

GROUP ACTIONS OF A_5 ON CONTRACTIBLE 2-COMPLEXES

IVÁN SADOFSCHI COSTA

ABSTRACT. We prove that every action of A_5 on a finite 2-dimensional contractible complex has a fixed point.

CONTENTS

1. Introduction	1
2. Fixed point free actions on acyclic 2-complexes	3
3. A reduction	6
4. Brown's short exact sequence	9
5. A moduli of representations	12
6. Quaternions	15
7. Group actions of A_5 on contractible 2-complexes	19
Appendix A. Equations over groups	20
Appendix B. Quaternion valued analytic functions	21
Appendix C. Alternative proof of Lemma 5.1	22
Appendix D. Alternative proofs using SAGE	23
References	25

1. INTRODUCTION

A well-known result of Jean-Pierre Serre [Ser80] states that every action of a finite group on a contractible 1-complex (i.e. a tree) has a fixed point. By Smith theory, every action of a p -group on the disk \mathbb{D}^n has a fixed point. The group A_5 acts simplicially and fixed point freely on the barycentric subdivision X of the 2-skeleton of the Poincaré homology sphere which is an acyclic 2-complex. By considering the join $X * A_5$, Edwin E. Floyd and Roger W. Richardson [FR59] proved that A_5 acts simplicially and fixed point freely on a contractible 3-complex. Moreover, by embedding $X * A_5$ in \mathbb{R}^{81} and taking a regular neighbourhood they proved that A_5 acts simplicially and fixed point freely on a triangulation of the disk \mathbb{D}^{81} . This was the only example known of this kind until Bob Oliver obtained a complete classification of the groups that act fixed point freely on a disk \mathbb{D}^n [Oli75]. The example of Floyd and Richardson makes clear that it is not possible to extend Serre's result to 3-complexes and is natural to wonder if it holds for 2-complexes. Carles Casacuberta and Warren Dicks [CD92] made the following conjecture (without requiring X to be finite) which was also posed by Michael Aschbacher and Yoav Segev as a question [AS93, Question 3] in the finite case.

2010 *Mathematics Subject Classification.* 57S17, 57M20, 57M60, 55M20, 55M25, 20F05, 20E06.

Key words and phrases. Group actions, contractible 2-complexes, moduli of group representations, mapping degree.

Researcher of CONICET. The author was partially supported by grants PICT-2017-2806, PIP 11220170100357CO and UBACyT 20020160100081BA.

Conjecture 1.1. *Let G be a finite group. If X is a 2-dimensional finite contractible G -complex then $X^G \neq \emptyset$.*

In [CD92] the conjecture is proved for solvable groups. Independently, Segev [Seg93] studied the question of which groups act without fixed points on an acyclic 2-complex and proved Conjecture 1.1 for solvable groups and the alternating groups A_n for $n \geq 6$. In [Seg94], Segev proved the conjecture for collapsible 2-complexes. Using the classification of the finite simple groups, Aschbacher and Segev proved that for many groups any action on a finite 2-dimensional acyclic complex has a fixed point [AS93]. Later, Oliver and Segev [OS02] gave a complete classification of the groups that act without fixed points on a finite acyclic 2-complex. Before [OS02], A_5 was the only group known to act without fixed points on an acyclic 2-complex. An excellent exposition on this topic is the one given by Alejandro Adem at the Séminaire Bourbaki [Ade03]. In [Cor01], J.M. Corson proved that Conjecture 1.1 holds for diagrammatically reducible complexes. The smallest group for which Conjecture 1.1 remained open is the alternating group A_5 . The main result of this paper is the following.

Theorem 7.2. *Every action of $A_5 \cong \mathrm{PSL}_2(2^2)$ on a finite, contractible 2-complex has a fixed point.*

From this, using the results of Oliver and Segev [OS02], we deduce the following.

Corollary 7.3. *Let G be one of the groups $\mathrm{PSL}_2(2^{2k})$, $\mathrm{PSL}_2(5^k)$ for $k \geq 1$ or $\mathrm{PSL}_2(q)$ for $q \equiv \pm 3 \pmod{8}$ and $q \equiv \pm 1 \pmod{5}$. Then every action of G on a finite contractible 2-complex has a fixed point.*

Our proof of Theorem 7.2 goes by constructing a nontrivial representation in $\mathrm{SO}(3, \mathbb{R})$ of the fundamental group of every fixed point free, 2-dimensional, finite acyclic A_5 -complex. Therefore we have the following.

Corollary 7.5. *Let X be a fixed point free 2-dimensional finite and acyclic A_5 -complex and let $\pi = \pi_1(X)$. Then π is infinite or there is an epimorphism $\pi \rightarrow A_5$.*

The paper is organized as follows. In Section 3 we prove Theorem 3.6 which says that to prove Theorem 7.2 it is enough to inspect the acyclic complexes of the type considered by Oliver and Segev in [OS02]. The necessary results from [OS02] are recalled in Section 2. In Section 3 we also prove Theorem 3.8, which describes a possible path towards settling Conjecture 1.1.

In Section 4 we establish the connection between Theorem 7.2 and the following group theoretic statement, using a result of Kenneth S. Brown [Bro84] in Bass–Serre theory.

Theorem 7.1. *There is no presentation of A_5 of the form*

$$\langle a, b, c, d, x_0, \dots, x_k \mid a^2, b^3, c^2, d^2, (ab)^3, (bc)^2, (cd)^5, x_0 a x_0^{-1} = d, w_0, \dots, w_k \rangle$$

with $w_0, \dots, w_k \in \ker(\phi)$, where $\phi: F(a, b, c, d, x_0, \dots, x_k) \rightarrow A_5$ is given by $a \mapsto (2, 5)(3, 4)$, $b \mapsto (3, 5, 4)$, $c \mapsto (1, 2)(3, 5)$, $d \mapsto (2, 5)(3, 4)$ and $x_i \mapsto 1$ for each $i = 0, \dots, k$.

In order to prove Theorem 7.1, in Section 5 we introduce a moduli of representations of the group

$$\Gamma_k = \langle a, b, c, d, x_0, \dots, x_k \mid a^2, b^3, c^2, d^2, (ab)^3, (bc)^2, (cd)^5, x_0 a x_0^{-1} = d \rangle$$

in $\mathrm{SO}(3)$. In Section 6 we view these rotations in $S^3 \subset \mathbb{H}$, enabling us to do a degree argument which completes the proof of Theorem 7.1. This proof is inspired by James Howie’s proof of the

Scott–Wiegold conjecture [How02]. Finally, in Section 7 we put everything together to complete the proof of Theorem 7.2.

Note. Some of the results presented here appeared originally in the author’s thesis [SC19].

Acknowledgements. I am grateful to my mentor, Jonathan Barmak, for his constant advice, in particular for suggesting me to work on this problem during my PhD. I would like to thank Bob Oliver for his feedback on my thesis. I would also like to thank Kevin Piterman for taking a look at a previous version of this article, and Yago Antolin, Ignacio Darago, Carlos Di Fiore, Fernando Martin and Pedro Tamaroff for valuable conversations.

2. FIXED POINT FREE ACTIONS ON ACYCLIC 2-COMPLEXES

In this section we review the results obtained by Bob Oliver and Yoav Segev in their article [OS02] that are needed later.

Throughout the paper, by G -complex we mean a G -CW complex. That is, a CW complex with a continuous G -action that is *admissible* (i.e. the action permutes the open cells of X , and maps a cell to itself only via the identity). For more details see [OS02, Appendix A]. We will frequently assume that the 2-cells in a G -complex are attached along closed edge paths, this will make no difference for the questions that we study. A *graph* is a 1-dimensional CW complex. By G -graph we always mean a 1-dimensional G -complex.

Definition 2.1 ([OS02]). A G -complex X is *essential* if there is no normal subgroup $1 \neq N \triangleleft G$ such that for each $H \subseteq G$, the inclusion $X^{HN} \rightarrow X^H$ induces an isomorphism on integral homology.

The main results of [OS02] are the following two theorems.

Theorem 2.2 ([OS02, Theorem A]). *For any finite group G , there is an essential fixed point free 2-dimensional (finite) acyclic G -complex if and only if G is isomorphic to one of the simple groups $\mathrm{PSL}_2(2^k)$ for $k \geq 2$, $\mathrm{PSL}_2(q)$ for $q \equiv \pm 3 \pmod{8}$ and $q \geq 5$, or $\mathrm{Sz}(2^k)$ for odd $k \geq 3$. Furthermore, the isotropy subgroups of any such G -complex are all solvable.*

Theorem 2.3 ([OS02, Theorem B]). *Let G be any finite group, and let X be any 2-dimensional acyclic G -complex. Let N be the subgroup generated by all normal subgroups $N' \triangleleft G$ such that $X^{N'} \neq \emptyset$. Then X^N is acyclic; X is essential if and only if $N = 1$; and the action of G/N on X^N is essential.*

The following fundamental result of Segev [Seg93, Theorem 3.4] will be used frequently, sometimes implicitly. We state the version given in [OS02].

Theorem 2.4 ([OS02, Theorem 4.1]). *Let X be any 2-dimensional acyclic G -complex (not necessarily finite). Then X^G is acyclic or empty, and is acyclic if G is solvable.*

We denote the set of subgroups of G by $\mathcal{S}(G)$.

Definition 2.5 ([OS02]). By a *family* of subgroups of G we mean any subset $\mathcal{F} \subseteq \mathcal{S}(G)$ which is closed under conjugation. A nonempty family is said to be *separating* if it has the following three properties: (a) $G \notin \mathcal{F}$; (b) if $H' \subseteq H$ and $H \in \mathcal{F}$ then $H' \in \mathcal{F}$; (c) for any $H \triangleleft K \subseteq G$ with K/H solvable, $K \in \mathcal{F}$ if $H \in \mathcal{F}$.

For any family \mathcal{F} of subgroups of G , a (G, \mathcal{F}) -complex is a G -complex all of whose isotropy subgroups lie in \mathcal{F} . A (G, \mathcal{F}) -complex is *universal* (resp. *H-universal*) if the fixed point set of each $H \in \mathcal{F}$ is contractible (resp. acyclic).

If G is not solvable, the separating family of solvable subgroups of G is denoted by \mathcal{SLV} . If G is perfect, then the family of proper subgroups of G is denoted by $\mathcal{MA}\mathcal{X}$.

Lemma 2.6 ([OS02, Lemma 1.2]). *Let X be any 2-dimensional acyclic G -complex without fixed points. Let \mathcal{F} be the set of subgroups $H \subseteq G$ such that $X^H \neq \emptyset$. Then \mathcal{F} is a separating family of subgroups of G , and X is an H -universal (G, \mathcal{F}) -complex.*

Proposition 2.7 ([OS02, Proposition 6.4]). *Assume that L is one of the simple groups $\mathrm{PSL}_2(q)$ or $\mathrm{Sz}(q)$, where $q = p^k$ and p is prime ($p = 2$ in the second case). Let $G \subseteq \mathrm{Aut}(L)$ be any subgroup containing L , and let \mathcal{F} be a separating family for G . Then there is a 2-dimensional acyclic (G, \mathcal{F}) -complex if and only if $G = L$, $\mathcal{F} = \mathcal{SLV}$, and q is a power of 2 or $q \equiv \pm 3 \pmod{8}$.*

Lemma 2.8. *Let G be one of the groups in Theorem 2.2 and let X be a fixed point free 2-dimensional acyclic G -complex. If $K \leq G$ is not solvable then the action of K on X is fixed point free.*

Proof. Let $\mathcal{F} = \{H \leq G : X^H \neq \emptyset\}$. Then, by Lemma 2.6, \mathcal{F} is a separating family and X is an H -universal (G, \mathcal{F}) -complex. By Proposition 2.7, we must have $\mathcal{F} = \mathcal{SLV}$ and therefore $X^K = \emptyset$. \square

If X is a poset, then $\mathcal{K}(X)$ denotes the *order complex* of X , that is, the simplicial complex with simplices the finite nonempty totally ordered subsets of X (the complex $\mathcal{K}(X)$ is also known as the *nerve* of X).

Definition 2.9 ([OS02, Definition 2.1]). For any family \mathcal{F} of subgroups of G define

$$i_{\mathcal{F}}(H) = \frac{1}{[N_G(H) : H]}(1 - \chi(\mathcal{K}(\mathcal{F}_{>H}))).$$

Recall that if $G \curvearrowright X$, the orbit $G \cdot x$ is said to be of *type* G/H if the stabilizer G_x is conjugate to H in G . In other words, if the action of G on $G \cdot x$ is the same as the action of G on G/H .

Lemma 2.10 ([OS02, Lemma 2.3]). *Fix a separating family \mathcal{F} , a finite H -universal (G, \mathcal{F}) -complex X , and a subgroup $H \subseteq G$. For each n , let $c_n(H)$ denote the number of orbits of n -cells of type G/H in X . Then $i_{\mathcal{F}}(H) = \sum_{n \geq 0} (-1)^n c_n(H)$.*

Proposition 2.11 ([OS02, Tables 2,3,4]). *Let G be one of the simple groups $\mathrm{PSL}_2(2^k)$ for $k \geq 2$, $\mathrm{PSL}_2(q)$ for $q \equiv \pm 3 \pmod{8}$ and $q \geq 5$, or $\mathrm{Sz}(2^k)$ for odd $k \geq 3$. Then $i_{\mathcal{SLV}}(1) = 1$.*

For each family of groups appearing in Theorem 2.2, Oliver and Segev describe an example. In what follows, D_{2m} is a dihedral group of order $2m$ and C_m is a cyclic group of order m .

Proposition 2.12 ([OS02, Example 3.4]). *Set $G = \mathrm{PSL}_2(q)$, where $q = 2^k$ and $k \geq 2$. Then there is a 2-dimensional acyclic fixed point free G -complex X , all of whose isotropy subgroups are solvable. More precisely X can be constructed to have three orbits of vertices with isotropy subgroups isomorphic to $B = \mathbb{F}_q \rtimes C_{q-1}$, $D_{2(q-1)}$, and $D_{2(q+1)}$; three orbits of edges with isotropy subgroups isomorphic to C_{q-1} , C_2 and C_2 ; and one free orbit of 2-cells.*

We have $A_5 = \mathrm{PSL}_2(2^2)$. The barycentric subdivision of the 2-skeleton of the Poincaré dodecahedral space is an A_5 -complex of the type given in Proposition 2.12 with fundamental group the binary icosahedral group $A_5^* \cong \mathrm{SL}(2, 5)$ which has order 120. The Poincaré dodecahedral space appears in many other natural ways, for more information see [KS79].

Proposition 2.13 ([OS02, Example 3.5]). *Assume that $G = \mathrm{PSL}_2(q)$, where $q = p^k \geq 5$ and $q \equiv \pm 3 \pmod{8}$. Then there is a 2-dimensional acyclic fixed point free G -complex X , all of whose isotropy subgroups are solvable. More precisely, X can be constructed to have four orbits of vertices with isotropy subgroups isomorphic to $B = \mathbb{F}_q \rtimes C_{(q-1)/2}$, D_{q-1} , D_{q+1} , and A_4 ; four orbits of edges with isotropy subgroups isomorphic to $C_{(q-1)/2}$, C_2^2 , C_3 and C_2 ; and one free orbit of 2-cells.*

Proposition 2.14 ([OS02, Example 3.7]). *Set $q = 2^{2k+1}$ for any $k \geq 1$. Then there is a 2-dimensional acyclic fixed point free $\mathrm{Sz}(q)$ -complex X , all of whose isotropy subgroups are solvable. More precisely, X can be constructed to have four orbits of vertices with isotropy subgroups isomorphic to $M(q, \theta)$, $D_{2(q-1)}$, $C_{q+\sqrt{2q}+1} \rtimes C_4$, $C_{q-\sqrt{2q}+1} \rtimes C_4$; four orbits of edges with isotropy subgroups isomorphic to C_{q-1} , C_4 , C_4 and C_2 ; and one free orbit of 2-cells.*

We also have $A_5 \cong \mathrm{PSL}_2(5)$, so this group is addressed in both Proposition 2.12 and Proposition 2.13. There is no other such exception.

Definition 2.15. If G is one of the groups in Theorem 2.2, the *Oliver–Segev G -graph* $\Gamma_{OS}(G)$ is the 1-skeleton of any 2-dimensional fixed point free acyclic G -complex without free orbits of 1-cells of the type constructed in Propositions 2.12 to 2.14. For this definition, we regard A_5 as $\mathrm{PSL}_2(2^2)$ rather than $\mathrm{PSL}_2(5)$.

Generally, there is more than one possible choice for the G -graph $\Gamma_{OS}(G)$. Even for $G = A_5$, thought of as $\mathrm{PSL}_2(2^2)$, the quotient graph $\Gamma_{OS}(G)/G$ is not unique. However in Proposition 3.10 we show that $\Gamma_{OS}(G)$ is unique up to G -homotopy equivalence. Moreover, Corollary 3.11 shows the particular choice of $\Gamma_{OS}(G)$ is irrelevant for our purposes.

Definition 2.16 (A construction of $\Gamma_{OS}(A_5)$). Here we give a construction of $\Gamma_{OS}(A_5)$ and we fix some notation in regard to this graph. Consider the following subgroups of A_5 :

$$\begin{aligned} H_1 &= \langle (2, 5)(3, 4), (3, 5, 4) \rangle \cong A_4, \\ H_2 &= \langle (3, 5, 4), (1, 2)(3, 5) \rangle \cong D_6 \text{ and} \\ H_3 &= \langle (1, 2)(3, 5), (2, 5)(3, 4) \rangle \cong D_{10}. \end{aligned}$$

The graph $\Gamma_{OS}(A_5)$ has three orbits of vertices whose representatives v_1, v_2, v_3 have stabilizers H_1, H_2, H_3 respectively. In addition, $\Gamma_{OS}(A_5)$ has three orbits of edges whose representatives $v_1 \xrightarrow{e_{12}} v_2$, $v_3 \xrightarrow{e_{31}} v_1$ and $v_2 \xrightarrow{e_{23}} v_3$ have stabilizers

$$\begin{aligned} H_{12} &= H_1 \cap H_2 = \langle (3, 5, 4) \rangle \cong \mathbb{Z}_3, \\ H_{13} &= H_1 \cap H_3 = \langle (2, 5)(3, 4) \rangle \cong \mathbb{Z}_2 \text{ and} \\ H_{23} &= H_2 \cap H_3 = \langle (1, 2)(3, 5) \rangle \cong \mathbb{Z}_2 \end{aligned}$$

respectively.

Attaching a free orbit of 2-cells to $\Gamma_{OS}(A_5)$ along the orbit of the closed edge path (e_{12}, e_{23}, e_{31}) we obtain an acyclic 2-dimensional fixed point free A_5 -complex of the type given in Proposition 2.12. This complex is, in fact, the barycentric subdivision of the 2-skeleton of the Poincaré dodecahedral space (a simplicial complex having $21 = 5 + 10 + 6$ vertices, $80 = 20 + 30 + 30$ edges and 60 faces). A concrete isomorphism can be produced by mapping v_3 to the barycentre of a pentagonal 2-cell $ABCDE$, v_1 to A and v_2 to the barycentre (midpoint) of AB . For more details on this see [OS02, pp. 20-21].

3. A REDUCTION

In this section we rely on the results of Oliver and Segev to prove Theorem 3.6, which allows us to reduce the proof of Theorem 7.2 to the study of acyclic complexes of the type considered in [OS02]. We also prove Theorem 3.8 which describes a possible path to establish Conjecture 1.1. We first prove some results which will be used to do equivariant modifications to our complexes.

Definition 3.1. If X, Y are G -spaces, a G -homotopy is an equivariant map $H: X \times I \rightarrow Y$. We say that $f_0(x) = H(x, 0)$ and $f_1(x) = H(x, 1)$ are G -homotopic and we denote this by $f_0 \simeq_G f_1$. An equivariant map $f: X \rightarrow Y$ is a G -homotopy equivalence if there is an equivariant map $g: Y \rightarrow X$ such that $fg \simeq_G 1_Y$ and $gf \simeq_G 1_X$. A G -invariant subspace A of X is a *strong G -deformation retract* of X if there is a retraction $r: X \rightarrow A$ such that there is a G -homotopy $H: ir \simeq 1_X$ relative to A , where $i: A \rightarrow X$ is the inclusion.

Remark 3.2. An equivariant map $f: X \rightarrow Y$ is a G -homotopy equivalence if and only if $f^H: X^H \rightarrow Y^H$ is a homotopy equivalence for each subgroup $H \leq G$ (see [tD87, Chapter II, (2.7) Proposition]). Thus, if $f: X \rightarrow Y$ is a G -homotopy equivalence, the action $G \curvearrowright X$ is fixed point free (resp. essential) if and only if the action $G \curvearrowright Y$ is fixed point free (resp. essential).

The following lemma allows us to do elementary expansions equivariantly.

Lemma 3.3. *Let X be an acyclic 2-dimensional G -complex. Let $H \leq G$ and $x_0, x_1 \in X^{(0)} \cap X^H$. Then there is a G -complex $Y \supset X$, such that X is a strong G -deformation retract of Y and Y is obtained from X by attaching an orbit of 1-cells of type G/H with endpoints $\{x_0, x_1\}$ and an orbit of 2-cells of type G/H .*

Proof. We attach an orbit of 1-cells of type G/H to X using the attaching map $\varphi: G/H \times S^0 \rightarrow X^{(0)}$ defined by $(gH, 1) \mapsto g \cdot x_0$, $(gH, -1) \mapsto g \cdot x_1$. Let e be the 1-cell of this new orbit corresponding to the coset H . Since X is acyclic, by Theorem 2.4 X^H is also acyclic. Let γ be an edge path in X^H starting at x_1 and ending at x_0 . Then we attach an orbit of 2-cells of type G/H in such a way that the 2-cell corresponding to the coset H is attached along the closed edge path given by e and γ . It is clear that X is a strong G -deformation retract of Y . \square

The following very natural definitions appear in [KLV01, Section 2].

Definition 3.4. A *forest* is a graph with trivial first homology. If a subcomplex Γ of a CW complex X is a forest, there is a CW complex Y obtained from X by shrinking each connected component of Γ to a point. The quotient map $q: X \rightarrow Y$ is a homotopy equivalence and we say Y is obtained from X by a *forest collapse*.

If X is a G -complex and $\Gamma \subset X$ is a forest which is G -invariant, the quotient map q is a G -homotopy equivalence and we say the G -complex Y is obtained from X by a *G -forest collapse*. We say that a G -graph is *reduced* if it has no edge e such that $G \cdot e$ is a forest.

Lemma 3.5. *Let X be a 2-dimensional acyclic G -complex. If $X^{(1)}$ is a reduced G -graph then stabilizers of different vertices are not comparable.*

Proof. Let $\mathcal{F} = \{G_x : x \in X^{(0)}\}$ and let $M = \{v \in X^{(0)} : G_v \text{ is maximal in } \mathcal{F}\}$. We first prove, by contradiction, that $X^{(0)} = M$. Consider $v \in X^{(0)} - M$ such that G_v is maximal in $\{G_x : x \in X^{(0)} - M\}$. Then since X^{G_v} contains v , by Theorem 2.4 it must be acyclic. Since $v \notin M$, there is a vertex $w \in X^{G_v} \cap M$. By connectivity there is an edge $e \in X^{G_v}$ whose

endpoints v' and w' satisfy $v' \notin M$ and $w' \in M$. Since $G_{v'} \geq G_v$ and $v' \notin M$, by our choice of v we have $G_v = G_{v'}$. Since $e \in X^{G_v}$ we have $G_v \leq G_e$ and since v' is an endpoint of e we have $G_e \leq G_{v'}$. Thus $G_e = G_{v'}$ and then the degree of v' in the graph $G \cdot e$ (which has vertex set $G \cdot w' \amalg G \cdot v'$) is 1. Thus $G \cdot e$ is a forest, contradiction. Therefore we must have $M = X^{(0)}$. To conclude we have to prove that different vertices $u, v \in M$ have different stabilizers. Suppose $G_u = G_v$ to get a contradiction. Since u, v are vertices of X^{G_u} which is connected, there is an edge $e \in X^{G_u}$ and by maximality we must have $G_e = G_u$. If u', v' are the endpoints of e , we have $G_{u'} = G_{v'}$. We have two cases and in any case we obtain a contradiction. If $G \cdot u' \neq G \cdot v'$ then $G \cdot e$ is a forest consisting of $|G/G_e|$ disjoint edges, contradiction. Otherwise, there is a nontrivial element $g \in G$ such that $g \cdot u' = v'$ and we have $G_{u'} = G_{v'} = gG_{u'}g^{-1}$. Thus $g \in N_G(G_{u'})$. Consider the action of $\langle g \rangle$ on $X^{G_{u'}}$, which is acyclic and thus has a fixed point by the Lefschetz fixed point theorem. But this cannot happen, since this would imply that $\langle G_{u'}, g \rangle \gneq G_{u'}$ fixes a point of X , which is a contradiction since $u' \in M$. \square

Now we prove the main results of the section.

Theorem 3.6. *Let G be one of the groups in Theorem 2.2. Let X be a fixed point free 2-dimensional finite acyclic G -complex. Then there is a fixed point free 2-dimensional finite acyclic G -complex X' obtained from the G -graph $\Gamma_{OS}(G)$ by attaching $k \geq 0$ free orbits of 1-cells and $k + 1$ free orbits of 2-cells and an epimorphism $\pi_1(X) \rightarrow \pi_1(X')$.*

Proof. Let $\mathcal{F} = \{H \leq G : X^H \neq \emptyset\}$. Then, by Lemma 2.6, \mathcal{F} is a separating family and X is an H -universal (G, \mathcal{F}) -complex. By Proposition 2.7, we must have $\mathcal{F} = \mathcal{SLV}$. By doing enough G -forest collapses we can assume that $X^{(1)}$ is a reduced G -graph. The stabilizers of the vertices of $\Gamma_{OS}(G)$ are precisely the maximal solvable subgroups of G . Therefore, since every solvable subgroup of G fixes a point of X , by Lemma 3.5, we may identify $X^{(0)} = \Gamma_{OS}(G)^{(0)}$. Applying Lemma 3.3 enough times to modify X , we may further assume $\Gamma_{OS}(G)$ is a subcomplex of X .

Finally we will modify X so that for every subgroup $1 \neq H \leq G$, we have $X^H = \Gamma_{OS}(G)^H$. We do this by reverse induction on $|H|$. Assume that we have X such that it holds for every subgroup K with $H \lneq K \leq G$. If H is not solvable, we have $X^H = \Gamma_{OS}(G)^H = \emptyset$ so we are done. If H is solvable, since $\Gamma_{OS}(G)^H$ is a tree (it is acyclic and 1-dimensional) and X^H is acyclic by Theorem 2.4, the inclusion $\Gamma_{OS}(G)^H \hookrightarrow X^H$ is an $N_G(H)$ -equivariant homology equivalence. Now since $\Gamma_{OS}(G)^H$ is a tree we can define an $N_G(H)$ -equivariant retraction $r_H: X^H \rightarrow \Gamma_{OS}(G)^H$. Then r_H is a homology equivalence. Moreover, the stabilizer of the cells in $X^H - \Gamma_{OS}(G)^H$ is H (the stabilizer cannot be bigger by the induction hypothesis). We define retractions $r_{Hg}: X^{Hg} \rightarrow \Gamma_{OS}(G)^{Hg}$ by $r_{Hg}(gx) = g \cdot r_H(x)$ which glue to give a G -equivariant homology equivalence

$$r: \Gamma_{OS}(G) \bigcup_{g \in G} X^{Hg} \rightarrow \Gamma_{OS}(G).$$

We may replace X by the pushout \tilde{X} given by the following diagram

$$\begin{array}{ccc} \Gamma_{OS}(G) \bigcup_{g \in G} X^{Hg} & \xrightarrow{r} & \Gamma_{OS}(G) \\ \downarrow & & \downarrow \\ X & \xrightarrow{\bar{r}} & \tilde{X} \end{array}$$

It follows that \bar{r} is a homology equivalence, so the resulting G -complex \tilde{X} is acyclic. Moreover since $\tilde{X}^{(1)}$ is a subcomplex of $X^{(1)}$ and the restriction $\bar{r}: X^{(1)} \rightarrow \tilde{X}^{(1)}$ is a retraction, \bar{r} induces an epimorphism on π_1 . This procedure removes the excessive orbits of cells of type G/H . By induction we obtain a complex X' such that $X'^{(1)}$ coincides with $\Gamma_{OS}(G)$ up to $k \geq 0$ free orbits of 1-cells and such that every orbit of 2-cells is free. By Lemma 2.6 X' is an H -universal (G, \mathcal{SLV}) -complex. Now by Lemma 2.10 and Proposition 2.11 there are exactly $k + 1$ orbits of 2-cells. \square

From [OS02, Propositions 3.3 and 3.6] we have:

Proposition 3.7. *Each of the groups in the statement of Theorem 2.2 has a subgroup isomorphic to one of the following groups:*

- $\mathrm{PSL}_2(2^p)$ for p prime;
- $\mathrm{PSL}_2(3^p)$ for an odd prime p ;
- $\mathrm{PSL}_2(q)$ for a prime $q > 3$ such that $q \equiv \pm 3 \pmod{5}$ and $q \equiv \pm 3 \pmod{8}$;
- $\mathrm{Sz}(2^p)$ for p an odd prime.

Moreover, every proper subgroup of a group in this list is solvable.

Theorem 3.8. *To prove Conjecture 1.1 it is enough to prove $P(G)$ for each group G listed in Proposition 3.7, where $P(G)$ denotes the following proposition: “there is a nontrivial representation in $\mathrm{SO}(n, \mathbb{R})$ of the fundamental group of every acyclic G -complex obtained from $\Gamma_{OS}(G)$ by attaching $k \geq 0$ free orbits of 1-cells and $(k + 1)$ free orbits of 2-cells”.*

Proof. Let G be a finite group and suppose that X is a finite, acyclic 2-dimensional fixed point free G -complex. Let N be the subgroup generated by all normal subgroups $N' \triangleleft G$ such that $X^{N'} \neq \emptyset$. By Theorem 2.3 we have that $Y = X^N$ is acyclic and the action of G/N on Y is essential and fixed point free. Then G/N must be one of the groups in Theorem 2.2. We take a subgroup K of G/N isomorphic to one of the groups listed in Proposition 3.7. Then by Lemma 2.8 the action of K on Y is fixed point free. Now, by Theorem 3.6 and by $P(K)$, it follows that $\pi_1(Y)$ admits a nontrivial representation in $\mathrm{SO}(n, \mathbb{R})$. Therefore, by Proposition A.3, X cannot be contractible. \square

Remark 3.9. In Theorem 7.4 we prove the group $A_5 \cong \mathrm{PSL}_2(2^2)$ satisfies the condition P in Theorem 3.8.

The following explains why our particular choice of $\Gamma_{OS}(G)$ and the way the free orbits of 1-cells are attached is not relevant.

Proposition 3.10. *Any two choices for $\Gamma_{OS}(G)$ are G -homotopy equivalent. Moreover, attaching $k \geq 0$ free orbits of 1-cells to any two choices for $\Gamma_{OS}(G)$ produces G -homotopy equivalent graphs.*

Proof. Since any choice of $\Gamma_{OS}(G)$ is a universal $(G, \mathcal{SLV} - \{1\})$ -complex, the first part follows from [OS02, Proposition A.6]. The second part follows easily from the first and the gluing theorem for adjunction spaces [Bro06, 7.5.7]. \square

Corollary 3.11. *Let Γ be a graph obtained from $\Gamma_{OS}(G)$ by attaching $k \geq 0$ free orbits of 1-cells. The set of G -homotopy equivalence classes of 2-dimensional acyclic fixed point free G -complexes with 1-skeleton Γ does not depend on the particular choice of $\Gamma_{OS}(G)$ or the way the k free orbits of 1-cells are attached. In particular, the set of isomorphism classes of groups that occur as the fundamental group of such spaces does not depend on such choices.*

Proof. Again, this is an easy application of [Bro06, 7.5.7]. \square

4. BROWN'S SHORT EXACT SEQUENCE

Using Bass–Serre theory, K.S. Brown gave a method to produce a presentation for a group G acting on a simply connected complex X [Bro84, Theorem 1]. When X is not simply connected, Brown describes a presentation for an extension \tilde{G}_X of G by $\pi_1(X)$ [Bro84, Theorem 2]. The group \tilde{G}_X has a description as a quotient of the fundamental group of a graph of groups. A similar result in the simply connected case was given by Corson [Cor92, Theorem 5.1] in terms of *complexes of groups* (higher dimensional analogues of graphs of groups).

Using Brown's result we translate the A_5 case of Conjecture 1.1 into a nice looking problem in combinatorial group theory. This translation can be done in general, but to obtain similar results for the rest of the groups G that appear in Theorem 3.8 we need a choice of $\Gamma_{OS}(G)$ and presentations for the stabilizers of its vertices.

In Brown's original formulation, the result deals with actions that need not to be admissible (Brown uses the term G – CW -complex in a different way than us). Since the actions we are interested in are admissible, we state Brown's result only in that case.

Let X be a connected G -complex. By admissibility of the action, the group G acts on the set of oriented edges. If e is an oriented edge, the same 1-cell with the opposite orientation is denoted by e^{-1} . Each oriented edge e has a *source* and *target*, denoted by $s(e)$ and $t(e)$ and for every $g \in G$ we have $g \cdot s(e) = s(g \cdot e)$ and $g \cdot t(e) = t(g \cdot e)$.

To obtain a description of the group \tilde{G}_X we need a number of choices that we now specify. For each 1-cell of X we choose a preferred orientation in such a way that these orientations are preserved by G . This determines a set P of oriented edges. We choose a *tree of representatives* for X/G . That is, a tree $T \subset X$ such that the vertex set V of T is a set of representatives of $X^{(0)}/G$. Such tree always exists and the 1-cells of T are inequivalent modulo G . We give an orientation to the 1-cells of T so that they are elements of P . We also choose a set of representatives E of P/G in such a way that $s(e) \in V$ for every $e \in E$ and such that each oriented edge of T is in E . If e is an oriented edge, the unique element of V that is equivalent to $t(e)$ modulo G will be denoted by $w(e)$. For every $e \in E$ we fix an element $g_e \in G$ such that $t(e) = g_e \cdot w(e)$. If $e \in T$, we specifically choose $g_e = 1$. For each orbit of 2-cells we choose a closed edge path τ based at a vertex of T and representing the attaching map for this orbit of 2-cells. Let F be the set given by these closed edge paths.

The group \tilde{G}_X is defined as a quotient of

$$\bigast_{v \in V} G_v \bigast_{e \in E} \mathbb{Z}$$

by certain relations. In order to define these relations we introduce some notation. If $v \in V$ and $g \in G_v$ we denote the copy of g in the free factor G_v by g_v . The generator of the copy of \mathbb{Z} that corresponds to e is denoted by x_e . The relations are the following:

- (i) $x_e = 1$ if $e \in T$.
- (ii) $x_e^{-1} g_{s(e)} x_e = (g_e^{-1} g g_e)_{w(e)}$ for every $e \in E$ and $g \in G_e$.
- (iii) $r_\tau = 1$ for every $\tau \in F$.

We state Brown's theorem before giving the definition of the element r_ω associated to a closed edge path ω .

Theorem 4.1 (Brown, [Bro84, Theorems 1 and 2]). *The group*

$$\tilde{G}_X = \frac{\bigstar_{v \in V} G_v \bigstar_{e \in E} \mathbb{Z}}{\langle\langle R \rangle\rangle}$$

where R consists of relations (i)-(iii) is an extension

$$1 \rightarrow \pi_1(X, x_0) \xrightarrow{i} \tilde{G}_X \xrightarrow{\bar{\phi}} G \rightarrow 1.$$

The map $\bar{\phi}$ is defined passing to the quotient the coproduct ϕ of the inclusions $G_v \rightarrow G$ and the mappings $\mathbb{Z} \rightarrow G$ given by $x_e \mapsto g_e$. The map i sends a closed edge path ω based at $x_0 \in V$ to the class of r_ω .

Now we explain how to obtain the elements r_ω . If α is an oriented edge, we define

$$\varepsilon(\alpha) = \begin{cases} 1 & \alpha \in P \\ -1 & \text{if } \alpha \notin P \end{cases}$$

and we can always take $e \in E$ and $g \in G$ such that $\alpha = ge^{\varepsilon(\alpha)}$. Note that e is unique but g is not. Moreover, if α starts at $v \in V$, we can write

$$\alpha = \begin{cases} he & \text{with } h \in G_{s(e)}, \text{ if } \alpha \in P \\ hg_e^{-1}e^{-1} & \text{with } h \in G_{w(e)}, \text{ if } \alpha \notin P \end{cases}$$

Again, h is not unique.

Now if $\omega = (\alpha_1, \dots, \alpha_n)$ is a closed edge path starting at a vertex $v_0 \in V$ we define the element $r_\omega \in \bigstar_{v \in V} G_v \bigstar_{e \in E} \mathbb{Z}$. Recursively, we define some sequences. Since the oriented edge α_1 starts at $v_0 \in V$, we can obtain an oriented edge e_1 and an element $h_1 \in G_{v_0}$ as above. We set $\varepsilon_1 = \varepsilon(\alpha_1)$ and $g_1 = h_1 g_{e_1}^{\varepsilon_1}$. Set $v_1 = w(e_1)$ if $\alpha_1 \in P$ and otherwise $v_1 = s(e_1)$. Now suppose we have defined e_1, \dots, e_k , h_1, \dots, h_k , $\varepsilon_1, \dots, \varepsilon_k$, g_1, \dots, g_k and v_1, \dots, v_k such that the oriented edge $(g_1 g_2 \cdots g_k)^{-1} \alpha_{k+1}$ starts at $v_k \in V$. We can obtain an oriented edge e_{k+1} and an element $h_{k+1} \in G_{v_k}$ as before. We set $\varepsilon_{k+1} = \varepsilon(\alpha_{k+1})$ and $g_{k+1} = h_{k+1} g_{e_{k+1}}^{\varepsilon_{k+1}}$. Set $v_{k+1} = w(e_{k+1})$ if $\alpha_{k+1} \in P$ and otherwise $v_{k+1} = s(e_{k+1})$. When we conclude, we have an element $g_1 g_2 \cdots g_n \in G_{v_0}$. Finally the word associated to ω is given by

$$r_\omega = (h_1)_{v_0} x_{e_1}^{\varepsilon_1} (h_2)_{v_1} x_{e_2}^{\varepsilon_2} \cdots (h_n)_{v_{n-1}} x_{e_n}^{\varepsilon_n} (g_1 g_2 \cdots g_n)_{v_0}^{-1}.$$

A closed edge path ω in X determines a conjugacy class $[\omega]$ of $\pi_1(X)$. The following describes the conjugation action of \tilde{G}_X on $\pi_1(X)$.

Proposition 4.2 ([Bro84, Proposition 1]). *Let ω be a closed edge path in X and $g \in G$. Then the conjugacy classes $i([\omega])$ and $i([g\omega])$ are contained in the same \tilde{G}_X -conjugacy class. Moreover for any element $\tilde{g} \in \bar{\phi}^{-1}(g)$ we have $\tilde{g}i([\omega])\tilde{g}^{-1} = i([g\omega])$.*

The following proposition summarizes many ideas of this section.

Proposition 4.3. *Let Γ be a G -graph and let $w_1, \dots, w_k \in \ker(\bar{\phi}: \tilde{G}_\Gamma \rightarrow G)$. Let X be a G -complex obtained by attaching orbits of 2-cells to Γ along closed edge paths τ_1, \dots, τ_k such that $r_{\tau_i} = w_i$. Then we have a diagram with exact rows and columns*

$$\begin{array}{ccccccc}
& & 1 & & 1 & & 1 \\
& & \downarrow & & \downarrow & & \downarrow \\
1 & \longrightarrow & \langle\langle G \cdot \tau_i \rangle\rangle & \xrightarrow{\sim} & \langle\langle w_i \rangle\rangle^{\tilde{G}_\Gamma} & \longrightarrow & 1 \\
& & \downarrow & & \downarrow & & \downarrow \\
1 & \longrightarrow & \pi_1(\Gamma) & \xrightarrow{i} & \tilde{G}_\Gamma & \xrightarrow{\bar{\phi}} & G \longrightarrow 1 \\
& & \downarrow i_* & & \downarrow & & \parallel \\
1 & \longrightarrow & \pi_1(X) & \xrightarrow{i} & \tilde{G}_X & \xrightarrow{\bar{\phi}} & G \longrightarrow 1 \\
& & \downarrow & & \downarrow & & \downarrow \\
& & 1 & & 1 & & 1
\end{array}$$

and we have $H_1(X) \cong \frac{N}{\langle\langle w_i \rangle\rangle^{\tilde{G}_\Gamma}[N, N]}$, where $N = \ker(\bar{\phi}: \tilde{G}_\Gamma \rightarrow G)$.

Remark 4.4. If X is a connected G -complex, the group \tilde{G}_X is isomorphic to the group formed by the pairs (g, \tilde{g}) such that $g \in G$ and \tilde{g} is a lift of $g: X \rightarrow X$ to the universal cover \tilde{X} of X (see [Bro84]). Suppose Y is another G -complex and $h: X \rightarrow Y$ is equivariant and a homotopy equivalence. Let $\tilde{h}: \tilde{X} \rightarrow \tilde{Y}$ be a lift of h to the universal covers. Then if $g \in G$, for each lift $\tilde{g}_X: \tilde{X} \rightarrow \tilde{X}$ of $g: X \rightarrow X$ there is a unique lift $\tilde{g}_Y: \tilde{Y} \rightarrow \tilde{Y}$ of $g: Y \rightarrow Y$ such that the following diagram commutes:

$$\begin{array}{ccc}
\tilde{X} & \xrightarrow{\tilde{h}} & \tilde{Y} \\
\tilde{g}_X \downarrow & & \downarrow \tilde{g}_Y \\
\tilde{X} & \xrightarrow{\tilde{h}} & \tilde{Y}
\end{array}$$

Then it is easy to check that there is an isomorphism $\tilde{G}_X \rightarrow \tilde{G}_Y$ given by $\tilde{g}_X \mapsto \tilde{g}_Y$. In particular, the isomorphism type of $\tilde{G}_{\Gamma_{OS}(G)}$ does not depend on any choice.

We now apply Brown's result for $G = A_5$. Recall the construction of $\Gamma_{OS}(A_5)$ given in Definition 2.16. Suppose that we have an acyclic 2-complex X obtained from $\Gamma_{OS}(A_5)$ by attaching a free A_5 -orbit of 2-cells. We want to apply Brown's method to obtain a presentation for the extension \tilde{G}_X . We take $T = \{e_{12}, e_{23}\}$. Thus $V = \{v_1, v_2, v_3\}$. We take $E = \{e_{12}, e_{23}, e_{31}\}$. Note that we have $w(e) = t(e)$ for every $e \in E$. We can take $g_e = 1$ for every $e \in E$.

Then Brown's result gives

$$\tilde{G}_X = \frac{(H_1 *_{H_{12}} H_2 *_{H_{23}} H_3) *_{H_{13}}}{\langle\langle w \rangle\rangle}$$

We explain this. First we amalgamate the groups H_1, H_2, H_3 identifying the copy of H_{12} in H_1 with the copy of H_{12} in H_2 and the copy of H_{23} in H_2 with the copy of H_{23} in H_3 . This comes from the relations of type (i) and (ii) for $e \in T$. Then we form an HNN extension with stable letter $x = x_{e_{31}}$ that corresponds to the relation of type (ii) coming from e_{31} . The associated subgroups of this HNN extension are the copies of H_{13} in H_1 and H_3 . The quotient by the word w comes from the only relation of type (iii).

Now we obtain an explicit presentation for \tilde{G}_X . We have $\langle a, b \mid a^2, b^3, (ab)^3 \rangle \cong H_1$ via $a \mapsto (2, 5)(3, 4)$, $b \mapsto (3, 5, 4)$. We have $\langle b, c \mid b^3, c^2, (bc)^2 \rangle \cong H_2$ via $b \mapsto (3, 5, 4)$, $c \mapsto (1, 2)(3, 5)$.

Finally $\langle c, d \mid c^2, d^2, (cd)^5 \rangle \cong H_3$ via $c \mapsto (1, 2)(3, 5)$, $d \mapsto (2, 5)(3, 4)$. Thus we have a presentation

$$\tilde{G}_X = \langle a, b, c, d, x \mid a^2, b^3, c^2, d^2, (ab)^3, (bc)^2, (cd)^5, xax^{-1} = d, w \rangle$$

where the word w depends on the attaching map. The mapping $\bar{\phi}: \tilde{G}_X \rightarrow A_5$ is given by $a \mapsto (2, 5)(3, 4)$, $b \mapsto (3, 5, 4)$, $c \mapsto (1, 2)(3, 5)$, $d \mapsto (2, 5)(3, 4)$ and $x \mapsto 1$. Note that $\pi_1(X)$ is trivial if and only if $\bar{\phi}: \tilde{G}_X \rightarrow A_5$ is an isomorphism. If we also take into account k additional free orbits of 1 and 2 cells and we recall Theorem 3.6, from Brown's result we obtain:

Theorem 4.5. *The following are equivalent.*

- (i) *Every finite, 2-dimensional contractible A_5 -complex has a fixed point.*
- (ii) *There is no presentation of A_5 of the form*

$$\langle a, b, c, d, x_0, \dots, x_k \mid a^2, b^3, c^2, d^2, (ab)^3, (bc)^2, (cd)^5, x_0ax_0^{-1} = d, w_0, \dots, w_k \rangle$$

with $w_0, \dots, w_k \in \ker(\phi)$, where $\phi: F(a, b, c, d, x_0, \dots, x_k) \rightarrow A_5$ is given by $a \mapsto (2, 5)(3, 4)$, $b \mapsto (3, 5, 4)$, $c \mapsto (1, 2)(3, 5)$, $d \mapsto (2, 5)(3, 4)$ and $x_i \mapsto 1$.

5. A MODULI OF REPRESENTATIONS

In order to prove Theorem 7.1 we define a moduli of representations of the group

$$\Gamma_k = \langle a, b, c, d, x_0, \dots, x_k \mid a^2, b^3, c^2, d^2, (ab)^3, (bc)^2, (cd)^5, x_0ax_0^{-1} = d \rangle$$

in $\text{SO}(3)$. Our argument is inspired by James Howie's proof of the Scott–Wiegold conjecture [How02].

Let $\bar{\phi}: \Gamma_k \rightarrow A_5$ be the homomorphism induced by $\phi: F(a, b, c, d, x_0, \dots, x_k) \rightarrow A_5$.

Lemma 5.1. *We have $\ker(\bar{\phi}) = \langle\langle x_0, \dots, x_k, (bac)^3 \rangle\rangle$.*

Proof. It is straightforward to verify that $(bac)^3 \in \ker(\bar{\phi})$, so it is enough to show the induced epimorphism

$$\bar{\phi}: \Gamma_k / \langle\langle x_0, \dots, x_k, (bac)^3 \rangle\rangle \rightarrow A_5$$

is in fact an isomorphism. Eliminating d and the x_i we see

$$\Gamma_k / \langle\langle x_0, \dots, x_k, (bac)^3 \rangle\rangle = \langle a, b, c \mid a^2, b^3, c^2, (ab)^3, (bc)^2, (ca)^5, (bac)^3 \rangle.$$

The quickest way to finish the proof is by using GAP [GAP19] to compute the order of this group:

```
gap> F:=FreeGroup("a","b","c");;
gap> AssignGeneratorVariables(F);;
#I Assigned the global variables [ a, b, c ]
gap> R:=[a^2,b^3,c^2,(a*b)^3,(b*c)^2,(c*a)^5,(b*a*c)^3];;
gap> Order(F/R);
60
```

In Appendix C we give an alternative proof by hand. □

Proposition 5.2. *Let $w_0, \dots, w_k \in \ker(\phi)$. If the group Γ_k admits a representation ρ such that*

- (i) $\rho(w_i) = 1$ for each $i = 0, \dots, k$ and
- (ii) *there exists $r \in \{x_0, \dots, x_k, (bac)^3\}$ such that $\rho(r) \neq 1$*

then $\Gamma_k / \langle\langle w_0, \dots, w_k \rangle\rangle \xrightarrow{\bar{\phi}} A_5$ is not an isomorphism.

Proof. This follows from Lemma 5.1. □

Remark 5.3. Note that in some cases (for example when $k = 0$ and $w_0 = x_0$) a representation of Γ_k with image isomorphic to A_5 may suffice to conclude that $\Gamma_k / \langle\langle w_0, \dots, w_k \rangle\rangle$ is not A_5 . This may seem counterintuitive.

If $\alpha, \beta \in \mathbb{C}$ we consider the matrix

$$R(\alpha, \beta) = \begin{pmatrix} \alpha & \beta & 0 \\ -\beta & \alpha & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

which lies in $\mathrm{SO}(3, \mathbb{C})$ whenever $\alpha^2 + \beta^2 = 1$. Recall that $\mathrm{SO}(n, \mathbb{C})$ is the group of matrices $M \in M_n(\mathbb{C})$ such that $M \cdot M^T = 1$ and $\det(M) = 1$. We now introduce our moduli of representations of Γ_k .

Theorem 5.4. *If $\mathbf{z} = (\alpha_1, \beta_1, \alpha_2, \beta_2, \alpha_3, \beta_3, X_1, \dots, X_k) \in \mathbb{C}^6 \times \mathrm{SO}(3, \mathbb{C})^k$ satisfies $\alpha_i^2 + \beta_i^2 = 1$ for $i = 1, 2, 3$ then there is a group representation*

$$\rho_{\mathbf{z}}: \Gamma_k \rightarrow \mathrm{SO}(3, \mathbb{C})$$

defined by the following matrices

$$\begin{aligned} A &= \begin{pmatrix} -1 & 0 & 0 \\ 0 & \frac{1}{3} & -\frac{2}{3}\sqrt{2} \\ 0 & -\frac{2}{3}\sqrt{2} & -\frac{1}{3} \end{pmatrix} \\ B &= \begin{pmatrix} -\frac{1}{2} & -\frac{\sqrt{3}}{2} & 0 \\ \frac{\sqrt{3}}{2} & -\frac{1}{2} & 0 \\ 0 & 0 & 1 \end{pmatrix} \\ C &= R(\alpha_1, \beta_1) S_0 R(\alpha_1, \beta_1)^T \\ D &= R(\alpha_1, \beta_1) S_1 R(\alpha_2, \beta_2) S_2 R(\alpha_2, \beta_2)^T S_1^T R(\alpha_1, \beta_1)^T \\ X_0 &= R(\alpha_1, \beta_1) S_1 R(\alpha_2, \beta_2) S_3 R(\alpha_3, \beta_3) S_4 \end{aligned}$$

where

$$\begin{aligned} S_0 &= \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, S_1 = \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & -1 & 0 \end{pmatrix}, S_2 = \begin{pmatrix} -\cos(\frac{2\pi}{5}) & 0 & -\sin(\frac{2\pi}{5}) \\ 0 & -1 & 0 \\ -\sin(\frac{2\pi}{5}) & 0 & \cos(\frac{2\pi}{5}) \end{pmatrix}, \\ S_3 &= \begin{pmatrix} 0 & \cos(\frac{\pi}{5}) & \sin(\frac{\pi}{5}) \\ 1 & 0 & 0 \\ 0 & \sin(\frac{\pi}{5}) & -\cos(\frac{\pi}{5}) \end{pmatrix} \text{ and } S_4 = \begin{pmatrix} 0 & -\frac{\sqrt{3}}{3} & -\frac{\sqrt{6}}{3} \\ 1 & 0 & 0 \\ 0 & -\frac{\sqrt{6}}{3} & \frac{\sqrt{3}}{3} \end{pmatrix}. \end{aligned}$$

Proof. The proof reduces to the case $k = 0$. We describe the computations needed to finish the proof. It is straightforward to prove $A^2 = 1$, $B^3 = 1$ and $(AB)^3 = 1$; $C^2 = 1$ and $D^2 = 1$ reduce to $S_0^2 = 1$ and $S_2^2 = 1$ respectively. Since $R(\alpha_1, \beta_1)$ commutes with B , to prove $(BC)^2 = 1$ it is enough to verify $(BS_0)^2 = 1$. To prove $(CD)^5 = 1$ it is enough to verify that $(S_1^T S_0 S_1 R(\alpha_2, \beta_2) S_2 R(\alpha_2, \beta_2)^T)^5 = 1$ and, since

$$S_1^T S_0 S_1 = \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

commutes with $R(\alpha_2, \beta_2)$, this reduces to proving $(S_1^T S_0 S_1 S_2)^5 = 1$ which follows from

$$S_1^T S_0 S_1 S_2 = \begin{pmatrix} \cos(\frac{2\pi}{5}) & 0 & \sin(\frac{2\pi}{5}) \\ 0 & 1 & 0 \\ -\sin(\frac{2\pi}{5}) & 0 & \cos(\frac{2\pi}{5}) \end{pmatrix}.$$

Finally, $X_0 A X_0^T = D$ reduces to $S_3 R(\alpha_3, \beta_3) S_4 A S_4^T R(\alpha_3, \beta_3)^T S_3^T = S_2$ which follows from

$$S_4 A S_4^T = S_3^T S_2 S_3 = \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

The function `check_rep` in Appendix D gives an alternative proof using SAGE [Sag19]. \square

Remark 5.5. We also regard A, B, C, D, X_0 as matrices with coefficients in the polynomial ring $\mathbb{C}[\alpha_1, \beta_1, \alpha_2, \beta_2, \alpha_3, \beta_3]$.

Remark 5.6. This family of representations was obtained in the following way. We first obtained a single representation of the group Γ_0 in $\text{SO}(3, \mathbb{R})$ by choosing reflections $\sigma_1, \sigma_2, \sigma_3, \sigma_4, \sigma_5$ with axes forming the appropriate angles so that $a \mapsto \sigma_1 \sigma_2$, $b \mapsto \sigma_2 \sigma_3$, $c \mapsto \sigma_3 \sigma_4$ and $d \mapsto \sigma_4 \sigma_5$ defines a representation of the (alternating Coxeter) group generated by a, b, c , and d . Since $\sigma_1 \sigma_2$ and $\sigma_4 \sigma_5$ are rotations of the same angle, they are conjugate, so it is possible to extend this to a representation of Γ_0 by mapping x_0 to a rotation r . Then we *twisted* this representation in the following way to obtain three degrees of freedom. If θ_1, θ_2 and θ_3 are rotations commuting with $\sigma_1 \sigma_2$, $\sigma_2 \sigma_3$, and $\sigma_3 \sigma_4$ respectively then $a \mapsto \sigma_1 \sigma_2$, $b \mapsto \sigma_2 \sigma_3$, $c \mapsto \theta_2 \sigma_3 \sigma_4 \theta_2^{-1}$, $d \mapsto \theta_2 \theta_3 \sigma_4 \sigma_5 \theta_3^{-1} \theta_2^{-1}$ and $x_0 \mapsto \theta_2 \theta_3 r \theta_1$ gives a representation of Γ_0 . After tidying up these computations we obtain the moduli in Theorem 5.4.

Remark 5.7. Given a family $\{w_i\}_{i \in I}$ of words in $F(a, b, c, d, x_0, \dots, x_k)$, the set of points $\mathbf{z} \in \mathbb{C}^6 \times \text{SO}(3, \mathbb{C})^k \subseteq \mathbb{C}^{6+9k}$ such that $\rho_{\mathbf{z}}(w_i) = 1$ for all $i \in I$ is an affine algebraic variety that we denote $Z(\{w_i : i \in I\})$. For $k = 0$ the variety $Z(w_0)$ can be described with only 6 equations. More generally, if we allow X_1, \dots, X_k to take values in $\text{O}(3, \mathbb{C})$ the variety $Z(w_0, \dots, w_k)$ can be described using $6 + 9k$ equations. This suggests that it may be possible to use a result such as Bézout's theorem to count points. We could not finish this approach so we took a different one.

Proposition 5.8. *There is exactly one choice of*

$$(\alpha_1, \beta_1, \alpha_2, \beta_2, \alpha_3, \beta_3, X_1, \dots, X_k) \in \mathbb{C}^6 \times \text{SO}(3, \mathbb{C})^k$$

with $\alpha_i^2 + \beta_i^2 = 1$ for $i = 1, 2, 3$ such that the matrices in Theorem 5.4 satisfy

$$X_0 = X_1 = \dots = X_k = (BAC)^3 = 1.$$

The unique solution

$$\mathbf{z}_u = (\alpha_1^u, \beta_1^u, \alpha_2^u, \beta_2^u, \alpha_3^u, \beta_3^u, 1, \dots, 1)$$

is real and its exact value is given by

$$\begin{aligned} \alpha_1^u &= -\frac{1}{4}\sqrt{3\sqrt{5}+9} & \alpha_2^u &= -\sqrt{-\frac{2}{15}\sqrt{5}+\frac{1}{3}} & \alpha_3^u &= -\sqrt{-\frac{1}{5}\sqrt{5}+\frac{1}{2}} \\ \beta_1^u &= \frac{1}{4}\sqrt{-3\sqrt{5}+7} & \beta_2^u &= \sqrt{\frac{2}{15}\sqrt{5}+\frac{2}{3}} & \beta_3^u &= \sqrt{\frac{1}{5}\sqrt{5}+\frac{1}{2}}. \end{aligned}$$

Proof. Again this reduces to the case $k = 0$. In Appendix D we give a proof using SAGE. We indicate here how to prove this by hand. We rewrite $(BAC)^3 = 1$ as

$$(BAC)^2 - (BAC)^T = 0 \quad (\text{i})$$

and $X_0 = 1$ as

$$S_4 R(\alpha_1, \beta_1) S_1 R(\alpha_2, \beta_2) - R(\alpha_3, \beta_3)^T S_3^T = 0. \quad (\text{ii})$$

We have

$$BAC = \begin{pmatrix} -\frac{\sqrt{3}}{3}\alpha_1\beta_1 - \frac{1}{2}(\alpha_1^2 - \beta_1^2) & \alpha_1\beta_1 - \frac{\sqrt{3}}{6}(\alpha_1^2 - \beta_1^2) & -\frac{\sqrt{6}}{3} \\ -\frac{1}{3}\alpha_1\beta_1 + \frac{\sqrt{3}}{2}(\alpha_1^2 - \beta_1^2) & -\sqrt{3}\alpha_1\beta_1 - \frac{1}{6}(\alpha_1^2 - \beta_1^2) & -\frac{\sqrt{2}}{3} \\ -\frac{4\sqrt{2}}{3}\alpha_1\beta_1 & -\frac{2\sqrt{2}}{3}(\alpha_1^2 - \beta_1^2) & \frac{1}{3} \end{pmatrix}.$$

Then the $(3, 3)$ entry of (i) gives $\frac{8\sqrt{3}}{9}\alpha_1\beta_1 + \frac{4}{9}(\alpha_1^2 - \beta_1^2) - \frac{2}{9} = 0$ and from $\alpha_1^2 + \beta_1^2 = 1$ we obtain

$$\frac{8\sqrt{3}}{9}\alpha_1\beta_1 + \frac{8}{9}\alpha_1^2 - \frac{2}{3} = 0. \quad (\text{iii})$$

To find the entries of (ii) it is useful to recall that $\cos(\frac{2\pi}{5}) = \frac{1}{4}(\sqrt{5} - 1)$, $\sin(\frac{2\pi}{5}) = \frac{1}{4}\sqrt{2\sqrt{5} + 10}$, $\cos(\frac{\pi}{5}) = \frac{1}{4}(\sqrt{5} + 1)$, and $\sin(\frac{\pi}{5}) = \frac{1}{4}\sqrt{-2\sqrt{5} + 10}$. We obtain

$$\begin{pmatrix} -\frac{\sqrt{3}}{3}\alpha_2\beta_1 - \frac{\sqrt{6}}{3}\beta_2 + \frac{\sqrt{5}+1}{4}\beta_3 & -\frac{\sqrt{3}}{3}\beta_1\beta_2 + \frac{\sqrt{6}}{3}\alpha_2 - \alpha_3 & \frac{\sqrt{3}}{3}\alpha_1 + \frac{\sqrt{-2\sqrt{5}+10}}{4}\beta_3 \\ -\alpha_1\alpha_2 - \frac{\sqrt{5}+1}{4}\alpha_3 & -\alpha_1\beta_2 - \beta_3 & -\beta_1 - \frac{\sqrt{-2\sqrt{5}+10}}{4}\alpha_3 \\ -\frac{\sqrt{6}}{3}\alpha_2\beta_1 + \frac{\sqrt{3}}{3}\beta_2 - \frac{\sqrt{-2\sqrt{5}+10}}{4} & -\frac{\sqrt{6}}{3}\beta_1\beta_2 - \frac{\sqrt{3}}{3}\alpha_2 & \frac{\sqrt{6}}{3}\alpha_1 + \frac{\sqrt{5}+1}{4} \end{pmatrix} = 0.$$

The $(3, 3)$ entry determines the value of α_1 . From (iii) we obtain the value of β_1 . The $(1, 3)$ entry allows to obtain the value of β_3 . Now the $(2, 2)$ entry gives the value of β_2 . The $(3, 2)$ entry gives the value of α_2 and finally the $(2, 3)$ entry determines the value of α_3 . Computing the remaining entries we see these values form a solution to $X_0 = (BAC)^3 = 1$ and satisfy $\alpha_i^2 + \beta_i^2 = 1$. \square

Remark 5.9. We say that \mathbf{z}_u is *universal* in the following sense: if $\{w_i\}_{i \in I} \subseteq \ker(\phi)$ then $\mathbf{z}_u \in Z(\{w_i : i \in I\})$.

The following result is proved in Section 6.

Theorem 5.10. *Let $w_0, \dots, w_k \in \ker(\phi)$. Let $N = \ker(\bar{\phi})$. If $N = \langle\langle w_0, \dots, w_k \rangle\rangle^{\Gamma_k}[N, N]$ then the variety $Z(w_0, \dots, w_k)$ has at least two different points.*

Note that, by Proposition 4.3, the condition $N = \langle\langle w_0, \dots, w_k \rangle\rangle^{\Gamma_k}[N, N]$ is equivalent to the acyclicity of the corresponding 2-complex. This is also the same as saying that w_0, \dots, w_k generate the A_5 -module $N/[N, N]$ (i.e. the *relation module* of $1 \rightarrow N \rightarrow \Gamma_k \xrightarrow{\bar{\phi}} A_5 \rightarrow 1$).

6. QUATERNIONS

To prove Theorem 5.10 we study the real part of the moduli, working with quaternions instead of orthogonal matrices. This is useful because representing a rotation as a quaternion allows to find the axis easily.

Recall that $S^3 = \{q \in \mathbb{H} : |q| = 1\}$ acts on $S^2 = \{b\mathbf{i} + c\mathbf{j} + d\mathbf{k} : b^2 + c^2 + d^2 = 1\}$ by conjugation. Recall that any element of S^3 can be written as $\cos(\theta/2) + \sin(\theta/2)q$ with $\theta \in [0, 2\pi]$ and $q = b\mathbf{i} + c\mathbf{j} + d\mathbf{k} \in S^2$. There is a homomorphism $p: S^3 \rightarrow \text{SO}(3, \mathbb{R})$ with $\ker(p) = \{1, -1\}$ and which sends $\cos(\frac{\theta}{2}) + \sin(\frac{\theta}{2})(b\mathbf{i} + c\mathbf{j} + d\mathbf{k})$ to the rotation matrix with angle θ and axis (b, c, d) . Note that $\tilde{R}(t) = \cos(\frac{t}{2}) + \mathbf{k}\sin(\frac{t}{2})$ is a lift of $R(\cos(t), \sin(t))$ by p .

Let $\psi: \mathbb{H} \rightarrow \mathbb{R}^3$ be given by $a + b\mathbf{i} + c\mathbf{j} + d\mathbf{k} \mapsto (b, c, d)$. Recall that if $q \in S^3$ and v is a pure quaternion we have $\psi(qvq^{-1}) = p(q) \cdot \psi(v)$. Let $\mathbb{D}^3 \subset \mathbb{R}^3$ be the unit disk. Let $\varphi: \mathbb{D}^3 \rightarrow \mathbb{H}$ be given by $(b, c, d) \mapsto \sqrt{1 - b^2 - c^2 - d^2} + b\mathbf{i} + c\mathbf{j} + d\mathbf{k}$. We denote the coordinates of $[0, 2\pi]^3 \times (\mathbb{D}^3)^k$ by $t_1, t_2, t_3, \dots, t_{3(k+1)}$.

Definition 6.1. Let $\tilde{A}, \tilde{B}, \tilde{S}_0, \tilde{S}_1, \tilde{S}_2, \tilde{S}_3, \tilde{S}_4$, be preimages by p of the matrices $A, B, S_0, S_1, S_2, S_3, S_4$ which appear in the statement of Theorem 5.4. We also define functions $\tilde{C}, \tilde{D}, \tilde{X}_0: [0, 2\pi]^3 \times (\mathbb{D}^3)^k \rightarrow \mathbb{H}$ by

$$\begin{aligned}\tilde{C}(\mathbf{t}) &= \tilde{R}(t_1) \tilde{S}_0 \tilde{R}(t_1)^{-1}, \\ \tilde{D}(\mathbf{t}) &= \tilde{R}(t_1) \tilde{S}_1 \tilde{R}(t_2) \tilde{S}_2 \tilde{R}(t_2)^{-1} \tilde{S}_1^{-1} \tilde{R}(t_1)^{-1}, \\ \tilde{X}_0(\mathbf{t}) &= \tilde{R}(t_1) \tilde{S}_1 \tilde{R}(t_2) \tilde{S}_3 \tilde{R}(t_3) \tilde{S}_4.\end{aligned}$$

For $i = 1, \dots, k$ we define $\tilde{X}_i(\mathbf{t}) = \varphi(t_{3i+1}, t_{3i+2}, t_{3i+3})$. Let $t_1^u, t_2^u, t_3^u \in [0, 2\pi]^3$ be the unique numbers such that $\cos(t_i^u) = \alpha_i^u$ and $\sin(t_i^u) = \beta_i^u$. Let $\mathbf{t}_u = (t_1^u, t_2^u, t_3^u, 0, \dots, 0) \in [0, 2\pi]^3 \times (\mathbb{D}^3)^k$. Note that we can arrange the signs of these preimages so that $(\tilde{B}\tilde{A}\tilde{C})^3(\mathbf{t}_u) = 1$ and $\tilde{X}_0(\mathbf{t}_u) = 1$.

If $w \in F(a, b, c, d, x_0, \dots, x_k)$ there is an induced map $\widetilde{W}: [0, 2\pi]^3 \times (\mathbb{D}^3)^k \rightarrow S^3$. Note that any two words w, w' which are equal in Γ_k induce maps $\widetilde{W}, \widetilde{W}'$ which are equal or differ on a sign. If $w_0, \dots, w_k \in \ker(\phi)$ we can consider

$$\widetilde{\mathbf{W}} = (\widetilde{W}_0, \dots, \widetilde{W}_k): [0, 2\pi]^3 \times (\mathbb{D}^3)^k \rightarrow (S^3)^{k+1}$$

which can be composed with

$$\Psi = (\psi, \dots, \psi): \mathbb{H}^{k+1} \rightarrow \mathbb{R}^{3(k+1)}$$

to obtain a map

$$\Psi \widetilde{\mathbf{W}}: [0, 2\pi]^3 \times (\mathbb{D}^3)^k \rightarrow (\mathbb{D}^3)^{k+1}.$$

The plan is to assume \mathbf{t}_u is the only zero in order to do a degree argument. We will get a contradiction by computing the degree in two different ways.

Lemma 6.2. Let $I = [-1, 1]$ and let $\mathbb{D}^3 \subset \mathbb{R}^3$ be the unit disk. Let

$$\mathbf{F} = (f_0, \dots, f_k): I^3 \times (\mathbb{D}^3)^k \rightarrow (\mathbb{D}^3)^{k+1}$$

be a continuous map which is nonzero on the boundary of $I^3 \times (\mathbb{D}^3)^k$ and satisfies the following parity condition:

- For $t_1, t_2, t_3 \in I, x_1, \dots, x_k \in \mathbb{D}^3$ we have

$$\begin{aligned}(f_0, f_1, \dots, f_k)((-1, t_2, t_3), x_1, \dots, x_k) &= (-f_0, f_1, \dots, f_k)((1, t_2, t_3), x_1, \dots, x_k) \\ (f_0, f_1, \dots, f_k)((t_1, -1, t_3), x_1, \dots, x_k) &= (-f_0, f_1, \dots, f_k)((t_1, 1, t_3), x_1, \dots, x_k) \\ (f_0, f_1, \dots, f_k)((t_1, t_2, -1), x_1, \dots, x_k) &= (-f_0, f_1, \dots, f_k)((t_1, t_2, 1), x_1, \dots, x_k).\end{aligned}$$
- For each $1 \leq i \leq k$ and for every $(x_0, \dots, x_k) \in I^3 \times (\mathbb{D}^3)^k$ with $x_i \in \partial\mathbb{D}^3$ we have

$$(f_0, f_1, \dots, f_k)(x_0, \dots, x_{i-1}, -x_i, x_{i+1}, \dots, x_k) = (f_0, \dots, f_{i-1}, -f_i, f_{i+1}, \dots, f_k)(x_0, \dots, x_k).$$

Then the restriction

$$\mathbf{F}: \partial(I^3 \times (\mathbb{D}^3)^k) \rightarrow (\mathbb{D}^3)^{k+1} - \{0\}$$

has even degree.

Proof. We fix cellular structures. For I we take the structure with two 0-cells and one 1-cell. For \mathbb{D}^3 we take the cell structure with two 0-cells, two 1-cells, two 2-cells and one 3-cell (the antipodal map interchanges the i -cells in each pair for $0 \leq i \leq 2$). We take the product cellular structure for I^3 , $I^3 \times (\mathbb{D}^3)^k$ and $(\mathbb{D}^3)^{k+1}$. Let $S = \partial(I^3 \times (\mathbb{D}^3)^k)$. Note that the $(3k+2)$ -cells of S can be divided into $3+k$ pairs of *opposite* cells in a natural way. Note that it is easy to define a cellular map $h_0: I^3 \rightarrow \partial \mathbb{D}^3$ which satisfies

$$\begin{aligned} h_0(-1, t_2, t_3) &= -h_0(1, t_2, t_3) \\ h_0(t_1, -1, t_3) &= -h_0(t_1, 1, t_3) \\ h_0(t_1, t_2, -1) &= -h_0(t_1, t_2, 1). \end{aligned}$$

Let $h_i: \mathbb{D}^3 \rightarrow \mathbb{D}^3$ be the identity for $1 \leq i \leq k$. Now we can define a homotopy between $\mathbf{F}|_S$ and a map $\mathbf{G}: S \rightarrow \partial(\mathbb{D}^3)^{k+1}$ that satisfies the parity condition and coincides with $\mathbf{H} = (h_0, \dots, h_k)$ on the $(3k+1)$ -skeleton of S . This is done skeleton by skeleton using that $\partial(\mathbb{D}^3)^{k+1}$ is $(3k+1)$ -connected. For each pair of opposite $(3k+2)$ -cells we can extend the homotopy so that the parity condition is also satisfied by \mathbf{G} . Clearly the degrees of $\mathbf{F}|_S$ and \mathbf{G} are equal. Now note that if e, e' is a pair of opposite $(3k+2)$ -cells then $\mathbf{G}_*(e), \mathbf{H}_*(e) \in C_{3k+2}(\partial(\mathbb{D}^3)^{k+1})$ differ on an element of $H_{3k+2}(\partial(\mathbb{D}^3)^{k+1})$. Moreover, by the parity condition, $\mathbf{G}_*(e')$ and $\mathbf{H}_*(e')$ differ on the same element. Thus the degree of $\mathbf{H}|_S$ and the degree of \mathbf{G} are equal modulo 2. To conclude, note that $\deg(\mathbf{H}|_S) = 0$ since $\mathbf{H}: I^3 \times (\mathbb{D}^3)^k \rightarrow \partial(\mathbb{D}^3)^{k+1}$ is an extension of $\mathbf{H}|_S$ to a contractible space. \square

Corollary 6.3. *Let $w_0, \dots, w_k \in F(a, b, c, d, x_0, \dots, x_k)$ be words and assume the total exponent of x_i in w_j is $\delta_{i,j}$. If $\Psi\tilde{\mathbf{W}}$ is nonzero on the boundary of $[0, 2\pi]^3 \times (\mathbb{D}^3)^k$, then the degree of the restriction $\Psi\tilde{\mathbf{W}}: \partial([0, 2\pi]^3 \times (\mathbb{D}^3)^k) \rightarrow (\mathbb{D}^3)^{k+1} - \{0\}$ is even.*

Proof. Since the total exponent of x_i in w_j is $\delta_{i,j}$, by looking at Definition 6.1 we see the parity condition of Lemma 6.2 is satisfied. \square

Recall that the degree can be computed in the following way

Lemma 6.4. *Let $f: \mathbb{R}^n \rightarrow \mathbb{R}^n$ be smooth and assume $f(0) = 0$. If $\det(Df_0) \neq 0$ then 0 is an isolated zero and the degree of f around 0 is given by $\deg(f, 0) = \text{sg}(\det(Df_0))$.*

We need some basic differentiation properties for quaternion valued analytic functions analogous to the usual ones (see Appendix B). Note that $\tilde{R}(t) = \cos(\frac{t}{2}) + \mathbf{k} \sin(\frac{t}{2}) = 1 + \frac{t}{2}\mathbf{k} + O(t^2)$.

Lemma 6.5. *Let $\tilde{\mathbf{X}} = (\tilde{X}_0, \dots, \tilde{X}_k)$. Then $D(\Psi\tilde{\mathbf{X}})_{\mathbf{t}_u}$ is invertible.*

Proof. Again this reduces to the case $k = 0$ by noting that

$$D(\Psi\tilde{\mathbf{X}})_{\mathbf{t}_u} = \begin{pmatrix} M & 0 \\ 0 & I \end{pmatrix}$$

where M is the 3×3 matrix we obtain in the $k = 0$ case. We now prove M is invertible. Recall that $\tilde{X}_0(\mathbf{t}_u) = 1$. Then

$$\begin{aligned} \tilde{X}_0(\mathbf{t}_u + \mathbf{t}) &= \tilde{R}(t_1^u) \tilde{R}(t_1) \tilde{S}_1 \tilde{R}(t_2^u) \tilde{R}(t_2) \tilde{S}_3 \tilde{R}(t_3^u) \tilde{R}(t_3) \tilde{S}_4 \\ &= \tilde{R}(t_1^u) \left(1 + \frac{t_1}{2} \mathbf{k}\right) \tilde{S}_1 \tilde{R}(t_2^u) \left(1 + \frac{t_2}{2} \mathbf{k}\right) \tilde{S}_3 \tilde{R}(t_3^u) \left(1 + \frac{t_3}{2} \mathbf{k}\right) \tilde{S}_4 + O(\mathbf{t}^2) \\ &= 1 + \frac{1}{2} \tilde{R}(t_1^u) \mathbf{k} \tilde{R}(t_1^u)^{-1} t_1 + \frac{1}{2} \left(\tilde{R}(t_1^u) \tilde{S}_1 \tilde{R}(t_2^u) \right) \mathbf{k} \left(\tilde{R}(t_1^u) \tilde{S}_1 \tilde{R}(t_2^u) \right)^{-1} t_2 \\ &\quad + \frac{1}{2} \tilde{S}_4^{-1} \mathbf{k} \tilde{S}_4 t_3 + O(\mathbf{t}^2) \end{aligned}$$

Now recalling that $q \mathbf{k} q^{-1} = (\mathbf{i}, \mathbf{j}, \mathbf{k}) \cdot p(q) \cdot (0, 0, 1)$ for any $q \in S^3$ we see that the columns of M are given by

$$\begin{aligned} \frac{1}{2} R(\alpha_1^u, \beta_1^u) \cdot (0, 0, 1) &= \left(0, 0, \frac{1}{2}\right) \\ \frac{1}{2} R(\alpha_1^u, \beta_1^u) S_1 R(\alpha_2^u, \beta_2^u) \cdot (0, 0, 1) &= \left(-\frac{1}{2} \beta_1^u, -\frac{1}{2} \alpha_1^u, 0\right) \\ \frac{1}{2} S_4^{-1} \cdot (0, 0, 1) &= \left(0, -\frac{1}{6} \sqrt{6}, \frac{1}{6} \sqrt{3}\right). \end{aligned}$$

Thus

$$M = \begin{pmatrix} 0 & -\frac{1}{2} \beta_1^u & 0 \\ 0 & -\frac{1}{2} \alpha_1^u & -\frac{1}{6} \sqrt{6} \\ \frac{1}{2} & 0 & \frac{1}{6} \sqrt{3} \end{pmatrix}$$

and therefore $\det(M) = \frac{1}{24} \sqrt{6} \beta_1^u \neq 0$. \square

Lemma 6.6. *Let $w \in \ker(\phi)$. Then $\frac{\partial \tilde{W}}{\partial t_i}(\mathbf{t}_u)$ is a pure quaternion for $i = 1, \dots, 3(k+1)$.*

Proof. Since w belongs to $\ker(\phi)$, in Γ_k it equals a product of conjugates of the x_i , $(bac)^3$ and their inverses. Recall that S^2 is invariant by the action of S^3 . By Proposition B.1, it is enough to prove that $\frac{\partial \tilde{X}_j}{\partial t_i}(\mathbf{t}_u)$ and $\frac{\partial (\tilde{B} \tilde{A} \tilde{C})^3}{\partial t_i}(\mathbf{t}_u)$ are pure quaternions.

For $i = 0$ the first claim follows from the computation in the proof of Lemma 6.5 and is easy to verify for $i > 0$. The second claim follows similarly by noting that $(\tilde{B} \tilde{A} \tilde{C})^3(\mathbf{t}_u) = 1$ and writing

$$(\tilde{B} \tilde{A} \tilde{C})^3(\mathbf{t}_u + \mathbf{t}) = \left(\tilde{B} \tilde{A} \left(1 + \frac{t_1}{2} \mathbf{k}\right) \tilde{S}_0 \left(1 - \frac{t_1}{2} \mathbf{k}\right) \right)^3 + O(\mathbf{t}^2).$$

\square

Lemma 6.7. *Let $N = \ker(\bar{\phi})$ and let $w_0, \dots, w_k \in \ker(\phi)$. If $N = \langle\langle w_0, \dots, w_k \rangle\rangle^{\Gamma_k} [N, N]$ then $D(\Psi \tilde{\mathbf{W}})_{\mathbf{t}_u}$ is invertible.*

Proof. We may assume without loss of generality that $\tilde{W}_j(\mathbf{t}_u) = 1$ for all j . For each j there are numbers $a_j, \ell_j \in \mathbb{N}_0$, words $v_{j,1}, \dots, v_{j,a_j}, u_{j,1}, \dots, v_{j,a_j} \in \ker(\phi)$, words $p_{j,1}, \dots, p_{j,\ell_j} \in F(a, b, c, d, x_0, \dots, x_k)$, indices $\alpha_{j,1}, \dots, \alpha_{j,\ell_j} \in \{0, \dots, k\}$ and signs $\varepsilon_{j,1}, \dots, \varepsilon_{j,\ell_j} \in \{1, -1\}$ such that in Γ_k we have

$$x_j = \prod_{s=1}^{\ell_j} p_{j,s} w_{\alpha_{j,s}}^{\varepsilon_{j,s}} p_{j,s}^{-1} \prod_{i=1}^{a_j} [u_{j,i}, v_{j,i}].$$

Then we have

$$\tilde{X}_j(\mathbf{t}_u + \mathbf{t}) = \left(\prod_{s=1}^{\ell_j} \tilde{P}_{j,s} \tilde{W}_{\alpha_{j,s}}^{\varepsilon_{j,s}} \tilde{P}_{j,s}^{-1} \prod_{i=1}^{a_j} [\tilde{U}_{j,i}, \tilde{V}_{j,i}] \right) (\mathbf{t}_u + \mathbf{t})$$

and using Proposition B.1 we obtain

$$\begin{aligned} \frac{\partial \tilde{X}_j}{\partial t_i}(\mathbf{t}_u) &= \sum_{s=1}^{\ell_j} \tilde{P}_{j,s}(\mathbf{t}_u) \frac{\partial \tilde{W}_{\alpha_{j,s}}^{\varepsilon_{j,s}}}{\partial t_i}(\mathbf{t}_u) \tilde{P}_{j,s}^{-1}(\mathbf{t}_u) \\ &= \sum_{s=1}^{\ell_j} \varepsilon_{j,s} \tilde{P}_{j,s}(\mathbf{t}_u) \frac{\partial \tilde{W}_{\alpha_{j,s}}}{\partial t_i}(\mathbf{t}_u) \tilde{P}_{j,s}^{-1}(\mathbf{t}_u) \end{aligned}$$

By Lemma 6.6, $D(\Psi \tilde{\mathbf{W}})_{\mathbf{t}_u}$ is invertible if and only if

$$\left\{ \left(\frac{\partial \tilde{W}_0}{\partial t_i}(\mathbf{t}_u), \dots, \frac{\partial \tilde{W}_k}{\partial t_i}(\mathbf{t}_u) \right) : 1 \leq i \leq 3(k+1) \right\}$$

is linearly independent over \mathbb{R} . If $\lambda_i \in \mathbb{R}$ satisfy

$$\sum_{i=1}^{3(k+1)} \lambda_i \frac{\partial \tilde{\mathbf{W}}}{\partial t_i}(\mathbf{t}_u) = 0$$

it follows that

$$\sum_{i=1}^{3(k+1)} \lambda_i \frac{\partial \tilde{\mathbf{X}}}{\partial t_i}(\mathbf{t}_u) = 0.$$

By Lemma 6.5, $D(\Psi \tilde{\mathbf{X}})_{\mathbf{t}_u}$ is invertible and, again by Lemma 6.6, the set

$$\left\{ \left(\frac{\partial \tilde{X}_0}{\partial t_i}(\mathbf{t}_u), \dots, \frac{\partial \tilde{X}_k}{\partial t_i}(\mathbf{t}_u) \right) : 1 \leq i \leq 3(k+1) \right\}$$

is linearly independent over \mathbb{R} . Thus $\lambda_1 = \dots = \lambda_{3(k+1)} = 0$ and we are done. \square

Proof of Theorem 5.10. We can assume that the total exponent of x_i in w_j is $\delta_{i,j}$. To prove this, consider the abelianization and note that it is possible to achieve this by using the following operations:

- replacing w_i by $w_i w_j$ (if $i \neq j$),
- replacing w_i by w_i^{-1} , and
- interchanging w_i and w_j .

By Lemma 6.7 and Lemma 6.4, the degree of $\Psi \tilde{\mathbf{W}}$ near \mathbf{t}_u is ± 1 . If $\Psi \tilde{\mathbf{W}}$ has a zero on $\partial([0, 2\pi]^3 \times (\mathbb{D}^3)^k)$ we are done. Otherwise, by Corollary 6.3, the degree of $\Psi \tilde{\mathbf{W}}$ restricted to the boundary of $[0, 2\pi]^3 \times (\mathbb{D}^3)^k$ is even. It follows that there must be a point $\mathbf{t} \neq \mathbf{t}_u$ such that $\Psi \tilde{\mathbf{W}}(\mathbf{t}) = 0$. This gives a second point in $Z(w_0, \dots, w_k)$. \square

7. GROUP ACTIONS OF A_5 ON CONTRACTIBLE 2-COMPLEXES

We can now prove the following.

Theorem 7.1. *There is no presentation of A_5 of the form*

$$\langle a, b, c, d, x_0, \dots, x_k \mid a^2, b^3, c^2, d^2, (ab)^3, (bc)^2, (cd)^5, x_0 a x_0^{-1} = d, w_0, \dots, w_k \rangle$$

with $w_0, \dots, w_k \in \ker(\phi)$, where $\phi: F(a, b, c, d, x_0, \dots, x_k) \rightarrow A_5$ is given by $a \mapsto (2, 5)(3, 4)$, $b \mapsto (3, 5, 4)$, $c \mapsto (1, 2)(3, 5)$, $d \mapsto (2, 5)(3, 4)$ and $x_i \mapsto 1$ for each $i = 0, \dots, k$.

Proof. This follows from Theorem 5.10, Proposition 5.8 and Proposition 5.2. \square

Now from Theorem 7.1 and Theorem 4.5 we deduce.

Theorem 7.2. *Every action of $A_5 \cong \mathrm{PSL}_2(2^2)$ on a finite, contractible 2-complex has a fixed point.*

Corollary 7.3. *Let G be one of the groups $\mathrm{PSL}_2(2^{2k})$, $\mathrm{PSL}_2(5^k)$ for $k \geq 1$ or $\mathrm{PSL}_2(q)$ for $q \equiv \pm 3 \pmod{8}$ and $q \equiv \pm 1 \pmod{5}$. Then every action of G on a finite contractible 2-complex has a fixed point.*

Proof. Let G be one of these groups and let X be a finite acyclic fixed point free 2-dimensional G -complex. By [OS02, Proposition 3.3], A_5 is a subgroup of G and by Lemma 2.8 the action of A_5 on X is fixed point free. By Theorem 7.2 X cannot be contractible. \square

Looking more carefully at the proof of Theorem 7.2 we obtain the following.

Theorem 7.4. *Let X be a fixed point free 2-dimensional finite, acyclic A_5 -complex. Then there is a nontrivial representation $\pi_1(X) \rightarrow \mathrm{SO}(3, \mathbb{R})$.*

Proof. By Theorem 3.6 we see that π surjects onto the fundamental group of an acyclic 2-dimensional A_5 complex X' which is obtained from $\Gamma_{OS}(A_5)$ by attaching $k \geq 0$ free orbits of 1-cells and $k + 1$ free orbits of 2-cells. Now note that the representation constructed to prove Theorem 7.1 restricted to $\pi_1(X')$ gives a nontrivial morphism into $\mathrm{SO}(3, \mathbb{R})$. \square

Since A_5 is the only finite perfect subgroup of $\mathrm{SO}(3, \mathbb{R})$ we deduce the following.

Corollary 7.5. *Let X be a fixed point free 2-dimensional finite and acyclic A_5 -complex and let $\pi = \pi_1(X)$. Then π is infinite or there is an epimorphism $\pi \rightarrow A_5$.*

Recall that $N = \ker(\bar{\phi})$ is a free group of rank $60(k + 1)$. We can restate Theorem 7.1 in the following way which highlights the connection with the relation gap problem (see [Har18, Har15]).

Corollary 7.6. *The extension*

$$1 \rightarrow N \rightarrow \Gamma_k \xrightarrow{\bar{\phi}} A_5 \rightarrow 1$$

has a relation gap. That is, the A_5 -module $N/[N, N]$ is free of rank $k + 1$. However N cannot be generated by $k + 1$ elements as a Γ_k -group.

Note that since Γ_k is not free this is not an example of a presentation with a relation gap.

APPENDIX A. EQUATIONS OVER GROUPS

Let G be a group. An *equation* over G in the variables x_1, \dots, x_n is an element $w \in G * F(x_1, \dots, x_n)$. We say that a system of equations

$$\begin{aligned} w_1(x_1, \dots, x_n) &= 1 \\ w_2(x_1, \dots, x_n) &= 1 \\ &\dots \\ w_m(x_1, \dots, x_n) &= 1 \end{aligned}$$

has a solution in an overgroup of G if the map $G \rightarrow G * F(x_1, \dots, x_m) / \langle\langle w_1, \dots, w_m \rangle\rangle$ is injective. Such a system of equations determines an $(m \times n)$ -matrix M where $M_{i,j}$ is given by the total

exponent of the letter x_j in the word w_i . A system is said to be *independent* if the rank of M is m .

One of the most important open problems in the theory of equations over groups is the Kervaire–Laudenbach–Howie conjecture [How81, Conjecture].

Conjecture A.1 (Kervaire–Laudenbach–Howie). *An independent system of equations over G has a solution in an overgroup of G .*

The Gerstenhaber–Rothaus theorem [GR62, Theorem 3] says that finitely generated subgroups of compact connected Lie groups satisfy Conjecture A.1.

Proposition A.2. *Let X be a finite acyclic 2-complex and let $A \subset X$ be an acyclic subcomplex. Then we can write*

$$\pi_1(X) = \pi_1(A) * F(x_1, \dots, x_n) / \langle\langle w_1, \dots, w_n \rangle\rangle$$

and the $(n \times n)$ -matrix M such that $M_{i,j}$ is the total exponent of x_j in w_i is invertible.

Proof. Take a maximal tree T for A and consider a maximal tree \bar{T} of X containing T . Then $A/T \simeq A$ is an acyclic subcomplex of the acyclic 2-complex $X/\bar{T} \simeq X$. As usual, from A/T we can read a presentation for $\pi_1(A)$ which is balanced since A/T is acyclic. Now we consider a variable x_i for each 1-cell of X/\bar{T} which is not in A/T and we read words from the attaching maps for the 2-cells of X/\bar{T} which are not part of A/T . In this way we obtain equations in these variables with coefficients in A which give the desired description of $\pi_1(X)$. Since X/\bar{T} is acyclic, there is an equal number of variables and equations and the matrix M is invertible. \square

Now from the Gerstenhaber–Rothaus theorem we deduce.

Proposition A.3. *Let X be a finite acyclic 2-complex. If $A \subset X$ is an acyclic subcomplex and there is a nontrivial representation $\rho: \pi_1(A) \rightarrow \mathrm{SO}(n, \mathbb{R})$ then $\pi_1(X)$ is nontrivial.*

Proof. By Proposition A.2 can write $\pi_1(X) = \pi_1(A) * F(x_1, \dots, x_n) / \langle\langle w_1, \dots, w_n \rangle\rangle$ and the system is independent. Let $G_0 = \rho(\pi_1(A))$. There is an induced map

$$\rho: \pi_1(A) * F(x_1, \dots, x_n) \rightarrow G_0 * F(x_1, \dots, x_n)$$

which induces an epimorphism $\pi_1(X) \rightarrow G_0 * F(x_1, \dots, x_n) / \langle\langle \rho(w_1), \dots, \rho(w_n) \rangle\rangle$. Finally from [GR62, Theorem 3] it follows that this group is nontrivial. \square

APPENDIX B. QUATERNION VALUED ANALYTIC FUNCTIONS

A *quaternion valued analytic function* is a function $f: U \rightarrow \mathbb{H}$ where $U \subset \mathbb{R}^n$ is open, such that its components are analytic, that is a function that can be written as

$$f = f_1 + f_i \mathbf{i} + f_j \mathbf{j} + f_k \mathbf{k}$$

with $f_1, f_i, f_j, f_k: U \rightarrow \mathbb{R}$ are analytic. For $i = 1, \dots, n$ we can define the partial derivative

$$\frac{\partial f}{\partial t_i} = \frac{\partial f_1}{\partial t_i} + \frac{\partial f_i}{\partial t_i} \mathbf{i} + \frac{\partial f_j}{\partial t_i} \mathbf{j} + \frac{\partial f_k}{\partial t_i} \mathbf{k}.$$

We define

$$Df_{\mathbf{t}} = \left(\frac{\partial f}{\partial t_1}(\mathbf{t}), \dots, \frac{\partial f}{\partial t_n}(\mathbf{t}) \right).$$

If each coordinate of $\mathbf{F} = (f_1, \dots, f_m): U \rightarrow \mathbb{H}^m$ is analytic then we use the notation

$$\frac{\partial \mathbf{F}}{\partial t_i} = \left(\frac{\partial f_1}{\partial t_i}, \dots, \frac{\partial f_m}{\partial t_i} \right).$$

The usual properties extend to this context. We need the following

Proposition B.1. *Let $f, g: U \rightarrow \mathbb{H}$ be analytic. Then*

(i) *We have the product rule*

$$\frac{\partial f \cdot g}{\partial t_i}(\mathbf{t}) = \frac{\partial f}{\partial t_i}(\mathbf{t})g(\mathbf{t}) + f(\mathbf{t})\frac{\partial g}{\partial t_i}(\mathbf{t}).$$

(ii) *Suppose f is nowhere zero and $g(\mathbf{t}_0) \in \mathbb{R}$ then*

$$\frac{\partial f \cdot g \cdot \frac{1}{f}}{\partial t_i}(\mathbf{t}_0) = f(\mathbf{t}_0)\frac{\partial g}{\partial t_i}(\mathbf{t}_0)f(\mathbf{t}_0)^{-1}.$$

(iii) *Suppose $f(\mathbf{t}_0) = \pm 1$ then*

$$\frac{\partial \frac{1}{f}}{\partial t_i}(\mathbf{t}_0) = -\frac{\partial f}{\partial t_i}(\mathbf{t}_0).$$

(iv) *Suppose that f, g are nowhere zero and $f(\mathbf{t}_0), g(\mathbf{t}_0) \in \{1, -1\}$. Then the commutator $[f, g] = f \cdot g \cdot \frac{1}{f} \cdot \frac{1}{g}$ satisfies $[f, g](\mathbf{t}_0) = 1$ and*

$$\frac{\partial [f, g]}{\partial t_i}(\mathbf{t}_0) = 0.$$

Proof. (i) is a straightforward computation, (ii) and (iii) follow from (i). Finally, (iv) follows from the previous properties. \square

As usual we have the Taylor series

$$f(\mathbf{t}_0 + \mathbf{t}) = f(\mathbf{t}_0) + \sum_{i=1}^n \frac{\partial f}{\partial t_i}(\mathbf{t}_0)t_i + O(\mathbf{t}^2).$$

From the product rule we see that we can multiply the Taylor series of two functions to obtain the Taylor series of the product.

APPENDIX C. ALTERNATIVE PROOF OF LEMMA 5.1

An alternative way to finish the proof of Lemma 5.1 goes by noting that $x = bc$ and $y = ca$ satisfy $x^2 = y^5 = (xy)^3 = 1$. Thus it would suffice to show the group

$$\langle a, b, c \mid a^2, b^3, c^2, (ab)^3, (bc)^2, (ca)^5, (bac)^3 \rangle$$

is generated by x, y , for it is well known that $\langle x, y \mid x^2, y^5, (xy)^3 \rangle$ is a presentation of A_5 . To do this, it is enough to show that $a \in \langle x, y \rangle$. The following computation proves this claim.

$$\begin{aligned}
xy^2xy^{-2}xy &= (bc)(ca)(ca)(bc)(a^{-1}c^{-1})(a^{-1}c^{-1})(bc)(ca) \\
&\quad (\text{using } a^2 = c^2 = 1) = bacabcacacba \\
&\quad (\text{replacing } acba \text{ by } b^{-1}cab^{-1}c) = bacabcacb^{-1}cab^{-1}c \\
&\quad (\text{replacing } cb^{-1}c \text{ by } b) = bacabcabab^{-1}c \\
&\quad (\text{replacing } aba \text{ by } b^2ab^2) = bacabcb^2abc \\
&\quad (\text{replacing } bcb \text{ by } c) = bacacbabc \\
&\quad (\text{replacing } acba \text{ by } b^{-1}cab^{-1}c) = bacb^{-1}cab^{-1}cbc \\
&\quad (\text{replacing } cb^{-1}c \text{ by } b) = babab^{-1}cbc \\
&\quad (\text{replacing } cbc \text{ by } b^{-1}) = babab^{-2} \\
&\quad (\text{using } b^3 = 1) = babab \\
&\quad (\text{using } (ab)^3 = 1) = a.
\end{aligned}$$

APPENDIX D. ALTERNATIVE PROOFS USING SAGE

The following SAGE code gives alternative proofs of Theorem 5.4 and Proposition 5.8 which are easier to verify. Note that SAGE computes exactly over the algebraic numbers so there is no numerical error. The function `check_rep`, shows A, B, C, D, X_0 satisfy the defining relations for Γ_0 in

$$M_3(\mathbb{C}[\alpha_1, \beta_1, \alpha_2, \beta_2, \alpha_3, \beta_3] / \langle \alpha_1^2 + \beta_1^2 - 1, \alpha_2^2 + \beta_2^2 - 1, \alpha_3^2 + \beta_3^2 - 1 \rangle).$$

The function `find_universal_representations` gives the exact value of the unique solution \mathbf{z}_u of $X_0 = (BAC)^3 = 1$ by solving the corresponding system of polynomial equations over the algebraic closure of \mathbb{Q} . Note that in this code we use x_i, y_i instead of α_i, β_i .

```

def R(x,y):
    return matrix([
        (x,y,0),
        (-y,x,0),
        (0,0,1),
    ]);

A = matrix([
    (-1,0,0),
    (0,1/3,-2/3*sqrt(2)),
    (0,-2/3*sqrt(2),-1/3),
]);

B = matrix([
    (-1/2,-sqrt(3)/2,0),
    (sqrt(3)/2, -1/2,0),
    (0,0,1),
]);

```

```

S0 = matrix([
    (-1,0,0),
    (0,1,0),
    (0,0,-1)
])

S1 = matrix([
    (-1,0,0),
    (0,0,-1),
    (0,-1,0),
]);

S2 = matrix([
    (-cos(2*pi/5),0,-sin(2*pi/5)),
    (0,-1,0),
    (-sin(2*pi/5),0,cos(2*pi/5)),
]);

S3 = matrix([
    (0,cos(pi/5),sin(pi/5)),
    (1,0,0),
    (0,sin(pi/5),-cos(pi/5)),
]);

S4 = matrix([
    (0, -sqrt(3)/3, -sqrt(6)/3),
    (1,0,0),
    (0,-sqrt(6)/3,sqrt(3)/3),
]);

def rep(x1,y1,x2,y2,x3,y3):
    C = R(x1,y1) * S0 * R(x1,y1).T;
    D = R(x1,y1) * S1 * R(x2,y2) * S2 * R(x2,y2).T * S1.T * R(x1,y1).T;
    X0 = R(x1,y1) * S1 * R(x2,y2) * S3 * R(x3,y3) * S4;
    return (A,B,C,D,X0);

I = matrix.identity(3);

def check_rep():
    R.<x1,y1,x2,y2,x3,y3> = QQbar[];
    A,B,C,D,X0 = rep(x1,y1,x2,y2,x3,y3);
    J = R.ideal([x1^2+y1^2-1, x2^2+y2^2-1, x3^2+y3^2-1]);
    S = R.quotient(J);
    f = S.cover(); # f: R -> S
    M3R = MatrixSpace(R,3,3);
    M3S = MatrixSpace(S,3,3);

```

```

M3f = M3R.hom(f,M3S); # M3f: M3R -> M3S
A,B,C,D,X0 = M3f(A), M3f(B), M3f(C), M3f(D), M3f(X0);
for M in [A,B,C,D,X0]:
    assert(M*transpose(M)==I);
    assert(M.det()==1);
relations = [ A**2, (A*B)**3, B**3, (B*C)**2, C**2,
              (C*D)**5, D**2, X0*A*transpose(X0)*transpose(D) ];
for r in relations:
    assert(r==I);
print("The construction defines a representation of Gamma0.");

def find_universal_representations():
    R.<x1,y1,x2,y2,x3,y3> = QQbar[]
    A,B,C,D,X0 = rep(x1,y1,x2,y2,x3,y3);
    equations = [
        x1^2+y1^2-1,
        x2^2+y2^2-1,
        x3^2+y3^2-1
    ] + [
        M[i][j] for i in range(3)
            for j in range(3)
            for M in [ X0-I, (B*A*C)**3-I ]
    ];
    J = R.ideal(equations);
    dim_Z = J.dimension()
    assert(dim_Z==0)
    print("The variety of universal representations has dimension "
          + str(dim_Z));
    Z = J.variety();
    assert(len(Z)==1);
    print("The number of universal representations is "+str(len(Z)));
    z_u = Z[0];
    print("The unique universal representation is given by:");
    for v in [x1,y1,x2,y2,x3,y3]:
        print(v, z_u[v].radical_expression(), z_u[v], z_u[v].minpoly());
    return z_u;

```

REFERENCES

- [Ade03] Alejandro Adem. Finite group actions on acyclic 2-complexes. *Astérisque*, (290):Exp. No. 894, vii, 1–17, 2003. Séminaire Bourbaki. Vol. 2001/2002.
- [AS93] Michael Aschbacher and Yoav Segev. A fixed point theorem for groups acting on finite 2-dimensional acyclic simplicial complexes. *Proc. London Math. Soc. (3)*, 67(2):329–354, 1993.
- [Bro84] Kenneth S. Brown. Presentations for groups acting on simply-connected complexes. *J. Pure Appl. Algebra*, 32(1):1–10, 1984.
- [Bro06] Ronald Brown. *Topology and groupoids*. BookSurge, LLC, Charleston, SC, 2006.

- [CD92] Carles Casacuberta and Warren Dicks. On finite groups acting on acyclic complexes of dimension two. *Publicacions Matemàtiques*, pages 463–466, 1992.
- [Cor92] Jon M. Corson. Complexes of groups. *Proc. London Math. Soc.* (3), 65(1):199–224, 1992.
- [Cor01] Jon M. Corson. On finite groups acting on contractible complexes of dimension two. *Geom. Dedicata*, 87(1-3):161–166, 2001.
- [FR59] Edwin E. Floyd and Roger W. Richardson. An action of a finite group on an n -cell without stationary points. *Bull. Amer. Math. Soc.*, 65:73–76, 1959.
- [GAP19] The GAP Group. *GAP – Groups, Algorithms, and Programming, Version 4.10.1*, 2019.
- [GR62] Murray Gerstenhaber and Oscar S. Rothaus. The solution of sets of equations in groups. *Proc. Nat. Acad. Sci. U.S.A.*, 48:1531–1533, 1962.
- [Har15] Jens Harlander. On the relation gap and relation lifting problem. In *Groups St Andrews 2013*, volume 422 of *London Math. Soc. Lecture Note Ser.*, pages 278–285. Cambridge Univ. Press, Cambridge, 2015.
- [Har18] Jens Harlander. The relation gap problem. In *Advances in two-dimensional homotopy and combinatorial group theory*, volume 446 of *London Math. Soc. Lecture Note Ser.*, pages 128–148. Cambridge Univ. Press, Cambridge, 2018.
- [How81] James Howie. On pairs of 2-complexes and systems of equations over groups. *J. Reine Angew. Math.*, 324:165–174, 1981.
- [How02] James Howie. A proof of the Scott-Wiegold conjecture on free products of cyclic groups. *J. Pure Appl. Algebra*, 173(2):167–176, 2002.
- [KLV01] Sava Krstić, Martin Lustig, and Karen Vogtmann. An equivariant Whitehead algorithm and conjugacy for roots of Dehn twist automorphisms. *Proc. Edinb. Math. Soc.* (2), 44(1):117–141, 2001.
- [KS79] Robion C. Kirby and Martin G. Scharlemann. Eight faces of the Poincaré homology 3-sphere. *Geometric topology (Proc. Georgia Topology Conf., Athens, Ga., 1977)*, pages 113–146, 1979.
- [Oli75] Robert Oliver. Fixed-point sets of group actions on finite acyclic complexes. *Comment. Math. Helv.*, 50:155–177, 1975.
- [OS02] Bob Oliver and Yoav Segev. Fixed point free actions on \mathbf{Z} -acyclic 2-complexes. *Acta Math.*, 189(2):203–285, 2002.
- [Sag19] The Sage Developers. *SageMath, the Sage Mathematics Software System (Version 8.9)*, 2019. <https://www.sagemath.org>.
- [SC19] Iván Sadofski Costa. *Fixed points of maps and actions on 2-complexes*. PhD thesis, Universidad de Buenos Aires, 2019. https://bibliotecadigital.exactas.uba.ar/collection/tesis/document/tesis_n6591_SadofskiCosta.
- [Seg93] Yoav Segev. Group actions on finite acyclic simplicial complexes. *Israel J. Math.*, 82(1-3):381–394, 1993.
- [Seg94] Yoav Segev. Some remarks on finite 1-acyclic and collapsible complexes. *J. Combin. Theory Ser. A*, 65(1):137–150, 1994.
- [Ser80] Jean-Pierre Serre. *Trees*. Springer-Verlag, Berlin-New York, 1980. Translated from the French by John Stillwell.
- [tD87] Tammo tom Dieck. *Transformation groups*, volume 8 of *De Gruyter Studies in Mathematics*. Walter de Gruyter & Co., Berlin, 1987.

DEPARTAMENTO DE MATEMÁTICA - IMAS, FCEyN, UNIVERSIDAD DE BUENOS AIRES. BUENOS AIRES, ARGENTINA.

Email address: `isadofski@dm.uba.ar`