

AN INFINITE PRESENTATION FOR THE TWIST SUBGROUP OF THE MAPPING CLASS GROUP OF A COMPACT NON-ORIENTABLE SURFACE

*RYOMA KOBAYASHI AND GENKI OMORI

ABSTRACT. A finite presentation for the subgroup of the mapping class group of a compact non-orientable surface generated by all Dehn twists was given by Stukow [26]. In this paper, we give an infinite presentation for this group, mainly using the presentation given by Stukow [26] and Birman exact sequences on mapping class groups of non-orientable surfaces.

1. INTRODUCTION

1.1. Background.

For $g \geq 1$ and $n \geq 0$, let $N_{g,n}$ denote a compact non-orientable surface of genus g with n boundary components, that is, $N_{g,n}$ is a surface obtained by removing n open disks from a connected sum of g real projective planes. For $g \geq 0$ and $n \geq 0$, let $\Sigma_{g,n}$ denote a compact orientable surface of genus g with n boundary components, that is, $\Sigma_{g,n}$ is a surface obtained by removing n open disks from a connected sum of g tori. As shown in Figure 1, we can regard $N_{g,n}$ as a surface obtained by attaching $g - 2h$ Möbius bands to $g - 2h$ boundary components of $\Sigma_{h,n+g-2h}$, for $0 \leq h < \frac{g}{2}$. We call these attached Möbius bands *crosscaps*.




FIGURE 1. A model of a non-orientable surface $N_{g,n}$.

The mapping class group of $N_{g,n}$, denoted by $\mathcal{M}(N_{g,n})$, is the group consisting of isotopy classes of all diffeomorphisms of $N_{g,n}$ which fix the boundary pointwise. The mapping class group of $\Sigma_{g,n}$, denoted by $\mathcal{M}(\Sigma_{g,n})$, is the group consisting of isotopy classes of all *orientation-preserving* diffeomorphisms of $\Sigma_{g,n}$ which fix the boundary pointwise. It is well known that $\mathcal{M}(\Sigma_{g,n})$ is generated by only *Dehn twists* (see [3, 17, 4]). On the other hand, $\mathcal{M}(N_{g,n})$ can not be generated by only

2010 *Mathematics Subject Classification.* Primary 20F05, Secondary 57M07.

Key words and phrases. mapping class group, twist subgroup, presentation.

The first author was supported by JSPS KAKENHI Grant Number JP19K14542 and 22K13920. The second author was supported by JSPS KAKENHI Grant Number JP19K23409 and JP21K13794.

Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

Dehn twists. As generators of $\mathcal{M}(N_{g,n})$, other than Dehn twists, *crosscap slides* or *crosscap transpositions* are needed (see [16, 18]). Let us consider the subgroup of $\mathcal{M}(N_{g,n})$ generated by all Dehn twists, denoted by $\mathcal{T}(N_{g,n})$. We call $\mathcal{T}(N_{g,n})$ the *twist subgroup* of $\mathcal{M}(N_{g,n})$.

We now explain about the history of studies on presentations for $\mathcal{M}(\Sigma_{g,n})$, $\mathcal{M}(N_{g,n})$ and $\mathcal{T}(N_{g,n})$. Finite presentations for $\mathcal{M}(\Sigma_{g,n})$ were given by Hatcher-Thurston [10] and Harer [9], and subsequently simplified by Wajnryb [28] and Matsumoto [20] for $n \leq 1$. Gervais [8] and Labruère-Paris [15] gave finite presentations of $\mathcal{M}(\Sigma_{g,n})$ for $n \geq 2$. Gervais [7] gave an infinite presentation for $\mathcal{M}(\Sigma_{g,n})$ by using the presentation for $\mathcal{M}(\Sigma_{g,n})$ given in [9, 28], and then Luo [19] simplified its presentation. Finite presentations for $\mathcal{M}(N_{2,0})$, $\mathcal{M}(N_{2,1})$, $\mathcal{M}(N_{3,0})$ and $\mathcal{M}(N_{4,0})$ were given by [16], [23], [2] and [27] respectively. Note that $\mathcal{M}(N_{1,0})$ and $\mathcal{M}(N_{1,1})$ are trivial (see [5]). Paris-Szepietowski [22] gave a finite presentation of $\mathcal{M}(N_{g,n})$ for $g+n > 3$ with $n \leq 1$. Stukow [25] gave another finite presentation of $\mathcal{M}(N_{g,n})$ for $g+n > 3$ with $n \leq 1$, applying Tietze transformations for the presentation of $\mathcal{M}(N_{g,n})$ given in [22]. The second author [21] gave an infinite presentation of $\mathcal{M}(N_{g,n})$ for $g \geq 1$ and $n \leq 1$, using the presentation of $\mathcal{M}(N_{g,n})$ given in [25], and then, following this work, the authors [12] gave an infinite presentation of $\mathcal{M}(N_{g,n})$ for $g \geq 1$ and $n \geq 2$. It is known that $\mathcal{T}(N_{g,n})$ is the index 2 subgroup of $\mathcal{M}(N_{g,n})$ (see [18]). Stukow [26] gave a finite presentation of $\mathcal{T}(N_{g,n})$ for $g+n > 3$ with $n \leq 1$, applying the Reidemeister-Schreier method for the presentation of $\mathcal{M}(N_{g,n})$ given in [25] (see Theorems 2.5 and 2.6).

In this paper, we give an infinite presentation of $\mathcal{T}(N_{g,n})$ for $g \geq 1$ and $n \geq 0$ (see Theorem 1.1), mainly using the presentation of $\mathcal{T}(N_{g,n})$ given in [26] and Birman exact sequences on mapping class groups of non-orientable surfaces.

Through this paper, the product gf of mapping classes f and g means that we apply f first and then g . Moreover we do not distinguish a loop from its isotopy class.

1.2. Main result.

For a simple closed curve c of $N_{g,n}$, a regular neighborhood of c is either an annulus or a Möbius band. We call c a *two sided* or a *one sided* simple closed curve respectively. For a two sided simple closed curve c , we can take two orientations $+_c$ and $-_c$ of a regular neighborhood of c . The *right handed Dehn twist* $t_{c;\theta}$ about a two sided simple closed curve c with respect to $\theta \in \{+_c, -_c\}$ is the isotopy class of the map described as shown in Figure 2. $t_{c;\theta}$ does not depend on a choice of a representative curve of the isotopy class of c and its regular neighborhood. We remark that although the Dehn twist was defined for an *oriented* simple closed curve in [21, 12], in this paper we do not consider an orientation of a simple closed curve in the definition of the Dehn twist. We write $t_{c;\theta} = t_c$ if the orientation θ is given explicitly. That is, the direction of the twist is indicated by an arrow written beside c as shown in Figure 2.




FIGURE 2. The Dehn twist $t_{c;\theta} = t_c$ about c with respect to $\theta \in \{+\alpha, -\alpha\}$.

We denote by $f_*(\theta)$ the orientation of a regular neighborhood of $f(c)$ induced from $\theta \in \{+c, -c\}$, for a two sided simple closed curve c of $N_{g,n}$ and $f \in \mathcal{M}(N_{g,n})$. Let $c_1, \dots, c_k, c_0, c'_0$ and $d_1, \dots, d_7$ be simple closed curves with arrows as shown in Figure 3. $\mathcal{M}(N_{g,n})$ admits following relations.

- (1) $t_{c;\theta} = 1$ if c bounds a disk or a Möbius band.
- (2) $t_{c;-c}^{-1} = t_{c;+c}$.
- (3) $ft_{c;\theta}f^{-1} = t_{f(c);f_*(\theta)}$ for $f \in \mathcal{M}(N_{g,n})$.
- (4) $(t_{c_1}t_{c_2} \dots t_{c_k})^{k+1} = t_{c_0}t_{c'_0}$ if k is odd,
 $(t_{c_1}t_{c_2} \dots t_{c_k})^{2k+2} = t_{c_0}$ if k is even.
- (5) $t_{d_1}t_{d_2}t_{d_3} = t_{d_4}t_{d_5}t_{d_6}t_{d_7}$.

We can check the relations (1) and (2) easily. The relations (3), (4) and (5) are called a *conjugation relation*, a *k-chain relation*, a *lantern relation* respectively. These are famous relations on mapping class groups. In the relation (3), if $f = t_{c';\theta'}$, $|c \cap c'| = 0$ or 1, and the orientations θ and θ' are compatible, then the relation can be rewritten as a *commutativity relation* $t_{c;\theta}t_{c';\theta'} = t_{c';\theta'}t_{c;\theta}$ or a *braid relation* $t_{c;\theta}t_{c';\theta'}t_{c;\theta} = t_{c';\theta'}t_{c;\theta}t_{c';\theta'}$ respectively.




FIGURE 3.

Our main result is as follows.

Theorem 1.1. *For $g \geq 1$ and $n \geq 0$, $\mathcal{T}(N_{g,n})$ admits a presentation with a generating set*

$$X = \left\{ t_{c;\theta} \mid \begin{array}{l} c \text{ is a two sided simple closed curve of } N_{g,n} \text{ and} \\ \theta \text{ is an orientation of a regular neighborhood of } c. \end{array} \right\}.$$

The defining relations are

- (1) $t_{c;\theta} = 1$ if c bounds a disk or a Möbius band,
- (2) $t_{c;-c}^{-1} = t_{c;+c}$,
- (3) all the conjugation relations $ft_{c;\theta}f^{-1} = t_{f(c);f_*(\theta)}$ for $f \in X$,
- (4) all the 2-chain relations,
- (5) all the lantern relations,

Remark 1.2. If a two sided simple closed curve c is a non-separating curve of $N_{2,0}$, then we can check that $t_{c;+c}$ and $t_{c;-c}$ are the same elements in X of Theorem 1.1. Hence by the relation (2) of Theorem 1.1, we have the relation $t_{c;\theta}^2 = 1$.

Remark 1.3. Applying Theorem in [19] to a regular neighborhood of $\cup_{i=1}^k c_i$ in Figure 3, it follows that any chain relation is obtained from the relations (1), (3), (4) and (5) of Theorem 1.1. We assign the label (4)' to all the chain relations.

Here is the outline of this paper. In Section 2, we explain what we need to prove our main result. In Section 3, we prove that the infinitely presented group with the presentation in Theorem 1.1 is isomorphic to $\mathcal{T}(N_{g,n})$ with a presentation which is already known, for $n \leq 1$. In Section 4, we complete the proof of Theorem 1.1, by induction on n .

2. PRELIMINARIES

In this section, we explain what we need to prove our main result Theorem 1.1. In Section 2.1, we define the *capping map*, the *point pushing map*, the *forgetful map* and the *crosscap pushing map* on mapping class groups for non-orientable surfaces. In Section 2.2, we introduce a finite presentation of $\mathcal{T}(N_{g,n})$ for $g \geq 1$ and $n \leq 1$.

2.1. Homomorphisms on mapping class groups of non-orientable surfaces.

We first define the *capping map*, the *point pushing map* and the *forgetful map* on mapping class groups of non-orientable surfaces. Take a point $*$ in the interior of $N_{g,n-1}$. Let $\mathcal{M}(N_{g,n-1}, *)$ denote the group consisting of isotopy classes of all diffeomorphisms of $N_{g,n-1}$ which fix $*$ and the boundary pointwise. We can regard $N_{g,n}$ as a complement of an open disk neighborhood of $*$ in $N_{g,n-1}$. The natural embedding $N_{g,n} \hookrightarrow N_{g,n-1}$ induces the homomorphism

$$\mathcal{C} : \mathcal{M}(N_{g,n}) \rightarrow \mathcal{M}(N_{g,n-1}, *)$$

which is called the *capping map*. The *point pushing map*

$$\mathcal{P}_* : \pi_1(N_{g,n-1}, *) \rightarrow \mathcal{M}(N_{g,n-1}, *)$$

is defined as follows. For any loop $x \in \pi_1(N_{g,n-1}, *)$, $\mathcal{P}_*(x)$ is described by pushing $*$ once along x . The *forgetful homomorphism*

$$\mathcal{F} : \mathcal{M}(N_{g,n-1}, *) \rightarrow \mathcal{M}(N_{g,n-1})$$

is defined naturally, that is, $\mathcal{F}(f)$ does not necessarily fix $*$.


Next, we consider exact sequences on mapping class groups of non-orientable surfaces. Let $\mathcal{M}^+(N_{g,n-1}, *)$ denote the subgroup of $\mathcal{M}(N_{g,n-1}, *)$ consisting of elements which preserve a local orientation of $*$, and $\pi_1^+(N_{g,n-1}, *)$ the subgroup of $\pi_1(N_{g,n-1}, *)$ generated by two sided simple loops. We have the exact sequences

$$(1) \quad 1 \rightarrow \mathbb{Z} \rightarrow \mathcal{M}(N_{g,n}) \xrightarrow{\mathcal{C}} \mathcal{M}^+(N_{g,n-1}, *) \rightarrow 1,$$

$$(2) \quad \pi_1^+(N_{g,n-1}, *) \xrightarrow{\mathcal{P}_*} \mathcal{M}^+(N_{g,n-1}, *) \xrightarrow{\mathcal{F}} \mathcal{M}(N_{g,n-1}) \rightarrow 1.$$

The second sequence is called the *Birman exact sequence*, introduced by Birman [1]. Note that $\mathcal{P}_*$ is injective except for the case $(g, n) = (2, 1)$. For details see [1, 14, 24, 6, 22].

Remark 2.1. For a simple loop $\alpha \in \pi_1^+(N_{g,n-1}, *)$, we take an orientation θ of a regular neighborhood of α . Let γ_1 and γ_2 be the right side boundary curve and the left side boundary curve of the regular neighborhood of α for the direction of α , respectively, where the left and right sides are determined by θ . Then we have $\mathcal{P}_*(\alpha) = t_{\gamma_1; \theta_1} t_{\gamma_2; \theta_2}^{-1}$, where θ_1 and θ_2 are the orientations compatible with θ (see Figure 4).

FIGURE 4. The boundary curves γ_1 and γ_2 of a regular neighborhood of α .

Remark 2.2. Let α , β and $\gamma \in \pi_1^+(N_{g,n-1}, *)$ be simple loops with $\alpha\beta = \gamma$. If α and β intersect transversally at only $*$ as shown in Figure 5, then the relation $\mathcal{P}_*(\gamma) = \mathcal{P}_*(\beta)\mathcal{P}_*(\alpha)$ is obtained from the relation (3) of Theorem 1.1 (see Lemma 5.4 in [12]). If α and β intersect tangentially at only $*$ as shown in Figure 5, then the relation $\mathcal{P}_*(\gamma) = \mathcal{P}_*(\beta)\mathcal{P}_*(\alpha)$ is obtained from the relation (5) of Theorem 1.1 and a relation $t_{c;\theta} = 1$, where c is a simple closed curve bounding a disk neighborhood of $*$ (see Lemma 5.2 in [12]). We call this relation the *extended lantern relation*.




FIGURE 5.

Finally, we define the *crosscap pushing map*. Let S be a surface obtained by shrinking some crosscap of $N_{g,n}$ into a point. We call this operation the *blowdown* with respect to the crosscap. Note that S is diffeomorphic to either $N_{g-1,n}$ or $\Sigma_{\frac{g-1}{2},n}$ (see Figure 6). Let $*$ be the shrunk point of S . Conversely, we can obtain $N_{g,n}$ from S , and call this operation the *blowup* with respect to $*$. The *crosscap pushing map*

$$\mathcal{P}_{\otimes} : \pi_1(S, *) \rightarrow \mathcal{M}(N_{g,n})$$

is defined as follows. For $x \in \pi_1(S, *)$, let $\tilde{x}$ be an oriented loop of $N_{g,n}$ induced from x by the blowup with respect to $*$. $\mathcal{P}_{\otimes}(x)$ is described by pushing the crosscap, which is obtained by the blowup with respect to $*$, once along $\tilde{x}$ (see Figure 7).





FIGURE 6.

FIGURE 7. The crosscap pushing map $\mathcal{P}_\otimes$.

Similar to Remarks 2.1 and 2.2, we have the followings.

Remark 2.3. For a two sided simple loop $\alpha \in \pi_1(S, *)$, we take an orientation θ of a regular neighborhood of α . Let γ_1 and γ_2 be the right side boundary curve and the left side boundary curve of the regular neighborhood of α for the direction of α , respectively, where the left and right sides are determined by θ . We also denote by $\tilde{\gamma}_1$ and $\tilde{\gamma}_2$ loops of $N_{g,n}$ induced from γ_1 and γ_2 by the blowup with respect to $*$, respectively. Then we have $\mathcal{P}_\otimes(\alpha) = t_{\tilde{\gamma}_1; \tilde{\theta}_1} t_{\tilde{\gamma}_2; \tilde{\theta}_2}^{-1}$, where $\tilde{\theta}_1$ and $\tilde{\theta}_2$ are the orientations of regular neighborhoods of $\tilde{\gamma}_1$ and $\tilde{\gamma}_2$ induced by the blowup with respect to $*$ from the orientations of regular neighborhoods of γ_1 and γ_2 which are compatible with θ , respectively.

Remark 2.4. Let α, β and $\gamma \in \pi_1(S, *)$ be two sided simple loops with $\alpha\beta = \gamma$. If α and β intersect transversally at only $*$ as shown in Figure 5, then the relation $\mathcal{P}_\otimes(\gamma) = \mathcal{P}_\otimes(\beta)\mathcal{P}_\otimes(\alpha)$ is obtained from the relations (3) of Theorem 1.1 (see Lemma 5.4 in [12]). If α and β intersect tangentially at only $*$ as shown in Figure 5, then the relation $\mathcal{P}_\otimes(\gamma) = \mathcal{P}_\otimes(\beta)\mathcal{P}_\otimes(\alpha)$ is obtained from the relations (1) and (5) of Theorem 1.1 (see Lemma 5.2 in [12]).

2.2. A finite presentation for $\mathcal{T}(N_{g,n})$.

For $g \geq 2$ and $n \geq 0$ we have the short exact sequence

$$(3) \quad 1 \rightarrow \mathcal{T}(N_{g,n}) \rightarrow \mathcal{M}(N_{g,n}) \rightarrow \mathbb{Z}/2\mathbb{Z} \rightarrow 1$$

(see [16, 18]). Using the Reidemeister Schreier method for a presentation of $\mathcal{M}(N_{g,n})$, we can obtain a presentation of $\mathcal{T}(N_{g,n})$.

For $n \leq 1$, let $\alpha_1, \dots, \alpha_{g-1}, \beta, \gamma, \varepsilon$ and ζ be simple closed curves of $N_{g,n}$ with arrows as shown in Figure 8, and if g is even, $\beta_0, \dots, \beta_{\frac{g-2}{2}}, \bar{\beta}_{\frac{g-6}{2}}, \bar{\beta}_{\frac{g-4}{2}}$ and $\bar{\beta}_{\frac{g-2}{2}}$ simple closed curves of $N_{g,n}$ with arrows as shown in Figure 8. Let α be an oriented simple loop of $N_{g-1,n}$ based at $*$ induced from α_1 by the blowdown with respect to the first crosscap, as shown in Figure 8. For simplicity, we denote $t_{\alpha_i} = a_i, t_\beta = b, t_\gamma = c, t_\varepsilon = e, t_\zeta = f, t_{\beta_i} = b_i, t_{\bar{\beta}_i} = \bar{b}_i$ and $\mathcal{P}_\otimes(\alpha) = y$. Note that $y^2 = t_\delta$, where δ is a simple close curve of $N_{g,n}$ with an arrow as shown in Figure 8. For $g \geq 1$ and $n \leq 1$, $\mathcal{T}(N_{g,n})$ admits a finite presentation given in Theorems 2.5, 2.6 or Lemma 2.7.

Theorem 2.5 (Theorem 3.1 in [26]). *If $g \geq 3$ is odd or $g = 4$, then $\mathcal{T}(N_{g,1})$ admits a presentation with generators $a_1, \dots, a_{g-1}, e, f, y^2$ and b, c for $g \geq 4$. The defining relations are*

- (A1) $a_i a_j = a_j a_i$ for $g \geq 4, |i - j| > 1$,
- (A2) $a_i a_{i+1} a_i = a_{i+1} a_i a_{i+1}$ for $i = 1, \dots, g - 2$,
- (A3) $a_i b = b a_i$ for $g \geq 4, i \neq 4$,
- (A4) $b a_4 b = a_4 b a_4$ for $g \geq 5$,




FIGURE 8.

- (A5) $(a_2a_3a_4b)^{10} = (a_1a_2a_3a_4b)^6$ for $g \geq 5$,
 (A6) $(a_2a_3a_4a_5a_6b)^{12} = (a_1a_2a_3a_4a_5a_6b)^9$ for $g \geq 7$,
 (A1₁) $ea_j = a_je$ for $g \geq 5$, $j \geq 4$,
 (A1₂) $fa_j = a_jf$ for $g \geq 5$, $j \geq 4$,
 (A2₁) $a_1ea_1 = ea_1e$,
 (A2₂) $a_3^{-1}ea_3^{-1} = ea_3^{-1}e$ for $g \geq 4$,
 (A2₃) $a_1fa_1 = fa_1f$,
 (A3₁) $a_1c = ca_1$ for $g = 4, 5$,
 (A3₂) $ec = ce$ for $g = 4, 5$,
 (A4) $ca_4c = a_4ca_4$ for $g = 5, 6$,
 (A5) $(e^{-1}a_3a_4c)^{10} = (a_1^{-1}e^{-1}a_3a_4c)^6$ for $g = 5, 6$,
 (A6) $(e^{-1}a_3a_4a_5a_6c)^{12} = (a_1^{-1}e^{-1}a_3a_4a_5a_6c)^9$ for $g = 7, 8$,
 (B1) $(a_2a_3a_1a_2ea_1a_3^{-1}e)(a_2a_3a_1a_2fa_1a_3^{-1}f) = 1$ for $g \geq 4$,
 (B2₁) $y^2 = a_2a_1ea_1a_2a_1a_2a_1a_2fa_1a_2$,
 (B2₂) $(a_2a_1ea_1a_2a_1a_2a_1a_2fa_1a_2)(a_2a_1fa_1a_2a_1a_2a_1a_2ea_1a_2) = 1$,
 (B3) $y^2a_3 = a_3y^2$ for $g \geq 4$,
 (B4₁) $ea_2 = a_2e$,
 (B4₂) $fa_2 = a_2f$,
 (B6₁) $bc = (a_1a_2a_3f^{-1}a_3^{-1}a_2^{-1}a_1^{-1})(a_2^{-1}a_3^{-1}e^{-1}a_3a_2)$ for $g \geq 4$,
 (B6₂) $c(y^2by^{-2}) = (a_1^{-1}e^{-1}a_3a_2a_3^{-1}ea_1)(ea_3^{-1}y^2a_2y^{-2}a_3e^{-1})$ for $g = 4, 5$,
 (B7₁) $(a_4a_5a_3a_4a_2a_3a_1a_2ea_1a_3^{-1}ea_4^{-1}a_3^{-1}a_5^{-1}a_4^{-1})c$
 $= b(a_4a_5a_3a_4a_2a_3a_1a_2ea_1a_3^{-1}ea_4^{-1}a_3^{-1}a_5^{-1}a_4^{-1})$ for $g \geq 6$,
 (B7₂) $(a_2^{-1}a_1^{-1}a_3^{-1}a_2^{-1}a_4^{-1}a_3^{-1}a_5^{-1}a_4^{-1})b(a_4a_5a_3a_4a_2a_3a_1a_2)y^2$
 $= y^2(a_2^{-1}a_1^{-1}a_3^{-1}a_2^{-1}a_4^{-1}a_3^{-1}a_5^{-1}a_4^{-1})b(a_4a_5a_3a_4a_2a_3a_1a_2)$ for $g \geq 6$,
 (B8₁) $(a_1ea_3^{-1}a_4^{-1}ca_4a_3e^{-1}a_1^{-1})(a_1^{-1}a_2^{-1}a_3^{-1}a_4^{-1}b^{-1}a_4a_3a_2a_1)$
 $= a_4^{-1}(a_3^{-1}a_2^{-1}e^{-1}a_3a_4a_3^{-1}ea_2a_3)a_2^{-1}e^{-1}$ for $g \geq 5$,
 (B8₂) $(a_1^{-1}a_2^{-1}a_3^{-1}a_4^{-1}ba_4a_3a_2a_1)(a_1fa_3^{-1}a_4^{-1}y^{-2}c^{-1}y^2a_4a_3f^{-1}a_1^{-1})$
 $= a_4^{-1}(a_3^{-1}fa_2a_3a_4a_3^{-1}a_2^{-1}f^{-1}a_3)fa_2$ for $g = 5, 6$.

If $g \geq 6$ is even, then $\mathcal{T}(N_{g,1})$ admits a presentation with generators $a_1, \dots, a_{g-1}$, e, f, y^2, b, c and additionally $b_0, \dots, b_{\frac{g-2}{2}}, \bar{b}_{\frac{g-6}{2}}, \bar{b}_{\frac{g-4}{2}}, \bar{b}_{\frac{g-2}{2}}$. The defining relations are relations (A1)-(A6), (A1₁)-(A6), (B1)-(B8₂) and additionally

- (A7) $b_0 = a_1, b_1 = b$,
 (A8) $b_{i+1} = (b_{i-1}a_2ia_{2i+1}a_{2i+2}a_{2i+3}b_i)^5(b_{i-1}a_2ia_{2i+1}a_{2i+2}a_{2i+3})^{-6}$ for $1 \leq i \leq \frac{g-4}{2}$,

- (A9a) $b_2b = bb_2$ for $g = 6$,
 (A9b) $b_{\frac{g-2}{2}}a_{g-5} = a_{g-5}b_{\frac{g-2}{2}}$ for $g \geq 8$,
 (A7a) $\bar{b}_0 = a_1^{-1}$, $\bar{b}_1 = c$ for $g = 6$,
 (A7b) $\bar{b}_1 = c$ for $g = 8$,
 (A7c) $\bar{b}_i = z_{g-1}b_iz_{g-1}^{-1}$ where $i = \frac{g-6}{2}, \frac{g-4}{2}, i \geq 2$ and $z_{g-1} = (a_{g-1}a_ga_{g-2}a_{g-1} \cdots a_3a_4e^{-1}a_3a_1^{-1}e^{-1})(a_2^{-1}a_1^{-1} \cdots a_{g-1}^{-1}a_{g-2}^{-1}a_g^{-1}a_{g-1}^{-1})$,
 (A8a) $\bar{b}_2 = (\bar{b}_0e^{-1}a_3a_4a_5\bar{b}_1)^5(\bar{b}_0e^{-1}a_3a_4a_5)^{-6}$ for $g = 6$,
 (A8b) $\bar{b}_{\frac{g-2}{2}} = (\bar{b}_{\frac{g-6}{2}}a_{g-4}a_{g-3}a_{g-2}a_{g-1}\bar{b}_{\frac{g-4}{2}})^5(\bar{b}_{\frac{g-6}{2}}a_{g-4}a_{g-3}a_{g-2}a_{g-1})^{-6}$ for $g \geq 8$,
 (A9a) $\bar{b}_2c = c\bar{b}_2$ for $g = 6$,
 (A9b) $\bar{b}_{\frac{g-2}{2}}a_{g-5} = a_{g-5}\bar{b}_{\frac{g-2}{2}}$ for $g \geq 8$.

Theorem 2.6 (Theorem 3.2 in [26]). *If $g \geq 5$ is odd, then the group $\mathcal{T}(N_{g,0})$ is isomorphic to the quotient of the group $\mathcal{T}(N_{g,1})$ with the presentation given in Theorem 2.5 obtained by adding a generator ϱ and relations*

- (C1a) $(a_1a_2 \cdots a_{g-1})^g = \varrho$,
 (C1a) $(a_1^{-1}e^{-1}a_3 \cdots a_{g-1})^g = y^2\varrho$,
 (C2) $a_i\varrho = \varrho a_i$ for $1 \leq i \leq g-1$,
 (C2) $\varrho e = f\varrho$,
 (C5₁) $\varrho y^2 = y^{-2}\varrho$,
 (C3) $\varrho^2 = 1$,
 (C4a) $(a_2a_3 \cdots a_{g-1}e^{-1}a_3 \cdots a_{g-1})^{\frac{g-1}{2}} = 1$.

Moreover, relations $(\overline{A1_2})$, $(\overline{B2_2})$, $(\overline{B4_2})$ are superfluous.

If $g \geq 4$ is even, then the group $\mathcal{T}(N_{g,0})$ is isomorphic to the quotient of the group $\mathcal{T}(N_{g,1})$ with the presentation given in Theorem 2.5 obtained by adding a generator $\bar{\varrho}$ and relations

- (C1b) $(a_1a_2 \cdots a_{g-1})^g = 1$,
 (C2₁) $\bar{\varrho}a_1 = a_1^{-1}\bar{\varrho}$,
 (C2₂) $\bar{\varrho}a_i = a_i\bar{\varrho}$ for $3 \leq i \leq g-1$,
 (C2₃) $\bar{\varrho}a_2 = e^{-1}\bar{\varrho}$,
 (C5₂) $\bar{\varrho}y^2 = y^{-2}\bar{\varrho}$,
 (C3) $\bar{\varrho}^2 = 1$,
 (C4) $(\bar{\varrho}a_2a_3 \cdots a_{g-1})^{g-1} = 1$.

Moreover, relations $(\overline{A1_1})$, $(\overline{A2_1})$, $(\overline{A2_2})$ are superfluous.

Lemma 2.7. (1) $\mathcal{T}(N_{1,0})$ and $\mathcal{T}(N_{1,1})$ are trivial.

(2) $\mathcal{T}(N_{2,0}) = \langle a_1 \mid a_1^2 \rangle$.

(3) $\mathcal{T}(N_{2,1}) = \langle a_1, y^2 \mid a_1y^2a_1^{-1}y^{-2} \rangle$.

(4) $\mathcal{T}(N_{3,0}) = \langle a_1, a_2 \mid a_1a_2a_1a_2^{-1}a_1^{-1}a_2^{-1}, (a_1a_2)^6 \rangle$.

Proof. Note that $\mathbb{Z}/2\mathbb{Z}$ is generated by the image of y , in the sequence (3) at the beginning of Section 2.2.

(1) $\mathcal{M}(N_{1,0})$ and $\mathcal{M}(N_{1,1})$ are trivial (see Theorem 3.4 in [5]). Thus $\mathcal{T}(N_{1,0})$ and $\mathcal{T}(N_{1,1})$ are also trivial.

(2) We have the presentation

$$\mathcal{M}(N_{2,0}) = \langle a_1, y \mid a_1^2, y^2, (a_1y)^2 \rangle$$

(see Lemma 5 in [16]). Using the Reidemeister Schreier method for this presentation, we obtain the presentation

$$\begin{aligned}\mathcal{T}(N_{2,0}) &= \langle a_1, y^2 \mid a_1^2, y^2, a_1^{-1}y^2a_1 \rangle \\ &= \langle a_1 \mid a_1^2 \rangle.\end{aligned}$$

(3) We have the presentation

$$\mathcal{M}(N_{2,1}) = \langle a_1, y \mid ya_1y^{-1}a_1 \rangle$$

(see Theorem A.7 in [23]). Using the Reidemeister Schreier method for this presentation, we obtain the presentation

$$\mathcal{T}(N_{2,1}) = \langle a_1, y^2 \mid a_1y^2a_1^{-1}y^{-2} \rangle.$$

(4) We have the presentation

$$\mathcal{M}(N_{3,0}) = \langle a_1, a_2, y \mid a_1a_2a_1a_2^{-1}a_1^{-1}a_2^{-1}, (a_1a_2)^6, y^2, (a_1y)^2, (a_2y)^2 \rangle$$

(see Theorem 3 in [2]). Using the Reidemeister Schreier method for this presentation, we obtain the presentation

$$\begin{aligned}\mathcal{T}(N_{3,0}) &= \langle a_1, a_2, y^2 \mid a_1a_2a_1a_2^{-1}a_1^{-1}a_2^{-1}, (a_1a_2)^6, y^2, a_1^{-1}y^2a_1, a_2^{-1}y^2a_2 \rangle \\ &= \langle a_1, a_2 \mid a_1a_2a_1a_2^{-1}a_1^{-1}a_2^{-1}, (a_1a_2)^6 \rangle.\end{aligned}$$

□

3. PROOF OF THEOREM 1.1 FOR $g \geq 1$ AND $n \leq 1$

In this section, we prove Theorem 1.1 for $g \geq 1$ and $n \leq 1$, using the presentation of $\mathcal{T}(N_{g,n})$ given in Theorems 2.5, 2.6 and Lemma 2.7.

Let $\langle X \mid Y \rangle$ be the infinitely presented group with the presentation given in Theorem 1.1, and $\langle X_0 \mid Y_0 \rangle$ the presentation for $\mathcal{T}(N_{g,n})$ given in Theorems 2.5, 2.6 or Lemma 2.7. Denote by $F(X_0)$ the free group freely generated by X_0 . Let $p : F(X_0) \rightarrow \langle X_0 \mid Y_0 \rangle$ be the natural projection and $\eta : F(X_0) \rightarrow \langle X \mid Y \rangle$ the homomorphism defined as $\eta(x) = x$ for any $x \in X_0$. We consider a correspondence

$$\psi : \langle X_0 \mid Y_0 \rangle \rightarrow \langle X \mid Y \rangle$$

satisfying $\psi \circ p = \eta$. Showing the following proposition, we will obtain Theorem 1.1 for $g \geq 1$ and $n \leq 1$.

Proposition 3.1. *For $g \geq 1$ and $n \leq 1$, ψ is an isomorphism.*

Let $\varphi : \langle X \mid Y \rangle \rightarrow \langle X_0 \mid Y_0 \rangle = \mathcal{T}(N_{g,n})$ be the homomorphism defined as $\varphi(t_{c;\theta}) = t_{c;\theta}$ for any $t_{c;\theta} \in X$. Since $\varphi(Y) = 1$ in $\mathcal{T}(N_{g,n})$ clearly, φ is well-defined. By the definitions of ψ and φ , it is clear that $\varphi \circ \psi$ is the identity map if ψ is a homomorphism. Hence in order to prove Proposition 3.1, what we need is to show well-definedness and surjectivity of ψ . In Sections 3.1 and 3.2, we see well-definedness and surjectivity of ψ respectively.

3.1. Well-definedness of ψ .

In order to see well-definedness of ψ , it suffices to show that $\psi(r) = 1$ in $\langle X \mid Y \rangle$ for any $r \in Y_0$. More precisely, we check that any relation and relator of $\mathcal{T}(N_{g,n})$ in Theorems 2.5, 2.6 and Lemma 2.7 are obtained from the relations (1)-(5) of $\mathcal{T}(N_{g,n})$ in Theorem 1.1 and the relation (4)' assigned in Remark 1.3.

First we notice that the relations (A7), $(\overline{A7a})$ and $(\overline{A7b})$ hold since each of the pairs (b_0, a_1) , (b_1, b) , $(\bar{b}_0, a_1^{-1})$ for $g = 6$, and $(\bar{b}_1, c)$ for $g = 6, 8$ coincides as mapping classes from the definition of loops in Figure 8. We may also treat the relation (C1a) as a definition of ρ . It is easy to check that any relator of $\mathcal{T}(N_{g,n})$ in Lemma 2.7 is obtained from the relations (1)-(4) in Theorem 1.1. In particular, for the relator a^2 of $\mathcal{T}(N_{2,0})$, recall Remark 1.2. The relations (A1), (A2), (A3), (A4), $(\overline{A1_1})$, $(\overline{A1_2})$, $(\overline{A2_1})$, $(\overline{A2_2})$, $(\overline{A2_3})$, $(\overline{A3_1})$, $(\overline{A3_2})$, $(\overline{A4})$, $(\overline{B3})$, $(\overline{B4_1})$, $(\overline{B4_2})$, $(\overline{B7_1})$, $(\overline{B7_2})$, (A9a), (A9b), (A7c), (A9a), (A9b) (C2), $(\overline{C2})$, $(\overline{C5_1})$, $(\overline{C2_1})$, $(\overline{C2_2})$, $(\overline{C2_3})$ and $(\overline{C5_2})$ are obtained by repeating the relation (3) in Theorem 1.1. The relations (A5), (A6), $(\overline{A5})$, $(\overline{A6})$, (A8), $(\overline{A8a})$ and $(\overline{A8b})$ come from relations of mapping class groups of orientable surfaces (see Theorem 1.3 in [20]), and hence these relations are obtained from the relations (3), (4) and (5) in Theorem 1.1 by Theorem in [19]. The relations (C3) and (C1b) are obtained from the relations (1) in Theorem 1.1 and (4)' assigned in Remark 1.3. Thus it suffices to show that the relations $(\overline{B1})$, $(\overline{B2_1})$, $(\overline{B2_2})$, $(\overline{B6_1})$, $(\overline{B6_2})$, $(\overline{B8_1})$, $(\overline{B8_2})$, $(\overline{C1a})$, $(\overline{C4a})$, $(\overline{C3})$ and $(\overline{C4})$ are satisfied in $\langle X \mid Y \rangle$.

Recall the simple closed curves and the simple loop as shown in Figure 8.

3.1.1. On the relation $(\overline{B1})$.

The relation $(\overline{B1})$ can be rewritten as follows.

$$\begin{aligned}
 & (a_2 a_3 a_1 a_2 e a_1 a_3^{-1} e) \underbrace{(a_2 a_3 a_1 a_2 f a_1 a_3^{-1} f)}_{(\overline{B4_1}) \text{ (A1)}} = 1 \\
 \iff & \underbrace{a_2 a_3 a_1}_{(\overline{B4_2}) \text{ (A1)}} (a_2 e a_1 a_3^{-1} a_2 e a_1 a_3 a_2 f a_1 a_3^{-1} f a_2 a_3 a_1) a_1^{-1} a_3^{-1} a_2^{-1} = 1 \\
 \iff & a_1 \{ (a_2 e a_1) a_3^{-1} (a_2 e a_1) a_3 (a_2 f a_1) a_3^{-1} (a_2 f a_1) a_3 \} a_1^{-1} = 1.
 \end{aligned}$$

Similarly, the relation $(\overline{B2_1})$ can be rewritten as follows.

$$\begin{aligned}
 y^2 &= \underbrace{a_2 a_1 e a_1}_{(\overline{A2_1})} a_2 a_1 a_2 a_1 \underbrace{a_2 f a_1 a_2}_{(\overline{B4_2})} \\
 &= \underbrace{a_2 e a_1 e a_2}_{(\overline{B4_1})} a_1 a_2 a_1 f a_2 a_1 a_2 \underbrace{a_2}_{(\overline{A2})} \\
 &= \underbrace{a_2 e a_1 a_2 e a_1 a_2 a_1 f a_1 a_2 a_1}_{(\overline{A2_3})} \\
 &= \underbrace{a_2 e a_1 a_2 e a_1 a_2 f a_1 f a_2 a_1}_{(\overline{B4_2})} \\
 &= (a_2 e a_1)^2 (a_2 f a_1)^2.
 \end{aligned}$$

Let A , B , C , D and E be simple closed curves with arrows as shown in Figure 9. Repeating the relation (3), we have $t_A = a_1 (a_2 e a_1) a_3^{-1} (a_2 e a_1)^{-1} a_1^{-1}$, $t_B = a_1 (a_2 e a_1)^2 a_3 (a_2 e a_1)^{-2} a_1^{-1}$ and $t_C = a_1 (a_2 f a_1)^{-1} a_3^{-1} (a_2 f a_1) a_1^{-1}$. Assuming

that $(\overline{B2}_1)$ holds in $\langle X \mid Y \rangle$, we calculate

$$\begin{aligned} a_1 \{ (a_2 e a_1) a_3^{-1} (a_2 e a_1) a_3 (a_2 f a_1) a_3^{-1} (a_2 f a_1) a_3 \} a_1^{-1} &= t_A t_B y^2 t_C (a_1 a_3 a_1^{-1}) \\ &\stackrel{(1),(3)}{=} t_A t_B y^2 t_C a_3 t_D t_E \\ &\stackrel{(5)}{=} 1. \end{aligned}$$

Note that $B \cup \alpha_3 \cup D \cup E$ bounds $\Sigma_{0,4}$. Thus the relation $(\overline{B1})$ is satisfied in $\langle X \mid Y \rangle$, by Section 3.1.2.




FIGURE 9.

3.1.2. *On the relations $(\overline{B2}_1)$ and $(\overline{B2}_2)$.*

Let A, B, C and D be simple closed curves with arrows as shown in Figure 10. Repeating the relation (3), we have $t_A = a_2 a_1 e a_1^{-1} a_2^{-1}$ and $t_B = a_2 a_1 f^{-1} a_1^{-1} a_2^{-1}$. In addition, by the relations (1) and (4)' we have $t_C = (f a_1 a_2)^4$. Hence we calculate

$$\begin{aligned} y^2 &\stackrel{(5)}{=} t_A t_B t_C t_D a_1 a_1^{-1} \\ &\stackrel{(1)}{=} (a_2 a_1 e a_1^{-1} a_2^{-1}) (a_2 a_1 f^{-1} a_1^{-1} a_2^{-1}) (f a_1 a_2)^4 \\ &= a_2 a_1 e f^{-1} a_1^{-1} \underbrace{a_2^{-1} f a_1 a_2 f a_1 a_2 f a_1 a_2}_{(\overline{B4}_2)} \underbrace{a_1 a_2 f a_1 a_2}_{(\overline{B4}_2)} \\ &= a_2 a_1 e \underbrace{f^{-1} a_1^{-1} f a_2^{-1}}_{(\overline{A2}_3)} a_1 a_2 \underbrace{f a_1 f a_2 a_1 a_2 f a_1 a_2}_{(\overline{A2}_3)} \\ &= a_2 a_1 e a_1 \underbrace{f^{-1} a_1^{-1} a_2^{-1} a_1 a_2 a_1 f a_1 a_2 a_1 a_2 f a_1 a_2}_{(\overline{A2}), (\overline{B4}_2)} \\ &= a_2 a_1 e a_1 a_2 a_1 a_2 a_1 a_2 f a_1 a_2. \end{aligned}$$

Note that $C \cup D \cup \alpha_1$ bounds $\Sigma_{0,4}$. Thus the relation $(\overline{B2}_1)$ is satisfied in $\langle X \mid Y \rangle$.




FIGURE 10.

Moreover, similar to the transformation of the relation $(\overline{B2}_1)$ of Section 3.1.1, the relation $(\overline{B2}_2)$ can be rewritten as

$$(a_2 e a_1)^2 (a_2 f a_1)^4 (a_2 e a_1)^2 = 1.$$

By the relations (1) and (4)', we have $t_C = (a_2 f a_1)^4$ and $t_C^{-1} = (a_2 e a_1)^4$. Hence we calculate

$$\begin{aligned}
 (a_2 e a_1)^2 (a_2 f a_1)^4 (a_2 e a_1)^2 &= (a_2 e a_1)^2 t_C (a_2 e a_1)^2 \\
 &\stackrel{(3)}{=} (a_2 e a_1)^4 t_C \\
 &= t_C^{-1} t_C \\
 &= 1.
 \end{aligned}$$

Thus the relation $(\overline{B2}_2)$ is satisfied in $\langle X \mid Y \rangle$.

3.1.3. *On the relations $(\overline{B6}_1)$ and $(\overline{B6}_2)$.*

Let A, B, C, D and E be simple closed curves with arrows as shown in Figure 11. Repeating the relation (3), we have $t_A = a_1 a_2 a_3 f^{-1} a_3^{-1} a_2^{-1} a_1^{-1}$ and $t_B = a_2^{-1} a_3^{-1} e^{-1} a_3 a_2$. Then we calculate

$$\begin{aligned}
 b c y^2 &\stackrel{(5)}{=} t_C a_1 a_1^{-1} a_3 \\
 &= t_C a_3, \\
 t_A t_B y^2 &\stackrel{(5)}{=} t_C t_D t_E a_3 \\
 &\stackrel{(1)}{=} t_C a_3,
 \end{aligned}$$

and hence $bc = t_A t_B$. Note that $C \cup \alpha_1 \cup \alpha_3$ and $C \cup D \cup E \cup \alpha_3$ bound $\Sigma_{0,4}$. Thus the relation $(\overline{B6}_1)$ is satisfied in $\langle X \mid Y \rangle$.




FIGURE 11.

In addition, repeating the relation (3), we have $y^2 b y^{-2} = t_{y(\gamma)}$, $a_1^{-1} e^{-1} a_3 a_2 a_3^{-1} e a_1 = t_{y(A)}$ and $e a_3^{-1} y^2 a_2 y^{-2} a_3 e^{-1} = t_{y(B)}$. Then we calculate

$$\begin{aligned}
 t_{y(\beta)} t_{y(\gamma)} y^2 &\stackrel{(5)}{=} t_C a_1 a_1^{-1} a_3 \\
 &= t_C a_3, \\
 t_{y(A)} t_{y(B)} y^2 &\stackrel{(5)}{=} t_C t_{y(D)} t_{y(E)} a_3 \\
 &\stackrel{(1)}{=} t_C a_3,
 \end{aligned}$$

and hence $t_{y(\beta)} t_{y(\gamma)} = t_{y(A)} t_{y(B)}$. Note that $C \cup y(D) \cup y(E) \cup \alpha_3 = y(C \cup D \cup E \cup \alpha_3)$ bounds $\Sigma_{0,4}$. Thus, since $c = t_{y(\beta)}$, the relation $(\overline{B6}_2)$ is satisfied in $\langle X \mid Y \rangle$.

3.1.4. *On the relations $(\overline{B8}_1)$ and $(\overline{B8}_2)$.*

Let A, B, C and D be simple closed curves with arrows as shown in Figure 12. Repeating the relation (3), we have $t_A = (a_1 e a_3^{-1} a_4^{-1}) c (a_4 a_3 e^{-1} a_1^{-1})$, $t_B = (a_1^{-1} a_2^{-1} a_3^{-1} a_4^{-1}) b^{-1} (a_4 a_3 a_2 a_1)$ and $t_C = (a_3^{-1} a_2^{-1} e^{-1} a_3) a_4 (a_3^{-1} e a_2 a_3)$. By the relation (5) we see

$$t_A t_D a_2 a_4 = t_C e^{-1} t_B^{-1},$$

and by the relations (1) and (3) we see

$$t_A t_B = a_4^{-1} t_C a_2^{-1} e^{-1}.$$

Note that $A \cup D \cup \alpha_2 \cup \alpha_4$ bounds $\Sigma_{0,4}$. Thus the relation $(\overline{\text{B8}}_1)$ is satisfied in $\langle X \mid Y \rangle$.




FIGURE 12.

In addition, repeating the relation (3), we have $(a_1^{-1} a_2^{-1} a_3^{-1} a_4^{-1}) b(a_4 a_3 a_2 a_1) = t_{y^{-1}(A)}$, $(a_1 f a_3^{-1} a_4^{-1}) y^{-2} c^{-1} y^2 (a_4 a_3 f^{-1} a_1^{-1}) = t_{y^{-1}(B)}$ and $(a_3^{-1} f a_2 a_3) a_4 (a_3^{-1} a_2^{-1} f^{-1} a_3) = t_{y^{-1}(C)}$. By the relation (5) we see

$$t_{y^{-1}(A)} t_{y^{-1}(D)} f^{-1} a_4 = t_{y^{-1}(C)} a_2 t_{y^{-1}(B)}^{-1},$$

and by the relations (1) and (3) we see

$$t_{y^{-1}(A)} t_{y^{-1}(B)} = a_4^{-1} t_{y^{-1}(C)} f a_2.$$

Note that $y^{-1}(A) \cup y^{-1}(D) \cup \zeta \cup \alpha_4 = y^{-1}(A \cup D \cup \alpha_2 \cup \alpha_4)$ bounds $\Sigma_{0,4}$. Thus the relation $(\overline{\text{B8}}_2)$ is satisfied in $\langle X \mid Y \rangle$.

3.1.5. On the relation $(\overline{\text{C1a}})$.

We calculate

$$\begin{aligned} \varrho &\stackrel{(\text{C1a})}{=} (a_1 a_2 \cdots a_{g-1})^g \\ &= a_1 a_2 \cdots a_{g-1} \underbrace{(a_1 a_2 \cdots a_{g-1})^{g-2} a_1 a_2 \cdots a_{g-1}}_{(\text{A1}), (\text{A2})} \\ &= a_1 a_2 \cdots a_{g-1} a_{g-1} (a_1 a_2 \cdots a_{g-1})^{g-2} a_2 \cdots a_{g-1} \\ &= a_1 a_2 \cdots a_{g-1} a_{g-1} \underbrace{(a_1 a_2 \cdots a_{g-1})^{g-3} a_1 (a_2 \cdots a_{g-1})^2}_{(\text{A1}), (\text{A2})} \\ &= a_1 a_2 \cdots a_{g-1} a_{g-1} a_{g-2} (a_1 a_2 \cdots a_{g-1})^{g-3} (a_2 \cdots a_{g-1})^2 \\ &\vdots \\ &= a_1 a_2 a_3 \cdots a_{g-1} a_{g-1} \cdots a_3 a_2 a_1 (a_2 a_3 \cdots a_{g-1})^{g-1} \\ &\stackrel{(\text{A1}), (\text{A2})}{=} (a_2 a_3 \cdots a_{g-1})^{g-1} a_1 a_2 a_3 \cdots a_{g-1} a_{g-1} \cdots a_3 a_2 a_1. \end{aligned}$$

Let $\varrho' = (a_1^{-1}e^{-1}a_3 \cdots a_{g-1})^g$. Similarly, we calculate

$$\begin{aligned}
\varrho' &= (a_1^{-1}e^{-1}a_3 \cdots a_{g-1})^g \\
&= a_1^{-1}e^{-1}a_3 \cdots a_{g-1} \frac{(a_1^{-1}e^{-1}a_3 \cdots a_{g-1})^{g-2}a_1^{-1}e^{-1}a_3 \cdots a_{g-1}}{(\overline{A1}), (\overline{A2}), (\overline{A1_1}), (\overline{A2_1}), (\overline{A2_2})} \\
&= a_1^{-1}e^{-1}a_3 \cdots a_{g-1}a_{g-1}(a_1^{-1}e^{-1}a_3 \cdots a_{g-1})^{g-2}e^{-1}a_3 \cdots a_{g-1} \\
&= a_1^{-1}e^{-1}a_3 \cdots a_{g-1}a_{g-1} \frac{(a_1^{-1}e^{-1}a_3 \cdots a_{g-1})^{g-3}a_1^{-1}(e^{-1}a_3 \cdots a_{g-1})^2}{(\overline{A1}), (\overline{A2}), (\overline{A1_1}), (\overline{A2_1}), (\overline{A2_2})} \\
&= a_1^{-1}e^{-1}a_3 \cdots a_{g-1}a_{g-1}a_{g-2}(a_1^{-1}e^{-1}a_3 \cdots a_{g-1})^{g-3}(e^{-1}a_3 \cdots a_{g-1})^2 \\
&\vdots \\
&= a_1^{-1}e^{-1}a_3 \cdots a_{g-1}a_{g-1} \cdots a_3e^{-1}a_1^{-1}(e^{-1}a_3 \cdots a_{g-1})^{g-1} \\
&\stackrel{(\overline{A1}), (\overline{A2})}{=} (a_2a_3 \cdots a_{g-1})^{g-1}a_1a_2a_3 \cdots a_{g-1}a_{g-1} \cdots a_3a_2a_1.
\end{aligned}$$

We show that $\varrho' = y^2\varrho$. Since a regular neighborhood of $\alpha_1 \cup \varepsilon \cup \alpha_3 \cup \cdots \cup \alpha_{g-1}$ is diffeomorphic to $\Sigma_{\frac{g-1}{2}, 1}$, by the relations (1) and (4)', we have $(\varrho')^2 = 1$. We calculate

$$\begin{aligned}
\varrho'\varrho^{-1} &= (\varrho')^{-1}\varrho^{-1} \\
&= a_1ea_3^{-1} \cdots a_{g-1}^{-1}a_{g-1}^{-1} \cdots a_3^{-1}ea_1 \frac{(e^{-1}a_3 \cdots a_{g-1})^{1-g}\varrho^{-1}}{(\overline{C2}), (\overline{C2})} \\
&= a_1ea_3^{-1} \cdots a_{g-1}^{-1}a_{g-1}^{-1} \cdots a_3^{-1}ea_1 \underline{\varrho^{-1}}(f^{-1}a_3 \cdots a_{g-1})^{1-g} \\
&= a_1ea_3^{-1} \cdots a_{g-1}^{-1}a_{g-1}^{-1} \cdots a_3^{-1} \underline{ea_1} \\
&\quad \frac{a_1^{-1}a_2^{-1}a_3^{-1} \cdots a_{g-1}^{-1}a_{g-1}^{-1} \cdots a_3^{-1}a_2^{-1}a_1^{-1}(a_2a_3 \cdots a_{g-1})^{1-g}}{(f^{-1}a_3 \cdots a_{g-1})^{1-g}} \\
&\stackrel{(\overline{B4_1})}{=} a_1\{ea_2(a_2^{-1}a_3^{-1} \cdots a_{g-1}^{-1}a_{g-1}^{-1} \cdots a_3^{-1}a_2^{-1})\}^2a_1^{-1} \\
&\quad (a_2a_3 \cdots a_{g-1})^{1-g}(f^{-1}a_3 \cdots a_{g-1})^{1-g}.
\end{aligned}$$

It suffices to show the equality

$$\begin{aligned}
&y^2(f^{-1}a_3 \cdots a_{g-1})^{g-1}(a_2a_3 \cdots a_{g-1})^{g-1} \\
&= a_1\{ea_2(a_2^{-1}a_3^{-1} \cdots a_{g-1}^{-1}a_{g-1}^{-1} \cdots a_3^{-1}a_2^{-1})\}^2a_1^{-1}.
\end{aligned}$$

Let γ_1 , γ_2 and γ_3 be oriented simple loops based at $*$ which is a point obtained by the blowdown with respect to the first crosscap in $N_{g,0}$ when g is odd, as shown in Figure 13 (a), and A and B simple closed curves with arrows as shown in Figure 13 (b). By the relations (1), (4)' and Remark 2.3, we have $y^2 = \mathcal{P}_{\otimes}(\alpha^2)$, $(a_2a_3 \cdots a_{g-1})^{g-1} = \mathcal{P}_{\otimes}(\gamma_1)$, $(f^{-1}a_3 \cdots a_{g-1})^{g-1} = \mathcal{P}_{\otimes}(\gamma_2)$ and $t_At_B = \mathcal{P}_{\otimes}(\gamma_3)$.

Hence by Remark 2.4, we calculate

$$\begin{aligned}
y^2(f^{-1}a_3 \cdots a_{g-1})^{g-1}(a_2a_3 \cdots a_{g-1})^{g-1} &= \mathcal{P}_{\otimes}(\alpha^2)\mathcal{P}_{\otimes}(\gamma_2)\mathcal{P}_{\otimes}(\gamma_1) \\
&= \mathcal{P}_{\otimes}(\alpha^2)\mathcal{P}_{\otimes}(\gamma_1\gamma_2) \\
&= \mathcal{P}_{\otimes}(\gamma_1\gamma_2\alpha^2) \\
&= \mathcal{P}_{\otimes}(\gamma_3) \\
&= t_A t_B \\
&\stackrel{(1)}{=} t_A.
\end{aligned}$$

On the other hand, we see

$$\begin{aligned}
&a_1\{ea_2(a_2^{-1}a_3^{-1} \cdots a_{g-1}^{-1}a_{g-1}^{-1} \cdots a_3^{-1}a_2^{-1})\}^2a_1^{-1} \\
&= a_1(ea_2)a_1^{-1} \cdot a_1(a_2a_3 \cdots a_{g-1}a_{g-1} \cdots a_3a_2)^{-1}ea_2(a_2a_3 \cdots a_{g-1}a_{g-1} \cdots a_3a_2)a_1^{-1} \\
&\quad a_1(a_2a_3 \cdots a_{g-1}a_{g-1} \cdots a_3a_2)^{-2}a_1^{-1}.
\end{aligned}$$

In addition, by the calculation similar to the beginning of Section 3.1.5, we see

$$\begin{aligned}
(a_2a_3 \cdots a_{g-1})^{2(g-1)} &= \{(a_2a_3 \cdots a_{g-1})^{g-1}\}^2 \\
&\stackrel{(A1),(A2)}{=} \{(a_3 \cdots a_{g-1})^{g-2}(a_2a_3 \cdots a_{g-1}a_{g-1} \cdots a_3a_2)\}^2 \\
&\stackrel{(A1),(A2)}{=} (a_3 \cdots a_{g-1})^{2(g-2)}(a_2a_3 \cdots a_{g-1}a_{g-1} \cdots a_3a_2)^2,
\end{aligned}$$

and hence

$$\begin{aligned}
&a_1(a_2a_3 \cdots a_{g-1}a_{g-1} \cdots a_3a_2)^{-2}a_1^{-1} \\
&= a_1(a_2a_3 \cdots a_{g-1})^{-2(g-1)}(a_3 \cdots a_{g-1})^{2(g-2)}a_1^{-1} \\
&\stackrel{(A1)}{=} a_1(a_2a_3 \cdots a_{g-1})^{-2(g-1)}a_1^{-1}(a_3 \cdots a_{g-1})^{2(g-2)}.
\end{aligned}$$

Let γ_4 , γ_5 , γ_6 and γ_7 be oriented simple loops based at $*$ which is a point obtained by the blowdown with respect to the second crosscap, as shown in Figure 13 (a), $\gamma_{5,1}$, $\gamma_{5,2}$ and $\gamma_{5,3}$ oriented simple loops based at $*$ which is a point obtained by the blowdown with respect to the first crosscap, as shown in Figure 13 (b), and C a simple closed curve with an arrow as shown in Figure 13 (c). By the relations (3), (4)' and Remark 2.3, we have $a_1(a_2a_3 \cdots a_{g-1})^{-(g-1)}a_1^{-1} = \mathcal{P}_{\otimes}(\gamma_4)$,

$$\begin{aligned}
&a_1(a_2a_3 \cdots a_{g-1}a_{g-1} \cdots a_3a_2)^{-1}ea_2(a_2a_3 \cdots a_{g-1}a_{g-1} \cdots a_3a_2)a_1^{-1} \\
&= a_1(a_2a_3 \cdots a_{g-1}a_{g-1} \cdots a_3a_2)^{-1}\mathcal{P}_{\otimes}(\gamma_{5,1})(a_2a_3 \cdots a_{g-1}a_{g-1} \cdots a_3a_2)a_1^{-1} \\
&= a_1(a_{g-1} \cdots a_3a_2)^{-1}\mathcal{P}_{\otimes}(\gamma_{5,2})(a_{g-1} \cdots a_3a_2)a_1^{-1} \\
&= a_1\mathcal{P}_{\otimes}(\gamma_{5,3})a_1^{-1} \\
&= \mathcal{P}_{\otimes}(\gamma_5),
\end{aligned}$$

$a_1(ea_2)a_1^{-1} = \mathcal{P}_\otimes(\gamma_6)$, $t_A t_C^{-1} = \mathcal{P}_\otimes(\gamma_7)$ and $(a_3 \cdots a_{g-1})^{2(g-2)} = t_C$. Hence by Remark 2.4, we calculate

$$\begin{aligned}
 a_1\{ea_2(a_2^{-1}a_3^{-1} \cdots a_{g-1}^{-1}a_{g-1}^{-1} \cdots a_3^{-1}a_2^{-1})\}^2 a_1^{-1} &= \mathcal{P}_\otimes(\gamma_6)\mathcal{P}_\otimes(\gamma_5)\mathcal{P}_\otimes(\gamma_4)^2 t_C \\
 &= \mathcal{P}_\otimes(\gamma_6)\mathcal{P}_\otimes(\gamma_4\gamma_5)\mathcal{P}_\otimes(\gamma_4) t_C \\
 &= \mathcal{P}_\otimes(\gamma_4\gamma_5\gamma_6)\mathcal{P}_\otimes(\gamma_4) t_C \\
 &= \mathcal{P}_\otimes(\gamma_4^2\gamma_5\gamma_6) t_C \\
 &= \mathcal{P}_\otimes(\gamma_7) t_C \\
 &= t_A.
 \end{aligned}$$

Thus the relation $(\overline{\text{C1a}})$ is satisfied in $\langle X \mid Y \rangle$.




FIGURE 13.

3.1.6. On the relation $(\overline{\text{C4a}})$.

For $1 \leq i \leq \frac{g-1}{2}$, let γ_i be an oriented simple loop based at $*$ which is a point obtained by the blowdown with respect to the first crosscap, as shown in Figure 14, and $\Phi = e^{-1}a_3 \cdots a_{g-1}$.

We now prove the following lemma.

Lemma 3.2. *In $\langle X \mid Y \rangle$, we have $\mathcal{P}_\otimes(\gamma_1)\Phi^2\mathcal{P}_\otimes(\gamma_i) = \mathcal{P}_\otimes(\gamma_{i+1})\Phi^2$ for $1 \leq i \leq \frac{g-3}{2}$.*

Proof. Let δ_i be an oriented simple loop based at $*$ which is a point obtained by the blowdown with respect to the first crosscap, as shown in Figure 14. By Remark 2.3, repeating the relation (3), we have $\Phi^2\mathcal{P}_\otimes(\gamma_i)\Phi^{-2} = \mathcal{P}_\otimes(\delta_i)$. In addition, by Remarks 2.3 and 2.4, we have $\mathcal{P}_\otimes(\gamma_1)\mathcal{P}_\otimes(\delta_i) = \mathcal{P}_\otimes(\gamma_{i+1})$, and hence the claim holds. $\square$




FIGURE 14.

By the relation (4)' and Remark 2.3, we have $a_2e = \mathcal{P}_\otimes(\gamma_1)$ and $\mathcal{P}_\otimes(\gamma_{\frac{g-1}{2}}) = \Phi^{1-g}$. Hence by Lemma 3.2, we calculate

$$\begin{aligned}
 (a_2a_3 \cdots a_{g-1}e^{-1}a_3 \cdots a_{g-1})^{\frac{g-1}{2}} &= (a_2e(e^{-1}a_3 \cdots a_{g-1})^2)^{\frac{g-1}{2}} \\
 &= (\mathcal{P}_\otimes(\gamma_1)\Phi^2)^{\frac{g-1}{2}} \\
 &= (\mathcal{P}_\otimes(\gamma_1)\Phi^2)^{\frac{g-5}{2}}\mathcal{P}_\otimes(\gamma_2)\Phi^4 \\
 &= (\mathcal{P}_\otimes(\gamma_1)\Phi^2)^{\frac{g-7}{2}}\mathcal{P}_\otimes(\gamma_3)\Phi^6 \\
 &\vdots \\
 &= \mathcal{P}_\otimes(\gamma_{\frac{g-1}{2}})\Phi^{g-1} \\
 &= 1.
 \end{aligned}$$

Thus the relation $(\overline{C4a})$ is satisfied in $\langle X \mid Y \rangle$.

3.1.7. On the relation $(\overline{C3})$.

In $\mathcal{M}(N_{g,0})$, $\bar{\varrho}$ is defined as $\bar{\varrho} = y\varrho$ and we have the relation $\varrho = (y^{-1}a_2a_3 \cdots a_{g-1}ya_2a_3 \cdots a_{g-1})^{\frac{g-2}{2}}y^{-1}a_2a_3 \cdots a_{g-1}$ (see Theorem 2.2 in [26]). In addition, note that Lemma 3.2 holds even if g is even. Hence we calculate

$$\begin{aligned}
 \bar{\varrho} &= y(y^{-1}a_2a_3 \cdots a_{g-1}ya_2a_3 \cdots a_{g-1})^{\frac{g-2}{2}}y^{-1}a_2a_3 \cdots a_{g-1} \\
 &= (a_2a_3 \cdots a_{g-1}ya_2a_3 \cdots a_{g-1}y^{-1})^{\frac{g-2}{2}}a_2a_3 \cdots a_{g-1} \\
 &= (a_2a_3 \cdots a_{g-1}e^{-1}a_3 \cdots a_{g-1})^{\frac{g-2}{2}}a_2a_3 \cdots a_{g-1} \\
 &= (\mathcal{P}_\otimes(\gamma_1)\Phi^2)^{\frac{g-2}{2}}\mathcal{P}_\otimes(\gamma_1)\Phi \\
 &= (\mathcal{P}_\otimes(\gamma_1)\Phi^2)^{\frac{g-4}{2}}\mathcal{P}_\otimes(\gamma_2)\Phi^3 \\
 &= (\mathcal{P}_\otimes(\gamma_1)\Phi^2)^{\frac{g-6}{2}}\mathcal{P}_\otimes(\gamma_3)\Phi^5 \\
 &\vdots \\
 &= \mathcal{P}_\otimes(\gamma_1)\Phi^2\mathcal{P}_\otimes(\gamma_{\frac{g-2}{2}})\Phi^{g-3} \\
 &= \mathcal{P}_\otimes(\gamma_1)(\Phi^2\mathcal{P}_\otimes(\gamma_{\frac{g-2}{2}})\Phi^{-2})\Phi^{g-1},
 \end{aligned}$$

where γ_i is defined in Section 3.1.6. Let δ_1 and δ_2 be oriented simple loops based at $*$ which is a point obtained by the blowdown with respect to the first crosscap, as shown in Figure 15 (a). Repeating the relation (3), we have $\Phi^2 \mathcal{P}_\otimes(\gamma_{\frac{g-2}{2}}) \Phi^{-2} = \mathcal{P}_\otimes(\delta_1)$. In addition, by Remark 2.4, we have $\mathcal{P}_\otimes(\gamma_1) \mathcal{P}_\otimes(\delta_1) = \mathcal{P}_\otimes(\delta_2)$. Let δ_3 and δ_4 be oriented simple loops based at $*$ which is a point obtained by the blowdown with respect to the first crosscap, as shown in Figure 15 (a). Repeating the relation (3), we have $\Phi^{g-1} \mathcal{P}_\otimes(\delta_2) \Phi^{1-g} = \mathcal{P}_\otimes(\delta_3)$. In addition, by Remark 2.4, we have $\mathcal{P}_\otimes(\delta_2) \mathcal{P}_\otimes(\delta_3) = \mathcal{P}_\otimes(\delta_4)$. Let A and B be simple closed curves with arrows as shown in Figure 15 (b). By the relation (4)' and Remark 2.3, we have $\Phi^{2g-2} = t_A$ and $\mathcal{P}_\otimes(\delta_4) = t_B t_A^{-1}$. Hence we calculate

$$\begin{aligned}
 \bar{\varrho}^2 &= (\mathcal{P}_\otimes(\delta_2) \Phi^{g-1})^2. \\
 &= \mathcal{P}_\otimes(\delta_2) \mathcal{P}_\otimes(\delta_3) \Phi^{2g-2} \\
 &= \mathcal{P}_\otimes(\delta_4) t_A \\
 &= t_B \\
 &\stackrel{(1)}{=} 1.
 \end{aligned}$$

Thus the relation $(\overline{C3})$ is satisfied in $\langle X \mid Y \rangle$.




FIGURE 15.

3.1.8. On the relation $(\overline{C4})$.

Note that the relation $a_2 \bar{\varrho} = \bar{\varrho} e^{-1}$ is obtained from the relations $(\overline{C2_3})$ and $(\overline{C3})$. We call this relation $(\overline{C2_4})$. By the equality $\bar{\varrho} = (a_2 a_3 \cdots a_{g-1} e^{-1} a_3 \cdots a_{g-1})^{\frac{g-2}{2}} a_2 a_3 \cdots a_{g-1}$ which appeared in Section 3.1.7, we

calculate

$$\begin{aligned}
(\bar{\rho}a_2a_3 \cdots a_{g-1})^{g-1} &= \frac{\bar{\rho}a_2a_3 \cdots a_{g-1}\bar{\rho}a_2a_3 \cdots a_{g-1}(\bar{\rho}a_2a_3 \cdots a_{g-1})^{g-3}}{(\overline{C2_2}),(\overline{C2_4})} \\
&= \frac{\bar{\rho}^2e^{-1}a_3 \cdots a_{g-1}a_2a_3 \cdots a_{g-1}\bar{\rho}a_2a_3 \cdots a_{g-1}(\bar{\rho}a_2a_3 \cdots a_{g-1})^{g-4}}{(\overline{C2_2}),(\overline{C2_3}),(\overline{C2_4})} \\
&= \bar{\rho}^3a_2a_3 \cdots a_{g-1}e^{-1}a_3 \cdots a_{g-1}a_2a_3 \cdots a_{g-1}(\bar{\rho}a_2a_3 \cdots a_{g-1})^{g-4} \\
&\vdots \\
&= \bar{\rho}^{g-1}(a_2a_3 \cdots a_{g-1}e^{-1}a_3 \cdots a_{g-1})^{\frac{g-2}{2}}a_2a_3 \cdots a_{g-1} \\
&= \bar{\rho}^g \\
&\stackrel{(\overline{C3})}{=} 1.
\end{aligned}$$

Thus the relation $(\overline{C4})$ is satisfied in $\langle X \mid Y \rangle$.

Therefore well-definedness of ψ follows.

3.2. Surjectivity of ψ .

For any simple closed curve d of $N_{g,n}$, we denote the complement of the interior of a regular neighborhood of d by $N_{g,n} \setminus d$. If $N_{g,n} \setminus d$ and $N_{g,n} \setminus d'$ are diffeomorphic, there exists $x \in \mathcal{M}(N_{g,n})$ with $x(d') = d$ for $n \leq 1$. In particular, if d and d' are two sided, we have $xt_{d',\theta'}x^{-1} = t_{d,\theta}$ for some orientation θ and θ' .

For any two sided simple closed curve d of $N_{g,n}$ where $n \leq 1$, $N_{g,n} \setminus d$ is diffeomorphic to either one of

- $N_{g-2,n+2}$, where $g \geq 3$,
- $\Sigma_{\frac{g-2}{2},n+2}$, where g is even,
- $\Sigma_{h,1} \sqcup N_{g-2h,n+1}$, where $0 \leq h < \frac{g}{2}$,
- $N_{i,1} \sqcup N_{g-i,n+1}$, where $1 \leq i \leq g-1$ and $g \geq 2$,
- $N_{i,1} \sqcup \Sigma_{\frac{g-i}{2},n+1}$, where $1 \leq i \leq g$ and $g-i$ is even.

For each case, we would like to find a word w on X_0 such that $\psi(w) = t_{d,\theta}$, where X_0 is defined at the beginning of Section 3.

3.2.1. The case where $N_{g,n} \setminus d$ is diffeomorphic to $N_{g-2,n+2}$ or $\Sigma_{\frac{g-2}{2},n+2}$.

Since $N_{g,n} \setminus d$ is diffeomorphic to either $N_{g,n} \setminus \alpha_1$ or $N_{g,n} \setminus \beta_{\frac{g-2}{2}}$, there exists $x \in \mathcal{M}(N_{g,n})$ such that $xt_{d',\theta'}x^{-1} = t_{d,\theta}$, where $t_{d',\theta'} = a_1$ or $b_{\frac{g-2}{2}}$ respectively. If $x \in \mathcal{T}(N_{g,n})$, there exists a word $x = x_1x_2 \cdots x_s$ on X_0 . Then we obtain $\psi((x_1x_2 \cdots x_s)t_{d',\theta'}(x_1x_2 \cdots x_s)^{-1}) = t_{d,\theta}$, repeating the relation (3). If $x \notin \mathcal{T}(N_{g,n})$, since $xy^{-1} \in \mathcal{T}(N_{g,n})$ by the sequence (3) in Section 2.2, there exists a word $xy^{-1} = x_1x_2 \cdots x_s$ on X_0 . Since $yt_{d',\theta'}y^{-1} = a_1^{-1}$ or $b_{\frac{g-2}{2}}^{-1}$, we obtain $\psi((x_1x_2 \cdots x_s)(yt_{d',\theta'}y^{-1})(x_1x_2 \cdots x_s)^{-1}) = t_{d,\theta}$, repeating the relation (3).

3.2.2. The case where $N_{g,n} \setminus d$ is diffeomorphic to $\Sigma_{h,1} \sqcup N_{g-2h,n+1}$.

When $h = 0$, since $t_{d,\theta} = 1$ by the relation (1), we obtain $\psi(1) = t_{d,\theta}$. When $h \geq 1$, there exists $x \in \mathcal{M}(N_{g,n})$ such that $xt_{d',\theta'}x^{-1} = t_{d,\theta}$, where d' is the boundary curve of a regular neighborhood of $\alpha_1 \cup \alpha_2 \cup \cdots \cup \alpha_{2h}$ which is diffeomorphic to $\Sigma_{h,1}$, as shown in Figure 16. Note that we have $t_{d',\theta'}^\epsilon = (a_1a_2 \cdots a_{2h})^{4h+2}$ for some $\epsilon = \pm 1$, by the relation (4)'. If $x \in \mathcal{T}(N_{g,n})$, there exists a word $x = x_1x_2 \cdots x_s$ on X_0 . Then we obtain $\psi((x_1x_2 \cdots x_s)(a_1a_2 \cdots a_{2h})^{\epsilon(4h+2)}(x_1x_2 \cdots x_s)^{-1}) = t_{d,\theta}$,

repeating the relations (3) and (4)'. If $x \notin \mathcal{T}(N_{g,n})$, there exists a word $xy^{-1} = x_1x_2 \cdots x_s$ on X_0 . Since $ya_1y^{-1} = a_1^{-1}$, $ya_2y^{-1} = e^{-1}$ and $ya_iy^{-1} = a_i$ for $i \geq 3$, we obtain $\psi((x_1x_2 \cdots x_s)(a_1^{-1}e^{-1}a_3 \cdots a_{2h})^{\epsilon(4h+2)}(x_1x_2 \cdots x_s)^{-1}) = t_{d;\theta}$, repeating the relations (3) and (4)'.




FIGURE 16.

3.2.3. *The case where $N_{g,n} \setminus d$ is diffeomorphic to $N_{i,1} \sqcup N_{g-i,n+1}$.*

When $i = 1$, since $t_{d;\theta} = 1$ by the relation (1), we obtain $\psi(1) = t_{d;\theta}$. When $i = 2$, there exists $x \in \mathcal{M}(N_{g,n})$ such that $xy^2x^{-1} = xt_\delta x^{-1} = t_{d;\theta}$, where δ appeared in Figure 8. If $x \in \mathcal{T}(N_{g,n})$, there exists a word $x = x_1x_2 \cdots x_s$ on X_0 . Then we obtain $\psi((x_1x_2 \cdots x_s)y^2(x_1x_2 \cdots x_s)^{-1}) = t_{d;\theta}$, repeating the relation (3). If $x \notin \mathcal{T}(N_{g,n})$, there exists a word $xy^{-1} = x_1x_2 \cdots x_s$ on X_0 . Since $y \cdot y^2 \cdot y^{-1} = y^2$, we have $\psi((x_1x_2 \cdots x_s)y^2(x_1x_2 \cdots x_s)^{-1}) = t_{d;\theta}$, repeating the relation (3). When $i \geq 3$, we take simple closed curves $d_1, \dots, d_6$ as shown in Figure 17. By induction on i , we can suppose that $t_{d_j;\theta_j}$ is described as a word on X_0 for $1 \leq j \leq 6$. Hence we obtain $\psi(t_{d_1;\theta_1}^{\epsilon_1} t_{d_2;\theta_2}^{\epsilon_2} t_{d_3;\theta_3}^{\epsilon_3} t_{d_4;\theta_4}^{\epsilon_4} t_{d_5;\theta_5}^{\epsilon_5} t_{d_6;\theta_6}^{\epsilon_6}) = t_{d;\theta}$ for some $\epsilon_j = \pm 1$, by the relations (2) and (5).




FIGURE 17.

3.2.4. *The case where $N_{g,n} \setminus d$ is diffeomorphic to $N_{i,1} \sqcup \Sigma_{\frac{g-i}{2},n+1}$.*

When $i = 1$, since $t_{d;\theta} = 1$ by the relation (1), we obtain $\psi(1) = t_{d;\theta}$. When $i = 2$, d is described as shown in Figure 18, for some model of $N_{g,n}$. Let γ_1 and γ_2 be oriented loops of $N_{g-1,n}$ based at $*$ as shown in Figure 18, where $*$ is the point obtained by the blowdown with respect to the crosscap M in Figure 18. Note that γ_1 and γ_2 are two sided since g is even when $i = 2$. In addition, when $\mathcal{P}_\otimes(\gamma_j) = t_{c_j} t_{c'_j}^{-1}$, $N_{g,n} \setminus (c_j \cup c'_j)$ is diffeomorphic to $N_{g,n} \setminus (\alpha_2 \cup \zeta)$ for $j = 1$ and 2 , where α_2 and ζ appeared in Figure 8. Hence by Remark 2.3, we can describe $\mathcal{P}_\otimes(\gamma_j) = x_j(a_2f)x_j^{-1}$ for some $x_j \in \mathcal{M}(N_{g,n})$, for $j = 1$ and 2 . If $x_j \in \mathcal{T}(N_{g,n})$, x_j is represented by a word on X_0 , and hence $\mathcal{P}_\otimes(\gamma_j)$ is a word on X_0 . If $x_j \notin \mathcal{T}(N_{g,n})$, x_jy^{-1} is represented by a word on X_0 . Since $y(a_2f)y^{-1} = e^{-1}a_2^{-1}$, $\mathcal{P}_\otimes(\gamma_j)$ is a word on X_0 . We denote the word presentation of $\mathcal{P}_\otimes(\gamma_j)$ on X_0 by w_j . Then we calculate $\psi(w_2w_1) = w_2w_1 \stackrel{(3)}{=} \mathcal{P}_\otimes(\gamma_2)\mathcal{P}_\otimes(\gamma_1) = \mathcal{P}_\otimes(\gamma_1\gamma_2) \stackrel{(1)}{=} t_{d;\theta}$, by

Remarks 2.3 and 2.4. When $i \geq 3$, we take simple closed curves $d_1, \dots, d_6$ as shown in Figure 17, similar to Section 3.2.3. Since $N_{g,n} \setminus d_j$ is diffeomorphic to either $N_{1,1} \sqcup N_{g-1,n+1}$, $N_{2,1} \sqcup N_{g-2,n+1}$, $N_{i-1,1} \sqcup N_{g-i+1,n+1}$ or $N_{i-2,1} \sqcup N_{g-i+2,n+1}$, by Section 3.2.3, $t_{d_j;\theta_j}$ is described as a word on X_0 for $1 \leq j \leq 6$. Hence we obtain $\psi(t_{d_1;\theta_1}^{\epsilon_1} t_{d_2;\theta_2}^{\epsilon_2} t_{d_3;\theta_3}^{\epsilon_3} t_{d_4;\theta_4}^{\epsilon_4} t_{d_5;\theta_5}^{\epsilon_5} t_{d_6;\theta_6}^{\epsilon_6}) = t_{d;\theta}$ for some $\epsilon_j = \pm 1$, by the relations (2) and (5).




FIGURE 18.

Therefore surjectivity of ψ follows, and hence the proof of Proposition 3.1 is completed.

4. PROOF OF THEOREM 1.1 FOR $g \geq 1$ AND $n \geq 2$

Recall the capping map $\mathcal{C}$, the point pushing map $\mathcal{P}_*$ and the forgetful map $\mathcal{F}$ defined in Section 2.1. Let $\mathcal{T}^+(N_{g,n-1}, *) = \mathcal{C}(\mathcal{T}(N_{g,n}))$, then for $g \geq 1$ and $n \geq 2$, from the sequences (1) and (2) in Section 2.1, we have the short exact sequences

$$(4) \quad 1 \rightarrow \mathbb{Z} \rightarrow \mathcal{T}(N_{g,n}) \xrightarrow{\mathcal{C}} \mathcal{T}^+(N_{g,n-1}, *) \rightarrow 1,$$

$$(5) \quad 1 \rightarrow \pi_1^+(N_{g,n-1}, *) \xrightarrow{\mathcal{P}_*} \mathcal{T}^+(N_{g,n-1}, *) \xrightarrow{\mathcal{F}} \mathcal{T}(N_{g,n-1}) \rightarrow 1.$$

Using these sequences and the presentation of $\mathcal{T}(N_{g,1})$ given in Section 3, we give the presentation of $\mathcal{T}(N_{g,n})$ for $g \geq 1$ and $n \geq 2$, by induction on n .

Note that, in general, as basics on combinatorial group theory, from presented groups $\langle G_1 \mid R_1 \rangle$ and $\langle G_3 \mid R_3 \rangle$ with a short exact sequence $1 \rightarrow \langle G_1 \mid R_1 \rangle \xrightarrow{i} G \xrightarrow{p} \langle G_3 \mid R_3 \rangle \rightarrow 1$, we can give a presentation for G as follows. Let $\tilde{g}$ be any lift of $g \in G_3$ with respect to p , and $\tilde{r}$ a word obtained from $r \in R_3$ by replacing each $g \in G_3$ to $\tilde{g}$. For $x \in \ker p$, denote by w_x a word on $i(G_1)$ corresponding to x . For $r \in R_1$, denote by $\tilde{r}$ a word on $i(G_1)$ obtained from r by replacing $h \in G_1$ to $i(h)$. Let $G_2 = \{i(h), \tilde{g} \mid h \in G_1, g \in G_3\}$ and $R_2 = \{\tilde{r}_1, \tilde{r}_3 w_{r_3}^{-1}, \tilde{g} i(h) \tilde{g}^{-1} w_{\tilde{g} i(h) \tilde{g}^{-1}}^{-1} \mid r_1 \in R_1, r_3 \in R_3, h \in G_1, g \in G_3\}$. Then we have a presentation $G = \langle G_2 \mid R_2 \rangle$. For details, for instance see Proposition 1 in 10.2 in [11].

We now show the following proposition, using the sequence (5). Remember the infinite presentation $\langle X \mid Y \rangle$ for the group presented in Theorem 1.1.

Proposition 4.1. *For $g \geq 1$ and $n \geq 2$, suppose $\mathcal{T}(N_{g,n-1}) = \langle X \mid Y \rangle$, then $\mathcal{T}^+(N_{g,n-1}, *)$ admits a presentation with a generating set*

$$\tilde{X} = \left\{ t_{\tilde{c}, \tilde{\theta}} \left| \begin{array}{l} \tilde{c} \text{ is a two sided simple closed curve of } N_{g,n-1} \setminus \{*\} \\ \text{which does not bound a disk neighborhood of } *, \text{ and} \\ \tilde{\theta} \text{ is an orientation of a regular neighborhood of } \tilde{c}. \end{array} \right. \right\}.$$

The defining relations are

- ($\tilde{1}$) $t_{\tilde{c}, \tilde{\theta}} = 1$ if $\tilde{c}$ bounds a disk or a Möbius band,
- ($\tilde{2}$) $t_{\tilde{c}; -\tilde{\theta}}^{-1} = t_{\tilde{c}; +\tilde{\theta}}$,

- ($\tilde{3}$) all the conjugation relations $ft_{\tilde{c},\tilde{\theta}}f^{-1} = t_{f(\tilde{c});f_*(\tilde{\theta})}$ for $f \in \tilde{X}$,
- ($\tilde{4}$) all the 2-chain relations,
- ($\tilde{5}$) all the lantern relations,
- ($\tilde{5}'$) all the extended lantern relations defined in Remark 2.2.

Proof. $\pi_1^+(N_{g,n-1}, *)$ is a finite rank free group for $n \geq 2$. However, we consider an infinite presentation for $\pi_1^+(N_{g,n-1}, *)$. Let π be the group generated by symbols S_α for a non-trivial simple loop $\alpha \in \pi_1^+(N_{g,n-1}, *)$, and with the defining relations

- $S_{\alpha^{-1}} = S_\alpha^{-1}$,
- $S_\alpha S_\beta = S_\gamma$ if $\alpha\beta = \gamma$,
- $S_\alpha S_\beta S_\alpha^{-1} = S_\gamma$ if $\alpha\beta\alpha^{-1} = \gamma$.

Then we have that π is isomorphic to $\pi_1^+(N_{g,n-1}, *)$ (see Theorem 1.2 in [13]). So π gives an infinite presentation for $\pi_1^+(N_{g,n-1}, *)$. By this we identify $\pi_1^+(N_{g,n-1}, *)$ with π .

We take a simple path P of $N_{g,n-1}$ between $*$ and the $(n-1)$ -st boundary component. For a simple closed curve c of $N_{g,n-1}$, let $\hat{c}$ be a simple closed curve of $N_{g,n-1} \setminus \{*\}$ corresponding to c which does not intersect P , as shown in Figure 19.




FIGURE 19. Examples of simple closed curves of $N_{g,n-1} \setminus \{*\}$ corresponding to simple closed curves of $N_{g,n-1}$ which do not intersect the path P .

By the sequence ($\tilde{5}$), $\mathcal{T}^+(N_{g,n-1}, *)$ is generated by

- $t_{\tilde{c},\tilde{\theta}}$ for any simple closed curve c of $N_{g,n-1}$ and
- $\mathcal{P}_*(S_\alpha)$ for any generator S_α of π .

The defining relations are

- $\tilde{r} = w_{\tilde{r}}$ for the lift of any $r \in Y$ with respect to $\mathcal{F}$,
- $\mathcal{P}_*(S_{\alpha^{-1}}) = \mathcal{P}_*(S_\alpha)^{-1}$,
- $\mathcal{P}_*(S_\beta)\mathcal{P}_*(S_\alpha) = \mathcal{P}_*(S_\gamma)$ for any S_α, S_β and S_γ satisfying $\alpha\beta = \gamma$,
- $\mathcal{P}_*(S_\alpha)^{-1}\mathcal{P}_*(S_\beta)\mathcal{P}_*(S_\alpha) = \mathcal{P}_*(S_\gamma)$ for any S_α, S_β and S_γ satisfying $\alpha\beta\alpha^{-1} = \gamma$ and
- $t_{\tilde{c},\tilde{\theta}}\mathcal{P}_*(S_\alpha)t_{\tilde{c},\tilde{\theta}}^{-1} = w_{t_{\tilde{c},\tilde{\theta}}\mathcal{P}_*(S_\alpha)t_{\tilde{c},\tilde{\theta}}^{-1}}$ for any $t_{\tilde{c},\tilde{\theta}}$ and S_α ,

where w_x is a word corresponding to x on $\{\mathcal{P}_*(S_\alpha)\}$. We denote this presentation by $\langle \tilde{X}_0 \mid \tilde{Y}_0 \rangle$. In addition, let $\langle \tilde{X} \mid \tilde{Y} \rangle$ be the infinitely presented group with the presentation given in Proposition 4.1. We show that $\langle \tilde{X} \mid \tilde{Y} \rangle$ is isomorphic to $\langle \tilde{X}_0 \mid \tilde{Y}_0 \rangle = \mathcal{T}^+(N_{g,n-1}, *)$.

Denote by $F(\tilde{X}_0)$ the free group freely generated by $\tilde{X}_0$. Let $\tilde{p} : F(\tilde{X}_0) \rightarrow \langle \tilde{X}_0 \mid \tilde{Y}_0 \rangle$ be the natural projection and $\tilde{\eta} : F(\tilde{X}_0) \rightarrow \langle \tilde{X} \mid \tilde{Y} \rangle$ the homomorphism defined as $\tilde{\eta}(x) = x$ for any $x \in \tilde{X}_0$. Note that, by Remark 2.1, $\mathcal{P}_*(S_\alpha)$ is also in $\langle \tilde{X} \mid \tilde{Y} \rangle$ for any generator S_α of π . Hence $\tilde{\eta}$ is well-defined. We consider a correspondence

$$\tilde{\psi} : \langle \tilde{X}_0 \mid \tilde{Y}_0 \rangle \rightarrow \langle \tilde{X} \mid \tilde{Y} \rangle$$

satisfying $\tilde{\psi} \circ \tilde{p} = \tilde{\eta}$. We show that $\tilde{\psi}$ is an isomorphism.

First we show well-definedness of $\tilde{\psi}$. In the first relation of $\tilde{Y}_0$, from the definition of $\tilde{c}$, it is clear that $w_{\tilde{r}} = 1$ for any $r \in Y$. Hence the first relation of $\tilde{Y}_0$ is either one of the relations (1)-(5). In the second relation of $\tilde{Y}_0$, the relation $\mathcal{P}_*(S_{\alpha-1}) = \mathcal{P}_*(S_{\alpha})^{-1}$ is trivial. The relation $\mathcal{P}_*(S_{\beta})\mathcal{P}_*(S_{\alpha}) = \mathcal{P}_*(S_{\gamma})$ is obtained from the relations (3) or (5)' by Remark 2.2. The relation $\mathcal{P}_*(S_{\alpha})^{-1}\mathcal{P}_*(S_{\beta})\mathcal{P}_*(S_{\alpha}) = \mathcal{P}_*(S_{\gamma})$ is the relation (3). In the third relation of $\tilde{Y}_0$, we have $w_{t_{\tilde{c};\tilde{\theta}}\mathcal{P}_*(S_{\alpha})t_{\tilde{c};\tilde{\theta}}^{-1}} = \mathcal{P}_*(S_{t_{\tilde{c};\tilde{\theta}}(\alpha)})$, and hence it is the relation (3). Therefore, any relation of $\tilde{Y}_0$ is satisfied in $\langle \tilde{X} \mid \tilde{Y} \rangle$. So $\tilde{\psi}$ is well-defined as a homomorphism.

Next we show bijectivity of $\tilde{\psi}$. Let $\tilde{\varphi} : \langle \tilde{X} \mid \tilde{Y} \rangle \rightarrow \langle \tilde{X}_0 \mid \tilde{Y}_0 \rangle = \mathcal{T}^+(N_{g,n-1}, *)$ be the homomorphism defined as $\tilde{\varphi}(x) = x$ for any $x \in \tilde{X}$. Since $\tilde{\varphi}(\tilde{Y}) = 1$ in $\mathcal{T}^+(N_{g,n-1}, *)$ clearly, $\tilde{\varphi}$ is well-defined. By the definitions of $\tilde{\psi}$ and $\tilde{\varphi}$, it is clear that $\tilde{\varphi} \circ \tilde{\psi}$ is the identity map, and hence $\tilde{\psi}$ is injective. For any $t_{\tilde{c};\tilde{\theta}} \in \tilde{X}$, if $\tilde{c}$ intersects P transversally at $l \geq 1$ points, there exist $t_{\tilde{c}';\tilde{\theta}'}$ and S_{α} such that $t_{\tilde{c};\tilde{\theta}} = \tilde{\psi}(\mathcal{P}_*(S_{\alpha})t_{\tilde{c}';\tilde{\theta}'})$, as shown in Figure 20, by Remark 2.3. We notice that $\tilde{c}'$ intersects P transversally at $l-1$ points. By induction on the intersection number l of $\tilde{c}$ and P , we see that $t_{\tilde{c};\tilde{\theta}}$ is a product of some $\tilde{\psi}(t_{\tilde{c};\tilde{\theta}})$ and some $\tilde{\psi}(\mathcal{P}_*(S_{\alpha}))$'s. Hence $\tilde{\psi}$ is surjective, and so bijective.




FIGURE 20. $\tilde{c}$ and $\tilde{c}'$ are the boundary curves of a regular neighborhood of α .

Therefore $\tilde{\psi}$ is the isomorphism. Thus the claim is obtained. $\square$

Finally we complete the proof of Theorem 1.1.

Proof of Theorem 1.1. From the case where $n = 1$ of Theorem 1.1, Proposition 4.1 and the sequence (4), by induction on n , $\mathcal{T}(N_{g,n})$ is generated by

- the natural lift $t_{c;\theta}$ of $t_{\tilde{c};\tilde{\theta}}$ with respect to $\mathcal{C}$ and
- the Dehn twist $t_{\partial_n;\theta_n}$ about the n -th boundary curve ∂_n ,

that is, X generate $\mathcal{T}(N_{g,n})$. In addition $\mathcal{T}(N_{g,n})$ has the relations

- $\tilde{r} = w_{\tilde{r}}$ for the lift of any relator r of $\mathcal{T}^+(N_{g,n-1}, *)$ with respect to $\mathcal{C}$ and
- $t_{c;\theta}t_{\partial_n;\theta_n}t_{c;\theta}^{-1} = w_{t_{c;\theta}t_{\partial_n;\theta_n}t_{c;\theta}^{-1}}$,

where $w_x = t_{\partial_n;\theta_n}^m$ for some integer m corresponding to x . In the first relation, it is clear that $w_{\tilde{r}} = 1$ if r is a relator corresponding to the relations (1)-(5) of

Proposition 4.1. On the other hand, if r is a relator corresponding to the relation $(\tilde{5})'$ of Proposition 4.1, then $w_{\tilde{r}} = t_{\partial_n; \theta_n}^\epsilon$ for some $\epsilon = \pm 1$. Hence the first relation is either one of the relation (1)-(5). In the second relation, it is clear that $w_{t_{c; \theta} t_{\partial_n; \theta_n} t_{c; \theta}^{-1}} = t_{\partial_n; \theta_n}$, and hence it is the relation (3).

Thus we complete the proof. $\square$

ACKNOWLEDGEMENT

The authors would like to express their gratitude to Susumu Hirose for his useful comments.

REFERENCES

- [1] J. S. Birman, *Mapping class groups and their relationship to braid groups*, Comm. Pure Appl. Math. **22** (1969), 213–238.
- [2] J. S. Birman and D. R. J. Chillingworth, *On the homeotopy group of a non-orientable surface*, Proc. Cambridge Philos. Soc. **71** (1972), 437–448. Erratum: Math. Proc. Cambridge Philos. Soc. **136** (2004), no. 2, 441.
- [3] M. Dehn, *Die Gruppe der Abbildungsklassen*, Das arithmetische Feld auf Flächen, Acta Math. **69** (1938), no. 1, 135–206.
- [4] M. Dehn, *Papers on group theory and topology*, Springer-Verlag, New York, 1987.
- [5] D. B. A. Epstein, *Curves on 2-manifolds and isotopies*, Acta Math. **115** (1966), 83–107.
- [6] B. Farb and D. Margalit, *A primer on mapping class groups*, Princeton Mathematical Series, **49**.
- [7] S. Gervais, *Presentation and central extensions of mapping class groups*, Trans. Amer. Math. Soc. **348** (1996), no. 8, 3097–3132.
- [8] S. Gervais, *A finite presentation of the mapping class group of a punctured surface*, Topology **40** (2001), no. 4, 703–725.
- [9] J. Harer, *The second homology group of the mapping class group of an orientable surface*, Invent. Math. **72** (1983), no. 2, 221–239.
- [10] A. Hatcher and W. Thurston, *A presentation for the mapping class group of a closed orientable surface*, Topology **19** (1980), no. 3, 221–237.
- [11] D. L. Johnson, *Presentations of groups*, Second edition. London Mathematical Society Student Texts, **15**. Cambridge University Press, Cambridge, 1997.
- [12] R. Kobayashi and G. Omori, *An infinite presentation for the mapping class group of a non-orientable surface with boundary*, Osaka J. Math., **59** (2) (2022), 269–314.
- [13] R. Kobayashi, *Infinite presentations for fundamental groups of surfaces*, Hiroshima Math. J., to appear.
- [14] M. Korkmaz, *Mapping class groups of nonorientable surfaces*, Geom. Dedicata **89** (2002), 109–133.
- [15] C. Labruère and L. Paris, *Presentations for the punctured mapping class groups in terms of Artin groups*, Algebr. Geom. Topol. **1** (2001), 73–114.
- [16] W. B. R. Lickorish, *Homeomorphisms of non-orientable two-manifolds*, Proc. Cambridge Philos. Soc. **59** (1963), 307–317.
- [17] W. B. R. Lickorish, *A finite set of generators for the homeotopy group of a 2-manifold*, Proc. Cambridge Philos. Soc. **60** (1964), 769–778.
- [18] W. B. R. Lickorish, *On the homeomorphisms of a non-orientable surface*, Proc. Cambridge Philos. Soc. **61** (1965), 61–64.
- [19] F. Luo, *A presentation of the mapping class groups*, Math. Res. Lett. **4** (1997), no. 5, 735–739.
- [20] M. Matsumoto, *A presentation of mapping class groups in terms of Artin groups and geometric monodromy of singularities*, Math. Ann. **316** (2000), no. 3, 401–418.
- [21] G. Omori, *An infinite presentation for the mapping class group nonorientable surface*, Algebr. Geom. Topol. **17** (2017), no. 1, 419–437.
- [22] L. Paris and B. Szepietowski, *A presentation for the mapping class group of a nonorientable surface*, Bull. Soc. Math. France **143** (2015), no. 3, 503–566.
- [23] M. Stukow, *Dehn twists on nonorientable surfaces*, Fund. Math. **189** (2006), no. 2, 117–147.

- [24] M. Stukow, *Commensurability of geometric subgroups of mapping class groups*, Geom. Dedicata **143** (2009), 117–142.
- [25] M. Stukow, *A finite presentation for the mapping class group of a nonorientable surface with Dehn twists and one crosscap slide as generators*, J. Pure Appl. Algebra **218** (2014), no. 12, 2226–2239.
- [26] M. Stukow, *A finite presentation for the twist subgroup of the mapping class group of a nonorientable surface*, Bull. Korean Math. Soc. **53** (2016), no. 2, 601–614.
- [27] B. Szepietowski, *A presentation for the mapping class group of the closed non-orientable surface of genus 4*, J. Pure Appl. Algebra **213** (2009), no. 11, 2001–2016.
- [28] B. Wajnryb, *A simple presentation for the mapping class group of an orientable surface*, Israel J. Math. **45** (1983), no. 2-3, 157–174.

(*R. Kobayashi) DEPARTMENT OF GENERAL EDUCATION,
 NATIONAL INSTITUTE OF TECHNOLOGY, ISHIKAWA COLLEGE,
 TSUBATA, ISHIKAWA, 929-0392, JAPAN
Email address: kobayashi_ryoma@ishikawa-nct.ac.jp

(G. Omori) DEPARTMENT OF MATHEMATICS,
 FACULTY OF SCIENCE AND TECHNOLOGY,
 TOKYO UNIVERSITY OF SCIENCE,
 NODA, CHIBA, 278-8510, JAPAN
Email address: omori_genki@ma.noda.tus.ac.jp