

MASS OF ASYMPTOTICALLY FLAT 3-MANIFOLDS WITH BOUNDARY

SVEN HIRSCH, PENGZI MIAO, AND TIN-YAU TSANG

ABSTRACT. We study the mass of asymptotically flat 3-manifolds with boundary using the method of Bray-Kazaras-Khuri-Stern [6]. More precisely, we derive a mass formula on the union of an asymptotically flat manifold and fill-ins of its boundary, and give new sufficient conditions guaranteeing the positivity of the mass. Motivation to such consideration comes from studying the quasi-local mass of the boundary surface. If the boundary isometrically embeds in the Euclidean space, we apply the formula to obtain convergence of the Brown-York mass along large surfaces tending to ∞ which include the scaling of any fixed coordinate-convex surface.

1. INTRODUCTION

Given a complete asymptotically flat 3-manifold (M^3, g) , Bray, Kazaras, Khuri and Stern [6] gave a new proof of the three dimensional Riemannian positive mass theorem [28, 37]. More precisely, their result shows

$$(1.1) \quad \mathfrak{m}(g) \geq \frac{1}{16\pi} \int_{M_{ext}} \left(\frac{|\nabla^2 u|^2}{|\nabla u|} + R|\nabla u| \right),$$

where $\mathfrak{m}(g)$ is the total mass [2] of (M, g) , M_{ext} is the exterior region in (M, g) , R denotes the scalar curvature of the metric, and u is a harmonic function, satisfying a Neumann boundary condition at ∂M_{ext} , and is asymptotic to one of the asymptotically flat coordinate functions at infinity.

The method in [6] was inspired by the level sets approach to study scalar curvature via harmonic maps initiated by Stern [34] (also see Bray-Stern [5]). The method was further explored by Kazaras, Khuri and the first named author [16] to prove the three dimensional spacetime positive mass theorem [29] and produced a new lower bound on the total energy-momentum of initial data sets.

In this paper, we apply the method of Bray-Kazaras-Khuri-Stern [6] to analyze the mass of asymptotically flat 3-manifolds (M^3, g) with nonempty boundary Σ . In the case of M being diffeomorphic to $\mathbb{R}^3$ minus a ball, the mass of such manifolds is often used in the study of the quasi-local mass of Σ (see [4, 30, 36] for instance).

The following formula identifies the boundary terms of the integral in [6] that relates the mass and harmonic functions. (See Proposition 2.3 and Remark 2.1 in Section 2.)

Proposition 1.1. *Let (M, g) be an asymptotically flat 3-manifold with boundary Σ . Let u be a harmonic function on M which is asymptotic to one of the asymptotically flat coordinate functions at infinity. Let $\Sigma_t = u^{-1}(t)$ be the level set of u and ν be the infinity pointing unit*

P. Miao acknowledges the support of NSF Grant DMS-1906423.

normal to Σ . Then

$$\begin{aligned}
 (1.2) \quad & 8\pi \mathbf{m}(g) + 2\pi \int_{-T}^T (\chi(\Sigma_t) - 1) dt \\
 & \geq \int_M \frac{1}{2} \left(\frac{|\nabla^2 u|^2}{|\nabla u|} + R|\nabla u| \right) \\
 & \quad - \int_{\Sigma} H|\nabla u| + \int_{\{\nabla_{\Sigma} u \neq 0\}} \left\langle \nabla_{\Sigma} \beta, \frac{\nabla_{\Sigma} u}{|\nabla_{\Sigma} u|} \right\rangle |\nabla u|.
 \end{aligned}$$

Here $T > 0$ is some large constant so that the Euler characteristic $\chi(\Sigma_t) = 1$ for $|t| \geq T$; H is the mean curvature of Σ with respect to ν ; β is the angle between ∇u and Σ , which equals zero if $\frac{\partial u}{\partial \nu} = 0$; and operators Δ_{Σ} , ∇_{Σ} and ∇_{Σ}^2 denote the corresponding Laplacian, gradient, and Hessian on Σ .

Using (1.2), we derive a formula of the mass of (M, g) via fill-ins of its boundary Σ . We say a compact Riemannian 3-manifold (Ω, g_{Ω}) is a fill-in of Σ if $\partial\Omega$ is isometric to Σ .




FIGURE 1. An asymptotically flat manifold and a fill-in of its boundary

Throughout the paper, M and Ω are assumed to be orientable and metrics g and g_{Ω} are assumed to be C^2 up to the boundary.

Theorem 1.2. *Let (M, g) be an asymptotically flat 3-manifold with boundary Σ . Let (Ω, g_{Ω}) be a fill-in of Σ . Then there exist harmonic functions u, u_{Ω} on $(M, g), (\Omega, g_{\Omega})$ respectively, so that u is asymptotic to one of the asymptotically flat coordinate functions near infinity on (M, g) , u and u_{Ω} are C^2 up to the boundary, and along the boundary,*

$$(1.3) \quad u = u_{\Omega}, \quad \frac{\partial u}{\partial \nu} = \frac{\partial u_{\Omega}}{\partial \nu}.$$

Here ν denotes the ∞ -pointing unit normal to Σ in (M, g) , and by abuse of notation also the outward normal to $\partial\Omega$ in (Ω, g_{Ω}) . The equation is to be understood as two sides being equal under the isometry between Σ and $\partial\Omega$. If the topological manifold obtained by gluing M and Ω along their boundary contains no non-separating 2-spheres, then

$$\begin{aligned}
 (1.4) \quad \mathbf{m}(g) & \geq \frac{1}{16\pi} \left[\int_M \left(\frac{|\nabla^2 u|}{|\nabla u|} + R|\nabla u| \right) + \int_{\Omega} \left(\frac{|\nabla^2 u_{\Omega}|}{|\nabla u_{\Omega}|} + R|\nabla u_{\Omega}| \right) \right] \\
 & \quad + \frac{1}{8\pi} \int_{\Sigma} (H_{\Omega} - H)|\nabla u|.
 \end{aligned}$$

Here H, H_{Ω} are the mean curvature of $\Sigma, \partial\Omega$ in $(M, g), (\Omega, g_{\Omega})$ respectively, and $|\nabla u|$ denotes the common value of the length of the gradient of u and u_{Ω} at Σ and $\partial\Omega$.

Remark 1.1. Given (M^3, g) with boundary Σ , a fill-in of Σ modulo the smoothness at a point can be produced using (M, g) itself (see Section 3.5 for details). More generally, it is shown by Shi, Wang and Wei [32] that fill-ins with nonnegative scalar curvature always

exist for Σ . Related study of fill-ins with or without mean curvature constraint can be found in [17, 18, 22, 23, 32, 33].

Formula (1.4) is not surprising in the sense that the mean curvature difference $(H_\Omega - H)$ measures the distributional scalar curvature of the manifold $(N, \hat{g}) = (M, g) \cup (\Omega, g_\Omega)$ across $\Sigma = \partial\Omega$ (see [25, 30]). A proof of (1.4) then can be given by approximating $\hat{g}$ by smooth metrics produced in [25] and applying formula (1.1) to the approximation (see the appendix). In Section 3 below, we choose a more direct approach toward Theorem 1.2 by working with harmonic functions defined on the singular manifold $(N, \hat{g})$. By analyzing their regularity at the corner surface Σ , in Proposition 3.1 we show these functions are necessarily smooth on Σ , and hence smooth up to Σ from the two sides in $(N, \hat{g})$. As a result, except the mean curvature term, boundary terms in (1.2) from (M, g) are compensated by the corresponding boundary terms from (Ω, g_Ω) , which gives (1.4). We think this type of tangential regularity result can be useful in other PDE settings as well.

Formula (1.4) leads to a sufficient condition that guarantees positivity of $\mathbf{m}(g)$.

Corollary 1.3. *Let (M, g) and (Ω, g_Ω) be given as in Theorem 1.2. Suppose there exist vector fields X, Y on Ω, M , respectively, such that*

- a) $R \geq C_1|X|^2 - 2 \operatorname{div} X$ on Ω ,
- b) $R \geq C_2|Y|^2 - 2 \operatorname{div} Y$ on M with $Y = O(|x|^{-1-2\tau})$ for some $\tau > \frac{1}{2}$ and
- c) $H - \langle Y, \nu \rangle \leq H_\Omega - \langle X, \nu \rangle$.

Here $C_1, C_2 \geq \frac{2}{3}$ are some constants. Then $\mathbf{m}(g) \geq 0$. If $\mathbf{m}(g) = 0$ and $C_1, C_2 > \frac{2}{3}$, then $X = 0, Y = 0, \Sigma$ and $\partial\Omega$ have the same second fundamental forms, and the manifold $(M, g) \cup (\Omega, g_\Omega)$ obtained by gluing along $\Sigma = \partial\Omega$ is isometric to $\mathbb{R}^3$.

Remark 1.2. Motivations for allowing such vector fields on Ω and M come from the use of Jang's equation [29] in the context of quasi-local mass (see [20, 21, 35, 36] for instance). It is known the scalar curvature R of the graph of solutions to Jang's equation satisfies $R \geq 2|X|^2 - 2 \operatorname{div} X$ for a vector field X . We are recently informed by Aghil Alaei that results similar to $\mathbf{m}(g) \geq 0$ in Corollary 1.3 may also be derived via the method in [1].

If we make use of a fill-in of Σ produced by (M, g) itself, we obtain a result that only depends on (M, g) .

Corollary 1.4. *Let (M, g) be an asymptotically flat 3-manifold with boundary Σ . Let G be the harmonic function on (M, g) satisfying $G = 1$ at Σ and $G \rightarrow 0$ at ∞ . Let w be the harmonic function satisfying $w = 0$ at Σ and w is asymptotic to an asymptotically flat coordinate function at ∞ . Then there exists a harmonic function v on (M, g) such that $v \rightarrow 0$ at ∞ and*

$$(1.5) \quad v \frac{\partial G}{\partial \nu} - 2 \frac{\partial v}{\partial \nu} = \frac{\partial w}{\partial \nu} \text{ at } \Sigma.$$

Moreover, if (M, g) is diffeomorphic to $\mathbb{R}^3$ minus a ball,

$$(1.6) \quad \mathbf{m}(g) \geq \frac{1}{16\pi} \int_M \left[\frac{|\nabla^2 u|^2}{|\nabla u|^2} + R|\nabla u|^2 \right] - \frac{1}{4\pi} \int_\Sigma (H + 2 \frac{\partial G}{\partial \nu}) |\nabla u|^2.$$

Here $u = v + w$.

Remark 1.3. If $R \geq 0$ and $H \leq -2 \frac{\partial G}{\partial \nu}$, (1.6) implies $\mathbf{m}(g) \geq 0$, which is a special case of results in [15].

Given a surface Σ with mean curvature H in a 3-manifold (M, g) , if Σ has positive Gauss curvature, the Brown-York mass ([7, 8]) of Σ is given by

$$(1.7) \quad \mathbf{m}_{BY}(\Sigma) = \frac{1}{8\pi} \int_{\Sigma} (H_0 - H).$$

Here H_0 denotes the mean curvature of the isometric embedding of Σ in the Euclidean space $(\mathbb{R}^3, g_0)$.

In an asymptotically flat 3-manifold (M, g) , let Σ denote a large 2-sphere near infinity. It is a natural question to compare the Brown-York mass of Σ and the mass of (M, g) .




FIGURE 2. Brown-York mass of large surfaces in (M, g)

If $\{\Sigma_r\}$ is a family of large coordinate spheres, it was demonstrated in [12] that

$$(1.8) \quad \lim_{r \rightarrow \infty} \mathbf{m}_{BY}(\Sigma_r) = \mathbf{m}(g).$$

The same convergence property was shown for nearly round surfaces in [31] and for surfaces of revolution in asymptotically Schwarzschild manifold in [10, 11].

Theorem 1.2 suggests a natural approach to analyze the convergence of $\mathbf{m}_{BY}(\Sigma_r)$ for a more general class of surfaces. Let Ω_0 be the domain enclosed by Σ in $\mathbb{R}^3$. Then (Ω_0, g_0) provides a natural fill-in of the asymptotically end (E, g) , where E is the exterior of Σ in M . Applying (1.4) to the manifold $(N, \hat{g}) = (E, g) \cup (\Omega_0, g_0)$, one expects to relate $\mathbf{m}(g)$ to $\mathbf{m}_{BY}(\Sigma)$ by estimating the corresponding harmonic function. (In this case it can be shown that (1.4) is indeed an equation.) By implementing this idea, we show that

Theorem 1.5. *Let (M, g) be an asymptotically flat 3-manifold with a coordinate chart $\{x_i\}$ in which the metric coefficients g_{ij} satisfies the asymptotically flat condition. Let $\{\Sigma_r\}_{r \geq r_0}$ be a 1-parameter family of surfaces approaching infinity, where $r = \min_{\Sigma_r} \{|x| \mid x \in \Sigma_r\}$. Suppose the rescaled surfaces*

$$\tilde{\Sigma}_r := \left\{ \frac{1}{r}x \mid x \in \Sigma_r \right\} \subset \mathbb{R}^3$$

satisfy the property

$$(1.9) \quad 0 < k_1 < \kappa_\alpha < k_2, \quad \alpha = 1, 2,$$

where $\{\kappa_\alpha\}$ are the principal curvatures of $\tilde{\Sigma}_r$ in $\mathbb{R}^3$ with respect to the background Euclidean metric and k_1, k_2 are two positive constants independent of r . Then

$$(1.10) \quad \lim_{r \rightarrow \infty} \mathbf{m}_{BY}(\Sigma_r) = \mathbf{m}(g).$$

In particular, Theorem 1.5 applies to any family of surfaces that are obtained by scaling a fixed convex surface in the background coordinate space.

We end this introduction with a connection of (1.4) to the positivity of Brown-York mass. Suppose $\Omega \subset \mathbb{R}^3$ is a bounded domain with smooth boundary Σ . Let g be a metric on Ω such

that g and g_0 induce the same metric γ on Σ . Applying Theorem 1.2 to $(\mathbb{R}^3 \setminus \Omega, g_0) \cup (\Omega, g)$, one has

$$(1.11) \quad \frac{1}{8\pi} \int_{\Sigma} (H_0 - H_{\Omega}) |\nabla u| \geq \frac{1}{16\pi} \left[\int_{\mathbb{R}^3 \setminus \Omega} \frac{|\nabla^2 u|}{|\nabla u|} + \int_{\Omega} \left(\frac{|\nabla^2 u_{\Omega}|}{|\nabla u_{\Omega}|} + R |\nabla u_{\Omega}| \right) \right].$$

Here u and u_{Ω} are given by Theorem 1.2. In particular, u is harmonic on $(\mathbb{R}^3 \setminus \Omega, g_0)$, asymptotic to a coordinate function at infinity, and satisfies $\int_{\Sigma} \frac{\partial u}{\partial \nu} = 0$ at Σ .

If g has nonnegative scalar curvature, it follows that

$$(1.12) \quad \frac{1}{8\pi} \int_{\Sigma} (H_0 - H_{\Omega}) |\nabla u| \geq 0.$$

Comparing to the positivity of Brown-York mass ([30]), (1.12) has the drawback of having a weight function $|\nabla u|$ that is not explicitly determined by the boundary data of $(\Sigma, \gamma, H_{\Omega})$. On the other hand, (1.12) does not require an assumption on the mean curvature H_{Ω} and the Gauss curvature of γ .

The rest of the paper is organized as follows. In Section 2, we derive the relevant formulae on manifolds with boundary in the spirit of [6, 34]. In Section 3, we study regularity of harmonic functions on manifolds with corner along a hypersurface and prove Theorem 1.2, Corollary 1.3 and 1.4. In Section 4, we analyze the convergence of Brown-York mass using Theorem 1.2 and prove Theorem 1.5.

2. THE BOUNDARY FORMULAE

We start with a lemma.

Lemma 2.1. *Let U be an open set with piecewise smooth boundary ∂U in a given manifold. Let S be a smooth portion of ∂U . Let u be a harmonic function on U . Suppose u is C^2 up to S . Let ν denote the outward unit normal to S . Given a constant $\epsilon > 0$, let $\phi = \sqrt{|\nabla u|^2 + \epsilon}$. Then, at $p \in S$,*

- (1) $\partial_{\nu} \phi = 0$, if $\nabla u(p) = 0$;
- (2) $\partial_{\nu} \phi = -\frac{|\nabla u|^2}{\phi} H - \frac{\partial_{\nu} u}{\phi} \Delta_S \eta$, if $\nabla u(p) \neq 0$ and $\nabla_S \eta(p) = 0$;
- (3) $\partial_{\nu} \phi = -\frac{|\nabla u|^2}{\phi} H + \frac{|\nabla u|}{\phi} \kappa |\nabla_S \eta| + \frac{1}{\phi} \langle \nabla_S \partial_{\nu} u, \nabla_S \eta \rangle - \frac{\partial_{\nu} u}{\phi} |\nabla_S \eta|^{-2} \nabla_S^2 \eta (\nabla_S \eta, \nabla_S \eta)$, if $\nabla_S \eta(p) \neq 0$.

Here $\eta = u|_S$ is the restriction of u to S ; ∇_S , ∇_S^2 and Δ_S denote the gradient, the Hessian and the Laplacian on S ; H is the mean curvature of S with respect to ν ; and κ is the geodesic curvature of the level curve of η in the level surface of u with respect to the outward direction.

Proof. At p , we have

$$(2.1) \quad \partial_{\nu} \phi = \frac{1}{\phi} \langle \nabla_{\nu} \nabla u, \nabla u \rangle = \frac{1}{\phi} \nabla^2 u(\nu, \nabla u).$$

Thus, $\partial_{\nu} \phi = 0$ if $\nabla u = 0$ at p . In what follows, we assume $\nabla u(p) \neq 0$. Near p in S , let $n = \frac{\nabla u}{|\nabla u|}$. Then

$$(2.2) \quad \partial_{\nu} \phi = \frac{|\nabla u|}{\phi} \nabla^2 u(\nu, n).$$

Let $\theta \in [0, \pi]$ be the angle between ν and n . We have two cases:

Case 1. $\nabla_S \eta = 0$ at p . In this case, $\nabla u = (\partial_\nu u) \nu$, and

$$(2.3) \quad \partial_\nu \phi = \frac{\partial_\nu u}{\phi} \nabla^2 u(\nu, \nu) = \frac{\partial_\nu u}{\phi} (-H \partial_\nu u - \Delta_S \eta).$$

Here we have used the fact that $\Delta u = 0$ implies

$$(2.4) \quad \nabla^2 u(\nu, \nu) = -H \partial_\nu u - \Delta_S \eta.$$

We rewrite (2.3) as

$$(2.5) \quad \partial_\nu \phi = -\frac{|\nabla u|^2}{\phi} H - \frac{\partial_\nu u}{\phi} \Delta_S \eta.$$

Case 2. $\nabla_S \eta \neq 0$ at p . In this case, we introduce the following notations near p :

- $\tau_t = \eta^{-1}(t)$ and $\Sigma_t = u^{-1}(t)$;
- τ'_t denotes a unit tangent vector to the curve τ_t ;
- $n_t = \frac{1}{|\nabla_S \eta|} \nabla_S \eta$, which is a unit normal to τ_t in S ;
- ν_t denotes the outward unit normal to τ_t within Σ_t .

Along the curve τ_t , $\{\nu, n_t\}$ and $\{\nu_t, n_t\}$ both form an orthonormal basis of the normal bundle $\tau'_t{}^\perp$, and

$$(2.6) \quad \cos \theta = \langle \nu, n_t \rangle = \frac{\partial_\nu u}{|\nabla u|} \quad \text{and} \quad |\nabla_S \eta| = |\nabla u| \sin \theta.$$

We have

$$(2.7) \quad \begin{aligned} \partial_\nu \phi &= \frac{|\nabla u|}{\phi} \nabla^2 u(\nu, n) \\ &= \frac{|\nabla u|}{\phi} \nabla^2 u(\nu, n_t \sin \theta + \nu \cos \theta) \\ &= \frac{|\nabla u|}{\phi} \sin \theta \nabla^2 u(\nu, n_t) + \frac{|\nabla u|}{\phi} \cos \theta \nabla^2 u(\nu, \nu). \end{aligned}$$

Let $\mathbb{I}\mathbb{I}(v, w) = \langle \nabla_v \nu, w \rangle$ denote the second fundamental form of S with respect to ν . Then

$$(2.8) \quad \begin{aligned} \nabla^2 u(\nu, n_t) &= \langle \nabla_{n_t} \nabla u, \nu \rangle \\ &= \langle \nabla_{n_t} \nabla_S \eta, \nu \rangle + \langle \nabla_{n_t} (\partial_\nu u) \nu, \nu \rangle \\ &= -\mathbb{I}\mathbb{I}(\nabla_S \eta, n_t) + n_t(\partial_\nu u). \end{aligned}$$

Therefore,

$$(2.9) \quad \sin \theta \nabla^2 u(\nu, n_t) = -\mathbb{I}\mathbb{I}(n_t, n_t) |\nabla_S \eta| \sin \theta + n_t(\partial_\nu u) \sin \theta.$$

Thus,

$$(2.10) \quad \begin{aligned} \partial_\nu \phi &= \frac{|\nabla u|}{\phi} [-\mathbb{I}\mathbb{I}(n_t, n_t) |\nabla_S \eta| \sin \theta + n_t(\partial_\nu u) \sin \theta \\ &\quad + (-H \partial_\nu u - \Delta_S \eta) \cos \theta]. \end{aligned}$$

Here we also have used (2.4).

Now let κ denote the geodesic curvature of $\tau_t = \Sigma_t \cap S$ in Σ_t with respect to ν_t , that is

$$(2.11) \quad \kappa = \langle \nabla_{\tau'_t} \nu_t, \tau'_t \rangle = -\langle \nabla_{\tau'_t} \tau'_t, \nu_t \rangle$$

Using the fact $\nu_t = -\cos \theta n_t + \sin \theta \nu$, we have

$$\begin{aligned}
 \kappa &= -\langle \nabla_{\tau'_t} \tau'_t, -\cos \theta n_t + \sin \theta \nu \rangle \\
 (2.12) \quad &= \sin \theta \text{III}(\tau'_t, \tau'_t) + \cos \theta \langle \nabla_{\tau'_t} \tau'_t, n_t \rangle \\
 &= \sin \theta [H - \text{III}(n_t, n_t)] + \cos \theta \langle \nabla_{\tau'} \tau', n_t \rangle.
 \end{aligned}$$

Or equivalently,

$$(2.13) \quad -\text{III}(n_t, n_t) \sin \theta = \kappa - \sin \theta H - \cos \theta \langle \nabla_{\tau'} \tau', n_t \rangle.$$

It follows from (2.10), (2.13) and (2.6) that

$$\begin{aligned}
 \partial_\nu \phi &= \frac{|\nabla u|}{\phi} [(\kappa - \sin \theta H - \cos \theta \langle \nabla_{\tau'} \tau', n_t \rangle) |\nabla_S \eta| + n_t(\partial_\nu u) \sin \theta \\
 &\quad + (-H \partial_\nu u - \Delta_S \eta) \cos \theta] \\
 (2.14) \quad &= \frac{|\nabla u|}{\phi} [\kappa |\nabla_S \eta| - H |\nabla u| + n_t(\partial_\nu u) \sin \theta \\
 &\quad - (\Delta_S \eta + \langle \nabla_{\tau'} \tau', \nabla_S \eta \rangle) \cos \theta].
 \end{aligned}$$

Observe that

$$(2.15) \quad |\nabla u| n_t(\partial_\nu u) \sin \theta = \langle \nabla_S \partial_\nu u, \nabla_S \eta \rangle$$

and

$$(2.16) \quad \Delta_S \eta + \langle \nabla_{\tau'} \tau', \nabla_S \eta \rangle = \nabla_S^2 \eta(n_t, n_t).$$

We can rewrite (2.14) as

$$(2.17) \quad \partial_\nu \phi = \frac{|\nabla u|}{\phi} \kappa |\nabla_S \eta| - \frac{|\nabla u|^2}{\phi} H + \frac{1}{\phi} \langle \nabla_S \partial_\nu u, \nabla_S \eta \rangle - \frac{\partial_\nu u}{\phi} \nabla_S^2 \eta(n_t, n_t).$$

This finishes the proof. $\square$

The next proposition gives an analogue of Proposition 4.2 in [6] with no boundary condition imposed.

Proposition 2.2. *Let (Ω, g) be a 3-dimensional compact, oriented Riemannian manifold with smooth boundary $\partial\Omega$. Here $\partial\Omega$ is not necessarily connected. Let u be a harmonic function on Ω . Then*

$$\begin{aligned}
 (2.18) \quad &\int_\Omega \frac{1}{2} \left[\frac{|\nabla^2 u|^2}{|\nabla u|} + R |\nabla u| \right] - 2\pi \int_{u_{\min}}^{u_{\max}} \chi(\Sigma_t) dt \\
 &\leq - \int_{\partial\Omega} H |\nabla u| + \int_{\{\nabla_{\partial\Omega} u \neq 0\}} \left\langle \nabla_{\partial\Omega} \partial_\nu u, \frac{\nabla_{\partial\Omega} u}{|\nabla u|} \right\rangle - \frac{\partial_\nu u}{|\nabla u|} \nabla_{\partial\Omega}^2 u \left(\frac{\nabla_{\partial\Omega} u}{|\nabla_{\partial\Omega} u|}, \frac{\nabla_{\partial\Omega} u}{|\nabla_{\partial\Omega} u|} \right).
 \end{aligned}$$

Here R is the scalar curvature of g ; $\Sigma_t = u^{-1}(t)$ is the level set of u ; $\chi(\Sigma_t)$ denotes the Euler characteristic of Σ_t if Σ_t is a regular surface; $u_{\min} = \min_\Omega u$, $u_{\max} = \max_\Omega u$; and H is the mean curvature of $\partial\Omega$ with respect to the outward unit normal ν .

Proof. Given $\epsilon > 0$, let $\phi = \sqrt{|\nabla u|^2 + \epsilon}$. Integrating $\Delta\phi$ on Ω and applying Lemma 2.1, we have

$$(2.19) \quad \int_\Omega \Delta\phi = \int_{\partial\Omega} \partial_\nu \phi = \int_{\partial\Omega \cap \{\nabla u \neq 0\}} \partial_\nu \phi = I_1 + I_2,$$

where

$$(2.20) \quad I_1 = \int_{\{\nabla_{\partial\Omega} u=0, \partial_\nu u \neq 0\}} \partial_\nu \phi = \int_{\{\nabla_{\partial\Omega} u=0, \partial_\nu u \neq 0\}} -\frac{|\nabla u|^2}{\phi} H - \frac{\partial_\nu u}{\phi} \Delta_{\partial\Omega} u,$$

and

$$(2.21) \quad \begin{aligned} I_2 &= \int_{\{\nabla_{\partial\Omega} u \neq 0\}} \partial_\nu \phi \\ &= \int_{\{\nabla_{\partial\Omega} u \neq 0\}} -\frac{|\nabla u|^2}{\phi} H + \frac{|\nabla u|}{\phi} \kappa |\nabla_{\partial\Omega} u| + \frac{1}{\phi} \langle \nabla_{\partial\Omega} \partial_\nu u, \nabla_{\partial\Omega} u \rangle \\ &\quad - \frac{\partial_\nu u}{\phi} |\nabla_{\partial\Omega} u|^{-2} \nabla_{\partial\Omega}^2 u (\nabla_{\partial\Omega} u, \nabla_{\partial\Omega} u). \end{aligned}$$

Here $|\nabla u|$, H , $\Delta_{\partial\Omega} u$, $|\nabla_{\partial\Omega} \partial_\nu u|$, and $|\nabla_{\partial\Omega}^2 u|$ are all bounded quantities on $\partial\Omega$, independent on ϵ . Moreover, by item 3 in Lemma 2.1,

$$(2.22) \quad \frac{|\nabla u|}{\phi} \kappa |\nabla_{\partial\Omega} u| = \partial_\nu \phi + \frac{|\nabla u|^2}{\phi} H - \frac{1}{\phi} \langle \nabla_{\partial\Omega} \partial_\nu u, \nabla_{\partial\Omega} u \rangle + \frac{\partial_\nu u}{\phi} |\nabla_{\partial\Omega} u|^{-2} \nabla_{\partial\Omega}^2 u (\nabla_{\partial\Omega} u, \nabla_{\partial\Omega} u)$$

on $\{\nabla_{\partial\Omega} u \neq 0\} \subset \partial\Omega$. This, together with $|\partial_\nu \phi| \leq |\nabla \phi| \leq |\nabla^2 u|$, shows $\frac{|\nabla u|}{\phi} \kappa |\nabla_{\partial\Omega} u|$ is bounded by a constant independent on ϵ as well. Therefore, by dominated convergence theorem,

$$(2.23) \quad \begin{aligned} \lim_{\epsilon \rightarrow 0} \int_{\partial\Omega} \partial_\nu \phi &= \int_{\{\nabla_{\partial\Omega} u=0, \partial_\nu u \neq 0\}} -H |\nabla u| - \frac{\partial_\nu u}{|\nabla u|} \Delta_{\partial\Omega} u \\ &\quad + \int_{\{\nabla_{\partial\Omega} u \neq 0\}} -H |\nabla u| + \kappa |\nabla_{\partial\Omega} u| + \frac{1}{|\nabla u|} \langle \nabla_{\partial\Omega} \partial_\nu u, \nabla_{\partial\Omega} u \rangle \\ &\quad - \frac{\partial_\nu u}{|\nabla u|} |\nabla_{\partial\Omega} u|^{-2} \nabla_{\partial\Omega}^2 u (\nabla_{\partial\Omega} u, \nabla_{\partial\Omega} u). \end{aligned}$$

Combining the terms involving H and applying the coarea formula to the term involving κ , we can rewrite the above as

$$(2.24) \quad \begin{aligned} \lim_{\epsilon \rightarrow 0} \int_{\partial\Omega} \partial_\nu \phi &= - \int_{\partial\Omega} H |\nabla u| + \int_{u_{min}}^{u_{max}} \left(\int_{\{\nabla_{\partial\Omega} u \neq 0\} \cap \Sigma_t} \kappa \right) dt \\ &\quad + \int_{\{\nabla_{\partial\Omega} u=0, \partial_\nu u \neq 0\}} -\frac{\partial_\nu u}{|\nabla u|} \Delta_{\partial\Omega} u \\ &\quad + \int_{\{\nabla_{\partial\Omega} u \neq 0\}} \langle \nabla_{\partial\Omega} \partial_\nu u, \frac{\nabla_{\partial\Omega} u}{|\nabla u|} \rangle - \frac{\partial_\nu u}{|\nabla u|} \nabla_{\partial\Omega}^2 u \left(\frac{\nabla_{\partial\Omega} u}{|\nabla_{\partial\Omega} u|}, \frac{\nabla_{\partial\Omega} u}{|\nabla_{\partial\Omega} u|} \right). \end{aligned}$$

Under an assumption u is C^2 on Σ , elementary arguments shows the set

$$\{x \in \Sigma \mid \nabla_{\partial\Omega} u = 0, \Delta_{\partial\Omega} u \neq 0\}$$

is a set of measure zero in Σ . Hence, the integral in the middle line above vanishes and (2.24) simplifies to

$$(2.25) \quad \lim_{\epsilon \rightarrow 0} \int_{\partial\Omega} \partial_\nu \phi = - \int_{\partial\Omega} H |\nabla u| + \int_{u_{min}}^{u_{max}} \left(\int_{\{\nabla_{\partial\Omega} u \neq 0\} \cap \Sigma_t} \kappa \right) dt \\ + \int_{\{\nabla_{\partial\Omega} u \neq 0\}} \langle \nabla_{\partial\Omega} \partial_\nu u, \frac{\nabla_{\partial\Omega} u}{|\nabla u|} \rangle - \frac{\partial_\nu u}{|\nabla u|} \nabla_{\partial\Omega}^2 u \left(\frac{\nabla_{\partial\Omega} u}{|\nabla_{\partial\Omega} u|}, \frac{\nabla_{\partial\Omega} u}{|\nabla_{\partial\Omega} u|} \right).$$

Next, we analyze the interior terms following the argument of Theorem 1.1 in [34]. First, by Bochner's formula,

$$(2.26) \quad \Delta \phi = \frac{1}{\phi} \left[|\nabla^2 u|^2 + \text{Ric}(\nabla u, \nabla u) - \frac{1}{4\phi^2} |\nabla |\nabla u|^2|^2 \right] \\ \geq \phi^{-1} \text{Ric}(\nabla u, \nabla u).$$

Let $\mathcal{C}$ be the union of critical values of both $u : \Omega \rightarrow \mathbb{R}$ and $u|_{\partial\Omega} : \partial\Omega \rightarrow \mathbb{R}$. By Sard's theorem, $\mathcal{C}$ is a set of measure zero. Let A be an open set of $[u_{min}, u_{max}]$ so that A contains $\mathcal{C}$. Let B be the complement of A in $[u_{min}, u_{max}]$. By formula (14) in [34], along a regular level set Σ_t of u ,

$$(2.27) \quad \Delta \phi \geq \frac{1}{2\phi} [|\nabla^2 u|^2 + |\nabla u|^2 (R - 2K_\Sigma)],$$

where K_Σ is the Gauss curvature of Σ_t . Thus,

$$(2.28) \quad \int_{\Omega} \Delta \phi \geq \int_{u^{-1}(B)} \frac{1}{2\phi} [|\nabla^2 u|^2 + |\nabla u|^2 (R - 2K_\Sigma)] \\ - \max_{\Omega} |\text{Ric}| \int_{u^{-1}(A)} |\nabla u| dt.$$

As K_Σ is bounded by a constant independent on ϵ on the compact set $u^{-1}(B)$, passing to the limit in (2.28) gives

$$(2.29) \quad \lim_{\epsilon \rightarrow 0} \int_{\Omega} \Delta \phi \geq \int_{u^{-1}(B)} \frac{1}{2|\nabla u|} [|\nabla^2 u|^2 + |\nabla u|^2 (R - 2K_\Sigma)] \\ - \max_{\Omega} |\text{Ric}| \int_{u^{-1}(A)} |\nabla u| dt.$$

By coarea formula, this can be written as

$$(2.30) \quad \lim_{\epsilon \rightarrow 0} \int_{\Omega} \Delta \phi \geq \int_B \left(\int_{\Sigma_t} \frac{1}{2|\nabla u|^2} [|\nabla^2 u|^2 + |\nabla u|^2 R] \right) dt \\ - \int_B \left(\int_{\Sigma_t} K_\Sigma \right) dt - \left(\max_{\Omega} |\text{Ric}| \right) \int_A |\Sigma_t| dt.$$

As $t \in B$ is a regular value for both u and $u|_{\partial\Omega}$, Σ_t is a compact surface with boundary satisfying

$$\partial \Sigma_t = \Sigma_t \cap \partial\Omega = \Sigma_t \cap \{\nabla_{\partial\Omega} u \neq 0\}.$$

By the Gauss-Bonnet theorem,

$$(2.31) \quad \lim_{\epsilon \rightarrow 0} \int_{\Omega} \Delta \phi \geq \int_B \left(\int_{\Sigma_t} \frac{1}{2|\nabla u|^2} [|\nabla^2 u|^2 + |\nabla u|^2 R] \right) dt \\ - \int_B \left(2\pi \chi(\Sigma_t) - \int_{\Sigma_t \cap \{\nabla_{\partial\Omega} u \neq 0\}} \kappa \right) dt - \left(\max_{\Omega} |\text{Ric}| \right) \int_A |\Sigma_t| dt.$$

Since $\int_{u_{\min}}^{u_{\max}} |\Sigma_t| dt = \int_{\Omega} |\nabla u| < \infty$, (2.18) now follows from (2.25) and (2.31) by letting the measure of A tend to zero. $\square$

Remark 2.1. We learn from Daniel Stern that, at points $\nabla_{\partial\Omega} u \neq 0$, the integrand in the second boundary integral in (2.18) can be written as

$$(2.32) \quad \left\langle \nabla_{\partial\Omega} \partial_{\nu} u, \frac{\nabla_{\partial\Omega} u}{|\nabla u|} \right\rangle - \frac{\partial_{\nu} u}{|\nabla u|} \nabla_{\partial\Omega}^2 u \left(\frac{\nabla_{\partial\Omega} u}{|\nabla_{\partial\Omega} u|}, \frac{\nabla_{\partial\Omega} u}{|\nabla_{\partial\Omega} u|} \right) \\ = \left\langle \frac{|\nabla_{\partial\Omega} u| \nabla_{\partial\Omega} \partial_{\nu} u - \partial_{\nu} u \nabla_{\partial\Omega} |\nabla_{\partial\Omega} u|}{|\nabla u|^2}, \frac{\nabla_{\partial\Omega} u}{|\nabla_{\partial\Omega} u|} \right\rangle |\nabla u| \\ = \left\langle \nabla_{\partial\Omega} \beta, \frac{\nabla_{\partial\Omega} u}{|\nabla_{\partial\Omega} u|} \right\rangle |\nabla u|,$$

where $\beta = \frac{\pi}{2} - \theta$ is the angle between ∇u and the boundary surface $\partial\Omega$. It is intriguing if this interpretation can yield relevant geometric boundary conditions for harmonic functions.

Next we consider the mass of an asymptotically flat 3-manifold with boundary. Adopting conventions from [3], we say a Riemannian manifold (M^3, g) is asymptotically flat if there is a compact set $K \subset M$ and a diffeomorphism $\Psi : M \setminus K \rightarrow \mathbb{R}^3 \setminus B_r$, where $B_r = \{|x| \leq r\}$ for some r , such that

$$(2.33) \quad (\Psi^{-1})^*(g)_{ij} - \delta_{ij} \in W_{-\tau}^{k,p}(\mathbb{R}^3 \setminus B_r),$$

where $k \geq 2$, $p > 3$ and $\tau \in (\frac{1}{2}, 1]$. Here $W_{\delta}^{k,p}(\mathbb{R}^3 \setminus B_r)$ denotes the weighted Sobolev space consisting of functions $u \in W_{loc}^{k,p}(\mathbb{R}^3 \setminus B_r)$ such that

$$(2.34) \quad \|u\|_{W_{\delta}^{k,p}(\mathbb{R}^3 \setminus B_r)} = \sum_{|\beta| \leq k} \|\partial^{\beta} u |x|^{|\beta| - \delta - \frac{3}{p}}\|_{L^p(\mathbb{R}^3 \setminus B_r)} < \infty.$$

If the scalar curvature R of g is integrable, the ADM mass [2] of (M, g) is defined and given by

$$(2.35) \quad \mathfrak{m} = \lim_{r \rightarrow \infty} \frac{1}{16\pi} \int_{S_r} (g_{ij,i} - g_{ii,j}) v^j.$$

Here $S_r = \{|x| = r\}$ and v denotes the outward unit normal to S_r . The fact that $\mathfrak{m}$ is a geometric invariant of (M, g) , independent on the choice of the coordinates, was shown by Bartnik [3] and by Chruściel [9] independently.

Proposition 2.3. *Let (M^3, g) be an asymptotically flat 3-manifold with boundary Σ . Given a function η on Σ , let u be the harmonic function on M such that $u = \eta$ on Σ and u is asymptotic to some coordinate function x_i at the infinity. Let $\Sigma_t = u^{-1}(t)$ be the level set of u . Let $\mathfrak{m}$ be the mass of (M, g) and ν be the infinity pointing unit normal to Σ . Then*

$$\begin{aligned}
(2.36) \quad & 8\pi \mathbf{m} + 2\pi \int_{-T}^T (\chi(\Sigma_t) - 1) dt \\
& \geq \int_M \frac{1}{2} \left[\frac{|\nabla^2 u|^2}{|\nabla u|} + R|\nabla u| \right] - \int_{\Sigma} H|\nabla u| \\
& \quad + \int_{\{\nabla_{\Sigma} \eta \neq 0\}} \left\langle \nabla_{\Sigma} \partial_{\nu} u, \frac{\nabla_{\Sigma} \eta}{|\nabla u|} \right\rangle - \frac{\partial_{\nu} u}{|\nabla u|} \nabla_{\Sigma}^2 \eta \left(\frac{\nabla_{\Sigma} \eta}{|\nabla_{\Sigma} \eta|}, \frac{\nabla_{\Sigma} \eta}{|\nabla_{\Sigma} \eta|} \right).
\end{aligned}$$

Here $T > 0$ is some constant satisfying $\chi(\Sigma_t) = 1$ for all $|t| \geq T$, R is the scalar curvature of g , and H is the mean curvature of Σ with respect to ν .

Formula (2.36) is a corollary of the result of Bray-Kazaras-Khuri-Stern (Section 6 of [6]) and the preceding Proposition 2.2.

Proof. Suppose u is asymptotic to x_1 . By constructing two other harmonic functions u_2 and u_3 , asymptotic to x_2 and x_3 , respectively, we may well assume $u = y_1$ outside a compact set K , where $\{y_1, y_2, y_3\}$ are harmonic coordinates on $M \setminus K$ (see [3] for instance). This, in particular, implies that there exists a constant $T > 0$ such that $\chi(\Sigma_t) = 1$ for all $|t| \geq T_0$. For instance, let $D \subset M$ be a compact set such that $M \setminus D = \mathbb{R}^3 \setminus \{(y_1, y_2, y_3) \mid |y_i| \leq a, i = 1, 2, 3\}$ for some constant $a > 0$. Then, for any t with $|t| > \max\{a, \max_D |u|\}$, Σ_t is precisely the entire coordinate plane $\{y_1 = t\}$, hence $\chi(\Sigma_t) = 1$ for such t .

For each large constant $L > 0$, let S denote the cylindrical surface given by

$$S = \{y_1 = \pm L, y_2^2 + y_3^2 \leq L\} \cup \{|y_1| \leq L, y_2^2 + y_3^2 = L\}.$$

Let Ω be the region bounded by S and Σ . Let $\phi = \sqrt{|\nabla u|^2 + \epsilon}$ for $\epsilon > 0$. Though Ω does not have a smooth boundary, Stoke's theorem still holds on this Ω . Thus,

$$(2.37) \quad \int_S \partial_{\nu} |\nabla u| = \lim_{\epsilon \rightarrow 0} \int_S \partial_{\nu} \phi = \lim_{\epsilon \rightarrow 0} \int_{\Sigma} \partial_{\nu} \phi + \lim_{\epsilon \rightarrow 0} \int_{\Omega} \Delta \phi.$$

Here, by abuse of notations, ν also denotes the infinity pointing unit normal to S and we use the fact $|\nabla u| > 0$ at S for large L . For the other boundary term $\lim_{\epsilon \rightarrow 0} \int_{\Sigma} \partial_{\nu} \phi$ and the interior term $\lim_{\epsilon \rightarrow 0} \int_{\Omega} \Delta \phi$, we can apply (2.25) and (2.31), respectively, to have

$$\begin{aligned}
(2.38) \quad & \lim_{\epsilon \rightarrow 0} \int_{\Sigma} \partial_{\nu} \phi + \lim_{\epsilon \rightarrow 0} \int_{\Omega} \Delta \phi \\
& \geq \int_{\Omega} \frac{1}{2} \left[\frac{|\nabla^2 u|^2}{|\nabla u|} + R|\nabla u| \right] dt - \int_{\min_{\Omega} u}^{\max_{\Omega} u} 2\pi \chi(\Sigma_t) dt + \int_{\min_{\Omega} u}^{\max_{\Omega} u} \int_{\Sigma_t \cap S} \kappa dt \\
& \quad - \int_{\Sigma} H|\nabla u| - \int_{\{\nabla_{\Sigma} u = 0, \partial_{\nu} u \neq 0\}} \frac{\partial_{\nu} u}{|\nabla u|} \Delta_{\Sigma} u \\
& \quad + \int_{\{\nabla_{\Sigma} u \neq 0\}} \left\langle \nabla_{\Sigma} \partial_{\nu} u, \frac{\nabla_{\Sigma} u}{|\nabla u|} \right\rangle - \frac{\partial_{\nu} u}{|\nabla u|} \nabla_{\Sigma}^2 u \left(\frac{\nabla_{\Sigma} u}{|\nabla_{\Sigma} u|}, \frac{\nabla_{\Sigma} u}{|\nabla_{\Sigma} u|} \right).
\end{aligned}$$

For large L , observe that $\max_{\Omega} u = \max_S u = L$ and $\min_{\Omega} u = \min_S u = -L$. Hence, by (2.37) and (2.38),

$$\begin{aligned}
 (2.39) \quad & \int_S \partial_{\nu} |\nabla u| + \int_{-L}^L \left(2\pi \chi(\Sigma_t) - \int_{\Sigma_t \cap S} \kappa \right) dt \\
 & \geq \int_{\Omega} \frac{1}{2} \left[\frac{|\nabla^2 u|^2}{|\nabla u|} + R |\nabla u| \right] dt - \int_{\Sigma} H |\nabla u| - \int_{\{\nabla_{\Sigma} u = 0, \partial_{\nu} u \neq 0\}} \frac{\partial_{\nu} u}{|\nabla u|} \Delta_{\Sigma} u \\
 & \quad + \int_{\{\nabla_{\Sigma} u \neq 0\}} \left\langle \nabla_{\Sigma} \partial_{\nu} u, \frac{\nabla_{\Sigma} u}{|\nabla u|} \right\rangle - \frac{\partial_{\nu} u}{|\nabla u|} \nabla_{\Sigma}^2 u \left(\frac{\nabla_{\Sigma} u}{|\nabla_{\Sigma} u|}, \frac{\nabla_{\Sigma} u}{|\nabla_{\Sigma} u|} \right).
 \end{aligned}$$

Now recall the mass formula derived in (6.27) of [6], which asserts

$$(2.40) \quad \lim_{L \rightarrow \infty} \left[\int_S \partial_{\nu} |\nabla u| + \int_{-L}^L \left(2\pi - \int_{\Sigma_t \cap S} \kappa \right) dt \right] = 8\pi m.$$

Letting $L \rightarrow \infty$, (2.36) follows from (2.39) and (2.40). $\square$

3. MASS OF MANIFOLDS WITH A FILL-IN OF THE BOUNDARY

In this section, we use (E, g) , (Ω, g) to denote an asymptotically flat 3-manifold with boundary Σ , a compact Riemannian 3-manifold with boundary $\partial\Omega$, respectively. By abuse of notation, metrics on E and Ω are both denoted by g . We say (Ω, g) is a fill-in of Σ if $\partial\Omega$ is isometric to Σ , both of which are equipped with the induced metric. By a recent result of Shi, Wang and Wei [32], which answered a question of Gromov [14], a fill-in of Σ with nonnegative scalar curvature always exists.

Suppose (Ω, g) is a fill-in of (E, g) . Let U_+ and U_- be the Gaussian tubular neighborhoods of Σ in E and Ω , respectively, so that U_+ is diffeomorphic to $\Sigma \times [0, \epsilon)$ and U_- is diffeomorphic to $\Sigma \times (-\epsilon, 0]$. Identifying $U_+ \cup U_-$ with $\Sigma \times (-\epsilon, \epsilon)$ by lining up geodesics that are normal to Σ , we obtain a manifold (M, g) on which g is a continuous metric, which is smooth up to Σ from both sides. In particular, g is Lipschitz across Σ . As a result, by (2.33), g is a $W_{loc}^{1,p}$ metric on M satisfying $g_{ij} - \delta_{ij} \in W_{-\tau}^{1,p}(\mathbb{R}^3 \setminus B_r)$ for $p > 3$ at the end of E . By Theorem 3.1 in [3], there exists a function $u \in W_{loc}^{2,p}(M)$ such that $u - x_1 \in W_{1-\tau}^{2,p}(\mathbb{R}^3 \setminus B_r)$ and $\Delta u = 0$.

We want to apply Proposition 2.2 and Proposition 2.3 to analyze the mass of (M, g) with this choice of u .

3.1. Regularity of harmonic functions on the corner surface Σ . By Sobolev embedding, $u \in W_{loc}^{2,p}(M) \subset C_{loc}^{1,\alpha}(M)$, and hence $u|_{\Sigma} \in C^{1,\alpha}(\Sigma)$. Standard elliptic theory shows u is smooth away from Σ . In what follows, we examine the regularity of u on Σ .

Proposition 3.1. *If g is C^2 up to Σ on its two sides in M , then $u|_{\Sigma}$, the restriction of u to Σ , is in $C^{2,\alpha}(\Sigma)$, and u is $C^{2,\alpha}$ up to Σ on its two sides. Similarly, if g is C^∞ up to Σ , then $u|_{\Sigma}$ is in $C^\infty(\Sigma)$ and u is C^∞ up to Σ on its two sides.*

Proof. Consider a point $p \in \Sigma$. Using Fermi coordinates, we identify a neighborhood V of p in M with $\Sigma \times (-\epsilon, \epsilon)$, where coordinates on Σ are (x_1, x_2) and a coordinate in $(-\epsilon, \epsilon)$ is $x_3 = t$. Given a small constant h , consider the difference quotient of u along the direction of $e_\alpha = \partial_\alpha$, that is

$$(3.1) \quad \delta_\alpha^h u(x) = \frac{1}{h} [u(x + h e_\alpha) - u(x)], \quad \alpha \in \{1, 2\}.$$

It follows from $\Delta u = 0$ that

$$\begin{aligned}
 \Delta \delta_\alpha^h u(x) &= \frac{1}{h} \left[g^{ij}(x) \partial_{ij}^2 u(x + he_\alpha) - g^{ij}(x) \Gamma_{ij}^k(x) \partial_k u(x + he_\alpha) \right] \\
 (3.2) \quad &= \frac{1}{h} \left[[g^{ij}(x) - g^{ij}(x + he_\alpha)] \partial_{ij}^2 u(x + he_\alpha) \right. \\
 &\quad \left. - [g^{ij}(x) \Gamma_{ij}^k(x) - g^{ij}(x + he_\alpha) \Gamma_{ij}^k(x + he_\alpha)] \partial_k u(x + he_\alpha) \right].
 \end{aligned}$$

Since $g_{ij} \in C_{loc}^{0,1}(V)$,

$$(3.3) \quad \frac{g^{ij}(x) - g^{ij}(x + he_\alpha)}{h}, \Gamma_{ij}^k(x), \Gamma_{ij}^k(x + he_\alpha) \in L_{loc}^\infty(V).$$

Since g is C^2 up to Σ from both sides of Σ , g_{ij} and ∂g_{ij} are locally Lipschitz along the ∂_α -direction in V , therefore

$$(3.4) \quad \frac{1}{h} \left[\Gamma_{ij}^k(x) - \Gamma_{ij}^k(x + he_\alpha) \right] \in L_{loc}^\infty(V).$$

These show that the right side of (3.2) is in $L_{loc}^p(V)$. By Theorem 9.11 in [13], $\|\delta_\alpha^h u\|_{W^{2,p}(\tilde{V})}$ is bounded by a constant independent on h , on any subdomain $\tilde{V} \subset\subset V$. This readily implies $\partial_\alpha u = \lim_{h \rightarrow 0} \delta_\alpha^h u \in W_{loc}^{2,p}(V)$. By Sobolev embedding, $\partial_\alpha u \in C_{loc}^{1,\alpha}(V)$. In particular, this shows $u|_\Sigma \in C^{2,\alpha}(\Sigma)$.

Next, we show that this argument can be iterated to show $u|_\Sigma$ is smooth in the case that g is smooth up to Σ from both sides. By integrating again a test function, (3.2) implies

$$(3.5) \quad \Delta \partial_\alpha u = -\partial_\alpha g^{ij} \partial_{ij}^2 u + \left(\partial_\alpha g^{ij} \Gamma_{ij}^k + g^{ij} \partial_\alpha \Gamma_{ij}^k \right) \partial_k u.$$

Therefore,

$$\begin{aligned}
 \Delta \delta_\beta^h \partial_\alpha u &= -\delta_\beta^h \partial_\alpha g^{ij} \partial_{ij}^2 u - \tau_\alpha^h \partial_\alpha g^{ij} \delta_\beta^h \partial_{ij}^2 u \\
 (3.6) \quad &+ \delta_\beta^h \left(\partial_\alpha g^{ij} \Gamma_{ij}^k + g^{ij} \partial_\alpha \Gamma_{ij}^k \right) \partial_k u \\
 &+ \tau_\beta^h \left(\partial_\alpha g^{ij} \Gamma_{ij}^k + g^{ij} \partial_\alpha \Gamma_{ij}^k \right) \delta_\beta^h \partial_k u.
 \end{aligned}$$

Here τ_β^h is the operator of shifting the coordinate along e_β by h , i.e. $\tau_\beta^h f(x) = f(x + he_\beta)$. Recall that, on any subdomain $\tilde{V} \subset\subset V$, the preceding step has shown $\|\delta_\beta^h u\|_{W^{2,p}(\tilde{V})} \leq C$, where C is independent on h . This, combined with the fact $\delta_\beta^h \partial_{ij}^2 u = \partial_{ij}^2 \delta_\beta^h u$ and the assumption that g is smooth up to Σ on both sides, then shows the right side of (3.6) is uniformly bounded in $L^p(\tilde{V})$. Hence, by Theorem 9.11 in [13], $\|\delta_\beta^h \partial_\alpha u\|_{W^{2,p}(\tilde{V})}$ is uniformly bounded and consequently $\partial_{\beta\alpha}^2 u \in W_{loc}^{2,p}(V)$. By Sobolev embedding, $\partial_{\beta\alpha}^2 u \in C_{loc}^{1,\alpha}(V)$, and this implies $u|_\Sigma \in C^{3,\alpha}(\Sigma)$. Repeating these arguments, we conclude $u|_\Sigma \in C^\infty(\Sigma)$.

The remaining claim that u is $C^{2,\alpha}$ (C^∞) up to Σ on the two sides of Σ follows directly from the standard elliptic boundary regularity theory, see Theorem 6.19 in [13]. $\square$

Remark 3.1. Proposition 3.1 is a local result that holds for any $W^{2,p}$ harmonic function in a neighborhood of the corner hypersurface Σ . It continues to hold in higher dimensions as the proof does not require Σ to be 2-dimensional. The proof technique also works for more general linear, elliptic PDEs on manifolds with corner type singularity.

Remark 3.2. Let H and H_Ω be the mean curvature of Σ in (E, g) and (Ω, g) , respectively. Evaluating $\Delta u = 0$ along Σ in its both sides in (M, g) , one has

$$0 = \Delta_\Sigma u + H \frac{\partial u}{\partial t} + \frac{\partial^2 u}{\partial t^2} \quad \text{and} \quad 0 = \Delta_\Sigma u + H_\Omega \frac{\partial u}{\partial t} + \frac{\partial^2 u}{\partial t^2},$$

where $\Delta_\Sigma u$ is the Laplacian of $u|_\Sigma$ on Σ and $\partial_t = \nu$ is the ∞ -pointing unit normal in (M, g) . This implies, if $H = H_\Omega$, u is C^2 across Σ in the given coordinates $\{x_\alpha, t\}$.

3.2. Mass of manifolds with corner surface. We analyze the mass of (M, g) following the method of Bray-Kazaras-Khrui-Stern [6]. For this purpose, we impose the following topological assumption on $M = E \cup \Omega$:

Condition (T): Assume M contains no non-separating two-spheres.

As $u \in W_{loc}^{2,p}(M) \subset C^1(M)$, $\Sigma_t = u^{-1}(t)$ is C^1 surface for each regular value t . Moreover, u satisfies the maximum principle (see Theorem 8.1 in [13] for instance). Thus, it follows from condition (T) and the maximum principle that Σ_t consists of a single non-compact connected component, together with some compact components (possibly empty) that are closed surfaces with positive genus. Therefore, $\chi(\Sigma_t) \leq 1$ (cf. Section 6 in [6]).

By Proposition 3.1, we can apply Proposition 2.3 and Proposition 2.2 to (E, g) and (Ω, g) , respectively, to obtain

$$\begin{aligned} (3.7) \quad & 8\pi\mathfrak{m} + 2\pi \int_{-T}^T (\chi(\Sigma_{t,E}) - 1) dt \\ & \geq \int_E \frac{1}{2} \left[\frac{|\nabla^2 u|^2}{|\nabla u|} + R|\nabla u| \right] - \int_\Sigma H_E |\nabla u| \\ & \quad + \int_{\{\nabla_\Sigma u \neq 0\}} \left\langle \nabla_\Sigma \partial_\nu u, \frac{\nabla_\Sigma u}{|\nabla u|} \right\rangle - \frac{\partial_\nu u}{|\nabla u|} \nabla_\Sigma^2 u \left(\frac{\nabla_\Sigma u}{|\nabla_\Sigma u|}, \frac{\nabla_\Sigma u}{|\nabla_\Sigma u|} \right) \end{aligned}$$

and

$$\begin{aligned} (3.8) \quad & 2\pi \int_{\min_\Omega u}^{\max_\Omega u} \chi(\Sigma_{t,\Omega}) dt \\ & + \int_{\{\nabla_\Sigma u \neq 0\}} \left\langle \nabla_\Sigma \partial_\nu u, \frac{\nabla_\Sigma u}{|\nabla u|} \right\rangle - \frac{\partial_\nu u}{|\nabla u|} \nabla_\Sigma^2 u \left(\frac{\nabla_\Sigma u}{|\nabla_\Sigma u|}, \frac{\nabla_\Sigma u}{|\nabla_\Sigma u|} \right) \\ & \geq \int_\Omega \frac{1}{2} \left[\frac{|\nabla^2 u|^2}{|\nabla u|} + R|\nabla u| \right] + \int_\Sigma H_\Omega |\nabla u|. \end{aligned}$$

Here t is chosen to be a regular value of u and $u|_\Sigma$, $\Sigma_{t,E} = \Sigma_t \cap E$, $\Sigma_{t,\Omega} = \Sigma_t \cap \Omega$. Since u is C^1 on M and C^2 on Σ , adding (3.7) and (3.8) gives

$$\begin{aligned} (3.9) \quad & 8\pi\mathfrak{m} + 2\pi \int_{-T}^T [\chi(\Sigma_{t,E}) + \chi(\Sigma_{t,\Omega}) - 1] dt \\ & \geq \int_E \frac{1}{2} \left[\frac{|\nabla^2 u|^2}{|\nabla u|} + R|\nabla u| \right] + \int_\Omega \frac{1}{2} \left[\frac{|\nabla^2 u|^2}{|\nabla u|} + R|\nabla u| \right] + \int_\Sigma (H_\Omega - H_E) |\nabla u|. \end{aligned}$$

Note that, for each chosen t , $\Sigma_{t,E} \cup \Sigma_{t,\Omega} = \Sigma_t$ if $\Sigma_t \cap \Omega \neq \emptyset$; and $\Sigma_{t,E} = \Sigma_t$ if $\Sigma_t \cap \Omega = \emptyset$. In either case, $\chi(\Sigma_{t,E}) + \chi(\Sigma_{t,\Omega}) = \chi(\Sigma_t) \leq 1$. Therefore, it follows from (3.9) that

$$(3.10) \quad 8\pi\mathfrak{m} \geq \int_E \frac{1}{2} \left[\frac{|\nabla^2 u|^2}{|\nabla u|} + R|\nabla u| \right] + \int_\Omega \frac{1}{2} \left[\frac{|\nabla^2 u|^2}{|\nabla u|} + R|\nabla u| \right] + \int_\Sigma (H_\Omega - H_E) |\nabla u|,$$

and equality holds only if $\chi(\Sigma_t) = 1$ for a.e. t . This proves Theorem 1.2.

Remark 3.3. If the gradient $|\nabla u|$ never vanishes on M , it is clear from its proof that equality holds in (3.10).

Remark 3.4. The proof allows E and Ω to have additional boundary components if those components have zero mean curvature. In this case, one can impose a Neumann boundary condition at the minimal boundary portion (see [6]) or impose a suitable Dirichlet boundary condition (see [16]).

3.3. Positivity of the mass. We give a proof of Corollary 1.3 using (1.4). We keep the same notations as in the previous subsection, that is E denotes the asymptotically flat piece, and metrics on both E and Ω are denoted by g .

Suppose there are vector fields X, Y on Ω, E , respectively, satisfying

$$(3.11) \quad R \geq \frac{2}{3}|X|^2 - 2 \operatorname{div} X \quad \text{and} \quad R \geq \frac{2}{3}|Y|^2 - 2 \operatorname{div} Y,$$

where $Y = O(|x|^{-1-2\tau})$ for some $\tau > \frac{1}{2}$. Here we consider the case $C_1 = C_2 = \frac{2}{3}$.

Let (M, g) , u be the glued manifold, the harmonic function as in the preceding proof of Theorem 1.2. On (Ω, g) , $|\nabla u|$ is Lipschitz, integrating by parts gives

$$(3.12) \quad \begin{aligned} & \int_{\Omega} \frac{1}{2} \left(\frac{1}{|\nabla u|} |\nabla^2 u|^2 + R |\nabla u| \right) \\ & \geq \int_{\Omega} \frac{1}{2|\nabla u|} |\nabla^2 u|^2 + \frac{1}{3} |X|^2 |\nabla u| + \langle X, \nabla |\nabla u| \rangle - \int_{\Sigma} \langle X, \nu \rangle |\nabla u|. \end{aligned}$$

To estimate the integral on Ω , we use a refined Kato inequality for harmonic functions (c.f. Lemma 3.1 in [16] for instance).

Lemma 3.2. *If u is a harmonic function on a Riemannian manifold (U^n, g) , then*

$$(3.13) \quad |\nabla^2 u|^2 \geq \frac{n}{n-1} |\nabla |\nabla u||^2 \quad \text{on } U \setminus D.$$

Here $D \subset \{\nabla u = 0\}$ is a set of measure zero at which $|\nabla u|$ is not differentiable. If equality holds at some $p \in U \setminus D$, then

- i) $\nabla^2 u(p) = 0$ if $\nabla u(p) = 0$; and
- ii) $\nabla^2 u(p) = \frac{n}{n-1} \nabla^2 u(\nu, \nu)(p) \left(\nu \otimes \nu - \frac{1}{n} g \right)(p)$, if $\nabla u(p) \neq 0$. Here $\nu = \frac{\nabla u(p)}{|\nabla u(p)|}$ and $\nu \otimes \nu$ is understood by $\nu \otimes \nu(v, w) = \langle \nu, v \rangle \langle \nu, w \rangle$ for any $v, w \in T_p U$.

Proof. Let $u_{;i} = \partial_i u$ and $Z = \frac{1}{2} \nabla |\nabla u|^2$, then

$$(3.14) \quad \begin{aligned} |Z|^2 &= \frac{1}{2} (u_{;j} Z_i + u_{;i} Z_j) (\nabla^2 u)^{ij} \\ &= \left[\frac{1}{2} (u_{;j} Z_i + u_{;i} Z_j) - \frac{1}{n} \nabla^2 u (\nabla u, \nabla u) g_{ij} \right] (\nabla^2 u)^{ij}, \end{aligned}$$

where $\Delta u = 0$ was used in the second step. Define

$$(3.15) \quad W_{ij} = \frac{1}{2} (u_{;j} Z_i + u_{;i} Z_j) - \frac{1}{n} \nabla^2 u (\nabla u, \nabla u) g_{ij},$$

then $g^{ij}W_{ij} = 0$ and

$$(3.16) \quad \begin{aligned} |W|^2 &= \frac{1}{2}|Z|^2|\nabla u|^2 + \left(\frac{1}{2} - \frac{1}{n}\right) \langle Z, \nabla u \rangle^2 \\ &\leq \frac{n-1}{n}|Z|^2|\nabla u|^2. \end{aligned}$$

By (3.14) and (3.16),

$$(3.17) \quad |Z|^2 = \langle W, \nabla^2 u \rangle \leq \sqrt{\frac{n-1}{n}}|Z||\nabla u||\nabla^2 u|,$$

which gives $|Z| \leq \sqrt{\frac{n-1}{n}}|\nabla u||\nabla^2 u|$. This readily shows

$$(3.18) \quad |\nabla|\nabla u||^2 \leq \frac{n-1}{n}|\nabla^2 u|^2, \quad \text{on } U \setminus D.$$

Suppose equality in (3.18) holds at some $p \in U \setminus D$. If $\nabla u(p) = 0$, then $\nabla|\nabla u|(p) = 0$ which shows $\nabla^2 u(p) = 0$.

Next suppose $\nabla u(p) \neq 0$. From the proof of (3.18), we have

$$(3.19) \quad \langle Z, \nabla u \rangle^2(p) = |Z(p)|^2|\nabla u(p)|^2 \quad \text{and} \quad \langle W, \nabla^2 u \rangle(p) = |W(p)||\nabla^2 u(p)|.$$

If $\nabla^2 u(p) = 0$, ii) holds automatically. Suppose $\nabla^2 u(p) \neq 0$. In this case, $Z(p) \neq 0$. For otherwise $\nabla|\nabla u|(p) = 0$ which would imply $\nabla^2 u(p) = 0$ by the equality in (3.18). Thus, (3.19) is equivalent to

$$(3.20) \quad Z(p) = \phi(p)\nabla u(p) \quad \text{and} \quad W(p) = \psi(p)\nabla^2 u(p),$$

where $\phi(p)$ and $\psi(p)$ are some numbers with $\phi(p) \neq 0$ and $\psi(p) \geq 0$.

By the definition of W and the fact $g^{ij}W_{ij} = 0$, (3.20) gives $\phi(p) = \nabla^2 u(\nu, \nu)(p)$ and

$$(3.21) \quad \phi(p) \left(du \otimes du - \frac{1}{n}|\nabla u|^2 g \right)(p) = \psi(p)\nabla^2 u(p).$$

Evaluating (3.21) along (ν, ν) shows $\psi(p) = \frac{n-1}{n}|\nabla u(p)|^2$. Part ii) follows. $\square$

Coming back to (3.12), by Lemma 3.2, we have

$$(3.22) \quad \begin{aligned} &\int_{\Omega} \frac{1}{2|\nabla u|}|\nabla^2 u|^2 + \frac{1}{3}|X|^2|\nabla u| + \langle X, \nabla|\nabla u| \rangle \\ &\geq \int_{\Omega} \left(\frac{\sqrt{3}}{2\sqrt{|\nabla u|}}|\nabla|\nabla u|| - \frac{1}{\sqrt{3}}|X|\sqrt{|\nabla u|} \right)^2 + |\nabla|\nabla u|||X| + \langle X, \nabla|\nabla u| \rangle \\ &\geq 0. \end{aligned}$$

Similarly, on E ,

$$(3.23) \quad \begin{aligned} &\int_E \frac{1}{2|\nabla u|}|\nabla^2 u|^2 + \frac{1}{3}|Y|^2|\nabla u| + \langle Y, \nabla|\nabla u| \rangle \\ &\geq \int_E \left(\frac{\sqrt{3}}{2\sqrt{|\nabla u|}}|\nabla|\nabla u|| - \frac{1}{\sqrt{3}}|Y|\sqrt{|\nabla u|} \right)^2 + |\nabla|\nabla u|||Y| + \langle Y, \nabla|\nabla u| \rangle \\ &\geq 0. \end{aligned}$$

Thus, by (3.10), (3.12), and assumption c),

$$(3.24) \quad 8\pi\mathfrak{m} \geq \int_{\Sigma} (H_{\Omega} - \langle X, \nu \rangle - H + \langle Y, \nu \rangle) |\nabla u| \geq 0.$$

3.4. The zero mass case. Suppose $\mathfrak{m} = 0$ in the above context. Tracing all inequalities, we have

$$(3.25) \quad \chi(\Sigma_t) = 1, \text{ a.e. } t \in \mathbb{R},$$

$$(3.26) \quad \left(R - \frac{2}{3}|X|^2 + 2 \operatorname{div} X \right) |\nabla u| = 0, \text{ on } \Omega,$$

$$(3.27) \quad \frac{1}{|\nabla u|} \left(|\nabla^2 u|^2 - \frac{3}{2} |\nabla |\nabla u||^2 \right) = 0, \text{ a.e. on } \Omega,$$

$$(3.28) \quad \frac{\sqrt{3}}{2\sqrt{|\nabla u|}} |\nabla |\nabla u|| = \frac{1}{\sqrt{3}} |X| \sqrt{|\nabla u|}, \text{ a.e. on } \Omega$$

and

$$(3.29) \quad |\nabla |\nabla u|| |X| + \langle X, \nabla |\nabla u| \rangle = 0, \text{ a.e. on } \Omega.$$

Analogous relations hold for R , u and Y on E .

To show the rigidity case in Corollary 1.3, we assume

$$R \geq C_1 |X|^2 - 2 \operatorname{div} X \text{ on } \Omega$$

for some constant $C_1 > \frac{2}{3}$. Under this assumption, (3.26) implies $|X||\nabla u| = 0$ on Ω . By (3.28), $\nabla |\nabla u| = 0$ a. e. on Ω , and hence $|\nabla u|$ is a constant on Ω . Similarly, if

$$R \geq C_2 |Y|^2 - 2 \operatorname{div} Y \text{ on } E$$

for some $C_2 > \frac{2}{3}$, then $|\nabla u|$ is a constant on E . Consequently, $|\nabla u|$ equals the same constant on E and Ω since $|\nabla u|$ is C^0 on M . As $|\nabla u| \rightarrow 1$ near infinity, we have $|\nabla u| = 1$ on M . As a result, $X = 0$, $Y = 0$, $R \geq 0$ and $H_{\Omega} \geq H$. Theorem 1.2 and $\mathfrak{m} = 0$ then imply $\nabla^2 u = 0$, $R = 0$ and $H_{\Omega} = H$.

Let u_i denote the harmonic function u with x_1 replaced by x_i , $i = 1, 2, 3$. The preceding proof shows that $\{\nabla u_1, \nabla u_2, \nabla u_3\}$, being linearly independent near infinity, are parallel on (E, g) and (Ω, g) . Hence, (E, g) and (Ω, g) are flat.

We claim that Σ has the same second fundamental form in (E, g) and (Ω, g) . To see this, on each side of Σ in (M, g) , we have

$$(3.30) \quad 0 = \nabla^2 u_i(\partial_{\alpha}, \partial_{\beta}) = \nabla_{\Sigma}^2 u_i(\partial_{\alpha}, \partial_{\beta}) + \mathbb{I}(\partial_{\alpha}, \partial_{\beta}) \frac{\partial u_i}{\partial \nu},$$

where ∇_{Σ}^2 is the Hessian on Σ and $\mathbb{I}$ is the corresponding second fundamental form of Σ . As $\{\nabla u_1, \nabla u_2, \nabla u_3\}$ form a basis for $T_p M$ at each $p \in \Sigma$, one of u_i satisfies

$$(3.31) \quad \frac{\partial u_i}{\partial \nu}(p) \neq 0.$$

It follows from (3.30) and (3.31) that $\mathbb{I}$ are the same from both sides of Σ in (M, g) . This combined with the fact g is flat near Σ implies that g is C^2 across Σ (see [30] for instance). On the other hand, $H_{\Omega} = H$ implies u is C^2 across Σ by Remark 3.2. The smoothness of u is further promoted to be $C^{2,\alpha}$ via the regularity of g . We then proceed as in [6] to conclude that the map $F = (u_1, u_2, u_3)$ is a $C^{2,\alpha}$ isometry between (M, g) and $\mathbb{R}^3$.

Remark 3.5. In the setting of Theorem 1.2, Corollary 1.3 establishes the positive mass theorem with corner along Σ (c.f. [25, 30, 24]). However, it does not seem to be a replacement of the later results due to the dimension restriction and the topological condition (T).

The preceding proof does not make use of (3.25), (3.27) and (3.29). Exploring these relations, we can prove a rigidity result for the case $C_1 = C_2 = \frac{2}{3}$, but under a stronger regularity assumption of g across Σ .

Theorem 3.3. *Let (M, g) be a smooth asymptotically flat 3-manifold, without boundary, which contains no non-separating 2-spheres. Let $\Omega \subset M$ be a bounded domain with smooth boundary Σ . Suppose there exist vector fields X, Y on $\bar{\Omega}, M \setminus \Omega$, respectively, such that*

- a) $R \geq \frac{2}{3}|X|^2 - 2 \operatorname{div} X$ on Ω ,
- b) $R \geq \frac{2}{3}|Y|^2 - 2 \operatorname{div} Y$ on $M \setminus \Omega$ with $Y = O(|x|^{-1-2\tau})$, and
- c) $\langle Y, \nu \rangle \geq \langle X, \nu \rangle$, where ν is the ∞ -pointing unit normal to Σ .

Then $\mathbf{m} = 0$ implies (M, g) is isometric to $\mathbb{R}^3$.

Proof. As we assume g is smooth, the function u is smooth. Hence each regular level set $\Sigma_t = u^{-1}(t)$ is a surface with a smooth metric induced from g .

Suppose $\mathbf{m} = 0$. It suffices to show $\nabla^2 u = 0$ on M . We first note

$$(3.32) \quad \nabla^2 u = 0, \text{ a.e. on } \{x \in M \mid \nabla u(x) = 0\}.$$

This is because (1.4) implies $\int_M \frac{|\nabla^2 u|}{|\nabla u|} < \infty$, as R is integrable and $|\nabla u|$ is bounded.

Next, consider a set

$$A = \{t \in \mathbb{R} \mid t \text{ is a regular value of both } u \text{ and } u|_{\Sigma}, \text{ and } \chi(\Sigma_t) = 1\}.$$

By (3.25), the complement of A in $\mathbb{R}$ is a set of measure zero. The following is the main assertion in this proof:

$$(3.33) \quad \nabla^2 u(x) = 0, \forall x \in \Sigma_t \text{ where } t \in A.$$

If (3.33) holds, then $\nabla^2 u = 0$ on M follows. To see this, consider an open set

$$B = \{x \in M \mid \nabla^2 u(x) \neq 0\} \cap \{x \in M \mid \nabla u(x) \neq 0\}.$$

Let $p \in B$ if $B \neq \emptyset$. Consider a curve $\tau : (-\epsilon, \epsilon) \rightarrow B$, for some small $\epsilon > 0$, with $\tau(0) = p$ and $\tau'(0) = \nabla u(p)$. As $\tau(s) \in B$ for $s \in (-\epsilon, \epsilon)$, (3.33) implies $u(\tau(s)) \notin A$. As a result, $\{u(\tau(s)) \mid s \in (-\epsilon, \epsilon)\}$ is a subset of measure zero in $\mathbb{R}$. However, this set contains an open neighborhood of $u(p)$ because $\frac{d}{ds}|_{s=0} u(\tau(s)) > 0$. This is a contradiction. Therefore $B = \emptyset$. Considering (3.32), we then conclude $\nabla^2 u = 0$ on M .

The rest of the proof is to show (3.33). We start with some local information near a connected surface S on which u is a constant with $\nabla u \neq 0$. By (3.27) and Lemma 3.2, along S , u satisfies

$$(3.34) \quad \nabla^2 u = \frac{3}{2} \nabla^2 u(\nu, \nu) \left(\nu \otimes \nu - \frac{1}{3} g \right), \text{ where } \nu = \frac{\nabla u}{|\nabla u|}.$$

This implies

- $|\nabla u|$ is a constant on S ; and
- $|\nabla u| \text{III} = -\frac{1}{2} \nabla^2 u(\nu, \nu) \gamma$, where III is the second fundamental form of S with respect to ν and γ is the metric on S induced from g . As a result, the mean curvature H of S satisfies $H|\nabla u| = -\nabla^2 u(\nu, \nu)$.

To get more information, we can consider the level set foliation of u near S . More precisely, let U be a bounded, connected open set in S and let W be an open neighborhood of U in M obtained by flowing U along the integral flow of $|\nabla u|^{-2}\nabla u$. W is diffeomorphic to $(-\delta, \delta) \times U$ for some $\delta > 0$ and the metric g on W can be written as

$$(3.35) \quad g = \frac{1}{f(s)^2} ds^2 + \gamma_s.$$

Here $f = |\nabla u|$ and γ_s is the induced metric on $\{s\} \times U$, which is part of the level set Σ_{t+s} . For convenience, we may assume $u = 0$ on S . Then $u = s$ on W , and

$$(3.36) \quad \nabla^2 u(\nu, \nu) = f^2 \nabla^2 s(\partial_s, \partial_s) = f f_{,s},$$

where $f_{,s} = \frac{df}{ds}$. Thus, III and H satisfy

$$(3.37) \quad \text{III} = -\frac{1}{2} f_{,s} \gamma \quad \text{and} \quad H = -f_{,s}.$$

On the other hand, (3.35) implies $\text{III} = \frac{1}{2} f \frac{d}{ds} \gamma_s$. Therefore, $f \frac{d}{ds} \gamma_s + f_{,s} \gamma = 0$ which shows

$$(3.38) \quad f \gamma_s = \sigma,$$

where σ is a fixed metric on U . As a result, on $W = (-\delta, \delta) \times U$,

$$(3.39) \quad g = \frac{1}{f(s)^2} ds^2 + \frac{1}{f(s)} \sigma.$$

Next, we incorporate the scalar curvature R and the vector fields X and Y . Suppose $U \subset\subset S \cap \Omega$. Then, for δ small, $W \subset \Omega$. By (3.26), (3.28) and (3.29), on W , we have

$$(3.40) \quad R = \frac{2}{3} |X|^2 + 2 \operatorname{div} X \quad \text{and} \quad X = -\frac{3}{2|\nabla u|} \nabla |\nabla u|.$$

This shows, in terms of $f = |\nabla u|$,

$$(3.41) \quad R = -\frac{3}{2} f_{,s}^2 + 3 f f_{,ss}.$$

On the other hand, (3.39) and direct calculation gives

$$(3.42) \quad R = -\frac{3}{2} f_{,s}^2 + 2 f f_{,ss} + 2 f K,$$

where K is the Gauss curvature of the metric σ on U . As a result,

$$(3.43) \quad f_{,ss} = 2K.$$

Consequently, σ is a metric of constant Gauss curvature on U and $f_{,ss}$ equals a constant that is independent on s . By using the vector field Y , we know the same conclusion holds on $U \subset\subset S \cap (M \setminus \bar{\Omega})$. As a result, if S intersects $\Sigma = \partial\Omega$ transversely, then the induced metric on S has constant Gauss curvature everywhere.

We proceed to get information on u near infinity. Since u is asymptotic to a coordinate function, we know there exists a compact set T , such that

$$(3.44) \quad |\nabla u| = 1 \quad \text{on} \quad M \setminus T.$$

This can be seen, for instance, by taking $u = u_1$ in $\{u_i\}_{1 \leq i \leq 3}$ where (u_1, u_2, u_3) forms a harmonic coordinate chart near ∞ . Then, outside a large coordinate ball B_r , each level set of u is $\{u_1 = c\} \setminus B_r$ for some constant c . As $|\nabla u| \rightarrow 1$ at ∞ and $|\nabla u|$ is locally a constant on each regular level set of u , we have $|\nabla u| = 1$ outside B_r .

It follows from (3.44), (3.36) and (3.37) that

$$(3.45) \quad \nabla^2 u(\nu, \nu) = 0 \text{ and } H = 0 \text{ on } \Sigma_t \cap (M \setminus T)$$

for any regular level set Σ_t . This, combined with (3.34), shows

$$(3.46) \quad \nabla^2 u = 0 \text{ on } M \setminus T.$$

We are now in a position to prove (3.33). Let S be a connected component of Σ_t with $t \in A$. By (3.37), H is locally a constant and hence a constant on S . If S is non-compact, then $S \cap (M \setminus T) \neq \emptyset$. By (3.45), $H = 0$ on $S \cap (M \setminus T)$. Hence $H = 0$ everywhere on S . By (3.36) and (3.37), $\nabla^2 u(\nu, \nu) = 0$ on S . Therefore, by (3.34), $\nabla^2 u = 0$ on S .

If S is compact, S must be a torus. This is because $\chi(\Sigma_t) = 1$ and any higher genus surface has negative Euler characteristics. Let σ be the induced metric on S . Since t is a regular value of both u and $u|_\Sigma$, S intersects Σ transversely. Hence σ has constant Gauss curvature K . By Gauss-Bonnet theorem, $K = 0$.

Now consider an open neighborhood W of S , diffeomorphic to $(-\delta, \delta) \times S$ for some $\delta > 0$, on which g takes the form of (3.39). By (3.43) and the fact $K = 0$, we have

$$(3.47) \quad \frac{d^2}{ds^2} f(s) = 0, \quad \forall s \in (-\delta, \delta).$$

Therefore, $f(s) = |\nabla u|(s) = as + b$ for some constants a and b , with $b = |\nabla u|$ on S .

If $a > 0$, the integral flow of $|\nabla u|^{-2} \nabla u$ starting from S exits on $[0, \infty)$. This means u goes to $+\infty$ along the flow. Hence the flow eventually enters $M \setminus T$. By (3.44), $|\nabla u|(s) = 1$ for large s . This contradicts with $|\nabla u|(s) = as + b$.

If $a < 0$, by consider the integral flow of $-|\nabla u|^{-2} \nabla u$ starting from S , we again have a contradiction.

Therefore, $a = 0$, i.e. $f_{,s} = 0$. This combined with (3.36) shows

$$\nabla^2 u(\nu, \nu) = 0 \text{ on } S.$$

We conclude $\nabla^2 u = 0$ on S by (3.34). This completes the proof of (3.33). $\square$

Remark 3.6. If the metric is merely $C^{0,1}$ across Σ , like that in Corollary 1.3, new difficulties seem to arise in analyzing rigidity for the case $C_1 = C_2 = \frac{2}{3}$. For instance, the function u is $W^{2,p}$ (thus $C^{1,\alpha}$) across Σ . A regular level set S of u , intersecting transversely with Σ at a curve $\mathcal{C}$, may not be C^2 across $\mathcal{C}$. In particular, the two Gauss curvature constants on both sides of $\mathcal{C}$ in S might not agree.

It is plausible to work with the condition $|\nabla u| = 1$, instead of $\nabla^2 u = 0$, in the claim (3.33) since $|\nabla u|$ is C^0 on M . By an argument similar to part of the proof of Theorem 3.3, it indeed can be shown that $|\nabla u| = 1$ along any non-compact component of Σ_t for $t \in A$. Thus, under a more stringent topological assumption that M contains no non-separating spheres and torus, the rigidity statement in Corollary 1.3 holds for $C_1 = C_2 = \frac{2}{3}$.

3.5. Fill-in of (M, g) by its conformal deformation. Given an asymptotically flat (E^3, g) with boundary Σ , consider its Green function G , which is determined by

$$(3.48) \quad \Delta_g G = 0 \text{ on } E, \quad G = 1 \text{ at } \Sigma, \quad \text{and } G \rightarrow 0 \text{ at } \infty.$$

Let $g_* = G^4 g$. Let $\Omega = E \cup \{q\}$ denote the one-point compactification of M by including a point q representing ∞ . It follows from the asymptotically flatness of g and the asymptotic expansion of G that g_* is continuously extendable across q , and moreover, $g_* \in W^{1,p}(B)$ for some $p > 6$ in a neighborhood B of q (see Lemma 6.1 in [23] for instance).

This (Ω, g_*) serves as a fill-in of (E, g) . The metric on $(E, g) \cup (\Omega, g_*)$, glued along Σ , is a $W_{loc}^{1,p}$ metric on $E \cup \Omega$. In particular, the function u in Section 3.1 exists on $(E, g) \cup (\Omega, g_*)$. Moreover, at most one regular level set of u passes through the point q . If E is topologically $\mathbb{R}^3$ minus a ball, $E \cup \Omega$ is homeomorphic to $\mathbb{R}^3$. Therefore, the proof of Theorem 1.2 applies to show

$$(3.49) \quad 8\pi \mathbf{m} \geq \int_E \frac{1}{2} \left[\frac{|\nabla^2 u|^2}{|\nabla u|} + R|\nabla u| \right] + \int_\Omega \frac{1}{2} \left[\frac{|\nabla^2 u|^2}{|\nabla u|} + R|\nabla u| \right] + \int_\Sigma (H_* - H)|\nabla u|.$$

Here H_* is the mean curvature of Σ in (Ω, g_*) with respect to the outward normal and is given by

$$(3.50) \quad H_* = -H - 4 \frac{\partial G}{\partial \nu},$$

where ν is the ∞ -pointing unit normal to Σ in (M, g) and H is the mean curvature of Σ in (M, g) with respect to ν . As a result,

$$(3.51) \quad H_* - H = -2H - 4 \frac{\partial G}{\partial \nu}.$$

Now let u denote the restriction of u on E and let u_Ω be the restriction of u to Ω . On $\Omega = M \cup \{q\}$, let $v = u_\Omega G$, then

$$(3.52) \quad \Delta_g v = 0, \quad v \rightarrow 0 \text{ at } \infty, \quad v = u \text{ at } \Sigma,$$

and

$$(3.53) \quad \frac{\partial v}{\partial \nu} = \frac{\partial u_\Omega}{\partial \nu} + u \frac{\partial G}{\partial \nu}.$$

As u is C^1 on $(E, g) \cup (\Omega, g_*)$,

$$(3.54) \quad \frac{\partial u_\Omega}{\partial \nu} = -\frac{\partial u}{\partial \nu}.$$

(Here ν points inward in the conformal fill-in (Ω, g_*) .) Set $w = u - v$, which satisfies

$$(3.55) \quad \Delta_g w = 0 \text{ on } E, \quad w = 0 \text{ at } \Sigma, \quad w \text{ is asymptotic to } x_i \text{ at } \infty.$$

By (3.53), (3.54) and the fact $u = v$ at Σ , we have

$$(3.56) \quad 2 \frac{\partial v}{\partial \nu} = v \frac{\partial G}{\partial \nu} - \frac{\partial w}{\partial \nu}.$$

Corollary 1.4 then follows from (3.49) and (3.56).

4. CONVERGENCE OF THE BROWN-YORK MASS

In this section, we apply Theorem 1.2 to study convergence of the Brown-York mass of surfaces approaching infinity in an asymptotically flat 3-manifold.

Recall that the Brown-York mass of a surface Σ with positive Gauss curvature, with mean curvature H in a 3-manifold (M, g) , is

$$(4.1) \quad \mathbf{m}_{BY}(\Sigma) = \frac{1}{8\pi} \int_\Sigma (H_0 - H) d\sigma,$$

where H_0 is the mean curvature of the isometric embedding of Σ in the Euclidean space $\mathbb{R}^3$.

The fact $\mathbf{m}_{BY}(\cdot)$ converges to the mass of an asymptotically flat manifold (M, g) along large coordinate spheres was demonstrated by Fan, Shi and Tam [12]. The convergence was later established for large nearly round surfaces by Shi, Wang and Wu [31]. Similar

results for surfaces of revolution were obtained by Fan and Kwong [10, 11] in asymptotically Schwarzschild manifolds.

Using Theorem 1.2, we demonstrate the convergence of $\mathbf{m}_{BY}(\cdot)$ for a large class of surfaces, which in particular includes surfaces obtained by scaling any fixed convex surface in the background Euclidean space.

Theorem 4.1. *Let (M^3, g) be an asymptotically flat manifold with a coordinate chart $\{x_i\}$ in which the metric coefficients g_{ij} satisfies the asymptotically flat condition (2.33) with $k \geq 4$ and $p > 3$. Let $\{\Sigma_r\}_{r \geq r_0}$ be a 1-parameter family of surfaces approaching the infinity with $r = \min_{\Sigma_r} \{|x| \mid x \in \Sigma_r\}$. Suppose the rescaled surfaces*

$$\tilde{\Sigma}_r := \left\{ \frac{1}{r}x \mid x \in \Sigma_r \right\} \subset \mathbb{R}^3$$

satisfy curvature condition

$$(4.2) \quad 0 < k_1 < \kappa_\alpha < k_2, \quad \alpha = 1, 2,$$

where $\{\kappa_\alpha\}$ are the principal curvatures of $\tilde{\Sigma}_r$ with respect to the background Euclidean metric in $\mathbb{R}^3$, and k_1, k_2 are two positive constants independent on r . Then

$$(4.3) \quad \lim_{r \rightarrow \infty} \mathbf{m}_{BY}(\Sigma_r) = \mathbf{m},$$

where $\mathbf{m}$ is the mass of (M, g) .

Note that condition (4.2) is equivalent to

$$(4.4) \quad \bar{K} > C_1 > 0 \quad \text{and} \quad 0 < \bar{H} < C_2$$

for some constants C_1, C_2 independent on r . Here $\bar{K}, \bar{H}$ denotes the Gauss curvature, the mean curvature of $\tilde{\Sigma}_r$ with respect to the background Euclidean metric in $\mathbb{R}^3$, respectively.

We first outline the idea behind the proof of Theorem 4.1. By condition (4.2) and the asymptotically flatness of g , Σ_r has positive Gauss curvature for large r . Thus Σ_r isometrically embeds in the Euclidean space $(\mathbb{R}^3, g_0)$ as a convex surface $\Sigma_r^{(0)}$ ([26, 27]). Let $\Omega_r^{(0)}$ be the domain enclosed by $\Sigma_r^{(0)}$ in $\mathbb{R}^3$ and E_r be the exterior region of Σ_r in M . The Euclidean domain $(\Omega_r^{(0)}, g_0)$ provides a natural fill-in of $\Sigma_r = \partial E_r$. Consider the manifold (N_r, g_r) obtained by gluing $(\Omega_r^{(0)}, g_0)$ and (E_r, g) along Σ_r . Intuitively, (N_r, g_r) approaches $(\mathbb{R}^3, g_0)$ in a suitable sense as $r \rightarrow \infty$. Hence applying Theorem 1.2 to (N_r, g_r) , we would expect the harmonic function u on (N_r, g_r) to approach a standard coordinate function on $(\mathbb{R}^3, g_0)$. In particular (1.4) would be an equation, the bulk integral on its right side would tend to zero, and the surface integral would be $\mathbf{m}_{BY}(\Sigma_r) + o(1)$.

To implement this idea, we need to have uniform control on the isometric embedding and to impose a uniform gauge when identifying $(E_r, g) \cup (\Omega_r^{(0)}, g_0)$ with $(\mathbb{R}^3, g_r)$ so that a fixed background is available when estimating u . We use condition (4.2) for achieving both these purposes.

Proof. For each large r , let $M_r = \{|x| \geq r\} \subset M$. Consider the diffeomorphism

$$\Phi_r : M_r \longrightarrow E = \{|x| \geq 1 \mid x \in \mathbb{R}^3\},$$

where $\Phi_r(x) = r^{-1}x$. Define a metric $\tilde{g}_r$ on E by $\tilde{g}_r = r^{-2}(\Phi_r^{-1})^*(g)$, then

$$(4.5) \quad \tilde{g}_{rij}(x) = g_{ij}(rx), \quad x \in E.$$

By definition, $\tilde{\Sigma}_r = \Phi_r(\Sigma_r) \subset E$. Let g_0 denote the background Euclidean metric on $\mathbb{R}^3$. Let $\bar{\sigma}_r$ be the induced metric on $\tilde{\Sigma}_r$ in $(\mathbb{R}^3, g_0)$ and $\bar{K}$ be the Gauss curvature of $(\tilde{\Sigma}_r, \bar{\sigma}_r)$. By condition (4.2), $\bar{K} \geq k_1^2 > 0$. This shows the intrinsic diameter of $(\tilde{\Sigma}_r, \bar{\sigma}_r)$ is bounded above by a constant independent on r . Thus there exists a constant $L \geq 1$ so that $\tilde{\Sigma}_r$ lies between $\{|x| = L\}$ and $\{|x| = 1\}$ for all large r .

In E , let T be the closed annular region bounded by $\{|x| = L\}$ and $\{|x| = 1\}$. By the asymptotically flatness condition (2.33) and weighted Sobolev inequalities,

$$(4.6) \quad \|\tilde{g}_r - g_0\|_{C^{3,\alpha}(T)} = O(r^{-\tau})$$

for some $0 < \alpha < 1$. As a result,

$$(4.7) \quad \|\tilde{\sigma}_r - \bar{\sigma}_r\|_{C^{3,\alpha}(\tilde{\Sigma}_r)} = O(r^{-\tau}),$$

$$(4.8) \quad \|\tilde{\nu} - \bar{\nu}\|_{C^3(\tilde{\Sigma}_r)} = O(r^{-\tau}),$$

and

$$(4.9) \quad \|\tilde{H} - \bar{H}\|_{C^0(\tilde{\Sigma}_r)} = O(r^{-\tau}).$$

Here $\tilde{\sigma}_r$, $\tilde{\nu}$, $\tilde{H}$ denote the induced metric, the inward unit normal, the mean curvature of $\tilde{\Sigma}_r$ in $(T, \tilde{g}_r)$, and $\bar{\nu}$, $\bar{H}$ denote the inward unit normal, the mean curvature of $\tilde{\Sigma}_r$ in (T, g_0) .

By the openness of solutions to the Weyl embedding problem [26, 19], it follows from the conditions $\tilde{\Sigma}_r \subset T$ and $k_1^2 < K < k_2^2$ that, for large r , there is an isometric embedding $\tilde{X}_r : (\tilde{\Sigma}_r, \tilde{\sigma}_r) \rightarrow \mathbb{R}^3$ so that

$$(4.10) \quad \|\tilde{X}_r - \text{Id}\|_{C^{3,\alpha}(\tilde{\Sigma}_r)} \leq C \|\tilde{\sigma}_r - \bar{\sigma}_r\|_{C^{3,\alpha}(\tilde{\Sigma}_r)}.$$

Here $\text{Id} : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ denotes the identity map and C is a constant independent on r . (This estimate was stated for $C^{2,\alpha}$ in [26], but examining sections 5-9 in [26] shows higher order derivatives estimate hold as well.) Moreover, $\tilde{X}_r$ has the same regularity as the metric $\tilde{\sigma}_r$ (see section 3 in [26].)

In what follows, we let $\tilde{\Sigma}_r^{(0)} = \tilde{X}_r(\tilde{\Sigma}_r)$ and let $\tilde{\Omega}_r^{(0)}$ be the region enclosed by $\tilde{\Sigma}_r^{(0)}$ in $\mathbb{R}^3$. Let $\tilde{\nu}_0$ be the unit inward normal to $\tilde{\Sigma}_r^{(0)}$ in $(\tilde{\Omega}_r^{(0)}, g_0)$. By (4.10),

$$(4.11) \quad \|\tilde{\nu}_0 - \bar{\nu}\|_{C^{2,\alpha}(\tilde{\Sigma}_r)} = O(r^{-\tau}),$$

and

$$(4.12) \quad \|\tilde{H}_0 - \bar{H}\|_{C^0(\Sigma)} = O(r^{-\tau}).$$

Here $\tilde{H}_0$ denote the mean curvature of $\tilde{\Sigma}_r^{(0)}$ in $(\mathbb{R}^3, g_0)$.

The Euclidean domains $(\tilde{\Omega}_r^{(0)}, g_0)$ provides a natural fill-in of $(\tilde{E}_r, \tilde{g}_r)$, where $\tilde{E}_r$ denotes the exterior of $\tilde{\Sigma}_r$ in E . To derive estimates on solutions to the corresponding Laplacian equation, we need a gauge to identify $\tilde{E}_r \cup \tilde{\Omega}_r^{(0)}$ with $\mathbb{R}^3$, which can be estimated via r . For this purpose, let $\tilde{\Omega}_r$ be the region enclosed by $\tilde{\Sigma}_r$ in $\mathbb{R}^3$. If we could extend $\tilde{X}_r : \tilde{\Sigma}_r \rightarrow \tilde{\Sigma}_r^{(0)}$ to a diffeomorphism $F_r : \tilde{\Omega}_r \rightarrow \tilde{\Omega}_r^{(0)}$, such that $(F_r)_*(\tilde{\nu}) = \tilde{\nu}_0$ and F_r remains close to Id , then $F_r^*(g_0)$ on $\tilde{\Omega}_r$ and $\tilde{g}_r$ on $\tilde{E}_r$ would form a locally Lipschitz metric $\hat{g}_r$ on $\mathbb{R}^3$, which is close to g_0 . We may then carry out analysis on this fixed $\mathbb{R}^3$ equipped with $\{\hat{g}_r\}$. We do not know how to construct such an map $F_r : \tilde{\Omega}_r \rightarrow \tilde{\Omega}_r^{(0)}$. Instead, we produce an alternative map $\tilde{F}_r : \tilde{\Omega}_r \rightarrow \mathbb{R}^3$ below, which suffices for the purpose of analyzing $\mathbf{m}_{BY}(\Sigma_r)$.

By condition (4.2), the second fundamental forms of $\tilde{\Sigma}_r$ are uniformly bounded in $(\mathbb{R}^3, g_0)$, and so are the second fundamental forms of $\tilde{\Sigma}_r^{(0)}$ in $(\mathbb{R}^3, g_0)$ by (4.10). It follows that $\tilde{\Sigma}_r, \tilde{\Sigma}_r^{(0)}$ have Gaussian tubular neighborhoods in $(\mathbb{R}^3, g_0)$ which have a fixed width independent on r . For small $s \geq 0$, consider the map

$$(4.13) \quad \Phi_r(\omega + s\tilde{\nu}) = \tilde{X}_r(\omega) + s\tilde{\nu}_0.$$

By (4.6), there exists a small $\epsilon > 0$, independent on r , such that

- $U_{r,\epsilon} = \{\omega + s\tilde{\nu} \mid \omega \in \tilde{\Sigma}_r, 0 \leq s < 2\epsilon\}$ is an open neighborhood of $\tilde{\Sigma}_r$ in $\tilde{\Omega}_r$, diffeomorphic to $\tilde{\Sigma}_r \times [0, 2\epsilon]$; and
- Φ_r is a diffeomorphism from $U_{r,\epsilon}$ to a Gaussian neighborhood of $\tilde{\Sigma}_r^{(0)}$ in $(\tilde{\Omega}_r^{(0)}, g_0)$.

Let η be a fixed smooth function on $[0, \infty)$, independent on r , satisfying $0 \leq \eta \leq 1$, $\eta = 1$ on $[0, \frac{1}{2}\epsilon]$, and $\eta = 0$ on $[\epsilon, \infty)$. Define $\tilde{F}_r : \tilde{\Omega}_r \rightarrow \mathbb{R}^3$, which interpolates Φ_r and Id, by

- $\tilde{F}_r(x) = \text{Id}(x)$, if $x \notin U_{r,\epsilon}$;
- $\tilde{F}_r(x) = \Phi_r(x)\eta(s) + \text{Id}(x)(1 - \eta(s))$, if $x = \omega + s\tilde{\nu}_r \in U_{r,\epsilon}$.

It follows from (4.8), (4.10) and (4.11) that

$$(4.14) \quad \|\tilde{F}_r - \text{Id}\|_{C^{2,\alpha}(\tilde{\Omega}_r)} \leq C(\|\tilde{X}_r - \text{Id}\|_{C^{2,\alpha}(\tilde{\Sigma}_r)} + \|\tilde{\nu} - \tilde{\nu}_0\|_{C^{2,\alpha}(\tilde{\Sigma}_r)}) = O(r^{-\tau}).$$

In particular, for large r , $\tilde{F}_r : \tilde{\Omega}_r \rightarrow \mathbb{R}^3$ is an immersion. Define $\tilde{g}_r = \tilde{F}_r^*(g_0)$ on $\tilde{\Omega}_r$. By construction, $(\tilde{\Omega}_r, \tilde{g}_r)$ satisfies

- a) the induced metric from $\tilde{g}_r$ on $\tilde{\Sigma}_r$ is $\tilde{\sigma}_r$;
- b) $\tilde{\nu}$ is the inward unit normal to $\tilde{\Sigma}_r$ in $(\tilde{\Omega}_r, \tilde{g}_r)$, since $(\tilde{F}_r)_*(\tilde{\nu}) = \tilde{\nu}_0$;
- c) $\|\tilde{g}_r - g_0\|_{C^{1,\alpha}(\tilde{\Omega}_r)} = O(r^{-\tau})$.

Moreover, the mean curvature of $\tilde{\Sigma}_r$ in $(\tilde{\Omega}_r, \tilde{g}_r)$ equals $\tilde{H}_0$, which is the mean curvature of $\tilde{\Sigma}_r^{(0)}$ in $(\mathbb{R}^3, g_0)$.

We now write $(\mathbb{R}^3, \hat{g}_r) = (\tilde{E}_r, \tilde{g}_r) \cup (\tilde{\Omega}_r, \tilde{g}_r)$, where $\hat{g}_r$ is a $C_{loc}^{0,1}$ metric on $\mathbb{R}^3$. Moreover, by properties a), b), c) above,

$$(4.15) \quad \|(\hat{g}_r - g_0)_{ij}\|_{W_{-\tau}^{2,p}(\tilde{E}_r)} = r^{-\tau} \|g_{ij} - \delta_{ij}\|_{W_{-\tau}^{2,p}(E_r)} = O(r^{-\tau}),$$

where E_r is the exterior of Σ_r in (M, g) , and

$$(4.16) \quad \|\hat{g}_r - g_0\|_{C^1(\tilde{\Omega}_r)} = O(r^{-\tau}).$$

Consider the operator

$$(4.17) \quad \Delta_{\hat{g}_r} : W_{1-\tau}^{2,p}(\mathbb{R}^3) \longrightarrow W_{-1-\tau}^{0,p}(\mathbb{R}^3).$$

Here $W_{\delta}^{k,p}(\mathbb{R}^3)$ is the weighted Sobolev space on $\mathbb{R}^3$, consisting of functions $v \in W_{loc}^{k,p}(\mathbb{R}^3)$ such that

$$(4.18) \quad \|v\|_{W_{\delta}^{k,p}(\mathbb{R}^3)} = \sum_{|\beta| \leq k} \|\partial^{\beta} u (1 + |x|^2)^{\frac{1}{2}(|\beta| - \delta - \frac{3}{p})}\|_{L^p(\mathbb{R}^3)} < \infty.$$

By (4.15) and (4.16), $\Delta_{\hat{g}_r} x_1 \in W_{-1-\tau}^{0,p}(\mathbb{R}^3)$ and

$$(4.19) \quad \|\Delta_{\hat{g}_r} x_1\|_{W_{-1-\tau}^{0,p}(\mathbb{R}^3)} = O(r^{-\tau}).$$

By the usual construction of harmonic coordinates (see Section 3 in [3] for instance), there exists $w_r \in W_{1-\tau}^{2,p}(\mathbb{R}^3)$ so that

$$(4.20) \quad \Delta_{\hat{g}_r} w_r = \Delta_{\hat{g}_r} x_1.$$

Let $u_r = x_1 - w_r$, then u_r satisfies $\Delta_{\hat{g}_r} u_r = 0$ and u_r is asymptotic to x_1 .

We estimate w_r by applying Corollary 1.16 of [3]. Shrinking τ if needed, we may assume $\tau \in (\frac{1}{2}, 1)$. By (4.15) and (4.16),

$$(4.21) \quad \lim_{r \rightarrow \infty} \|\hat{g}_r - g_0\|_{W_0^{1,p}(\mathbb{R}^3)} = 0.$$

By Corollary 1.16 in [3],

$$(4.22) \quad \|w_r - \text{Ker}(\Delta_{\hat{g}_r})\|_{W_{1-\tau}^{2,p}(\mathbb{R}^3)} \leq C \|\Delta_{\hat{g}_r} w_r\|_{W_{-1-\tau}^{0,p}(\mathbb{R}^3)}.$$

Here $\text{Ker}(\Delta_{\hat{g}_r})$ is the Kernel of $\Delta_{\hat{g}_r} : W_{1-\tau}^{2,p}(\mathbb{R}^3) \rightarrow W_{-1-\tau}^{0,p}(\mathbb{R}^3)$, and C is some constant independent on r . Since $1 - \tau \in (0, \frac{1}{2})$, we have

$$(4.23) \quad \dim \text{Ker}(\Delta_{\hat{g}_r}) = \dim \text{Ker}(\Delta_{g_0}) = 1$$

(see Proposition 1.15 and Corollary 1.9 of [3]). Thus,

$$(4.24) \quad \text{Ker}(\Delta_{\hat{g}_r}) = \{\text{constant functions}\}.$$

This, combined with (4.22), gives

$$(4.25) \quad \inf_{c \in \mathbb{R}} \|w_r - c\|_{W_{1-\tau}^{2,p}(\mathbb{R}^3)} \leq C \|\Delta_{\hat{g}_r} w_r\|_{W_{-1-\tau}^{0,p}(\mathbb{R}^3)} \\ = C \|\Delta_{\hat{g}_r} x_1\|_{W_{-1-\tau}^{0,p}(\mathbb{R}^3)}.$$

We pick $c_r \in \mathbb{R}$ as follows: if $\|\Delta_{\hat{g}_r} x_1\|_{W_{-1-\tau}^{0,p}(\mathbb{R}^3)} > 0$, pick $c_r \in \mathbb{R}$ so that

$$(4.26) \quad \|w_r - c_r\|_{W_{1-\tau}^{2,p}} \leq 2C \|\Delta_{\hat{g}_r} x_1\|_{W_{-1-\tau}^{0,p}(\mathbb{R}^3)}.$$

If $\|\Delta_{\hat{g}_r} x_1\|_{W_{-1-\tau}^{0,p}(\mathbb{R}^3)} = 0$, we simply pick $c_r = 0$, in which case we also let $w_r = 0$. For these choices of w_r and c_r , we have

$$(4.27) \quad \|w_r - c_r\|_{W_{1-\tau}^{2,p}(\mathbb{R}^3)} \leq 2C \|\Delta_{\hat{g}_r} x_1\|_{W_{-1-\tau}^{0,p}}.$$

Together with (4.19), this shows

$$(4.28) \quad \|\partial w_r\|_{W_{-\tau}^{1,p}(\mathbb{R}^3)} = O(r^{-\tau}).$$

We now estimate the gradient of $u_r = x_1 - w_r$. By (4.28) and Sobolev embedding,

$$(4.29) \quad |\nabla u_r - \nabla x_1|_{\hat{g}_r} = |\nabla w_r|_{\hat{g}_r} = O(r^{-\tau}).$$

As $|\nabla x_1|_{\hat{g}_r} = 1 + O(r^{-\tau})$, this gives

$$(4.30) \quad |\nabla u_r|_{\hat{g}_r} = 1 + O(r^{-\tau}).$$

In particular, $|\nabla u_r|_{\hat{g}_r} > 0$ on $\mathbb{R}^3$, for large r . By Theorem 1.2 and Remark 3.3, we have

$$(4.31) \quad \mathbf{m}(\hat{g}_r) = \frac{1}{16\pi} \int_{\mathbb{R}^3} \left[\frac{|\nabla^2 u_r|_{\hat{g}_r}}{|\nabla u_r|_{\hat{g}_r}} + R_{\hat{g}_r} |\nabla u_r|_{\hat{g}_r} \right] dV_{\hat{g}_r} + \frac{1}{8\pi} \int_{\tilde{\Sigma}_r} (\tilde{H}_0 - \tilde{H}) |\nabla u_r|_{\hat{g}_r} d\hat{\sigma}_r,$$

where

$$(4.32) \quad \mathbf{m}(\hat{g}_r) = r^{-1} \mathbf{m}(g) \quad \text{and} \quad \frac{1}{8\pi} \int_{\Sigma} (\tilde{H}_0 - \tilde{H}) d\hat{\sigma}_r = r^{-1} \mathbf{m}_{BY}(\Sigma_r).$$

We estimate the other terms in the right side of (4.31). By (4.30),

$$(4.33) \quad \int_{\tilde{\Sigma}_r} (\tilde{H}_0 - \tilde{H})(|\nabla u_r|_{\hat{g}_r} - 1) d\hat{\sigma}_r = O(r^{-2\tau}).$$

Here we used (4.9) and (4.12), which together imply

$$(4.34) \quad |\tilde{H} - \tilde{H}_0| = O(r^{-\tau}).$$

For the integral involving $R_{\hat{g}_r}$, we have

$$(4.35) \quad \int_{\tilde{\Omega}_r} R_{\hat{g}_r} |\nabla u_r|_{\hat{g}_r} dV_{\hat{g}_r} = 0$$

because, $\hat{g}_r = \tilde{F}_r^*(g_0)$ on $\tilde{\Omega}_r$, which is flat. On $\tilde{E}_r$, by (4.30),

$$(4.36) \quad \begin{aligned} \int_{\tilde{E}_r} R_{\hat{g}_r} |\nabla u_r|_{\hat{g}_r} dV_{\hat{g}_r} &\leq C \int_{\tilde{E}_r} R_{\hat{g}_r} dV_{\hat{g}_r} = Cr^{-1} \int_{M_r} R_g dV_g \\ &= o(r^{-1}), \end{aligned}$$

where in the last step, we used the assumption $R(g)$ is integrable on (M, g) .

For the integral involving $\nabla^2 u_r$, we have

$$(4.37) \quad \int_{\mathbb{R}^3} |\nabla^2 u_r|_{\hat{g}_r}^2 dV_{\hat{g}_r} \leq C \left(\int_{\mathbb{R}^3} |\partial^2 u_r|_{g_0}^2 dV_{g_0} + \int_{\mathbb{R}^3} |\hat{\Gamma}_{ij}^k|^2 |\nabla u_r|_{g_0}^2 dV_{g_0} \right).$$

Here $\hat{\Gamma}_{ij}^k$ is the connection coefficients of $\hat{g}_r$. The right side can be handled by Hölder's inequality. Let $\rho = \sqrt{1 + |x|^2}$, then

$$(4.38) \quad \begin{aligned} \int_{\mathbb{R}^3} |\hat{\Gamma}_{ij}^k|^2 dV_{g_0} &\leq C \int_{\mathbb{R}^3} |\partial \hat{g}_r|^2 dV_{g_0} \\ &= C \int_{\mathbb{R}^3} |\partial \hat{g}_r|^2 \rho^{2(\tau+1) - \frac{6}{q}} \rho^{-2(\tau+1) + \frac{6}{q}} dV_{g_0} \\ &\leq C \|\partial \hat{g}_r\|_{W_{-1-\tau}^{0,p}(\mathbb{R}^3)}^2 \left(\int_{\mathbb{R}^3} \rho^{[-2(\tau+1) + \frac{6}{q}] \frac{q}{q-2}} dx \right)^{1-\frac{2}{q}} \\ &= O(r^{-2\tau}), \end{aligned}$$

where we used (4.15), (4.16) and the assumption $\tau > \frac{1}{2}$.

Similarly, using the fact $\partial^2 u_r = \partial^2 w_r$, we have

$$(4.39) \quad \begin{aligned} \int_{\mathbb{R}^3} |\partial^2 u_r|^2 dV_{g_0} &= \int_{\mathbb{R}^3} |\partial^2 w_r|^2 \rho^{2(\tau+1) - \frac{6}{q}} \rho^{-2(\tau+1) + \frac{6}{q}} dV_{g_0} \\ &\leq \|\partial^2 w_r\|_{W_{-1-\tau}^{0,p}(\mathbb{R}^3)}^2 \left(\int_{\mathbb{R}^3} \rho^{[-2(\tau+1) + \frac{6}{q}] \frac{q}{q-2}} dV_{g_0} \right)^{1-\frac{2}{q}} \\ &= O(r^{-2\tau}), \end{aligned}$$

where we used (4.28) in the last step.

It follows from (4.31) – (4.39) that

$$(4.40) \quad \mathbf{m}(g) = \frac{1}{8\pi} \int_{\Sigma_r} (H_0 - H) d\sigma_r + O(r^{1-2\tau}) + o(1),$$

which proves (4.3). □

Remark 4.1. In the statement of Theorem 4.1, we formulate the asymptotically flatness of g via weighted Sobolev spaces $W_{-\tau}^{k,p}$ with $k \geq 4$ and $p > 3$. It is clear from the proof that one can also use a weighted Hölder space $C_{-\tau}^{k,\alpha}$ (see [3]), with $k \geq 3$, to specify the asymptotic of g . The fact we need a Hölder decay condition on the third order derivative of g comes from estimate (4.10) which is needed for the estimate of $\tilde{\nu} - \tilde{\nu}_0$ in (4.14).

APPENDIX A. THE DESINGULARIZATION APPROACH

We give another proof of (1.4) by combining the mass formula in [6] with the mollification method in [25]. Let (E, g) , (Ω, g) , (M, g) and Σ be given as in the beginning of Section 3. Applying Proposition 3.1 in [25] (and its proof therein) to (M, g) near Σ , we have

Proposition A.1. *There exists a family of C^2 metrics $\{g_\delta\}_{\delta>0}$ on M with the properties:*

- (1) $g_\delta = g$ outside $\Sigma \times (-\delta, \delta)$; g_δ converges to g in C^0 on M ;
- (2) $\|g_\delta\|_{C^{0,1}(M)}$ and thus $\|\Gamma_\delta\|_{L^\infty(M)}$ are bounded uniformly with respect to δ . Here Γ_δ denotes the Christoffel symbol of g_δ ;
- (3) $R_\delta(x, t) = O(1)$ in $\Sigma \times \{\delta^2 < |t| < \delta\}$, and
- (4) $R_\delta(x, t) = O(1) + 2(H_\Omega - H)(x) \frac{1}{\delta^2} \phi\left(\frac{t}{\delta^2}\right)$ in $\Sigma \times (-\delta^2, \delta^2)$.

Here $\Sigma \times (-\delta, \delta)$ denotes a Gaussian tubular neighborhood of Σ in (M, g) which consists of points that are less than δ distance to Σ ; t is the coordinate in $(-\delta, \delta)$ such that ∂_t points to the infinity; $R_\delta(x, t)$ is the scalar curvature of g_δ at a point $(x, t) \in \Sigma \times (-\delta, \delta)$; $O(1)$ is a bounded quantity that is independent on δ , and $\phi(t)$ is a standard mollifier on $\mathbb{R}^1$.

Applying equation (6.28) from [6] to (M, g_δ) , using the fact $g_\delta = g$ near infinity, we have

Proposition A.2. *Let u_δ be the harmonic function on (M, g_δ) which is asymptotic to x_1 . Then*

$$(A.1) \quad \mathfrak{m}(g) \geq 16\pi \int_M \frac{|\nabla_\delta^2 u_\delta|^2}{|\nabla_\delta u_\delta|} + R_\delta |\nabla_\delta u_\delta|.$$

Here $\nabla_\delta, \nabla_\delta^2$ denote the operators with respect to g_δ .

Proof of (1.4) using Propositions A.1 and A.2.

On (M, g) , let $g_0 = g$ and let $\{g_\delta\}$ be given in Proposition A.1. Let u_0 be the harmonic function on (M, g_0) , asymptotic to x_1 near infinity. (u_0 is precisely the function u considered in Section 3.) Similar to the start of the proof of Proposition 2.3, we may view this u_0 as one of the asymptotically flat coordinate functions near infinity. In what follows, we define $\bar{x}_1 = u_0$.

For each $\delta > 0$, we recall how one finds the harmonic function u_δ that is asymptotic to $\bar{x}_1$. First, we compute $\Delta_\delta \bar{x}_1$, the Laplacian of $\bar{x}_1$ with respect to g_δ , which satisfies $\Delta_\delta \bar{x}_1 = 0$ outside $\Sigma \times (-\delta, \delta)$. This shows $\Delta_\delta \bar{x}_1 \in W_{-\eta-2}^{0,p}(M)$, $\forall \eta > 0$. Consider the operator

$$(A.2) \quad \Delta_\delta : W_{-\eta}^{2,p}(M) \rightarrow W_{-\eta-2}^{0,p}(M).$$

By Proposition 2.2 in [3], Δ_δ is an isomorphism provided $0 < \eta < 1$. Therefore, for a given $\eta \in (0, 1)$, there exists a $w_\delta \in W_{-\eta}^{2,p}(M)$ such that $\Delta_\delta w_\delta = \Delta_\delta \bar{x}_1$ and

$$(A.3) \quad \begin{aligned} \|w_\delta\|_{W_{-\eta}^{2,p}(M)} &\leq C \|\Delta_\delta w_\delta\|_{W_{-\eta-2}^{0,p}(M)} \\ &= C \|\Delta_\delta \bar{x}_1\|_{W_{-\eta-2}^{0,p}(M)}. \end{aligned}$$

Here C can be taken to be independent on δ by (i) and (ii) of Proposition A.1.

Since $\bar{x}_1 = u_0 \in W_{loc}^{2,p}(M)$ and $\{g_\delta\}$ is uniformly bounded in $C^{0,1}(M)$, we have

$$\int_U |\Delta_\delta \bar{x}_1|^p < \Lambda,$$

where U is a fixed bounded open set containing Σ and Λ is a constant independent on δ . This combined with (i) of Proposition A.1 shows, as $\delta \rightarrow 0$,

$$(A.4) \quad \|\Delta_\delta \bar{x}_1\|_{W_{-\eta-2}^{0,p}(M)} = o(1),$$

Therefore,

$$(A.5) \quad \|w_\delta\|_{W_{-\eta}^{2,p}(M)} = o(1).$$

By Sobolev embedding, w_δ also satisfies $\|w_\delta\|_{C_{-\eta}^{1,\alpha}(M)} = o(1)$. Set $u_\delta = u_0 - w_\delta$, u_δ is the harmonic function on (M, g_δ) that is asymptotic to $\bar{x}_1$.

By Propositions A.1 and A.2, we have

$$(A.6) \quad \mathfrak{m}(g) \geq \underbrace{\int_M \frac{|\nabla^2 u_\delta|^2}{|\nabla u_\delta|}}_I + \underbrace{\int_{M \setminus \{\Sigma \times (-\delta, \delta)\}} R_\delta |\nabla u_\delta|}_{II} + \underbrace{\int_\Sigma \int_{-\delta}^\delta R_\delta |\nabla u_\delta|}_{III}.$$

Letting $\delta \rightarrow 0$, by (A.5) and Fatou's Lemma,

$$(A.7) \quad \liminf_{\delta \rightarrow 0} I \geq \liminf_{\delta \rightarrow 0} \int_{M \setminus \{\Sigma \times (-\delta, \delta)\}} \frac{|\nabla^2 u_\delta|^2}{|\nabla u_\delta|} \geq \int_M \frac{|\nabla^2 u_0|^2}{|\nabla u_0|}.$$

By (A.5) and Proposition A.1,

$$(A.8) \quad \lim_{\delta \rightarrow 0} II = \int_M R |\nabla u_0|, \quad \text{and} \quad \lim_{\delta \rightarrow 0} III = 2 \int_\Sigma (H_\Omega - H) |\nabla u_0|.$$

Thus (1.4) follows from (A.6) – (A.8). $\square$

We remark that though this approach established (1.4), it does not give the regularity of u on Σ which is a main component of Theorem 1.2.

Acknowledgements. SH thanks Hubert Bray for insightful discussions relating to the level set method. PM is grateful to Danial Stern for giving the geometric interpretation of the boundary term in (2.32). PM is also indebted to Siyuan Lu for explaining estimates on isometric embedding pertinent to Theorem 1.5. TT thanks Man-Chun Lee and Richard Schoen for helpful discussions on the desingularisation approach.

The authors want to thank the anonymous referee for helpful suggestions, specially for the insightful comment on the zero mass case in relation to Corollary 1.3.

REFERENCES

- [1] A. Alaei, M. Khuri and S.-T. Yau, *Geometric inequalities for quasi-local masses*, Commun. Math. Phys., **378** (2020), 467–505.
- [2] R. Arnowitt, S. Deser, and C. Misner, *Energy and the criteria for radiation in general relativity*, Physical Review **118** (1960), no. 4, 1100.
- [3] R. Bartnik, *The mass of an asymptotically flat manifold*, Comm. Pure Appl. Math., **39** (1986), no. 5, 661–693.
- [4] R. Bartnik, *New definition of quasilocal mass*, Phys. Rev. Lett. **62** (1989), 2346–2348.

- [5] H. Bray and D. Stern, *Scalar curvature and harmonic one-forms on three-manifolds with boundary*, to appear in Comm. Anal. Geom., arXiv:1911.06803 (2019).
- [6] H. Bray, D. Kazaras, M. Khuri, and D. Stern, *Harmonic functions and the mass of 3-dimensional asymptotically flat Riemannian manifolds.*, arXiv preprint arXiv:1911.06754 (2019).
- [7] J. D. Brown and J. W. York, Jr., *Quasilocal energy in general relativity*, in Mathematical aspects of classical field theory (Seattle, WA, 1991), volume 132 of Contemp. Math., pages 129–142. Amer. Math. Soc., Providence, RI, 1992, MR1188439.
- [8] J. D. Brown and J. W. York, Jr., *Quasilocal energy and conserved charges derived from the gravitational action*, Phys. Rev. D (3) **47** (1993), no. 4, 1407–1419, MR1211109.
- [9] P. Chruściel, *Boundary conditions at spatial infinity from a Hamiltonian point of view*, Topological properties and global structure of space-time, Bergmann P.G., De Sabbata V. (eds), NATO Adv. Sci. Inst. Ser. B Phys., (1986), 49–59.
- [10] X.-Q. Fan and K.-K. Kwong, *The Brown-York mass of revolution surface in asymptotically Schwarzschild manifold*, J. Geom. Anal., **21** (2011), no. 3, 527–542.
- [11] X.-Q. Fan and K.-K. Kwong, *A property of the Brown-York mass in Schwarzschild manifolds*, J. Math. Anal. and Appl., **400** (2013), no. 2, 615–623.
- [12] X.-Q. Fan, Y. Shi and L.-F. Tam, *Large-sphere and small-sphere limits of the Brown-York mass*, Comm. Anal. Geom. **17** (2009), no. 1, 37–72.
- [13] D. Gilbarg and N. Trudinger, *Elliptic partial differential equations of second order*. Springer, 2015.
- [14] M. Gromov, *Four lectures on scalar curvature*, arXiv:1908.10612, 2019.
- [15] S. Hirsch and P. Miao, *A positive mass theorem for manifolds with boundary*, Pacific J. Math., **306** (2020), no. 1, 185–201.
- [16] S. Hirsch, D. Kazaras and M. Khuri, *Spacetime harmonic functions and the mass of 3-dimensional asymptotically flat initial data for the Einstein equations*, arXiv preprint arXiv:2002.01534 (2020).
- [17] J. Jauregui, *Fill-ins of nonnegative scalar curvature, static metrics, and quasi-local mass*, Pacific J. Math., **261** (2013), 417–444.
- [18] J. Jauregui, P. Miao and L.-F. Tam, *Extensions and fill-ins with non-negative scalar curvature*, Class. Quantum Grav., **30** (2013), 195007.
- [19] C. Li and Z. Wang, *The Weyl problem in warped product spaces*, J. Differential Geom., **114** (2020), no. 2, 243–304.
- [20] C.-C. M. Liu and S.-T. Yau, *Positivity of quasi-local mass*, Phys. Rev. Lett., **90** (2003), no. 23, 231102.
- [21] C.-C. M. Liu and S.-T. Yau, *Positivity of quasi-local mass II*, J. Amer. Math. Soc., **19** (2006), no. 1, 181–204.
- [22] C. Mantoulidis and P. Miao, *Total mean curvature, scalar curvature, and a variational analog of Brown-York mass*, Commun. Math. Phys. **352** (2017), no. 2, 703–718.
- [23] C. Mantoulidis, P. Miao and L.-F. Tam, *Capacity, quasi-local mass, singular fill-ins*, Journal für die reine und angewandte Mathematik, Crelle's Journal, DOI: <https://doi.org/10.1515/crelle-2019-0040>.
- [24] D. McFeron and G. Székelyhidi, *On the positive mass theorem for manifolds with corners*, Commun. Math. Phys. **313** (2012), no. 2, 425–443.
- [25] P. Miao, *Positive mass theorem on manifolds admitting corners along a hypersurface.*, Adv. Theor. Math. Phys., **6** (2002). no.6, 1163–1182.
- [26] L. Nirenberg, *The Weyl and Minkowski problem in differential geometry in the large*, Commun. Pure Appl. Math. **6** (1953), no. 3, 337–394.
- [27] A. V. Pogorelov, *Regularity of a convex surface with given Gaussian curvature*, (Russian) Mat. Sbornik N.S. **31**(73), (1952), no. 1, 88–103, MR0052807.
- [28] R. Schoen and S.-T. Yau, *On the proof of the positive mass conjecture in general relativity*. Comm. Math. Phys., **65** (1979), no.1, 45–76.
- [29] R. Schoen and S.-T. Yau, *Proof of the positive mass theorem II*. Comm. Math. Phys., **79** (1981), no.2, 231–260.
- [30] Y. Shi and L.-F. Tam, *Positive mass theorem and the boundary behaviors of compact manifolds with nonnegative scalar curvature*, Journal of Differential Geometry, **62** (2002), no.1, 79–125.
- [31] Y. Shi, G. Wang and J. Wu, *On the behavior of quasi-local mass at the infinity along nearly round surfaces*, Ann. Global Anal. Geom., **36** (2009), no. 4, 419–441.
- [32] Y. Shi, W. Wang and G. Wei, *Total mean curvature of the boundary and nonnegative scalar curvature fill-ins*, arXiv:2007.06756.

- [33] Y. Shi, W. Wang, G. Wei and J. Zhu, *On the fill-in of nonnegative scalar curvature metrics*, Math. Ann., **379** (2021), 235-270.
- [34] D. Stern, *Scalar curvature and harmonic maps to S^1* , to appear in J. Diff. Geom., arXiv:1908.09754 (2019).
- [35] M.-T. Wang and S.-T. Yau, *A generalization of Liu-Yau's quasi-local mass*, Comm. Anal. Geom., **15** (2007), no.2, 249-282.
- [36] M.-T. Wang and S.-T. Yau, *Isometric embeddings into the Minkowski space and new quasi-local mass*, Commun. Math. Phys., **288** (2009), no.3, 919-942.
- [37] E. Witten, *A new proof of the positive energy theorem*, Commun. Math. Phys., **80** (1981), no.3, 381-402.

DEPARTMENT OF MATHEMATICS, DUKE UNIVERSITY, DURHAM, NC 27708, USA
Email address: `sven.hirsch@duke.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MIAMI, CORAL GABLES, FL 33146, USA
Email address: `pengzim@math.miami.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, IRVINE, CA 92697
Email address: `tytsang@uci.edu`