

MATRIX REARRANGEMENT INEQUALITIES REVISITED

Victoria M Chayes
Department of Mathematics
Rutgers University
Piscataway, NJ 08854
vc362@math.rutgers.edu

November 1, 2021

ABSTRACT

Let $\|X\|_p = \text{Tr}[(X^*X)^{p/2}]^{1/p}$ denote the p -Schatten norm of a matrix $X \in M_{n \times n}(\mathbb{C})$, and $\sigma(X)$ the singular values with $\uparrow \downarrow$ indicating its increasing or decreasing rearrangements. We wish to examine inequalities between $\|A+B\|_p^p + \|A-B\|_p^p$, $\|\sigma_\downarrow(A) + \sigma_\downarrow(B)\|_p^p + \|\sigma_\downarrow(A) - \sigma_\downarrow(B)\|_p^p$, and $\|\sigma_\uparrow(A) + \sigma_\downarrow(B)\|_p^p + \|\sigma_\uparrow(A) - \sigma_\downarrow(B)\|_p^p$ for various values of $1 \leq p < \infty$. It was conjectured in [6] that a universal inequality $\|\sigma_\downarrow(A) + \sigma_\downarrow(B)\|_p^p + \|\sigma_\downarrow(A) - \sigma_\downarrow(B)\|_p^p \leq \|A+B\|_p^p + \|A-B\|_p^p \leq \|\sigma_\uparrow(A) + \sigma_\downarrow(B)\|_p^p + \|\sigma_\uparrow(A) - \sigma_\downarrow(B)\|_p^p$ might hold for $1 \leq p \leq 2$ and reverse at $p \geq 2$, potentially providing a stronger inequality to the generalization of Hanner's Inequality to complex matrices $\|A+B\|_p^p + \|A-B\|_p^p \geq (\|A\|_p + \|B\|_p)^p + \||A\|_p - \|B\|_p|^p$. We extend some of the cases in which the inequalities of [6] hold, but offer counterexamples to any general rearrangement inequality holding. We simplify the original proofs of [6] with the technique of majorization. This also allows us to characterize the equality cases of all of the inequalities considered. We also address the commuting, unitary, and $\{A, B\} = 0$ cases directly, and expand on the role of the anticommutator. In doing so, we extend Hanner's Inequality for self-adjoint matrices to the $\{A, B\} = 0$ case for all ranges of p .

Keywords Matrix Inequality · Hanner's Inequality · p -Schatten Norm · Majorization

1 Introduction

It has been of great interest to extend Hanner's Inequality for L^p spaces

$$\|f+g\|_p^p + \|f-g\|_p^p \geq (\|f\|_p + \|g\|_p)^p + \||f\|_p - \|g\|_p|^p \quad (1.1)$$

for $1 \leq p \leq 2$ to the non-communative analogue in C^p

$$\|A+B\|_p^p + \|A-B\|_p^p \geq (\|A\|_p + \|B\|_p)^p + \||A\|_p - \|B\|_p|^p. \quad (1.2)$$

In [6], Carlen and Lieb proposed the following two conjectures for their potential pertinence to proving 1.2:

Conjecture 1.1. For all $1 \leq p \leq 2$, and all complex-valued $n \times n$ matrices A and B ,

$$\|A+B\|_p^p + \|A-B\|_p^p \geq \|\sigma_\downarrow(A) + \sigma_\downarrow(B)\|_p^p + \|\sigma_\downarrow(A) - \sigma_\downarrow(B)\|_p^p. \quad (1.3)$$

For $p > 2$, the inequality reverses.

Conjecture 1.2. For all $1 \leq p \leq 2$, and all complex-valued $n \times n$ matrices A and B ,

$$\|A+B\|_p^p + \|A-B\|_p^p \leq \|\sigma_\uparrow(A) + \sigma_\downarrow(B)\|_p^p + \|\sigma_\uparrow(A) - \sigma_\downarrow(B)\|_p^p. \quad (1.4)$$

For $p > 2$, the inequality reverses.

For these, the authors proved Conjecture 1.1 in the case $A \geq B \geq 0$ and $1 \leq p \leq 2$; and proved Conjecture 1.2 in the case $A \geq |B| \geq 0$ and $1 \leq p \leq 2$. We note a missing requirement in [6] used in the proof for Conjecture 1.2 in

those conditions is also that $\sigma_n(A) \geq \sigma_1(B)$. Lemma 2.1 in [15] proves that Conjecture 1.1 holds for all matrices and $p = 2k$, $k \in \mathbb{N}$. To the best of our knowledge, no further work has been done on the subject.

If Conjecture 1.1 were true in general, with an additional application of Hanner's Inequality on the sequences of singular values, the non-commutative Hanner's Inequality for matrices would be proven in general; currently, it is only known for $A + B$, $A - B \geq 0$ for all p , or general A and B in the ranges $1 \leq p \leq \frac{4}{3}$ and $p \geq 4$ [3].

In this paper we extend the range of Conjecture 1.1 with the requirements $A \geq B \geq 0$ to $2 \leq p \leq 3$, and we prove Conjecture 1.2 in the $A + B$, $A - B \geq 0$, $\sigma_n(A) \geq \sigma_1(B)$ case for the range $1 \leq p \leq 3$. We prove both conjectures for the full range of p in the commuting case. We prove Conjecture 1.1 in the case that A and B are both unitary, and in the case when A and B are self-adjoint and $\{A, B\} = 0$. However, we demonstrate that both conjectures are false in general. Section 2 gives a background to majorization, which is the primary technique that we use in our proofs. Section 3 presents the extensions of the conjectures' requirements and ranges, and general counterexamples.

The key observation as to why the conjectures cannot hold in general is that if the matrix B is taken to be unitary, all its singular values are equal to 1, and therefore there is no distinction between the "aligned" and the "up-down" rearrangements. If both conjectures were true, this would imply equality everywhere when B is unitary, which can easily be numerically confirmed as false.

The fact that these re-arrangement inequalities do not hold in general is notable, because the analogue to Conjecture 1.1 with complex functions and the spherically symmetric decreasing rearrangement is shown in [6] to hold. Therefore, we see directly that the non-commutativity of the matrices ruins a commutative identity. In disproving Conjecture 1.1 in general, we also rule it out as a method to attempt to extend Hanner's inequality to C^p .

We will use the following notation throughout this paper: $\sigma(X)$ denotes the vector of singular values of a matrix X , assumed to be in descending order unless $\sigma_{\uparrow}(X)$ is specified; $\sigma_{\downarrow}(X)$ may then be used for emphasis. The norm $\|\cdot\|_p$ may either indicate the vector p -norm or the p -Schatten norm dependent on context. We use for a vector \mathbf{v} the notation $[\mathbf{v}]$ to indicate the matrix $[\text{Diag}(\mathbf{v})]$.

2 Majorization

Let $\mathbf{a}, \mathbf{b} \in \mathbb{R}^n$ with components labeled in descending order $a_1 \geq \dots \geq a_n$ and $b_1 \geq \dots \geq b_n$. Then \mathbf{b} weakly majorizes \mathbf{a} , written $\mathbf{a} \prec_w \mathbf{b}$, when

$$\sum_{i=1}^k a_i \leq \sum_{i=1}^k b_i, \quad 1 \leq k \leq n \quad (2.1)$$

and majorizes $\mathbf{a} \prec \mathbf{b}$ when the final inequality is an equality. Weak log majorization $\mathbf{a} \prec_{w(\log)} \mathbf{b}$ is similarly defined for non-negative vectors as

$$\prod_{i=1}^k a_i \leq \prod_{i=1}^k b_i, \quad 1 \leq k \leq n \quad (2.2)$$

with log majorization $\mathbf{a} \prec_{(\log)} \mathbf{b}$ when the final inequality is an equality. An important fact is that log majorization and weak log majorization both imply weak majorization [2] [Lemma 1.8].

Note that it is not necessary that the vectors \mathbf{a} and \mathbf{b} be in descending order—majorization is explicitly defined with respect to the rearrangements of the values in descending order. We define all of the above majorization for matrices, ie $A \prec B$ and all variations, when the singular values considered as a vector are majorized $\sigma(A) \prec \sigma(B)$. All operators stated for majorization (ie $f(\mathbf{a})$ or $\mathbf{a}\mathbf{b}$) should be considered to be applied entrywise to the vectors (ie $(f(a_1), \dots, f(a_n))$ or $(a_1 b_1, \dots, a_n b_n)$).

Majorization holds the following vital property:

Theorem 2.1. (Hardy, Littlewood, and Pólya [9] [10]; Tomić, Weyl [16] [17]) Suppose $\mathbf{a} \prec_w \mathbf{b}$. Then for any function $f : \mathbb{R} \rightarrow \mathbb{R}$ that is increasing and convex on the domain containing all elements of \mathbf{a} and \mathbf{b} ,

$$\sum_{i=1}^n f(a_i) \leq \sum_{i=1}^n f(b_i). \quad (2.3)$$

If $\mathbf{a} \prec \mathbf{b}$, the 'increasing' requirement can be dropped.

An immediate yet highly useful lemma follows:

Lemma 2.2. Let $\mathbf{a}, \mathbf{b} \in \mathbb{R}_n^+$. Suppose $\mathbf{a} \prec_w \mathbf{b}$. Then $\mathbf{a}^s \prec_w \mathbf{b}^s$ for all $s \geq 1$.

Log majorization also allows us to characterize equality cases:

Lemma 2.3. (Hiai [11] [Lemma 2.2]) *Let $\Phi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ be a strictly convex increasing function. Then $\mathbf{a} \prec_{(\log)} \mathbf{b}$ and $\sum_{i=1}^n \Phi(a_i) = \sum_{i=1}^n \Phi(b_i)$ implies $\mathbf{a} = \Theta \mathbf{b}$ for some permutation matrix Θ .*

As exponentiating is strictly convex, an immediate corollary is $\mathbf{a} \prec_{(\log)} \mathbf{b}$ and $\mathbf{a} \prec \mathbf{b}$ implies $\mathbf{a} = \Theta \mathbf{b}$.

Majorization is an incredibly powerful technique in matrix analysis used to prove numerous inequalities about eigenvalues and singular values of matrices, powers of products of positive semidefinite matrices, Golden-Thompson-like inequalities, and more. A good overview of the techniques and important results can be found in [12], [13]. The two results that we will need for this paper will regard the eigenvalues of the sums of Hermitian matrices:

Theorem 2.4. (Fan [7]) *Let $A, B \in M_{n \times n}(\mathbb{C})$ be self-adjoint. Then*

$$\lambda(A + B) \prec \lambda(A) + \lambda(B). \quad (2.4)$$

and the singular values of products of general matrices:

Theorem 2.5. (Gel'fand and Naimark [8]) *Let $A, B \in M_{n \times n}(\mathbb{C})$. Then*

$$\sigma(AB) \prec_{(\log)} \sigma(A)\sigma(B). \quad (2.5)$$

We will also need a fairly intuitive lemma that to our knowledge has not yet been addressed in existing literature, characterizing the concatenation of majorized vectors:

Lemma 2.6. *Let $\mathbf{x} \prec_w \mathbf{y}$, and $\mathbf{a} \prec_w \mathbf{b}$ be non-negative vectors labeled in descending order. Then $\mathbf{x}\mathbf{a} \prec_w \mathbf{y}\mathbf{b}$.*

Proof. We can write the components of \mathbf{y} as $y_{n-1} = y_n + \epsilon_1$, $y_{n-2} = y_n + \epsilon_1 + \epsilon_2$, \dots , $y_1 = y_n + \epsilon_1 + \dots + \epsilon_{n-1}$ where $\epsilon_i \geq 0$. Then applying $\mathbf{a} \prec_w \mathbf{b}$

$$\epsilon_{n-1}a_1 \leq \epsilon_{n-1}b_1 \quad (2.6)$$

$$\epsilon_{n-2} \left(\sum_{i=1}^2 a_i \right) \leq \epsilon_{n-2} \left(\sum_{i=1}^2 b_i \right) \quad (2.7)$$

$$\vdots \quad (2.8)$$

$$y_n \left(\sum_{i=1}^n a_i \right) \leq y_n \left(\sum_{i=1}^n b_i \right) \quad (2.9)$$

and summing them all together,

$$\sum_{i=1}^n y_i a_i \leq \sum_{i=1}^n y_i b_i. \quad (2.10)$$

Applying the same splitting argument to a_i with $\mathbf{x} \prec_w \mathbf{y}$ gives

$$\sum_{i=1}^n x_i a_i \leq \sum_{i=1}^n y_i a_i, \quad (2.11)$$

and stringing the two inequalities together

$$\sum_{i=1}^n x_i a_i \leq \sum_{i=1}^n y_i b_i. \quad (2.12)$$

Finally, nothing that when $\mathbf{a} \prec_w \mathbf{b}$, the first k th components maintain the weak majorization relationship $(a_1, \dots, a_k) \prec_w (b_1, \dots, b_k)$, applying the argument to the k th components gives the desired result. \square

Note that the above technique can be expressed compactly as the weighted sum of Ky-Fan norms for matrices $[\mathbf{a}]$, $[\mathbf{b}]$, $[\mathbf{x}]$, and $[\mathbf{y}]$, and leveraging the matrix majorization result that $A \prec_w B$ implies $|||A||| \leq |||B|||$ for every unitarily invariant norm $||| \cdot |||$.

3 Extensions And Counterexamples

First, we address rearrangement of commuting matrices:

Theorem 3.1. *Let $A, B \in M_{n \times n}(\mathbb{C})$ be two self-adjoint matrices that commute. Then*

$$\|\sigma(A) + \sigma(B)\|_p^p + \|\sigma(A) - \sigma(B)\|_p^p \leq \|A + B\|_p^p + \|A - B\|_p^p \leq \|\sigma_\uparrow(A) + \sigma_\downarrow(B)\|_p^p + \|\sigma_\uparrow(A) - \sigma_\downarrow(B)\|_p^p \quad (3.1)$$

for $1 \leq p \leq 2$, and the inequality reverses for $p \geq 2$. Furthermore, there is equality in either inequality for $p \neq 1, 2$ if and only if A and B are aligned in the extremized arrangement.

Proof. In the mutually diagonalizable basis, we can write

$$A = \begin{bmatrix} \lambda_1(A) & & \\ & \ddots & \\ & & \lambda_n(A) \end{bmatrix}, \quad B = \begin{bmatrix} \lambda_{i_1}(B) & & \\ & \ddots & \\ & & \lambda_{i_n}(B) \end{bmatrix}. \quad (3.2)$$

Then we note that the singular values of $A + B$ and $A - B$ can be grouped as

$$\{|\lambda_j(A) \pm \lambda_{i_j}(B)|\} = \{|\lambda_j(A)| \pm |\lambda_{i_j}(B)|\} = \{|\sigma_i(A) \pm \sigma_{k_i}(B)|\}. \quad (3.3)$$

Re-labeling to preserve the pairings above, we consider the functions

$$f(x) = \sum_{i=1}^n \sigma_i(A) \chi_{[i-1, i)}(x) \quad (3.4)$$

$$g(x) = \sum_{i=1}^n \sigma_{k_i}(B) \chi_{[i-1, i)}(x) \quad (3.5)$$

and let f^* and g^* denote the spherically symmetric decreasing rearrangements.

We will need an extension of the Riesz rearrangement inequality:

Lemma 3.2. (Almgren, Lieb [1] Theorem 2.2). *Let $F : \mathbb{R}^+ \times \mathbb{R}^+ \rightarrow \mathbb{R}^+$ be a continuous function such that $F(0, 0) = 0$ and*

$$F(u_2, v_2) + F(u_1, v_1) \geq F(u_2, v_1) + F(u_1, v_2) \quad (3.6)$$

whenever $u_1 \geq u_2 \geq 0$ and $v_1 \geq v_2 \geq 0$. Then for f, g in the class \mathcal{C}_0 (ie f^*, g^* are well-defined), the inequality

$$\int F(f(x), g(x)) d\mathcal{L}^n x \leq \int F(f^*(x), g^*(x)) d\mathcal{L}^n x \quad (3.7)$$

holds. If condition 3.6 is reversed, the inequality 3.7 is reversed.

The following technique is inspired by [6] [Lemma 1.1], which in fact proves a more general theorem on symmetric decreasing arrangements of general complex functions. For the left half of our inequality, we choose

$$F(x, y) = |x + y|^p + |x - y|^p. \quad (3.8)$$

We see that $\partial^2 F(x, y) / \partial x \partial y \leq 0$ when $1 < p \leq 2$, with the inequality switching at $p = 2$, satisfying the condition of Equation 3.6. Then

$$\|f + g\|_p^p + \|f - g\|_p^p \geq \|f^* + g^*\|_p^p + \|f^* - g^*\|_p^p \quad (3.9)$$

for $1 < p \leq 2$ (and taking the limit for $p = 1$), with the inequality switching for $p \geq 2$. As $\|f \pm g\|_p^p = \|A \pm B\|_p^p$ and $\|f^* \pm g^*\|_p^p = \|\sigma(A) \pm \sigma(B)\|_p^p$, the left half of Equation 3.1 is proven.

For the right half, without loss of generality, we can assume that B is invertible; otherwise, we could consider a limit of perturbations. As the inequality for matrices A, B holds if and only if it holds for cA, cB for some scaling constant c , we can further assume that the largest singular value of B is equal to 1. We define piecewise functions such as

$$F(x, y) = \begin{cases} \left|x + \frac{1}{y}\right|^p + \left|x - \frac{1}{y}\right|^p & x \geq 0, y \geq 1 \\ e^{1-\frac{1}{y}} \left(\left|x + \frac{1}{y}\right|^p + \left|x - \frac{1}{y}\right|^p\right) & x \geq 0, 0 \leq y < 1 \end{cases} \quad (3.10)$$

for $1 \leq p \leq 2$ and

$$F(x, y) = \begin{cases} \left| x + \frac{1}{y} \right|^p + \left| x - \frac{1}{y} \right|^p & x \geq 0, y \geq 1 \\ e^{p(1-y)} (|x + y|^p + |x - y|^p) & x \geq 0, 0 \leq y < 1 \end{cases} \quad (3.11)$$

for $p \geq 2$. The values of the function F that we care about will be in the $y \geq 1$ range; these are merely examples of functions existing that satisfy the necessary conditions to apply Theorem 3.2.

It can be readily confirmed that $F(x, y)$ is continuous, and by exponential domination in the limit $F(0, 0) = 0$. We therefore calculate the partial derivative on each piece and see that $\partial^2 F(x, y)/\partial x \partial y \geq 0$ when $1 < p \leq 2$, with the inequality reversing at $p = 2$, satisfying the condition of Equation (3.6). Then letting

$$f(x) = \sum_{i=1}^n \sigma_i(A) \chi_{[i-1, i)}(x) \quad (3.12)$$

$$g(x) = \sum_{i=1}^n (\sigma_{k_i}(B))^{-1} \chi_{[i-1, i)}(x) \quad (3.13)$$

and comparing $\int F(f(x), g(x)) dx$ and $\int F(f^*(x), g^*(x)) dx$ (and taking the limit for $p = 1$), the full inequality is proven.

To characterize the equality cases, we can consider F as the limit of twice-differentiable functions with $W(x - y) = \epsilon^{-1} \exp(|x - y|/\epsilon)$ and take the limit as $\epsilon \rightarrow 0$. One can express

$$\begin{aligned} & \int \int F(f(x), g(y)) W(x - y) d\mathcal{L}^n x d\mathcal{L}^n y \\ &= \int \int F_{12}(s, t) \left[\int \int \chi_{\{f > s\}}(x) \chi_{\{g > t\}}(y) W(x - y) d\mathcal{L}^n x d\mathcal{L}^n y \right] d\mathcal{L}^1 s d\mathcal{L}^1 t \end{aligned} \quad (3.14)$$

and apply the Riesz rearrangement inequality to the interior integral. Note that for piecewise f and g as defined in Equations (3.4) and (3.5) or (3.12) and (3.13), there is strict inequality in application of the Riesz rearrangement inequality if the functions are not aligned [5]. When we take the limit, W converges to the delta distribution, and we have

$$\begin{aligned} & \int \int F_{12}(s, t) \left[\int \int \chi_{\{f > s\}}(x) \chi_{\{g > t\}}(y) \delta(x - y) d\mathcal{L}^n x d\mathcal{L}^n y \right] d\mathcal{L}^1 s d\mathcal{L}^1 t \\ & \geq \int \int F_{12}(s, t) \left[\int \int \chi_{\{f^* > s\}}(x) \chi_{\{g^* > t\}}(y) \delta(x - y) d\mathcal{L}^n x d\mathcal{L}^n y \right] d\mathcal{L}^1 s d\mathcal{L}^1 t \end{aligned} \quad (3.15)$$

Suppose, as in our case, that $F_{12}(s, t) > 0$ for $s, t > 0$, and that the matrices A and B are unaligned. Then there exists at least one interval $[i - 1, i)$ and values $0 < c_0 < c_1$ such for $x \in [i - 1, i)$ and $c_0 < c < c_1$ where $f^*(x), g^*(x) > c$, but either $f(x) > c$ and $g(x) \leq c$ or $f(x) \leq c$ and $g(x) > c$; otherwise, the matrices are in fact aligned to the equality case. Then $\chi_{\{f > s\}}(x) \chi_{\{g > t\}}(x) = 0$ and $\chi_{\{f^* > s\}}(x) \chi_{\{g^* > t\}}(x) = 1$ for $s, t \in [c_0, c_1]$. This means that the interior integral will be strictly greater when f and g are in decreasing rearrangements, and we conclude that the inequality (3.15) must be strict. \square

Next, we address the case when anticommutator $\{A, B\} = 0$.

Theorem 3.3. *Let $A, B \in M_{n \times n}(\mathbb{C})$ be self-adjoint such that $AB + BA = 0$. Then*

$$\|A + B\|_p^p + \|A - B\|_p^p \geq \|\sigma(A) + \sigma(B)\|_p^p + \|\sigma(A) - \sigma(B)\|_p^p \quad (3.16)$$

for $1 \leq p \leq 2$, with the inequality reversing for $p \geq 2$.

Proof. We note that as $\lambda(X^2) = \sigma(X^2)$ for self-adjoint X ,

$$\|A + B\|_p^p = \sum_{i=1}^n \lambda_i((A + B)^2)^{p/2} = \sum_{i=1}^n \lambda_i(A^2 + B^2)^{p/2} = \sum_{i=1}^n \lambda_i((A - B)^2)^{p/2} = \|A - B\|_p^p \quad (3.17)$$

When $1 \leq p \leq 2$, we make use of the majorization identity of Theorem 2.4 of $\lambda(A+B) \prec \lambda(A) + \lambda(B)$ and the fact that $f(x) = x^{p/2}$ is concave to conclude that

$$\|A+B\|_p^p + \|A-B\|_p^p = 2 \sum_{i=1}^n \lambda_i(A^2 + B^2)^{p/2} \quad (3.18)$$

$$\geq 2 \sum_{i=1}^n (\lambda_i(A^2) + \lambda_i(B^2))^{p/2} \quad (3.19)$$

$$= 2 \sum_{i=1}^n (\sigma_i(A)^2 + \sigma_i(B)^2)^{p/2} \quad (3.20)$$

$$= 2 \sum_{i=1}^n \left(\frac{(\sigma_i(A) + \sigma_i(B))^2}{2} + \frac{(\sigma_i(A) - \sigma_i(B))^2}{2} \right)^{p/2} \quad (3.21)$$

$$\geq \sum_{i=1}^n ((\sigma_i(A) + \sigma_i(B))^2)^{p/2} + ((\sigma_i(A) - \sigma_i(B))^2)^{p/2} \quad (3.22)$$

$$= \|\sigma(A) + \sigma(B)\|_p^p + \|\sigma(A) - \sigma(B)\|_p^p \quad (3.23)$$

An identical argument for $p \geq 2$ with reversed inequalities can be made now leveraging convexity of $x^{p/2}$.

Note that this proof extends to general A, B when $AB^* + BA^* = 0$. \square

The unitary case gives some insight to the role of the anticommutator.

Theorem 3.4. *Let $U, V \in M^{n \times n}(\mathbb{C})$ unitary. Then*

$$\|U+V\|_p^p + \|U-V\|_p^p \geq 2^p n \quad (3.24)$$

for $1 \leq p \leq 2$, with the inequality switching for $p \geq 2$. There is equality for $p \neq 2$ if and only if $U = V$. The extremization of the inequality is directly dependent on $\sigma(UV + VU)$, with greatest difference when $\{U, V\} = 0$.

Proof. Note that Equation 3.24 can be directly derived from [14] the Clarkson type inequalities

$$2(\|A\|_p^p + \|B\|_p^p) \leq \|A+B\|_p^p + \|A-B\|_p^p \leq 2^{p-1}(\|A\|_p^p + \|B\|_p^p) \quad (3.25)$$

for $p \geq 2$ and reversing for $1 \leq p \leq 2$; and in fact can be seen from direct matrix inequalities of Theorems 2.1 and 2.5 of [4]. However, we can use majorization to examine this inequality on the level of the eigenvalues to see the direct role of the anticommutator.

We can assume without loss of generality that U and V are also self-adjoint; otherwise, consider the unitary matrices

$$\widehat{U} = \begin{bmatrix} 0 & U \\ U^* & 0 \end{bmatrix}, \quad \widehat{V} = \begin{bmatrix} 0 & V \\ V^* & 0 \end{bmatrix} \quad (3.26)$$

then the inequality holds for U, V if and only if it holds for \widehat{U}, \widehat{V} by dividing by the appropriate factor of 2.

Once more we will make use of $\lambda((U \pm V)^2) = \sigma((U \pm V)^2)$. We note that $UV + VU$ is a Hermitian matrix, and as $\|UV + VU\| \leq \|UV\| + \|VU\| \leq 2\|U\|\|V\| = 2$, the eigenvalues of $UV + VU$ must be within the interval $[-2, 2]$, and can be written as $2\cos(\theta_j)$. Then

$$\|U+V\|_p^p + \|U-V\|_p^p = \|(U+V)^2\|_{p/2}^{p/2} + \|(U-V)^2\|_{p/2}^{p/2} \quad (3.27)$$

$$= \|\lambda(2I + UV + VU)\|_{p/2}^{p/2} + \|\lambda(2I - UV - VU)\|_{p/2}^{p/2} \quad (3.28)$$

$$= \|2 + \lambda(UV + VU)\|_{p/2}^{p/2} + \|2 - \lambda(UV + VU)\|_{p/2}^{p/2} \quad (3.29)$$

$$= \sum_{j=1}^n 2^{p/2} |1 + \cos(\theta_j)|^{p/2} + 2^{p/2} |1 - \cos(\theta_j)|^{p/2} \quad (3.30)$$

The function $f(\theta) = (1 + \cos(\theta))^s + (1 - \cos(\theta))^s$ can be examined on the interval $(0, \frac{\pi}{2})$. It has derivative $s \sin(\theta)[(1 - \cos(\theta))^{s-1} - (1 + \cos(\theta))^{s-1}]$, which can only be 0 at $\theta = 0, \frac{\pi}{2}$. It is immediately confirmed that the

function monotone for all $s \geq 0$ is minimized at $\theta = 0$ and maximized at $\theta = \frac{\pi}{2}$ for $0 \leq s \leq 1$, with the maximum and minimum reversing for $s \geq 1$. Therefore,

$$\sum_{j=1}^n 2^{p/2} |1 + \cos(\theta_j)|^{p/2} + 2^{p/2} |1 - \cos(\theta_j)|^{p/2} \geq \sum_{j=1}^n 2^{p/2} |2|^{p/2} = n2^p \quad (3.31)$$

and the rearrangement inequality holds as desired for $1 \leq p \leq 2$, with the inequality reversing for $p \geq 2$. As the desired extrema are reached only at $\theta = 0$, then if there is equality for $p > 2$, we must have $\theta_j = 0$ for all j , and hence $UV = VU = I$. As U is self-adjoint and unitary, we know that $U^{-1} = U$, and hence we conclude $V = U$. The alternative extrema are reached when $\theta_j = \frac{\pi}{2}$ for all j , and hence $UV + VU = 0$. \square

We finally expand upon the ranges of Conjectures 1.1 and 1.2 as originally seen in [6], and comment on how this can lead to counterexamples.

Theorem 3.5. *Let $A, B \in M_{n \times n}(\mathbb{C})$ be self-adjoint with $A \geq B \geq 0$. Then*

$$\|A + B\|_p^p + \|A - B\|_p^p \geq \|\sigma(A) + \sigma(B)\|_p^p + \|\sigma(A) - \sigma(B)\|_p^p \quad (3.32)$$

with the inequality reversing for $2 \leq p \leq 3$. There is equality for $p \neq 1, 2$ if and only if there is equality in the entire range $1 \leq p \leq 3$.

Proof. For a positive matrix C and $1 < p < 2$, for positive normalization constant k_p we have

$$C^p = k_p \int_0^\infty \left(\frac{C}{t^2} - \frac{1}{t} + \frac{1}{t+C} \right) t^p dt. \quad (3.33)$$

We can therefore express the difference between sides in Equation 3.32 for $1 < p < 2$ by the integral representation after cancellation as

$$k_p \operatorname{Tr} \left[\int_0^\infty \left(\frac{1}{A+B+t} + \frac{1}{A-B+t} - \frac{1}{\sigma(A)+\sigma(B)+t} - \frac{1}{\sigma(A)-\sigma(B)+t} \right) t^p dt \right]. \quad (3.34)$$


In [6] it is proven that when $A \geq B \geq 0$, this integrand is always positive semidefinite. Therefore, the integral is zero if and only if it is zero everywhere, if and only if Equation 3.34 is zero. This would happen independent of p , and hence if there is equality for some $1 < p < 2$, there must be equality for all $1 \leq p \leq 2$.

To extend the range to $2 \leq p \leq 3$, we see that

$$C^p = k_p C \int_0^\infty \left(\frac{C}{t^2} - \frac{1}{t} + \frac{1}{t+C} \right) t^p dt = k_p \int_0^\infty \left(\frac{C^2}{t^2} - \frac{C}{t} + \frac{C}{t+C} \right) t^p dt \quad (3.35)$$

$$= k_p \int_0^\infty \left(\frac{C^2}{t^3} - \frac{C}{t^2} + \frac{1}{t} - \frac{1}{t+C} \right) t^{p+1} dt \quad (3.36)$$

The first three terms of the integral cancel completely between each side of Equation 3.32, and now as the sign of the final term is reversed, the argument for $1 < p < 2$ is reversed. \square



The obvious question is whether or not it is possible to relax the requirement that $A \geq B \geq 0$, perhaps even to $A + B, A - B \geq 0$. The answer is: it is not.

Counterexample 3.6. The matrices

$$A = \begin{bmatrix} 6 & 0 \\ 0 & 5 \end{bmatrix}, \quad B = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \quad (3.37)$$

have the property $A + B, A - B \geq 0$, and

$$\|A+B\|_p^p + \|A-B\|_p^p \leq \|\sigma(A)+\sigma(B)\|_p^p + \|\sigma(A)-\sigma(B)\|_p^p \quad (3.38)$$

for $1 \leq p \leq 2$ and $p \geq 3$, with the inequality reversing between $2 \leq p \leq 3$.

A plot of Counterexample 3.6 can be seen in Figure 1. This counterexample hinges on the fact that we chose B to be unitary, so the “up-down” rearrangement and the “aligned” rearrangements were the same. In this case, as A and B satisfied the requirements of Theorem 3.7 (the extension of Conjecture 1.2) but not of Theorem 3.5, $\|\sigma(A) \pm \sigma(B)\|_p^p$ was treated as the “up-down” and not the “aligned” case.

Our proof of Theorem 3.7 is very similar to our proof of Theorem 3.5 which drew heavy inspiration from the proofs in [6]. However, it diverges from [6] in a very important manner: in [6], the rearrangement inequalities in the integral representation required both $A, B \geq 0$. Therefore for Conjecture 1.2, they first proved $\|A + B\|_p^p + \|A - B\|_p^p \leq \|A + |B|\|_p^p + \|A - |B|\|_p^p$ for $1 \leq p \leq 2$, then working with positive matrices A and $|B|$ addressed the rearrangement. As monotonicity of X^p was required, this does not extend as easily to $2 \leq p \leq 3$ as the proof of Theorem 3.5 did. We instead use majorization in the integral representation, removing the need to consider $|B|$ at all, which then allows us to extend the range without trouble:

Theorem 3.7. *Let $A, B \in M_{n \times n}(\mathbb{C})$ be self-adjoint with $A + B, A - B \geq 0$ and $\sigma_n(A) \geq \sigma_1(B)$. Then*

$$\|A + B\|_p^p + \|A - B\|_p^p \leq \|\sigma_\uparrow(A) + \sigma_\downarrow(B)\|_p^p + \|\sigma_\uparrow(A) - \sigma_\downarrow(B)\|_p^p \quad (3.39)$$

with the inequality reversing for $2 \leq p \leq 3$. There is equality for $p \neq 1, 2$ if and only if A and B commute and they have simultaneous diagonalizations with diagonals $\sigma_\uparrow(A)$ and $\sigma_\downarrow(B)$, and hence there is equality in the entire range $1 \leq p \leq 3$.

Proof. Once more, we use the integral representation. We can express the difference between sides in Equation 3.39 for $1 < p < 2$ by the integral representation after cancellation as

$$k_p \operatorname{Tr} \left[\int_0^\infty \left(\frac{1}{A + B + t} + \frac{1}{A - B + t} - \frac{1}{\sigma_\uparrow(A) + \sigma_\downarrow(B) + t} - \frac{1}{\sigma_\uparrow(A) - \sigma_\downarrow(B) + t} \right) t^p dt \right] \quad (3.40)$$

We will show that the integrand is always negative. We make the substitution $H = A + t$, $K = H^{-1/2} B H^{-1/2}$, then

$$(A \pm B + t)^{-1} = H^{-1/2} (I \pm K)^{-1} H^{-1/2} = H^{-1/2} \left(\sum_{n=0}^\infty (-1)^n (\pm K)^n \right) H^{-1/2} \quad (3.41)$$

and

$$\frac{1}{A + B + t} + \frac{1}{A - B + t} = 2H^{-1/2} \left(\sum_{n=0}^\infty K^{2n} \right) H^{-1/2} \quad (3.42)$$




Figure 2: $\|\sigma_{\uparrow}(C) + \sigma_{\downarrow}(D)\|_p^p + \|\sigma_{\uparrow}(C) - \sigma_{\downarrow}(D)\|_p^p - \|C + D\|_p^p - \|C - D\|_p^p$ for $1 \leq p \leq 3$, demonstrating the expected behavior on the interval $1 \leq p \leq 2$, and contrary behavior within $2 \leq p \leq 3$.




Figure 3: $\|\sigma(C) + \sigma(D)\|_p^p + \|\sigma(C) - \sigma(D)\|_p^p - \|C + D\|_p^p - \|C - D\|_p^p$ with contrary behavior within $1 \leq p \leq 2$.

For each m , we notice that K^{2m} is a positive matrix, and hence $H^{-1/2}K^{2m}H^{-1/2}$ is positive, and hence the eigenvalues and singular values are the same. Therefore,

$$\text{Tr}[H^{-1/2}K^{2m}H^{-1/2}] = \sum_{i=1}^n \sigma_i(H^{-1/2}K^{2m}H^{-1/2}) \quad (3.43)$$

$$= \sum_{i=1}^n \sigma_i(H^{-1/2}(H^{-1/2}BH^{-1/2})^{2m}H^{-1/2}) \quad (3.44)$$

$$= \sum_{i=1}^n \sigma_i(H^{-1/2}(H^{-1/2}BH^{-1/2})^m)^2 \quad (3.45)$$

$$\leq \sum_{i=1}^n \sigma_i(H^{-1/2})^2 \sigma_i((H^{-1/2}BH^{-1/2})^m)^2 \quad (3.46)$$

$$\leq \sum_{i=1}^n \sigma_i(H^{-1}) \sigma_i(H^{-1/2})^{2m} \sigma_i(B)^{2m} \sigma_i(H^{-1/2})^{2m} \quad (3.47)$$

$$= \sum_{i=1}^n \sigma_{n+1-i}(H)^{-2m-1} \sigma_i(B)^{2n} \quad (3.48)$$

This string makes repeated use of the majorization inequalities from Theorem 2.5 and Lemmas 2.2 and 2.6. Furthermore, there is equality for $p \neq 1, 2$ if and only if the integrand is always 0, and there is equality throughout. As we made use of log majorization $\sigma(AB) \prec_{(\log)} \sigma(A)\sigma(B)$, by Lemma 2.3 this must imply that $\sigma(AB) = \sigma(A)\sigma(B)$, which happens if and only if A and B commute with singular values aligned. Reversing the expansion trick from line 3.48 gives $\frac{1}{\sigma_{\uparrow}(A) \pm \sigma_{\downarrow}(B) + t}$ as desired, completing our proof for $1 < p < 2$. The same integral representation for $2 \leq p \leq 3$ as in Theorem 3.5 now extends the range. \square

An obvious counterexample to Conjecture 1.2 for all ranges are any pair of unitary matrices, as shown by Theorem 3.4. However, there are matrices that hold in the range $1 \leq p \leq 2$, but not in the range $2 \leq p \leq 3$, as demonstrated by Counterexample 3.8 and Figure 2. In fact, these matrices C and D also provide a counterexample to Conjecture 1.1, as seen in Figure 3.

Counterexample 3.8. The matrices

$$C = \begin{bmatrix} 6 & 0 \\ 0 & -1 \end{bmatrix}, \quad D = \begin{bmatrix} -1.97035 & 1.72243 \\ 1.72243 & 1.79035 \end{bmatrix} \quad (3.49)$$

are a counterexample for both Conjecture 1.1 and Conjecture 1.2, with contrary behavior for Conjecture 1.1 within the interval $1 \leq p \leq 2$; and contrary behavior for Conjecture 1.2 within the interval $2 \leq p \leq 3$.

Acknowledgements

This research was funded by the NDSEG Fellowship, Class of 2017. Thank you to my advisor, Professor Eric Carlen, for bringing my attention to the problem and providing me with a background to the subject.

References

- [1] Almgren Jr, F.J., Lieb, E.H.: Symmetric decreasing rearrangement is sometimes continuous. *Journal of the American Mathematical Society* pp. 683–773 (1989)
- [2] Ando, T.: Majorization, doubly stochastic matrices, and comparison of eigenvalues. *Linear Algebra and its Applications* **118**, 163 – 248 (1989). DOI [https://doi.org/10.1016/0024-3795\(89\)90580-6](https://doi.org/10.1016/0024-3795(89)90580-6). URL <http://www.sciencedirect.com/science/article/pii/0024379589905806>
- [3] Ball, K., Carlen, E.A., Lieb, E.H.: Sharp uniform convexity and smoothness inequalities for trace norms. *Inventiones mathematicae* **115**(1), 463–482 (1994). DOI 10.1007/BF01231769. URL <https://doi.org/10.1007/BF01231769>
- [4] Bourin, J.C., Lee, E.Y.: Clarkson-mccarthy inequalities with unitary and isometry orbits. *Linear Algebra and its Applications* **601**, 170 – 179 (2020). DOI <https://doi.org/10.1016/j.laa.2020.04.019>. URL <http://www.sciencedirect.com/science/article/pii/S0024379520302135>
- [5] Burchard, A.: Cases of equality in the riesz rearrangement inequality. *Annals of Mathematics* **143**(3), 499–527 (1996). URL <http://www.jstor.org/stable/2118534>
- [6] Carlen, E., Lieb, E.H.: Some matrix rearrangement inequalities. *Annali di Matematica Pura ed Applicata* **185**(5), S315–S324 (2006). DOI 10.1007/s10231-004-0147-z. URL <https://doi.org/10.1007/s10231-004-0147-z>
- [7] Fan, K.: Maximum properties and inequalities for the eigenvalues of completely continuous operators. *Proceedings of the National Academy of Sciences of the United States of America* **37**(11), 760–766 (1951). DOI 10.1073/pnas.37.11.760. URL <https://www.ncbi.nlm.nih.gov/pubmed/16578416>
- [8] Gel’fand, I.M., Naimark, M.A.: The relation between the unitary representations of the complex unimodular group and its unitary subgroup. *Izv. Akad. Nauk SSSR Ser. Mat.* **14**(3), 239–260 (1950)
- [9] Hardy, G.H., Littlewood, J.E., Pólya, G.: Some simple inequalities satisfied by convex functions. *Messenger Math.* **58**, 145–152 (1929). URL <https://ci.nii.ac.jp/naid/10009422169/en/>
- [10] Hardy, G.H., Polya, G.: *Inequalities*. Cambridge : Cambridge University Press (1934). Bibliography: p. [300]-314
- [11] Hiai, F.: Equality cases in matrix norm inequalities of golden-thompson type. *Linear and Multilinear Algebra* **36**(4), 239–249 (1994). DOI 10.1080/03081089408818297. URL <https://doi.org/10.1080/03081089408818297>
- [12] Hiai, F., Petz, D.: *Introduction To Matrix Analysis And Applications*, 1 edn., chap. 6, pp. 227–271. Springer International Publishing, Cham (2014)
- [13] Marshall, A.W., Olkin, I., Arnold, B.C.: *Inequalities: Theory of Majorization and Its Applications*, 2 edn. Springer, New York (2011)
- [14] McCarthy, C.: c_p cp. *Isr. J. Math.* **5**, 249–271 (1967)
- [15] Tomczak-Jaegermann, N.: The moduli of smoothness and convexity and the Rademacher averages of the trace classes $S_p(1 \leq p \leq \infty)^*$. *Studia Mathematica* **50**(2), 163–182 (1974). URL <http://eudml.org/doc/217886>
- [16] Tomić, M.: Théoreme de gauss relatif au centre de gravité et son application. *Bull. Soc. Math. Phys. Serbie* **1**, 31–40 (1949)
- [17] Weyl, H.: Inequalities between two types of eigenvalues of a linear transformation. *Proceedings of the National Academy of Sciences of the United States of America* **35**(7), 408–411 (1949)