

On Bilinear Hardy Inequality and Corresponding Geometric Mean Inequality

Amiran Gogatishvili, Pankaj Jain and Saikat Kanjilal

Abstract. The main aim of this paper to provide several scales of equivalent conditions for the bilinear Hardy inequalities in the case $1 < q, p_1, p_2 < \infty$ with $q \geq \max(p_1, p_2)$.

2010 AMS Subject Classification. 26D10, 46E35.

Key words and Phrases: Hardy inequality, bilinear Hardy inequality, integral conditions, equivalent conditions, geometric mean inequality.

1 Introduction

Let \mathfrak{M} denote the set of all Lebesgue measurable functions on (a, b) , $-\infty \leq a < b \leq \infty$, $\mathfrak{M}^+ \subset \mathfrak{M}$ is the subset of all non-negative functions.

Let $u, v, \in \mathfrak{M}^+$, $0 < p, q \leq \infty$, $p \geq 1$. Denote $p' := \frac{p}{p-1}$ and write

$$U(x) = \int_x^b u(t) dt, \quad V(x) = \int_a^x v^{1-p'}(t) dt, \quad (1.1)$$

and assume that $U(x) < \infty$, $V(x) < \infty$ for almost everywhere (a.e.) $x \in (a, b)$.

Consider the one dimensional Hardy inequality

$$\left(\int_a^b \left(\int_a^x f(t) dt \right)^q u(x) dx \right)^{\frac{1}{q}} \leq C \left(\int_a^b f^p(x)v(x) dx \right)^{\frac{1}{p}}, \quad f \in \mathfrak{M}^+. \quad (1.2)$$

It is known that the inequality (1.2) is characterized by the Muckenhoupt condition [9] in the case $1 < p \leq q < \infty$ which is given by

$$A_M := \sup_{a < x < b} A_M(x) = \sup_{a < x < b} U^{\frac{1}{q}}(x)V^{\frac{1}{p'}}(x) < \infty. \quad (1.3)$$

It is noted that the Muckenhoupt condition $A_M < \infty$ is not unique. It can be replaced by several scales of equivalent conditions (see [2], [3]). Precisely, the following result was proved in [3]:

Theorem A. Let $-\infty \leq a < b \leq \infty$, α, β, s be positive numbers and $f, g, h \in \mathfrak{M}^+$. Denote

$$F(x) := \int_x^b f(t) dt, \quad G(x) := \int_a^x g(t) dt, \quad (1.4)$$

and suppose that $F(x) < \infty$, $G(x) < \infty$ for every $x \in (a, b)$. Furthermore, denote

$$\begin{aligned} B_1(x; \alpha, \beta) &:= F^\alpha(x) G^\beta(x); \\ B_2(x; \alpha, \beta, s) &:= \left(\int_x^b f(t) G^{\frac{\beta-s}{\alpha}}(t) dt \right)^\alpha G^s(x); \\ B_3(x; \alpha, \beta, s) &:= \left(\int_a^x g(t) F^{\frac{\alpha-s}{\beta}}(t) dt \right)^\beta F^s(x); \\ B_4(x; \alpha, \beta, s) &:= \left(\int_a^x f(t) G^{\frac{\beta+s}{\alpha}}(t) dt \right)^\alpha G^{-s}(x); \\ B_5(x; \alpha, \beta, s) &:= \left(\int_x^b g(t) F^{\frac{\alpha+s}{\beta}}(t) dt \right)^\beta F^{-s}(x); \\ B_6(x; \alpha, \beta, s) &:= \left(\int_x^b f(t) G^{\frac{\beta}{\alpha+s}}(t) dt \right)^{\alpha+s} F^{-s}(x); \\ B_7(x; \alpha, \beta, s) &:= \left(\int_a^x g(t) F^{\frac{\alpha}{\beta+s}}(t) dt \right)^{\beta+s} G^{-s}(x); \\ B_8(x; \alpha, \beta, s) &:= \left(\int_a^x f(t) G^{\frac{\beta}{\alpha-s}}(t) dt \right)^{\alpha-s} F^s(x), \quad \alpha > s; \\ B_9(x; \alpha, \beta, s) &:= \left(\int_x^b f(t) G^{\frac{\beta}{\alpha-s}}(t) dt \right)^{\alpha-s} F^s(x), \quad \alpha < s; \\ B_{10}(x; \alpha, \beta, s) &:= \left(\int_x^b g(t) F^{\frac{\alpha}{\beta-s}}(t) dt \right)^{\beta-s} G^s(x), \quad \beta > s; \\ B_{11}(x; \alpha, \beta, s) &:= \left(\int_a^x g(t) F^{\frac{\alpha}{\beta-s}}(t) dt \right)^{\beta-s} G^s(x), \quad \beta < s; \\ B_{12}(x; \alpha, \beta, s; h) &:= \left(\int_x^b f(t) h^{\frac{\beta-s}{\alpha}}(t) dt \right)^\alpha \left(h(x) + G(x) \right)^s, \quad \beta < s; \\ B_{13}(x; \alpha, \beta, s; h) &:= \left(\int_a^x g(t) h^{\frac{\alpha-s}{\beta}}(t) dt \right)^\beta \left(h(x) + F(x) \right)^s, \quad \alpha < s; \\ B_{14}(x; \alpha, \beta, s; h) &:= \left(\int_a^x f(t) \left(h(t) + G(t) \right)^{\frac{\beta+s}{\alpha}} dt \right)^\alpha h^{-s}(x); \\ B_{15}(x; \alpha, \beta, s; h) &:= \left(\int_x^b g(t) \left(h(t) + F(t) \right)^{\frac{\alpha+s}{\beta}} dt \right)^\beta h^{-s}(x). \end{aligned}$$

The numbers $B_1(\alpha, \beta) := \sup_{a < x < b} B_1(x; \alpha, \beta)$, $B_i(\alpha, \beta, s) := \sup_{a < x < b} B_i(x; \alpha, \beta, s)$ ($i =$

2, 3, ..., 11) and $B_i(\alpha, \beta, s) := \inf_{h \geq 0} \sup_{a < x < b} B_i(x; \alpha, \beta, s; h)$ ($i = 12, 13, 14, 15$) are mutually equivalent.

For $\alpha = \frac{1}{q}$, $\beta = \frac{1}{p'}$, $F(x) = \int_x^b u(t) dt$ and $G(x) = \int_a^x v^{1-p'}(t) dt$, we find that

$$B_1\left(x; \frac{1}{q}, \frac{1}{p'}\right) = A_M(x)$$

so that the condition

$$\sup_{a < x < b} B_1\left(x; \frac{1}{q}, \frac{1}{p'}\right) < \infty$$

in Theorem A is the Muckenhoupt condition (1.3). Consequently, the other conditions in Theorem A are equivalent to (1.3). Let us mention that similar equivalent conditions for the inequality (1.2) for the case $1 < q < p < \infty$ were obtained by Persson, Stepanov and Wall [12].

Towards the first aim of this paper, we provide a stronger version of Theorem A by proving that, in Theorem A, the supremum over the interval (a, b) can be considered in certain truncated intervals. This is done in Section 2.

Recently, Cañestro *et al.* [1] (see also [8]) considered the weighted bilinear Hardy inequality

$$\left(\int_a^b \left(\int_a^x f(t) dt\right)^q \left(\int_a^x g(t) dt\right)^q u(x) dx\right)^{\frac{1}{q}} \leq C \left(\int_a^b f^{p_1}(x) v_1(x) dt\right)^{\frac{1}{p_1}} \times \left(\int_a^b g^{p_2}(x) v_2(x) dt\right)^{\frac{1}{p_2}} \quad (1.5)$$

and proved the following:

Theorem B. *Let $1 < q, p_1, p_2 < \infty$ with $q \geq \max(p_1, p_2)$. Let $u, v_1, v_2 \in \mathfrak{M}^+$. Then there exists a positive constant C such that the inequality (1.5) holds for all $f, g \in \mathfrak{M}^+$ if and only if $\mathcal{D} < \infty$, where*

$$\mathcal{D} := \sup_{a < x < b} \left(\int_x^b u(t) dt\right)^{\frac{1}{q}} \left(\int_a^x v_1^{1-p'_1}(t) dt\right)^{\frac{1}{p'_1}} \left(\int_a^x v_2^{1-p'_2}(t) dt\right)^{\frac{1}{p'_2}}. \quad (1.6)$$

The next aim of this paper is to prove a result of the type of Theorem A in respect of the bilinear Hardy inequality (1.5). Some of these equivalent conditions have recently been proved in [7]. In this paper, we give different proofs of some of the conditions proved in [7] and moreover, we provide several new equivalent conditions. This is done in Section 3.

Finally, in Section 4, we give a characterization for the bilinear type geometric mean inequality

$$\left(\int_0^\infty (Tf)^q(x) (Tg)^q(x) u(x) dx\right)^{\frac{1}{q}} \leq C \left(\int_0^\infty f^{p_1}(x) v_1(x) dx\right)^{\frac{1}{p_1}}$$

$$\times \left(\int_0^\infty f^{p_2}(x) v_2(x) dx \right)^{\frac{1}{p_2}},$$

where

$$(Tf)(x) := \exp \left(\frac{1}{x} \int_0^x \ln(f(t)) dt \right), \quad f \in \mathfrak{M}^+.$$

2 Improvement of the Theorem A

We prove the following theorem:

Theorem 2.1. *Under the setting of Theorem A, for $x \in (a, b)$, the following hold:*

- (i) $\sup_{x < y < b} B_1(y; \alpha, \beta) \approx \sup_{x < y < b} B_2(y; \alpha, \beta, s);$
- (ii) $\sup_{x < y < b} B_1(y; \alpha, \beta) \approx \sup_{x < y < b} \left(\int_x^y f(t) G^{\frac{\beta+s}{\alpha}}(t) dt \right)^\alpha G^{-s}(y);$
- (iii) $\sup_{x < y < b} B_1(y; \alpha, \beta) \approx \sup_{x < y < b} B_6(y; \alpha, \beta, s);$
- (iv) $\sup_{x < y < b} B_1(y; \alpha, \beta) \approx \sup_{x < y < b} \left(\int_x^y f(t) G^{\frac{\beta}{\alpha-s}}(t) dt \right)^{\alpha-s} F^s(y), \quad \alpha > s;$
- (v) $\sup_{x < y < b} B_1(y; \alpha, \beta) \approx \sup_{x < y < b} B_9(y; \alpha, \beta, s), \quad \alpha < s;$
- (vi) $\sup_{x < y < b} B_1(y; \alpha, \beta) \approx \inf_{h \geq 0} \sup_{x < y < b} B_{12}(y; \alpha, \beta, s; h), \quad \beta < s;$
- (vii) $\sup_{x < y < b} B_1(y; \alpha, \beta) \approx \inf_{h \geq 0} \sup_{x < y < b} h(y)^{-s} \left(\int_x^y f(z) (h(z) + G(z))^{\frac{\beta+s}{\alpha}} dz \right)^\alpha;$
- (viii) $\sup_{a < y < x} B_1(y; \alpha, \beta) \approx \sup_{a < y < x} B_3(y; \alpha, \beta, s);$
- (ix) $\sup_{a < y < x} B_1(y; \alpha, \beta) \approx \sup_{a < y < x} \left(\int_y^x g(t) F^{\frac{\alpha+s}{\beta}}(t) dt \right)^\beta F^{-s}(y);$
- (x) $\sup_{a < y < x} B_1(y; \alpha, \beta) \approx \sup_{a < y < x} B_7(y; \alpha, \beta, s);$
- (xi) $\sup_{a < y < x} B_1(y; \alpha, \beta) \approx \sup_{a < y < x} \left(\int_y^x g(t) F^{\frac{\alpha}{\beta-s}}(t) dt \right)^{\beta-s} G^s(y), \quad \beta > s;$
- (xii) $\sup_{a < y < x} B_1(y; \alpha, \beta) \approx \sup_{a < y < x} B_{11}(y; \alpha, \beta, s), \quad \beta < s;$
- (xiii) $\sup_{a < y < x} B_1(y; \alpha, \beta) \approx \inf_{h \geq 0} \sup_{a < y < x} B_{13}(y; \alpha, \beta, s; h), \quad \alpha < s;$

$$(xiv) \sup_{a < y < x} B_1(y; \alpha, \beta) \approx \inf_{h \geq 0} \sup_{a < y < x} h(y)^{-s} \left(\int_y^x g(z) (h(z) + F(z))^{\frac{\alpha+s}{\beta}} dz \right)^\beta.$$

Proof. (i) Let us consider the function

$$f_x(y) := \chi_{(x,b)}(y)f(y).$$

By using the equivalence $B_1(\alpha, \beta) \approx B_2(\alpha, \beta, s)$ from Theorem A, we get

$$\begin{aligned} \sup_{x < y < b} B_1(y; \alpha, \beta) &= \sup_{x < y < b} F^\alpha(y)G^\beta(y) \\ &= \sup_{a < y < b} \left(\int_y^b f_x(t) dt \right)^\alpha \left(\int_a^y g(t) dt \right)^\beta \\ &\approx \sup_{a < y < b} \left(\int_y^b f_x(t) G^{\frac{\beta-s}{\alpha}}(t) dt \right)^\alpha G^s(y) \\ &= \sup_{x < y < b} \left(\int_y^b f(t) G^{\frac{\beta-s}{\alpha}}(t) dt \right)^\alpha G^s(y) \\ &= \sup_{x < y < b} B_2(y; \alpha, \beta, s). \end{aligned}$$

(ii) Let us consider the function

$$f_x(y) := \chi_{(x,b)}(y)f(y).$$

By using the equivalence $B_1(\alpha, \beta) \approx B_4(\alpha, \beta, s)$ from Theorem A, we get

$$\begin{aligned} \sup_{x < y < b} B_1(y; \alpha, \beta) &= \sup_{x < y < b} F^\alpha(y)G^\beta(y) \\ &= \sup_{a < y < b} \left(\int_y^b f_x(t) dt \right)^\alpha \left(\int_a^y g(t) dt \right)^\beta \\ &\approx \sup_{a < y < b} \left(\int_a^y f_x(t) G^{\frac{\beta+s}{\alpha}}(t) dt \right)^\alpha G^{-s}(y) \\ &= \sup_{x < y < b} \left(\int_x^y f(t) G^{\frac{\beta+s}{\alpha}}(t) dt \right)^\alpha G^{-s}(y). \end{aligned}$$

(iii) Let us consider the function

$$f_x(y) := \chi_{(x,b)}(y)f(y).$$

By using the equivalence $B_1(\alpha, \beta) \approx B_6(\alpha, \beta, s)$ from Theorem A, we get

$$\begin{aligned} \sup_{x < y < b} B_1(y; \alpha, \beta) &= \sup_{x < y < b} F^\alpha(y)G^\beta(y) \\ &= \sup_{a < y < b} \left(\int_y^b f_x(t) dt \right)^\alpha \left(\int_a^y g(t) dt \right)^\beta \end{aligned}$$

$$\begin{aligned}
&\approx \sup_{a < y < b} \left(\int_y^b f_x(t) G^{\frac{\beta}{\alpha+s}}(t) dt \right)^{\alpha+s} F^{-s}(y) \\
&= \sup_{x < y < b} \left(\int_y^b f(t) G^{\frac{\beta}{\alpha+s}}(t) dt \right)^{\alpha+s} F^{-s}(y) \\
&= \sup_{x < y < b} B_6(y; \alpha, \beta, s).
\end{aligned}$$

(iv) Let us consider the function

$$f_x(y) := \chi_{(x,b)}(y) f(y).$$

By using the equivalence $B_1(\alpha, \beta) \approx B_8(\alpha, \beta, s)$ from Theorem A, we get

$$\begin{aligned}
\sup_{x < y < b} B_1(y; \alpha, \beta) &= \sup_{x < y < b} F^\alpha(y) G^\beta(y) \\
&= \sup_{a < y < b} \left(\int_y^b f_x(t) dt \right)^\alpha \left(\int_a^y g(t) dt \right)^\beta \\
&\approx \sup_{a < y < b} \left(\int_a^y f_x(t) G^{\frac{\beta}{\alpha-s}}(t) dt \right)^{\alpha-s} F^s(y) \\
&= \sup_{x < y < b} \left(\int_x^y f(t) G^{\frac{\beta}{\alpha-s}}(t) dt \right)^{\alpha-s} F^s(y).
\end{aligned}$$

(v) Let us consider the function

$$f_x(y) := \chi_{(x,b)}(y) f(y).$$

By using the equivalence $B_1(\alpha, \beta) \approx B_9(\alpha, \beta, s)$ from Theorem A, we get

$$\begin{aligned}
\sup_{x < y < b} B_1(y; \alpha, \beta) &= \sup_{x < y < b} F^\alpha(y) G^\beta(y) \\
&= \sup_{a < y < b} \left(\int_y^b f_x(t) dt \right)^\alpha \left(\int_a^y g(t) dt \right)^\beta \\
&\approx \sup_{a < y < b} \left(\int_y^b f_x(t) G^{\frac{\beta}{\alpha-s}}(t) dt \right)^{\alpha-s} F^s(y) \\
&= \sup_{x < y < b} \left(\int_y^b f(t) G^{\frac{\beta}{\alpha-s}}(t) dt \right)^{\alpha-s} F^s(y) \\
&= \sup_{x < y < b} B_9(y; \alpha, \beta, s).
\end{aligned}$$

(vi) Let us consider the function

$$f_x(y) := \chi_{(x,b)}(y) f(y).$$

By using the equivalence $B_1(\alpha, \beta) \approx B_{12}(\alpha, \beta, s)$ from Theorem A, we get

$$\begin{aligned}
\sup_{x < y < b} B_1(y; \alpha, \beta) &= \sup_{x < y < b} F^\alpha(y) G^\beta(y) \\
&= \sup_{a < y < b} \left(\int_y^b f_x(t) dt \right)^\alpha \left(\int_a^y g(t) dt \right)^\beta \\
&\approx \inf_{h \geq 0} \sup_{a < y < b} \left(h(y) + G(y) \right)^s \left(\int_y^b f_x(z) h(z)^{\frac{\beta-s}{\alpha}} dz \right)^\alpha \\
&= \inf_{h \geq 0} \sup_{x < y < b} \left(h(y) + G(y) \right)^s \left(\int_y^b f(z) h(z)^{\frac{\beta-s}{\alpha}} dz \right)^\alpha \\
&= \inf_{h \geq 0} \sup_{x < y < b} B_{12}(y; \alpha, \beta, s; h).
\end{aligned}$$

(vii) Let us consider the function

$$f_x(y) := \chi_{(x,b)}(y) f(y).$$

By using the equivalence $B_1(\alpha, \beta) \approx B_{14}(\alpha, \beta, s)$ from Theorem A, we get

$$\begin{aligned}
\sup_{x < y < b} B_1(y; \alpha, \beta) &= \sup_{x < y < b} F^\alpha(y) G^\beta(y) \\
&= \sup_{a < y < b} \left(\int_y^b f_x(t) dt \right)^\alpha \left(\int_a^y g(t) dt \right)^\beta \\
&\approx \inf_{h \geq 0} \sup_{a < y < b} h(y)^{-s} \left(\int_a^y f_x(z) \left(h(z) + G(z) \right)^{\frac{\beta+s}{\alpha}} dz \right)^\alpha \\
&= \inf_{h \geq 0} \sup_{x < y < b} h(y)^{-s} \left(\int_x^y f(z) \left(h(z) + G(z) \right)^{\frac{\beta+s}{\alpha}} dz \right)^\alpha.
\end{aligned}$$

(viii) Let us consider the function

$$g_x(y) := \chi_{(a,x)}(y) g(y).$$

By using the equivalence $B_1(\alpha, \beta) \approx B_3(\alpha, \beta, s)$ from Theorem A, we get

$$\begin{aligned}
\sup_{a < y < x} B_1(y; \alpha, \beta) &= \sup_{a < y < x} F^\alpha(y) G^\beta(y) \\
&= \sup_{a < y < b} \left(\int_y^b f(t) dt \right)^\alpha \left(\int_a^y g_x(t) dt \right)^\beta \\
&\approx \sup_{a < y < b} \left(\int_a^y g_x(t) F^{\frac{\alpha-s}{\beta}}(t) dt \right)^\beta F^s(y) \\
&= \sup_{a < y < x} \left(\int_a^y g(t) F^{\frac{\alpha-s}{\beta}}(t) dt \right)^\beta F^s(y) \\
&= \sup_{a < y < x} B_3(y; \alpha, \beta, s).
\end{aligned}$$

(ix) Let us consider the function

$$g_x(y) := \chi_{(a,x)}(y)g(y).$$

By using the equivalence $B_1(\alpha, \beta) \approx B_5(\alpha, \beta, s)$ from Theorem A, we get

$$\begin{aligned} \sup_{a < y < x} B_1(y; \alpha, \beta) &= \sup_{a < y < x} F^\alpha(y)G^\beta(y) \\ &= \sup_{a < y < b} \left(\int_y^b f(t) dt \right)^\alpha \left(\int_a^y g_x(t) dt \right)^\beta \\ &\approx \sup_{a < y < b} \left(\int_y^b g_x(t) F^{\frac{\alpha+s}{\beta}}(t) dt \right)^\beta F^{-s}(y) \\ &= \sup_{a < y < x} \left(\int_y^x g(t) F^{\frac{\alpha+s}{\beta}}(t) dt \right)^\beta F^{-s}(y). \end{aligned}$$

(x) Let us consider the function

$$g_x(y) := \chi_{(a,x)}(y)g(y).$$

By using the equivalence $B_1(\alpha, \beta) \approx B_7(\alpha, \beta, s)$ from Theorem A, we get

$$\begin{aligned} \sup_{a < y < x} B_1(y; \alpha, \beta) &= \sup_{a < y < x} F^\alpha(y)G^\beta(y) \\ &= \sup_{a < y < b} \left(\int_y^b f(t) dt \right)^\alpha \left(\int_a^y g_x(t) dt \right)^\beta \\ &\approx \sup_{a < y < b} \left(\int_a^y g_x(t) F^{\frac{\alpha}{\beta+s}}(t) dt \right)^{\beta+s} G^{-s}(y) \\ &= \sup_{a < y < x} \left(\int_a^y g(t) F^{\frac{\alpha}{\beta+s}}(t) dt \right)^{\beta+s} G^{-s}(y) \\ &= \sup_{a < y < x} B_7(y; \alpha, \beta, s). \end{aligned}$$

(xi) Let us consider the function

$$g_x(y) := \chi_{(a,x)}(y)g(y).$$

By using the equivalence $B_1(\alpha, \beta) \approx B_{10}(\alpha, \beta, s)$ from Theorem A, we get

$$\begin{aligned} \sup_{a < y < x} B_1(y; \alpha, \beta) &= \sup_{a < y < x} F^\alpha(y)G^\beta(y) \\ &= \sup_{a < y < b} \left(\int_y^b f(t) dt \right)^\alpha \left(\int_a^y g_x(t) dt \right)^\beta \\ &\approx \sup_{a < y < b} \left(\int_y^b g_x(t) F^{\frac{\alpha}{\beta-s}}(t) dt \right)^{\beta-s} G^s(y) \end{aligned}$$

$$= \sup_{a < y < x} \left(\int_y^x g(t) F^{\frac{\alpha}{\beta-s}}(t) dt \right)^{\beta-s} G^s(y).$$

(xii) Let us consider the function

$$g_x(y) := \chi_{(a,x)}(y)g(y).$$

By using the equivalence $B_1(\alpha, \beta) \approx B_{11}(\alpha, \beta, s)$ from Theorem A, we get

$$\begin{aligned} \sup_{a < y < x} B_1(y; \alpha, \beta) &= \sup_{a < y < x} F^\alpha(y)G^\beta(y) \\ &= \sup_{a < y < b} \left(\int_y^b f(t) dt \right)^\alpha \left(\int_a^y g_x(t) dt \right)^\beta \\ &\approx \sup_{a < y < b} \left(\int_a^y g_x(t) F^{\frac{\alpha}{\beta-s}}(t) dt \right)^{\beta-s} G^s(y) \\ &= \sup_{a < y < x} \left(\int_a^y g(t) F^{\frac{\alpha}{\beta-s}}(t) dt \right)^{\beta-s} G^s(y) \\ &= \sup_{a < y < x} B_{11}(y; \alpha, \beta, s). \end{aligned}$$

(xiii) Let us consider the function

$$g_x(y) := \chi_{(a,x)}(y)g(y).$$

By using the equivalence $B_1(\alpha, \beta) \approx B_{13}(\alpha, \beta, s)$ from Theorem A, we get

$$\begin{aligned} \sup_{a < y < x} B_1(y; \alpha, \beta) &= \sup_{x < y < b} F^\alpha(y)G^\beta(y) \\ &= \sup_{a < y < b} \left(\int_y^b f(t) dt \right)^\alpha \left(\int_a^y g_x(t) dt \right)^\beta \\ &\approx \inf_{h \geq 0} \sup_{a < y < b} \left(h(y) + F(y) \right)^s \left(\int_a^y g_x(z) h(z)^{\frac{\alpha-s}{\beta}} dz \right)^\beta \\ &= \inf_{h \geq 0} \sup_{a < y < x} \left(h(y) + F(y) \right)^s \left(\int_a^y g(z) h(z)^{\frac{\alpha-s}{\beta}} dz \right)^\beta \\ &= \inf_{h \geq 0} \sup_{a < y < x} B_{13}(y; \alpha, \beta, s; h). \end{aligned}$$

(xiv) Let us consider the function

$$g_x(y) := \chi_{(a,x)}(y)g(y).$$

By using the equivalence $B_1(\alpha, \beta) \approx B_{15}(\alpha, \beta, s)$ from Theorem A, we get

$$\sup_{a < y < x} B_1(y; \alpha, \beta) = \sup_{a < y < x} F^\alpha(y)G^\beta(y)$$

$$\begin{aligned}
&= \sup_{a < y < b} \left(\int_y^b f(t) dt \right)^\alpha \left(\int_a^y g_x(t) dt \right)^\beta \\
&\approx \inf_{h \geq 0} \sup_{a < y < b} h(y)^{-s} \left(\int_y^b g_x(z) \left(h(z) + F(z) \right)^{\frac{\alpha+s}{\beta}} dz \right)^\beta \\
&= \inf_{h \geq 0} \sup_{a < y < x} h(y)^{-s} \left(\int_y^x g(z) \left(h(z) + F(z) \right)^{\frac{\alpha+s}{\beta}} dz \right)^\beta.
\end{aligned}$$

□

Remark 2.2. Theorem 2.1 suggests that the equivalence is independent of x .

Remark 2.3. Theorem 2.1 demonstrates that, in Theorem A, the supremum over the interval (a, b) can be considered in certain truncated intervals. This consideration will be useful in the results of next section. Moreover, Theorem A can be obtained by Theorem 2.1 by taking $x = a$ or $x = b$ appropriately.

3 Equivalence Theorem for Bilinear Hardy Inequality

We prove the following:

Theorem 3.1. Let $-\infty \leq a < b \leq \infty$, $\alpha, \beta, \gamma, s, s_1, s_2$ be positive numbers and $f, g, h, h_1, h_2 \in \mathfrak{M}^+$. Let F, G be as in (1.4) and denote

$$H(x) := \int_a^x h(t) dt. \quad (3.1)$$

Assume that $F(x)$, $G(x)$ and $H(x)$ are finite for all $x \in (a, b)$. Consider

$$\tilde{B}_1(x; \alpha, \beta, \gamma) := F^\alpha(x) G^\beta(x) H^\gamma(x);$$

$$\tilde{B}_2(x; \alpha, \beta, \gamma, s_1, s_2) := \left(\int_x^b f(t) G^{\frac{\beta-s_1}{\alpha}}(t) H^{\frac{\gamma-s_2}{\alpha}}(t) dt \right)^\alpha G^{s_1}(x) H^{s_2}(x);$$

$$\tilde{B}_3(x; \alpha, \beta, \gamma, s_1, s_2) := \left(\int_a^x f(t) G^{\frac{\beta+s_1}{\alpha}}(t) H^{\frac{\gamma+s_2}{\alpha}}(t) dt \right)^\alpha G^{-s_1}(x) H^{-s_2}(x);$$

$$\tilde{B}_4(x; \alpha, \beta, \gamma, s_1, s_2) := \left(\int_a^x f(t) G^{\frac{\beta}{\alpha+s_1}}(t) H^{\frac{\gamma-s_2}{\alpha+s_1}}(t) dt \right)^{\alpha+s_1} F^{-s_1}(x) H^{s_2}(x);$$

$$\tilde{B}_5(x; \alpha, \beta, \gamma, s_1, s_2) := \left(\int_a^x f(t) G^{\frac{\beta-s_2}{\alpha+s_1}}(t) H^{\frac{\gamma}{\alpha+s_1}}(t) dt \right)^{\alpha+s_1} F^{-s_1}(x) G^{s_2}(x);$$

$$\tilde{B}_6(x; \alpha, \beta, \gamma, s_1, s_2) := \left(\int_a^x f(t) G^{\frac{\beta}{\alpha-s_1}}(t) H^{\frac{\gamma+s_2}{\alpha-s_1}}(t) dt \right)^{\alpha-s_1} F^{s_1}(x) H^{-s_2}(x), \quad \alpha > s_1;$$

$$\tilde{B}_7(x; \alpha, \beta, \gamma, s_1, s_2) := \left(\int_a^x f(t) G^{\frac{\beta+s_2}{\alpha-s_1}}(t) H^{\frac{\gamma}{\alpha-s_1}}(t) dt \right)^{\alpha-s_1} F^{s_1}(x) G^{-s_2}(x), \quad \alpha > s_1;$$

$$\tilde{B}_8(x; \alpha, \beta, \gamma, s_1, s_2) := \left(\int_x^b f(t) G^{\frac{\beta}{\alpha-s_1}}(t) dt \right)^{\alpha-s_1} H^{s_2}(x) \left(\int_x^b f(t) H^{\frac{\gamma-s_2}{s_1}}(t) dt \right)^{s_1}, \quad \alpha < s_1;$$

$$\begin{aligned}
\tilde{B}_9(x; \alpha, \beta, \gamma, s_1, s_2) &:= \left(\int_x^b f(t) H^{\frac{\gamma}{\alpha-s_1}}(t) dt \right)^{\alpha-s_1} G^{s_2}(x) \left(\int_x^b f(t) G^{\frac{\beta-s_2}{s_1}}(t) dt \right)^{s_1}, \quad \alpha < s_1; \\
\tilde{B}_{10}(x; \alpha, \beta, \gamma, s_1, s_2) &:= \left(\int_a^x g(t) F^{\frac{\alpha/2}{\beta+s_1}}(t) dt \right)^{\beta+s_1} \left(\int_a^x h(t) F^{\frac{\alpha/2}{\gamma+s_2}}(t) dt \right)^{\gamma+s_2} G^{-s_1}(x) H^{-s_2}(x); \\
\tilde{B}_{11}(x; \alpha, \beta, \gamma, s) &:= \left(\int_x^b f(t) G^{\frac{\beta(1-s)}{\alpha}}(t) H^{\frac{\gamma(1-s)}{\alpha}}(t) dt \right)^{\alpha} G^{\beta s}(x) H^{\gamma s}(x); \\
\tilde{B}_{12}(x; \alpha, \beta, \gamma, s) &:= \left(\int_a^x f(t) G^{\frac{\beta(1+s)}{\alpha}}(t) H^{\frac{\gamma(1+s)}{\alpha}}(t) dt \right)^{\alpha} G^{-\beta s}(x) H^{-\gamma s}; \\
\tilde{B}_{13}(x; \alpha, \beta, \gamma, s) &:= \left(\int_x^b f(t) G^{\frac{\beta}{\alpha+s}}(t) H^{\frac{\gamma}{\alpha+s}}(t) dt \right)^{\alpha+s} F^{-s}(x); \\
\tilde{B}_{14}(x; \alpha, \beta, \gamma, s) &:= \left(\int_a^x f(t) G^{\frac{\beta}{\alpha-s}}(t) H^{\frac{\gamma}{\alpha-s}}(t) dt \right)^{\alpha-s} F^s(x), \quad \alpha > s; \\
\tilde{B}_{15}(x; \alpha, \beta, \gamma, s) &:= \left(\int_x^b f(t) G^{\frac{\beta}{\alpha-s}}(t) H^{\frac{\gamma}{\alpha-s}}(t) dt \right)^{\alpha-s} F^s(x), \quad \alpha < s; \\
\tilde{B}_{16}(x; \alpha, \beta, \gamma, s; h_1) &:= \left(\int_x^b f(t) h_1^{\frac{\gamma-s}{\alpha}}(t) dt \right)^{\alpha} G^{\beta}(x) (h_1(x) + H(x))^s, \quad \gamma < s; \\
\tilde{B}_{17}(x; \alpha, \beta, \gamma, s; h_1) &:= \left(\int_x^b f(t) h_1^{\frac{\beta-s}{\alpha}}(t) dt \right)^{\alpha} H^{\gamma}(x) (h_1(x) + G(x))^s, \quad \beta < s; \\
\tilde{B}_{18}(x; \alpha, \beta, \gamma, s_1, s_2; h_1, h_2) &:= \left(\int_x^b f(t) h_1^{\frac{\beta-s_1}{\alpha}}(t) h_2^{\frac{\gamma-s_2}{\alpha}}(t) dt \right)^{\alpha} \left\{ \sup_{a < y < x} (h_1(y) + G(y))^{s_1} \right\} \times \\
&\quad \times (h_2(x) + H(x))^{s_2}, \quad \beta < s_1, \gamma < s_2; \\
\tilde{B}_{19}(x; \alpha, \beta, \gamma, s_1, s_2; h_1, h_2) &:= \left(\int_x^b f(t) h_1^{\frac{\beta-s_1}{\alpha}}(t) h_2^{\frac{\gamma-s_2}{\alpha}}(t) dt \right)^{\alpha} (h_1(x) + G(x))^{s_1} \times \\
&\quad \times \left\{ \sup_{a < y < x} (h_2(y) + H(y))^{s_2} \right\}, \quad \beta < s_1, \gamma < s_2; \\
\tilde{B}_{20}(x; \alpha, \beta, \gamma, s_1, s_2; h_1, h_2) &:= \left(\int_x^b f(t) h_1^{\frac{\beta-s_1}{\alpha}}(t) h_2^{\frac{\gamma-s_2}{\alpha}}(t) dt \right)^{\alpha} \left\{ \sup_{a < y < x} (h_1(y) + G(y))^{s_1} \right\} \times \\
&\quad \times \left\{ \sup_{a < y < x} (h_2(y) + H(y))^{s_2} \right\}, \quad \beta < s_1, \gamma < s_2; \\
\tilde{B}_{21}(x; \alpha, \beta, \gamma, s_1, s_2; h_1, h_2) &:= \left(\int_a^x f(t) (h_1(t) + G(t))^{\frac{\beta+s_1}{\alpha}} (h_2(t) + H(t))^{\frac{\gamma+s_2}{\alpha}} dt \right)^{\alpha} \times \\
&\quad \times \left\{ \sup_{x < y < b} h_1^{-s_1}(y) \right\} h_2^{-s_2}(x); \\
\tilde{B}_{22}(x; \alpha, \beta, \gamma, s_1, s_2; h_1, h_2) &:= \left(\int_a^x f(t) (h_1(t) + G(t))^{\frac{\beta+s_1}{\alpha}} (h_2(t) + H(t))^{\frac{\gamma+s_2}{\alpha}} dt \right)^{\alpha} \times
\end{aligned}$$

$$\begin{aligned} \tilde{B}_{23}(x; \alpha, \beta, \gamma, s_1, s_2; h_1, h_2) &:= \left(\int_a^x f(t) (h_1(t) + G(t))^{\frac{\beta+s_1}{\alpha}} (h_2(t) + H(t))^{\frac{\gamma+s_2}{\alpha}} dt \right)^\alpha \\ &\quad \times h_1^{-s_1}(x) \left\{ \sup_{x < y < b} h_2^{-s_2}(y) \right\}; \\ &\quad \times \left\{ \sup_{x < y < b} h_1^{-s_1}(y) \right\} \left\{ \sup_{x < y < b} h_2^{-s_2}(y) \right\}. \end{aligned}$$

The numbers

$$\begin{aligned} \tilde{B}_1(\alpha, \beta, \gamma) &:= \sup_{a < x < b} \tilde{B}_1(x; \alpha, \beta, \gamma), \\ \tilde{B}_i(\alpha, \beta, \gamma, s_1, s_2) &:= \sup_{a < x < b} \tilde{B}_i(x; \alpha, \beta, \gamma, s_1, s_2); \quad (i = 2, 3, 4, 5, 6, 7, 8, 9, 10), \\ \tilde{B}_i(\alpha, \beta, \gamma, s) &:= \sup_{a < x < b} \tilde{B}_i(x; \alpha, \beta, \gamma, s); \quad (i = 11, 12, 13, 14, 15), \\ \tilde{B}_i(\alpha, \beta, \gamma, s) &:= \inf_{h_1 \geq 0} \sup_{a < x < b} \tilde{B}_i(x; \alpha, \beta, \gamma, s; h_1); \quad (i = 16, 17) \end{aligned}$$

and

$$\tilde{B}_i(\alpha, \beta, \gamma, s_1, s_2) := \inf_{h_1, h_2 \geq 0} \sup_{a < x < b} \tilde{B}_i(x; \alpha, \beta, \gamma, s_1, s_2; h_1, h_2); \quad (i = 18, 19, 20, 21, 22, 23)$$

are mutually equivalent in the sense that if one number is finite then the others are so.

Proof. The equivalence $\tilde{B}_1(\alpha, \beta, \gamma) \approx \tilde{B}_i(\alpha, \beta, \gamma, s_1, s_2) \approx \tilde{B}_j(\alpha, \beta, \gamma, s)$, ($i = 2, 3, 10$), ($j = 13, 14, 15$) has been proved in [7]. Here, we give different proofs for the equivalence

$$\tilde{B}_1(\alpha, \beta, \gamma) \approx \tilde{B}_k(\alpha, \beta, \gamma, s_1, s_2), \quad (k = 2, 3)$$

and prove the remaining ones.

$$(i) \quad \tilde{B}_1(\alpha, \beta, \gamma) \approx \tilde{B}_2(\alpha, \beta, \gamma, s_1, s_2)$$

By using the equivalence $B_1(\alpha, \beta) \approx B_2(\alpha, \beta, s)$ from Theorem A and in view of Theorem 2.1 (i)

$$\sup_{x < y < b} F^\alpha(y) G^\beta(y) \approx \sup_{x < y < b} \left(\int_y^b f(t) G^{\frac{\beta-s_1}{\alpha}}(t) dt \right)^\alpha G^{s_1}(y),$$

we get

$$\begin{aligned} \tilde{B}_1(\alpha, \beta, \gamma) &= \sup_{a < x < b} \tilde{B}_1(x; \alpha, \beta, \gamma) \\ &= \sup_{a < x < b} F^\alpha(x) G^\beta(x) H^\gamma(x) \end{aligned}$$

$$\begin{aligned}
&= \sup_{a < x < b} H^\gamma(x) \sup_{x < y < b} F^\alpha(y) G^\beta(y) \\
&\approx \sup_{a < x < b} H^\gamma(x) \sup_{x < y < b} G^{s_1}(y) \left(\int_y^b f(t) G^{\frac{\beta-s_1}{\alpha}}(t) dt \right)^\alpha \\
&= \sup_{a < x < b} G^{s_1}(x) \sup_{x < y < b} H^\gamma(y) \left(\int_y^b f(t) G^{\frac{\beta-s_1}{\alpha}}(t) dt \right)^\alpha \\
&\approx \sup_{a < x < b} G^{s_1}(y) \sup_{x < y < b} H^{s_2}(y) \left(\int_y^b f(t) G^{\frac{\beta-s_1}{\alpha}}(t) H^{\frac{\gamma-s_2}{\alpha}}(t) dt \right)^\alpha \\
&= \sup_{a < x < b} \left(\int_x^b f(t) G^{\frac{\beta-s_1}{\alpha}}(t) H^{\frac{\gamma-s_2}{\alpha}}(t) dt \right)^\alpha G^{s_1}(x) H^{s_2}(x) \\
&= \sup_{a < x < b} \tilde{B}_2(x; \alpha, \beta, \gamma, s_1, s_2) \\
&= \tilde{B}_2(\alpha, \beta, \gamma, s_1, s_2).
\end{aligned}$$

$$(ii) \quad \tilde{B}_1(\alpha, \beta, \gamma) \approx \tilde{B}_3(\alpha, \beta, \gamma, s_1, s_2)$$

By using the equivalence $B_1(\alpha, \beta) \approx B_4(\alpha, \beta, s)$ from Theorem A and in view of Theorem 2.1 (ii)

$$\sup_{x < y < b} F^\alpha(y) G^\beta(y) \approx \sup_{x < y < b} \left(\int_x^y f(t) G^{\frac{\beta+s_1}{\alpha}}(t) dt \right)^\alpha G^{-s_1}(y),$$

we get

$$\begin{aligned}
\tilde{B}_1(\alpha, \beta, \gamma) &= \sup_{a < x < b} \tilde{B}_1(x; \alpha, \beta, \gamma) \\
&= \sup_{a < x < b} F^\alpha(x) G^\beta(x) H^\gamma(x) \\
&= \sup_{a < x < b} H^\gamma(x) \sup_{x < y < b} F^\alpha(y) G^\beta(y) \\
&\approx \sup_{a < x < b} H^\gamma(x) \sup_{x < y < b} G^{-s_1}(y) \left(\int_x^y f(t) G^{\frac{\beta+s_1}{\alpha}}(t) dt \right)^\alpha \\
&= \sup_{a < y < b} G^{-s_1}(y) \sup_{a < x < y} H^\gamma(x) \left(\int_x^y f(t) G^{\frac{\beta+s_1}{\alpha}}(t) dt \right)^\alpha \\
&\approx \sup_{a < y < b} G^{-s_1}(y) \sup_{a < x < y} H^{-s_2}(x) \left(\int_a^x f(t) G^{\frac{\beta+s_1}{\alpha}}(t) H^{\frac{\gamma+s_2}{\alpha}}(t) dt \right)^\alpha \\
&= \sup_{a < y < b} \left(\int_a^y f(t) G^{\frac{\beta+s_1}{\alpha}}(t) H^{\frac{\gamma+s_2}{\alpha}}(t) dt \right)^\alpha G^{-s_1}(y) H^{-s_2}(y) \\
&= \sup_{a < x < b} \left(\int_a^x f(t) G^{\frac{\beta+s_1}{\alpha}}(t) H^{\frac{\gamma+s_2}{\alpha}}(t) dt \right)^\alpha G^{-s_1}(x) H^{-s_2}(x)
\end{aligned}$$

$$\begin{aligned}
&= \sup_{a < x < b} \tilde{B}_3(x; \alpha, \beta, \gamma, s_1, s_2) \\
&= \tilde{B}_3(\alpha, \beta, \gamma, s_1, s_2).
\end{aligned}$$

$$(iii) \quad \tilde{B}_1(\alpha, \beta, \gamma) \approx \tilde{B}_4(\alpha, \beta, \gamma, s_1, s_2)$$

By using the equivalences $B_1(\alpha, \beta) \approx B_6(\alpha, \beta, s)$, $B_1(\alpha, \beta) \approx B_2(\alpha, \beta, s)$ from Theorem A and in view of Theorem 2.1 (iii)

$$\sup_{x < y < b} F^\alpha(y)G^\beta(y) \approx \sup_{x < y < b} \left(\int_y^b f(t)G^{\frac{\beta}{\alpha+s_1}}(t) dt \right)^{\alpha+s_1} F^{-s_1}(y),$$

we get

$$\begin{aligned}
\tilde{B}_1(\alpha, \beta, \gamma) &= \sup_{a < x < b} \tilde{B}_1(x; \alpha, \beta, \gamma) \\
&= \sup_{a < x < b} F^\alpha(x)G^\beta(x)H^\gamma(x) \\
&= \sup_{a < x < b} H^\gamma(x) \sup_{x < y < b} F^\alpha(y)G^\beta(y) \\
&\approx \sup_{a < x < b} H^\gamma(x) \sup_{x < y < b} \left(\int_y^b f(t)G^{\frac{\beta}{\alpha+s_1}}(t) dt \right)^{\alpha+s_1} F^{-s_1}(y) \\
&= \sup_{a < x < b} F^{-s_1}(x) \sup_{x < y < b} \left(\int_y^b f(t)G^{\frac{\beta}{\alpha+s_1}}(t) dt \right)^{\alpha+s_1} H^\gamma(y) \\
&\approx \sup_{a < x < b} F^{-s_1}(x) \sup_{x < y < b} \left(\int_y^b f(t)G^{\frac{\beta}{\alpha+s_1}}(t)H^{\frac{\gamma-s_2}{\alpha+s_1}}(t) dt \right)^{\alpha+s_1} H^{s_2}(y) \\
&= \sup_{a < x < b} F^{-s_1}(x)H^{s_2}(x) \left(\int_x^b f(t)G^{\frac{\beta}{\alpha+s_1}}(t)H^{\frac{\gamma-s_2}{\alpha+s_1}}(t) dt \right)^{\alpha+s_1} \\
&= \sup_{a < x < b} \tilde{B}_4(x; \alpha, \beta, \gamma, s_1, s_2) \\
&= \tilde{B}_4(\alpha, \beta, \gamma, s_1, s_2).
\end{aligned}$$

$$(iv) \quad \tilde{B}_1(\alpha, \beta, \gamma) \approx \tilde{B}_5(\alpha, \beta, \gamma, s_1, s_2)$$

By using the equivalences $B_1(\alpha, \beta) \approx B_6(\alpha, \beta, s)$, $B_1(\alpha, \beta) \approx B_2(\alpha, \beta, s)$ from Theorem A and in view of Theorem 2.1 (iii)

$$\sup_{x < y < b} F^\alpha(y)H^\gamma(y) \approx \sup_{x < y < b} \left(\int_y^b f(t)H^{\frac{\gamma}{\alpha+s_1}}(t) dt \right)^{\alpha+s_1} F^{-s_1}(y),$$

we get

$$\begin{aligned}
\tilde{B}_1(\alpha, \beta, \gamma) &= \sup_{a < x < b} \tilde{B}_1(x; \alpha, \beta, \gamma) \\
&= \sup_{a < x < b} F^\alpha(x) G^\beta(x) H^\gamma(x) \\
&= \sup_{a < x < b} G^\beta(x) \sup_{x < y < b} F^\alpha(y) H^\gamma(y) \\
&\approx \sup_{a < x < b} G^\beta(x) \sup_{x < y < b} \left(\int_y^b f(t) H^{\frac{\gamma}{\alpha+s_1}}(t) dt \right)^{\alpha+s_1} F^{-s_1}(y) \\
&= \sup_{a < x < b} F^{-s_1}(x) \sup_{x < y < b} \left(\int_y^b f(t) H^{\frac{\gamma}{\alpha+s_1}}(t) dt \right)^{\alpha+s_1} G^\beta(y) \\
&\approx \sup_{a < x < b} F^{-s_1}(x) \sup_{x < y < b} \left(\int_y^b f(t) H^{\frac{\gamma}{\alpha+s_1}}(t) G^{\frac{\beta-s_2}{\alpha+s_1}}(t) dt \right)^{\alpha+s_1} G^{s_2}(y) \\
&= \sup_{a < x < b} F^{-s_1}(x) G^{s_2}(x) \left(\int_x^b f(t) H^{\frac{\gamma}{\alpha+s_1}}(t) G^{\frac{\beta-s_2}{\alpha+s_1}}(t) dt \right)^{\alpha+s_1} \\
&= \sup_{a < x < b} \tilde{B}_5(x; \alpha, \beta, \gamma, s_1, s_2) \\
&= \tilde{B}_5(\alpha, \beta, \gamma, s_1, s_2).
\end{aligned}$$

$$(v) \quad \tilde{B}_1(\alpha, \beta, \gamma) \approx \tilde{B}_6(\alpha, \beta, \gamma, s_1, s_2)$$

By using the equivalences $B_1(\alpha, \beta) \approx B_8(\alpha, \beta, s)$, $B_1(\alpha, \beta) \approx B_4(\alpha, \beta, s)$ from Theorem A and in view of Theorem 2.1 (iv)

$$\sup_{x < y < b} G^\beta(y) F^\alpha(y) \approx \sup_{x < y < b} \left(\int_x^y f(t) G^{\frac{\beta}{\alpha-s_1}}(t) dt \right)^{\alpha-s_1} F^{s_1}(y); \quad \alpha > s_1,$$

we have

$$\begin{aligned}
\tilde{B}_1(\alpha, \beta, \gamma) &= \sup_{a < x < b} \tilde{B}_1(x; \alpha, \beta, \gamma) \\
&= \sup_{a < x < b} F^\alpha(x) G^\beta(x) H^\gamma(x) \\
&= \sup_{a < x < b} H^\gamma(x) \sup_{x < y < b} F^\alpha(y) G^\beta(y) \\
&\approx \sup_{a < x < b} H^\gamma(x) \sup_{x < y < b} \left(\int_x^y f(t) G^{\frac{\beta}{\alpha-s_1}}(t) dt \right)^{\alpha-s_1} F^{s_1}(y) \\
&= \sup_{a < x < b} F^{s_1}(x) \sup_{a < y < x} H^\gamma(y) \left(\int_y^x f(t) G^{\frac{\beta}{\alpha-s_1}}(t) dt \right)^{\alpha-s_1} \\
&\approx \sup_{a < x < b} F^{s_1}(x) \sup_{a < y < x} H^{-s_2}(y) \left(\int_a^y f(t) H^{\frac{\gamma+s_2}{\alpha-s_1}}(t) G^{\frac{\beta}{\alpha-s_1}}(t) dt \right)^{\alpha-s_1}
\end{aligned}$$

$$\begin{aligned}
&= \sup_{a < x < b} F^{s_1}(x) H^{-s_2}(x) \left(\int_a^x f(t) H^{\frac{\gamma+s_2}{\alpha-s_1}}(t) G^{\frac{\beta}{\alpha-s_1}}(t) dt \right)^{\alpha-s_1} \\
&= \sup_{a < x < b} \tilde{B}_6(x; \alpha, \beta, \gamma, s_1, s_2) \\
&= \tilde{B}_6(\alpha, \beta, \gamma, s_1, s_2).
\end{aligned}$$

$$(vi) \quad \tilde{B}_1(\alpha, \beta, \gamma) \approx \tilde{B}_7(\alpha, \beta, \gamma, s_1, s_2)$$

By using the equivalences $B_1(\alpha, \beta) \approx B_8(\alpha, \beta, s)$, $B_1(\alpha, \beta) \approx B_4(\alpha, \beta, s)$ from Theorem A and in view of Theorem 2.1 (iv)

$$\sup_{x < y < b} H^\gamma(y) F^\alpha(y) \approx \sup_{x < y < b} \left(\int_x^y f(t) H^{\frac{\gamma}{\alpha-s_1}}(t) dt \right)^{\alpha-s_1} F^{s_1}(y); \quad \alpha > s_1,$$

we have

$$\begin{aligned}
\tilde{B}_1(\alpha, \beta, \gamma) &= \sup_{a < x < b} \tilde{B}_1(x; \alpha, \beta, \gamma) \\
&= \sup_{a < x < b} F^\alpha(x) G^\beta(x) H^\gamma(x) \\
&= \sup_{a < x < b} G^\beta(x) \sup_{x < y < b} F^\alpha(y) H^\gamma(y) \\
&\approx \sup_{a < x < b} G^\beta(x) \sup_{x < y < b} \left(\int_x^y f(t) H^{\frac{\gamma}{\alpha-s_1}}(t) dt \right)^{\alpha-s_1} F^{s_1}(y) \\
&= \sup_{a < x < b} F^{s_1}(x) \sup_{a < y < x} G^\beta(y) \left(\int_y^x f(t) H^{\frac{\gamma}{\alpha-s_1}}(t) dt \right)^{\alpha-s_1} \\
&\approx \sup_{a < x < b} F^{s_1}(x) \sup_{a < y < x} G^{-s_2}(y) \left(\int_a^y f(t) G^{\frac{\beta+s_2}{\alpha-s_1}}(t) H^{\frac{\gamma}{\alpha-s_1}}(t) dt \right)^{\alpha-s_1} \\
&= \sup_{a < x < b} F^{s_1}(x) G^{-s_2}(x) \left(\int_a^x f(t) G^{\frac{\beta+s_2}{\alpha-s_1}}(t) H^{\frac{\gamma}{\alpha-s_1}}(t) dt \right)^{\alpha-s_1} \\
&= \sup_{a < x < b} \tilde{B}_7(x; \alpha, \beta, \gamma, s_1, s_2) \\
&= \tilde{B}_7(\alpha, \beta, \gamma, s_1, s_2).
\end{aligned}$$

$$(vii) \quad \tilde{B}_1(\alpha, \beta, \gamma) \approx \tilde{B}_8(\alpha, \beta, \gamma, s_1, s_2)$$

By using the equivalences $B_1(\alpha, \beta) \approx B_9(\alpha, \beta, s)$, $B_1(\alpha, \beta) \approx B_2(\alpha, \beta, s)$ from Theorem A and in view of Theorem 2.1 (v)

$$\sup_{x < y < b} G^\beta(y) F^\alpha(y) \approx \sup_{x < y < b} \left(\int_y^b f(t) G^{\frac{\beta}{\alpha-s_1}}(t) dt \right)^{\alpha-s_1} F^{s_1}(y); \quad \alpha < s_1,$$

we have

$$\begin{aligned}
\tilde{B}_1(\alpha, \beta, \gamma) &= \sup_{a < x < b} \tilde{B}_1(x; \alpha, \beta, \gamma) \\
&= \sup_{a < x < b} F^\alpha(x) G^\beta(x) H^\gamma(x) \\
&= \sup_{a < x < b} H^\gamma(x) \sup_{x < y < b} F^\alpha(y) G^\beta(y) \\
&\approx \sup_{a < x < b} H^\gamma(x) \sup_{x < y < b} \left(\int_y^b f(t) G^{\frac{\beta}{\alpha-s_1}}(t) dt \right)^{\alpha-s_1} F^{s_1}(y) \\
&= \sup_{a < x < b} \left(\int_x^b f(t) G^{\frac{\beta}{\alpha-s_1}}(t) dt \right)^{\alpha-s_1} F^{s_1}(x) H^\gamma(x) \\
&= \sup_{a < x < b} \left(\int_x^b f(t) G^{\frac{\beta}{\alpha-s_1}}(t) dt \right)^{\alpha-s_1} \sup_{x < y < b} F^{s_1}(y) H^\gamma(y) \\
&\approx \sup_{a < x < b} \left(\int_x^b f(t) G^{\frac{\beta}{\alpha-s_1}}(t) dt \right)^{\alpha-s_1} \sup_{x < y < b} H^{s_2}(y) \left(\int_y^b f(t) H^{\frac{\gamma-s_2}{s_1}}(t) dt \right)^{s_1} \\
&= \sup_{a < x < b} \left(\int_x^b f(t) G^{\frac{\beta}{\alpha-s_1}}(t) dt \right)^{\alpha-s_1} H^{s_2}(x) \left(\int_x^b f(t) H^{\frac{\gamma-s_2}{s_1}}(t) dt \right)^{s_1} \\
&= \sup_{a < x < b} \tilde{B}_8(x; \alpha, \beta, \gamma, s_1, s_2) \\
&= \tilde{B}_8(\alpha, \beta, \gamma, s_1, s_2).
\end{aligned}$$

$$(viii) \quad \tilde{B}_1(\alpha, \beta, \gamma) \approx \tilde{B}_9(\alpha, \beta, \gamma, s_1, s_2)$$

By using the equivalences $B_1(\alpha, \beta) \approx B_9(\alpha, \beta, s)$, $B_1(\alpha, \beta) \approx B_2(\alpha, \beta, s)$ from Theorem A and in view of Theorem 2.1 (v)

$$\sup_{x < y < b} H^\gamma(y) F^\alpha(y) \approx \sup_{x < y < b} \left(\int_y^b f(t) H^{\frac{\gamma}{\alpha-s_1}}(t) dt \right)^{\alpha-s_1} F^{s_1}(y); \quad \alpha < s_1,$$

we have

$$\begin{aligned}
\tilde{B}_1(\alpha, \beta, \gamma) &= \sup_{a < x < b} \tilde{B}_1(x; \alpha, \beta, \gamma) \\
&= \sup_{a < x < b} F^\alpha(x) G^\beta(x) H^\gamma(x) \\
&= \sup_{a < x < b} G^\beta(x) \sup_{x < y < b} F^\alpha(y) H^\gamma(y) \\
&\approx \sup_{a < x < b} G^\beta(x) \sup_{x < y < b} \left(\int_y^b f(t) H^{\frac{\gamma}{\alpha-s_1}}(t) dt \right)^{\alpha-s_1} F^{s_1}(y)
\end{aligned}$$

$$\begin{aligned}
&= \sup_{a < x < b} G^\beta(x) F^{s_1}(x) \left(\int_x^b f(t) H^{\frac{\gamma}{\alpha-s_1}}(t) dt \right)^{\alpha-s_1} \\
&= \sup_{a < x < b} \left(\int_x^b f(t) H^{\frac{\gamma}{\alpha-s_1}}(t) dt \right)^{\alpha-s_1} \sup_{x < y < b} F^{s_1}(y) G^\beta(y) \\
&\approx \sup_{a < x < b} \left(\int_x^b f(t) H^{\frac{\gamma}{\alpha-s_1}}(t) dt \right)^{\alpha-s_1} \sup_{x < y < b} G^{s_2}(y) \left(\int_y^b f(t) G^{\frac{\beta-s_2}{s_1}}(t) dt \right)^{s_1} \\
&= \sup_{a < x < b} \left(\int_x^b f(t) H^{\frac{\gamma}{\alpha-s_1}}(t) dt \right)^{\alpha-s_1} G^{s_2}(x) \left(\int_x^b f(t) G^{\frac{\beta-s_2}{s_1}}(t) dt \right)^{s_1} \\
&= \sup_{a < x < b} \tilde{B}_9(x; \alpha, \beta, \gamma, s_1, s_2) \\
&= \tilde{B}_9(\alpha, \beta, \gamma, s_1, s_2).
\end{aligned}$$

$$(ix) \quad \tilde{B}_1(\alpha, \beta, \gamma) \approx \tilde{B}_{11}(\alpha, \beta, \gamma, s)$$

Easily follows from $B_1(\alpha, \beta) \approx B_2(\alpha, \beta, s)$.

$$(x) \quad \tilde{B}_1(\alpha, \beta, \gamma) \approx \tilde{B}_{12}(\alpha, \beta, \gamma, s)$$

Similarly from $B_1(\alpha, \beta) \approx B_4(\alpha, \beta, s)$.

$$(xi) \quad \tilde{B}_1(\alpha, \beta, \gamma) \approx \tilde{B}_{16}(\alpha, \beta, \gamma, s)$$

By using the equivalences $B_1(\alpha, \beta) \approx B_{12}(\alpha, \beta, s)$ from Theorem A and in view of Theorem 2.1 (vi)

$$\sup_{x < y < b} H^\gamma(y) F^\alpha(y) \approx \inf_{h_1 \geq 0} \sup_{x < y < b} \left(h_1(y) + H(y) \right)^s \left(\int_y^b f(t) h_1^{\frac{\gamma-s}{\alpha}}(t) dt \right)^\alpha; \quad \gamma < s,$$

we have

$$\begin{aligned}
\tilde{B}_1(\alpha, \beta, \gamma) &= \sup_{a < x < b} \tilde{B}_1(x; \alpha, \beta, \gamma) \\
&= \sup_{a < x < b} F^\alpha(x) G^\beta(x) H^\gamma(x) \\
&= \sup_{a < x < b} G^\beta(x) \sup_{x < y < b} F^\alpha(y) H^\gamma(y) \\
&\approx \inf_{h_1 \geq 0} \sup_{a < x < b} G^\beta(x) \sup_{x < y < b} \left(h_1(y) + H(y) \right)^s \left(\int_y^b f(t) h_1^{\frac{\gamma-s}{\alpha}}(t) dt \right)^\alpha \\
&= \inf_{h_1 \geq 0} \sup_{a < x < b} \left(\int_x^b f(t) h_1^{\frac{\gamma-s}{\alpha}}(t) dt \right)^\alpha G^\beta(x) \left(h_1(x) + H(x) \right)^s \\
&= \inf_{h_1 \geq 0} \sup_{a < x < b} \tilde{B}_{16}(x; \alpha, \beta, \gamma, s; h_1)
\end{aligned}$$

$$=\tilde{B}_{16}(\alpha, \beta, \gamma, s).$$

$$(xii) \quad \tilde{B}_1(\alpha, \beta, \gamma) \approx \tilde{B}_{17}(\alpha, \beta, \beta, s)$$

By using the equivalences $B_1(\alpha, \beta) \approx B_{12}(\alpha, \beta, s)$ from Theorem A and in view of Theorem 2.1 (vi)

$$\sup_{x < y < b} G^\beta(y) F^\alpha(y) \approx \inf_{h_1 \geq 0} \sup_{x < y < b} \left(h_1(y) + G(y) \right)^s \left(\int_y^b f(t) h_1^{\frac{\beta-s}{\alpha}}(t) dt \right)^\alpha; \quad \beta < s,$$

we have

$$\begin{aligned} \tilde{B}_1(\alpha, \beta, \gamma) &= \sup_{a < x < b} \tilde{B}_1(x; \alpha, \beta, \gamma) \\ &= \sup_{a < x < b} F^\alpha(x) G^\beta(x) H^\gamma(x) \\ &= \sup_{a < x < b} H^\gamma(x) \sup_{x < y < b} F^\alpha(y) G^\beta(y) \\ &\approx \inf_{h_1 \geq 0} \sup_{a < x < b} H^\gamma(x) \sup_{x < y < b} \left(h_1(y) + G(y) \right)^s \left(\int_y^b f(t) h_1^{\frac{\beta-s}{\alpha}}(t) dt \right)^\alpha \\ &= \inf_{h_1 \geq 0} \sup_{a < x < b} \left(\int_x^b f(t) h_1^{\frac{\beta-s}{\alpha}}(t) dt \right)^\alpha H^\gamma(x) \left(h_1(x) + G(x) \right)^s \\ &= \inf_{h_1 \geq 0} \sup_{a < x < b} \tilde{B}_{17}(x; \alpha, \beta, \gamma, s; h_1) \\ &= \tilde{B}_{17}(\alpha, \beta, \gamma, s). \end{aligned}$$

$$(xiii) \quad \tilde{B}_1(\alpha, \beta, \gamma) \approx \tilde{B}_{18}(\alpha, \beta, \gamma, s_1, s_2)$$

By using the equivalence $B_1(\alpha, \beta) \approx B_{12}(\alpha, \beta, s)$ from Theorem A and in view of Theorem 2.1 (vi)

$$\sup_{x < y < b} F^\alpha(y) G^\beta(y) \approx \inf_{h_1 > 0} \sup_{x < y < b} \left(h_1(y) + G(y) \right)^s \left(\int_y^b f(z) h_1^{\frac{\beta-s}{\alpha}}(z) dz \right)^\alpha; \quad \beta < s$$

and also the fact that $H^\gamma(x) \geq 0$ as well as $\left(h_1(x) + G(x) \right)^{s_1} \geq 0$, we get

$$\begin{aligned} \tilde{B}_1(\alpha, \beta, \gamma) &= \sup_{a < x < b} \tilde{B}_1(x; \alpha, \beta, \gamma) \\ &= \sup_{a < x < b} F^\alpha(x) G^\beta(x) H^\gamma(x) \\ &= \sup_{a < x < b} H^\gamma(x) \sup_{x < y < b} F^\alpha(y) G^\beta(y) \end{aligned}$$

$$\begin{aligned}
&\approx \sup_{a < x < b} H^\gamma(x) \inf_{h_1 \geq 0} \sup_{x < y < b} \left(\int_y^b f(t) h_1^{\frac{\beta-s_1}{\alpha}}(t) dt \right)^\alpha \left(h_1(y) + G(y) \right)^{s_1} \\
&= \inf_{h_1 \geq 0} \sup_{a < x < b} H^\gamma(x) \sup_{x < y < b} \left(\int_y^b f(t) h_1^{\frac{\beta-s_1}{\alpha}}(t) dt \right)^\alpha \left(h_1(y) + G(y) \right)^{s_1} \\
&\leq \inf_{h_1 \geq 0} \sup_{a < x < b} \left\{ \sup_{a < y < x} \left(h_1(y) + G(y) \right)^{s_1} \right\} \left\{ \sup_{x < y < b} \left(\int_y^b f(t) h_1^{\frac{\beta-s_1}{\alpha}}(t) dt \right)^\alpha H^\gamma(y) \right\} \\
&\approx \inf_{h_1 \geq 0} \sup_{a < x < b} \left\{ \sup_{a < y < x} \left(h_1(y) + G(y) \right)^{s_1} \right\} \times \\
&\quad \times \left\{ \inf_{h_2 \geq 0} \sup_{x < y < b} \left(\int_y^b f(t) h_1^{\frac{\beta-s_1}{\alpha}}(t) h_2^{\frac{\gamma-s_2}{\alpha}}(t) dt \right)^\alpha \left(h_2(y) + H(y) \right)^{s_2} \right\} \\
&= \inf_{h_1 \geq 0} \inf_{h_2 \geq 0} \sup_{a < x < b} \left\{ \sup_{a < y < x} \left(h_1(y) + G(y) \right)^{s_1} \right\} \times \\
&\quad \times \left\{ \sup_{x < y < b} \left(\int_y^b f(t) h_1^{\frac{\beta-s_1}{\alpha}}(t) h_2^{\frac{\gamma-s_2}{\alpha}}(t) dt \right)^\alpha \left(h_2(y) + H(y) \right)^{s_2} \right\} \\
&= \inf_{h_1, h_2 \geq 0} \sup_{a < x < b} \left(\int_x^b f(t) h_1^{\frac{\beta-s_1}{\alpha}}(t) h_2^{\frac{\gamma-s_2}{\alpha}}(t) dt \right)^\alpha \times \\
&\quad \times \left\{ \sup_{a < y < x} \left(h_1(y) + G(y) \right)^{s_1} \right\} \left(h_2(x) + H(x) \right)^{s_2} \\
&= \inf_{h_1, h_2 \geq 0} \sup_{a < x < b} \tilde{B}_{18}(x; \alpha, \beta, \gamma, s_1, s_2; h_1, h_2) \\
&= \tilde{B}_{18}(\alpha, \beta, \gamma, s_1, s_2).
\end{aligned}$$

Thus

$$\tilde{B}_1(\alpha, \beta, \gamma) \lesssim \tilde{B}_{18}(\alpha, \beta, \gamma, s_1, s_2). \quad (3.2)$$

Since for $h_1(x) = G(x)$ and $h_2(x) = H(x)$, we have

$$\begin{aligned}
\tilde{B}_{18}(x; \alpha, \beta, \gamma, s_1, s_2; G, H) &= 2^{s_1+s_2} \left(\int_x^b f(t) G^{\frac{\beta-s_1}{\alpha}}(t) H^{\frac{\gamma-s_2}{\alpha}}(t) dt \right)^\alpha G^{s_1}(x) H^{s_2}(x) \\
&= 2^{s_1+s_2} \tilde{B}_2(x; \alpha, \beta, \gamma, s_1, s_2) \\
&\leq 2^{s_1+s_2} \sup_{a < x < b} \tilde{B}_2(x; \alpha, \beta, \gamma, s_1, s_2) \\
&= 2^{s_1+s_2} \tilde{B}_2(\alpha, \beta, \gamma, s_1, s_2) \\
&\approx 2^{s_1+s_2} \tilde{B}_1(\alpha, \beta, \gamma).
\end{aligned}$$

The last relation holds by using $\tilde{B}_1(\alpha, \beta, \gamma) \approx \tilde{B}_2(\alpha, \beta, \gamma, s_1, s_2)$. Clearly we get that

$$\tilde{B}_{18}(\alpha, \beta, \gamma, s_1, s_2) \lesssim \tilde{B}_1(\alpha, \beta, \gamma). \quad (3.3)$$

In view of (3.2) and (3.3), we obtain

$$\tilde{B}_1(\alpha, \beta, \gamma) \approx \tilde{B}_{18}(\alpha, \beta, \gamma, s_1, s_2).$$

$$(xiv) \quad \tilde{B}_1(\alpha, \beta, \gamma) \approx \tilde{B}_{19}(\alpha, \beta, \gamma, s_1, s_2)$$

Using the similar reasoning as in (xiii).

$$(xv) \quad \tilde{B}_1(\alpha, \beta, \gamma) \approx \tilde{B}_{20}(\alpha, \beta, \gamma, s_1, s_2)$$

Using the similar reasoning as in (xiii).

$$(xvi) \quad \tilde{B}_1(\alpha, \beta, \gamma) \approx \tilde{B}_{21}(\alpha, \beta, \gamma, s_1, s_2)$$

By using the equivalence $B_1(\alpha, \beta) \approx B_{14}(\alpha, \beta, s)$ from Theorem A and in view of Theorem 2.1 (vii)

$$\sup_{x < y < b} F^\alpha(y) G^\beta(y) \approx \inf_{h_1 > 0} \sup_{x < y < b} h_1^{-s_1}(y) \left(\int_x^y f(z) (h_1(z) + G(z))^{\frac{\beta+s_1}{\alpha}} dz \right)^\alpha$$

and also the fact that $H^\gamma(x) \geq 0$ as well as $h_1^{-s_1}(x) \geq 0$, we get

$$\begin{aligned} \tilde{B}_1(\alpha, \beta, \gamma) &= \sup_{a < x < b} \tilde{B}_1(x; \alpha, \beta, \gamma) \\ &= \sup_{a < x < b} F^\alpha(x) G^\beta(x) H^\gamma(x) \\ &= \sup_{a < x < b} H^\gamma(x) \sup_{x < y < b} F^\alpha(y) G^\beta(y) \\ &\approx \sup_{a < x < b} H^\gamma(x) \inf_{h_1 \geq 0} \sup_{x < y < b} \left(\int_x^y f(t) (h_1(t) + G(t))^{\frac{\beta+s_1}{\alpha}} dt \right)^\alpha h_1^{-s_1}(y) \\ &= \inf_{h_1 \geq 0} \sup_{a < x < b} H^\gamma(x) \sup_{x < y < b} h_1^{-s_1}(y) \left(\int_x^y f(t) (h_1(t) + G(t))^{\frac{\beta+s_1}{\alpha}} dt \right)^\alpha \\ &\leq \inf_{h_1 \geq 0} \sup_{a < y < b} \left\{ \sup_{y < x < b} h_1^{-s_1}(x) \right\} \left\{ \sup_{a < x < y} \left(\int_x^y f(t) (h_1(t) + G(t))^{\frac{\beta+s_1}{\alpha}} dt \right)^\alpha H^\gamma(x) \right\} \\ &\approx \inf_{h_1 \geq 0} \sup_{a < y < b} \left\{ \sup_{y < x < b} h_1^{-s_1}(x) \right\} \times \\ &\quad \times \left\{ \inf_{h_2 \geq 0} \sup_{a < x < y} h_2^{-s_2}(x) \left(\int_a^x f(t) (h_1(t) + G(t))^{\frac{\beta+s_1}{\alpha}} (h_2(t) + H(t))^{\frac{\gamma+s_2}{\alpha}} dt \right)^\alpha \right\} \\ &= \inf_{h_1 \geq 0} \inf_{h_2 \geq 0} \sup_{a < y < b} \left\{ \sup_{y < x < b} h_1^{-s_1}(x) \right\} \times \end{aligned}$$

$$\begin{aligned}
& \times \left\{ \sup_{a < x < y} h_2^{-s_2}(x) \left(\int_a^x f(t) (h_1(t) + G(t))^{\frac{\beta+s_1}{\alpha}} (h_2(t) + H(t))^{\frac{\gamma+s_2}{\alpha}} dt \right)^\alpha \right\} \\
& = \inf_{h_1, h_2 \geq 0} \sup_{a < x < b} \left(\int_a^x f(t) (h_1(t) + G(t))^{\frac{\beta+s_1}{\alpha}} (h_2(t) + H(t))^{\frac{\gamma+s_2}{\alpha}} dt \right)^\alpha \times \\
& \quad \times \left\{ \sup_{x < y < b} h_1^{-s_1}(y) \right\} h_2^{-s_2}(x) \\
& = \inf_{h_1, h_2 \geq 0} \sup_{a < x < b} \tilde{B}_{21}(x; \alpha, \beta, \gamma, s_1, s_2; h_1, h_2) \\
& = \tilde{B}_{21}(\alpha, \beta, \gamma, s_1, s_2).
\end{aligned}$$

Thus

$$\tilde{B}_1(\alpha, \beta, \gamma) \lesssim \tilde{B}_{21}(\alpha, \beta, \gamma, s_1, s_2). \quad (3.4)$$

Since for $h_1(x) = G(x)$ and $h_2(x) = H(x)$, we have

$$\begin{aligned}
\tilde{B}_{21}(x; \alpha, \beta, \gamma, s_1, s_2; G, H) & = 2^{\beta+\gamma+s_1+s_2} \left(\int_a^x f(t) G^{\frac{\beta+s_1}{\alpha}}(t) H^{\frac{\gamma+s_2}{\alpha}}(t) dt \right)^\alpha G^{-s_1}(x) H^{-s_2}(x) \\
& = 2^{\beta+\gamma+s_1+s_2} \tilde{B}_3(x; \alpha, \beta, \gamma, s_1, s_2) \\
& \leq 2^{\beta+\gamma+s_1+s_2} \sup_{a < x < b} \tilde{B}_3(x; \alpha, \beta, \gamma, s_1, s_2) \\
& = 2^{\beta+\gamma+s_1+s_2} \tilde{B}_3(\alpha, \beta, \gamma, s_1, s_2) \\
& \approx 2^{\beta+\gamma+s_1+s_2} \tilde{B}_1(\alpha, \beta, \gamma).
\end{aligned}$$

The last relation holds by using $\tilde{B}_1(\alpha, \beta, \gamma) \approx \tilde{B}_3(\alpha, \beta, \gamma, s_1, s_2)$. Clearly we get that

$$\tilde{B}_{21}(\alpha, \beta, \gamma, s_1, s_2) \lesssim \tilde{B}_1(\alpha, \beta, \gamma). \quad (3.5)$$

Therefore, in view of (3.4) and (3.5), we find

$$\tilde{B}_1(\alpha, \beta, \gamma) \approx \tilde{B}_{21}(\alpha, \beta, \gamma, s_1, s_2).$$

$$(xvii) \quad \tilde{B}_1(\alpha, \beta, \gamma) \approx \tilde{B}_{22}(\alpha, \beta, \gamma, s_1, s_2)$$

Using the similar reasoning as in (xvi).

$$(xviii) \quad \tilde{B}_1(\alpha, \beta, \gamma) \approx \tilde{B}_{23}(\alpha, \beta, \gamma, s_1, s_2)$$

Using the similar reasoning as in (xvi). \square

We apply Theorem 3.1 and provide equivalent conditions for the bilinear inequality (1.5) to hold.

Theorem 3.2. *Let $1 < \max(p_1, p_2) \leq q < \infty$ with s, s_1, s_2 be positive numbers. Define U as in (1.1) and*

$$V_i(x) = \int_a^x v_i^{1-p'_i}(t) dt, \quad i = 1, 2.$$

Moreover define

$$\begin{aligned} \tilde{A}_1(s_1, s_2) &:= \sup_{a < x < b} \left(\int_x^b u(t) V_1^{\left(\frac{\frac{1}{p_1} - s_1}{\frac{1}{q}}\right)}(t) V_2^{\left(\frac{\frac{1}{p_2} - s_2}{\frac{1}{q}}\right)}(t) dt \right)^{\frac{1}{q}} V_1^{s_1}(x) V_2^{s_2}(x); \\ \tilde{A}_2(s_1, s_2) &:= \sup_{a < x < b} \left(\int_a^x u(t) V_1^{\left(\frac{\frac{1}{p_1} + s_1}{\frac{1}{q}}\right)}(t) V_2^{\left(\frac{\frac{1}{p_2} + s_2}{\frac{1}{q}}\right)}(t) dt \right)^{\frac{1}{q}} V_1^{-s_1}(x) V_2^{-s_2}(x); \\ \tilde{A}_3(s_1, s_2) &:= \sup_{a < x < b} \left(\int_a^x u(t) V_1^{\left(\frac{\frac{1}{p_1}}{\frac{1}{q} + s_1}\right)}(t) V_2^{\left(\frac{\frac{1}{p_2} - s_2}{\frac{1}{q} + s_1}\right)}(t) dt \right)^{\frac{1}{q} + s_1} U^{-s_1}(x) V_2^{s_2}(x); \\ \tilde{A}_4(s_1, s_2) &:= \sup_{a < x < b} \left(\int_a^x u(t) V_1^{\left(\frac{\frac{1}{p_1} - s_2}{\frac{1}{q} + s_1}\right)}(t) V_2^{\left(\frac{\frac{1}{p_2}}{\frac{1}{q} + s_1}\right)}(t) dt \right)^{\frac{1}{q} + s_1} U^{-s_1}(x) V_1^{s_2}(x); \\ \tilde{A}_5(s_1, s_2) &:= \sup_{a < x < b} \left(\int_a^x u(t) V_1^{\left(\frac{\frac{1}{p_1}}{\frac{1}{q} - s_1}\right)}(t) V_2^{\left(\frac{\frac{1}{p_2} + s_2}{\frac{1}{q} - s_1}\right)}(t) dt \right)^{\frac{1}{q} - s_1} U^{s_1}(x) V_2^{-s_2}(x), \quad \frac{1}{q} > s_1; \\ \tilde{A}_6(s_1, s_2) &:= \sup_{a < x < b} \left(\int_a^x u(t) V_1^{\left(\frac{\frac{1}{p_1} + s_2}{\frac{1}{q} - s_1}\right)}(t) V_2^{\left(\frac{\frac{1}{p_2}}{\frac{1}{q} - s_1}\right)}(t) dt \right)^{\frac{1}{q} - s_1} U^{s_1}(x) V_1^{-s_2}(x), \quad \frac{1}{q} > s_1; \\ \tilde{A}_7(s_1, s_2) &:= \sup_{a < x < b} \left(\int_x^b u(t) V_1^{\left(\frac{\frac{1}{p_1}}{\frac{1}{q} - s_1}\right)}(t) dt \right)^{\frac{1}{q} - s_1} V_2^{s_2}(x) \left(\int_x^b u(t) V_2^{\left(\frac{\frac{1}{p_2} - s_2}{s_1}\right)}(t) dt \right)^{s_1}, \quad \frac{1}{q} < s_1; \end{aligned}$$

$$\tilde{A}_8(s_1, s_2) := \sup_{a < x < b} \left(\int_x^b u(t) V_2^{\left(\frac{\frac{1}{p_2}}{\frac{1}{q} - s_1}\right)}(t) dt \right)^{\frac{1}{q} - s_1} V_1^{s_2}(x) \left(\int_x^b u(t) V_1^{\left(\frac{\frac{1}{p_1} - s_2}{s_1}\right)}(t) dt \right)^{s_1}, \quad \frac{1}{q} < s_1;$$

$$\tilde{A}_9(s_1, s_2) := \sup_{a < x < b} \left(\int_a^x v_1^{1-p_1'}(t) U^{\left(\frac{\frac{1}{2q}}{p_1' + s_1}\right)}(t) dt \right)^{\frac{1}{p_1'} + s_1} \left(\int_a^x v_2^{1-p_2'}(t) U^{\left(\frac{\frac{1}{2q}}{p_2' + s_2}\right)}(t) dt \right)^{\frac{1}{p_2'} + s_2} \times \\ \times V_1^{-s_1}(x) V_2^{-s_2}(x);$$

$$\tilde{A}_{10}(s) := \sup_{a < x < b} \left(\int_x^b u(t) V_1^{\left(\frac{\frac{1}{p_1}(1-s)}{\frac{1}{q}}\right)}(t) V_2^{\left(\frac{\frac{1}{p_2}(1-s)}{\frac{1}{q}}\right)}(t) dt \right)^{\frac{1}{q}} V_1^{\frac{s}{p_1'}}(x) V_2^{\frac{s}{p_2'}}(x);$$

$$\tilde{A}_{11}(s) := \sup_{a < x < b} \left(\int_a^x u(t) V_1^{\left(\frac{\frac{1}{p_1}(1+s)}{\frac{1}{q}}\right)}(t) V_2^{\left(\frac{\frac{1}{p_2}(1+s)}{\frac{1}{q}}\right)}(t) dt \right)^{\frac{1}{q}} V_1^{-\frac{s}{p_1'}}(x) V_2^{-\frac{s}{p_2'}}(x);$$

$$\tilde{A}_{12}(s) := \sup_{a < x < b} \left(\int_x^b u(t) V_1^{\left(\frac{\frac{1}{p_1}}{\frac{1}{q} + s}\right)}(t) V_2^{\left(\frac{\frac{1}{p_2}}{\frac{1}{q} + s}\right)}(t) dt \right)^{\frac{1}{q} + s} U^{-s}(x);$$

$$\tilde{A}_{13}(s) := \sup_{a < x < b} \left(\int_a^x u(t) V_1^{\left(\frac{\frac{1}{p_1}}{\frac{1}{q} - s}\right)}(t) V_2^{\left(\frac{\frac{1}{p_2}}{\frac{1}{q} - s}\right)}(t) dt \right)^{\frac{1}{q} - s} U^s(x), \quad \frac{1}{q} > s;$$

$$\tilde{A}_{14}(s) := \sup_{a < x < b} \left(\int_x^b u(t) V_1^{\left(\frac{\frac{1}{p_1}}{\frac{1}{q} - s}\right)}(t) V_2^{\left(\frac{\frac{1}{p_2}}{\frac{1}{q} - s}\right)}(t) dt \right)^{\frac{1}{q} - s} U^s(x), \quad \frac{1}{q} < s;$$

$$\tilde{A}_{15}(s) := \inf_{h_1 \geq 0} \sup_{a < x < b} \left(\int_x^b u(t) h_1^{\left(\frac{\frac{1}{p_2} - s}{\frac{1}{q}}\right)}(t) dt \right)^{\frac{1}{q}} V_1^{\frac{1}{p_1'}}(x) \left(h_1(x) + V_2(x) \right)^s, \quad \frac{1}{p_2'} < s;$$

$$\tilde{A}_{16}(s) := \inf_{h_1 \geq 0} \sup_{a < x < b} \left(\int_x^b u(t) h_1^{\left(\frac{\frac{1}{p'_1} - s}{\frac{1}{q}}\right)}(t) dt \right)^{\frac{1}{q}} V_2^{\frac{1}{p'_2}}(x) \left(h_1(x) + V_1(x) \right)^s, \quad \frac{1}{p'_1} < s;$$

$$\begin{aligned} \tilde{A}_{17}(s_1, s_2) := & \inf_{h_1, h_2 \geq 0} \sup_{a < x < b} \left(\int_x^b u(t) h_1^{\left(\frac{\frac{1}{p'_1} - s_1}{\frac{1}{q}}\right)}(t) h_2^{\left(\frac{\frac{1}{p'_2} - s_2}{\frac{1}{q}}\right)}(t) dt \right)^{\frac{1}{q}} \times \\ & \times \left\{ \sup_{a < y < x} \left(h_1(y) + V_1(y) \right)^{s_1} \right\} \left(h_2(x) + V_2(x) \right)^{s_2}, \quad \frac{1}{p'_1} < s_1, \frac{1}{p'_2} < s_2; \end{aligned}$$

$$\begin{aligned} \tilde{A}_{18}(s_1, s_2) := & \inf_{h_1, h_2 \geq 0} \sup_{a < x < b} \left(\int_x^b u(t) h_1^{\left(\frac{\frac{1}{p'_1} - s_1}{\frac{1}{q}}\right)}(t) h_2^{\left(\frac{\frac{1}{p'_2} - s_2}{\frac{1}{q}}\right)}(t) dt \right)^{\frac{1}{q}} \times \\ & \times \left(h_1(x) + V_1(x) \right)^{s_1} \left\{ \sup_{a < y < x} \left(h_2(y) + V_2(y) \right)^{s_2} \right\}, \quad \frac{1}{p'_1} < s_1, \frac{1}{p'_2} < s_2; \end{aligned}$$

$$\begin{aligned} \tilde{A}_{19}(s_1, s_2) := & \inf_{h_1, h_2 \geq 0} \sup_{a < x < b} \left(\int_x^b u(t) h_1^{\left(\frac{\frac{1}{p'_1} - s_1}{\frac{1}{q}}\right)}(t) h_2^{\left(\frac{\frac{1}{p'_2} - s_2}{\frac{1}{q}}\right)}(t) dt \right)^{\frac{1}{q}} \times \\ & \times \left\{ \sup_{a < y < x} \left(h_1(y) + V_1(y) \right)^{s_1} \right\} \left\{ \sup_{a < y < x} \left(h_2(y) + V_2(y) \right)^{s_2} \right\}, \quad \frac{1}{p'_1} < s_1, \frac{1}{p'_2} < s_2; \end{aligned}$$

$$\begin{aligned} \tilde{A}_{20}(s_1, s_2) := & \inf_{h_1, h_2 \geq 0} \sup_{a < x < b} \left(\int_a^x u(t) \left(h_1(t) + V_1(t) \right)^{\left(\frac{\frac{1}{p'_1} + s_1}{\frac{1}{q}}\right)} \left(h_2(t) + V_2(t) \right)^{\left(\frac{\frac{1}{p'_2} + s_2}{\frac{1}{q}}\right)} dt \right)^{\frac{1}{q}} \times \\ & \times \left\{ \sup_{x < y < b} h_1^{-s_1}(y) \right\} h_2^{-s_2}(x); \end{aligned}$$

$$\begin{aligned} \tilde{A}_{21}(s_1, s_2) := & \inf_{h_1, h_2 \geq 0} \sup_{a < x < b} \left(\int_a^x u(t) \left(h_1(t) + V_1(t) \right)^{\left(\frac{\frac{1}{p'_1} + s_1}{\frac{1}{q}}\right)} \left(h_2(t) + V_2(t) \right)^{\left(\frac{\frac{1}{p'_2} + s_2}{\frac{1}{q}}\right)} dt \right)^{\frac{1}{q}} \times \\ & \times h_1^{-s_1}(x) \left\{ \sup_{x < y < b} h_2^{-s_2}(y) \right\}; \end{aligned}$$

$$\begin{aligned} \tilde{A}_{22}(s_1, s_2) := & \inf_{h_1, h_2 \geq 0} \sup_{a < x < b} \left(\int_a^x u(t) \left(h_1(t) + V_1(t) \right)^{\left(\frac{\frac{1}{p_1} + s_1}{q} \right)} \left(h_2(t) + V_2(t) \right)^{\left(\frac{\frac{1}{p_2} + s_2}{q} \right)} dt \right)^{\frac{1}{q}} \times \\ & \times \left\{ \sup_{x < y < b} h_1^{-s_1}(y) \right\} \left\{ \sup_{x < y < b} h_2^{-s_2}(y) \right\}. \end{aligned}$$

Then the inequality (1.5) holds for all $f, g \in \mathfrak{M}^+$ if and only if any of the $\tilde{A}_i(s_1, s_2)$, $\tilde{A}_j(s)$, ($i = 1, 2, \dots, 9, 17, 18, \dots, 22$), ($j = 10, 11, \dots, 16$) is finite. Also the best constant C in (1.5) satisfies $C \approx \tilde{A}_i(s_1, s_2) \approx \tilde{A}_j(s)$, ($i = 1, 2, \dots, 9, 17, 18, \dots, 22$), ($j = 10, 11, \dots, 16$).

Proof. By putting $f(x) = u(x)$, $g(x) = v_1^{1-p_1'}(x)$ in (1.4) and $h(x) = v_2^{1-p_2'}(x)$ in (3.1), we get

$$F(x) = U(x), G(x) = V_1(x), H(x) = V_2(x).$$

Consequently for $\alpha = \frac{1}{q}$, $\beta = \frac{1}{p_1'}$ and $\gamma = \frac{1}{p_2'}$, we have

$$\begin{aligned} \tilde{A}_i(s_1, s_2) &= \tilde{B}_{i+1} \left(\frac{1}{q}, \frac{1}{p_1'}, \frac{1}{p_2'}, s_1, s_2 \right); \quad (i = 1, 2, \dots, 9, 17, 18, \dots, 22), \\ \tilde{A}_j(s) &= \tilde{B}_{j+1} \left(\frac{1}{q}, \frac{1}{p_1'}, \frac{1}{p_2'}, s \right); \quad (j = 10, 11, \dots, 16). \end{aligned}$$

Now by Theorem 3.1, these are all equivalent to $\tilde{B}_1 \left(\frac{1}{q}, \frac{1}{p_1'}, \frac{1}{p_2'} \right) \stackrel{(1.6)}{=} \mathcal{D}$ and \mathcal{D} is necessary and sufficient condition for the inequality (1.5) to hold.

Since the constant C in (1.5) verifies $C \approx \mathcal{D} = \tilde{B}_1 \left(\frac{1}{q}, \frac{1}{p_1'}, \frac{1}{p_2'} \right)$, it follows that $C \approx \tilde{A}_i(s_1, s_2) \approx \tilde{A}_j(s)$, ($i = 1, 2, \dots, 9, 17, 18, \dots, 22$), ($j = 10, 11, \dots, 16$). \square

4 Bilinear Type Geometric Mean Operator

Consider the Hardy averaging operator A :

$$(Af)(x) := \frac{1}{x} \int_0^x f(t) dt$$

and the geometric mean operator T which is defined by

$$(Tf)(x) := \exp \left(\frac{1}{x} \int_0^x \ln(f(t)) dt \right), \quad f \in \mathfrak{M}^+.$$

Note that

$$\lim_{\alpha \rightarrow 0} \left\{ (Af^\alpha)(x) \right\}^{\frac{1}{\alpha}} = Tf(x). \quad (4.1)$$

Therefore, it seems reasonable to study the inequality

$$\left(\int_0^\infty (Tf)^q(x)u(x) dx \right)^{\frac{1}{q}} \leq C \left(\int_0^\infty f^p(x)v(x) dx \right)^{\frac{1}{p}} \quad (4.2)$$

via (1.2) with suitable modifications. However, it was observed in [5], [6] that the Muckenhoupt condition (1.5) is not suitable for these modifications. To this end, Persson and Stepanov [11] obtained an alternate condition for the inequality (1.2) and by suitable modifications studied the inequality (4.2). Precisely, they proved the following

Theorem C. *Let $0 < p \leq q < \infty$. Then the inequality (4.2) with the best constant C holds for $f \in \mathfrak{M}^+$ if and only if following holds*

$$\mathcal{B} := \sup_{t>0} t^{-\frac{1}{p}} \left(\int_0^t \left[T \left(\frac{1}{v(x)} \right) \right]^{\frac{q}{p}} u(x) dx \right)^{\frac{1}{q}} < \infty. \quad (4.3)$$

This was perhaps one of the motivations to obtain several equivalent conditions for the inequality (1.2). Let us mention that the inequality (4.2) has also independently been studied in [4], [10], [13].

Our aim, in this section, is to study the following bilinear type inequality involving the geometric mean operator T :

$$\begin{aligned} \left(\int_0^\infty (Tf)^q(x)(Tg)^q(x)u(x) dx \right)^{\frac{1}{q}} &\leq C \left(\int_0^\infty f^{p_1}(x)v_1(x) dx \right)^{\frac{1}{p_1}} \\ &\times \left(\int_0^\infty f^{p_2}(x)v_2(x) dx \right)^{\frac{1}{p_2}}. \end{aligned} \quad (4.4)$$

We prove the following:

Theorem 4.1. *Let $1 < \max(p_1, p_2) < q < \infty$. Then the inequality (4.4) holds for all $f, g \in \mathfrak{M}^+$ if and only if*

$$\tilde{\mathcal{B}} := \sup_{t>0} t^{-\left(\frac{1}{p_1} + \frac{1}{p_2}\right)} \left(\int_0^t \left[T \left(\frac{1}{v_1(x)} \right) \right]^{\frac{q}{p_1}} \left[T \left(\frac{1}{v_2(x)} \right) \right]^{\frac{q}{p_2}} u(x) dx \right)^{\frac{1}{q}} < \infty.$$

Proof. By using (4.3) from Theorem C, we get

$$\begin{aligned} C &= \sup_{g \in \mathfrak{M}^+} \sup_{f \in \mathfrak{M}^+} \frac{\left(\int_0^\infty (Gf)^q(x)(Gg)^q(x)u(x) dx \right)^{\frac{1}{q}}}{\|f\|_{p_1, v_1} \|g\|_{p_2, v_2}} \\ &\approx \sup_{g \in \mathfrak{M}^+} \|g\|_{p_2, v_2}^{-1} \sup_{t_1 > 0} t_1^{-\frac{1}{p_1}} \left(\int_0^{t_1} \left[G \left(\frac{1}{v_1(x)} \right) \right]^{\frac{q}{p_1}} (Gg)^q(x)u(x) dx \right)^{\frac{1}{q}} \end{aligned}$$

$$\begin{aligned}
&= \sup_{t_1 > 0} t_1^{-\frac{1}{p_1}} \sup_{g \in \mathfrak{M}^+} \|g\|_{p_2, v_2}^{-1} \left(\int_0^{t_1} \left[G \left(\frac{1}{v_1(x)} \right) \right]^{\frac{q}{p_1}} (Gg)^q(x) u(x) dx \right)^{\frac{1}{q}} \\
&\approx \sup_{t_1 > 0} t_1^{-\frac{1}{p_1}} \sup_{0 < t_2 < t_1} t_2^{-\frac{1}{p_2}} \left(\int_0^{t_2} \left[G \left(\frac{1}{v_1(x)} \right) \right]^{\frac{q}{p_1}} \left[G \left(\frac{1}{v_2(x)} \right) \right]^{\frac{q}{p_2}} u(x) dx \right)^{\frac{1}{q}} \\
&= \sup_{t > 0} t^{-\left(\frac{1}{p_1} + \frac{1}{p_2}\right)} \left(\int_0^t \left[G \left(\frac{1}{v_1(x)} \right) \right]^{\frac{q}{p_1}} \left[G \left(\frac{1}{v_2(x)} \right) \right]^{\frac{q}{p_2}} u(x) dx \right)^{\frac{1}{q}} \\
&= \tilde{\mathcal{B}}.
\end{aligned}$$

□

5 Acknowledgement

The work of first author, has been partially supported by Shota Rustaveli National Science Foundation of Georgia (SRNSFG) [grant number FR17-589], by Grant 18-00580S of the Czech Science Foundation and RVO:67985840.

The second author acknowledges the MATRICS Research Grant No. MTR/2017/000126 of SERB, Department of Science and Technology, India.

References

- [1] M.I. Aguilar Cañestro, P.Ortega Salvador and C.Ramírez Torreblanca, *Weighted bilinear Hardy inequalities*, J. Math. Anal. Appl., 387 (2012), 320–334.
- [2] A. Gogatishvili, A. Kufner, L.-E. Persson, and A. Wedestig, *An equivalence theorem for some scales of integral conditions related to Hardy’s inequality with applications*, Real Anal. Exchange, 29 (2004), 867–880.
- [3] A. Gogatishvili, A. Kufner and L.E. Persson, *Some new scales of characterization of Hardy’s inequality*, Proc. Est. Acad. Sci., 59 (2010), 7–18.
- [4] H.P. Heinig, R. Kerman and M. Krbec, *Weighted exponential inequalities*, Georgian Math. J., 8 (2001), 69–86.
- [5] P. Jain, L.-E. Persson and A. Wedestig, *From Hardy to Carleman and general mean-type inequalities*, Function Spaces and Applications, CRC Press (New York)/Narosa Publishing House (New Delhi)/Alpha Science (Pangbourne) (2000), 117–130 .
- [6] P. Jain, L.-E. Persson and A. Wedestig, *Carleman-Knopp type inequalities via Hardy inequalities*, Math. Inequal. Appl., 4 (2001), 343–355.

- [7] S. Kanjilal, L.-E. Persson and G.E. Shambilova, *Equivalent integral conditions related to bilinear Hardy-type inequalities*, Math. Inequal. Appl., 22 (2019), 1535–1548.
- [8] M. Křepela, *Iterating bilinear Hardy inequalities*, Proc. Edinburgh Math. Soc., 60 (2017), 955–971.
- [9] B. Muckenhoupt, *Hardy's inequality with weights*, Studia Math., 44 (1972), 31–38.
- [10] B. Opic and P. Gurka, *Weighted inequalities for geometric means*, Proc. Amer. Math. Soc., 3 (1994), 771–779.
- [11] L.-E. Persson, V.D. Stepanov, *Weighted integral inequalities with the geometric mean operator*, J. Inequal. Appl., 7 (2002), 727–746.
- [12] L.-E. Persson, V.D. Stepanov, P. Wall, *Some scale of equivalent weight characterizations of Hardy's inequality: the case $q < p$* , Math. Inequal. Appl., 10 (2007), 267–279.
- [13] L. Pick and B. Opic, *On the geometric mean operator*, J. Math. Anal. Appl., 183 (1994), 652–662.

Amiran Gogatishvili

Institute of Mathematics
of the Czech Academy of Sciences
Žitná 25, 115 67 Prague 1
Czech Republic
Email: gogatish@math.cas.cz

Pankaj Jain

Department of Mathematics
South Asian University
Chanakya Puri, New Delhi-110021
India
Email: pankaj.jain@sau.ac.in & pankajkrjain@hotmail.com

Saikat Kanjilal

Department of Mathematics
South Asian University
Chanakya Puri, New Delhi-110021
India
Email: saikatkanjilal@students.sau.ac.in & saikat.kanjilal.07@gmail.com