

Kim-type APN functions are affine equivalent to Gold functions

Benjamin Chase

Petr Lisoněk*

Department of Mathematics

Simon Fraser University

Burnaby, BC

Canada V5A 1S6

plisonek@sfu.ca

Abstract

The problem of finding APN permutations of \mathbb{F}_{2^n} where n is even and $n > 6$ has been called the Big APN Problem. Li, Li, Helleseht and Qu recently characterized APN functions defined on \mathbb{F}_{q^2} of the form $f(x) = x^{3q} + a_1x^{2q+1} + a_2x^{q+2} + a_3x^3$, where $q = 2^m$ and $m \geq 4$. We will call functions of this form Kim-type functions because they generalize the form of the Kim function that was used to construct an APN permutation of \mathbb{F}_{2^6} . We extend the result of Li, Li, Helleseht and Qu by proving that if a Kim-type function f is APN and $m \geq 4$, then f is affine equivalent to one of two Gold functions $G_1(x) = x^3$ or $G_2(x) = x^{2^{m-1}+1}$. Combined with the recent result of Gölöğlu and Langevin who proved that, for even n , Gold APN functions are never CCZ equivalent to permutations, it follows that for $m \geq 4$ Kim-type APN functions on $\mathbb{F}_{2^{2m}}$ are never CCZ equivalent to permutations.

1 Background

Let \mathbb{F}_{2^k} denote the finite field with 2^k elements and let $\mathbb{F}_{2^k}^*$ be the set of its nonzero elements. Let Tr_k denote the absolute trace from \mathbb{F}_{2^k} to \mathbb{F}_2 .

*Research of both authors was supported in part by the Natural Sciences and Engineering Research Council of Canada (NSERC).

We say that a function $f : \mathbb{F}_{2^n} \rightarrow \mathbb{F}_{2^n}$ is *almost perfect nonlinear (APN)* if for any $a \in \mathbb{F}_{2^n}^*$ and any $b \in \mathbb{F}_{2^n}$ the equation

$$f(x+a) + f(x) = b$$

has at most two solutions $x \in \mathbb{F}_{2^n}$. We say that functions $f : \mathbb{F}_{2^n} \rightarrow \mathbb{F}_{2^n}$ and $g : \mathbb{F}_{2^n} \rightarrow \mathbb{F}_{2^n}$ are *extended affine (EA) equivalent* if there exist affine permutations L_1, L_2 of \mathbb{F}_{2^n} and an affine mapping $L_3 : \mathbb{F}_{2^n} \rightarrow \mathbb{F}_{2^n}$ such that $g(x) = L_1(f(L_2(x))) + L_3(x)$. If $L_3(x) = 0$ we will say that f and g are *affine equivalent*; this is the kind of equivalence that we will be using in the present paper. EA equivalence is a special case of *Carlet-Charpin-Zimoviev (CCZ) equivalence* [5]. CCZ equivalence preserves the property whether the function is APN or not.

Many infinite families of APN functions are known. A recent account of the known families is given in [3]. In the present paper we will encounter the *Gold* APN functions $G(x) = x^{2^k+1}$ where $\gcd(k, n) = 1$.

APN functions are of great interest in cryptography as components of S-boxes in block ciphers, where they provide optimal protection against differential cryptanalysis. They also have connections to combinatorial objects such as finite geometries and combinatorial designs [10].

In some designs of block ciphers, such as substitution-permutation networks, it is desirable that the APN function is also a *permutation* of \mathbb{F}_{2^n} . While many APN permutations are known for odd n , the situation is very different when n is even. It is known that no APN permutations of \mathbb{F}_{2^n} exist when $n = 2, 4$. An APN permutation of \mathbb{F}_{2^6} was announced at the Fq9 conference in 2009 [2]. Its construction is based on the APN function

$$\kappa(x) = x^3 + x^{10} + ux^{24}$$

where u is a suitable primitive element of \mathbb{F}_{2^6} . The function κ is known as *Kim function*. APN permutations of \mathbb{F}_{2^6} are then constructed by applying a suitable CCZ equivalence transformation to the Kim function [2]; all examples constructed in [2] are mutually CCZ equivalent.

The question of existence of APN permutations of \mathbb{F}_{2^n} for even $n > 6$ was posed as the *Big APN Problem* in [2], and it has attracted lot of interest since then.

By generalizing the form of the Kim function, Carlet in [4] posed the following open problem:

Problem 1.1. [4, Section 3.7] Find more APN functions or, better, infinite classes of APN functions of the form $X^3 + aX^{2+q} + bX^{2q+1} + cX^{3q}$ where

$q = 2^{n/2}$ with n even, or more generally of the form $X^{2^k+1} + aX^{2^k+q} + bX^{2^kq+1} + cX^{2^kq+q}$, where $\gcd(k, n) = 1$.

The motivation for solving this problem is given by possibly applying a suitable CCZ transformation to the resulting APN functions to obtain APN permutations, in analogy to the approach of [2]. There have been several lines of attack to approach Problem 1.1 and similar problems; we refer to the Introduction section of [9] for a detailed account of them.

Most recently, Li, Li, Hellesest and Qu [9] solved the first part of Problem 1.1 by proving Theorem 1.4 stated below, in which they completely characterize APN functions on \mathbb{F}_{2^n} of the form $f(x) = x^{3q} + a_1x^{2q+1} + a_2x^{q+2} + a_3x^3$, where $q = 2^m$ and $m = n/2$, $m \geq 4$. We will call functions of this form *Kim-type functions* in the present paper. We extend the result of [9] by proving that if a Kim-type function f is APN and $m \geq 4$, then f is affine equivalent to one of two Gold functions $G_1(x) = x^3$ or $G_2(x) = x^{2^{m-1}+1}$. Combined with the recent result of Göloğlu and Langevin [7] which states that, for even n , Gold APN functions are never CCZ equivalent to permutations, it follows that for $m \geq 4$ Kim-type functions are never CCZ equivalent to permutations.

1.1 Preliminaries

Throughout the paper we assume that $m \geq 4$ is an integer and $q = 2^m$. For $z \in \mathbb{F}_{q^2}$ we denote $\bar{z} = z^q$. The set

$$U = \{z \in \mathbb{F}_{q^2} \mid z^{q+1} = 1\}$$

will play an important role. Since $z^{q+1} = z\bar{z}$ is the norm of $z \in \mathbb{F}_{q^2}$ with respect to \mathbb{F}_q , the set U is sometimes called the “unit circle” in \mathbb{F}_{q^2} . A well known fact is that every $z \in \mathbb{F}_{q^2}^*$ can be written *uniquely* as $z = xy$ where $x \in \mathbb{F}_q$ and $y \in U$. Of course, for $z \in U$ we have $z^q = \frac{1}{z}$. This computational rule will come handy in many proofs in this paper. Some of the calculations in this paper, in particular factorization of polynomials and the computation of a resultant of polynomials, were done with the support of the computer algebra system Magma [1].

We will need a condition for solubility of certain quadratic equations by elements of U distinct from 1.

Lemma 1.2. *Assume that $A, B, C \in \mathbb{F}_q$ and the equation $Az^2 + Bz + C = 0$ has a solution $z \in U \setminus \{1\}$. Then $A = C$.*

Proof. Suppose $Az^2 + Bz + C = 0$ and $z \in U \setminus \{1\}$. After raising the equation to the q -th power and simplifying we get $A + Bz + Cz^2 = 0$. Adding the equations yields $(A+C)(z^2+1) = 0$. Since $z \neq 1$, the conclusion follows. \square

1.2 Characterizations of Kim-type APN functions

In Section 1.2 we summarize the most important results of the recent preprints [9] and [6].

We start with an observation which is not stated as a numbered item in [9] but we want to give it a label for our future reference, and we also state it in a slightly different form.

Lemma 1.3. *Any function $f : \mathbb{F}_{q^2} \rightarrow \mathbb{F}_{q^2}$ given by $f(x) = x^{3q} + a_1x^{2q+1} + a_3x^3$ where $a_1, a_3 \in \mathbb{F}_{q^2}$ is affine equivalent to a function $f'(x) = x^{3q} + a'_1x^{2q+1} + a'_3x^3$ where $a'_1 \in \mathbb{F}_q$ and $a'_3 \in \mathbb{F}_{q^2}$.*

Proof. This is proved at the bottom of page 2 in [9]. The proof given there applies to the general case when $f(x)$ also contains the term a_2x^{q+2} . Denote the function obtained in the proof by $f'(x) = x^{3q} + a'_1x^{2q+1} + a'_2x^{q+2} + a'_3x^3$. It follows from the proof that if $a_2 = 0$, then $a'_2 = 0$. \square

The following constants are introduced in [9]:

$$\begin{cases} \theta_1 = 1 + a_1^2 + a_2\bar{a}_2 + a_3\bar{a}_3 \\ \theta_2 = a_1 + \bar{a}_2a_3 \\ \theta_3 = \bar{a}_2 + a_1\bar{a}_3 \\ \theta_4 = a_1^2 + a_2\bar{a}_2. \end{cases} \quad (1)$$

The following theorem is the main result of [9].

Theorem 1.4. [9, Theorem 1.2] *Let $n = 2m$ with $m \geq 4$ and $f(x) = \bar{x}^3 + a_1\bar{x}^2x + a_2\bar{x}x^2 + a_3x^3$, where $a_1 \in \mathbb{F}_{2^m}$, $a_2, a_3 \in \mathbb{F}_{2^n}$. Let θ_i 's be defined as in (1) and define*

$$\Gamma_1 = \left\{ (a_1, a_2, a_3) \mid \theta_1 \neq 0, \operatorname{Tr}_m \left(\frac{\theta_2\bar{\theta}_2}{\theta_1^2} \right) = 0, \theta_1^2\theta_4 + \theta_1\theta_2\bar{\theta}_2 + \theta_2^2\theta_3 + \bar{\theta}_2^2\bar{\theta}_3 = 0 \right\} \quad (2)$$

and

$$\Gamma_2 = \left\{ (a_1, a_2, a_3) \mid \theta_1 \neq 0, \operatorname{Tr}_m \left(\frac{\theta_2\bar{\theta}_2}{\theta_1^2} \right) = 0, \theta_1^2\theta_3 + \theta_1\bar{\theta}_2^2 + \theta_2^2\theta_3 + \bar{\theta}_2^2\bar{\theta}_3 = 0 \right\}. \quad (3)$$

Then f is APN over \mathbb{F}_{2^n} if and only if

(1) m is even, $(a_1, a_2, a_3) \in \Gamma_1 \cup \Gamma_2$; or

(2) m is odd, $(a_1, a_2, a_3) \in \Gamma_1$.

Prior to the appearance of the manuscript [9], a characterization of the Kim-type APN functions for the special case when the coefficients a_i lie in the subfield \mathbb{F}_q was given in [8], and then the following result on equivalence with Gold functions was proved in [6].

Theorem 1.5. [6] Suppose that $m \geq 4$ is an integer and let $q = 2^m$. Let $f : \mathbb{F}_{q^2} \rightarrow \mathbb{F}_{q^2}$ be given by $f(x) = x^{3q} + a_1x^{2q+1} + a_2x^{q+2} + a_3x^3$ where $a_1, a_2, a_3 \in \mathbb{F}_q$. If f is APN, then f is affine equivalent to $G_1(x) = x^3$ or f is affine equivalent to $G_2(x) = x^{2^{m-1}+1}$.

In order to make the present paper self-contained, in the remainder of Section 1.2 we include the proof of Theorem 1.5 using Theorem 1.4 and some ideas of [6].

Assume that $a_1, a_2, a_3 \in \mathbb{F}_q$. Then $\theta_1 = (1 + a_1 + a_2 + a_3)^2$ and equations (2) and (3) factorize as $\theta_1 S^2 = 0$ and $\theta_1 ST = 0$ respectively, where

$$S = a_1^2 + a_1a_3 + a_2^2 + a_2$$

and

$$T = a_1a_3 + a_2 + a_3^2 + 1.$$

If f is APN, then $\theta_1 \neq 0$, thus we must have $S = 0$ for equation (2) to hold, and we must have $ST = 0$ for equation (3) to hold.

Recall that a linear map from \mathbb{F}_{q^2} to \mathbb{F}_{q^2} is a bijection if and only if it has no nonzero roots in \mathbb{F}_{q^2} . In the following proof of Theorem 1.5 we will often use linear maps of the form $L(x) = x^q + tx$ which are bijections if and only if $t \notin U$. For the case of $t \in \mathbb{F}_q$ we only need to check that $t \neq 1$ for L to be a bijection.

First we will show that if the function f given in Theorem 1.5 is APN, then f is affine equivalent to a function that has one of the following forms:

$$(i) \quad f_1(x) = x^{3q} + c_1x^{2q+1} + c_2x^{q+2},$$

$$(ii) \quad f_2(x) = x^{3q} + c_2x^{q+2} + x^3,$$

with $c_1, c_2 \in \mathbb{F}_q$.

If $a_3 = 0$ then we are done. Assume $a_3 = 1$. If $a_1 = 0$ then we are done. If $a_1 = a_2$ then $f(x) \in \mathbb{F}_q$ for all $x \in \mathbb{F}_{q^2}$, which contradicts f being APN. If $a_1 \notin \{0, a_2\}$ then let $r = a_2/a_1$, and let

$$L(u) = \frac{u^q + ru}{r + 1}.$$

The function $f_2(x) = L(f(x))$ is affine equivalent to f , and f_2 is of the form (ii) given above.

Now assume $a_3 \notin \{0, 1\}$. Let

$$L(u) = \frac{a_3 u^q + u}{a_3^2 + 1}.$$

The function $f_1(x) = L(f(x))$ is affine equivalent to f , and f_1 is of the form (i) given above. Therefore, if f is APN, it must be affine equivalent to a function of the form (i) or (ii).

Now we will show that APN functions of the form f_1 or f_2 are affine equivalent to G_1 or G_2 . First consider f_1 . Suppose

$$S = c_1^2 + c_2^2 + c_2 = 0.$$

If $c_2 = 0$ then $c_1 = 0$ and $f_1(x)^q = G_1(x)$. If $c_2 = 1$ then $S = c_1^2 = 0$, but by Theorem 1.4, we require that $\theta_1 = c_1^2 \neq 0$. So assume $c_2 \notin \{0, 1\}$. Let $b \in \mathbb{F}_q \setminus \mathbb{F}_2$ be such that $c_2 = (b+1)^2$. Then it follows from $S = 0$ that $c_1 = b^2 + b$. So f_1 is of the form

$$f_1(x) = x^{3q} + (b^2 + b)x^{2q+1} + (b+1)^2 x^{q+2}. \quad (4)$$

Let $r \in \mathbb{F}_q \setminus \mathbb{F}_4$ so that $r^3 \neq 1$ and the linear maps $L_1(x) = x^q + rx$ and $L_2(x) = r^3 x^q + x$ are bijections. We have

$$L_2(G_1(L_1(x))) = (r^6 + 1)x^{3q} + (r^5 + r)x^{2q+1} + (r^4 + r^2)x^{q+2}.$$

Dividing by $(r^6 + 1)$, we see that G_1 is affine equivalent to

$$F_1(x) = x^{3q} + \frac{r^5 + r}{r^6 + 1} x^{2q+1} + \frac{r^4 + r^2}{r^6 + 1} x^{q+2}.$$

Letting

$$b' = \frac{(r+1)^3}{r^3 + 1},$$

we have

$$F_1(x) = x^{3q} + (b'^2 + b')x^{2q+1} + (b' + 1)^2 x^{q+2}.$$

From the trace condition of Theorem 1.4 applied to function f_1 in equation (4) we have

$$\text{Tr}_m \left(\frac{\theta_2^{q+1}}{\theta_1^2} \right) = \text{Tr}_m \left(\frac{\theta_2}{\theta_1} \right) = \text{Tr}_m \left(\frac{b^2 + b}{b^2} \right) = 0,$$

equivalently

$$\mathrm{Tr}_m\left(\frac{1}{b}\right) = \mathrm{Tr}_m(1).$$

To show that f_1 is affine equivalent to G_1 , we must show that b' takes all values from $\mathbb{F}_q \setminus \mathbb{F}_2$ such that $\mathrm{Tr}_m(b'^{-1}) = \mathrm{Tr}_m(1)$ as r runs through $\mathbb{F}_q \setminus \mathbb{F}_4$. Let $r' = r + 1$. Then as r runs through $\mathbb{F}_q \setminus \mathbb{F}_4$, so does r' , and

$$\begin{aligned} \mathrm{Tr}_m(b'^{-1}) &= \mathrm{Tr}_m\left(\frac{r^3 + 1}{(r + 1)^3}\right) = \mathrm{Tr}_m\left(\frac{r'^3 + r'^2 + r'}{r'^3}\right) \\ &= \mathrm{Tr}_m(1) + \mathrm{Tr}_m\left(\frac{1}{r'}\right) + \mathrm{Tr}_m\left(\frac{1}{r'^2}\right) = \mathrm{Tr}_m(1). \end{aligned}$$

Therefore, if $S = 0$, then f_1 is affine equivalent to G_1 .

Now suppose $S \neq 0$, that is, m is even and $T = c_2 + 1 = 0$. Then $c_2 = 1$ and

$$f_1(x) = x^{3q} + c_1 x^{2q+1} + x^{q+2}.$$

By Theorem 1.4, we require $\theta_1 = c_1^2 \neq 0$, that is, $c_1 \neq 0$. We also require the trace condition

$$\mathrm{Tr}_m\left(\frac{\theta_2^{q+1}}{\theta_2^2}\right) = \mathrm{Tr}_m\left(\frac{1}{c_1}\right) = 0.$$

Let $r \in \mathbb{F}_q \setminus \mathbb{F}_2$ and let $L(x) = x^q + rx$. We have

$$L(G_2(L(x)^{2q})) = (r^3 + r)x^{3q} + (r + 1)^4 x^{2q+1} + (r^3 + r)x^{q+2}.$$

After dividing by $r^3 + r$, we see that G_2 is affine equivalent to

$$F_1(x) = x^{3q} + dx^{2q+1} + x^{q+2}$$

where

$$d = \frac{(r + 1)^2}{r}.$$

Let $r' = r + 1$ so that r' is also in $\mathbb{F}_q \setminus \mathbb{F}_2$. We have

$$\mathrm{Tr}_m\left(\frac{1}{d}\right) = \mathrm{Tr}_m\left(\frac{r}{(r + 1)^2}\right) = \mathrm{Tr}_m\left(\frac{r' + 1}{r'^2}\right) = \mathrm{Tr}_m\left(\frac{1}{r'}\right) + \mathrm{Tr}_m\left(\frac{1}{r'^2}\right) = 0$$

which means that d takes all values in \mathbb{F}_q^* such that $\mathrm{Tr}_m(d^{-1}) = 0$. Thus f_1 is affine equivalent to G_2 when $S \neq 0$.

Considering f_2 , we note that $\theta_1 = c_2^2$ must be nonzero. Since $T = c_2$ we must also have $S = c_2^2 + c_2 = 0$. Therefore $c_2 = 1$. From the trace condition, we have

$$\text{Tr}_m \left(\frac{\theta_2^{q+1}}{\theta_1^2} \right) = \text{Tr}_m \left(\frac{\theta_2}{\theta_1} \right) = \text{Tr}_m(1) = 0.$$

Therefore m must be even, so \mathbb{F}_q contains a primitive cube root of unity ω . We have

$$L(G_1(L(x))) = (1 + \omega)f_2(x)$$

where $L(x) = x^q + \omega x$ which is a bijection since $\omega \notin U$ when m is even. This completes the proof of Theorem 1.5.

2 Main result

Theorem 2.1. *Suppose that $m \geq 4$ is an integer and let $q = 2^m$. Let $f : \mathbb{F}_{q^2} \rightarrow \mathbb{F}_{q^2}$ be given by $f(x) = x^{3q} + a_1x^{2q+1} + a_2x^{q+2} + a_3x^3$ where $a_1, a_2, a_3 \in \mathbb{F}_{q^2}$. If f is APN, then f is affine equivalent to $G_1(x) = x^3$ or f is affine equivalent to $G_2(x) = x^{2^{m-1}+1}$.*

Proof. The proof of the theorem is obtained by combining the results of Lemma 1.3, Theorem 1.4, and Propositions 2.2 through 2.8. \square

Proposition 2.2. *Let $f : \mathbb{F}_{q^2} \rightarrow \mathbb{F}_{q^2}$ be given by $f(x) = x^{3q} + a_1x^{2q+1} + a_2x^{q+2} + a_3x^3$ where $a_1 \in \mathbb{F}_q$, $a_2 \in \mathbb{F}_{q^2}^*$ and $a_3 \in \mathbb{F}_{q^2}$. If $a_1/a_2 \notin U$ and $a_1/a_2 \neq a_3^q$, then f is affine equivalent to $h(x) = x^{3q} + a'_1x^{2q+1} + a'_3x^3$ where $a'_1 \in \mathbb{F}_q$ and $a'_3 \in \mathbb{F}_{q^2}$.*

Proof. Let $t = a_1/a_2$ and let $h_0(x) = L(f(x))$ where $L(z) = z^q + tz$. Then

$$h_0(x) = \left(\frac{a_1}{a_2} + a_3^q \right) x^{3q} + \left(\frac{a_1^2}{a_2} + a_2^q \right) x^{2q+1} + \left(\frac{a_1 a_3}{a_2} + 1 \right) x^3.$$

Since $a_1/a_2 \notin U$, functions f and h_0 are affine equivalent. Now let

$$h_1(x) = \left(\frac{a_1}{a_2} + a_3^q \right)^{-1} \cdot h_0(x)$$

and apply Lemma 1.3 to function h_1 . \square

We first deal with the special cases when two coefficients of the function vanish.

Proposition 2.3. *Let $f : \mathbb{F}_{q^2} \rightarrow \mathbb{F}_{q^2}$ be given by $f(x) = x^{3q} + a_1x^{2q+1} + a_3x^3$ where $a_1, a_3 \in \mathbb{F}_{q^2}$ and $a_1 = 0$ or $a_3 = 0$. If f is APN, then f is affine equivalent to $G_1(x) = x^3$.*

Proof. Assume $a_1 = a_2 = 0$. Then $\theta_1 = a_3^{q+1} + 1$. If $a_3 \in U$, then $\theta_1 = 0$ and f is not APN. If $a_3 \notin U$, let $L(z) = z^q + tz$ where $t = a_3^q$. Then L is a permutation of \mathbb{F}_{q^2} and $L(f(x)) = (a_3^{q+1} + 1)x^3$.

Next assume $a_2 = a_3 = 0$. Then $\theta_1 = (a_1 + 1)^2$. After plugging into (2) we get $a_1^4(a_1 + 1)^2 = 0$, and after plugging into (3) we get $a_1^2(a_1 + 1)^2 = 0$. If f is APN, then $\theta_1 \neq 0$, hence $a_1 = 0$ and $f(x) = x^{3q}$ which is affine equivalent to $G_1(x) = x^3$. \square

Proposition 2.4. *Let $f : \mathbb{F}_{q^2} \rightarrow \mathbb{F}_{q^2}$ be given by $f(x) = x^{3q} + a_1x^{2q+1} + a_3x^3$ where $a_1 \in \mathbb{F}_q$ and $a_3 \in \mathbb{F}_{q^2}$. If $(a_1, 0, a_3) \in \Gamma_1$ as defined in equation (2) then f is affine equivalent to $G_1(x) = x^3$ or f is affine equivalent to $G_2(x) = x^{2^{m-1}+1}$.*

Proof. We will show that $a_3 \in \mathbb{F}_q$ or $a_1 = 0$ and then the conclusion will follow by applying Theorem 1.5 or Proposition 2.3 accordingly.

If $a_1 = 0$ or $a_3 = 0$, then the conclusion follows immediately. Hence assume that $a_1 \neq 0$ and $a_3 \neq 0$. Let $a_3 = yz$ where $y \in \mathbb{F}_q^*$ and $z \in U$. After plugging this along with $a_2 = 0$ into equation (2), dividing by a_1^2 and simplifying we get

$$a_1yz^2 + (a_1^4 + a_1^2 + y^4 + y^2)z + a_1y = 0. \quad (5)$$

By the definition of set Γ_1 we have

$$\text{Tr}_m \left(\frac{\theta_2^{q+1}}{\theta_1^2} \right) = \text{Tr}_m \left(\frac{a_1^2}{(a_1^2 + y^2 + 1)^2} \right) = \text{Tr}_m \left(\frac{a_1}{a_1^2 + y^2 + 1} \right) = 0.$$

Hence there exist $t \in \mathbb{F}_q$ such that

$$(t^2 + t)(a_1^2 + y^2 + 1) + a_1 = 0. \quad (6)$$

The resultant of the left-hand sides of (5) and (6) with respect to y is $a_1^2(Az^2 + B)(Bz^2 + A)$ where $A = (a_1t + t + 1)t^3$ and $B = (a_1t + a_1 + t)(t + 1)^3$.

If $z = 1$, then $a_3 \in \mathbb{F}_q$. Otherwise, we have $A, B \in \mathbb{F}_q$ but $z \notin \mathbb{F}_q$. Then the resultant can vanish only if $A = B = 0$. This occurs only if $(a_1, t) = (0, 0)$ or $(a_1, t) = (0, 1)$. This completes the proof. \square

Proposition 2.5. *Let $f : \mathbb{F}_{q^2} \rightarrow \mathbb{F}_{q^2}$ be given by $f(x) = x^{3q} + a_1x^{2q+1} + a_3x^3$ where $a_1 \in \mathbb{F}_q$ and $a_3 \in \mathbb{F}_{q^2}$. If $(a_1, 0, a_3) \in \Gamma_2$ as defined in equation (3) then f is affine equivalent to $G_1(x) = x^3$ or f is affine equivalent to $G_2(x) = x^{2^{m-1}+1}$.*

Proof. We will show that $a_3 \in \mathbb{F}_q$ and then the conclusion will follow by applying Theorem 1.5.

Again let $a_3 = yz$ where $y \in \mathbb{F}_q$ and $z \in U$. After plugging this along with $a_2 = 0$ into equation (3) and simplifying we get

$$(a_1^2y)z^2 + (a_1^3 + a_1y^2 + a_1)z + a_1^4y + a_1^2y + y^5 + y = 0.$$

If $z = 1$ then $a_3 \in \mathbb{F}_q$ and we are done. Otherwise, by Lemma 1.2 we must have

$$(a_1^2y) + (a_1^4y + a_1^2y + y^5 + y) = y(a_1 + y + 1)^4 = 0.$$

If $y = 0$, then $a_3 \in \mathbb{F}_q$. If $(a_1 + y + 1)^4 = 0$, then $\theta_1 = (a_1 + y + 1)^2 = 0$ and $(a_1, 0, a_3) \notin \Gamma_2$ by the first condition of equation (3). \square

Proposition 2.6. *Let $f : \mathbb{F}_{q^2} \rightarrow \mathbb{F}_{q^2}$ be given by $f(x) = x^{3q} + a_1x^{2q+1} + a_2x^{q+2} + a_3x^3$ where $a_1 \in \mathbb{F}_q$, $a_2 \in \mathbb{F}_{q^2}^*$ and $a_3 \in \mathbb{F}_{q^2}$, and assume that $a_1/a_2 \in U$. If f is APN, then m is even, and furthermore $a_1, a_2, a_3 \in \mathbb{F}_q$ or there exist $u, z \in U$ such that*

$$\begin{aligned} a_1 &= \frac{(u^3 + z)^2}{u(u^2 + z)^2} \\ a_2 &= \frac{(u^3 + z)^2}{(u^2 + z)^2} \\ a_3 &= \frac{uz^2(u + 1)^2}{(u^2 + z)^2}. \end{aligned}$$

Proof. Under the assumptions listed in the proposition, we first prove that $(a_1, a_2, a_3) \notin \Gamma_1$, and then we prove that $(a_1, a_2, a_3) \in \Gamma_2$ implies the conclusions given in the statement.

From $a_1/a_2 \in U$ it follows that $a_1 \neq 0$.

Let $a_2 = a_1u$ where $u \in U$, and let $a_3 = yz$ where $y \in \mathbb{F}_q$ and $z \in U$. After substituting this into (2), simplifying and clearing denominators we get

$$a_1^2(a_1u^4y + a_1u^3z + u^2y^2z + a_1yz^2 + a_1uz + u^2z)(yz + u)(uy + z) = 0.$$

Since $u/z \in U$, any of the last two factors can vanish only if $y = 1$, but then $\theta_1 = y^2 + 1 = 0$ and $(a_1, a_2, a_3) \notin \Gamma_1$. Since $a_1 \neq 0$, the second factor must vanish, hence

$$a_1(u^4y + u^3z + yz^2 + uz) = u^2z(y^2 + 1).$$

The right-hand side can not vanish, hence we have

$$a_1 = \frac{u^2z(y^2 + 1)}{u^4y + u^3z + yz^2 + uz}. \quad (7)$$

We will obtain the desired conclusion by invoking the condition $\text{Tr}_m \left(\frac{\theta_2^{q+1}}{\theta_1^2} \right) = 0$ from the definition of set Γ_1 in equation (2). After substituting the formula (7) for a_1 into θ_2 we get

$$\frac{\theta_2^{q+1}}{\theta_1^2} = \frac{u^3z(yz + u)(uy + z)}{(u^4y + u^3z + yz^2 + uz)^2}.$$

For the trace value condition to hold there must exist $w \in \mathbb{F}_q$ such that

$$(w^2 + w)(u^4y + u^3z + yz^2 + uz)^2 + u^3z(yz + u)(uy + z) = 0.$$

This equation factors as

$$(Dw + z(yz + u))(Dw + u^3(uy + z)) = 0$$

where $D = u^4y + u^3z + yz^2 + uz$ is the denominator in (7). We already know that $D \neq 0$. Hence the values of w are

$$w_1 = \frac{z(yz + u)}{D}, \quad w_2 = \frac{u^3(uy + z)}{D}.$$

Let

$$r = \frac{a_1}{w_1} = \frac{u^2(y^2 + 1)}{yz + u}.$$

Since $a_1, w_1 \in \mathbb{F}_q$, we must have $r \in \mathbb{F}_q$. However

$$r + r^q = \frac{(y + 1)^2 D}{u(yz + u)(uy + z)} \neq 0$$

which finishes the proof that $(a_1, a_2, a_3) \notin \Gamma_1$.

Thus, if f is APN, then we must have $(a_1, a_2, a_3) \in \Gamma_2$ and m must be even. We will analyze this case now.

Again assume that $a_2 = a_1u$ and $a_3 = yz$ where $y \in \mathbb{F}_q$ and $u, z \in U$. By Proposition 2.3 we can assume $a_3 \neq 0$, hence $y \neq 0$. After plugging the expressions for a_2 and a_3 into (3) and simplifying we get

$$a_1(uy + z)(Aa_1^2 + Ba_1 + C) = 0$$

where $A = D(yz + u)$, $D = (u^2 + z)^2y + uz(u^2 + 1)$, $B = (y + 1)^2u^3(uy + z)$ and $C = (y + 1)^4u^2z$. Since $\theta_1 = 1 + y^2$, we have $y \neq 1$, hence $uy + z \neq 0$ and $yz + u \neq 0$.

We will distinguish two cases: $A = 0$ and $A \neq 0$, equivalently, $D = 0$ and $D \neq 0$.

First let us assume that $D = 0$. If $u^2 = z$, then $u = z = 1$ and $a_1, a_2, a_3 \in \mathbb{F}_q$. Otherwise

$$y = \frac{uz(u^2 + 1)}{(u^2 + z)^2}$$

and

$$a_1 = \frac{C}{B} = \frac{(y + 1)^2z}{u(uy + z)} = \frac{(u^3 + z)^2}{u(u^2 + z)^2}.$$

Then

$$a_2 = a_1u = \frac{(u^3 + z)^2}{(u^2 + z)^2}$$

and

$$a_3 = yz = \frac{uz^2(u + 1)^2}{(u^2 + z)^2}.$$

With these values of a_1, a_2, a_3 we get after simplifications $\theta_2^{q+1}/\theta_1^2 = 1$, hence $\text{Tr}_m(\theta_2^{q+1}/\theta_1^2) = 0$ since m is even, and $(a_1, a_2, a_3) \in \Gamma_2$.

We complete the proof by analyzing the case $A \neq 0$. We note that a_1 is also a root of

$$Aa_1^2 + Ba_1 + C + \frac{A}{A^q}(A^qa_1^2 + B^qa_1 + C^q) = 0,$$

which after simplifications becomes $u(Ra_1 + S) = 0$ where

$R = (u + z)^4(y + 1)^2y^2$ and $S = (y + 1)^4(uyz^2 + u^2z + uy + z)uz$. If $u = z$, then it must be $S = z^3(y + 1)^5(z + 1)^2 = 0$. Since $\theta_1 = (y + 1)^2$, we get $u = z = 1$ and $a_1, a_2, a_3 \in \mathbb{F}_q$. If $y = 0$, then we must have $S = z^2u(u + 1)^2 = 0$, hence $u = 1$ and $a_1, a_2, a_3 \in \mathbb{F}_q$. Otherwise we get

$$a_1 = \frac{S}{R} = \frac{(y + 1)^2(uyz^2 + u^2z + uy + z)uz}{(u + z)^4y^2}. \quad (8)$$

After plugging this into (3) we get

$$\frac{z^2(y+1)^4u^2(u+1)^2(u+yz)((u+z^2)y+z(u+1))^2D}{(y(u+z)^2)^4} = 0. \quad (9)$$

If $u = 1$ then

$$a_1 = \frac{z(y+1)^2}{y(z+1)^2}$$

and

$$\frac{\theta_2^{q+1}}{\theta_1^2} = \frac{(yz+1)(y+z)z}{y^2(z+1)^4}.$$

For the trace condition to be satisfied there must exist $w \in \mathbb{F}_q$ such that

$$(w^2 + w)(y^2(z+1)^4) + (yz+1)(y+z)z = 0 \quad (10)$$

which factors as

$$(wyz^2 + wy + y + z)(wyz^2 + wy + yz^2 + z) = 0$$

and the root of the first factor is

$$w_1 = \frac{y+z}{y(z^2+1)}$$

for which we get

$$w_1 + w_1^q = 1.$$

Hence $w_1 \notin \mathbb{F}_q$. Of course for the other root of (10) we have $w_2 = w_1 + 1 \notin \mathbb{F}_q$. Therefore the case $u = 1$ can not occur.

Otherwise, $u \neq 1$ and since we are also assuming $D \neq 0$, we get from (9)

$$y = \frac{z(u+1)}{u+z^2}$$

and after plugging this into (8) we get

$$a_1 = \frac{u(z+1)^2}{(u+1)(u+z^2)}$$

and after plugging the last two formulas into $\theta_2^{q+1}/\theta_1^2$ and simplifying we get

$$\frac{\theta_2^{q+1}}{\theta_1^2} = \frac{u}{(u+1)^2}.$$

Since we require $\text{Tr}_m(\theta_2^{q+1}/\theta_1^2) = 0$, there must exist $w \in \mathbb{F}_q$ such that $(w^2 + w)(u + 1)^2 + u = 0$. This factors as

$$(uw + w + 1)(uw + u + w) = 0$$

and the roots are $w_1 = 1/(u + 1)$ and $w_2 = u/(u + 1)$. However $w_1 + w_1^q = w_2 + w_2^q = 1$ which means that $w_1, w_2 \notin \mathbb{F}_q$ and $\text{Tr}_m(\theta_2^{q+1}/\theta_1^2) \neq 0$. Hence the case $A \neq 0$ does not produce APN functions, and the entire proof of the proposition is now completed. \square

Proposition 2.7. *Let $f : \mathbb{F}_{q^2} \rightarrow \mathbb{F}_{q^2}$ be given by $f(x) = x^{3q} + a_1x^{2q+1} + a_2x^{q+2} + a_3x^3$ where*

$$\begin{aligned} a_1 &= \frac{(u^3 + z)^2}{u(u^2 + z)^2} \\ a_2 &= \frac{(u^3 + z)^2}{(u^2 + z)^2} \\ a_3 &= \frac{uz^2(u + 1)^2}{(u^2 + z)^2} \end{aligned}$$

for some $u, z \in U$ such that $u^2 \neq z$, and m is even. If f is APN, then f is affine equivalent to $G_1(x) = x^3$ or f is affine equivalent to $G_2(x) = x^{2^{m-1}+1}$.

Proof. Since m is even, \mathbb{F}_q contains a primitive cube root of unity, which we will denote ω , that is, $\omega^2 + \omega + 1 = 0$.

Let $D = (\omega^2u^3 + (u + \omega)z)^2$. Let $L_1(x) = x^q + tx$ where

$$t = \frac{(\omega u^3 + (u + \omega^2)z)^2}{uD} \tag{11}$$

and let $L_2(x) = x^{2q} + sx^2$ where $s = \omega u^2$.

Let us consider the cases when the denominator of (11) may vanish. This happens when $z_1 \in U$ where

$$z_1 = \frac{w^2u^3}{u + \omega}.$$

We have

$$\frac{1}{z_1} + z_1^q = \frac{\omega(u + 1)^2}{u^3(\omega + u + 1)}$$

hence $z_1 \in U$ only if $u = 1$, but then $z_1 = 1$ and $u^2 = z_1$, hence this case can not occur.

Let $G_2(x) = x^{2^{m-1}+1}$ be a Gold APN function. Denote

$$F(x) = L_1(G_2(L_2(x))) = c_0x^{3q} + c_1x^{2q+1} + c_2x^{q+2} + c_3x^3. \quad (12)$$

A calculation shows that

$$\begin{aligned} c_0 &= \frac{(u^2 + z)^2}{D} \\ c_1 &= \frac{(u^3 + z)^2}{uD} \\ c_2 &= \frac{(u^3 + z)^2}{D} \\ c_3 &= \frac{uz^2(u+1)^2}{D}. \end{aligned}$$

Since $u^2 \neq z$, we have $c_0 \neq 0$. Since $c_i/c_0 = a_i$ for all $i = 1, 2, 3$, we get that $F(x) = c_0f(x)$.

We need to show that L_1 and L_2 are permutations of \mathbb{F}_{q^2} . For L_1 this is the case if and only if $t \notin U$. We can assume without loss of generality that $t \neq 0$ and compute

$$t^q + \frac{1}{t} = \frac{u^3(u^3 + z)^2(u + z)^2}{\omega(u^3 + \omega^2uz + \omega z)^2(u^3 + \omega^2u^2 + \omega z)^2}.$$

Hence $t^{q+1} \neq 1$ unless $u^3 = z$ or $u = z$. If $u^3 = z$ then $a_1 = a_2 = 0$ and f is affine equivalent to $G_1(x) = x^3$ by Proposition 2.3. If $u = z$ then $\theta_1 = 0$ and f is not APN.

To show that L_2 is a permutation of \mathbb{F}_{q^2} we need to show that the only root of $L_2(x) = x^2(x^{2q-2} + s)$ in \mathbb{F}_{q^2} is 0, equivalently, that there does not exist $x \in \mathbb{F}_{q^2}$ such that $x^{2q-2} = s$. After raising this equation to power $q+1$ the left-hand side becomes $(x^{2q-2})^{q+1} = (x^{q^2-1})^2 = 1$ but the right-hand side becomes $s^{q+1} = (\omega u^2)^{q+1} = \omega^{q+1} = \omega^2$ since m is even. Hence the equation can not hold, and L_2 is a permutation of \mathbb{F}_{q^2} . \square

Proposition 2.8. *Let $f : \mathbb{F}_{q^2} \rightarrow \mathbb{F}_{q^2}$ be given by $f(x) = x^{3q} + a_1x^{2q+1} + a_2x^{q+2} + a_3x^3$ where $a_1 \in \mathbb{F}_q$, $a_2 \in \mathbb{F}_{q^2}^*$ and $a_3 \in \mathbb{F}_{q^2}$, and assume that $a_1/a_2 = a_3^q$. If f is APN, then f is affine equivalent to $G_1(x) = x^3$.*

Proof. Assume $a_1 = a_2a_3^q$. There exist $v, y \in \mathbb{F}_q$ and $z \in U$ such that $a_1 = vy$, $a_2 = vz$ and $a_3 = yz$. After simplifications we get $\theta_1 = (v+1)^2(y+1)^2$. After plugging into (2) and simplifications we get

$$v^2(v+1)^4(y+1)^6 = 0.$$

Since $\theta_1 \neq 0$, we must have $v = 0$. Then $a_1 = a_2 = 0$ and the conclusion follows from Proposition 2.3. Finally after plugging into (3) and simplifications we get

$$\frac{v(v+1)^4(y+1)^6}{z} = 0$$

and the conclusion is the same as in the previous case. \square

References

- [1] W. Bosma, J. Cannon and C. Playoust, The Magma algebra system. I. The user language. *J. Symbolic Comput.* **24** (1997), 235–265.
- [2] K.A. Browning, J.F. Dillon, M.T. McQuistan, A.J. Wolfe, An APN permutation in dimension six. *Finite fields: theory and applications*, 33–42, *Contemp. Math.*, 518, Amer. Math. Soc., Providence, RI, 2010.
- [3] L. Budaghyan, M. Calderini, I. Villa, On equivalence between known families of quadratic APN functions. *Finite Fields and Their Applications* **66** (2020), 101704.
- [4] C. Carlet, Open questions on nonlinearity and on APN functions. *Arithmetic of finite fields*, 83–107, *Lecture Notes in Comput. Sci.*, 9061, Springer, Cham, 2015.
- [5] C. Carlet, P. Charpin, V. Zinoviev, Codes, bent functions and permutations suitable for DES-like cryptosystems. *Des. Codes Cryptogr.* **15** (1998), 125–156.
- [6] F. Göloğlu, D. Krasnayová, P. Lisoněk, Generalized Kim functions are not equivalent to permutations. Preprint, 2020.
- [7] F. Göloğlu, P. Langevin, Almost perfect nonlinear families which are not equivalent to permutations. *Finite Fields Appl.* **67** (2020), 101707.
- [8] D. Krasnayová, Constructions of APN permutations. Diploma Thesis. Charles University, Prague, 2016.
<https://dspace.cuni.cz/handle/20.500.11956/83075> (accessed on 12 September 2020)
- [9] K. Li, C. Li, T. Helleseht, L. Qu, A complete characterization of the APN property of a class of quadrimonomials. arXiv:2007.03996 [cs.IT] (accessed on 19 August 2020)

- [10] S. Li, W. Meidl, A. Polujan, A. Pott, C. Riera, P. Stănică, Vanishing flats: a combinatorial viewpoint on the planarity of functions and their application. To appear in IEEE Transactions on Information Theory. arXiv:2006.01941 [cs.IT] (accessed on 10 September 2020)