

Continuous time approximation of Nash equilibria

Romeo Awi*, Ryan Hynd†, and Henok Mawi‡

September 15, 2020

Abstract

We consider the problem of approximating Nash equilibria of N functions f_1, \dots, f_N of N variables. In particular, we deduce conditions under which systems of the form

$$\dot{u}_j(t) = -\nabla_{x_j} f_j(u(t))$$

($j = 1, \dots, N$) are well posed and in which the large time limits of their solutions $u(t) = (u_1(t), \dots, u_N(t))$ are Nash equilibria for f_1, \dots, f_N . To this end, we will invoke the theory of maximal monotone operators. We will also identify an application of these ideas in game theory and show how to approximate equilibria in some game theoretic problems in function spaces.

1 Introduction

Let us first recall the notion of a Nash equilibrium. Consider a collection of N sets X_1, \dots, X_N and define

$$X := X_1 \times \dots \times X_N.$$

For a given $x = (x_1, \dots, x_N) \in X$, $j \in \{1, \dots, N\}$, and $y_j \in X_j$, we will use the notation (y_j, x_{-j}) for the point in X in which y_j replaces x_j in the coordinates of x . That is,

$$(y_j, x_{-j}) := (x_1, \dots, x_{j-1}, y_j, x_{j+1}, \dots, x_N).$$

A collection of functions $f_1, \dots, f_N : X \rightarrow \mathbb{R}$ has a *Nash equilibrium* at $x \in X$ provided

$$f_j(x) \leq f_j(y_j, x_{-j})$$

for all $y_j \in X_j$ and $j = 1, \dots, N$.

*Department of Mathematics, Hampton University.

†Department of Mathematics, University of Pennsylvania.

‡Department of Mathematics, Howard University.

Nash recognized that an equilibrium is a fixed point of the set valued mapping

$$X \ni x \mapsto \arg \min_{y \in X} \left\{ \sum_{j=1}^N f_j(y_j, x_{-j}) \right\}.$$

In particular, he applied Kakutani's fixed point theorem [19] to show that if X_1, \dots, X_N are nonempty, convex, compact subsets of Euclidean space and each f_1, \dots, f_N is multilinear, then f_1, \dots, f_N has a Nash equilibrium at some $x \in X$ [24, 25]. Nash was interested in multilinear f_1, \dots, f_N as they correspond to the expected cost of players assuming mixed strategies in noncooperative games. More generally, his existence theorem holds if each $f_j : X \rightarrow \mathbb{R}$ is continuous and

$$X \ni y \mapsto \sum_{j=1}^N f_j(y_j, x_{-j}) \text{ is convex for each } x \in X. \quad (1.1)$$

As the existence of a Nash equilibrium is due to a nonconstructive fixed point theorem, it seems unlikely that there are good ways to approximate these points. Indeed, it has been established that the approximation of Nash equilibria is computationally challenging [9, 10, 21]. Nevertheless, we contend that there is a nontrivial class of functions f_1, \dots, f_N for which the approximation of Nash equilibria is at least theoretically feasible. We will explain below that a sufficient condition for the approximation of Nash equilibria is that f_1, \dots, f_N satisfy

$$\sum_{j=1}^N (\nabla_{x_j} f_j(x) - \nabla_{x_j} f_j(y)) \cdot (x_j - y_j) \geq 0 \quad (1.2)$$

for each $x, y \in X$. Here we are considering each X_j as a subset of a Euclidean space and use ' \cdot ' to denote the dot products on any of these spaces.

Our approach starts with the observation that if (1.1) holds and X_1, \dots, X_N are convex, then x is a Nash equilibrium if and only if

$$\sum_{j=1}^N \nabla_{x_j} f_j(x) \cdot (y_j - x_j) \geq 0 \quad (1.3)$$

for each $y \in X$. As a result, it is natural to consider the differential inequalities

$$\sum_{j=1}^N (\dot{u}_j(t) + \nabla_{x_j} f_j(u(t))) \cdot (y_j - u_j(t)) \geq 0 \quad (1.4)$$

for $t \geq 0$ and $y \in X$. Here

$$u : [0, \infty) \rightarrow X; t \mapsto (u_1(t), \dots, u_N(t))$$

is the unknown. We will below argue that for appropriately chosen initial condition $u(0) \in X$, the *Cesàro mean* of u

$$t \mapsto \frac{1}{t} \int_0^t u(s) ds$$

will converge to a Nash equilibrium of f_1, \dots, f_N as $t \rightarrow \infty$ provided that (1.2) holds.

1.1 A few concrete examples

An elementary example which illustrates this approach may be observed when $X_1 = X_2 = [-1, 1]$, $X = [-1, 1]^2$,

$$f_1(x_1, x_2) = x_1x_2, \quad \text{and} \quad f_2(x_1, x_2) = -x_1x_2.$$

It is not hard to check that the origin $(0, 0)$ is the unique Nash equilibrium of f_1, f_2 and that f_1, f_2 satisfy (1.2) (with equality holding) for each $x, y \in [-1, 1]^2$. We now seek an absolutely continuous path

$$u : [0, \infty) \rightarrow [-1, 1]^2$$

such that

$$(\dot{u}_1(t) + u_2(t))(y_1 - u_1(t)) + (\dot{u}_2(t) - u_1(t))(y_2 - u_2(t)) \geq 0 \quad (1.5)$$

holds for almost every $t \geq 0$ and each $(y_1, y_2) \in [-1, 1]$.

It turns out that if

$$u_1(0)^2 + u_2(0)^2 \leq 1,$$

then the unique solution of these differential inequalities is

$$\begin{cases} u_1(t) = u_1(0) \cos(t) - u_2(0) \sin(t) \\ u_2(t) = u_2(0) \cos(t) + u_1(0) \sin(t). \end{cases}$$

This solution simply parametrizes the circle of radius $\sqrt{u_1(0)^2 + u_2(0)^2}$. In particular,

$$\lim_{t \rightarrow \infty} \frac{1}{t} \int_0^t u(s) ds = (0, 0).$$

For general initial conditions $(u_1(0), u_2(0)) \in [-1, 1]^2$, it can be shown that $u_1(s)^2 + u_2(s)^2 \leq 1$ in a finite time $s \geq 0$. Refer to Figure 1 for a schematic. It then follows that the same limit above holds for this solution.

The method we present can be extended in certain cases when X_1, \dots, X_N are not compact. For example, suppose $X_1 = X_2 = \mathbb{R}$,

$$f_1(x_1, x_2) = \frac{1}{2}x_1^2 - x_1x_2 - x_1, \quad \text{and} \quad f_2(x_1, x_2) = \frac{1}{2}x_2^2 + x_1x_2 - 2x_2.$$

It is easy to check that f_1, f_2 satisfy (1.2). Furthermore,

$$\begin{cases} \partial_{x_1} f_1(x_1, x_2) = x_1 - x_2 - 1 = 0 \\ \partial_{x_2} f_2(x_1, x_2) = x_2 + x_1 - 2 = 0 \end{cases}$$




Figure 1: Plot of a solution u of (1.5) where the initial position of u is indicated in red. This solution starts out on a circle centered at the origin traversing it counterclockwise until it hits the boundary line segment $(x_1 = 1, -1 \leq x_2 \leq 1)$. Then it proceeds upwards along this boundary line segment until it arrives at some time s at the orange marker which is located at $(1, 0)$. For times $t \geq s$, the position $u(t)$ as shown in purple remains on the unit circle traversing it counterclockwise.

provided

$$x_1 = \frac{3}{2} \quad \text{and} \quad x_2 = \frac{1}{2}.$$

As a result, this is the unique Nash equilibrium of f_1, f_2 .

We note that the solution of

$$\begin{cases} \dot{u}_1(t) = -(u_1(t) - u_2(t) - 1) \\ \dot{u}_2(t) = -(u_2(t) + u_1(t) - 2) \end{cases}$$

is given by

$$\begin{cases} u_1(t) = \frac{3}{2} + \left(u_1(0) - \frac{3}{2}\right) e^{-t} \cos(t) + \left(u_2(0) - \frac{1}{2}\right) e^{-t} \sin(t) \\ u_2(t) = \frac{1}{2} + \left(u_2(0) - \frac{1}{2}\right) e^{-t} \cos(t) + \left(\frac{3}{2} - u_1(0)\right) e^{-t} \sin(t). \end{cases}$$

It is now clear that

$$\lim_{t \rightarrow \infty} (u_1(t), u_2(t)) = \left(\frac{3}{2}, \frac{1}{2} \right).$$

In particular, we do not need to employ the Cesàro mean of u in order to approximate the Nash equilibrium of f_1, f_2 .

1.2 A general setting

It just so happens that our method does not rely on the spaces X_1, \dots, X_N being finite dimensional. Consequently, we will consider the following version of our approximation problem. Let V be a reflexive Banach space over \mathbb{R} with norm $\|\cdot\|$ and continuous dual V^* . We further suppose

$$X_1, \dots, X_N \subset V \text{ are closed and convex with nonempty interiors}$$

and consider N functions

$$f_1, \dots, f_N : X \rightarrow \mathbb{R}$$

which satisfy

$$\left\{ \begin{array}{l} f_1, \dots, f_N \text{ are weakly lowersemicontinuous,} \\ X \ni y \mapsto \sum_{j=1}^N f_j(y_j, x_{-j}) \text{ is convex for each } x \in X, \\ X \ni x \mapsto \sum_{j=1}^N f_j(y_j, x_{-j}) \text{ is weakly continuous for each } y \in X, \text{ and} \\ \sum_{j=1}^N (\partial_{x_j} f_j(x) - \partial_{x_j} f_j(y), x_j - y_j) \geq 0 \text{ for each } x, y \in X. \end{array} \right. \quad (1.6)$$

In the last condition listed above, we mean

$$\sum_{j=1}^N (p_j - q_j, x_j - y_j) \geq 0$$

for each

$$p_j \in \partial_{x_j} f_j(x) := \{\zeta \in V^* : f_j(z, x_{-j}) \geq f_j(x) + (\zeta, z - x_j), z \in X_j\}$$

and $q_j \in \partial_{x_j} f_j(y)$ for $j = 1, \dots, N$.

When V is finite dimensional, we naturally identify V and V^* with Euclidean space of the same dimension. Alternatively, when V is infinite dimensional, we will suppose there is a real Hilbert space H for which

$$V \subset H \subset V^*$$

and $V \subset H$ is continuously embedded. It is with respect to this space H in which we consider the following initial value problem: for a given $u^0 \in X$, find an absolutely continuous $u : [0, \infty) \rightarrow H^N$ such that

$$\begin{cases} u(0) = u^0, \text{ and} \\ \sum_{j=1}^N (\dot{u}_j(t) + \partial_{x_j} f_j(u(t)), y_j - u_j(t)) \geq 0 \text{ for a.e. } t \geq 0 \text{ and each } y \in X. \end{cases} \quad (1.7)$$

Our central result involving this initial value problem is as follows. In this statement, we will make use of the set

$$\mathcal{D} = \left\{ x \in X : \bigcap_{j=1}^N (\partial_{x_j} f_j(x) + n_{X_j}(x_j)) \cap H \neq \emptyset \right\}. \quad (1.8)$$

Here $n_{X_j} : V \rightarrow 2^{V^*}$ is the normal cone of X_j defined as

$$n_{X_j}(z) := \{ \zeta \in V^* : (\zeta, y - z) \leq 0 \text{ all } y \in X_j \}$$

for $z \in V$ ($j = 1, \dots, N$).

Theorem 1.1. *Assume f_1, \dots, f_N satisfy (1.6).*

(i) *For any $u^0 \in \mathcal{D}$ there is a unique Lipschitz continuous $u : [0, \infty) \rightarrow H^N$ satisfying $u(t) \in \mathcal{D}$ for each $t \geq 0$ and the initial value problem (1.7).*

(ii) *If f_1, \dots, f_N has a Nash equilibrium, then*

$$\frac{1}{t} \int_0^t u(s) ds$$

converges weakly in H to a Nash equilibrium of f_1, \dots, f_N as $t \rightarrow \infty$.

We note that if $X_1 = \dots = X_N = V$, then $n_{X_1} \equiv \dots \equiv n_{X_N} \equiv \{0\}$. In this case

$$\mathcal{D} = \left\{ x \in X : \bigcap_{j=1}^N (\partial_{x_j} f_j(x) \cap H) \neq \emptyset \right\},$$

and the initial value problem reduces to

$$\begin{cases} u(0) = u^0, \text{ and} \\ \dot{u}_j(t) + \partial_{x_j} f_j(u(t)) \ni 0 \text{ for a.e. } t \geq 0 \text{ and each } j = 1, \dots, N. \end{cases}$$

We will also explain that if in addition $(\partial_{x_1} f_1, \dots, \partial_{x_N} f_N)$ is single-valued, everywhere-defined, monotone, and hemicontinuous we can obtain the same result as above without assuming (1.6). This follows directly from a theorem of Browder and Minty.

We will phrase our initial value problem (1.7) as the evolution generated by a monotone operator on H^N with domain \mathcal{D} . According to pioneering work of Brézis [5], the crucial task will be to verify the maximality of this operator. Regarding the large time behavior of solutions, the convergence to equilibria of maximal monotone operators by the Cesàro means of semigroups they generate originates in the work of Brézis and Baillon [3]. We also refer the reader to Chapter 3 of [2] which gives a detailed discussion of this phenomenon.

Of course, the asymptotic statements we made above are predicated on the existence of a Nash equilibrium. This is only guaranteed to be the case if X is weakly compact or if there is some $\theta > 0$ such that

$$\sum_{j=1}^N (\partial_{x_j} f_j(x) - \partial_{x_j} f_j(y), x_j - y_j) \geq \theta \|x - y\|^2$$

for each $x, y \in X$. When such coercivity holds, f_1, \dots, f_N has a unique Nash equilibrium and solutions of the initial value problem (1.7) converge exponentially fast in H^N to this equilibrium point.

We will prove Theorem 1.1 in the following section. Then we will apply this result to flows in Euclidean spaces of the form (1.4) in section 3. Finally, we will consider examples in Lebesgue and Sobolev spaces in section 4. A prototypical collection of functionals we will study is

$$f_j(v) = \int_{\Omega} \frac{1}{2} |\nabla v_j|^2 + F_j(v) dx$$

for $j = 1, \dots, N$, where $v = (v_1, \dots, v_N) \in H_0^1(\Omega)^N$ and $\Omega \subset \mathbb{R}^d$ is a bounded domain. For this particular example, we will argue that if $F_1, \dots, F_N : \mathbb{R}^N \rightarrow \mathbb{R}$ satisfy suitable growth and monotonicity conditions, then f_1, \dots, f_N has a unique Nash equilibrium which can be approximated by solutions $u_1, \dots, u_N : \Omega \times [0, \infty) \rightarrow \mathbb{R}$ of the parabolic initial/boundary value problem

$$\begin{cases} \partial_t u_j = \Delta u_j - \partial_{z_j} F_j(u) & \text{in } \Omega \times (0, \infty) \\ u_j = 0 & \text{on } \partial\Omega \times (0, \infty) \\ u_j = u_j^0 & \text{on } \Omega \times \{0\} \end{cases}$$

for appropriately chosen initial conditions $u_1^0, \dots, u_N^0 : \Omega \rightarrow \mathbb{R}$.

We also remark that in finite dimensions, similar results were known to Flåm [15]. In particular, Flåm seems to be the first author to identify the important role of the monotonicity

condition (1.2) in approximating Nash equilibrium. However, Rosen appears to be the first to consider using equations such as (1.4) to approximate equilibrium points [26]. In addition, we would like to point out that the finite dimensional variational inequality (1.3) and its relation to Nash equilibrium is studied in depth in the monograph by Facchinei and Pang [13]. In infinite dimensions, there have also been several recent works which discuss discrete time approximations for variational inequalities that correspond to Nash equilibrium [1,6,7,11,18].

Acknowledgements: This material is based upon work supported by the National Science Foundation under Grants DMS-1440140, DMS-1554130, and HRD-1700236, the National Security Agency under Grant No. H98230-20-1-0015, and the Sloan Foundation under Grant No. G-2020-12602 while the authors participated in a program hosted by the Mathematical Sciences Research Institute in Berkeley, California, during the summer of 2020.

2 Approximation theorem

In this section, we will briefly recall the notion of a maximal monotone operator on a Hilbert space and state a few key results for these operators. Then we will apply these results to prove Theorem 1.1 and a few related corollaries.

2.1 Maximal monotone operators on a Hilbert space

Let H be Hilbert space with inner product $\langle \cdot, \cdot \rangle$ and norm $|\cdot|$. We will denote 2^H as the power set or collection of all subsets of H . We recall that $B : H \rightarrow 2^H$ is monotone if

$$\langle p - q, x - y \rangle \geq 0$$

for all $x, y \in H$, $p \in Bx$, and $q \in By$. Moreover, B is *maximally monotone* if the only monotone $C : H \rightarrow 2^H$ such that $Bx \subset Cx$ for all $x \in H$ is $C = B$. Minty's well known maximality criterion is as follows [22].

Theorem (Minty's Lemma). *A monotone operator $B : H \rightarrow 2^H$ is maximal if and only if for each $y \in H$, there is a unique $x \in H$ such that*

$$x + Bx \ni y.$$

Let us now recall a fundamental theorem for the initial value problem

$$\begin{cases} \dot{u}(t) + Bu(t) \ni 0, \text{ a.e. } t \geq 0 \\ u(0) = u^0. \end{cases} \quad (2.1)$$

As discussed in Chapter II of [5] and Chapter 3 of [2], the initial value problem is well-posed provided that B is maximally monotone and u^0 is an element of the domain of B

$$D(B) = \{x \in H : Bx \neq \emptyset\}.$$

Theorem (Brézis's Theorem). *Suppose $B : H \rightarrow 2^H$ is maximally monotone and $u^0 \in D(B)$. Then there is a unique Lipschitz continuous*

$$u : [0, \infty) \rightarrow H$$

solution of (2.1) such that $u(t) \in D(B)$ for all $t \geq 0$. Moreover,

$$|\dot{u}(t)| \leq \min\{|p| : p \in Bu^0\} \quad (2.2)$$

for almost every $t \geq 0$.

Remark 2.1. The full statement of Brézis's Theorem (Theorem 3.1 of [2]) is more extensive than what is written above.

Remark 2.2. We note that the right hand side of (2.2) is finite as the images of maximal monotone operators are closed and convex subsets of H .

We also note that the operator B generates a contraction. Let u be a path as described in Brézis' Theorem, and suppose $v : [0, \infty) \rightarrow H$ is any other Lipschitz continuous path with $v(t) \in D(B)$ for $t \geq 0$ and

$$\dot{v}(t) + Bv(t) \ni 0, \text{ a.e. } t \geq 0.$$

Then

$$|u(t) - v(t)| \leq |u(0) - v(0)|, \quad t \geq 0. \quad (2.3)$$

If, in addition, there is some $\lambda > 0$ such that

$$\langle p - q, x - y \rangle \geq \lambda|x - y|^2 \quad (2.4)$$

for all $x, y \in H$, $p \in Bx$, and $q \in By$, then (2.3) can be improved to

$$|u(t) - v(t)| \leq e^{-\lambda t}|u(0) - v(0)|, \quad t \geq 0. \quad (2.5)$$

Remarkably, it is also possible to use solutions of the initial value problem (as described in Brézis's Theorem) to approximate equilibria of B . These are points $y \in H$ such that

$$By \ni 0.$$

The following theorem was proved in [3].

Theorem (The Baillon-Brézis Theorem). *Suppose B has an equilibrium point and u is a solution of the initial value problem (2.1). Then*

$$\frac{1}{t} \int_0^t u(s) ds$$

converges weakly in H to some equilibrium point of B .

Remark 2.3. If B satisfies (2.4), it is plain to see that $By \ni 0$ can have at most one solution. In this case,

$$|u(t) - y| \leq e^{-\lambda t}|u(0) - y|, \quad t \geq 0$$

for any solution of the initial value problem (2.1). This follows from (2.5) as $v(t) = y$ is a solution of the initial value problem with $v(0) = y$.

2.2 Proof of Theorem 1.1

Let us now suppose the spaces V , X_1, \dots, X_N , and H are as described in the subsection 1.2 of the introduction. We will further assume $f_1, \dots, f_N : X \rightarrow \mathbb{R}$ are given functions which satisfy (1.6). An elementary but important observation we will use is that a given $y \in H$ induces a linear form in V^* by the formula

$$(y, x) := \langle y, x \rangle, \quad (x \in V).$$

That this linear form is continuous is due to the continuity of the embedding $V \subset H$. For convenience, we will use $\|\cdot\|$ to denote both the norm on V and the associated norm on V^N . Namely, we'll write

$$\|x\| = \left(\sum_{j=1}^N \|x_j\|^2 \right)^{1/2}$$

for $x = (x_1, \dots, x_N) \in V^N$. We will also use this convention for the inner product $\langle \cdot, \cdot \rangle$ and norm $|\cdot|$ on H and on H^N and the induced norm $\|\cdot\|_*$ on V^* and on $(V^*)^N$.

Let us specify $A : H^N \rightarrow 2^{H^N}$ via

$$Ax = \begin{cases} (\partial_{x_1} f_1(x) + n_{X_1}(x_1)) \cap H \times \cdots \times (\partial_{x_N} f_N(x) + n_{X_N}(x_N)) \cap H, & x \in \mathcal{D} \\ \emptyset, & x \notin \mathcal{D} \end{cases}$$

for $x \in H^N$. Here \mathcal{D} is defined in (1.8).

Lemma 2.4. *The domain of A is \mathcal{D} , and A is monotone. Moreover, an absolutely continuous $u : [0, \infty) \rightarrow H^N$ solves*

$$\begin{cases} \dot{u}(t) + Au(t) \ni 0, & \text{a.e. } t \geq 0 \\ u(0) = u^0 \end{cases} \quad (2.6)$$

if and only it solves (1.7).

Proof. By definition, the domain of A is \mathcal{D} . Suppose $x, y \in \mathcal{D}$,

$$\zeta_j \in (\partial_{x_j} f_j(x) + n_{X_j}(x_j)) \cap H, \text{ and } \xi_j \in (\partial_{x_j} f_j(y) + n_{X_j}(y_j)) \cap H$$

for $j = 1, \dots, N$. In view of the last condition listed in (1.6) and the fact that each n_{X_j} is monotone,

$$\sum_{j=1}^N \langle \zeta_j - \xi_j, x_j - y_j \rangle = \sum_{j=1}^N (\zeta_j - \xi_j, x_j - y_j) \geq 0.$$

Thus, A is monotone.

Note that if u solves (2.6), then

$$-\dot{u}_j(t) \in (\partial_{x_j} f_j(u(t)) + n_{X_j}(u_j(t))) \cap H \subset \partial_{x_j} f_j(u(t)) + n_{X_j}(u_j(t))$$

for almost every $t \geq 0$ and each $j = 1, \dots, N$. It now follows from the definition of the normal cone that u is a solution of (1.7). Conversely, if u solves (1.7) then

$$-\dot{u}_j(t) \in \partial_{x_j} f_j(u(t)) + n_{X_j}(u_j(t))$$

for almost every $t \geq 0$ and each $j = 1, \dots, N$. As u is absolutely continuous, $-\dot{u}(t) \in H$ for almost every $t \geq 0$, and therefore, $-\dot{u}(t) \in Au(t)$ for almost every $t \geq 0$. \square

In view of Brézis' Theorem and the Baillon-Brézis Theorem, we can conclude Theorem 1.1 once we verify that A is maximal. To this end, we will verify the hypotheses of Minty's Lemma.

Proof of Theorem 1.1. For a given $y \in H^N$, it suffices to show there is $x \in X$ such that

$$\sum_{j=1}^N (x_j + \partial_{x_j} f_j(x) - y_j, z_j - x_j) \geq 0 \quad (2.7)$$

for each $z \in X$. In this case, $y_j - x_j \in H$ and

$$y_j - x_j \in \partial_{x_j} f_j(x) + n_{X_j}(x_j)$$

for $j = 1, \dots, N$ so that $x \in \mathcal{D}$ and $y - x \in Ax$.

In order to solve (2.7), we will employ the auxiliary functions defined by

$$g_j(x) := \frac{1}{2} \|x_j\|^2 - (y_j, x_j) + f_j(x)$$

for $x \in X$ and $j = 1, \dots, N$. In view of (1.6),

$$\left\{ \begin{array}{l} g_1, \dots, g_N \text{ are weakly lowersemicontinuous,} \\ X \ni y \mapsto \sum_{j=1}^N g_j(y_j, x_{-j}) \text{ is convex for each } x \in X, \\ X \ni x \mapsto \sum_{j=1}^N g_j(y_j, x_{-j}) \text{ is weakly continuous for each } y \in X, \text{ and} \\ \sum_{j=1}^N (\partial_{x_j} g_j(x) - \partial_{x_j} g_j(y), x_j - y_j) \geq \|x - y\|^2 \text{ for each } x, y \in X. \end{array} \right. \quad (2.8)$$

Note that $x \in X$ is a Nash equilibrium of g_1, \dots, g_N if and only if x solves (2.7). Since we are assuming that each X_j has nonempty interior, this follows from the Pshenichnii–Rockafellar conditions for the minimum of a proper, lowersemicontinuous, convex function on a closed

subset of a Banach space (Theorem 4.3.6 of [4]). Consequently, we now aim to show that g_1, \dots, g_N has a Nash equilibrium.

For each $r > 0$, set

$$X^r := \{x \in X : \|x\| \leq r\}.$$

It is clear that X^r is convex and weakly closed. And as each X_1, \dots, X_N is nonempty, X^r is nonempty for all r greater or equal to some fixed $s > 0$. Let us consider the map $\Phi^r : X^r \rightarrow 2^{X^r}$ specified as

$$\Phi^r(x) := \arg \min_{y \in X^r} \left\{ \sum_{j=1}^N g_j(y_j, x_{-j}) \right\}$$

($x \in X^r$) for $r > s$.

The first and second properties of g_1, \dots, g_N listed in (2.8) imply that $\Phi^r(x) \neq \emptyset$ and convex for any $x \in X^r$ and $r > s$; and the first and third properties imply that the graph of Φ^r is weakly closed. Indeed let us suppose $(x^k)_{k \in \mathbb{N}}, (y^k)_{k \in \mathbb{N}} \subset X$, $x^k \rightharpoonup x$, $y^k \rightharpoonup y$, and

$$\sum_{j=1}^N g_j(y_j^k, x_{-j}^k) \leq \sum_{j=1}^N g_j(z_j, x_{-j}^k)$$

for each $z \in X$ with $\|z\| \leq r$ and all $k \in \mathbb{N}$. We can then send $k \rightarrow \infty$ to get

$$\sum_{j=1}^N g_j(y_j, x_{-j}) \leq \sum_{j=1}^N g_j(z_j, x_{-j}).$$

The Kakutani-Glicksberg-Fan theorem [14, 16, 19] then implies there is $x^r \in \Phi^r(x^r)$. In particular,

$$\sum_{j=1}^N g_j(x^r) \leq \sum_{j=1}^N g_j(z_j, x_{-j}^r) \tag{2.9}$$

for each $z \in X$ with $\|z\| \leq r$. It follows that there are $p_j^r \in \partial_{x_j} g_j(x^r)$ for each $j = 1, \dots, N$ such that

$$\sum_{j=1}^N (p_j^r, z_j - x_j^r) \geq 0$$

for all $z \in X$ with $\|z\| \leq r$. Since $u^0 \in \mathcal{D}$, there are

$$w_j \in \partial_{x_j} g_j(u^0)$$

for $j = 1, \dots, N$.

Note that the fourth listed property of g_1, \dots, g_N in (2.8) gives

$$\|x^r - u^0\|^2 \leq \sum_{j=1}^N (p_j^r - w_j, x_j^r - u_j^0)$$

$$\begin{aligned}
&\leq \sum_{j=1}^N (-w_j, x_j^r - u_j^0) \\
&\leq \|w\|_* \|x^r - u^0\|
\end{aligned}$$

for all r sufficiently large. As a result, there is a sequence $(r_k)_{k \in \mathbb{N}}$ which increases to infinity such that $(x^{r_k})_{k \in \mathbb{N}} \subset X$ converges weakly to some $x \in X$. Upon sending $r = r_k \rightarrow \infty$ in (2.9), we find that x is the desired Nash equilibrium of g_1, \dots, g_N . \square

Corollary 2.5. *If X is weakly compact, then f_1, \dots, f_N has a Nash equilibrium.*

Proof. If X is weakly compact, we can repeat the argument given in the proof above used to show $g_1, \dots, g_N : X^r \rightarrow \mathbb{R}$ has a Nash equilibrium. All we would need to do is to replace X^r with X and g_1, \dots, g_N with f_1, \dots, f_N . \square

Corollary 2.6. *Suppose there is $\theta > 0$ such that*

$$\sum_{j=1}^N (\partial_{x_j} f_j(x) - \partial_{x_j} f_j(y), x_j - y_j) \geq \theta \|x - y\|^2$$

for each $x, y \in X$. Then f_1, \dots, f_N has a unique Nash equilibrium $z \in X$. Moreover, if $u : [0, \infty) \rightarrow H^N$ is a solution of the initial value problem as described in Theorem 1.1, there is $\lambda > 0$ such that

$$|u(t) - z| \leq e^{-\lambda t} |u(0) - z| \tag{2.10}$$

for each $t \geq 0$.

Proof. We can repeat the argument given in the proof of Theorem 1.1 that shows g_1, \dots, g_N has a Nash equilibrium to conclude that f_1, \dots, f_N has a Nash equilibrium.

If x is a Nash equilibrium of f_1, \dots, f_N , then

$$\sum_{j=1}^N (\partial_{x_j} f_j(x), z_j - x_j) \geq 0$$

for each $z \in X$. Likewise,

$$\sum_{j=1}^N (\partial_{x_j} f_j(y), w_j - y_j) \geq 0$$

for each $w \in X$ if y is another Nash equilibrium. Choosing $z = y$ and $w = x$ and adding these inequalities give

$$\theta \|x - y\|^2 \leq \sum_{j=1}^N (\partial_{x_j} f_j(x) - \partial_{x_j} f_j(y), x_j - y_j) \leq 0.$$

As $V^N \subset H^N$ is continuously embedded, there is a constant $C > 0$ such that

$$|x| \leq C \|x\|, \quad x \in V^N.$$

Setting

$$\lambda := \frac{\theta}{C^2},$$

gives

$$\langle Ax - Ay, x - y \rangle \geq \theta \|x - y\|^2 \geq \lambda |x - y|^2$$

for $x, y \in D(A)$. The inequality (2.10) now follows from Remark 2.3. \square

We now consider the special case mentioned in the introduction. In the statement below, we will suppose that V and H are as above. However, we will not assume f_1, \dots, f_N satisfy (1.6). We also note that the proof essentially follows from an observation made by Brézis in Remark 2.3.7 of [5].

Theorem 2.7. *Suppose $f_1, \dots, f_N : V^N \rightarrow \mathbb{R}$ satisfy*

$$\left\{ \begin{array}{l} \partial_{x_j} f_j(x) = \{\nabla_{x_j} f_j(x)\} \text{ for each } x \in V^N, \\ \nabla_{x_j} f_j(x + ty) \rightharpoonup \nabla_{x_j} f_j(x) \text{ as } t \rightarrow 0^+ \text{ for each } x, y \in V^N, \text{ and} \\ \sum_{j=1}^N (\nabla_{x_j} f_j(x) - \nabla_{x_j} f_j(y), x_j - y_j) \geq 0 \text{ for each } x, y \in V^N, \end{array} \right.$$

and define

$$\mathcal{D} = \{x \in V^N : \nabla_{x_1} f_1(x), \dots, \nabla_{x_N} f_N(x) \in H\}.$$

Then for each $u^0 \in \mathcal{D}$, there is a unique Lipschitz continuous $u : [0, \infty) \rightarrow H^N$ such that $u(t) \in \mathcal{D}$ for each $t \geq 0$ and

$$\left\{ \begin{array}{l} u(0) = u^0, \text{ and} \\ \dot{u}_j(t) + \nabla_{x_j} f_j(u(t)) = 0 \text{ for a.e. } t \geq 0 \text{ and each } j = 1, \dots, N. \end{array} \right.$$

Proof. It suffices to show that $A : H^N \rightarrow 2^{H^N}$ defined via

$$Ax := \begin{cases} \{(\nabla_{x_1} f_1(x), \dots, \nabla_{x_N} f_N(x))\}, & x \in \mathcal{D} \\ \emptyset, & x \notin \mathcal{D} \end{cases}$$

($x \in H^N$) is maximal. To this end, we first consider the operator $B : V^N \rightarrow (V^*)^N$ defined via

$$Bx := (x_1 + \nabla_{x_1} f_1(x), \dots, x_N + \nabla_{x_N} f_N(x))$$

for $x \in V^N$. Notice that

$$B(x + ty) \rightharpoonup Bx$$

and

$$(Bx - By, x - y) \geq \|x - y\|^2$$

for all $x, y \in V^N$.

By a theorem due independently to Browder [8] and Minty [23], B is surjective; see also Corollary 1.8 of [20]. In particular, if $y \in H^N$, there is $x \in V^N$ such that

$$Bx = y.$$

That is,

$$x_j + \nabla_{x_j} f_j(x) = y_j$$

for $j = 1, \dots, N$. Then $\nabla_{x_j} f_j(x) = y_j - x_j \in H$ for $j = 1, \dots, N$, which implies $x \in \mathcal{D}$ and

$$x + Ax \ni y.$$

□

Remark 2.8. If f_1, \dots, f_N additionally satisfy

$$\sum_{j=1}^N (\nabla_{x_j} f_j(x) - \nabla_{x_j} f_j(y), x_j - y_j) \geq \theta \|x - y\|^2$$

($x, y \in V^N$), then f_1, \dots, f_N has a unique Nash equilibrium and (2.10) holds for any $u(0) \in \mathcal{D}$.

3 Finite dimensional flows

In this section, we will study a few implications of Theorem 1.1 in the particular case

$$V = H = \mathbb{R}^d$$

equipped with the standard Euclidean dot product and norm. As there is just one topology to consider, the statements we'll make will be simpler than in the infinite dimensional setting. We will also identify a potential application to noncooperative games.

3.1 Compact domains

Let $X = X_1 \times \dots \times X_N$, where $X_1, \dots, X_N \subset \mathbb{R}^d$ are convex, compact subsets with nonempty interior. Further assume

$$\left\{ \begin{array}{l} f_1, \dots, f_N : X \rightarrow \mathbb{R} \text{ are continuous,} \\ X \ni y \mapsto \sum_{j=1}^N f_j(y_j, x_{-j}) \text{ is convex for each } x \in X, \text{ and} \\ \sum_{j=1}^N (\partial_{x_j} f_j(x) - \partial_{x_j} f_j(y)) \cdot (x_j - y_j) \geq 0 \text{ for each } x, y \in X. \end{array} \right.$$

As before we set

$$\mathcal{D} = \left\{ x \in X : \bigcap_{j=1}^N (\partial_{x_j} f_j(x) + n_{X_j}(x)) \neq \emptyset \right\},$$

where

$$n_{X_j}(z) = \{\zeta \in \mathbb{R}^d : \zeta \cdot (y - z) \leq 0 \text{ for all } y \in X_j\}$$

($j = 1, \dots, N$). Theorem 1.1 implies the following.

Proposition 3.1. *For each $u^0 \in \mathcal{D}$, there is a unique Lipschitz continuous $u : [0, \infty) \rightarrow \mathbb{R}^{Nd}$ with $u(t) \in \mathcal{D}$ for $t \geq 0$ and which satisfies*

$$\begin{cases} u(0) = u^0, \text{ and} \\ \sum_{j=1}^N (\dot{u}_j(t) + \partial_{x_j} f_j(u(t))) \cdot (y_j - u_j(t)) \geq 0 \text{ for a.e. } t \geq 0 \text{ and each } y \in X. \end{cases}$$

Furthermore, the limit

$$\lim_{t \rightarrow \infty} \frac{1}{t} \int_0^t u(s) ds$$

exists and equals a Nash equilibrium of f_1, \dots, f_N .

Remark 3.2. In view of Corollary 2.6, if

$$\sum_{j=1}^N (\partial_{x_j} f_j(x) - \partial_{x_j} f_j(y)) \cdot (x_j - y_j) \geq \theta |x - y|^2$$

for $x, y \in X$, f_1, \dots, f_N has a unique Nash equilibrium and the solution u in the proposition above converges exponentially fast to this equilibrium point.

3.2 An application to game theory

Again suppose $X_1, \dots, X_N \subset \mathbb{R}^d$ are convex, compact subsets with nonempty interior. Another interesting case to consider is when $f_1, \dots, f_N : X \rightarrow \mathbb{R}$ are each N -linear. That is,

$$X_j \ni y_j \mapsto f_j(y_j, x_{-j}) \text{ is linear} \quad (3.1)$$

for each $j = 1, \dots, N$ and $x \in X$. These are precisely the types of functions Nash considered in his celebrated work on noncooperative games [24, 25]. Note that (3.1) implies

$$y_j \cdot \nabla_{x_j} f_j(x) = f_j(y_j, x_{-j}) \quad (3.2)$$

for $y_j \in X_j$, $x \in \mathbb{R}^{Nd}$, and $j = 1, \dots, N$. Consequently,

$$\sum_{j=1}^N (\nabla_{x_j} f_j(x) - \nabla_{x_j} f_j(y)) \cdot (x_j - y_j) = \sum_{j=1}^N (f_j(x) - f_j(y_j, x_{-j}) + f_j(y) - f_j(x_j, y_{-j})) \geq 0$$

if and only if

$$\sum_{j=1}^N (f_j(x) + f_j(y)) \geq \sum_{j=1}^N (f_j(y_j, x_{-j}) + f_j(x_j, y_{-j})). \quad (3.3)$$

With these assumptions, Theorem 1.1 implies the following statement.

Proposition 3.3. *Suppose f_1, \dots, f_N are N -linear and satisfy (3.3). Then for each $u^0 \in X$, there is a unique Lipschitz $u : [0, \infty) \rightarrow X$ such that*

$$\begin{cases} u(0) = u^0 \\ \sum_{j=1}^N (\dot{u}_j(t) + \nabla_{x_j} f_j(u(t))) \cdot (y_j - u_j(t)) \geq 0, \text{ for a.e. } t \geq 0 \text{ and all } y \in X. \end{cases} \quad (3.4)$$

Moreover,

$$\frac{1}{t} \int_0^t u(s) ds$$

converges as $t \rightarrow \infty$ to a Nash equilibrium of f_1, \dots, f_N .

Remark 3.4. Using (3.2), we can rewrite the inequality in (3.4) as

$$\sum_{j=1}^N (\dot{u}_j(t) \cdot (y_j - u_j(t)) + f_j(y_j, u_{-j}(t)) - f_j(u(t))) \geq 0.$$

3.3 The whole space

Finally, we consider the case where f_1, \dots, f_N are defined on the whole space \mathbb{R}^{Nd} . We will assume

$$\begin{cases} f_1, \dots, f_N \text{ are continuous,} \\ \mathbb{R}^{Nd} \ni y \mapsto \sum_{j=1}^N f_j(y_j, x_{-j}) \text{ is convex for each } x \in \mathbb{R}^{Nd}, \text{ and} \\ \sum_{j=1}^N (\partial_{x_j} f_j(x) - \partial_{x_j} f_j(y)) \cdot (x_j - y_j) \geq 0 \text{ for each } x, y \in \mathbb{R}^{Nd}. \end{cases}$$

Let us also set

$$\mathcal{D} := \left\{ x \in \mathbb{R}^{Nd} : \bigcap_{j=1}^N \partial_{x_j} f_j(x) \neq \emptyset \right\}.$$

Theorem 1.1 gives us the following proposition.

Proposition 3.5. For each $u^0 \in \mathcal{D}$, there is a unique Lipschitz continuous $u : [0, \infty) \rightarrow \mathbb{R}^{Nd}$ such that $u(t) \in \mathcal{D}$ for each $t \geq 0$ and which satisfies

$$\begin{cases} u(0) = u^0 \\ \dot{u}_j(t) + \partial_{x_j} f_j(u(t)) \ni 0, \text{ for a.e. } t \geq 0 \text{ and } j = 1, \dots, N. \end{cases}$$

If f_1, \dots, f_N has a Nash equilibrium, then

$$\lim_{t \rightarrow \infty} \frac{1}{t} \int_0^t u(s) ds$$

exists and is also Nash equilibrium.

Remark 3.6. If

$$\sum_{j=1}^N (\partial_{x_j} f_j(x) - \partial_{x_j} f_j(y)) \cdot (x_j - y_j) \geq \theta |x - y|^2$$

for all $x, y \in \mathbb{R}^{Nd}$ and some $\theta > 0$, then f_1, \dots, f_N has a unique Nash equilibrium at some $z \in \mathbb{R}^{Nd}$ and

$$|u(t) - z| \leq e^{-\theta t} |u^0 - z|, \quad t \geq 0.$$

Remark 3.7. A basic example we had in mind when considering Proposition 3.5 was

$$f_j(x) = \sum_{k, \ell=1}^N \frac{1}{2} A_{k, \ell}^j x_k \cdot x_\ell$$

for $x = (x_1, \dots, x_N) \in \mathbb{R}^{Nd}$ and $j = 1, \dots, N$. Here $A_{k, \ell}^j$ are $d \times d$ symmetric matrices which additionally satisfy

$$A_{k, \ell}^j = A_{\ell, k}^j$$

($j, k, \ell = 1, \dots, N$). The required monotonicity condition on f_1, \dots, f_N is satisfied provided

$$\sum_{j, k=1}^N A_{k, j}^j y_j \cdot y_k \geq 0$$

for each $y \in \mathbb{R}^{Nd}$.

4 Examples in function spaces

In this final section, we will apply the ideas we have developed to consider functionals defined on Lebesgue and Sobolev spaces. As we have done above, we will consider both the existence of Nash equilibria and their approximation by continuous time flows. Throughout, we will assume $\Omega \subset \mathbb{R}^d$ is a bounded domain with smooth boundary. We will also change notation by using v for the variables in which our functionals are defined. This allows us to reserve x for points in Ω . Finally, we will use $|\cdot|$ for any norm on a finite dimensional space.

4.1 A few remarks on Sobolev spaces

Before discussing examples, let us recall a few facts about Sobolev spaces and establish some notation. A good reference for this material is Chapter 5 and 6 of [12]. First we note

$$H_0^1(\Omega) \subset L^2(\Omega) \subset H^{-1}(\Omega)$$

where each inclusion is compact. Here $L^2(\Omega)$ is the space of Lebesgue measurable functions on Ω which are square integrable equipped with the standard inner product. The space $H_0^1(\Omega)$ is the closure of all smooth functions having compact support in Ω in the norm

$$u \mapsto \left(\int_{\Omega} |Du|^2 dx \right)^{1/2}.$$

The topological dual of $H_0^1(\Omega)$ is denoted $H^{-1}(\Omega)$. Recall that the Dirichlet Laplacian $-\Delta : H_0^1(\Omega) \rightarrow H^{-1}(\Omega)$ is defined via

$$(-\Delta u, v) = \int_{\Omega} Du \cdot Dv dx$$

for $u, v \in H_0^1(\Omega)$. Here we are using (\cdot, \cdot) as the natural pairing between $H^{-1}(\Omega)$ and $H_0^1(\Omega)$. Moreover, $-\Delta$ is an isometry; in particular, this map is invertible. This allows us to define the following inner product on $H^{-1}(\Omega)$

$$(f, g) \mapsto (f, (-\Delta)^{-1}g).$$

We also note that if $f \in L^2(\Omega)$, the inner product between f and g simplifies to

$$(f, (-\Delta)^{-1}g) = \int_{\Omega} f(-\Delta)^{-1}g dx.$$

In addition, we will consider the space $H^2(\Omega) \cap H_0^1(\Omega)$ consisting of $H_0^1(\Omega)$ functions for which $-\Delta u \in L^2(\Omega)$. This is a Hilbert space endowed with the inner product

$$(u, v) \mapsto \int_{\Omega} \Delta u \Delta v dx.$$

4.2 $H = L^2(\Omega)$

For $v \in H_0^1(\Omega)^N$, define

$$f_j(v) = \int_{\Omega} H_j(\nabla v_j) + F_j(v) - h_j v_j dx$$

for $j = 1, \dots, N$. Here each $h_j \in L^2(\Omega)$, and $H_j : \mathbb{R}^d \rightarrow \mathbb{R}$ is assumed to be smooth and satisfy

$$\theta |p - q|^2 \leq (\nabla H_j(p) - \nabla H_j(q)) \cdot (p - q) \leq \frac{1}{\theta} |p - q|^2 \quad (4.1)$$

for all $p, q \in \mathbb{R}^d$ and some $\theta \in (0, 1]$. We'll also suppose $F_1, \dots, F_N : \mathbb{R}^N \rightarrow \mathbb{R}$ are smooth,

$$\sum_{j=1}^N (\partial_{z_j} F_j(z) - \partial_{z_j} F_j(w))(z_j - w_j) \geq 0$$

for $z, w \in \mathbb{R}^N$, and there is C such that

$$|F_j(z)| \leq C(1 + |z|^2) \text{ and } |\partial_{z_j} F_j(z)| \leq C(1 + |z|) \quad (4.2)$$

for $z \in \mathbb{R}^N$ and $j = 1, \dots, N$.

Using these assumptions, it is straightforward to verify that f_1, \dots, f_N fulfills (1.6) with $X_1 = \dots = X_N = V = H_0^1(\Omega)$. The following lemma also follows by routine computations.

Lemma 4.1. *For each $v, w \in H_0^1(\Omega)^N$ and $j = 1, \dots, N$,*

$$\partial_{v_j} f_j(v) = \{\nabla_{v_j} f_j(v)\}$$

with

$$(\nabla_{v_j} f_j(v), w_j) = \int_{\Omega} \nabla H_j(\nabla v_j) \cdot \nabla w_j + \partial_{z_j} F_j(v) w_j - h_j w_j dx$$

and

$$\sum_{j=1}^N (\nabla_{v_j} f_j(v) - \nabla_{v_j} f_j(w), v_j - w_j) \geq \theta \int_{\Omega} |\nabla v - \nabla w|^2 dx.$$

Moreover, v is a Nash equilibrium of f_1, \dots, f_N if and only if it is a weak solution of

$$\begin{cases} -\operatorname{div}(\nabla H_j(\nabla v_j)) + \partial_{z_j} F_j(v) = h_j & \text{in } \Omega \\ v_j = 0 & \text{on } \partial\Omega \end{cases}$$

($j = 1, \dots, N$).

Remark 4.2. In this statement, (\cdot, \cdot) denotes the natural pairing between $H^{-1}(\Omega)$ and $H_0^1(\Omega)$.

We now can state a result involving the approximation of Nash equilibrium for f_1, \dots, f_N .

Proposition 4.3. *Let $u^0 \in (H^2(\Omega) \cap H_0^1(\Omega))^N$. There is a unique Lipschitz continuous*

$$u : [0, \infty) \rightarrow L^2(\Omega)^N; t \mapsto u(\cdot, t)$$

such that $u(\cdot, t) \in (H^2(\Omega) \cap H_0^1(\Omega))^N$ for each $t \geq 0$ and

$$\begin{cases} \partial_t u_j = \operatorname{div}(\nabla H_j(\nabla u_j)) - \partial_{z_j} F_j(u) + h_j & \text{in } \Omega \times (0, \infty) \\ u_j = 0 & \text{on } \partial\Omega \times (0, \infty) \\ u_j = u_j^0 & \text{on } \Omega \times \{0\} \end{cases} \quad (4.3)$$

($j = 1, \dots, N$). Furthermore, there is $\lambda > 0$ such that

$$\int_{\Omega} |u(x, t) - v(x)|^2 dx \leq e^{-2\lambda t} \int_{\Omega} |u^0(x) - v(x)|^2 dx$$

for $t \geq 0$. Here v is the unique Nash equilibrium of f_1, \dots, f_N .

Remark 4.4. For almost every $t \geq 0$, the PDE in (4.3) holds almost everywhere in Ω ; the boundary condition holds in the trace sense; and $u(\cdot, 0) = u^0$ as $L^2(\Omega)^N$ functions.

Proof. Suppose that $\zeta \in L^2(\Omega)^N$ and $v \in H_0^1(\Omega)^N$ is a weak solution of

$$\begin{cases} -\operatorname{div}(\nabla H_j(\nabla v_j)) + \partial_{z_j} F_j(v) = h_j + \zeta_j & \text{in } \Omega \\ v_j = 0 & \text{on } \partial\Omega. \end{cases}$$

As $\partial\Omega$ is assumed to be smooth and in view of the hypothesis (4.1), elliptic regularity (Theorem 1, section 8.3 of [12]) implies $v \in (H^2(\Omega) \cap H_0^1(\Omega))^N$. Therefore,

$$\{v \in H_0^1(\Omega)^N : \nabla_{v_1} f_1(v), \dots, \nabla_{v_N} f_N(v) \in L^2(\Omega)\} = (H^2(\Omega) \cap H_0^1(\Omega))^N.$$

Theorem 1.1 and Corollary 2.6 now allow us to conclude. □

4.3 Another example with $H = L^2(\Omega)$

Define

$$f_j(v) = \int_{\Omega} L_j(\nabla v_1, \dots, \nabla v_N) - h_j v_j \, dx$$

for $v \in H_0^1(\Omega)^N$. Here $h_1, \dots, h_N \in L^2(\Omega)$ and each $L_j : \mathbb{R}^{Nd} \rightarrow \mathbb{R}$ is smooth with

$$\theta |p - q|^2 \leq \sum_{j=1}^N (\nabla_{p_j} L_j(p) - \nabla_{p_j} L_j(q)) \cdot (p_j - q_j) \leq \frac{1}{\theta} |p - q|^2 \quad (4.4)$$

for all $p, q \in \mathbb{R}^{Nd}$ and some $\theta \in (0, 1]$. We'll also assume there is $C > 0$ such that

$$|\nabla_{p_j} L_j(p)| \leq C(1 + |p|)$$

for each $p \in \mathbb{R}^{Nd}$ and $j = 1, \dots, N$.

Direct computation gives the following lemma.

Lemma 4.5. *For each $v, w \in H_0^1(\Omega)^N$ and $j = 1, \dots, N$,*

$$\partial_{v_j} f_j(v) = \{\nabla_{v_j} f_j(v)\}$$

with

$$(\nabla_{v_j} f_j(v), w_j) = \int_{\Omega} \nabla_{p_j} L_j(\nabla v_1, \dots, \nabla v_N) \cdot \nabla w_j - h_j w_j \, dx$$

and

$$\sum_{j=1}^N (\nabla_{v_j} f_j(v) - \nabla_{v_j} f_j(w), v_j - w_j) \geq \theta \int_{\Omega} |\nabla v - \nabla w|^2 \, dx.$$

Moreover,

$$\nabla_{v_j} f_j(v + tw) \rightarrow \nabla_{v_j} f_j(v)$$

as $t \rightarrow 0^+$, and v is a Nash equilibrium of f_1, \dots, f_N if and only if it is a weak solution of

$$\begin{cases} -\operatorname{div}(\nabla_{p_j} L_j(\nabla v_1, \dots, \nabla v_N)) = h_j & \text{in } \Omega \\ v_j = 0 & \text{on } \partial\Omega \end{cases}$$

($j = 1, \dots, N$).

With this lemma, we can now establish the subsequent assertion about the Nash equilibrium of f_1, \dots, f_N .

Proposition 4.6. *Let $u^0 \in (H^2(\Omega) \cap H_0^1(\Omega))^N$. There is a unique Lipschitz continuous*

$$u : [0, \infty) \rightarrow L^2(\Omega)^N; t \mapsto u(\cdot, t)$$

such that $u(\cdot, t) \in (H^2(\Omega) \cap H_0^1(\Omega))^N$ for each $t \geq 0$ and

$$\begin{cases} \partial_t u_j = \operatorname{div}(\nabla_{p_j} L_j(\nabla u_1, \dots, \nabla u_N)) + h_j & \text{in } \Omega \times (0, \infty) \\ u_j = 0 & \text{on } \partial\Omega \times (0, \infty) \\ u_j = u_j^0 & \text{on } \Omega \times \{0\} \end{cases}$$

($j = 1, \dots, N$). Furthermore, there is $\lambda > 0$ such that

$$\int_{\Omega} |u(x, t) - v(x)|^2 dx \leq e^{-2\lambda t} \int_{\Omega} |u^0(x) - v(x)|^2 dx$$

for $t \geq 0$. Here v is the unique Nash equilibrium of f_1, \dots, f_N .

Proof. Suppose that $\xi \in L^2(\Omega)^N$ and $v \in H_0^1(\Omega)^N$ is a weak solution of

$$\begin{cases} -\operatorname{div}(\nabla_{p_j} L_j(\nabla v_1, \dots, \nabla v_N)) = h_j + \xi_j & \text{in } \Omega \\ v_j = 0 & \text{on } \partial\Omega. \end{cases}$$

By the uniform estimate (4.4), elliptic regularity implies $v \in (H^2(\Omega) \cap H_0^1(\Omega))^N$ (see the remark at the end of section 9.1 in [12]). As a result,

$$\{v \in H_0^1(\Omega)^N : \nabla_{v_1} f_1(v), \dots, \nabla_{v_N} f_N(v) \in L^2(\Omega)\} = (H^2(\Omega) \cap H_0^1(\Omega))^N.$$

We then conclude by Theorem 2.7 and Remark 2.8. □

4.4 $H = H^{-1}(\Omega)$

For a given $v \in L^2(\Omega)^N$, set

$$f_j(v) = \int_{\Omega} F_j(v) dx - \langle h_j, v_j \rangle$$

for $j = 1, \dots, N$. Here $h_1, \dots, h_N \in H^{-1}(\Omega)$, and $\langle \cdot, \cdot \rangle$ denotes the inner product in $H^{-1}(\Omega)$. We will assume $F_1, \dots, F_N : \mathbb{R}^N \rightarrow \mathbb{R}$ are smooth and satisfy (4.2) and

$$\begin{cases} |\partial_{z_k} \partial_{z_j} F_j(z)| \leq C \\ \theta |z - w|^2 \leq \sum_{i=1}^N (\partial_{z_i} F_i(z) - \partial_{z_i} F_i(w))(z_i - w_i) \end{cases} \quad (4.5)$$

for $z, w \in \mathbb{R}^N$, $j, k = 1, \dots, N$, and some $C, \theta > 0$. For convenience, we will also assume

$$\partial_{z_j} F_j(0) = 0 \quad (4.6)$$

for $j = 1, \dots, N$.

The following observations can be verified by routine computations.

Lemma 4.7. *For each $v, w \in L^2(\Omega)^N$ and $j = 1, \dots, N$,*

$$\partial_{v_j} f_j(v) = \{\nabla_{v_j} f_j(v)\}$$

with

$$\nabla_{v_j} f_j(v) = \partial_{z_j} F_j(v) - (-\Delta)^{-1} h_j.$$

Moreover, $v \in H_0^1(\Omega)^N$ is a Nash equilibrium of f_1, \dots, f_N if and only if it is a weak solution of

$$\begin{cases} -\Delta(\partial_{z_j} F_j(v)) = h_j & \text{in } \Omega \\ v_j = 0 & \text{on } \partial\Omega \end{cases} \quad (4.7)$$

($j = 1, \dots, N$).

By Proposition A.1 in the appendix, there are smooth functions G_1, \dots, G_N such that

$$\begin{cases} \partial_{z_j} F_j(z) = y_j \text{ if and only if } \partial_{y_j} G_j(y) = z_j \\ |G_j(y)| \leq B(1 + |y|^2) \text{ and } |\partial_{y_j} G_j(y)| \leq B(1 + |y|) \\ 0 \leq \sum_{i=1}^N (\partial_{y_i} G_i(y) - \partial_{y_i} G_i(z)) (y_i - z_i) \\ |\partial_{y_k} \partial_{y_j} G_j(y)| \leq \frac{1}{\theta} \end{cases} \quad (4.8)$$

for all $y, z \in \mathbb{R}^N$, $j, k = 1, \dots, N$ and some constant B . Notice that in view of (4.6)

$$\partial_{y_j} G_j(0) = 0$$

for $j = 1, \dots, N$. Thus, it is possible to solve (4.7) by choosing

$$w_j = (-\Delta)^{-1}h_j \in H_0^1(\Omega)$$

for $j = 1, \dots, N$ and then setting

$$v_j = \partial_{y_j} G_j(w)$$

for $j = 1, \dots, N$. In particular, we have just shown that f_1, \dots, f_N has a Nash equilibrium, and it is not hard to see it is unique.

Let us now see how to approximate this Nash equilibrium.

Proposition 4.8. *Let $u^0 \in H_0^1(\Omega)^N$. There is a unique Lipschitz continuous*

$$u : [0, \infty) \rightarrow H^{-1}(\Omega)^N; t \mapsto u(\cdot, t)$$

such that $u(\cdot, t) \in H_0^1(\Omega)^N$ for each $t \geq 0$ and

$$\begin{cases} \partial_t u_j = \Delta(\partial_{z_j} F_j(u)) - h_j & \text{in } \Omega \times (0, \infty) \\ u_j = 0 & \text{on } \partial\Omega \times (0, \infty) \\ u_j = u_j^0 & \text{on } \Omega \times \{0\} \end{cases}$$

($j = 1, \dots, N$). Furthermore, there is $\lambda > 0$ such that

$$\|u(\cdot, t) - v\| \leq e^{-\lambda t} \|u^0 - v\|$$

for $t \geq 0$. Here $\|\cdot\|$ denotes the norm on $H^{-1}(\Omega)^N$ and v is the Nash equilibrium of f_1, \dots, f_N .

Proof. Define $A : H^{-1}(\Omega)^N \rightarrow 2^{H^{-1}(\Omega)^N}$ as

$$Av = \begin{cases} \{(-\Delta(\partial_{z_1} F_1(v)) + h_1, \dots, -\Delta(\partial_{z_N} F_N(v)) + h_N)\}, & v \in H_0^1(\Omega)^N \\ \emptyset, & \text{otherwise} \end{cases}$$

for $v \in H^{-1}(\Omega)^N$. We note that $D(A) = H_0^1(\Omega)^N$, and $0 \in Av$ if and only if $v \in H_0^1(\Omega)^N$ is a Nash equilibrium for f_1, \dots, f_N . By the second inequality listed in (4.5),

$$\begin{aligned} \langle Av - Aw, v - w \rangle &= \int_{\Omega} \sum_{j=1}^N (\partial_{z_j} F_j(v) - \partial_{z_j} F_j(w))(v_j - w_j) dx \\ &\geq \theta \int_{\Omega} |v - w|^2 dx \\ &\geq \lambda \|v - w\|^2 \end{aligned}$$

for $v, w \in H_0^1(\Omega)^N$ and some $\lambda > 0$.

As a result, it suffices to check that A is maximal. This amounts to finding a weak solution $v \in H_0^1(\Omega)^N$ of

$$\begin{cases} v_j - \Delta(\partial_{z_j} F_j(v)) = h_j + \zeta_j & \text{in } \Omega \\ v_j = 0 & \text{on } \partial\Omega \end{cases} \quad (4.9)$$

for a given $\zeta \in H^{-1}(\Omega)^N$. To this end, we consider the auxiliary problem of finding $w \in H_0^1(\Omega)^N$ such that

$$\begin{cases} \partial_{y_j} G_j(w) - \Delta w_j = h_j + \zeta_j & \text{in } \Omega \\ w_j = 0 & \text{on } \partial\Omega. \end{cases} \quad (4.10)$$

If we can find w , then $v \in H_0^1(\Omega)^N$ defined by

$$v_j = \partial_{y_j} G_j(w)$$

($j = 1, \dots, N$) is a solution of (4.9).

Notice that (4.10) has a solution if and only if

$$g_j(w) = \int_{\Omega} G_j(w) + \frac{1}{2} |\nabla w_j|^2 dx - (h_j + \zeta_j, w_j) \quad (4.11)$$

($w \in H_0^1(\Omega)^N$, $j = 1, \dots, N$) has a Nash equilibrium. Here (\cdot, \cdot) is the natural pairing between $H^{-1}(\Omega)$ and $H_0^1(\Omega)$. Using (4.8), it is straightforward to check that g_1, \dots, g_N satisfies the properties listed in (1.6) with $X_1 = \dots = X_N = V = H_0^1(\Omega)$. As a result, Corollary 2.6 implies that g_1, \dots, g_N has a Nash equilibrium. \square

4.5 $H = H_0^1(\Omega)$

Set

$$f_j(v) = \int_{\Omega} F_j(\Delta v) - \nabla h_j \cdot \nabla v_j dx$$

for $v \in (H^2(\Omega) \cap H_0^1(\Omega))^N$ and $j = 1, \dots, N$. Here $h_1, \dots, h_N \in H_0^1(\Omega)$ and

$$\Delta v = (\Delta v_1, \dots, \Delta v_N).$$

We will suppose $F_1, \dots, F_N : \mathbb{R}^N \rightarrow \mathbb{R}$ are smooth and satisfy (4.5) and (4.6). A few key observations regarding f_1, \dots, f_N are as follows.

Lemma 4.9. *For each $v, w \in (H^2(\Omega) \cap H_0^1(\Omega))^N$ and $j = 1, \dots, N$,*

$$\partial_{v_j} f_j(v) = \{\nabla_{v_j} f_j(v)\}$$

with

$$(\nabla_{v_j} f_j(v), w_j) = \int_{\Omega} \partial_{z_j} F_j(\Delta v) \Delta w_j - \nabla h_j \cdot \nabla w_j dx.$$

Moreover, v is a Nash equilibrium of f_1, \dots, f_N if and only if it is a solution of

$$\begin{cases} -\partial_{z_j} F_j(\Delta v) = h_j & \text{in } \Omega \\ v_j = 0 & \text{on } \partial\Omega \end{cases} \quad (4.12)$$

($j = 1, \dots, N$).

Remark 4.10. Above, (\cdot, \cdot) denotes the natural pairing between $(H^2(\Omega) \cap H_0^1(\Omega))^*$ and $H^2(\Omega) \cap H_0^1(\Omega)$.

Using the functions G_1, \dots, G_N from the previous subsection, we can solve

$$\begin{cases} -\Delta v_j = -\partial_{y_j} G_j(-h) & \text{in } \Omega \\ v_j = 0 & \text{on } \partial\Omega, \end{cases}$$

for $j = 1, \dots, N$ to obtain a solution of (4.12). It follows that f_1, \dots, f_N has a unique Nash equilibrium v , which belongs to the space

$$\mathcal{D} = \{v \in (H^2(\Omega) \cap H_0^1(\Omega))^N : \Delta v \in H_0^1(\Omega)^N\}.$$

As with our previous examples, we can approximate this Nash equilibrium with a continuous time flow.

Proposition 4.11. *Let $u^0 \in \mathcal{D}$. There is a unique Lipschitz continuous*

$$u : [0, \infty) \rightarrow H_0^1(\Omega)^N; t \mapsto u(\cdot, t)$$

such that $u(\cdot, t) \in \mathcal{D}$ for each $t \geq 0$ and

$$\begin{cases} \partial_t u_j = \partial_{z_j} F_j(\Delta u) + h_j & \text{in } \Omega \times (0, \infty) \\ u_j = 0 & \text{on } \partial\Omega \times (0, \infty) \\ u_j = u_j^0 & \text{on } \Omega \times \{0\} \end{cases}$$

($j = 1, \dots, N$). Furthermore, there is $\lambda > 0$ such that

$$\int_{\Omega} |\nabla u(x, t) - \nabla v(x)|^2 dx \leq e^{-2\lambda t} \int_{\Omega} |\nabla u^0(x) - \nabla v(x)|^2 dx$$

for $t \geq 0$. Here v is the unique Nash equilibrium of f_1, \dots, f_N .

Proof. Define $A : H_0^1(\Omega)^N \rightarrow 2^{H_0^1(\Omega)^N}$ by

$$Av = \begin{cases} \{(-\partial_{z_1} F_1(\Delta v) - h_1, \dots, -\partial_{z_N} F_N(\Delta v) - h_N)\}, & v \in \mathcal{D} \\ \emptyset, & \text{otherwise} \end{cases}$$

for $v \in H_0^1(\Omega)^N$. Note that $D(A) = \mathcal{D}$, and $0 \in Av$ if and only if $v \in \mathcal{D}$ is a Nash equilibrium for f_1, \dots, f_N . In view of (4.5),

$$\begin{aligned} \langle Av - Aw, v - w \rangle &= \int_{\Omega} \sum_{j=1}^N (\partial_{z_j} F_j(\Delta v) - \partial_{z_j} F_j(\Delta w)) (\Delta v_j - \Delta w_j) dx \\ &\geq \theta \int_{\Omega} |\Delta v - \Delta w|^2 dx \\ &\geq \lambda \int_{\Omega} |\nabla v - \nabla w|^2 dx. \end{aligned}$$

for $v, w \in \mathcal{D}$ and some $\lambda > 0$.

Therefore, it suffices to show

$$\begin{cases} v_j - \partial_{z_j} F_j(\Delta v) = h_j + \zeta_j & \text{in } \Omega \\ v_j = 0 & \text{on } \partial\Omega \end{cases}$$

has a solution $v \in (H^2(\Omega) \cap H_0^1(\Omega))^N$ for a given $\zeta \in H_0^1(\Omega)^N$. We note that any such solution v will automatically belong to \mathcal{D} . Moreover, this is equivalent to finding a weak solution $v \in H_0^1(\Omega)^N$ of

$$\begin{cases} \partial_{y_j} G_j(v - h - \zeta) - \Delta v_j = 0 & \text{in } \Omega \\ v_j = 0 & \text{on } \partial\Omega. \end{cases} \quad (4.13)$$

We emphasize that elliptic regularity implies that any such weak solution actually belongs to $(H^2(\Omega) \cap H_0^1(\Omega))^N$.

Any $v \in H_0^1(\Omega)^N$ which solves (4.13) is a Nash equilibrium of g_1, \dots, g_N defined by

$$g_j(v) = \int_{\Omega} G_j(v - h - \zeta) + \frac{1}{2} |\nabla v_j|^2 dx.$$

As we did when considering the functions defined in (4.11), we can apply Corollary 2.6 and conclude that g_1, \dots, g_N has a unique Nash equilibrium. \square

A Dual functions

In this appendix, we will verify a technical assertion that was used to establish the existence of a Nash equilibrium in a few of the examples we considered above.

Proposition A.1. *Suppose $F_1, \dots, F_N : \mathbb{R}^N \rightarrow \mathbb{R}$ are smooth and satisfy*

$$|F_j(z)| \leq C(1 + |z|^2) \quad (A.1)$$

for $z \in \mathbb{R}^N$ and $j = 1, \dots, N$ and

$$\theta |z - w|^2 \leq \sum_{j=1}^N (\partial_{z_j} F_j(z) - \partial_{z_j} F_j(w)) (z_j - w_j) \quad (A.2)$$

for $z, w \in \mathbb{R}^N$ and some $\theta > 0$. There are smooth functions $G_1, \dots, G_N : \mathbb{R}^N \rightarrow \mathbb{R}$ with the following properties.

(i) $\partial_{z_j} F_j(z) = y_j$ if and only if $\partial_{y_j} G_j(y) = z_j$.

(ii) For all $j, k = 1, \dots, N$ and $y \in \mathbb{R}^N$,

$$|\partial_{y_k} \partial_{y_j} G_j(y)| \leq \frac{1}{\theta}.$$

(iii) For each $y, w \in \mathbb{R}^N$,

$$0 \leq \sum_{j=1}^N (\partial_{y_j} G_j(y) - \partial_{y_j} G_j(w)) (y_j - w_j).$$

(iv) There is a constant B such that

$$|G_j(y)| \leq B(1 + |y|^2) \text{ and } |\partial_{y_j} G_j(y)| \leq B(1 + |y|)$$

for $y \in \mathbb{R}^N$ and $j = 1, \dots, N$.

Proof. First we note that the mapping of \mathbb{R}^N

$$z \mapsto (\partial_{z_1} F_1(z), \dots, \partial_{z_N} F_N(z)) \tag{A.3}$$

is invertible. This follows from the theorem due to Browder [8] and Minty [23] as (A.2) implies that this map is both monotone and coercive; invertibility is a consequence of Hadamard's global invertibility theorem (Theorem A of [17]), as well. The condition (A.2) also gives

$$\theta |v|^2 \leq \sum_{j,k=1}^N \partial_{z_k} \partial_{z_j} F_j(z) v_k v_j \tag{A.4}$$

for each $z, v \in \mathbb{R}^N$.

Now fix $y \in \mathbb{R}^N$ and let $z \in \mathbb{R}^N$ be the unique Nash equilibrium of

$$\mathbb{R}^N \ni x \mapsto F_j(x) - y_j x_j$$

($j = 1, \dots, N$). Such a Nash equilibrium exists for these N functions by Corollary 2.6. As a result,

$$\partial_{z_j} F_j(z) = y_j, \quad j = 1, \dots, N. \tag{A.5}$$

Since (A.3) is invertible and smooth, (A.5) determines z as a smooth function of y . Let us now define

$$G_j(y) := z_j y_j - F_j(z) \tag{A.6}$$

for $j = 1, \dots, N$. Differentiating with respect to y_j and using (A.5), we find

$$\partial_{y_j} G_j(y) = z_j.$$

This proves (i) and that

$$y \mapsto (\partial_{y_1} G_1(y), \dots, \partial_{y_N} G_N(y))$$

is the inverse mapping of (A.3).

It follows from part (i) and the inverse function theorem that whenever $\partial_{y_j} G_j(y) = z_j$ and $\partial_{z_j} F_j(z) = y_j$ for $j = 1, \dots, N$, the matrices

$$(\partial_{z_k} \partial_{z_j} F_j(z))_{j,k=1}^N \text{ and } (\partial_{y_k} \partial_{y_j} G_j(y))_{j,k=1}^N$$

are inverses. Consequently, for a given $w \in \mathbb{R}^N$, we can choose $v \in \mathbb{R}^N$ defined as

$$v_j = \sum_{k=1}^N \partial_{y_k} \partial_{y_j} G_j(y) w_k$$

in (A.4) to get

$$\begin{aligned} 0 &\leq \theta \sum_{j=1}^N \left(\sum_{k=1}^N \partial_{y_k} \partial_{y_j} G_j(y) w_k \right)^2 \\ &\leq \sum_{j,k=1}^N \partial_{y_k} \partial_{y_j} G_j(y) w_k w_j \\ &= \sum_{j=1}^N \left(\sum_{k=1}^N \partial_{y_k} \partial_{y_j} G_j(y) w_k \right) w_j \\ &\leq \sqrt{\sum_{j=1}^N \left(\sum_{k=1}^N \partial_{y_k} \partial_{y_j} G_j(y) w_k \right)^2} \cdot |w|. \end{aligned} \tag{A.7}$$

As a result,

$$\sum_{j=1}^N \left(\sum_{k=1}^N \partial_{y_k} \partial_{y_j} G_j(y) w_k \right)^2 \leq \frac{1}{\theta^2} |w|^2.$$

Upon selecting $w = e_k$, the k th standard unit vector in \mathbb{R}^N , we find

$$(\partial_{y_k} \partial_{y_j} G_j(y))^2 = \left(\sum_{\ell=1}^N \partial_{y_\ell} \partial_{y_j} G_j(y) w_\ell \right)^2 \leq \sum_{i=1}^N \left(\sum_{\ell=1}^N \partial_{y_\ell} \partial_{y_i} G_i(y) w_\ell \right)^2 \leq \frac{1}{\theta^2}$$

for $j, k = 1, \dots, N$. We conclude assertion (ii).

Part (iii) follows directly from (A.7). Indeed, for any $y, w \in \mathbb{R}^N$, we can use the fundamental theorem of calculus to derive

$$\begin{aligned} & \sum_{j=1}^N (\partial_{y_j} G_j(y) - \partial_{y_j} G_j(w)) (y_j - w_j) \\ &= \int_0^1 \left(\sum_{j,k=1}^N \partial_{y_k} \partial_{y_j} G_j(w + t(y-w))(y_k - w_k)(y_j - w_j) \right) dt \\ &\geq 0. \end{aligned}$$

We are left to verify assertion (iv). First note that by part (ii),

$$\begin{aligned} |\partial_{y_j} G_j(y)| &\leq |\partial_{y_j} G_j(y) - \partial_{y_j} G_j(0)| + |\partial_{y_j} G_j(0)| \\ &= \left| \int_0^1 \sum_{k=1}^N \partial_{y_k} \partial_{y_j} G_j(ty) y_k dt \right| + |\partial_{y_j} G_j(0)| \\ &\leq \sum_{k=1}^N \left(\int_0^1 |\partial_{y_k} \partial_{y_j} G_j(ty)| dt \right) |y_k| + |\partial_{y_j} G_j(0)| \\ &\leq \frac{1}{\theta} \sum_{k=1}^N |y_k| + |\partial_{y_j} G_j(0)| \\ &\leq D(1 + |y|) \end{aligned}$$

for some $D > 0$ independent of $j = 1, \dots, N$ and y .

For a given $y \in \mathbb{R}^N$ there is a unique z such that $\partial_{z_j} F_j(z) = y_j$ and $\partial_{y_j} G_j(y) = z_j$ for $j = 1, \dots, N$. We just showed above that

$$|z_j| = |\partial_{y_j} G_j(y)| \leq D(1 + |y|).$$

In view of hypothesis, (A.1) and our definition of G_j (A.6), we additionally have

$$\begin{aligned} |G_j(y)| &= |z_j y_j - F_j(z)| \\ &\leq |z_j| |y_j| + |F_j(z)| \\ &\leq D(1 + |y|) |y| + C(1 + |z|^2) \\ &\leq D(1 + |y|) |y| + C(1 + ND^2(1 + |y|)^2) \\ &\leq B(1 + |y|^2) \end{aligned}$$

for an appropriately chosen constant B . □

References

- [1] Abdullah Alotaibi, Patrick L. Combettes, and Naseer Shahzad. Solving coupled composite monotone inclusions by successive Fejér approximations of their Kuhn-Tucker set. *SIAM J. Optim.*, 24(4):2076–2095, 2014.

- [2] Jean-Pierre Aubin and Arrigo Cellina. *Differential inclusions*, volume 264 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1984. Set-valued maps and viability theory.
- [3] J. B. Baillon and H. Brézis. Une remarque sur le comportement asymptotique des semigroupes non linéaires. *Houston J. Math.*, 2(1):5–7, 1976.
- [4] Jonathan M. Borwein and Qiji J. Zhu. *Techniques of variational analysis*, volume 20 of *CMS Books in Mathematics/Ouvrages de Mathématiques de la SMC*. Springer-Verlag, New York, 2005.
- [5] H. Brézis. *Opérateurs maximaux monotones et semi-groupes de contractions dans les espaces de Hilbert*. North-Holland Publishing Co., Amsterdam-London; American Elsevier Publishing Co., Inc., New York, 1973. North-Holland Mathematics Studies, No. 5. Notas de Matemática (50).
- [6] Luis M. Briceño Arias. A Douglas-Rachford splitting method for solving equilibrium problems. *Nonlinear Anal.*, 75(16):6053–6059, 2012.
- [7] Luis M. Briceño Arias and Patrick L. Combettes. Monotone operator methods for Nash equilibria in non-potential games. In *Computational and analytical mathematics*, volume 50 of *Springer Proc. Math. Stat.*, pages 143–159. Springer, New York, 2013.
- [8] Felix E. Browder. Nonlinear elliptic boundary value problems. *Bull. Amer. Math. Soc.*, 69:862–874, 1963.
- [9] Constantinos Daskalakis, Paul W. Goldberg, and Christos H. Papadimitriou. The complexity of computing a Nash equilibrium. *SIAM J. Comput.*, 39(1):195–259, 2009.
- [10] Constantinos Daskalakis, Aranyak Mehta, and Christos Papadimitriou. A note on approximate Nash equilibria. *Theoret. Comput. Sci.*, 410(17):1581–1588, 2009.
- [11] Jonathan Eckstein and B. F. Svaiter. General projective splitting methods for sums of maximal monotone operators. *SIAM J. Control Optim.*, 48(2):787–811, 2009.
- [12] Lawrence C. Evans. *Partial differential equations*, volume 19 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, second edition, 2010.
- [13] Francisco Facchinei and Jong-Shi Pang. *Finite-dimensional variational inequalities and complementarity problems. Vol. I*. Springer Series in Operations Research. Springer-Verlag, New York, 2003.
- [14] Ky Fan. Fixed-point and minimax theorems in locally convex topological linear spaces. *Proc. Nat. Acad. Sci. U.S.A.*, 38:121–126, 1952.
- [15] Sjur Didrik Flam. Paths to constrained Nash equilibria. *Appl. Math. Optim.*, 27(3):275–289, 1993.

- [16] I. L. Glicksberg. A further generalization of the Kakutani fixed theorem, with application to Nash equilibrium points. *Proc. Amer. Math. Soc.*, 3:170–174, 1952.
- [17] W. B. Gordon. On the diffeomorphisms of Euclidean space. *Amer. Math. Monthly*, 79:755–759, 1972.
- [18] Patrick R. Johnstone and Jonathan Eckstein. Convergence rates for projective splitting. *SIAM J. Optim.*, 29(3):1931–1957, 2019.
- [19] Shizuo Kakutani. A generalization of Brouwer’s fixed point theorem. *Duke Math. J.*, 8:457–459, 1941.
- [20] David Kinderlehrer and Guido Stampacchia. *An introduction to variational inequalities and their applications*, volume 31 of *Classics in Applied Mathematics*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2000. Reprint of the 1980 original.
- [21] Richard J. Lipton, Evangelos Markakis, and Aranyak Mehta. Playing large games using simple strategies. In *Proceedings of the 4th ACM Conference on Electronic Commerce, EC’03*, pages 36–41, New York, NY, USA, 2003. Association for Computing Machinery.
- [22] George J. Minty. Monotone (nonlinear) operators in Hilbert space. *Duke Math. J.*, 29:341–346, 1962.
- [23] George J. Minty. on a “monotonicity” method for the solution of non-linear equations in Banach spaces. *Proc. Nat. Acad. Sci. U.S.A.*, 50:1038–1041, 1963.
- [24] John Nash. Non-cooperative games. *Ann. of Math. (2)*, 54:286–295, 1951.
- [25] John F. Nash, Jr. Equilibrium points in n -person games. *Proc. Nat. Acad. Sci. U.S.A.*, 36:48–49, 1950.
- [26] J. B. Rosen. Existence and uniqueness of equilibrium points for concave n -person games. *Econometrica*, 33:520–534, 1965.