

HARDY INEQUALITIES IN FRACTIONAL ORLICZ-SOBOLEV SPACES

ARIEL M. SALORT

ABSTRACT. In this article we prove both norm and modular Hardy inequalities for a class functions in one-dimensional fractional Orlicz-Sobolev spaces.

1. INTRODUCTION

In the early 1920's, in the seminal article [15], G.H. Hardy obtained inequalities of the form

$$(1.1) \quad \int_0^\infty \frac{|u(x)|^p}{x^p} dx \leq \left(\frac{p}{p-1} \right)^p \int_0^\infty |u'(x)|^p dx$$

where $p > 1$ and u is a nonnegative measurable function defined on $(0, \infty)$.

Throughout the decades that followed, many authors contributed to characterize the family of weights v, w and powers p, q for which inequalities of the type

$$\left(\int_0^\infty |u(x)|^p v(x) dx \right)^{\frac{1}{p}} \leq \mathbf{C}_H \left(\int_0^\infty |u'(x)|^q w(x) dx \right)^{\frac{1}{q}}$$

are valid for a suitable positive constant $\mathbf{C}_H$ independent of u . See for instance the pioneering works [4, 30, 31]. We also refer to the seminal books [22, 26].

Orlicz spaces play a fundamental role when describing phenomena with non-standard growth. See [19, 21, 28]. Generalizations of Hardy type inequalities to the Orlicz space structure provide for a family of inequalities admitting behaviors more general than powers. In the last years many authors leaded the task of characterizing the class of admissible weights v, w, a, b and Young functions P, Q for which is valid an inequality as follows

$$Q^{-1} \left(\int_0^\infty Q(a(x)|Tu(x)|)v(x) dx \right) \leq \mathbf{C}_H P^{-1} \left(\int_0^\infty P(b(x)|u(x)|)w(x) dx \right),$$

where $Tu(x) = \int_0^x \mathcal{K}(x, y)u(y) dy$ is the generalized Hardy operator and $\mathbf{C}_H$ is a positive constant independent of u . For more details we refer to [6, 7, 16, 23] and references therein. In contrast with the L^p case, Hardy inequalities in integral form may differ from norm inequalities. See [6, 8].

Nonlocal Hardy inequalities were object of study in the last years. The nonlocal counterpart of (1.1), for suitable values of $p > 1$ and $s \in (0, 1)$, takes the form

$$\int_0^\infty \frac{|u(x)|^p}{x^{sp}} dx \leq \mathbf{C}_H \int_0^\infty \int_0^\infty \frac{|u(x) - u(y)|^p}{|x - y|^{1+sp}} dx dy$$

for functions in an appropriated Sobolev space. Hardy inequalities for the fractional p -Laplacian date back to the early 1960's and were derived independently in [14] and [18]. For the case $p = 2$ see also [1, 3]. See also [20] for a different approach.

Up to our knowledge, the only research on Hardy inequalities in a non-local framework with non-standard growth was made in [2] for values of $s \in (0, 1)$ close

2010 *Mathematics Subject Classification.* 46E30, 35R11, 45G05.

Key words and phrases. Fractional order Sobolev spaces, nonlocal eigenvalues, g -laplace operator, nonlocal Hardy inequalities.

to 0. See also [24]. Therefore, the main scope of this paper is to study the validity of these inequalities in the fractional order Orlicz-Sobolev spaces introduced in [2, 12], in both integral and norm form for *big* values of $s \in (0, 1)$, that is, when $s > 1/p^-$, where $p^- > 1$ is a fixed constant.

In order to state our results, recall that a *Young function* G is a continuous, nonnegative, strictly increasing and convex function on $\mathbb{R}_+ := [0, \infty)$ (see Section 2), for which we assume the growth condition

$$(L) \quad 1 < p^- \leq \frac{tg(t)}{G(t)} \leq p^+ < \infty \quad \text{for all } t > 0$$

for fixed constants $p^\pm$, where $g = G'$. Given a Young function G , we define

$$L^G(\mathbb{R}_+) := \{u: \mathbb{R}_+ \rightarrow \mathbb{R} \text{ measurable: } \Phi_G(u) < \infty\}.$$

The *modular* and the *Luxemburg norm* of $u \in L^G(\mathbb{R}_+)$, respectively, are defined as

$$\Phi_G(u) = \int_0^\infty G(|u(x)|) dx \quad \text{and} \quad \|u\|_G := \inf \left\{ \lambda > 0: \Phi_G\left(\frac{u}{\lambda}\right) \leq 1 \right\}.$$

Given $s \in (0, 1)$, we define the *fractional Orlicz-Sobolev space* as (see [12])

$$W^{s,G}(\mathbb{R}_+) := \{u \in L^G(\mathbb{R}_+) \text{ such that } \Phi_{s,G}(u) < \infty\}.$$

Here the *fractional modular* of $u \in W^{s,G}(\mathbb{R}_+)$ is defined as

$$\Phi_{s,G}(u) := \int_0^\infty \int_0^\infty G\left(\frac{|u(x) - u(y)|}{|x - y|^s}\right) \frac{dx dy}{|x - y|}.$$

These spaces are endowed with the so-called *Luxemburg norm* defined as

$$\|u\|_{s,G} := \|u\|_G + [u]_{s,G},$$

where the (s, G) -Gagliardo semi-norm reads as $[u]_{s,G} := \inf \left\{ \lambda > 0: \Phi_{s,G}\left(\frac{u}{\lambda}\right) \leq 1 \right\}$.

We describe now our results. As we will see, the proof of our non-local modular inequality is based on an astute application of the following modular Hardy inequality for the local Orlicz-Sobolev space $W^{1,G}(\mathbb{R}_+)$ (see Lemma 3.4 for details)

$$(1.2) \quad \int_0^\infty G\left(\left|\frac{1}{x^s} \int_0^x \frac{u(t)}{t} dt\right|\right) dx \leq \mathbf{c}_H \int_0^\infty G\left(\frac{|u(x)|}{x^s}\right) dx.$$

Following [5], inequality (1.2) can be obtained from the local norm inequalities studied in [25, 27].

Finally, for simplicity in our notation, given a Young function G satisfying (L) we define the functions

$$(1.3) \quad \psi_G(x) = \begin{cases} x^{p^+} & \text{if } x \geq 1 \\ x^{p^-} & \text{if } x < 1 \end{cases} \quad \text{and} \quad \phi_G(x) = \begin{cases} x^{1/p^-} & \text{if } x \geq 1 \\ x^{1/p^+} & \text{if } x < 1 \end{cases}.$$

With these preliminaries, our first result reads as follows.

Theorem 1.1. *Let G be a Young function satisfying (L) and let $s \in (0, 1)$ be such that $sp^- > 1$. Then for all $u \in W^{s,G}(\mathbb{R}_+)$ such that $\lim_{x \rightarrow 0} \frac{1}{x} \int_0^x u(x) dx = u_0$, the following inequality holds*

$$\int_0^\infty G\left(\frac{|u(x) - u(x_0)|}{x^s}\right) dx \leq \mathbf{C}_H \Phi_{s,G}(u), \quad \mathbf{C}_H := \mathbf{C}(1 + \mathbf{c}_H)$$

where $\mathbf{c}_H$ is given in (1.2) and $\mathbf{C} := 2^{p^+}$ is the doubling constant for G .

Remark 1.2. The constant $\mathbf{C}_H$ in Theorem 1.1 can be computed explicitly as

$$\mathbf{C}_H = \mathbf{C}(1 + \mathbf{c}_H) = 2^{p^+} \left(1 + \psi_G \left(\frac{p^-}{sp^- - 1} \right) \right).$$

Remark 1.3. Observe that $u \in W^{s,G}(\mathbb{R}_+)$ is in fact continuous in a neighborhood $\mathcal{O}$ of the origin when $sp^- > 1$ due to the embedding of $W^{s,G}(\mathcal{O})$ into the space $C^{0,s-1/p^-}(\mathcal{O})$ of Hölder continuous functions (see Proposition 2.2).

From the modular inequality it is easy to deduce a norm inequality.

Corollary 1.4. *With the same assumptions of Theorem 1.1,*

$$\left\| \frac{u - u_0}{x^s} \right\|_G \leq \phi_G(\mathbf{C}_H)[u]_{s,G}$$

where ϕ_G is given in (1.3) and $\mathbf{C}_H$ is the constant given in Theorem 1.1.

From Theorem 1.1 and Remark 1.3 it can be easily deduced the following consequence.

Corollary 1.5. *Let G be a Young function satisfying (L) and let $s \in (0, 1)$ be such that $sp^- > 1$. Then for all $u \in W^{s,G}(\mathbb{R}_+)$ such that $u(0) = 0$ it holds that*

$$\int_0^\infty G \left(\frac{|u(x)|}{x^s} \right) dx \leq \mathbf{C}_H \Phi_{s,G}(u), \quad \left\| \frac{u}{x^s} \right\|_G \leq \phi_G(\mathbf{C}_H)[u]_{s,G},$$

where ϕ_G is given in (1.3) and $\mathbf{C}_H$ is the constant given in Theorem 1.1.

Although a norm inequality can be deduced from Theorem 1.1, it can also be obtained independently of the modular inequality. In fact, the following result provides for a more accurate constant.

Theorem 1.6. *Let G be a Young function satisfying (L) and let $s \in (0, 1)$ be such that $sp^- > 1$. Then*

$$\left\| \frac{u}{x^s} \right\|_G \leq \frac{(1+s)p^- - 1}{sp^- - 1} [u]_{s,G}$$

for all $u \in W^{s,G}(\mathbb{R}_+)$.

Sharpness of the Hardy constant is known both in the local and nonlocal case when G is a power, see [13, 17] for instance. However, in the Orlicz setting it is unknown even in the local case.

2. PRELIMINARY RESULTS

2.1. Young functions. A *Young function* is an application $G: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which is continuous, nonnegative, strictly increasing, convex on $[0, \infty)$ and admits the integral formulation $G(t) = \int_0^t g(s) ds$. For some fixed constants $p^\pm$ we assume that G satisfies the growth condition (L).

The *complementary Young function* $\tilde{G}$ of a Young function G is defined as

$$\tilde{G}(t) = \sup\{tw - G(w) : w > 0\}.$$

We introduce some well-known results on Young function. See [21, 28] for details.

Lemma 2.1. *Let G be a Young function satisfying (L) and $a, t \geq 0$. Then*

$$\begin{aligned} (G_1) \quad & \min\{a^{p^-}, a^{p^+}\}G(t) \leq G(at) \leq \max\{a^{p^-}, a^{p^+}\}G(t), \\ (G_2) \quad & G(a+t) \leq \mathbf{C}(G(a) + G(t)) \quad \text{with } \mathbf{C} := 2^{p^+}. \end{aligned}$$

Condition (G₂) is known as the Δ_2 condition or doubling condition. It can be proved that (L) implies that both G and $\tilde{G}$ satisfy (G₂). See [21, Theorem 3.4.4 and Theorem 3.13.9].

Young functions includes for instance powers (when $g(t) = t^{p-1}$, $p^\pm = p > 1$ and hence $G(t) = \frac{t^p}{p}$) and logarithmic perturbations of powers (when $g(t) = t \log(b+ct)$, where $p^- = 1 + a$, $p^+ = 2 + a$). See [21] for more examples.

2.2. Fractional Orlicz-Sobolev spaces. Given a Young function G , a fractional parameter $s \in (0, 1)$ and an open interval $\Omega \subseteq \mathbb{R}$, we have already defined the fractional Orlicz-Sobolev space $W^{s,G}(\Omega)$ in the introduction. We also define the following related space

$$W_0^{s,G}(\Omega) := \{u \in W^{s,G}(\mathbb{R}) : u = 0 \text{ a.e. in } \mathbb{R} \setminus \Omega\},$$

which coincides with the closure of C_c^∞ functions with respect to the $\|\cdot\|_{s,G}$ norm, and it is the natural space for the well-posedness of Dirichlet problems.

For a further generalization of these spaces we refer to [9].

The following result characterizes continuous functions in fractional Orlicz-Sobolev spaces.

Proposition 2.2. *Let G be a Young function satisfying (L) and let $s \in (0, 1)$ be such that $sp^- > 1$. Then, given an open and bounded interval $\Omega \subset \mathbb{R}$, it holds that $W^{s,G}(\Omega) \subset C^{0,s-1/p^-}(\Omega)$.*

Proof. From [11, Proposition 2.9 and Proposition 2.7], for any $u \in W^{s,G}(\Omega)$ we get

$$[u]_{W_{reg}^{s,p^-}(\Omega)} + \|u\|_{L^{p^-}(\Omega)} \leq C([u]_{W^{s,G}(\Omega)} + \|u\|_{L^G(\Omega)}).$$

where

$$[u]_{W_{reg}^{s,p^-}(\Omega)} = \left(\iint_{\Omega \times \Omega} \frac{|u(x) - u(y)|^{p^-}}{|x - y|^{n+sp^-}} dx dy \right)^{\frac{1}{p^-}}.$$

Moreover, by [10, Theorem 8.2], $\|u\|_{C^{0,s-1/p^-}(\Omega)} \leq C([u]_{W_{reg}^{s,p^-}(\Omega)} + \|u\|_{L^{p^-}(\Omega)})$ and the result follows. $\square$

3. ONE-DIMENSIONAL HARDY INEQUALITIES

In this section we prove our main results. First, following [5], we link norm inequalities for linear operators with integral inequalities.

Proposition 3.1. *Let G be a Young function satisfying (L). Suppose that the inequality $\|Tu\|_{\varepsilon G} \leq C\|u\|_{\varepsilon G}$ holds for all $\varepsilon > 0$ with C independent of ε , where*

$\|u\|_{\varepsilon G} = \inf \left\{ \lambda: \int_0^\infty G\left(\frac{|u(x)|}{\lambda}\right) \varepsilon dx \leq 1 \right\}$ and T is a linear operator. Then it holds that

$$\int_0^\infty G(|Tu(x)|) dx \leq \int_0^\infty G(C|u(x)|) dx.$$

Proof. Given $u \in L^G(\mathbb{R}_+)$, define the number $\varepsilon = \left(\int_0^\infty G(|u|) dx\right)^{-1}$ and observe that $\|u\|_{\varepsilon G} \leq 1$. Therefore, $\|Tu\|_{\varepsilon G} \leq C\|u\|_{\varepsilon G} \leq C$, and then, by definition of the Luxemburg norm we get

$$\int_0^\infty G\left(\frac{|Tu|}{C}\right) dx \leq \frac{1}{\varepsilon} \int_0^\infty G\left(\frac{|Tu|}{\|Tu\|_{\varepsilon G}}\right) \varepsilon dx \leq \frac{1}{\varepsilon} = \int_0^\infty G(|u|) dx.$$

Finally, since T is linear, replacing u with Cu the result follows. $\square$

In order to apply Proposition 3.1 we use the following inequality due to [27] (cf. also [25, Corollary 4]).

Proposition 3.2. *Given $I = (0, \ell)$, $0 < \ell \leq \infty$, if $\theta \in \mathbb{R}$ is such that $\theta < 1/(p^-)'$, then, for $x^\theta u(x) \in L^G(I)$,*

$$\left\| x^{\theta-1} \int_0^x u(t) dt \right\|_{L^G(I)} \leq \frac{(p^-)'}{1 - \theta(p^-)'} \|x^\theta u(x)\|_{L^G(I)}.$$

where $(p^-)'$ is the conjugated exponent of p^- .

Corollary 3.3. *Given $I = (0, \ell)$, $0 < \ell \leq \infty$ and $s \in (0, 1)$,*

(i) *if $\theta = 1 - s$ and $x^{1-s}u(x) \in L^G(I)$, for $sp^- > 1$ it holds that*

$$(3.1) \quad \left\| \frac{1}{x^s} \int_0^x u(t) dt \right\|_{L^G(I)} \leq \frac{p^-}{sp^- - 1} \|x^{1-s}u(x)\|_{L^G(I)},$$

(ii) *if $\theta = -s$ and $x^{-s}u(x) \in L^G(I)$, for $(1+s)p^- > 1$ it holds that*

$$(3.2) \quad \left\| \frac{1}{x^{1+s}} \int_0^x u(t) dt \right\|_{L^G(I)} \leq \frac{p^-}{(1+s)p^- - 1} \|x^{-s}u(x)\|_{L^G(I)}.$$

We prove now a key lemma for our arguments.

Lemma 3.4. *Let G be a Young function satisfying (L) and let $s \in (0, 1)$ be such that $sp^- > 1$. Given u such that $x^{-s}u(x) \in L^G(\mathbb{R}_+)$, then we have that*

$$(3.3) \quad \int_0^\infty G\left(\left|\frac{1}{x^s} \int_0^x \frac{u(t)}{t} dt\right|\right) dx \leq \mathbf{c}_H \int_0^\infty G\left(\frac{|u(x)|}{x^s}\right) dx.$$

The constant $\mathbf{c}_H$ is given by $\mathbf{c}_H = \psi_G\left(\frac{p^-}{sp^- - 1}\right)$, where ψ_G is given in (1.3).

Proof. Given G satisfying (L), we define the Young function $G_\varepsilon := \varepsilon G$ for some $\varepsilon > 0$. It is immediate that G_ε satisfies (L) with the same constants that G .

Let u be a fixed function such that $x^{-s}u(x) \in L^G(\mathbb{R}_+)$, then also $x^{-s}u(x) \in L^{G_\varepsilon}(\mathbb{R}_+)$. Hence, by applying Corollary 3.3 item (3.1) to $x^{-1}u(x)$ we find that

$$(3.4) \quad \left\| \frac{1}{x^s} \int_0^x \frac{u(t)}{t} dt \right\|_{G_\varepsilon} \leq \frac{p^-}{sp^- - 1} \left\| \frac{u}{x^s} \right\|_{G_\varepsilon} \quad \forall \varepsilon > 0.$$

Since (3.4) holds with constant independent of ε , from Proposition 3.1 we get that

$$\int_0^\infty G\left(\left|\frac{1}{x^s} \int_0^x \frac{u(t)}{t} dt\right|\right) dx \leq \int_0^\infty G\left(\frac{p^-}{sp^- - 1} \frac{|u(x)|}{x^s}\right) dx$$

and the result follows by using condition (G_1) . $\square$

We are ready to prove our main result.

Proof of Theorem 1.1. We assume without loss of generality that

$$\lim_{x \rightarrow 0} \frac{1}{x} \int_0^x u(x) dx = 0.$$

Then, the general case will follow by applying Theorem 1.1 to $u - u_0$.

Given $u \in W^{s,G}(\mathbb{R}_+)$ we consider the auxiliary function

$$v(x) = u(x) - \frac{1}{x} \int_0^x u(t) dt.$$

For $0 < a < b < \infty$, using integration by parts it is straightforward to see that

$$\begin{aligned} \int_a^b \frac{v(x)}{x} dx &= \int_a^b \frac{u(x)}{x} dx - \int_a^b \frac{1}{x^2} \int_0^x u(s) ds dx \\ &= \int_a^b \frac{u(x)}{x} dx + \frac{1}{x} \int_0^x u(s) ds \Big|_a^b - \int_a^b \frac{u(x)}{x} dx \\ &= \frac{1}{b} \int_0^b u(x) dx - \frac{1}{a} \int_0^a u(x) dx. \end{aligned}$$

Taking $b = t$ and $a \rightarrow 0$ in the last expression we get $\int_0^t \frac{v(x)}{x} dx = \frac{1}{t} \int_0^t u(x) dx$, and

$$(3.5) \quad u(x) = v(x) + \int_0^x \frac{v(t)}{t} dt.$$

We prove now that $x^{-s}v(x) \in L^G(\mathbb{R}_+)$. Indeed, by definition of v we have that

$$\int_0^\infty G\left(\frac{|v(x)|}{x^s}\right) dx = \int_0^\infty G\left(\frac{|u(x) - \frac{1}{x} \int_0^x u(t) dt|}{x^s}\right) dx.$$

Now, by using Jensen's inequality we get

$$\begin{aligned} \int_0^\infty G\left(\frac{|u(x) - \frac{1}{x} \int_0^x u(t) dt|}{x^s}\right) dx &\leq \int_0^\infty G\left(\frac{1}{x} \int_0^x \frac{|u(x) - u(t)|}{x^s} dt\right) dx \\ &\leq \int_0^\infty \frac{1}{x} \int_0^x G\left(\frac{|u(x) - u(t)|}{x^s}\right) dt dx, \end{aligned}$$

and the last term in the inequality above can be bounded as

$$\begin{aligned} \int_0^\infty \frac{1}{x} \int_0^x G\left(\frac{|u(x) - u(t)|}{x^s}\right) dt dx &\leq \int_0^\infty \int_0^x G\left(\frac{|u(x) - u(y)|}{|x - y|^s}\right) \frac{dy dx}{|x - y|} \\ &\leq \int_0^\infty \int_0^\infty G\left(\frac{|u(x) - u(y)|}{|x - y|^s}\right) \frac{dy dx}{|x - y|}. \end{aligned}$$

Since $x^{-s}v(x) \in L^G(\mathbb{R}_+)$, we are in position to apply Lemma 3.4 to v , obtaining

$$(3.6) \quad \int_0^\infty G\left(\left|\frac{1}{x^s} \int_0^x \frac{v(t)}{t} dt\right|\right) dx \leq \mathbf{c}_H \int_0^\infty G\left(\frac{|v(x)|}{x^s}\right) dx.$$

Therefore, from (3.5), property (G₂) and (3.6) we find that

$$\begin{aligned} \int_0^\infty G\left(\frac{|u(x)|}{x^s}\right) dx &\leq \mathbf{C} \int_0^\infty G\left(\frac{|v(x)|}{x^s}\right) dx + \mathbf{C} \int_0^\infty G\left(\frac{1}{x^s} \left|\int_0^x \frac{v(t)}{t} dt\right|\right) dx \\ &\leq \mathbf{C}(1 + \mathbf{c}_H) \int_0^\infty G\left(\frac{|v(x)|}{x^s}\right) dx \\ &\leq \mathbf{C}(1 + \mathbf{c}_H) \Phi_{s,G}(u) \end{aligned}$$

giving the desired inequality. $\square$

As a corollary, we obtain the Hardy inequality for norms stated in Corollary 1.4.

Proof of Corollary 1.4. We assume without loss of generality that

$$\lim_{x \rightarrow 0} \frac{1}{x} \int_0^x u(x) dx = 0.$$

Then, the general case will follow by applying the inequality to $u - u_0$.

By using Theorem 1.1 together with (G₁) one gets that $\Phi_G(\frac{u}{x^s}) \leq \Phi_{s,G}(\tilde{\mathbf{C}}_H u)$ for $u \in W^{s,G}(\mathbb{R}_+)$, where $\tilde{\mathbf{C}}_H = \max\{\mathbf{C}_H^{\frac{1}{p^-}}, \mathbf{C}_H^{\frac{1}{p^+}}\}$. Then, given $u \in W^{s,G}(\mathbb{R}_+)$, from the last expression we find that $\Phi_G(u/\tilde{\mathbf{C}}_H[u]_{s,G}x^s) \leq \Phi_{s,G}(u/[u]_{s,G}) \leq 1$. Hence, by definition of the Luxemburg norm we get

$$\left\| \frac{u}{x^s} \right\|_G = \inf \left\{ \lambda : \Phi_G\left(\frac{u}{\lambda x^s}\right) \leq 1 \right\} \leq \tilde{\mathbf{C}}_H [u]_{s,G}$$

and the proof concludes. $\square$

Now, we provide for the proof of the norm inequality given in Theorem 1.6.

Proof of Theorem 1.6. Given $u \in W^{s,G}(\mathbb{R}_+)$, by using the triangular inequality for the Luxemburg norm we obtain that

$$\left\| \frac{u}{x^s} \right\|_G \leq \left\| \frac{u - \frac{1}{x} \int_0^x u(y) dy}{x^s} \right\|_G + \left\| \frac{1}{x^{1+s}} \int_0^x u(y) dy \right\|_G := (i) + (ii).$$

Let us find a bound for (i). By using Jensen's inequality we get

$$\begin{aligned} \int_0^\infty G\left(\frac{|u(x) - \frac{1}{x} \int_0^x u(t) dt|}{x^s}\right) dx &\leq \int_0^\infty G\left(\frac{1}{x} \int_0^x \frac{|u(x) - u(t)|}{x^s} dt\right) dx \\ &\leq \int_0^\infty \frac{1}{x} \int_0^x G\left(\frac{|u(x) - u(t)|}{x^s}\right) dt dx \leq \int_0^\infty \int_0^x G\left(\frac{|u(x) - u(y)|}{|x-y|^s}\right) \frac{dy dx}{|x-y|} \\ &\leq \Phi_{s,G}(u). \end{aligned}$$

The inequality above applied to $u/[u]_{s,G}$ gives that

$$\int_0^\infty G\left(\frac{|u(x) - \frac{1}{x} \int_0^x u(y) dy|}{x^s [u]_{s,G}}\right) dx \leq \Phi_{s,G}\left(\frac{u}{[u]_{s,G}}\right) \leq 1,$$

from where, from the definition of the Luxemburg norm, we obtain that (i) is bounded by $[u]_{s,G}$. Expression (3.2) from Corollary 3.3 gives that (ii) is less than $\frac{p^-}{(1+s)p^- - 1} \left\| \frac{u}{x^s} \right\|_G$. From these two relations we get that

$$\left\| \frac{u}{x^s} \right\|_G \leq [u]_{s,G} + \frac{p^-}{(1+s)p^- - 1} \left\| \frac{u}{x^s} \right\|_G.$$

Finally, the condition $sp^- > 1$ leads to

$$\mathbf{C}_H = \left(1 - \frac{p^-}{(1+s)p^- - 1} \right)^{-1} = \frac{(1+s)p^- - 1}{sp^- - 1} > 0$$

and the desired inequality is obtained. $\square$

4. APPLICATIONS AND EXAMPLES

4.1. Lower bound of Dirichlet eigenvalues. Given a Young function G satisfying (L) and a fractional parameter $s \in (0, 1)$ such that $sp^- > 1$, consider the eigenvalue problem for the fractional g -Laplacian operator in an open and bounded interval $\Omega \subset \mathbb{R}^+$ (see [12, 29]):

$$(4.1) \quad \begin{cases} (-\Delta_g)^s u = \lambda g(u)u & \text{in } \Omega, \\ u = 0 & \text{on } \mathbb{R} \setminus \Omega. \end{cases}$$

This operator is well defined between $W^{s,G}(\mathbb{R})$ and its dual, and acts as

$$(-\Delta_g)^s u(x) := 2p.v. \int_{-\infty}^{\infty} g(|D_s u|) \frac{D_s u}{|D_s u|} \frac{dy}{|x-y|^{1+s}},$$

where $D_s u(x, y) = \frac{u(x) - u(y)}{|x-y|^s}$. The following representation formula holds

$$\langle (-\Delta_g)^s u, v \rangle = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} g(|D_s u|) \frac{D_s u}{|D_s u|} D_s v \frac{dx dy}{|x-y|} \quad \forall v \in W^{s,G}(\mathbb{R}).$$

The natural space for solutions of (4.1) is $W_0^{s,G}(\Omega)$, i.e., functions in $W^{s,G}(\mathbb{R})$ such that $u = 0$ in $\mathbb{R} \setminus \Omega$.

Due to the possible lack of homogeneity of the problem, eigenpairs depend strongly on the normalization: for each $\alpha > 0$, $u_\alpha \in \mathcal{X}_\alpha := \left\{ W_0^{s,G}(\Omega) : \Phi_G(u) = \alpha \right\}$ is an eigenfunction of (4.1) with eigenvalue λ_α if it holds that

$$\langle (-\Delta_g)^s u_\alpha, v \rangle = \lambda_\alpha \int_{\Omega} g(u_\alpha) u_\alpha \frac{u_\alpha}{|u_\alpha|} v dx \quad \forall v \in W_0^{s,G}(\Omega).$$

On the other hand, for each $\alpha > 0$ we define the minimizer (see [11, Theorem 2.12])

$$(4.2) \quad \Lambda_\alpha := \inf \{ \Phi_{s,G}(u) / \Phi_G(u) : u \in \mathcal{X}_\alpha \}.$$

In [29] it is proved that for each $\alpha > 0$ the infimum in (4.2) is attained for some function $u_\alpha \in \mathcal{X}_\alpha$, and then, by a Lagrange's multiplier argument it is deduced the existence of an eigenvalue λ_α of (4.1) with eigenfunction u_α . However, in contrast with the case of powers, in general λ_α does not admit a variational characterization and $\lambda_\alpha \neq \Lambda_\alpha$. Both constants are comparable each other. Indeed, $\frac{p^-}{p^+} \Lambda_\alpha \leq \lambda_\alpha \leq \frac{p^+}{p^-} \Lambda_\alpha$.

Using Lemma 2.1 it is easy to see that for any function $u \in \mathcal{X}_\alpha$ it holds that

$$\int_{\Omega} G(|u|) dx \leq \int_{\Omega} G\left(\frac{|u(x)|}{x^s} \text{diam}(\Omega)\right) dx \leq \psi_G(\text{diam}(\Omega)) \int_{\Omega} G\left(\frac{|u(x)|}{x^s}\right) dx,$$

where ψ_G is given in (1.3). Hence, from Theorems 1.1 we get that

$$(\psi_G(\text{diam}(\Omega))\mathbf{C}_H)^{-1} \leq \Lambda_\alpha \leq \frac{p^+}{p^-} \lambda_\alpha.$$

4.2. Lower bound of weighted eigenvalues. With the same ideas of [29], we can consider eigenvalues λ_α and minimizers Λ_α of the weighted problem

$$(4.3) \quad \begin{cases} (-\Delta_g)^s u = \frac{\lambda}{|x|^s} g\left(\frac{|u|}{x^s}\right) \frac{u}{|u|} & \text{in } \Omega, \\ u = 0 & \text{on } \mathbb{R} \setminus \Omega. \end{cases}$$

where $\Omega \subset \mathbb{R}^+$ is an open and bounded interval. Given $\alpha > 0$, a number λ_α is an eigenvalue of (4.3) with eigenfunction $u_\alpha \in \mathcal{X}_\alpha := \left\{ W_0^{s,G}(\Omega) : \int_{\Omega} G\left(\frac{|u(x)|}{x^s}\right) dx = \alpha \right\}$ if it holds that

$$\langle (-\Delta_g)^s u_\alpha, v \rangle = \lambda_\alpha \int_{\Omega} g\left(\frac{|u_\alpha(x)|}{|x|^s}\right) \frac{u_\alpha(x)}{|u_\alpha(x)|} \frac{v(x)}{x^s} dx \quad \forall v \in W_0^{s,G}(\Omega).$$

We also define the number $\Lambda_\alpha := \inf \left\{ \Phi_{s,G}(u) / \int_{\Omega} G\left(\frac{|u(x)|}{x^s}\right) dx : u \in \mathcal{X}_\alpha \right\}$.

We have again in this case that $\frac{p^-}{p^+} \Lambda_\alpha \leq \lambda_\alpha \leq \frac{p^+}{p^-} \Lambda_\alpha$. As a direct consequence of Theorem 1.1 we get $\frac{1}{\mathbf{C}_H} \leq \Lambda_\alpha \leq \frac{p^+}{p^-} \lambda_\alpha$.

ACKNOWLEDGEMENTS

This paper is partially supported by grants UBACyT 20020130100283BA, CONICET PIP 11220150100032CO and ANPCyT PICT 2012-0153.

REFERENCES

- [1] Adams, R., Aronszajn, N. and Smith, K., *Theory of Bessel potentials*, Part II, Ann. Inst. Fourier (Grenoble) 17 (1967), 1-135. [1](#)
- [2] Alberico, A., Cianchi, A., Pick, L. and Slavkova, L. *Fractional Orlicz-Sobolev embeddings*, arxiv preprint 2001.05565 (2020) [1](#), [2](#)
- [3] Aronszajn, N. and Smith, K., *Theory of Bessel potentials*, Technical report 26, Department of Mathematics, University of Kansas, USA, 1961. [1](#)
- [4] Bradley, J.S., *Hardy's inequality with weights*, Can. Math. Bull., 21, No. 4, 405-408 (1978). [1](#)
- [5] Bloom S. and Kerman, R., *Weighted L_Φ integral inequalities for operators of Hardy type*. Studia Mathematica, 1(110), 35-52, (1994). [2](#), [4](#)
- [6] Bloom, S. and Kerman, R., *Weighted norm inequalities for operators of Hardy type*, Proc. Amer. Math. Soc. 113, 135141, (1991). [1](#)
- [7] Bloom, S. and Kerman, R., *Weighted Orlicz space integral inequalities for the Hardy-Littlewood maximal operator*, Studia Math. 110, 149167, (1994). [1](#)
- [8] Cianchi, A., *Hardy inequalities in Orlicz spaces*. Transactions of the American Mathematical Society, 351(6), 2459-2478, (1999). [1](#)
- [9] De Nápoli, P., Fernández Bonder, J. and Salort, A. (2019). *A Pólya-Szegő principle for general fractional Orlicz-Sobolev spaces*. Complex Variables and Elliptic Equations, 2020, p. 1-23. [4](#)
- [10] Di Nezza, E., Palatucci, G., and Valdinoci, E. *Hitchhiker's guide to the fractional Sobolev spaces*. Bulletin des Sciences Mathématiques, 136(5), 521-573. (2012). [4](#)

- [11] Fernández Bonder, J., Pérez Llanos, M. and Salort, A., *A Hölder Infinity Laplacian obtained as limit of Orlicz Fractional Laplacians*, preprint arXiv:1807.01669 [4](#), [8](#)
- [12] Fernández Bonder, J. and Salort, A., *Fractional order Orlicz-Sobolev spaces*, Journal of Functional Analysis, 2019, vol. 277, no 2, p. 333-367. [2](#), [8](#)
- [13] Frank, R. L., and Seiringer, R. *Non-linear ground state representations and sharp Hardy inequalities*. Journal of Functional Analysis, 255(12), 3407-3430. (2008) [3](#)
- [14] Grisvard, P., *Espaces intermediaires entre espaces de Sobolev avec poids*, Ann. Scuola Norm. Sup. Pisa 17 (1963), 255-296. [1](#)
- [15] Hardy, G. H., *Note on a theorem of Hilbert*, Math. Z. 6, 314317, (1920). [1](#)
- [16] Heinig, H. and Maligranda, L., *Interpolation with weights in Orlicz spaces*, Boll. Un. Mat. Ital. (7) 8, 3755, (1994). [1](#)
- [17] Herbst, I., *Spectral Theory of the Operator $(p^2 + m^2)^{1/2} - Ze^2/r$* . Comm. Math. Phys. 53(1977), no. 3, 285-294. [3](#)
- [18] Jakovlev, G., *Boundary properties of a certain class of functions (Russian)*. Trudy Mat. Inst. Steklov 60, (1961). [1](#)
- [19] Krasnoselskii, M. and Rutitskii, I., *Convex functions and Orlicz spaces*. (1961). [1](#)
- [20] Krugljak, N., Maligranda, L., and Persson, L., *On an elementary approach to the fractional Hardy inequality*. Proc. Amer. Math. Soc., 128(3), 727-734. (2000). [1](#)
- [21] Kufner, A., John, O. and Fucik, S., *Function spaces*(Vol. 3). Springer Science Business Media. (1979). [1](#), [3](#), [4](#)
- [22] Kufner, A., Persson, L. and Samko, N. *Weighted inequalities of Hardy type*. World Scientific Publishing Company. (2017). [1](#)
- [23] Lai, Q., *Two weight mixed Φ -inequalities for the Hardy operator and the Hardy-Littlewood maximal operator*, J. London Math. Soc. 48, 301318, (1992). [1](#)
- [24] Maione, A., Salort, A. and Vecchi, E., *Maz'ya-Shaposhnikova formula in Magnetic Fractional Orlicz-Sobolev spaces*. arXiv preprint arXiv:2005.04662 (2020). [2](#)
- [25] Maligranda, L., *Generalized Hardy inequalities in rearrangement invariant spaces*. Journal des Mathematiques Pures et Appliques, 59(4), 405-415, (1980). [2](#), [5](#)
- [26] Opic, B., and Kufner, A. *Hardy-type inequalities* (Vol. 219). Halsted Press. (1990). [1](#)
- [27] Palmieri, P., *Un approccio alla teoria degli spazi di traccia relativi agli spazi di Orlicz-Sobolev*, Bolletino U. M. I. 5, 16B, (1979). [2](#), [5](#)
- [28] Rao, M. and Ren, Z., *Applications of Orlicz spaces* (Vol. 250). CRC Press. (2002). [1](#), [3](#)
- [29] Salort, A. *Eigenvalues and minimizers for a non-standard growth non-local operator*, Journal of Differential Equations, 2020, vol. 268, no 9, p. 5413-5439. [8](#), [9](#)
- [30] Talenti, G., *Osservazioni sopra una classe di disuguaglianze*, Rend. Sem. Mat. Fis. Milano, 39, 171-185 (1969). [1](#)
- [31] Tomaselli, G., *A class of inequalities*, Boll. Un. Mat. Ital., Set. IV, No. 6, 622-631 (1969). [1](#)

DEPARTAMENTO DE MATEMÁTICA, FCEyN - UNIVERSIDAD DE BUENOS AIRES AND
 IMAS - CONICET - CIUDAD UNIVERSITARIA, PABELLÓN I (1428) AV. CANTILLO S/N.
 BUENOS AIRES, ARGENTINA.

E-mail address, A.M. Salort: asalort@dm.uba.ar

URL: <http://mate.dm.uba.ar/~asalort>