

SHARP NONUNIQUENESS FOR THE NAVIER-STOKES EQUATIONS

ALEXEY CHESKIDOV AND XIAOYUTAO LUO

ABSTRACT. In this paper, we prove a sharp nonuniqueness result for the incompressible Navier-Stokes equations in the periodic setting. In any dimension $d \geq 2$ and given any $p < 2$, we show the nonuniqueness of weak solutions in the class $L_t^p L^\infty$, which is sharp in view of classical uniqueness results. The proof is based on the construction of a class of non-Leray-Hopf weak solutions. More specifically, for any $p < 2$, $q < \infty$, and $\varepsilon > 0$, we construct non-Leray-Hopf weak solutions $u \in L_t^p L^\infty \cap L_t^1 W^{1,q}$ that are locally smooth outside a singular set in time of Hausdorff dimension less than ε . As a byproduct, examples of anomalous dissipation in the class $L_t^{3/2-\varepsilon} C^{1/3}$ for the Euler equations are given.

1. INTRODUCTION

1.1. The incompressible Navier-Stokes equations. The Navier-Stokes equations are a fundamental mathematical model of incompressible viscous fluid flow, which consists of a system of equations

$$\begin{cases} \partial_t u - \Delta u + \operatorname{div}(u \otimes u) + \nabla p = 0 \\ \operatorname{div} u = 0, \end{cases} \quad (1.1)$$

posed on a spatial domain $\Omega \subset \mathbb{R}^d$ with suitable boundary conditions. In (1.1), $u : [0, T] \times \Omega \rightarrow \mathbb{R}^d$ is the unknown velocity, and $p : [0, T] \times \Omega \rightarrow \mathbb{R}$ is a scalar pressure. We consider the Cauchy problem of (1.1) on a time interval $[0, T]$ for some initial data u_0 and $T > 0$.

In the paper we confine ourselves to the periodic case $\Omega = \mathbb{T}^d = \mathbb{R}^d / \mathbb{Z}^d$ in dimension $d \geq 2$ and consider solutions with zero spacial mean

$$\int_{\mathbb{T}^d} u(t, x) dx = 0,$$

which is propagated under the evolution of the equation (1.1).

In this paper, the notion of weak solutions refers to that of distributional solutions that solve (1.1) in the sense space-time distribution, cf. [FJR72, Che11].

Definition 1.1. Denote by \mathcal{D}_T the space of divergence-free test function $\varphi \in C^\infty(\mathbb{R} \times \mathbb{T}^d)$ such that $\varphi = 0$ if $t \geq T$.

Let $u_0 \in L^2(\mathbb{T}^d)$ be weakly divergence-free¹. A vector field $u \in L^2([0, T] \times \mathbb{T}^d)$ is a weak solution of (1.1) with initial data u_0 if the following hold:

- (1) For a.e. $t \in [0, T]$, u is weakly divergence-free;
- (2) For any $\varphi \in \mathcal{D}_T$,

$$\int_{\mathbb{T}^d} u_0(x) \cdot \varphi(0, x) dx = - \int_0^T \int_{\mathbb{T}^d} u \cdot (\partial_t \varphi + \Delta \varphi + u \cdot \nabla \varphi) dx dt. \quad (1.2)$$

In the literature, such solutions are sometimes called “very weak solutions” [Ama00, Gal00] due to the minimal regularity assumptions. Remarkably, by [FJR72, Theorem 2.1], up to possibly redefining u on a set of measure zero in space-time, the above weak formulation is equivalent to the integral equation

$$u = e^{t\Delta} u_0 + \int_0^t e^{(t-s)\Delta} \mathbb{P} \operatorname{div}(u \otimes u)(s) ds, \quad (1.3)$$

where $e^{t\Delta}$ is the heat semigroup and \mathbb{P} is the Leray projection onto the divergence-free vector fields. Note that formulation (1.3) was also used in a variety of works [FK64, Kat84, KT01] to construct unique solutions (called mild solutions) of (1.1) when the initial data u_0 is critical or subcritical, starting from the work of Fujita and Kato [FK64].

Date: September 15, 2020.

¹It is possible to consider more general initial data, such as L^p for some $1 \leq p \leq \infty$ as in [FJR72].

A more physical class of weak solutions, introduced by Leray [Ler34] and constructed by Leray [Ler34] in \mathbb{R}^3 and Hopf [Hop51] in general domains in $d \geq 2$, is also considered in the literature.

Definition 1.2. A weak solution u of (1.1) is called *Leray-Hopf weak solution* if $u \in C_w([0, T; L^2] \cap L^2(0, T; H^1))$ and

$$\frac{1}{2} \|u(t)\|_2^2 + \int_0^t \|\nabla u(s)\|_2^2 ds \leq \frac{1}{2} \|u(0)\|_2^2, \quad (1.4)$$

for all $t \in [0, T]$

The Leray-Hopf weak solutions encode the natural conservation law of (1.1) and satisfy much better properties than general weak solutions, especially in the most relevant case of 3D, such as Leray's structure theorem [Ler34], weak-strong uniqueness [Ser62, Ger06, Che11], partial regularity [Str88, CKN82], integrability of higher Sobolev norms [CV14], estimates of potential blowup rates [ESŠ03, GKP16, CZ16]. In fact, it is well-known that Leray-Hopf solutions are smooth and unique in 2D. These nice properties, much desirable from a regularity stand point, make it significantly harder to construct nonunique Leray-Hopf solutions in $d \geq 3$, though partial results [Lad69, JŠ15] and numerical evidence [GŠ17] are available.

The main focus of the present paper is to study the question of uniqueness/nonuniqueness for general weak solutions of (1.1) in the sense of Definition 1.1. Even though the solutions constructed in this paper live on a borderline of a class of Leray-Hopf solutions, they do not have the regularity to justify (1.4) and are not Leray-Hopf solutions; whether Leray-Hopf solutions are unique in dimension $d \geq 3$ remains a challenging open question.

1.2. Uniqueness vs. nonuniqueness. We first discuss the uniqueness results. For brevity, we do not emphasize the underlying spatial domain of the results that we mention. Since we mainly work with the scales of mixed Lebesgue norms $L^p L^q$, to simplify notation, given $p, q \in [1, \infty]$ and $\Omega \subset \mathbb{R}^d$ let us denote the Banach space

$$X^{p,q}([0, T]; \Omega) = \begin{cases} L^p([0, T]; L^q(\Omega)) & \text{if } p \neq \infty, \\ C([0, T]; L^q(\Omega)) & \text{if } p = \infty. \end{cases}$$

In the setting of Leray-Hopf solutions, Prodi [Pro59], Serrin [Ser62], and Ladyzhenskaya [Lad67] proved that if a Leray-Hopf solution u satisfies

$$u \in X^{p,q} \quad \text{for some } p < \infty \text{ and } q > d \text{ such that } \frac{2}{p} + \frac{d}{q} \leq 1,$$

then *all* Leray-Hopf solutions with the same initial data must coincide. Remarkably, the endpoint case $p = \infty$ can be strengthened to $L^\infty L^d$ [LM01, ESŠ03]. This type of results are often referred to as *weak-strong uniqueness* in the sense that if there exists a strong solution, then any weak solution with the same initial data coincides with it. In fact, a membership in such functional classes implies the regularity of Leray-Hopf solutions as well, though we will not go into details in this direction and simply refer interested readers to [Ser62, ESŠ03, GKP16] and references therein.

In the setting of Definition 1.1, one generally loses the property of weak-strong uniqueness due to the lack of the energy inequality (1.2). For such weak solutions, one can instead study the uniqueness issue within certain functional classes, such as $X^{p,q}$. The general scheme is to write (1.3) as the following abstract formulation

$$u = e^{t\Delta} u_0 + B(u, u), \quad (1.5)$$

and study the continuity of the bilinear operator B in the various underlying functional spaces, see for instance [LR16] and references therein. The first result in this direction dates back to Fabes, Jones, and Rivière [FJR72] who proved that weak solutions in the class $X^{p,q}$ are unique if $\frac{2}{p} + \frac{d}{q} \leq 1$ and $d < q < \infty$. The limit case $p = \infty$ and $q = d$ was later covered first by [FLRT00] in $d \geq 2$, and then [Mey97, Mon99, LM01] via different methods and for different spatial domains.

Based on a scaling analysis, when $\frac{2}{p} + \frac{d}{q} = 1$, the space $X^{p,q}$ is invariant under the parabolic scaling of the equation $u \mapsto u_\lambda := \lambda u(\lambda^2 t, \lambda x)$. In the literature, the space $X^{p,q}$ is called *sub-critical* when $\frac{2}{p} + \frac{d}{q} < 1$, *critical* when $\frac{2}{p} + \frac{d}{q} = 1$ and, *super-critical* when $\frac{2}{p} + \frac{d}{q} > 1$. Therefore, the uniqueness condition $u \in X^{p,q}$ for $\frac{2}{p} + \frac{d}{q} \leq 1$ seems to be sharp. In fact, thanks to [FJR72, Kat84, FLRT00, LM01] when $\frac{2}{p} + \frac{d}{q} \leq 1$, all weak solutions (in the

sense of Definition 1.1) belonging to the class $X^{p,q}$ are automatically Leray-Hopf² and hence regular. In other words, sub-critical or critical weak solutions are classical solutions.

We can summarize these uniqueness results as follows. Since these results were originally stated for \mathbb{R}^d , we also include a proof applicable to our specific setup in the appendix for readers' convenience.

Theorem 1.3 ([FJR72, Kat84, FLRT00, LM01]). *Let $d \geq 2$ and u be a weak solution of (1.1) such that $u \in X^{p,q}([0, T]; \mathbb{T}^d)$ for some $p, q \in [1, \infty]$ such that $\frac{2}{p} + \frac{d}{q} \leq 1$. Then*

- (1) u is unique in the class $X^{p,q}$,
- (2) u is a Leray-Hopf solution.

So far, the positive results suggest $\frac{2}{p} + \frac{d}{q} = 1$ as the critical regularity threshold for uniqueness/nonuniqueness of the weak solutions. One would naturally ask what would happen in the super-critical regime $\frac{2}{p} + \frac{d}{q} > 1$, or more specifically, whether the following conjecture is valid.

Conjecture 1.4. *Let $d \geq 2$ and $p, q \in [1, \infty]$ such that $\frac{2}{p} + \frac{d}{q} > 1$. Then*

- (1) *There exist two weak solutions $u, v \in X^{p,q}([0, T]; \mathbb{T}^d)$ of (1.1) such that*

$$u(0) = v(0) \text{ but } v \neq u.$$
- (2) *There exists a weak solution $u \in X^{p,q}([0, T]; \mathbb{T}^d)$ of (1.1) such that u is not Leray-Hopf.*

In stark contrast to the positive result of Theorem 1.3, which has been known for quite some time, Conjecture 1.4 was completely open until very recently due to the groundbreaking work [BV19]. Even though the regularity was very far from the threshold $\frac{2}{p} + \frac{d}{q} = 1$, the nonuniqueness of weak solutions has been shown in dimension $d \geq 3$: $C_t L^{2+}$ [BV19, BCV18] in dimension $d = 3$ and $H^{1/200-}$ [Luo19] in dimension $d \geq 4$. In fact, these works used a unified approach to tackle both parts of Conjecture 1.4 at the same time: one perturbs a given smooth solution of (1.1) to obtain a “wild” solution with certain regularity $X^{p,q}$ and then the existence of such a wild solution implies both the nonuniqueness of weak solutions and the existence of non-Leray-Hopf solutions in the said class.

Unfortunately, the strategy of [BV19], which was in turn developed from a series of works [BDLS15, BDL16, Ise18, BLJV18], breaks down in 2D. The reason is that, roughly speaking, the framework of [BV19] is L^2 critical, in the sense that the mechanism can produce finite energy wild solutions if the system is L^2 super-critical. This heuristic has been confirmed in [BCV18, LT20, LQ20] for the generalized Navier-Stokes equations with fractional dissipation.

Since the 2D Navier-Stokes equations is L^2 -critical, there are no nonuniqueness results for the 2D case to date. In fact, a direct corollary of [BV19] is false in 2D: any $C_t L^2$ weak solution of the 2D Navier-Stokes equations is Leray-Hopf, and hence smooth and unique. One of the main results in this paper is to show that the nonuniqueness of weak solutions holds even in 2D.

Theorem 1.5 (Strong nonuniqueness in 2D). *Let $d = 2$ be the dimension. Every weak solution of (1.1) is not unique: for any weak solution $v : [0, T] \times \mathbb{T}^2 \rightarrow \mathbb{R}^2$, there exists a different weak solution u with the same initial data.*

It is worth noting that the nonuniqueness is proved here in a stronger sense than [BV19, Luo19], namely that *every* solution is nonunique in the class of weak solutions. We can classify different types of nonuniqueness results as follows. Here, X denotes different functional classes of weak solutions.

- “Weak nonuniqueness”: there exists a nonunique weak solution in the class X .
- “Strong nonuniqueness”: any weak solution in the class X is nonunique.

Under this classification, currently the only strong nonuniqueness available is [BCV18] for 3D, where X can be taken as $C_t H^\varepsilon$ weak solutions with intervals of regularity for a small $\varepsilon > 0$. In fact, the main results of [BCV18] imply the strong uniqueness of weak solutions on \mathbb{T}^d for dimension $d = 3, 4$ since Leray-Hopf solutions in $d \leq 4$ have intervals of regularity.

In this spirit, we prove a strong nonuniqueness result in a class of $L_t^p L^\infty$ weak solutions, for any $p < 2$ and in any dimension $d \geq 2$, which is sharp in view of Theorem 1.3. A detailed list of the properties of constructed solutions can be found in Theorem 1.7 below. In particular, the below theorem settles Conjecture 1.4 in the case $q = \infty$.

Theorem 1.6 (Sharp nonuniqueness in $d \geq 2$). *Let $d \geq 2$ be the dimension and $1 \leq p < 2$.*

²In contrast to [FJR72, Kat84, FLRT00] where one has to assume L^2 initial data, our setting is on the d -dimensional torus and the initial data is always L^2 .

- (1) A weak solution $u \in L^p(0, T; L^\infty(\mathbb{T}^d))$ of (1.1) is not unique in the class $L^p(0, T; L^\infty(\mathbb{T}^d))$ if u has at least one interval of regularity³.
- (2) There exist non-Leray-Hopf weak solutions $u \in L^p(0, T; L^\infty(\mathbb{T}^d))$.

In view of Theorem 1.3, nonunique solutions cannot live in the class $L_t^2 L^\infty$. However, Theorem 1.6 shows the existence of nonunique solutions on the borderline of this Leray-Hopf class and raises the question of whether such constructions can be extended to the Leray-Hopf solutions.

One clearly sees that Theorem 1.6 only implies the sharpness of [FJR72] near $q = \infty$ and the rest of the borderline regime remains open, see the discussion at the end of the introduction. Alternatively, we can present the result as: for any smooth initial data, there are infinitely many weak solutions with regularity $L_t^p L^\infty$ emerging from the same data and agrees with each other for a short time, see the proof of Theorem 1.10 at the end of Section 2.

1.3. The main theorem and intervals of regularity. We now present the main theorem of the paper, which implies immediately Theorem 1.5 and 1.6 above as we will show in Section 2. One of the most interesting features of our constructed weak solutions is that they possess intervals of regularity, i.e. they are classical solutions on many sub-intervals whose union occupies a majority of the time axis, cf. [Ler34] and [BCV18], which we will discuss in detail towards the end of the introduction.

Theorem 1.7. *Let $d \geq 2$ be the dimension and $1 \leq p < 2$, $q < \infty$, and $\varepsilon > 0$. For any smooth, divergence-free vector field $v \in C^\infty([0, T] \times \mathbb{T}^d)$ with zero spatial mean for each $t \in [0, T]$, there exists a weak solution u of (1.1) and a set*

$$\mathcal{I} = \bigcup_{i=1}^{\infty} (a_i, b_i) \subset [0, T],$$

such that the following holds.

- (1) The solution u satisfies

$$u \in L^p(0, T; L^\infty(\mathbb{T}^d)) \cap L^1(0, T; W^{1,q}(\mathbb{T}^d)).$$

- (2) u is a smooth solution on (a_i, b_i) for every i . Namely,

$$u|_{\mathcal{I} \times \mathbb{T}^d} \in C^\infty(\mathcal{I} \times \mathbb{T}^d).$$

In addition, u agrees with the smooth solution emerging from the initial data $v(0)$ near $t = 0$.

- (3) The Hausdorff dimension of the residue set $\mathcal{S} = [0, T] \setminus \mathcal{I}$ satisfies

$$d_{\mathcal{H}}(\mathcal{S}) \leq \varepsilon.$$

- (4) The solution u and the given vector field v are ε -close in $L_t^p L^\infty \cap L_t^1 W^{1,q}$:

$$\|u - v\|_{L^p(0, T; L^\infty(\mathbb{T}^d)) \cap L^1(0, T; W^{1,q}(\mathbb{T}^d))} \leq \varepsilon.$$

Remark 1.8. We list a few remarks here concerning the above result.

- (1) In terms of the scaling, u also lies on the borderline of the Beale-Kato-Majda criterion [BKM84] which scales as $L_t^1 W^{1,\infty}$.
- (2) The residue set $\mathcal{S} = [0, T] \setminus \mathcal{I}$ is a singular set in the sense that for any $t \notin \mathcal{S}$, there is $\delta > 0$ such that $u \in C^\infty((t - \delta, t + \delta) \times \mathbb{T}^d)$.
- (3) Since the solution $u(t)$ is smooth on (a_i, b_i) , the energy equality is satisfied

$$\frac{1}{2} \|u(t_1)\|_2^2 + \int_{t_0}^{t_1} \|\nabla u(s)\|_2^2 ds = \frac{1}{2} \|u(t_0)\|_2^2 \quad \text{for all } t_0, t_1 \in (a_i, b_i).$$

- (4) The driving mechanism of nonuniqueness is a result of large chunks of mass emerging from/escaping to finer time scales. There is no blowup on each intervals of regularity (a_i, b_i) but norms do blow up as $i \rightarrow \infty$.

³In particular, it can be applied to a smooth solution to obtain a nonunique weak solution in the class $L_t^p L^\infty$.

1.4. Applications to the Euler equations. Our results also apply to the inviscid case with no changes in contrast to a recent work [BCV18], which relies heavily on the parabolic regularization.

Theorem 1.9. *Theorem 1.7 also holds for the Euler equations. Namely, under the same assumptions, there exists a weak solution u of the Euler equations satisfying the same properties.*

As a byproduct of the construction, we provide an improvement to the Onsager’s conjecture in 2D in the negative direction, where the best result currently stands at $L_t^\infty C^{1/5-\varepsilon}$ [BDLIS15], see also [BSV19, pp. 1817] and [Nov20].

Theorem 1.10. *Let $d \geq 2$ be the dimension and $\varepsilon > 0$. There exist infinitely many non-conserving weak solutions $u \in L_t^{\frac{3}{2}-\varepsilon} C^{\frac{1}{3}} \cap L_t^1 C^{1-\varepsilon}$ with the same initial data.*

This result appears to establish the first non-conserving solutions with an exact “ $\frac{1}{3}$ -Hölder regularity” in space, albeit with a non-optimal $L^{3/2-}$ exponent in time, cf. positive results [CET94, CCFS08]. We also note that previous constructions [BDLIS15, BDLS16, Ise18, BLJV18] are continuous in space-time whereas our solutions are not, and in fact, the kinetic energy of our solutions becomes unbounded in a piece-wise constant fashion.

1.5. Main ideas of the construction. The construction in Theorem 1.7 is based on an iterative scheme to obtain suitable approximate solutions to (1.1) that consists of two main steps. The first step is to concentrate the stress error of the approximate solutions to many smaller sub-intervals, allowing us to achieve a small Hausdorff dimension in time of the singular set. In particular, this ensures the final approximate solution is an exact smooth solution on many small intervals. Crucially, this is done by adding a very small corrector to the existing solution while keeping the size of the stress error unchanged L^1 in time up to a constant multiple. After this concentration procedure, the stress error is zero on a large subset of the time interval and thus the size of the concentrated stress error is much larger on its support set.

The second step uses a convex integration scheme to add another perturbation to the concentrated solution, reducing the size of the stress error. This convex integration technique has been developed over the last decades, see for instance [DLS09, DLS13, BDLIS15, BLJV18, Ise18, BV19, Luo19] and references therein, since its inception to fluid dynamics in [DLS09]. In particular, its latest iteration for the transport equation in [CL20] allows us to achieve a very high level of temporal concentration of the perturbation in the sense that higher Sobolev norms blow up while their time averages remain bounded. The introduction of temporal concentration in the convex integration scheme allows us to trade temporal integrability for spatial regularity, answering a question raised in [BV21, Problem 4.4]. To avoid a dimensional loss, the “building blocks” used in the scheme are almost spatially homogeneous, in stark contrast to [BV19, Luo19, BCV18].

The most difficult and important part of the iterative scheme is ensuring that the perturbation w satisfies the regularity $w \in L_t^p L^\infty \cap L_t^1 W^{1,q}$, while at the same time successfully reducing the size of the stress error. This boils down to balancing four different aspects of the perturbation: temporal and spatial oscillation/concentration. In the present work, the leading order effects are temporal concentration and spatial oscillation, whereas temporal oscillation and spatial concentration effectively play no role and are kept to a minimum.

1.6. Comparison with previous works. In the last part of the introduction, we compare our main results to the previous works and list a few open questions. We divide the discussion into three topics as follows.

Regularity threshold for uniqueness/nonuniqueness. The first nonuniqueness result for the Navier-Stokes system was established in [BV19] by Buckmaster and Vicol, where finite energy nonunique weak solutions were constructed in 3D. Even though the nonuniqueness is only proved in $C_t L^2$, the iteration scheme in [BV19] allows for a very small regularity H^ε for $\varepsilon \ll 1$, which was then used in [BCV18] to show nonuniqueness at such a regularity. The work [Luo19] built upon the observation that in higher dimension, weak solutions can be less intermittent and thus the regularity of nonuniqueness was improved to $H^{1/200-}$ for $d \geq 4$. In fact, as noted in [Tao19], in very high dimension one can show nonuniqueness in $C_t H^{1/2-}$ or $H^{1/2-}$ in the stationary case, although the regularity $H^{1/2-}$ is still very far from the critical scale $H^{\frac{d-2}{2}}$ or L^d .

Below we compare different results using the scales of space-time Lebesgue spaces $X^{p,q}$.

Results	Category	Scaling	Range
Leray-Hopf solutions	Existence	$\frac{2}{p} + \frac{d}{q} = \frac{d}{2}$	$q \geq 2$
[BV19]	Nonuniqueness	$\frac{2}{p} + \frac{d}{q} = \frac{d}{2}$	$q = 2$ and $d = 3$
[BCV18]	Nonuniqueness	$\frac{2}{p} + \frac{d}{q} = \frac{d}{2} - \varepsilon$	$q = 2+$ and $d = 3$
[Luo19]	Nonuniqueness	$\frac{2}{p} + \frac{d}{q} = \frac{d}{2} - \frac{1}{200}$	$q = 2+$ and $d \geq 4$
Theorem 1.7	Nonuniqueness	$\frac{2}{p} + \frac{d}{q} = 1 + \varepsilon$	$q = \infty$
Theorem 1.3	Uniqueness	$\frac{2}{p} + \frac{d}{q} = 1$	$q \leq \infty$

In light of the current state, we expect the nonuniqueness of weak solutions continue to hold in the full range of the super-critical regime $\frac{2}{p} + \frac{d}{q} > 1$. Unfortunately, the method developed in this paper heavily relies on the constraint $p < 2$ ($q = \infty$) and is not able to achieve the nonuniqueness of weak solutions in $X^{p,q}$ for $p \geq 2$ and $q \geq 2$.

Size of the potential singular set. Here we discuss our result in the context of partial regularity, more specifically, the size of the singular set in time or in space-time. By singular times we mean the union of times at which the solution is not locally smooth, while singular points in space-time refer to points (t, x) where the solution is not locally bounded (in the sense of ess sup).

By the classical results of Leray, in 3D the Hausdorff dimension of possible singular times of a Leray-Hopf solution is bounded by $1/2^4$. A key step in understanding the (possible) singular set of weak solutions was made by Scheffer [Sch76a, Sch77, Sch80] where the notion of suitable weak solutions was introduced. It was proved in [Sch80] that the singular sets of these suitable weak solutions have finite $\frac{5}{3}$ -dimensional Hausdorff measure in space-time. The theory of partial regularity culminated with the work [CKN82] by Caffarelli, Kohn, and Nirenberg where they show that $\mathcal{P}^1(S) = 0$, i.e., the 1-dimensional parabolic Hausdorff measure of the singular set in space-time is zero. Note that these partial regularity results only provide upper bounds on the potential singular sets.

While convincing evidence [Sch85, Sch87, Oza20] suggests that the upper bound of 1-dimensional parabolic singularities in 3D is likely to be sharp for suitable weak solutions, it was unknown whether there are weak solutions with a nontrivial⁵ singular set until the work [BCV18] where the authors constructed wild solutions with a nonempty singular set with a dimension strictly less than 1. As in [BCV18], solutions constructed here are not Leray-Hopf; however, they constitute the first example of 3D weak solutions that surpass the $1/2$ upper bound (with a nonempty singular set).

In dimension $d \geq 4$, the existence of partially regular (in space-time or in time) weak solutions becomes highly nontrivial. In fact, Leray's structure theorem only holds up to $d = 4$ and the local energy inequality, a key ingredient in the partial regularity theory, remains absent in $d \geq 4$ [DD07, Remark 1.1]. Despite such a difficulty, the existence of partially regular weak solutions in space-time was established in 4D [Sch77] by Scheffer and also a recent preprint [Wu20] by Wu. In dimension $d \geq 5$ the existence of partially regular weak solutions (in space-time or in time) was unknown to our knowledge and Theorem 1.7 appears to be the first example of weak solutions with partial regularity in time in dimension $d \geq 5$.

In relation to the partial regularity in space-time, the singular set of our solutions is the whole spatial domain at each singular time, as with all the other constructions exploiting a convex integration scheme. It might be possible to construct wild solutions that enjoy a certain space-time partial regularity by a space-time variant of the concentration procedure used here.

Anomalous dissipation of the Euler equations. A recent milestone in incompressible fluid dynamics is the resolution of the Onsager conjecture [Ons49] which states that $\frac{1}{3}$ -Hölder is the critical threshold for energy conservation for the 3D Euler equations. While the positive direction was settled in the 90s in [CET94] following the first attempt by [Eyi94] and then later refined in [CCFS08], the negative part was significantly harder and the regularity of counterexamples [Sch93, Shn97] was far below the threshold. Advances in the negative direction really took off with the modern convex integration approach starting with the seminal paper of De Lellis and Székelyhidi Jr. [DLS09]. The approach of using convex integration was refined and improved in a series of works [DLS13, DLS14, BDLIS15, BDLS16]. Building upon these works, the threshold $C_t C^{1/3-}$ was finally reached by Isett [Ise18], see also [BLJV18].

⁴This interpretation was made explicit in [Sch76b].

⁵Here by nontrivial we mean that the singular set is not empty or full since smooth solutions have no singularity while the singular set of the solutions in [BV19] is the whole space-time domain.

So far, constructed anomalous weak solutions have a limited regularity on the whole time axis, namely, the Hölder regularity in space is always below $\frac{1}{3}$. The works [CET94, CCFS08] suggest that insisting on the exact “ $\frac{1}{3}$ -Hölder regularity” in space leads to L^3 being the right critical scale in time for the energy conservation. This exact “ $\frac{1}{3}$ -Hölder regularity” seems to be out of reach for the previous Euler schemes, an issue that has been investigated recently by Isett [Ise17].

Even though our inviscid solutions have a worse global-in-time regularity, they are smooth solutions on a “large” portion of the time axis, and hence the kinetic energy is conserved locally in time. The mechanisms of the failure of the energy conservation are completely different: fast spatial oscillations play a key role in the previous Euler examples, whereas a strong temporal concentration here causes the breakdown at small time scales. It would be very interesting to combine the previous Euler results with the current paper to show that there exist “wild solutions” in $C_t C^{\frac{1}{3}-}$ or $L_t^{3-} C^{\frac{1}{3}}$ that are locally smooth in time away from a small singular set.

1.7. Notations. For reader’s convenience, we collect the notations used throughout the manuscript.

- $\mathbb{T}^d = \mathbb{R}^d / \mathbb{Z}^d$ is the d -dimensional torus and is identified with $[0, 1]^d$. For any function $f : \mathbb{T}^d \rightarrow \mathbb{R}$ we denote by $f(\sigma \cdot)$ the $\sigma^{-1}\mathbb{T}^d$ -periodic function $f(\sigma x)$. The space $C_0^\infty(\mathbb{T}^d)$ is the set of periodic smooth functions with zero mean and $C_0^\infty(\mathbb{T}^d, \mathbb{R}^d)$ is the set of periodic smooth vector fields with zero mean.
- The Lebesgue space is denoted by L^p . For any $f \in L^1(\mathbb{T}^d)$, its spacial average is

$$\int_{\mathbb{T}^d} f dx = \int_{\mathbb{T}^d} f dx.$$

For any function $f : [0, T] \times \mathbb{T}^d \rightarrow \mathbb{R}$, denote by $\|f(t)\|_p$ the Lebesgue norm on \mathbb{T}^d (in space only) at a fixed time t . If the norm is taken in space-time, we use $\|f\|_{L_{t,x}^p}$.

- The tensor divergence $\operatorname{div} A = \partial_j A_{ij}$ for any matrix-valued function $A : \mathbb{T}^d \rightarrow \mathbb{R}^{d \times d}$ and the tensor product $f \otimes g = f_i g_j$ for any two vectors $f, g \in \mathbb{R}^d$. The notion ∇ indicates full differentiation in space only, and space-time gradient is denoted by $\nabla_{t,x}$.
- For any Banach space X , the Bochner space $L^p(0, T; X)$ is equipped with the norm

$$\left(\int_0^T \|\cdot\|_X^p dt \right)^{\frac{1}{p}},$$

and we often use the short notations $L_t^p X$ and $\|\cdot\|_{L_t^p X}$.

- We write $X \lesssim Y$ if there exists a constant $C > 0$ independent of X and Y such that $X \leq CY$. If the constant C depends on quantities a_1, a_2, \dots, a_n we will write $X \lesssim_{a_1, \dots, a_n} Y$ or $X \leq C_{a_1, \dots, a_n} Y$.

1.8. Organization of the paper. The organization of the rest of the paper is as follows.

The outline of construction is given in Section 2, where main theorems will be proved assuming the main proposition of the paper, proposition 2.2. The proof of the main proposition is the content of the rest of the paper: we concentrate the stress error to many small sub-intervals in Section 3, design a velocity perturbation using convex integration to obtain a new solution pair (u_1, R_1) in Section 4, and finally estimate the perturbation along with the new stress error to conclude the proof in Section 5.

Appendix A includes a proof of Theorem 1.3. Appendix B contains some technical tools used in the paper, namely an improved Hölder’s inequality and antidivergence operators on \mathbb{T}^d .

2. OUTLINE OF THE PROOF

The proof of Theorem 1.7 consists of a iterative scheme, which is achieved by repeatedly applying the main proposition of this paper, Proposition 2.2 to obtain a sequence of solutions (u_n, R_n) to (2.1). The proof mainly consists of three goals:

- (1) The convergence of $u_n \rightarrow u$ in $L_{t,x}^2$ and $R_n \rightarrow 0$ in $L_{t,x}^1$ so that u is a weak solution of (1.1).
- (2) Ensuring the final solution verifies $u \in L_t^p L^\infty \cap L_t^1 W^{1,q}$ for $p < 2$ and $q < \infty$.
- (3) Achieving a small dimension of the singular set of u in time.

To this end, we employ a two-step approach:

- Step 1: $(u_n, R_n) \xrightarrow{\text{concentrating the stress error}} (\bar{u}_n, \bar{R}_n)$

- Step 2: $(\bar{u}_n, \bar{R}_n) \xrightarrow{\text{space-time convex integration}} (u_{n+1}, R_{n+1})$

where the first step is mainly for achieving a small singular set in time and the second step is to ensure the convergence of R_n .

2.1. The Navier-Stokes-Reynolds system. Let us first introduce the approximate equations of (1.1) for our approximate solutions. These approximate solutions solved the so-called Navier-Stokes-Reynolds systems

$$\begin{cases} \partial_t u - \Delta u + \operatorname{div}(u \otimes u) + \nabla p = \operatorname{div} R \\ \operatorname{div} u = 0 \end{cases} \quad (2.1)$$

where $R : [0, T] \times \mathbb{T}^d \rightarrow \mathcal{S}_0^{d \times d}$ is a traceless symmetric matrix called Reynolds stress.

Since the associated pressure p can be uniquely determined by the elliptic equation:

$$\Delta p = \operatorname{div} \operatorname{div} R - \operatorname{div} \operatorname{div}(u \otimes u) = \partial_i \partial_j (R_{ij} - u_i u_j),$$

throughout the paper, we denote the solution of (2.1) by (u, R) .

This system (2.1) arises naturally when studying weak solutions of the Navier-Stokes equations. The Reynolds stress R emerges as the noncommutativity between average ensembles and the quadratic nonlinearity.

In the inviscid case, we can just drop the Laplacian term in (2.1) and the system becomes the so-called Euler-Reynolds equations, which was widely used in constructing non-conserving weak solutions of the Euler equations in the context of Onsager's conjecture [DLS13, DLS14, BDLIS15, BDLS16, Ise18, BLJV18]. Since the Laplacian plays no role in our construction, in what follows we simply use (2.1).

2.2. Concentrating the stress error. As stated in the introduction, we proceed with two steps to prove this main proposition. The first step is a procedure that concentrates the stress error into many smaller sub-intervals.

Given (u_{n-1}, R_{n-1}) , we divide the time interval $[0, T]$ into smaller sub-intervals I_i of length $\tau^\varepsilon > 0$, where $\tau > 0$ will be chosen to be very small depending on (u_{n-1}, R_{n-1}) . So the total number of sub-intervals is $\sim \tau^{-\varepsilon}$.

On each sub-interval I_i , we solve a generalized Navier-Stokes equations linearized around (u_{n-1}, R_{n-1}) to obtain a corrector v_i on I_i . More precisely, $v_i : I_i \times \mathbb{T}^d \rightarrow \mathbb{R}^d$ solves

$$\begin{cases} \partial_t v_i - \Delta v_i + \operatorname{div}(v_i \otimes v_i) + \operatorname{div}(v_i \otimes u) + \operatorname{div}(u \otimes v_i) + \nabla q_i = -\operatorname{div} R \\ \operatorname{div} v_i = 0 \\ v_i(t_i) = 0. \end{cases} \quad (2.2)$$

so that $u_{n-1} + v_i$ is an exact solution (of the Navier-Stokes equations) on I_i .

To concentrate the error and obtain a solution on $[0, T]$, we apply a sharp cutoff χ_i to the corrector v_i and obtain the glued solution \bar{u}_{n-1} defined by

$$\bar{u}_{n-1} := u_{n-1} + \sum_i \chi_i v_i.$$

Specifically, each χ_i equals 1 on a majority of the sub-interval I_i , but $\chi = 0$ near endpoints of each I_i of scale $\sim \tau$. Since $\varepsilon \ll 1$, the cutoff χ_i is very sharp when comparing to the length of the sub-interval I_i .

On one hand, due to the sharp cutoff χ_i , the stress error \bar{R}_n associated with \bar{u}_n will only be supported near the endpoints of I_i of time scale τ . In other words, the temporal support of \bar{u}_n can be covered by $\sim \tau^{-\varepsilon}$ many intervals of size $\sim \tau$, from which one can already see a small dimension of the singular set of the final solution.

On the other hand, the corrector v_i is very small, say in $L_t^\infty H^d$, since it starts with initial data 0 and we can choose time scale $\tau^\varepsilon = |I_i|$ to be sufficiently small. More importantly, the new stress error \bar{R}_{n-1} associated with \bar{u}_{n-1} satisfies the estimate

$$\|\bar{R}_{n-1}\|_{L_t^1 L^r} \lesssim \|R_{n-1}\|_{L_t^1 L^r} \quad \text{for } 1 < r < \infty,$$

with an implicit constant independent of the time scale $\tau > 0$. In other words, concentrating the stress error R_{n-1} to \bar{R}_{n-1} cost a loss of a constant multiple when measuring in L^1 norm in time.

2.3. Space-time convex integration. The next step is to use a convex integration technique to reduce to size of \overline{R}_{n-1} by adding further a perturbation w_n to \overline{u}_{n-1} to obtain a new solution (u_n, R_n) of (2.1). The perturbation w_n and the new stress R_n satisfies the equation

$$\operatorname{div} R_n = \operatorname{div} \overline{R}_{n-1} + \operatorname{div}(w_n \otimes w_n) + \partial_t w_n - \Delta w_n + \operatorname{div}(\overline{u}_{n-1} \otimes w_n) + \operatorname{div}(w_n \otimes \overline{u}_{n-1}) + \nabla P_n,$$

for a suitable pressure P_n . The heuristic is that the high-high to low cascade in space-time of $w_n \otimes w_n$ can balance the old stress error \overline{R}_{n-1} in the sense that

$$\operatorname{div}(\overline{R}_{n-1} + w_n \otimes w_n) = \text{High Spacial Freq. Term} + \text{High Temporal Freq. Term} + \text{Lower Order Terms}, \quad (2.3)$$

where the ‘‘High Temporal Freq. Term’’ above will further be balanced by a part of $\partial_t w_n$, as in [BV19, BCV18] and [CL20]. However, one of the fundamental differences to [BV19, BCV18] is that this additional ‘‘convex integration in time’’ requires no additional constraint of oscillation and concentration and is basically free, which is crucial to obtain the sharp regularity $L_t^p L^\infty \cap L_t^1 W^{1,q}$.

In particular, executing the scheme of [CL20] requires two ingredients:

- (1) Suitable stationary flows as the spatial building blocks that can achieve some level of spatial concentration.
- (2) The use of intermittent temporal functions to oscillate the spatial building blocks in time.

Once (1) is available, it is relatively straightforward to implement (2). On the technical side, we use the stationary Mikado flows introduced in [DS17] as the spatial building blocks. These are periodic pipe flows that can be arranged to supported on periodic cylinders with a small radius. In other words, Mikado flows can achieved a $d-1$ -dimensional concentration on \mathbb{T}^d , which is more than enough in view of (1). It is worth noting that in the framework of [BV19], stationary Mikado flows are not sufficiently intermittent to be used for the Navier-Stokes equations in dimension $d \leq 3$, cf. [Luo19].

The space-time cascade (2.3) imposed a relation between the perturbation w_n and the stress error \overline{R}_{n-1} as

$$\|w_n\|_{L_{t,x}^2} \sim \|\overline{R}_{n-1}\|_{L_{t,x}^1}. \quad (2.4)$$

The relation (2.4) will imply the convergence in $L_{t,x}^2$ of the approximate solutions u_n as long as one can successfully reduce the size of the stress error

$$\|R_n\|_{L_{t,x}^1} \ll \|\overline{R}_{n-1}\|_{L_{t,x}^1}. \quad (2.5)$$

In particular, special attention will be paid to estimating the temporal derivative part of the new stress error

$$\operatorname{div} R_{\text{tem}} = \partial_t w_n, \quad (2.6)$$

and achieving the regularity of the perturbation

$$\|w_n\|_{L^p L^\infty} + \|w_n\|_{L^1 W^{1,q}} \ll 1. \quad (2.7)$$

These two constraints (2.6) and (2.7) require a very delicate choice of parameters when designing the perturbation w_n . On one hand, (2.6) implies the temporal frequency can not be too large, relative to the spatial frequency, otherwise the time derivative will dominate. On the other hand, (2.7) requires a large temporal frequency so that temporal concentration can offset the loss caused by going from L^2 to L^∞ or $W^{1,q}$ in space in relation to (2.4).

Nevertheless, it turns out that the scheme in [CL20] is flexible enough to accommodate (2.6) and (2.7). We could somehow explain why this is possible. Roughly speaking, the method in [CL20] is $L_{t,x}^2$ -critical. While it is difficult for w_n to go above the $L_{t,x}^2$ regularity, we trade L^2 for L^p (resp. L^1) in time to obtain an improvement of L^2 to L^∞ (resp. $W^{1,q}$) in space. We provide a scaling analysis below.

2.4. Oscillation and concentration. We do this computation in general dimension $d \geq 2$ and $D \in [0, d]$ denotes the spatial intermittency, cf. [Man76, Fri95].

We start with a velocity perturbation in $L_{t,x}^2$ with a certain decay given by the previous stress error,

$$\|w_n\|_{L_{t,x}^2} \lesssim 1.$$

Denote the spacial frequency by λ and the temporal frequency by κ , namely

$$\|\partial_t^m \nabla^n w\|_{L_{t,x}^2} \lesssim \kappa^m \lambda^n.$$

The intermittency parameter D in space dictates the concentration level of w_n and the scaling law

$$\|w_n(t)\|_{L^p} \lesssim \|w_n(t)\|_{L^2} \lambda^{(d-D)(\frac{1}{2}-\frac{1}{p})}. \quad (2.8)$$

As for the temporal scaling, we assume for simplicity a full concentration in time:

$$\|w_n\|_{L^q L^p} \lesssim \|w_n\|_{L^2 L^p} \kappa^{\frac{1}{2}-\frac{1}{q}} \sim \kappa^{\frac{1}{2}-\frac{1}{q}} \lambda^{(d-D)(\frac{1}{2}-\frac{1}{p})}. \quad (2.9)$$

With such scaling laws, we effectively assume a negligible amount of temporal oscillation and the goal then boils down to finding a working choice of D in terms of the given parameters d, p, q . In other words, we need to find a balance between spatial oscillation and spacial concentration.

By the scaling relations (2.8) and (2.9), the stress error contributed by the time derivative (2.6) satisfies

$$\|\operatorname{div}^{-1}(\partial_t w_n)\|_{L^1_{t,x}} \lesssim \kappa^{\frac{1}{2}} \lambda^{-1} \lambda^{-\frac{d-D}{2}} \quad (2.10)$$

where we assume div^{-1} gains one full derivative in space. The regularity condition (2.7) becomes

$$\|w_n\|_{L^p L^\infty} \sim \kappa^{\frac{1}{2}-\frac{1}{p}} \lambda^{\frac{d-D}{2}} \ll 1, \quad (2.11)$$

and

$$\|w_n\|_{L^1 W^{1,q}} \sim \kappa^{-\frac{1}{2}} \lambda^{1+\frac{d-D}{2}-\frac{d-D}{q}} \ll 1. \quad (2.12)$$

In particular, (2.10) and (2.12) imply that

$$\lambda^{1+\frac{d-D}{2}-\frac{d-D}{q}} \ll \kappa^{\frac{1}{2}} \ll \lambda^{1+\frac{d-D}{2}}, \quad (2.13)$$

which always has some room since $q < \infty$. Then all we need to is to choose D to ensure (2.11) and (2.13). One can already see that D should be very close to d , which means we need much more spatial oscillation than spatial concentration. Indeed, solutions to (2.11) and (2.13) do exist and we refers to Section 5.1 for the exact choice used.

2.5. The main iteration proposition. We are ready to introduce the main iteration proposition of the paper that materializes the above discussion. To simplify presentation, let us introduce the notion of well-preparedness of solutions to (2.1), which encodes the small Hausdorff dimension of the singular set in time. Throughout the paper, we take $T = 1$ and assume $0 < \varepsilon < 1$ without loss of generality.

Definition 2.1. *We say a smooth solution (u, R) of (2.1) on $[0, 1]$ is well-prepared if there exist a set I and a length scale $\tau > 0$ such that I is a union of at most $\tau^{-\varepsilon}$ many closed intervals of length 5τ and*

$$R(t, x) = 0 \quad \text{if} \quad \operatorname{dist}(t, I^c) \leq \tau.$$

With this definition, to ensure the solution u has intervals of regularity with a small residue set of Hausdorff dimension $\lesssim \varepsilon$, it suffices to construct approximate solutions (u_n, R_n) that are well-prepared for some I_n and τ_n such that

$$I_n \subset I_{n-1} \quad \text{and} \quad \tau_n \rightarrow 0.$$

The main proposition of this paper states as follows.

Proposition 2.2 (Main iteration). *For any $\varepsilon > 0$ and $p < 2$, there exists a universal constant $M = M(\varepsilon, p) > 0$ and $r > 1$ depending only on p and q such that the following holds.*

Let $\delta > 0$ and (u, R) be a well-prepared smooth solution of (2.1) for some set \tilde{I} and a length scale $\tilde{\tau} > 0$. Then there exists another well-prepared smooth solution (u_1, R_1) of (2.1) for some set $I \subset \tilde{I}$ with $0, 1 \notin I$ and $\tau < \tilde{\tau}/2$ such that

$$\|R_1\|_{L^1(0,1;L^r(\mathbb{T}^d))} \leq \delta.$$

Moreover, the velocity perturbation $w := u_1 - u$ satisfies

(1) *The temporal support of w is contained in I , i.e.*

$$\operatorname{supp} w \subset I \times \mathbb{T}^d.$$

(2) *The $L^2 L^2$ estimate,*

$$\|w\|_{L^2([0,1] \times \mathbb{T}^d)} \leq M \|R\|_{L^1([0,1] \times \mathbb{T}^d)};$$

(3) *The $L^p L^\infty \cap L^1 W^{1,q}$ estimate,*

$$\|w\|_{L^p(0,1;L^\infty(\mathbb{T}^d))} + \|w\|_{L^1(0,1;W^{1,q}(\mathbb{T}^d))} \leq \delta;$$

Remark 2.3. *We list a few comments concerning the main proposition.*

(1) *It will be clear that the construction adapts no change for the Euler equations, except dropping the Laplacian.*

- (2) The parameter $r > 1$ is used to ensure the L^r boundedness of the Calderón-Zygmund singular integral and is very close to 1.
- (3) Due to the local well-preparedness, on large portions of the time axis, the solutions are exact solutions of the Navier-Stokes equations (or the Euler equations) and we do not touch them in the future.

We will break down Proposition 2.2 into two separate propositions, whose proof will be the context of Section 3 and respectively Section 4 and Section 5.

2.6. Proof of main theorems. We first deduce Theorem 1.5 and Theorem 1.6 from Theorem 1.7.

Proof of Theorem 1.5. Given a weak solution v (in the sense of Definition 1.1) of the 2D Navier-Stokes equations on $[0, T]$, we need to construct a different weak solution u with the same initial data. For brevity, we take $T = 1$.

Case 1: v is a Leray-Hopf solution.

Since v is Leray-Hopf on $[0, 1]$, v is in fact smooth on $(0, 1]$. Let $\tilde{v} : [1/2, 1] \times \mathbb{T}^2 \rightarrow \mathbb{R}^2$ be a smooth divergence-free vector field that coincide with v on $[1/2, 3/4]$ but

$$\|\tilde{v} - v\|_{L^p([1/2, 1]; L^\infty(\mathbb{T}^d))} \geq 1. \quad (2.14)$$

Then we can apply Theorem 1.7 for \tilde{v} and $\varepsilon < 1$ to obtain a weak solution \tilde{u} on $[1/2, 1] \times \mathbb{T}^2$. The conclusion follows once we define the weak solution $u : [0, 1] \times \mathbb{T}^2 \rightarrow \mathbb{R}^2$ by

$$u = \begin{cases} v & \text{if } t \in [0, 1/2] \\ \tilde{u} & \text{if } t \in [1/2, 1]. \end{cases}$$

Indeed, the new glued solution u is still a weak solution of (1.1) and moreover, by (2.14)

$$\|u - v\|_{L^p([1/2, 1]; L^\infty)} \geq \|\tilde{v} - v\|_{L^p([1/2, 1]; L^\infty)} - \|\tilde{u} - \tilde{v}\|_{L^p([1/2, 1]; L^\infty)} \geq 1 - \varepsilon > 0,$$

which implies $u \neq v$.

Case 2: v is not Leray-Hopf.

Since the initial data $v_0 \in L^2(\mathbb{T}^2)$, we may just use v_0 as initial data to obtain a Leray-Hopf solution u on $[0, 1]$ which is different from v . □

Proof of Theorem 1.6. To prove both points, it suffices to show that given a weak solution $v \in L^p(0, 1; L^\infty(\mathbb{T}^d))$ of (1.1) with at least one interval of regularity, there exists a non-Leray-Hopf weak solution $u \in L^p(0, 1; L^\infty(\mathbb{T}^d))$ having intervals of regularity and with the same initial data.

Let $[a, b]$ be an interval of regularity of the weak solution v . We first choose a smooth, divergence-free vector field $\tilde{v} : [a, 1] \times \mathbb{T}^d \rightarrow \mathbb{R}^d$ such that

$$\tilde{v}|_{[a, b] \times \mathbb{T}^d} \equiv v|_{[a, b] \times \mathbb{T}^d},$$

but

$$\|\tilde{v} - v\|_{L^p([a, 1]; L^\infty(\mathbb{T}^d))} \geq 1 \quad \text{and} \quad \|\tilde{v}\|_{L^p([a, 1]; L^2(\mathbb{T}^d))} \geq 1 + \|v(0)\|_{L^2(\mathbb{T}^d)}, \quad (2.15)$$

As in the proof of Theorem 1.5, we apply Theorem 1.7 to \tilde{v} and $\varepsilon < 1$ to obtain a weak solution $\tilde{u} \in L^p(a, 1; L^\infty(\mathbb{T}^d))$ such that

$$\|\tilde{u} - \tilde{v}\|_{L^p([a, 1]; L^\infty(\mathbb{T}^d))} \leq \varepsilon. \quad (2.16)$$

We then define a new solution $u : [0, 1] \times \mathbb{T}^d \rightarrow \mathbb{R}^d$ by

$$u = \begin{cases} v & \text{if } t \in [0, a] \\ \tilde{u} & \text{if } t \in [a, 1]. \end{cases}$$

The glued solution $u \in L^p L^\infty$ is still a weak solution due the smoothness of both v and \tilde{u} near $t = a$. Next, u and v are different since

$$\|u - v\|_{L^p L^\infty} = \|\tilde{u} - v\|_{L^p([a, 1]; L^\infty)} \geq \|\tilde{v} - v\|_{L^p([a, 1]; L^\infty)} - \|\tilde{u} - \tilde{v}\|_{L^p([a, 1]; L^\infty)} \geq 1 - \varepsilon,$$

where we have used (2.15) and (2.16). Finally, u can not be a Leray-Hopf solution since

$$\|u\|_{L_t^p L^2} \geq \|\tilde{u}\|_{L^p([a, 1]; L^2)} \geq \|\tilde{v}\|_{L^p([a, 1]; L^2)} - \|\tilde{u} - \tilde{v}\|_{L^p([a, 1]; L^\infty)} > \|v(0)\|_2$$

and Leray-Hopf solutions must have non-increasing L^2 norm. □

Next, we prove Theorem 1.10 in the case of the Euler equations.

Proof of Theorem 1.10. We first choose an infinite sequence of smooth divergence-free vector fields $v_n \in C_0^\infty([0, 1] \times \mathbb{T}^d)$ such that:

- (1) On $[0, \frac{1}{2}]$, every v_n coincides and is equal to an exact solution of the Euler equations;
- (2) Each v_n satisfies

$$\|v_n\|_{L^p([0,1];L^\infty)} = 2^n.$$

Since Theorem 1.7 holds for the Euler equations as well, we can apply it for v_n and some $\varepsilon < \frac{1}{2}$. Then we obtain infinite many weak solution $u_n \in L^p L^2 \cap L^1 W^{1,q}$ of the Euler equations.

On one hand, since each v_n we used agrees and solves the Euler equations on $[0, 1/2]$, by Theorem 1.7, for any $n \geq 1$, there exists $\tau_n > 0$ such that u_n coincides with v_n on $[0, \tau_n]$ ⁶. So every weak solution u_n has the same initial data.

On the other hand, these weak solutions u_n are different since for any $n > m$,

$$\|u_n - u_m\|_{L^p L^\infty} \geq \|u_n - v_n\|_{L^p L^\infty} - \|u_m - v_m\|_{L^p L^\infty} \geq 2^n - 1.$$

where we have used that $\varepsilon < \frac{1}{2}$.

It remains to show the regularity $u_n \in L^{3/2-\varepsilon} C^{1/3} \cap L^1 C^{1-\varepsilon}$. This can be done by standard interpolations. Since for any $\varepsilon > 0$ there exists $q < \infty$ such that the embedding

$$W^{1,q}(\mathbb{T}^d) \hookrightarrow C^{1-\varepsilon}(\mathbb{T}^d)$$

holds, we get $L^1 C^{1-\varepsilon}$. The bound $u \in L^{3/2-\varepsilon} C^{1/3}$ can then be obtained by interpolating $L^p L^\infty$ with $L^1 C^{1-\varepsilon}$ (with different ε). \square

Finally, we prove Theorem 1.7 and Theorem 1.9 assuming Proposition 2.2.

Proof of Theorem 1.7 and Theorem 1.9. Let $u_0 = v$ and

$$R_0 = \mathcal{R}\left(\partial_t u_0 - \Delta u_0 + \operatorname{div}(u_0 \otimes u_0)\right)$$

where \mathcal{R} is an inverse divergence operator on \mathbb{T}^d defined in Appendix B.

Since the given vector field v has zero spatial mean for each $t \in [0, 1]$, (u_0, R_0) solves (2.1) trivially and is well-prepared for $I = [0, 1]$ and $\tau = 1$. We construct a sequence of solution (u_n, R_n) for $n \in \mathbb{N}$ as follows.

Given (u_{n-1}, R_{n-1}) , we apply Proposition 2.2 with the parameter

$$\delta_n := 2^{-n} \min \{\|R_{n-1}\|_{L^1 L^r}, \varepsilon\}$$

to obtain (u_n, R_n) . Denote the perturbations by $w_n := u_n - u_{n-1}$ for $n \geq 1$.

As a result, we have

$$\|R_n\|_{L_t^1 L_x^1} \leq \|R_n\|_{L_t^1 L_x^r} \leq \delta_n,$$

and

$$\|w_n\|_{L_t^p L^\infty} + \|w_n\|_{L_t^1 W^{1,q}} \leq \delta_n,$$

for any $n \geq 1$. Also,

$$\|w_n\|_{L_t^2 L^2} \leq M \|R_{n-1}\|_{L_t^1 L^1} \leq M \delta_{n-1},$$

for any $n \geq 2$.

Since u_n is Cauchy in $L_{t,x}^2 \cap L_t^p L^\infty \cap L_t^1 W^{1,q}$, there exists $u \in L_{t,x}^2 \cap L_t^p L^\infty \cap L_t^1 W^{1,q}$ such that

$$u_n \rightarrow u \quad \text{in} \quad L_{t,x}^2 \cap L^p L^\infty \cap L^1 W^{1,q}.$$

To show that u is a weak solution of (1.1), we need to verify the *a.e.* in time divergence-free condition and the weak formulation. As we will show below, the Lebesgue measure of the singular set in time is zero, so u is *a.e.* in time divergence-free by construction.

Now we show that the weak formulation holds. Indeed, take any $\varphi \in \mathcal{D}_T$, then using weak formulation of (2.1) or integrating by parts we have

$$\int_{\mathbb{T}^d} u_n(0, x) \varphi(0, x) dx = - \int_{[0,1] \times \mathbb{T}^d} (u_n \cdot \Delta \varphi + u_n \otimes u_n : \nabla \varphi + u_n \cdot \partial_t \varphi) dx dt - \int_{[0,1] \times \mathbb{T}^d} R_n : \nabla \varphi dx dt.$$

⁶In fact, τ_n can be taken to be $1/2$ if we use Proposition 2.2 instead of Theorem 1.7.

Since $u_n \rightarrow u$ in $L_{t,x}^2$, $R_n \rightarrow 0$ in $L_{t,x}^1$, and $u_n(0, x) \equiv v(0, x)$ for all $n \geq 1$, it follows that all terms above converge to their natural limits and hence u is a weak solution of (1.1).

Moreover,

$$\|u - v\|_{L_t^p L^\infty \cap L_t^1 W^{1,q}} \leq \sum_{n \geq 1} \left[\|w_n\|_{L_t^p L^\infty} + \|w_n\|_{L_t^1 W^{1,q}} \right] \leq \sum_{n \geq 1} 2^{-n} \varepsilon \leq \varepsilon.$$

Finally, we show the structure of the intervals of regularity of u . Recall that each solution (u_n, R_n) is well-prepared, so let us denote by $\mathcal{B}_n \subset [0, 1]$ and $\tau_n > 0$ the set and length scale of the well-preparedness of (u_n, R_n) .

Let

$$\mathcal{I} = \bigcup_{n \geq 0} \mathcal{B}_n^c \setminus \{0, 1\},$$

where the complement is taken in $[0, 1]$. Note that $\mathcal{B}_n^c \subset (\text{supp } w_n)^c$ for all $n \geq 1$, and $\mathcal{B}_n^c \setminus \{0, 1\}$ are monotonically increasing open sets. Therefore $w_k(t) \equiv 0$ on \mathcal{B}_n^c for all $k \geq n$, and hence $u(t) \equiv u_n(t)$ on \mathcal{B}_n^c for each n . Since u_n is smooth, this proves that $u|_{\mathcal{I} \times \mathbb{T}^d} \in C^\infty(\mathcal{I} \times \mathbb{T}^d)$. By construction, each u_n also agrees with each other for $0 \leq t \leq \tau_1/2$, and hence u agree with the unique smooth solution with initial data $v(0)$ near $t = 0$.

Finally, since each \mathcal{B}_n is a finite union of closed intervals, we can write \mathcal{I} as a union of countably many open intervals:

$$\mathcal{I} = \bigcup_{i \geq 0} (a_i, b_i).$$

For the Hausdorff dimension bound of $[0, 1] \setminus \mathcal{I}$, we notice that

$$(0, 1) \setminus \mathcal{I} = \bigcap_{n \geq 0} \mathcal{B}_n = \limsup_n \mathcal{B}_n.$$

Since each \mathcal{B}_n is covered by at most $\tau_n^{-\varepsilon}$ many balls of radius $5\tau_n$, we have

$$d_{\mathcal{H}}(\limsup_n \mathcal{B}_n) \leq \varepsilon.$$

□

3. CONCENTRATING THE STRESS ERROR

The goal of this section is to prove Proposition 3.1 below. The idea is that given a solution (u, R) of (2.1), we can add a small correction term to the existing solution (u, R) so that all of the stress error R concentrates to a set I , the union of small intervals of length τ , and thus obtain a new solution (\bar{u}, \bar{R}) . The key is that the procedure

$$(u, R) \rightarrow (\bar{u}, \bar{R})$$

leaves the size of the stress error R invariant L^1 in time, up to a cost of a constant multiple, namely

$$\|\bar{R}\|_{L^1 L^r} \leq C \|R\|_{L^1 L^r} \quad \text{for any } 1 < r < \infty, \quad (3.1)$$

where $C = C(r, \varepsilon)$ is a universal constant that only depends on r and $\varepsilon > 0$ in the well-preparedness.

Proposition 3.1. *Let $0 < \varepsilon < 1$ and (u, R) be a well-prepared smooth solution of (2.1) for some set and a length scale \tilde{I} and $\tilde{\tau} > 0$. For any $1 < r < \infty$, there exists a universal constant $C = C(r, \varepsilon) > 0$ such that the following holds.*

For any $\delta > 0$, there exists another well-prepared smooth solution (\bar{u}, \bar{R}) of (2.1) some set $I \subset \tilde{I}$ with $0, 1 \notin I$ and $\tau < \tilde{\tau}/2$ satisfying the following.

(1) *The new stress error \bar{R} satisfies*

$$\bar{R}(t, x) = 0 \quad \text{if } \text{dist}(t, I^c) \leq \frac{3\tau}{2},$$

and

$$\|\bar{R}\|_{L^1(0,1; L^r(\mathbb{T}^d))} \leq C \|R\|_{L^1(0,1; L^r(\mathbb{T}^d))};$$

(2) *The velocity perturbation $\bar{w} := \bar{u} - u$ satisfies*

$$\text{supp } \bar{w} \subset \tilde{I} \times \mathbb{T}^d,$$

and

$$\|\bar{w}\|_{L^\infty(0,1; H^d(\mathbb{T}^d))} \leq \delta.$$

Note the slightly stricter bound $\text{dist}(t, I^c) \leq \frac{3\tau}{2}$ versus the definition of well-preparedness is to leave room for the future convex integration scheme in the next section.

3.1. Subdividing the time interval. We first introduce a subdivision of the time interval $[0, 1]$. Then on each sub-interval $[t_i, t_{i+1}]$, we solve a generalized Navier-Stokes (or Euler in the inviscid case) equations and obtain a solution v_i so that $u + v_i$ is a exact solution of the Navier-Stokes equations on $[t_i, t_{i+1}]$.

Let $\tau > 0$ be a small length scale to be fixed in the end of this section and define for $0 \leq i \leq \lfloor \tau^{-\varepsilon} \rfloor$

$$t_i = i\tau^\varepsilon.$$

Without loss of generality, we assume $\tau^{-\varepsilon}$ is always an integer so that the time interval $[0, 1]$ is perfectly divided.

For $0 \leq i \leq \tau^{-\varepsilon} - 1$, let $v_i : [t_i, t_{i+1}] \times \mathbb{T}^d \rightarrow \mathbb{R}^d$ and $q_i : [t_i, t_{i+1}] \times \mathbb{T}^d \rightarrow \mathbb{R}$ be the solution of the following generalized Navier-Stokes system

$$\begin{cases} \partial_t v_i - \Delta v_i + \text{div}(v_i \otimes v_i) + \text{div}(v_i \otimes u) + \text{div}(u \otimes v_i) + \nabla q_i = -\text{div} R \\ \text{div} v_i = 0 \\ v_i(t_i) = 0. \end{cases} \quad (3.2)$$

Since the initial data for v_i is zero and u and R are smooth on $[0, 1] \times \mathbb{T}^d$, thanks to the general local wellposedness theory of the Navier-Stoke equations (or the Euler equations in the inviscid case), for all sufficiently small $\tau > 0$, we may solve equation (3.2) on intervals $t \in [t_i, t_{i+1}]$ to obtain a unique smooth solution v_i .

We shall focus on estimating each v_i on the associated interval $[t_i, t_{i+1}]$. The solution v_i serves as an ‘‘accumulator’’ of the stress error on $[t_i, t_{i+1}]$, and it will provide the major contribution to the new stress error \overline{R} once we use a gluing procedure.

Recall that $\mathcal{R} : C^\infty(\mathbb{T}^d, \mathbb{R}^d) \rightarrow C^\infty(\mathbb{T}^d, \mathcal{S}_0^{d \times d})$ is an inverse divergence operator on \mathbb{T}^d defined in Appendix B. The below result quantifies the size of the corrector v_i in relation to the time scale τ and the forcing $-\text{div} R$.

Proposition 3.2. *Let $d \geq 2$ and (u, R) be a smooth solution of (2.1). There exists a universal constant $C_r > 0$ depending on $1 < r < \infty$ so that the following holds.*

For any $\delta > 0$, if $\tau > 0$ is sufficiently small, then the unique smooth solutions v_i to (3.2) on $[t_i, t_{i+1}]$ satisfies

$$\|v_i\|_{L^\infty([t_i, t_{i+1}]; H^d(\mathbb{T}^d))} \leq \delta,$$

and

$$\|\mathcal{R}v_i\|_{L^\infty([t_i, t_{i+1}]; L^r(\mathbb{T}^d))} \leq C_r \int_{[t_i, t_{i+1}]} \|R(t)\|_{L^r} dt + C_u \delta \tau^\varepsilon,$$

where C_u is a sufficiently large constant depending on u but not δ or τ .

Proof. The first estimate follows directly from the uniform equicontinuity of solutions on bounded sets of H^d and the fact that $\|u\|_{H^d}$ is bounded on $[0, 1]$. Assuming $\delta > 0$ is sufficiently small, we prove the second one as follows.

To reduce notations, we simply write v for v_i and denote $z := \mathcal{R}v$. Note that (3.2) preserves the zero-mean condition.

Denote by \mathbb{P} the Leray projection onto the divergence-free vector fields on \mathbb{T}^d . Once the pressure is eliminated by projecting (3.2), the evolution of z is governed by

$$\partial_t z - \Delta z = F,$$

where

$$F = -\mathcal{R}\mathbb{P} \text{div}(v \otimes v + u \otimes v + v \otimes u) - \mathcal{R}\mathbb{P} \text{div} R.$$

Since $1 < r < \infty$ and $\mathcal{R}\mathbb{P} \text{div}$ is a Calderón-Zygmund operator on $L^r(\mathbb{T}^d)$, a standard energy method yields

$$\|z(t)\|_{L^r(\mathbb{T}^d)} \leq C_r \int_{t_i}^t (\|R(s)\|_r + \|v \otimes v\|_r + \|u \otimes v\|_r + \|v \otimes u\|_r) ds \quad \text{for all } t \in [t_i, t_{i+1}],$$

where the constant C_r depends only on $r > 1$ and dimension d . Using the obtained estimate on v and the embedding $H^d(\mathbb{T}^d) \subset L^\infty(\mathbb{T}^d)$, we get

$$\|z\|_{L^\infty([t_i, t_{i+1}]; L^r(\mathbb{T}^d))} \leq C_r \int_{t_i}^{t_{i+1}} \left(\|R(t)\|_r + \|v\|_{L^\infty([t_i, t_{i+1}]; L^r(\mathbb{T}^d))}^2 + \|u\|_{L_{i,x}^\infty} \|v\|_{L^\infty([t_i, t_{i+1}]; L^r(\mathbb{T}^d))} \right) dt$$

Since $t_{i+1} - t_i = \tau^\varepsilon$, it follows that

$$\|z\|_{L^\infty([t_i, t_{i+1}]; L^r(\mathbb{T}^d))} \leq C_r \int_{t_i}^{t_{i+1}} \|R(t)\|_r dt + C_r \tau^\varepsilon \delta (\delta + \|u\|_{L_{t,x}^\infty}).$$

□

3.2. Temporal concentration by sharp gluing. Since $u + v_i$ is an exact solution of the Navier-Stokes equations on each interval $[t_i, t_{i+1}]$, $0 \leq i \leq \tau^{-\varepsilon} - 1$, the next step is to suitably glue each v_i together so that the glued solution $u + \sum \chi_i v_i$ is still an exact solution on a majority of the time interval $[0, 1]$, with an error supported on many small disjoint sub-intervals.

We first choose cutoff functions that will be used to glue together v_i . We define $\chi_i \in C_c^\infty(\mathbb{R})$ be a smooth cutoff such that when $1 \leq i \leq \tau^{-\varepsilon} - 2$,

$$\chi_i = \begin{cases} 1 & \text{if } t_i + \tau \leq t \leq t_{i+1} - \tau \\ 0 & \text{if } t_i + \tau/2 \geq t \text{ or } t \geq t_{i+1} - \tau/2, \end{cases} \quad (3.3)$$

and when $i = 0$,

$$\chi_i = \begin{cases} 1 & \text{if } 0 \leq t \leq t_{i+1} - \tau \\ 0 & \text{if } t \geq t_{i+1} - \tau/2, \end{cases} \quad (3.4)$$

and when $i = \tau^{-\varepsilon} - 1$,

$$\chi_i = \begin{cases} 1 & \text{if } t_i + \tau \leq t \leq 1 \\ 0 & \text{if } t \leq t_i + \tau/2. \end{cases} \quad (3.5)$$

In other words, we do not cut near the endpoints $t = 0$ and $t = 1$, and the glued solution \bar{u} is an exact solution for a short time near $t = 0$ and $t = 1$. It is worth noting that in the iteration scheme v_i for $i = 0$ or $i = \tau^{-\varepsilon} - 1$ will be zero after the first step (since it is already an exact solution of (1.1) there), so (3.4) and (3.5) are only used once.

Furthermore, we require that the bounds

$$|\nabla^m \chi_i| \lesssim_m \tau^{-m},$$

hold uniformly in τ and i .

Note that for sub-intervals $[t_i, t_{i+1}]$, $1 \leq i \leq \tau^{-\varepsilon} - 2$, we cut near both the left and the right point of $[t_i, t_{i+1}]$. The left cutoff is to ensure smoothness near t_i since each v_i only has a limited amount of time regularity at $t = t_i$ whereas the right cutoff is where the intended gluing takes place. With χ_i in hand, we can simply define the glued solution as

$$\bar{u} := u + \sum_i \chi_i v_i = u + \bar{w}.$$

It is clear that $\bar{u} : [0, 1] \times \mathbb{T}^d \rightarrow \mathbb{R}^d$ has zero spatial mean and is also divergence-free. It remains to show that \bar{u} satisfies the properties listed in Proposition 3.1.

Heuristically, \bar{u} should be an exact solution with a stress error supported on smaller intervals of size τ . To confirm this claim, we must compute the stress error \bar{R} associated with \bar{u} . Since supports of χ_i are disjoint, we can compute

$$\begin{aligned} \partial_t \bar{u} - \Delta \bar{u} + \operatorname{div}(\bar{u} \otimes \bar{u}) &= \operatorname{div} R + (\partial_t - \Delta) \sum_i \chi_i v_i \\ &\quad + \sum_i \chi_i \operatorname{div}(u \otimes v_i) + \sum_i \chi_i \operatorname{div}(v_i \otimes u) \\ &\quad + \sum_i \chi_i^2 \operatorname{div}(v_i \otimes v_i). \end{aligned}$$

Thus, using the fact that v_i solves (3.2) on $[t_i, t_{i+1}]$ and u solves (2.1) on $[0, 1]$, we have

$$\begin{aligned} \partial_t \bar{u} - \Delta \bar{u} + \operatorname{div}(\bar{u} \otimes \bar{u}) + \nabla p &= \operatorname{div} R + \sum_i \partial_t \chi_i v_i + \sum_i (\chi_i^2 - \chi_i) \operatorname{div}(v_i \otimes v_i) \\ &\quad + \sum_i \chi_i (\partial_t v_i - \Delta v_i + \operatorname{div}(v_i \otimes v_i) + \operatorname{div}(u \otimes v_i) + \operatorname{div}(v_i \otimes u)) \\ &= (1 - \sum_i \chi_i) \operatorname{div} R + \sum_i \partial_t \chi_i v_i + \sum_i (\chi_i^2 - \chi_i) \operatorname{div}(v_i \otimes v_i) - \sum_i \chi_i \nabla q_i. \end{aligned}$$

Now let

$$\bar{R} := (1 - \sum_i \chi_i)R + \mathcal{R} \sum_i \partial_t \chi_i v_i + \sum_i (\chi_i^2 - \chi_i) v_i \otimes v_i, \quad (3.6)$$

where \otimes denotes a traceless tensor product, i.e. $f \otimes g = f_i g_j - \frac{1}{d} \delta_{ij} f_k g_k$. Since each v_i has zero spacial mean, $\operatorname{div} \mathcal{R} v_i = v_i$, and we can conclude that

$$\partial_t \bar{u} - \Delta \bar{u} + \operatorname{div}(\bar{u} \otimes \bar{u}) + \nabla \bar{p} = \operatorname{div} \bar{R},$$

where the pressure $\bar{p} : [0, 1] \times \mathbb{T}^d \rightarrow \mathbb{R}$ is defined by

$$\bar{p} = p + \sum_i \chi_i q_i - \sum_i (\chi_i^2 - \chi_i) \frac{|v_i|^2}{d}.$$

The last step is then to show that the new Reynolds stress \bar{R} is comparable to the original one when measured in L_t^1 . It is clear that \bar{R} is much more ‘‘turbulent’’ than the original R as its value changes much more drastically due to the sharp cutoff χ near the endpoints of each interval $[t_i, t_{i+1}]$.

The heuristic is that if τ is small enough, then v_i behaves linearly with a rate $\sim \operatorname{div} R$, and thus gluing together v_i only counts the input from the stress forcing $\operatorname{div} R$. More precisely, the leading order term in (3.6) is the second term, where $\mathcal{R} v_i$ is proportional to R thanks to Proposition 3.2.

Proposition 3.3. *For any $1 < r < \infty$, there exists a universal constant C_r depending on r and ε such that for all sufficiently small $\tau > 0$, the glued solution (\bar{u}, \bar{R}) satisfies*

$$\|\bar{R}\|_{L_t^1 L^r} \leq C_r \|R\|_{L_t^1 L^r}.$$

Proof. By the triangle inequality, we need to estimate

$$\|\bar{R}\|_{L_t^1 L^r} \leq \left\| (1 - \sum_i \chi_i)R \right\|_{L_t^1 L^r} + \sum_i \|\partial_t \chi_i \mathcal{R} v_i\|_{L_t^1 L^r} + \left\| \sum_i (\chi_i - \chi_i^2) v_i \otimes v_i \right\|_{L_t^1 L^r}.$$

The idea is to treat the first and the last terms as small errors. By Hölder’s inequality, we get

$$\|\bar{R}\|_{L_t^1 L^r} \leq \left\| 1 - \sum_i \chi_i \right\|_{L^1([0,1])} \|R\|_{L_t^\infty L^r} + \sum_i \|\partial_t \chi_i\|_{L^1([0,1])} \|\mathcal{R} v_i\|_{L_t^1 L^r} + \sum_i \|(\chi_i - \chi_i^2)\|_{L^1([0,1])} \|v_i \otimes v_i\|_{L_t^\infty L^r}.$$

By the definition of the cutoff χ_i , we have the following trivial bounds (with implicit constants depending on ε) in time:

$$\left\| 1 - \sum_i \chi_i \right\|_{L^1([0,1])} \lesssim \tau^{1-\varepsilon},$$

and for any $0 \leq i \leq \tau^{-\varepsilon} - 1$,

$$\begin{aligned} \|\partial_t \chi_i\|_{L^1([0,1])} &\lesssim 1, \\ \|\chi_i - \chi_i^2\|_{L^1([0,1])} &\lesssim \tau. \end{aligned}$$

In addition, it follows from the bounds for v_i in Proposition 3.2 that

$$\|v_i \otimes v_i\|_{L_t^\infty L^r} \lesssim \|v_i\|_{L_t^\infty H^d}^2 \leq \delta^2.$$

Combining these together and using the bound on $\mathcal{R} v_i$ from Proposition 3.2, we get

$$\begin{aligned} \|\bar{R}\|_{L_t^1 L^r} &\lesssim \tau^{1-\varepsilon} \|R\|_{L_t^\infty L^r} + \sum_i \left(\int_{t_i}^{t_{i+1}} \|R(t)\|_{L^r} dt + C_u \delta \tau^\varepsilon \right) + \delta^2 \sum_i \tau \\ &\lesssim \tau^{1-\varepsilon} \|R\|_{L_t^\infty L^r} + \|R\|_{L_t^1 L^r} + C_u \delta, \end{aligned}$$

where we have used the fact that $\sum_i 1 \leq \tau^{-\varepsilon} \leq \tau^{-1}$.

The conclusion follows once we choose $\tau > 0$ sufficiently small such that

$$C_u \delta \leq \|R\|_{L^1 L^r} \quad \text{and} \quad \tau^{1-\varepsilon} \|R\|_{L_t^\infty L^r} \leq \|R\|_{L^1 L^r}.$$

□

3.3. Proof of Proposition 3.1. We conclude this section with the last step in the proof of Proposition 3.1. Since all the estimates have been obtained, we only need to verify that the temporal support of $\bar{w} = \sum_i \chi_i v_i$ is contained in \tilde{I} and show the well-preparedness of (\bar{u}, \bar{R}) .

Note that (u, R) is well-prepared for \tilde{I} and $\tilde{\tau}$, and it follows from (3.2) that

$$v_i \equiv 0 \quad \text{for } 0 \leq i \leq \tau^{-\varepsilon} - 1 \text{ such that } R \equiv 0 \text{ on } [t_i, t_{i+1}].$$

Hence, if $\tau^\varepsilon = \lfloor [t_i, t_{i+1}] \rfloor$ is sufficiently smaller than $\tilde{\tau}$, the definition of well-preparedness of (u, R) implies that

$$\bigcup_i \text{supp}_t \chi_i v_i \subset \tilde{I}.$$

Thus we have proved that $\text{supp}_t \bar{w} \subset \tilde{I}$.

Let us now show the well-preparedness of (\bar{u}, \bar{R}) . Define an index set

$$E = \{i \in \mathbb{Z} : 1 \leq i \leq \tau^{-\varepsilon} - 1 \text{ and } v_i \neq 0\}$$

satisfying a trivial estimate

$$|E| \leq \tau^{-\varepsilon}.$$

The idea is that the concentrated stress \bar{R} is supported around each t_i , for $i \in E$. Therefore, we can define a set on the time axis

$$I := \bigcup_{i \in E} \left[t_i - \frac{5\tau}{2}, t_i + \frac{5\tau}{2} \right],$$

where as before $t_i = i\tau^\varepsilon$. Note that each interval in I has length 5τ and the total number of intervals is at most $\tau^{-\varepsilon}$, consistent with the well-preparedness, and $0, 1 \notin I$ due to (3.4)

Now take any $t \in [0, 1]$ such that $\text{dist}(t, I^c) \leq \frac{3\tau}{2}$. Then by (3.3) and (3.4),

$$\sum_i \chi_i(t) = 1.$$

Moreover, $\partial_t \chi_i(t) = 0$ and $\chi_i(t) \in \{0, 1\}$ for any i . Consequently,

$$\bar{R}(t) = (1 - \sum_i \chi_i)R + \mathcal{R} \sum_i \partial_t \chi_i v_i + \sum_i (\chi_i^2 - \chi_i) v_i \otimes v_i = 0,$$

for every t such that $\text{dist}(t, I^c) \leq \frac{3\tau}{2}$. In particular, (\bar{u}, \bar{R}) is also well-prepared, which concludes the proof.

4. CONVEX INTEGRATION IN SPACE-TIME

In this section, we will use a convex integration scheme to reduce the size of the Reynolds stress. The goal is to design a suitable velocity perturbation w to the glued solution (\bar{u}, \bar{R}) so that

$$u_1 := \bar{u} + w$$

solves the equation (2.1) with a much smaller Reynolds stress R_1 .

The main goal of the current and the following section is summarized in the following proposition.

Proposition 4.1. *There exists a universal constant $M > 0$ such that for any $p < 2$ and $q < \infty$, there exists $r > 1$ depending only on p and q such that the following holds.*

Let $\delta > 0$ and (\bar{u}, \bar{R}) be a well-prepared smooth solution of (2.1) given by Proposition 3.1 for the set I and time scale τ . Then there exists another well-prepared smooth solution (u_1, R_1) of (2.1) for the same set I and time scale τ such that

$$\|R_1\|_{L^1(0,1;L^r(\mathbb{T}^d))} \leq \delta.$$

Moreover, the velocity perturbation $w := u_1 - \bar{u}$ satisfies

(1) *The temporal support of w is contained in I , i.e.*

$$\text{supp } w \subset I \times \mathbb{T}^d;$$

(2) *The $L^2 L^2$ estimate,*

$$\|w\|_{L^2([0,1] \times \mathbb{T}^d)} \leq M \|\bar{R}\|_{L^1([0,1] \times \mathbb{T}^d)};$$

(3) *The $L^p L^\infty \cap L^1 W^{1,q}$ estimate,*

$$\|w\|_{L^p(0,1;L^\infty(\mathbb{T}^d))} + \|w\|_{L^1(0,1;W^{1,q}(\mathbb{T}^d))} \leq \delta.$$

It is clear that Proposition 2.2 follows from Proposition 3.1 and Proposition 4.1. In the remainder of this section, we will construct the velocity perturbation w and define its associated Reynolds stress R_1 and pressure p_1 . The well-preparedness of (u_1, R_1) will be an easy consequence of the definition of w , whereas all the estimates will be proven in the next section.

4.1. Stationary Mikado flows for the convex integration. The main building blocks of the convex integration scheme is the Mikado flows $\mathbf{W}_k : \mathbb{T}^d \rightarrow \mathbb{R}^d$ introduced in [DS17]. In other contexts [MS18, Luo19], they are called concentrated Mikado flows, or Mikado flows with concentration since they are supported on periodic cylinders with a small radius. Here for brevity, we simply refer to them as the Mikado flows.

We start with a geometric lemma that dates back to the work of Nash [Nas54]. A proof of the following version, which is essentially due to De Lellis and Székelyhidi Jr., can be found in [Sze13, Lemma 3.3]. This lemma is the key reason to use Mikado flows in the convex integration.

Recall that $\mathcal{S}_+^{d \times d}$ is the set of positive definite symmetric $d \times d$ matrices and $\mathbf{e}_k = \frac{k}{|k|}$ for any $k \in \mathbb{Z}^d$.

Lemma 4.2. *For any compact subset $\mathcal{N} \subset \mathcal{S}_+^{d \times d}$, there exists a finite set $\Lambda \subset \mathbb{Z}^d$ and smooth functions $\Gamma_k \in C^\infty(\mathcal{N}; \mathbb{R})$ for any $k \in \Lambda$ such that*

$$R = \sum_{k \in \Lambda} \Gamma_k^2(R) \mathbf{e}_k \otimes \mathbf{e}_k, \quad \text{for all } R \in \mathcal{N}.$$

We apply the lemma $\mathcal{N} = B_{1/2}(\text{Id})$ where $B_{1/2}(\text{Id})$ denotes the metric ball of radius $1/2$ around the identity Id in the space $\mathcal{S}_+^{d \times d}$ to obtain smooth functions Γ_k for any $k \in \Lambda \subset \mathbb{Z}^d$. Throughout the paper, the direction set Λ is fixed, and we construct the Mikado flows as follows.

We choose points $p_k \in (0, 1)^d$ such that $p_k \neq p_{-k}$ if both $k, -k \in \Lambda$. For each $k \in \Lambda$, we denote by $l_k \subset \mathbb{T}^d$ the periodic line passing through p_k in direction k , namely

$$l_k = \{tk + p_k \in \mathbb{T}^d : t \in \mathbb{R}\}.$$

Since Λ is a finite lattice set and we identify \mathbb{T}^d with a periodic box $[0, 1]^d$, there exists a geometric constant $C_\Lambda \in \mathbb{N}$ depending on the set Λ such that

$$|l_k \cap l_{k'}| \leq C_\Lambda \quad \text{for any } k, k' \in \Lambda,$$

where we note that $l_k \cap l_{-k} = \emptyset$ due to $p_k \neq p_{-k}$.

Here we do not even require nonparallel periodic lines to be disjoint in $d \geq 3$ in contrast to [Luo19]. Since nonparallel periodic lines have to intersect in the 2D case, we would rather present a unified approach based on the fact that the intersection parts are very small due to the concentration parameter $\mu > 0$, see Theorem 4.3 below.

Let $\mu > 0$ be the spatial concentration parameter whose value will be fixed in the end of the proof. Let $\phi, \psi \in C_c^\infty([1/2, 1])$ be such that if we define $\psi_k, \phi_k : \mathbb{T}^d \rightarrow \mathbb{R}$ by

$$\psi_k := \mu^{\frac{d-1}{2}} \psi(\mu \text{dist}(l_k, x)) \tag{4.1}$$

and

$$\phi_k := \mu^{\frac{d-1}{2}-2} \phi(\mu \text{dist}(l_k, x)) \tag{4.2}$$

then

$$\Delta \phi_k = \psi_k \quad \text{on } \mathbb{T}^d \quad \text{and} \quad \int_{\mathbb{T}^d} \psi_k^2 dx = 1. \tag{4.3}$$

Note that

$$\text{supp } \psi_k \cap \text{supp } \psi_{k'} \subset \{x \in \mathbb{T}^d : \text{dist}(x, l_k \cap l_{k'}) \leq M_\Lambda \mu^{-1}\} \tag{4.4}$$

for a sufficiently large constant M_Λ depending on Λ .

Finally, the stationary Mikado flows $\mathbf{W}_k : \mathbb{T}^d \rightarrow \mathbb{R}^d$ are defined by

$$\mathbf{W}_k = \psi_k \mathbf{e}_k,$$

where the constant vector $\mathbf{e}_k = \frac{k}{|k|}$. Using the gradient field $\nabla \phi_k$, we may write \mathbf{W}_k as a divergence of a skew-symmetric tensor $\Omega_k \in C_0^\infty(\mathbb{T}^d, \mathbb{R}^{d \times d})$,

$$\Omega_k := \mathbf{e}_k \otimes \nabla \phi_k - \nabla \phi_k \otimes \mathbf{e}_k.$$

Indeed, Ω_k is skew-symmetric by definition, and by a direct computation

$$\text{div } \Omega_k = \text{div}(\nabla \phi_k) \mathbf{e}_k - (\mathbf{e}_k \cdot \nabla) \nabla \phi_k = \Delta \phi_k \mathbf{e}_k - 0 = \mathbf{W}_k.$$

We summarize the main properties of the Mikado flows \mathbf{W}_k in the following theorem.

Theorem 4.3. *Let $d \geq 2$ be the dimension. The stationary Mikado flows $\mathbf{W}_k : \mathbb{T}^d \rightarrow \mathbb{R}^d$ satisfy the following.*

(1) *Each $\mathbf{W}_k \in C_0^\infty(\mathbb{T}^d)$ is divergence-free, satisfies*

$$\mathbf{W}_k = \operatorname{div} \boldsymbol{\Omega}_k,$$

and solves the pressureless Euler equations

$$\operatorname{div}(\mathbf{W}_k \otimes \mathbf{W}_k) = 0.$$

(2) *For any $1 \leq p \leq \infty$, the estimates*

$$\begin{aligned} \mu^{-m} \|\nabla^m \mathbf{W}_k\|_{L^p(\mathbb{T}^d)} &\lesssim_m \mu^{\frac{d-1}{2} - \frac{d-1}{p}}, \\ \mu^{-m} \|\nabla^m \boldsymbol{\Omega}_k\|_{L^p(\mathbb{T}^d)} &\lesssim_m \mu^{-1 + \frac{d-1}{2} - \frac{d-1}{p}}, \end{aligned}$$

hold uniformly in μ .

(3) *For any $k \in \Lambda$, there holds*

$$\int_{\mathbb{T}^d} \mathbf{W}_k \otimes \mathbf{W}_k = \mathbf{e}_k \otimes \mathbf{e}_k,$$

and for any $1 \leq p \leq \infty$

$$\|\mathbf{W}_k \otimes \mathbf{W}_{k'}\|_{L^p(\mathbb{T}^d)} \lesssim \mu^{(d-1) - \frac{d}{p}} \quad \text{if } k \neq k'.$$

Proof. We only prove the last claim as the rest of them are standard and can easily be deduced from (4.1)-(4.3). It suffices to assume $k \neq -k'$. Using the L^∞ bound in (2), we obtain

$$\|\mathbf{W}_k \otimes \mathbf{W}_{k'}\|_{L^p(\mathbb{T}^d)} \lesssim \mu^{(d-1)} \left(\int_{\operatorname{supp} \psi_k \cap \operatorname{supp} \psi_{k'}} dx \right)^{\frac{1}{p}}.$$

Note that due to (4.4) $\operatorname{supp} \psi_k \cap \operatorname{supp} \psi_{k'}$ is contained in a union of finitely many balls of radius $\sim \mu^{-1}$. Thus

$$\|\mathbf{W}_k \otimes \mathbf{W}_{k'}\|_{L^p(\mathbb{T}^d)} \lesssim \mu^{(d-1) - \frac{d}{p}}.$$

□

4.2. Implementation of temporal concentration. Since Mikado flows are stationary, the velocity perturbation will be homogeneous in time if we simply use Lemma 4.2 to define the coefficients. To obtain $L^p L^\infty$ and $L^1 W^{1,q}$ estimates, it is necessary to introduce temporal concentration in the perturbation.

To this end, we choose temporal functions g_κ and h_κ to oscillate the building blocks \mathbf{W}_k intermittently in time. Specifically, g_κ will be used to oscillate \mathbf{W}_k so that the space-time cascade balances the low temporal frequency part of the old stress error \overline{R} , whereas h_κ is used to define a temporal corrector whose time derivative will further balance the high temporal frequency part of the old stress error \overline{R} .

Let $g \in C_c^\infty([0, 1])$ be such that

$$\int_0^1 g^2(t) dt = 1.$$

To add in temporal concentration, let $\kappa > 0$ be a large constant whose value will be specified later and define $g_\kappa : [0, 1] \rightarrow \mathbb{R}$ as the 1-periodic extension of $\kappa^{1/2} g(\kappa t)$ so that

$$\|g_\kappa\|_{L^p([0,1])} \lesssim \kappa^{\frac{1}{2} - \frac{1}{p}} \quad \text{for all } 1 \leq p \leq \infty.$$

The value of κ will be specified later and the function g_κ will be used in the definition of the velocity perturbation. As we will see in Lemma 4.5, the nonlinear term can only balance a portion of the stress error \overline{R} and there is a leftover term which is of high temporal frequency. This motivates us to consider the following temporal corrector.

Let $h_\kappa : [0, 1] \rightarrow \mathbb{R}$ be defined by

$$h_\kappa(t) = \int_0^t g_\kappa^2(s) - 1 ds.$$

In view of the zero-mean condition for $g_\kappa^2(t) - 1$, the function $h_\kappa : [0, 1] \rightarrow \mathbb{R}$ is well-defined and periodic, and we have

$$\|h_\kappa\|_{L^\infty([0,1])} \leq 1, \tag{4.5}$$

uniformly in κ .

We remark that for any $\nu \in \mathbb{N}$, the periodically rescaled function $g_\kappa(\nu \cdot) : [0, 1] \rightarrow \mathbb{R}$ also verifies the bound

$$\|g_\kappa(\nu \cdot)\|_{L^p([0,1])} \lesssim \kappa^{\frac{1}{2} - \frac{1}{p}} \quad \text{for all } 1 \leq p \leq \infty. \quad (4.6)$$

Moreover, we have the identity

$$\partial_t (\nu^{-1} h_\kappa(\nu t)) = g_\kappa^2(\nu t) - 1, \quad (4.7)$$

which will imply the smallness of the corrector, cf (4.15).

4.3. Space-time cutoffs. Before introducing the velocity perturbation, we need to define two important cutoff functions, one to ensure Lemma 4.2 applies and the other to ensure the well-preparedness of the new solution (u_1, R_1) .

Since Lemma 4.2 is stated for a fix compact set in $\mathcal{S}_+^{d \times d}$, we need to introduce a cutoff for the stress \bar{R} . Let $\chi : \mathbb{R}^{d \times d} \rightarrow \mathbb{R}^+$ be a positive smooth function such that χ is monotonically increasing with respect to $|x|$ and

$$\chi(x) = \begin{cases} 1 & \text{if } 0 \leq |x| \leq 1 \\ |x| & \text{if } |x| \geq 2. \end{cases} \quad (4.8)$$

With this cutoff χ , we may define a divisor for the stress \bar{R} so that Lemma 4.2 applies. Indeed, define $\rho \in C^\infty([0, 1] \times \mathbb{T}^d)$ by

$$\rho = \chi(\bar{R}). \quad (4.9)$$

Then immediately

$$\text{Id} - \frac{\bar{R}}{\rho} \in B_{\frac{1}{2}}(\text{Id}) \quad \text{for any } (t, x) \in [0, 1] \times \mathbb{T}^d,$$

which means we can use $\text{Id} - \frac{\bar{R}}{\rho}$ as the argument in the smooth functions Γ_k given by Lemma 4.2.

Next, we need another cutoff to take care of the well-preparedness of the new solution (u_1, R_1) as the perturbation has to be supported within I . Let $\theta \in C_c^\infty(\mathbb{R})$ be a smooth temporal cutoff function such that

$$\theta(t) = \begin{cases} 1 & \text{if } \text{dist}(t, I^c) \geq \frac{3\tau}{2} \\ 0 & \text{if } \text{dist}(t, I^c) \leq \tau, \end{cases} \quad (4.10)$$

where $I \subset [0, 1]$ and $\tau > 0$ are given by Proposition 3.1. Note that this cutoff ensures that the new solution will still be well-prepared.

4.4. The velocity perturbation. We recall that we have defined four parameters for the perturbation so far:

- (1) Temporal oscillation $\nu \in \mathbb{N}$ and temporal concentration $\kappa > 0$;
- (2) Spatial oscillation $\sigma \in \mathbb{N}$ and spatial concentration $\mu \in \mathbb{N}$.

These four parameters are assumed to be sufficiently large for the moment and will be taken to be explicit fixed powers of a frequency parameter $\lambda > 0$ in the next section, where we also fix the value of $r > 1$ appearing in the main proposition.

With all the ingredients in hand, we are ready to define the velocity perturbation. In summary, the velocity perturbation $w : [0, 1] \times \mathbb{T}^d \rightarrow \mathbb{R}^d$ consists of three parts,

$$w = w^{(p)} + w^{(c)} + w^{(t)}.$$

The principle part of the perturbation $w^{(p)}$ consists of super-positions of the building blocks \mathbf{W}_k oscillating with period σ^{-1} on \mathbb{T}^d and period ν^{-1} on $[0, 1]$:

$$w^{(p)}(t, x) := \sum_{k \in \Lambda} a_k(t, x) \mathbf{W}_k(\sigma x), \quad (4.11)$$

where the amplitude function $a_k : [0, 1] \times \mathbb{T}^d \rightarrow \mathbb{R}$ is given by

$$a_k = \theta g_\kappa(\nu t) \rho^{1/2} \Gamma_k \left(\text{Id} - \frac{\bar{R}}{\rho} \right). \quad (4.12)$$

Note that (4.11) is not divergence-free. To fix this, we introduce a divergence-free corrector using the tensor potential Ω_k

$$w^{(c)}(t, x) := \sigma^{-1} \sum_{k \in \Lambda} \nabla a_k(t, x) : \Omega_k(\sigma x). \quad (4.13)$$

Indeed, we can rewrite $w^{(p)} + w^{(c)}$ as

$$\begin{aligned} w^{(p)} + w^{(c)} &= \sigma^{-1} \sum_{k \in \Lambda} a_k(t, x) \operatorname{div} \mathbf{\Omega}_k(\sigma x) + \sigma^{-1} \sum_{k \in \Lambda} \nabla a_k(t, x) : \mathbf{\Omega}_k(\sigma x) \\ &= \sigma^{-1} \operatorname{div} \sum_{k \in \Lambda} a_k(t, x) \mathbf{\Omega}_k(\sigma x), \end{aligned} \quad (4.14)$$

where each $a_k \mathbf{\Omega}_k$ is skew-symmetric and hence $\operatorname{div}(w^{(p)} + w^{(c)}) = 0$.

Finally, we define a temporal corrector to balance the high temporal frequency part of the interaction. This ansatz was first introduced in [BV19] and also used in [BCV18]. The heart of the argument is to ensure that

$$\partial_t w^{(t)} + \operatorname{div}(w^{(p)} \otimes w^{(p)}) = \text{Pressure gradient} + \text{Terms with high spacial frequencies} + \text{Lower order terms.}$$

However, the key difference between [BV19, BCV18] and the current scheme is that here the smallness of the corrector is free and it does not require much temporal oscillation, which is the reason we must use stationary spatial building blocks.

Specifically, the temporal corrector $w^{(t)}$ is defined as

$$w^{(t)} = \sigma^{-1} h_\kappa(\nu t) (\operatorname{div}(\overline{R}) - \nabla \Delta^{-1} \operatorname{div} \operatorname{div}(\overline{R})), \quad (4.15)$$

where we note that Δ^{-1} is well-defined on \mathbb{T}^d since $\operatorname{div} \operatorname{div}(\overline{R}) = \partial_i \partial_j \overline{R}_{ij}$ has zero spatial mean.

It is easy to check $\operatorname{supp}_t w^{(t)} \subset \operatorname{supp}_t \overline{R}$ and $w^{(t)}$ is divergence-free. Indeed,

$$\operatorname{div} w^{(t)} = \sigma^{-1} h_\kappa(\nu t) (\partial_i \partial_j \overline{R}_{ij} - \partial_k \partial_k \Delta^{-1} \partial_i \partial_j \overline{R}_{ij}) = 0.$$

In the lemma below, we show that the leading order interaction of the principle part $w^{(p)}$ is able to balance the low temporal frequency part of the stress error \overline{R} , which motivates the choice of the corrector $w^{(t)}$, cf. (4.21).

Lemma 4.4. *The coefficients a_k satisfy*

$$a_k = 0 \quad \text{if} \quad \operatorname{dist}(t, I^c) \leq \tau,$$

and

$$\sum_{k \in \Lambda} a_k^2 \int_{\mathbb{T}^d} \mathbf{W}_k \otimes \mathbf{W}_k dx = \theta^2 g_\kappa^2(\nu t) \rho \operatorname{Id} - g_\kappa^2(\nu t) \overline{R}.$$

Proof. The first property follows from (4.10).

Now since $\int \mathbf{W}_k \otimes \mathbf{W}_k = \mathbf{e}_k \otimes \mathbf{e}_k$, a direct computation and Lemma 4.2 give

$$\sum_{k \in \Lambda} a_k^2 \mathbf{e}_k \otimes \mathbf{e}_k = \sum_{k \in \Lambda} \theta^2 g_\kappa^2(\nu t) \rho \Gamma_k^2 \mathbf{e}_k \otimes \mathbf{e}_k = \theta^2 g_\kappa^2(\nu t) \rho \sum_{k \in \Lambda} \Gamma_k^2 \left(\operatorname{Id} - \frac{\overline{R}}{\rho} \right) \mathbf{e}_k \otimes \mathbf{e}_k = \theta^2 g_\kappa^2(\nu t) \rho \operatorname{Id} - \theta^2 g_\kappa^2(\nu t) \overline{R}.$$

The identity follows from the fact that

$$\operatorname{supp} \overline{R} \subset \{t : \theta(t) = 1\}.$$

□

4.5. The new Reynolds stress. In this subsection, our goal is to design a suitable stress tensor $R_1 : [0, 1] \times \mathbb{T}^d \rightarrow \mathcal{S}_0^{d \times d}$ such that the pair (u_1, R_1) is a smooth solution of (2.1) for a suitable smooth pressure p_1 .

We first compute the nonlinear term and isolate nonlocal interactions:

$$\operatorname{div}(w^{(p)} \otimes w^{(p)} + \overline{R}) = \operatorname{div} \left[\sum_{k \in \Lambda} a_k^2 \mathbf{W}_k(\sigma x) \otimes \mathbf{W}_k(\sigma x) + \overline{R} \right] + \operatorname{div} R_{\text{far}}, \quad (4.16)$$

where R_{far} denotes the nonlocal interactions between Mikado flows of different directions

$$R_{\text{far}} = \sum_{k \neq k'} a_k a_{k'} \mathbf{W}_k(\sigma x) \otimes \mathbf{W}_{k'}(\sigma x). \quad (4.17)$$

And then we proceed to examine the first term in (4.16), for which by Lemma 4.4 we have

$$\begin{aligned}
& \operatorname{div} \left[\sum_{k \in \Lambda} a_k^2 \mathbf{W}_k(\sigma x) \otimes \mathbf{W}_k(\sigma x) + \bar{R} \right] \\
&= \operatorname{div} \left[\sum_{k \in \Lambda} a_k^2 (\mathbf{W}_k(\sigma x) \otimes \mathbf{W}_k(\sigma x) - \int \mathbf{W}_k \otimes \mathbf{W}_k) + \int \mathbf{W}_k \otimes \mathbf{W}_k + \bar{R} \right] \\
&= \operatorname{div} \sum_{k \in \Lambda} a_k^2 (\mathbf{W}_k(\sigma x) \otimes \mathbf{W}_k(\sigma x) - \int \mathbf{W}_k \otimes \mathbf{W}_k) + \nabla(\theta^2 g_k^2 \rho) + (1 - g_\kappa^2(\sigma t)) \operatorname{div} \bar{R}. \tag{4.18}
\end{aligned}$$

Finally, using the product rule, we compute the divergence term as

$$\begin{aligned}
& \operatorname{div} \sum_{k \in \Lambda} a_k^2 (\mathbf{W}_k(\sigma x) \otimes \mathbf{W}_k(\sigma x) - \int \mathbf{W}_k \otimes \mathbf{W}_k) \\
&= \sum_{k \in \Lambda} \nabla(a_k^2) \cdot (\mathbf{W}_k(\sigma x) \otimes \mathbf{W}_k(\sigma x) - \int \mathbf{W}_k \otimes \mathbf{W}_k). \tag{4.19}
\end{aligned}$$

Typical in the convex integration, we can gain a factor of σ^{-1} in (4.19) by inverting the divergence. To this end, let us use the bilinear anti-divergence operator \mathcal{B} defined in Appendix B. Since (4.19) has zero spatial mean, by (B.4) it is equal to $\operatorname{div} R_{\text{osc},x}$, where

$$R_{\text{osc},x} = \sum_{k \in \Lambda} \mathcal{B} \left(\nabla(a_k^2), \mathbf{W}_k(\sigma x) \otimes \mathbf{W}_k(\sigma x) - \int \mathbf{W}_k \otimes \mathbf{W}_k \right). \tag{4.20}$$

Combining (4.16), (4.18), (4.19), and (4.20) we have

$$\operatorname{div}(w^{(p)} \otimes w^{(p)} + \bar{R}) = \operatorname{div} R_{\text{osc},x} + \operatorname{div} R_{\text{far}} + \nabla(\theta^2 g_k^2 \rho) + (1 - g_\kappa^2(\sigma t)) \operatorname{div} \bar{R}. \tag{4.21}$$

In view of the above computations, we define a temporal oscillation error

$$R_{\text{osc},t} = \nu^{-1} h_\kappa(\nu t) \operatorname{div} \partial_t \bar{R}, \tag{4.22}$$

so that the following decomposition holds.

Lemma 4.5. *Let the space-time oscillation error R_{osc} be*

$$R_{\text{osc}} = R_{\text{osc},x} + R_{\text{osc},t} + R_{\text{far}}.$$

Then

$$\partial_t w^{(t)} + \operatorname{div}(w^{(p)} \otimes w^{(p)} + \bar{R}) + \nabla P = \operatorname{div} R_{\text{osc}}.$$

where the pressure term P is defined by

$$P = \theta^2 g_k^2(\nu t) \rho + \nu^{-1} \Delta^{-1} \operatorname{div} \operatorname{div} \partial_t (\bar{R} h_\kappa(\nu t)). \tag{4.23}$$

Proof. By the definition of $w^{(t)}$, we have

$$\begin{aligned}
\partial_t w^{(t)} &= \partial_t (\nu^{-1} h_\kappa(\nu t) \operatorname{div} \bar{R}) + \partial_t (\nu^{-1} h_\kappa(\nu t) \nabla \Delta^{-1} \operatorname{div} \operatorname{div} \bar{R}) \\
&= (g_\kappa^2(\nu t) - 1) \operatorname{div} \bar{R} + \nu^{-1} h_\kappa(\nu t) \operatorname{div} \partial_t \bar{R} + \nabla \Delta^{-1} \operatorname{div} \operatorname{div} \partial_t (\bar{R} h_\kappa(\nu t)) \\
&= (g_\kappa^2(\nu t) - 1) \operatorname{div} \bar{R} + R_{\text{osc},t} + \nabla \Delta^{-1} \operatorname{div} \operatorname{div} \partial_t (\bar{R} h_\kappa(\nu t)),
\end{aligned}$$

where we used identity (4.7) for the time derivative of h_κ . The conclusion follows immediately from (4.21). \square

Finally, we can define the correction error and the linear error as usual:

$$R_{\text{cor}} = \mathcal{R}(\operatorname{div}(w^{(c)} + w^{(t)}) \otimes w + w^{(p)} \otimes (w^{(c)} + w^{(t)})) \tag{4.24}$$

and

$$R_{\text{lin}} = \mathcal{R}(\partial_t(w^{(p)} + w^{(c)}) - \Delta u + \bar{u} \otimes w + w \otimes \bar{u}). \tag{4.25}$$

To conclude, we summarize the main results in this section below.

Lemma 4.6. *Define the new Reynolds stress by*

$$R_1 = R_{\text{lin}} + R_{\text{cor}} + R_{\text{osc}},$$

and the new pressure by

$$p_1 = \bar{p} + P.$$

Then (u_1, R_1) is a well-prepared solution to (2.1),

$$\partial_t u_1 - \Delta u_1 + \operatorname{div}(u_1 \otimes u_1) + \nabla p_1 = \operatorname{div} R_1,$$

and the velocity perturbation $w = u_1 - \bar{u}$ satisfies

$$\operatorname{supp} w \subset I \times \mathbb{T}^d.$$

Proof. A direct computation of the left-hand side gives

$$\begin{aligned} \partial_t u_1 - \Delta u_1 + \operatorname{div}(u_1 \otimes u_1) + \nabla p_1 &= \partial_t \bar{u} - \Delta \bar{u} + \operatorname{div}(\bar{u} \otimes \bar{u}) + \nabla p \\ &\quad + \partial_t w - \Delta w + \operatorname{div}(\bar{u} \otimes w) + \operatorname{div}(w \otimes \bar{u}) + \operatorname{div}(w \otimes w) + \nabla P \\ &= \operatorname{div} \bar{R} + \partial_t w - \Delta w + \operatorname{div}(\bar{u} \otimes w) + \operatorname{div}(w \otimes \bar{u}) + \operatorname{div}(w \otimes w) + \nabla P, \end{aligned}$$

where we have use the fact that (\bar{u}, \bar{R}) solves (2.1) with pressure \bar{p} .

From the definitions (4.24), (4.25) and Lemma 4.5 we can conclude that (u_1, R_1) solves (2.1).

The claim that (u_1, R_1) is well-prepared and $\operatorname{supp}_t w \subset I$ follows from the fact that (\bar{u}, \bar{R}) is given by Proposition 3.1 and the perturbations $w^{(p)}$, $w^{(c)}$ and $w^{(t)}$ satisfy

$$w^{(p)} = w^{(c)} = 0 \quad \text{if } \operatorname{dist}(t, I^c) \leq \tau$$

and

$$\operatorname{supp}_t w^{(t)} \subset \operatorname{supp}_t \bar{R}.$$

□

5. PROOF OF PROPOSITION 4.1

In this section we will show that the velocity perturbation w and the new Reynolds stress R_1 derived in Section 4 satisfy the claimed properties in Proposition 4.1.

As a general note, we use a constant C_u for dependency on the previous solution (\bar{u}, \bar{R}) throughout this section. Unless otherwise indicated, in the statement of below lemmas and propositions the exponents p, q and r refer to the ones given by Proposition 2.2, cf. (5.1) and (5.2).

5.1. Choice of parameters. We fix two parameters $0 < \gamma < 1$ and $1 < r < 2$ as follows.

(1) First, we choose $\gamma > 0$ small enough such that

$$10d\gamma \leq \min \left\{ \frac{1}{p} - \frac{1}{2}, \frac{1}{q} \right\}. \quad (5.1)$$

(2) Once γ is fixed, choose $r > 1$ such that

$$d - \frac{d}{r} \leq \gamma. \quad (5.2)$$

It is clear that $r > 1$ only depends on p, q , as claimed in Proposition 4.1.

Let λ be a sufficiently large number whose value will be fixed in the end. We choose the parameters σ, κ along with ν, μ in the building blocks as explicit powers of λ as follows.

(1) Temporal oscillation $\nu \in \mathbb{N}$ and spatial oscillation $\sigma \in \mathbb{N}$:

$$\begin{aligned} \nu &= \lceil \lambda^\gamma \rceil, \\ \sigma &= \lceil \lambda^{\frac{1}{\gamma}} \rceil. \end{aligned}$$

(2) Temporal concentration $\kappa > 0$ and spatial concentration $\mu > 0$:

$$\begin{aligned} \kappa &= \lambda^{\frac{2}{\gamma} + d + 1 - 6\gamma}, \\ \mu &= \lambda. \end{aligned}$$

Note that we have the hierarchy of parameters

$$\nu \ll \mu \ll \sigma \ll \kappa^{1/2}.$$

More precisely, we have the following useful lemma that will be used throughout the next section.

Lemma 5.1. *For any $\lambda > 0$, there hold*

$$\nu \kappa^{\frac{1}{2}} \sigma^{-1} \mu^{-1} \mu^{\frac{d-1}{2} - \frac{d-1}{r}} \leq \lambda^{-\gamma} \quad (5.3)$$

$$\kappa^{\frac{1}{2} - \frac{1}{p}} \mu^{\frac{d-1}{2}} \leq \lambda^{-\gamma} \quad (5.4)$$

$$\kappa^{-\frac{1}{2}} \sigma \mu \mu^{\frac{d-1}{2} - \frac{d-1}{q}} \leq \lambda^{-\gamma}. \quad (5.5)$$

Proof. The first inequality (5.3) is equivalent to

$$\gamma + \left(\frac{1}{\gamma} + \frac{d+1}{2} - 3\gamma \right) - \frac{1}{\gamma} - 1 + \left(\frac{d-1}{2} - \frac{d-1}{r} \right) \leq -\gamma$$

It can be simplified to

$$d-1 - \frac{d-1}{r} \leq \gamma$$

which holds due to (5.2).

For the second one (5.4), thanks to (5.1) it suffices to show that

$$\kappa^{-d\gamma} \mu^{\frac{d-1}{2}} \leq \lambda^{-\gamma}.$$

It is equivalent to

$$-2d + \gamma d(d+1 - 6\gamma) + \frac{d-1}{2} \leq -\gamma$$

which holds trivially since $10d\gamma \leq 1$.

The third one (5.5), looking again at the exponents, reads as

$$-\left(\frac{1}{\gamma} + \frac{d+1}{2} - 3\gamma \right) + \frac{1}{\gamma} + 1 + \left(\frac{d-1}{2} - \frac{d-1}{q} \right) \leq -\gamma.$$

Simplifying, we obtain

$$-\frac{d-1}{q} \leq -4\gamma,$$

which also holds due to (5.1). □

5.2. Estimates on velocity perturbation. We first estimate the coefficient a_k of the perturbation w .

Lemma 5.2. *The coefficients a_k are smooth on $[0, 1] \times \mathbb{T}^d$ and*

$$\|\partial_t^n \nabla^m a_k\|_{L^p(0,1; L^\infty(\mathbb{T}^d))} \leq C_{u,m,n} (\nu \kappa)^n \kappa^{\frac{1}{2} - \frac{1}{p}} \quad \text{for any } p \in [1, \infty].$$

In addition, the bound

$$\|a_k(t)\|_{L^2(\mathbb{T}^d)} \lesssim \theta(t) g_\kappa(\sigma t) \left(\int_{\mathbb{T}^d} \rho(t, x) dx \right)^{\frac{1}{2}}$$

holds for all time $t \in [0, 1]$.

Proof. It follows from definition (4.12) that a_k is smooth. Since the implicit constant is allowed to depend on (\bar{u}, \bar{R}) , it suffices to consider only the time differentiation.

We have that

$$\begin{aligned} \|\partial_t^n [\theta(\cdot) g_\kappa(\nu \cdot)]\|_{L^p([0,1])} &\leq \|\theta\|_{C^n([0,1])} \|\partial_t^n g_\kappa(\nu \cdot)\|_{L^p([0,1])} \\ &\lesssim (\sigma \kappa)^n \kappa^{\frac{1}{2} - \frac{1}{p}}, \end{aligned}$$

which implies the first bound. The second bound follows immediately from the definition of a_k :

$$\|a_k(t)\|_{L^2(\mathbb{T}^d)} \lesssim \theta g_\kappa \left(\int_{\mathbb{T}^d} \rho \Gamma_k \left(\text{Id} - \frac{\bar{R}}{\rho} \right) dx \right)^{\frac{1}{2}} \lesssim \theta g_\kappa \left(\int_{\mathbb{T}^d} \rho dx \right)^{\frac{1}{2}}.$$

□

With the estimates of a_k in hand, we start estimating the velocity perturbation. As expected, the principle part $w^{(p)}$ is the largest among all parts in w .

Proposition 5.3. *The principle part $w^{(p)}$ satisfies*

$$\|w^{(p)}\|_{L^2([0,1]\times\mathbb{T}^d)} \lesssim \|\overline{R}\|_{L^1([0,1]\times\mathbb{T}^d)}^{\frac{1}{2}} + C_u \sigma^{-\frac{1}{2}},$$

and

$$\|w^{(p)}\|_{L^p(0,1;L^\infty(\mathbb{T}^d))} + \|w^{(p)}\|_{L^1(0,1;W^{1,q}(\mathbb{T}^d))} \leq C_u \lambda^{-\gamma}.$$

In particular, for sufficiently large λ ,

$$\begin{aligned} \|w^{(p)}\|_{L^2([0,1]\times\mathbb{T}^d)} &\lesssim \|\overline{R}\|_{L^1([0,1]\times\mathbb{T}^d)}^{\frac{1}{2}}, \\ \|w^{(p)}\|_{L^p(0,1;L^\infty(\mathbb{T}^d))} &\leq \frac{\delta}{4}. \end{aligned}$$

Proof. We first show the estimate for $L^2_{t,x}$ and then for $L^p L^\infty$.

$L^2_{t,x}$ estimate:

Taking L^2 norm in space and appealing to Lemma B.1, we have

$$\|w^{(p)}(t)\|_{L^2(\mathbb{T}^d)} \lesssim \sum_{k \in \Lambda} \|a_k(t)\|_2 \|\mathbf{W}_k\|_2 + \sigma^{-\frac{1}{2}} C_u.$$

Recall that $\|\mathbf{W}_k\|_2 \lesssim 1$. Then using Lemma 5.2 and taking L^2 norm in time gives

$$\|w^{(p)}\|_{L^2([0,1]\times\mathbb{T}^d)} \lesssim \sum_{k \in \Lambda} \left(\int_0^1 g_\kappa^2(\nu t) \int_{\mathbb{T}^d} \rho(t, x) dx dt \right)^{\frac{1}{2}} + \sigma^{-\frac{1}{2}} C_u. \quad (5.6)$$

Notice that

$$t \mapsto \int_{\mathbb{T}^d} \rho(t, x) dx$$

is a smooth map on $[0, 1]$. Thus, we may apply Lemma B.1 once again (with $p = 1$) to obtain that

$$\int_0^1 g_\kappa^2(\nu t) \int_{\mathbb{T}^d} \rho(t, x) dx dt \lesssim \|\overline{R}\|_{L^1([0,1]\times\mathbb{T}^d)} + C_u \nu^{-1}, \quad (5.7)$$

where we have used the fact that $\int g_\kappa^2 = 1$ and the bound

$$\int_{\mathbb{T}^d} \rho(t, x) dx \lesssim \|\overline{R}(t)\|_{L^1(\mathbb{T}^d)}.$$

Hence, combining (5.6) and (5.7) gives

$$\|w^{(p)}\|_{L^2([0,1]\times\mathbb{T}^d)} \lesssim \|\overline{R}\|_{L^1([0,1]\times\mathbb{T}^d)}^{\frac{1}{2}} + C_u \sigma^{-\frac{1}{2}}.$$

$L^p L^\infty$ estimate:

Taking L^∞ norm in space and using Hölder's inequality give

$$\|w^{(p)}(t)\|_{L^\infty(\mathbb{T}^d)} \lesssim \sum_{k \in \Lambda} \|a_k(t)\|_\infty \|\mathbf{W}_k\|_\infty.$$

We can now take L^p in time and apply the estimates in Lemma 5.2 and Theorem 4.3 to obtain

$$\begin{aligned} \|w^{(p)}\|_{L^p(0,1;L^\infty(\mathbb{T}^d))} &\lesssim \mu^{\frac{d-1}{2}} \sum_{k \in \Lambda} \|a_k\|_{L^p(0,1;L^\infty(\mathbb{T}^d))} \\ &\leq C_u \kappa^{\frac{1}{2} - \frac{1}{p}} \mu^{\frac{d-1}{2}}, \end{aligned}$$

which by Lemma 5.1 implies that

$$\|w^{(p)}\|_{L^p(0,1;L^\infty(\mathbb{T}^d))} \leq C_u \lambda^{-\gamma}.$$

$L^1 W^{1,q}$ estimate:

This part is similar to the $L^p L^\infty$. We first take $W^{1,q}$ norm in space to obtain

$$\|w^{(p)}(t)\|_{W^{1,q}(\mathbb{T}^d)} \lesssim \sum_{k \in \Lambda} \|a_k(t)\|_{C^1} \|\mathbf{W}_k(\sigma \cdot)\|_{W^{1,q}}.$$

Integrating in time, by Theorem 4.3, Lemma 5.2 we have that

$$\begin{aligned} \|w^{(p)}(t)\|_{L^1(0,1;W^{1,q}(\mathbb{T}^d))} &\lesssim \sum_{k \in \Lambda} \|a_k(t)\|_{L^1 C^1} \|\mathbf{W}_k(\sigma \cdot)\|_{W^{1,q}} \\ &\lesssim \sigma \mu \mu^{\frac{d-1}{2} - \frac{d-1}{q}} \sum_{k \in \Lambda} \|a_k(t)\|_{L^1 C^1} \\ &\lesssim \sigma \kappa^{-\frac{1}{2}} \mu \mu^{\frac{d-1}{2} - \frac{d-1}{q}}, \end{aligned}$$

which implies the desired bound thanks to Lemma 5.1. □

Next, we estimate the corrector $w^{(c)}$, which is expected to be much smaller than $w^{(p)}$ due to the derivative gains from both the fast oscillation σ and the tensor potential Ω_k .

Proposition 5.4. *The divergence-free corrector $w^{(c)}$ satisfies*

$$\|w^{(c)}\|_{L^2(0,1;L^\infty(\mathbb{T}^d))} \leq C_u \lambda^{-\gamma},$$

and

$$\|w^{(c)}\|_{L^1(0,1;W^{1,q}(\mathbb{T}^d))} \leq C_u \lambda^{-\gamma}.$$

In particular, for sufficiently large λ ,

$$\|w^{(c)}\|_{L^2([0,1] \times \mathbb{T}^d)} \leq \|\bar{R}\|_{L^1([0,1] \times \mathbb{T}^d)}^{\frac{1}{2}},$$

and

$$\|w^{(c)}\|_{L^p(0,1;L^\infty(\mathbb{T}^d))} + \|w^{(c)}\|_{L^1(0,1;W^{1,q}(\mathbb{T}^d))} \leq \frac{\delta}{4}.$$

Proof. $L^2 L^\infty$ **estimate:**

From the definition, we have

$$\begin{aligned} \|w^{(c)}(t)\|_{L^\infty(\mathbb{T}^d)} &\leq \sigma^{-1} \left\| \sum_{k \in \Lambda} \nabla a_k(t) : \Omega_k(\sigma \cdot) \right\|_{L^\infty(\mathbb{T}^d)} \\ &\lesssim \sigma^{-1} \sum_{k \in \Lambda} \|\nabla a_k(t)\|_{L^\infty(\mathbb{T}^d)} \|\Omega_k(\sigma \cdot)\|_{L^\infty(\mathbb{T}^d)}. \end{aligned}$$

Now, thanks to Lemma 5.2 and (5.2), we take L^2 in time to obtain

$$\begin{aligned} \|w^{(c)}\|_{L^2(0,1;L^\infty(\mathbb{T}^d))} &\lesssim \sigma^{-1} \mu^{-1 + \frac{d-1}{2}} \sum_{k \in \Lambda} \|\nabla a_k\|_{L^2(0,1;L^\infty(\mathbb{T}^d))} \\ &\leq C_u \sigma^{-1} \mu^{-1 + \frac{d-1}{2}}, \end{aligned}$$

which by the definition of γ implies that

$$\|w^{(c)}\|_{L^2(0,1;L^\infty(\mathbb{T}^d))} \leq C_u \lambda^{-\gamma}.$$

$L^1 W^{1,q}$ **estimate:** This part is very similar to the estimation of $w^{(p)}$. We first take $W^{1,q}$ in space to obtain that

$$\begin{aligned} \|w^{(c)}(t)\|_{W^{1,q}(\mathbb{T}^d)} &\leq \sigma^{-1} \left\| \sum_{k \in \Lambda} \nabla a_k(t) : \Omega_k(\sigma \cdot) \right\|_{W^{1,q}(\mathbb{T}^d)} \\ &\lesssim \sigma^{-1} \sum_{k \in \Lambda} \|a_k(t)\|_{C^2(\mathbb{T}^d)} \|\Omega_k(\sigma \cdot)\|_{W^{1,q}(\mathbb{T}^d)}. \end{aligned}$$

Integrating in space and using Lemma 5.2 and Theorem 4.3 we have

$$\begin{aligned} \|w^{(c)}\|_{L^1(0,1;W^{1,q}(\mathbb{T}^d))} &\lesssim \sigma^{-1} \sum_{k \in \Lambda} \|a_k(t)\|_{L^1(0,1;C^2(\mathbb{T}^d))} \|\Omega_k(\sigma \cdot)\|_{W^{1,q}(\mathbb{T}^d)} \\ &\lesssim \kappa^{-\frac{1}{2}} \mu^{\frac{d-1}{2} - \frac{d-1}{q}}, \end{aligned}$$

which differs the estimate of $\|w^{(p)}\|_{L^1W^{1,q}}$ by a factor of $\sigma\mu$ and hence

$$\|w^{(c)}\|_{L^1(0,1;W^{1,q}(\mathbb{T}^d))} \lesssim C_u \lambda^{-\gamma}.$$

□

Finally, we estimate the temporal corrector $w^{(t)}$. From its definition (4.15), one can see that the spatial frequency of $w^{(t)}$ is independent from the parameters σ , τ and μ . As a result, this term poses no constraints to the choice of temporal and spatial oscillation/concentration at all and is small for basically any choice of parameters (as long as temporal oscillation ν is present). This is one of the main technical differences from [BV19, BCV18] where the leading order effect is temporal oscillation.

Proposition 5.5. *The temporal corrector $w^{(t)}$ satisfies*

$$\|w^{(t)}\|_{L^\infty(0,1;W^{1,\infty}(\mathbb{T}^d))} \leq C_u \nu^{-1}.$$

In particular, for sufficiently large λ ,

$$\|w^{(t)}\|_{L^2([0,1] \times \mathbb{T}^d)} \leq \|\overline{R}\|_{L^1([0,1] \times \mathbb{T}^d)}^{\frac{1}{2}},$$

and

$$\|w^{(t)}\|_{L^p(0,1;L^\infty(\mathbb{T}^d))} + \|w^{(t)}\|_{L^1(0,1;W^{1,q}(\mathbb{T}^d))} \leq \frac{\delta}{4}.$$

Proof. It follows directly from the definition of $w^{(t)}$ that

$$\|w^{(t)}\|_{L^\infty(0,1;W^{1,\infty}(\mathbb{T}^d))} \lesssim \nu^{-1} \|h\|_{L^\infty([0,1])} \|\overline{R}\|_{L^\infty(0,1;W^{2,\infty}(\mathbb{T}^d))} \leq \nu^{-1} C_u.$$

□

5.3. Estimates on the new Reynolds stress. The last step of the proof is to estimate R_1 . We proceed with the decomposition in Lemma 4.6. More specifically, we will prove that for all sufficiently large λ , each part of the stress R_1 is less than $\frac{\delta}{4}$.

5.3.1. *Linear error.*

Lemma 5.6. *For sufficiently large λ ,*

$$\|R_{\text{lin}}\|_{L^1(0,1;L^r(\mathbb{T}^d))} \leq \frac{\delta}{4}.$$

Proof. We split the linear error into three parts:

$$\|R_{\text{lin}}\|_{L^1(0,1;L^r(\mathbb{T}^d))} \leq \underbrace{\|\mathcal{R}(\Delta w)\|_{L_t^1 L^r}}_{:=L_1} + \underbrace{\|\mathcal{R}(\partial_t(w^{(p)} + w^{(c)}))\|_{L_t^1 L^r}}_{:=L_2} + \underbrace{\|\mathcal{R}(\text{div}(w \otimes \bar{u}) + \text{div}(\bar{u} \otimes w))\|_{L_t^1 L^r}}_{:=L_3}.$$

Estimate of L_1 :

By (B.3) or boundedness of Riesz transform we have

$$L_1 \lesssim \|w\|_{L^1(0,1;W^{1,r}(\mathbb{T}^d))}.$$

Note that we have estimated w in $L^1W^{1,q}$ and thus by Proposition 5.3–5.5 we can conclude that

$$L_1 \leq C_u \lambda^{-\gamma}. \tag{5.8}$$

Estimate of L_2 :

By (4.14), we have

$$\partial_t(w^{(p)} + w^{(c)}) = \sigma^{-1} \sum_k \text{div}(\partial_t a_k \Omega_k(\sigma \cdot)),$$

and hence

$$\begin{aligned} L_2 &\leq \|\mathcal{R}\partial_t(w^{(p)} + w^{(c)})\|_{L^1(0,1;L^r(\mathbb{T}^d))} \\ &\lesssim \sigma^{-1} \sum_k \|\mathcal{R} \operatorname{div}(\partial_t a_k \Omega_k(\sigma \cdot))\|_{L^1(0,1;L^r(\mathbb{T}^d))}. \end{aligned}$$

Since $\mathcal{R} \operatorname{div}$ is a Calderón-Zygmund operator on \mathbb{T}^d , we have

$$L_2 \lesssim \sigma^{-1} \sum_k \|\partial_t a_k\|_{L^1 L^\infty} \|\Omega_k\|_r.$$

Appealing to Lemma 5.2 and estimates of Ω_k listed in Theorem 4.3 we have

$$\begin{aligned} L_2 &\leq C_u (\sigma \kappa) \kappa^{-\frac{1}{2}} \mu^{\frac{d-1}{2} - \frac{d-1}{r}} \\ &\leq C_u \lambda^{-\gamma}. \end{aligned} \tag{5.9}$$

Estimate of L_3 : For the last term we simply use L^r boundedness of \mathcal{R} to obtain

$$L_3 \lesssim \|w \otimes \bar{w}\|_{L^1_t L^r}.$$

Here we use a crude bound

$$\|w \otimes \bar{w}\|_{L^1_t L^r} \lesssim \|w\|_{L^p L^\infty} \|\bar{w}\|_{L^\infty_{t,x}},$$

and apply the obtained estimates in Proposition 5.3, 5.4 and 5.5 to conclude

$$L_3 \leq C_u \lambda^{-\gamma}. \tag{5.10}$$

From (5.8), (5.9), and (5.10), we can conclude that for all sufficiently large λ , there holds

$$\|R_{\text{lin}}\|_{L^1(0,1;L^r(\mathbb{T}^d))} \leq \frac{\delta}{4}.$$

□

5.3.2. Correction error.

Lemma 5.7. *For sufficiently large λ ,*

$$\|R_{\text{cor}}\|_{L^1(0,1;L^r(\mathbb{T}^d))} \leq \frac{\delta}{4}.$$

Proof. By the boundedness of $\mathcal{R} \operatorname{div}$ in L^r and Hölder's inequality,

$$\begin{aligned} \|R_{\text{cor}}\|_{L^1(0,1;L^r(\mathbb{T}^d))} &\lesssim \|(w^{(c)} + w^{(t)}) \otimes w\|_{L^1 L^r} + \|w^{(p)} \otimes (w^{(c)} + w^{(t)})\|_{L^1 L^r} \\ &\lesssim \left(\|w^{(c)}\|_{L^2 L^\infty} + \|w^{(t)}\|_{L^2 L^\infty} \right) \|w\|_{L^2 L^2} + \|w^{(p)}\|_{L^2 L^2} \left(\|w^{(c)}\|_{L^2 L^\infty} + \|w^{(t)}\|_{L^2 L^\infty} \right). \end{aligned}$$

By Propositions 5.3,

$$\begin{aligned} \|w\|_{L^2 L^2} &\lesssim \|w^{(p)}\|_{L^2 L^2} + \|w^{(c)}\|_{L^2 L^2} + \|w^{(t)}\|_{L^2 L^2} \\ &\lesssim \|\bar{R}\|_{L^1([0,1] \times \mathbb{T}^d)}^{\frac{1}{2}}, \end{aligned}$$

and by Proposition 5.4 and 5.5,

$$\|w^{(c)}\|_{L^2 L^\infty} + \|w^{(t)}\|_{L^2 L^\infty} \leq C_u \lambda^{-\gamma}.$$

So for all λ sufficiently large, we can conclude that

$$\|R_{\text{cor}}\|_{L^1(0,1;L^r(\mathbb{T}^d))} \leq \frac{\delta}{4}.$$

□

5.3.3. Oscillation error.

Lemma 5.8. For sufficiently large λ ,

$$\|R_{\text{osc}}\|_{L^1(0,1;L^r(\mathbb{T}^d))} \leq \frac{\delta}{4}.$$

Proof. We will use the decomposition from Lemma 4.5

$$R_{\text{osc}} = R_{\text{osc},x} + R_{\text{osc},t} + R_{\text{far}}.$$

Estimate of $R_{\text{osc},x}$:

Denote $\mathbf{T}_k : [0, 1] \times \mathbb{T}^d \rightarrow \mathbb{R}^{d \times d}$ by

$$\mathbf{T}_k = \mathbf{W}_k \otimes \mathbf{W}_k - \int \mathbf{W}_k \otimes \mathbf{W}_k,$$

so that

$$R_{\text{osc},x} = \sum_{k \in \Lambda} \mathcal{B}(\nabla(a_k^2), \mathbf{T}_k(\sigma \cdot)).$$

Using Theorem B.4, we can estimate the L^r norm of $R_{\text{osc},x}$ as follows.

$$\begin{aligned} \|R_{\text{osc},x}\|_{L^r(\mathbb{T}^d)} &= \left\| \sum_{k \in \Lambda} \mathcal{B}(\nabla(a_k^2), \mathbf{T}_k(\sigma \cdot)) \right\|_{L^r} \\ &\lesssim \sum_{k \in \Lambda} \|\nabla(a_k^2)\|_{C^1} \|\mathcal{R}(\mathbf{T}_k(\sigma \cdot))\|_{L^r} \\ &\lesssim \sigma^{-1} \sum_{k \in \Lambda} \|\nabla(a_k^2)\|_{C^1} \|\mathbf{T}_k\|_{L^r}, \end{aligned}$$

where we have used the fact that $\mathbf{T}_k(\sigma \cdot)$ has zero spatial mean in the last inequality. Thanks to Theorem 4.3, for any $k \in \Lambda$

$$\|\mathbf{T}_k\|_{L^r} \lesssim \|\mathbf{W}_k \otimes \mathbf{W}_k\|_{L^r} \lesssim \|\mathbf{W}_k\|_{L^{2r}}^2 \lesssim \mu^{d-1-\frac{d-1}{r}}.$$

Therefore, by Lemma 5.2,

$$\|R_{\text{osc},x}\|_{L^1(0,1;L^r(\mathbb{T}^d))} \leq C_u \sigma^{-1} \mu^{d-1-\frac{d-1}{r}}.$$

Estimate of $R_{\text{osc},t}$:

Using the bound on g_k (4.5), we infer

$$\begin{aligned} \|R_{\text{osc},t}\|_{L^1(0,1;L^r(\mathbb{T}^d))} &= \|\sigma^{-1} h_k(\sigma t) \operatorname{div} \partial_t \bar{R}\|_{L^1 L^r} \\ &\lesssim \sigma^{-1} \|h_k(\sigma \cdot)\|_{L^1 C_u} \\ &\leq C_u \sigma^{-1}. \end{aligned}$$

Estimate of R_{far} :

We can use Theorem 4.3 and Lemma 5.2 to obtain

$$\begin{aligned} \|R_{\text{far}}\|_{L^1(0,1;L^r(\mathbb{T}^d))} &= \left\| \sum_{k \neq k'} a_k a_{k'} \mathbf{W}_k(\sigma \cdot) \otimes \mathbf{W}_{k'}(\sigma \cdot) \right\|_{L^1(0,1;L^r(\mathbb{T}^d))} \\ &\lesssim \sum_{k \neq k'} \|a_k\|_{L^2(0,1;L^\infty(\mathbb{T}^d))} \|a_{k'}\|_{L^2(0,1;L^\infty(\mathbb{T}^d))} \|\mathbf{W}_k \otimes \mathbf{W}_{k'}\|_{L^r} \\ &\leq C_u \mu^{d-1-\frac{d}{r}}. \end{aligned}$$

Now we can combine all the estimates and conclude

$$\|R_{\text{osc}}\|_{L^1(0,1;L^r(\mathbb{T}^d))} \leq C_u \left(\sigma^{-1} \mu^{d-1-\frac{d-1}{r}} + \sigma^{-1} + \mu^{d-1-\frac{d}{r}} \right).$$

Thanks to (5.2), we have

$$\|R_{\text{osc}}\|_{L^1(0,1;L^r(\mathbb{T}^d))} \leq C_u \lambda^{-\gamma}.$$

And thus for λ large enough, the desire bound holds:

$$\|R_{\text{osc}}\|_{L^1(0,1;L^r(\mathbb{T}^d))} \leq \frac{\delta}{4}.$$

□

APPENDIX A. $X^{p,q}$ WEAK SOLUTIONS ON THE TORUS

In this section, we show that sub-critical and critical weak solutions in $X^{p,q}([0, T]; \mathbb{T}^d)$ are in fact Leray-Hopf. In particular, by the weak-strong uniqueness results of Ladyzhenskaya-Prodi-Serrin, this implies the uniqueness part of Theorem 1.3. The content of Theorem A.1 is classical [FJR72, Kat84, LM01] and we include a proof in our specific setting for the convenience of the readers. Note that the proof applies to the entire range of exponents p, q including the case $q = \infty$ which is most relevant to the results of this paper, but was omitted in [FJR72].

Theorem A.1. *Let $d \geq 2$ be the dimension and $u \in X^{p,q}([0, T]; \mathbb{T}^d)$ be a weak solution of (1.1) with $\frac{2}{p} + \frac{d}{q} = 1$, $d \leq q \leq \infty$. Then u is a Leray-Hopf solution.*

To prove Theorem A.1, we follow the duality approach in [LM01] and use only classical ingredients.

The first ingredient is an existence result for a linearized Navier-Stokes equation.

Theorem A.2. *Let $u \in X^{p,q}([0, T]; \mathbb{T}^d)$ be a weak solution of (1.1) with $\frac{2}{p} + \frac{d}{q} = 1$. For any divergence-free $v_0 \in L^2(\mathbb{T}^d)$, there exists a weak solution $v \in C_w L^2 \cap L^2 H^1$ to the linearized Navier-Stokes equation:*

$$\begin{cases} \partial_t v - \Delta v + u \cdot \nabla v + \nabla p = 0 \\ \operatorname{div} v = 0, \end{cases} \quad (\text{A.1})$$

satisfying the energy inequality

$$\frac{1}{2} \|v(t)\|_2^2 + \int_{t_0}^t \|\nabla v(s)\|_2^2 ds \leq \frac{1}{2} \|v(t_0)\|_2^2,$$

for all $t \in [t_0, T]$, a.e. $t_0 \in [0, T]$ (including $t_0 = 0$).

Proof. This follows by a standard Galerkin method and can be found in many textbooks. See [RRS16, Chapter 4] or [Gal94] for details. \square

Let v be the weak solution given by Theorem A.2 with initial data $u(0)$. The goal is to show $u \equiv v$. Setting $w = u - v$, the equation for w reads

$$\partial_t w - \Delta w + u \cdot \nabla w + \nabla q = 0,$$

and its weak formulation

$$\int_0^T \int_{\mathbb{T}^d} w \cdot (\partial_t \varphi + \Delta \varphi + u \cdot \nabla \varphi) dx dt = 0 \quad \text{for any } \varphi \in \mathcal{D}_T. \quad (\text{A.2})$$

Fix $F \in C_c^\infty([0, T] \times \mathbb{T}^d)$. Let $\Phi : [0, T] \times \mathbb{T}^d \rightarrow \mathbb{R}^d$ and $\chi : [0, T] \times \mathbb{T}^d \rightarrow \mathbb{R}$ satisfy the system of equations

$$\begin{cases} -\partial_t \Phi - \Delta \Phi - u \cdot \nabla \Phi + \nabla \chi = F \\ \operatorname{div} \Phi = 0 \\ \Phi(T) = 0. \end{cases} \quad (\text{A.3})$$

Note that the equation of Φ is ‘‘backwards in time’’ and by a change of variable one can convert (A.3) into a more conventional form.

If we can use $\varphi = \Phi$ in the weak formulation (A.2), then immediately

$$\int_{[0, T] \times \mathbb{T}^d} w \cdot F dx dt = 0.$$

Since $F \in C_c^\infty([0, T] \times \mathbb{T}^d)$ is arbitrary, we have

$$w = 0 \quad \text{for a.e. } (t, x) \in [0, T] \times \mathbb{T}^d.$$

So the question of whether $u \equiv v$ reduces to showing a certain regularity of Φ . More specifically, we can prove the following theorem.

Theorem A.3. *Let $d \geq 2$ be the dimension and $u \in X^{p,q}([0, T]; \mathbb{T}^d)$ be a weak solution of (1.1) with $\frac{2}{p} + \frac{d}{q} = 1$, $d \leq q \leq \infty$. For any $F \in C_c^\infty([0, T] \times \mathbb{T}^d)$, the system (A.3) has a weak solution $\Phi \in L^\infty L^2 \cap L^2 H^1$ such that*

$$\partial_t \Phi, \Delta \Phi, \nabla \chi \in L^s([0, T] \times \mathbb{T}^d) \quad \text{for some } s > 1.$$

Proof. The Galerkin method is based on solving the following finite-dimensional approximation

$$\begin{cases} -\partial_t \Phi_n - \Delta \Phi_n - P_n [u_n \cdot \nabla \Phi_n] + \nabla \chi_n = F_n \\ \operatorname{div} \Phi_n = 0 \\ \Phi_n(T) = 0, \end{cases} \quad (\text{A.4})$$

where Φ_n, u_n, χ_n, F_n are restricted to the first n Fourier modes, and P_n is the projection operator on those modes.

It suffices to verify *a priori* estimates, as they are preserved in the limit as $n \rightarrow \infty$.

Step 1: Energy bounds $L^\infty L^2 \cap L^2 H^1$.

The solutions to the Galerkin approximation (A.4) enjoy the energy estimate

$$-\frac{1}{2} \frac{d}{dt} \|\Phi\|_2^2 + \|\nabla \Phi\|_2^2 \leq \int_{\mathbb{T}^d} |F \cdot \Phi| dx,$$

which implies the desired energy bounds.

Step 2: Higher bounds $L_t^\infty H^1 \cap L_t^2 H^2$ when $2 < q \leq \infty$.

We now take the L^2 inner product of (A.4) with $\Delta \Phi$ to obtain

$$-\frac{1}{2} \frac{d}{dt} \|\nabla \Phi\|_2^2 + \|\Delta \Phi\|_2^2 \leq \int_{\mathbb{T}^d} |F \cdot \Delta \Phi| dx + \int_{\mathbb{T}^d} |u \cdot \nabla \Phi \cdot \Delta \Phi| dx.$$

- **Case 1:** $p < \infty$ and $d < q \leq \infty$.

Thanks to Proposition A.4,

$$\int_{\mathbb{T}^d} |u \cdot \nabla \Phi \cdot \Delta \Phi| dx \lesssim \|u\|_q \|\nabla \Phi\|_2^{\frac{q-d}{q}} \|\Delta \Phi\|_2^{\frac{q+d}{q}}. \quad (\text{A.5})$$

Then Hölder's, Poincaré's, and Young's inequalities yield

$$-\frac{1}{2} \frac{d}{dt} \|\nabla \Phi\|_2^2 + \frac{1}{2} \|\Delta \Phi\|_2^2 \lesssim \|u\|_q^{\frac{2q}{q-d}} \|\nabla \Phi\|_2^2 + \|F\|_2^2.$$

Thanks to the integrability of $\|u\|_q^{\frac{2q}{q-d}}$ ($\|u\|_\infty^2$ when $q = \infty$), Gronwall's inequality immediately imply that $\Phi \in L^\infty H^1 \cap L^2 H^2$.

- **Case 2:** $p = \infty$ and $q = d > 2$.

Since $u \in C_t L^d$, for any $\varepsilon > 0$ there exists a decomposition $u = u_1 + u_2$ such that

$$\|u_1\|_{C_t L^d} \leq \varepsilon \quad \text{and} \quad u_2 \in L_{t,x}^\infty,$$

and then the u_1 portion of the nonlinear term in (A.5) can be absorbed by $\|\Delta \Phi\|_2^2$, so we arrive at the same conclusion.

Step 2 continue: Higher bounds $L_t^\infty H^{\frac{1}{2}} \cap L_t^2 H^{\frac{3}{2}}$ when $q = d = 2$.

To fix the lack of embedding when $d = q = 2$, we take the L^2 inner product of (A.4) with $\Lambda \Phi$, where $\Lambda^s := (-\Delta)^{\frac{s}{2}}$ is the fractional Laplacian on \mathbb{T}^d for zero mean functions, and obtain

$$-\frac{1}{2} \frac{d}{dt} \|\Lambda^{\frac{1}{2}} \Phi\|_2^2 + \|\Lambda^{\frac{3}{2}} \Phi\|_2^2 \leq \int_{\mathbb{T}^d} |F \cdot \Lambda \Phi| dx + \int_{\mathbb{T}^d} |u \cdot \nabla \Phi \cdot \Lambda \Phi| dx.$$

Then the Sobolev embedding $H^{\frac{1}{2}}(\mathbb{T}^2) \hookrightarrow L^4(\mathbb{T}^2)$ gives

$$\int_{\mathbb{T}^d} |u \cdot \nabla \Phi \cdot \Lambda \Phi| dx \lesssim \|u\|_2 \|\nabla \Phi\|_4^2 \lesssim \|u\|_2 \|\Lambda^{\frac{3}{2}} \Phi\|_2^2. \quad (\text{A.6})$$

With (A.6) in hand, the above decomposition $u = u_1 + u_2$ and Gronwall's inequality imply that $\Phi \in L^\infty H^{\frac{1}{2}} \cap L^2 H^{\frac{3}{2}}$.

Step 3: Conclusion from maximal regularity of the heat equation.

Taking Leary's projection \mathbb{P} onto the divergence-free vector fields, the equation for Φ can be rewritten as

$$\begin{cases} -\partial_t \Phi - \Delta \Phi = \mathbb{P}(u \cdot \nabla \Phi) + \mathbb{P}F \\ \Phi(T) = 0. \end{cases}$$

Therefore, we may use the maximal $L_t^p L^q$ regularity of the heat equation to conclude the proof.

- **Case 1:** $q \neq 2$: $\Phi \in L^\infty H^1 \cap L^2 H^2$.
Since $u \in L^p L^q$, $\frac{2}{p} + \frac{d}{q} = 1$ and $\nabla \Phi \in L^\infty L^2$,

$$u \cdot \nabla \Phi \in L^s([0, T] \times \mathbb{T}^d) \quad \text{for some } s > 1.$$

Thus by the maximal regularity of the heat equation

$$\partial_t \Phi, \Delta \Phi \in L^s([0, T] \times \mathbb{T}^d) \quad \text{for some } s > 1.$$

- **Case 2:** $q = 2$: $\Phi \in L^\infty H^{\frac{1}{2}} \cap L^2 H^{\frac{3}{2}}$.

By the argument in Case 1, it suffices to show that $\nabla \Phi \in L^r_{t,x}$ for some $r > 2$. By a Sobolev interpolation $H^1 = [H^{\frac{1}{2}}, H^{\frac{3}{2}}]$, we first infer that $\nabla \Phi \in L^4 L^2$. Since $\nabla \Phi \in L^2 H^{\frac{1}{2}}$, by the Sobolev embedding $H^{\frac{1}{2}}(\mathbb{T}^2) \hookrightarrow L^4(\mathbb{T}^2)$, we have $\nabla \Phi \in L^2 L^4$. From $\nabla \Phi \in L^4 L^2 \cap L^2 L^4$ it follows by a Hölder interpolation that $\nabla \Phi \in L^r_{t,x}$ for $r = \frac{8}{3} > 2$.

Now that we have shown the estimates for $\partial_t \Phi$ and $\Delta \Phi$, the same holds true for $\nabla \chi$ thanks to the fact that

$$\Delta \chi = \operatorname{div}(u \cdot \nabla \Phi) + \operatorname{div} F.$$

□

The last result is a classical estimate of the nonlinear term, cf. [RRS16, pp. 172]. Notice that one of the embedding fails when $q = 2$ which is the reason we prove Theorem A.3 for $q > 2$ and $q = 2$ separately.

Proposition A.4. *Let $d \geq 2$ be the dimension and $d \leq q \leq \infty$ such that $q > 2$. For any smooth vector fields $u, v \in C_0^\infty(\mathbb{T}^d)$,*

$$\int_{\mathbb{T}^d} |u \cdot \nabla v \cdot \Delta v| dx \lesssim \|u\|_q \|\nabla v\|_2^{\frac{q-d}{q}} \|\Delta v\|_2^{\frac{q+d}{q}}.$$

Proof. We apply Hölder's inequality with exponents $\frac{1}{q} + \frac{1}{r} + \frac{1}{2} = 1$, $r = \frac{2q}{q-2} \in [2, \frac{2d}{d-2})$,

$$\int_{\mathbb{T}^d} |u \cdot \nabla v \cdot \Delta v| dx \lesssim \|u\|_q \|\nabla v\|_r \|\Delta v\|_2.$$

Since $2 \leq r < \infty$, by the Sobolev embedding $H^s(\mathbb{T}^d) \hookrightarrow L^r(\mathbb{T}^d)$, $s = d(\frac{1}{2} - \frac{1}{r})$,

$$\|\nabla v\|_r \lesssim \|\nabla v\|_{H^s}.$$

Finally, by a standard Sobolev interpolation and the L^2 -boundedness of Riesz transform, we have

$$\|\nabla v\|_{H^s} \lesssim \|\nabla v\|_2^{\frac{q-d}{q}} \|\Delta v\|_2^{\frac{d}{q}},$$

which concludes the proof. □

APPENDIX B. SOME TECHNICAL TOOLS

B.1. Improved Hölder's inequality on \mathbb{T}^d . We recall the following result due to Modena and Székelyhidi [MS18], which was inspired by [BV19, Lemma 3.7]. This lemma allows us to quantify the decorrelation in the usual Hölder's inequality when we increase the oscillation of one function.

Lemma B.1. *Let $p \in [1, \infty]$ and $a, f : \mathbb{T}^d \rightarrow \mathbb{R}$ be smooth functions. Then for any $\sigma \in \mathbb{N}$,*

$$\left| \|af(\sigma \cdot)\|_p - \|a\|_p \|f\|_p \right| \lesssim \sigma^{-\frac{1}{p}} \|a\|_{C^1} \|f\|_p. \quad (\text{B.1})$$

The proof is based on the interplay between the Poincaré's inequality and the fast oscillation of $f(\sigma \cdot)$ and can be found in [MS18, Lemma 2.1].

B.2. Tensor-valued antidivergence \mathcal{R} . For any $f \in C^\infty(\mathbb{T}^d)$, there exists a $v \in C_0^\infty(\mathbb{T}^d)$ such that

$$\Delta v = f - \int_{\mathbb{T}^d} f.$$

And we denote v by $\Delta^{-1}f$. Note that if $f \in C_0^\infty(\mathbb{T}^d)$, then by rescaling we have

$$\Delta^{-1}(f(\sigma \cdot)) = \sigma^{-2}v(\sigma \cdot) \quad \text{for } \sigma \in \mathbb{N}.$$

We recall the following antidivergence operator \mathcal{R} introduced in [DLS13].

Definition B.2. $\mathcal{R} : C^\infty(\mathbb{T}^d, \mathbb{R}^d) \rightarrow C^\infty(\mathbb{T}^d, \mathcal{S}_0^{d \times d})$ is defined by

$$(\mathcal{R}v)_{ij} = \mathcal{R}_{ijk}v_k \tag{B.2}$$

where

$$\mathcal{R}_{ijk} = -\frac{2-d}{d-1}\Delta^{-2}\partial_i\partial_j\partial_k - \frac{1}{d-1}\Delta^{-1}\partial_k\delta_{ij} + \Delta^{-1}\partial_i\delta_{jk} + \Delta^{-1}\partial_j\delta_{ik}.$$

It is clear that \mathcal{R} is well-defined since \mathcal{R}_{ijk} is symmetric in i, j and taking the trace gives

$$\begin{aligned} \text{Tr } \mathcal{R}v &= \frac{2-d}{d-1}\Delta^{-1}\partial_k v_k + \frac{-d}{d-1}\Delta^{-1}\partial_k v_k + \Delta^{-1}\partial_k v_k + \Delta^{-1}\partial_k v_k \\ &= \left(\frac{2-d}{d-1} + \frac{-d}{d-1} + 2\right)\Delta^{-1}\partial_k v_k = 0. \end{aligned}$$

By a direct computation, one can also show that

$$\text{div}(\mathcal{R}v) = v - \int_{\mathbb{T}^d} v \quad \text{for any } v \in C^\infty(\mathbb{T}^d, \mathbb{R}^d)$$

and

$$\text{div}(\mathcal{R}\Delta v) = \nabla v + \nabla v^T \quad \text{for any divergence-free } v \in C^\infty(\mathbb{T}^d, \mathbb{R}^d). \tag{B.3}$$

We can show that \mathcal{R} is bounded on $L^p(\mathbb{T}^d)$ for any $1 \leq p \leq \infty$.

Theorem B.3. Let $1 \leq p \leq \infty$. For any $f \in C_0^\infty(\mathbb{T}^d)$, there holds

$$\|\mathcal{R}f\|_{L^p(\mathbb{T}^d)} \lesssim \|f\|_{L^p(\mathbb{T}^d)}.$$

Proof. We only show that the operator

$$\Delta^{-1}\Delta^{-1}\partial_i\partial_j\partial_k$$

is bounded on $L^p(\mathbb{T}^d)$ for $1 \leq p \leq \infty$ since the argument applies also to $\Delta^{-1}\partial_i$.

When $1 < p < \infty$, this follows from the boundedness of the Riesz transforms and the Poincare inequality.

When $p = \infty$, the Sobolev embedding $W^{1, d+1}(\mathbb{T}^d) \hookrightarrow L^\infty(\mathbb{T}^d)$ implies that

$$\|\Delta^{-1}\Delta^{-1}\partial_i\partial_j\partial_k f\|_{L^\infty(\mathbb{T}^d)} \lesssim \|\Delta^{-1}\Delta^{-1}\partial_i\partial_j\partial_k f\|_{W^{1, d+1}(\mathbb{T}^d)} \lesssim \|f\|_{L^{d+1}(\mathbb{T}^d)} \leq \|f\|_{L^\infty(\mathbb{T}^d)}$$

where we have used the boundedness of the Riesz transforms once again.

When $p = 1$, it suffices to consider $f \in L^1(\mathbb{T}^d)$ with zero mean. Then one can use a duality approach and use the boundedness in L^∞ since integrating by parts yields

$$\langle \Delta^{-1}\Delta^{-1}\partial_i\partial_j\partial_k f, \varphi \rangle = -\langle f, \Delta^{-1}\Delta^{-1}\partial_i\partial_j\partial_k \varphi \rangle \quad \text{if } \varphi \in C_0^\infty(\mathbb{T}^d).$$

□

B.3. Bilinear antidivergence \mathcal{B} . We can also introduce the bilinear version $\mathcal{B} : C^\infty(\mathbb{T}^d, \mathbb{R}^d) \times C^\infty(\mathbb{T}^d, \mathcal{S}_0^{d \times d}) \rightarrow C_0^\infty(\mathbb{T}^d, \mathcal{S}_0^{d \times d})$ of \mathcal{R} . Let

$$(\mathcal{B}(v, A))_{ij} = v_l \mathcal{R}_{ijk} A_{lk} - \mathcal{R}(\partial_i v_l \mathcal{R}_{ijk} A_{lk})$$

or by a slight abuse of notations

$$\mathcal{B}(v, A) = v\mathcal{R}A - \mathcal{R}(\nabla v \mathcal{R}A).$$

Theorem B.4. Let $1 \leq p \leq \infty$. For any $v \in C^\infty(\mathbb{T}^d)$ and $A \in C_0^\infty(\mathbb{T}^d, \mathbb{R}^d)$,

$$\text{div}(\mathcal{B}(v, A)) = vA - \int_{\mathbb{T}^d} vA, \tag{B.4}$$

and

$$\|\mathcal{B}(a, v)\|_{L^p(\mathbb{T}^d)} \lesssim \|v\|_{C^1(\mathbb{T}^d)} \|f\|_{L^p(\mathbb{T}^d)}.$$

Proof. A direct computation gives

$$\begin{aligned} \operatorname{div}(\mathcal{B}(v, A)) &= \partial_j v_l \mathcal{R}_{ijk} A_{lk} + v_l \partial_j \mathcal{R}_{ijk} A_{lk} - \operatorname{div} \mathcal{R}(\partial_i v_l \mathcal{R}_{ijk} A_{lk}) \\ &= v_l A_{il} + \int \partial_i v_l \mathcal{R}_{ijk} A_{lk} \end{aligned}$$

where we have used the fact that A has zero mean and \mathcal{R} is symmetric.

Integrating by parts, we have

$$\int \partial_i v_l \mathcal{R}_{ijk} A_{lk} = - \int v_l \partial_i \mathcal{R}_{ijk} A_{lk} = - \int v_l A_{lj},$$

which implies that

$$\operatorname{div}(\mathcal{B}(v, A)) = vA - \int vA.$$

The second estimate follows immediately from the definition of \mathcal{B} and Theorem B.3. □

Acknowledgement. AC was partially supported by the NSF grant DMS–1909849.

REFERENCES

- [Ama00] Herbert Amann. On the strong solvability of the Navier-Stokes equations. *J. Math. Fluid Mech.*, 2(1):16–98, 2000.
- [BCV18] Tristan Buckmaster, Maria Colombo, and Vlad Vicol. Wild solutions of the Navier-Stokes equations whose singular sets in time have Hausdorff dimension strictly less than 1. *arXiv:1809.00600*, 2018.
- [BDLIS15] Tristan Buckmaster, Camillo De Lellis, Philip Isett, and László Székelyhidi, Jr. Anomalous dissipation for 1/5-Hölder Euler flows. *Ann. of Math. (2)*, 182(1):127–172, 2015.
- [BDLS16] Tristan Buckmaster, Camillo De Lellis, and László Székelyhidi, Jr. Dissipative Euler flows with Onsager-critical spatial regularity. *Comm. Pure Appl. Math.*, 69(9):1613–1670, 2016.
- [BKM84] J. T. Beale, T. Kato, and A. Majda. Remarks on the breakdown of smooth solutions for the 3-D Euler equations. *Comm. Math. Phys.*, 94(1):61–66, 1984.
- [BLJV18] T. Buckmaster, C. De Lellis, L. Székelyhidi Jr., and V. Vicol. Onsager’s conjecture for admissible weak solutions. *Comm. Pure Appl. Math.*, to appear, 2018.
- [BSV19] Tristan Buckmaster, Steve Shkoller, and Vlad Vicol. Nonuniqueness of weak solutions to the SQG equation. *Comm. Pure Appl. Math.*, 72(9):1809–1874, 2019.
- [BV19] Tristan Buckmaster and Vlad Vicol. Nonuniqueness of weak solutions to the Navier-Stokes equation. *Ann. of Math. (2)*, 189(1):101–144, 2019.
- [BV21] Tristan Buckmaster and Vlad Vicol. Convex integration constructions in hydrodynamics. *Bull. Am. Math. Soc.*, to appear, 2021.
- [CCFS08] A. Cheskidov, P. Constantin, S. Friedlander, and R. Shvydkoy. Energy conservation and Onsager’s conjecture for the Euler equations. *Nonlinearity*, 21(6):1233–1252, 2008.
- [CET94] P. Constantin, W. E, and E. S. Titi. Onsager’s conjecture on the energy conservation for solutions of Euler’s equation. *Comm. Math. Phys.*, 165(1):207–209, 1994.
- [Che11] Jean-Yves Chemin. About weak-strong uniqueness for the 3D incompressible Navier-Stokes system. *Comm. Pure Appl. Math.*, 64(12):1587–1598, 2011.
- [CKN82] L. Caffarelli, R. Kohn, and L. Nirenberg. Partial regularity of suitable weak solutions of the Navier-Stokes equations. *Comm. Pure Appl. Math.*, 35(6):771–831, 1982.
- [CL20] Alexey Cheskidov and Xiaoyutao Luo. Nonuniqueness of weak solutions for the transport equation at critical space regularity. *arXiv:2004.09538*, 2020.
- [CV14] Kyudong Choi and Alexis F. Vasseur. Estimates on fractional higher derivatives of weak solutions for the Navier-Stokes equations. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 31(5):899–945, 2014.
- [CZ16] Jean-Yves Chemin and Ping Zhang. On the critical one component regularity for 3-D Navier-Stokes systems. *Ann. Sci. Éc. Norm. Supér. (4)*, 49(1):131–167, 2016.
- [DD07] Hongjie Dong and Dapeng Du. Partial regularity of solutions to the four-dimensional Navier-Stokes equations at the first blow-up time. *Comm. Math. Phys.*, 273(3):785–801, 2007.
- [DLS09] C. De Lellis and L. Székelyhidi, Jr. The Euler equations as a differential inclusion. *Ann. of Math. (2)*, 170(3):1417–1436, 2009.
- [DLS13] Camillo De Lellis and László Székelyhidi, Jr. Dissipative continuous Euler flows. *Invent. Math.*, 193(2):377–407, 2013.
- [DLS14] Camillo De Lellis and László Székelyhidi, Jr. Dissipative Euler flows and Onsager’s conjecture. *J. Eur. Math. Soc. (JEMS)*, 16(7):1467–1505, 2014.
- [DS17] Sara Daneri and László Székelyhidi, Jr. Non-uniqueness and h-principle for Hölder-continuous weak solutions of the Euler equations. *Arch. Ration. Mech. Anal.*, 224(2):471–514, 2017.
- [ESŠ03] L. Escauriaza, G. A. Seregin, and V. Šverák. $L_{3,\infty}$ -solutions of Navier-Stokes equations and backward uniqueness. *Uspekhi Mat. Nauk*, 58(2(350)):3–44, 2003.
- [Eyi94] Gregory L. Eyink. Energy dissipation without viscosity in ideal hydrodynamics. I. Fourier analysis and local energy transfer. *Phys. D*, 78(3-4):222–240, 1994.

- [FJR72] E. B. Fabes, B. F. Jones, and N. M. Rivière. The initial value problem for the Navier-Stokes equations with data in L^p . *Arch. Rational Mech. Anal.*, 45:222–240, 1972.
- [FK64] Hiroshi Fujita and Tosio Kato. On the Navier-Stokes initial value problem. I. *Arch. Rational Mech. Anal.*, 16:269–315, 1964.
- [FLRT00] Giulia Furioli, Pierre G. Lemarié-Rieusset, and Elide Terraneo. Unicité dans $L^3(\mathbb{R}^3)$ et d’autres espaces fonctionnels limites pour Navier-Stokes. *Rev. Mat. Iberoamericana*, 16(3):605–667, 2000.
- [Fri95] Uriel Frisch. *Turbulence*. Cambridge University Press, Cambridge, 1995. The legacy of A. N. Kolmogorov.
- [Gal94] Giovanni P. Galdi. *An introduction to the mathematical theory of the Navier-Stokes equations. Vol. II*, volume 39 of *Springer Tracts in Natural Philosophy*. Springer-Verlag, New York, 1994. Nonlinear steady problems.
- [Gal00] Giovanni P. Galdi. An introduction to the Navier-Stokes initial-boundary value problem. In *Fundamental directions in mathematical fluid mechanics*. Adv. Math. Fluid Mech., pages 1–70. Birkhäuser, Basel, 2000.
- [Ger06] P. Germain. Multipliers, paramultipliers, and weak-strong uniqueness for the Navier-Stokes equations. *J. Differential Equations*, 226(2):373–428, 2006.
- [GKP16] Isabelle Gallagher, Gabriel S. Koch, and Fabrice Planchon. Blow-up of critical Besov norms at a potential Navier-Stokes singularity. *Comm. Math. Phys.*, 343(1):39–82, 2016.
- [GŠ17] J. Guillod and V. Šverák. Numerical investigations of non-uniqueness for the Navier-Stokes initial value problem in borderline spaces. [arXiv:1704.00560](https://arxiv.org/abs/1704.00560), 2017.
- [Hop51] E. Hopf. Über die anfangswertaufgabe für die hydrodynamischen grundgleichungen. erhard schmidt zu seinem 75. geburtstag gewidmet. *Math. Nachr.*, 4(1-6):213–231, 1951.
- [Ise17] Philip Isett. On the Endpoint Regularity in Onsager’s Conjecture. [arXiv:1706.01549](https://arxiv.org/abs/1706.01549), 2017.
- [Ise18] Philip Isett. A proof of Onsager’s conjecture. *Ann. of Math. (2)*, 188(3):871–963, 2018.
- [JŠ15] Hao Jia and Vladimír Šverák. Are the incompressible 3d Navier-Stokes equations locally ill-posed in the natural energy space? *J. Funct. Anal.*, 268(12):3734–3766, 2015.
- [Kat84] Tosio Kato. Strong L^p -solutions of the Navier-Stokes equation in \mathbf{R}^m , with applications to weak solutions. *Math. Z.*, 187(4):471–480, 1984.
- [KT01] Herbert Koch and Daniel Tataru. Well-posedness for the Navier-Stokes equations. *Adv. Math.*, 157(1):22–35, 2001.
- [Lad67] O. A. Ladyženskaja. Uniqueness and smoothness of generalized solutions of Navier-Stokes equations. *Zap. Naučn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)*, 5:169–185, 1967.
- [Lad69] O A Ladyženskaja. Example of nonuniqueness in the hopf class of weak solutions for the Navier-Stokes equations. *Mathematics of the USSR-Izvestiya*, 3(1):229–236, feb 1969.
- [Ler34] J. Leray. Sur le mouvement d’un liquide visqueux emplissant l’espace. *Acta Math.*, 63(1):193–248, 1934.
- [LM01] P.-L. Lions and N. Masmoudi. Uniqueness of mild solutions of the Navier-Stokes system in L^N . *Comm. Partial Differential Equations*, 26(11-12):2211–2226, 2001.
- [LQ20] Tianwen Luo and Peng Qu. Non-uniqueness of weak solutions to 2D hypoviscous Navier-Stokes equations. *J. Differential Equations*, 269(4):2896–2919, 2020.
- [LR16] Pierre Gilles Lemarié-Rieusset. *The Navier-Stokes problem in the 21st century*. CRC Press, Boca Raton, FL, 2016.
- [LT20] Tianwen Luo and Edriss S. Titi. Non-uniqueness of weak solutions to hyperviscous Navier-Stokes equations: on sharpness of J.-L. Lions exponent. *Calc. Var. Partial Differential Equations*, 59(3):Paper No. 92, 15, 2020.
- [Luo19] Xiaoyutao Luo. Stationary solutions and nonuniqueness of weak solutions for the Navier-Stokes equations in high dimensions. *Arch. Ration. Mech. Anal.*, 233(2):701–747, 2019.
- [Man76] Benoit Mandelbrot. Intermittent turbulence and fractal dimension: kurtosis and the spectral exponent $5/3 + B$. In *Turbulence and Navier-Stokes equations (Proc. Conf., Univ. Paris-Sud, Orsay, 1975)*, pages 121–145. Lecture Notes in Math., Vol. 565. Springer, Berlin, 1976.
- [Mey97] Yves Meyer. Wavelets, paraproducts, and Navier-Stokes equations. In *Current developments in mathematics, 1996 (Cambridge, MA)*, pages 105–212. Int. Press, Boston, MA, 1997.
- [Mon99] Sylvie Monniaux. Uniqueness of mild solutions of the Navier-Stokes equation and maximal L^p -regularity. *C. R. Acad. Sci. Paris Sér. I Math.*, 328(8):663–668, 1999.
- [MS18] Stefano Modena and László Székelyhídi, Jr. Non-uniqueness for the transport equation with Sobolev vector fields. *Ann. PDE*, 4(2):Art. 18, 38, 2018.
- [Nas54] John Nash. C^1 isometric imbeddings. *Ann. of Math. (2)*, 60:383–396, 1954.
- [Nov20] Matthew Novack. Non-uniqueness of weak solutions to the 3D Quasi-Geostrophic Equations. *SIAM Journal on Mathematical Analysis*, to appear, 2020.
- [Ons49] L. Onsager. Statistical hydrodynamics. *Nuovo Cimento (9)*, 6(Suppl, Supplemento, no. 2 (Convegno Internazionale di Meccanica Statistica)):279–287, 1949.
- [Oža20] Wojciech S. Ożański. Weak Solutions to the Navier–Stokes Inequality with Arbitrary Energy Profiles. *Comm. Math. Phys.*, 374(1):33–62, 2020.
- [Pro59] G. Prodi. Un teorema di unicità per le equazioni di navier-stokes. *Ann. Mat. Pura ed Appl.*, 48(1):173–182, 1959.
- [RRS16] James C. Robinson, José L. Rodrigo, and Witold Sadowski. *The three-dimensional Navier-Stokes equations*, volume 157 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2016. Classical theory.
- [Sch76a] Vladimir Scheffer. Partial regularity of solutions to the Navier-Stokes equations. *Pacific J. Math.*, 66(2):535–552, 1976.
- [Sch76b] Vladimir Scheffer. Turbulence and Hausdorff dimension. In *Turbulence and Navier-Stokes equations (Proc. Conf., Univ. Paris-Sud, Orsay, 1975)*, pages 174–183. Lecture Notes in Math., Vol. 565, 1976.
- [Sch77] Vladimir Scheffer. Hausdorff measure and the Navier-Stokes equations. *Comm. Math. Phys.*, 55(2):97–112, 1977.
- [Sch80] Vladimir Scheffer. The Navier-Stokes equations on a bounded domain. *Comm. Math. Phys.*, 73(1):1–42, 1980.
- [Sch85] Vladimir Scheffer. A solution to the Navier-Stokes inequality with an internal singularity. *Comm. Math. Phys.*, 101(1):47–85, 1985.

- [Sch87] Vladimir Scheffer. Nearly one-dimensional singularities of solutions to the Navier-Stokes inequality. *Comm. Math. Phys.*, 110(4):525–551, 1987.
- [Sch93] Vladimir Scheffer. An inviscid flow with compact support in space-time. *J. Geom. Anal.*, 3(4):343–401, 1993.
- [Ser62] J. Serrin. On the interior regularity of weak solutions of the Navier-Stokes equations. *Arch. Rational Mech. Anal.*, 9:187–195, 1962.
- [Shn97] A. Shnirelman. On the nonuniqueness of weak solution of the Euler equation. *Comm. Pure Appl. Math.*, 50(12):1261–1286, 1997.
- [Str88] M. Struwe. On partial regularity results for the Navier-Stokes equations. *Comm. Pure Appl. Math.*, 41(4):437–458, 1988.
- [Sze13] László Székelyhidi, Jr. From isometric embeddings to turbulence. In *HCDTE lecture notes. Part II. Nonlinear hyperbolic PDEs, dispersive and transport equations*, volume 7 of *AIMS Ser. Appl. Math.*, page 63. Am. Inst. Math. Sci. (AIMS), Springfield, MO, 2013.
- [Tao19] Terence Tao. 255B, Notes 2: Onsager’s conjecture. <https://terrytao.wordpress.com/2019/01/08/255b-notes-2-onsagers-conjecture/>, 2019.
- [Wu20] Bian Wu. Partially regular weak solutions of the Navier-Stokes equations in $\mathbb{R}^4 \times [0, \infty[$. *arXiv:2008.05802*, 2020.

DEPARTMENT OF MATHEMATICS, STATISTICS AND COMPUTER SCIENCE, UNIVERSITY OF ILLINOIS AT CHICAGO, CHICAGO, ILLINOIS 60607

E-mail address: acheskid@uic.edu

DEPARTMENT OF MATHEMATICS, DUKE UNIVERSITY, DURHAM, NC 27708

E-mail address: xiaoyutao.luo@duke.edu