

HARDY AND POINCARÉ INEQUALITIES IN FRACTIONAL ORLICZ-SOBOLEV SPACES

KAUSHIK BAL, KAUSHIK MOHANTA, PROSENJIT ROY, AND FIROJ SK

ABSTRACT. We provide sufficient conditions for boundary Hardy inequality to hold in bounded Lipschitz domains, complement of a point (the so called point Hardy inequality), domain above the graph of a Lipschitz function, complement of a bounded Lipschitz domain in fractional Orlicz-Sobolev setting. As a consequence we get sufficient conditions for regional fractional Orlicz Poincaré inequality in bounded Lipschitz domains. Necessary conditions for fractional Orlicz Hardy and regional fractional Orlicz Poincaré inequalities are also given for bounded Lipschitz domains. Various sufficient conditions on open sets are provided for fractional Orlicz Poincaré inequality and regional fractional Orlicz Poincaré inequality to hold.

1. INTRODUCTION

The aim of this article is to study two very well known inequalities, the Poincaré inequality and the Hardy inequality on fractional Orlicz-Sobolev setting. The classical Poincaré inequality [Eva10, Chapter 5.8.1] states that for any bounded domain $D \subseteq \mathbb{R}^N$, $q \geq 1$ there exists $c = c(q, N, D) > 0$ such that for any $f \in C_c^\infty(D)$,

$$(1.1) \quad \|f\|_{L^q(D)} \leq c \|\nabla f\|_{L^q(D)}.$$

For bounded domains it is a standard fact that the best constant is attained. Although generalized in many different direction as can be seen in [Beb03, J⁺86, KZ08, LSY03], for the purpose of this paper we would like to refer Gossez [Gos74, Lemma 5.7], who generalized eq. (1.1) in Orlicz-Sobolev setting (defined below). The standard boundary Hardy inequality states that if D is a bounded Lipschitz domain in \mathbb{R}^N , $q \geq 1$, then there exists a constant $c = c(q) > 0$ such that for $f \in C_c^\infty(D)$,

$$(1.2) \quad \|f/\delta_D\|_{L^q(D)} \leq c \|\nabla f\|_{L^q(D)},$$

where $\delta_D(x) := \inf\{|x-y| : y \in D^c\}$. We shall frequently denote $\delta_D(x)$ by δ_x . This has been further developed in [BM97, BMS00, KMP07, MMP98] and several other works. For the generalization to Orlicz version of local Hardy inequality we refer to [Hei95, KaPPu09, MMOS14, MNOS11].

We shall primarily be concerned with the “Orlicz space” $L^A(D)$ and the “fractional Orlicz-Sobolev space” $W^{s,A}(D)$. We start by defining these spaces. A continuous, convex function $A : [0, \infty) \rightarrow [0, \infty)$ such that $\lim_{t \rightarrow 0} \frac{A(t)}{t} = 0$ and $\lim_{t \rightarrow \infty} \frac{A(t)}{t} = \infty$ is called an **N -function** or an **Young function** [AF03, Chapter 8.2].

Definition 1 ([AF03, Chapter 8.2]). *We say that A satisfies the Δ_2 condition globally or simply the Δ_2 condition ($A \in \Delta_2$) if there exists constant $p > 0$ such that*

$$A(2t) \leq pA(t), \quad \forall t > 0.$$

The set

$$L^A(D) := \left\{ f : D \rightarrow \mathbb{R} \text{ measurable} \mid \exists \lambda > 0 \text{ such that } M_{L^A(D)}\left(\frac{f}{\lambda}\right) < \infty \right\}$$

INDIAN INSTITUTE OF TECHNOLOGY KANPUR, INDIA

E-mail addresses: kaushik@iitk.ac.in, kmohanta@iitk.ac.in, prosenjit@iitk.ac.in, firoj@iitk.ac.in.

2020 *Mathematics Subject Classification.* 35B27; 46E35; 49J52.

Key words and phrases. fractional Hardy inequality; regional fractional Poincaré inequality; fractional Poincaré inequality; fractional Orlicz Sobolev spaces; unbounded domains.

is called the **Orlicz space** and the set

$$W^{s,A}(D) := \left\{ f \in L^A(D) \mid \exists \lambda > 0 \text{ such that } M_{W^{s,A}(D)}\left(\frac{f}{\lambda}\right) < \infty \right\}$$

is called the **fractional Orlicz-Sobolev space**, where

$$M_{L^A(D)}(f) := \int_D A(|f(x)|) dx \quad \text{and} \quad M_{W^{s,A}(D)}(f) := \iint_{D \times D} A\left(\frac{|f(x) - f(y)|}{|x - y|^s}\right) \frac{dxdy}{|x - y|^N}.$$

In the case $A(t) = t^q$ for some $q > 1$, $L^A(D)$ and $W^{s,A}(D)$ are well known Lebesgue space $L^q(D)$ and the fractional Sobolev space $W^{s,q}(D)$ respectively [DNPV12, p. 524].

For $s \in (0, 1)$, we start by defining the following important quotients that will be used frequently throughout this article:

$$H_{N,s,A}(D) := \inf_{\substack{f \in C_c^\infty(D) \\ f \neq 0}} \frac{M_{W^{s,A}(D)}(f)}{\int_D A\left(\frac{|f(x)|}{\delta_x^s}\right) dx}, \quad P_{N,s,A}^1(D) := \inf_{\substack{f \in C_c^\infty(D) \\ f \neq 0}} \frac{M_{W^{s,A}(D)}(f)}{\int_D A(|f(x)|) dx}$$

and

$$P_{N,s,A}^2(D) := \inf_{\substack{f \in C_c^\infty(D) \\ f \neq 0}} \frac{M_{W^{s,A}(\mathbb{R}^N)}(f)}{\int_D A(|f(x)|) dx}.$$

We shall say that

- fractional Orlicz Hardy inequality (denoted as $FOHI(s, A)$) holds if $H_{N,s,A}(D) > 0$;
- regional fractional Orlicz Poincaré inequality ($RFOPI(s, A)$) holds if $P_{N,s,A}^1(D) > 0$;
- fractional Orlicz Poincaré inequality ($FOPI(s, A)$) holds if $P_{N,s,A}^2(D) > 0$.

We start by recalling some literature on fractional Orlicz Hardy inequality for the case when $A(t) = t^q$, $q > 1$. Kufner [KT78] proved $FOHI(s, t^q)$ in one dimension when $D = (0, \infty)$. For bounded Lipschitz domain D , Dyda [Dyd04, Theorem 1.1 and Section 2] proved the following results:

Theorem 1.1 (Dyda). *Let $\beta > 0$ and $q > 1$. The Hardy inequality,*

$$(1.3) \quad \int_D \frac{|u(x)|^q}{\delta_D(x)^\beta} dx \leq c \iint_{D \times D} \frac{|u(x) - u(y)|^q}{|x - y|^{N+\beta}} dxdy, \quad \text{for all } u \in C_c(D),$$

where $c = c(D, \beta, N, q) < \infty$ is a constant, holds true in each of the following cases:

- (T1) D is a bounded Lipschitz domain and $\beta > 1$;
- (T2) D is a complement of a bounded Lipschitz domain, $\beta \neq 1$ and $\beta \neq N$;
- (T3) D is a domain above the graph of a Lipschitz function $\mathbb{R}^{N-1} \rightarrow \mathbb{R}$ and $\beta \neq 1$;
- (T4) D is a complement of a point and $\beta \neq N$.

The question of best constant in fractional Hardy inequality, that is the exact value of $FOHI(s, t^q)$, was first addressed in [BD11, Theorem 1] for upper half space. Heinig *et al.* [HKP97, Theorem 3.1] and Kufner *et al.* [KP03, Theorem 5.23] studied one dimensional $FOHI(s, t^q)$ between two weighted L^q spaces. Reader may refer to [BC18, CS03, Dyd11, DF12, DV14, EHSV14, FL12, FS08, HKP97, HSV15, ILTV14, LS10] and references therein for more information related to fractional Hardy inequality.

Concerning $RFOPI(s, A)$ and $FOPI(s, A)$, we start with the trivial observation that $P_{N,s,A}^2(D) \geq P_{N,s,A}^1(D)$, that is $FOPI(s, A)$ holds whenever $RFOPI(s, A)$ is true. For a bounded domain D , $RFOPI(s, t^2)$ is true if and only if $2s > 1$ [Che18, Proposition 3.2] whereas $FOPI(s, t^2)$ holds for all $s \in (0, 1)$ [CR20, Theorem 1.2]. It was first established in [Yer14, Lemma 1] that $FOPI(s, t^2)$ is true for all values of s , if the domain is an infinite strip i.e. $D = (0, 1) \times \mathbb{R}^{N-1}$, though the best constant was not established. The best constant for the above case is obtained in

[CR20, Theorem 1.3]. $RFOPI(s, t^q)$ for strip like domain is further studied in [CCRS21, Theorem 1.1], where it is established that $RFOPI(s, t^2)$ is true if and only if $2s > 1$. Sufficient criteria on domains for $RFOPI(s, t^2)$ and $FOPI(s, t^2)$ to hold, is provided in [CCRS21, Theorem 1.1-1.4] and [CR20, Theorem 1.2] respectively. For other related works on fractional Poincaré inequality and its applications, we refer the reader to [CS03, CR20, CR17], and the references therein.

As of now the theory of fractional Orlicz-Sobolev spaces is still at an early stage of development. Bonder-Salort [FBS19, Section 2.2] defined the space $W^{s,A}(\mathbb{R}^N)$ and established basic properties of the space. For an open bounded set $D \subseteq \mathbb{R}^N$, the space $\{f \in L^A(D) \mid M_{W^{s,A}(\mathbb{R}^N)}(f) < \infty\}$ and corresponding fractional Laplacian have been considered in [SV20, Sal20a]. The fractional Orlicz-Sobolev space $W^{s,A}(D)$ has appeared in Bahrouni *et al.* [BBX20, Section 2] for any open bounded set $D \subseteq \mathbb{R}^N$. For other works on fractional Orlicz-Sobolev spaces one may refer to [ACPS20b, ACPS20a, ABS18, BO20, BOT20, BS21, CdP20, FHR20, MSV]. Salort [Sal20b, Theorem 1.1] proved $FOHI$ in one dimension. In [ACPS21, Theorem 7.4], the authors proved $FOHI$ in the special case where the domain is \mathbb{R}^N . Fractional Orlicz Hardy inequality and regional fractional Orlicz Poincaré inequality for any domain or fractional Orlicz Poincaré inequality for unbounded domains have not yet appeared in the literature (apart from [Sal20b, ACPS21] as mentioned above). For $A \in \Delta_2$, an open bounded set D , $s \in (0, 1)$, the following version of $FOPI(s, A)$ is proved in [FBS19, Theorem 6.1 and Corollary 6.2]:

$$M_{L^A(D)}(f) \leq cM_{W^{s,A}(\mathbb{R}^N)}(f), \quad f \in L^A(D)$$

where $c = c(A, s, N) > 0$. Similar results are also proved in [SV20, Sal20a] for general N -function (without Δ_2 condition).

Primary aim of this article is to generalize theorem 1.1 in the fractional Orlicz setting. This is achieved via theorems 1.2, 1.4 and 1.5. Theorem 1.2 also provides sufficient condition on A and s for $RFOPI(s, A)$ to hold on any bounded Lipschitz domain, whereas theorem 1.3 gives sufficient condition, on bounded Lipschitz domains, for $FOHI(s, A)$ and $RFOPI(s, A)$ not to hold. As an application of theorems 1.2 and 1.3, at the end of section 3 (see table 1), we give several examples of $A \in \Delta_2$ and respective ranges of s for which Hardy and regional Poincaré inequality is true or false. In particular for N -functions t^q , $q > 1$ and $(1+t)\log(1+t) - t$ a complete answer, for all values of $s \in (0, 1)$ can be provided for $FOHI(s, A)$. At this point, we would like to mention that the line of argument in the proofs of theorems 1.2, 1.4 and 1.5 are adapted from [Dyd04] where Dyda, in fact, predicted the possibility of his methods being generalized. Secondary objective of this article is to study $RFOPI(s, A)$ and $FOPI(s, A)$ for unbounded domains in \mathbb{R}^N . In this direction our results are proposition 5.1 and theorem 1.6. In proposition 5.1 we give complete characterization of domains in 1-dimension for which $RFOPI(s, A)$ holds, provided $\lim_{\lambda \rightarrow 0+} \alpha_{s,A}(\lambda) = 0$ (see eq. (1.4)).

Theorem 1.6 provides different sufficient criteria on domains for $FOPI(s, A)$ and $RFOPI(s, A)$ to hold. As an application of theorem 1.6, at the end of section 5 we provide several non-trivial examples of domains for which $RFOPI(s, A)$ and $FOPI(s, A)$ holds. An adapted version of a change of variable formula introduced in [LS10, Lemma 2.4] is the key ingredient in the proof of theorem 1.6 which can be regarded as the fractional Orlicz analogue of the results obtained in [CR20, Theorem 1.2] and [CCRS21, Theorem 1.3].

Our main results are stated below.

Theorem 1.2. *Let $D \subseteq \mathbb{R}^N$ be a bounded Lipschitz domain, $A \in \Delta_2$, $s \in (0, 1)$ and $\liminf_{\lambda \rightarrow 0+} \alpha_{s,A}(\lambda) = 0$, where $\alpha_{s,A} : [0, \infty) \rightarrow \mathbb{R}$ is defined by*

$$(1.4) \quad \alpha_{s,A}(\lambda) := \sup_{t \in [0, \infty)} \frac{A(\lambda t)}{\lambda^{\frac{1}{s}} A(t)}.$$

Then $FOHI(s, A)$ and $RFOPI(s, A)$ holds, that is $H_{N,s,A}(D) > 0$ and $P_{N,s,A}^1(D) > 0$.

Theorem 1.3. Let $D \subseteq \mathbb{R}^N$ be a bounded Lipschitz domain, A be an N -function and

$$(1.5) \quad \lim_{\varepsilon \rightarrow 0+} \varepsilon \int_0^{\varepsilon^{-s}} \frac{A(z)}{z} dz = \beta \in \mathbb{R}.$$

Then

- (1) if $\beta = 0$, then both $FOHI(s, A)$ and $RFOPI(s, A)$ do not hold,
- (2) if $\beta \in (0, \infty)$, then $FOHI(s, A)$ does not hold.

Theorem 1.4. Let $s \in (0, 1)$, $A \in \Delta_2$ and $\alpha_{s,A}$ be as in eq. (1.4). Assume $\liminf_{\lambda \rightarrow 0+} \lambda^{\frac{1-N}{s}} \alpha_{s,A}(\lambda) = 0$ or $\liminf_{\lambda \rightarrow \infty} \lambda^{\frac{1-N}{s}} \alpha_{s,A}(\lambda) = 0$. Then $FOHI(s, A)$ holds in $\mathbb{R}^N \setminus \{0\}$, that is $H_{N,s,A}(\mathbb{R}^N \setminus \{0\}) > 0$.

Theorem 1.5. Suppose $D \subseteq \mathbb{R}^N$ be an open set, $A \in \Delta_2$, $s \in (0, 1)$ and $\alpha_{s,A}$ be as in eq. (1.4). Then $FOHI(s, A)$ holds true, that is $H_{N,s,A}(D) > 0$, in each of the following cases:

- (1) $D = \{(x', x_N) \in \mathbb{R}^N \mid x' \in \mathbb{R}^{N-1}, x_N > \Phi(x')\}$, where $\Phi : \mathbb{R}^{N-1} \rightarrow \mathbb{R}$ is a Lipschitz map and s, A are such that $\liminf_{\lambda \rightarrow 0} \alpha_{s,A}(\lambda) = 0$ or $\liminf_{\lambda \rightarrow \infty} \alpha_{s,A}(\lambda) = 0$;
- (2) D^c is closure of some bounded Lipschitz domain and s, A are such that $\liminf_{\lambda \rightarrow \infty} \lambda^{\frac{1-N}{s}} \alpha_{s,A}(\lambda) = 0 = \liminf_{\lambda \rightarrow 0+} \alpha_{s,A}(\lambda)$ or $\liminf_{\lambda \rightarrow 0+} \lambda^{\frac{1-N}{s}} \alpha_{s,A}(\lambda) = 0$ or $\liminf_{\lambda \rightarrow \infty} \alpha_{s,A}(\lambda) = 0$.

Before giving the statement of our last result, that is [theorem 1.6](#), we define some terminologies that will be required to formulate it in a precise manner.

Definition 2. We say that a set $D \subseteq \mathbb{R}^N$ satisfies **the finite ball condition** if D does not contain arbitrarily large balls, that is if

$$BC(D) := \sup\{r : B(x, r) \subseteq D, x \in D\} < \infty.$$

Definition 3. Let $\{D_\beta\}_\beta$ be a family of sets in \mathbb{R}^N , where $\beta \in \Lambda$ (some indexing set). We say that the **FOPI(s, A)** holds uniformly for $\{D_\beta\}_\beta$ if $\inf_\beta P_{N,s,A}^2(D_\beta) > 0$.

Let $\omega \in \mathbb{S}^{N-1}$ and $x \in \omega^\perp$, define $L_D(x, \omega) := \{t \mid x + t\omega \in D\} \subseteq \mathbb{R}$.

Definition 4. We say an open set $D \subseteq \mathbb{R}^N$ is of **type LS(s, A)** if there exists $\Sigma \subseteq \mathbb{S}^{N-1}$ with positive $(N-1)$ -dimensional Hausdorff measure, such that uniform $FOPI(s, A)$ holds for the family $\{L_D(x, \omega)\}_{\omega \in \Sigma, x \in \omega^\perp}$.

Theorem 1.6. Let $D \subseteq \mathbb{R}^N$ be an open set, $s \in (0, 1)$, A be an N -function and $\alpha_{s,A}$ be as in eq. (1.4).

- (1) Assume that $A \in \Delta_2$ and $\lim_{\lambda \rightarrow 0+} \alpha_{s,A}(\lambda) = 0$. Let there exist $\Sigma \subseteq \mathbb{S}^{N-1}$ with positive $(N-1)$ -dimensional Hausdorff measure such that $\sup_{\omega \in \Sigma, x \in \omega^\perp} BC(L_D(x, \omega)) < \infty$. Then the $RFOPI(s, A)$ holds true in D .
- (2) Assume that there exist $R, c_1 > 0$ such that $\mathcal{L}^N(B(x, R) \cap D^c) > c_1$ for any $x \in D$. Then $FOPI(s, A)$ holds in $D \forall s \in (0, 1)$.
- (3) $FOPI(s, A)$ holds if D is an $LS(s, A)$ domain.

This paper is arranged in the following way: In [section 2](#), some necessary preliminaries are discussed. Proof of [theorems 1.2](#) and [1.3](#) followed by some applications are given in [section 3](#). [Theorems 1.4](#) and [1.5](#) are proved in [section 4](#). In [section 5](#) we prove [proposition 5.1](#) and [theorem 1.6](#).

2. NOTATIONS AND PRELIMINARIES

Throughout the paper the following conventions and notations will be followed, unless mentioned otherwise explicitly:

D will denote an open set in \mathbb{R}^N , $s \in (0, 1)$, A will denote an N -function, \mathcal{L}^k will denote the Lebesgue measure on \mathbb{R}^k , \mathbb{S}^{k-1} will denote the unit sphere in \mathbb{R}^k , c will denote a generic constant which may change from line to line, X^c will stand for the complement of the set X in appropriate universal set (to be understood from the context), for any real valued function f or for any Lipschitz domain D , $\text{Lip}(f)$ or $\text{Lip}(D)$ will denote the Lipschitz constant, $\alpha_{s,A}$ will be as in [eq. \(1.4\)](#), p will be as in [definition 1](#).

We start with some basic facts about N -functions. One may refer to [\[AF03, KR61\]](#) for detailed discussion on the topic.

Lemma 2.1 ([\[AF03, Chapter 8.2\]](#)). *A is an N -function if and only if there exists a non-decreasing, right continuous function $a : [0, \infty) \rightarrow [0, \infty)$ satisfying $a(0) = 0$, $a(t) > 0$ for $t > 0$, $\lim_{t \rightarrow 0^+} a(t) = \infty$ such that*

$$A(x) = \int_0^x a(t) dt.$$

The following two lemmas will be used frequently in the rest of the article.

Lemma 2.2 ([\[AF03, Chapter 8.2\]](#)). *Let A be an N -function, then A and $t \mapsto \frac{A(t)}{t}$ both are strictly increasing function on $(0, \infty)$.*

Lemma 2.3. *Let $A \in \Delta_2$. Then*

$$A(\lambda t) \leq \lambda^p A(t) \text{ for } t \in [0, \infty), \quad \forall \lambda \geq 1,$$

where p is the constant in [definition 1](#). The above inequality is equivalent to

$$A(\lambda t) \geq \lambda^p A(t) \text{ for } t \in [0, \infty), \quad \forall 0 \leq \lambda \leq 1.$$

Proof. Since $A \in \Delta_2$, there exists $p > 0$ such that $A(2t) \leq pA(t)$, $\forall t > 0$. Now for a as in [lemma 2.1](#), using the non decreasing property of a ,

$$pA(t) \geq A(2t) = \int_0^{2t} a(\tau) d\tau > \int_t^{2t} a(\tau) d\tau > ta(t).$$

This implies for any $\lambda > 1$,

$$\log \left(\frac{A(\lambda t)}{A(t)} \right) = \int_t^{\lambda t} \frac{a(\tau)}{A(\tau)} d\tau < \int_t^{\lambda t} \frac{p}{\tau} d\tau = p \log \left(\frac{\lambda t}{t} \right) = \log(\lambda^p).$$

The lemma follows. □

Lemma 2.4. *Let $D \subseteq \mathbb{R}^N$ be a bounded domain, $A \in \Delta_2$. Then for some constant $c = c(D, A) > 0$,*

$$P_{N,s,A}^1(D) \geq cH_{N,s,A}(D).$$

Proof. Let $f \in C_c^\infty(D)$. First assume $\text{diam}(D) \leq 1$. By the monotonicity of A ,

$$\int_D A \left(\frac{|f(x)|}{\delta_x^s} \right) dx \geq \int_D A(|f(x)|) dx.$$

In the case $\text{diam}(D) > 1$, we exploit the Δ_2 condition of A and use [lemma 2.3](#) to get

$$\int_D A \left(\frac{|f(x)|}{\delta_x^s} \right) dx \geq \int_D A(\text{diam}(D)^{-s} |f(x)|) dx \geq \text{diam}(D)^{-sp} \int_D A(|f(x)|) dx.$$

Hence the lemma follows. □

Proposition 2.1. *Let $s \in (0, 1)$ and $p > 1$ be as in [lemma 2.3](#).*

- (1) *If $D_1 \subseteq D_2 \subseteq \mathbb{R}^N$, then $P_{N,s,A}^2(D_2) \leq P_{N,s,A}^2(D_1)$.*
- (2) *Let $D \subseteq \mathbb{R}^N$ be an open set and $u \in W^{s,A}(D)$. Assume that $A \in \Delta_2$. For $t > 0$, define $v_t \in W^{s,A}(D)$ by $v_t(x) = u(tx)$. Then*

$$\iint_{D \times D} A \left(\frac{|v_t(x) - v_t(y)|}{|x - y|^s} \right) \frac{dxdy}{|x - y|^N} \leq \begin{cases} t^{s-N} \iint_{tD \times tD} A \left(\frac{|u(x) - u(y)|}{|x - y|^s} \right) \frac{dxdy}{|x - y|^N}, & t < 1, \\ t^{sp-N} \iint_{tD \times tD} A \left(\frac{|u(x) - u(y)|}{|x - y|^s} \right) \frac{dxdy}{|x - y|^N}, & t \geq 1, \end{cases}$$

and also

$$\iint_{D \times D} A \left(\frac{|v_t(x) - v_t(y)|}{|x - y|^s} \right) \frac{dxdy}{|x - y|^N} \geq \begin{cases} t^{s-N} \iint_{tD \times tD} A \left(\frac{|u(x) - u(y)|}{|x - y|^s} \right) \frac{dxdy}{|x - y|^N}, & t \geq 1, \\ t^{sp-N} \iint_{tD \times tD} A \left(\frac{|u(x) - u(y)|}{|x - y|^s} \right) \frac{dxdy}{|x - y|^N}, & t < 1. \end{cases}$$

Furthermore,

$$\begin{aligned} \frac{P_{N,s,A}^1(D)}{t^s} &\leq P_{N,s,A}^1(tD) \leq \frac{P_{N,s,A}^1(D)}{t^{sp}} \quad \text{if } t < 1, \\ \frac{P_{N,s,A}^1(D)}{t^{sp}} &\leq P_{N,s,A}^1(tD) \leq \frac{P_{N,s,A}^1(D)}{t^s} \quad \text{if } t \geq 1. \end{aligned}$$

- (3) *Let $D \subseteq \mathbb{R}^N$ be an open set and $t > 0$. Assume that $A \in \Delta_2$. Then*

$$\begin{aligned} \frac{P_{N,s,A}^2(D)}{t^s} &\leq P_{N,s,A}^2(tD) \leq \frac{P_{N,s,A}^2(D)}{t^{sp}} \quad \text{if } t < 1, \\ \frac{P_{N,s,A}^2(D)}{t^{sp}} &\leq P_{N,s,A}^2(tD) \leq \frac{P_{N,s,A}^2(D)}{t^s} \quad \text{if } t \geq 1. \end{aligned}$$

- (4) *Let $D \subseteq \mathbb{R}^N$ be such that $BC(D) = \infty$, then $P_{N,s,A}^1(D) = 0 = P_{N,s,A}^2(D)$.*

Proof. (1) Directly follows from the definition.

- (2) For the first inequality, take the change of variable $X = tx, Y = ty$ and in case of $t < 1$, use [lemma 2.2](#) to infer $\frac{A(t^s f(x,y))}{t^s f(x,y)} < \frac{A(f(x,y))}{f(x,y)}$. In case of $t \geq 1$ use [lemma 2.3](#). For the second inequality replace t by $\frac{1}{t}$.

- (3) Similar as case (2).

- (4) $BC(D) = \infty$ implies there exist a positive sequence $\{r_n\}_n$ and a sequence $\{x_n\}_n$ such that $r_n \rightarrow \infty$ and $B(x_n, r_n) \subseteq D$. Note that $P_{N,s,A}^1(D) \leq P_{N,s,A}^2(D)$. Then by (1) and (3) we infer that

$$P_{N,s,A}^2(D) \leq P_{N,s,A}^2(B(x_n, r_n)) = P_{N,s,A}^2(B(0, r_n)) \leq \frac{P_{N,s,A}^2(B(0, 1))}{r_n^s} \rightarrow 0.$$

This finishes the proof. \square

Lemma 2.5. *Let $D \subsetneq \mathbb{R}$ be an open set with $BC(D) < \infty$ and $D = \cup_{k=1}^{\infty} I_k$, where I_k 's are disjoint intervals. Then for any $k \in \mathbb{N}$, $P_{1,s,A}^1(I_k) \geq \min\{BC(D)^{-sp}, 1\} P_{1,s,A}^1((0, 1))$, where $p > 1$ is as in [lemma 2.3](#).*

Proof. Using (2) of [proposition 2.1](#), we have $P_{1,s,A}^1(I_k) \geq \text{diam}(I_k)^{-\beta} P_{1,s,A}^1((0, 1))$, where

$$\beta = \begin{cases} s, & \text{if } \text{diam}(I_k) < 1, \\ sp, & \text{if } \text{diam}(I_k) \geq 1. \end{cases}$$

If $\text{diam}(I_k) < 1$, $P_{1,s,A}^1(I_k) \geq P_{1,s,A}^1(0, 1)$. On the other hand if $\text{diam}(I_k) \geq 1$,

$$P_{1,s,A}^1(I_k) \geq \text{diam}(I_k)^{-sp} P_{1,s,A}^1((0, 1)) \geq BC(D)^{-sp} P_{1,s,A}^1((0, 1)).$$

Combining the two cases we finally get the desired result. \square

3. $FOHI(s, A)$ AND $RFOPI(s, A)$ ON BOUNDED LIPSCHITZ DOMAINS

In this section we shall prove [theorems 1.2](#) and [1.3](#). Our proof of [theorem 1.2](#) is motivated by [\[Dyd04\]](#). We start this section by proving some technical lemmas which will be used in the proof of [theorem 1.2](#). Let $D \subsetneq \mathbb{R}^N$ be a non-empty open set. Throughout the section we shall assume, unless stated otherwise, $\Omega \subseteq D$, $A \in \Delta_2$ and p is as in [definition 1](#).

We now fix a function $f \in C_c^\infty(D)$ and define

$$(3.1) \quad G = G(f, \Omega; l_1, l_2) := \left\{ x \in \Omega \mid A\left(\frac{|f(x)|}{\delta_x^s}\right) > \frac{2^{p+1}}{l_2 \delta_x^N} \int_{B(x, l_1 \delta_x) \cap \Omega} A\left(\frac{|f(x) - f(y)|}{\delta_x^s}\right) dy \right\};$$

where $l_1 > 1$, l_2 are positive numbers, independent of f , whose values are given later.

Lemma 3.1. *Let f, l_1, l_2 be as above. Then*

$$\int_{\Omega \setminus G} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx \leq \frac{2^{p+1} l_1^{N+sp}}{l_2} \int_{x \in \Omega \setminus G} \int_{y \in \Omega} A\left(\frac{|f(x) - f(y)|}{|x - y|^s}\right) \frac{dy dx}{|x - y|^N}.$$

Proof. For $x \in \Omega \setminus G$, using [lemmas 2.2](#) and [2.3](#), we have

$$\begin{aligned} \int_{\Omega} A\left(\frac{|f(x) - f(y)|}{|x - y|^s}\right) \frac{dy}{|x - y|^N} &\geq \int_{B(x, l_1 \delta_x) \cap \Omega} A\left(\frac{|f(x) - f(y)|}{|x - y|^s}\right) \frac{dy}{|x - y|^N} \\ &\geq \int_{B(x, l_1 \delta_x) \cap \Omega} A\left(\frac{|f(x) - f(y)|}{(l_1 \delta_x)^s}\right) \frac{dy}{(l_1 \delta_x)^N} \\ &\geq \frac{1}{l_1^{N+sp} \delta_x^N} \int_{B(x, l_1 \delta_x) \cap \Omega} A\left(\frac{|f(x) - f(y)|}{\delta_x^s}\right) dy \geq \frac{l_2}{l_1^{N+sp} 2^{p+1}} A\left(\frac{|f(x)|}{\delta_x^s}\right). \end{aligned}$$

Integrating over $\Omega \setminus G$, we get the desired result. \square

In the view of the above lemma, we can say that the set $\Omega \setminus G$ is a good set for fractional Orlicz Hardy inequality in the sense that the required inequality holds in it.

For any $x \in G$, let us define the set $E^*(x) = \left\{ y \in E \mid \frac{|f(x)|}{2} \leq |f(y)| \leq \frac{3}{2}|f(x)| \right\}$.

Lemma 3.2. *Let $x \in G$ and $E \subseteq B(x, l_1 \delta_x) \cap \Omega$. If $\mathcal{L}^N(E) \geq l_2 \delta_x^N$, then*

$$\frac{\mathcal{L}^N(E)}{2} \leq \mathcal{L}^N(E) - \frac{l_2 \delta_x^N}{2} \leq \mathcal{L}^N(E^*(x)).$$

Proof. Since $x \in G$, $E \subseteq B(x, l_1 \delta_x) \cap \Omega$, by [eq. \(3.1\)](#) we have

$$A\left(\frac{|f(x)|}{\delta_x^s}\right) \geq \frac{2^{p+1}}{l_2 \delta_x^N} \int_{B(x, l_1 \delta_x) \cap \Omega} A\left(\frac{|f(x) - f(y)|}{\delta_x^s}\right) dy \geq \frac{2^{p+1}}{l_2 \delta_x^N} \int_{E \setminus E^*(x)} A\left(\frac{|f(x) - f(y)|}{\delta_x^s}\right) dy.$$

Note that, $y \in E \setminus E^*(x)$ implies $|f(x) - f(y)| \geq ||f(x)| - |f(y)|| \geq \frac{|f(x)|}{2}$. Thus by using [lemmas 2.2](#) and [2.3](#) we get

$$A\left(\frac{|f(x)|}{\delta_x^s}\right) \geq \frac{2}{l_2 \delta_x^N} A\left(\frac{|f(x)|}{\delta_x^s}\right) \{ \mathcal{L}^N(E) - \mathcal{L}^N(E^*(x)) \}.$$

Using the hypothesis on measure of E , we get $\frac{\mathcal{L}^N(E)}{2} \leq \mathcal{L}^N(E) - \frac{l_2 \delta_x^N}{2} \leq \mathcal{L}^N(E^*(x))$. \square

Lemma 3.3. *Let $E_1 \subseteq \Omega$ and $E_2 \subseteq B(x, l_1 \delta_x) \cap \Omega$ be such that $\mathcal{L}^N(E_2) \geq l_2 \delta_x^N$ for all $x \in E_1$. Then*

$$\int_{E_1 \cap G} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx \leq \frac{2^{p+1} \mathcal{L}^N(E_1)}{\mathcal{L}^N(E_2)} \int_{E_2} A\left(\frac{\sup\{\delta_x^s \mid x \in E_2\} |f(y)|}{\inf\{\delta_x^s \mid x \in E_1\} \delta_y^s}\right) dy.$$

Proof. Assume that $E_1 \cap G$ is non empty. First, we fix $\eta > 1$ and pick $x_0 \in E_1 \cap G$ such that $\sup_{x \in E_1 \cap G} |f(x)| \leq \eta |f(x_0)|$. Also $|f(x_0)| \leq 2|f(y)|$ for any $y \in E_2^*(x_0)$. Hence, for any $y \in E_2^*(x_0)$, using [lemmas 2.2](#) and [2.3](#),

$$\begin{aligned} \int_{E_1 \cap G} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx &\leq \mathcal{L}^N(E_1 \cap G) A\left(\frac{\sup_{x \in E_1 \cap G} |f(x)|}{\inf\{\delta_x^s \mid x \in E_1 \cap G\}}\right) \\ &\leq \mathcal{L}^N(E_1 \cap G) A\left(\frac{2\eta |f(y)|}{\inf\{\delta_x^s \mid x \in E_1 \cap G\}}\right) \leq 2^p \eta^p \mathcal{L}^N(E_1) A\left(\frac{\sup\{\delta_y^s \mid y \in E_2^*(x_0)\} |f(y)|}{\inf\{\delta_x^s \mid x \in E_1\} \delta_y^s}\right). \end{aligned}$$

Integrating over $y \in E_2^*(x_0)$, using [lemma 3.2](#) and the fact that $E_2^*(x_0) \subseteq E_2$, we obtain

$$\int_{E_1 \cap G} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx \leq \frac{2^{p+1} \eta^p \mathcal{L}^N(E_1)}{\mathcal{L}^N(E_2)} \int_{E_2} A\left(\frac{\sup\{\delta_x^s \mid x \in E_2\} |f(y)|}{\inf\{\delta_x^s \mid x \in E_1\} \delta_y^s}\right) dy.$$

Letting $\eta \rightarrow 1$ the proof follows. \square

Lemma 3.4. Let $\Omega \subseteq D \subseteq \mathbb{R}^N$ be two open sets, $f \in C_c^\infty(D)$, $l_1, l_2 > 0$, $G = G(f, \Omega; l_1, l_2)$, $0 < \gamma < 1$ and $m \in \mathbb{N}$. Assume $\Omega = \cup_{j=0}^\infty A_j$, $\mathcal{L}^N(A_i \cap A_j) = 0 \ \forall j \neq i$ and $\exists n_0 \in \mathbb{N}$ such that $f \equiv 0$ on A_j for $j \geq n_0$.

(1) If $\forall j \in \mathbb{N}$,

$$\int_{G \cap A_j} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx \leq \gamma \int_{A_{j+m}} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx,$$

then there exists a constant $c > 0$ such that

$$\int_{\Omega} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx \leq c \iint_{\Omega \times \Omega} A\left(\frac{|f(x) - f(y)|}{|x - y|^s}\right) \frac{dx dy}{|x - y|^N}.$$

(2) If $\forall j \geq m$,

$$\int_{G \cap A_j} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx \leq \gamma \int_{A_{j-m}} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx,$$

then there exists a constant $c > 0$ such that

$$\int_{\Omega} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx \leq c \iint_{\Omega \times \Omega} A\left(\frac{|f(x) - f(y)|}{|x - y|^s}\right) \frac{dx dy}{|x - y|^N} + c \int_{A_0 \cup \dots \cup A_{m-1}} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx.$$

Proof. (1)

$$\begin{aligned} \int_{\Omega \cap G} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx &= \sum_{j=0}^\infty \int_{A_j \cap G} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx \leq \sum_{j=0}^\infty \gamma \int_{A_{j+m}} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx \\ &= \sum_{j=0}^\infty \gamma \left[\int_{A_{j+m} \cap G} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx + \int_{A_{j+m} \setminus G} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx \right] \\ &\leq \sum_{j=0}^\infty \gamma \left[\gamma \int_{A_{j+2m}} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx + \int_{A_{j+m} \setminus G} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx \right] \\ &\leq \dots \leq \sum_{j=0}^\infty \sum_{k=1}^\infty \gamma^k \int_{A_{j+km} \setminus G} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx. \end{aligned}$$

Here we used the fact that $f \equiv 0$ for large j forcing the iterative process to terminate. So we have

$$\int_{\Omega \cap G} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx \leq \sum_{k=1}^{\infty} \gamma^k \int_{\Omega \setminus G} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx.$$

Hence by using above estimate and [lemma 3.1](#) we get

$$\begin{aligned} \int_{\Omega} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx &= \int_{\Omega \cap G} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx + \int_{\Omega \setminus G} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx \\ &\leq \sum_{k=0}^{\infty} \gamma^k \int_{\Omega \setminus G} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx \leq c \int_{\Omega} \int_{\Omega} A\left(\frac{|f(x) - f(y)|}{|x - y|^s}\right) \frac{dx dy}{|x - y|^N}. \end{aligned}$$

This completes the proof of (1).

(2) Let for each $j \geq 0$, k_j be the largest nonnegative integer such that $j - k_j m \geq 0$, so that $0 \leq j - k_j m \leq m - 1$. Then k_j -many repeated applications of the inequality in the hypothesis gives

$$\begin{aligned} \int_{A_j \cap G} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx &\leq \gamma \int_{A_{j-m}} A\left(\frac{|f(y)|}{\delta_y^s}\right) dy \\ &= \gamma \left[\int_{A_{j-m} \setminus G} A\left(\frac{|f(y)|}{\delta_y^s}\right) dy + \int_{A_{j-m} \cap G} A\left(\frac{|f(y)|}{\delta_y^s}\right) dy \right] \\ &\leq \dots \leq \gamma^{k_j} \int_{A_{j-k_j m} \cap G} A\left(\frac{|f(y)|}{\delta_y^s}\right) dy + \sum_{k=1}^{k_j} \gamma^k \int_{A_{j-km} \setminus G} A\left(\frac{|f(y)|}{\delta_y^s}\right) dy \\ &\leq \gamma^{k_j} \int_{A_0 \cup \dots \cup A_{m-1}} A\left(\frac{|f(y)|}{\delta_y^s}\right) dy + \sum_{k=1}^{k_j} \gamma^k \int_{A_{j-km} \setminus G} A\left(\frac{|f(y)|}{\delta_y^s}\right) dy. \end{aligned}$$

Summing over j ,

$$\int_{\Omega \cap G} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx \leq \frac{\gamma}{1 - \gamma} \left[\int_{A_0 \cup \dots \cup A_{m-1}} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx + \int_{\Omega \setminus G} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx \right].$$

Proceeding as in part (1), the lemma follows. \square

Before proving [theorem 1.2](#), we need a geometric decomposition of a bounded Lipschitz domain D , given in [[Dyd04](#), p. 581]. We outline the construction for the sake of completeness.

Let us denote $x = (x_1, x_2, \dots, x_{N-1}, x_N) = (\tilde{x}, x_N) \in \mathbb{R}^N$ with $\tilde{x} \in \mathbb{R}^{N-1}$, $x_N \in \mathbb{R}$. $D \subsetneq \mathbb{R}^N$ shall be assumed to be a bounded Lipschitz domain throughout the rest of the section. For any $z \in \partial D$ there are linear isometry $L_z : \mathbb{R}^N \rightarrow \mathbb{R}^N$ and Lipschitz function $\phi_z : \mathbb{R}^{N-1} \rightarrow \mathbb{R}$ such that $\text{Lip}(\phi_z) \leq \text{Lip}(D)$ and

$$L_z(D) \cap B(L_z(z), r_0) = \{x \in \mathbb{R}^N \mid x_N > \phi_z(\tilde{x})\} \cap B(L_z(z), r_0),$$

for some positive r_0 which depends only on D (existence of r_0 is guaranteed because ∂D is compact). Without loss of generality, we can assume L_z to be the identity map. Because otherwise we can work with the Lipschitz domain $L_z(D)$ and the point $L_z(z)$, then pull back the construction to the original domain D and the point z via L_z^{-1} . For $x \in \mathbb{R}^N$ we set

$$V_z(x) := |x_N - \phi_z(\tilde{x})|.$$

For $E \subseteq \mathbb{R}^{N-1}$ and $r > 0$ define

$$Q_z(E, r) := \{x \in D \mid \tilde{x} \in E, 0 < V_z(x) \leq r\}.$$

Set

$$K_r := \left\{x \in \mathbb{R}^{N-1} \mid |x_\ell - z_\ell| \leq \frac{r}{2}, \ell = 1, 2, \dots, N-1\right\}.$$

That is K_r is an $N-1$ dimensional square of side-length r . $Q_z(K_\rho, r)$ is referred to as a Lipschitz box and is denoted by $Q_z(r)$. We now choose $\rho > 0$ small enough such that $Q_z(\rho) \subseteq D \cap B(z, r_0/2)$. Let $x \in Q_z(\rho)$. Then using the fact that $\phi_z(w) = w_N$ for any $w \in \partial D \cap B(z, r_0)$,

$$V_z(x) = |x_N - \phi_z(\tilde{x})| \leq |x_N - w_N| + |\phi_z(\tilde{w}) - \phi_z(\tilde{x})| \leq |x_N - w_N| + \text{Lip}(D)|\tilde{w} - \tilde{x}| \leq (1 + \text{Lip}(D))|w - x|.$$

Since w is arbitrary, we get

$$(3.2) \quad \frac{V_z(x)}{(1 + \text{Lip}(D))} \leq \delta_x \leq V_z(x), \quad \forall x \in Q_z(\rho).$$

For $j \in \mathbb{N} \cup \{0\}$ consider the dyadic decomposition of K_ρ (by dissecting sides first and then decomposing the cube into smaller cubes and proceeding by induction) into the union of $(N-1)$ -dimensional cubes K_j^i for $1 \leq i \leq 2^{j(N-1)}$. Now for each j , K_j^i 's have disjoint interiors and have sides-length $\frac{\rho}{2^j}$. Set $Q_j^i := Q_z(K_j^i, \rho)$ and define

$$(3.3) \quad A_k = Q_z(K_\rho, \rho/2^{k+1}) \setminus Q_z(K_\rho, \rho/2^k), \text{ for } k \in \mathbb{N} \cup \{0\}.$$

Then A_k 's are mutually disjoint sets with $\cup_{k=0}^\infty A_k = Q_z(\rho)$. Thus the Lipschitz box $Q_z(\rho)$ is the union of sets $A_j \cap Q_j^i$, $i, j \in \mathbb{N}$ whose pairwise intersections are measure zero sets. Moreover using eqs. (3.2) and (3.3), for $k \geq j$, we have

$$(3.4) \quad \frac{\rho}{2^{k+1}(1 + \text{Lip}(D))} \leq \delta_x \leq \frac{\rho}{2^k}, \text{ for } x \in A_k \cap Q_j^i$$

and

$$(3.5) \quad \mathcal{L}^N(A_k \cap Q_j^i) = \left(\frac{\rho}{2^k} - \frac{\rho}{2^{k+1}}\right) \left(\frac{\rho}{2^j}\right)^{N-1} = \frac{\rho^N}{2^{k+1+jN-j}}.$$

Let $x \in A_k \cap Q_j^i$ and $y \in A_j \cap Q_j^i$ where $k \geq j$. Then $\frac{\rho}{2^{k+1}} \leq |x_N - \phi_z(\tilde{x})| \leq \frac{\rho}{2^k}$ and $\frac{\rho}{2^{j+1}} \leq |y_N - \phi_z(\tilde{y})| \leq \frac{\rho}{2^j}$. Moreover both x and y lie 'above' the same dyadic cube Q_j^i which has side length $\frac{\rho}{2^j}$. So for $1 \leq \ell \leq N-1$, $|x_\ell - y_\ell| \leq \frac{\rho}{2^j}$. Therefore $|\tilde{x} - \tilde{y}| \leq \sqrt{N-1} \frac{\rho}{2^j}$. Again, using these


$$\begin{aligned} |x_N - y_N| &= |x_N - \phi_z(\tilde{x}) + \phi_z(\tilde{x}) - \phi_z(\tilde{y}) + \phi_z(\tilde{y}) - y_N| \leq \frac{\rho}{2^k} + \text{Lip}(D)|\tilde{x} - \tilde{y}| + \frac{\rho}{2^j} \\ &\leq \frac{\rho}{2^{j-1}} + \text{Lip}(D)\sqrt{N-1} \frac{\rho}{2^j}. \end{aligned}$$

So we finally get

$$\begin{aligned} |x - y|^2 &\leq \left(\sqrt{N-1} \frac{\rho}{2^j}\right)^2 + \left(\frac{\rho}{2^{j-1}} + \text{Lip}(D)\sqrt{N-1} \frac{\rho}{2^j}\right)^2 \\ &= \left(\frac{\rho}{2^j}\right)^2 \left(N-1 + \left(2 + \text{Lip}(D)\sqrt{N-1}\right)^2\right). \end{aligned}$$

This gives us for $k \geq j$,

$$(3.6) \quad |x - y| \leq \frac{\rho}{2^j} \sqrt{N-1 + (\text{Lip}(D)\sqrt{N-1} + 2)^2} \quad \text{for } x \in A_k \cap Q_j^i \text{ and } y \in A_j \cap Q_j^i.$$

FIGURE 1. Lipschitz box $Q_z(\rho)$, $N = 2$.

For the next result we consider $Q = Q_z(\rho)$ defined above.

Lemma 3.5. *Let $\lim_{\lambda \rightarrow 0+} \alpha_{s,A}(\lambda) = 0$, where $\alpha_{s,A}$ is as in eq. (1.4). Then there exists a constant $c = c(D, N, A, s) > 0$ such that for all $f \in C_c^\infty(D)$ we have*

$$\int_{Q_z(\rho)} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx \leq c \iint_{Q_z(\rho) \times Q_z(\rho)} A\left(\frac{|f(x) - f(y)|}{|x - y|^s}\right) \frac{dxdy}{|x - y|^N}.$$

Proof. From the hypothesis on s, A it follows that there exists $m \in \mathbb{N}$ such that

$$2^{2+p}(1 + \text{Lip}(D))\alpha_{s,A}(2^{s-ms}(1 + \text{Lip}(D))^s) < \frac{1}{2}.$$

Set

$$l_1 = 2(1 + \text{Lip}(D))\sqrt{N - 1 + (\text{Lip}(D)\sqrt{N - 1} + 2)^2} \text{ and } l_2 = \frac{1}{2^{m+1}}.$$

Let $f \in C_c^\infty(D)$ and $G = G(f, Q_z(\rho); l_1, l_2)$. Set $E_1 := A_j \cap Q_j^i$ and $E_2 := A_{j+m} \cap Q_j^i$. For these choices of l_1 and l_2 , from eqs. (3.4) and (3.5), we get $\mathcal{L}^N(E_2) = \frac{\rho^N}{2^{1+m+jN}}$ and $\delta_x \leq \frac{\rho}{2^{j+m}}$ for $x \in E_2$. Also utilizing eqs. (3.4) and (3.6), $E_2 \subseteq B(x, l_1\delta_x)$. Thus the sets E_1 and E_2 satisfy the hypotheses of lemma 3.3. Thus we obtain, using lemma 3.3, eq. (3.4) and the choice of m ,

$$\begin{aligned} \int_{(A_j \cap Q_j^i) \cap G} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx &\leq 2^{1+p+m} \int_{A_{j+m} \cap Q_j^i} A\left(2^{s-ms}(1 + \text{Lip}(D))^s \frac{|f(y)|}{\delta_y^s}\right) dy \\ &= 2^{2+p}(1 + \text{Lip}(D)) \int_{A_{j+m} \cap Q_j^i} \frac{A\left(2^{s-ms}(1 + \text{Lip}(D))^s \frac{|f(y)|}{\delta_y^s}\right)}{2^{1-m}(1 + \text{Lip}(D))A\left(\frac{|f(y)|}{\delta_y^s}\right)} A\left(\frac{|f(y)|}{\delta_y^s}\right) dy \\ &\leq 2^{2+p}(1 + \text{Lip}(D))\alpha_{s,A}(2^{s-ms}(1 + \text{Lip}(D))^s) \int_{A_{j+m} \cap Q_j^i} A\left(\frac{|f(y)|}{\delta_y^s}\right) dy \leq \frac{1}{2} \int_{A_{j+m} \cap Q_j^i} A\left(\frac{|f(y)|}{\delta_y^s}\right) dy. \end{aligned}$$

Summing over $i = 1, \dots, 2^{j(N-1)}$ we obtain

$$(3.7) \quad \int_{A_j \cap G} A \left(\frac{|f(x)|}{\delta_x^s} \right) dx \leq \frac{1}{2} \int_{A_{j+m}} A \left(\frac{|f(x)|}{\delta_x^s} \right) dx.$$

The lemma follows from [lemma 3.4](#). □

Proof of Theorem 1.2. Consider the following two sets

$$D_1 = \{x \in D \mid \delta_x \geq \tilde{\rho}\}, \text{ and } D_2 = \{x \in D \mid \delta_x < \tilde{\rho}\},$$

where $\tilde{\rho} > 0$ is sufficiently small. Then by [lemma 3.5](#) and compactness of ∂D , we have

$$(3.8) \quad \int_{D_2} A \left(\frac{|f(x)|}{\delta_x^s} \right) dx \leq \sigma c \int_{D_2} \int_{D_2} A \left(\frac{|f(x) - f(y)|}{|x - y|^s} \right) \frac{dxdy}{|x - y|^N},$$

since D_2 may be covered by sets of the form $Q_{z_k}(\rho)$ such that every $x \in D_2$ belongs to at-most $\sigma = \sigma(D, N) \in \mathbb{N}$ sets of type $Q_{z_k}(\rho)$. This is possible for sufficiently small $\tilde{\rho} < \rho$, and such a $\tilde{\rho}$ may be chosen to depend only on $\text{Lip}(D)$, N , r_0 . We now take $G = G(f, D; l_1, l_2)$, where $l_1 = \frac{\text{diam}(D)}{\rho}$ and $l_2 = \frac{\mathcal{L}^N(D_2)}{\text{diam}(D)^N}$. Set $E_1 = D_1$ and $E_2 = D_2$. Then for any $x \in E_1$,

$$l_1 \delta_x = \text{diam}(D) \frac{\delta_x}{\rho} \geq \text{diam}(D),$$

which implies $E_2 \subseteq B(x, l_1 \delta_x)$ and

$$\mathcal{L}^N(E_2) = \text{diam}(D)^N l_2 \geq \delta_x^N l_2.$$

We can now apply [lemma 3.3](#) to get

$$(3.9) \quad \int_{D_1 \cap G} A \left(\frac{|f(x)|}{\delta_x^s} \right) dx \leq c(E_1, E_2) \int_{D_2} A \left(\frac{|f(x)|}{\delta_x^s} \right) dx,$$

and also from [lemma 3.1](#) we get

$$(3.10) \quad \int_{D_1 \setminus G} A \left(\frac{|f(x)|}{\delta_x^s} \right) dx \leq c \int_{D_1 \setminus G} \int_D A \left(\frac{|f(x) - f(y)|}{|x - y|^s} \right) \frac{dydx}{|x - y|^N}.$$

Combining the estimates [eq. \(3.8\)](#), [eq. \(3.9\)](#) and [eq. \(3.10\)](#) we conclude $H_{N,s,A}(D) > 0$. If $A \in \Delta_2$, then using [lemma 2.4](#), $P_{N,s,A}^1(D) > 0$. This completes the proof. □

Proof of Theorem 1.3. For any $\varepsilon \in (0, 1)$ denote $D_\varepsilon := \{x \in D \mid \text{dist}(x, \partial D) < \varepsilon\}$. Let $f_\varepsilon \in C_c^\infty(D)$ be such that $f_\varepsilon \equiv 1$ on $D \setminus D_\varepsilon$, $0 \leq f_\varepsilon \leq 1$ on D and $|\nabla f_\varepsilon| < \frac{c_1}{\varepsilon}$ on D for some constant $c_1 > 0$. Now

$$\begin{aligned} \iint_{D \times D} A \left(\frac{|f_\varepsilon(x) - f_\varepsilon(y)|}{|x - y|^s} \right) \frac{dxdy}{|x - y|^N} &= \iint_{D_\varepsilon \times D_\varepsilon} A \left(\frac{|f_\varepsilon(x) - f_\varepsilon(y)|}{|x - y|^s} \right) \frac{dxdy}{|x - y|^N} \\ &+ 2 \iint_{D_\varepsilon^c \times D_\varepsilon} A \left(\frac{|f_\varepsilon(x) - f_\varepsilon(y)|}{|x - y|^s} \right) \frac{dxdy}{|x - y|^N} \leq 3 \iint_{D \times D_\varepsilon} A \left(\frac{|f_\varepsilon(x) - f_\varepsilon(y)|}{|x - y|^s} \right) \frac{dxdy}{|x - y|^N} \\ &\leq 3 \int_{D_\varepsilon} \left\{ \int_{y \in D, |x-y| < \varepsilon} A \left(\frac{|f_\varepsilon(x) - f_\varepsilon(y)|}{|x - y|^s} \right) \frac{dy}{|x - y|^N} \right. \\ &\quad \left. + \int_{y \in D, |x-y| > \varepsilon} A \left(\frac{|f_\varepsilon(x) - f_\varepsilon(y)|}{|x - y|^s} \right) \frac{dy}{|x - y|^N} \right\} dx = 3 \int_{D_\varepsilon} (I_1(x) + I_2(x)) dx. \end{aligned}$$

Using $|\nabla f_\varepsilon| < \frac{c_1}{\varepsilon}$ on D , we get

$$\begin{aligned} I_1(x) &= \int_{y \in D, |x-y| < \varepsilon} A\left(\frac{|f_\varepsilon(x) - f_\varepsilon(y)|}{|x-y|^s}\right) \frac{dy}{|x-y|^N} \leq \int_{|x-y| < \varepsilon} A\left(\frac{c_1|x-y|^{1-s}}{\varepsilon}\right) \frac{dy}{|x-y|^N} \\ &= \int_{|y| < \varepsilon} A\left(\frac{c_1|y|^{1-s}}{\varepsilon}\right) \frac{dy}{|y|^N} \leq c \int_0^\varepsilon A\left(\frac{c_1 r^{1-s}}{\varepsilon}\right) \frac{dr}{r} = c \int_0^{c_1 \varepsilon^{-s}} \frac{A(z)}{z} dz. \end{aligned}$$

The fact that $0 \leq f_\varepsilon(x) \leq 1$ on D gives

$$\begin{aligned} I_2(x) &= \int_{y \in D, |x-y| > \varepsilon} A\left(\frac{|f_\varepsilon(x) - f_\varepsilon(y)|}{|x-y|^s}\right) \frac{dy}{|x-y|^N} \leq \int_{|x-y| > \varepsilon} A(2|x-y|^{-s}) \frac{dy}{|x-y|^N} \\ &= c \int_\varepsilon^\infty \frac{A(2r^{-s})}{r} dr \leq c \int_0^{2\varepsilon^{-s}} \frac{A(z)}{z} dz. \end{aligned}$$

The hypothesis on D that it has bounded Lipschitz boundary implies that $\mathcal{L}^N(D_\varepsilon)$ is bounded above as well as bounded below by a constant multiple of ε . Let $\frac{1}{c_2} := \max\{c_1, 2\}$. Then we have, as $\varepsilon \rightarrow 0$,

$$(3.11) \quad \iint_{D \times D} A\left(\frac{|f_\varepsilon(x) - f_\varepsilon(y)|}{|x-y|^s}\right) \frac{dx dy}{|x-y|^N} \leq c \mathcal{L}^N(D_\varepsilon) \int_0^{(c_2 \varepsilon)^{-s}} \frac{A(z)}{z} dz \leq c (c_2 \varepsilon) \int_0^{(c_2 \varepsilon)^{-s}} \frac{A(z)}{z} dz \rightarrow \beta.$$

Now $f_\varepsilon \in C_c^\infty(D)$, $f_\varepsilon \rightarrow 1$ pointwise a.e. in D and A is continuous. Therefore by Fatou's lemma,

$$(3.12) \quad \lim_{\varepsilon \rightarrow 0} \int_D A\left(\frac{|f_\varepsilon(x)|}{\delta_x^s}\right) dx \geq \int_D A\left(\frac{1}{\delta_x^s}\right) dx > 0$$

and

$$(3.13) \quad \lim_{\varepsilon \rightarrow 0} \int_D A(|f_\varepsilon(x)|) dx \geq A(1) \mathcal{L}^N(D) > 0.$$

Now to prove (1) assume $\beta = 0$. Then LHS of eq. (3.11), which is also the numerator in the definition of both $H_{N,s,A}$ and $P_{N,s,A}^1$, converges to 0. We now use eqs. (3.12) and (3.13) to conclude $H_{N,s,A}(D) = 0$ and $P_{N,s,A}^1(D) = 0$ respectively. This proves (1).

Now we prove (2). Applying L'hospital rule to eq. (1.5), we get $\lim_{\varepsilon \rightarrow 0} \varepsilon A(\varepsilon^{-s}) = \beta$. So there exists $n_0 \in \mathbb{N}$ such that for $n \geq n_0$, $A(n^s) \geq \frac{\beta n}{2}$. Using eq. (3.12) we have

$$\begin{aligned} \lim_{\frac{1}{n} \rightarrow 0} \int_D A\left(\frac{|f_{\frac{1}{n}}(x)|}{\delta_x^s}\right) dx &\geq \int_D A\left(\frac{1}{\delta_x^s}\right) dx \geq \sum_{n=n_0}^\infty \int_{D_{\frac{1}{n}} \setminus D_{\frac{1}{n+1}}} A\left(\frac{1}{\delta_x^s}\right) dx \\ &\geq \sum_{n=n_0}^\infty \int_{D_{\frac{1}{n}} \setminus D_{\frac{1}{n+1}}} A(n^s) dx = \sum_{n=n_0}^\infty \frac{\beta n}{2} \mathcal{L}^N\left(D_{\frac{1}{n}} \setminus D_{\frac{1}{n+1}}\right) = \sum_{n=n_0}^\infty \frac{\beta n}{2} \left(\frac{1}{n} - \frac{1}{n+1}\right) = \infty. \end{aligned}$$

The proof follows after observing that $\beta > 0$ in eq. (3.11). \square

The conclusions we can draw as an application of theorems 1.2 and 1.3, for any bounded Lipschitz domain D and for any $q > 1$ are shown below in table 1.

TABLE 1. Conclusions we can draw from [theorems 1.2](#) and [1.3](#)

$A(t)$	$H_{N,s,A}(D) > 0$	$H_{N,s,A}(D) = 0$	$P_{N,s,A}^1(D) > 0$	$P_{N,s,A}^1(D) = 0$
t^q	$s \in (\frac{1}{q}, 1)$	$s \in (0, \frac{1}{q}]$	$s \in (\frac{1}{q}, 1)$	$s \in (0, \frac{1}{q})$
$t^q(1 + \log t)$	$s \in (\frac{1}{q}, 1)$	$s \in (0, \frac{1}{q})$	$s \in (\frac{1}{q}, 1)$	$s \in (0, \frac{1}{q})$
$\frac{t^q}{\log(e+t)}$	$s \in (\frac{1}{q}, 1)$	$s \in (0, \frac{1}{q}]$	$s \in (\frac{1}{q}, 1)$	$s \in (0, \frac{1}{q}]$
$(1+t)\log(1+t) - t$	NA	$s \in (0, 1)$	NA	$s \in (0, 1)$.

4. PROOF OF [THEOREMS 1.4](#) AND [1.5](#)

In this section we prove [theorems 1.4](#) and [1.5](#). The notations and conventions followed in this section will be the same as that in [section 3](#). We need [lemmas 4.1](#) and [4.2](#) to prove [theorem 1.4](#) and to prove [theorem 1.5](#) we further need [lemma 4.3](#).

Lemma 4.1. *Suppose $\liminf_{\lambda \rightarrow \infty} \lambda^{\frac{1-N}{s}} \alpha_{s,A}(\lambda) = 0$. Then there exists a constant $c = c(s, A, N) > 0$ such that for any $r > 0$ and any $f \in C_c^\infty(\mathbb{R}^N)$,*

$$\int_{B(0,r)^c} A\left(\frac{|f(x)|}{|x|^s}\right) dx \leq c \iint_{B(0,r)^c \times B(0,r)^c} A\left(\frac{|f(x) - f(y)|}{|x - y|^s}\right) \frac{dy dx}{|x - y|^N}.$$

Proof. We can assume $f \in C_c^\infty(\mathbb{R}^N \setminus \{0\})$, because for the sake of proving the above inequality the value of f inside $B(0, r)$ does not matter. Let $D := \mathbb{R}^N \setminus \{0\}$ so that $\delta_x = |x|$, $\Omega := B(0, r)^c$, $l_1 = 2^{m+1}$, $l_2 = \mathcal{L}^N(B(0, 1))2^{mN}(2^N - 1)$, $G := G(f, \Omega; l_1, l_2)$ and $A_k = B(0, 2^{k+1}r) \setminus B(0, 2^k r)$ for $k \in \mathbb{N} \cup \{0\}$. For a fixed $j \in \mathbb{N} \cup \{0\}$, set $E_1 = A_j$ and $E_2 = A_{j+m}$. Then for $x \in E_1$, $2^j r \leq \delta_x \leq 2^{j+1}r$, which gives $\text{dist}(x, A_{j+m}) \leq 2^{j+m+1}r \leq l_1 \delta_x$. Again $\mathcal{L}^N(A_k) = \mathcal{L}^N(B(0, 1))2^{kN}r^N(2^N - 1)$. So E_1, E_2 satisfy the hypotheses of [lemma 3.3](#). We can then conclude

$$\begin{aligned} \int_{A_j \cap G} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx &\leq \frac{2^{p+1}\mathcal{L}^N(A_j)}{\mathcal{L}^N(A_{j+m})} \int_{A_{j+m}} A\left(\frac{\sup\{\delta_x^s \mid x \in A_{j+m}\} |f(y)|}{\inf\{\delta_x^s \mid x \in A_j\} \delta_y^s}\right) dy \\ &= 2^{p+1-mN} \int_{A_{j+m}} A\left(\frac{(2^{j+m+1}r)^s |f(y)|}{(2^j r)^s \delta_y^s}\right) dy = 2^{p+1-mN} \int_{A_{j+m}} A\left(\frac{2^{(m+1)s} |f(y)|}{\delta_y^s}\right) dy \\ &= 2^{p+1+N} \int_{A_{j+m}} \frac{A\left(\frac{2^{(m+1)s} |f(y)|}{\delta_y^s}\right)}{2^{(m+1)N} A\left(\frac{|f(y)|}{\delta_y^s}\right)} A\left(\frac{|f(y)|}{\delta_y^s}\right) dy \\ &= 2^{p+1+N} 2^{(m+1)(1-N)} \alpha_{s,A}(2^{(m+1)s}) \int_{A_{j+m}} A\left(\frac{|f(y)|}{\delta_y^s}\right) dy. \end{aligned}$$

Now we use the hypothesis and choose large enough $m = m(s, A, N) \in \mathbb{N}$, so that

$$\int_{A_j \cap G} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx \leq \frac{1}{2} \int_{A_{j+m}} A\left(\frac{|f(y)|}{\delta_y^s}\right) dy.$$

Now we can apply [lemma 3.4](#) to complete the proof. \square

Lemma 4.2. Suppose $\liminf_{\lambda \rightarrow 0+} \lambda^{\frac{1-N}{s}} \alpha_{s,A}(\lambda) = 0$. Then there exists a constant $c = c(s, A, N) > 0$ and $m = m(s, A, N) \in \mathbb{N}$ such that for any $r > 0$ and any $f \in C_c^\infty(\mathbb{R}^N)$,

$$\begin{aligned} & \int_{B(0, 2^m r)^c} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx \\ & \leq c \left[\iint_{B(0, 2^m r)^c \times B(0, 2^m r)^c} A\left(\frac{|f(x) - f(y)|}{|x - y|^s}\right) \frac{dy dx}{|x - y|^N} + \int_{B(0, 2^m r) \setminus B(0, r)} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx \right]. \end{aligned}$$

Proof. From the hypothesis on s and A , there exists $m \in \mathbb{N}$ such that

$$2^{p+1+N} 2^{(1-m)(1-N)} \alpha_{s,A} \left(2^{(1-m)s} \right) < \frac{1}{2}.$$

We fix this m . Let $D = C_c^\infty(\mathbb{R}^N \setminus \{0\})$, $\Omega = B(0, r)^c$, $l_1 = 2$, $l_2 = \mathcal{L}^N(B(0, 1)) 2^{-mN} (2^N - 1)$, $G = G(f, \Omega; l_1, l_2)$ and $A_k = B(0, 2^{k+1}r) \setminus B(0, 2^k r)$, then $\mathcal{L}^N(A_k) = \mathcal{L}^N(B(0, 1)) 2^{kN} r^N (2^N - 1)$. As in the previous lemma, we assume $f \in C_c^\infty(D)$. For $j \geq m$, set $E_1 = A_j$ and $E_2 = A_{j-m}$. So for $x \in A_j$, $\text{dist}(x, A_{j-m}) \leq 2^{j+1}r \leq l_1 \delta_x$. Also $\frac{\mathcal{L}^N(A_{j-m})}{\delta_x^N} \geq \mathcal{L}^N(B(0, 1)) 2^{-mN} (2^N - 1) = l_2$. Hence, for any $j \geq m$, we can apply lemma 3.3 to get

$$\begin{aligned} & \int_{A_j \cap G} A\left(\frac{|f(x)|}{|x|^s}\right) dx \leq 2^{p+1+mN} \int_{A_{j-m}} A\left(\frac{2^{(1-m)s} |f(y)|}{|y|^s}\right) dy \\ & = 2^{p+1+N} \int_{A_{j-m}} \frac{A\left(\frac{2^{(1-m)s} |f(y)|}{|y|^s}\right) dy}{2^{(1-m)N} A\left(\frac{|f(y)|}{|y|^s}\right) dy} A\left(\frac{|f(y)|}{|y|^s}\right) dy \\ & = 2^{p+1+N} 2^{(1-m)(1-N)} \alpha_{s,A} \left(2^{(1-m)s} \right) \int_{A_{j-m}} A\left(\frac{|f(y)|}{|y|^s}\right) dy < \frac{1}{2} \int_{A_{j-m}} A\left(\frac{|f(y)|}{|y|^s}\right) dy. \end{aligned}$$

Finally we apply lemma 3.4 to complete the proof. \square

Proof of Theorem 1.4. Set $D = \mathbb{R}^N \setminus \{0\}$, take $f \in C_c^\infty(D)$. Then $f \equiv 0$ near 0. First let us assume $\liminf_{\lambda \rightarrow \infty} \lambda^{\frac{1-N}{s}} \alpha_{s,A}(\lambda) = 0$. Choose $r > 0$ small enough, such that, $\text{supp}(f) \subseteq B(0, r)^c$. Then lemma 4.1 implies $H_{N,s,A}(D) > 0$.

If, however, $\liminf_{\lambda \rightarrow 0+} \lambda^{\frac{1-N}{s}} \alpha_{s,A}(\lambda) = 0$, we choose $r > 0$ so small that $\text{supp}(f) \subseteq B(0, 2^m r)^c$. Proceed exactly same as in the previous case and apply lemma 4.2. The restriction on $\text{supp}(f)$ ensures that the last term in the inequality is zero. Hence $H_{N,s,A}(\mathbb{R}^N \setminus \{0\}) > 0$. \square

We shall the notations as in the geometric decomposition of D , done after lemma 3.3, for lemma 4.3.

Lemma 4.3. For a fixed $z \in \partial D$, define $Q_k := \cup_{\ell=k}^\infty A_\ell$, where A_ℓ are as in eq. (3.3). Let $\liminf_{\lambda \rightarrow \infty} \alpha_{s,A}(\lambda) = 0$. Then there exists $m = m(\text{Lip}(D), s, A)$ and $c = c(\text{Lip}(D), s, A)$ such that for any $f \in C_c^\infty(D)$,

$$\int_{Q_m} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx \leq c \left[\iint_{Q_m \times Q_m} A\left(\frac{|f(x) - f(y)|}{|x - y|^s}\right) \frac{dy dx}{|x - y|^N} + \int_{Q_z(\rho) \setminus Q_m} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx \right].$$

Proof. From the hypothesis on s and A , there exists $m \in \mathbb{N}$ such that

$$2^{2+p}(1 + \text{Lip}(D))\alpha_{s,A}(2^{s+ms}(1 + \text{Lip}(D))^s) < \frac{1}{2}.$$

We fix this m . Let $l_1 = 2(1 + \text{Lip}(D))\sqrt{N-1 + (\text{Lip}(D)\sqrt{N-1} + 2)^2}$, $l_2 = 2^{m-1}$ and $G = G(f, Q_z(\rho); l_1, l_2)$. For $j \geq m$, set $E_1 = A_j \cap Q_j^i$ and $E_2 = A_{j-m} \cap Q_j^i$. Equations (3.4) and (3.5) imply that we can apply lemma 3.3 on E_1, E_2 . Thus we get, utilizing the choice of m ,

$$\begin{aligned} \int_{(A_j \cap Q_j^i) \cap G} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx &\leq 2^{1+p-m} \int_{A_{j-m} \cap Q_j^i} A\left(2^{s+ms}(1 + \text{Lip}(D))^s \frac{|f(y)|}{\delta_y^s}\right) dy \\ &= 2^{2+p}(1 + \text{Lip}(D)) \int_{A_{j-m} \cap Q_j^i} \frac{A\left(2^{s+ms}(1 + \text{Lip}(D))^s \frac{|f(y)|}{\delta_y^s}\right)}{2^{1+m}(1 + \text{Lip}(D))A\left(\frac{|f(y)|}{\delta_y^s}\right)} A\left(\frac{|f(y)|}{\delta_y^s}\right) dy \\ &\leq 2^{2+p}(1 + \text{Lip}(D))\alpha_{s,A}(2^{s+ms}(1 + \text{Lip}(D))^s) \int_{A_{j-m} \cap Q_j^i} A\left(\frac{|f(y)|}{\delta_y^s}\right) dy < \frac{1}{2} \int_{A_{j-m} \cap Q_j^i} A\left(\frac{|f(y)|}{\delta_y^s}\right) dy. \end{aligned}$$

Summing over $1 \leq i \leq 2^{j(N-1)}$, we get

$$\int_{A_j \cap G} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx \leq \frac{1}{2} \int_{A_{j-m}} A\left(\frac{|f(y)|}{\delta_y^s}\right) dy.$$

Applying lemma 3.4 the lemma follows. \square

Proof of Theorem 1.5. (1) Without loss of generality, we can assume $D = \mathbb{R}^{N-1} \times \{0\}$. Let $z \in D$. First assume the case $\liminf_{\lambda \rightarrow 0} \alpha_{s,A}(\lambda) = 0$. Take $f \in C_c^\infty(D)$. Note that for the choice of D in this case, we are free to choose ρ as large as we want and we still have $Q_z(\rho) \subset D$. Choose $\rho > 0$ so that $\text{supp}(f) \subseteq Q_z(\rho)$. Then applying lemma 3.5 we get

$$\begin{aligned} \int_D A\left(\frac{|f(x)|}{\delta_x^s}\right) dx &= \int_{Q_z(\rho)} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx \leq c \iint_{Q_z(\rho) \times Q_z(\rho)} A\left(\frac{|f(x) - f(y)|}{|x - y|^s}\right) \frac{dxdy}{|x - y|^N} \\ &\leq c \iint_{D \times D} A\left(\frac{|f(x) - f(y)|}{|x - y|^s}\right) \frac{dxdy}{|x - y|^N}. \end{aligned}$$

Now assume $\liminf_{\lambda \rightarrow \infty} \alpha_{s,A}(\lambda) = 0$. Lemma 4.3 implies

$$\int_{0 < x_N < \frac{\rho}{2^m}} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx \leq c \iint_{0 < x_N, y_N < \frac{\rho}{2^m}} A\left(\frac{|f(x) - f(y)|}{|x - y|^s}\right) \frac{dxdy}{|x - y|^N} + \int_{x_N > \frac{\rho}{2^m}} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx,$$

where m, ρ is the integer considered in lemma 4.3. Note that m does not depend on ρ . So we choose ρ big enough, so that $\text{supp}(f) \subseteq \{x \in \mathbb{R}^N \mid x_N < \frac{\rho}{2^m}\}$ forcing the remainder term in the above equation to vanish.

(2) Let $f \in C_c^\infty(D)$. We choose $R > 0$ large enough so that $D^c \subseteq B(0, R)$. Now $D \cap B(0, R)$ is a bounded Lipschitz domain. A careful observation of the proof of theorem 1.2 reveals that, under the assumption $\liminf_{\lambda \rightarrow 0+} \alpha_{s,A}(\lambda) = 0$, we can use the same technique to get a constant $c = c(A, s, N, D)$ such that

$$(4.1) \quad \int_{D \cap B(0, R)} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx \leq c \iint_{D \cap B(0, R) \times D \cap B(0, R)} A\left(\frac{|f(x) - f(y)|}{|x - y|^s}\right) \frac{dxdy}{|x - y|^N},$$

even if $f \notin C_c^\infty(D \cap B(0, R))$.

First we assume $\liminf_{\lambda \rightarrow \infty} \lambda^{\frac{1-N}{s}} \alpha_{s,A}(\lambda) = \liminf_{\lambda \rightarrow 0+} \alpha_{s,A}(\lambda) = 0$.

We apply [lemma 4.1](#) to get, for any $r = R$,

$$(4.2) \quad \int_{B(0,R)^c} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx \leq c \iint_{B(0,R)^c \times B(0,R)^c} A\left(\frac{|f(x) - f(y)|}{|x - y|^s}\right) \frac{dy dx}{|x - y|^N}.$$

Adding [eqs. \(4.1\)](#) and [\(4.2\)](#), the claim follows.

Now we assume $\liminf_{\lambda \rightarrow 0+} \lambda^{\frac{1-N}{s}} \alpha_{s,A}(\lambda) = 0$. Note that in this case $\liminf_{\lambda \rightarrow 0+} \alpha_{s,A}(\lambda) = 0$ is implied. Consider the extension of f by 0 to whole of \mathbb{R}^N . Applying [lemma 4.2](#) with $r = \frac{R}{2^m}$ and then proceeding similarly as above, the claim follows.

Finally, assume $\liminf_{\lambda \rightarrow \infty} \alpha_{s,A}(\lambda) = 0$. Denote $\forall \eta > 0$, $D_\eta := \{x \in D \mid \text{dist}(x, \partial D) < \eta\}$. An application of [lemma 4.3](#), as [lemma 3.5](#) was used in the proof of [theorem 1.2](#), gives

$$\begin{aligned} \int_{D_\varepsilon} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx &\leq c \left[\iint_{D_\varepsilon \times D_\varepsilon} A\left(\frac{|f(x) - f(y)|}{|x - y|^s}\right) \frac{dy dx}{|x - y|^N} + \int_{D_{2\varepsilon} \setminus D_\varepsilon} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx \right] \\ &\leq c \left[\iint_{D_\varepsilon \times D_\varepsilon} A\left(\frac{|f(x) - f(y)|}{|x - y|^s}\right) \frac{dy dx}{|x - y|^N} + \int_{D \setminus D_\varepsilon} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx \right], \end{aligned}$$

for some fixed $\varepsilon > 0$ and a constant $c = c(s, N, A, D) > 0$ and or any $f \in C_c^\infty(D)$. Now set $\Omega := D \setminus D_\varepsilon$ and let $R > 0$ be such that $\Omega^c \subseteq B(0, R)$. Before proceeding further note that if, for some constant $c = c(s, N, A, D) > 0$ and for all $f \in C_c^\infty(D)$, we can show

$$\int_{\Omega} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx \leq c \iint_{\Omega \times \Omega} A\left(\frac{|f(x) - f(y)|}{|x - y|^s}\right) \frac{dy dx}{|x - y|^N},$$

then the proof will be complete. Set $A_j := B(0, 2^j R) \setminus B(0, 2^{j-1} R)$ and $A_0 := B(0, R) \cap \Omega$. Then for $k \in \mathbb{N}$, $\mathcal{L}^N(A_k) = \mathcal{L}^N(B(0, 1)) R^N 2^{(k-1)N} (2^N - 1)$ and for any $x \in A_k$, $k > 1$, $(2^{k-1} - 1)R \leq \delta_x \leq 2^k R$. Whereas $\mathcal{L}^N(A_0) = \mathcal{L}^N(B(0, R)) - \mathcal{L}^N(\Omega^c)$, for $x \in A_0 \cup A_1$, $\varepsilon \leq \delta_x \leq R$ and for $x \in A_1$, $\varepsilon \leq \delta_x \leq 2R$. Now let $x \in A_j$, $y \in A_{j+m}$. Then for any $j \geq 0$, $\text{dist}(x, y) \leq 2^{j+m} R = 2^m \frac{2^j}{2^{j-1}-1}$. So it is clear that there exists $l_1, l_2 > 0$ such that the hypotheses of [lemma 3.3](#) are satisfied. So, we get

$$\begin{aligned} \int_{A_j \cap G} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx &\leq \frac{2^{p+1} \mathcal{L}^N(A_j)}{\mathcal{L}^N(A_{j+m})} \int_{A_{j+m}} A\left(\frac{\sup\{\delta_x^s \mid x \in A_{j+m}\} |f(y)|}{\inf\{\delta_x^s \mid x \in A_j\} \delta_y^s}\right) dy \\ &\leq c(R, \mathcal{L}^N(\Omega^c), N) 2^{-mN} \int_{A_{j+m}} A\left(\frac{\sup\{\delta_x^s \mid x \in A_{j+m}\} |f(y)|}{\inf\{\delta_x^s \mid x \in A_j\} \delta_y^s}\right) dy \\ &\leq c 2^{-mN} \int_{A_{j+m}} A\left(\frac{c(\varepsilon, R) 2^{ms} |f(y)|}{\delta_y^s}\right) dy. \end{aligned}$$

Note that $\liminf_{\lambda \rightarrow \infty} \alpha_{s,A}(\lambda) = 0$ implies $\liminf_{\lambda \rightarrow \infty} \lambda^{\frac{1-N}{s}} \alpha_{s,A}(\lambda) = 0$. Choosing m sufficiently large and then applying this, we get

$$\int_{A_j \cap G} A\left(\frac{|f(x)|}{\delta_x^s}\right) dx \leq \int_{A_{j+m}} A\left(\frac{|f(y)|}{\delta_y^s}\right) dy.$$

The proof follows as an application of [lemma 3.4](#). \square

5. $RFOPI(s, A)$ AND $FOPI(s, A)$ ON UNBOUNDED DOMAINS

In this section we study $RFOPI(s, A)$ and $FOPI(s, A)$ for general domains in \mathbb{R}^N .

Proposition 5.1. *Let $D \subseteq \mathbb{R}^N$ be an open set and $\lim_{\lambda \rightarrow 0} \alpha_{s,A}(\lambda) = 0$. Then $P_{N,s,A}^1(D) > 0$ if and only if $BC(D) < \infty$.*

Proof. Suppose $BC(D) < \infty$. There exists a countable number of disjoint open intervals, say I_k , such that $D = \cup_{k=1}^{\infty} I_k$. Let $f \in C_c^\infty(D) \setminus \{0\}$. Using [theorem 1.2](#) followed by [lemma 2.5](#), we get

$$\begin{aligned} \iint_{D \times D} A\left(\frac{|f(x) - f(y)|}{|x - y|^s}\right) \frac{dxdy}{|x - y|} &\geq \sum_{k=1}^{\infty} \iint_{I_k \times I_k} A\left(\frac{|f(x) - f(y)|}{|x - y|^s}\right) \frac{dxdy}{|x - y|} \\ &\geq \sum_{k=1}^{\infty} P_{1,s,A}^1(I_k) \int_{I_k} A(|f(x)|) dx \geq \min\{BC(D)^{-sp}, 1\} P_{1,s,A}^1((0, 1)) \sum_{k=1}^{\infty} \int_{I_k} A(|f(x)|) dx \\ &= \min\{BC(D)^{-sp}, 1\} P_{1,s,A}^1((0, 1)) \int_D A(|f(x)|) dx, \end{aligned}$$

which implies $P_{1,s,A}^1(D) \geq c P_{1,s,A}^1((0, 1)) > 0$. The other part follows from [proposition 2.1](#). \square

The next lemma is an important change of variable formula for the fractional Orlicz seminorm.

Lemma 5.1. *Let $0 < s < 1$ and $D \subseteq \mathbb{R}^N$ be a measurable set. Then for any $f \in C_c^\infty(D)$,*

$$\begin{aligned} 2 \iint_{D \times D} A\left(\frac{|f(x) - f(y)|}{|x - y|^s}\right) \frac{dxdy}{|x - y|^N} \\ = \int_{\omega \in \mathbb{S}^{N-1}} \int_{x \in \omega^\perp} \int_{\ell \in L_D(x, \omega)} \int_{t \in L_D(x, \omega)} A\left(\frac{|f(x + \ell\omega) - f(x + t\omega)|}{|\ell - t|^s}\right) \frac{dtd\ell d\sigma(x) d\sigma(\omega)}{|\ell - t|}, \end{aligned}$$

where $L_D(x, \omega) := \{t \mid x + t\omega \in D\}$.

Proof. For a fixed $f \in C_c^\infty(D)$ and $x \in D$, consider the change of variable $y = x + z$ followed by another change of variable to polar coordinates. We then get

$$\begin{aligned} 2 \iint_{D \times D} A\left(\frac{|f(x) - f(y)|}{|x - y|^s}\right) \frac{dxdy}{|x - y|^N} &= 2 \int_{x \in D} \int_{z \in D - x} A\left(\frac{|f(x) - f(x + z)|}{|z|^s}\right) \frac{dzdx}{|z|^N} \\ &= 2 \int_{x \in D} \int_{\omega \in \mathbb{S}^{N-1}} \int_{r \in \{r > 0 \mid x + r\omega \in D\}} A\left(\frac{|f(x) - f(x + r\omega)|}{r^s}\right) \frac{dr d\sigma(\omega) dx}{r} \\ &= \int_{\omega \in \mathbb{S}^{N-1}} \int_{x \in D} \int_{h \in L_D(x, \omega)} A\left(\frac{|f(x) - f(x + h\omega)|}{|h|^s}\right) \frac{dh dx d\sigma(\omega)}{|h|}. \end{aligned}$$

Now, for a fixed $\omega \in \mathbb{S}^{N-1}$ we can write $x = (x - (x \cdot \omega)\omega) + (x \cdot \omega)\omega$. Using this we can split the integral over $x \in D$ by considering the change of variable $\ell = x \cdot \omega$ and $z = x - (x \cdot \omega)\omega$, where $\ell \in L_D(z, \omega)$ and $z \in \omega^\perp$. So we get

$$\begin{aligned}
& 2 \iint_{D \times D} A \left(\frac{|f(x) - f(y)|}{|x - y|^s} \right) \frac{dx dy}{|x - y|^N} \\
&= \int_{\omega \in \mathbb{S}^{N-1}} \int_{z \in \omega^\perp} \int_{\ell \in L_D(z, \omega)} \int_{h \in L_D(x, \omega)} A \left(\frac{|f(z + \ell\omega) - f(z + (\ell + h)\omega)|}{|h|^s} \right) \frac{dh d\ell d\sigma(z) d\sigma(\omega)}{|h|} \\
&= \int_{\omega \in \mathbb{S}^{N-1}} \int_{z \in \omega^\perp} \int_{\ell \in L_D(z, \omega)} \int_{t \in L_D(z + \ell\omega, \omega) + \ell} A \left(\frac{|f(z + \ell\omega) - f(z + t\omega)|}{|\ell - t|^s} \right) \frac{dh d\ell d\sigma(z) d\sigma(\omega)}{|\ell - t|}.
\end{aligned}$$

We conclude the proof of this lemma with the observation that $L_D(z, \omega) = L_D(z + \ell\omega, \omega) + \ell$. This finishes the proof of the theorem. \square

Proof of Theorem 1.6. Let $f \in C_c^\infty(D)$. (1) Applying lemma 5.1 we obtain

$$\begin{aligned}
& 2 \iint_{D \times D} A \left(\frac{|f(x) - f(y)|}{|x - y|^s} \right) \frac{dx dy}{|x - y|^N} \\
&= \int_{\omega \in \mathbb{S}^{N-1}} \int_{x \in \omega^\perp} \int_{\ell \in L_D(x, \omega)} \int_{t \in L_D(x, \omega)} A \left(\frac{|f(x + \ell\omega) - f(x + t\omega)|}{|\ell - t|^s} \right) \frac{dt d\ell d\sigma(x) d\sigma(\omega)}{|\ell - t|} \\
&\geq \int_{\omega \in \Sigma} \int_{x \in \omega^\perp} \int_{\ell \in L_D(x, \omega)} \int_{t \in L_D(x, \omega)} A \left(\frac{|f(x + \ell\omega) - f(x + t\omega)|}{|\ell - t|^s} \right) \frac{dt d\ell d\sigma(x) d\sigma(\omega)}{|\ell - t|} \\
&= \sum_{k=1}^{\infty} \int_{\omega \in \Sigma} \int_{x \in \omega^\perp} \int_{\ell \in L_D(x, \omega)} \int_{t \in I_k} A \left(\frac{|f(x + \ell\omega) - f(x + t\omega)|}{|\ell - t|^s} \right) \frac{dt d\ell d\sigma(x) d\sigma(\omega)}{|\ell - t|} \\
&\geq \sum_{k=1}^{\infty} \int_{\omega \in \Sigma} \int_{x \in \omega^\perp} P_{1,s,A}^1(I_k) \int_{t \in I_k} A(|f(x + t\omega)|) dt d\sigma(x) d\sigma(\omega),
\end{aligned}$$

where $L_D(x, \omega) = \cup_{k=1}^{\infty} I_k$, for mutually disjoint family of intervals $\{I_k\}_k$. From lemma 2.5 we have, for any k ,

$$P_{1,s,A}^1(I_k) \geq \min\{BC(L_D(x, \omega))^{-sp}, 1\} P_{1,s,A}^1((0, 1)).$$

We can now use the hypothesis of the lemma to get a constant $c = c(D) > 0$ such that $c < \min\{BC(L_D(x, \omega))^{-sp}, 1\}$. Again as an application of theorem 1.2 $P_{1,s,A}^1((0, 1)) > 0$. Using these we get

$$\begin{aligned}
\iint_{D \times D} A \left(\frac{|f(x) - f(y)|}{|x - y|^s} \right) \frac{dx dy}{|x - y|^N} &\geq c \int_{\omega \in \Sigma} \int_{x \in \omega^\perp} \int_{t \in L_D(x, \omega)} A(|f(x + t\omega)|) dt d\sigma(x) d\sigma(\omega) \\
&\geq c \int_{\omega \in \Sigma} \int_{x \in D} A(|f(x)|) dx d\sigma(\omega) = c \int_D A(|f(x)|) dx.
\end{aligned}$$

which implies that $P_{N,s,A}^1(D) > 0$. This completes the first part of the proof.

(2) We use the positivity, increasing property of A , the fact that $f \equiv 0$ on D^c and suitable change of variable in the following calculation.

$$\begin{aligned}
& \iint_{\mathbb{R}^N \times \mathbb{R}^N} A \left(\frac{|f(x) - f(y)|}{|x - y|^s} \right) \frac{dy dx}{|x - y|^N} = \frac{1}{R^N} \iint_{\mathbb{R}^N \times \mathbb{R}^N} A \left(\frac{R^s |f(\frac{x}{R}) - f(\frac{y}{R})|}{|x - y|^s} \right) \frac{dy dx}{|x - y|^N} \\
&\geq \frac{1}{R^N} \int_{x \in \mathbb{R}^N} \int_{y \in B(x, R) \cap D^c} A \left(\frac{R^s |f(\frac{x}{R})|}{|x - y|^s} \right) \frac{dy dx}{|x - y|^N} \geq \frac{\mathcal{L}^N(B(x, R) \cap D^c)}{R^N} \int_{\mathbb{R}^N} A \left(|f\left(\frac{x}{R}\right)| \right) dx
\end{aligned}$$

$$= \frac{c_1}{R^N} \int_{\mathbb{R}^N} A(|f(x)|) dx = \frac{c_1}{R^N} \int_D A(|f(x)|) dx.$$

This shows $P_{N,s,A}^2(D) > 0$ and the proof of the second part follows.

(3) This is a straightforward use of [lemma 5.1](#) as in the first part of the proof. \square

As an application of [theorem 1.6](#) we discuss here some examples of some unbounded domains for which Poincaré inequality holds true, provided $\lim_{\lambda \rightarrow 0^+} \alpha_{s,A}(\lambda) = 0$ for the first condition (see also [[CCRS21](#), [CR20](#)]). Although verification of the claims in these examples is straight forward, we work it out in the second example.

Example 5.1 (Domain between graphs of two functions). *Let $f_i : \mathbb{R}^{N-1} \rightarrow [m, M]$ ($i = 1, 2$) be two bounded continuous function such that $f_1 < f_2$. We then define D as*

$$D = \{(\tilde{x}, x_N) \in \mathbb{R}^N \mid f_1(\tilde{x}) < x_N < f_2(\tilde{x})\}.$$

It is easy to verify that all the hypotheses of [theorem 1.6](#) are true.

Example 5.2 (Countable union of parallel strips). *Let $D := I \times \mathbb{R} \subseteq \mathbb{R}^2$, where $I := \cup_{k=1}^{\infty} I_k$ and I_k 's are disjoint intervals with $\text{dist}(I_i, I_j) \geq \gamma$ for a constant $\gamma > 0$, where I_i, I_j are any two distinct intervals. Also assume that $\sup_k \mathcal{L}^1(I_k) = M < \infty$. Then it is easy to check that the second condition of [theorem 1.6](#) holds. This gives $P_{1,s,A}^2(I) > 0$, for any $s \in (0, 1)$. Now choose $\Sigma = \{\omega_\theta := (\cos\theta, \sin\theta) \mid \theta \in (0, \pi/4)\}$. Then for $x \in \omega_\theta^\perp$, the set $L_D(x, \omega_\theta)$ can be expressed as the countable union of disjoint intervals, i.e. $L_D(x, \omega_\theta) = \cup_{k=1}^{\infty} J_k(x, \omega_\theta)$. Observe that $\mathcal{L}^1(J_k(x, \omega_\theta)) = \frac{\mathcal{L}^1(I_k)}{\cos\theta}$ for any k , which is independent of x . Thus, we have for some constant $c = c(x, \omega_\theta)$,*

$$(5.1) \quad L_D(x, \omega_\theta) = c + (\sec\theta)I.$$

This gives $\sup_{\omega_\theta \in \Sigma, x \in \omega_\theta^\perp} BC(L_D(x, \omega_\theta)) \leq M\sqrt{2}$ and which shows that the first condition holds. Consequently $RFOPI(s, A)$ holds in D when $\lim_{\lambda \rightarrow 0^+} \alpha_{s,A}(\lambda) = 0$. For third condition, Using (3) of [proposition 2.1](#) and [eq. \(5.1\)](#) we obtain, since $\theta \in (0, \pi/4)$,

$$P_{1,s,A}^2(L_D(x, \omega_\theta)) \geq \frac{P_{1,s,A}^2(I)}{(\cos\theta)^{sp}} \geq 2^{\frac{sp}{2}} P_{1,s,A}^2(I).$$

So, uniform $FOPI(s, A)$ holds for the collection $\{L_D(x, \omega_\theta)\}_{\omega_\theta \in \Sigma, x \in \omega_\theta^\perp}$. Hence the domain D is of type $LS(s, A)$ for any $s \in (0, 1)$. Consequently $FOPI(s, A)$ holds in D for any $s \in (0, 1)$.

Example 5.3 (Concentric annulus). *The following domain satisfies condition 2 for all $s \in (0, 1)$ of [theorem 1.6](#):*

$$D = \bigcup_{k=1}^{\infty} B(0, 2k) \setminus B(0, 2k-1).$$

Example 5.4 (Domain with holes around points of $\mathbb{Z} \times \mathbb{Z}$). *The following domain satisfies the condition 2 of [theorem 1.6](#):*

$$D = \mathbb{R}^2 \setminus \left(\bigcup_{x \in \mathbb{Z}^2} B(x, 1/10) \right).$$

ACKNOWLEDGEMENT

Research work of first author is funded by Matrics grant (MTR/2020/000594). The work of the second author is supported by CSIR(09/092(0940)/2015-EMR-I). Research work of third author is funded by Matrics grant (MTR/2019/000585) and Inspire grant (IFA14-MA43) of Department of Science and Technology (DST).

REFERENCES

- [ABS18] Elhoussine Azroul, Abdelmoujib Benkirane, and Mohammed Sрати. Introduction to fractional orlicz-sobolev spaces. *arXiv preprint arXiv:1807.11753*, 2018.
- [ACPS20a] Angela Alberico, Andrea Cianchi, Luboš Pick, and Lenka Slavíková. On the limit as $s \rightarrow 1^-$ of possibly non-separable fractional Orlicz-Sobolev spaces. *Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl.*, 31(4):879–899, 2020.
- [ACPS20b] Angela Alberico, Andrea Cianchi, Luboš Pick, and Lenka Slavíková. On the limit as $s \rightarrow 0^+$ of fractional Orlicz-Sobolev spaces. *J. Fourier Anal. Appl.*, 26(6):Paper No. 80, 19, 2020.
- [ACPS21] Angela Alberico, Andrea Cianchi, Luboš Pick, and Lenka Slavíková. Fractional Orlicz-Sobolev embeddings. *J. Math. Pures Appl. (9)*, 149:216–253, 2021.
- [AF03] Robert A. Adams and John J. F. Fournier. *Sobolev spaces*, volume 140 of *Pure and Applied Mathematics (Amsterdam)*. Elsevier/Academic Press, Amsterdam, second edition, 2003.
- [BBX20] Anouar Bahrouni, Sabri Bahrouni, and Mingqi Xiang. On a class of nonvariational problems in fractional Orlicz-Sobolev spaces. *Nonlinear Anal.*, 190:111595, 13, 2020.
- [BC18] Lorenzo Brasco and Eleonora Cinti. On fractional Hardy inequalities in convex sets. *Discrete Contin. Dyn. Syst.*, 38(8):4019–4040, 2018.
- [BD11] Krzysztof Bogdan and Bartłomiej Dyda. The best constant in a fractional Hardy inequality. *Math. Nachr.*, 284(5-6):629–638, 2011.
- [Beb03] M. Bebendorf. A note on the Poincaré inequality for convex domains. *Z. Anal. Anwendungen*, 22(4):751–756, 2003.
- [BM97] Haïm Brezis and Moshe Marcus. Hardy’s inequalities revisited. *Annali della Scuola Normale Superiore di Pisa-Classe di Scienze*, 25(1-2):217–237, 1997.
- [BMS00] Haim Brezis, Moshe Marcus, and Itai Shafir. Extremal functions for Hardy’s inequality with weight. *Journal of Functional Analysis*, 171(1):177–191, 2000.
- [BO20] Sabri Bahrouni and Hichem Ounaies. Embedding theorems in the fractional Orlicz-Sobolev space and applications to non-local problems. *Discrete Contin. Dyn. Syst.*, 40(5):2917–2944, 2020.
- [BOT20] Sabri Bahrouni, Hichem Ounaies, and Leandro S. Tavares. Basic results of fractional Orlicz-Sobolev space and applications to non-local problems. *Topol. Methods Nonlinear Anal.*, 55(2):681–695, 2020.
- [BS21] Sabri Bahrouni and Ariel M. Salort. Neumann and Robin type boundary conditions in fractional Orlicz-Sobolev spaces. *ESAIM Control Optim. Calc. Var.*, 27(suppl.):Paper No. S15, 23, 2021.
- [CCRS21] Indranil Chowdhury, Gyula Csátó, Prosenjit Roy, and Firoj Sk. Study of fractional Poincaré inequalities on unbounded domains. *Discrete Contin. Dyn. Syst.*, 41(6):2993–3020, 2021.
- [CdP20] Ernesto Correa and Arturo de Pablo. Remarks on a nonlinear nonlocal operator in Orlicz spaces. *Adv. Nonlinear Anal.*, 9(1):305–326, 2020.
- [Che18] Huyuan Chen. The Dirichlet elliptic problem involving regional fractional Laplacian. *J. Math. Phys.*, 59(7):071504, 19, 2018.
- [CR17] Indranil Chowdhury and Prosenjit Roy. On the asymptotic analysis of problems involving fractional Laplacian in cylindrical domains tending to infinity. *Commun. Contemp. Math.*, 19(5):1650035, 21, 2017.
- [CR20] Indranil Chowdhury and Prosenjit Roy. Fractional Poincaré inequality for unbounded domains with finite ball condition: Counter example. *arXiv preprint arXiv:2001.04441*, 2020.
- [CS03] Zhen-Qing Chen and Renming Song. Hardy inequality for censored stable processes. *Tohoku Math. J. (2)*, 55(3):439–450, 2003.
- [DF12] Bartłomiej Dyda and Rupert L. Frank. Fractional Hardy-Sobolev-Maz’ya inequality for domains. *Studia Math.*, 208(2):151–166, 2012.
- [DNPV12] Eleonora Di Nezza, Giampiero Palatucci, and Enrico Valdinoci. Hitchhiker’s guide to the fractional Sobolev spaces. *Bull. Sci. Math.*, 136(5):521–573, 2012.
- [DV14] Bartłomiej Dyda and Antti V. Vähäkangas. A framework for fractional Hardy inequalities. *Ann. Acad. Sci. Fenn. Math.*, 39(2):675–689, 2014.
- [Dyd04] Bartłomiej Dyda. A fractional order Hardy inequality. *Illinois J. Math.*, 48(2):575–588, 2004.
- [Dyd11] Bartłomiej Dyda. Fractional Hardy inequality with a remainder term. *Colloq. Math.*, 122(1):59–67, 2011.
- [EHSV14] David E. Edmunds, Ritva Hurri-Syrjänen, and Antti V. Vähäkangas. Fractional Hardy-type inequalities in domains with uniformly fat complement. *Proc. Amer. Math. Soc.*, 142(3):897–907, 2014.
- [Eva10] Lawrence C. Evans. *Partial differential equations*, volume 19 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, second edition, 2010.

- [FBS19] Julián Fernández Bonder and Ariel M. Salort. Fractional order Orlicz-Sobolev spaces. *J. Funct. Anal.*, 277(2):333–367, 2019.
- [FHR20] Rita Ferreira, Peter Hästö, and Ana Margarida Ribeiro. Characterization of generalized Orlicz spaces. *Commun. Contemp. Math.*, 22(2):1850079, 25, 2020.
- [FL12] Rupert L. Frank and Michael Loss. Hardy-Sobolev-Maz’ya inequalities for arbitrary domains. *J. Math. Pures Appl. (9)*, 97(1):39–54, 2012.
- [FS08] Rupert L. Frank and Robert Seiringer. Non-linear ground state representations and sharp Hardy inequalities. *J. Funct. Anal.*, 255(12):3407–3430, 2008.
- [Gos74] Jean-Pierre Gossez. Nonlinear elliptic boundary value problems for equations with rapidly (or slowly) increasing coefficients. *Trans. Amer. Math. Soc.*, 190:163–205, 1974.
- [Hei95] Hans P. Heinig. A note on Hardy’s inequality in Orlicz spaces. In *Fourier analysis and partial differential equations (Miraflores de la Sierra, 1992)*, Stud. Adv. Math., pages 185–194. CRC, Boca Raton, FL, 1995.
- [HKP97] Hans P. Heinig, Alois Kufner, and Lars-Erik Persson. On some fractional order Hardy inequalities. *J. Inequal. Appl.*, 1(1):25–46, 1997.
- [HSV15] Ritva Hurri-Syrjänen and Antti V. Vähäkangas. Fractional Sobolev-Poincaré and fractional Hardy inequalities in unbounded John domains. *Mathematika*, 61(2):385–401, 2015.
- [ILTV14] Lizaveta Ihnatsyeva, Juha Lehrbäck, Heli Tuominen, and Antti V. Vähäkangas. Fractional Hardy inequalities and visibility of the boundary. *Studia Math.*, 224(1):47–80, 2014.
- [J⁺86] David Jerison et al. The Poincaré inequality for vector fields satisfying Hörmander’s condition. *Duke Mathematical Journal*, 53(2):503–523, 1986.
- [KaPPu09] Agnieszka Kał amajska and Katarzyna Pietruska-Pał uba. On a variant of the Hardy inequality between weighted Orlicz spaces. *Studia Math.*, 193(1):1–28, 2009.
- [KMP07] Alois Kufner, Lech Maligranda, and Lars-Erik Persson. *The Hardy inequality*. Vydavatelský Servis, Plzeň, 2007. About its history and some related results.
- [KP03] Alois Kufner and Lars-Erik Persson. *Weighted inequalities of Hardy type*. World Scientific Publishing Co., Inc., River Edge, NJ, 2003.
- [KR61] M. A. Krasnosel’kiĭ and Ja. B. Rutickiĭ. *Convex functions and Orlicz spaces*. Translated from the first Russian edition by Leo F. Boron. P. Noordhoff Ltd., Groningen, 1961.
- [KT78] Alois Kufner and Hans Triebel. Generalizations of Hardy’s inequality. *Confer. Sem. Mat. Univ. Bari*, (156):21 pp. (1979), 1978.
- [KZ08] Stephen Keith and Xiao Zhong. The Poincaré inequality is an open ended condition. *Annals of Mathematics*, pages 575–599, 2008.
- [LS10] Michael Loss and Craig Sloane. Hardy inequalities for fractional integrals on general domains. *J. Funct. Anal.*, 259(6):1369–1379, 2010.
- [LSY03] Elliott H Lieb, Robert Seiringer, and Jakob Yngvason. Poincaré inequalities in punctured domains. *Annals of mathematics*, pages 1067–1080, 2003.
- [MMOS14] Fumi-Yuki Maeda, Yoshihiro Mizuta, Takao Ohno, and Tetsu Shimomura. Hardy’s inequality in Musielak-Orlicz-Sobolev spaces. *Hiroshima Math. J.*, 44(2):139–155, 2014.
- [MMP98] Moshe Marcus, Victor J. Mizel, and Yehuda Pinchover. On the best constant for Hardy’s inequality in \mathbb{R}^n . *Trans. Amer. Math. Soc.*, 350(8):3237–3255, 1998.
- [MNOS11] Yoshihiro Mizuta, Eiichi Nakai, Takao Ohno, and Tetsu Shimomura. Hardy’s inequality in Orlicz-Sobolev spaces of variable exponent. *Hokkaido Math. J.*, 40(2):187–203, 2011.
- [MSV] Alberto Maione, Ariel Martin Salort, and Eugenio Vecchi. Maz’ya-Shaposhnikova formula in magnetic fractional Orlicz-Sobolev spaces. *Asymptotic Analysis (To appear)*, <https://doi.org/10.3233/ASY-211677>.
- [Sal20a] Ariel M Salort. Eigenvalues and minimizers for a non-standard growth non-local operator. *Journal of Differential Equations*, 268(9):5413–5439, 2020.
- [Sal20b] Ariel M Salort. Hardy inequalities in fractional Orlicz-Sobolev spaces. *arXiv preprint arXiv:2009.06431*, 2020.
- [SV20] Ariel M Salort and Hernán Vivas. Fractional eigenvalues in Orlicz spaces with no Δ_2 condition. *arXiv preprint arXiv:2005.01847*, 2020.
- [Yer14] Karen Yeressian. Asymptotic behavior of elliptic nonlocal equations set in cylinders. *Asymptot. Anal.*, 89(1-2):21–35, 2014.