

On Exotic Consistent Anomalies in $(1+1)d$: A Ghost Story

Chi-Ming Chang^a, Ying-Hsuan Lin^{b,c}

^a*Yau Mathematical Sciences Center, Tsinghua University, Beijing, 10084, China*

^b*Walter Burke Institute for Theoretical Physics,
California Institute of Technology, Pasadena, CA 91125, USA*

^c*Jefferson Physical Laboratory, Harvard University, Cambridge, MA 02138, USA*

cmchang@tsinghua.edu.cn, yinhslin@gmail.com

Abstract

We revisit 't Hooft anomalies in $(1+1)d$ non-spin quantum field theory, starting from the consistency and locality conditions, and find that consistent $U(1)$ and gravitational anomalies cannot always be canceled by properly quantized $(2+1)d$ classical Chern-Simons actions. On the one hand, we prove that certain exotic anomalies can only be realized by non-reflection-positive or non-compact theories; on the other hand, without insisting on reflection-positivity, the exotic anomalies present a caveat to the inflow paradigm. For the mixed $U(1)$ gravitational anomaly, we propose an inflow mechanism involving a mixed $U(1) \times SO(2)$ classical Chern-Simons action with a boundary condition that matches the $SO(2)$ gauge field with the $(1+1)d$ spin connection. Furthermore, we show that this mixed anomaly gives rise to an isotopy anomaly of $U(1)$ topological defect lines. The isotopy anomaly can be canceled by an extrinsic curvature improvement term, but at the cost of potentially uplifting $U(1)$ to a multifold cover. We end with a survey of the holomorphic bc ghost system, which realizes all the exotic consistent anomalies.

Contents

1	Introduction	1
1.1	Consistency and locality	3
2	Pure anomalies	4
2.1	Perturbative pure anomalies	5
2.2	Global gravitational anomaly	8
2.3	Global U(1) anomaly	12
3	Mixed gravitational anomaly	14
3.1	Perturbative mixed gravitational anomaly	14
3.2	Topological defects and isotopy anomaly	18
4	Holomorphic bc ghost system	21
5	Concluding remarks	24
A	Bardeen-Zumino counter-terms	26
A.1	Pure gravitational	26
A.2	Mixed gravitational	27
B	No covariant stress tensor for mixed anomaly	28

1 Introduction

't Hooft anomaly is a controlled breaking of symmetries in quantum field theory (QFT). Let Φ collectively denote the background gauge fields and metric, and Λ collectively denote diffeomorphisms and background gauge transformations. Under Λ , the partition function on Φ transforms as

$$Z[\Phi^\Lambda] = Z[\Phi] e^{i\alpha[\Phi, \Lambda]}, \tag{1.1}$$

The anomalous phase $\alpha[\Phi, \Lambda]$ is a functional that satisfies the consistency and locality conditions. Consistency — or finite Wess-Zumino consistency [1] — of an 't Hooft anomaly

requires the background gauge transformation (1.1) to respect the group multiplication law, which amounts to the commutativity of the diagram

$$\begin{array}{ccc}
 & Z[\Phi^{\Lambda_1}] & \\
 \Lambda_1 \nearrow & & \searrow \Lambda_2 \\
 Z[\Phi] & \xrightarrow{\Lambda_2 \Lambda_1} & Z[\Phi^{\Lambda_2 \Lambda_1}]
 \end{array} \tag{1.2}$$

The anomalous phases generated by the two routes differ by $2\pi\mathbb{Z}$. Locality of an 't Hooft anomaly is expected because anomaly is a short distance effect, *i.e.* it originates in the ultraviolet. The consistency and locality conditions led to the old cohomological classification of perturbative anomalies – the 't Hooft anomaly of a semi-simple Lie algebra \mathcal{G} in D spacetime dimensions is classified by the Lie algebra cohomology $H^{D+1}(\mathcal{G}, \mathbb{R})$ through the descent equations [2–13].

A more modern perspective on 't Hooft anomalies is the *inflow paradigm*: a D -dimensional anomalous QFT should be viewed as the boundary theory of a $(D + 1)$ -dimensional bulk classical action, also called a symmetry protected topological phase or an invertible field theory, such that the coupled system exhibits no anomaly [14–28]. From this perspective, the classification of boundary 't Hooft anomalies amounts to the classification of bulk classical actions. One recent triumph has been the classification of *reflection-positive* invertible topological field theories in $D + 1$ spacetime dimensions by cobordism groups [20, 25, 28].¹

For a discrete internal symmetry group G in a $(1+1)d$ non-spin QFT, the inflow paradigm suggests that the 't Hooft anomalies have the same $H^3(G, U(1))$ classification as the $(2+1)d$ Dijkgraaf-Witten theories [29]. The same classification can also be deduced from a purely $(1+1)d$ perspective [30, 31].² According to the inflow paradigm, the chiral central charge $c_- \equiv c - \bar{c}$ of a $(1+1)d$ non-spin CFT must be a multiple of eight, because only then can the gravitational anomaly be canceled by a properly quantized $(2+1)d$ gravitational Chern-Simons action. Similarly, the level $k_- \equiv k - \bar{k}$ of a $U(1)$ internal symmetry in a non-spin CFT must be an even integer for the $U(1)$ anomaly to be canceled by a $(2+1)d$ $U(1)$ Chern-Simons.

The above quantization conditions are violated by the holomorphic bc ghost system. Recall that b and c are left-moving anti-commuting free fields with weights λ and $1 - \lambda$. For

¹Reflection positivity of a QFT in Euclidean spacetime is equivalent to the unitarity of time evolutions in Lorentzian spacetime. However, in this paper we always call this property reflection-positivity, to avoid confusion of QFT unitarity with the unitarity of symmetry representations.

²The $(1+1)d$ classification is achieved by the pentagon identity, which arises as the consistency condition for the fusion category of symmetry defect lines, or equivalently from the finite Wess-Zumino consistency condition applied to patch-wise background gauge transformations.

integer λ , the holomorphic bc ghost system is a non-spin CFT, but has $c_- = 1 - 3(2\lambda - 1)^2 \in 2\mathbb{Z}$ and $k_- = 1$, suggesting that the gravitational and $U(1)$ anomalies cannot be canceled by inflow of familiar Chern-Simons actions. On the other hand, the consistency and locality conditions lead to weaker quantization conditions $c_- \in 2\mathbb{Z}$ and $k_- \in \mathbb{Z}$ that are precisely satisfied by the holomorphic bc ghost system.

The bc ghost system has a $U(1)$ ghost number symmetry, which exhibits a mixed gravitational anomaly: On any Riemann surface, it is conserved up to a background charge proportional to the Euler characteristic. In [32], it was pointed out that the mixed gravitational anomaly, albeit consistent, cannot be canceled by the inflow of a relativistic classical action if the boundary $(1+1)d$ spin connection is to be matched with the bulk $(2+1)d$ spin connection. However, a non-relativistic inflow is possible using the renowned Wen-Zee topological term [33,34]. In this paper, we propose a relativistic inflow that matches the boundary $(1+1)d$ spin connection with a bulk $SO(2)$ gauge field.

Another slightly bizarre feature of the mixed gravitational anomaly is the non-existence of an improved stress tensor with covariant anomalous conservation. Recall that a consistent anomaly requires the current to be defined via the variation of background fields, and the resulting anomalous conservation equations are generally not gauge-covariant. By adding Bardeen-Zumino currents [35], the consistent current can often be improved to a covariant one, *i.e.* with covariant anomalous conservation equations. The covariant currents are no longer equal to the variation of background fields, and do not satisfy the Wess-Zumino consistency condition. We show that this improvement is not possible for the mixed gravitational anomaly at hand.

The rest of this paper is organized as follows. Section 1.1 defines the consistency and locality conditions. Section 2 concerns pure anomalies, by first reviewing the anomaly descent and inflow of perturbative pure anomalies, and then examining the finite Wess-Zumino condition for their global versions. Section 3 explores the mixed $U(1)$ -gravitational anomaly and its connection to the isotopy anomaly of topological defect lines. Section 4 surveys the bc ghost system and finds it to realize every exotic consistent anomaly discussed in this paper. Appendix A reviews the Bardeen-Zumino counter-terms for pure gravitational anomaly, and constructs its counterpart for the mixed anomaly. Appendix B proves that the consistent mixed gravitational anomaly does not have covariant counterpart.

1.1 Consistency and locality

't Hooft anomalies satisfy two conditions: (finite Wess-Zumino) consistency and locality. Consistency amounts to the commutativity of the diagram (1.2) up to $2\pi\mathbb{Z}$ phase differences.

Condition 1.1 (Consistency). *For two arbitrary background diffeomorphism/gauge transformations Λ_1 and Λ_2 , the anomalous phases satisfy*

$$\alpha[\Phi, \Lambda_2 \Lambda_1] - \alpha[\Phi^{\Lambda_1}, \Lambda_2] - \alpha[\Phi, \Lambda_1] \in 2\pi\mathbb{Z}. \quad (1.3)$$

Locality amounts to the following two properties:

1. Under general background diffeomorphism/gauge transformations Λ , the anomalous phase $\alpha[\Phi, \Lambda]$ is a local functional of Φ .
2. Under infinitesimal background diffeomorphism/gauge transformations Λ , the anomalous phase $\alpha[\Phi, \Lambda]$ is a local functional of Φ and Λ , and vanishes when $\Phi = 0$. For the gravitational background, $\Phi = 0$ means that the spin connection (or Levi-Civita connection) vanishes, with no further constraint on the vielbein.

An argument for the second locality property can be made as follows. For continuous symmetries, the divergence of the Noether current J^μ should vanish in correlation functions up to contact terms,

$$\langle \nabla_\mu J^\mu(x) \cdots \rangle \Big|_{\Phi=0} = \text{contact terms}. \quad (1.4)$$

Had the second locality property been false, this contact structure would be violated by the anomalous Ward identities. The first locality property can be viewed as an extension of the second locality property to large background diffeomorphism/gauge transformations. The two locality properties above can be stated in more precise terms by the following locality condition.

Condition 1.2 (Locality). *Let \mathcal{G} be the space of all background diffeomorphism/gauge transformations, with connected components \mathcal{G}_n for $n = 0, 1, 2, \dots$, and with \mathcal{G}_0 containing the trivial transformation. The anomalous phase $\alpha[\Phi, \Lambda]$ takes the form*

$$\alpha[\Phi, \Lambda] = \sum_i \kappa_i(n) \mathcal{A}_i[\Phi, \Lambda] + \theta(n), \quad (1.5)$$

where $\mathcal{A}_i[\Phi, \Lambda]$ is a basis of independent local functionals that vanish in the trivial background $\Phi = 0$, and $\theta(0) = 0$.

2 Pure anomalies

This section first reviews the perturbative pure gravitational and $U(1)$ anomalies in non-spin QFT, and then examines the finite Wess-Zumino (fWZ) consistency condition for global anomalies. We derive a weaker quantization condition on the anomaly coefficients than that of inflow. A comparison can be found in Table 1.

2.1 Perturbative pure anomalies

We begin by reviewing the well-known perturbative pure anomalies. Consider a (1+1) d non-spin QFT with U(1) internal symmetry coupled to a background metric $g_{\mu\nu}$ and a background U(1) gauge field A . We parameterize the background metric by the zweibein e_μ^a and write $g_{\mu\nu} = e_\mu^a e_\nu^b \delta_{ab}$. We use μ, ν, \dots to denote spacetime indices, and a, b, \dots to denote frame indices. Under diffeomorphisms (ξ), local frame rotations (θ), and U(1) gauge transformations (λ) the background zweibein and the background U(1) gauge field transform as

$$\delta e^a = -\theta^a_b e^b + \mathcal{L}_\xi e^a, \quad \delta A = d\lambda + \mathcal{L}_\xi A, \quad (2.1)$$

where \mathcal{L}_ξ denotes the Lie-derivative.

The effective action $W[e, A] = -\log Z[e, A]$ is a complex-valued functional of the background fields, and is in general non-local. The infinitesimal part of the anomalous phase is a local functional linear in the gauge parameters,

$$\alpha[e, A, \theta, \xi, \lambda] = \mathcal{A}_\theta[e, A, \theta] + \mathcal{A}_\xi[e, A, \xi] + \mathcal{A}_\lambda[e, A, \lambda]. \quad (2.2)$$

The effective action shifts by

$$W[e + \delta e, A + \delta A] = W[e, A] - i(\mathcal{A}_\theta[e, A, \theta] + \mathcal{A}_\xi[e, A, \xi] + \mathcal{A}_\lambda[e, A, \lambda]). \quad (2.3)$$

The anomalous phases \mathcal{A}_θ , \mathcal{A}_ξ and \mathcal{A}_λ are constrained by the Wess-Zumino consistency condition [36]

$$\delta_{\chi_1} \mathcal{A}_{\chi_2} - \delta_{\chi_2} \mathcal{A}_{\chi_1} = \mathcal{A}_{[\chi_2, \chi_1]}, \quad \text{for } \chi = \theta, \xi, \lambda. \quad (2.4)$$

Descent equations

A large class of solutions to the Wess-Zumino consistency condition are obtained by the descent equations

$$\mathcal{I}^{(4)} = d\mathcal{I}^{(3)}, \quad \delta\mathcal{I}^{(3)} = d\mathcal{I}^{(2)}, \quad \mathcal{A} = 2\pi \int_{\mathcal{M}_2} \mathcal{I}^{(2)}, \quad (2.5)$$

where $\mathcal{I}^{(3)}$ and $\mathcal{I}^{(4)}$ are formal 3- and 4-forms. The 4-form anomaly polynomial responsible for the pure gravitational and U(1) anomalies is

$$\mathcal{I}^{(4)} = \frac{1}{(2\pi)^2} \left[\frac{\kappa R^2}{48} \text{tr} (R \wedge R) + \frac{\kappa F^2}{2} F \wedge F \right], \quad (2.6)$$

where $R_{ab} = \frac{1}{2} e_a^\mu e_b^\nu R_{\mu\nu\rho\sigma} dx^\rho dx^\sigma$ and $F = dA$. The descent 3-form is

$$\mathcal{I}^{(3)} = \frac{1}{(2\pi)^2} \left[\frac{\kappa R^2}{48} \text{CS}(\omega) + \frac{\kappa F^2}{2} A \wedge F \right], \quad (2.7)$$

and the anomalous phases are

$$\mathcal{A}_\theta = \frac{\kappa_{R^2}}{96\pi} \int_{\mathcal{M}_2} \theta^{ab} R_{ba}, \quad A_\xi = 0, \quad \mathcal{A}_\lambda = \frac{\kappa_{F^2}}{4\pi} \int_{\mathcal{M}_2} \lambda F. \quad (2.8)$$

Inflow mechanism

An anomaly that solves the descent equations has a natural bulk classical action. Consider

$$S_{\text{bulk}} = \frac{ik_{R^2}}{192\pi} \int_{\mathcal{M}_3} \text{CS}(\omega) + \frac{ik_{F^2}}{4\pi} \int_{\mathcal{M}_3} \text{CS}(A), \quad (2.9)$$

which, to be well-defined, must have quantized levels³

$$\frac{k_{R^2}}{8}, \quad k_{F^2} \in 2\mathbb{Z}. \quad (2.11)$$

If \mathcal{M}_3 is a three-manifold with boundary $\partial\mathcal{M}_3 = \mathcal{M}_2$, then the classical action on \mathcal{M}_3 contributes the following amount of anomaly to the $(1+1)d$ non-spin QFT on \mathcal{M}_2 ,

$$\Delta\kappa_{R^2} = -\frac{1}{2}k_{R^2}, \quad \Delta\kappa_{F^2} = -k_{F^2}. \quad (2.12)$$

For the coupled system to be free of anomalies, the quantization conditions (2.11) on the Chern-Simons levels k_{R^2} and k_{F^2} translate to

$$\frac{1}{8}\kappa_{R^2}, \quad \frac{1}{2}\kappa_{F^2} \in \mathbb{Z}. \quad (2.13)$$

Bardeen-Zumino counter-term

The Bardeen-Zumino counter-term provides a trade-off between the frame rotation anomaly and the diffeomorphism anomaly [37]. The conventional choice eliminates the former in favor of the latter. The counter-term is constructed from the zweibein $e^a{}_\mu$, with the explicit form given in (A.1). The modified effective action is

$$W'[e, A] \equiv W[e, A] + S_{\text{BZ}}[e], \quad (2.14)$$

³On a closed manifold \mathcal{M}_3 , the Chern-Simons action (2.9) is required to be invariant under background diffeomorphism and $U(1)$ gauge transformations. One way to manifest the invariance property is to rewrite the action as

$$S = \frac{ik_{R^2}}{192\pi} \int_{\mathcal{M}_4} \text{tr} R \wedge R + \frac{ik_{F^2}}{4\pi} \int_{\mathcal{M}_4} F \wedge F, \quad (2.10)$$

where \mathcal{M}_4 is a four manifold such that $\partial\mathcal{M}_4 = \mathcal{M}_3$. For (2.10) to be independent of the choice of \mathcal{M}_4 , the levels k_{R^2} and k_{F^2} must be quantized as in (2.11).

such that under local frame rotations,

$$\delta_\theta S_{\text{BZ}} = i\mathcal{A}_\theta. \quad (2.15)$$

Hence, the new effective action $W'[e, A]$ transforms as

$$W'[e + \delta e, A + \delta A] = W'[e, A] - i(\mathcal{A}_\lambda[e, A, \lambda] + \mathcal{A}'_\xi[e, A, \xi]), \quad (2.16)$$

with a nonzero anomalous phase $\mathcal{A}'_\xi[e, A, \xi]$ under diffeomorphism,

$$\mathcal{A}'_\xi = i\delta_\xi S_{\text{BZ}} = \frac{\kappa_{R^2}}{96\pi} \int_{\mathcal{M}_2} \partial_\mu \xi^\nu d\Gamma^\mu{}_\nu. \quad (2.17)$$

Anomalous conservation and covariant improvement

The anomalous phases (2.8) imply the anomalous conservation equations

$$\begin{aligned} \langle \nabla_\mu T^{\mu\nu}(x) \rangle &= -\frac{2\pi i}{\sqrt{g}} \frac{\delta \mathcal{A}'_\xi[e, A, \xi]}{\delta \xi_\nu} = \frac{i\kappa_{R^2}}{48} \frac{1}{\sqrt{g}} g^{\nu\lambda} \partial_\mu (\sqrt{g} \varepsilon^{\rho\sigma} \partial_\rho \Gamma^\mu{}_{\lambda\sigma}), \\ \langle \nabla^\mu J_\mu(x) \rangle &= -\frac{2\pi}{\sqrt{g}} \frac{\delta \mathcal{A}_\lambda[e, A, \lambda]}{\delta \lambda(x)} = -\frac{\kappa_{F^2}}{4} \varepsilon^{\mu\nu} F_{\mu\nu}. \end{aligned} \quad (2.18)$$

Note that the first equation is not covariant. In technical terms, these are consistent anomalies and not covariant anomalies [35]. To arrive at the latter, the stress tensor $T^{\mu\nu}$ must be improved by

$$\mathcal{T}^{\mu\nu} = T^{\mu\nu} - \frac{i\kappa_{R^2}}{48} \nabla_\lambda (\Gamma^{(\mu\lambda}{}_{\sigma} \varepsilon^{\nu)\sigma} - \Gamma^{\lambda(\mu}{}_{\sigma} \varepsilon^{\nu)\sigma} - \Gamma^{(\mu\nu)}{}_{\sigma} \varepsilon^{\lambda\sigma}), \quad (2.19)$$

The anomalous conservation equation for the improved stress tensor $\mathcal{T}^{\mu\nu}$ takes the covariant form

$$\langle \nabla_\mu \mathcal{T}^{\mu\nu}(x) \rangle = \frac{i\kappa_{R^2}}{48} \nabla_\mu (R^{\mu\nu}{}_{\rho\sigma} \varepsilon^{\rho\sigma}). \quad (2.20)$$

Operator product in CFT

On flat space, the two-point functions of the stress tensor $T_{\mu\nu}$ and the conserved current J_μ are constrained by conformal symmetry to be

$$\langle T_{zz}(z, \bar{z}) T_{zz}(0) \rangle = \frac{c}{2z^4}, \quad \langle T_{\bar{z}\bar{z}}(z, \bar{z}) T_{\bar{z}\bar{z}}(0) \rangle = \frac{\bar{c}}{2\bar{z}^4}, \quad (2.21)$$

and

$$\langle J_z(z, \bar{z}) J_z(0) \rangle = \frac{k}{z^2}, \quad \langle J_{\bar{z}}(z, \bar{z}) J_{\bar{z}}(0) \rangle = \frac{\bar{k}}{\bar{z}^2}. \quad (2.22)$$

The remaining components

$$\langle T_{zz}(z, \bar{z})T_{z\bar{z}}(0) \rangle, \langle T_{\bar{z}\bar{z}}(z, \bar{z})T_{z\bar{z}}(0) \rangle, \langle T_{zz}(z, \bar{z})T_{\bar{z}\bar{z}}(0) \rangle, \langle T_{z\bar{z}}(z, \bar{z})T_{z\bar{z}}(0) \rangle, \langle J_z(z, \bar{z})J_{\bar{z}}(0) \rangle$$

are contact terms with coefficients related to the anomalies. The above two point functions can be obtained from the Ward identities implied by the anomalous conservation equations (3.14), giving

$$c_- \equiv c - \bar{c} = \kappa_{R^2}, \quad k_- \equiv k - \bar{k} = \kappa_{F^2}. \quad (2.23)$$

The discussion of the $\langle TJ \rangle$ two-point functions is deferred to Section 3.1.

2.2 Global gravitational anomaly

Let us now study the pure anomaly of large diffeomorphisms. Since we do not assume time-reversal symmetry, orientation-reversing operations such as reflections are excluded. For concreteness, consider a $(1+1)d$ non-spin CFT on a torus with complex moduli τ and a flat metric

$$ds^2 = |dx^1 + \tau dx^2|^2, \quad x^\mu \cong x^\mu + 2\pi\mathbb{Z}. \quad (2.24)$$

The orientation-preserving large diffeomorphisms that respect the periodicity of the coordinates x^μ are

$$\begin{pmatrix} x^1 \\ x^2 \end{pmatrix} \rightarrow \begin{pmatrix} x'^1 \\ x'^2 \end{pmatrix} = \begin{pmatrix} a & -b \\ -c & d \end{pmatrix} \begin{pmatrix} x^1 \\ x^2 \end{pmatrix} \quad \text{for } ad - bc = 1 \quad \text{and } a, b, c, d \in \mathbb{Z}, \quad (2.25)$$

and form the mapping class group $\text{SL}(2, \mathbb{Z})$. It is generated by

$$S = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad T = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}, \quad (2.26)$$

which satisfy the relations

$$S^4 = 1, \quad (ST)^3 = S^2. \quad (2.27)$$

The form of the metric (2.24) is preserved, modulo Weyl transformations, by $\text{SL}(2, \mathbb{Z})$, with the complex moduli τ and the complex coordinate $w = x^1 + \tau x^2$ transformed as

$$\tau \rightarrow \tau' = \frac{a\tau + b}{c\tau + d}, \quad w \rightarrow w' = x'^1 + \tau' x'^2 = \frac{w}{c\tau + d}. \quad (2.28)$$

Torus partition function

Suppose the partition function on a flat torus does not vanish identically over all moduli.⁴ Under $\text{SL}(2, \mathbb{Z})$, the only possible dependence on the flat background geometry that is compatible with locality is through the volume integral $\int d^2x \sqrt{g}$. However, an anomalous phase

⁴The usual reason for a partition function to vanish identically is the existence of anti-commuting zero modes.

proportional to the volume violates the fWZ consistency condition 1.1, with Λ_1 an $\text{SL}(2, \mathbb{Z})$ transformation, and Λ_2 a Weyl transformation. Hence, the torus partition function must be invariant under $\text{SL}(2, \mathbb{Z})$ up to τ -independent anomalous phases⁵

$$Z\left(\frac{a\tau + b}{c\tau + d}, \frac{a\bar{\tau} + b}{c\bar{\tau} + d}\right) = Z(\tau, \bar{\tau}) e^{i\theta(a,b,c,d)}. \quad (2.30)$$

By the fWZ consistency condition 1.1, the general phases $\theta(a, b, c, d)$ are determined from the phases θ_S and θ_T of the S and T generators, *i.e.*

$$Z\left(-\frac{1}{\tau}, -\frac{1}{\bar{\tau}}\right) = Z(\tau, \bar{\tau}) e^{i\theta_S}, \quad Z(\tau + 1, \bar{\tau} + 1) = Z(\tau, \bar{\tau}) e^{i\theta_T}. \quad (2.31)$$

The chiral central charge is related to the T anomalous phase by $2\pi c_- = -24\theta_T$. Under the relations (2.27), fWZ constrains⁶

$$2\theta_S \in 2\pi\mathbb{Z}, \quad \theta_S + 3\theta_T \in 2\pi\mathbb{Z}. \quad (2.33)$$

There are two scenarios:

$$\begin{aligned} \text{(i)} \quad \theta_S, 3\theta_T \in 2\pi\mathbb{Z} &\Rightarrow Z(\tau, \bar{\tau}) = Z(-1/\tau, -1/\bar{\tau}), \quad c_- \in 8\mathbb{Z}, \\ \text{(ii)} \quad \theta_S, 3\theta_T \in 2\pi\left(\mathbb{Z} + \frac{1}{2}\right) &\Rightarrow Z(\tau, \bar{\tau}) = -Z(-1/\tau, -1/\bar{\tau}), \quad c_- \in 8\mathbb{Z} + 4. \end{aligned} \quad (2.34)$$

In scenario (ii), $Z(\tau = i, \bar{\tau} = -i)$ on the square torus must either blow up or vanish. The former means that the spectrum exhibits Hagedorn growth, which violates our expectation of QFT in finite volume.⁷ The latter violates reflection-positivity. See Figure 1. Hence, a reflection-positive CFT must fall into scenario (i).⁸

⁵On the flat torus (in Cartesian coordinates (2.24)) where the Christoffel symbols all vanish, no local integral term can contribute. Therefore, the fWZ consistency condition 1.1 modulo phase redefinitions defines the first group cohomology with $\text{U}(1)$ coefficients. This subsection is essentially an exercise computing

$$H^1(\text{PSL}(2, \mathbb{Z}), \text{U}(1)) = \mathbb{Z}_6, \quad H^1(\text{SL}(2, \mathbb{Z}), \text{U}(1)) = \mathbb{Z}_{12}. \quad (2.29)$$

⁶Note that the anomalous phases form a representations of $\text{PSL}(2, \mathbb{Z})$, defined by the relations

$$S^2 = (ST)^3 = 1. \quad (2.32)$$

This is physically expected because S^2 is charge conjugation, and acts trivially on a torus with no operator insertions.

⁷See [38] for a discussion. In the following we always assume that the torus partition function (for non-compact CFTs normalized by the volume) does not blow up.

⁸Many non-reflection-positive CFTs such as the $c < 0$ minimal models still have positive torus partition functions. They must also fall into scenario (i).




Figure 1: The square torus $\tau = i$, $\bar{\tau} = -1$ is symmetric under 90 degree rotations (modular S) and reflections. The partition function on the square torus transforms with a phase θ_S under the former, and must be positive in a reflection-positive theory due to the later.

More generally, an ST^n transformation produces a phase factor

$$e^{i(\theta_S + n\theta_T)} = \begin{cases} 1 & c_- \in 24\mathbb{Z}, \\ \omega^{\pm n} & c_- \in 24\mathbb{Z} \pm 8, \\ -(-)^n & c_- \in 24\mathbb{Z} + 12, \\ -(-\omega)^{\pm n} & c_- \in 24\mathbb{Z} \pm 4, \end{cases} \quad (2.35)$$

where $\omega = e^{\frac{2}{3}\pi i}$. An immediate consequence is that the partition function $Z(\tau, \bar{\tau})$ must vanish at the S -invariant point $\tau = i$ and/or the ST -invariant point $\tau = \omega$ whenever $c_- \notin 24\mathbb{Z}$. More specifically, the vanishing points in the standard fundamental domain are

$$\tau = \begin{cases} \omega & c_- \in 24\mathbb{Z} \pm 8, \\ i & c_- \in 24\mathbb{Z} + 12. \\ i, \omega & c_- \in 24\mathbb{Z} \pm 4. \end{cases} \quad (2.36)$$

As a check, the chiral half of the E_8 WZW model has $c_- = 8$, and its torus partition function $Z(\tau) = J(\tau)^{\frac{1}{3}}$ indeed vanishes at $\tau = \omega$.

Torus one-point function

One can derive similar conditions by looking at the torus one-point function

$$G(\tau, \bar{\tau}) = \langle \mathcal{O}_{h, \bar{h}}(w, \bar{w}) \rangle_{T_\tau^2}. \quad (2.37)$$

of a local operator $\mathcal{O}_{h, \bar{h}}$ that has definite holomorphic and anti-holomorphic weights h and \bar{h} but is not required to be a primary. By translational invariance, the torus one-point function does not depend on the coordinate w of the operator insertion. Under S , it transforms as

$$\langle \mathcal{O}_{h, \bar{h}}(w, \bar{w}) \rangle_{T_\tau^2} = e^{-i\theta_S} \langle \mathcal{O}_{h, \bar{h}}(w, \bar{w}) \rangle_{T_{-1/\tau}^2} = e^{-i\theta_S} \tau^{-h} \bar{\tau}^{-\bar{h}} \langle \mathcal{O}'_{h, \bar{h}}(w/\tau, \bar{w}/\bar{\tau}) \rangle_{T_{-1/\tau}^2}. \quad (2.38)$$

Under T , there is no conformal factor. In summary,

$$G\left(-\frac{1}{\tau}, -\frac{1}{\bar{\tau}}\right) = e^{i\theta_S \tau^h \bar{\tau}^{\bar{h}}} G(\tau, \bar{\tau}), \quad G(\tau + 1, \bar{\tau} + 1) = e^{i\theta_T} G(\tau, \bar{\tau}). \quad (2.39)$$

The anomalous phases θ_S and θ_T satisfy the quantization conditions

$$2\theta_S \in 2\pi\left(\mathbb{Z} + \frac{\ell}{2}\right), \quad \theta_S + 3\theta_T \in 2\pi\mathbb{Z}, \quad (2.40)$$

where $\ell = h - \bar{h}$ is the spin of the operator \mathcal{O} .

In a given theory, the torus one-point functions for different operators can have different θ_S , but they must have the same θ_T , which is related to the chiral central charge by $2\pi c_- = -24\theta_T$. A CFT with a non-vanishing torus one-point function of an operator of odd spin necessarily contains anti-commuting fields, *i.e.* ghosts if the QFT is non-spin. This is because $\mathcal{O}_{h, \bar{h}}$ must appear in the OPE of some real operator $\mathcal{O}_{h', \bar{h}'}$ with itself,

$$\mathcal{O}_{h', \bar{h}'}(z_1, \bar{z}_1) \mathcal{O}_{h', \bar{h}'}(z_2, \bar{z}_2) \ni C(z_1 - z_2)^{h-2h'} (\bar{z}_1 - \bar{z}_2)^{\bar{h}-2\bar{h}'} \mathcal{O}_{h, \bar{h}}\left(\frac{z_1 + z_2}{2}, \frac{\bar{z}_1 + \bar{z}_2}{2}\right). \quad (2.41)$$

Exchanging z_1 and z_2 produces a sign since

$$(-)^{h-\bar{h}-2(h'-\bar{h}')} = -1. \quad (2.42)$$

For the OPE coefficient C to be non-zero, the operator $\mathcal{O}_{h', \bar{h}'}$ must therefore be anti-commuting (Grassmann-valued) to produce a compensating sign.

- If the torus one-point function for at least one operator of even spin does not vanish identically over all torus moduli, then we recover the previous condition (2.34), hence $c_- \in 4\mathbb{Z}$.
- If the torus one-point functions for at least one operator of odd spin does not vanish identically over all torus moduli — which can only happen in the presence of anti-commuting fields, *i.e.* ghosts if the CFT is non-spin — then (2.40) leads to $c_- \in 4\mathbb{Z} + 2$.

Quantization of the chiral central charge

The preceding results can be summarized as follows.

Lesson 2.1. *The chiral central charge of a non-spin CFT satisfies $c_- \in 2\mathbb{Z}$ if at least one torus one-point function does not vanish identically over all moduli of the torus. If the torus partition function itself does not vanish identically, then $c_- \in 4\mathbb{Z}$. If the partition function is positive on the square torus (true if reflection-positive), then $c_- \in 8\mathbb{Z}$.⁹*

⁹The condition $c_- \in 2\mathbb{Z}$ was also found in the classification of $(2+1)d$ non-spin invertible topological orders by BF categories [39]. There is no known non-spin invertible topological order that realizes the minimal chiral central charge $c_- = \pm 2$. We thank Xiao-Gang Wen for pointing this out to us.

Note that a $(2+1)d$ bulk gravitational Chern-Simons action can cancel the global gravitational anomaly if $c_- \in 8\mathbb{Z}$, which is guaranteed for reflection-positive CFTs. If not reflection-positive and $c_- \notin 8\mathbb{Z}$, then the global gravitational anomaly is consistent but more exotic. The holomorphic bc ghost system realizes $c_- \in -2 + 24\mathbb{Z}$.

2.3 Global U(1) anomaly

Consider a $(1+1)d$ non-spin QFT with U(1) global symmetry on a genus- g Riemann surface Σ . Let \mathcal{C}_i for $i = 1, \dots, 2g$ be a basis of non-contractable cycles on the Riemann surface Σ , with intersection matrix Ω . The winding numbers of the gauge transformation λ are

$$\vec{m}[\lambda] = \frac{1}{2\pi} \int_{\vec{\mathcal{C}}} d\lambda. \quad (2.43)$$

The locality condition 1.2 dictates that the anomalous phase takes the form

$$\alpha[A, \lambda] = -\frac{\kappa(\vec{m}[\lambda])}{4\pi} \int_{\Sigma} d\lambda A + \sum_i \frac{\kappa'_i(\vec{m}[\lambda])}{2\pi} \int_{\Sigma} f_i(\lambda) F + \theta(\vec{m}[\lambda]), \quad (2.44)$$

where f_i is a basis of periodic functions,

$$f_i(\lambda + 2\pi) = f_i(\lambda), \quad (2.45)$$

and $\kappa, \kappa'_i, \theta$ are functions that satisfy

$$\kappa(0) = \kappa_{F^2}, \quad \theta(0) = 0. \quad (2.46)$$

Let us focus on background gauge orbits that are flat, so that κ'_i does not appear. Consider two large background gauge transformations λ_1 and λ_2 with nontrivial windings. For shorthand, we write

$$\vec{m}_1 \equiv \vec{m}[\lambda_1], \quad \vec{m}_2 \equiv \vec{m}[\lambda_2], \quad \vec{m}_{12} \equiv \vec{m}[\lambda_1 + \lambda_2] = \vec{m}_1 + \vec{m}_2. \quad (2.47)$$

The fWZ consistency condition 1.1 requires that

$$\begin{aligned} & \left[-\frac{\kappa(\vec{m}_{12})}{4\pi} \int_{\Sigma} d(\lambda_1 + \lambda_2) A + \theta(\vec{m}_{12}) \right] - \left[\frac{\kappa(\vec{m}_2)}{4\pi} \int_{\Sigma} d\lambda_2 (A + d\lambda_2) + \theta(\vec{m}_2) \right] \\ & - \left[\frac{\kappa(\vec{m}_1)}{4\pi} \int_{\Sigma} d\lambda_1 A + \theta(\vec{m}_1) \right] \equiv 0 \pmod{2\pi}. \end{aligned} \quad (2.48)$$

The above can be reorganized into

$$\begin{aligned} & [-\pi\kappa(\vec{m}_2) \vec{m}_1 \cdot \Omega \cdot \vec{m}_2 + \theta(\vec{m}_{12}) - \theta(\vec{m}_1) - \theta(\vec{m}_2)] - \left[\frac{\kappa(\vec{m}_{12}) - \kappa(\vec{m}_1)}{4\pi} \int_{\Sigma} d\lambda_1 A \right] \\ & - \left[\frac{\kappa(\vec{m}_{12}) - \kappa(\vec{m}_2)}{4\pi} \int_{\Sigma} d\lambda_2 A \right] \equiv 0 \pmod{2\pi}, \end{aligned} \quad (2.49)$$

where we used

$$\frac{1}{4\pi^2} \int_{\Sigma} d\lambda_1 d\lambda_2 = \vec{m}_1 \cdot \Omega \cdot \vec{m}_2. \quad (2.50)$$

Because A is an arbitrary flat connection and λ_1, λ_2 are independent and arbitrary, the coefficients in second and third brackets must separately vanish. Hence,

$$\kappa(\vec{m}[\lambda]) = \kappa(0) = \kappa_{F^2} \quad (2.51)$$

is a constant.

We left with

$$- \pi \kappa_{F^2} \vec{m}_1 \cdot \Omega \cdot \vec{m}_2 + \theta(\vec{m}_{12}) - \theta(\vec{m}_1) - \theta(\vec{m}_2) \equiv 0 \pmod{2\pi}, \quad (2.52)$$

For concreteness, let Σ be a torus, and choose a basis of cycles \mathcal{C}_i with intersection matrix

$$\Omega = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}. \quad (2.53)$$

With

$$\vec{m}_1 = (1, 0) \quad \vec{m}_2 = (-1, 0), \quad (2.54)$$

and separately

$$\vec{m}_1 = (0, 1), \quad \vec{m}_2 = (0, -1), \quad (2.55)$$

together with (2.46), we find

$$\theta(1, 0) + \theta(-1, 0) \equiv \theta(0, 1) + \theta(0, -1) \equiv 0 \pmod{2\pi}. \quad (2.56)$$

With

$$\vec{m}_1 = (m-1, n), \quad \vec{m}_2 = (1, 0), \quad (2.57)$$

and separately

$$\vec{m}_1 = (m, n-1), \quad \vec{m}_2 = (0, 1), \quad (2.58)$$

we find recurrence relations on $\theta(m, n)$ for (m, n) in the first quadrant,

$$\begin{aligned} \theta(m, n) &\equiv \theta(m-1, n) + \theta(1, 0) - \pi \kappa_{F^2} n \pmod{2\pi}, \\ \theta(m, n) &\equiv \theta(m, n-1) + \theta(0, 1) + \pi \kappa_{F^2} m \pmod{2\pi}. \end{aligned} \quad (2.59)$$

Similarly, there are recurrence relations for (m, n) in the three other quadrants. The solution in all quadrants is

$$\theta(m, n) = \theta(1, 0)m + \theta(0, 1)n - \pi \kappa_{F^2} mn. \quad (2.60)$$

Plugging this solution back into (2.52), we find the quantization condition

$$\kappa_{F^2} \in \mathbb{Z}. \quad (2.61)$$

The quantization condition (2.61) is weaker than the quantization condition (2.13) expected from the inflow of (2+1) d bulk U(1) Chern-Simons.

Mixing with the modular transforms

Let $P_{m,n}$ denote a background $U(1)$ gauge transformation with winding numbers (m, n) . From the fWZ consistency condition 1.1 for the relations

$$P_{1,0} S = S P_{0,1}, \quad T P_{1,0} = P_{1,1} T, \quad (2.62)$$

one deduces

$$\theta(1, 0) = \theta(0, 1) = \pi \kappa_{F^2}. \quad (2.63)$$

Quantization of the level

In CFT, the anomaly coefficient and the level are related by $\kappa_{F^2} = k_-$.

Lesson 2.2. *The level k_- of a $U(1)$ current algebra in a non-spin CFT must satisfy $k_- \in \mathbb{Z}$ if the flavored torus partition function does not vanish identically over all moduli of the torus and all flat gauge backgrounds.*

Note that a $(2+1)d$ bulk $U(1)$ Chern-Simons action can cancel the anomaly if k_- is even. The holomorphic bc ghost system realizes $k_- = 1$.

3 Mixed gravitational anomaly

This section examines the mixed $U(1)$ -gravitational anomaly in a $(1+1)d$ non-spin QFT. In the first part of this section, we characterize the mixed gravitational anomaly by descent and inflow, examine the possibility of a covariant improvement, and study the imprint of the anomaly on local operator products in CFT. In the second part, we study the mixed gravitational anomaly from the perspective of topological defects, and show that the mixed gravitational anomaly gives rise to an isotopy anomaly.

3.1 Perturbative mixed gravitational anomaly

In the following, A and F denote the $U(1)$ connection and field strength, and ω denotes the spin connection, with R its field strength. We use a, b, \dots to denote frame indices.

Descent equations

The mixed gravitational anomaly is described by the anomaly polynomial,

$$\mathcal{I}^{(4)} = \frac{\kappa_{FR}}{(2\pi)^2} F \wedge (\varepsilon^{ab} R_{ba}) . \quad (3.1)$$

The descent 3-form is¹⁰

$$\mathcal{I}^{(3)} = \frac{1}{(2\pi)^2} \left[\frac{\kappa_{FR}}{2} A \wedge \varepsilon^{ab} R_{ba} + \frac{\kappa_{FR}}{2} F \wedge \varepsilon^{ab} \omega_{ba} + sd (A \wedge \varepsilon^{ab} \omega_{ba}) \right] , \quad (3.2)$$

where the ambiguity s is related to the freedom of adding the Bardeen counter-term

$$S_B = -\frac{is'}{2\pi} \int A \wedge (\varepsilon^{ab} \omega_{ba}) . \quad (3.3)$$

Its addition to the action shifts the ambiguity s to $s + s'$. The anomalous phases are

$$\mathcal{A}_\lambda = \frac{1}{2\pi} \left(\frac{\kappa_{FR}}{2} - s \right) \int_{\mathcal{M}_2} \lambda \varepsilon^{ab} R_{ba} , \quad \mathcal{A}_\theta = \frac{1}{2\pi} \left(\frac{\kappa_{FR}}{2} + s \right) \int_{\mathcal{M}_2} \theta^{ab} \varepsilon_{ba} F , \quad \mathcal{A}_\xi = 0 . \quad (3.4)$$

Inflow mechanism

Can the mixed gravitational anomaly of a $(1+1)d$ non-spin QFT be canceled by coupling to a $(2+1)d$ classical action? When the $(2+1)d$ spacetime is a product manifold $\mathcal{M}_3 = \mathcal{M}_2 \times [0, \infty)$, one could consider the $(2+1)d$ classical action of the renowned Wen-Zee topological term [33, 34] relevant for the Hall viscosity in non-relativistic quantum Hall systems (see [40–42] for the connection)

$$\frac{ik_{FR}}{16\pi} \int_{\mathcal{M}_2 \times [0, \infty)} \varepsilon^{ab} \omega_{ab} \wedge F , \quad (3.5)$$

where \mathcal{M}_2 is the spatial manifold, and the anomaly coefficient k_{FR} is also called the spin vector.¹¹ The above inflow action explicitly breaks $(2+1)d$ Lorentz invariance, and thus requires non-relativistic geometry to generalize to non-product manifolds.

We propose a slightly different inflow mechanism that preserves $(2+1)d$ Lorentz invariance. Consider the mixed Chern-Simons term

$$\frac{ik_{FR}}{4\pi} \int_{\mathcal{M}_3} A \wedge F_R , \quad (3.6)$$

where \mathcal{M}_3 is a three-dimensional manifold whose boundary is \mathcal{M}_2 , and $F_R = dA_R$ is the field strength of a background $\text{SO}(2)$ gauge field on \mathcal{M}_3 . The matching condition at $\partial\mathcal{M}_3 = \mathcal{M}_2$

¹⁰The descent equation $\mathcal{I}^{(4)} = d\mathcal{I}^{(3)}$ is insensitive to the addition of exact terms (total derivatives).

¹¹We thank Xiao-Gang Wen and Juven Wang for bringing our attention to [33] and [34].

is such that the normal component of A_R vanishes, and the tangent components of A_R are identified with the boundary $(1+1)d$ spin connection by

$$A_R|_{\mathcal{M}_2} = \frac{1}{\zeta} \varepsilon^{ab} \omega_{ba}, \quad (3.7)$$

with a proportionality constant ζ to be fixed by flux quantization. The flux of $\varepsilon^{ab} \omega_{ba}$ can be computed as

$$\int_{\mathcal{M}_2} \varepsilon^{ab} R_{ba} = - \int_{\mathcal{M}_2} d^2x \sqrt{g} R = -4\pi\chi. \quad (3.8)$$

Depending on whether the theory is defined only on orientable Riemann surfaces, for instance when there is no time-reversal symmetry, or on general Riemann surfaces, the Euler characteristic is quantized as $\chi \in 2\mathbb{Z}$ or $\chi \in \mathbb{Z}$, respectively. Hence, flux quantization determines

$$\zeta = \begin{cases} 4 & \mathcal{M}_2 \text{ orientable,} \\ 2 & \mathcal{M}_2 \text{ general.} \end{cases} \quad (3.9)$$

To cancel the mixed gravitational anomaly of the $(1+1)d$ QFT, the Chern-Simons level is chosen to be

$$k_{FR} = 2\zeta\kappa_{FR}. \quad (3.10)$$

The quantization condition for the Chern-Simons level is

$$k_{FR} \in 2\mathbb{Z} \quad (3.11)$$

translates to a quantization condition on the mixed gravitational anomaly coefficient

$$\kappa_{FR} \in \begin{cases} \frac{1}{4}\mathbb{Z} & \mathcal{M}_2 \text{ orientable,} \\ \frac{1}{2}\mathbb{Z} & \mathcal{M}_2 \text{ general.} \end{cases} \quad (3.12)$$

We will see in Section 4 that the holomorphic bc ghost system realizes $\kappa_{FR} \in \frac{1}{4} + \frac{1}{2}\mathbb{Z}$. It is in principle possible to derive a quantization condition on κ_{FR} from fWZ alone without the need of inflow. However, to probe κ_{FR} requires considering curved Riemann surfaces and is beyond the scope of this paper.

Bardeen-Zumino counter-terms

By adding a mixed Bardeen-Zumino counter-term $S_{\text{BZ}}^{\text{mixed}}$ which we construct in Appendix A.2, the anomalous phase \mathcal{A}_θ under frame rotations can be completely canceled, while generating

an extra contribution to the anomalous phase A'_ξ under diffeomorphisms. In summary, the new anomalous phases are

$$\begin{aligned}
\mathcal{A}'_\theta &= \mathcal{A}_\theta + i\delta_\theta S_{\text{BZ}}^{\text{mixed}} = 0, \\
\mathcal{A}'_\xi &= i\delta_\xi S_{\text{BZ}}^{\text{mixed}} = \frac{1}{2\pi} \left(\frac{\kappa_{FR}}{2} + s \right) \int_{\mathcal{M}_2} \partial_\mu \xi^\nu d(\varepsilon^\mu{}_\nu A), \\
\mathcal{A}'_\lambda &= \mathcal{A}_\lambda + i\delta_\lambda S_{\text{BZ}}^{\text{mixed}} = \frac{1}{2\pi} \int_{\mathcal{M}_2} \lambda \left[\kappa_{FR} \varepsilon^{ab} R_{ba} - \left(\frac{\kappa_{FR}}{2} + s \right) d(\varepsilon^\nu{}_\mu \Gamma^\mu{}_\nu) \right].
\end{aligned} \tag{3.13}$$

Anomalous conservation and covariant improvement

The anomalous phases (3.13) give the non-covariant anomalous conservation equations for the consistent currents,

$$\begin{aligned}
\langle \nabla_\mu T^{\mu\nu}(x) \rangle &= -\frac{2\pi i}{\sqrt{g}} \frac{\delta \mathcal{A}'_\xi[e, A, \xi]}{\delta \xi_\nu} = i \left(\frac{\kappa_{FR}}{2} + s \right) \frac{1}{\sqrt{g}} g^{\nu\lambda} \partial_\mu [\sqrt{g} \varepsilon^{\rho\sigma} \partial_\rho (\varepsilon^\mu{}_\lambda A_\sigma)], \\
\langle \nabla^\mu J_\mu(x) \rangle &= -\frac{2\pi}{\sqrt{g}} \frac{\delta \mathcal{A}'_\lambda[e, A, \lambda]}{\delta \lambda(x)} = \kappa_{FR} R + \left(\frac{\kappa_{FR}}{2} + s \right) \nabla_\rho (\varepsilon^{\rho\sigma} \varepsilon^\nu{}_\mu \Gamma^\mu{}_\nu).
\end{aligned} \tag{3.14}$$

The conservation of U(1) can be covariantized by improving the consistent current J^μ with Bardeen-Zumino currents, which are terms that depend only on the background fields and vanish when $A_\mu = 0$ and $g_{\mu\nu} = \delta_{\mu\nu}$. More precisely, the improved current is

$$\mathcal{J}^\mu = J^\mu - \left(\frac{\kappa_{FR}}{2} + s \right) \varepsilon^{\mu\nu} \varepsilon^\rho{}_\sigma \Gamma^\sigma{}_{\rho\nu}, \tag{3.15}$$

which has a covariant form of the anomalous conservation equation

$$\langle \nabla^\mu \mathcal{J}_\mu(x) \rangle = -\frac{\kappa_{FR}}{4} \varepsilon^{\mu\nu} F_{\mu\nu} + \kappa_{FR} R. \tag{3.16}$$

However, by an explicit computation in Appendix B, we show that no covariant improvement of the stress tensor exists.

Operator product in CFT

In flat space CFT, the two-point functions between the U(1) current J_μ and the stress tensor $T_{\mu\nu}$ must take the form

$$\langle T_{zz}(z) J_z(0) \rangle = \frac{\alpha}{z^3}, \quad \langle T_{\bar{z}\bar{z}}(\bar{z}) J_{\bar{z}}(0) \rangle = \frac{\bar{\alpha}}{\bar{z}^3}, \tag{3.17}$$

which imply the commutation relations¹²

$$[L_m, J_n] = -nJ_{m+n} + \frac{m(m+1)}{2}\alpha\delta_{m+n}, \quad [\bar{L}_m, \bar{J}_n] = -n\bar{J}_{m+n} + \frac{m(m+1)}{2}\bar{\alpha}\delta_{m+n}. \quad (3.18)$$

One also has the contact terms

$$\begin{aligned} \langle T_{zz}(z, \bar{z})J_{\bar{z}}(0) \rangle &= 2\pi\beta\partial\delta^{(2)}(z, \bar{z}), & \langle T_{\bar{z}\bar{z}}(z, \bar{z})J_z(0) \rangle &= 2\pi\bar{\beta}\bar{\partial}\delta^{(2)}(z, \bar{z}), \\ \langle T_{z\bar{z}}(z, \bar{z})J_{\bar{z}}(0) \rangle &= 2\pi\gamma\bar{\partial}\delta^{(2)}(z, \bar{z}), & \langle T_{\bar{z}z}(z, \bar{z})J_z(0) \rangle &= 2\pi\bar{\gamma}\partial\delta^{(2)}(z, \bar{z}). \end{aligned} \quad (3.19)$$

Matching the above with the anomalous Ward identities implied by the anomalous conservation equations (3.14), we arrive at the relations

$$\begin{aligned} \alpha + 2\beta &= 4\left(\frac{\kappa_{FR}}{2} - s\right), & \bar{\alpha} + 2\bar{\beta} &= 4\left(\frac{\kappa_{FR}}{2} - s\right), & \gamma + \bar{\gamma} &= -2\left(\frac{\kappa_{FR}}{2} - s\right), \\ \beta + \gamma &= -\left(\frac{\kappa_{FR}}{2} + s\right), & \bar{\beta} + \bar{\gamma} &= -\left(\frac{\kappa_{FR}}{2} + s\right), \\ \alpha + 2\bar{\gamma} &= 2\left(\frac{\kappa_{FR}}{2} + s\right), & \bar{\alpha} + 2\gamma &= 2\left(\frac{\kappa_{FR}}{2} + s\right). \end{aligned} \quad (3.20)$$

In particular,

$$\alpha + \bar{\alpha} = 4\kappa_{FR} \quad (3.21)$$

is insensitive to the coefficient s of the Bardeen counter-term.

When α or $\bar{\alpha}$ is nonzero, the operator J_z or $J_{\bar{z}}$ is not a Virasoro primary operator, respectively. In a compact reflection-positive CFT, an operator must be either primary or descendant (see for example [44]). Therefore, there must exist an operator \mathcal{O} of dimension $(h, \bar{h}) = (0, 0)$ such that $L_{-1}\mathcal{O} = J_z$ or $\bar{L}_{-1}\mathcal{O} = J_{\bar{z}}$. However, in a compact reflection-positive CFT, the only dimension zero operator is the identity which is annihilated by the Virasoro generators L_{-1} and \bar{L}_{-1} . We have learned the following.

Lesson 3.1. *A 2d CFT with mixed $U(1)$ -gravitational anomaly cannot be compact and reflection-positive.*

3.2 Topological defects and isotopy anomaly

An invertible topological defect line (TDL) can be constructed from a covariant current \mathcal{J}_μ that is conserved up to covariant anomalies,

$$\mathcal{L}_\eta(\mathcal{C}) = : \exp \left[i\eta \oint_{\mathcal{C}} ds n_\mu \mathcal{J}^\mu \right] :, \quad (3.22)$$

¹²In particular, $[L_0, J_0] = \alpha$. In the vertex operator algebra (VOA) language, $[L_1, J(0)] \neq 0$ means that the VOA is “not of strong CFT type”. For a strongly rational holomorphic VOA (which requires it to be of strong CFT type), it was proven by [43] that the central charge must be a multiple of 8.




Figure 2: Deforming a symmetry defect line from the curve \mathcal{C} across the domain \mathcal{D} to the new curve \mathcal{C}' .

where n_μ is the normal vector to the curve \mathcal{C} . The defect is topological up to an isotopy anomaly: When the curve \mathcal{C} is deformed across a domain \mathcal{D} , as shown in Figure 2, the defect \mathcal{L}_η is modified by a phase factor determined by the divergence theorem,

$$: \exp \left[i\eta \oint_{\partial\mathcal{D}} ds n_\mu \mathcal{J}^\mu \right] := : \exp \left[i\eta \int_{\mathcal{D}} d^2x \sqrt{g} \nabla_\mu \mathcal{J}^\mu \right] := \exp \left[i\eta \kappa_{FR} \int_{\mathcal{D}} d^2x \sqrt{g} R \right]. \quad (3.23)$$

The isotopy anomaly generalizes the mixed gravitational anomaly to discrete groups and non-invertible topological defects [45]. For discrete groups, there is no analog of Lesson 3.1. In particular, an anomalous \mathbb{Z}_2 symmetry line in compact reflection-positive CFTs has isotopy anomaly. Note that a defect line defined using the consistent current J_μ is not topological. Even on flat space, its anomalous conservation depends on the choice of coordinate system. The consistent and covariant currents agree only in Cartesian coordinates on flat space.

An uplifting discussion

On flat space, for every each $\eta \in \mathbb{Z}$, the defect \mathcal{L}_η commutes with all local operators and is therefore identified with the trivial line, reflecting the periodicity of $U(1)$. On curved space, this family of lines differ by their isotopy anomaly, and the periodicity of $U(1)$ is ruined.¹³

¹³In particular, a point of localized curvature can carry an arbitrary real amount of charge.

A remedy is to modify the topological defect by a local improvement term¹⁴

$$\tilde{\mathcal{L}}_\eta(\mathcal{C}) = \mathcal{L}_\eta(\mathcal{C}) \exp \left[-i\eta\kappa_{FR} \oint_{\mathcal{C}} ds K \right], \quad (3.24)$$

such that the isotopy anomaly is precisely canceled via the Gauss-Bonnet theorem. However, only when $\kappa_{FR} \in \frac{\mathbb{Z}}{2}$ does this fully restore the periodicity of U(1). Otherwise, the distinction between $\tilde{\mathcal{L}}_{\eta=0}$ and $\tilde{\mathcal{L}}_{\eta=1}$ can be detected by the loop expectation value on the plane,¹⁵

$$\langle \tilde{\mathcal{L}}_\eta(\mathcal{C}) \rangle_{\mathbb{R}^2} = \exp[-4\pi i\eta\kappa_{FR}]. \quad (3.26)$$

If the mixed gravitational anomaly is such that $\kappa_{FR} \in \frac{\mathbb{Z}}{2}$, then $\tilde{\mathcal{L}}_\eta$ defines a U(1) symmetry with the same periodicity as U(1), and is truly topological on curved backgrounds, *i.e.* free of isotopy anomaly, unlike \mathcal{L}_η . Hence $\tilde{\mathcal{L}}_\eta$ is in all respects better than \mathcal{L}_η . However, if $\kappa_{FR} \notin \frac{\mathbb{Z}}{2}$, then we are faced with a dilemma.

Lesson 3.2. *If the mixed gravitational anomaly is such that $\kappa_{FR} \notin \frac{\mathbb{Z}}{2}$, then*

1. *The abelian symmetry implemented by \mathcal{L}_η is U(1) on flat space, but is non-compact \mathbb{R} on curved background.*
2. *The abelian symmetry implemented by $\tilde{\mathcal{L}}_\eta$ is a multifold cover $\widetilde{\text{U}(1)}$ on any space, flat or curved.*

If the quantization condition (3.12), $\kappa_{FR} \in \frac{\mathbb{Z}}{4}$, obtained from inflow considerations is universally true, then $\widetilde{\text{U}(1)}$ is at most a two-fold lift.

Isotopy anomaly as contact term

On flat space, the isotopy anomaly can be detected by the contact terms in the OPE between the stress tensor and the symmetry defect \mathcal{L}_η . Using the two-point functions (3.17) and (3.19), we find

$$\begin{aligned} \langle T_{zz}(z, \bar{z}) \mathcal{L}_\eta \rangle &= \eta \left\langle : \oint_{\mathcal{C}} \left[\frac{\alpha}{(z-w)^3} dw - 2\pi\beta \partial_z \delta^{(2)}(z-w, \bar{z}-\bar{w}) d\bar{w} \right] \mathcal{L}_\eta : \right\rangle \\ &= -i\pi(\alpha + 2\beta)\eta \partial_z^2 \theta(z \in D) \langle \mathcal{L}_\eta \rangle, \end{aligned} \quad (3.27)$$

¹⁴In [45], this term was called an extrinsic curvature “counter-term” by the present authors. However, from a purely (1+1)*d* point of view, it is more appropriately regarded as an improvement term for a defect operator.

¹⁵The loop expectation value of a TDL \mathcal{L} on the plane was denoted by $R(\mathcal{L})$ in [45]. If \mathcal{C} is the unit circle on flat space, then

$$\oint_{\mathcal{C}} ds K = 4\pi. \quad (3.25)$$

where we have assumed that TDL \mathcal{L}_η is located on the boundary of a compact region D , i.e. $\mathcal{C} = \partial D$. Similarly, we also have

$$\begin{aligned}\langle T_{\bar{z}\bar{z}}(z, \bar{z})\mathcal{L}_\eta \rangle &= -i\pi(\bar{\alpha} + 2\bar{\beta})\alpha\partial_{\bar{z}}^2\theta(z \in D)\langle \mathcal{L}_\eta \rangle, \\ \langle T_{z\bar{z}}(z, \bar{z})\mathcal{L}_\eta \rangle &= -2\pi i(\gamma + \bar{\gamma})\alpha\partial_z\partial_{\bar{z}}\theta(z \in D)\langle \mathcal{L}_\eta \rangle.\end{aligned}\tag{3.28}$$

4 Holomorphic bc ghost system

The anomalies of the previous sections will now find life in a specific theory — the holomorphic bc ghost system, which is a CFT of anti-commuting complex free fields b and c , with weights

$$h_b = \lambda, \quad h_c = 1 - \lambda\tag{4.1}$$

and OPE

$$b(z)c(0) \sim \frac{1}{z}.\tag{4.2}$$

The stress tensor and the $U(1)$ current for the ghost number symmetry that assigns charges ± 1 to c and b are

$$T_{zz} = (1 - \lambda) : (\partial b)c : - \lambda : b\partial c : , \quad J_z = : bc : .\tag{4.3}$$

The anomalies coefficients, computed from the TT , jj , and Tj OPEs, are

$$c_- = \kappa_{R^2} = 1 - 3(2\lambda - 1)^2, \quad k_- = \kappa_{F^2} = 1, \quad \kappa_{FR} = \frac{2\lambda - 1}{4}.\tag{4.4}$$

To be a non-spin CFT, the spins of b and c must be integers, hence

$$\lambda \in \mathbb{Z}.\tag{4.5}$$

The $U(1)$ anomaly coefficient κ_F^2 is not an even integer, so it cannot be canceled by a bulk $(2+1)d$ $U(1)$ Chern-Simons. Likewise, the gravitational anomaly coefficient $\kappa_{R^2} \in 8\mathbb{Z} - 2$ is an even integer but not a multiple of eight. Hence, the gravitational anomaly cannot be canceled by a bulk $(2+1)d$ gravitational Chern-Simons. They do, however, satisfy and saturate the quantization conditions derived from the finite Wess-Zumino consistency conditions.

Torus one-point function of the ghost number current

Let us consider the holomorphic bc ghost system on a flat torus with complex moduli τ , with periodic boundary conditions around both the space and Euclidean time cycles. The torus

partition function vanishes due to the zero modes of the b and c fields. Consider instead the torus one-point function of the current J_z ,

$$G(\tau, \bar{\tau}) = \langle J_z(0) \rangle_{T\bar{\tau}} = \eta(\tau)^2. \quad (4.6)$$

Under modular S and T transformations, the anomalous phases are

$$\theta_S = \frac{3\pi}{2}, \quad \theta_T = \frac{\pi}{6}, \quad (4.7)$$

which satisfy the quantization condition (2.40) for $\ell = 1$.

Flavored torus partition function

Consider a flat torus with metric¹⁶

$$ds^2 = (d\sigma^1)^2 + (d\sigma^2)^2, \quad \sigma^1 \cong \sigma^1 + 2\pi\mathbb{Z}, \quad \sigma^2 \cong \sigma^2 + 2\pi\mathbb{Z}, \quad (4.8)$$

where $\tau = \tau_1 + \tau_2 i$ is the complex moduli, and let us compute the partition function of the bc system on this torus with constant background gauge field

$$A = A_1 d\sigma^1 + A_2 d\sigma^2. \quad (4.9)$$

A natural thing to evaluate is the trace

$$Z_H(\tau, z) = \text{Tr} \left(q^{L_0 - \frac{c}{24}} e^{2\pi i(z - \frac{1}{2})J_0} \right) = \frac{\theta_1(\tau|z)}{\eta(\tau)}, \quad (4.10)$$

where the chemical potential z is related to the constant background gauge field A by

$$z = -i\tau_2(A_1 + iA_2). \quad (4.11)$$

However, $Z_H(\tau, z)$ does not satisfy the transformation law (1.1). The resolution is an extra term B that comes from carefully taking the Legendre transformation that relates the Lagrangian to the Hamiltonian [46, 47], resulting in

$$Z(\tau, \bar{\tau}, z, \bar{z}) = Z_H(\tau, z) e^{\pi B(\tau, \bar{\tau}, z, \bar{z})}. \quad (4.12)$$

The function B a quadratic function in z and \bar{z} (by nature of the Legendre transform), that vanishes when $A_1 = 0$, and transforms under $\text{SL}(2, \mathbb{Z})$ as

$$B \left(\frac{a\tau + b}{c\tau + d}, \frac{a\bar{\tau} + b}{c\bar{\tau} + d}, \frac{z}{c\tau + d}, \frac{\bar{z}}{c\bar{\tau} + d} \right) = B(\tau, \bar{\tau}, z, \bar{z}) - \frac{icz^2}{c\tau + d} \quad (4.13)$$

¹⁶This is a different coordinate system from (2.24).

so that the flavored partition function $Z(\tau, \bar{\tau}, z, \bar{z})$ is invariant under $\text{SL}(2, \mathbb{Z})$ up to anomalous phases. It is fixed to be

$$B(\tau, \bar{\tau}, z, \bar{z}) = \frac{z(z - \bar{z})}{2\tau_2}. \quad (4.14)$$

Under the modular S and T transformations, the flavored partition function transforms as

$$\begin{aligned} Z(\tau + 1, \bar{\tau} + 1, z, \bar{z}) &= e^{\frac{\pi i}{6}} Z(\tau, \bar{\tau}, z, \bar{z}), \\ Z\left(-\frac{1}{\tau}, -\frac{1}{\bar{\tau}}, \frac{z}{\tau}, \frac{\bar{z}}{\bar{\tau}}\right) &= e^{\frac{3\pi i}{2}} Z(\tau, \bar{\tau}, z, \bar{z}), \end{aligned} \quad (4.15)$$

which agrees with the anomalous phases (4.7). Under a large gauge transformation

$$A \rightarrow A + d\lambda, \quad \lambda = m \left(\sigma^1 - \frac{\tau_1}{\tau_2} \sigma_2 \right) + n \frac{\sigma^2}{\tau_2}, \quad (4.16)$$

the flavored partition function transforms as

$$\begin{aligned} Z(\tau, \bar{\tau}, A + d\lambda) &= Z(\tau, \bar{\tau}, A) \exp[-\pi i(m\tau_2 A_2 - (n - m\tau_1)A_1) - (mn + m + n)\pi i] \\ &= Z(\tau, \bar{\tau}, A) \exp\left(-\frac{i}{4\pi} \int d\lambda A - (mn + m + n)\pi i\right), \end{aligned} \quad (4.17)$$

which also agrees with (2.44) with $\kappa_{F^2} = 1$.

Another uplifting discussion

From Lesson (3.2), we learned that when the mixed gravitational anomaly is such that $\kappa_{FR} \in \frac{1}{4} + \frac{\mathbb{Z}}{2}$, which is the case for the holomorphic bc ghost system with integer λ , the improved topological defect line $\tilde{\mathcal{L}}_\eta$ generates a symmetry group $\widetilde{\text{U}}(1)$ that is a two-fold lift of the standard ghost number $\text{U}(1)$. Correspondingly, the pure $\widetilde{\text{U}}(1)$ anomaly has coefficient

$$\kappa_{\widetilde{F}^2} = 4\kappa_{F^2} = 4, \quad (4.18)$$

and thus can be the inflow of a properly quantized $\widetilde{\text{U}}(1)$ classical Chern-Simons action.

From a purely $(1+1)d$ perspective, the fact that the pure anomaly coefficient $\kappa_{F^2} = 1 \notin 2\mathbb{Z}$ for the $\text{U}(1)$ internal symmetry (quantized such that the local operators span integer charges) makes the mapping of the anomaly to discrete subgroups subtle and confusing. When $\kappa_{F^2} \in 2\mathbb{Z}$, the \mathbb{Z}_N subgroup of the $\text{U}(1)$ has anomaly

$$\frac{\kappa_{F^2}}{2} \bmod N \in H^3(\mathbb{Z}_N, \text{U}(1)). \quad (4.19)$$

However, when $\kappa_{F^2} \in 2\mathbb{Z} + 1$, the anomaly of the \mathbb{Z}_N subgroup does not fall in the classification by group cohomology. This phenomenon can be described more concretely as follows. Consider say three parallel \mathbb{Z}_3 symmetry defect lines wrapped around the time direction on a torus. By the standard axioms of topological defect lines [45], they should fuse to the trivial line *with no extra phase*. Now, if we embed this configuration into $U(1)$, it is described by the gauge background

$$A = \frac{2\pi}{3}\delta(x) + \frac{2\pi}{3}\delta(x - \frac{2\pi}{3}) + \frac{2\pi}{3}\delta(x - \frac{4\pi}{3}), \quad (4.20)$$

where x is the spatial direction. The key point here is that this $U(1)$ background gauge configuration is a pure gauge that has winding number one around the spatial circle, *i.e.*

$$A = d\lambda, \quad \lambda = \frac{2\pi}{3}\theta(x) + \frac{2\pi}{3}\theta(x - \frac{2\pi}{3}) + \frac{2\pi}{3}\theta(x - \frac{4\pi}{3}). \quad (4.21)$$

It is thus related to the trivial configuration by a large background gauge transformation, which can produce an anomalous phase via the $\theta(\vec{m}[\lambda])$ term in (2.44). When $\kappa_{F^2} \in 2\mathbb{Z} + 1$, the anomalous phase is nontrivial, and thus the fusion of three \mathbb{Z}_3 symmetry defect lines on the torus produces the trivial defect line but *with a sign*, violating the standard axioms. None of these issues would arise for $\widetilde{U(1)}$.

The preceding discussion suggests that perhaps $\widetilde{U(1)}$ is more sensible than $U(1)$ as the symmetry group of the holomorphic bc ghost system, despite the charge quantization of local operators being even integers.

5 Concluding remarks

Starting with the finite Wess-Zumino consistency condition (1.3), we derived quantization conditions on the pure gravitational and $U(1)$ anomaly coefficients κ_{R^2} and κ_{F^2} in $(1+1)d$ non-spin quantum field theory. The quantization conditions turned out to be weaker than those predicted by the inflow of properly quantized classical Chern-Simons actions. We also examined the mixed $U(1)$ -gravitational anomaly, proposed an inflow mechanism, and from inflow derived a quantization condition on κ_{FR} . It may be possible to derive a quantization condition on κ_{FR} from the finite Wess-Zumino consistency alone without invoking inflow, but this requires going beyond the flat torus background to *e.g.* a genus-two Riemann surface, and is beyond the scope of this paper. The quantization conditions are summarized in Table 1. A survey of the holomorphic bc ghost system found the theory to realize the minimal quantization condition for all three anomalies.

We called an anomaly exotic if

$$\kappa_{R^2} \notin 8\mathbb{Z} \quad \text{or} \quad \kappa_{F^2} \notin 2\mathbb{Z} \quad \text{or} \quad \kappa_{FR} \neq 0, \quad (5.1)$$

	Inflow	fWZ
$\kappa_{R^2} = c_-$	$8\mathbb{Z}$	$2\mathbb{Z}$
$\kappa_{F^2} = k_-$	$2\mathbb{Z}$	\mathbb{Z}
κ_{FR}	$\frac{1}{4}\mathbb{Z}$?

Table 1: Quantization of anomaly coefficients predicted by inflow of classical Chern-Simons actions versus the finite Wess-Zumino consistency condition (1.3).

and proved that certain exotic anomalies cannot be realized in any compact reflection-positive non-spin conformal field theory. Non-reflective-positivity is no reason to dismiss these exotic anomalies.¹⁷ Besides the central role Faddeev-Popov ghosts play in gauge theories, ghost fields also appear in interesting holographic contexts, including a purported holographic dual of dS₄ higher spin gravity [51–53], and supergroup gauge theories [54–56].

The mixed gravitational anomaly discussed in Section 3 has a natural even D -dimensional generalization, described by an anomalous phase \mathcal{A}_λ that involves the D -dimensional Euler form E_D ,

$$\mathcal{A}_\lambda = \kappa_{FR} \int_{\mathcal{M}_D} \lambda E_D, \quad E_D = \frac{1}{(2\pi)^{\frac{D}{2}}} \varepsilon^{a_1, \dots, a_D} R_{a_1 a_2} \wedge \dots \wedge R_{a_{D-1} a_D}, \quad (5.2)$$

where we ignored the ambiguity from Bardeen counter-terms. An inflow mechanism of this anomaly involves a mixed classical Chern-Simons action of a background U(1) gauge field and a background SO(D) gauge field that matches with the D -dimensional spin-connection by a boundary matching condition analogous to (3.7). On product manifolds $\mathcal{M}_D \times [0, 1)$ with a distinguished time direction, a higher-dimensional generalization of the Wen-Zee topological term [33, 34] can also provide the inflow. Further studies of the higher-dimensional mixed gravitational anomaly is left for future work.

Acknowledgements

We are grateful to Po-Shen Hsin, Chao-Ming Jian, Shu-Heng Shao, Ryan Thorngren, Yifan Wang and Xiao-Gang Wen for helpful discussions and comments. CC thanks the hospitality of National Taiwan University. YL is supported by the Sherman Fairchild Foundation, by the U.S. Department of Energy, Office of Science, Office of High Energy Physics, under Award Number DE-SC0011632, and by the Simons Collaboration Grant on the Non-Perturbative Bootstrap.

¹⁷Lattice models at criticality need not be reflection-positive. Non-reflection-positive CFTs are known to be important landmarks in RG space that in fact influence reflection-positive RG flows [48, 49]. Quantum field theory realizing non-integer “ $O(N)$ ” symmetry is necessarily non-reflection-positive [50].

A Bardeen-Zumino counter-terms

In this appendix, we review the pure Bardeen-Zumino counter-term of [37], and construct a mixed Bardeen-Zumino counter-term for the mixed U(1)-gravitational anomaly of Section 3.

A.1 Pure gravitational

Let us treat the vielbein $e^a{}_\mu$ as a matrix and denote it by E . The two-dimensional pure Bardeen-Zumino term is

$$S_{\text{BZ}} = -\frac{i}{2\pi} \int_{\mathcal{M}_2} \int_0^1 dt \frac{\kappa R^2}{48} \text{tr}(Hd\Gamma_t) = -\frac{i}{2\pi} \int_{\mathcal{M}_2} \int_0^1 d\tau \frac{\kappa R^2}{48} \text{tr}(Hd\omega_\tau), \quad (\text{A.1})$$

where the H is

$$E = e^H, \quad (\text{A.2})$$

and the Γ_t and the ω_t are defined by

$$\Gamma_t = E^t \Gamma E^{-t} + E^t dE^{-t} = E^{-1+t} \omega E^{1-t} + E^{-1+t} dE^{1-t} = \omega_\tau, \quad (\text{A.3})$$

where $\tau = 1-t$. The matrix valued Christoffel one-form Γ and spin connection ω are defined by

$$\Gamma^\mu{}_\nu \equiv \Gamma^\mu{}_{\nu\rho} dx^\rho, \quad \omega^a{}_b \equiv \omega^a{}_{b\mu} dx^\mu. \quad (\text{A.4})$$

In the matrix notation, diffeomorphisms and local frame rotations act on the vielbein E , Christoffel one-form Γ , and spin connection ω as

$$\begin{aligned} \delta_\xi E &= (\mathcal{L}_\xi + T_\Lambda)E, & \delta_\xi \Gamma &= (\mathcal{L}_\xi + T_\Lambda)\Gamma, \\ T_\Lambda E &\equiv E\Lambda, & T_\Lambda \Gamma &\equiv d\Lambda + [\Gamma, \Lambda], \\ \delta_\theta E &= -\theta E, & \delta_\theta \omega &= d\theta + [\omega, \theta]. \end{aligned} \quad (\text{A.5})$$

where \mathcal{L}_ξ is the Lie derivative.¹⁸ The gauge parameter Λ is related to the diffeomorphism parameter ξ by

$$\Lambda^\rho{}_\mu = \partial_\mu \xi^\rho. \quad (\text{A.7})$$

The Γ_t transforms under T_Λ as

$$T_\Lambda \Gamma_t = d\Lambda_t + [\Gamma_t, \Lambda_t] \equiv T_{\Lambda_t} \Gamma_t, \quad \Lambda_t \equiv E^t \Lambda E^{-t} + E^t (T_\Lambda E^{-t}). \quad (\text{A.8})$$

¹⁸An useful identity between the Lie derivative \mathcal{L}_ξ , exterior derivative d and interior product ι_ξ is

$$\mathcal{L}_\xi = d\iota_\xi + \iota_\xi d. \quad (\text{A.6})$$

We also have the identities

$$\begin{aligned}\frac{\partial \Lambda_t}{\partial t} &= [H, \Lambda_t] - T_\Lambda H, \\ \frac{\partial \Gamma_t}{\partial t} &= -dH + [H, \Gamma_t].\end{aligned}\tag{A.9}$$

Using the above, we compute the diffeomorphism variation of the Bardeen-Zumino action to be

$$\delta_\xi S_{\text{BZ}} = -\frac{i}{2\pi} \int_{\mathcal{M}_2} \frac{\kappa_{R^2}}{48} \text{tr}(\Lambda d\Gamma).\tag{A.10}$$

By a similar computation, we find

$$\delta_\theta S_{\text{BZ}} = \frac{i}{2\pi} \int_{\mathcal{M}_2} \frac{\kappa_{R^2}}{48} \text{tr}(\theta d\omega).\tag{A.11}$$

Hence, adding the Bardeen-Zumino counter-term S_{BZ} to the effective action $W[e, A]$, we cancel the pure frame rotation anomaly, *i.e.* \mathcal{A}_θ in (2.8), while introducing a pure diffeomorphism anomaly (2.17).

A.2 Mixed gravitational

We introduce a mixed Bardeen-Zumino action

$$S_{\text{BZ}}^{\text{mixed}} = -\frac{i}{2\pi} \int_0^1 dt \int_{\mathcal{M}_2} \left(\frac{\kappa_{FR}}{2} + s \right) \text{tr} (Hd(\mathcal{E}_t \mathbb{A})).\tag{A.12}$$

The matrix \mathcal{E}_t is defined by

$$\mathcal{E}_t = E^t \mathcal{E} E^{-t},\tag{A.13}$$

where the matrix \mathcal{E} is the $(1+1)d$ Levi-Civita tensor $\varepsilon^\mu{}_\nu$. Note that the matrix $\mathcal{E} \equiv \mathcal{E}_1$ is the Levi-Civita symbol $\varepsilon^a{}_b$ with local Lorentz indices. The \mathcal{E}_t transforms under T_Λ as

$$T_\Lambda \mathcal{E}_t = [\mathcal{E}_t, \Lambda_t].\tag{A.14}$$

We also have identity

$$\frac{\partial \mathcal{E}_t}{\partial t} = H E^t \mathcal{E} E^{-t} - E^t \mathcal{E} E^{-t} H = [H, \mathcal{E}_t].\tag{A.15}$$

Using the above, we obtain the diffeomorphism variation of the mixed Bardeen-Zumino action to be

$$\delta_\xi S_{\text{BZ}}^{\text{mixed}} = -\frac{i}{2\pi} \int_{\mathcal{M}_2} \left(\frac{\kappa_{FR}}{2} + s \right) \text{tr}(\Lambda d(\mathcal{E} A)).\tag{A.16}$$

Similarly, we have the variation of the mixed Bardeen-Zumino term under local frame rotations,

$$\delta_\theta S_{\text{BZ}}^{\text{mixed}} = \frac{i}{2\pi} \int_{\mathcal{M}_2} \left(\frac{\kappa_{FR}}{2} + s \right) \text{tr}(\theta \mathcal{E}) dA.\tag{A.17}$$

To derive the variation of the mixed Bardeen-Zumino term under the background U(1) gauge transformation, let us first rewrite the mixed Bardeen-Zumino term by integrating out the auxiliary variable t in (A.12) as

$$S_{\text{BZ}}^{\text{mixed}} = -\frac{i}{2\pi} \int_{\mathcal{M}_2} \left(\frac{\kappa_{FR}}{2} + s \right) \text{tr} (\omega \mathcal{E} - \Gamma \mathcal{E}) A. \quad (\text{A.18})$$

Under background U(1) gauge transformations, the mixed Bardeen-Zumino term becomes

$$\delta_\lambda S_{\text{BZ}}^{\text{mixed}} = -\frac{i}{2\pi} \int_{\mathcal{M}_2} \left(\frac{\kappa_{FR}}{2} + s \right) \lambda \text{tr} (\mathcal{E} d\omega - d(\Gamma \mathcal{E})). \quad (\text{A.19})$$

B No covariant stress tensor for mixed anomaly

Let us examine the possibility of improving the stress tensor such that the mixed gravitational anomaly becomes covariant. The most general improvement terms linear in derivatives and linear in A come in two forms, ∂A and ΓA . For the first form, it is clear that there are two possibilities

$$\partial^{(\mu} A^{\nu)}, \quad g^{\mu\nu} \partial^\sigma A_\sigma. \quad (\text{B.1})$$

For the second form, if A takes a μ, ν index, then we have

$$g^{\rho\sigma} \Gamma_{\rho\sigma}^\mu A^\nu, \quad g^{\mu\rho} \Gamma_{\rho\sigma}^\sigma A^\nu, \quad (\text{B.2})$$

and if A takes a dummy index that is contracted, then we have

$$g^{\mu\rho} \Gamma_{\rho\sigma}^\nu A^\sigma, \quad g^{\mu\rho} g^{\nu\sigma} \Gamma_{\rho\sigma}^\tau A_\tau, \quad g^{\mu\nu} \Gamma_{\rho\sigma}^\rho A^\sigma, \quad g^{\mu\nu} g^{\rho\sigma} \Gamma_{\rho\sigma}^\tau A_\tau. \quad (\text{B.3})$$

Hence, the most general improvement takes the form

$$\begin{aligned} Y_2^{\mu\nu} = & c_1 \Gamma^{(\mu\nu)\rho} A_\rho + c_2 \Gamma^{\rho\mu\nu} A_\rho + c_3 \Gamma^{(\underline{\mu}\rho\sigma} g_{\rho\sigma} A^{\underline{\nu})} + c_4 g_{\rho\sigma} \Gamma^{\rho\sigma(\mu} A^{\nu)} \\ & + c_5 g^{\mu\nu} \Gamma^\sigma_{\sigma\rho} A^\rho + c_6 g^{\mu\nu} g_{\sigma\lambda} \Gamma^{\rho\sigma\lambda} A_\rho + c_7 g^{(\underline{\mu}\rho} g^{\underline{\nu})\sigma} \partial_\rho A_\sigma + c_8 g^{\mu\nu} g^{\rho\sigma} \partial_\rho A_\sigma. \end{aligned} \quad (\text{B.4})$$

The only possible covariant form of the conservation equation is

$$\langle \nabla_\mu \mathcal{T}^{\mu\nu}(x) \rangle \supset \nabla_\mu F^{\mu\nu}. \quad (\text{B.5})$$

Using the MathGR package [57], it is straightforward to evaluate $\nabla_\mu Y_2^{\mu\nu}$, $\nabla_\mu F^{\mu\nu}$ and the consistent mixed gravitational anomaly in $\nabla_\mu T^{\mu\nu}$ in conformal gauge. The results can be decomposed with respect to a basis (with the overall conformal factor e^{-4w} stripped off)

$$\begin{aligned} (A\partial)w\partial_\nu w, \quad A_\nu \partial^2 w, \quad (A\partial)\partial_\nu w, \quad \partial_\nu w(\partial A), \\ (\partial_\nu A_\rho)\partial_\rho w, \quad \partial_\nu(\partial A), \quad \partial_\rho w \partial_\rho A_\nu, \quad \partial^2 A_\nu, \end{aligned} \quad (\text{B.6})$$

where $\partial A = \partial_\rho A_\rho$ and $A\partial = A_\rho\partial_\rho$. In this basis, the eight terms in $\nabla_\mu Y_2^{\mu\nu}$ can be represented by a coefficient matrix

$$\begin{pmatrix} 2 & -1 & -1 & -2 & 0 & \frac{1}{2} & -1 & \frac{1}{2} \\ 4 & 0 & -2 & -2 & -2 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -2 & 1 & 1 & 1 & 0 & 0 & 1 & 0 \\ -2 & 0 & 1 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & -1 & 0 & 1 & 0 \\ -4 & 0 & 2 & 0 & 2 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}. \quad (\text{B.7})$$

The covariant anomaly $\nabla_\mu F^{\mu\nu}$ is represented by

$$(0 \ 0 \ 0 \ 0 \ -2 \ 1 \ 2 \ -1), \quad (\text{B.8})$$

and the consistent anomaly in $\nabla_\mu T^{\mu\nu}$ is represented by

$$(0 \ 0 \ 0 \ 0 \ 0 \ -1 \ 0 \ 1). \quad (\text{B.9})$$

No combination of (B.8) with the rows of (B.7) produces (B.9). Hence, no covariant stress tensor exists.

References

- [1] J. A. d. Azcárraga and J. M. Izquierdo, *Lie Groups, Lie Algebras, Cohomology and some Applications in Physics*. Cambridge University Press, 4, 2011.
- [2] R. Stora, *Continuum Gauge Theories, Conf. Proc. C* **7607121** (1976) 201.
- [3] B. Zumino, *CHIRAL ANOMALIES AND DIFFERENTIAL GEOMETRY: LECTURES GIVEN AT LES HOUCHES, AUGUST 1983*, in *Les Houches Summer School on Theoretical Physics: Relativity, Groups and Topology*, pp. 1291–1322, 10, 1983.
- [4] B. Zumino, Y.-S. Wu, and A. Zee, *Chiral Anomalies, Higher Dimensions, and Differential Geometry, Nucl. Phys. B* **239** (1984) 477–507.
- [5] R. Stora, *ALGEBRAIC STRUCTURE AND TOPOLOGICAL ORIGIN OF ANOMALIES, NATO Sci. Ser. B* **115** (1984).
- [6] L. Faddeev and S. L. Shatashvili, *Algebraic and Hamiltonian Methods in the Theory of Nonabelian Anomalies, Theor. Math. Phys.* **60** (1985) 770–778.

- [7] J. Manes, R. Stora, and B. Zumino, *Algebraic Study of Chiral Anomalies*, *Commun. Math. Phys.* **102** (1985) 157.
- [8] R. Stora, *ALGEBRAIC STRUCTURE OF CHIRAL ANOMALIES*, in *XVI GIFT International Seminar on Theoretical Physics: New Perspectives in Quantum Field Theory*, pp. 339–371, 9, 1985.
- [9] B. Zumino, *Cohomology of Gauge Groups: Cocycles and Schwinger Terms*, *Nucl. Phys. B* **253** (1985) 477–493.
- [10] J. Manes and B. Zumino, *NONTRIVIALITY OF GAUGE ANOMALIES*, in *Nuffield Workshop on Supersymmetry and its Applications*, p. 0003, 12, 1985.
- [11] D. Kastler and R. Stora, *A DIFFERENTIAL GEOMETRIC SETTING FOR BRS TRANSFORMATIONS AND ANOMALIES. 1.*, *J. Geom. Phys.* **3** (1986) 437.
- [12] D. Kastler and R. Stora, *A DIFFERENTIAL GEOMETRIC SETTING FOR BRS TRANSFORMATIONS AND ANOMALIES. 2.*, *J. Geom. Phys.* **3** (1986) 483–505.
- [13] A. Alekseev, Y. Madaichik, L. Faddeev, and S. L. Shatashvili, *Derivation of Anomalous Commutators in Functional Integral Formalism*, *Theor. Math. Phys.* **73** (1988) 1149–1151.
- [14] J. Callan, Curtis G. and J. A. Harvey, *Anomalies and Fermion Zero Modes on Strings and Domain Walls*, *Nucl. Phys. B* **250** (1985) 427–436.
- [15] S. Ryu, J. E. Moore, and A. W. Ludwig, *Electromagnetic and gravitational responses and anomalies in topological insulators and superconductors*, *Phys. Rev. B* **85** (2012) 045104, [[arXiv:1010.0936](#)].
- [16] X. Chen, Z.-C. Gu, Z.-X. Liu, and X.-G. Wen, *Symmetry protected topological orders and the group cohomology of their symmetry group*, *Phys. Rev. B* **87** (2013), no. 15 155114, [[arXiv:1106.4772](#)].
- [17] X.-G. Wen, *Classifying gauge anomalies through symmetry-protected trivial orders and classifying gravitational anomalies through topological orders*, *Phys. Rev. D* **88** (2013), no. 4 045013, [[arXiv:1303.1803](#)].
- [18] A. Kapustin and R. Thorngren, *Anomalies of discrete symmetries in various dimensions and group cohomology*, [arXiv:1404.3230](#).
- [19] D. S. Freed, *Anomalies and Invertible Field Theories*, *Proc. Symp. Pure Math.* **88** (2014) 25–46, [[arXiv:1404.7224](#)].
- [20] A. Kapustin, *Symmetry Protected Topological Phases, Anomalies, and Cobordisms: Beyond Group Cohomology*, [arXiv:1403.1467](#).
- [21] J. C. Wang, Z.-C. Gu, and X.-G. Wen, *Field theory representation of gauge-gravity symmetry-protected topological invariants, group cohomology and beyond*, *Phys. Rev. Lett.* **114** (2015), no. 3 031601, [[arXiv:1405.7689](#)].

- [22] D. V. Else and C. Nayak, *Classifying symmetry-protected topological phases through the anomalous action of the symmetry on the edge*, *Phys. Rev. B* **90** (2014), no. 23 235137, [arXiv:1409.5436].
- [23] C.-T. Hsieh, G. Y. Cho, and S. Ryu, *Global anomalies on the surface of fermionic symmetry-protected topological phases in (3+1) dimensions*, *Phys. Rev. B* **93** (2016), no. 7 075135, [arXiv:1503.01411].
- [24] E. Witten, *Fermion Path Integrals And Topological Phases*, *Rev. Mod. Phys.* **88** (2016), no. 3 035001, [arXiv:1508.04715].
- [25] D. S. Freed and M. J. Hopkins, *Reflection positivity and invertible topological phases*, arXiv:1604.06527.
- [26] E. Witten, *The "Parity" Anomaly On An Unorientable Manifold*, *Phys. Rev. B* **94** (2016), no. 19 195150, [arXiv:1605.02391].
- [27] M. Guo, P. Putrov, and J. Wang, *Time reversal, $SU(N)$ Yang–Mills and cobordisms: Interacting topological superconductors/insulators and quantum spin liquids in 3+1D*, *Annals Phys.* **394** (2018) 244–293, [arXiv:1711.11587].
- [28] K. Yonekura, *On the cobordism classification of symmetry protected topological phases*, *Commun. Math. Phys.* **368** (2019), no. 3 1121–1173, [arXiv:1803.10796].
- [29] R. Dijkgraaf and E. Witten, *Topological Gauge Theories and Group Cohomology*, *Commun. Math. Phys.* **129** (1990) 393.
- [30] D. Gaiotto, A. Kapustin, N. Seiberg, and B. Willett, *Generalized Global Symmetries*, *JHEP* **02** (2015) 172, [arXiv:1412.5148].
- [31] L. Bhardwaj and Y. Tachikawa, *On finite symmetries and their gauging in two dimensions*, *JHEP* **03** (2018) 189, [arXiv:1704.02330].
- [32] Y. Nakayama, *Realization of impossible anomalies*, *Phys. Rev. D* **98** (2018), no. 8 085002, [arXiv:1804.02940].
- [33] X. Wen and A. Zee, *Shift and spin vector: New topological quantum numbers for the Hall fluids*, *Phys. Rev. Lett.* **69** (1992) 953–956. [Erratum: *Phys.Rev.Lett.* 69, 3000 (1992)].
- [34] B. Han, H. Wang, and P. Ye, *Generalized Wen-Zee Terms*, *Phys. Rev. B* **99** (2019), no. 20 205120, [arXiv:1807.10844].
- [35] W. A. Bardeen, *Anomalous ward identities in spinor field theories*, *Physical Review* **184** (1969), no. 5 1848–1859.
- [36] J. Wess and B. Zumino, *Consequences of anomalous Ward identities*, *Phys. Lett. B* **37** (1971) 95–97.

- [37] W. A. Bardeen and B. Zumino, *Consistent and Covariant Anomalies in Gauge and Gravitational Theories*, *Nucl. Phys. B* **244** (1984) 421–453.
- [38] D. Simmons-Duffin, *Tasi lectures on the conformal bootstrap*, *arXiv preprint arXiv:1602.07982* (2016).
- [39] L. Kong and X.-G. Wen, *Braided fusion categories, gravitational anomalies, and the mathematical framework for topological orders in any dimensions*, *arXiv:1405.5858*.
- [40] N. Read, *Non-Abelian adiabatic statistics and Hall viscosity in quantum Hall states and $p(x) + ip(y)$ paired superfluids*, *Phys. Rev. B* **79** (2009) 045308, [[arXiv:0805.2507](#)].
- [41] N. Read and E. Rezayi, *Hall viscosity, orbital spin, and geometry: paired superfluids and quantum Hall systems*, *Phys. Rev. B* **84** (2011) 085316, [[arXiv:1008.0210](#)].
- [42] B. Bradlyn and N. Read, *Low-energy effective theory in the bulk for transport in a topological phase*, *Phys. Rev. B* **91** (2015), no. 12 125303, [[arXiv:1407.2911](#)]. [Erratum: *Phys.Rev.B* 93, 239902 (2016)].
- [43] Y. Zhu, *Modular invariance of characters of vertex operator algebras*, *Journal of the American Mathematical Society* **9** (1996), no. 1 237–302.
- [44] D. Simmons-Duffin, *The Conformal Bootstrap*, in *Theoretical Advanced Study Institute in Elementary Particle Physics: New Frontiers in Fields and Strings*, pp. 1–74, 2017. [arXiv:1602.07982](#).
- [45] C.-M. Chang, Y.-H. Lin, S.-H. Shao, Y. Wang, and X. Yin, *Topological Defect Lines and Renormalization Group Flows in Two Dimensions*, *JHEP* **01** (2019) 026, [[arXiv:1802.04445](#)].
- [46] P. Kraus, *Lectures on black holes and the $AdS(3) / CFT(2)$ correspondence*, *Lect. Notes Phys.* **755** (2008) 193–247, [[hep-th/0609074](#)].
- [47] E. Dyer, A. L. Fitzpatrick, and Y. Xin, *Constraints on Flavored 2d CFT Partition Functions*, *JHEP* **02** (2018) 148, [[arXiv:1709.01533](#)].
- [48] V. Gorbenko, S. Rychkov, and B. Zan, *Walking, weak first-order transitions, and complex cfts*, *Journal of High Energy Physics* **2018** (Oct, 2018).
- [49] S. Giombi, R. Huang, I. R. Klebanov, S. S. Pufu, and G. Tarnopolsky, *$o(n)$ model in $4 \leq d \leq 6$: Instantons and complex cfts*, *Phys. Rev. D* **101** (Feb, 2020) 045013.
- [50] D. J. Binder and S. Rychkov, *Deligne Categories in Lattice Models and Quantum Field Theory, or Making Sense of $O(N)$ Symmetry with Non-integer N* , *JHEP* **04** (2020) 117, [[arXiv:1911.07895](#)].
- [51] D. Anninos, T. Hartman, and A. Strominger, *Higher Spin Realization of the dS/CFT Correspondence*, *Class. Quant. Grav.* **34** (2017), no. 1 015009, [[arXiv:1108.5735](#)].

- [52] G. S. Ng and A. Strominger, *State/Operator Correspondence in Higher-Spin dS/CFT*, *Class. Quant. Grav.* **30** (2013) 104002, [[arXiv:1204.1057](#)].
- [53] C.-M. Chang, A. Pathak, and A. Strominger, *Non-Minimal Higher-Spin DS₄/CFT₃*, [arXiv:1309.7413](#).
- [54] T. Okuda and T. Takayanagi, *Ghost D-branes*, *JHEP* **03** (2006) 062, [[hep-th/0601024](#)].
- [55] C. Vafa, *Non-Unitary Holography*, [arXiv:1409.1603](#).
- [56] R. Dijkgraaf, B. Heidenreich, P. Jefferson, and C. Vafa, *Negative Branes, Supergroups and the Signature of Spacetime*, *JHEP* **02** (2018) 050, [[arXiv:1603.05665](#)].
- [57] Y. Wang, *MathGR: a tensor and GR computation package to keep it simple*, [arXiv:1306.1295](#).