

A note on the closed-form solution for the longest head run problem of Abraham de Moivre

Yaakov Malinovsky *

Department of Mathematics and Statistics
University of Maryland, Baltimore County, Baltimore, MD 21250, USA

December 29, 2021

Abstract

The problem of the longest head run was introduced and solved by Abraham de Moivre in the second edition of his book *Doctrine of Chances* (de Moivre, 1738). The closed-form solution as a finite sum involving binomial coefficients was provided in Uspensky (1937). Since then, the problem and its variations and extensions have found broad interest and diverse applications. Surprisingly, a very simple closed form can be obtained, which we present in this note.

Keywords: longest run problem; generating functions; history of probability

1 Introduction

In a series of n independent trials, an event E has a probability p of occurrence for each trial. If, in these trials, event E occurs *at least* r times without interruption, then we have a *run* of size r . What is the probability $y_{n,r}$ of having a run of size r in n trials? This problem was formulated and solved by Abraham de Moivre in the second edition of his book *The Doctrine of Chances: or, A Method of Calculating the Probabilities of Events in Play* (de Moivre (1738), Problem LXXXVIII, p. 243). Although more than 280 years have passed since then, de Moivre's problem and its variations remain of great interest in probability and statistics; see for example Novak (2017) and references therein.

*email: yaakovm@umbc.edu

Formally, let X_1, \dots, X_n be a sequence of independent and identically distributed Bernoulli random variables with the probability of success $p = P(X_1 = 1) = P(E)$ and the probability of failure $1 - p = P(X_1 = 0)$. We denote by

$$L_n = \max \{k : X_{i+1} = \dots = X_{i+k} = 1, i = 0, 1, \dots, n - k\}$$

the length of the longest consecutive series of trials where event E occurs. Then, recalling that having a run of size r is defined as the occurrence of at least r consecutive successes in n iid Bernoulli trials with corresponding probability $y_{r,n}$, we obtain

$$y_{n,r} = P(L_n \geq r).$$

de Moivre did not provide a proof, but demonstrated a method of finding $y_{n,r}$. Reviewing that method, one can see that he used the method of generating functions. He demonstrated the method with an example of ten trials having $p = 1/2$, in which the probability of a run of size 3 equals $65/128$.

The closed-form solution of $y_{n,r}$ was given by Uspensky (1937) as a polynomial with binomial coefficients, arrived at by first obtaining a difference equation and then using a method of generating functions to solve it. Surprisingly, a simple closed form of $y_{n,r}$ can be obtained as a corollary from the difference equation given by Uspensky. This closed-form solution seems to have never been reported in the literature. In this note, we present it along with Uspensky's original derivations.

2 Uspensky's solution

We present Uspensky's solution (Uspensky (1937), pages 77-79) while keeping his original notations. This solution demonstrates the power of the use of ordinary linear difference equations along with the generating functions. Let's consider $n + 1$ trials with the corresponding probability $y_{n+1,r}$. A run of size r in $n + 1$ trials can happen in two mutually-exclusive ways:

(W1) : the run is obtained in the first n trials or

(W2) : the run is obtained as of trial $n + 1$.

(W2) means that among the first $n - r$ trials, there is no run of size r ; event E^C occurred at trial $n - r + 1$; and event E occurred in the trials $n - r + 2, \dots, n + 1$. Combining (W1) and (W2), we obtain a linear difference equation of order $r + 1$,

$$y_{n+1,r} = y_{n,r} + (1 - y_{n-r,r})qp^r, \quad (1)$$

with initial conditions

$$y_{0,r} = y_{1,r} = \dots = y_{r-1,r} = 0, \quad y_{r,r} = p^r,$$

where $q = 1 - p$.

Using a method of generating functions, Uspensky obtained the closed-form solution of (1)

$$y_{n,r} = 1 - \beta_{n,r} + p^r \beta_{n-r,r}$$

$$\beta_{n,r} = \sum_{l=0}^{\lfloor \frac{n}{r+1} \rfloor} (-1)^l \binom{n-lr}{l} (qp^r)^l, \quad (2)$$

where $\lfloor x \rfloor$ is defined as the greatest integer less than or equal to x .

Going back to de Moivre's original example, where $n = 10, r = 3$ and $p = 1/2$, and using (2), we obtain $y_{n,r} = 65/128$, which coincides with de Moivre's answer on page 245 of his *Doctrine of Chances* (de Moivre, 1738).

3 Closed-form solution for the case $r \geq n/2$

Surprisingly, a simple closed-form solution follows from Uspensky's equation (1) as follows: Replacing $n + 1$ with n , the difference equation (1) becomes

$$y_{n,r} = y_{n-1,r} + (1 - y_{n-1-r,r})qp^r, \quad (3)$$

with initial conditions $y_{0,r} = \dots = y_{r-1,r} = 0, y_{r,r} = p^r$.

If $n - 1 - r \leq r - 1$ (i.e. $n/2 \leq r$), then $y_{n-1-r,r} = 0$ and we obtain from (3)

$$y_{n,r} = y_{n-1,r} + qp^r \quad \text{for } n/2 \leq r, \quad (4)$$

with initial conditions $y_{0,r} = \dots = y_{r-1,r} = 0, y_{r,r} = p^r$.

Iterating (4), we obtain the following corollary:

Corollary 1. *If $n/2 \leq r \leq n$, where r is an integer, then*

$$y_{n,r} = p^r + (n - r)p^r q. \quad (5)$$

An alternative explanation for (5) is that, when $n/2 \leq r \leq n$, we cannot have two separate runs of size r . Therefore, a run of size r in n trials either begins at position 1 (this happens with probability p^r) or at the positions $2, \dots, n - (r - 1)$ (at each such position, this happens with probability qp^r). Then, denote by R_i an event in which a run of size r begins at position $i, i = 1, 2, \dots, n - (r - 1)$. Since the events $R_1, R_2, \dots, R_{n-(r-1)}$ are disjoint, we obtain that $y_{n,r} = P(R_1) + P(R_2) + \dots + P(R_{n-(r-1)}) = p^r + (n - r)qp^r$.

4 Comments

There are a number of interesting problems discussed in the Uspensky (1937) book, many of which have roots in the classics of probability, their origins tracing back to founders of modern-probability such as Pascal, Fermat, Huygens, Bernoulli, de Moivre, Laplace, Markoff, Bernstein, and others. For example, Uspensky (1937) considered another problem of de Moivre's that was later discussed and extended by Diaconis and Zabell (1991). A large collection of classic problems in probability with historical comments and original citations are nicely presented in the book by Gorroochurn (2012).

There are many follow-ups on and extensions of de Moivre's longest head run problem. An interesting recursive solution of the problem in the case of the fair coin was given by Székely and Tusnády (1979). That problem was then extended to the Markov chain setting, where Uspensky's generation function was generalized to the case of dependent observations (Novak, 1989). The problem has also found applications in numerous fields. Among which are reliability (Derman *et al.*, 1982), computational biology (Schbath, 2000), and finance where time dependent-sequences naturally occur (see Novak (2011) and references therein).

Acknowledgement

I thank Serguei Novak for comments on the early version of this article, and I thank Danny Segev for following the early version of this article on arXiv and suggesting the alternative

proof that is now presented after Corollary 1. I also would like to thank the referee for insightful and helpful comments that led to significant improvements in the paper.

References

- de Moivre, A. (1738). The doctrine of chances. 2d ed. H Woodfall, London.[Publicly available on website of Linda Hall Library <http://lhldigital.lindahall.org/cdm/ref/collection/math/id/16415>].
- Derman, C., Liberman, G. J., and Ross, S. M.(1982). On the consecutive-k-out-of-n:F systems. *IEEE Trans. Reliab.* **31**, 57–63.
- Diaconis, P., Zabell, S. (1991). Closed form summation for classical distributions: Variations on a theme of de Moivre. *Statist. Sci.* **6**, 284–302.
- Gorroochurn, P. (2012). Classic problems of probability. Wiley, Hoboken, NJ.
- Novak, S.Y. (1989). Asymptotic expansions in the problem of the length of the longest head-run for Markov chain with two states. *Trudy Inst. Math. (Novosibirsk)* **13**, 136–147. In Russian.
- Novak, S.Y. (2011). Extreme value methods with application to Finance. Chapman&Hall/CRC press, London.
- Novak, S.Y. (2017). On the length of the longest head run. *Stat Probab Lett.* **130**, 111–114.
- Schbath S. (2000). An overview on the distribution of word counts in Markov chains. *J. Comput. Biology* **7**, 193–201.
- Székel, G., Tusnády, G. (1979). Generalized Fibonacci numbers, and the number of "pure heads". *Matematikai Lapok* **27**, 147–151. In Hungarian.
- Uspensky, J. V.(1937). Introduction to mathematical probability. McGraw-Hill Book Company.