

Exponential decay for semilinear wave equations with viscoelastic damping and delay feedback

Alessandro Paolucci*

Cristina Pignotti†

Abstract

In this paper we study a class of semilinear wave type equations with viscoelastic damping and delay feedback with time variable coefficient. By combining semigroup arguments, careful energy estimates and an iterative approach we are able to prove, under suitable assumptions, a well-posedness result and an exponential decay estimate for solutions corresponding to *small* initial data. This extends and concludes the analysis initiated in [16] and then developed in [13, 17].

1 Introduction

Let H be a Hilbert space and let A be a positive self-adjoint operator with dense domain $D(A)$ in H and compact inverse in H . Let us consider the system:

$$\begin{aligned} u_{tt}(t) + Au(t) - \int_0^{+\infty} \mu(s)Au(t-s)ds + k(t)BB^*u_t(t-\tau) &= \nabla\psi(u(t)), \quad t \in (0, +\infty) \\ u(t) &= u_0(t), \quad t \in (-\infty, 0], \\ u_t(0) &= u_1 \\ B^*u_t(t) &= g(t), \quad t \in (-\tau, 0), \end{aligned} \tag{1.1}$$

where $\tau > 0$ represents the time delay, B is a bounded linear operator of H into itself, B^* denotes its adjoint, and $(u_0(\cdot), u_1, g(\cdot))$ are the initial data taken in suitable spaces. Moreover, the delay damping coefficient $k : [0, +\infty) \rightarrow \mathbb{R}$ is a function in $L^1_{loc}([0, +\infty))$ such that

$$\int_{t-\tau}^t |k(s)|ds < C^*, \quad \forall t \in (0, +\infty), \tag{1.2}$$

for a suitable constant C^* , and the memory kernel $\mu : [0, +\infty) \rightarrow [0, +\infty)$ satisfies the following assumptions:

- (i) $\mu \in C^1(\mathbb{R}^+) \cap L^1(\mathbb{R}^+)$;
- (ii) $\mu(0) = \mu_0 > 0$;

*Dipartimento di Ingegneria e Scienze dell'Informazione e Matematica, Università di L'Aquila, Via Vetoio, Loc. Coppito, 67010 L'Aquila Italy (alessandro.paolucci2@graduate.univaq.it).

†Dipartimento di Ingegneria e Scienze dell'Informazione e Matematica, Università di L'Aquila, Via Vetoio, Loc. Coppito, 67010 L'Aquila Italy (pignotti@univaq.it).

$$(iii) \int_0^{+\infty} \mu(t) dt = \tilde{\mu} < 1;$$

$$(iv) \mu'(t) \leq -\delta \mu(t), \text{ for some } \delta > 0.$$

Furthermore, $\psi : D(A^{\frac{1}{2}}) \rightarrow \mathbb{R}$ is a functional having Gâteaux derivative $D\psi(u)$ at every $u \in D(A^{\frac{1}{2}})$. Moreover, in the spirit of [2], we assume the following hypotheses:

(H1) For every $u \in D(A^{\frac{1}{2}})$, there exists a constant $c(u) > 0$ such that

$$|D\psi(u)(v)| \leq c(u) \|v\|_H \quad \forall v \in D(A^{\frac{1}{2}}).$$

Then, ψ can be extended to the whole H and we denote by $\nabla\psi(u)$ the unique vector representing $D\psi(u)$ in the Riesz isomorphism, i.e.

$$\langle \nabla\psi(u), v \rangle_H = D\psi(u)(v), \quad \forall v \in H;$$

(H2) for all $r > 0$ there exists a constant $L(r) > 0$ such that

$$\|\nabla\psi(u) - \nabla\psi(v)\|_H \leq L(r) \|A^{\frac{1}{2}}(u - v)\|_H,$$

for all $u, v \in D(A^{\frac{1}{2}})$ satisfying $\|A^{\frac{1}{2}}u\|_H \leq r$ and $\|A^{\frac{1}{2}}v\|_H \leq r$.

(H3) $\psi(0) = 0$, $\nabla\psi(0) = 0$ and there exists a strictly increasing continuous function h such that

$$\|\nabla\psi(u)\|_H \leq h(\|A^{\frac{1}{2}}u\|_H) \|A^{\frac{1}{2}}u\|_H, \quad (1.3)$$

for all $u \in D(A^{\frac{1}{2}})$.

We are interested in studying well-posedness and stability results, for small initial data, for the above model. Our results extend the ones of [16, 17] where abstract evolution equations are analyzed and, in the specific case of memory damping, exponential decay is obtained essentially only in the linear case. Indeed, in the nonlinear setting, an extra standard frictional damping, not delayed, was needed in order to obtain existence and uniqueness of global solutions with exponentially decaying energy for suitably small initial data. Moreover, in [16, 17] the delay damping coefficient $k(t)$ is assumed to be constant and the results there obtained require a smallness assumption on $\|k\|_\infty$. The analysis of [16, 17] has been extended in [13] by considering a time variable delay damping coefficient $k(t)$ as in the present paper. However, also in [13] an extra frictional not delayed damping was needed, in the case of wave type equation with memory damping, when a locally Lipschitz continuous nonlinear term is included into the equation.

Then, here, we focus on wave type equations with viscoelastic damping, delay feedback and source term, obtaining well-posedness and stability results for small initial data without adding extra frictional not delayed damping. So, we here improve and conclude the analysis developed in [16, 17, 13] for the class of models at hand. Other models with viscoelastic damping and time delay are studied in recent literature. The first result is due to [12], in the linear setting. In that paper a standard frictional damping, not delayed, is included into the model in order to compensate the destabilizing effect of the delay feedback. Actually, at least in the linear case, the viscoelastic damping alone can counter the destabilizing delay effect, under suitable assumptions, without needing other dampings. This has been shown, e.g., in [11, 3, 9, 23]. The case of viscoelastic wave equation with intermittent delay feedback has been studied in [20]

while [14] deals with a model for plate equation with memory, source term, delay feedback and standard not delayed frictional damping.

More extended is the literature in case of a frictional/structural damping, instead of a viscoelastic term, which compensates the destabilizing effect of a time delay and, for specific models, mainly in the linear setting, several stability results have been quite recently obtained under appropriate assumptions (see e.g. [1, 4, 5, 7, 15, 18, 21, 22]).

The paper is organized as follows. In Section 2 we give some preliminaries, writing system (1.1) in an abstract way. In Section 3 we prove the exponential decay of the energy associated to (1.1). Finally, in Section 4 some examples are illustrated.

2 Preliminaries

As in Dafermos [8], we define the function

$$\eta^t(s) := u(t) - u(t-s), \quad s, t \in (0, +\infty), \quad (2.4)$$

so that we can rewrite (1.1) in the following way:

$$\begin{aligned} u_{tt}(t) + (1 - \tilde{\mu})Au(t) + \int_0^{+\infty} \mu(s)A\eta^t(s)ds + k(t)BB^*u_t(t-\tau) \\ = \nabla\psi(u(t)), \quad t \in (0, +\infty), \\ \eta_t^t(s) = -\eta_s^t(s) + u_t(t), \quad t, s \in (0, +\infty), \\ u(0) = u_0(0), \\ u_t(0) = u_1, \\ B^*u_t(t) = g(t), \quad t \in (-\tau, 0), \\ \eta^0(s) = \eta_0(s) = u_0(0) - u_0(-s) \quad s \in (0, +\infty). \end{aligned} \quad (2.5)$$

Let $L_\mu^2((0, +\infty); D(A^{\frac{1}{2}}))$ be the Hilbert space of the $D(A^{\frac{1}{2}})$ -valued functions in $(0, +\infty)$ endowed with the scalar product

$$\langle \varphi, \psi \rangle_{L_\mu^2((0, +\infty); D(A^{\frac{1}{2}}))} = \int_0^{+\infty} \mu(s) \langle A^{\frac{1}{2}}\varphi, A^{\frac{1}{2}}\psi \rangle_H ds$$

and denote by \mathcal{H} the Hilbert space

$$\mathcal{H} = D(A^{\frac{1}{2}}) \times H \times L_\mu^2((0, +\infty); D(A^{\frac{1}{2}})),$$

equipped with the inner product

$$\left\langle \begin{pmatrix} u \\ v \\ w \end{pmatrix}, \begin{pmatrix} \tilde{u} \\ \tilde{v} \\ \tilde{w} \end{pmatrix} \right\rangle_{\mathcal{H}} := (1 - \tilde{\mu}) \langle A^{\frac{1}{2}}u, A^{\frac{1}{2}}\tilde{u} \rangle_H + \langle v, \tilde{v} \rangle_H + \int_0^{+\infty} \mu(s) \langle A^{\frac{1}{2}}w, A^{\frac{1}{2}}\tilde{w} \rangle_H ds. \quad (2.6)$$

Setting $U = (u, u_t, \eta^t)$ we can restate (1.1) in the abstract form

$$\begin{aligned} U'(t) &= \mathcal{A}U(t) - k(t)\mathcal{B}U(t-\tau) + F(U(t)), \\ \mathcal{B}U(t-\tau) &= \tilde{g}(t) \quad \text{for } t \in [0, \tau], \\ U(0) &= U_0, \end{aligned} \quad (2.7)$$

where the operator \mathcal{A} is defined by

$$\mathcal{A} \begin{pmatrix} u \\ v \\ w \end{pmatrix} = \begin{pmatrix} v \\ -(1 - \tilde{\mu})Au - \int_0^{+\infty} \mu(s)Aw(s)ds \\ -w_s + v \end{pmatrix}$$

with domain

$$D(\mathcal{A}) = \{(u, v, w) \in D(A^{\frac{1}{2}}) \times D(A^{\frac{1}{2}}) \times L^2_\mu((0, +\infty); D(A^{\frac{1}{2}})) : (1 - \tilde{\mu})u + \int_0^{+\infty} \mu(s)w(s)ds \in D(A), w_s \in L^2_\mu((0, +\infty); D(A^{\frac{1}{2}}))\}, \quad (2.8)$$

in the Hilbert space \mathcal{H} , and the operator $\mathcal{B} : \mathcal{H} \rightarrow \mathcal{H}$ is defined by

$$\mathcal{B} \begin{pmatrix} u \\ v \\ w \end{pmatrix} := \begin{pmatrix} 0 \\ BB^*v \\ 0 \end{pmatrix}.$$

Moreover, $\tilde{g}(t) = (0, Bg(t - \tau), 0)$, $U_0 = (u_0(0), u_1, \eta_0)$ and $F(U) := (0, \nabla\psi(u), 0)^T$. From (H2) and (H3) we deduce that the function F satisfies:

(F1) $F(0) = 0$;

(F2) for each $r > 0$ there exists a constant $L(r) > 0$ such that

$$\|F(U) - F(V)\|_{\mathcal{H}} \leq L(r)\|U - V\|_{\mathcal{H}} \quad (2.9)$$

whenever $\|U\|_{\mathcal{H}} \leq r$ and $\|V\|_{\mathcal{H}} \leq r$.

3 Stability result

In this section we want to prove an exponential stability result for the system (1.1) for *small* initial data. It's well-known (see e.g. [10]) that the operator \mathcal{A} in the problem's formulation (2.7) generates an exponentially stable semigroup $\{S(t)\}_{t \geq 0}$, namely there exist two constants $M, \omega > 0$ such that

$$\|S(t)\|_{\mathcal{L}(H)} \leq Me^{-\omega t}. \quad (3.10)$$

Denoting

$$\|B\|_{\mathcal{L}(H)} = \|B^*\|_{\mathcal{L}(H)} = b, \quad (3.11)$$

then $\|\mathcal{B}\|_{\mathcal{L}(\mathcal{H})} = b^2$. Our result will be obtained under an assumption on the coefficient $k(t)$ of the delay feedback which includes as particular cases k integrable and k in L^∞ with $\|k\|_\infty$ sufficiently small. More precisely, we assume (cf. [13]) that there exist two constants $\omega' \in [0, \omega)$ and $\gamma \in \mathbb{R}$ such that

$$b^2 Me^{\omega\tau} \int_0^t |k(s + \tau)|ds \leq \gamma + \omega' t, \quad \text{for all } t \geq 0. \quad (3.12)$$

Theorem 3.1. *Assume (3.12). Moreover, suppose that*

(W) there exist $\rho > 0$, $C_\rho > 0$, with $L(C_\rho) < \frac{\omega - \omega'}{M}$ such that if $U_0 \in \mathcal{H}$ and if $\tilde{g} \in C([0, \tau]; \mathcal{H})$ satisfy

$$\|U_0\|_{\mathcal{H}}^2 + \int_0^\tau |k(s)| \cdot \|\tilde{g}(s)\|_{\mathcal{H}}^2 ds < \rho^2, \quad (3.13)$$

then the system (2.7) has a unique solution $U \in C([0, +\infty); \mathcal{H})$ satisfying $\|U(t)\|_{\mathcal{H}} \leq C_\rho$ for all $t > 0$.

Then, for every solution U of (2.7), with initial datum U_0 satisfying (3.13),

$$\|U(t)\|_{\mathcal{H}} \leq \tilde{M} \left(\|U_0\|_{\mathcal{H}} + \int_0^\tau e^{\omega s} |k(s)| \cdot \|\tilde{g}(s)\|_{\mathcal{H}} ds \right) e^{-(\omega - \omega' - ML(C_\rho))t}, \quad t \geq 0, \quad (3.14)$$

with $\tilde{M} = Me^\gamma$.

Proof. By Duhamel's formula, using (3.10), we have

$$\begin{aligned} \|U(t)\|_{\mathcal{H}} &\leq Me^{-\omega t} \|U_0\|_{\mathcal{H}} + Me^{-\omega t} \int_0^t e^{\omega s} |k(s)| \cdot \|\mathcal{B}U(s - \tau)\|_{\mathcal{H}} ds + ML(C_\rho) e^{-\omega t} \int_0^t e^{\omega s} \|U(s)\|_{\mathcal{H}} ds \\ &\leq Me^{-\omega t} \|U_0\|_{\mathcal{H}} + Me^{-\omega t} \int_0^{\tau} e^{\omega s} |k(s)| \cdot \|\mathcal{B}U(s - \tau)\|_{\mathcal{H}} ds \\ &\quad + Me^{-\omega t} \int_\tau^t e^{\omega s} |k(s)| \cdot \|\mathcal{B}U(s - \tau)\|_{\mathcal{H}} ds + ML(C_\rho) e^{-\omega t} \int_0^t e^{\omega s} \|U(s)\|_{\mathcal{H}} ds. \end{aligned}$$

Hence, we obtain

$$\begin{aligned} e^{\omega t} \|U(t)\|_{\mathcal{H}} &\leq M \|U_0\|_{\mathcal{H}} + M \int_0^\tau e^{\omega s} |k(s)| \cdot \|\mathcal{B}U(s - \tau)\|_{\mathcal{H}} ds \\ &\quad + \int_0^t \left(Me^{\omega \tau} |k(s + \tau)| \cdot \|\mathcal{B}\|_{\mathcal{L}(H)} + ML(C_\rho) \right) e^{\omega s} \|U(s)\|_{\mathcal{H}} ds, \end{aligned}$$

and then

$$\begin{aligned} e^{\omega t} \|U(t)\|_{\mathcal{H}} &\leq M \|U_0\|_{\mathcal{H}} + M \int_0^\tau e^{\omega s} |k(s)| \cdot \|\tilde{g}(s)\|_{\mathcal{H}} ds \\ &\quad + \int_0^t \left(Mb^2 e^{\omega \tau} |k(s + \tau)| + ML(C_\rho) \right) e^{\omega s} \|U(s)\|_{\mathcal{H}} ds. \end{aligned}$$

Therefore, using Gronwall's inequality,

$$e^{\omega t} \|U(t)\|_{\mathcal{H}} \leq M \left(\|U_0\|_{\mathcal{H}} + \int_0^\tau e^{\omega s} |k(s)| \cdot \|\tilde{g}(s)\|_{\mathcal{H}} ds \right) e^{Mb^2 e^{\omega \tau} \int_0^t |k(s + \tau)| ds + ML(C_\rho)t}$$

and so, from (3.12),

$$\|U(t)\|_{\mathcal{H}} \leq Me^\gamma \left(\|U_0\|_{\mathcal{H}} + \int_0^\tau e^{\omega s} |k(s)| \cdot \|\tilde{g}(s)\|_{\mathcal{H}} ds \right) e^{-(\omega - \omega' - ML(C_\rho))t}.$$

This gives (3.14) with \tilde{M} as in the statement. ■

In order to prove the stability result we need then to show that the well-posedness assumption (W) of Theorem 3.1 is satisfied for problem (1.1). For this, let us define the energy of the model (1.1) as

$$E(t) := E(u(t)) = \frac{1}{2} \|u_t(t)\|_H^2 + \frac{1-\tilde{\mu}}{2} \|A^{\frac{1}{2}}u(t)\|_H^2 - \psi(u) + \frac{1}{2} \int_{t-\tau}^t |k(s+\tau)| \cdot \|B^*u_t(s)\|_H^2 ds + \frac{1}{2} \int_0^{+\infty} \mu(s) \|A^{\frac{1}{2}}\eta^t(s)\|_H^2 ds. \quad (3.15)$$

The following lemma holds.

Lemma 3.2. *Let $u : [0, T) \rightarrow \mathbb{R}$ be a solution of (1.1). Assume that $E(t) \geq \frac{1}{4} \|u_t(t)\|_H^2$ for all $t \in [0, T)$. Then,*

$$E(t) \leq \bar{C}(t)E(0), \quad (3.16)$$

for all $t \in [0, T)$, where

$$\bar{C}(t) = e^{2b^2 \int_0^t (|k(s)| + |k(s+\tau)|) ds}. \quad (3.17)$$

Proof. Differentiating $E(t)$, we obtain

$$\begin{aligned} \frac{dE}{dt} &= \langle u_t, u_{tt} \rangle_H + (1-\tilde{\mu}) \langle A^{\frac{1}{2}}u, A^{\frac{1}{2}}u_t \rangle_H - \langle \nabla \psi(u), u_t \rangle_H + \frac{1}{2} |k(t+\tau)| \cdot \|B^*u_t(t)\|_H^2 \\ &\quad - \frac{1}{2} |k(t)| \cdot \|B^*u_t(t-\tau)\|_H^2 + \int_0^{+\infty} \mu(s) \langle A^{\frac{1}{2}}\eta^t(s), A^{\frac{1}{2}}\eta_t^t(s) \rangle_H ds. \end{aligned}$$

Then, from (1.1),

$$\begin{aligned} \frac{dE}{dt} &= - \int_0^{+\infty} \mu(s) \langle u_t(t), A\eta^t(s) \rangle_H ds - k(t) \langle u_t, BB^*u_t(t-\tau) \rangle_H + \frac{1}{2} |k(t+\tau)| \cdot \|B^*u_t(t)\|_H^2 \\ &\quad - \frac{1}{2} |k(t)| \cdot \|B^*u_t(t-\tau)\|_H^2 + \int_0^{+\infty} \mu(s) \langle A\eta^t(s), \eta_t^t(s) \rangle_H ds. \end{aligned}$$

Using the second equation of (2.5), we have that

$$\begin{aligned} \frac{dE}{dt} &= -k(t) \langle u_t, BB^*u_t(t-\tau) \rangle_H + \frac{1}{2} |k(t+\tau)| \cdot \|B^*u_t\|_H^2 - \frac{1}{2} |k(t)| \cdot \|B^*u_t(t-\tau)\|_H^2 \\ &\quad - \int_0^{+\infty} \mu(s) \langle A\eta^t(s), \eta_s^t(s) \rangle_H ds. \end{aligned}$$

Now, we claim that

$$\int_0^{+\infty} \mu(s) \langle \eta_s^t, A\eta^t(s) \rangle_H ds \geq 0.$$

Indeed, integrating by parts and recalling assumption (iv) on $\mu(\cdot)$, we deduce

$$\int_0^{+\infty} \mu(s) \langle \eta_s^t, A\eta^t(s) \rangle_H ds = -\frac{1}{2} \int_0^{+\infty} \mu'(s) \|A^{\frac{1}{2}}\eta^t(s)\|_H^2 ds \geq 0.$$

Therefore, we have that

$$\frac{dE(t)}{dt} \leq -k(t) \langle B^*u_t, B^*u_t(t-\tau) \rangle_H + \frac{1}{2} |k(t+\tau)| \cdot \|B^*u_t(t)\|_H^2 - \frac{1}{2} |k(t)| \cdot \|B^*u_t(t-\tau)\|_H^2.$$

Now, using Cauchy-Schwarz inequality, we obtain the following estimate:

$$\begin{aligned}
\frac{dE(t)}{dt} &\leq \frac{1}{2}(|k(t)| + |k(t+\tau)|) \|B^*u_t(t)\|_H^2 \\
&\leq \frac{1}{2}(|k(t)| + |k(t+\tau)|) b^2 \|u_t\|_H^2 \\
&= 2b^2(|k(t)| + |k(t+\tau)|) \frac{1}{4} \|u_t\|_H^2 \\
&\leq 2b^2(|k(t)| + |k(t+\tau)|) E(t).
\end{aligned}$$

Hence, the Gronwall Lemma concludes the proof. ■

Before proving the well-posedness assumption (W) for solutions to (2.7), we need the following two lemmas.

Lemma 3.3. *Let us consider the system (2.7) with initial data $U_0 \in \mathcal{H}$ and $\tilde{g} \in C([0, \tau]; \mathcal{H})$. Then, there exists a unique local solution $U(\cdot)$ defined on a time interval $[0, \delta)$, with $\delta \leq \tau$.*

Proof. Since $t \in [0, \tau]$, we can rewrite the abstract system (2.7) as an undelayed problem:

$$\begin{aligned}
U'(t) &= \mathcal{A}U(t) - k(t)\tilde{g}(t) + F(U(t)), \quad t \in (0, \tau), \\
U(0) &= U_0.
\end{aligned}$$

Then, we can apply the classical theory of nonlinear semigroups (see e.g. [19]) obtaining the existence of a unique solution on a set $[0, \delta)$, with $\delta \leq \tau$. ■

Lemma 3.4. *Let $U(t) = (u(t), u_t(t), \eta^t)$ be a solution to (2.7) defined on the interval $[0, \delta)$, with $\delta \leq \tau$. Then,*

1. *if $h(\|A^{\frac{1}{2}}u_0(0)\|_H) < \frac{1-\tilde{\mu}}{2}$, then $E(0) > 0$;*
2. *if $h(\|A^{\frac{1}{2}}u_0(0)\|_H) < \frac{1-\tilde{\mu}}{2}$ and $h\left(\frac{2}{(1-\tilde{\mu})^{\frac{1}{2}}}\bar{C}^{\frac{1}{2}}(\tau)E^{\frac{1}{2}}(0)\right) < \frac{1-\tilde{\mu}}{2}$, with $\bar{C}(\tau)$ defined in (3.17), then*

$$\begin{aligned}
E(t) &> \frac{1}{4}\|u_t\|_H^2 + \frac{1-\tilde{\mu}}{4}\|A^{\frac{1}{2}}u\|_H^2 \\
&\quad + \frac{1}{4}\int_{t-\tau}^t |k(s+\tau)| \cdot \|B^*u_t(s)\|_H^2 ds + \frac{1}{4}\int_0^{+\infty} \mu(s)\|A^{\frac{1}{2}}\eta^t(s)\|_H^2 ds,
\end{aligned} \tag{3.18}$$

for all $t \in [0, \delta)$. In particular,

$$E(t) > \frac{1}{4}\|U(t)\|_{\mathcal{H}}^2, \quad \text{for all } t \in [0, \delta). \tag{3.19}$$

Proof. We first deduce by assumption (H3) on ψ that

$$\begin{aligned}
|\psi(u)| &\leq \int_0^1 |\langle \nabla \psi(su), u \rangle| ds \\
&\leq \|A^{\frac{1}{2}}u\|_H^2 \int_0^1 h(s\|A^{\frac{1}{2}}u\|_H) s ds \leq \frac{1}{2}h(\|A^{\frac{1}{2}}u\|_H)\|A^{\frac{1}{2}}u\|_H^2.
\end{aligned} \tag{3.20}$$

Hence, under the assumption $h(\|A^{\frac{1}{2}}u_0(0)\|_H) < \frac{1-\tilde{\mu}}{2}$, we have that

$$\begin{aligned} E(0) &= \frac{1}{2}\|u_1\|_H^2 + \frac{1-\tilde{\mu}}{2}\|A^{\frac{1}{2}}u_0(0)\|_H^2 - \psi(u_0(0)) + \frac{1}{2}\int_{-\tau}^0 |k(s+\tau)| \cdot \|B^*u_t(s)\|_H^2 ds \\ &\quad + \frac{1}{2}\int_0^{+\infty} \mu(s)\|A^{\frac{1}{2}}\eta_0(s)\|_H^2 ds \\ &\geq \frac{1}{2}\|u_1\|_H^2 + \frac{1-\tilde{\mu}}{4}\|A^{\frac{1}{2}}u_0(0)\|_H^2 + \frac{1}{2}\int_{-\tau}^0 |k(s+\tau)| \cdot \|B^*u_t(s)\|_H^2 ds \\ &\quad + \frac{1}{2}\int_0^{+\infty} \mu(s)\|A^{\frac{1}{2}}\eta_0(s)\|_H^2 ds > 0, \end{aligned}$$

obtaining 1.

In order to prove the second statement, we argue by contradiction. Let us denote

$$r := \sup\{s \in [0, \delta) : (3.18) \text{ holds } \forall t \in [0, s)\}.$$

We suppose by contradiction that $r < \delta$. Then, by continuity, we have

$$\begin{aligned} E(r) &= \frac{1}{4}\|u_t(r)\|_H^2 + \frac{1-\tilde{\mu}}{4}\|A^{\frac{1}{2}}u(r)\|_H^2 + \frac{1}{4}\int_{r-\tau}^r |k(s+\tau)| \cdot \|B^*u_t(s)\|_H^2 ds \\ &\quad + \frac{1}{4}\int_0^{+\infty} \mu(s)\|A^{\frac{1}{2}}\eta^r(s)\|_H^2 ds. \end{aligned} \tag{3.21}$$

Now, from (3.21) and Lemma 3.2 we can infer that

$$\begin{aligned} h(\|A^{\frac{1}{2}}u(r)\|_H) &\leq h\left(\frac{2}{(1-\tilde{\mu})^{\frac{1}{2}}}E^{\frac{1}{2}}(r)\right) \\ &< h\left(\frac{2}{(1-\tilde{\mu})^{\frac{1}{2}}}\bar{C}^{\frac{1}{2}}(\tau)E^{\frac{1}{2}}(0)\right) < \frac{1-\tilde{\mu}}{2}. \end{aligned}$$

Hence, we have that

$$\begin{aligned} E(r) &= \frac{1}{2}\|u_t(r)\|_H^2 + \frac{1-\tilde{\mu}}{2}\|A^{\frac{1}{2}}u(r)\|_H^2 - \psi(u(r)) + \frac{1}{2}\int_{r-\tau}^r |k(s+\tau)| \cdot \|B^*u_t(s)\|_H^2 ds \\ &\quad + \frac{1}{2}\int_0^{+\infty} \mu(s)\|A^{\frac{1}{2}}\eta^r(s)\|_H^2 ds \\ &> \frac{1}{4}\|u_t(r)\|_H^2 + \frac{1-\tilde{\mu}}{4}\|A^{\frac{1}{2}}u(r)\|_H^2 + \frac{1}{4}\int_{r-\tau}^r |k(s+\tau)| \cdot \|B^*u_t(s)\|_H^2 ds \\ &\quad + \frac{1}{4}\int_0^{+\infty} \mu(s)\|A^{\frac{1}{2}}\eta^r(s)\|_H^2 ds, \end{aligned}$$

which contradicts the maximality of r . Hence, $r = \delta$ and this concludes the proof. \blacksquare

Theorem 3.5. *Problem (2.7), with initial data $U_0 \in \mathcal{H}$ and $\tilde{g} \in C([0, \tau]; \mathcal{H})$, satisfies the well-posedness assumption (W). Then, for solutions of (2.7) corresponding to sufficiently small initial data the exponential decay estimate (3.14) holds.*

Proof. Let us fix $N \in \mathbb{N}$ such that

$$2M^2\left(1 + e^{2\omega\tau}C^*\right)e^{2\gamma}e^{-(\omega-\omega')(N-1)\tau} < \frac{1}{1 + e^{\omega\tau}b^2C^*}, \tag{3.22}$$

where C^* is the constant defined in (1.2). Then, let ρ be a positive constant such that

$$\rho \leq \frac{(1 - \tilde{\mu})^{\frac{1}{2}}}{2\bar{C}^{\frac{1}{2}}(N\tau)} h^{-1} \left(\frac{1 - \tilde{\mu}}{2} \right). \quad (3.23)$$

Now, let us assume that the initial data $(u_0(0), u_1, \eta_0)$ and $B^*u_t(s)$, $s \in [-\tau, 0]$, satisfy the smallness assumption

$$\begin{aligned} (1 - \tilde{\mu}) \|A^{\frac{1}{2}}u_0(0)\|_H^2 + \|u_1\|_H^2 + \int_{-\tau}^0 |k(s + \tau)| \|B^*u_t(s)\|_H^2 ds \\ + \int_0^{+\infty} \mu(s) \|A^{\frac{1}{2}}\eta_0(s)\|_H^2 ds < \rho^2. \end{aligned} \quad (3.24)$$

Note that (3.24) is equivalent to

$$\|U_0\|_{\mathcal{H}}^2 + \int_0^\tau |k(s)| \cdot \|\tilde{g}(s)\|_{\mathcal{H}}^2 ds < \rho^2. \quad (3.25)$$

From Lemma 3.3 we know that there exists a local solution defined on a time interval $[0, \delta)$, with $\delta \leq \tau$. From (3.24) and (3.23) we have that

$$h(\|A^{\frac{1}{2}}u_0(0)\|_H) < h\left(\frac{\rho}{(1 - \tilde{\mu})^{\frac{1}{2}}}\right) \leq h\left(\frac{1}{2\bar{C}^{\frac{1}{2}}(N\tau)} h^{-1}\left(\frac{1 - \tilde{\mu}}{2}\right)\right) < \frac{1 - \tilde{\mu}}{2}, \quad (3.26)$$

where we have used the fact that $\bar{C}(N\tau) > 1$. This implies, from Lemma 3.4, $E(0) > 0$. Furthermore, from (3.20) and (3.26) we get

$$\begin{aligned} E(0) &\leq \frac{1}{2} \|u_1\|_H^2 + \frac{3}{4} (1 - \tilde{\mu}) \|A^{\frac{1}{2}}u_0(0)\|_H^2 + \frac{1}{2} \int_{-\tau}^0 |k(s + \tau)| \cdot \|B^*u_t(s)\|_H^2 ds \\ &\quad + \frac{1}{2} \int_0^{+\infty} \mu(s) \|A^{\frac{1}{2}}\eta_0(s)\|_H^2 ds < \rho^2, \end{aligned}$$

which gives, recalling (3.23),

$$h\left(\frac{2}{(1 - \tilde{\mu})^{\frac{1}{2}}} \bar{C}^{\frac{1}{2}}(N\tau) E^{\frac{1}{2}}(0)\right) < h\left(\frac{2}{(1 - \tilde{\mu})^{\frac{1}{2}}} \bar{C}^{\frac{1}{2}}(N\tau) \rho\right) \leq h\left(h^{-1}\left(\frac{1 - \tilde{\mu}}{2}\right)\right) = \frac{1 - \tilde{\mu}}{2}. \quad (3.27)$$

Since $\bar{C}(N\tau) \geq \bar{C}(\tau)$, then

$$h\left(\frac{2}{(1 - \tilde{\mu})^{\frac{1}{2}}} \bar{C}^{\frac{1}{2}}(\tau) E^{\frac{1}{2}}(0)\right) \leq h\left(\frac{2}{(1 - \tilde{\mu})^{\frac{1}{2}}} \bar{C}^{\frac{1}{2}}(N\tau) E^{\frac{1}{2}}(0)\right) < \frac{1 - \tilde{\mu}}{2}. \quad (3.28)$$

So, we can apply Lemma 3.4 and we obtain

$$\begin{aligned} E(t) &> \frac{1}{4} \|u_t(t)\|_H^2 + \frac{1 - \tilde{\mu}}{4} \|A^{\frac{1}{2}}u(t)\|_H^2 \\ &\quad + \frac{1}{4} \int_{t-\tau}^t |k(s + \tau)| \cdot \|B^*u_t(s)\|_H^2 ds + \frac{1}{4} \int_0^{+\infty} \mu(s) \|A^{\frac{1}{2}}\eta^t(s)\|_H^2 ds, \end{aligned}$$

for all $t \in [0, \delta)$. In particular we have that

$$E(t) > \frac{1}{4} \|u_t(t)\|_H^2, \quad \text{for } t \in [0, \delta).$$

Therefore, we can apply Lemma 3.2, obtaining

$$E(t) \leq \bar{C}(\tau)E(0) < \bar{C}(\tau)\rho^2,$$

for any $t \in [0, \delta]$. Since

$$\begin{aligned} 0 &< \frac{1}{4} \|u_t(t)\|_H^2 + \frac{1-\tilde{\mu}}{4} \|A^{\frac{1}{2}}u(t)\|_H^2 \\ &+ \frac{1}{4} \int_{t-\tau}^t |k(s+\tau)| \cdot \|B^*u_t(s)\|_H^2 ds + \frac{1}{4} \int_0^{+\infty} \mu(s) \|A^{\frac{1}{2}}\eta^t(s)\|_H^2 ds \leq E(t) \leq \bar{C}(\tau)E(0), \end{aligned} \quad (3.29)$$

for all $t \in [0, \delta]$, then we can extend the solution to the entire interval $[0, \tau]$.

Now, observe that from (3.29) and (3.28) we have

$$h(\|A^{\frac{1}{2}}u(\tau)\|_H) \leq h\left(\frac{2}{(1-\tilde{\mu})^{\frac{1}{2}}} \bar{C}^{\frac{1}{2}}(\tau)E^{\frac{1}{2}}(0)\right) < \frac{1-\tilde{\mu}}{2}. \quad (3.30)$$

By continuity, (3.30) implies that there exists $\delta' > 0$ such that

$$h(\|A^{\frac{1}{2}}u(t)\|_H) < \frac{1-\tilde{\mu}}{2}, \quad \forall t \in [\tau, \tau + \delta').$$

From this, arguing as before, we deduce

$$\begin{aligned} E(t) &> \frac{1}{4} \|u_t(t)\|_H^2 + \frac{1-\tilde{\mu}}{4} \|A^{\frac{1}{2}}u(t)\|_H^2 + \frac{1}{4} \int_{t-\tau}^t |k(s+\tau)| \cdot \|B^*u_t(s)\|_H^2 ds \\ &+ \frac{1}{4} \int_0^{+\infty} \mu(s) \|A^{\frac{1}{2}}\eta^t(s)\|_H^2 ds, \end{aligned}$$

for any $t \in [\tau, \tau + \delta')$. In particular, also in such an interval we have $E(t) > \frac{1}{4} \|u_t(t)\|_H^2$. Hence, we can apply again Lemma 3.2 on the time interval $[0, \tau + \delta')$ obtaining

$$0 < E(t) \leq \bar{C}(2\tau)E(0).$$

As before, we can then extend the solution the whole interval $[0, 2\tau]$. At time $t = 2\tau$ we have that

$$h(\|A^{\frac{1}{2}}u(2\tau)\|_H) \leq h\left(\frac{2}{(1-\tilde{\mu})^{\frac{1}{2}}} E^{\frac{1}{2}}(2\tau)\right) \leq h\left(\frac{2}{(1-\tilde{\mu})^{\frac{1}{2}}} \bar{C}^{\frac{1}{2}}(2\tau)E^{\frac{1}{2}}(0)\right) < \frac{1-\tilde{\mu}}{2},$$

where we have used (3.28). Moreover, if $3 \leq N$,

$$h\left(\frac{2}{(1-\tilde{\mu})^{\frac{1}{2}}} \bar{C}^{\frac{1}{2}}(3\tau)E^{\frac{1}{2}}(0)\right) \leq h\left(\frac{2}{(1-\tilde{\mu})^{\frac{1}{2}}} \bar{C}^{\frac{1}{2}}(N\tau)E^{\frac{1}{2}}(0)\right) < \frac{1-\tilde{\mu}}{2}.$$

Thus, one can repeat again the same argument. By iteration, we then find a unique solution to the problem (2.7) on the interval $[0, N\tau]$, where N is the natural number fixed at the beginning of the proof. Moreover, the definition (3.23) of ρ , ensures that

$$h(\|A^{\frac{1}{2}}u(N\tau)\|_H) \leq h\left(\frac{2}{(1-\tilde{\mu})^{\frac{1}{2}}} E^{\frac{1}{2}}(N\tau)\right) \leq h\left(\frac{2}{(1-\tilde{\mu})^{\frac{1}{2}}} \bar{C}^{\frac{1}{2}}(N\tau)E^{\frac{1}{2}}(0)\right) < \frac{1-\tilde{\mu}}{2},$$

where we have used (3.27). Note that, by construction, (3.18) and (3.19) are satisfied in the whole $[0, N\tau]$. Then, from (3.19),

$$\|U(t)\|_{\mathcal{H}}^2 \leq 4E(t) \leq 4\bar{C}(N\tau)E(0) < 4\bar{C}(N\tau)\rho^2$$

and so

$$\|U(t)\|_{\mathcal{H}} \leq 2\bar{C}^{\frac{1}{2}}(N\tau)\rho, \quad \forall t \in [0, N\tau].$$

Thus, we have proved that, under the assumption (3.25) on the initial data, there exists a unique solution $U(\cdot)$ to problem (2.7) defined on the time interval $[0, N\tau]$. Moreover,

$$\|U(t)\|_{\mathcal{H}} \leq C_\rho := 2\bar{C}^{\frac{1}{2}}(N\tau)\rho.$$

So far we have fixed ρ satisfying (3.23); now, eventually choosing a smaller ρ , we assume that ρ satisfies the additional assumption

$$L(C_\rho) = L(\bar{C}^{\frac{1}{2}}(N\tau)\rho) < \frac{\omega - \omega'}{2M}.$$

Then, the well-posedness assumption (W) of Theorem 3.1 is satisfied on $[0, N\tau]$. Therefore, we obtain that U satisfies the exponential decay estimate (3.14) and then

$$\|U(t)\|_{\mathcal{H}} \leq M \left(\|U(0)\|_{\mathcal{H}} + \int_0^t |k(s)| e^{\omega s} \|\tilde{g}(s)\|_{\mathcal{H}} ds \right) e^{\gamma} e^{-\frac{\omega - \omega'}{2}t}, \quad \forall t \in [0, N\tau]. \quad (3.31)$$

In particular,

$$\|U(N\tau)\|_{\mathcal{H}} \leq M \left(\|U(0)\|_{\mathcal{H}} + \int_0^{N\tau} e^{\omega s} |k(s)| \cdot \|\tilde{g}(s)\|_{\mathcal{H}} ds \right) e^{\gamma} e^{-\frac{\omega - \omega'}{2}N\tau}. \quad (3.32)$$

Now, observe that

$$\int_0^{\tau} e^{\omega s} |k(s)| \cdot \|\tilde{g}(s)\|_{\mathcal{H}}^2 ds \leq e^{\omega\tau} \left(\int_0^{\tau} |k(s)| ds \right)^{1/2} \left(\int_0^{\tau} |k(s)| \cdot \|\tilde{g}(s)\|_{\mathcal{H}}^2 ds \right)^{1/2}.$$

Hence,

$$\|U(t)\|_{\mathcal{H}} \leq M\rho \left(1 + e^{\omega\tau} C^{*1/2} \right) e^{\gamma} e^{-\frac{\omega - \omega'}{2}t}, \quad \forall t \in [0, N\tau],$$

and then

$$\|U(t)\|_{\mathcal{H}}^2 \leq 2M^2\rho^2 \left(1 + e^{2\omega\tau} C^* \right) e^{2\gamma} e^{-(\omega - \omega')t}, \quad \forall t \in [0, N\tau], \quad (3.33)$$

where C^* is the constant defined in (1.2). From (3.33) we deduce

$$\begin{aligned} & \|U(N\tau)\|_{\mathcal{H}}^2 + \int_{N\tau}^{(N+1)\tau} e^{\omega(s-N\tau)} |k(s)| \cdot \|B^* u_t(s - \tau)\|_{\mathcal{H}}^2 ds \\ & \leq 2M^2\rho^2 \left(1 + e^{2\omega\tau} C^* \right) e^{2\gamma} e^{-(\omega - \omega')N\tau} + e^{\omega\tau} b^2 \int_{N\tau}^{(N+1)\tau} |k(s)| \cdot \|U(s - \tau)\|_{\mathcal{H}}^2 ds. \end{aligned} \quad (3.34)$$

Now, observe that, for $s \in [N\tau, (N+1)\tau]$, it results $s - \tau \in [(N-1)\tau, N\tau]$, then from (3.33) we deduce

$$\|U(s - \tau)\|_{\mathcal{H}}^2 \leq 2M^2\rho^2 \left(1 + e^{2\omega\tau} C^* \right) e^{2\gamma} e^{-(\omega - \omega')(N-1)\tau}, \quad \forall s \in [N\tau, (N+1)\tau].$$

This last estimate, used in (3.34), gives

$$\begin{aligned} & \|U(N\tau)\|_{\mathcal{H}}^2 + \int_{N\tau}^{(N+1)\tau} e^{\omega(s-N\tau)} |k(s)| \cdot \|B^* u_t(s-\tau)\|_H^2 ds \\ & \leq 2M^2 \rho^2 (1 + e^{2\omega\tau} C^*) e^{2\gamma} e^{-(\omega-\omega')(N-1)\tau} (1 + e^{\omega\tau} b^2 C^*). \end{aligned} \quad (3.35)$$

From (3.35) and (3.22), we then deduce

$$\|U(N\tau)\|_{\mathcal{H}}^2 + \int_{N\tau}^{(N+1)\tau} |k(s)| \cdot \|B^* u_t(s-\tau)\|_H^2 ds < \rho^2.$$

Thus (cf. with (3.25)), one can argue as before on the interval $[N\tau, 2N\tau]$ obtaining a solution on $[0, 2N\tau]$. Iterating this procedure we get a global solution satisfying

$$\|U(t)\|_{\mathcal{H}} < C_\rho.$$

Therefore, we have showed that the problem (2.7) satisfies the well-posedness assumption (W) of Theorem 3.1. ■

We have then proved that, for suitably small data, solutions to problem (2.7) are globally defined and their energies satisfy an exponential decay estimate. Therefore, one can state the following theorem.

Theorem 3.6. *Let us consider (2.5). Then, there exists $\delta > 0$ such that if*

$$(1 - \tilde{\mu}) \|A^{\frac{1}{2}} u_0\|_H^2 + \|u_1\|_H^2 + \int_0^{+\infty} \mu(s) \|A^{\frac{1}{2}} \eta_0\|_H^2 ds + \int_0^\tau |k(s)| \cdot \|Bg(s-\tau)\|_H^2 ds < \delta, \quad (3.36)$$

then the solution u is globally defined and it satisfies

$$E(t) \leq C e^{-\beta t}, \quad (3.37)$$

where C is a constant depending only on the initial data and $\beta > 0$.

Proof. By a direct application of Theorem 3.1 and Theorem 3.5 we have that there exist $K, \gamma > 0$ such that

$$\|U(t)\|_{\mathcal{H}}^2 \leq K e^{-\gamma t}, \quad (3.38)$$

for all $t \geq 0$, if the initial data are suitably small. Now, observe that there exists a constant $\overline{C} > 0$ such that

$$\int_{t-\tau}^t |k(s)| \cdot \|B^* u_t(s)\|_H^2 ds \leq \overline{C} C^* e^{-\gamma(t-\tau)}. \quad (3.39)$$

Then, from (3.38) and (3.39) we obtain (3.37). ■

4 Examples

4.1 The wave equation with memory and source term

Let Ω be a non-empty bounded set in \mathbb{R}^n , with boundary Γ of class C^2 . Moreover, let $\mathcal{O} \subset \Omega$ be a nonempty open subset of Ω . We consider the following wave equation:

$$\begin{aligned} u_{tt}(x, t) - \Delta u(x, t) + \int_0^{+\infty} \mu(s) \Delta u(x, t-s) ds + k(t) \chi_{\mathcal{O}} u_t(x, t-\tau) \\ = |u(x, t)|^\sigma u(x, t), \quad \text{in } \Omega \times (0, +\infty), \\ u(x, t) = 0, \quad \text{in } \Gamma \times (0, +\infty), \\ u(x, t) = u_0(x, t) \quad \text{in } \Omega \times (-\infty, 0], \\ u_t(x, 0) = u_1(x), \quad \text{in } \Omega, \\ u_t(x, t) = g(x, t), \quad \text{in } \Omega \times (-\tau, 0], \end{aligned} \quad (4.40)$$

where $\tau > 0$ is the time-delay, $\mu : (0, +\infty) \rightarrow (0, +\infty)$ is a locally absolutely continuous memory kernel, which satisfies the assumptions (i)-(iv), $\sigma > 0$ and the damping coefficient $k(\cdot)$ is a function in $L^1_{loc}([0, +\infty))$ satisfying (1.2). Then, system (4.40) falls in the form (1.1) with $A = -\Delta$ and $D(A) = H^2(\Omega) \cap H^1_0(\Omega)$.

Defining η_s^t as in (2.4), we can rewrite the system (4.40) in the following way:

$$\begin{aligned} u_{tt}(x, t) - (1 - \tilde{\mu}) \Delta u(x, t) - \int_0^{+\infty} \mu(s) \Delta \eta^t(x, s) ds + k(t) \chi_{\mathcal{O}} u_t(x, t-\tau) \\ = |u(x, t)|^\sigma u(x, t), \quad \text{in } \Omega \times (0, +\infty), \\ \eta_t^t(x, s) = -\eta_s^t(x, s) + u_t(x, t), \quad \text{in } \Omega \times (0, +\infty) \times (0, +\infty), \\ u(x, t) = 0, \quad \text{in } \Gamma \times (0, +\infty), \\ \eta^t(x, s) = 0, \quad \text{in } \Gamma \times (0, +\infty), \quad \text{for } t \geq 0, \\ u(x, 0) = u_0(x) := u_0(x, 0), \quad \text{in } \Omega, \\ u_t(x, 0) = u_1(x) := \frac{\partial u_0}{\partial t}(x, t) \Big|_{t=0}, \quad \text{in } \Omega, \\ \eta^0(x, s) = \eta_0(x, s) := u_0(x, 0) - u_0(x, -s), \quad \text{in } \Omega \times (0, +\infty), \\ u_t(x, t) = g(x, t), \quad \text{in } \Omega \times (-\tau, 0). \end{aligned} \quad (4.41)$$

As before, we introduce the Hilbert space $L^2_\mu((0, +\infty); H^1_0(\Omega))$ endowed with the inner product

$$\langle \phi, \psi \rangle_{L^2_\mu((0, +\infty); H^1_0(\Omega))} := \int_\Omega \left(\int_0^{+\infty} \mu(s) \nabla \phi(x, s) \nabla \psi(x, s) dx \right) ds,$$

and consider the Hilbert space

$$\mathcal{H} = H^1_0(\Omega) \times L^2(\Omega) \times L^2_\mu((0, +\infty); H^1_0(\Omega)),$$

equipped with the inner product

$$\left\langle \begin{pmatrix} u \\ v \\ w \end{pmatrix}, \begin{pmatrix} \tilde{u} \\ \tilde{v} \\ \tilde{w} \end{pmatrix} \right\rangle_{\mathcal{H}} := (1 - \tilde{\mu}) \int_\Omega \nabla u \nabla \tilde{u} dx + \int_\Omega v \tilde{v} dx + \int_\Omega \int_0^{+\infty} \mu(s) \nabla w \nabla \tilde{w} ds dx.$$

Setting $U = (u, u_t, \eta^t)$, we can rewrite (4.43) in the form (2.7), where

$$\mathcal{A} \begin{pmatrix} u \\ v \\ w \end{pmatrix} = \begin{pmatrix} u \\ (1 - \tilde{\mu}) \Delta u + \int_0^{+\infty} \mu(s) \Delta w(s) ds \\ -w_s + v \end{pmatrix},$$

with domain

$$D(A) = \{(u, v, w) \in H_0^1(\Omega) \times H_0^1(\Omega) \times L_\mu^2((0, +\infty); H_0^1(\Omega)) : \\ (1 - \tilde{\mu})u + \int_0^{+\infty} \mu(s)w(s)ds \in H^2(\Omega) \cap H_0^1(\Omega), w_s \in L_\mu^2((0, +\infty); H_0^1(\Omega))\},$$

$\mathcal{B}(u, v, \eta^t)^T := (0, \chi_{\mathcal{O}}v, 0)^T$ and $F(U(t)) = (0, |u(t)|^\sigma u(t), 0)^T$. For any $u \in H_0^1(\Omega)$ consider the functional

$$\psi(u) := \frac{1}{\sigma + 2} \int_\Omega |u(x)|^{\sigma+2} dx.$$

By Sobolev's embedding theorem, we know that if $0 < \sigma < \frac{4}{n-2}$, then ψ is well-defined, and Gâteaux differentiable at any point $u \in H_0^1(\Omega)$, with Gâteaux derivative given by

$$D\psi(u)(v) = \int_\Omega |u(x)|^\sigma u(x)v(x)dx,$$

for any $v \in H_0^1(\Omega)$. Moreover, as in [2], if $0 < \sigma \leq \frac{2}{n-2}$, then ψ satisfies the assumptions (H1), (H2), (H3). Define the energy as follows:

$$E(t) := \frac{1}{2} \int_\Omega |u_t(x, t)|^2 dx + \frac{1 - \tilde{\mu}}{2} \int_\Omega |\nabla u(x, t)|^2 dx - \psi(u(x, t)) \\ + \frac{1}{2} \int_{t-\tau}^t \int_{\mathcal{O}} |k(s + \tau)| \cdot |u_t(x, s)|^2 dx ds + \frac{1}{2} \int_0^{+\infty} \mu(s) \int_\Omega |\nabla \eta^t(x, s)|^2 dx ds.$$

Theorem 3.1 applies to this model giving well-posedness and exponential decay of the energy for suitably small initial data, provided that the condition (3.12) on the time delay holds for every $t \geq 0$.

4.2 The plate system with memory and source term

Let Ω be a non-empty bounded set in \mathbb{R}^n , with boundary Γ of class C^2 . Let us denote $\nu(x)$ the outward unit normal vector at any point $x \in \Gamma$. Moreover, let $\mathcal{O} \subset \Omega$ be a nonempty open subset of Ω . We consider the following viscoelastic plate system:

$$u_{tt}(x, t) + \Delta^2 u(x, t) - \int_0^{+\infty} \mu(s) \Delta^2 u(x, t - s) ds + k(t) \chi_{\mathcal{O}} u_t(x, t - \tau) \\ = |u(x, t)|^\sigma u(x, t), \quad \text{in } \Omega \times (0, +\infty), \\ u(x, t) = \frac{\partial u}{\partial \nu}(x, t) = 0, \quad \text{in } \Gamma \times (0, +\infty), \\ u(x, t) = u_0(x, t) \quad \text{in } \Omega \times (-\infty, 0], \\ u_t(x, 0) = u_1(x), \quad \text{in } \Omega, \\ u_t(x, t) = g(x, t), \quad \text{in } \Omega \times (-\tau, 0], \tag{4.42}$$

where $\tau > 0$ is the time-delay, $\mu : (0, +\infty) \rightarrow (0, +\infty)$ is a locally absolutely continuous memory kernel, which satisfies the assumptions (i)-(iv), $\sigma > 0$ and the damping coefficient $k(\cdot)$ is a function in $L_{loc}^1([0, +\infty))$ satisfying (1.2). This system again falls in (1.1) for $A = \Delta^2$ with domain $D(A) = H^4(\Omega) \cap H_0^1(\Omega)$. Defining η_s^t as in (2.4), we can rewrite the system (4.42) in the

following way:

$$\begin{aligned}
& u_{tt}(x, t) + (1 - \tilde{\mu})\Delta^2 u(x, t) + \int_0^{+\infty} \mu(s)\Delta^2 \eta^t(x, s)ds + k(t)\chi_{\mathcal{O}}u_t(x, t - \tau) \\
& \quad = |u(x, t)|^\sigma u(x, t), \quad \text{in } \Omega \times (0, +\infty), \\
& \eta_t^t(x, s) = -\eta_s^t(x, s) + u_t(x, t), \quad \text{in } \Omega \times (0, +\infty) \times (0, +\infty), \\
& u(x, t) = \frac{\partial u}{\partial \nu}(x, t) = 0, \quad \text{in } \Gamma \times (0, +\infty), \\
& \eta^t(x, s) = 0, \quad \text{in } \Gamma \times (0, +\infty), \quad \text{for } t \geq 0, \\
& u(x, 0) = u_0(x) := u_0(x, 0), \quad \text{in } \Omega, \\
& u_t(x, 0) = u_1(x) := \frac{\partial u_0}{\partial t}(x, t) \Big|_{t=0}, \quad \text{in } \Omega, \\
& \eta^0(x, s) = \eta_0(x, s) := u_0(x, 0) - u_0(x, -s), \quad \text{in } \Omega \times (0, +\infty), \\
& u_t(x, t) = g(x, t), \quad \text{in } \Omega \times (-\tau, 0).
\end{aligned} \tag{4.43}$$

Then, arguing analogously to the previous example, one can recast (4.43) in the form (2.7). Moreover, for $(n-4)\sigma \leq 4$ (cf. e.g. [14]) the nonlinear source satisfies the required assumptions.

Theorem 3.1 applies then to this model giving well-posedness and exponential decay of the energy for suitably small initial data, provided that the condition (3.12) on the time delay holds for every $t \geq 0$.

References

- [1] E.M. AIT BENHASSI, K. AMMARI, S. BOULITE AND L. MANIAR. Feedback stabilization of a class of evolution equations with delay. *J. Evol. Equ.*, 9:103–121, 2009.
- [2] F. ALABAU-BOUSSOUIRA, P. CANNARSA AND D. SFORZA. Decay estimates for second order evolution equations with memory. *J. Funct. Anal.*, 254:1342–1372, 2008.
- [3] F. ALABAU-BOUSSOUIRA, S. NICAISE AND C. PIGNOTTI. Exponential stability of the wave equation with memory and time delay. *New prospects in direct, inverse and control problems for evolution equations*, 1–22, Springer INdAM Ser., 10:1–22, Springer, Cham, 2014.
- [4] K. AMMARI AND S. GERBI. Interior feedback stabilization of wave equations with dynamic boundary delay. *Z. Anal. Anwend.*, 36:297–327, 2017.
- [5] T.A. APALARA AND S.A. MESSAOUDI. An exponential stability result of a Timoshenko system with thermoelasticity with second sound and in the presence of delay. *Appl. Math. Optim.*, 71:449–472, 2015.
- [6] C. BARDOS, G. LEBEAU AND J. RAUCH. Sharp sufficient conditions for the observation, control and stabilization of waves from the boundary. *SIAM J. Control Optim.*, 30:1024–1065, 1992.
- [7] B. CHENTOUF. Compensation of the interior delay effect for a rotating disk-beam system. *IMA J. Math. Control Inform.*, 33:963–978, 2016.
- [8] C.M. DAFERMOS. Asymptotic stability in viscoelasticity. *Arch. Rational Mech. Anal.*, 37:297–308, 1970.

- [9] Q. DAI AND Z. YANG. Global existence and exponential decay of the solution for a viscoelastic wave equation with a delay. *Z. Angew. Math. Phys.*, 65:885–903, 2014.
- [10] C. GIORGI, J.E. MUÑOZ RIVERA AND V. PATA. Global attractors for a semilinear hyperbolic equation in viscoelasticity. *J. Math. Anal. Appl.*, 260:83–99, 2001.
- [11] A. GUESMIA. Well-posedness and exponential stability of an abstract evolution equation with infinite memory and time delay. *IMA J. Math. Control Inform.*, 30:507–526, 2013.
- [12] M. KIRANE AND B. SAID-HOUARI. Existence and asymptotic stability of a viscoelastic wave equation with a delay. *Z. Angew. Math. Phys.*, 62:1065–1082, 2011.
- [13] V. KOMORNIK AND C. PIGNOTTI. Energy decay for evolution equations with delay feedbacks. *Math. Nachr.*, to appear.
- [14] M. I. MUSTAFA AND M. KAFINI. Energy decay for viscoelastic plates with distributed delay and source term. *Z. Angew. Math. Phys.*, 67, Art. 36, 18 pp., 2016.
- [15] S. NICAISE AND C. PIGNOTTI. Stability and instability results of the wave equation with a delay term in the boundary or internal feedbacks. *SIAM J. Control Optim.*, 45:1561–1585, 2006.
- [16] S. NICAISE AND C. PIGNOTTI. Exponential stability of abstract evolution equations with time delay. *J. Evol. Equ.*, 15:107–129, 2015.
- [17] S. NICAISE AND C. PIGNOTTI. Well-posedness and stability results for nonlinear abstract evolution equations with time delays. *J. Evol. Equ.*, 18:947–971, 2018.
- [18] R.L. OLIVEIRA AND H.P. OQUENDO. Stability and instability results for coupled waves with delay term. *J. Math. Phys.*, 61, Art. 071505, 2020.
- [19] A. PAZY. Semigroups of linear operators and applications to partial differential equations. *Vol. 44 of Applied Math. Sciences. Springer-Verlag*, New York, 1983.
- [20] C. PIGNOTTI. Stability results for second-order evolution equations with memory and switching time-delay. *J. Dynam. Differential Equations* 29:1309–1324, 2017.
- [21] B. SAID-HOUARI AND A. SOUFYANE. Stability result of the Timoshenko system with delay and boundary feedback. *IMA J. Math. Control Inform.*, 29:383–398, 2012.
- [22] G.Q. XU, S.P. YUNG AND L.K. LI. Stabilization of wave systems with input delay in the boundary control. *ESAIM Control Optim. Calc. Var.*, 12:770–785, 2006.
- [23] Z. YANG. Existence and energy decay of solutions for the Euler-Bernoulli viscoelastic equation with a delay. *Z. Angew. Math. Phys.*, 66:727–745, 2015.