

ArielRad: the *Ariel* Radiometric Model

Lorenzo V. Mugnai · Enzo Pascale ·
Billy Edwards · Andreas Papageorgiou ·
Subhajit Sarkar

Received: date / Accepted: date

Abstract ArielRad, the *Ariel* radiometric model, is a simulator developed to address the challenges in optimising the space mission science payload and to demonstrate its compliance with the performance requirements. *Ariel*, the Atmospheric Remote-Sensing Infrared Exoplanet Large-survey, has been selected by ESA as the M4 mission in the Cosmic Vision programme and, during its 4 years primary operation, will provide the first unbiased spectroscopic survey of a large and diverse sample of transiting exoplanet atmospheres. To allow for an accurate study of the mission, ArielRad uses a physically motivated noise model to estimate contributions arising from stationary processes, and includes margins for correlated and time-dependent noise sources. We show that the measurement uncertainties are dominated by the photon statistic, and that an observing programme with about 1000 exoplanetary targets can be completed during the primary mission lifetime.

Keywords *Ariel* · exoplanet · simulated science

1 Introduction

In the past 20 years more than 4000 exoplanets have been detected using space and ground based surveys, and many more are expected to be discovered in

This work has been supported by ASI grant n. 2018.22.HH.O.

L. V. Mugnai · E. Pascale
Dipartimento di Fisica, La Sapienza Università di Roma, Piazzale Aldo Moro 2, 00185 Roma,
Italy E-mail: lorenzo.mugnai@uniroma1.it

B. Edwards
Department of Physics and Astronomy, University College London, Gower Street, London,
WC1E 6BT, UK

A. Papageorgiou · S. Sarkar
School of Physics and Astronomy, Cardiff University, Queens Buildings, The Parade, Cardiff,
CF24 3AA, UK

the coming years thanks to space missions such as *TESS* (Ricker et al., 2016), *CHEOPS* (Cessa et al., 2017), *PLATO* (Rauer et al., 2014) and *GAIA* (Gaia Collaboration et al., 2016), and to ground instrumentation such as *HARPS* (Mayor et al., 2003), *HATnet* (Bakos, 2018), *WASP* (Pollacco et al., 2006), *KELT* (Pepper et al., 2018), *OGLE* (Udalski et al., 2015), *NGTS* (Wheatley et al., 2013) and many others.

Planets have been found to be ubiquitous in our Galaxy, have been detected around almost every type of star and Cassan et al. (2012) infer that on average every star in our Galaxy hosts one planetary companion. The exoplanets detected thus-far show a diversity in their masses, sizes, orbits, and, presumably, physical and chemical conditions unseen among the planets in our own Solar System.

However, the essential nature of these exoplanets remains elusive. We have little idea whether the planet chemistry is linked to the formation environment or whether the type of host star drives the physics and chemistry of the planets birth, and evolution (Tinetti et al., 2018).

Atmospheric spectroscopy holds the key to unlock the mysteries of the chemical and physical conditions of these alien worlds as well as their formation and evolutionary histories. Multi-band photometry and spectroscopy of transiting exoplanets (Seager and Sasselov, 2000) is currently one of the most effective observational techniques for revealing the chemistry and thermodynamics of exoplanet atmospheres (Charbonneau et al., 2005; Tinetti et al., 2007; Sing et al., 2016; Madhusudhan et al., 2012; Huitson et al., 2012; Kreidberg et al., 2014). Photometric and spectroscopic light-curves of transiting exoplanets provide a measurement of the transmission (transit) or emission (eclipse) spectrum of an exo-atmosphere, and can be used to reveal chemical constituents, as well as the pressure and temperature profile, using retrieval techniques originally developed for the study of the Earth and Solar System planets, and adapted to the new field of investigation (Irwin et al., 2008; Line et al., 2013; Waldmann et al., 2015; Gandhi and Madhusudhan, 2017, e.g.).

Existing astronomical instrumentation has allowed to study spectroscopically a few tens of exoplanets, selected among those that are more easily observable based on their sizes and temperatures, and over a limited spectral range (e.g. Sing et al., 2016; Tsiaras et al., 2018). A significantly larger population study is now required in order to decipher the secrets of the exoplanets and their diversity.

The Atmospheric Remote-Sensing Infrared Exoplanet Large-survey, *Ariel*, has been selected by ESA as the next medium class mission of the Cosmic Vision programme to spectroscopically characterise the atmospheres of a large and diverse sample of exoplanets. *Ariel* will largely focus on warm and hot exoplanets, taking advantage from their well mixed atmospheres that show minimal condensation and sequestration of high atomic weight metals such as C, O, N, S, Si. The *Ariel* science payload uses a 1-m class telescope to feed a multi-band photometer and spectrometers covering the wavelength range from $0.5\,\mu\text{m}$ to $7.8\,\mu\text{m}$, to sample both the peak thermal emission of the exoplanet

atmospheres, and the spectral signature of all major atmospheric gases (e.g. H_2O , CO , CO_2 , NH_3 , CH_4 , HCN , H_2S , TiO , VO) and condensed species.

The *Ariel* payload design is investigated using detailed simulations of the astrophysical detection, that take into account mission design parameters such as flight duration and sky availability, and payload and astrophysical uncertainties. Margins are used on each estimate to ensure all performance predictions are derived under reasonably conservative assumptions.

In this work we describe ArielRad, the *Ariel* radiometric model simulator used to assess the payload science performance and to demonstrate its compliance with the science requirements. ArielRad is the third simulation tool developed to assess the mission performance and follows ExoSim (Sarkar et al., 2020), and the *Ariel* ESA Radiometric Model (AERM), developed by the Space Agency (Puig et al., 2015) to support the flow down of science requirements to instrument requirements the radiometric model during the *Ariel* phase/A study (Tinetti et al., 2018).

ExoSim is a end-to-end, time-domain simulator of *Ariel* observations. It evaluates photometric and spectroscopic light curves implementing a detailed description of the instrument design, source of uncertainties, and systematics of instrument and astrophysical origin. As such, the simulated observations produced by ExoSim are similar to those expected with *Ariel* and require a full data reduction pipeline to detrend the observations and reconstruct the planet spectrum. ExoSim has been fully validated using noise modelling as well as real measurements obtained with Hubble Space Telescope. ExoSim simulations allow us to study effects of spatially and temporally correlated noise processes such as the photometric uncertainties arising from the jitter of the line of sight, or from the activity of the host star. However, ExoSim analyses are computationally intensive and it is currently impractical to conduct studies on more than a few targets, until a fully automated data reduction pipeline is developed and validated in the next phase of the project. AERM overcomes this limitation implementing a simplified approach based on a radiometric modelling of the detection and of the uncertainties. These simplifications mean it is capable of assessing the confidence limit on the detection of emission and transmission spectra of hundreds of exoplanet targets. AERM implements a noise model calibrated using ExoSim estimates and delivers compatible estimates on test targets. However, the noise model implemented in AERM is a two parameters model that falls short in capturing the complexity of the *Ariel* payload design. Consequently, AERM provides an overly pessimistic prediction on some targets which makes this simulator unsuitable in assessing instrument design solutions.

ArielRad has been written to derive payload requirements from science requirements through detailed error budgeting, to validate the compliance of the payload design with the science requirements, and to optimise the payload design evaluating instrument design solutions over a proposed target list comprising about 1000 exoplanets. ArielRad overcomes the limitations of AERM by implementing a detailed payload model, similar to that used by ExoSim, capable of describing all major instrument components. ExoSim computing

limitations are overcome in ArielRad by implementing radiometric estimates of the uncertainties of the detection. Noise contributions that need to be estimated in the time domain, such as the photometric noise arising from the jitter of the line-of-sight, are imported in ArielRad from pre-computed ExoSim estimates. ArielRad is used to create and maintain the top level payload performance error budgets, allowing a balanced allocation of resources across the payload.

In this work we describe ArielRad, the models implemented, their validation, and provide examples of how ArielRad can be used to support the *Ariel* mission development, leaving to a future work a detailed assessment of the *Ariel* mission performance.

2 *Ariel* instrument design and observational strategy

The *Ariel* payload design is briefly described in this section with more details available in Tinetti et al. (2016); Tinetti et al. (2018); Pascale et al. (2018), and in the *Ariel* Assessment Study Reports¹. The telescope is an off-axis $0.63m^2$ Cassegrain with an elliptical primary mirror, that provides diffraction limited performance at wavelengths longer than $3\mu m$; there is no need for imaging capabilities and the telescope is cooled to less than $70K$. A refocusing mechanism actuates the secondary mirror to correct possible misalignments, that can occur at launch and during thermalisation.

The flux collected by the primary aperture feeds two separated instrument modules. A dichroic mirror splits the light into two beams, one at wavelengths shorter than $1.95\mu m$ and the second at wavelengths between $1.95\mu m$ and $7.8\mu m$. The first beam is fed to an instrument module containing three photometers (VISPhot, $0.5\mu m - 0.6\mu m$; FGS-1, $0.6\mu m - 0.80\mu m$; FGS-2, $0.80\mu m - 1.1\mu m$) and a slit-less prism spectrometer (NIRSpec, $1.1\mu m - 1.95\mu m$) with spectral resolving power > 15 . The two photometers, FGS1 and FGS2, operate as Fine Guidance Sensors (FGS), providing both photometric and pointing information for the attitude and orbital control system (AOCS). The second instrument module, fed by the longer wavelength beam, hosts the *Ariel* Infrared Spectrometer (AIRS), which consists of two prism-dispersed channels: Channel 0 (CH0) covering the $1.95\mu m - 3.9\mu m$ band with a spectral resolving power larger than 100, and Channel 1 (CH1) covering the $3.9\mu m - 7.8\mu m$ band with a spectral resolving power larger than 30. The spectrometers have field stops (slits) at an intermediate image plane, that are wider than the telescope Point Spread Function (PSF) and are used to limit the stray-light and the diffuse emission from reaching the focal plane.

During its four years primary mission, *Ariel* will observe ~ 1000 planets. To maximise the scientific return, the mission implements a three-tier observational and data analysis approach, where three different sample spectra are analysed with optimised spectral resolutions, wavelength intervals and Signal-to-Noise ratios (SNR). A summary of this observational strategy is given in

¹ <https://arielmission.space/ariel-publications/>.

Table 1 Summary of the science addressed in each tier.

Tier Name	Observational Strategy	Science case
Tier 1: Reconnaissance survey	Low Spectral Resolution observation of ~ 1000 planets in the VIS and IR, with SNR ~ 7	– What fraction of planets are covered by clouds? – What fraction of small planets have still retained H/He? – Classification through colour-colour diagrams – Constraining/removing degeneracies in the interpretation of mass-radius diagrams – Albedo, bulk temperature and energy balance for sub-sample
Tier 2: Deep survey	Higher Spectral Resolution observations of a sub-sample in the VIS-IR	– Main atmospheric component for small planets – Chemical abundances of trace gases – Atmospheric thermal structure (vertical/horizontal) – Cloud characterisation – Elemental composition
Tier 3: Benchmark planets	Very best planets, re-observed multiple times with all techniques	– Very detailed knowledge of the planetary chemistry dynamics – Weather, spatial and temporal variability

Tab. 1 with the first tier (Tier 1) being a low spectral resolution reconnaissance survey of ~ 1000 planets to address science questions on a large population of targets, such as to the fraction of planets covered by clouds or of small planets that have retained a H/He atmosphere. The second tier, Tier 2, which consists of $\sim 50\%$ of the planets from Tier 1, are studied with a higher spectral resolution, merging the data collected in two juxtaposed spectral bins. Tier 2 analysis searches for potential correlations between atmospheric chemistry and basic parameters such as planetary size, density, temperature, stellar type and metallicity. It allows investigations of chemical abundances, cloud characterisation and elemental composition. Finally, $\sim 10\%$ of Tier 1 planets are re-observed multiple times in Tier 3, and the data are analysed using the full spectral resolution provided by the payload to gain detailed knowledge of the planetary chemistry dynamics and temporal variability of the exoplanet atmospheres.

3 The ArielRad simulator

ArielRad is a radiometric simulator of the *Ariel* payload, it is implemented in Python and it is maintained by the *Ariel* Consortium. The software package comes with an exhaustive documentation and the primary inputs are contained in a XML configuration file describing the payload, an XML configuration file describing the mission parameters, and a CSV or spreadsheet table containing




Fig. 1 Simulator work flow. The simulation starts with two input files: a payload configuration file and a candidate planet list. ArielRad propagates the target host star signal through the payload, then evaluates the noise. Finally, the simulator estimates the transit or eclipse observation and the resultant SNR.

a list of target exoplanetary systems with their parameters. An instrument independent version of the radiometric simulator called ExoRad is publicly available on GitHub².

The simulator evaluates the payload science performance by estimating the expected experimental uncertainties on measured exoplanet atmospheric spectra in emission and transmission. Each simulation starts with the generation of the source signal from the star. The signal is then propagated through the instrument to the detector focal planes (assuming no field stars), accounting for the transmission of each optical component and the dispersion of the prism spectrometers.

Uncertainty estimates account for detector noise (readout noise and gain noise, the latter arising from variations in acquisition electronic chain), dark current, photon noise from the star, the zodiacal background and includes instrument emission, and jitter noise. Margins are included to account for uncertainties in the current noise estimates and instrument performances. The simulation continues by estimating the planet atmospheric signal and thus the SNR of the detection when observing the target in transit or eclipse. Every candidate target observation is considered to last 2.5 times the planet transit time, i.e. the time between the first and the last contact between the planetary and the stellar disks, T_{14} . This strategy allows the collection of data both in and out of transit for the light curve fit and the transit depth estimation. This parameter can be configured by the user as well as all other parameters described later. The data is then binned according to the tier resolution, as will be done in *Ariel* data analysis. Tier 1 uses a low resolution spectroscopy (4 spectral resolution elements covering $1.1 - 7.80\mu m$); Tier 2 has spectral resolution $R \sim 10$ for $1.1 < \lambda < 1.90\mu m$, $R \sim 50$ for $1.95 < \lambda < 3.90\mu m$ and $R \sim 15$ for $3.90 < \lambda < 7.80\mu m$; Tier 3 uses the full R , that means $R = 100, 30$ for AIRS-CH0 and AIRS-CH1 respectively, and for NIRSpec, as the requirement is $R > 15$, we use $R = 20$. Finally, ArielRad estimates the number of observations required to achieve a desired SNR for each planet (e.g. $SNR = 7$) and from this the total observing time required on target. This process is summarised graphically in Fig. 1) and each stage described in detail in the following sections.

² <https://github.com/ExObsSim/ExoRad2-public>

3.1 Source model

ArielRad models the target star using a grid of synthetic Phoenix spectra from Baraffe et al. (2015). Given a target, ArielRad selects the Phoenix spectrum that refers to a source with the most similar temperature, surface gravity and metallicity to the target star. The Spectral Energy Density (SED) of the target star at the telescope input is evaluated as

$$S(\lambda_k) = \frac{1}{\lambda_k - \lambda_{k-1}} \frac{R_\star^2}{D^2} \int_{\lambda_{k-1}}^{\lambda_k} S_{Ph}(\lambda) d\lambda \quad (1)$$

where $\lambda_k = \left(1 + \frac{1}{6000}\right)^k \lambda_{min}$ is a logarithmic spaced wavelength grid, determined starting from the sample spectral resolution of 6000 and the minimum wavelength λ_{min} . This binning reduces the spectral resolution of the input spectra for computational efficiency, while preserving the total power. $R_\star$ is the target star radius, D is the distance and S_{ph} the input Phoenix spectrum. The units of $S(\lambda_k)$ are those of a spectral energy density.

3.1.1 Diffuse radiation

ArielRad includes contributions from the Zodiacal background, and instrument thermal emission.

The Zodiacal background emission is modelled using a modified version of the JWST-MIRI Zodiacal model (Glasse et al., 2010), scaled according to the target position in the sky and the Zodi model of Kelsall et al. (1998):

$$I_{zodi}(\lambda) = A \cdot 3.510^{-14} \cdot BB(\lambda, 5500K) + B \cdot 3.5810^{-8} \cdot BB(\lambda, 270K) \quad (2)$$

where BB is the Planck law, and A and B are the fitted coefficients. Typically, A and B evaluate to ~ 0.9 at ecliptic poles, ~ 8 at a Solar elongation angle of 55 deg and at an ecliptic latitude of 0 deg, and ~ 2.5 on average. Therefore $A = B = 2.5$ is assumed when target position coordinates are not available (Pascale et al., 2015).

Instrument thermal emission, I_{inst} , is estimated by modelling each of the optical element as a Planck law at the element operating temperature and modified by a wavelength dependent emissivity. All the reflective surfaces are made in aluminium, and we conservatively assume a 3% emissivity. Same emissivity is assumed for the refractive components. All optical elements are passively cooled to 60K. Because of the almost isothermal *Ariel* design, the total instrument emission from optics is computed from the emission of one component, multiplied by the number of optical components along the light path.

The exception is the contribution of the detector box, I_{Inner} , as the flux is not coming from the instrument field of view (FoV) but instead directly irradiates the pixels from all directions. This contribution is referred to as the “Inner Sanctum” and it is estimated as the radiation emitted from a black body cavity at the detector operating temperature (i.e. unit emissivity is assumed).




Fig. 2 Total photon conversion efficiency, $\Phi_Y(\lambda)QE_Y(\lambda)$, as used in ArielRad.

3.2 Payload configuration

The payload is modelled using the current best estimates from the instrument system engineers. The transmission $\Phi_Y(\lambda)$, where Y is one of VISPhot, FGS1, FGS2, NIRSpec, AIRS-CH0 or AIRS-CH1, is obtained by simulating the light path from the telescope to the detectors through the optics. The detector quantum efficiency $QE_Y(\lambda)$ is also dependent on wavelength and defined for each channel. The photon conversion efficiency (PCE) is the product between transmission and quantum efficiency and it is shown in Fig. 2. The lower PCE observed in VISPhot and FGS1 with respect to FGS2 and NIRSpec is caused by a lower detector QE at short wavelengths, while a similar PCE reduction at AIRS wavelengths is mainly a consequence of the refractive materials used in the mid-infrared. The Point Spread Functions (PSF) are estimated as a function of the wavelength using external software and included, allowing for wavelength interpolations.

3.3 Signal model

The Point Response Function (PRF) is the PSF, normalised to unit volume and convolved with the pixel response, which is assumed to be a top-hat function, $H_{pix}(x, y)$. The effects of a non-flat pixel response function are only relevant for pointing jitter. As discussed later, this effect is estimated with ExoSim and imported in ArielRad as an additional noise contribution (see sec 3.4.3). Hence,

$$PRF(x_l, y_m, \lambda) = \int \int PSF(x, y, \lambda) H_{pix}(x_l - x, y_m - y) dx dy \quad (3)$$

where l and m are detector pixel indices, and $\sum_{l,m} PRF(x_l, y_m, \lambda) = 1$.

For each photometric channel, the signal of the target star T is modelled as the sum of the signals sampled by all detector pixels within an elliptical aperture, the size of which depends on the PRF. Hence, this is estimated as

$$S_{T,phot}(\lambda_s) = A_{tel} \sum_{x_l, y_m \in A_p} PRF(x_l, y_m, \lambda_s) \int QE_Y(\lambda) \Phi_Y(\lambda) S(\lambda) \frac{\lambda}{hc} d\lambda \quad (4)$$

where λ_s is the central wavelength of the photometric band (e.g. 0.55, 0.70 and 0.90 μm for VISPhot, FGS1 and FGS2 respectively). A_{tel} is the telescope effective collecting area. A_p is an elliptical aperture used for aperture photometry. Typical aperture sizes are chosen to encircle either 83.8% or 91.0% of the energy in the PSF as ExoSim models indicate a reduction of jitter noise to negligible levels compared to other noise sources for these aperture sizes (Sarkar and Pascale, 2015). The aperture, A_p , is defined over x,y pixels with coordinate indices l, m .

The case of a spectroscopic channel is similar and the signal sampled by the detector is estimated as

$$S_{T,spec}(\lambda_s) = A_{tel} \sum_{x_l, y_m \in A_s} PRF(x_l, y_m) QE_Y(\lambda) \Phi_Y(\lambda) S(y_m) \frac{\lambda}{hc} \Delta\lambda_m \quad (5)$$

where A_s is now a rectangular box aperture of the size of the spectral bin along the y-axis, assumed to be parallel to the dispersion direction. Apertures are spaced such as to obtain the desired spectral binning R , which is 20, 100, 30 for NIRSpec, AIRS-CH0 and AIRS-CH1 respectively. $\Delta\lambda_m$ is the wavelength interval sampled by the pixel at coordinate y_m , and λ_s is the wavelength at the centre of the spectral bin sampled by the spectrometer, defined as

$$\lambda_s = \frac{1}{2} (\lambda_j + \lambda_{j+1}) \quad (6)$$

where λ_j is the wavelength at the bin edge defined by the recursive relation

$$\lambda_{j+1} = \lambda_j \left(1 + \frac{1}{R} \right) \quad (7)$$

and the focal plane wavelength solution maps pixel coordinates to wavelength ($\lambda_{j+1}, \lambda_j \rightarrow y_{i+1}, y_i$).

For the Zodiacal light contribution, the effect is proportional to the pixel solid angle, $\Omega = \left(\frac{\Delta_{pix}}{f_{eff}} \right)^2$, where Δ_{pix} is the pixel size and f_{eff} is the effective focal length. For the photometers and the slit-less NIRSpec spectrometer this contribution is given by

$$S_{zodi,phot}(\lambda_s) = A_{tel} \Omega \int QE_Y(\lambda) \Phi_Y(\lambda) I_{zodi}(\lambda) \frac{\lambda}{hc} d\lambda \quad (8)$$

However, the AIRS instrument includes a field stop that being wider than the input PSF has no effects on a point source target, but acts as a slit for

diffuse radiation. Therefore, Zodiacal light must be modelled differently for AIRS channels. ArielRad simulates the signal incoming to the detector as the convolution between the Zodiacal light and the field stop/slit. If the slit width expressed in number of pixels at the focal plane is L , and the spectral resolving power computed at a certain λ_0 is $R(\lambda_0)$, the detector receives diffuse radiation over the wavelength range $\left(\lambda_j - \frac{L\lambda_0}{4R(\lambda_0)}, \lambda_j + \frac{L\lambda_0}{4R(\lambda_0)}\right)$, and not over the full range of wavelengths accepted by the filter, so

$$S_{zodi,spec}(\lambda_s) = A_{tel}\Omega \int_{\lambda_c - \frac{L\lambda_0}{4R(\lambda_0)}}^{\lambda_c + \frac{L\lambda_0}{4R(\lambda_0)}} QE_Y(\lambda)\Phi_Y(\lambda)I_{zodi}(y_m)\frac{\lambda}{hc}\Delta\lambda_m \quad (9)$$

where λ_0 is 1.95 and 3.90 μm and $R(\lambda_0)$ is $R(1.95) = 100$ and $R(3.90) = 30$ for AIRS-CH0 and AIRS-CH1 respectively.

ArielRad models the diffuse light coming from instrument emission by substituting $I_{zodi}(\lambda)$ with $n\epsilon I_{instr}(\lambda)$ in Eq. 8 and 9, where we are assuming that all the n optical elements (lenses, mirrors, etc.) have the same emissivity, ϵ . If the components are modelled with different emissivity, $\sum_{i=1}^n \epsilon_i I_{instr}(\lambda)$ is used instead.

The Inner Sanctum contribution is the same for all detector focal planes, as it originates from the emission of their enclosures, and it is proportional to the pixel surface:

$$S_{Inner}(\lambda_s) = \Delta_{pix}^2 (\pi - \Omega) \int QE_Y(\lambda)I_{Inner}(\lambda)\frac{\lambda}{hc}d\lambda \quad (10)$$

Diffuse backgrounds add a DC offset in the measured stellar signal. ArielRad assumes that these are removed using standard techniques, e.g. aperture photometry, and only their (uncorrelated) contribution to the noise budget is considered later in this work.

The resultant total signal in one spectral bin is the sum of all the previous contributions

$$S_Y(\lambda_s) = S_{T,Y}(\lambda_s) + S_{zodi,Y}(\lambda_s) + S_{instr,Y}(\lambda_s) + S_{Inner}(\lambda_s) \quad (11)$$

where in our notation Y can be one of VISPhot, FGS1, FGS2, NIRSpec, AIRS-CH0 or AIRS-CH1. $S_Y(\lambda_s)$ has units of counts (electrons) per second.

3.4 Noise model

ArielRad simulates the noise for each spectral bin from the signal estimate. In a real instrument, noise sources act at every stage of the detection chain and can be stationary and non-stationary. ArielRad estimates the contributions of noise components that are stationary random processes, such as Poisson noise and detector noise. It also includes Jitter noise (provided from an ExoSim simulation), margins for other noise contributions, and a noise floor. Fig. 3 shows a noise tree including all the noise sources considered.




Fig. 3 Noise tree diagram listing all noise contributors considered in Eq. 12. The Poisson Noise term depends on signals collected by the telescope (in yellow the target and the Zodiacal background photon noise), and on signals of instrument origin (in blue, photon noise from instrument and inner sanctum self emission). The detector noise contributors are from dark current, gain and read noise. The Jitter noise term is discussed separately from the rest of the noise terms as it is imported as a model from ExoSim simulations. The payload noise floor prevents noise from integrating down indefinitely.

The variance of each spectral bin is modelled as

$$\begin{aligned}
 \text{VAR}[S_Y(\lambda_s)] = & \begin{cases} (1 + \chi_Y) \text{VAR}_P[S_Y(\lambda_s)] \frac{1}{\Delta t_{int}} + & \text{(Poisson noise term)} \\ + \text{VAR}_D[S_Y(\lambda_s)] \frac{1}{\Delta t_{int}} + & \text{(Detector noise term)} \\ + \text{VAR}_J[S_Y(\lambda_s)] \frac{1}{\Delta t_{int}} + & \text{(Jitter noise term)} \\ + [p_0 S_{T,Y}(\lambda_s)]^2 & \text{(Payload noise floor)} \end{cases} \\
 & (12)
 \end{aligned}$$

where all quantities except p_0 are estimated at a 1 hr integration time and at longer time scales are assumed to decrease as the inverse of Δt_{int} , the integration time in hours. Time scales longer than 1 hr are expected to be above the correlation time scale of coloured noise processes considered. χ_Y is a margin added to include noise sources that cannot be simulated with ArielRad, such as correlated noise, time-dependent effects, and unknown unknowns. Typically, $\chi = 0.4$ is assumed, allowing for approximately 20% margin on top the photon noise that, from one side, it is conservatively larger than what has been achieved with *Hubble*/WFC3 and *Spitzer*/IRAC ($\sim 15\%$, Zellem et al., 2019), but from the other it is adequate for the current stage of the mission design.

The *Payload noise floor*, p_0 , models low frequencies instabilities (e.g. Brownian or pink noise processes) that prevents noise to integrate down indefinitely with time. Following Greene et al. (2016), we use $p_0 = 20$ ppm of the incoming signal $S_{T,Y}(\lambda_s)$. See also Beichman et al. (2014); Mollière et al. (2017) for an example of this approach.

3.4.1 Poisson Noise

The Poisson noise term includes all the photon noise contributions: target source, zodiacal background, inner sanctum and instrument emission. Hence, the Poisson variance at 1 hr time scale is

$$VAR_P[S_Y(\lambda_s)] = \frac{1 \text{ hr}}{3600 \text{ s}} \frac{1}{\eta} \frac{6(n^2 + 1)}{5n(n + 1)} S_Y(\lambda_s) \quad (13)$$

The factor 3600 s normalises the variance at 1 hr time scale and the parameter $\eta \lesssim 1$ is an efficiency accounting for detector reset events needed to prevent saturation and includes the temporal gap occurring between the end of the reset event and the first read. As the detector is read in n consecutive non-destructive reads (NDRs) during each exposure, Eq. 13 is multiplied by $\frac{6(n^2+1)}{5n(n+1)}$ as discussed in Rauscher et al. (2007) (MULTIACCUM readout strategy, see their Equation 1). This is appropriate as the number of group elements is 1 in the *Ariel* baseline payload design. The minimum value of n is 2, i.e. correlated double sampling (CDS) required to remove the detector KTC noise; $\frac{6(n^2+1)}{5n(n+1)}$ evaluates to 1 when $n = 2$. NDRs are read at a constant cadence and n depends from the detector saturation time

$$T_{sat} = \frac{f_{WD} \cdot WD}{S_{Y,max}} \quad (14)$$

where WD is the detector well depth, f_{WD} is the fraction of well depth used (e.g. $f_{WD} = 0.75$, Berta et al., 2012), and $S_{Y,max}$ is the largest pixel signal in the channel Y , that is estimated by ArielRad, but is not explicitly discussed here for the sake of conciseness. $VAR_P[S_Y(\lambda_s)]$ has units of s^{-2}hr .

3.4.2 Detector Noise

The detector noise variance can be divided in three terms, read noise, gain noise and dark current. The dark current signal depends on the detector pixel dark current, I_{dark} , and on the number of pixels, N_{pix} , included in the photometric apertures A_p or A_s defined earlier, with $S_{dark curr} = N_{pix} I_{dark}$. The detector noise variance at 1 hr is

$$\begin{aligned} VAR_D[S_Y(\lambda_s)] = & \frac{1 \text{ hr}}{3600 \text{ s}} \frac{12(n-1)}{n(n+1)} \frac{N_{pix} \sigma_{rd,Y}^2}{\eta^2 T_{sat}} + \\ & + \sigma_{gain,Y}^2 S_Y^2(\lambda_s) + \\ & + \frac{1 \text{ hr}}{3600 \text{ s}} \frac{1}{\eta} \frac{6(n^2 + 1)}{5n(n + 1)} S_{dark curr} \end{aligned} \quad (15)$$

where $\sigma_{rd,Y}^2$ is the noise variance on each individual NDR and has no units. Following Rauscher et al. (2007), the factor $\frac{12(n-1)}{n(n+1)}$ accounts for a line fit to the NDR ramp in one exposure, and decreases as n increases.

The gain noise, $\sigma_{gain,Y}$ in units of $\sqrt{\text{hr}}$, accounts for instabilities of the electronic acquisition chain (amplifiers, digitisers, etc.), assumed post-processing, i.e. after common modes are removed using housekeeping information. With T_{sat} in units of seconds, $VAR_D[S_Y(\lambda_s)]$ has units of s^{-2}hr .

3.4.3 Jitter Noise

Pointing drifts and jitter of the line-of-sight (LoS) manifest themselves in the observed data product via two mechanisms: 1) the drifting of the spectrum along the spectral dispersion axis of the detector; 2) the drift of the spectrum along the cross-dispersion (spatial) direction. The effect of jitter on the observed time series is the introduction of noise, characterised by the power-spectrum of the telescope pointing (usually not stationary). It is correlated in time, as the power-spectrum is not constant in frequency. The amplitude of the resultant photometric scatter depends on the amount of spectral/spatial displacement of the spectrum, the monochromatic PSF of the instruments, the detector pixel response function (intra-pixel response) and the amplitude of the inter-pixel variations (i.e. QE variations across the focal plane detector array). Modelling the complexity of the jitter noise effect is beyond the capabilities of a radiometric model, therefore ArielRad imports jitter noise models from ExoSim simulations that provides variance vs wavelength at a timescale of 1 hr, i.e. $VAR_J([S_Y(\lambda_s)]$ in Eq. 12. At longer timescales, jitter noise is to good approximation time uncorrelated and can be therefore scaled at any desired observing time longer than 1 hr.

3.4.4 Additional noise contributions

As ArielRad is a radiometric model of the *Ariel* payload performance, it cannot simulate non-stationary noise processes, or processes that require a more sophisticated simulation strategy, i.e. time-domain. One example of the latter is the already discussed pointing jitter, that is accounted for in ArielRad using external modelling with ExoSim.

Detector persistence is an additional potential systematic that cannot be modelled with ArielRad, but *HST* observations have shown that it can be effectively corrected in data analysis (Zhou et al., 2017). Further, this effect is expected to be negligible in *Ariel* observations compared to *HST* as *Ariel* stares continuously at a target from an L2 orbit ensuring that detectors reach a steady state in a relatively short amount of time (few minutes at most).

Detector non-linear behaviour is well characterised in detector testing activities before launch and during commissioning. Coupled with a stable line-of-sight provided by the AOCS at the sub-pixel level, detectors pixels sample the same optical power level during an observation, therefore dwarfing, relatively




Fig. 4 ArielRad noise budget example of the *Ariel* payload at 1 hr integration time, using Tier 3 binning. Two targets are considered: GJ 1214, a faint target for the *Ariel* mission, and HD 209458, a typical bright target. The total noise estimate is in black and includes a 20% margin in excess of the target photon noise (green). Also shown are the detector dark current and read noise (purple and yellow, respectively), photon noise from the zodiacal background (blue), gain noise (brown), and pointing jitter noise (gray). A noise floor is indicated by the horizontal dashed line. The pointing jitter noise model is estimated using ExoSim (Sarkar and Pascale, 2015) simulations with identical payload parametrization. Gain noise is assumed to be equal to 40 ppm $\sqrt{\text{hr}}$. Photon noise from instrument and inner sanctum emission are omitted because are negligibly small and out of the scale of the diagram. The *Ariel* payload model used in this budget is that at the mid of phase B1, it includes a noise floor of 20ppm and identical detector median dark current noise across all channels, that might result in an overestimate of up to a factor of 3 in all channels, but CH1.

to other sources of experimental uncertainties, this type of systematic that is also known to be amendable (de Jong, 2006).

Throughput variations can be caused by thermoelastic deformations that can also induce temporal variations in the shape of the PSF. To minimise this effect, the *Ariel* payload design uses aluminium structures, including the telescope mirrors. Along with the thermal stability enjoyed by spacecraft in L2 orbits, this prevents driving any significant thermal gradient on the payload structure and subsystems. It is therefore expected by design that thermoelastic throughput variations during *Ariel* observations will be significantly below any detection limit.

3.4.5 Output format

ArielRad outputs the noise integrated over 1 hr of observation relative to the target signal:

$$\sigma_{Y,1hr}(\lambda_s) = \frac{\sqrt{(1 + \chi_Y)VAR_P + VAR_D + VAR_J}}{S_{T,Y}(\lambda_s)} \quad (16)$$

The units are $\text{hr}^{1/2}$. The relative noise achieved during one observation is

$$\sigma_{Y,\Delta t}^2(\lambda_s) = \frac{\sigma_{Y,1\text{hr}}^2(\lambda_s)}{\Delta t_{\text{int}}} + p_0^2 \quad (17)$$

where Δt_{int} is the observing time in hours, and the payload noise floor p_0 is added in quadrature to prevent the noise to integrate down indefinitely as discussed earlier.

The payload (relative) noise budget is shown in Fig. 4 for a typical bright target star (HD 209458) and for a faint target star (GJ 1214). These stars are representative of respectively the bright and faint flux limits of the payload design requirements. The noise budget has been estimated at 1 hr integration assuming that detectors are analysed in CDS samples, i.e. $n = 2$. The budget is sampled at Tier 3 resolution and it allows comparison of noise contributions to the total noise in any individual *Ariel* channel. A comparison of sensitivity among channels for any given target needs to take into account the different spectral binning used that is higher in CH0 ($R = 100$) compared to NIRSpec ($R = 20$) and CH1 ($R = 30$). A strong increase with wavelength is seen in the read and dark current noise components as a consequence of a decreasing stellar SED toward the red.

As it is evident from the noise budget, the *Ariel* payload design reaches photon noise limited performances, allowing observations of even dimmer targets.

3.4.6 Signal-to-Noise ratio

In a transit or eclipse observation, the observable is the wavelength-dependent contrast-ratio that is the difference between the flux incoming from the extra-solar system star plus planet when the planet is moving in front (transit) or behind (secondary transit, i.e. eclipse) the star and when it is not:

$$CR(\lambda) = \frac{S_{OOT}(\lambda) - S_{IT}(\lambda)}{S_{OOT}(\lambda)} \quad (18)$$

where the labels *OOT* and *IT* identify signals measured respectively out of transit and in transit.

For a primary transit the contrast ratio is evaluated from the comparison between the SED of the star $S_\star(\lambda)$ plus that of the planet $S_p(\lambda)$, $S_{OOT}(\lambda) = S_\star(\lambda) + S_p(\lambda)$, and the SED measured during the transit, that up to the first order depends on the radii ratio (Mandel and Agol, 2002; Seager and Mallén-Ornelas, 2002). ArielRad simulates the contrast ratio in each spectral bin for the primary transit, considering also the portion of the star light passing through the planet atmosphere, as described in Seager and Sasselov (2000) and Brown (2001):

$$CR_{\text{tot}}(\lambda) = \left(\frac{R_p}{R_\star}\right)^2 + 2\Delta z(\lambda) \frac{R_p}{R_\star^2} \quad (19)$$

where R_p and $R_\star$ are the planet and star radii respectively and $\Delta z(\lambda)$ is the atmospheric height. Because the interest is on the detection of the exoplanet atmosphere, the “signal” is just the rightmost quantity (Louie et al., 2018; Zellem et al., 2019) in the expression above, i.e.

$$CR(\lambda) = 2\Delta z(\lambda) \frac{R_p}{R_\star^2} \quad (20)$$

The atmospheric height is proportional to the scale height $H = k_B T / \mu g$ where k_B is the Boltzmann constant, T is the temperature of the atmosphere, μ is the mean molecular weight and g is the gravitational acceleration. The wavelength dependent constant of proportionality can be provided by atmospheric models or, following Tinetti et al. (2013), set to 5 independently from the wavelength for a simple, yet representative performance estimate.

For the eclipse case, the contrast ratio is simply $CR(\lambda) = S_p(\lambda) / S_\star(\lambda)$ (Charbonneau et al., 2005; Deming et al., 2005), and the planetary SED is part due to the planet thermal emission, $S_{em}(\lambda)$, and part due to the reflected star light, $A_I(\lambda)$. Thus,

$$CR(\lambda) = \frac{S_{em}(\lambda) + A_I(\lambda)}{S_\star(\lambda)} \quad (21)$$

The simulator estimates the contrast ratio in each spectral bin modelling the planet emission as a Black Body at the planet temperature, and computes the reflected light component according to Charbonneau et al. (1999) as $A_I(\lambda) = \alpha(\lambda) \left(\frac{R_p}{a}\right)^2 S_\star(\lambda)$ where $\alpha(\lambda)$ is the geometric albedo and a is the semi-major orbital axis.

ArielRad can estimate these contrast-ratios considering the observation of S_{IT} lasting the time between the first and last contact, T_{14} , and S_{OOT} lasting γT_{14} , with $\gamma = 1.5$, typically, from current *Ariel* science requirements. Using Eq. 17 and 18, the noise variance estimate on a contrast ratio measurement is

$$VAR(CR_Y, \lambda_s) = \sigma_{Y,1hr}^2(\lambda_s) \left[1 + \frac{1}{\gamma} \right] \frac{1}{T_{14}} + p_0^2 \quad (22)$$

and the Signal-to-Noise Ratio in each spectral bin is

$$SNR_Y(\lambda_s) = \frac{CR_Y(\lambda_s)}{\sqrt{VAR(CR_Y, \lambda_s)}} \quad (23)$$

where we can substitute $CR(\lambda_s)$ with the estimated contrast ratio for transit or eclipse observations. The label Y indicates that quantities are integrated over the photometric band or spectral bin of interest. Therefore ArielRad estimates three sets of SNR, one for each of the three *Ariel* tier discussed in Section 2.

The SNR achieved in multiple, N_{obs} observations of the same target extra-solar system is assumed to scale as $\sqrt{N_{obs}}$, under the assumption that disturbances originate from stochastic processes that are uncorrelated over the time scale separating two observations, and longer. If this were not the case,

it would be more appropriate to assume that the noise floor is not reduced by averaging multiple observations. However, there is currently no evidence supporting this as *Hubble*/WFC3 and *Spitzer*/IRAC observations have not yet reached a noise floor (Zellem et al., 2019) .

4 Validation

ArielRad has been validated by comparing its estimates to those of ExoSim and AERM introduced in Section 1.

As ExoSim has been extensively validated against real astrophysical observations (Sarkar et al., 2020), the comparison between ArielRad and ExoSim estimates are the most interesting to investigate. For this, we chose to compare the predictions made of T_{sat} (Eq. 14). A consistent result between ExoSim and ArielRad on this parameter implies that ArielRad captures the complexity of the payload design as thoroughly as ExoSim does, and validates the implementation of the radiometric algorithms. The comparison is therefore done implementing the same baseline *Ariel* model in the two simulators, and T_{sat} is evaluated for the three target stars GJ 1214 (M4.5, $\text{mag}_K \simeq 8.8$), HD 209458 (G0V, $\text{mag}_K \simeq 6.3$) and HD 219134 (K3V, $\text{mag}_K \simeq 3.3$) that cover a range in brightness and temperature representative of potential *Ariel* targets (e.g. Edwards et al., 2019). The comparison is given in Table 2 For the visible photometer and the infrared spectrometer; it is found that the two models agree to better than 5% on all targets.

Table 2 Comparison between T_{sat} estimates with ExoSim and ArielRad.

Channel	GJ 1214	HD 209458	HD 219134
	Percent variation		
VISPhot	-0.8	-0.5	-0.4
AIRS-CH0	-2.9	1.0	1.2
AIRS-CH1	4.4	2.5	2.8

The comparison between ArielRad and AERM provides validation of the SNR calculations summarised by Eq. 23, that means a validation of all calculations implemented concerning the estimate of the exoplanet atmospheric signatures observable during a transit or an eclipse and uncertainties. Puig et al. (2015), and reference therein, detail the algorithms implemented by AERM, that estimates the noise during 1 s of integration as

$$\sigma_{Y,Total} = \sqrt{[S_{T,Y}(\lambda_s) + S_{zodi,Y}(\lambda_s)] \cdot (1 + \chi_Y) + N_{min,Y}(\lambda_s)} \quad (24)$$

The main difference with the ArielRad implementations is the $N_{min}(\lambda_s)$ term that combines the noise variance in a spectral bin or photometric band from detector dark current and instrument emissions. For the photometric channels, $N_{min,Y}(\lambda_s)$ is set to 400 s^{-1} . For the prism spectrometers, $N_{min,Y}(\lambda_s) \propto \lambda_s^3$,




Fig. 5 Comparison between ArielRad and AERM SNR estimates in one transit (*left*) and one eclipse (*right*) observation of a sample of 2500 candidate exoplanet atmospheres. Each datapoint is the average SNR across the NIRSpec (green), AIRC-CH0 (red) and AIRS-CH1 (blue) bands, binned at the *Ariel* tier 3 spectral resolution. Trend lines are shown with slope and intercept given in the text annotation in each panel. A black dotted line with unity slope is shown. While estimates for CH0 and CH1 are in good agreement among both models, AERM provides SNR estimates in NIRSpec that systematically and unrealistically larger than ArielRad. This is due to a limitation in the AERM algorithms implemented and it is further discussed in Section 4.

and it is set to $17\text{ s}^{-1}\mu\text{m}^{-3}$, $20\text{ s}^{-1}\mu\text{m}^{-3}$, and $5\text{ s}^{-1}\mu\text{m}^{-3}$ at the blue end of NIRSpec, AIRS-CH0 and AIRS-CH1, respectively. There is no provision for other noise sources in AERM, including detector readout and gain noise, nor pointing jitter. These effects are accounted for in the term χ_Y that is set to 0.2 for the photometers and NIRSpec, and to 0.3 for both AIRS channels. AERM further assume a photon conversion efficiency that, in each photometric or spectroscopic channel is wavelength independent. For this validation exercise, the ArielRad input configuration is adapted to match that in AERM, in terms of photon conversion efficiency, and matching dark currents to the equivalent $N_{\min,Y}(\lambda_s)$. Furthermore, the parameters in Eq. 13 are chosen such that $n = 2$ (i.e. AERM assumes CDS), $\eta = 1$ (i.e. AERM assumes 100% sampling efficiency) and the photon noise margin parameter χ are set to those in use in AERM. VAR_D , VAR_J , and p_0 are set to zero in Eq. 12. With this, AERM and ArielRad implement effectively the same instrument model, and are run to evaluate the SNR achieved on one transit or one eclipse over 2500 candidate *Ariel* targets (Edwards et al., 2019). Figure 5 shows the comparison of the two model estimates of the average SNR over spectral bins in each spectrometer. We find that, on average, the SNR estimates agree to better than 2 or 3% for AIRS channels. For NIRSpec we find that on average ArielRad predicts SNR that are up to 8% smaller than AERM's predictions. A further investigation has revealed the root cause of this discrepancy in the way AERM estimates the signals: while ArielRad integrates SEDs over wavelengths in a spectral bin, AERM uses the SED estimated at the blue end of each spectral bin. This always implies that signal in AERM are systematically larger resulting

is smaller uncertainties. The effect is more evident in NIRSpec because the larger spectral bin width ($R = 10$, as implemented in ESARad) compared to AIRS channels ($R=100$ and $R = 30$ in CH0 and CH1, respectively).

5 Use of ArielRad

ArielRad has been developed to support the *Ariel* phase B study, leading to Mission Adoption by ESA in the Autumn 2020. ArielRad allows to evaluate the payload science performance over a large target list of thousands of potential exoplanetary targets and to assess the compliance of the payload design with the science requirements briefly discussed in Section 2, with a more detailed discussion in Tinetti et al. (2018) and in the *Ariel* yellow book³. ArielRad is the main tool used to assess the *Ariel* payload design solutions and provides a guide to optimise the payload to achieve requirements, and to maximise the *Ariel* science return beyond requirements, when possible.

The observed diversity of exoplanets can only be investigated by surveying a large parameter space in planetary radii and masses, thermodynamic conditions, chemical properties and host star types. *Ariel* is designed to provide the first large survey of the atmospheres of about 1000 diverse planets and ArielRad is the tool used to craft a target list that is compliant with this science mandate. Edwards et al. (2019) used ArielRad simulations to provide a preliminary mission reference sample (MRS) of 1000 planets. While the MRS is expected to evolve during the next phases of the project until launch in 2028, the ArielRad performance analysis demonstrates that the atmospheres of planets in the MRS can be characterised with a $SNR > 7$ during the 4 year nominal mission lifetime.

As discussed in Section 3.4, ArielRad provides a detailed description of the *Ariel* noise budget (Figure 4) on individual targets. The analysis can be extended to provide a comprehensive description of the payload performance over all targets in the MRS to show how *Ariel* achieves a photon noise limited performance on all targets, as shown in Figure 6. The MRS of Edwards et al. (2019) is used. It lists both exoplanets already discovered and expected TESS yields. At AIRS wavelengths ($\lambda_s > 1.95 \mu\text{m}$) photon noise is the dominant source of uncertainty, dwarfing all other noise contributions. At shorter wavelength, photon noise is less important for a small, but significant number of targets, with detector gain noise playing a larger role in the noise budget. These are largely TESS targets around M-type (cold) host stars, demonstrating the power of the *Ariel* IR bands for these type of targets. The *Ariel* telescope is diffraction limited at a wavelength of $3 \mu\text{m}$ and significant optical aberrations degrade the image quality at shorter wavelengths. However, *Ariel* works as a light bucket, and image quality is not relevant to achieve its performance requirements. This aspect is also investigated in Figure 6 where the analysis is done for both a diffraction limited instrument and using estimates of aberrated PSF from engineering optical modelling, corresponding to a wave front

³ <https://sci.esa.int/s/8zPrb9w>.




Fig. 6 Total noise to photon noise ratio in one hour integration for all *Ariel* MRS targets. Solid and dashed lines mark median values across all targets evaluated for diffraction limited (red) and aberrated (blue) optics. Blue area represents the dispersion of 95% of simulated planets for the aberrated optics configuration. As the MRS contains exoplanets already discovered as well as expected TESS yields, these are separated respectively in the left and right panels. As discussed in the text, the apparent excess noise above photon noise at visible-nearIR wavelengths is due to the presence of cold M-type stellar targets, that are more numerous among TESS targets, and the assumed gain noise contribution. For these type of targets, observations in the nearIR-midIR are more sensitive showing the power of *Ariel*’s IR bands that are always shot noise limited.

error of 250 nm RMS at the VISPhot, FGS1, FGS2 and NIRSpec focal planes, and 280 nm RMS at the AIRS focal planes. The differences are negligible and it can be noted from the figure that an aberrated PSF behaves slightly better than a diffraction limited PSF, despite the former requires a larger number of pixel in each photometric or spectral bin aperture. However, because the aberrated PSF dilutes the signal more, pixels take longer to saturate, there are fewer exposure in a given observing time, hence read noise has overall a smaller impact.

ArielRad uncertainties estimates support the the science community to optimise the science of *Ariel*. For instance Changeat et al. (2019) used ArielRad estimates to investigate the capability of *Ariel* in retrieving pressure-dependent chemical profiles from predictions of observed atmospheric spectra. Science analyses using ArielRad performance estimates are ongoing, include aspects related to atmospheric retrieval, phase curve detection, transit timing variations, etc., and will be reported in *Ariel* phase B study report later in 2020 ahead of Mission Adoption: the *Ariel* ”Red Book”.

6 Conclusions

In this work we have discussed the algorithmic implementation of ArielRad, the *Ariel* radiometric simulator, used for the optimisation of the payload design and to evaluate the science performance of the ESA M4 space mission.

ArielRad accounts for all relevant sources of uncertainties on the detection of exoplanetary atmospheres with *Ariel*, that are: photon noise (of astrophysical origin and from the instrument self emission), detector noise and electronic noise, and jitter noise. All other potential systematic of instrument origin are expected to be made negligible by a careful instrument design as detailed by Tinetti et al. (2018) and further discussed in the *Ariel* Yellow Book.

ArielRad has been extensively validated against two alternative models, ExoSim and AERM, always showing excellent agreement at a few percent level across all *Ariel* bands.

Ariel will perform the first statistical survey of the atmospheres of a large and diverse sample, observing about 1000 exoplanets during its life time, as discussed in Edwards et al. (2019) with the help of ArielRad. For all exoplanet observations, the photon noise of their host stars is the dominant source of uncertainties, as revealed by an ArielRad assessment.

ArielRad performance estimates, in the form of noise vs wavelength achieved on a given set of exoplanetary targets, are a product distributed upon request. As the phase-B continues, the payload design is optimised in an iterative process that aims at building the most performant space mission within the envelope provided. An advancement in payload design does not however imply the need to modify the algorithmic implementation of ArielRad thanks to its parametric description of the payload model. As a consequence, the simulator itself is planned to be released at the freeze-out of the payload design, to occur after mission adoption.

Acknowledgements We thank G. Pilbratt (ESA-ESTEC) for comments that greatly improved the development of ArielRad.

Appendices

Acronyms and symbols

References

- Bakos GÁ (2018) The HATNet and HATSouth Exoplanet Surveys, p 111. DOI 10.1007/978-3-319-55333-7_111
- Baraffe I, Homeier D, Allard F, Chabrier G (2015) New evolutionary models for pre-main sequence and main sequence low-mass stars down to the hydrogen-burning limit. *Astronomy & Astrophysics* 577:A42, DOI 10.1051/0004-6361/201425481
- Beichman C, Benneke B, Knutson H, Smith R, Lagage PO, Dressing C, Latham D, Lunine J, Birkmann S, Ferruit P, Giardino G, Kempton E, Carey S, Krick J, Deroo PD, Mandell A, Ressler ME, Shporer A, Swain M, Vasisht G, Ricker G, Bouwman J, Crossfield I, Greene T, Howell S, Christiansen J, Ciardi D, Clampin M, Greenhouse M, Sozzetti A, Goudfrooij P, Hines D, Keyes T, Lee J, McCullough P, Robberto M, Stansberry J, Valenti J, Rieke M, Rieke G, Fortney J, Bean J, Kreidberg L, Ehrenreich D, Deming D, Albert L, Doyon R, Sing D (2014) Observations of Transiting Exoplanets with the James Webb Space Telescope (JWST). *pasp* 126(946):1134, DOI 10.1086/679566

Table 3 Symbols used in the main text ordered by comparison

Symbol	meaning
T_{14}	time between first and fourth contact in transit
S_{Ph}	Phoenix Spectral Energy Density
S	stellar Spectral Energy Density
R_*	Stellar radius
D	stellar distance
I_{zodi}	zodiacal Background Spectral Energy Density
BB	black body Spectral Energy Density
I_{inst}	payload instrument Spectral Energy Density
I_{Inner}	detector box Spectral Energy Density
Φ	optical transmission
Y	channel identifier
QE	quantum efficiency
PRF	point response function
H_{pix}	pixel response function
x, y	pixels in the detector
l, m	pixel indices in x and y axes
$S_{T,Y}$	target signal in channel
λ_s	wavelength at spectral bin center
A_{tel}	telescope area
A_p	elliptical aperture in pixel used for aperture photometry in photometric channels
A_s	rectangular box aperture of the size of spectral bin used in spectrometers
R	spectral resolution
Ω	field of view
Δ_{pix}	pixel size
f_{eff}	telescope effective focal length
$S_{zodi,Y}$	zodiacal signal in channel
R_0	spectral resolution at $\lambda = 3.90\mu m$
L	slit width expressed in number of pixel on the focal plane

- Berta ZK, Charbonneau D, Désert JM, Miller-Ricci Kempton E, McCullough PR, Burke CJ, Fortney JJ, Irwin J, Nutzman P, Homeier D (2012) The Flat Transmission Spectrum of the Super-Earth GJ1214b from Wide Field Camera 3 on the Hubble Space Telescope. *ApJ* 747(1):35, DOI 10.1088/0004-637X/747/1/35, 1111.5621
- Brown TM (2001) Transmission Spectra as Diagnostics of Extrasolar Giant Planet Atmospheres. *The Astrophysical Journal* 553(2):57, DOI 10.1086/320950, 0101307
- Cassan A, Kubas D, Beaulieu JP, Dominik M, Horne K, Greenhill J, Wambsganss J, Menzies J, Williams A, Jørgensen UG, Udalski A, Bennett DP, Albrow MD, Batista V, Brilliant S, Caldwell JAR, Cole A, Coutures C, Cook KH, Dieters S, Dominis Prester D, Donatowicz J, Fouqué P, Hill K, Kains N, Kane S, Marquette JB, Martin R, Pollard KR, Sahu KC, Vinter C, Warren D, Watson B, Zub M, Sumi T, Szymański MK, Kubiak M, Poleski R, Soszynski I, Ulaczyk K, Pietrzyński G, Wyrzykowski L (2012) One or more bound planets per Milky Way star from microlensing observations. *Nature* 481:167–169, DOI 10.1038/nature10684, 1202.0903
- Cessa V, Beck T, Benz W, Broeg C, Ehrenreich D, Fortier A, Peter G, Magrin D, Pagano I, Plesseria JY, Steller M, Szoke J, Thomas N, Ragazzoni R, Wildi F (2017) CHEOPS: a space telescope for ultra-high precision photometry of exoplanet transits. In: Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, vol 10563, p 105631L, DOI 10.1117/12.2304164
- Changeat Q, Edwards B, Waldmann I, Tinetti G (2019) Towards a more complex description of chemical profiles in exoplanets retrievals: A 2-layer parameterisation. *arXiv e-prints* arXiv:1903.11180, 1903.11180

Table 4 Table 3 (*continued*)

Symbol	meaning
S_{Inner}	detector box signal in spectroscopic channels
S_Y	total incoming signal in photometric or spectroscopic channels
$S_{instr,Y}$	instrument signal in photometric or spectroscopic channels
$VAR()$	total variance
$VAR_P()$	Poisson noise contribution to the variance
$VAR_D()$	Detector noise contribution to the variance
$VAR_J()$	Jitter noise contribution to the variance
Δt_{int}	integration time expressed in hours
χ_Y	margin added to Poisson noise
N_{pix}	number of pixel included in spectral bin
I_{dark}	detector dark current
n	number of Non Destructive Reads
T_{sat}	detector saturation time
WD	pixel well depth
f_{WD}	coefficient describing the fraction of pixel well depth
σ_{rd}	pixel read noise
CR	contrast ratio
S_{OOT}	flux observed out-of-transit
S_{IT}	flux observed in-transit
S_P	flux from the planet
$S_\star$	flux from the star
Δz	atmosphere height
R_p	planetary radius
S_{em}	flux from the planet thermal emission
A_l	fraction of starlight reflected by the planet
α	geometrical albedo
a	semi-major axis in planet orbit
N_{min}	minimum noise considered in the channel in ESA radiometric model for <i>Ariel</i>

- Charbonneau D, Noyes RW, Korzennik SG, Nisenson P, Jha S, Vogt SS, Kibrick RI (1999) An upper limit on the reflected light from the planet orbiting the star τ bootis. The Astrophysical Journal 522(2):L145–L148, DOI 10.1086/312234
- Charbonneau D, Allen LE, Megeath ST, Torres G, Alonso R, Brown TM, Gilliland RL, Latham DW, Mandushev G, O'Donovan FT, Sozzetti A (2005) Detection of Thermal Emission from an Extrasolar Planet. Astrophysical Journal 626(1):523–529, DOI 10.1086/429991, [astro-ph/0503457](#)
- de Jong RS (2006) Correcting the NICMOS count-rate dependent non-linearity. Tech. rep.
- Deming D, Seager S, Richardson LJ, Harrington J (2005) Infrared radiation from an extrasolar planet. Nature 434(7034):740–743, DOI 10.1038/nature03507, [astro-ph/0503554](#)
- Edwards B, Mugnai L, Tinetti G, Pascale E, Sarkar S (2019) An Updated Study of Potential Targets for Ariel. AJ 157(6):242, DOI 10.3847/1538-3881/ab1cb9, [1905.04959](#)
- Gaia Collaboration, Prusti T, de Bruijne JHJ, Brown AGA, Vallenari A, Babusiaux C, Bailer-Jones CAL, Bastian U, Biermann M, Evans DW, Eyer L, Jansen F, Jordi C, Klioner SA, Lammers U, Lindegren L, Luri X, Mignard F, Milligan DJ, Panem C, Poinsignon V, Pourbaix D, Randich S, Sarri G, Sartoretti P, Siddiqui HI, Soubiran C, Valette V, van Leeuwen F, Walton NA, Aerts C, Arenou F, Cropper M, Drimmel R, Høg E, Katz D, Lattanzi MG, O'Mullane W, Grebel EK, Holland AD, Huc C, Passot X, Bramante L, Cacciari C, Castañeda J, Chaoul L, Cheek N, De Angeli F, Fabricius C, Guerra R, Hernández J, Jean-Antoine-Piccolo A, Masana E, Messineo R, Mowlavi N, Nienartowicz K, Ordóñez-Blanco D, Panuzzo P, Portell J, Richards PJ, Riello M, Seabroke GM, Tanga P, Thévenin F, Torra J, Els SG, Gracia-Abril G, Comoretto G, Garcia-Reinaldos M, Lock T, Mercier E, Altmann M, Andrae R, Astraatmadja TL, Bellas-Velidis I, Benson K,

Berthier J, Blomme R, Busso G, Carry B, Cellino A, Clementini G, Cowell S, Creevey O, Cuyppers J, Davidson M, De Ridder J, de Torres A, Delchambre L, Dell'Oro A, Ducourant C, Frémat Y, García-Torres M, Gosset E, Halbwachs JL, Hambly NC, Harrison DL, Hauser M, Hestroffer D, Hodgkin ST, Huckle HE, Hutton A, Jasiewicz G, Jordan S, Kontizas M, Korn AJ, Lanzafame AC, Manteiga M, Moitinho A, Muinonen K, Osinde J, Pancino E, Pauwels T, Petit JM, Recio-Blanco A, Robin AC, Sarro LM, Siopis C, Smith M, Smith KW, Sozzetti A, Thuillot W, van Reeve W, Viala Y, Abbas U, Abreu Aramburu A, Accart S, Aguado JJ, Allan PM, Allasia W, Altavilla G, Álvarez MA, Alves J, Anderson RI, Andrei AH, Anglada Varela E, Antiche E, Antoja T, Antón S, Arcay B, Atzei A, Ayache L, Bach N, Baker SG, Balaguer-Núñez L, Barache C, Barata C, Barbier A, Barblan F, Baroni M, Barrado y Navascués D, Barros M, Barstow MA, Becciani U, Bellazzini M, Bellei G, Bello García A, Belokurov V, Bendjoya P, Berihuete A, Bianchi L, Bienaymé O, Billebaud F, Blagorodnova N, Blanco-Cuaresma S, Boch T, Bombrun A, Borrachero R, Bouquillon S, Bourda G, Bouy H, Bragaglia A, Breddels MA, Brouillet N, Brüsemeister T, Bucciarelli B, Budnik F, Burgess P, Burgon R, Burlacu A, Busonero D, Buzzi R, Caffau E, Cambras J, Campbell H, Cancelliere R, Cantat-Gaudin T, Carlucci T, Carrasco JM, Castellani M, Charlot P, Charnas J, Charvet P, Chassat F, Chiavassa A, Clotet M, Coccozza G, Collins RS, Collins P, Costigan G, Crifo F, Cross NJG, Crosta M, Crowley C, Dafonte C, Damerdjy Y, Dapergolas A, David P, David M, De Cat P, de Felice F, de Laverny P, De Luise F, De March R, de Martino D, de Souza R, Debosscher J, del Pozo E, Delbo M, Delgado A, Delgado HE, di Marco F, Di Matteo P, Diakite S, Distefano E, Dolding C, Dos Anjos S, Drazinos P, Durán J, Dzigan Y, Ecale E, Edvardsson B, Enke H, Erdmann M, Escolar D, Espina M, Evans NW, Eynard Bontemps G, Fabre C, Fabrizio M, Faigler S, Falcão AJ, Farràs Casas M, Faye F, Federici L, Fedorets G, Fernández-Hernández J, Fernique P, Fienga A, Figueras F, Filippi F, Findeisen K, Fonti A, Fouesneau M, Fraile E, Fraser M, Fuchs J, Furnell R, Gai M, Galletti S, Galluccio L, Garabato D, García-Sedano F, Garé P, Garofalo A, Garralda N, Gavras P, Gerssen J, Geyer R, Gilmore G, Girona S, Giuffrida G, Gomes M, González-Marcos A, González-Núñez J, González-Vidal JJ, Granvik M, Guerrier A, Guillout P, Guiraud J, Gúrpide A, Gutiérrez-Sánchez R, Guy LP, Haigron R, Hatzidimitriou D, Haywood M, Heiter U, Helmi A, Hobbs D, Hofmann W, Holl B, Holland G, Hunt JAS, Hypki A, Icardi V, Irwin M, Jevardat de Fombelle G, Jofré P, Jonker PG, Jorissen A, Julbe F, Karamelas A, Kochoska A, Kohley R, Kolenberg K, Kontizas E, Koposov SE, Kordopatis G, Koubsky P, Kowalczyk A, Krone-Martins A, Kudryashova M, Kull I, Bachchan RK, Lacoste-Seris F, Lanza AF, Lavigne JB, Le Poncin-Lafitte C, Lebreton Y, Lebzelter T, Leccia S, Leclerc N, Lecoœur-Taïbi I, Lemaitre V, Lenhardt H, Leroux F, Liao S, Licata E, Lindstrøm HEP, Lister TA, Livanou E, Lobel A, Löffler W, López M, Lopez-Lozano A, Lorenz D, Loureiro T, MacDonald I, Magalhães Fernandes T, Managau S, Mann RG, Mantelet G, Marchal O, Marchant JM, Marconi M, Marie J, Marinoni S, Marrese PM, Marschalló G, Marshall DJ, Martín-Fleitas JM, Martino M, Mary N, Matijević G, Mazeh T, McMillan PJ, Messina S, Mestre A, Michalik D, Millar NR, Miranda BMH, Molina D, Molinaro R, Molinaro M, Molnár L, Moniez M, Montegriffo P, Monteiro D, Mor R, Mora A, Morbidelli R, Morel T, Morgenthaler S, Morley T, Morris D, Mulone AF, Muraveva T, Musella I, Narbonne J, Nelemans G, Nicastro L, Noval L, Ordénovic C, Ordieres-Meré J, Osborne P, Pagani C, Pagano I, Pailler F, Palacin H, Palaversa L, Parsons P, Paulsen T, Pecoraro M, Pedrosa R, Pentikäinen H, Pereira J, Pichon B, Piersimoni AM, Pineau FX, Plachy E, Plum G, Poujoulet E, Prša A, Pulone L, Ragaini S, Rago S, Rambaux N, Ramos-Lerate M, Ranalli P, Rauw G, Read A, Regibo S, Renk F, Reylé C, Ribeiro RA, Rimoldini L, Ripepi V, Riva A, Rixon G, Roelens M, Romero-Gómez M, Rowell N, Royer F, Rudolph A, Ruiz-Dern L, Sadowski G, Sagristà Sellés T, Sahlmann J, Salgado J, Salguero E, Sarasso M, Savietto H, Schnorhk A, Schultheis M, Sciacca E, Segol M, Segovia JC, Segransan D, Serpell E, Shih IC, Smareglia R, Smart RL, Smith C, Solano E, Solitro F, Sordo R, Soria Nieto S, Souchay J, Spagna A, Spoto F, Stampa U, Steele IA, Steidelmüller H, Stephenson CA, Stoev H, Suess FF, Süveges M, Surdej J, Szabados L, Szegedi-Elek E, Tapiador D, Taris F, Tauran G, Taylor MB, Teixeira R, Terrett D, Tingley B, Trager SC, Turon C, Ulla A, Utrilla E, Valentini G, van Elteren A, Van Hemelryck E, van Leeuwen M, Varadi M, Vecchiato A, Veljanoski J, Via T, Vicente D, Vogt S, Voss H, Votruba V, Voutsinas S, Walmsley G, Weiler M, Weingrill K, Werner D, Wevers T, Whitehead G, Wyrzykowski

- L, Yoldas A, Žerjal M, Zucker S, Zurbach C, Zwitter T, Alecu A, Allen M, Allende Prieto C, Amorim A, Anglada-Escudé G, Arsenijevic V, Azaz S, Balm P, Beck M, Bernstein HH, Bigot L, Bijaoui A, Blasco C, Bonfigli M, Bono G, Boudreault S, Bressan A, Brown S, Brunet PM, Bunclark P, Buonanno R, Butkevich AG, Carret C, Carrion C, Chemin L, Chéreau F, Corcione L, Darmigny E, de Boer KS, de Teodoro P, de Zeeuw PT, Delle Luche C, Domingues CD, Dubath P, Fodor F, Frézouls B, Fries A, Fustes D, Fyfe D, Gallardo E, Gallegos J, Gardiol D, Gebran M, Gomboc A, Gómez A, Grux E, Gueguen A, Heyrovsky A, Hoar J, Iannicola G, Isasi Parache Y, Janotto AM, Joliet E, Jonckheere A, Keil R, Kim DW, Klagyivik P, Klar J, Knude J, Kochukhov O, Kolka I, Kos J, Kutka A, Lainey V, LeBouquin D, Liu C, Loreggia D, Makarov VV, Marseille MG, Martayan C, Martinez-Rubi O, Massart B, Meynadier F, Mignot S, Munari U, Nguyen AT, Nordlander T, Ocvirk P, O’Flaherty KS, Olias Sanz A, Ortiz P, Osorio J, Oszkiewicz D, Ouzounis A, Palmer M, Park P, Pasquato E, Peltzer C, Peralta J, Péturaud F, Pieniluoma T, Pigozzi E, Poels J, Prat G, Prod’homme T, Raison F, Rebordao JM, Riskey D, Rocca-Volmerange B, Rosen S, Ruiz-Fuertes MI, Russo F, Sembay S, Serraller Vizcaino I, Short A, Siebert A, Silva H, Sinachopoulos D, Slezak E, Soffel M, Sosnowska D, Straizys V, ter Linden M, Terrell D, Theil S, Tiede C, Troisi L, Tsalmantza P, Tur D, Vaccari M, Vachier F, Valles P, Van Hamme W, Veltz L, Virtanen J, Wallut JM, Wichmann R, Wilkinson MI, Ziaeeppour H, Zschocke S (2016) The Gaia mission. *aap* 595:A1, DOI 10.1051/0004-6361/201629272, 1609.04153
- Gandhi S, Madhusudhan N (2017) GENESIS: new self-consistent models of exoplanetary spectra. *mnras* 472(2):2334–2355, DOI 10.1093/mnras/stx1601, 1706.02302
- Glasse ACH, Bauwens E, Bouwman J, Detre Ö, Fischer S, Garcia-Marin M, Justannont K, Labiano A, Nakos T, Ressler M, Rieke G, Scheithauer S, Wells M, Wright GS (2010) The throughput and sensitivity of the JWST mid-infrared instrument. In: *Space Telescopes and Instrumentation 2010: Optical, Infrared, and Millimeter Wave*, Proc. SPIE, vol 7731, p 77310K, DOI 10.1117/12.856975
- Greene TP, Line MR, Montero C, Fortney JJ, Lustig-Yaeger J, Luther K (2016) Characterizing Transiting Exoplanet Atmospheres with JWST. *APJ* 817(1):17, DOI 10.3847/0004-637X/817/1/17, 1511.05528
- Huitson CM, Sing DK, Vidal-Madjar A, Ballester GE, Lecavelier des Etangs A, Désert JM, Pont F (2012) Temperature-pressure profile of the hot Jupiter HD 189733b from HST sodium observations: detection of upper atmospheric heating. *mnras* 422(3):2477–2488, DOI 10.1111/j.1365-2966.2012.20805.x, 1202.4721
- Irwin PGJ, Teanby NA, de Kok R, Fletcher LN, Howett CJA, Tsang CCC, Wilson CF, Calcutt SB, Nixon CA, Parrish PD (2008) The NEMESIS planetary atmosphere radiative transfer and retrieval tool. *jsrt* 109:1136–1150, DOI 10.1016/j.jsrt.2007.11.006
- Kelsall T, Weiland JL, Franz BA, Reach WT, Arendt RG, Dwek E, Freudenreich HT, Hauser MG, Moseley SH, Odegard NP, Silverberg RF, Wright EL (1998) The COBE Diffuse Infrared Background Experiment Search for the Cosmic Infrared Background. II. Model of the Interplanetary Dust Cloud. *APJ* 508(1):44–73, DOI 10.1086/306380, astro-ph/9806250
- Kreidberg L, Bean JL, Désert JM, Benneke B, Deming D, Stevenson KB, Seager S, Berta-Thompson Z, Seifahrt A, Homeier D (2014) Clouds in the atmosphere of the super-Earth exoplanet GJ1214b. *Nature* 505(7481):69–72, DOI 10.1038/nature12888, 1401.0022
- Line MR, Wolf AS, Zhang X, Knutson H, Kammer JA, Ellison E, Deroo P, Crisp D, Yung YL (2013) A Systematic Retrieval Analysis of Secondary Eclipse Spectra. I. A Comparison of Atmospheric Retrieval Techniques. *APJ* 775(2):137, DOI 10.1088/0004-637X/775/2/137, 1304.5561
- Louie DR, Deming D, Albert L, Bouma LG, Bean J, Lopez-Morales M (2018) Simulated JWST/NIRISS Transit Spectroscopy of Anticipated TESS Planets Compared to Select Discoveries from Space-based and Ground-based Surveys. *PASP* 130(986):044401, DOI 10.1088/1538-3873/aaa87b, 1711.02098
- Madhusudhan N, Lee KKM, Mousis O (2012) A Possible Carbon-rich Interior in Super-Earth 55 Cancri e. *APJ* 759(2):L40, DOI 10.1088/2041-8205/759/2/L40, 1210.2720
- Mandel K, Agol E (2002) Analytic Light Curves for Planetary Transit Searches. *The Astrophysical Journal* 580(2):L171–L175, DOI 10.1086/345520, 0210099v1

- Mayor M, Pepe F, Queloz D, Bouchy F, Rupprecht G, Lo Curto G, Avila G, Benz W, Bertaux JL, Bonfils X, Dall T, Dekker H, Delabre B, Eckert W, Fleury M, Gilliotte A, Gojak D, Guzman JC, Kohler D, Lizon JL, Longinotti A, Lovis C, Megevand D, Pasquini L, Reyes J, Sivan JP, Sosnowska D, Soto R, Udry S, van Kesteren A, Weber L, Weilenmann U (2003) Setting New Standards with HARPS. *The Messenger* 114:20–24
- Mollière P, van Boekel R, Bouwman J, Henning T, Lagage PO, Min M (2017) Observing transiting planets with JWST. Prime targets and their synthetic spectral observations. *aap* 600:A10, DOI 10.1051/0004-6361/201629800, 1611.08608
- Pascale E, Waldmann IP, MacTavish CJ, Papageorgiou A, Amaral-Rogers A, Varley R, Coudé du Foresto V, Griffin MJ, Ollivier M, Sarkar S, Spencer L, Swinyard BM, Tessenyi M, Tinetti G (2015) EChOSim: The Exoplanet Characterisation Observatory software simulator. *Experimental Astronomy* 40:601–619, DOI 10.1007/s10686-015-9471-0, 1406.3984
- Pascale E, Bezawada N, Barstow J, Beaulieu JP, Bowles N, Coudé Du Foresto V, Coustenis A, Decin L, Drossart P, Eccleston P, Encrenaz T, Forget F, Griffin M, Güdel M, Hartogh P, Heske A, Lagage PO, Leconte J, Malaguti P, Micela G, Middleton K, Min M, Moneti A, Morales J, Mugnai L, Ollivier M, Pace E, Papageorgiou A, Pilbratt G, Puig L, Rataj M, Ray T, Ribas I, Rocchetto M, Sarkar S, Selsis F, Taylor W, Tennyson J, Tinetti G, Turrini D, Vandenbussche B, Venot O, Waldmann I, Wolkenberg P, Wright G, Zapatero Osorio MR, Zingales T (2018) The ariel space mission. *Proceedings of SPIE - The International Society for Optical Engineering* 10698
- Pepper J, Stassun KG, Gaudi BS (2018) KELT: The Kilodegree Extremely Little Telescope, a Survey for Exoplanets Transiting Bright, Hot Stars, p 128. DOI 10.1007/978-3-319-55333-7_128
- Pollacco D, Skillen I, Collier Cameron A, Christian D, Irwin J, Lister T, Street R, West R, Clarkson W, Evans N, Fitzsimmons A, Haswell C, Hellier C, Hodgkin S, Horne K, Jones B, Kane S, Keenan F, Norton A, Wheatley P (2006) The wasp project and superwasp camera. *Astrophysics and Space Science* 304, DOI 10.1007/s10509-006-9124-x
- Puig L, Isaak K, Linder M, Escudero I, Crouzet PE, Walker R, Ehle M, Hübner J, Timm R, de Vogeleer B, Drossart P, Hartogh P, Lovis C, Micela G, Ollivier M, Ribas I, Snellen I, Swinyard B, Tinetti G, Eccleston P (2015) The phase 0/A study of the ESA M3 mission candidate EChO. *Experimental Astronomy* 40(2-3):393–425, DOI 10.1007/s10686-014-9419-9
- Rauer H, Catala C, Aerts C, Appourchaux T, Benz W, Brandeker A, Christensen-Dalsgaard J, Deleuil M, Gizon L, Goupil MJ, Güdel M, Janot-Pacheco E, Mas-Hesse M, Pagano I, Piotto G, Pollacco D, Santos C, Smith A, Suárez JC, Szabó R, Udry S, Adibekyan V, Alibert Y, Almenara JM, Amaro-Seoane P, Eiff MAV, Asplund M, Antonello E, Barnes S, Baudin F, Belkacem K, Bergemann M, Bihain G, Birch AC, Bonfils X, Boisse I, Bonomo AS, Borsa F, Brandão IM, Brocato E, Brun S, Burleigh M, Burston R, Cabrera J, Cassisi S, Chaplin W, Charpinet S, Chiappini C, Church RP, Csizmadia S, Cunha M, Damasso M, Davies MB, Deeg HJ, Díaz RF, Dreizler S, Dreyer C, Eggenberger P, Ehrenreich D, Eigmüller P, Erikson A, Farmer R, Feltzing S, de Oliveira Fialho F, Figueira P, Forveille T, Fridlund M, García RA, Giommi P, Giuffrida G, Godolt M, Gomes da Silva J, Granzer T, Grenfell JL, Grottsch-Noels A, Günther E, Haswell CA, Hatzes AP, Hébrard G, Hekker S, Helled R, Heng K, Jenkins JM, Johansen A, Khodachenko ML, Kislyakova KG, Kley W, Kolb U, Krivova N, Kupka F, Lammer H, Lanza AF, Lebreton Y, Magrin D, Marcos-Arenal P, Marrese PM, Marques JP, Martins J, Mathis S, Mathur S, Messina S, Miglio A, Montalbán J, Montalto M, Monteiro MJPF, Moradi H, Moravveji E, Mordasini C, Morel T, Mortier A, Nascimbeni V, Nelson RP, Nielsen MB, Noack L, Norton AJ, Ofir A, Oshagh M, Ouazzani RM, Pápics P, Parro VC, Petit P, Plez B, Poretti E, Quirrenbach A, Ragazzoni R, Raimondo G, Rainer M, Reese DR, Redmer R, Reffert S, Rojas-Ayala B, Roxburgh IW, Salmon S, Santerne A, Schneider J, Schou J, Schuh S, Schunker H, Silva-Valio A, Silvotti R, Skillen I, Snellen I, Sohl F, Sousa SG, Sozzetti A, Stello D, Strassmeier KG, Švanda M, Szabó GM, Tkachenko A, Valencia D, Van Grootel V, Vauclair SD, Ventura P, Wagner FW, Walton NA, Weingrill J, Werner SC, Wheatley PJ, Zwintz K (2014) The PLATO 2.0 mission. *Experimental Astronomy* 38(1-2):249–330, DOI 10.1007/s10686-014-9383-4, 1310.0696

- Rauscher BJ, Fox O, Ferruit P (2007) Detectors for the James Webb Space Telescope Near-Infrared Spectrograph I: Readout Mode, Noise Model, and Calibration Considerations. *Publications of the Astronomical Society of the Pacific* 119:768–786, DOI 10.1086/520887, 0706.2344
- Ricker GR, Vanderspek R, Winn J, Seager S, Berta-Thompson Z, Levine A, Villaseñor J, Latham D, Charbonneau D, Holman M, Johnson J, Sasselov D, Szentgyorgyi A, Torres G, Bakos G, Brown T, Christensen-Dalsgaard J, Kjeldsen H, Clampin M, Rinehart S, Deming D, Doty J, Dunham E, Ida S, Kawai N, Sato B, Jenkins J, Lissauer J, Jernigan G, Kaltenegger L, Laughlin G, Lin D, McCullough P, Narita N, Pepper J, Stassun K, Udry S (2016) The Transiting Exoplanet Survey Satellite. In: *Space Telescopes and Instrumentation 2016: Optical, Infrared, and Millimeter Wave*, Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, vol 9904, p 99042B, DOI 10.1117/12.2232071
- Sarkar S, Pascale E (2015) ExoSim: a novel simulator of exoplanet spectroscopic observations. In: *European Planetary Science Congress*, pp EPSC2015–187
- Sarkar S, Pascale E, Papageorgiou A, Johnson LJ, Waldmann I (2020) ExoSim: the Exoplanet Observation Simulator. *arXiv e-prints* arXiv:2002.03739, 2002.03739
- Seager S, Mallén-Ornelas G (2002) On the Unique Solution of Planet and Star Parameters from an Extrasolar Planet Transit Light Curve. *Tech. rep.*
- Seager S, Sasselov DD (2000) Theoretical transmission spectra during extrasolar giant planet transits. *The Astrophysical Journal* 537(2):916–921, DOI 10.1086/309088
- Sing DK, Fortney JJ, Nikolov N, Wakeford HR, Kataria T, Evans TM, Aigrain S, Ballester GE, Burrows AS, Deming D, Désert JM, Gibson NP, Henry GW, Huitson CM, Knutson HA, Lecavelier Des Etangs A, Pont F, Showman AP, Vidal-Madjar A, Williamson MH, Wilson PA (2016) A continuum from clear to cloudy hot-Jupiter exoplanets without primordial water depletion. *Nature* 529(7584):59–62, DOI 10.1038/nature16068, 1512.04341
- Tinetti G, Vidal-Madjar A, Liang MC, Beaulieu JP, Yung Y, Carey S, Barber RJ, Tennyson J, Ribas I, Allard N, Ballester GE, Sing DK, Selsis F (2007) Water vapour in the atmosphere of a transiting extrasolar planet. *Nature* 448(7150):169–171, DOI 10.1038/nature06002
- Tinetti G, Encrenaz T, Coustenis A (2013) Spectroscopy of planetary atmospheres in our galaxy. *A&A Rev.* 21:63, DOI 10.1007/s00159-013-0063-6
- Tinetti G, Drossart P, Eccleston P, Hartogh P, Heske A, Leconte J, Micela G, Ollivier M, Pilbratt G, Puig L, Turrini D, Vandenbussche B, Wolkenberg P, Pascale E, Beaulieu JP, Güdel M, Min M, Rataj M, Ray T, Ribas I, Barstow J, Bowles N, Coustenis A, Coudé du Foresto V, Decin L, Encrenaz T, Forget F, Friswell M, Griffin M, Lagage PO, Malaguti P, Moneti A, Morales JC, Pace E, Rocchetto M, Sarkar S, Selsis F, Taylor W, Tennyson J, Venot O, Waldmann IP, Wright G, Zingales T, Zapatero-Osorio MR (2016) The science of ARIEL (Atmospheric Remote-sensing Infrared Exoplanet Large-survey). In: *Space Telescopes and Instrumentation 2016: Optical, Infrared, and Millimeter Wave*, Proc. SPIE, vol 9904, p 99041X, DOI 10.1117/12.2232370
- Tinetti G, Drossart P, Eccleston P, Hartogh P, Heske A, Leconte J, Micela G, Ollivier M, Pilbratt G, Puig L, Turrini D, Vandenbussche B, Wolkenberg P, Beaulieu JP, Buchave LA, Ferus M, Griffin M, Guedel M, Justtanont K, Lagage PO, Machado P, Malaguti G, Min M, Nørgaard-Nielsen HU, Rataj M, Ray T, Ribas I, Swain M, Szabo R, Werner S, Barstow J, Burleigh M, Cho J, du Foresto VC, Coustenis A, Decin L, Encrenaz T, Galand M, Gillon M, Helled R, Morales JC, Muñoz AG, Moneti A, Pagano I, Pascale E, Piccioni G, Pinfield D, Sarkar S, Selsis F, Tennyson J, Triaud A, Venot O, Waldmann I, Waltham D, Wright G, Amiaux J, Augères JL, Berthé M, Bezawada N, Bishop G, Bowles N, Coffey D, Colomé J, Crook M, Crouzet PE, Da Peppo V, Sanz IE, Focardi M, Frericks M, Hunt T, Kohley R, Middleton K, Morgante G, Ottensamer R, Pace E, Pearson C, Stamper R, Symonds K, Rengel M, Renotte E, Ade P, Affer L, Alard C, Allard N, Altieri F, André Y, Arena C, Argyriou I, Aylward A, Baccani C, Bakos G, Banaszekiewicz M, Barlow M, Batista V, Bellucci G, Benatti S, Bernardi P, Bézard B, Blecka M, Bolmont E, Bonfond B, Bonito R, Bonomo AS, Brucato JR, Brun AS, Bryson I, Bujwan W, Casewell S, Charnay B, Pestellini CC, Chen G, Ciaravella A, Claudi R, Clédassou R, Damasso M, Damiano M, Danielski C, Deroo P, Di Giorgio AM, Dominik C, Doublier V, Doyle S, Doyon R, Drummond B, Duong B, Eales S, Edwards B, Farina

- M, Flaccomio E, Fletcher L, Forget F, Fossey S, Fränz M, Fujii Y, García-Piquer Á, Gear W, Geoffray H, Gérard JC, Gesa L, Gomez H, Graczyk R, Griffith C, Grodent D, Guarcello MG, Gustin J, Hamano K, Hargrave P, Hello Y, Heng K, Herrero E, Hornstrup A, Hubert B, Ida S, Ikoma M, Iro N, Irwin P, Jarchow C, Jaubert J, Jones H, Julien Q, Kameda S, Kerschbaum F, Kervella P, Koskinen T, Krijger M, Krupp N, Lafarga M, Landini F, Lellouch E, Leto G, Luntzer A, Rank-Lüftinger T, Maggio A, Maldonado J, Maillard JP, Mall U, Marquette JB, Mathis S, Maxted P, Matsuo T, Medvedev A, Miguel Y, Minier V, Morello G, Mura A, Narita N, Nascimbeni V, Nguyen Tong N, Noe V, Oliva F, Palte E, Palmer P, Pancrazzi M, Papageorgiou A, Parmentier V, Perger M, Petralia A, Pezzuto S, Pierrehumbert R, Pillitteri I, Piotto G, Pisano G, Prisinzano L, Radioti A, Réess JM, Rezac L, Rocchetto M, Rosich A, Sanna N, Santerne A, Savini G, Scandariato G, Sicardy B, Sierra C, Sindoni G, Skup K, Snellen I, Sobiecki M, Soret L, Sozzetti A, Stiepen A, Strugarek A, Taylor J, Taylor W, Terenzi L, Tessenyi M, Tsiasas A, Tucker C, Valencia D, Vasisht G, Vazan A, Vilardell F, Vinatier S, Viti S, Waters R, Wawer P, Wawrzaszek A, Whitworth A, Yung YL, Yurchenko SN, Osorio MRZ, Zellem R, Zingales T, Zwart F (2018) A chemical survey of exoplanets with ariel. *Experimental Astronomy* 46(1):135–209, DOI 10.1007/s10686-018-9598-x
- Tsiasas A, Waldmann IP, Zingales T, Rocchetto M, Morello G, Damiano M, Karpouzas K, Tinetti G, McKemmish LK, Tennyson J, Yurchenko SN (2018) A Population Study of Gaseous Exoplanets. *aj* 155:156, DOI 10.3847/1538-3881/aaaf75, 1704.05413
- Udalski A, Szymański MK, Szymański G (2015) OGLE-IV: Fourth Phase of the Optical Gravitational Lensing Experiment. *acta* 65(1):1–38, 1504.05966
- Waldmann IP, Tinetti G, Rocchetto M, Barton EJ, Yurchenko SN, Tennyson J (2015) Tau-REx I: A Next Generation Retrieval Code for Exoplanetary Atmospheres. *apj* 802(2):107, DOI 10.1088/0004-637X/802/2/107, 1409.2312
- Wheatley PJ, Pollacco DL, Queloz D, Rauer H, Watson CA, West RG, Chazelas B, Loudon TM, Walker S, Bannister N, Bento J, Burleigh M, Cabrera J, Eigmüller P, Erikson A, Genolet L, Goad M, Grange A, Jordán A, Lawrie K, McCormac J, Neveu M (2013) The Next Generation Transit Survey (NGTS). In: *European Physical Journal Web of Conferences*, European Physical Journal Web of Conferences, vol 47, p 13002, DOI 10.1051/epjconf/20134713002, 1302.6592
- Zellem RT, Swain MR, Cowan NB, Bryden G, Komacek TD, Colavita M, Ardila D, Roudier GM, Fortney JJ, Bean J, Line MR, Griffith CA, Shkolnik EL, Kreidberg L, Moses JJ, Showman AP, Stevenson KB, Wong A, Chapman JW, Ciardi DR, Howard AW, Kataria T, Kempton EMR, Latham D, Mahadevan S, Meléndez J, Parmentier V (2019) Constraining Exoplanet Metallicities and Aerosols with the Contribution to ARIEL Spectroscopy of Exoplanets (CASE). *PASP* 131(1003):094401, DOI 10.1088/1538-3873/ab2d54, 1906.02820
- Zhou Y, Apai D, Lew BWP, Schneider G (2017) A Physical Model-based Correction for Charge Traps in the Hubble Space Telescope’s Wide Field Camera 3 Near-IR Detector and Its Applications to Transiting Exoplanets and Brown Dwarfs. *AJ* 153(6):243, DOI 10.3847/1538-3881/aa6481, 1703.01301