

MAPPING CLASS GROUP OF MANIFOLDS WHICH LOOK LIKE 3-DIMENSIONAL COMPLETE INTERSECTIONS

MATTHIAS KRECK AND YANG SU

ABSTRACT. In this paper we compute the mapping class group of closed simply-connected 6-manifolds M which look like complete intersections, i. e. $H^2(M; \mathbb{Z}) = \mathbb{Z} \cdot x$ with $x^3 \neq 0$. We prove some algebraic properties of the mapping class group, like for example we compute the abelianization and the center. We show that modulo the center the mapping class group is residually finite and virtually torsion-free. We also study low dimensional homology groups. The results are very similar to the computation of the mapping class group of Riemann surfaces. We give generators of the mapping class group, and generators and relations for the subgroup acting trivially on $\pi_3(M)$.

1. INTRODUCTION

The mapping class group of Riemann surfaces plays a big role in different areas of mathematics in particular in algebraic geometry in the study of moduli spaces. The mapping class group of complex algebraic surfaces is at present inaccessible due to the mysteries of smooth 4-manifolds. In this paper we will determine the mapping class group of certain 6-manifolds including all complex 3-dimensional complete intersections.

The approach is similar to one of the methods for Riemann surfaces through the comparison with the Torelli group, the subgroup acting trivially on the cohomology ring. For the Torelli group a central tool is a homomorphism into an abelian group which for Riemann surfaces was defined by Johnson. In our situation we define a similar homomorphism. Surprisingly the kernel of this homomorphism which for Riemann surfaces is still not understood can be completely computed. So one has a complete picture in terms of extensions of groups.

Knowing this one can ask some obvious questions concerning the nature of the mapping class group. It turns out that the results for Riemann surfaces essentially generalize to our situation. We apply this to get some information about the cohomology of the mapping class group.

To work with groups generators and relations are very helpful. We give explicit generators in form of generalized Dehn twists, and for the Torelli-type subgroup acting trivially on $\pi_3(M)$ relations, too.

Recently Oscar Randal-Williams applied our results to the moduli space of hypersurfaces in $\mathbb{C}P^4$ and determined the image of the monodromy representation [RW23].

2. RESULTS

2.1. The setting. All manifolds are assumed to be smooth and oriented.

Definition 2.1. The group of isotopy classes of orientation preserving diffeomorphisms of a manifold M is denoted by $\text{MCG}(M)+$:

$$\text{MCG}(M) := \pi_0 \text{Diff}^+(M).$$

This is called the (oriented) **mapping class group**.

The kernel of the homomorphism $\rho : \text{MCG}(M) \rightarrow \text{Aut}(H^*(M; \mathbb{Z}))$ induced by the action on the cohomology ring is denoted by

$$\mathcal{I}(M) := \ker\{\rho : \text{MCG}(M) \rightarrow \text{Aut}(H^*(M; \mathbb{Z}))\}.$$

This is called the **Torelli group**.

These groups play a central role in the study of surfaces and related moduli spaces. In higher dimensions there is not very much known about these groups except for highly connected manifolds [Kre78]. In this paper we study the mapping class group of the following manifolds.

Definition 2.2. A **manifold which looks like a 3-dimensional complete intersection** is an oriented simply-connected closed 6-manifold M with $H_2(M; \mathbb{Z}) \cong \mathbb{Z}$ and $x^3 \neq 0$ for a generator $x \in H^2(M; \mathbb{Z})$.

This notion is motivated by the fact that all complex 3-dimensional complete intersections have this property. This follows from the Lefschetz Hyperplane Theorem which states that an n -dimensional complete intersection $X \subset \mathbb{C}P^{n+k}$ is simply-connected and the inclusion into $\mathbb{C}P^{n+k}$ induces an isomorphism in homology below degree n and is surjective in degree n .

From now on all manifolds are assumed to be of this type unless otherwise said.

The **degree** of M is defined by

$$d(M) := \langle x^3, [M] \rangle$$

where x is the integral generator of $H^2(M)$ that makes the degree positive.

Since x^3 is non-zero and we consider orientation preserving diffeomorphisms the action on $H^2(M; \mathbb{Z}) \cong \mathbb{Z}$ is trivial and so the Torelli group is the kernel of the action on $H^3(M; \mathbb{Z})$ equipped with the intersection form. Since the intersection form is skew symmetric it is isometric to the skew symmetric hyperbolic form. Once and for all we choose a symplectic basis for $H^3(M; \mathbb{Z})$ and identify the group of isometries with the symplectic group $\text{Sp}(b, \mathbb{Z})$, where $b = b_3(M)$, the third Betti number of M . So ρ is a homomorphism from $\text{MCG}(M)$ to $\text{Sp}(b, \mathbb{Z})$.

In addition to the cohomology ring characteristic classes are attached to a smooth manifold. In our case the second Stiefel Whitney class $w_2(M)$

shows up and is equivalent to M being spin ($w_2(M) = 0$) or non-spin. And the first Pontrjagin $p_1(M) \in H^4(M; \mathbb{Z})$ class has to be considered. By Poincaré duality $H^4(M; \mathbb{Z}) \cong \mathbb{Z}$ and since x^2 is non zero, there is a unique generator y , such that $x^2 = d(M)y$. It can be shown, when M is spin, $p_1(M)$ is divisible by 4, and we define $k(M)$ to be the integer such that $p_1(M) = 4k(M)y$. When M is non-spin, we define $k(M)$ to be the integer such that $p_1(M) = (4k(M) - d(M))y$. For details see section 3.1. Finally we abbreviate $H^3(M; \mathbb{Z})$ by H .

Following the study of the mapping class group of Riemann surfaces (see for example [FaMa12]) we determine the mapping class group in terms of two types of short exact sequences. The first type of exact sequences starts with the action on the cohomology ring, whereas the second begins instead with the action on $\pi_3(M)$.

2.2. Computation of the mapping class group via the Torelli group.

The Torelli group $\mathcal{I}(M)$ is the kernel of ρ and we prove that ρ is surjective. Denote by $\Delta_{d(M), k(M)}$ the subgroup $\{(d(M)u, k(M)u) \mid u \in H\} \subset H \times H$. Next we construct a homomorphism

$$v_{x,p} : \mathcal{I}(M) \rightarrow (H \times H) / \Delta_{d(M), k(M)},$$

which we call the **variation of the cup product and the Pontrjagin class**, and denote its kernel by $\mathcal{K}(M)$. We read off $v_{x,p}(f)$ from the Pontrjagin class and the ring structure of the mapping torus M_f . For details of the construction we refer to section 4.2. We note that the variation of the cup product and the Pontrjagin class plays the role of the Johnson homomorphism for the mapping class group of surfaces.

Theorem 2.3. *Let M be a manifold which looks like a 3-dimensional complete intersection. Then there are short exact sequences*

$$(2.1) \quad 1 \rightarrow \mathcal{I}(M) \rightarrow \text{MCG}(M) \xrightarrow{\rho} \text{Sp}(b, \mathbb{Z}) \rightarrow 1$$

$$(2.2) \quad 1 \rightarrow \mathcal{K}(M) \rightarrow \mathcal{I}(M) \xrightarrow{v_{x,p}} (H \times H) / \Delta_{d(M), k(M)} \rightarrow 0.$$

The group $\mathcal{K}(M)$ only depends on whether M is spin or not and $d(M)$ and $p(M)$. In Theorem 5.1 we give tables of the groups $\mathcal{K}(M)$ in terms of $d(M)$ and $p(M)$. In the spin case it is isomorphic to a subgroup of $\mathbb{Z}/2 \times \mathbb{Z}/4 \times \mathbb{Z}/84$, all subgroups except the trivial subgroup and $(\mathbb{Z}/2)^3$, $(\mathbb{Z}/2)^3 \times \mathbb{Z}/3$, $(\mathbb{Z}/2)^3 \times \mathbb{Z}/7$, $(\mathbb{Z}/2)^3 \times \mathbb{Z}/21$ appear; in the non-spin case it is a subgroup of $\mathbb{Z}/168$, and all subgroups except the trivial one appear.

The group $\mathcal{K}(M)$ is the center of $\text{MCG}(M)$.

The determination of $\mathcal{K}(M)$ is the heart of our work.

From Theorem 2.3 we see that the Torelli group $\mathcal{I}(M)$ is an infinite group when $b_3(M) > 0$. This applies especially to 3-dimensional complex complete intersections and hence contradicts [Ver13, Theorem 3.4]. Another class of counterexamples with second Betti number ≥ 2 was found by R. Hain [Hai19]. For more general results on this issue see [KrSu19].

Looking at the exact sequences (2.1) and (2.2) above, one might wonder if one can combine them into one. In the case of surfaces Morita [Mor93] has made a step into that direction. He has combined the action on the first homology and Johnson homomorphism to a homomorphism

$$\mathrm{MCG}(F_g) \rightarrow \left(\frac{1}{2} \wedge^3 H^1(F_g)/H^1(F_g)\right) \rtimes \mathrm{Sp}(2g, \mathbb{Z})$$

whose kernel is equal to the kernel of the Johnson homomorphism and its image has finite index. In our situation we have a similar but stronger result.

Theorem 2.4. *Let M be a manifold which looks like a 3-dimensional complete intersection with 3rd Betti number b . When $b = 2$, or when $b \geq 4$ with M nonspin and $d(M)/2$ odd¹, there is a surjective homomorphism*

$$\tau: \mathrm{MCG}(M) \rightarrow (H \times H)/\Delta_{d(M), k(M)} \rtimes \mathrm{Sp}(b, \mathbb{Z})$$

whose kernel is $\mathcal{K}(M)$.

In the other cases there is a homomorphism

$$\tau: \mathrm{MCG}(M) \rightarrow (H \times H)/\Delta_{d(M), 2k(M)} \rtimes \mathrm{Sp}(b, \mathbb{Z})$$

whose image is

$$\{([a, b], A) \mid a, b \in H, A \in \mathrm{Sp}(b, \mathbb{Z}), b \equiv s(A) \pmod{2}\},$$

whose kernel is $\mathcal{K}(M)$, and where $s: \mathrm{Sp}(b, \mathbb{Z}) \rightarrow H \otimes \mathbb{Z}/2$ is a 1-cocycle which we give in Theorem 4.7.

For example we determine the mapping class group of the quintic X_5 in \mathbb{CP}^4 defined by the Fermat equation

$$z_0^5 + z_1^5 + \cdots + z_n^5 = 0,$$

which is of particular interest, last not least because it is a Calabi Yau manifold. Note that the 3rd Betti number of X_5 is 204.

Theorem 2.5. *Let X_5 be the quintic hypersurface in \mathbb{CP}^4 . The center of $\mathrm{MCG}(X_5)$ is $\mathcal{K}(X_5)$, which is isomorphic to $\mathbb{Z}/2$. The group $\mathrm{MCG}(X_5)/\mathcal{K}(X_5)$ is a subgroup of the semi-direct product $(H \times (H/5)) \rtimes \mathrm{Sp}(204, \mathbb{Z})$ of index 2^{204} , where the action of symplectic group on the abelian group is the tautological one on both factors. More precisely,*

$$\mathrm{MCG}(X_5)/\mathcal{K}(X_5) \cong \{((a, b), A) \mid a \equiv s(A) \pmod{2}\} \subset (H \times (H/5)) \rtimes \mathrm{Sp}(204, \mathbb{Z})$$

where $s: \mathrm{Sp}(204, \mathbb{Z}) \rightarrow H/2$ is a 1-cocycle given in Theorem 4.7.

The nontrivial element in the center is represented by a diffeomorphism supported in a closed disk D^6 . We will give an explicit formula of this diffeomorphism in §2.5.

A celebrated theorem of Sullivan [Sul77, Theorem 13.3] says that the mapping class group of a simply-connected closed smooth manifold of dimension > 5 is commensurable to an arithmetic group. In our situation one may see this in view of Theorem 2.4 and 2.3.

¹The degree of a non-spin manifold is always even, see the footnote in section 3.1.

2.3. Computation of the mapping class group via the special Torelli group. The Torelli group is the kernel of the action on $H_3(M)$ (which by Poincaré duality is isomorphic to $H^3(M)$). Instead one can consider the action on $\pi_3(M)$. In our situation the Hurewicz homomorphism h is surjective with kernel isomorphic to $\mathbb{Z}/d(M)$ (see section 3.1)

$$0 \rightarrow \mathbb{Z}/d(M) \rightarrow \pi_3(M) \xrightarrow{h} H_3(M) \rightarrow 0.$$

Via the Hurewicz homomorphism we can define the intersection form λ on $\pi_3(M)$, it's radical is the kernel of the Hurewicz map.

An orientation preserving diffeomorphism of M induces an isometry of $\pi_3(M)$ which is the identity on the kernel of the Hurewicz homomorphism (see the paragraph below (3.2)). Denote the group of such isometries by $\text{Sp}(\pi_3(M))$ and the kernel of this action by $\mathcal{J}(M)$, a subgroup of $\mathcal{I}(M)$, which we call the **special Torelli group**:

$$(2.3) \quad \mathcal{J}(M) := \ker\{\text{MCG}(M) \rightarrow \text{Sp}(\pi_3(M))\}.$$

It is easy to see that the group $\text{Sp}(\pi_3(M))$ is a split extension

$$0 \rightarrow H \otimes \mathbb{Z}/d(M) \rightarrow \text{Sp}(\pi_3(M)) \rightarrow \text{Sp}(b, \mathbb{Z}) \rightarrow 0.$$

On $\mathcal{J}(M)$ we define an invariant v_p landing in H , which is derived from the first Pontrjagin class of the mapping torus and called the **variation of the Pontrjagin class** and in this context plays the role of the homomorphism $v_{x,p}$. The kernel of this invariant is again $\mathcal{K}(M)$.

So altogether we have a filtration of subgroups of $\text{MCG}(M)$:

$$\text{MCG}(M) \supset \mathcal{I}(M) \supset \mathcal{J}(M) \supset \mathcal{K}(M)$$

Theorem 2.6. *Let M be a manifold which looks like a 3-dimensional complete intersection. Then there are short exact sequences*

$$(2.4) \quad 1 \rightarrow \mathcal{J}(M) \rightarrow \text{MCG}(M) \rightarrow \text{Sp}(\pi_3(M)) \rightarrow 1$$

$$(2.5) \quad 1 \rightarrow \mathcal{K}(M) \rightarrow \mathcal{J}(M) \xrightarrow{v_p} H \rightarrow 0$$

We can give an explicit finite presentation of $\mathcal{J}(M)$. We will return to this point in §2.5.

Combine the exact sequences (2.2) and (2.5) we have a commutative diagram of exact sequences

$$\begin{array}{ccccccc}
 & 0 & & 0 & & & \\
 & \downarrow & & \downarrow & & & \\
 & \mathcal{K}(M) & \xrightarrow{=} & \mathcal{K}(M) & & & \\
 & \downarrow & & \downarrow & & & \\
 0 \longrightarrow & \mathcal{J}(M) & \longrightarrow & \mathcal{I}(M) & \longrightarrow & H/d(M) & \longrightarrow 0 \\
 & \downarrow v_p & & \downarrow v_{x,p} & & \downarrow = & \\
 0 \longrightarrow & H & \xrightarrow{i_2} & (H \times H)/\Delta_{d(M),k(M)} & \longrightarrow & H/d(M) & \longrightarrow 0 \\
 & \downarrow & & \downarrow & & & \\
 & 0 & & 0 & & &
 \end{array}$$

2.4. Properties of the mapping class group. Given Theorem 2.3 one can try to generalize some of the fundamental properties of the mapping class group of surfaces to our situation. In many cases this works, sometimes with obvious modifications which often involve the center. It's known that the mapping class group of surfaces is centerless when $g \geq 3$. In our situation, as stated in Theorem 2.3, the center $Z(\text{MCG}(M))$ of $\text{MCG}(M)$ equals to $\mathcal{K}(M)$.

The mapping class group of surfaces is perfect when $g \geq 3$. The same holds in our context in most cases with some exceptional cases where the abelianization is $\mathbb{Z}/2$.

Theorem 2.7. *When $b > 0$ the homomorphism $\text{MCG}(M) \rightarrow \text{Sp}(b, \mathbb{Z})$ induces an isomorphism $H_1(\text{MCG}(M)) \rightarrow H_1(\text{Sp}(b, \mathbb{Z}))$, except for the case M spin and d even, where there is a split short exact sequence*

$$0 \rightarrow \mathbb{Z}/2 \rightarrow H_1(\text{MCG}(M)) \rightarrow H_1(\text{Sp}(b, \mathbb{Z})) \rightarrow 0.$$

The group $H_1(\text{Sp}(b, \mathbb{Z}))$ is isomorphic to (c. f. [BCRR18, Appendix])

$b = 2$	$b = 4$	$b \geq 6$
$\mathbb{Z}/12$	$\mathbb{Z}/2$	0

Therefore by Theorem 2.7 and the above table $H_1(\text{MCG}(M))$ is known.

For the mapping class group of surfaces, the homomorphism $\text{MCG}(F_g) \rightarrow \text{Sp}(2g, \mathbb{Z})$ induces an isomorphism $H_2(\text{MCG}(F_g)) \rightarrow H_2(\text{Sp}(2g, \mathbb{Z}))$ and the latter is isomorphic to \mathbb{Z} when $g \geq 4$ ([FaMa12]). In our situation we are able to compute the second homology group with \mathbb{Q} -coefficients.

Theorem 2.8. *When $b > 0$ the homomorphism $\text{MCG}(M) \rightarrow \text{Sp}(b, \mathbb{Z})$ induces a surjective homomorphism $H_2(\text{MCG}(M)) \rightarrow H_2(\text{Sp}(b, \mathbb{Z}))$. When $b \geq 6$ we have $H_2(\text{MCG}(M); \mathbb{Q}) \cong \mathbb{Q}^2$.*

The study of the moduli spaces of high dimensional manifolds is currently an active research field. As an example, for M a hypersurface in \mathbb{CP}^4 , it's shown in [GaRW18, §5.3] that $H_2(B\text{Diff}(M); \mathbb{Q}) \cong \mathbb{Q}^4$. For manifolds which look like complete intersection, as a corollary of Theorem 2.8 we can provide some integral information.

Corollary 2.9. *When $b > 0$ there is a surjective homomorphism*

$$H_2(B\text{Diff}(M)) \rightarrow H_2(\text{Sp}(b, \mathbb{Z})).$$

Let $\text{Diff}_0(M)$ be the identity component of $\text{Diff}(M)$. Combining the computation in [GaRW18] with Theorem 2.8 we have an estimation of the second homology of $B\text{Diff}_0(M)$

Corollary 2.10. $\dim H_2(B\text{Diff}_0(M); \mathbb{Q}) \geq 2$.

The mapping class group of surfaces is virtually torsion free, i. e., there exists a torsion free subgroup of finite index, and is residually finite, which can be characterized by the property that for every non-trivial element there is a homomorphism to a finite group which maps this element non-trivially. In our situation, after dividing the center (which in the case of surfaces of genus ≥ 3 doesn't change anything) we obtain the same result:

Theorem 2.11. $\text{MCG}(M)/Z(\text{MCG}(M))$ is virtually torsion free and residually finite.

It's shown by Randal-Williams [RW23] that $\text{MCG}(M)$ is not always residually finite.

2.5. Generators and presentations. The celebrated theorem of Sullivan [Sul77, Theorem 13.3] says that the mapping class group of simply-connected closed smooth manifolds of dimension > 5 is finitely presented. An obvious question is to describe the generators and relators of $\text{MCG}(M)$. In the case of surfaces generators are given by a finite number of Dehn twists and this is a very useful fact, since these are very concrete diffeomorphisms. To describe generators of $\text{MCG}(M)$ in our situation we need a generalization of Dehn twists about 3-spheres.

2.5.1. Dehn twists. A Dehn twist on a surface is given by an embedding of $S^1 \times [0, 1]$ into the surface and the corresponding Dehn twist is the identity outside this embedding and there it is given by rotating the circle $S^1 \times t$ by the angle $2\pi t$. Passing from $S^1 \times [0, 1]$ to thickenings of higher dimensional spheres one obtains an obvious generalization of Dehn twist: let $S \subset M$ be an embedded k -sphere with trivial normal bundle in an n -dimensional manifold M , identify a tubular neighborhood of S with $S^k \times D^{n-k}$ using a trivialization of the normal bundle; let $\alpha \in \pi_{n-k}SO(k+1)$ be represented by a smooth map $f: D^{n-k} \rightarrow SO(k+1)$, which maps to I near ∂D^{n-k} . Then we define the **(generalized) Dehn twist**

$$\mathbf{t}_{S,\alpha}: M \rightarrow M$$

as the diffeomorphism which maps the complement of $S^k \times D^{n-k}$ by the identity and $(x, y) \in S^k \times D^{n-k}$ to $(f(y)x, y)$.

For Dehn twists about 3-spheres in 6-manifolds, the action on the 3rd homotopy group is given as follows.

Lemma 2.12. (*Picard-Lefschetz formula*) *Let S_1 and S_2 be embedded 3-spheres in M , S_1 intersects with S_2 transversely at one point. Denote $e_i = [S_i] \in \pi_3(M)$. Let $\alpha \in \pi_3 SO(4)$, $\mathbf{t}_{S_1, \alpha}$ be the corresponding Dehn twist. Then*

$$\mathbf{t}_{S_1, \alpha*}(e_1) = e_1, \quad \mathbf{t}_{S_1, \alpha*}(e_2) = \langle e(\alpha), [S^4] \rangle \cdot e_1 + e_2,$$

where $e(\alpha)$ is the Euler class of the 4-dimensional vector bundle over S^4 represented by α .

Proof. Note that the evaluation $\langle e(\alpha), [S^4] \rangle$ equals the image of α under

$$p_*: \pi_3 SO(4) \rightarrow \pi_3(S^3) = \mathbb{Z}$$

where $p: SO(4) \rightarrow S^3$ is the natural fiber projection. Then the behaviour of e_1 and e_2 under $\mathbf{t}_{S_1, \alpha*}$ is clear by inspection. \square

2.5.2. Generators. We distinguish generators of $\pi_3(SO(4))$ as follows. The fibration $SO(3) \rightarrow SO(4) \rightarrow S^3$ gives a split short exact sequence

$$0 \rightarrow \pi_3(SO(3)) \rightarrow \pi_3(SO(4)) \rightarrow \pi_3(S^3) \rightarrow 0$$

We denote the class given by the inclusion of the unit quaternions S^3 to $SO(4)$ by ϵ

$$\epsilon: S^3 \rightarrow SO(4)$$

and the element given by the projection $S^3 \rightarrow SO(3)$ composed with the inclusion by ρ

$$\rho: S^3 \rightarrow SO(3) \rightarrow SO(4).$$

The Euler class of vector bundle over S^4 with clutching function ϵ is 1, whereas the Euler class for ρ is 0, and the first Pontrjagin class for ρ is 4 (the minimal value with trivial Euler class).

In the decomposition $M = N \# (S^3 \times S^3)^{\#r}$ as in (3.1), let S_i and T_i correspond to $S^3 \times \{0\}$ and $\{0\} \times S^3$ in the i -th copy of $S^3 \times S^3$; their homology classes form a symplectic basis of the intersection form on $H_{\mathbb{Z}}$. Let Z be an embedded 3-sphere representing a generator of the torsion subgroup of $\pi_3(M)$ (see (3.2)), disjoint from S_i and T_i . Now we define the Dehn twists for $1 \leq i \leq r$

$$\mathbf{a}_i = \mathbf{t}_{S_i, \rho}, \quad \mathbf{b}_i = \mathbf{t}_{T_i, \rho}, \quad \mathbf{e}_i = \mathbf{t}_{S_i, \epsilon}, \quad \mathbf{f}_i = \mathbf{t}_{T_i, \epsilon}, \quad \mathbf{z} = \mathbf{t}_{Z, \rho}.$$

Let $T_i \# T_{i+1}$ ($1 \leq i \leq r-1$), $S_i \# Z$ and $T_i \# Z$ be the connected sum of the corresponding spheres, define

$$\mathbf{f}_{i, i+1} = \mathbf{t}_{T_i \# T_{i+1}, \epsilon}, \quad \mathbf{g}_i = \mathbf{t}_{S_i \# Z, \epsilon}, \quad \mathbf{h}_i = \mathbf{t}_{T_i \# Z, \epsilon}$$

Theorem 2.13. *Let M be a manifold which looks like a 3-dimensional complete intersection with $b_3(M) > 0$. When M is spin and $d(M)$ is even, the mapping class group $\text{MCG}(M)$ is generated by $7r$ Dehn twists*

$$\mathbf{z}, \mathbf{a}_i, \mathbf{b}_i, \mathbf{e}_i, \mathbf{f}_i, \mathbf{g}_i, \mathbf{h}_i, \mathbf{f}_{i,i+1}.$$

In the other cases $\text{MCG}(M)$ is generated by $7r - 1$ Dehn twists

$$\mathbf{a}_i, \mathbf{b}_i, \mathbf{e}_i, \mathbf{f}_i, \mathbf{g}_i, \mathbf{h}_i, \mathbf{f}_{i,i+1}.$$

When $b_3(M) = 0$, the case is a little different. We need a generator of different type. Fix an embedding of the 6-disc D^6 in M , we may extend a diffeomorphism of D^6 which is the identity near the boundary to a diffeomorphism of M by the identity in the complement. We call these **disk supported diffeomorphisms** — diffeomorphisms which are the identity outside a disc in M . Denote the group of isotopy classes of diffeomorphisms on D^6 which are the identity near the boundary by $\text{MCG}(D^6, \partial)$, then above construction gives a homomorphism $\text{MCG}(D^6, \partial) \rightarrow \text{MCG}(M)$. The group $\text{MCG}(D^6, \partial)$ is isomorphic to $\text{MCG}(S^6)$, which in turn by the glueing construction is isomorphic to the group Θ_7 of oriented homotopy 7-spheres. The group Θ_7 is isomorphic to $\mathbb{Z}/28$ [KeMi63], a generator can be taken as the boundary of the E_8 -plumbing manifold. A diffeomorphism $f_c: D^6 \rightarrow D^6$ corresponding to this generator is constructed as follows [Las65].

Represent a generator of $\pi_3 SO(3)$ by a smooth compactly supported function $\alpha: \mathbb{R}^3 \rightarrow SO(3)$ and define the following diffeomorphisms of $\mathbb{R}^6 = \mathbb{R}^3 \times \mathbb{R}^3$,

$$\begin{aligned} F_1: \mathbb{R}^3 \times \mathbb{R}^3 &\rightarrow \mathbb{R}^3 \times \mathbb{R}^3, (x, y) \mapsto (x, \alpha(x)y) \\ F_2: \mathbb{R}^3 \times \mathbb{R}^3 &\rightarrow \mathbb{R}^3 \times \mathbb{R}^3, (x, y) \mapsto (\alpha(y)x, y) \end{aligned}$$

and

$$f_c = F_1 \circ F_2 \circ F_1^{-1} \circ F_2^{-1}.$$

It follows that f_c is compactly supported and so can be viewed as a diffeomorphism of D^6 which is the identity near the boundary. We denote by \mathbf{f}_c the extended diffeomorphism of M .

Recall that Z is an embedded 3-sphere representing a generator of $\pi_3(M) \cong \mathbb{Z}/d$, $\mathbf{t}_{Z,\epsilon}$ and $\mathbf{t}_{Z,\rho}$ are Dehn twists about Z with parameter ϵ and ρ respectively.

Theorem 2.14. *Let M be a manifold which looks like a 3-dimensional complete intersection with $b_3(M) = 0$. When M is spin, $\text{MCG}(M)$ is generated by \mathbf{f}_c , $\mathbf{t}_{Z,\epsilon}$ and $\mathbf{t}_{Z,\rho}$; when M is non-spin, $\text{MCG}(M)$ is generated by $\mathbf{t}_{Z,\rho}$.*

We don't have a finite list of relations between the generators. But we can do this for the group $\mathcal{J}(M)$, which is the kernel of the representation of $\text{MCG}(M)$ on $\text{Sp}(\pi_3(M))$ (see (2.3)). Let $\mathbf{u} = \mathbf{t}_{Z,\rho}$ be the Dehn twists about Z with parameter ρ .

Theorem 2.15. *A presentation of $\mathcal{J}(M)$ is given as follows:*

When M is spin, $\mathcal{J}(M)$ has generators \mathbf{f}_c , \mathbf{u} , \mathbf{z} , \mathbf{a}_i and \mathbf{b}_i ($1 \leq i \leq r$) and relations:

- (1) *all the commutators are trivial except for $[\mathbf{a}_i, \mathbf{b}_i] = \mathbf{f}_c$, $1 \leq i \leq r$;*
- (2) *$\mathbf{f}_c^r = \mathbf{u}^p = \mathbf{z}^q = 1$, where r , p and q is the order of the image of \mathbf{f}_c , \mathbf{u} and \mathbf{z} in $\mathcal{K}(M)$ respectively.*

When M is non-spin, $\mathcal{J}(M)$ has generators \mathbf{z} , \mathbf{a}_i and \mathbf{b}_i ($1 \leq i \leq r$) and relations:

- (1) *all the commutators are trivial except for $[\mathbf{a}_i, \mathbf{b}_i] = \mathbf{z}^6$, $1 \leq i \leq r$;*
- (2) *$\mathbf{z}^r = 1$, where r is the order of the image of \mathbf{z} in $\mathcal{K}(M)$.*

Structure of the paper. In section 3 we collect background knowledge of the diffeomorphism classification of manifolds which look like a 3-dimensional complete intersection, and of the mapping class group of $\#_r(S^3 \times S^3)$. In section 4 we prove the exact sequences in Theorem 2.3 and Theorem 2.4. Sections 5-8 are the core of the paper, in which we prove our main theorems about the kernel $\mathcal{K}(M)$ in the spin case: In section 5 we define the generalized Kreck-Stolz invariant on $\mathcal{K}(M, D)$. In section 6 we compute the mapping class group of $S^2 \times S^4$ and $(S^2 \times D^4, \partial)$ in terms of the generalized Kreck-Stolz invariant. In section 7 we show that the extension homomorphism $\text{MCG}(S^2 \times D^4, \partial) \rightarrow \mathcal{K}(M, D)$ is surjective and use it to compute the group $\mathcal{K}(M, D)$, leading to the table in Theorem 5.1. In section 8 we show that the forgetful homomorphism $\mathcal{K}(M, D) \rightarrow \mathcal{K}(M)$ is an isomorphism, finishing the proof of Theorem 5.1. In section 9 we explain the modifications needed for the non-spin case. The proofs of the group theoretic properties of $\text{MCG}(M)$ are in section 10. The computation of relevant cobordism groups is in the appendix.

Acknowledgements. We would like to thank Benson Farb, Daniel Huybrechts, Manuel Krannich, and Eduard Looijenga for helpful comments and Richard Hain and Oscar Randal-Williams for pointing out errors and suggestions how to correct them, and a referee for very valuable suggestions concerning the presentation.

3. PRELIMINARIES

3.1. Diffeomorphism classification. Based on the work of Wall [Wal66] and Jupp [Jup73], there is a complete diffeomorphism classification of manifolds which look like 3-dimensional complete intersections. This classification is the starting point of our computation of their mapping class groups. In this subsection we summarize the classification and invariants of these manifolds.

A manifold which looks like 3-dimensional complete intersections has a connected sum decomposition

$$(3.1) \quad M = N \# (S^3 \times S^3)^{\#r},$$

where N is simply connected with $H^*(N)$ isomorphic to $H^*(\mathbb{CP}^3)$ as groups, and $2r = b_3(M)$ is the 3rd Betti number of M . By standard algebraic

topology there is an exact sequence (of $\text{MCG}(M)$ -modules)

$$(3.2) \quad 0 \rightarrow \mathbb{Z}/d(M) \rightarrow \pi_3(M) \xrightarrow{h} H_3(M) \rightarrow 0.$$

where h is the Hurewicz homomorphism. A generator of the subgroup \mathbb{Z}/d is represented by the composite of the Hopf map $S^3 \rightarrow S^2$ and a map $S^2 \rightarrow M$ representing a generator of $\pi_2(M) \cong H_2(M)$. Hence \mathbb{Z}/d is a trivial $\text{MCG}(M)$ -module.

By Poincaré duality, $y := x^2/d(M)$ is an integral generator of $H^4(M)$. When M is spin, the first Pontrjagin class $p_1(M)$ of M can be written as

$$(3.3) \quad p_1(M) = 4k(M)y$$

with $k = 6l(M) + d(M)$, and $l(M)$ can take all integers. When M is non-spin, d is an even integer² and we can write

$$(3.4) \quad p_1(M) = (4k(M) - d)y$$

with $p = 12l(M) + d(M)/2$, $l(M)$ takes all integers. The diffeomorphism type of M is determined by the property M being spin/nonspin and the integers r (or b), $d(M)$ and $k(M)$ (or $l(M)$).

For a 3-dimensional complete intersection X of multi-degree (d_1, \dots, d_m) , all these invariants can be calculated from the equation of vector bundles

$$TX \oplus \gamma^{\otimes d_1} \oplus \dots \oplus \gamma^{\otimes d_m} = i^*T\mathbb{CP}^{3+m}$$

where γ is the tautological line bundle over \mathbb{CP}^{3+m} and $i: X \rightarrow \mathbb{CP}^{3+m}$ the inclusion map. For example, if X is a hypersurface of degree d , then

$$(3.5) \quad \begin{aligned} d(X) &= d, \quad w_2(X) \equiv 5 - d \pmod{2}, \quad p_1(X) = (5 - d^2)dy \\ r &= (d^4 - 5d^3 + 10d^2 - 10d + 4)/2 \end{aligned}$$

3.2. Mapping class group of $\#_r(S^3 \times S^3)$. In [Kre78] the mapping class group of $(k-1)$ -connected almost-parallelizable $2k$ -manifolds is studied. For our purpose in this section we recall the result for the connected sum of r copies of $S^3 \times S^3$, denoted by $\#_r(S^3 \times S^3)$.

The intersection form

$$I: H_3(\#_r(S^3 \times S^3)) \times H_3(\#_r(S^3 \times S^3)) \rightarrow \mathbb{Z}$$

is a skew-symmetric unimodular bilinear form over \mathbb{Z} . After taking a symplectic basis, the isometry group of this intersection form is identified with $\text{Sp}(2r, \mathbb{Z})$. An orientation preserving diffeomorphism $f: \#_r(S^3 \times S^3) \rightarrow \#_r(S^3 \times S^3)$ induces an isometry of I . Therefore there is a homomorphism

$$\text{MCG}(\#_r(S^3 \times S^3)) \rightarrow \text{Sp}(2r, \mathbb{Z}).$$

Denote the kernel of this homomorphism by $\mathcal{I}(\#_r(S^3 \times S^3))$, we define a homomorphism

$$\chi: \mathcal{I}(\#_r(S^3 \times S^3)) \rightarrow H^3(\#_r(S^3 \times S^3))$$

²Let x_2 and y_2 be the mod 2 reduction of x and y respectively, if d is odd, then $y_2 = x_2^2$ and by the Wu formula $w_2(M) \cup y_2 = \text{Sq}^2 y_2 = \text{Sq}^2(x_2^2) = 0$. This shows that M is spin.

as follows. Let $[f] \in \mathcal{I}(\#_r(S^3 \times S^3))$, denote the mapping torus by $(\#_r(S^3 \times S^3))_f$, consider the collapsing map

$$c: (\#_r(S^3 \times S^3))_f \rightarrow (\#_r(S^3 \times S^3))_f / \{0\} \times \#_r(S^3 \times S^3) = \Sigma(\#_r(S^3 \times S^3)^+)$$

If $f_* = \text{id}$ on homology, then

$$c^*: H^4(\Sigma(\#_r(S^3 \times S^3)^+)) \rightarrow H^4(\#_r(S^3 \times S^3))_f$$

is an isomorphism. Then $\chi(f)$ is defined to be $\frac{1}{4}p'(f)$, where $p'(f)$ is the desuspension of $(c^*)^{-1}(p_1(\#_r(S^3 \times S^3))_f)$. The geometric meaning of this invariant is the following: let $x \in H_3(\#_r(S^3 \times S^3))$ be represented by an embedded S^3 , which is fixed pointwisely by f , since the stable normal bundle of the embedded S^3 is trivial, the differential df induces a map $S^3 \rightarrow SO(4)$, giving an element in $\pi_3 SO(4)$. This element lies in the image of the suspension map $\pi_3 SO(3) \rightarrow \pi_3 SO(4)$. Identify this group with \mathbb{Z} , then this element equals $\langle \chi(f), x \rangle$ (see [Kre78, Lemma 2], and a correction of a factor 2 by [Cro11, Lemma 1.6]).

Theorem 3.1. [Kre78, Theorem 2] *There are exact sequences*

$$1 \rightarrow \mathcal{I}(\#_r(S^3 \times S^3)) \rightarrow \text{MCG}(\#_r(S^3 \times S^3)) \rightarrow \text{Sp}(2r, \mathbb{Z}) \rightarrow 1$$

$$0 \rightarrow \Theta_7 \rightarrow \mathcal{I}(\#_r(S^3 \times S^3)) \xrightarrow{\chi} H^3(\#_r(S^3 \times S^3)) \rightarrow 1$$

Remark 3.2. If we consider the quotient by the subgroup Θ_7 , then we get a group extension with abelian kernel

$$0 \rightarrow H^3(\#_r(S^3 \times S^3)) \rightarrow \text{MCG}(\#_r(S^3 \times S^3)) / \Theta_7 \rightarrow \text{Sp}(2r, \mathbb{Z}) \rightarrow 1$$

in which the action of $\text{Sp}(2r, \mathbb{Z})$ on $H^3(\#_r(S^3 \times S^3))$ is the standard linear action. This extension problem is studied in [Cro11]: it is split if and only if $r = 1$; for $r \geq 2$ the cohomology class in $H^2(\text{Sp}(2r, \mathbb{Z}), H^3(\#_r(S^3 \times S^3)))$ corresponding to this extension is described in [Cro11, Corollary 3.5].

4. THE TORELLI GROUP $\mathcal{I}(M)$

Convention. Recall that in section 3.1 we attach numerical invariants $d(M)$, $k(M)$ and $l(M)$ to the manifold M . From now on we simply write these numbers as d , k and l , respectively.

4.1. surjectivity of ρ . Recall we have a decomposition (3.1)

$$M = N \# (S^3 \times S^3)^{\#r}.$$

Given a diffeomorphism of $\#_r(S^3 \times S^3)$ which is the identity on an embedded disk D , we extend it to a diffeomorphism of M via the identity on the complement. Denote the group of isotopy classes of diffeomorphisms of $\#_r(S^3 \times S^3)$ which are the identity on D by $\text{MCG}(\#_r(S^3 \times S^3), D)$, we have a homomorphism $\text{MCG}(\#_r(S^3 \times S^3), D) \rightarrow \text{MCG}(M)$. By [Kre78, Lemma 3] the natural homomorphism $\text{MCG}(\#_r(S^3 \times S^3), D) \rightarrow \text{MCG}(\#_r(S^3 \times S^3))$ is

an isomorphism. Therefore we have a homomorphism $\text{MCG}(\#_r(S^3 \times S^3)) \rightarrow \text{MCG}(M)$ which clearly fits into the following commutative diagram

$$(4.1) \quad \begin{array}{ccc} \text{MCG}(\#_r(S^3 \times S^3)) & \longrightarrow & \text{Sp}(b, \mathbb{Z}) \\ \downarrow & & \downarrow = \\ \text{MCG}(M) & \xrightarrow{\rho} & \text{Sp}(b, \mathbb{Z}) \end{array}$$

By Theorem 3.1 the homomorphism in the upper row is surjective, therefore $\text{MCG}(M) \rightarrow \text{Sp}(b, \mathbb{Z})$ is surjective. This is the proof of the exact sequence (2.1) in Theorem 2.3.

4.2. variation of the cup product and the Pontrjagin class.

4.2.1. *definition of the variation map.* Next we define a homomorphism

$$\bar{v}_{x,p}: \mathcal{I}(M) \rightarrow (H^3(M) \times H^3(M))/\Delta_{d,2k}$$

where $\Delta_{d,2k} = \{(du, 2ku) \mid u \in H^3(M)\}$ is the diagonal of slope $(d, 2k)$, as follows.

Let $f: M \rightarrow M$ be a diffeomorphism such that $[f] \in \mathcal{I}(M)$,

$$M_f = M \times [0, 1]/(x, 0) \sim (f(x), 1)$$

be the mapping torus of f . Then M_f is an M -bundle over the circle. Let $i: M \rightarrow M_f$ be the inclusion of a fiber. Since f acts trivially on homology, the Wang sequence of the fiber bundle implies

- (1) $i^*: H^2(M_f) \rightarrow H^2(M)$ is an isomorphism;
- (2) there is a short exact sequence

$$(4.2) \quad 0 \rightarrow H^3(M) \xrightarrow{\delta} H^4(M_f) \xrightarrow{i^*} H^4(M) \rightarrow 0$$

Let $x \in H^2(M)$, $y \in H^4(M)$ be generators, such that $x^2 = dy$ and $d > 0$. Identifying $H^2(M)$ with $H^2(M_f)$ via i^* , we still denote the corresponding generator of $H^2(M_f)$ by x ; let $\bar{y} \in H^4(M_f)$ be a preimage of y .

The homomorphism $i^*: H^2(M_f; \mathbb{Z}/2) \rightarrow H^2(M; \mathbb{Z}/2)$ is an isomorphism, and $i^*w_2(M_f) = w_2(M)$. Therefore M_f is spin if and only if M is spin. Recall that for a spin vector bundle ξ , there is a spin first Pontrjagin class, denoted by $\frac{p_1}{2}(\xi)$, such that $2 \cdot \frac{p_1}{2}(\xi) = p_1(\xi)$. Therefore if M is spin, we have (see (3.3))

$$\frac{p_1}{2}(M) = 2ky;$$

if M is non-spin, then $TM \oplus \eta$ is a spin vector bundle, where η is the complex line bundle over M with $c_1(\eta) = x$. Since (see (3.4))

$$p_1(TM \oplus \eta) = p_1(M) + x^2 = p_1(M) + dy = 4ky$$

we have

$$\frac{p_1}{2}(TM \oplus \eta) = 2ky.$$

Therefore if M is spin, in the homomorphism $i^*: H^4(M_f) \rightarrow H^4(M)$, we have

$$(4.3) \quad i^*(x^2 - d\bar{y}) = x^2 - dy = 0, \quad i^*\left(\frac{p_1}{2}(M_f) - 2k\bar{y}\right) = \frac{p_1}{2}(M) - 2ky = 0$$

Therefore $x^2 - d\bar{y}$ and $\frac{p_1}{2}(M_f) - 2k\bar{y}$ have unique preimages under δ in $H^3(M)$

$$\delta^{-1}(x^2 - d\bar{y}) \in H^3(M), \quad \delta^{-1}\left(\frac{p_1}{2}(M_f) - 2k\bar{y}\right) \in H^3(M)$$

Note that \bar{y} is not unique, we may change it by $\bar{y} \rightarrow \bar{y} + \delta(u)$, $u \in H^3(M)$. The corresponding change of the preimages under δ follows from the equations

$$\begin{aligned} \delta^{-1}(x^2 - d(\bar{y} + \delta(u))) &= \delta^{-1}(x^2 - d\bar{y}) + du, \\ \delta^{-1}\left(\frac{p_1}{2}(M_f) - 2k(\bar{y} + \delta(u))\right) &= \delta^{-1}\left(\frac{p_1}{2}(M_f) - 2k\bar{y}\right) + 2ku \end{aligned}$$

Thus we have a well-defined map

$$\bar{v}_{x,p}: \mathcal{I}(M) \rightarrow (H^3(M) \times H^3(M))/\Delta_{d,2k}.$$

If M is non-spin, we just need to replace $\frac{p_1}{2}(M_f)$ in the second equation in (4.3) by $\frac{p_1}{2}(TM_f \oplus \bar{\eta})$, where $\bar{\eta}$ is the complex line bundle with $c_1(\bar{\eta}) = x$. Now since

$$i^*\left(\frac{p_1}{2}(TM_f \oplus \bar{\eta}) - 2k\bar{y}\right) = \frac{p_1}{2}(TM \oplus i^*\bar{\eta}) - 2ky = 0$$

we have an element $\delta^{-1}\left(\frac{p_1}{2}(TM_f \oplus \bar{\eta}) - 2k\bar{y}\right) \in H^3(M)$.

Lemma 4.1. *The map $\bar{v}_{x,p}: \mathcal{I}(M) \rightarrow (H^3(M) \times H^3(M))/\Delta_{d,2k}$ is a homomorphism.*

Proof. The exact sequence (4.2) is identified with

$$0 \rightarrow H^4(M_f, M) \rightarrow H^4(M_f) \rightarrow H^4(M) \rightarrow 0.$$

We construct M_{fg} from M_f by cutting $I \times M$ along $1/2 \times M$ and regluing it via g from the left to the right. Denote the two fibres over $1/4$ and $3/4$ by M_1 and M_2 . Consider the restriction map

$$H^4(M_{fg}, M_1 \cup M_2) \rightarrow H^4(M_{fg}, M_1).$$

We have an isomorphism

$$H^4(M_f, M) \oplus H^4(M_g, M) \xrightarrow{\cong} H^4(M_{fg}, M_1 \cup M_2).$$

If we identify $H^4(M_{fg}, M_1)$, $H^4(M_f, M)$ and $H^4(M_g, M)$ with $H^3(M)$, the homomorphism

$$H^4(M_f, M) \oplus H^4(M_g, M) \xrightarrow{\cong} H^4(M_{fg}, M_1 \cup M_2) \rightarrow H^4(M_{fg}, M_1)$$

corresponds to the homomorphism $H^3(M) \oplus H^3(M) \rightarrow H^3(M)$ given by the sum, and the corresponding cohomology classes for M_f and M_g are mapped to that of M_{fg} . This shows that $\bar{v}_{x,p}$ is a homomorphism. \square

4.2.2. *image of the variation map.* Next we determine the image of $\bar{v}_{x,p}$. Consider the Hurewicz homomorphism $h: \pi_3(M) \rightarrow H_3(M)$. Let $M \rightarrow \mathbb{CP}^\infty$ be the classifying map of the chosen generator $x \in H^2(M)$, denote the homotopy fiber by F . Then $\pi_3(M) \cong \pi_3(F) \cong H_3(F)$. The Gysin sequence of the fibration $S^1 \rightarrow F \rightarrow M$ gives an exact sequence

$$(4.4) \quad 0 \rightarrow H_4(M) \xrightarrow{\cap x} H_2(M) \rightarrow H_3(F) \rightarrow H_3(M) \rightarrow 0$$

From this we have a short exact sequence

$$0 \rightarrow \mathbb{Z}/d \rightarrow \pi_3(M) \xrightarrow{h} H_3(M) \rightarrow 0$$

This exact sequence is compatible with the action of f_* , and for $f \in \mathcal{I}(M)$, $f_* = \text{id}$ on $H_3(M)$ and \mathbb{Z}/d . The homomorphism $f_* - \text{id}: \pi_3(M) \rightarrow \pi_3(M)$ gives rise to a homomorphism $\beta: H_3(M) \rightarrow \mathbb{Z}/d$, which we view as an element in $H^3(M; \mathbb{Z}/d)$.

Lemma 4.2. *Under the composition*

$$\mathcal{I}(M) \xrightarrow{\bar{v}_{x,p}} (H^3(M) \times H^3(M))/\Delta_{d,2k} \xrightarrow{\text{pr}_1} H^3(M)/d = H^3(M; \mathbb{Z}/d)$$

the image of f is β . And this map is surjective.

Proof. We have a commutative diagram

$$\begin{array}{ccccc} F & \longrightarrow & M & \longrightarrow & \mathbb{CP}^\infty \\ \downarrow & & \downarrow & & \downarrow = \\ F_f & \longrightarrow & M_f & \longrightarrow & \mathbb{CP}^\infty \\ \downarrow & & \downarrow & & \\ S^1 & \xrightarrow{=} & S^1 & & \end{array}$$

with horizontal maps and vertical maps are fibrations. The Gysin sequences and Wang sequences fit together to commutative diagrams

$$\begin{array}{ccccccc} 0 & \longrightarrow & H^3(M) & \xrightarrow{\delta} & H^4(M_f) & \longrightarrow & H^4(M) \longrightarrow 0 \\ & & \downarrow i^* & & \downarrow & & \downarrow \\ 0 & \longrightarrow & H^3(F) & \xrightarrow{\delta} & H^4(F_f) & \longrightarrow & H^4(F) \longrightarrow 0 \end{array}$$

and

$$\begin{array}{ccccc}
H_5(M_f) & \longrightarrow & H_3(M_f) & \longrightarrow & H_4(F_f) \\
\downarrow & & \downarrow & \searrow & \downarrow \\
H_4(M) & \longrightarrow & H_2(M) & \xrightarrow{j_*} & H_3(F) \\
\downarrow & & \downarrow & \searrow & \downarrow f_* - \text{id} \\
H_4(M) & \longrightarrow & H_2(M) & \xrightarrow{j_*} & H_3(F) \\
\downarrow & & \downarrow \cong & \searrow & \downarrow \\
H_4(M_f) & \xrightarrow{x \cap -} & H_2(M_f) & \longrightarrow & H_3(F_f) \\
\downarrow \partial & \swarrow & & & \\
H_3(M) & & & &
\end{array}$$

In the above diagram the columns and the zigzagged rows are exact sequences.

Let $\alpha \in H^3(M)$ be the preimage of $x^2 - d\bar{y} \in H^4(M_f)$ under δ . For any $a \in H_3(M)$, let $\bar{a} \in H_3(F)$ be a preimage of a , since $(f_* - \text{id})(a) = 0$, there is a preimage $b \in H_2(M)$ of $(f_* - \text{id})(\bar{a})$ under j_* , view b as a class in $H_2(M_f)$, let $c \in H_4(M_f)$ be a preimage of b under $x \cap -$, then $\partial c = a$. We have

$$\langle \alpha, a \rangle = \langle \alpha, \partial c \rangle = \langle \delta \alpha, c \rangle = \langle x^2 - d\bar{y}, c \rangle \equiv \langle x^2, c \rangle = \langle x, x \cap c \rangle = \langle x, b \rangle \pmod{d}$$

This implies that the homomorphism $\beta: H_3(M) \rightarrow \mathbb{Z}/d$ coincides with the cohomology class $\delta^{-1}(x^2 - d\bar{y}) \in H^3(M; \mathbb{Z}/d)$.

Next we show the surjectivity. In the decomposition $M = N \#_r (S^3 \times S^3)$, let $N_0 = N - \text{int} D^6$, then it's easy to see that $\pi_3(N_0) \cong \pi_3(N) \cong \mathbb{Z}/d$, and $\pi_3(N_0) \rightarrow \text{tors} \pi_3(M)$ is an isomorphism, where $\text{tors} \pi_3(M)$ is the torsion subgroup of $\pi_3(M)$. Let $Z \subset N_0$ represent a generator of $\pi_3(N_0) \cong \mathbb{Z}/d$, $S_i = S^3 \times \{x_0\}$ and $T_i = \{x_0\} \times S^3$ in the i -th copy of $S^3 \times S^3$. Denote

$$[Z] = e_0, \quad [S_1] = e_1, \quad [T_1] = e_2 \in \pi_3(M)$$

Consider the Dehn twists $\mathbf{t}_{Z \# S_1, \alpha}$ and $\mathbf{t}_{S_1, -\alpha}$, where $\alpha \in \pi_3 SO(4)$ such that $\langle e(\alpha), [S^4] \rangle = 1$, where $e(\alpha)$ is the Euler class of the vector bundle over S^4 with clutching function α (for the notations of Dehn twists see section 2.5.1). Then by Lemma 2.12

$$\mathbf{t}_{Z \# S_1, \alpha*}(e_2) = e_0 + e_1 + e_2, \quad \mathbf{t}_{S_1, -\alpha*}(e_2) = e_2 - e_1.$$

The other generators of $\pi_3(M)$ are not changed. Then $(\mathbf{t}_{S_1, -\alpha} \circ \mathbf{t}_{Z \# S_1, \alpha})_* - \text{id}$ maps e_2 to e_0 , and other generators of $H_3(M)$ to 0. This finishes the proof of Lemma 4.2. \square

Remark 4.3. From Lemma 4.2 we see that the restriction of $\bar{v}_{x,p}$ on the subgroup of \mathcal{J}_M lifts to a homomorphism to $H^3(M; \mathbb{Z})$. In view of [Sul77, Corollary 13.3], in our situation $D = H^3(M; \mathbb{Q})$ and $I = 0$, and the lattice is $H^3(M; \mathbb{Z})$.

Next we consider the composition

$$\mathcal{I}(\#_r(S^3 \times S^3)) \rightarrow \mathcal{I}(M) \xrightarrow{\bar{v}_{x,p}} (H \times H)/\Delta_{d,2k}$$

If $f: M \rightarrow M$ is a diffeomorphism coming from $\#_r(S^3 \times S^3)$, then the mapping torus is

$$M_f = (N \times S^1) \#_{S^1} (\#_r(S^3 \times S^3))_f,$$

where $\#_{S^1}$ denotes the fiber connected-sum along S^1 . Under the decomposition

$$H^4(M_f) = H^4(N) \oplus H^4((\#_r(S^3 \times S^3))_f)$$

we may choose $\bar{y} = (y, 0)$, where $y = x^2/d \in H^4(N)$ is the canonical generator. Therefore in $H^4(M_f)$ we have the equation

$$x^2 - d\bar{y} = 0.$$

For the Pontrjagin class, in the spin case we have

$$\frac{p_1}{2}(M_f) = \left(\frac{p_1}{2}(N), \frac{p_1}{2}((\#_r(S^3 \times S^3))_f)\right),$$

therefore

$$\frac{p_1}{2}(M_f) - 2k\bar{y} = \frac{p_1}{2}((\#_r(S^3 \times S^3))_f).$$

In the non-spin case we have

$$\frac{p_1}{2}(TM_f \oplus \bar{\eta}) - 2k\bar{y} = \frac{p_1}{2}((\#_r(S^3 \times S^3))_f).$$

The coboundary map $\delta: H^3(M) \rightarrow H^4(M_f)$ in the Wang sequence (4.2) can be identified with

$$H^3(M) \xrightarrow{\Sigma} H^4(\Sigma(M^+)) \xrightarrow{c^*} H^4(M_f).$$

Identify $H^3(\#_r(S^3 \times S^3))$ with $H^3(M)$ and compare the homomorphism $\bar{v}_{x,p}$ with the invariant χ defined in §3.2, we have the following

Lemma 4.4. *There is a commutative diagram*

$$\begin{array}{ccc} \mathcal{I}_{\#_r(S^3 \times S^3)} & \xrightarrow{\chi} & H \\ \downarrow & & \downarrow 2 \cdot i_2 \\ \mathcal{I}(M) & \xrightarrow{\bar{v}_{x,p}} & (H \times H)/\Delta_{d,2k} \end{array}$$

where $2 \cdot i_2: \alpha \mapsto [0, 2\alpha]$.

Moreover, we have the following

Lemma 4.5. $\text{Im}(\bar{v}_{x,p}) \cap i_2(H) = (0 \times 2H)/\Delta_{d,2k}$

Proof. That the first component of $\bar{v}_{x,p}(f)$ equals 0 is equivalent to that f induces the identity on $\pi_3(M)$ (Lemma 4.2).

We first assume M is spin. Recall the second component of $\bar{v}_{x,p}(f)$ is $\delta^{-1}(\frac{p_1}{2}(M_f) - 2k\bar{y})$. It suffices to show that $p_1(M_f)$ is divisible by 4. This is done by showing that the evaluation of $p_1(M_f)$ on a basis of $H_4(M_f)$ is

divisible by 4. Choose embeddings $S_i^3 \subset M$ representing a basis of $H_3(M)$. By Haefliger's theorem [Hae61, Theorem 1b], S_i^3 and $f(S_i^3)$ are isotopic in $M \times [0, 1]$. We choose an isotopy and obtain a smooth map $g_i: S_i^3 \times S^1 \rightarrow M_f$, which is an embedding on each $S_i^3 \times \{t\}$. Then from the exact sequence

$$0 \rightarrow H_4(M) \rightarrow H_4(M_f) \rightarrow H_3(M) \rightarrow 0$$

a basis of $H_4(M_f)$ is given by a generator of $H_4(M)$ and $g_{i*}[S_i^3 \times S^1]$. The evaluation of $p_1(M_f)$ on the generator of $H_4(M)$ equals $4k$. Next we consider the evaluation $\langle p_1(M_f), g_{i*}[S_i^3 \times S^1] \rangle$. We drop the subscript i in the sequel.

We form the map $g': S^3 \times S^1 \rightarrow M_f \times S^1$, $(x, t) \mapsto (g(x, t), t)$, which is an embedding of a 4-manifold into an 8-manifold. Then

$$\langle p_1(M_f), g_*[S^3 \times S^1] \rangle = \langle p_1(M_f \times S^1), g'_*[S^3 \times S^1] \rangle = \langle p_1(\nu), g'_*[S^3 \times S^1] \rangle$$

where ν is the normal bundle of $g'(S^3 \times S^1)$ in $M_f \times S^1$. Note that g' is homotopic to $g'': S^3 \times S^1 \rightarrow M_f \times S^1$ with $g''(x, t) = (g(x, t), -t)$, and the intersection of g' and g'' is empty. This shows that the Euler class of ν is trivial, hence ν is reduced to a 3-dimensional vector bundle. For such vector bundles, the first Pontrjagin class is divisible by 4.

In the non-spin case we need to replace $p_1(M_f)$ by $p_1(TM_f \oplus \bar{\eta})$. But since any complex line bundle over $S^3 \times S^1$ is trivial, the above computation still works. □

By Lemma 4.4 and 4.5, we may conclude that $\text{Im } \bar{v}_{x,p} = (H \times 2H)/\Delta_{d,2k}$. Now we may remove the factor 2 by modifying the definition of the variation map. Namely, in the spin case we define

$$v_{x,p}(f) = (\delta^{-1}(x^2 - d\bar{y}), \delta^{-1}(\frac{1}{4}(p_1(M_f) - 4k\bar{y}))) \in (H \times H)/\Delta_{d,k};$$

in the non-spin case

$$v_{x,p}(f) = (\delta^{-1}(x^2 - d\bar{y}), \delta^{-1}(\frac{1}{4}(p_1(TM_f \oplus \bar{\eta}) - 4k\bar{y}))) \in (H \times H)/\Delta_{d,k}.$$

Then we have the following

Proposition 4.6. $v_{x,p}: \mathcal{I}(M) \rightarrow (H \times H)/\Delta_{d,k}$ is a surjective homomorphism.

This gives the proof of the exact sequence (2.2) in Theorem 2.3.

Theorem 2.4 is a consequence of the following stronger statement.

Theorem 4.7. Taking the quotient by $\mathcal{K}(M)$ we have a short exact sequence

$$(4.5) \quad 0 \rightarrow (H \times H)/\Delta_{d,k} \rightarrow \text{MCG}(M)/\mathcal{K}(M) \rightarrow \text{Sp}(b, \mathbb{Z}) \rightarrow 1$$

in which as a $\text{Sp}(b, \mathbb{Z})$ -module $(H \times H)/\Delta_{d,k}$ fits into an exact sequence of $\text{Sp}(b, \mathbb{Z})$ -modules

$$0 \rightarrow H \xrightarrow{i_2} (H \times H)/\Delta_{d,k} \xrightarrow{pr_1} H \otimes \mathbb{Z}/d \rightarrow 0$$

where the $\text{Sp}(b, \mathbb{Z})$ -actions on H and $H \otimes \mathbb{Z}/d$ are the standard linear actions.

When $b = 2$, or when $b \geq 4$ and M is nonspin with $d/2$ odd, the exact sequence (4.5) splits, therefore

$$\mathrm{MCG}(M)/\mathcal{K}(M) = ((H \times H)/\Delta_{d,k}) \rtimes \mathrm{Sp}(b, \mathbb{Z}).$$

In the other cases the exact sequence (4.5) does not split. There is an injective homomorphism

$$\mathrm{MCG}(M)/\mathcal{K}(M) \rightarrow ((H \times H)/\Delta_{d,2k}) \rtimes \mathrm{Sp}(b, \mathbb{Z})$$

whose image has index 2^b . More precisely, let $\psi: H \rightarrow \mathbb{Z}/2$ be any quadratic refinement of the standard symplectic form on H , and let

$$s: \mathrm{Sp}(b, \mathbb{Z}) \rightarrow H \otimes \mathbb{Z}/2, \quad A \mapsto \psi \cdot A - \psi$$

be the 1-cocycle associated with ψ , then the image is the subgroup

$$\{([a, b], A) \mid b \equiv s(A) \pmod{2}\} \subset ((H \times H)/\Delta_{d,2k}) \rtimes \mathrm{Sp}(b, \mathbb{Z}).$$

Proof. The exact sequence (4.5) follows from the exact sequences in Theorem 2.3. We examine the action of $\mathrm{Sp}(b, \mathbb{Z})$ on $(H \times H)/\Delta_{d,k}$. There is a commutative diagram

$$(4.6) \quad \begin{array}{ccccccc} 0 & \longrightarrow & H & \longrightarrow & \mathrm{MCG}(\#_r(S^3 \times S^3))/\Theta_7 & \longrightarrow & \mathrm{Sp}(b, \mathbb{Z}) \longrightarrow 1 \\ & & \downarrow i_2 & & \downarrow & & \downarrow = \\ 0 & \longrightarrow & (H \times H)/\Delta_{d,k} & \longrightarrow & \mathrm{MCG}(M)/\mathcal{K}(M) & \longrightarrow & \mathrm{Sp}(b, \mathbb{Z}) \longrightarrow 1 \end{array}$$

in which the inclusion $i_2: H \rightarrow (H \times H)/\Delta_{d,k}$ is a $\mathrm{Sp}(b, \mathbb{Z})$ -module homomorphism, and the action of $\mathrm{Sp}(b, \mathbb{Z})$ on H is the standard linear action [Kre78]. In the exact sequence

$$0 \rightarrow H \xrightarrow{i_2} (H \times H)/\Delta_{d,k} \xrightarrow{\mathrm{pr}_1} H \otimes \mathbb{Z}/d \rightarrow 0$$

the $\mathrm{Sp}(b, \mathbb{Z})$ -module structure on $H \otimes \mathbb{Z}/d$ can be obtained as follows: for any $f \in \mathcal{I}(M)$, $\varphi \in \mathrm{MCG}(M)$, we consider $(\varphi^{-1}f\varphi)_* - \mathrm{id}$ as a homomorphism from H to \mathbb{Z}/d . For any $a \in H$, let $\bar{a} \in \pi_3(M)$ be a pre-image of a , then

$$(\varphi^{-1}f\varphi)_*(\bar{a}) - \bar{a} = \varphi_*^{-1}(f_*(\varphi_*(\bar{a})) - \varphi_*(\bar{a})) = (f_* - \mathrm{id})(\varphi_*(\bar{a}))$$

where the second equality follows from the facts that $f_*(\varphi_*(\bar{a})) - \varphi_*(\bar{a}) \in \mathbb{Z}/d$ and φ_*^{-1} is trivial on \mathbb{Z}/d . This shows that the induced action of $\mathrm{Sp}(b, \mathbb{Z})$ on $H \otimes \mathbb{Z}/d$ is the standard linear action.

Let $\alpha \in H^2(\mathrm{Sp}(b, \mathbb{Z}), H)$ be the cohomology class of the extension of the upper exact sequence in diagram (4.6), then it is a standard fact that $i_{2*}(\alpha) \in H^2(\mathrm{Sp}(b, \mathbb{Z}), (H \times H)/\Delta_{d,k})$ is the cohomology class of the extension of the lower exact sequence (see [Bro82, p.94 Ex. 1(b)]).

When $b = 2$ we have $\alpha = 0$ (by [Kry02] or the fact $H^2(\mathrm{Sp}(2, \mathbb{Z}), \mathbb{Z}^2) = 0$ [Kra19, Lemma A.3]), hence $i_{2*}(\alpha) = 0$. Therefore

$$\mathrm{MCG}(M)/\mathcal{K}(M) \cong ((H \times H)/\Delta_{d,k}) \rtimes \mathrm{Sp}(b, \mathbb{Z}).$$

When $b \geq 4$, the following commutative diagram of exact sequences

$$\begin{array}{ccccccc} 0 & \longrightarrow & H & \xrightarrow{\cdot d} & H & \longrightarrow & H \otimes \mathbb{Z}/d \longrightarrow 0 \\ & & \downarrow \cdot k & & \downarrow i_1 & & \downarrow = \\ 0 & \longrightarrow & H & \xrightarrow{i_2} & (H \times H)/\Delta_{d,k} & \longrightarrow & H \otimes \mathbb{Z}/d \longrightarrow 0 \end{array}$$

induces a commutative diagram of exact sequence of cohomology groups

$$\begin{array}{ccccccc} H^1(\mathrm{Sp}(b, \mathbb{Z}), H \otimes \mathbb{Z}/d) & \longrightarrow & H^2(\mathrm{Sp}(b, \mathbb{Z}), H) & \xrightarrow{\cdot d} & H^2(\mathrm{Sp}(b, \mathbb{Z}), H) & & \\ \downarrow = & & \downarrow \cdot k & & \downarrow i_{1*} & & \\ H^1(\mathrm{Sp}(b, \mathbb{Z}), H \otimes \mathbb{Z}/d) & \xrightarrow{\delta} & H^2(\mathrm{Sp}(b, \mathbb{Z}), H) & \xrightarrow{i_{2*}} & H^2(\mathrm{Sp}(b, \mathbb{Z}), (H \times H)/\Delta_{d,k}) & & \end{array}$$

The cohomology groups $H^*(\mathrm{Sp}(b, \mathbb{Z}), H)$ are annihilated by 2 ([Kra19, Lemma A.3]). When M is spin, d and $p/2$ have the same parity (see (3.3)), therefore $\delta = 0$. By [Cro11, Theorem 1.1] $\alpha \neq 0$, therefore $i_{2*}\alpha \neq 0$. When M is nonspin, k and $d/2$ have the same parity. When k is even, $\delta = 0$; when k is odd, $i_{2*} = 0$. This shows the statements about the splitting of the exact sequence (4.5).

The cocycle $\alpha \in H^2(\mathrm{Sp}(b, \mathbb{Z}), H)$ corresponding to the extension problem of the upper line of diagram (4.6) is described in [Cro11, Corollary 3.5]. As a consequence, the embedding $2 \cdot : H \rightarrow H$ induces an embedding of $\mathcal{I}(\#_r(S^3 \times S^3))/\Theta_7$ into the semi-direct product $H \rtimes \mathrm{Sp}(b, \mathbb{Z})$ whose image is

$$\{(b, A) \mid b \equiv s(A) \pmod{2}\}$$

where $s : \mathrm{Sp}(b, \mathbb{Z}) \rightarrow H \otimes \mathbb{Z}/2$, $A \mapsto \psi \cdot A - \psi$ is a 1-cocycle associated with an arbitrary quadratic refinement $\psi : H \rightarrow \mathbb{Z}/2$ of the standard symplectic form (c. f. [Cro11, Proposition 2.3]). Therefore the embedding

$$(H \times H)/\Delta_{d,k} = (H \times 2H)/\Delta_{d,2k} \subset (H \times H)/\Delta_{d,2k}$$

induces an embedding of $\mathrm{MCG}(M)/\mathcal{K}(M)$ into the semi-direct product $(H \times H)/\Delta_{d,2k} \rtimes \mathrm{Sp}(b, \mathbb{Z})$, and the image is

$$\{([a, b], A) \mid a, b \in H, A \in \mathrm{Sp}(b, \mathbb{Z}), b \equiv s(A) \pmod{2}\}.$$

□

5. GENERALIZED KRECK-STOLZ INVARIANT

Now we come to the problem of determining the kernel

$$\mathcal{K}(M) = \mathrm{Ker}\{v_{x,p} : \mathcal{I}(M) \rightarrow H \times H/\Delta_{d,k}\}.$$

This task will be done by defining an invariant of the mapping torus. This invariant is not only crucial for the computation of \mathcal{K}_M , but also important for the study of the properties of $\mathrm{MCG}(M)$. We first present the results of $\mathcal{K}(M)$ in the following theorem. Recall from section 3.1 that the first Pontrjagin class of M is determined by an integer l .

Theorem 5.1. *The group $\mathcal{K}(M)$ is the center of $\text{MCG}(M)$. There is an isomorphism*

$$S: \mathcal{K}(M) \rightarrow \mathbb{Z}^3/L_M,$$

where L_M is a lattice of rank 3. The group $\mathcal{K}(M)$ is given as follows

(1) When M is spin

$$\mathcal{K}(M) \cong (\mathcal{K}(M))_{(2)} \times \mathbb{Z}/3^b \times \mathbb{Z}/7^c$$

where

$$b = \begin{cases} 1 & d \equiv 0 \pmod{3} \\ 0 & \text{otherwise} \end{cases} \quad c = \begin{cases} 1 & d - l \equiv 0 \pmod{7} \\ 0 & \text{otherwise} \end{cases}$$

The group $(\mathcal{K}(M))_{(2)}$ is given in the following table

	d odd	$d \equiv 2 \pmod{4}$	$d \equiv 4 \pmod{8}$	$d \equiv 0 \pmod{8}$
l odd	$\mathbb{Z}/2$	$\mathbb{Z}/2 \times \mathbb{Z}/4$	$(\mathbb{Z}/2)^2$	$(\mathbb{Z}/2)^2$
l even	$\mathbb{Z}/4$	$\mathbb{Z}/2 \times \mathbb{Z}/4$	$(\mathbb{Z}/2)^2 \times \mathbb{Z}/4$	$\mathbb{Z}/2 \times \mathbb{Z}/a \times \mathbb{Z}/4$

where

$$a = \begin{cases} 2 & l \equiv 2 \pmod{4} \\ 4 & l \equiv 0 \pmod{4} \end{cases}$$

(2) When M is non-spin

$$\mathcal{K}(M) \cong \mathbb{Z}/2^a \times \mathbb{Z}/3^b \times \mathbb{Z}/7^c,$$

where

$$b = \begin{cases} 1 & d \equiv 0 \pmod{3} \\ 0 & \text{otherwise} \end{cases} \quad c = \begin{cases} 1 & d \equiv l \equiv 0 \pmod{7} \\ 0 & \text{otherwise} \end{cases}$$

The value of a is given in the following table (where $\delta = d/2$)

	δ odd	$\delta \equiv 2 \pmod{4}$	$\delta \equiv 4 \pmod{8}$	$\delta \equiv 0 \pmod{8}$
l odd	0	1	2	2
l even	0	1	2	3

We first prove Theorem 5.1 in the spin case in §5-7. The treatment of the non-spin case is similar but with slight modifications. This is put in §9.

For the mapping class group of surfaces, the Birman-Craggs-Johnson invariants are defined using the relative mapping class group $\text{MCG}_{(F_g, D)}$ which is the mapping class group of diffeomorphisms fixing an embedded disk. In our situation we have to do the same and pass to $\text{MCG}(M, D)$. The group $\text{MCG}(M, D)$ is the group of isotopy classes of diffeomorphisms of M which are the identity on an embedded 6-dimensional disk $D \subset M$. The groups $\text{MCG}(M)$ and $\text{MCG}(M, D)$ are related by the following exact sequence ([Wal63, p.265])

$$(5.1) \quad \pi_1 GL(n, \mathbb{R}) \rightarrow \text{MCG}(M, D) \rightarrow \text{MCG}(M) \rightarrow 1.$$

The exact sequences (2.1) and (2.2) in Theorem 2.3 remain valid if we replace $\text{MCG}(M)$ by $\text{MCG}(M, D)$. We denote the corresponding kernel of $v_{x,p}$ by

$$\mathcal{K}(M, D) = \ker\{v_{x,p}: \mathcal{I}(M, D) \rightarrow H \times H/\Delta_{d,k}\}.$$

Then the above exact sequence gives rise to an exact sequence

$$(5.2) \quad \pi_1 GL(n, \mathbb{R}) \rightarrow \mathcal{K}(M, D) \rightarrow \mathcal{K}(M) \rightarrow 1.$$

In this section, based on the recent development of modified surgery theory in [Kre18], we define a Kreck-Stolz type invariant for elements in $\mathcal{K}(M, D)$. This invariant defines an injective homomorphism

$$S: \mathcal{K}(M, D) \hookrightarrow \mathbb{Z}^3 / \bar{L}(M).$$

In §6 we use this invariant to study $\text{MCG}(S^2 \times D^4, \partial)$ and then in §7 we determine the image of S by comparing $\mathcal{K}(M, D)$ with $\text{MCG}(S^2 \times D^4, \partial)$.

Let M be a spin 6-manifold which looks like a complete intersection, $f: (M, D) \rightarrow (M, D)$ be a diffeomorphism such that $[f] \in \mathcal{K}(M, D)$. Let M_f be the mapping torus. Let

$$B = \mathbb{CP}^\infty \times K(\mathbb{Z}, 4) \times B\text{Spin},$$

and let $p: B \rightarrow B\text{Spin} \rightarrow BSO$ be the canonical projection. Then the normal Gauss map $M_f \rightarrow BSO$ is lifted to B as follows: Fix a generator $x \in H^2(M) = H^2(M_f)$, there is a unique map $M_f \rightarrow \mathbb{CP}^\infty$ representing x . For $d \neq 0$, let $y = x^2/d \in H^4(M)$ be the generator; for $d = 0$ but $p_1(M) \neq 0$, let $y = p_1(M)/4k \in H^4(M)$; for $d = 0 = p_1(M)$, take $y \in H^4(M)$ an arbitrary generator³. For $[f] \in \mathcal{K}(M, D)$, since $\tau(f) = 0$, there is a unique $y' \in H^4(M_f)$ which is a preimage of $y \in H^4(M)$ under $H^4(M_f) \rightarrow H^4(M)$, such that $x^2 - dy' = 0$ and $\frac{p_1}{2}(M_f) - 2ky' = 0$. There is a unique map $M_f \rightarrow K(\mathbb{Z}, 4)$ representing y' . Finally choose the spin structure on the stable normal bundle of M_f , which restricts to the trivial spin structure on the preferred embedding $D \times S^1 \subset M_f$. This induces a unique map $M_f \rightarrow B\text{Spin}$. Put all these maps together we get $g: M_f \rightarrow B$ which is a lift of the normal Gauss map $M_f \rightarrow BSO$.

The pair (M_f, g) represents a bordism class in the B -bordism group $\Omega_7(B, p) = \Omega_7^{\text{spin}}(\mathbb{CP}^\infty \times K(\mathbb{Z}, 4))$, which is zero by an easy computation with the Atiyah-Hirzebruch spectral sequence, c. f. [Kre18, Theorem 6]. Therefore there exists a normal B -bordism $G: W^8 \rightarrow B$ such that $\partial(W, G) = (M_f, g)$. After taking connected sum with quaternion projective planes \mathbb{HP}^2 's (or $\overline{\mathbb{HP}^2}$'s) we may assume that the signature of W is zero.

Let $\bar{x} \in H^2(B)$ be the pull-back of the canonical generator of $H^2(\mathbb{CP}^\infty)$, $\bar{y} \in H^4(B)$ be the pull-back of the canonical generator of $H^4(K(\mathbb{Z}, 4))$ and $\frac{p_1}{2} \in H^4(B)$ be the pull-back of the universal spin Pontrjagin class in $H^4(B\text{Spin})$. Let $\alpha = G^*(\bar{x}^2 - d\bar{y})$ and $\beta = G^*(\frac{p_1}{2} + 2k\bar{y})$. Then by the construction above, the restriction of α and β on $\partial W = M_f$ are zero, we have relative cohomology classes $\bar{\alpha}, \bar{\beta} \in H^4(W, \partial W)$ which maps to α and

³Because the case of $S^2 \times S^4$ is crucial for the computation, here we need to include the possibility $d = 0$.

β respectively. Now define characteristic numbers

$$\begin{aligned} s_1(f) &= \langle \bar{\beta}^2, [W, \partial W] \rangle \\ s_2(f) &= \langle \bar{\alpha}^2 + \bar{\alpha} \cup \bar{\beta}, [W, \partial W] \rangle \\ s_3(f) &= \langle \bar{\alpha} \cup \bar{\beta}, [W, \partial W] \rangle \end{aligned}$$

Elementary algebraic topology shows that these numbers don't depend on the choices of the cohomology classes $\bar{\alpha}$ and $\bar{\beta}$. To get an invariant we also need to take into account of the choices of W . This amounts to a lattice $\bar{L}(M) \subset \mathbb{Z}^3$ consisting of characteristic numbers of closed 8-dimensional B -manifolds with signature 0. Therefore we have a well-defined invariant $S(f) = [(s_1(f), s_2(f), s_3(f))] \in \mathbb{Z}^3 / \bar{L}(M)$. We call it the generalized Kreck-Stolz invariant of f .

Proposition 5.2. *The generalized Kreck-Stolz invariant defines an injective homomorphism*

$$S: \mathcal{K}(M, D) \rightarrow \mathbb{Z}^3 / \bar{L}(M)$$

Proof. We first show that S is a group homomorphism. Given diffeomorphisms f_1 and f_2 , with normal B -bordism (W_1, G_1) and (W_2, G_2) for the corresponding mapping tori, we may take $W = W_1 \cup_{M \times [0,1]} W_2$ to be the normal B -bordism for the mapping torus of $f_1 \circ f_2$. It's easy to see that in Wall's formula of the additivity of the signature ([Wal69, Theorem]) the correction term $\sigma(V; A, B, C) = 0$, therefore $\text{sign}(W) = \text{sign}(W_1) + \text{sign}(W_2) = 0$. Therefore W can be taken as the null-bordism for computing the S -invariant of $f_1 \circ f_2$. The Mayer-Vietoris sequences form a commutative diagram

$$\begin{array}{ccc} H^4(W, \partial W) & \longrightarrow & H^4(W_1, \partial W_1) \oplus H^4(W_2, \partial W_2) \\ \downarrow & & \downarrow \\ H^4(W) & \longrightarrow & H^4(W_1) \oplus H^4(W_2) \end{array}$$

therefore $\bar{\alpha}, \bar{\beta} \in H^4(W, \partial W)$ are the pre-image of $(\bar{\alpha}_1, \bar{\alpha}_2)$ and $(\bar{\beta}_1, \bar{\beta}_2)$ respectively. The S -invariant of $f_1 \circ f_2$ is defined via characteristic numbers of W , which are now clearly the sum of the characteristic numbers of W_1 and W_2 , i. e. $s_i(f_1 \circ f_2) = s_i(f_1) + s_i(f_2)$ for $i = 1, 2, 3$.

To show the injectivity of S , notice that (W, G) can be viewed as a B -bordism between $M \times [0, 1]$ and $M \times [0, 1]$, where the boundaries $M \times 0$ and $M \times 1$ are identified by id_M and f respectively. By modified surgery theory [Kre99], the surgery obstruction $\theta(W, G)$ to (W, G) being B -bordant to an h -cobordism relative boundary is in $l_8(\mathbb{Z})$. It is shown in [Kre18, Theorem 5b] that $\theta(W, G)$ is elementary if

$$S(f) = [(s_1(f), s_2(f), s_3(f))] = 0 \in \mathbb{Z}^3 / \bar{L}(M)$$

In this situation one may replace W by an h -cobordism ([Kre99, Theorem 3]), still denoted by W . The embedding $D \times S^1 \subset M_f$ extends to an

embedding $D \times D^2 \subset W$, which can be viewed as an h -cobordism between $D \times I$ and $D \times I$. By the h -cobordism theorem, there is a diffeomorphism

$$(M \times [0, 1], D \times [0, 1]) \rightarrow (M \times [0, 1], D \times [0, 1])$$

which can be viewed as a pseudo-isotopy between f and $\text{id}_M \text{ rel } D$. Hence by Cerf's pseudo-isotopy theorem [Cer70], f is isotopic to $\text{id}_M \text{ rel } D$. This shows the injectivity of S . \square

Now we determine the lattice $\bar{L}(M)$. For this we need to compute the subgroup of the 8-dimensional B -bordism group consisting of manifolds with signature zero,

$$\Omega_8(B)_0 = \Omega_8^{\text{spin}}(\mathbb{CP}^\infty \times K(\mathbb{Z}, 4))_0$$

Computation with the Atiyah-Hirzebruch spectral sequence (c. f. [Kre18, Theorem 10]) shows that this group is isomorphic to \mathbb{Z}^6 , with generators

$$\begin{aligned} e_1 &= [B^8] - 8 \cdot 28[\mathbb{HP}^2] \\ e_2 &= [S^2 \times S^2 \times S^2 \times S^2 \xrightarrow{\sum_{i=1}^4 x_i} \mathbb{CP}^\infty \xrightarrow{i} \mathbb{CP}^\infty \times K(\mathbb{Z}, 4)] \\ e_3 &= [V(2) \hookrightarrow \mathbb{CP}^\infty \xrightarrow{i} \mathbb{CP}^\infty \times K(\mathbb{Z}, 4)] - 2[\mathbb{HP}^2] \\ e_4 &= [\mathbb{HP}^2 \xrightarrow{y} K(\mathbb{Z}, 4) \rightarrow \mathbb{CP}^\infty \times K(\mathbb{Z}, 4)] - [\mathbb{HP}^2] \\ e_5 &= [S^4 \times S^4 \xrightarrow{y_1 + y_2} K(\mathbb{Z}, 4)] \\ e_6 &= \frac{1}{2}[S^2 \times S^2 \times S^4 \xrightarrow{(x_1 + x_2, y)} \mathbb{CP}^\infty \times K(\mathbb{Z}, 4)] \end{aligned} \quad (5.3)$$

where B^8 is the Bott manifold, $V(2)$ is a degree 2 hypersurface in \mathbb{CP}^5 , $x_i \in H^2(S_i^2)$, $y_i \in H^4(S_i^4)$ and $y \in H^4(\mathbb{HP}^2)$ are generators.

The characteristic numbers of these generators are given in the following table, and the lattice $\bar{L}(M)$ is generated by column vectors.

$$(5.4) \quad \begin{array}{c|cccccc} & e_1 & e_2 & e_3 & e_4 & e_5 & e_6 \\ \hline s_1 & -8 \cdot 28 & 0 & 0 & 4k^2 - 4k & 8k^2 & 0 \\ s_2 & 0 & 24 & 0 & d^2 - d(2k - 1) & 2d^2 - 4dk & 2k - 2d \\ s_3 & 0 & 0 & -2 & -d(2k - 1) & -4dk & 2k \end{array}$$

6. THE MAPPING CLASS GROUP OF $(S^2 \times D^4, \partial)$

In this section we determine the mapping class group of $(S^2 \times D^4, \partial)$, denoted by $\text{MCG}(S^2 \times D^4, \partial)$. This is the group of isotopy classes of diffeomorphisms of $S^2 \times D^4$, which are the identity near the boundary. This is crucial for the determination of the image of S , and the generators of the mapping class group.

6.1. Simply-connected homology $S^2 \times S^5$. Let N^7 be a closed, simply-connected spin 7-manifold with $H_*(N) \cong H_*(S^2 \times S^5)$. Let \mathcal{N} be the set of equivalence classes of oriented 7-manifolds of this type. The oriented diffeomorphism type of these manifolds is determined by the Kreck-Stolz invariant [KrSt88]. We recall the definition here.

The normal 2-type of N is

$$p: B = \mathbb{CP}^\infty \times B\text{Spin} \rightarrow BSO$$

where p is the canonical projection on the second factor. Choose a generator of $H^2(N)$, there is a unique lifting $\bar{\nu}: N \rightarrow B$ of the normal Gauss map $\nu: N \rightarrow BSO$. Since the corresponding B -bordism group $\Omega_7(B, p) = \Omega_7^{\text{spin}}(\mathbb{CP}^\infty) = 0$, there exists a normal B -bordism $G: W^8 \rightarrow B$ such that $\partial(W, G) = (N, \bar{\nu})$. After taking connected sum with \mathbb{HP}^2 's (or $\overline{\mathbb{HP}^2}$'s) we may assume that the signature of W is zero.

Let $x \in H^2(B)$ be a generator, $\frac{p_1}{2} \in H^4(B)$ be the pull-back of the universal spin Pontrjagin class in $H^4(B\text{Spin})$, $\alpha = G^*(x^2)$, $\beta = G^*(\frac{p_1}{2})$. Since $H^3(N) = H^4(N) = 0$, α and β restrict to 0 on $\partial W = N$, therefore we have unique cohomology classes $\bar{\alpha}, \bar{\beta} \in H^4(W, \partial W)$ which are the preimage of α and β respectively. Now we define characteristic numbers

$$\begin{aligned} s_1(N) &= \langle \bar{\beta}^2, [W, \partial W] \rangle \\ s_2(N) &= \langle \bar{\alpha}^2 + \bar{\alpha} \cup \bar{\beta}, [W, \partial W] \rangle \\ s_3(N) &= \langle \bar{\alpha} \cup \bar{\beta}, [W, \partial W] \rangle \end{aligned}$$

It's easy to see that these numbers don't depend on the choice of the generator of $H^2(N)$. Taking into account the choices of W , we obtain a well-defined invariant $S(N) = [(s_1(N), s_2(N), s_3(N))] \in \mathbb{Z}^3/L$, where $L \subset \mathbb{Z}^3$ is the lattice generated by the corresponding characteristic numbers of closed 8-dimensional B -manifolds with signature 0. The generators of the bordism group are e_1, e_2, e_3 in 5.3 and from table 5.4 we see that L is generated by

$$(6.1) \quad \begin{array}{c|ccc} & e_1 & e_2 & e_3 \\ \hline s_1 & -8 \cdot 28 & 0 & 0 \\ s_2 & 0 & 24 & 0 \\ s_3 & 0 & 0 & -2 \end{array}$$

Therefore $\mathbb{Z}^3/L \cong \mathbb{Z}/(8 \cdot 28) \oplus \mathbb{Z}/24 \oplus \mathbb{Z}/2$,

Let (W, G) be a B -bordism between $(N_1, \bar{\nu}_1)$ and $(N_2, \bar{\nu}_2)$, then the surgery obstruction $\theta(W, G)$ to (W, G) being B -bordant to an h -cobordism is in $l_8(\mathbb{Z})$. By [Kre99, Theorem 6], $\theta(W, G)$ is elementary if $S(N_1) = S(N_2)$. Therefore $S(N)$ is a complete invariant of manifolds in \mathcal{N} .

Lemma 6.1. *There is an injective map between sets*

$$S = [(s_1, s_2, s_3)]: \mathcal{N} \rightarrow \mathbb{Z}^3/L = \mathbb{Z}/(8 \cdot 28) \oplus \mathbb{Z}/24 \oplus \mathbb{Z}/2$$

Remark 6.2. In [KrSt88] the Kreck-Stolz s -invariants are defined by the defect of (twisted) Dirac operators. An explicit formula expressing the s -invariants by characteristic numbers is given in [KrSt91, (2.4)]. If we turn the above defined S -invariant into \mathbb{Q}/\mathbb{Z} -valued invariants $s_1/(8 \cdot 28)$, $s_2/24$ and $s_3/2$, then it's easy to see from these expressions that

$$s_1/(8 \cdot 28) = s'_1, \quad s_2/24 = s'_2, \quad s_3/2 = s'_3 - 2^4 s'_2$$

where s'_i denotes the original Kreck-Stolz invariants.

We construct certain manifolds in \mathcal{N} and compute their S -invariant, which will be used in the next subsection. Let ξ_m be a 4-dimensional oriented real vector bundle over \mathbb{CP}^2 , with

$$w_2(\xi_m) \neq 0, \quad e(\xi_m) = x^2, \quad p_1(\xi_m) = (3 + 4m)x^2, \quad m \in \mathbb{Z}$$

where $x \in H^2(\mathbb{CP}^2)$ is a generator. (These bundles can be constructed as follows: let $p: \mathbb{CP}^2 \rightarrow \mathbb{CP}^2 \vee S^4$ be the pinch map, $\xi = p^*(\eta \oplus \varepsilon^2 \vee \xi')$, where η is the tautological line bundle over \mathbb{CP}^2 and ξ' is a 4-dimensional vector bundle over S^4 . Then $w_2(\xi) \neq 0$, $e(\xi) = p^*e(\xi')$ and $p_1(\xi) = p^*(p_1(\xi')) + x^2$. Note that 4-dimensional vector bundles over S^4 can have $e(\xi') = 2a + b$ and $p_1(\xi') = -2b$ for all $a, b \in \mathbb{Z}$.)

Let $N_m = S(\xi_m)$ be the sphere bundle of ξ_m . Since the Euler class of ξ_m is x^2 , from the Gysin sequence it is easy to see $N_m \in \mathcal{N}$. We may compute the S -invariant of N_m using $W = D(\xi_m) \# \overline{\mathbb{HP}^2}$. Straightforward computation gives

$$\begin{aligned} s_1(N_m) &= 4m(m+3) \equiv 0 \pmod{8} \\ s_2(N_m) &= -2(m+1) \\ s_3(N_m) &= 2m+3 \equiv 1 \pmod{2} \end{aligned} \tag{6.2}$$

6.2. Computation of $\text{MCG}(S^2 \times D^4, \partial)$. Let $\mathcal{I}(S^2 \times S^4)$ be the subgroup of the mapping class group $\text{MCG}_{S^2 \times S^4}$, consisting of isotopy classes of diffeomorphisms of $S^2 \times S^4$ which act trivially on homology. We define a map

$$\Psi: \mathcal{I}(S^2 \times S^4) \rightarrow \mathcal{N}$$

as follows: given a diffeomorphism $f: S^2 \times S^4 \rightarrow S^2 \times S^4$ which acts trivially on homology, let $N = S^2 \times D^5 \cup_f S^2 \times D^5$. It's easy to see that $N \in \mathcal{N}$. Clearly isotopic diffeomorphisms produce diffeomorphic 7-manifolds, therefore Ψ is a well-define map.

Lemma 6.3. *Ψ is surjective.*

Proof. Given $N \in \mathcal{N}$, there is an embedding $\varphi_1: S^2 \rightarrow N$ representing a generator of $H_2(N)$. The normal bundle of this embedding is trivial since N is spin, thus φ_1 extends to an embedding $\varphi_1: S^2 \times D^5 \hookrightarrow N$. The complement $N - \varphi_1(\text{int}(S^2 \times D^5))$ is simply-connected, spin, and has the same homology as S^2 . Therefore there is a second embedding $\varphi_2: S^2 \times D^5 \hookrightarrow N - \varphi_1(\text{int}(S^2 \times D^5))$ and it's easy to see that $N - (\varphi_1(\text{int}(S^2 \times D^5)) \cup \varphi_2(\text{int}(S^2 \times D^5)))$ is an h -cobordism. Therefore there exists a diffeomorphism $f: S^2 \times S^4 \rightarrow S^2 \times S^4$ such that $N = S^2 \times D^5 \cup_f S^2 \times D^5$. It's easy to see that f acts trivially on homology. \square

Recall that we have the exact sequence (5.1)

$$\pi_1 GL(6, \mathbb{R}) \rightarrow \mathcal{I}(S^2 \times S^4, D) \rightarrow \mathcal{I}(S^2 \times S^4) \rightarrow 1.$$

In this situation we may identify $\mathcal{I}(S^2 \times S^4)$ with $\mathcal{I}(S^2 \times S^4, D)$ by the following lemma.

Lemma 6.4. *Let M^n ($n \geq 3$) be a closed smooth manifold with a non-trivial S^1 -action. Let $p \in M$ be a fixed-point, such that the isotropy representation at p represents the non-trivial element in $\pi_1 GL(n, \mathbb{R})$. Then the boundary diffeomorphism is isotopic to the identity in $\text{MCG}(M, D)$*

Proof. An explicit isotopy is given by the S^1 -action. \square

Therefore by Proposition 5.2 and table 5.4, the generalized Kreck-Stolz S -invariant defines an injective homomorphism

$$S: \mathcal{I}(S^2 \times S^4) \rightarrow \mathbb{Z}^3/L$$

where L is generated by table 6.1. Comparing the definition of this invariant and that of the Kreck-Stolz invariant of manifolds in \mathcal{N} we have a commutative diagram

$$(6.3) \quad \begin{array}{ccc} \mathcal{I}(S^2 \times S^4) & \xrightarrow{\Psi} & \mathcal{N} \\ \downarrow S & & \downarrow S \\ \mathbb{Z}^3/L & \xrightarrow{=} & \mathbb{Z}^3/L \end{array}$$

(*Proof.* Let V be a coboundary of the mapping torus T_f , let

$$W = V \cup_{S^2 \times S^4 \times [0,1]} S^2 \times D^5 \times [0, 1],$$

then $\partial W = (S^2 \times D^5) \cup_f (S^2 \times D^5) = N$, we use W to compute the S -invariants of N , the $S^2 \times D^5 \times [0, 1]$ part doesn't contribute to the characteristic numbers.)

For any self-diffeomorphism f of $S^2 \times S^4$ which acts trivially on homology, since $\pi_2 SO(4) = 0$, after isotopy we may assume that f is the identity on a copy of $S^2 \times D^4$ in the standard decomposition $S^2 \times S^4 = S^2 \times D^4 \cup S^2 \times D^4$. This means, the natural homomorphism $\text{MCG}(S^2 \times D^4, \partial) \rightarrow \mathcal{I}(S^2 \times S^4)$ defined by extending a diffeomorphism of $(S^2 \times D^4, \partial)$ to $S^2 \times S^4$ by the identity on the other copy of $S^2 \times D^4$ is an epimorphism.

Recall that from the discussion after Theorem 2.13, we have a homomorphism $\text{MCG}(D^6, \partial) \rightarrow \text{MCG}(M)$, which is given by extending a diffeomorphism of (D^6, ∂) to the complement by the identity. The image are called disk supported diffeomorphisms. Clearly the image of this homomorphism is in the Torelli group. Now we study this homomorphism for $M = S^2 \times S^4$,

$$\text{MCG}(D^6, \partial) \rightarrow \text{MCG}(S^2 \times D^4, \partial) \rightarrow \mathcal{I}(S^2 \times S^4),$$

here we take D^6 in the interior of $S^2 \times D^4$. To this end we compute the S -invariant of disk supported diffeomorphisms in $\mathcal{I}(S^2 \times S^4)$. By diagram 6.3 this is equivalent to computing the S -invariant of $S^2 \times S^5 \# \Sigma$, where Σ is an exotic 7-sphere. Especially, let Σ be the boundary of the E_8 -plumbing manifold E_8 , which is a generator of Θ_7 . Then we may take

$$W = (S^2 \times D^6) \natural E_8 \# (8 \cdot \overline{\text{HIP}^2}) \rightarrow \mathbb{CP}^\infty \times B\text{Spin}$$




FIGURE 1.

as the null B -bordism, where \natural denotes the boundary connected-sum. From this it's easy to get

$$s_1 = -8, \quad s_2 = 0, \quad s_3 = 0.$$

Therefore the homomorphism $\text{MCG}(D^6, \partial) \rightarrow \mathcal{I}(S^2 \times S^4)$ is injective. The same holds for the homomorphism $\text{MCG}(D^6, \partial) \rightarrow \text{MCG}(S^2 \times D^4, \partial)$.

The next step to the computation of $\text{MCG}(S^2 \times D^4, \partial)$ is to determine the cokernel of $\text{MCG}(D^6, \partial) \rightarrow \text{MCG}(S^2 \times D^4, \partial)$, denoted by $\overline{\text{MCG}}(S^2 \times D^4, \partial)$. An upper bound of the order of $\overline{\text{MCG}}(S^2 \times D^4, \partial)$ is obtained as follows.

Let $P_3(S^2)$ be the 3rd stage Postnikov tower of S^2 , we define a homomorphism

$$\Phi: \Omega_7^{\text{string}}(P_3(S^2)) \rightarrow \overline{\text{MCG}}(S^2 \times D^4, \partial)$$

as follows. The normal 3-type of $S^2 \times D^4$ is

$$B = P_3(S^2) \times B\text{String} \xrightarrow{p} BSO$$

where p is the canonical projection on the second factor. Fix a generator of $H^2(S^2 \times D^4)$, there is a unique (up to homotopy) normal 3-smoothing $\overline{\nu}: S^2 \times D^4 \rightarrow B$. The corresponding B -bordism group $\Omega_*(B, p)$ is the string bordism group $\Omega_*^{\text{string}}(P_3(S^2))$. Given $[N^7, g] \in \Omega_7^{\text{string}}(P_3(S^2))$, consider the bordism class $[S^2 \times D^4 \times I, \overline{\nu}] + [N, g]$, where $I = [0, 1]$. View this as a B -bordism between $(S^2 \times D^4, \overline{\nu})$ and $(S^2 \times D^4, \overline{\nu})$, where the boundaries are identified by the identity. By [Kre99] the surgery obstruction to this B -bordism being B -bordant rel ∂ to an h -cobordism rel ∂ is in $L_7(\mathbb{Z}) = 0$. Thus we obtain an h -cobordism rel ∂ between $S^2 \times D^4$ and $S^2 \times D^4$, and henceforth a diffeomorphism $f: (S^2 \times D^4, \partial) \xrightarrow{\cong} (S^2 \times D^4, \partial)$.

To consider the dependence of this construction on the choices of representatives of a given B -bordism class, let (N, g) and (N', g') be two such representatives, then the h -cobordisms W and W' obtained by surgery are also B -bordant, say, through V^8 (see Figure 1). Then the surgery obstruction $\theta(V)$ to V being an h -cobordism rel ∂ lies in $L_8(\mathbb{Z}) \cong \mathbb{Z}$. After making a boundary connected sum of V with copies of E_8 along W , the surgery obstruction vanishes, and W is replaced by $W \natural \Sigma$ for some $\Sigma \in \Theta_7$. This means that we obtain a pseudo-isotopy between the diffeomorphism produced by




FIGURE 2.

(N, g) composed with the local diffeomorphism corresponding to Σ and the diffeomorphism produced by (N', g') . This shows that Φ is well-defined.

Lemma 6.5. $\Phi: \Omega_7^{\text{string}}(P_3(S^2)) \rightarrow \overline{\text{MCG}}(S^2 \times D^4, \partial)$ is a surjective homomorphism.

Proof. Φ is a homomorphism: clearly we have

$$\begin{aligned} & [S^2 \times D^4 \times I, \bar{\nu}] + [N_1, g_1] + [N_2, g_2] \\ &= [((S^2 \times D^4 \times I, \bar{\nu}) + (N_1, g_1)) \cup_{S^2 \times D^4} ((S^2 \times D^4 \times I, \bar{\nu}) + (N_2, g_2))] \end{aligned}$$

After surgery we obtain an h -cobordism rel ∂ , which is the union of the h -cobordisms obtained from (N_1, g_2) and (N_2, g_2) along $S^2 \times D^4$. Therefore the diffeomorphism induced by this h -cobordism is the composition of the diffeomorphisms induced by (N_1, g_1) and (N_2, g_2) .

Φ is surjective: given a diffeomorphism $f: (S^2 \times D^4, \partial) \rightarrow (S^2 \times D^4, \partial)$, the normal 3-smoothings $\bar{\nu}$ and $\bar{\nu} \circ f$ are homotopic. Choose a homotopy $g: S^2 \times D^4 \times I \rightarrow B$, extend g to a map $S^2 \times S^4 \times I \rightarrow B$, where on the other copy of $S^2 \times D^4 \times I$ we use $\bar{\nu}$. Still call this map g . We also extend f to a diffeomorphism of $S^2 \times S^4$, where on the other copy of $S^2 \times D^4$ we use the identity diffeomorphism. Still call this diffeomorphism f . Glue the two boundary components of $S^2 \times S^4 \times I$ via f , the map g induces a map $\bar{g}: (S^2 \times S^4)_f \rightarrow B$ which represents an element in $\Omega_7^{\text{string}}(P_3(S^2))$. It is easy to see that $[S^2 \times D^4 \times I, \bar{\nu}] + [(S^2 \times S^4)_f, \bar{g}]$ is B -bordant to $S^2 \times D^4 \times I \cup_f S^2 \times D^4 \times I$ (Figure 2). Therefore the image of $[S^2 \times D^4 \times I, \bar{\nu}] + [(S^2 \times S^4)_f, \bar{g}]$ under Φ is just f . This shows the surjectivity of Φ . \square

By a standard computation using the Atiyah-Hirzebruch spectral sequence one can get an upper bound of the order of $\Omega_7^{\text{string}}(P_3(S^2))$. We put the calculation in the appendix (Lemma 11.1).

Lemma 6.6. $|\Omega_7^{\text{string}}(P_3(S^2))| \leq 24$.

With this and Lemma 6.5 we see that the order of $\overline{\text{MCG}}(S^2 \times D^4, \partial) \leq 24$. Indeed the equality holds. To see this, denote the quotient of \mathcal{N} by the action

of Θ_7 via connected-sum with homotopy spheres by $\overline{\mathcal{N}}$ and recall that we have homomorphisms

$$\overline{\text{MCG}}(S^2 \times D^4, \partial) \xrightarrow{\Psi} \overline{\mathcal{N}} \xrightarrow{S} \mathbb{Z}/8 \oplus \mathbb{Z}/24 \oplus \mathbb{Z}/2$$

with the first one being surjective and the second one injective. Let $a, b \in \overline{\text{MCG}}(S^2 \times D^4, \partial)$ be pre-images of N_{-1} and N_{-2} under Ψ respectively, then by formula 6.2 and Lemma 6.6, a has order 2, b has order 12 and $a \neq 6b$. This proves the following lemma.

Lemma 6.7. $\overline{\text{MCG}}(S^2 \times D^4, \partial)$ is isomorphic to $\mathbb{Z}/12 \oplus \mathbb{Z}/2$, detected by $s_2/2$ and s_3 .

Theorem 6.8. *There are isomorphisms*

$$\text{MCG}(S^2 \times D^4, \partial) \xrightarrow{\cong} \mathcal{I}(S^2 \times S^4) \xrightarrow{\cong} \mathbb{Z}/28 \oplus \mathbb{Z}/12 \oplus \mathbb{Z}/2$$

where the second isomorphism is given by the invariants $(s_1/8, s_2/2, s_3)$.

Remark 6.9. It's shown in [Sat69, Theorem II] that $\mathcal{I}(S^2 \times S^4)$ is isomorphic to $FC_4^3 \oplus \Theta_7$, where FC_4^3 is the group of isotopy classes of framed 4-knots in S^7 . By [Hae66] there is an exact sequence

$$\pi_4 SO_3 \rightarrow FC_4^3 \rightarrow C_4^3 \rightarrow \pi_3 SO_3$$

with $C_4^3 \cong \mathbb{Z}/12$. It's easy to see that there is a splitting $FC_4^3 \rightarrow \pi_4(S^2) \cong \pi_4 SO_3$ given by the linking number of the image of $S^4 \times \{0\}$ and $S^4 \times \{x_0\}$ in S^7 , where $x_0 \in S^2$ is a fixed point. Therefore $FC_4^3 \cong \mathbb{Z}/2 \oplus \mathbb{Z}/12$. In Theorem 6.8 we get the same result using a different method, and moreover, we have found explicit invariants detecting these summands. We will use these invariants to give explicit generators of $\mathcal{I}(S^2 \times S^4)$ in the next subsection.

6.3. Generators of $\mathcal{I}(S^2 \times S^4)$. Let $S^3 \times D^3 \hookrightarrow S^2 \times S^4$ be an embedding such that $S^3 \times 0$ represents a generator of $\pi_3(S^2 \times S^4) \cong \mathbb{Z}$, $\mathbf{t}_{Z,\alpha}: S^2 \times S^4 \rightarrow S^2 \times S^4$ be the Dehn twist with parameter $\alpha \in \pi_3(SO(4))$ (see §2.5). Assume that the corresponding \mathbb{R}^4 -bundle over S^4 has Euler class $e(\alpha) = \chi$ and spin Pontrjagin class $\frac{1}{2}p_1(\alpha) = \chi - 2b$ with $\chi, b \in \mathbb{Z}$. We are going to compute the S -invariants of $\mathbf{t}_{Z,\alpha}$ in terms of χ and b .

First we have the following observation on the mapping torus of a Dehn twist.

Lemma 6.10. *Let $S^3 \times D^3 \subset M$ be an embedding, $\alpha \in \pi_3 SO(4)$, f be the Dehn twist in $S^3 \times D^3$ parametrized by α . Then the mapping torus is*

$$M_f = (M \times S^1 - S^3 \times D^4) \cup_{\varphi} S^3 \times D^4$$

where $\varphi: S^3 \times S^3 \rightarrow S^3 \times S^3$, $(x, y) \mapsto (\alpha(y)x, y)$. Let E_{α} be the D^4 -bundle over S^4 corresponding to α , with the standard decomposition $\partial E_{\alpha} = S^3 \times D^4 \cup S^3 \times D^4$. Then $M_f = \partial(M \times D^2 \cup_{S^3 \times D^4} E_{\alpha})$.

To compute the S -invariant of \mathbf{t}_{α} , we use the coboundary of the mapping torus constructed in this lemma, $W = S^2 \times S^4 \times D^2 \cup_{S^3 \times D^4} E_{\alpha}$. Then $H_4(W) \cong \mathbb{Z}^3$ is generated by the base S^4 of E_{α} , denoted by $[S_b^4]$, a fiber D^4

of E_α , denoted by u , and $pt \times S^4 \times pt \subset S^2 \times S^4 \times D^2$, denoted by $[S_0^4]$. The intersection form with respect to these basis is the matrix S

$$S = \begin{pmatrix} \chi & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

Also it's easy to see from the geometric construction

$$\langle \frac{1}{2}p_1(W), [S_b^4] \rangle = \chi - 2b, \quad \langle \frac{1}{2}p_1(W), u \rangle = \langle \frac{1}{2}p_1(W), [S_0^4] \rangle = 0$$

therefore $\frac{1}{2}p_1(W) = (\chi - 2b)[S_b^4]^*$.

Let $x \in H^2(W)$ be a generator, a representing cocycle X of $Dx \in H_6(W, \partial W)$ is constructed as follows: fix $p \in S^2$, consider $p \times S^4 \times D^2 \subset S^2 \times S^4 \times D^2$, which is the Poincaré dual of a generator of $H^2(S^2 \times S^4 \times D^2)$. We need to extend this to E_α . First notice that the intersection of $S^3 \times D^4 \subset S^2 \times S^4 \times S^1$ with $p \times S^4 \times D^2$ is $S_p^1 \times D^4 \subset S^3 \times D^4$, where $S_p^1 \subset S^3$ is the Hopf fiber over $p \in S^2$. We cone this $S_p^1 \times D^4$ in $D^4 \times D^4 \subset E_\alpha$ and get $D_p^2 \times D^4 \subset D^4 \times D^4$. The intersection of $D_p^2 \times D^4$ with $D^4 \times S^3$ is $D_p^2 \times S^3$. The clutching function of the bundle E_α is $\varphi_\alpha: D^4 \times S^3 \rightarrow D^4 \times S^3$, $(x, y) \mapsto (\alpha(y) \cdot x, y)$. Thus the image $\varphi_\alpha(D_p^2 \times S^3) = \{(\alpha(y) \cdot x, y) \mid x \in D_p^2, y \in S^3\}$. We take the cone of this over the S^3 factor and get $D^2 \times D^4 \subset D^4 \times D^4$. Now let

$$X_p = p \times S^4 \times D^2 \cup_{S_p^1 \times D^4} D_p^2 \times D^4 \cup_{\varphi_\alpha(D_p^2 \times S^3)} D^2 \times D^4$$

Then by the construction X_p is the Poincaré dual of $x \in H^2(W)$. Now we compute the self-intersection of X_p . We take another point $p' \in S^2$ and obtain $X_{p'}$ by the same procedure. The circles S_p^1 and $S_{p'}^1$ form a Hopf link in S^3 and $D_p^2 \cap D_{p'}^2 = 0 \in D^4$. Therefore from the construction $X_p \cap X_{p'} = S_b^4$, the base sphere of the bundle E_α . Let $\bar{\alpha}, \bar{\beta} \in H^4(W, \partial W)$ be the cohomology classes in the definition of the S -invariant, with these geometric constructions we have

$$D(\bar{\alpha}) = [S_b^4], \quad D(\bar{\beta}) = (\chi - 2b)u$$

Therefore

$$\begin{aligned} \frac{1}{8}s_1 &= \frac{1}{8}\langle \bar{\beta}^2, [W, \partial W] \rangle = \frac{1}{8}(\chi - 2b)\langle \frac{1}{2}p_1(W), u \rangle = 0 \\ \frac{1}{2}s_2 &= \frac{1}{2}\langle (\bar{\alpha}^2 + \bar{\alpha}\bar{\beta}), [W, \partial W] \rangle = \frac{1}{2}(\chi + \chi - 2b) = \chi - b \in \mathbb{Z}/12 \\ s_3 &= \langle \bar{\alpha}\bar{\beta}, [W, \partial W] \rangle = \chi - 2b = \chi \in \mathbb{Z}/2 \end{aligned}$$

Thus we have the following explicit geometric description of $\mathcal{I}(S^2 \times S^4)$.

Theorem 6.11. *In the mapping class group $\mathcal{I}(S^2 \times S^4) \cong \mathbb{Z}/28 \times \mathbb{Z}/12 \times \mathbb{Z}/2$, the $\mathbb{Z}/28$ -factor is generated by the disk supported diffeomorphism \mathbf{f}_c , the $\mathbb{Z}/12$ -factor is generated by the Dehn twists $\mathbf{t}_{Z,p}$ (with $\chi = 0, b = 1$) and the $\mathbb{Z}/2$ -factor is generated by the Dehn twist $\mathbf{t}_{Z,\epsilon}$ (with $\chi = 1, b = 0$).*

Note that Theorem 2.14 for spin manifolds is a corollary of this theorem and the fact that there is a surjective homomorphism $\text{MCG}(S^2 \times D^4, \partial) \rightarrow \text{MCG}(M)$ (see Lemma 7.5).

Remark 6.12. There is another natural Dehn twist, namely the Dehn twist along the embedded $S^2 \times D^4 \subset S^2 \times S^4$ parametrized by the non-trivial element of $\pi_4(SO(3)) \cong \mathbb{Z}/2$. Using similar construction as above it's easy to see that the coboundary W has trivial intersection form $H_4(W) \times H_4(W) \rightarrow \mathbb{Z}$, therefore the S -invariant of this Dehn twist is 0. Thus this Dehn twist is isotopic to the identity, which is not geometrically obvious.

7. DETERMINATION OF $\mathcal{K}(M)$

In this section we show that all diffeomorphisms in $\mathcal{K}(M, D)$ come from $(S^2 \times D^4, \partial)$. Together with the S -invariant we are able to determine the value of $\mathcal{K}(M)$, as given in the tables in Theorem 5.1.

Recall that for a 6-manifold M which looks like a complete intersection we have a decomposition 3.1

$$M = N \# (S^3 \times S^3)^{\#r}$$

with $N = (S^2 \times D^4) \cup (S^2 \times D^4)$. A diffeomorphism f of $(S^2 \times D^4, \partial)$ extends to a diffeomorphism of M by the identity on the complement, and such a diffeomorphism is clearly in $\mathcal{K}(M, D)$.

Let $f: (M, D) \rightarrow (M, D)$ be a diffeomorphism such that $[f] \in \mathcal{K}(M, D)$, since f induces the identity on $H_2(M)$, and $\pi_2 SO(4) = 0$, after isotopy we may assume that f is the identity on the first copy of $S^2 \times D^4 \subset M$. Let $M_0 = (S^2 \times D^4) \# (S^3 \times S^3)^{\#r}$ be the connected-sum of the second copy of $S^2 \times D^4$ with $\#_r(S^3 \times S^3)$, then we get a self-diffeomorphism f of M_0 which is the identity on the boundary. M_0 is stably parallelizable, choosing a framing of the stable normal bundle, we may compare it with the induced framing by the differential of f and get a map $df: M_0 \rightarrow SO$. By obstruction theory $[M_0, SO] \cong H^3(M_0; \mathbb{Z}) \cong H$, thus we may regard the differential df as an element in H . Also it's easy to see that $\pi_3(M_0) \cong \mathbb{Z} \oplus \mathbb{Z}^{2r}$ and $i_*: \pi_3(M_0) \rightarrow \pi_3(M) \cong \mathbb{Z}/d \oplus \mathbb{Z}^{2r}$ is the mod d reduction on the first summand. Since by lemma 4.2 the induced isomorphism f_* on $\pi_3(M)$ is the identity, on $\pi_3(M_0)$ f_* must have the form

$$\begin{pmatrix} I & 0 \\ B & I \end{pmatrix}$$

where $B: \mathbb{Z}^{2r} \rightarrow d\mathbb{Z}$ is a homomorphism. Conversely for any diffeomorphism of M_0 rel ∂ with this property, we extend it by the identity on the first copy of $S^2 \times D^4$ to a diffeomorphism of M whose action on π_3 is the identity.

Proposition 7.1. *Any class in $\mathcal{K}(M, D)$ can be obtained by a diffeomorphism of M_0 rel ∂ , whose action on $\pi_3(M_0)$ is the identity, and whose differential is trivial.*

Proof. We use notations for Dehn twists given in section 2.5.1. Let φ be the composition of the Dehn twists $\mathbf{t}_{Z\#S_1,\alpha}$ and $\mathbf{t}_{S_1,-\alpha}$, where the Euler characteristic of α is d ,

$$\varphi = \mathbf{t}_{S_1,-\alpha} \circ \mathbf{t}_{Z\#S_1,\alpha}.$$

Denote $e_0 = [Z]$ and $e_2 = [T_1] \in \pi_3(M_0)$. Then by Lemma 2.12, $\varphi_*(e_2) = d \cdot e_0 + e_2$, and φ_* fixes the other generators of $\pi_3(M_0)$. As a diffeomorphism of M_0 the differential of φ is trivial. Extend φ to a diffeomorphism of M , still denoted by φ . Then clearly φ^* is trivial on $H^3(M)$. By Lemma 4.2 $\varphi_* = \text{id}$ on $\pi_3(M)$. This implies $v_{x,p}(\varphi) = 0$, therefore $[\varphi] \in \mathcal{K}(M, D)$.

By Lemma 6.10, the mapping torus of φ is $M_\varphi = \partial(M \times D^2 \cup E_\alpha \cup E_{-\alpha})$. The signature of E_α and $E_{-\alpha}$ is 1 and -1 respectively. We may compute the S -invariant of φ as in §6.3, the result is

$$s_1 \in 8\mathbb{Z}/8 \cdot 28, \quad s_2 \in 2\mathbb{Z}/24, \quad s_3 = d \in \mathbb{Z}/2$$

By Theorem 6.11 we may realize these values as the S -invariant of some diffeomorphism φ' of $S^2 \times D^4$ rel ∂ . Extend φ' to a diffeomorphism of M , still denoted by φ' . Then $\varphi' \in \mathcal{K}(M, D)$. Let $\varphi_1 = \varphi'^{-1} \circ \varphi$, then φ_1 is isotopic to id_M .

Given a class in $\mathcal{K}(M, D)$ with a representing diffeomorphism f , the composition $\varphi_1 \circ f$ represents the same class, but has a simpler action on $\pi_3(M_0)$. Iterate this process, finally we will get a representing diffeomorphism f whose action on $\pi_3(M_0)$ is trivial.

Let $i^*: H^4(M_f) \rightarrow H^4((M_0)_f)$ be the induced homomorphism. Since $v_{x,p}(f) = 0$, there is a $y' \in H^4(M_f)$ such that $x^2 - dy' = 0$ and $\frac{1}{4}(p_1(M_f) - 2ky') = 0$. From this we have $i^*(y') = 0$ and $p_1((M_0)_f) = i^*(p_i(M_f)) = 0$. Similar to the proof of [Kre78, Lemma 2], this shows that the differential $df: M_0 \rightarrow SO$ is homotopic to the constant map. \square

Let $\text{MCG}_0(M_0, \partial)$ be the mapping class group of self-diffeomorphisms of M_0 rel ∂ , whose action on $\pi_3(M_0)$ is the identity, and whose differential is trivial. Then Proposition 7.1 states that $\text{MCG}_0(M_0, \partial) \rightarrow \mathcal{K}(M, D)$ is surjective. Let $\overline{\text{MCG}}_0(M_0, \partial)$ be the cokernel of $\Theta_7 \rightarrow \text{MCG}_0(M_0, \partial)$. By similar constructions and proofs to that of Lemma 6.5 we have

Lemma 7.2. *There is a surjective homomorphism*

$$\Phi_1: \Omega_7^{\text{string}}(P_3(S^2) \times K(\mathbb{Z}^{2r}, 3)) \rightarrow \overline{\text{MCG}}_0(M_0, \partial)$$

Combining this lemma with Lemma 6.5 and Proposition 7.1, we have the following

Proposition 7.3. *There is a commutative diagram*

$$\begin{array}{ccc} \Omega_7^{\text{string}}(P_3(S^2)) & \xrightarrow{\Phi} & \overline{\text{MCG}}(S^2 \times D^4, \partial) \\ \downarrow & & \downarrow \\ \Omega_7^{\text{string}}(P_3(S^2) \times K(\mathbb{Z}^{2r}, 3)) & \xrightarrow{\Phi_1} & \overline{\text{MCG}}_0(M_0, \partial) \longrightarrow \mathcal{K}(M, D)/\Theta_7 \end{array}$$

in which the horizontal maps are surjective.

A straightforward computation by the Atiyah-Hirzebruch spectral sequence shows that the natural inclusions of both factors induce an isomorphism (a sketch of the proof is in the Appendix (Lemma 11.3))

$$\Omega_7^{\text{string}}(P_3(S^2)) \oplus \Omega_7^{\text{string}}(K(\mathbb{Z}^{2r}, 3)) \rightarrow \Omega_7^{\text{string}}(P_3(S^2) \times K(\mathbb{Z}^{2r}, 3))$$

But elements in the direct summand $\Omega_7^{\text{string}}(K(\mathbb{Z}^{2r}, 3))$ correspond to diffeomorphisms of $\#_r(S^3 \times S^3)$ whose action on π_3 is the identity, and whose differential is the identity on the normal bundle of any embedded S^3 , hence is the identity by Theorem 3.1. Therefore we have

Lemma 7.4. *The homomorphism $\text{MCG}(S^2 \times D^4, \partial) \rightarrow \mathcal{K}(M, D)$ is surjective.*

Lemma 7.5. *There is a commutative diagram*

$$\begin{array}{ccc} \text{MCG}(S^2 \times D^4, \partial) & \xrightarrow{S} & \mathbb{Z}^3/L \\ \downarrow & & \downarrow \\ \mathcal{K}(M, D) & \xrightarrow{S} & \mathbb{Z}^3/\bar{L}(M) \end{array}$$

in which the vertical maps are surjective and the horizontal maps are injective.

Proof. Let f be a self-diffeomorphism of $S^2 \times D^4 \text{ rel } \partial$, $N = (S^2 \times D^5) \cup_f (S^2 \times D^5) \in \mathcal{N}$ be the corresponding 7-manifold and $W \rightarrow \mathbb{CP}^\infty \times B\text{Spin}$ be the null-bordism for computing the S -invariant of N . Assume

$$M = (S^2 \times D^4) \cup_\varphi (S^2 \times D^4 \#_r(S^3 \times S^3))$$

with glueing diffeomorphism $\varphi: S^2 \times S^3 \rightarrow S^2 \times S^3$. Extend f to a diffeomorphism of M by the identity on $S^2 \times D^4 \#_r(S^3 \times S^3)$. Let $\bar{\nu}: M \rightarrow B = \mathbb{CP}^\infty \times K(\mathbb{Z}, 4) \times B\text{Spin}$ be the lifting of the normal Gauss map of M , as in the definition of the S -invariant of f in section 5. We may assume that $\bar{\nu}$ maps $S^2 \times D^4$ to $\mathbb{CP}^\infty \times B\text{Spin}$. Now define (see Figure 3)

$$V = W \cup_{S^2 \times S^3 \times I \times I} ((S^2 \times D^4 \#_r(S^3 \times S^3)) \times I \times I) \rightarrow B$$

where W maps to $\mathbb{CP}^\infty \times B\text{Spin}$ and the map on $S^2 \times D^4 \times I \times I$ is $\bar{\nu}|_{S^2 \times D^4} \times \text{id} \times \text{id}$. Again in Wall's formula of the non-additivity of the signature the correcting term vanishes, therefore $\text{sign}(V) = 0$. Then V can be taken as the null B -bordism for computing the S -invariant of f . Similar to the proof of diagram 6.3, the $(S^2 \times D^4 \#_r(S^3 \times S^3)) \times I^2$ part doesn't contribute to the characteristic numbers. This proves the commutativity of the diagram. \square

Proof of Theorem 5.1 (spin case). In the next section we show that the forgetful homomorphism $\mathcal{K}(M, D) \rightarrow \mathcal{K}(M)$ is an isomorphism (Proposition 8.4). Therefore it suffices to compute the group $\mathcal{K}(M, D)$. By Theorem




FIGURE 3.

6.8 and Lemma 7.5, $\mathcal{K}(M, D)$ is isomorphic to the image of the surjective homomorphism

$$\mathbb{Z}/28 \oplus \mathbb{Z}/12 \oplus \mathbb{Z}/2 \rightarrow \mathbb{Z}^3/L(M)$$

where $L(M)$ is the lattice generated by the column vectors of the following table

$$(7.1) \quad \begin{array}{c|cccccc} & e_1 & e_2 & e_3 & e_4 & e_5 & e_6 \\ \hline s_1/8 & -28 & 0 & 0 & (k^2 - k)/2 & k^2 & 0 \\ s_2/2 & 0 & 12 & 0 & (d^2 - d(2k - 1))/2 & d^2 - 2dk & k - d \\ s_3 & 0 & 0 & 2 & d(2k - 1) & 4dk & -2k \end{array}$$

The finite abelian group $\mathbb{Z}^3/L(M)$ can have only 2-, 3- and 7-torsions, of exponents at most 2, 1, and 1 respectively. Therefore we may determine the structure of $\mathbb{Z}^3/L(M)$ by tensorizing it with $\mathbb{Z}/4$, $\mathbb{Z}/3$ and $\mathbb{Z}/7$ respectively. After solving this elementary arithmetic problem we get the tables in Theorem 5.1.

- (1) modulo 3: only the image of $s_2/2$ involves. Since

$$\gcd((d^2 - d(2k - 1))/2, d^2 - 2dk, k - d)$$

is divisible by 3 if and only if $d \equiv 0 \pmod{3}$, We have

$$\mathbb{Z}^3/L'(M) \otimes \mathbb{Z}/3 = \begin{cases} \mathbb{Z}/3 & \text{if } d \equiv 0 \pmod{3} \\ 0 & \text{if } d \not\equiv 0 \pmod{3} \end{cases}$$

- (2) modulo 7: only the image of $s_1/8$ involves. Since $\gcd((k^2 - k)/2, k^2)$ is divisible by 7 if and only if $d - l \equiv 0 \pmod{7}$, we have

$$\mathbb{Z}^3/L'(M) \otimes \mathbb{Z}/7 = \begin{cases} \mathbb{Z}/7 & \text{if } d - l \equiv 0 \pmod{7} \\ 0 & \text{if } d - l \not\equiv 0 \pmod{7} \end{cases}$$

- (3) modulo 4: the entries in table (7.1) are 8-periodic with respect to d . Therefore we only need to compute the lattice for $d = 0, \dots, 7$. The results are in the table in Theorem 5.1.

□

8. THE BOUNDARY DIFFEOMORPHISM IS TRIVIAL

In this section we prove that in the spin case the forgetful homomorphism $\mathcal{K}(M, D) \rightarrow \mathcal{K}(M)$ is an isomorphism. Recall that the relative mapping class group $\mathcal{K}(M, D)$ is related to the mapping class group $\mathcal{K}(M)$ by an exact sequence (see (5.2))

$$\pi_1(GL_6(\mathbb{R})) \rightarrow \mathcal{K}(M, D) \rightarrow \mathcal{K}(M) \rightarrow 1.$$

The image of the generator of $\pi_1(GL_6(\mathbb{R}))$ in $\mathcal{K}(M, D)$ is called the **boundary diffeomorphism**, it is given as follows: let $\alpha: [0, 1] \rightarrow SO(n)$ be a loop representing a generator of $\pi_1 SO(n)$, identify a collar neighborhood of $\partial(M - \mathring{D})$ with $S^{n-1} \times [0, 1]$, define

$$\delta: S^{n-1} \times [0, 1] \rightarrow S^{n-1} \times [0, 1], \quad (x, t) \mapsto (\alpha(t) \cdot x, t)$$

and extend it to a diffeomorphism of $M - \mathring{D}$ by identity on the complement of the collar neighborhood. The forgetful map $\mathcal{K}(M, D) \rightarrow \mathcal{K}(M)$ is an isomorphism if and only if the boundary diffeomorphism represents the trivial element in $\mathcal{K}(M, D)$.

The mapping torus of the boundary diffeomorphism of M is $S^1 \times M$, but the preferred spin structure is the twisted one when restricted to $S^1 \times D$. Although we know that the normal map $S^1 \times M \rightarrow \mathbb{CP}^\infty \times K(\mathbb{Z}, 4) \times B\text{Spin}$ is a boundary in corresponding spin bordism group, there isn't an explicit construction for such a spin coboundary. Instead we will make use of a spin^c -coboundary V constructed as follows: Let E be the non-trivial linear D^6 -bundle over S^2 , i. e.

$$E = (D^6 \times D^2)_1 \cup_g (D^6 \times D^2)_2$$

where $g: D^6 \times S^1 \rightarrow D^6 \times S^1$ is given by $(x, t) \mapsto (\alpha(t) \cdot x, t)$, with $\alpha: S^1 \rightarrow SO(6)$ representing the non-trivial element of the fundamental group. Let $S^5 \times D^2 \subset (D^6 \times D^2)_1$ be the standard inclusion. Also we have $S^5 = \partial D \subset M \subset S^1 \times M$, a tubular neighborhood of this embedding is diffeomorphic to $S^5 \times D^2$, let

$$(8.1) \quad V = (M \times D^2) \cup_{S^5 \times D^2} E,$$

then $\partial V = (M \times S^1 - S^5 \times D^2) \cup_g S^5 \times D^2$ is diffeomorphic to $S^1 \times M$, and the map $S^1 \times M \rightarrow \mathbb{CP}^\infty \times K(\mathbb{Z}, 4)$ extends to V , where on $M \times D^2$ it is the natural extension, and on E it represents a generator of $H^2(E)$. Let $z \in H^2(E)$ be a generator, L be the complex line bundle over V with $c_1(L) = (0, z) \in H^2(V) = H^2(M) \oplus H^2(E)$. Then the spin-structure on $\nu V \oplus L$ restricts to the non-trivial spin-structure on $S^1 \times D$. It's easy to see that the intersection pairing $H_4(V) \times H_4(V) \rightarrow \mathbb{Z}$ is trivial.

Theorem 8.1. *Let M be a simply-connected spin homology \mathbb{CP}^3 , $D \subset M$ be an embedded 6-disk, $N = M - \mathring{D}$, $f: N \rightarrow N$ be the boundary diffeomorphism. Then f is isotopic to the identity rel ∂N .*

Proof. The idea of the proof is the following (compare the proof of [Kre78, Proposition 3]): Do surgery on $S^1 \times D \subset S^1 \times M$ with the twisted framing we get $X = (N \times I)_{f \cup \text{id}}(N \times I)$. If X bounds a compact manifold U such that $(U, N \times I)$ is a relative h -cobordism, then f is isotopic to id_N rel ∂N . Now $H_3(X) \cong \mathbb{Z}$ and the homology classes are spherical. Choose an embedding $S^3 \times D^4 \subset X$ representing a generator of $H_3(X)$ and do surgery, we get a simply-connected spin homology $S^2 \times S^5$, denoted by Y . If Y is diffeomorphic to $S^2 \times S^5$, then we may take U to be the union of the trace of the second surgery and $S^2 \times D^6$ along Y . Thus in this case f is isotopic to id_N rel ∂N . If Y is diffeomorphic to $S^2 \times S^5 \# \Sigma$, where Σ is an exotic 7-sphere, then f is isotopic rel ∂N to a disk supported diffeomorphism corresponding to Σ . Then we determine Σ and look at the image of the corresponding disk supported diffeomorphism in $\mathcal{K}(M, D)$ in table (7.1).

The diffeomorphism type of Y is determined by the Kreck-Stolz S -invariants $S_i(Y)$ ($i = 1, 2, 3$). These invariants can be computed from a spin^c -coboundary of the map $Y \rightarrow \mathbb{CP}^\infty$ representing a generator of $H^2(Y)$ (c. f. [KrSt91, (2.7)]). We construct such a coboundary as follows: Let $S^1 \times M \rightarrow \mathbb{CP}^\infty$ be a map representing a generator of $H^2(S^1 \times M)$, this map extends to the trace W' of the above surgeries where on the attached 2-handle and 4-handle it is a constant map. There is a unique spin structure on W' , which restricts to the non-trivial spin structure on $S^1 \times M$. Take the spin^c -coboundary V described above and let $W = W' \cup V$. Then $W \rightarrow \mathbb{CP}^\infty$ is a spin^c -coboundary of $Y \rightarrow \mathbb{CP}^\infty$.

The S -invariants of Y are

$$S_i(Y) = \langle e^{z/2} \text{ch}(E_i) \hat{A}(W), [W, \partial W] \rangle \in \mathbb{Q}/\mathbb{Z} \quad (i = 1, 2, 3)$$

where E_i is a (virtual) complex vector bundle pulled back from a vector bundle over \mathbb{CP}^∞ . From the construction of V it is easy to see $z^2 = 0$, therefore

$$S_i(Y) = \langle \text{ch}(E_i) \hat{A}(W), [W, \partial W] \rangle,$$

which are linear combinations of

$$\langle x^4, [W, \partial W] \rangle, \quad \langle x^2 \frac{p_1}{2}, [W, \partial W] \rangle, \quad \langle (\frac{p_1}{2})^2, [W, \partial W] \rangle.$$

Explicit formulas are given in [KrSt91, (2.4)].

Now $H_4(W) \cong \mathbb{Z}^2$ with basis $\{u, v\}$, where u is a generator of $H_4(M)$, v is represented by a cycle C such that $C \cap V = D^2 \times S^2 \subset D^2 \times M$, where $S^2 \subset M$ represents a generator of $H_2(M)$. The intersection pairings of W are

$$u \cdot u = 0, \quad u \cdot v = 1, \quad v \cdot v = \lambda$$

(thus $\text{sign}(W) = 0$) and we have the following evaluations

$$\langle x^2, u \rangle = d, \quad \langle x^2, v \rangle = 0, \quad \langle \frac{p_1}{2}, u \rangle = 2k, \quad \langle \frac{p_1}{2}, v \rangle = \mu$$

where λ and μ are integers depending on the framing of the 3-surgery. From this we have

$$\begin{aligned}\langle x^4, [W, \partial W] \rangle &= -d^2\lambda \\ \langle x^2 \frac{p_1}{2}, [W, \partial W] \rangle &= d(\mu - 2k\lambda) \\ \langle (\frac{p_1}{2})^2, [W, \partial W] \rangle &= 2k(\mu - 2\lambda k) + 2k\mu.\end{aligned}$$

And the S -invariants of Y are

$$\begin{aligned}S_1(Y) &= \frac{1}{2^5 \cdot 7}(2k(\mu - 2\lambda k) + 2k\mu) \\ S_2(Y) &= \frac{-1}{2^3 \cdot 3}(d^2\lambda + d(\mu - 2\lambda k)) \\ S_3(Y) &= \frac{-1}{2 \cdot 3}(4d^2\lambda + d(\mu - 2\lambda k)).\end{aligned}$$

We analyse these values case by case. Since the boundary diffeomorphism is an element of order 2, we only need to consider the 2-primary part. When d is odd, notice that $S_2(Y)$ takes values in $\mathbb{Z}/12$, we must have $\lambda + \mu \equiv 0 \pmod{2}$. Changing the framing of the embedding $S^3 \times D^4 \subset X$, μ will be changed to $\mu + b$ and λ will be changed to $\lambda + 2a + b$, where a and b are arbitrary integers. Take $b = -\mu$ and $a = (\mu - \lambda)/2$, we get $\lambda = \mu = 0$ and hence all the $S_i(Y)$'s are 0.

When $d \equiv 0 \pmod{4}$, $S_3(Y)$ is automatically 0. We may choose a and b such that $\mu = 0$ and $\lambda = 0$ or 1. Thus $S_2(Y) = 0$, $S_1(Y) = 0$ or $k^2/2^3 \cdot 7$. In the latter case, when l is even, $k^2 \equiv 0 \pmod{2^3}$ hence $S_1(Y) = 0$. When l is odd, from the simplification of the lattice $L(M)$ in the proof of Theorem 5.1, disk supported diffeomorphisms don't contribute to $\mathcal{K}(M, D)$. Therefore the boundary diffeomorphism is trivial in $\mathcal{K}(M, D)$.

When $d = 4r + 2$, $S_3(Y)$ is again automatically 0. For λ odd we set $b = \mu + 2$, for λ even we set $b = \mu$. This makes $S_2(Y) = 0$, $S_1(Y) = (2k(\mu + 1) - \lambda k^2)/2^3 \cdot 7$ when λ is odd, or $S_1(Y) = (2k\mu - \lambda k^2)/2^3 \cdot 7$ when λ is even. Choosing appropriate a such that $\lambda = 0$ or 1 according to the parity of λ and μ , we will get $S_1(Y) = 0$. For example when λ is odd and μ is even

$$2k(\mu + 1) - \lambda k^2 \equiv (2k - k^2) \equiv 2^2(2r + 3l)(2r + 3l + 1) \equiv 0 \pmod{2^3}.$$

Therefore $S_1(Y) = 0$. The other cases are similar. \square

Lemma 8.2. *Let M be a closed n -manifold ($n \geq 3$), $D, D' \subset M$ be closed disjointly embedded n -disks. Let $N = M - \mathring{D}$, f be the boundary diffeomorphism in a neighborhood of ∂D ; $N' = N - \mathring{D}'$, f' be the boundary diffeomorphism in a neighborhood of $\partial D'$. Assume f is isotopic to id_N rel ∂N . Then f' is isotopic to either $\text{id}_{N'}$ or f' rel $\partial N'$.*

Proof. We have the following observation from the proof of [Wal63, Hilfsatz]: Let $h: N \times I \rightarrow N \times I$ be an isotopy between f and id_N , x be the center of D' , then $h(x \times I)$ is isotopic to $x \times I$. By the ambient isotopy theorem,

we may modify h such that $h(x \times I) = x \times I$. Now $h(D' \times I)$ is a tubular neighborhood of $x \times I$, by the tubular neighborhood theorem we may assume that $h(D' \times I) = D' \times I$ and it is a bundle map, thus $h|_{\partial D' \times I}$ is either the identity or the boundary diffeomorphism. Reparametrize the boundary of $\partial N' \times I$, we may view h as an isotopy between f and $\text{id}_{N'}$ or $f' \text{ rel } \partial N'$. \square

Theorem 8.3. *Let M_1, M_2 be closed oriented n -manifolds, $N_i = M_i - D_i$. Assume that the boundary diffeomorphism is isotopic to $\text{id}_{N_i} \text{ rel } \partial N_i$ for $i = 1, 2$. Then for $N = (M_1 \# M_2) - D$, the boundary diffeomorphism is also isotopic to $\text{id}_N \text{ rel } \partial N$.*

Proof. Let $D' \subset N_1$ be an embedded disk disjoint from D_1 , $N'_1 = N_1 - D'$, then the boundary diffeomorphism f in a neighborhood of ∂D_1 is isotopic rel $\partial N'_1$ to either $\text{id}_{N'}$ or the boundary diffeomorphism f' in a neighborhood of $\partial D'$. In the second case f' is isotopic to $\text{id}_{N_2} \text{ rel } \partial D'$. Put these two isotopies together we have an isotopy from f to $\text{id}_N \text{ rel } \partial D_1$ \square

Combining Theorem 8.1, Theorem 8.3 and the fact that for $\#r(S^3 \times S^3)$ the boundary diffeomorphism is isotopic to $\text{id} \text{ rel } D$ ([Kre78, Lemma 3]), we have proved

Proposition 8.4. *The forgetful homomorphism $\mathcal{K}(M, D) \rightarrow \mathcal{K}(M)$ is an isomorphism.*

9. THE NON-SPIN CASE

9.1. Determination of $\mathcal{K}(M, D)$. In this section we determine the kernel $\mathcal{K}(M, D)$ in the non-spin case, along the lines of section 5–7.

First of all the definition of the S -invariant. In the non-spin case the normal type is

$$p: B = \mathbb{CP}^\infty \times K(\mathbb{Z}, 4) \times B\text{Spin} \rightarrow BSO$$

where the map on \mathbb{CP}^∞ is the classifying map of the canonical complex line bundle η over \mathbb{CP}^∞ . For a diffeomorphism $f \in \mathcal{K}(M, D)$, there is a unique lifting $g: M_f \rightarrow B$ of the normal Gauss map. The corresponding bordism group is $\Omega_*(B, p) = \Omega_*^{\text{spin}}(\mathbb{CP}^\infty \times K(\mathbb{Z}, 4); \eta)$, the twisted spin cobordism group of $\mathbb{CP}^\infty \times K(\mathbb{Z}, 4)$. A standard computation using the Atiyah-Hirzebruch spectral sequence shows that $\Omega_7^{\text{spin}}(\mathbb{CP}^\infty \times K(\mathbb{Z}, 4); \eta) = 0$ (c. f. [Kre18, Theorem 6]). Now $H^4(B; \mathbb{Z})$ is generated by $x^2 \in H^4(\mathbb{CP}^\infty)$, $y \in H^4(K(\mathbb{Z}, 4))$ and $\frac{p_1}{2} \in H^4(B\text{Spin})$. Note that

$$g^*\left(\frac{p_1}{2}\right) = \frac{p_1}{2}(\nu M_f - g^*\eta) = \frac{1}{2}(-d - 48l - d)g^*y = -(d + 24l)g^*y$$

Therefore $\ker g^*$ is generated by $x^2 - dy$ and $\frac{p_1}{2} + (d + 24l)y$. Let $G: W^8 \rightarrow B$ be a null-cobordism of $g: M_f \rightarrow B$, $G^*(x^2 - dy) = \alpha$, $G^*\left(\frac{p_1}{2} + (d + 24l)y\right) = \beta$,

there are cohomology classes $\overline{\alpha}$ and $\overline{\beta} \in H^4(W, \partial W)$. We may define

$$\begin{cases} \sigma_1 &= \langle \overline{\alpha} \cup \overline{\alpha}, [W, \partial W] \rangle \\ \sigma_2 &= \langle \overline{\alpha} \cup \overline{\beta}, [W, \partial W] \rangle \\ \sigma_3 &= \langle \overline{\beta} \cup \overline{\beta}, [W, \partial W] \rangle \end{cases}$$

and the S -invariant $S(f) = [(\sigma_1, \sigma_2, \sigma_3)] \in \mathbb{Z}^3 / \bar{L}(M)$. Similar to Proposition 5.2 we have

Proposition 9.1. *The generalized Kreck-Stolz invariant defines an injective homomorphism*

$$S: \mathcal{K}(M, D) \rightarrow \mathbb{Z}^3 / \bar{L}(M)$$

In the non-spin case, we are able to show that the S -invariant of the boundary diffeomorphism is always zero (see §9.3). Therefore in this case we also have an isomorphism $\mathcal{K}(M, D) \xrightarrow{\cong} \mathcal{K}(M)$.

The lattice $\bar{L}(M)$ is given by the values of σ_i on the generators of the subgroup of $\Omega_8^{\text{spin}}(\mathbb{CP}^\infty \times K(\mathbb{Z}, 4); \eta)$ consisting of bordism classes of signature 0. These generators are given as follows: (a sketch of the computation is given in the Appendix)

$$(9.1) \quad \begin{aligned} e_1 &= B^8 - 8 \cdot 28\text{HP}^2 \\ e_2 &= [\mathbb{CP}^2 \times \mathbb{CP}^2 \rightarrow \mathbb{CP}^\infty] - \text{HP}^2 \\ e_3 &= [\mathbb{CP}^4 \rightarrow \mathbb{CP}^\infty] - \text{HP}^2 \\ e_4 &= [\text{HP}^2 \rightarrow K(\mathbb{Z}, 4)] - \text{HP}^2 \\ e_5 &= [S^4 \times S^4 \rightarrow K(\mathbb{Z}, 4)] \\ e_6 &= [\mathbb{CP}^2 \times S^4 \rightarrow \mathbb{CP}^\infty \times K(\mathbb{Z}, 4)] \end{aligned}$$

The lattice $\bar{L}(M)$ is generated by the column vectors in the following table (9.2)

	e_1	e_2	e_3	e_4	e_5	e_6
σ_1	0	6	1	d^2	$2d^2$	$-2d$
σ_2	0	-6	-3	$-d(d+24l-1)$	$-2d(d+24l)$	$3d+24l$
σ_3	$-8 \cdot 28$	8	8	$(d+24l-1)^2 - 1$	$2(d+24l)^2$	$-4(d+24l)$

In the non-spin case, M can be decomposed as a union

$$M = (D^4 \widetilde{\times} S^2) \cup (D^4 \widetilde{\times} S^2) \# r(S^3 \times S^3)$$

where $D^4 \widetilde{\times} S^2$ is the nontrivial D^4 -bundle over S^2 .

Next we compute $\text{MCG}(D^4 \widetilde{\times} S^2, \partial)$. Let \mathcal{N}' denote the set of equivalence classes of simply-connected oriented non-spin 7-manifolds N with $H_*(N) \cong H_*(S^2 \times S^5)$. Manifolds of this type are determined by the Kreck-Stolz S -invariant [KrSt88]. Let $S^4 \widetilde{\times} S^2$ be the non-trivial linear S^4 -bundle over S^2 , $\mathcal{I}(S^4 \widetilde{\times} S^2)$ be the group consisting of isotopy classes of diffeomorphisms acting trivially on homology, there is a surjective map $\Psi: \mathcal{I}(S^4 \widetilde{\times} S^2) \rightarrow \mathcal{N}'$,

which fits in a commutative diagram

$$\begin{array}{ccccc} \mathrm{MCG}(D^4 \widetilde{\times} S^2, \partial) & \longrightarrow & \mathcal{I}(S^4 \widetilde{\times} S^2) & \xrightarrow{\Psi} & \mathcal{N}' \\ & & \downarrow S & & \downarrow S \\ & & \mathbb{Z}^3/L & \xrightarrow{=} & \mathbb{Z}^3/L \end{array}$$

where the two maps in the first row are surjective and the two vertical homomorphisms are injective.

The lattice L is generated by the column vectors of the following table.

$$(9.3) \quad \begin{array}{c|ccc} & e_1 & e_2 & e_3 \\ \hline x^4 & 0 & 6 & 1 \\ x^2 \frac{p_1}{2} & 0 & -6 & -3 \\ (\frac{p_1}{2})^2 & -8 \cdot 28 & 8 & 8 \end{array}$$

After simplification we see $\mathbb{Z}^3/L \cong \mathbb{Z}/4 \oplus \mathbb{Z}/3 \cdot 8 \cdot 28$, the summands are determined by the invariants

$$\lambda = x^4 + 3x^2 \frac{p_1}{2} + \left(\frac{p_1}{2}\right)^2 \in \mathbb{Z}/4$$

$$\mu = 6x^4 + 10x^2 \frac{p_1}{2} + 3\left(\frac{p_1}{2}\right)^2 \in \mathbb{Z}/3 \cdot 8 \cdot 28$$

Next we consider the action of Θ_7 . Let $\Sigma = \partial E_8$, a generator of Θ_7 . Let W be the null-cobordism we use to compute the invariants of a diffeomorphism f , then for the composition of f with the local diffeomorphism induced by Σ , we may use $(W \natural E_8) \# (-8\mathbb{H}P^2)$ as the null-cobordism. It turns out that the value of λ is unchanged, and the value of μ is changed to $\mu(f) - 3 \cdot 8$. Therefore the natural homomorphism $\Theta_7 \rightarrow \mathrm{MCG}(D^4 \widetilde{\times} S^2, \partial)$ is injective. Denote the cokernel by $\overline{\mathrm{MCG}}(D^4 \widetilde{\times} S^2, \partial)$, let $\overline{\mathcal{N}'}$ be the quotient of the action of Θ_7 on \mathcal{N}' , then there are homomorphisms

$$\overline{\mathrm{MCG}}(D^4 \widetilde{\times} S^2, \partial) \rightarrow \overline{\mathcal{N}'} \rightarrow \mathbb{Z}/4 \oplus \mathbb{Z}/24$$

where the first one is surjective and the second one is injective.

An upper bound of $\overline{\mathrm{MCG}}(D^4 \widetilde{\times} S^2, \partial)$ is obtained as follows. Let η be the pull back of the canonical complex line bundle over S^2 to the 3rd stage Postnikov tower $P_3(S^2)$. Then similar to Lemma 6.5, there is a surjective homomorphism

$$\Omega_7^{\mathrm{string}}(P_3(S^2); \eta) \twoheadrightarrow \overline{\mathrm{MCG}}(D^4 \widetilde{\times} S^2, \partial)$$

with $\Omega_7^{\mathrm{string}}(P_3(S^2); \eta) \cong \mathbb{Z}/6$ (computation in the Appendix, see Lemma 11.2).

Let ξ_m be an oriented 4-dimensional real vector bundle over $\mathbb{C}P^2$ with

$$w_2(\xi_m) = 0, \quad e(\xi_m) = 1, \quad p_1(\xi_m) = 4m + 2 \quad (m \in \mathbb{Z}).$$

These bundles can be obtained by pulling back suitable vector bundles over S^4 . Let $W_m = D(\xi_m)$ be the disk bundle and $N_m = \partial W_m = S(\xi_m)$. Then

we may use $W_m \# (-\mathbb{H}P^2)$ to compute the invariants of N_m and get

$$\lambda(N_m) = 4m^2 + 6m \in \mathbb{Z}/4, \quad \mu(N_m) = 12m^2 + 16m \in \mathbb{Z}/24.$$

For $m = -1$, $(\lambda(N_{-1}), \mu(N_{-1})) = (-2, -4)$, which has order 6.

Theorem 9.2. *There are isomorphisms*

$$\mathrm{MCG}(D^4 \widetilde{\times} S^2, \partial) \xrightarrow{\cong} \mathcal{I}(S^4 \widetilde{\times} S^2) \xrightarrow{\cong} \mathbb{Z}/6 \cdot 28$$

where the second isomorphism is given by $\mu/4$.

Lemma 9.3. *There is a commutative diagram*

$$\begin{array}{ccc} \mathrm{MCG}(D^4 \widetilde{\times} S^2, \partial) & \xrightarrow{S} & \mathbb{Z}^3/L \\ \downarrow & & \downarrow \\ \mathcal{K}(M) & \xrightarrow{S} & \mathbb{Z}^3/\bar{L}(M) \end{array}$$

in which the vertical homomorphisms are surjective and the horizontal ones are injective.

Simplify table (9.2) in the same way as in the simplification of table (9.3), using the linear combination

$$\lambda = \sigma_1 + 3\sigma_2 + \sigma_3$$

$$\mu = 6\sigma_1 + 10\sigma_2 + 3\sigma_3$$

we have the following table (in which $\delta = d/2$, recall that in the non-spin case d is even)

$$\begin{array}{c|cccccc} \mu/4 & 6 \cdot 28 & 0 & -\delta^2 - 48\delta l + 2\delta + 432l^2 - 36l & -2(\delta^2 + 96\delta l - 864l^2) & 3(\delta - 4l) \\ \lambda & 0 & -4 & -4\delta^2 - 96\delta l + 2\delta + 576l^2 - 48l & -2(4\delta^2 + 48\delta l - 576l^2) & 6(\delta - 4l) \end{array}$$

and $\mathbb{Z}^3/\bar{L}(M) \cong \mathbb{Z}^2/L(M)$ with $L(M)$ determined by this table.

Now we have a homomorphism $\mathbb{Z}/6 \cdot 28 \rightarrow \mathbb{Z}^2/L(M)$, with $1 \mapsto [(-1, -2)]$ (since $(\lambda(N_{-1}), \mu(N_{-1})) = (-2, -4)$). We summarize the above calculations in the following commutative diagram

$$\begin{array}{ccc} \Theta_7 & & \\ \downarrow & \searrow^{1 \mapsto 6} & \\ \mathrm{MCG}(D^4 \widetilde{\times} S^2, \partial) & \xrightarrow[\cong]{\mu/4} & \mathbb{Z}/6 \cdot 28 \\ \downarrow & & \downarrow^{1 \mapsto [(-1, -2)]} \\ \mathcal{K}(M) & \xrightarrow{(\mu/4, \lambda)} & \mathbb{Z}^2/L(M) \end{array}$$

Therefore to determine the group $\mathcal{K}(M)$ we only need to compute the order of $[(1, 2)]$ in $\mathbb{Z}^2/L(M)$. Since the order is necessarily a factor of $6 \cdot 28$, we only need to calculate modulo 3, 7 and 8 respectively. The result is presented in the table in Theorem 5.1 (non-spin case).

9.2. Generators of $\text{MCG}(S^4 \widetilde{\times} S^2)$. Now we show that the Dehn twist $\mathbf{t}_{Z,\rho}$ is a generator of $\mathcal{I}(S^4 \widetilde{\times} S^2)$ by computing the invariant $\mu/4$ and hence prove Theorem 2.14 in the non-spin case. The computation is similar to that in §6.3. Let $\mathbf{t}_{Z,\alpha}$ be the Dehn twist in $S^3 \times D^3 \subset S^4 \widetilde{\times} S^2$ with parameter $\alpha \in \pi_3 SO(4)$, $W = E_\alpha \cup_{S^3 \times D^4} S^4 \widetilde{\times} S^2 \times D^2$ be the coboundary of the mapping torus of $\mathbf{t}_{Z,\alpha}$ constructed in Lemma 6.10. Then $H_4(W) \cong \mathbb{Z}^3$ is generated by the base S^4 of E_α , denoted by $[S_b^4]$, a fiber D^4 of E_α , denoted by u , and $S^4 \times pt \times pt$, denoted by $[S_0^4]$. The intersection matrix with respect to this basis is

$$S = \begin{pmatrix} \chi & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

Note that the pull-back of $\frac{1}{2}p_1$ equals to

$$\frac{1}{2}(p_1(\nu W) - p_1(\eta)) = -(\chi - 2b)[S_b^4]^*.$$

Let $x \in H^2(W)$ be a generator, then by a similar construction as in §6.3 the Poincaré dual of x^2 is $[S_b^4]$. With these data it is easy to compute

$$\frac{1}{4}\mu(\mathbf{t}_{Z,\alpha}) = -\chi + 5b \in \mathbb{Z}/6 \cdot 28.$$

Especially, the Dehn twist $\mathbf{t}_{Z,\rho}$ (with $\chi = 0$, $b = 1$) is a generator of $\mathcal{I}(S^4 \widetilde{\times} S^2)$.

9.3. The boundary diffeomorphism is trivial. In the non-spin case the S -invariant of the boundary diffeomorphism can be calculated directly from a spin^c -coboundary V whose construction is almost the same as the spin case (see equation (8.1)), the only modification is that now the complex line bundle L over V has $c_1(L) = (x, z) \in H^2(V) = H^2(M) \oplus H^2(E)$, where x and z are generators of corresponding cohomology groups. The spin-structure on $\nu V \oplus L$ restricts to the non-trivial spin-structure on $S^1 \times D$. Thus V can be used as the coboundary in the definition of the S -invariant. It's easy to see that the intersection pairing $H_4(V) \times H_4(V) \rightarrow \mathbb{Z}$ is trivial, therefore all the characteristic numbers $\sigma_i = 0$, ($i = 1, 2, 3$). This shows that the S -invariant of the boundary diffeomorphism is zero, hence it is isotopic to the identity rel D . This finishes the proof of Theorem 5.1 in the non-spin case.

10. GROUP THEORETICAL PROPERTIES

10.1. Generators of $\text{MCG}(M)$. In this subsection we write down explicit generators of $\text{MCG}(M)$ for $r \geq 1$, hence prove Theorem 2.13. For the notations of the Dehn twists we stick to the description before Theorem 2.13.

First of all, similar to the case of surfaces, from a presentation of the symplectic group $\text{Sp}(2r, \mathbb{Z})$ (c.f. [Bir71, Theorem 1]) it's easy to see that the

set of Dehn twists $\mathbf{e}_i, \mathbf{f}_j$ ($1 \leq i, j \leq r$) and $\mathbf{f}_{i,i+1}$ ($1 \leq i \leq r-1$) maps to a set of generators of $\mathrm{Sp}(2r, \mathbb{Z})$.

Next we give generators of the Torelli group $\mathcal{I}(M)$. Recall that there is a central extension

$$0 \rightarrow \mathcal{K}(M) \rightarrow \mathcal{I}(M) \xrightarrow{\tau} (H \times H)/\Delta_{d,k} \rightarrow 0$$

Let a_i, b_j and a'_i, b'_j ($1 \leq i, j \leq r$) be the standard symplectic basis of $0 \times H$ and $H \times 0$ respectively. Let

$$\mathbf{a}_i = \mathbf{t}_{S_i, \rho}, \quad \mathbf{b}_j = \mathbf{t}_{T_j, \rho}, \quad \mathbf{a}'_i = (\mathbf{t}_{S_i, \epsilon})^{-1} \circ \mathbf{g}_i, \quad \mathbf{b}'_j = (\mathbf{t}_{T_j, \epsilon})^{-1} \circ \mathbf{h}_j,$$

by the definition of τ , these are pre-images of a_i, b_j and a'_i, b'_j respectively. The commutators of these diffeomorphisms are in $\mathcal{K}(M)$. Now we compute the S -invariants of the commutators.

We only need to consider the commutators of these diffeomorphisms when the corresponding embedded 3-spheres of the Dehn twists have non-empty intersection. First of all it is shown in [Kry02, Theorem 2] that $[\mathbf{a}_i, \mathbf{b}_i] = \mathbf{f}_c$, a generator of the disk supported diffeomorphisms. Next we compute the S -invariant of the commutator

$$[\mathbf{a}'_1, \mathbf{b}_1] = \mathbf{a}'_1 \mathbf{b}_1 \mathbf{a}'_1{}^{-1} \mathbf{b}_1{}^{-1} = (\mathbf{t}_{S_1, \epsilon})^{-1} \circ \mathbf{g}_1 \circ \mathbf{t}_{T_1, \rho} \circ (\mathbf{g}_1)^{-1} \circ \mathbf{t}_{S_1, \epsilon} \circ (\mathbf{t}_{T_1, \rho})^{-1}.$$

As in §6.3 we take

$$W = M \times D^2 \cup E_{1-1} \cup E_{01} \cup E_2 \cup E_{01-1} \cup E_1 \cup E_{2-1}$$

where the E 's are corresponding D^4 -bundles over S^4 . From the geometric construction it's seen that $H_4(W) \cong \mathbb{Z}^{11}$, with basis given by the base 4-spheres of E 's, denoted by e_i , and u_1, \dots, u_4 , represented by the unions of suitable fibers D^4 of the bundles, and a generator of $H_4(M)$. The signature of W is 0.

Let $x \in H^2(W)$ be a generator, its Poincaré dual X is constructed as in §6.3. From this we have $D(\bar{\alpha}) = X \cap X = e_{01} + e_{01-1}$. From the evaluation of $p_1(W)$ on the basis and the intersection matrix of W we have

$$D(\bar{\beta}) = e_{01} + e_{-1} - e_{01-1} - e_{1-1} + 2u_1 - e_2 + 2u_4$$

Therefore $D\bar{\alpha} \cdot D\bar{\alpha} = \chi(E_{01}) + \chi(E_{01-1}) = 0$, $D\bar{\alpha} \cdot D\bar{\beta} = 2$.

In the spin case we have

$$\begin{aligned} \frac{1}{2}s_2 &= \frac{1}{2}(D\bar{\alpha} \cdot D\bar{\alpha} - D\bar{\alpha} \cdot D\bar{\beta}) = -1 \\ s_3 &= D\bar{\alpha} \cdot D\bar{\beta} = 2 = 0 \in \mathbb{Z}/2 \end{aligned}$$

(The s_1 -invariant is also 0, but this is not relevant, since disk supported diffeomorphisms will produce all possible values of the s_1 -invariant.) Similar computations of the other commutators show that the s_3 -invariant is always 0. From the simplification of the lattices in the proof of Theorem 5.1, we see when d is odd the kernel $\mathcal{K}(M)$ is generated by commutators of these Dehn twists; when d is even, we need an additional generator $\mathbf{z} = \mathbf{t}_{Z, \rho}$ to produce diffeomorphisms with nontrivial s_3 -invariant (see Theorem 2.14).

In the non-spin case we have

$$\mu/4 = \frac{1}{4}(6D\bar{\alpha} \cdot D\bar{\alpha} + 10D\bar{\alpha} \cdot D\bar{\beta} + 3D\bar{\beta} \cdot D\bar{\beta}) = 5$$

which is a unit in $\mathcal{K}(M) = \mathbb{Z}/2^a \oplus \mathbb{Z}/3^b \oplus \mathbb{Z}/7^c$.

Therefore the Torelli group $\mathcal{I}(M)$ is generated by the Dehn twists $\mathbf{a}_i, \mathbf{b}_j, (\mathbf{e}_i)^{-1} \circ \mathbf{g}_i, (\mathbf{f}_j)^{-1} \circ \mathbf{h}_j$ ($1 \leq i, j \leq r$), and \mathbf{z} when M is spin and d is even. Together with the Dehn twists $\mathbf{e}_i, \mathbf{f}_j$ ($1 \leq i, j \leq r$) and $\mathbf{f}_{i,i+1}$ ($1 \leq i \leq r-1$) they form a set of generators of $\text{MCG}(M)$. This proves Theorem 2.13.

The presentation of $\mathcal{J}(M)$ in Theorem 2.15 can be directly deduced from the above computation.

10.2. Lower homology groups. In this subsection we compute the lower homology groups of $\text{MCG}(M)$. The abelianization $(\text{MCG}(M))^{\text{ab}}$ equals to the first homology group $H_1(\text{MCG}(M))$. When $r = 0$ the group $\text{MCG}(M)$ itself is an abelian group. So we assume $r \geq 1$. Notice that from the commuting relations of the generators of $\mathcal{I}(M)$ described in the previous subsection, one also obtains the abelianization of $\mathcal{I}(M)$. We summarize the results in the following proposition. Recall that $H = H^3(M; \mathbb{Z})$.

Proposition 10.1. *When M is spin*

$$(\mathcal{I}(M))^{\text{ab}} = \begin{cases} (H \times H)/\Delta_{d,k} & d \text{ odd} \\ (H \times H)/\Delta_{d,k} \oplus \mathbb{Z}/2 & d \text{ even} \end{cases}$$

When M is non-spin $(\mathcal{I}(M))^{\text{ab}} = (H \times H)/\Delta_{d,k}$.

Proof of Theorem 2.7, Theorem 2.8, Corollary 2.9 and Corollary 2.10. Denote $\Gamma_r = \text{Sp}(2r, \mathbb{Z})$, we compute $(\text{MCG}(M))^{\text{ab}} = H_1(\text{MCG}(M))$ using the Hochschild-Serre spectral sequence of the exact sequence

$$1 \rightarrow \mathcal{I}(M) \rightarrow \text{MCG}(M) \rightarrow \Gamma_r \rightarrow 1$$

The relevant E_2 -terms are as follows: first of all (c. f. [BCRR18, Appendix])

$$E_{1,0}^2 = H_1(\Gamma_r) = \begin{cases} \mathbb{Z}/12 & r = 1 \\ \mathbb{Z}/2 & r = 2 \\ 0 & r \geq 3 \end{cases}, \quad E_{2,0}^2 = H_2(\Gamma_r) = \begin{cases} 0 & r = 1 \\ \mathbb{Z} \oplus \mathbb{Z}/2 & r = 2 \\ \mathbb{Z} & r \geq 3 \end{cases}$$

Secondly from the exact sequence of Γ_r -modules

$$0 \rightarrow H \rightarrow (H \times H)/\Delta_{d,k} \rightarrow H/d \rightarrow 0$$

and the fact (c. f. [Kra19, Lemma A.2]) that for an abelian group A the co-invariants $(\mathbb{Z}^{2g} \otimes A)_{\Gamma_r} = 0$, we have $E_{0,1}^2 = H_1(\mathcal{I}(M))_{\Gamma_r} = 0$ except for the case M spin and d even, where $E_{0,1}^2 = \mathbb{Z}/2$.

To determine the differential $d: E_{2,0}^2 \rightarrow E_{0,1}^2$ (in the case M spin, d even and $r \geq 2$), we compare the spectral sequences of the upper and lower lines

of the following commutative diagram

$$\begin{array}{ccccc}
 \mathcal{I}(\#_r(S^3 \times S^3)) & \longrightarrow & \text{MCG}(\#_r(S^3 \times S^3)) & \longrightarrow & \Gamma_r \\
 \downarrow & & \downarrow & & \downarrow = \\
 \mathcal{I}(M) & \longrightarrow & \text{MCG}(M) & \longrightarrow & \Gamma_r
 \end{array}$$

where $\mathcal{I}(\#_r(S^3 \times S^3))^{\text{ab}} = H$ is the standard Γ_r -module [Kry02], we see that the differential $d: E_{2,0}^2 = H_2(\Gamma_r) \rightarrow E_{0,1}^2$ factors through $H_2(\Gamma_r) \rightarrow H_{\Gamma_r} \rightarrow E_{0,1}^2$. But $H_{\Gamma_r} = 0$. Therefore the differential $d: E_{2,0}^2 \rightarrow E_{0,1}^2$ is trivial.

So we have a short exact sequence

$$0 \rightarrow E_{0,1}^2 \rightarrow H_1(\text{MCG}(M)) \rightarrow H_1(\Gamma_r) \rightarrow 0$$

When $r \geq 3$, $H_1(\Gamma_r) = 0$ therefore $H_1(\text{MCG}(M)) = E_{0,1}^2$. When $r = 1$ or 2 , there is a splitting $H_1(\Gamma_r) = H_1(\text{MCG}(\#_r(S^3 \times S^3))) \rightarrow H_1(\text{MCG}(M))$, therefore $H_1(\text{MCG}(M)) = H_1(\Gamma_r) \oplus E_{0,1}^2$. This proves Theorem 2.7.

Since the differential on $E_{2,0}^2$ is trivial, this term survives to infinity and therefore there is a surjective homomorphism $H_2(\text{MCG}(M)) \rightarrow H_2(\Gamma_r)$. The short exact sequence of groups $\text{Diff}_0(M) \rightarrow \text{Diff}(M) \rightarrow \text{MCG}(M)$ induces a fiber bundle $B\text{Diff}_0(M) \rightarrow B\text{Diff}(M) \rightarrow B\text{MCG}(M)$ with simply-connected fiber $B\text{Diff}_0(M)$. There is a surjective homomorphism $H_2(B\text{Diff}(M)) \rightarrow H_2(\text{MCG}(M))$. Composed with the surjective homomorphism $H_2(\text{MCG}(M)) \rightarrow H_2(\Gamma_r)$ we get a surjective homomorphism $H_2(B\text{Diff}(M)) \rightarrow H_2(\Gamma_r)$. This proves Corollary 2.9.

To compute $H_2(\text{MCG}(M); \mathbb{Q})$ ($r \geq 3$) we may use the short exact sequence

$$1 \rightarrow (H \times H)/\Delta_{d,k} \rightarrow \text{MCG}(M)/\mathcal{K}(M) \rightarrow \Gamma_r \rightarrow 1$$

since $\mathcal{K}(M)$ is a finite group. The relevant terms in the Hochschild-Serre spectral sequence are

- (1) $E_{2,0}^2 = H_2(\Gamma_r; \mathbb{Q}) \cong \mathbb{Q}$;
- (2) $E_{1,1}^2 = H_1(\Gamma_r; (H \times H)/\Delta_{d,k} \otimes \mathbb{Q}) = 0$ (cf. [Kra19, Lemma A.3]);
- (3)

$$\begin{aligned}
 E_{0,2}^2 &= H_0(\Gamma_r; H_2((H \times H)/\Delta_{d,k}; \mathbb{Q})) \\
 &= (\wedge^2 H \otimes \mathbb{Q}) \otimes_{\Gamma_r} \mathbb{Q} = \wedge^2 \mathbb{Q}^{2r} \otimes_{\text{Sp}(2r, \mathbb{Q})} \mathbb{Q} \cong \mathbb{Q}
 \end{aligned}$$

where the reason for the last isomorphism is that the second exterior power $\wedge^2 \mathbb{Q}^{2r}$ of the standard representation of $\text{Sp}(2r, \mathbb{Q})$ is isomorphic to the direct sum of the trivial representation \mathbb{Q} and a non-trivial irreducible representation.

- (4) $E_{2,1}^2 = H_2(\Gamma_r; (H \times H)/\Delta_{d,k} \otimes \mathbb{Q}) = 0$ (cf. [Kra19, Lemma A.3]);
- (5) $E_{3,0}^2 = H_3(\Gamma_r; \mathbb{Q}) = 0$ when $r \geq 3$ (cf. [Bor74]).

For this data it's easy to see $H_2(\text{MCG}(M); \mathbb{Q}) \cong \mathbb{Q}^2$.

Corollary 2.10 is a direct consequence of the Leray-Serre spectral sequence of the fiber bundle $B\text{Diff}_0(M) \rightarrow B\text{Diff}(M) \rightarrow \text{BMCG}(M)$, and the facts $H_2(\text{MCG}(M); \mathbb{Q}) \cong \mathbb{Q}^2$ and $H_2(B\text{Diff}(M); \mathbb{Q}) \cong \mathbb{Q}^4$ ([GaRW18, §5.3]). \square

10.3. Group theoretical properties.

Theorem 10.2. *The center of $\text{MCG}(M)$ is $\mathcal{K}(M)$.*

Proof. Recall that elements in $\mathcal{K}(M)$ are represented by diffeomorphisms in $S^2 \times D^4$ or $D^4 \tilde{\times} S^2$, hence they commute with the generators of $\text{MCG}(M)$ represented by the Dehn twists described in Theorem 2.13. Therefore $\mathcal{K}(M)$ is in the center of $\text{MCG}(M)$. We will show that $\text{MCG}(M)/\mathcal{K}(M)$ is centerless. By equations (1.11) and (1.12), there is a short exact sequence

$$0 \rightarrow H \rightarrow \text{MCG}(M)/\mathcal{K}(M) \rightarrow \text{Sp}(\pi_3(M)) \rightarrow 1$$

where $\text{Sp}(\pi_3(M)) = (\mathbb{Z}/d)^{2r} \rtimes \text{Sp}(2r, \mathbb{Z})$. It's easy to compute the center of the semi-direct product $(\mathbb{Z}/d)^{2r} \rtimes \text{Sp}(2r, \mathbb{Z})$ — it is trivial when $d \neq 2$, and is $\{(0, \pm I)\}$ when $d = 2$ (cf. [Kra19, Lemma A.2]).

When the center of $\text{Sp}_3(\pi_3(M))$ is trivial, the center of $\text{MCG}(M)/\mathcal{K}(M)$ is contained in H . The action of $\text{Sp}(2r, \mathbb{Z})$ on H is the standard linear action, with 0 being the only fixed element. This shows the center of $\text{MCG}(M)/\mathcal{K}(M)$ is trivial. When the center of $\text{Sp}_3(\pi_3(M))$ is $(0, \pm I)$, let $a \in \text{MCG}(M)/\mathcal{K}(M)$ be an element whose image in $\text{Sp}(2r, \mathbb{Z})$ is $-I$, then for any $b \in H$, the commutator $[a, b] = (-I)(b) - b = -2b$. Therefore a is not in the center. \square

Proof of Theorem 2.11. To show that $\text{MCG}(M)/\mathcal{K}(M)$ is virtually torsion free we construct torsion free subgroups of $\text{MCG}(M)/\mathcal{K}(M)$ of finite index as follows: Let $\text{Sp}(2r, \mathbb{Z})[m]$ be the principal congruence subgroup of $\text{Sp}(2r, \mathbb{Z})$ of level m ($m \geq 3$), then it is a torsion-free subgroup of finite index of $\text{Sp}(\pi_3(M))$. Let $(\text{MCG}(M)/\mathcal{K}(M))[m]$ be the preimage of $\text{Sp}(2r, \mathbb{Z})[m]$ in $\text{MCG}(M)/\mathcal{K}(M)$, it fits in the exact sequence

$$0 \rightarrow H \rightarrow (\text{MCG}(M)/\mathcal{K}(M))[m] \rightarrow \text{Sp}(2r, \mathbb{Z})[m] \rightarrow 1$$

Therefore $(\text{MCG}(M)/\mathcal{K}(M))[m]$ is a torsion-free subgroup of finite index in $\text{MCG}(M)/\mathcal{K}(M)$.

Finally we show that $\text{MCG}(M)/\mathcal{K}(M)$ is residually finite. From the commutative diagram of exact sequences 4.6 we see $\text{MCG}(\#r(S^3 \times S^3))/\Theta_7$ is a subgroup of $\text{MCG}(M)/\mathcal{K}(M)$ of finite index. It suffices to show that $\text{MCG}(\#r(S^3 \times S^3))/\Theta_7$ is residually finite. This group is a subgroup of $H \rtimes \Gamma_r$ of index 2^{2r} . Both H and Γ_r are residually finite, then the semi-direct product $H \rtimes \Gamma_r$ is also residually finite (e. g. by Theorem II of [Hew94] which says that an extension of a residually finite group by a residually finite group is residually finite if and only if the extension is residually virtually splittable). Therefore $\text{MCG}(\#r(S^3 \times S^3))/\Theta_7$ is residually finite, hence also $\text{MCG}(M)/\mathcal{K}(M)$. \square

11. APPENDIX: COMPUTATION OF THE BORDISM GROUPS

In the appendix we sketch the computation of the bordism groups used in the paper.

Let $P = P_3(S^2)$ be the 3rd stage Postnikov tower of S^2 , η be the complex line bundle over P , whose first Chern class equals to a generator of $H^2(P)$. We compute the string bordism groups $\Omega_7^{\text{string}}(P)$ and $\Omega_7^{\text{string}}(P; \eta)$ using Atiyah-Hirzebruch spectral sequence.

We first compute the (co)-homology groups of P . P is a fibration over $K(\mathbb{Z}, 2)$ with fiber $K(\mathbb{Z}, 3)$ and k -invariant $x^2 \in H^4(K(\mathbb{Z}, 2))$, where $x \in H^2(K(\mathbb{Z}, 2))$ is a generator. The homology groups of $K(\mathbb{Z}, 3)$ were computed in the classical paper [EiMa54, Theorem 23.1-23.4]

	3	4	5	6	7	8
$H_*(K(\mathbb{Z}, 3))$	\mathbb{Z}	0	$\mathbb{Z}/2$	0	$\mathbb{Z}/3$	$\mathbb{Z}/2$
$H_*(K(\mathbb{Z}, 3); \mathbb{Z}/2)$	$\mathbb{Z}/2$	0	$\mathbb{Z}/2$	$\mathbb{Z}/2$	0	$\mathbb{Z}/2$

Now we apply the Leray-Serre spectral sequence to the fibration $K(\mathbb{Z}, 3) \rightarrow P \rightarrow K(\mathbb{Z}, 2)$, it's easy compute the homology groups of P in low dimensions

	0	1	2	3	4	5	6	7	8
$H_*(P)$	\mathbb{Z}	0	\mathbb{Z}	0	0	$\mathbb{Z}/2$	0	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	$\mathbb{Z}/2$

From this we can deduce the cohomology groups with $\mathbb{Z}/2$ -coefficients

	0	1	2	3	4	5	6	7	8
$H^*(P; \mathbb{Z}/2)$	$\mathbb{Z}/2$	0	$\mathbb{Z}/2$	0	0	$\mathbb{Z}/2$	$\mathbb{Z}/2$	$\mathbb{Z}/2$	$(\mathbb{Z}/2)^2$

Now if we look at the Leray-Serre spectral sequence for the cohomology with $\mathbb{Z}/2$ -coefficients, the differential $d: E_2^{0,6} \rightarrow E_2^{2,5}$ must be zero. From this we obtain the cup product structure of $H^*(P; \mathbb{Z}/2)$ as follows: let $x \in H^2(P; \mathbb{Z}/2)$, $a \in H^5(P; \mathbb{Z}/2)$, $b \in H^6(P; \mathbb{Z}/2)$ be generators, then $ax \in H^7(P; \mathbb{Z}/2)$ is a generator, and $H^8(P; \mathbb{Z}/2)$ has generators bx and c , where c restricts to a generator of $H^8(K(\mathbb{Z}, 3); \mathbb{Z}/2)$.

We also need the information on the Steenrod operations in $H^*(P; \mathbb{Z}/2)$. This is obtained as follows. Let P' be the restriction of P on $S^2 \subset K(\mathbb{Z}, 2)$. Then from the above spectral sequence we see that $H^*(P; \mathbb{Z}/2) \rightarrow H^*(P'; \mathbb{Z}/2)$ are isomorphisms for $*$ = 6, 7, 8; and $H^5(P; \mathbb{Z}/2) \rightarrow H^5(P'; \mathbb{Z}/2)$ is injective. Since the k -invariant of P' is trivial, P' is homotopy equivalent to $S^2 \times K(\mathbb{Z}, 3)$. Let $x \in H^2(K(\mathbb{Z}, 2); \mathbb{Z}/2)$, $y \in H^3(K(\mathbb{Z}, 3); \mathbb{Z}/2)$ be generators, then $Sq^2 y \in H^5(K(\mathbb{Z}, 3); \mathbb{Z}/2)$ and $y^2 \in H^6(K(\mathbb{Z}, 3); \mathbb{Z}/2)$ are generators, and $H^5(P'; \mathbb{Z}/2)$ has generators xy and $Sq^2 y$. The image of $H^5(P; \mathbb{Z}/2) \rightarrow H^5(P'; \mathbb{Z}/2)$ is generated by $Sq^2 y$. By the Adem relation $Sq^2 Sq^2 y = Sq^3 Sq^1 y = 0$. Also $Sq^2 y^2 = 0$. Therefore $Sq^2: H^5(P; \mathbb{Z}/2) \rightarrow H^7(P; \mathbb{Z}/2)$ and $Sq^2: H^6(P; \mathbb{Z}/2) \rightarrow H^8(P; \mathbb{Z}/2)$ are trivial.

Now we are ready to apply the Atiyah-Hirzebruch spectral sequence to compute $\Omega_*^{\text{string}}(P)$ and $\Omega_*^{\text{string}}(P; \eta)$. The string cobordism groups up to dimension 7 are (the first six groups are the same as the stable stems and

the 7-th by [Gia71])

	0	1	2	3	4	5	6	7
Ω_*^{string}	\mathbb{Z}	$\mathbb{Z}/2$	$\mathbb{Z}/2$	$\mathbb{Z}/24$	0	0	$\mathbb{Z}/2$	0

In the Atiyah-Hirzebruch spectral sequence of $\Omega_*^{\text{string}}(P)$, the differentials d_2 are dual to Sq^2 . Thus $d_2: E_{7,0}^2 \rightarrow E_{5,1}^2$, $d_2: E_{7,1}^2 \rightarrow E_{5,2}^2$ and $d_2: E_{8,0}^2 \rightarrow E_{6,1}^2$ are trivial. Therefore we have exact sequences

$$\begin{aligned} 0 \rightarrow G \rightarrow \Omega_7^{\text{string}}(P) \rightarrow \mathbb{Z}/2 \oplus \mathbb{Z}/3 \rightarrow 0 \\ \mathbb{Z}/2 \rightarrow G \rightarrow \mathbb{Z}/2 \rightarrow 0 \end{aligned}$$

Therefore we have

Lemma 11.1. $|\Omega_7^{\text{string}}(P_3(S^2))| \leq 24$.

In the Atiyah-Hirzebruch spectral sequence of $\Omega_*^{\text{string}}(P; \eta)$, the differentials d_2 are dual to $Sq^2 + x$, where $x \in H^2(P; \mathbb{Z}/2)$ is the nontrivial element. Thus $d_2: E_{7,1}^2 \rightarrow E_{5,2}^2$ is an isomorphism, $d_2: E_{7,0}^2 \rightarrow E_{5,1}^2$ is surjective and $d_2: E_{8,0}^2 \rightarrow E_{6,1}^2$ is trivial. Therefore we have

Lemma 11.2. $\Omega_7^{\text{string}}(P_3(S^2); \eta) \cong \mathbb{Z}/2 \oplus \mathbb{Z}/3$.

Now we apply the Atiyah-Hirzebruch spectral sequence to compute $\Omega_*^{\text{string}}(P_3(S^2) \times K(\mathbb{Z}^{2r}, 3))$ and $\Omega_*^{\text{string}}(P_3(S^2) \times K(\mathbb{Z}^{2r}, 3); \eta)$. For simplicity denote $K(\mathbb{Z}^{2r}, 3)$ by K . Let $x \in H^2(P; \mathbb{Z}/2)$, $b_i \in H^3(K; \mathbb{Z}/2)$ ($i = 1, \dots, 2r$) be generators. Then $H^5(K; \mathbb{Z}/2)$ is generated by $Sq^2 b_i$ ($i = 1, \dots, 2r$). In the spin case in the Atiyah-Hirzebruch spectral sequence the differential $d_2: E_{7,1}^2 \rightarrow E_{5,2}^2$ is dual to $Sq^2: H^5(P \times K; \mathbb{Z}/2) \rightarrow H^7(P \times K; \mathbb{Z}/2)$. The mixed term in $H^5(P \times K; \mathbb{Z}/2)$ is $H^2(P; \mathbb{Z}/2) \otimes H^3(K; \mathbb{Z}/2)$, with generators $a \otimes b_i$. The mixed term at $E_{7,0}^2 = H_7(P \times K)$ is $H_2(P) \otimes H_5(K) \cong (\mathbb{Z}/2)^{2r}$, which reduces to $H_2(P; \mathbb{Z}/2) \otimes H_5(K; \mathbb{Z}/2)$ isomorphically. Note that $Sq^2(x \otimes b_i) = x \otimes Sq^2 b_i$ are the generators of $H^2(P; \mathbb{Z}/2) \otimes H^5(K; \mathbb{Z}/2)$. And at $E_{6,1}^2 = H_6(P \times K; \mathbb{Z}/2)$ there is no mixed term. Therefore no mixed terms survive in the spectral sequence. In the non-spin case, in the Atiyah-Hirzebruch spectral sequence for $\Omega_7^{\text{string}}(P \times K; \eta)$, the differential $d_2: E_{7,1}^2 \rightarrow E_{5,2}^2$ is dual to $Sq^2 + x \cup$. Since $x^2 = 0 \in H^4(P; \mathbb{Z}/2)$, again we have $(Sq^2 + x \cup)(x \otimes b_i) = x \otimes Sq^2 b_i$. Therefore we have

Lemma 11.3. *There are isomorphisms*

$$\begin{aligned} \Omega_7^{\text{string}}(P_3(S^2)) \oplus \Omega_7^{\text{string}}(K(\mathbb{Z}^{2r}, 3)) &\rightarrow \Omega_7^{\text{string}}(P_3(S^2) \times K(\mathbb{Z}^{2r}, 3)) \\ \Omega_7^{\text{string}}(P_3(S^2); \eta) \oplus \Omega_7^{\text{string}}(K(\mathbb{Z}^{2r}, 3)) &\rightarrow \Omega_7^{\text{string}}(P_3(S^2) \times K(\mathbb{Z}^{2r}, 3); \eta) \end{aligned}$$

Finally we compute the twisted spin bordism group $\Omega_8^{\text{spin}}(\mathbb{CP}^\infty \times K(\mathbb{Z}, 4); \eta)$, where η is the nontrivial complex line bundle over \mathbb{CP}^∞ . This group can be identified with

$$\pi_{10}^S(M\text{Spin} \wedge \text{Th}(\eta)) = \widetilde{\Omega}_{10}^{\text{spin}}(\text{Th}(\eta)) = \Omega_{10}^{\text{spin}}(\mathbb{CP}^\infty \times K(\mathbb{Z}, 4)) / \Omega_{10}^{\text{spin}}(\text{pt} \times K(\mathbb{Z}, 4)).$$

We apply the Atiyah-Hirzebruch-Leray-Serre spectral sequence to the fiber bundle $K(\mathbb{Z}, 4) \rightarrow \mathbb{CP}^\infty \times K(\mathbb{Z}, 4) \rightarrow \mathbb{CP}^\infty$. The groups $\Omega_*^{\text{spin}}(K(\mathbb{Z}, 4))$ can be computed straightforwardly by the Atiyah-Hirzebruch spectral sequence. We have $\widetilde{\Omega}_i^{\text{spin}}(K(\mathbb{Z}, 4)) = 0$ for $i \leq 7$ and $i \neq 4$, $\widetilde{\Omega}_4^{\text{spin}}(K(\mathbb{Z}, 4)) \cong H_4(K(\mathbb{Z}, 4)) \cong \mathbb{Z}$, $\widetilde{\Omega}_8^{\text{spin}}(K(\mathbb{Z}, 4)) \cong \mathbb{Z} \oplus \mathbb{Z}$ (c. f. [Kre18, Theorem 8]). From this data, it's easy to see that

$$\begin{aligned} & \Omega_8^{\text{spin}}(\mathbb{CP}^\infty \times K(\mathbb{Z}, 4); \eta) \\ & \cong \Omega_8^{\text{spin}} \oplus \widetilde{\Omega}_8^{\text{spin}}(\mathbb{CP}^\infty; \eta) \oplus \widetilde{\Omega}_8^{\text{spin}}(K(\mathbb{Z}, 4)) \oplus H_4(\mathbb{CP}^\infty) \otimes H_4(K(\mathbb{Z}, 4)) \end{aligned}$$

where the last factor $H_4(\mathbb{CP}^\infty) \otimes H_4(K(\mathbb{Z}, 4)) \cong \mathbb{Z}$ is generated by the natural inclusion $\mathbb{CP}^2 \times S^4 \rightarrow \mathbb{CP}^\infty \times K(\mathbb{Z}, 4)$.

REFERENCES

- [BCRR18] D. Benson, C. Campagnolo, A. Ranicki, C. Rovi, *Cohomology of symplectic groups and Meyer's signature theorem*, Algebraic and Geometric Topology 18 (2018), 4069-4091.
- [Bir71] J. Birman, *On Siegel's modular group*, Math. Ann. 191, 59-68 (1971).
- [Bor74] A. Borel, *Stable real cohomology of arithmetic groups*, Ann. Sci. École Norm. Sup. (4) 7 (1974), 235-272.
- [BrLe66] W. Browder, J. Levine, *Fibering manifolds over a circle*, Comment. Math. Helv. 40 (1966), 153-160.
- [Bro82] K. Brown, *Cohomology of Groups*, Graduate Texts in Mathematics, 87. Springer-Verlag, New York-Berlin, 1982.
- [Bru71] G. Brumfiel, *Homotopy equivalences of almost smooth manifolds*, Comment. Math. Helv. 46 (1971), 381-407.
- [Cer70] J. Cerf, *The pseudo-isotopy theorem for simply connected differentiable manifolds*, Manifolds- Amsterdam (1970), 76-82, Lect. Notes Math. 197, Springer, Berlin (1970).
- [Cro11] D. Crowley, *On the mapping class groups of $\#_r S^p \times S^p$ for $p = 3, 7$* , Math. Z. (2011) 269:1189-1199.
- [EiMa54] S. Eilenberg and S. MacLane, *On the groups $H(\Pi, n)$, II: Methods of computation*, Annals of Mathematics Second Series, Vol. 60 (1954), 40-139.
- [FaMa12] B. Farb and D. Margalit, *A Primer on Mapping Class Groups*, Princeton Mathematical Series, 49. Princeton University Press, Princeton, NJ, 2012.
- [GaRW18] S. Galatius and O. Randal-Williams, *Moduli spaces of manifolds: a user's guide*, Handbook of homotopy theory, 443-485, CRC Press/Chapman Hall Handb. Math. Ser., CRC Press, Boca Raton, FL, 2020.
- [Gia71] V. Giambalvo, *On $O\langle 8 \rangle$ -cobordism*, Illinois J. Math. 15 (1971), 533-541.
- [Hae61] A. Haefliger, *Differentiable embeddings*, Bull. Amer. Math. Soc. 67 (1961), 109-112.
- [Hae66] A. Haefliger, *Differentiable embeddings of S^m in S^{m+q} for $q \geq 2$* , Ann. of Math. (2) 83 (1966), No.3, 402-436.
- [Hai19] R. Hain, *Mapping class groups of simply-connected Kähler manifolds*, arXiv:2304.01410.
- [Hew94] P. R. Hewitt, *Extensions of residually finite groups*, J. Algebra 163 (1994), No.3, 757-772.
- [Jup73] P. E. Jupp, *Classification of certain 6-manifolds*, Proc. Camb. Phil. Soc. (1973), 73, 293-300.
- [KeMi63] M. Kervaire and J. Milnor, *Groups of homotopy spheres. I*, Ann. of Math. (2), 77, 1963, 504-537.

- [Kor02] M. Korkmaz, *Low-dimensional homology groups of mapping class groups: a survey*, Turkish J. Math. 26 (2002), no. 1, 101-114.
- [Kra19] M. Krannich, *Mapping class group of highly-connected $(4k+2)$ -manifolds*, Selecta Math. (N. S.) 26 (2020), no. 5, Paper No. 81.
- [Kre99] M. Kreck, *Surgery and duality*, Annals of Mathematics, 149 (1999), 707-754.
- [Kre18] M. Kreck, *On the classification of 1-connected 7-manifolds with torsion free second homology*, J. Topology 11 (2018) No. 3, 720-738.
- [Kre78] M. Kreck, *Isotopy classes of diffeomorphisms of $(k-1)$ -connected almost parallelizable $2k$ -manifolds*, Algebraic topology, Aarhus 1978 (Proc. Sympos., Univ. Aarhus, Aarhus, 1978), pp. 643-C663, Lecture Notes in Math., 763, Springer, Berlin, 1979.
- [KrSt88] M. Kreck and S. Stolz, *A diffeomorphism classification of 7-dimensional homogeneous Einstein manifolds with $SU(3) \times SU(2) \times U(1)$ -symmetry*, Annals of Mathematics, 127 (1988), 373-388.
- [KrSt91] M. Kreck and S. Stolz, *Some nondiffeomorphic heomorphic homogeneous 7-manifolds with positive sectional curvature*, J. Differential Geometry, 33 (1991), 465-486.
- [KrSu19] M. Kreck and Y. Su, *Finiteness and infiniteness results for Torelli groups of (Hyper)-Kähler manifolds*, Math. Ann. 381 (2021), no. 1-2, 841-852.
- [Kry02] N. Krylov, *On the Jacobi group and the mapping class group of $S^3 \times S^3$* , Trans. A. M. S., Vol. 355 No. 1 (2002), 99-117.
- [Las65] R. Lashof, *Problems in differential and algebraic topology*, Seattle Conference, 1963, Ann. of Math. (2) 81 (1965), 565-591.
- [Mor93] S. Morita, *The extension of Johnsons homomorphism from the Torelli group to the mapping class group*, Invent. Math. 111 (1993), no. 1, 197-224.
- [RW23] O. Randal-Williams, *Monodromy and mapping class groups of 3-dimensional hypersurfaces*, preprint
- [Sat69] H. Sato, *Diffeomorphism groups of $S^p \times S^q$ and exotic spheres*, Quart. J. Math. Oxford (2), 20 (1969), 255-276.
- [Sul77] D. Sullivan, *Infinitesimal computations in topology*, Publications mathématiques de l'I.H.É.S., tome 47 (1977), 269-331.
- [Ver13] M. Verbitsky, *Mapping class group and a global Torelli theorem for hyperkähler manifolds*, Appendix A by Eyal Markman. Duke Math. J. 162 (2013), no. 15, 2929-2986.
- [Wal63] C. T. C. Wall, *Classification problems in differential topology II: Diffeomorphisms of Handlebodies*, Topology 2 (1963), 263-272.
- [Wal66] C. T. C. Wall, *Classification problems in differential topology V*, Invent. Math. 1 (1966), 355-374.
- [Wal69] C. T. C. Wall, *Non-additivity of the signature*, Invent. Math. 7 (1969), 269-274.

MATHEMATISCHES INSTITUT, UNIVERSITÄT BONN AND MATHEMATISCHES INSTITUT,
UNIVERSITÄT FRANKFURT

Email address: kreck@math.uni-bonn.de

HLM, ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE, CHINESE ACADEMY OF
SCIENCES AND SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF CHINESE ACADEMY
OF SCIENCES

Email address: suyang@math.ac.cn