

## DOUBLY NONLINEAR STOCHASTIC EVOLUTION EQUATIONS II

LUCA SCARPA AND ULISSE STEFANELLI

ABSTRACT. Complementing the analysis in [41], we investigate the well-posedness of SPDEs problems of doubly nonlinear type. These arise ubiquitously in the modelization of dissipative media and correspond to generalized balance laws between conservative and nonconservative dynamics. We extend the reach of the classical deterministic case by allowing for stochasticity. The existence of martingale solutions is proved via a regularization technique, hinging on the validity of a Itô formula in a minimal regularity setting. Under additional assumptions, the well-posedness of stochastically strong solutions is also shown.

## 1. INTRODUCTION

This is the second paper of a series devoted to the existence of solutions to stochastic doubly nonlinear evolution equations. Complementing the analysis of [41], we focus here on equations which, in the *deterministic* case, take the form

$$A(\partial_t u(t)) + \partial \hat{B}(u(t)) \ni F(t, u(t)). \quad (1)$$

Here,  $t \in [0, T] \mapsto u(t) \in H$  represents the state of a physical systems at time  $t$ , seen as an element in the separable Hilbert space  $H$ ,  $T > 0$  is some final reference time, and  $\partial_t$  is the time derivative. The functional  $\hat{B} : H \rightarrow (-\infty, +\infty]$  denotes the *energy* of the system, here assumed to be convex, and  $\partial \hat{B}$  is the corresponding subdifferential. The term  $t \mapsto F(t, u(t)) \in H$  represents further external actions. The possibly multivalued, maximal monotone operator  $A : H \rightarrow 2^H$  with  $0 \in A(0)$  corresponds to *dissipation* instead. More precisely, the scalar  $(A(\partial_t u), \partial_t u)_H \geq 0$  represents the instantaneous dissipation of the system corresponding to the rate  $\partial_t u$ , where  $(\cdot, \cdot)_H$  is the scalar product in  $H$ .

Given these provisions, relation (1) is nothing but the balance between *conservative* actions, modelled by  $\partial \hat{B}(u)$ , *dissipative* actions, encoded in  $A(\partial_t u)$ , and *external* actions, namely  $F(t, u)$ . By additionally letting  $F(t, u) = \partial_u \hat{F}(t, u)$ , where  $\hat{F}(t, u)$  denotes the work of external actions at time  $t$  on the state  $u \in H$ , and assuming enough smoothness, one can check in particular the *dissipation inequality*

$$\frac{d}{dt}(\hat{B}(u) - \hat{F}(\cdot, u)) + \partial_t \hat{F}(\cdot, u) = (\partial \hat{B}(u) - F(\cdot, u), \partial_t u)_H = -(A(\partial_t u), \partial_t u)_H \leq 0. \quad (2)$$

The left-hand side of this relation features the sum of the variation of the free energy of the system  $\hat{B}(u) - \hat{F}(t, u)$ , namely the stored part of the energy  $\hat{B}$  which is not contributed from the exterior, plus the power of external actions. In particular, (2) entails that this balance is negative along all trajectories, expressing indeed the dissipativity of the system. Indeed, the generalized balance (1) is prototypical of dissipative systems and arises ubiquitously in

2010 *Mathematics Subject Classification.* 35K55, 35R60, 60H15.

*Key words and phrases.* Doubly nonlinear stochastic equations, strong and martingale solutions, existence, maximal monotone operators, generalized Itô's formula.

applications, especially in relation with Thermomechanics and dissipative media: see [30, 31] and [18] for justifications. Note that the existence of a potential  $\widehat{F}(t, u)$  is instrumental to obtaining relation (2), being however not required for the existence theory.

The linear case where  $A$  is chosen to be the identity in  $H$  (not specifically in focus here, however), corresponds to perturbed gradient flows and is rather classical [6]. For existence results for  $A$  nonlinear and coercive the reader is referred to COLLI & VISINTIN [14] and COLLI [13] (see also [8], [7], and [45] among others). Besides existence, structural stability [1], perturbations and long-time behavior [2, 3, 37, 43, 44], and variational characterization of solutions [4, 5, 47] have also been considered. The case  $A$  nonlinear and 0-homogeneous (not included in our analysis) is related with the modeling of rate-independent evolution: see [32] for a recent comprehensive collection of results.

Our aim is to study a *stochastic* version of equation (1), namely (4) below, taking into account possible random disturbances occurring in the evolution and affecting the energy–dissipation balance. Let a filtered probability space  $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in [0, T]}, \mathbb{P})$  and a cylindrical Wiener process  $W$  on a separable Hilbert space  $U$  be given. Solutions  $(\omega, t) \mapsto u(\omega, t) \in H$  are assumed to be Itô processes of the form

$$u(t) = u^d(t) + \int_0^t u^s(s) dW(s), \quad t \in [0, T], \quad (3)$$

where  $u^d$  is an absolutely continuous process and  $u^s$  is a  $W$ -stochastically integrable process. For processes in the form (3), the averaged variation rate is given exactly by the contribution  $\mathbb{E} \partial_t u^d$ , where  $\mathbb{E}$  denotes expectation. It is then natural to consider the differentiable process  $u^d$  as the one accounting for dissipation, whereas the stochastic integral of  $u^s$  has to be interpreted as the random perturbation of the trajectory of  $u$ . Consequently, as we are interested in nonlinear dissipation evolutions, we prescribe the action of the dissipation  $A$  directly on the rate  $\partial_t u^d$ , by keeping the contribution  $u^s dW$  as a separate random disturbance affecting the trajectory of  $u$ . Bearing in mind these considerations, we are interested in the following doubly nonlinear stochastic evolution equation

$$A(\partial_t u^d) dt + u^s dW + \partial \widehat{B}(u) dt \ni F(\cdot, u) dt + G(\cdot, u) dW, \quad (4)$$

where  $G$  is a suitable stochastically integrable operator, possibly depending on  $u$ . This may be equivalently seen as a system

$$\begin{cases} A(\partial_t u^d) + \partial \widehat{B}(u) \ni F(\cdot, u), \\ u^s = G(\cdot, u), \end{cases} \quad (5)$$

under the constraint that  $u$  is an Itô process in the form (3). By taking  $G = 0$ , the latter entails  $u^s = 0$ , so that  $u = u^d$ , and the equation reduces to the deterministic case of (1). In the special case where  $A$  is the identity in  $H$ , the doubly nonlinear problem (4) reduces instead to the classical well-understood stochastic evolution equation

$$du + \partial \widehat{B}(u) dt \ni F(\cdot, u) dt + G(\cdot, u) dW.$$

More generally, equation (4), or equivalently (5), features a possibly nonquadratic dissipation in terms of  $\partial_t u^d$ , and is the natural stochastic version of the deterministic evolution (1).

The consistency of the doubly nonlinear stochastic evolution equation (4) is particularly evident at the level of the energy–dissipation balance. Indeed, in Proposition 3.3 below we prove a refined Itô formula which would allow us to argue along the lines of (2) and obtain the

averaged dissipativity inequality

$$\frac{d}{dt} \mathbb{E} \left( \widehat{B}(u) - \widehat{F}(\cdot, u) - \frac{1}{2} \mathbb{E}(\text{Tr } L(\cdot, u)) \right) + \mathbb{E} \partial_t \widehat{F}(\cdot, u) = - \mathbb{E}(A(\partial_t u), \partial_t u)_H \leq 0. \quad (6)$$

Here,  $L(\cdot, u) = G(\cdot, u)G^*(\cdot, u)D_G \partial \widehat{B}(\cdot)$ , where the symbol  $D_G$  represents the Gâteaux differential. With respect to the deterministic dissipation inequality (2), its stochastic version (6) features the additional contribution of the noise, which effectively enters the energy balance via the term in  $L$ . The dissipativity inequality (6) will be crucial, as it allows to obtain uniform estimates on the solutions of the doubly nonlinear stochastic problem (4).

The main result of this note is to prove that the Cauchy problem for equation (4) admits an analytically strong martingale solution if  $A$  is coercive and linearly bounded and the sublevels of  $\widehat{B}$  are compact, see Theorem 2.1. The key step is to prove a Itô formula under minimal regularity assumptions, see Proposition 3.3. This in turn allows us to perform a regularization procedure and identify the corresponding nonlinear limits by a lower semicontinuity argument. This technique is new and can be expected to be useful elsewhere. Eventually, if  $A$  is strongly monotone and  $\partial \widehat{B}$  is linear and self-adjoint, solutions are actually strong in probability and unique, see Theorem 2.2.

The main mathematical difficulty in tackling the doubly nonlinear problem (4) is duplex, and consists on the one hand in proving some uniform estimates on the solutions and on the other in identifying the two nonlinearities  $A(\partial_t u^d)$  and  $\partial \widehat{B}(u)$ . These two issues are indeed implicitly related to one another, and their proof hinges on the availability of an accurately refined Itô's formula in Hilbert spaces. First of all, uniform estimates on suitably approximated solutions of (4) can be obtained, provided to show the dissipativity inequality for a regularized equation, which is easier to handle. Unlike the deterministic setting, the dissipativity inequality cannot be achieved by simply testing the equation by  $\partial_t u^d$ , due to the presence of the noise terms. Instead, it is obtained by proving an Itô formula for a suitable regularization of the energy  $\widehat{B}$  and by comparison in the equation itself. As this first step is carried out on (relatively) arbitrary regularized problems, one has enough regularity on the approximated energy at disposal and uniform estimates on the solutions can be obtained. The delicate point, however, consists in passing to the limit. Indeed, as the problem exhibits two nonlinearities, the identification procedure is absolutely nontrivial. The main idea is to identify  $\partial B(u)$  by means of (stochastic) compactness arguments. Once this is done, we characterize  $A(\partial_t u^d)$  using lower semicontinuity techniques (the so-called lim sup argument): this however strongly relies on the availability of an energy–dissipation *equality* at the limit. From the mathematical point of view, we are then forced to prove first an *exact* Itô formula for the energy  $\widehat{B}$  (not regularized). While Itô *inequalities* are available in literature in connection with existence of strong solutions to SPDEs for example [19], to the best of our knowledge Itô *equalities* associated to a general energy  $\widehat{B}$  have not received attention elsewhere, the main problem being the fact that  $\widehat{B}$  is not everywhere defined in  $H$ , having compact sublevels. For these reasons, the analytical preliminaries contained here have independent relevance on their own.

To our knowledge, this paper delivers the first existence theory in the doubly nonlinear frame of (4). In the case of a linear and nondegenerate operator  $A$ , existence results can be traced back to the classical theory by PARDOUX [34, 35] and KRYLOV & ROZOVSKIĬ, [23]: see the monographs [15, 36] for a general overview. In particular, the Cauchy problem for (4) with  $A$  linear and  $\partial \widehat{B}$  subhomogeneous has been proved to admit strong solutions by GESS [19]. Well-posedness under more relaxed growth conditions are also available, see [25, 27, 28] for semilinear

equations, [26, 29, 39] for equations in divergence form, and [11, 33, 40] for porous-media, Allen-Cahn, and Cahn-Hilliard equations.

In the first paper [41] of this series on existence for doubly nonlinear stochastic evolution equations, we focused on relations of the form

$$\mathrm{d}A(u) + \partial\widehat{B}(u) \, \mathrm{d}t \ni F(\cdot, u) \, \mathrm{d}t + G(\cdot, u) \, \mathrm{d}W$$

instead. Here, the operator  $A$  is applied directly to  $u$  and then differentiated. Such equations are also very relevant from the applicative viewpoint and arise in connection with different nonlinear diffusion situations, including phase transitions, porous-media, the Hele-Shaw cell, non-Newtonian fluids, and the Mean-Curvature flow. Existence for the latter has been obtained in [38] (for  $A$  bi-Lipschitz) and in [41] (for  $A$  multivalued). The case of  $A$  nonlinear and  $\partial\widehat{B}$  linear has been originally discussed in [10] in the setting of the two-phase stochastic Stefan problem, see also [9, 22].

Let us also point out that in the case where  $A = \partial\widehat{A}$  is a subdifferential of a certain convex lower semicontinuous function  $\widehat{A}$  on  $H$ , the doubly nonlinear equation (4) fits well in the framework of the stochastic Weighted Energy–Dissipation approach, and can be thus seen as limit of convex minimization problems. Such technique has been shown to be effective when  $A$  is the identity [42]. The study of the genuinely doubly nonlinear case can also be considered.

We give the detail of our setting and state the existence results in Section 2. Section 3 is devoted to the proof of the above-mentioned generalized Itô formula. The proof of Theorem 2.1, namely the existence of martingale solutions, is detailed in Section 4. Finally, probabilistically strong solutions are discussed in Section 5. We conclude in Section 6 by presenting an example of a concrete PDE to which the abstract analysis can be applied.

## 2. SETTING AND MAIN RESULTS

In this section we introduce the setting and assumptions on the data of the problem (Subsections 2.1–2.3) and we present the main results of the work (Subsection 2.4).

**2.1. Setting and notation.** Let  $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in [0, T]}, \mathbb{P})$  be a filtered probability space satisfying the usual conditions, where  $T > 0$  is a fixed final reference time, and let also  $W$  be a cylindrical Wiener process on a separable Hilbert space  $U$ . We fix once and for all a complete orthonormal system  $(u_j)_j$  of  $U$ .

For every Banach spaces  $E_1$  and  $E_2$ , the symbol  $\mathcal{L}(E_1, E_2)$  denotes the space of linear continuous operator from  $E_1$  to  $E_2$  endowed with the norm-topology. The space  $\mathcal{L}(E_1, E_2)$  endowed with the strong and weak operator topology is denoted by  $\mathcal{L}_s(E_1, E_2)$  and  $\mathcal{L}_w(E_1, E_2)$ , respectively. If  $E_1$  and  $E_2$  are also Hilbert spaces, the space of trace-class and Hilbert-Schmidt operators from  $E_1$  to  $E_2$  are denoted by  $\mathcal{L}^1(E_1, E_2)$  and  $\mathcal{L}^2(E_1, E_2)$ , respectively.

In the paper,  $V$  is separable reflexive Banach space and  $H$  is a Hilbert space such that  $V \hookrightarrow H$  continuously, densely, and compactly. As usual, we identify  $H$  with its dual, so that we have the Gelfand triple

$$V \hookrightarrow H \hookrightarrow V^*.$$

Without loss of generality, we suppose that there exists a family of regularizing linear operators  $(R_\lambda)_{\lambda > 0} \subset \mathcal{L}(H, V)$  such that  $R_\lambda \rightarrow I$  in  $\mathcal{L}_s(V, V)$  as  $\lambda \searrow 0$ : this is trivially satisfied in the majority of interesting examples, for example when  $V$  is a Sobolev space.

The progressive sigma algebra on  $\Omega \times [0, T]$  is denoted by  $\mathscr{P}$ . We use the classical symbols  $L^s(\Omega; E)$  and  $L^r(0, T; E)$  for the spaces of strongly measurable Bochner-integrable functions on  $\Omega$  and  $(0, T)$ , respectively, for all  $s, r \in [1, +\infty]$  and for every Banach space  $E$ . If  $s, r \in [1, +\infty)$  we use  $L^s_{\mathscr{P}}(\Omega; L^r(0, T; E))$  to indicate that measurability is intended with respect to  $\mathscr{P}$ . In the case that  $s \in (1, +\infty)$ ,  $r = +\infty$ , and  $E$  is a separable, we set

$$L^s_{\mathscr{P}}(\Omega; L^\infty(0, T; E^*)) := \left\{ v : \Omega \rightarrow L^\infty(0, T; E^*) \text{ weakly}^* \text{ progressively measurable} \right. \\ \left. \text{with } \mathbb{E} \|v\|_{L^\infty(0, T; E^*)}^s < \infty \right\},$$

and recall that by [16, Thm. 8.20.3] we have the identification

$$L^s_{\mathscr{P}}(\Omega; L^\infty(0, T; E^*)) = \left( L^{\frac{s}{s-1}}_{\mathscr{P}}(\Omega; L^1(0, T; E)) \right)^*.$$

**2.2. Spaces of Itô processes.** We introduce here some specific notation for classes of Itô-type processes. Let  $E, F$  be two separable reflexive Banach spaces and  $K$  be a separable Hilbert space, such that  $E, K \hookrightarrow F$  continuously. For every  $r_1, r_2 \in [1, +\infty)$  we use the notation

$$\mathcal{I}^{r_1, r_2}(E, K) := L^{r_1}_{\mathscr{P}}(\Omega; W^{1, r_1}(0, T; E)) \oplus L^{r_2}_{\mathscr{P}}(\Omega; L^2(0, T; \mathcal{L}^2(U, K))) \cdot W.$$

In other words,  $\mathcal{I}^{r_1, r_2}(E, K)$  is the space of all processes  $u$  that can be written as

$$u(t) = u^d(t) + \int_0^t u^s(r) dW(r), \quad t \in [0, T],$$

for some  $u^d \in L^{r_1}_{\mathscr{P}}(\Omega; W^{1, r_1}(0, T; E))$  and  $u^s \in L^{r_2}_{\mathscr{P}}(\Omega; L^2(0, T; \mathcal{L}^2(U, K)))$ . Clearly, in such case we have

$$u(t) = u^d(0) + \int_0^t \partial_t u^d(s) ds + \int_0^t u^s(r) dW(r), \quad t \in [0, T],$$

Note that the sum is actually a direct sum, and the representation  $u = u^d + u^s \cdot W$  is unique.

**2.3. Assumptions.** We now introduce and comment the assumptions on operators and data. These will be assumed throughout, without further specific mention.

- **Assumption U0:**  $u_0 \in V$ .
- **Assumption A:**  $A : H \rightarrow 2^H$  is maximal monotone and there exist two constants  $c_A, C_A > 0$  such that
$$\|y\|_H \leq C_A(1 + \|x\|_H) \quad \forall x \in H, \quad \forall y \in A(x),$$

$$(y, x)_H \geq c_A \|x\|_H^2 - c_A^{-1} \quad \forall x \in D(A), \quad \forall y \in A(x).$$
- **Assumption B1:**  $\widehat{B} : H \rightarrow [0, +\infty]$  is convex lower semicontinuous with  $\widehat{B}(0) = 0$  and  $B := \partial \widehat{B} : H \rightarrow 2^H$  is its subdifferential. We ask that

$$V \subseteq D(\widehat{B}),$$

and  $\widehat{B}$  is continuous at some point of  $V$ : this implies that  $\partial(\widehat{B}|_V) : V \rightarrow 2^{V^*}$  coincides with  $B$  on  $D(B)$ , hence  $B$  can be also seen as a maximal monotone operator from  $V$  to  $2^{V^*}$ . We assume that  $B$  is single-valued, and that there exist constants  $c_B > 0$  and  $p \geq 2$  such that

$$(B(x_1) - B(x_2), x_1 - x_2)_H \geq c_B \|x_1 - x_2\|_V^p \quad \forall x_1, x_2 \in D(B).$$

- **Assumption B2:**  $B : V \rightarrow V^*$  is Gâteaux-differentiable with

$$D_{\mathcal{G}}B \in C^0(V; \mathcal{L}_w(V, V^*))$$

and there exists a constant  $C_B > 0$  such that

$$\|D_{\mathcal{G}}B(x)\|_{\mathcal{L}(V, V^*)} \leq C_B \left(1 + \|x\|_V^{p-2}\right) \quad \forall x \in V.$$

- **Assumption F:**  $F : [0, T] \times V \rightarrow H$  is measurable, and there exist a constant  $C_F > 0$  and  $h_F \in L^1(0, T)$  such that

$$\begin{aligned} \|F(\cdot, x_1) - F(\cdot, x_2)\|_H^2 &\leq C_F \left(1 + \|x_1\|_V^{p-2} + \|x_2\|_V^{p-2}\right) \|x_1 - x_2\|_V^2 & \forall x_1, x_2 \in V, \\ \|F(\cdot, x)\|_H^2 &\leq h_F(\cdot) + C_F \|x\|_V^p & \forall x \in V. \end{aligned}$$

- **Assumption G:**  $G : [0, T] \times H \rightarrow \mathcal{L}^2(U, H)$  is measurable,  $G([0, T] \times V) \subset \mathcal{L}(U, V)$ , and there exist a constant  $C_G > 0$  such that

$$\begin{aligned} \|G(\cdot, x_1) - G(\cdot, x_2)\|_{\mathcal{L}^2(U, H)} &\leq C_G \|x_1 - x_2\|_H & \forall x_1, x_2 \in H, \\ \|G(\cdot, x)\|_{\mathcal{L}^2(U, H)} &\leq C_G (1 + \|x\|_V^\nu) & \forall x \in V, \end{aligned}$$

where  $\nu = 1$  if  $p > 2$  and  $\nu \in (0, 1)$  if  $p = 2$ . Moreover, we assume that there exists  $L : [0, T] \times V \rightarrow \mathcal{L}^1(H, H)$  and  $h_G \in L^1(0, T)$  such that

$$L(\cdot, x) = G(\cdot, x)G(\cdot, x)^* D_{\mathcal{G}}B(x) \quad \forall x \in V,$$

and

$$\begin{aligned} L(t, \cdot) &\in C^0(V; \mathcal{L}^1(H, H)) \quad \text{for a.e. } t \in (0, T), \\ \|L(\cdot, x)\|_{\mathcal{L}^1(H, H)} &\leq h_G(\cdot) + C_G \|x\|_V^p \quad \forall x \in V. \end{aligned}$$

Let us comment on this last requirement. Note that by the regularity of  $G$  and  $B$ , for every  $t \in (0, T)$  and  $x \in V$  we have that  $G(t, x) \in \mathcal{L}(U, V)$  and  $D_{\mathcal{G}}B(x) \in \mathcal{L}(V, V^*)$ , so that  $G(t, x)G(t, x)^* D_{\mathcal{G}}B(x) \in \mathcal{L}(V, V)$ . In the last condition, we are assuming that actually such operator extends to a trace class operator on  $H$ . Clearly, any such extension is necessarily unique due to the density of  $V$  in  $H$ . This behaviour is very natural and actually occurs in several relevant situations: roughly speaking, it means that the composition with  $GG^*$  has a regularizing effect on  $D_{\mathcal{G}}B(x)$ , due for example to suitable compensations of the singular terms.

**2.4. Main results.** We are now in the position of stating the main results of the paper. Our first result concerns existence of analytically strong martingale solutions to (4).

**Theorem 2.1** (Existence of martingale solutions). *Under the assumptions of Subsection 2.3, there exists a probability space  $(\hat{\Omega}, \hat{\mathcal{F}}, \hat{\mathbb{P}})$ , a  $U$ -cylindrical Wiener process  $\hat{W}$  on it, and a triple  $(\hat{u}, \hat{u}^d, \hat{v})$ , with*

$$\begin{aligned} \hat{u} &\in L^p_{\hat{\mathcal{F}}}(\hat{\Omega}; C^0([0, T]; H)) \cap L^p_{\hat{\mathcal{F}}}(\hat{\Omega}; L^\infty(0, T; V)), & \hat{u}^d &\in L^2_{\hat{\mathcal{F}}}(\hat{\Omega}; H^1(0, T; H)), \\ \hat{u} &\in D(B) \quad \text{a.e. in } \hat{\Omega} \times (0, T), & B(\hat{u}) &\in L^2_{\hat{\mathcal{F}}}(\hat{\Omega}; L^2(0, T; H)), \\ \hat{v} &\in L^2_{\hat{\mathcal{F}}}(\hat{\Omega}; L^2(0, T; H)), \end{aligned}$$

such that

$$\begin{aligned} \hat{u}(t) &= u_0 + \int_0^t \partial_t \hat{u}^d(s) \, ds + \int_0^t G(s, \hat{u}(s)) \, d\hat{W}(s) & \forall t \in [0, T], \quad \hat{\mathbb{P}}\text{-a.s.}, \\ \hat{v}(t) + B(\hat{u}(t)) &= F(t, \hat{u}(t)) & \text{for a.e. } t \in (0, T), \quad \hat{\mathbb{P}}\text{-a.s.}, \\ \hat{v}(t) &\in A(\partial_t \hat{u}^d(t)) & \text{for a.e. } t \in (0, T), \quad \hat{\mathbb{P}}\text{-a.s.} \end{aligned}$$

Furthermore, the following energy equality holds:

$$\begin{aligned} &\widehat{B}(\hat{u}(t)) + \int_0^t (\hat{v}(s), \partial_t \hat{u}^d(s))_H \, ds \\ &= \widehat{B}(u_0) + \int_0^t (F(s, \hat{u}(s)), \partial_t \hat{u}^d(s))_H \, ds + \frac{1}{2} \int_0^t \text{Tr}[L(s, \hat{u}(s))] \, ds \\ &\quad + \int_0^t (B(\hat{u}(s)), G(s, \hat{u}(s)) \, dW(s))_H \quad \forall t \in [0, T], \quad \hat{\mathbb{P}}\text{-a.s.} \end{aligned}$$

Our second result gives a sufficient condition on the nonlinearities in order to obtain existence (and uniqueness) of strong solutions both in the probabilistic and analytical sense.

**Theorem 2.2** (Well posedness in the strong probabilistic sense). *Under the assumptions of Subsection 2.3, suppose further that  $A$  is strongly monotone,  $B$  is linear self-adjoint,  $G$  is linear in its second variable, and  $h_G = 0$ . Then, there exists a unique triple  $(u, u^d, v)$ , with*

$$\begin{aligned} u &\in L^p_{\mathcal{F}}(\Omega; C^0([0, T]; H)) \cap L^p_{\mathcal{F}}(\Omega; L^\infty(0, T; V)), \quad u^d \in L^2_{\mathcal{F}}(\Omega; H^1(0, T; H)), \\ B(u) &\in L^2_{\mathcal{F}}(\Omega; L^2(0, T; H)), \\ v &\in L^2_{\mathcal{F}}(\Omega; L^2(0, T; H)), \end{aligned}$$

such that

$$\begin{aligned} u(t) &= u_0 + \int_0^t \partial_t u^d(s) \, ds + \int_0^t G(s, u(s)) \, dW(s) & \forall t \in [0, T], \quad \mathbb{P}\text{-a.s.}, \\ v(t) + B(u(t)) &= F(t, u(t)) & \text{for a.e. } t \in (0, T), \quad \mathbb{P}\text{-a.s.}, \\ v(t) &\in A(\partial_t u^d(t)) & \text{for a.e. } t \in (0, T), \quad \mathbb{P}\text{-a.s.} \end{aligned}$$

Furthermore, there exists a constant  $K > 0$ , such that, for every set  $\{u_{0,i}\}_{i=1,2} \subset V$  of initial data, their respective solutions  $\{(u_i, u_i^d, v_i)\}_{i=1,2}$  satisfy

$$\|u_1 - u_2\|_{L^\infty(0, T; L^2(\Omega; V))} + \|u_1^d - u_2^d\|_{L^2(\Omega; H^1(0, T; H))} \leq K \|u_{0,1} - u_{0,2}\|_V.$$

### 3. AN EXTENDED ITÔ'S FORMULA

In this section we prove a general version of Itô's formula for  $\widehat{B}$ , under minimal differentiability assumptions, see Proposition 3.3. In particular, we recall that the second derivative of  $\widehat{B}$  is supposed to be defined only in the sense of Gâteaux in  $V$ : hence,  $\widehat{B}$  may even be not twice-differentiable on  $H$ , and, in fact, not even well-defined on the whole space  $H$ .

We find such Itô's formula very interesting on its own, as it widely generalizes the classical result to the case where the second derivatives are not necessarily well-defined and continuous.



For every  $\lambda > 0$ , we recall that the Moreau-Yosida regularization of  $\widehat{B}$ , the resolvent of  $B$ , and the Yosida approximation of  $B$ , are defined respectively as

$$\begin{aligned}\widehat{B}_\lambda : H &\rightarrow [0, +\infty), & \widehat{B}_\lambda(x) &:= \inf_{y \in H} \left\{ \widehat{B}(y) + \frac{\|x - y\|_H^2}{2\lambda} \right\}, \quad x \in H, \\ J_\lambda^B : H &\rightarrow D(B) \subset V, & J_\lambda^B(x) &:= (I + \lambda B)^{-1}(x), \quad x \in H, \\ B_\lambda : H &\rightarrow H, & B_\lambda(x) &:= \frac{x - J_\lambda^B(x)}{\lambda}, \quad x \in H,\end{aligned}$$

where  $I : H \rightarrow H$  is the identity in  $H$ . It is well-known that  $J_\lambda^B$  is non-expansive on  $H$  and that  $B_\lambda$  is  $1/\lambda$ -Lipschitz-continuous on  $H$ .

First of all, we prove some technical properties of the resolvent.

**Lemma 3.1.** *For every  $\lambda > 0$ , the resolvent  $J_\lambda^B$  uniquely extends to a  $\frac{1}{p-1}$ -Hölder-continuous Gâteaux-differentiable operator  $J_\lambda^B : V^* \rightarrow V$ , such that  $D_{\mathcal{G}} J_\lambda^B \in C^0(H; \mathcal{L}_w(H, H))$ . Moreover, as  $\lambda \searrow 0$  it holds that*

$$\begin{aligned}J_\lambda^B(x) &\rightarrow x && \text{in } H && \forall x \in H, \\ J_\lambda^B(x) &\rightharpoonup x && \text{in } V && \forall x \in V, \\ D_{\mathcal{G}} J_\lambda^B(x) &\rightarrow I && \text{in } \mathcal{L}_s(H, H) && \forall x \in H.\end{aligned}$$

In particular, there exists a constant  $M > 0$ , independent of  $\lambda$ , such that

$$\|J_\lambda(x)\|_V \leq M(1 + \|x\|_V) \quad \forall x \in V.$$

*Proof.* STEP 1. Let  $\lambda > 0$  be fixed. The operator

$$I + \lambda B : V \rightarrow V^*$$

is maximal monotone and coercive, hence surjective. Its inverse  $(I + \lambda B)^{-1} : V^* \rightarrow V$  is well-defined, and coincides with  $J_\lambda^B$  on  $H$ . In the sequel, we will use the same symbol  $J_\lambda^B$  to denote  $(I + \lambda B)^{-1} : V^* \rightarrow V$ . Note that actually  $J_\lambda^B : V^* \rightarrow V$  is  $\frac{1}{p-1}$ -Hölder-continuous by the strong monotonicity of  $B$ . Indeed, for every  $x_1, x_2 \in V^*$ , it is immediate to see that

$$\|J_\lambda^B(x_1) - J_\lambda^B(x_2)\|_H^2 + \lambda c_B \|J_\lambda^B(x_1) - J_\lambda^B(x_2)\|_V^p \leq \langle x_1 - x_2, J_\lambda^B(x_1) - J_\lambda^B(x_2) \rangle_{V^*, V},$$

from which it follows that

$$\|J_\lambda^B(x_1) - J_\lambda^B(x_2)\|_V \leq \left( \frac{1}{\lambda c_B} \right)^{\frac{1}{p-1}} \|x_1 - x_2\|_{V^*}^{\frac{1}{p-1}}.$$

Hence,  $(I + \lambda B)^{-1} : V^* \rightarrow V$  is Hölder-continuous, and the extension of  $J_\lambda^B$  to  $V^*$  is unique. Now, by **B1**–**B2** we know that  $I + \lambda B : V \rightarrow V^*$  is Gâteaux-differentiable with differential given by

$$T_\lambda(x) := I + \lambda D_{\mathcal{G}} B(x) \in \mathcal{L}(V, V^*), \quad x \in V.$$

It follows that for every  $x \in V$ , the linear continuous operator  $T_\lambda(x) : V \rightarrow V^*$  is monotone (hence maximal monotone), injective, and coercive on  $V$  (due to the strong monotonicity of  $B$ ): this implies that  $T_\lambda(x)$  is an isomorphism for all  $x \in V$ . Consequently, as we have already proved that  $I + \lambda B : V \rightarrow V^*$  is invertible with Hölder-continuous inverse, we deduce that  $J_\lambda^B : V^* \rightarrow V$  is Gâteaux-differentiable with

$$D_{\mathcal{G}} J_\lambda^B(x) = T_\lambda(J_\lambda^B(x))^{-1} \in \mathcal{L}(V^*, V), \quad x \in V^*.$$



Clearly, this implies in particular that  $J_\lambda^B : H \rightarrow H$  is Gâteaux-differentiable, and

$$D_G J_\lambda^B(y)h + \lambda D_G B(J_\lambda^B(y))D_G J_\lambda^B(y)h = h \quad \forall h, y \in H. \quad (7)$$

As a by-product, we deduce also that  $B_\lambda : H \rightarrow H$  is Gâteaux-differentiable with

$$D_G B_\lambda(y) = D_G B(J_\lambda^B(y))D_G J_\lambda^B(y) \quad \forall y \in H.$$

Let us show that actually  $D_G J_\lambda^B \in C^0(H; \mathcal{L}_w(H, H))$ . To this end, let  $(x_j)_j \subset H$  and  $x \in H$  such that  $x_j \rightarrow x$  in  $H$  as  $j \rightarrow \infty$ . For what we have just proved, it holds that  $J_\lambda^B(x_j) \rightarrow J_\lambda^B(x)$  in  $V$ . Taking  $y = x_j$  and  $h \in H$  in (7), and testing by  $D_G J_\lambda^B(x_j)h$  yields, by the monotonicity of  $B$  and the Young inequality,

$$\frac{1}{2} \|D_G J_\lambda^B(x_j)h\|_H^2 \leq \frac{1}{2} \|h\|_H^2 \quad \forall j \in \mathbb{N}.$$

Hence, there exists  $I_\lambda(x, h) \in H$  such that, as  $j \rightarrow \infty$ ,

$$D_G J_\lambda^B(x_j)h \rightharpoonup I_\lambda(x, h) \quad \text{in } H, \quad \forall h \in H.$$

Let now  $k \in H$  be arbitrary, and consider the double sequence

$$a_{j,i} := \left( \frac{J_\lambda^B(x_j + h/i) - J_\lambda^B(x_j)}{1/i}, k \right)_H, \quad (j, i) \in \mathbb{N}^2.$$

Since  $J_\lambda^B$  is non-expansive on  $H$ , the sequence  $\{a_{j,i}\}_{j,i}$  is uniformly bounded in  $(j, i)$ . Hence, there is  $a_\lambda \in \mathbb{R}$  such that, possibly on a non-relabelled subsequence of  $(j, i)$ , it holds

$$\lim_{j,i \rightarrow \infty} a_{j,i} = a.$$

Now, note also that by continuity and Gâteaux-differentiability of  $J_\lambda^B$  we have

$$\lim_{j \rightarrow \infty} a_{j,i} = \left( \frac{J_\lambda^B(x + h/i) - J_\lambda^B(x)}{1/i}, k \right)_H \quad \forall i \in \mathbb{N},$$

and

$$\lim_{i \rightarrow \infty} a_{j,i} = (D_G J_\lambda^B(x_j)h, k)_H \quad \forall j \in \mathbb{N}.$$

Consequently, since the partial limits are finite and the double sequence converges also jointly, it holds that

$$a = \lim_{j,i \rightarrow \infty} a_{j,i} = \lim_{j \rightarrow \infty} \lim_{i \rightarrow \infty} a_{j,i} = \lim_{i \rightarrow \infty} \lim_{j \rightarrow \infty} a_{j,i}.$$

Putting this information together, we deduce that

$$\lim_{j \rightarrow \infty} (D_G J_\lambda^B(x_j)h, k)_H = (D_G J_\lambda^B(x)h, k)_H = a \quad \forall h, k \in H.$$

Since the limit  $a$  is independent of the specific subsequence, this argument can be replicated on all subsequences of  $(j, i)$ , and the convergences holds actually along the entire sequence. By the arbitrariness of  $k \in H$ , it follows that  $I_\lambda(x, h) = D_G J_\lambda^B(x)h$  for all  $h \in H$ . This shows that  $D_G J_\lambda^B \in C^0(H; \mathcal{L}_w(H, H))$ , as required.

STEP 2. Let us show the behaviour as  $\lambda \searrow 0$ : let  $x \in H$  be fixed. First of all, recalling that

$$J_\lambda^B(x) + \lambda B(J_\lambda^B(x)) = x,$$

testing by  $J_\lambda^B(x)$  and using the strong monotonicity of  $B$ , the fact that  $0 \in B(0)$ , and the Young inequality we get

$$\frac{1}{2} \|J_\lambda^B(x)\|_H^2 + \lambda c_B \|J_\lambda^B(x)\|_V^p \leq \frac{1}{2} \|x\|_H^2. \quad (8)$$

It follows that  $(J_\lambda^B(x))_\lambda$  is relatively weakly compact in  $H$ . Moreover, by assumption **B2** and (8) we have that, for a positive constant  $M$  independent of  $\lambda$ ,

$$\lambda \|B(J_\lambda^B(x))\|_{V^*} \leq M\lambda \left(1 + \|J_\lambda^B(x)\|_V^{p-1}\right) \leq M\lambda + M\lambda(2c_B\lambda)^{-\frac{p-1}{p}} \|x\|_H^{\frac{2p-1}{p}} \rightarrow 0.$$

It follows then that  $J_\lambda^B(x) \rightharpoonup x$  in  $H$ , hence also, again by a classical lim sup-argument, that  $J_\lambda^B(x) \rightarrow x$  in  $H$ . Secondly, taking  $h \in H$  and  $y = x$  in (7), testing by  $D_G J_\lambda^B(x)h$ , using the monotonicity of  $B$  and the Young inequality yield

$$\frac{1}{2} \|D_G J_\lambda^B(x)h\|_H^2 \leq \frac{1}{2} \|h\|_H^2. \quad (9)$$

Hence, there exists  $\tilde{I}(x, h) \in H$  such that, as  $\lambda \searrow 0$ ,

$$D_G J_\lambda^B(x)h \rightharpoonup \tilde{I}(x, h) \quad \forall h \in H.$$

We show that  $\tilde{I}(x, h) = h$  using again an argument with double sequences. Let  $k \in H$  be arbitrary and consider

$$b_{j,i} := \left( \frac{J_{\lambda_j}^B(x + h/i) - J_{\lambda_j}^B(x)}{1/i}, k \right)_H, \quad (j, i) \in \mathbb{N}^2,$$

where  $(\lambda_j)_j$  is an arbitrary infinitesimal real sequence. Since  $J_\lambda^B$  is non-expansive on  $H$ , the sequence  $\{b_{j,i}\}_{j,i}$  is uniformly bounded in  $(j, i)$ , and there is  $b \in \mathbb{R}$  such that, possibly on a non-relabelled subsequence of  $(j, i)$ ,

$$\lim_{j,i \rightarrow \infty} b_{j,i} = b.$$

As before, by the convergence of  $J_\lambda$  as  $\lambda \searrow 0$  and the Gâteaux-differentiability of  $J_\lambda^B$  we have

$$\lim_{j \rightarrow \infty} b_{j,i} = (h, k)_H \quad \forall i \in \mathbb{N},$$

and

$$\lim_{i \rightarrow \infty} b_{j,i} = \left( D_G J_{\lambda_j}^B(x)h, k \right)_H \quad \forall j \in \mathbb{N}.$$

Consequently, since the partial limits are finite and the double sequence converges also jointly, the double limits can be computed in any order, so that it holds that

$$\lim_{j \rightarrow \infty} \left( D_G J_{\lambda_j}^B(x)h, k \right)_H = (h, k)_H \quad \forall h, k \in H.$$

Since the limit is independent of the specific subsequence, this argument can be replicated on all subsequences of  $(j, i)$ , and the convergences holds actually along the entire sequence. By the arbitrariness of  $k \in H$ , it follows that  $\tilde{I}(x, h) = h$  for all  $h \in H$ . Furthermore, from (9) we also have

$$\limsup_{\lambda \searrow 0} \|D_G J_\lambda^B(x)h\|_H^2 \leq \|h\|_H^2,$$

hence actually it holds that  $D_G J_\lambda^B(x)h \rightarrow h$  in  $H$ . This shows that  $D_G J_\lambda \rightarrow I$  in  $\mathcal{L}_s(H, H)$ , as required.

**STEP 3.** Lastly, let us suppose that  $x \in V$ : from the relation

$$J_\lambda^B(x) + \lambda B(J_\lambda^B(x)) = x,$$

testing by  $B(J_\lambda^B(x))$  and using **B1–B2** together with the Young inequality we get

$$\begin{aligned} c_B \|J_\lambda^B(x)\|_V^p + \lambda \|B(J_\lambda^B(x))\|_H^2 &= (B(J_\lambda^B(x)), x)_H \leq C_B \|x\|_V \left(1 + \|J_\lambda^B(x)\|_V^{p-1}\right) \\ &\leq C_B \|x\|_V + \frac{1}{p} C_B^p c_B^{1-p} \|x\|_V^p + \frac{p-1}{p} c_B \|J_\lambda^B(x)\|_V^p, \end{aligned}$$

from which

$$\frac{C_B}{p} \|J_\lambda^B(x)\|_V^p + \lambda \|B(J_\lambda^B(x))\|_H^2 \leq C_B \|x\|_V + \frac{1}{p} C_B^p c_B^{1-p} \|x\|_V^p. \quad (10)$$

It follows that  $J_\lambda^B(x) \rightarrow x$  in  $V$  as  $\lambda \searrow 0$ , and the proof is concluded.  $\square$

Lemma 3.1 implies in particular that the Yosida approximation  $B_\lambda : H \rightarrow H$  is Gâteaux-differentiable, but not necessarily in the sense of Fréchet. In the next lemma we show that it is possible to write Itô's formula for  $\widehat{B}_\lambda$ , even if its second derivative is not well-defined everywhere in the sense of Fréchet.

**Lemma 3.2.** *Let the processes  $v$ ,  $f$ , and  $C$  satisfy*

$$v \in L^2_{\mathcal{D}}(\Omega; C^0([0, T]; H)), \quad f \in L^2_{\mathcal{D}}(\Omega; L^2(0, T; H)), \quad C \in L^2_{\mathcal{D}}(\Omega; L^2(0, T; \mathcal{L}^2(U, H))),$$

and

$$v(t) + \int_0^t f(s) ds = v(0) + \int_0^t C(s) dW(s) \quad \forall t \in [0, T], \quad \mathbb{P}\text{-a.s.}$$

Then, for all  $\lambda > 0$  it holds that

$$\begin{aligned} \widehat{B}_\lambda(v(t)) + \int_0^t (f(s), B_\lambda(v(s)))_H ds &= \widehat{B}_\lambda(v(0)) + \frac{1}{2} \int_0^t \text{Tr}[C^*(s) D_{\mathcal{G}} B_\lambda(v(s)) C(s)] ds \\ &\quad + \int_0^t (B_\lambda(v(s)), C(s) dW(s))_H \quad \forall t \in [0, T], \quad \mathbb{P}\text{-a.s.} \end{aligned}$$

*Proof.* Let  $\{e_k\}_{k \in \mathbb{N}}$  be a complete orthonormal system of  $H$  included in  $V$  and set

$$H_n := \text{span}\{e_0, \dots, e_{n-1}\}, \quad n \in \mathbb{N}, \quad H_\infty := \bigcup_{n \in \mathbb{N}} H_n.$$

Clearly,  $H_n$  is a closed linear subspace of  $H$  for all  $n \in \mathbb{N}$ , and  $H_\infty$  is dense in  $H$ . For every  $n \in \mathbb{N}$ , it is well-defined the orthogonal projection  $P_n : H \rightarrow H_n$  on  $H_n$ , with respect to the scalar product of  $H$ . Clearly,  $P_n$  is linear, 1-Lipschitz continuous, and

$$P_n x \rightarrow x \quad \text{in } H \quad \forall x \in H.$$

Setting now  $v_n := P_n v$ ,  $f_n := P_n f$ , and  $C_n := P_n C$ , we have that, for every  $n \in \mathbb{N}$ ,

$$v_n(t) + \int_0^t f_n(s) ds = v_n(0) + \int_0^t C_n(s) dW(s) \quad \text{in } H_n, \quad \forall t \in [0, T], \quad \mathbb{P}\text{-a.s.} \quad (11)$$

Now, it is immediate to see that the restriction  $\widehat{B}_{\lambda|_{H_n}} \in C^1(H_n)$  and its differential is given by

$$D(\widehat{B}_{\lambda|_{H_n}}) = P_n \circ B_{\lambda|_{H_n}} : H_n \rightarrow H_n.$$

Furthermore, from Lemma 3.1 we have that  $B_\lambda : H \rightarrow H$  is Gâteaux-differentiable, which yields directly that  $P_n \circ B_{\lambda|_{H_n}} : H_n \rightarrow H_n$  is Gâteaux-differentiable in turn with

$$D_{\mathcal{G}}(P_n \circ B_{\lambda|_{H_n}})(x)h = P_n D_{\mathcal{G}} B_\lambda(x)h, \quad x \in H_n, \quad h \in H_n.$$

Let us show that actually  $D_{\mathcal{G}}^2 \widehat{B}_\lambda \in C^0(H_n; \mathcal{L}(H_n, H_n))$ . To this end, let  $\{x_j\}_j \subset H_n$  and  $x \in H_n$  such that  $x_j \rightarrow x$  in  $H_n$  as  $j \rightarrow \infty$ . Then we have

$$\begin{aligned} \|D_{\mathcal{G}}^2 \widehat{B}_\lambda(x_j) - D_{\mathcal{G}}^2 \widehat{B}_\lambda(x)\|_{\mathcal{L}(H_n, H_n)} &= \sup_{\|h\|_{H_n} \leq 1} \left\{ \|D_{\mathcal{G}}^2 \widehat{B}_\lambda(x_j)h - D_{\mathcal{G}}^2 \widehat{B}_\lambda(x)h\|_{H_n} \right\} \\ &= \sup_{\|h\|_{H_n} \leq 1} \sup_{\|k\|_{H_n} \leq 1} \left\{ ((D_{\mathcal{G}} B_\lambda(x_j) - D_{\mathcal{G}} B_\lambda(x))h, k)_{H_n} \right\}. \end{aligned}$$

Now, for any such arbitrary  $h, k \in H_n$ , we can write

$$h = \sum_{i=1}^n a_i^h e_i, \quad k = \sum_{i=1}^n a_i^k e_i,$$

where

$$a_i^h := (h, e_i)_H, \quad a_i^k := (k, e_i)_H, \quad i = 1, \dots, n.$$

Clearly,  $\|h\|_{H_n} \leq 1$  and  $\|k\|_{H_n} \leq 1$  if and only if

$$\sum_{i=1}^n |a_i^h|^2 \leq 1, \quad \sum_{i=1}^n |a_i^k|^2 \leq 1.$$

Consequently, setting  $B_n$  as the closed unit ball in  $\mathbb{R}^n$ , we have  $a^h := (a_1^h, \dots, a_n^h) \in B_n$  and  $a^k := (a_1^k, \dots, a_n^k) \in B_n$ . We deduce that

$$\begin{aligned} \left\| D_{\mathcal{G}}^2 \widehat{B}_\lambda(x_j) - D_{\mathcal{G}}^2 \widehat{B}_\lambda(x) \right\|_{\mathcal{L}(H_n, H_n)} &= \sup_{a^h, a^k \in B_n} \left\{ \sum_{i,l=1}^n a_i^h a_l^k ((D_{\mathcal{G}} B_\lambda(x_j) - D_{\mathcal{G}} B_\lambda(x)) e_i, e_l)_{H_n} \right\} \\ &\leq \sum_{i,l=1}^n |((D_{\mathcal{G}} B_\lambda(x_j) - D_{\mathcal{G}} B_\lambda(x)) e_i, e_l)_H|, \end{aligned}$$

where the right-hand side converges to 0 when  $j \rightarrow \infty$  because  $D_{\mathcal{G}} B_\lambda \in C^0(H; \mathcal{L}_w(H, H))$  by Lemma 3.1. Hence, we have proved that  $\widehat{B}_{\lambda|_{H_n}} \in C^2(H_n)$  for every  $n \in \mathbb{N}$ , and the finite dimensional Itô formula yields then

$$\begin{aligned} \widehat{B}_\lambda(v_n(t)) + \int_0^t (f_n(s), B_\lambda(v_n(s)))_H ds &= \widehat{B}_\lambda(v_n(0)) + \frac{1}{2} \int_0^t \text{Tr} [C_n^*(s) D_{\mathcal{G}} B_\lambda(v_n(s)) C_n(s)] ds \\ &\quad + \int_0^t (B_\lambda(v_n(s)), C_n(s) dW(s))_H \quad \forall t \in [0, T], \quad \mathbb{P}\text{-a.s.} \end{aligned}$$

We want to pass now to the limit as  $n \rightarrow \infty$ . To this end, note that by the properties of  $P_n$  we have that

$$\begin{aligned} v_n &\rightarrow v && \text{in } L^2_{\mathcal{D}}(\Omega; L^\ell(0, T; H)) \quad \forall \ell \geq 1, \\ v_n(t) &\rightarrow v(t) && \text{in } L^2(\Omega, \mathcal{F}_t; H), \quad \forall t \in [0, T], \\ f_n &\rightarrow f && \text{in } L^2_{\mathcal{D}}(\Omega; L^2(0, T; H)), \\ C_n &\rightarrow C && \text{in } L^2_{\mathcal{D}}(\Omega; L^2(0, T; \mathcal{L}^2(U, H))). \end{aligned}$$

Moreover, by the Lipschitz-continuity of  $B_\lambda$ , the quadratic growth of  $\widehat{B}_\lambda$ , and the Vitali Convergence Theorem ensure that

$$\begin{aligned} B_\lambda(v_n) &\rightarrow B_\lambda(v) && \text{in } L^2_{\mathcal{D}}(\Omega; L^\ell(0, T; H)) \quad \forall \ell \geq 1, \\ \widehat{B}_\lambda(v_n(t)) &\rightarrow \widehat{B}_\lambda(v(t)) && \text{in } L^1(\Omega, \mathcal{F}_t), \quad \forall t \in [0, T]. \end{aligned}$$

Furthermore, Lemma 3.1 and the definition of Yosida approximation imply that

$$D_{\mathcal{G}} B_\lambda \in C^0(H; \mathcal{L}_w(H, H)),$$

so that by the Dominated Convergence Theorem we also infer that

$$\text{Tr} [C_n^* D_{\mathcal{G}} B_\lambda(v_n) C_n] \rightarrow \text{Tr} [C^* D_{\mathcal{G}} B_\lambda(v) C] \quad \text{in } L^1_{\mathcal{D}}(\Omega; L^1(0, T)).$$

Eventually, noting that by the dominated convergence theorem

$$B_\lambda(v_n(s)) C_n \rightarrow B_\lambda(v_n) C \quad \text{in } L^2_{\mathcal{D}}(\Omega; L^2(0, T; \mathcal{L}^2(U, \mathbb{R}))),$$

the Burkholder-Davis-Gundy inequality yields

$$\int_0^\cdot (B_\lambda(v_n(s)), C_n(s) dW(s))_H \rightarrow \int_0^\cdot (B_\lambda(v(s)), C(s) dW(s))_H \quad \text{in } L^2_{\mathcal{P}}(\Omega; C^0([0, T])).$$

Taking this information into account, we can let  $n \rightarrow \infty$  in (11), and deduce that Itô's formula holds for a suitable  $d\mathbb{P} \otimes dt$ -version of  $v$ , denoted by the same symbol for brevity of notation.  $\square$

We are now ready to prove the most general version of Itô formula for  $\widehat{B}$ . Let us stress again that this is not obvious at all, as the second derivative of  $\widehat{B}$  is not everywhere defined and is intended only in the sense of Gâteaux.

**Proposition 3.3** (Itô formula for  $\widehat{B}$ ). *Let the processes  $v$  and  $f$  satisfy*

$$\begin{aligned} v &\in L^2_{\mathcal{P}}(\Omega; C^0([0, T]; H)), & \widehat{B}(v(t)) &\in L^1(\Omega, \mathcal{F}_t) \quad \forall t \in [0, T], \\ v &\in D(B) \quad \text{a.e. in } \Omega \times (0, T), & B(v) &\in L^2_{\mathcal{P}}(\Omega; L^2(0, T; H)), \\ f &\in L^2_{\mathcal{P}}(\Omega; L^2(0, T; H)), \end{aligned}$$

and

$$v(t) + \int_0^t f(s) ds = v(0) + \int_0^t G(s, v(s)) dW(s) \quad \forall t \in [0, T], \quad \mathbb{P}\text{-a.s.}$$

Then, it holds that

$$\begin{aligned} \widehat{B}(v(t)) + \int_0^t (f(s), B(v(s)))_H ds &= \widehat{B}(v(0)) + \frac{1}{2} \int_0^t \text{Tr}[L(s, v(s))] ds \\ &\quad + \int_0^t (B(v(s)), G(s, v(s)) dW(s))_H \quad \forall t \in [0, T], \quad \mathbb{P}\text{-a.s.} \end{aligned}$$

*Proof.* For every  $\lambda > 0$ , there exists a unique  $v_\lambda \in L^2_{\mathcal{P}}(\Omega; C^0([0, T]; H))$  such that

$$v_\lambda(t) + \int_0^t f(s) ds = v(0) + \int_0^t G(s, J_\lambda^B(v_\lambda(s))) dW(s) \quad \forall t \in [0, T], \quad \mathbb{P}\text{-a.s.}$$

Hence, it holds that

$$(v - v_\lambda)(t) = \int_0^t (G(s, v(s)) - G(s, J_\lambda^B(v_\lambda(s)))) dW(s) \quad \forall t \in [0, T], \quad \mathbb{P}\text{-a.s.}$$

Itô's isometry, the Lipschitz-continuity of  $G$ , and the properties of  $J_\lambda^B$  and  $B_\lambda$  yield then, for every  $t \in [0, T]$ ,

$$\begin{aligned} \mathbb{E} \|(v - v_\lambda)(t)\|_H^2 &= \mathbb{E} \int_0^t \|G(s, v(s)) - G(s, J_\lambda^B(v_\lambda(s)))\|_{\mathcal{L}^2(U, H)}^2 ds \\ &\leq C_G \int_0^t \mathbb{E} \|(v - J_\lambda^B(v_\lambda))(s)\|_H^2 ds \\ &\leq 2C_G \int_0^t \mathbb{E} \|(v - J_\lambda^B(v))(s)\|_H^2 ds + 2C_G \int_0^t \mathbb{E} \|(J_\lambda^B(v) - J_\lambda^B(v_\lambda))(s)\|_H^2 ds \\ &\leq 2C_G \lambda^2 \int_0^t \mathbb{E} \|B_\lambda(v(s))\|_H^2 ds + 2C_G \int_0^t \mathbb{E} \|(v - v_\lambda)(s)\|_H^2 ds \\ &\leq 2C_G \lambda^2 \|B(v)\|_{L^2_{\mathcal{P}}(\Omega; L^2(0, T; H))}^2 + 2C_G \int_0^t \mathbb{E} \|(v - v_\lambda)(s)\|_H^2 ds. \end{aligned}$$

The Gronwall lemma implies that there exists  $M > 0$ , independent of  $\lambda$ , such that

$$\|v_\lambda - v\|_{C^0([0,T];L^2(\Omega;H))} \leq M\lambda.$$

In particular, it holds that  $v_\lambda \rightarrow v$  in  $C^0([0,T];L^2(\Omega;H))$ . A classical argument involving the Burkholder-David-Gundy inequality allows then to show that

$$v_\lambda \rightarrow v \quad \text{in } L^2_{\mathcal{D}}(\Omega;C^0([0,T];H)). \quad (12)$$

Moreover, since  $B_\lambda$  is  $\frac{1}{\lambda}$ -Lipschitz-continuous on  $H$ , by the estimate just proved we also have that

$$\begin{aligned} \|B_\lambda(v_\lambda)\|_{L^2_{\mathcal{D}}(\Omega;L^2(0,T;H))} &\leq \|B_\lambda(v)\|_{L^2_{\mathcal{D}}(\Omega;L^2(0,T;H))} + \|B_\lambda(v_\lambda) - B_\lambda(v)\|_{L^2_{\mathcal{D}}(\Omega;L^2(0,T;H))} \\ &\leq \|B(v)\|_{L^2_{\mathcal{D}}(\Omega;L^2(0,T;H))} + \frac{1}{\lambda} \|v_\lambda - v\|_{L^2_{\mathcal{D}}(\Omega;L^2(0,T;H))} \\ &\leq \|B(v)\|_{L^2_{\mathcal{D}}(\Omega;L^2(0,T;H))} + M. \end{aligned}$$

It follows that  $(B_\lambda(v_\lambda))_\lambda$  is uniformly bounded in  $L^2_{\mathcal{D}}(\Omega;L^2(0,T;H))$ : as we already know that  $v_\lambda \rightarrow v$  in  $L^2_{\mathcal{D}}(\Omega;L^2(0,T;H))$ , by strong-weak closure of  $B$  we also have that

$$B_\lambda(v_\lambda) \rightharpoonup B(v) \quad \text{in } L^2_{\mathcal{D}}(\Omega;L^2(0,T;H)). \quad (13)$$

As a consequence, by the strong monotonicity of  $B$  in **B1**, we get

$$c_B \|J_\lambda^B(v_\lambda) - v\|_{L^p_{\mathcal{D}}(\Omega;L^p(0,T;V))}^p \leq \mathbb{E} \int_0^T (B_\lambda(v_\lambda(s)) - B(v(s)), J_\lambda^B(v_\lambda(s)) - v(s))_H \, ds \rightarrow 0,$$

so that

$$J_\lambda^B(v_\lambda) \rightarrow v \quad \text{in } L^p_{\mathcal{D}}(\Omega;L^p(0,T;V)). \quad (14)$$

Now, by Lemma 3.2, for every  $\lambda > 0$  we have,

$$\begin{aligned} &\widehat{B}_\lambda(v_\lambda(t)) + \int_0^t (f(s), B_\lambda(v_\lambda(s)))_H \, ds \\ &= \widehat{B}_\lambda(v(0)) + \frac{1}{2} \int_0^t \text{Tr} [G^*(s, J_\lambda^B(v_\lambda(s))) D_G B_\lambda(v_\lambda(s)) G(s, J_\lambda^B(v_\lambda(s)))] \, ds \\ &\quad + \int_0^t (B_\lambda(v_\lambda(s)), G(s, J_\lambda^B(v_\lambda(s))) \, dW(s))_H \quad \forall t \in [0, T], \quad \mathbb{P}\text{-a.s.} \end{aligned} \quad (15)$$

By definition of  $\widehat{B}_\lambda$  and  $B_\lambda$ , we have that

$$\widehat{B}_\lambda(v_\lambda) = \widehat{B}(J_\lambda(v_\lambda)) + \frac{\lambda}{2} \|B_\lambda(v_\lambda)\|_H^2 = (B_\lambda(v_\lambda), J_\lambda(v_\lambda))_H - \widehat{B}^*(B_\lambda(v_\lambda)) + \frac{\lambda}{2} \|B_\lambda(v_\lambda)\|_H^2,$$

from which, recalling (13) and that  $\widehat{B}^*$  is weakly lower semicontinuous on  $H$ ,

$$\begin{aligned} &\limsup_{\lambda \searrow 0} \mathbb{E} \int_0^T \widehat{B}_\lambda(v_\lambda(s)) \, ds \\ &\leq \lim_{\lambda \searrow 0} \mathbb{E} \int_0^T (B_\lambda(v_\lambda(s)), J_\lambda(v_\lambda(s)))_H \, ds - \liminf_{\lambda \searrow 0} \mathbb{E} \int_0^T \widehat{B}^*(B_\lambda(v_\lambda(s))) \, ds \\ &\leq \mathbb{E} \int_0^T (B(v(s)), v(s))_H \, ds - \mathbb{E} \int_0^T \widehat{B}^*(B(v(s))) \, ds = \mathbb{E} \int_0^T \widehat{B}(v(s)) \, ds. \end{aligned}$$

As the  $\liminf$  inequality is immediate due to the lower semicontinuity of  $\widehat{B}$ , we infer that

$$\widehat{B}_\lambda(v_\lambda) \rightarrow \widehat{B}(v) \quad \text{in } L^1_{\mathcal{D}}(\Omega;L^1(0,T)). \quad (16)$$

Furthermore, by the Lipschitz continuity of  $G$  we get

$$\begin{aligned} & \|B_\lambda(v_\lambda)G(\cdot, J_\lambda^B(v_\lambda)) - B(v)G(\cdot, v)\|_{\mathcal{L}^2(U, \mathbb{R})} \\ & \leq \|B_\lambda(v_\lambda)(G(\cdot, J_\lambda^B(v_\lambda)) - G(\cdot, v))\|_{\mathcal{L}^2(U, \mathbb{R})} + \|(B_\lambda(v_\lambda) - B(v))G(\cdot, v)\|_{\mathcal{L}^2(U, \mathbb{R})} \\ & \leq C_G \|B_\lambda(v_\lambda)\|_H \|J_\lambda^B(v_\lambda) - v\|_H + \|(B_\lambda(v_\lambda) - B(v))G(\cdot, v)\|_{\mathcal{L}^2(U, \mathbb{R})}, \end{aligned}$$

so that from (12)–(13) and the Dominated Convergence Theorem we deduce that

$$B_\lambda(v_\lambda)G(\cdot, J_\lambda^B(v_\lambda)) \rightarrow B(v)G(\cdot, v) \quad \text{in } L^2_{\mathcal{P}}(\Omega; L^2(0, T; \mathcal{L}^2(U, \mathbb{R}))).$$

It follows, thanks to the Burkholder-Davis-Gundy inequality, that

$$\begin{aligned} & \int_0^\cdot (B_\lambda(v_\lambda(s)), G(s, J_\lambda^B(v_\lambda(s))) dW(s))_H \\ & \rightarrow \int_0^\cdot (B(v(s)), G(s, v(s))) dW(s))_H \quad \text{in } L^2_{\mathcal{P}}(\Omega; C^0([0, T])). \end{aligned} \quad (17)$$

It remains to show that we can let  $\lambda \searrow 0$  in the trace term. To this end, note that by Lemma 3.1 and the definition of  $L$  in  $\mathbf{G}$  we have

$$\begin{aligned} \text{Tr} [G^*(\cdot, J_\lambda^B(v_\lambda)) D_{\mathcal{G}} B_\lambda(v_\lambda) G(\cdot, J_\lambda^B(v_\lambda))] &= \text{Tr} [G(\cdot, J_\lambda^B(v_\lambda)) G^*(\cdot, J_\lambda^B(v_\lambda)) D_{\mathcal{G}} B_\lambda(v_\lambda)] \\ &= \text{Tr} [L(\cdot, J_\lambda^B(v_\lambda)) D_{\mathcal{G}} J_\lambda^B(v_\lambda)], \end{aligned}$$

where, by the strong convergence (14), the continuity of  $L$  in assumption  $\mathbf{G}$ , and again Lemma 3.1,

$$\begin{aligned} L(\cdot, J_\lambda^B(v_\lambda)) &\rightarrow L(\cdot, v) \quad \text{in } \mathcal{L}^1(H, H), \quad \text{a.e. in } \Omega \times (0, T), \\ D_{\mathcal{G}} J_\lambda^B(v_\lambda) &\rightarrow I \quad \text{in } \mathcal{L}_s(H, H), \quad \text{a.e. in } \Omega \times (0, T). \end{aligned}$$

It follows then that

$$\text{Tr} [L(\cdot, J_\lambda^B(v_\lambda)) D_{\mathcal{G}} J_\lambda^B(v_\lambda)] \rightarrow \text{Tr} [L(\cdot, v)] \quad \text{a.e. in } \Omega \times (0, T).$$

Moreover, by assumption  $\mathbf{G}$  we have that

$$\begin{aligned} |\text{Tr} [L(\cdot, J_\lambda^B(v_\lambda)) D_{\mathcal{G}} J_\lambda^B(v_\lambda)]| &\leq \|L(\cdot, J_\lambda^B(v_\lambda))\|_{\mathcal{L}^1(H, H)} \|D_{\mathcal{G}} J_\lambda^B(v_\lambda)\|_{\mathcal{L}(H, H)} \\ &\leq \|L(\cdot, J_\lambda^B(v_\lambda))\|_{\mathcal{L}^1(H, H)} \\ &\leq h_G(\cdot) + C_G \|J_\lambda^B(v_\lambda)\|_V^p. \end{aligned}$$

From (14) the right-hand converges in  $L^1_{\mathcal{P}}(\Omega; L^1(0, T))$ , hence it is uniformly integrable, and so is by comparison the left-hand side. Putting all this information together, the Vitali Convergence Theorem yields then

$$\text{Tr} [G^*(\cdot, J_\lambda^B(v_\lambda)) D_{\mathcal{G}} B_\lambda(v_\lambda) G(\cdot, J_\lambda^B(v_\lambda))] \rightarrow \text{Tr} [L(\cdot, v)] \quad \text{in } L^1_{\mathcal{P}}(\Omega; L^1(0, T)). \quad (18)$$

Letting then  $\lambda \searrow 0$  in (15) and using the convergences (12)–(14) and (16)–(18) we conclude.  $\square$

#### 4. EXISTENCE OF MARTINGALE SOLUTIONS

This section is devoted to the proof of Theorem 2.1. The proof is based on a Yosida-type approximation on the operators  $A$  and  $B$ . The passage to the limit hinges then on a lower-semicontinuity arguments, which in turn makes uses of the Itô formula for  $\hat{B}$  from Proposition 3.3. For the sake of clarity, we subdivide the argument in subsequent steps in the coming subsections.



**4.1. Approximation.** For every  $\lambda > 0$  let  $A_\lambda, B_\lambda : H \rightarrow H$  be the Yosida approximations of  $A$  and  $B$ , respectively, which we recall to be a maximal monotone  $\frac{1}{\lambda}$ -Lipschitz-continuous operators on  $H$ . The respective resolvents are denoted by  $J_\lambda^A, J_\lambda^B : H \rightarrow H$ . Let us also recall that  $R_\lambda \in \mathcal{L}(H, V)$  is a regularizing operator converging to the identity in  $\mathcal{L}_s(V, V)$ .

The approximated reads as follows:

$$\begin{aligned} & \text{find } u_\lambda \in \mathcal{I}^{2,2}(H, H) \text{ such that} \\ & \begin{cases} \lambda \partial_t u_\lambda^d + A_\lambda(\partial_t u_\lambda^d) + B_\lambda(u_\lambda) = F(\cdot, R_\lambda J_\lambda^B(u_\lambda)), \\ u_\lambda^d(0) = u_0, \\ u_\lambda^s = G(\cdot, J_\lambda^B(u_\lambda)). \end{cases} \end{aligned} \quad (19)$$

More precisely, this means that we look for an  $H$ -valued continuous process  $u_\lambda$  such that

$$u_\lambda = u_0 + \int_0^\cdot \partial_t u_\lambda^d(s) ds + \int_0^\cdot G(s, J_\lambda^B(u_\lambda(s))) dW(s), \quad (20)$$

where  $u_\lambda^d \in L^2_{\mathcal{D}}(\Omega; H^1(0, T; H))$  is such that

$$\lambda \partial_t u_\lambda^d + A_\lambda(\partial_t u_\lambda^d) + B_\lambda(u_\lambda) = F(\cdot, R_\lambda J_\lambda^B(u_\lambda)) \quad \text{in } L^2_{\mathcal{D}}(\Omega; L^2(0, T; H)). \quad (21)$$

By definition of  $A_\lambda$ , it is readily seen that  $\lambda I + A_\lambda : H \rightarrow H$  is Lipschitz-continuous and strongly monotone: hence, the inverse operator  $(\lambda I + A_\lambda)^{-1} : H \rightarrow H$  is well-defined and Lipschitz-continuous as well. It follows that we can equivalently rewrite the differential relation above as

$$\partial_t u_\lambda^d = (\lambda I + A_\lambda)^{-1}(F(\cdot, R_\lambda J_\lambda^B(u_\lambda)) - B_\lambda(u_\lambda)).$$

Then, the approximated problem (20)–(21) can be written in so-called normal form as

$$du_\lambda = (\lambda I + A_\lambda)^{-1}(F(\cdot, R_\lambda J_\lambda^B(u_\lambda)) - B_\lambda(u_\lambda)) dt + G(\cdot, J_\lambda^B(u_\lambda)) dW, \quad u_\lambda(0) = u_0.$$

Now, it is clear that by assumption **G** and the fact that  $J_\lambda^B : H \rightarrow H$  is Lipschitz-continuous, the operator  $G(\cdot, J_\lambda^B) : [0, T] \times H \rightarrow \mathcal{L}^2(U, H)$  is Lipschitz-continuous and linearly bounded in  $H$ , uniformly on  $[0, T]$ . Moreover, recall also that  $(\lambda I + A_\lambda)^{-1}$  and  $B_\lambda$  are Lipschitz-continuous on  $H$ , and that  $R_\lambda : H \rightarrow V$  is linear continuous (so in particular Lipschitz-continuous). Hence, bearing in mind that by Lemma 3.1 we have

$$\|J_\lambda^B(x_1) - J_\lambda^B(x_2)\|_V^p \leq \frac{1}{\lambda c_B} \|x_1 - x_2\|_H^2 \quad \forall x_1, x_2 \in H,$$

we deduce that the operator

$$S_\lambda(\cdot, x) := (\lambda I + A_\lambda)^{-1}(F(\cdot, R_\lambda J_\lambda^B(x)) - B_\lambda(x)), \quad x \in H,$$

is locally Lipschitz-continuous and linearly bounded on  $H$ . Indeed, by assumption **F** we have

$$\begin{aligned} \|S_\lambda(\cdot, x_1) - S_\lambda(\cdot, x_2)\|_H & \leq M_\lambda \left( 1 + \|J_\lambda^B(x_1)\|_V^{\frac{p-2}{2}} + \|J_\lambda^B(x_2)\|_V^{\frac{p-2}{2}} \right) \|x_1 - x_2\|_H \\ & \leq M'_\lambda \left( 1 + \|x_1\|_H^{\frac{p-2}{p}} + \|x_2\|_H^{\frac{p-2}{p}} \right) \|x_1 - x_2\|_H \quad \forall x_1, x_2 \in H \end{aligned}$$

and

$$\|S_\lambda(\cdot, x)\|_H \leq M'_\lambda (1 + |h_F(\cdot)| + \|x\|_H) \quad \forall x \in H,$$

for certain constants  $M_\lambda, M'_\lambda > 0$ . Since  $\frac{p-2}{p} \leq 1$ , we can apply the classical existence-uniqueness results for SPDEs with locally Lipschitz coefficients (see [24]), and infer that the approximated problem (20)–(21) admits a unique global solution

$$u_\lambda \in \mathcal{I}^{2,2}(H, H).$$

**4.2. Uniform estimates.** We prove here some estimates on the approximated solutions uniformly in  $\lambda$ . To this end, from (20), we can write Itô's formula for  $\widehat{B}_\lambda$  by using Lemma 3.2, getting

$$\begin{aligned}\widehat{B}_\lambda(u_\lambda(t)) &= \widehat{B}_\lambda(u_0) + \int_0^t (\partial_t u_\lambda^d(s), B_\lambda(u_\lambda(s)))_H \, ds \\ &\quad + \frac{1}{2} \int_0^t \text{Tr} [G(s, J_\lambda^B(u_\lambda(s)))^* D_{\mathcal{G}} B_\lambda(u_\lambda(s)) G(s, J_\lambda^B(u_\lambda(s)))] \, ds \\ &\quad + \int_0^t (B_\lambda(u_\lambda(s)), G(s, J_\lambda^B(u_\lambda(s))) \, dW(s))_H \quad \forall t \in [0, T], \quad \mathbb{P}\text{-a.s.}\end{aligned}$$

Taking equation (21) into account, this leads to

$$\begin{aligned}\widehat{B}_\lambda(u_\lambda(t)) &+ \lambda \int_0^t \|\partial_t u_\lambda^d(s)\|_H^2 \, ds + \int_0^t (A_\lambda(\partial_t u_\lambda^d(s)), \partial_t u_\lambda^d(s))_H \, ds \\ &= \widehat{B}_\lambda(u_0) + \int_0^t (F(s, R_\lambda J_\lambda^B(u_\lambda(s))), \partial_t u_\lambda^d(s))_H \, ds + \int_0^t (B_\lambda(u_\lambda(s)), G(s, J_\lambda^B(u_\lambda(s))) \, dW(s))_H \\ &\quad + \frac{1}{2} \int_0^t \text{Tr} [G(s, J_\lambda^B(u_\lambda(s)))^* D_{\mathcal{G}} B_\lambda(u_\lambda(s)) G(s, J_\lambda^B(u_\lambda(s)))] \, ds \quad \forall t \in [0, T], \quad \mathbb{P}\text{-a.s.} \quad (22)\end{aligned}$$

On the left-hand side, the coercivity of  $A$  in assumption **A** and the definition of  $\widehat{B}_\lambda$  and  $A_\lambda$  give

$$\begin{aligned}\widehat{B}_\lambda(u_\lambda(t)) &+ \int_0^t (A_\lambda(\partial_t u_\lambda^d(s)), \partial_t u_\lambda^d(s))_H \, ds \\ &\geq \widehat{B}(J_\lambda^B(u_\lambda(t))) + \int_0^t (A_\lambda(\partial_t u_\lambda^d(s)), J_\lambda^A(\partial_t u_\lambda^d(s)))_H \, ds + \lambda \int_0^t \|A_\lambda(\partial_t u_\lambda^d(s))\|_H^2 \, ds \\ &\geq \widehat{B}(J_\lambda^B(u_\lambda(t))) + c_A \int_0^t \|J_\lambda^A(\partial_t u_\lambda^d(s))\|_H^2 \, ds - c_A^{-1} T + \lambda \int_0^t \|A_\lambda(\partial_t u_\lambda^d(s))\|_H^2 \, ds.\end{aligned}$$

On the right-hand side, first of all it is clear that

$$\widehat{B}_\lambda(u_0) \leq \widehat{B}(u_0) \in L^1(\Omega).$$

Secondly, assumption **F**, the weighted Young inequality, and the uniform boundedness of  $(R_\lambda)_\lambda$  in  $\mathcal{L}(V, V)$  imply that, for every  $\delta > 0$ ,

$$\begin{aligned}\int_0^t (F(s, R_\lambda J_\lambda^B(u_\lambda(s))), \partial_t u_\lambda^d(s))_H \, ds &\leq \int_0^t \|F(s, R_\lambda J_\lambda^B(u_\lambda(s)))\|_H \|\partial_t u_\lambda^d(s)\|_H \, ds \\ &\leq \delta \int_0^t \|\partial_t u_\lambda^d(s)\|_H^2 \, ds + \frac{1}{4\delta} \left( \|h_F\|_{L^1(0, T)} + C_F \int_0^t \|R_\lambda J_\lambda^B(u_\lambda(s))\|_V^p \, ds \right) \\ &\leq \delta \int_0^t \|\partial_t u_\lambda^d(s)\|_H^2 \, ds + M_\delta \left( 1 + \int_0^t \|J_\lambda^B(u_\lambda(s))\|_V^p \, ds \right),\end{aligned}$$

where  $M_\delta > 0$  is a positive constant independent of  $\lambda$ . Noting further that

$$\int_0^t \|\partial_t u_\lambda^d(s)\|_H^2 \, ds \leq 2 \int_0^t \|J_\lambda^A(\partial_t u_\lambda^d(s))\|_H^2 \, ds + 2\lambda^2 \int_0^t \|A_\lambda(\partial_t u_\lambda^d(s))\|_H^2 \, ds,$$

taking  $\lambda \in (0, 1)$  and choosing  $\delta$  sufficient small, rearranging the terms in (22) we infer that there exists a constant  $M > 0$  independent of  $\lambda$  such that

$$\begin{aligned} & \widehat{B}(J_\lambda^B(u_\lambda(t))) + \frac{c_A}{2} \int_0^t \|J_\lambda^A(\partial_t u_\lambda^d(s))\|_H^2 ds + \frac{\lambda}{2} \int_0^t \|\partial_t u_\lambda^d(s)\|_H^2 ds \\ & \leq M \left( 1 + \int_0^t \|J_\lambda^B(u_\lambda(s))\|_V^p ds \right) + \int_0^t (B_\lambda(u_\lambda(s)), G(s, J_\lambda^B(u_\lambda(s))) dW(s))_H \\ & \quad + \frac{1}{2} \int_0^t \text{Tr} [G(s, J_\lambda^B(u_\lambda(s)))^* D_G B_\lambda(u_\lambda(s)) G(s, J_\lambda^B(u_\lambda(s)))] ds \quad \forall t \in [0, T], \quad \mathbb{P}\text{-a.s.} \end{aligned}$$

At this point, the trace term can be handled using assumption **G**, Lemma 3.1, and the fact that  $\|D_G J_\lambda^B(u_\lambda)\|_{\mathcal{L}(H, H)} \leq 1$  as

$$\begin{aligned} & \int_0^t \text{Tr} [G(s, J_\lambda^B(u_\lambda(s)))^* D_G B_\lambda(u_\lambda(s)) G(s, J_\lambda^B(u_\lambda(s)))] ds \\ & = \int_0^t \text{Tr} [G(s, J_\lambda^B(u_\lambda(s))) G(s, J_\lambda^B(u_\lambda(s)))^* D_G B(J_\lambda^B(u_\lambda(s))) D_G J_\lambda^B(u_\lambda(s))] ds \\ & = \int_0^t \text{Tr} [L(s, J_\lambda^B(u_\lambda(s))) D_G J_\lambda^B(u_\lambda(s))] ds \\ & \leq \int_0^t \|L(s, J_\lambda^B(u_\lambda(s)))\|_{\mathcal{L}^1(H, H)} \|D_G J_\lambda^B(u_\lambda(s))\|_{\mathcal{L}(H, H)} ds \\ & \leq \|h_G\|_{L^1(0, T)} + C_G \int_0^t \|J_\lambda^B(u_\lambda(s))\|_V^p ds \end{aligned}$$

Furthermore, let  $x \in V$  be arbitrary and note that by **B1** we have

$$\begin{aligned} \widehat{B}(x) & \geq \widehat{B}_\varepsilon(x) = \int_0^1 (B_\varepsilon(rx), x)_H dr = \int_0^1 r^{-1} (B_\varepsilon(rx), J_\varepsilon^B(rx))_H dr + \varepsilon \int_0^1 r^{-1} \|B_\varepsilon(rx)\|_H^2 dr \\ & \geq c_B \int_0^1 r^{-1} \|J_\varepsilon^B(rx)\|_V^p dr \quad \forall \varepsilon > 0. \end{aligned}$$

Letting  $\varepsilon \rightarrow 0$ , by Lemma 3.1 we have  $J_\varepsilon^B(rx) \rightharpoonup rx$  in  $V$  for every  $r \in [0, 1]$ , so that by the Fatou Lemma it follows that

$$\widehat{B}(x) \geq \frac{c_B}{p} \|x\|_V^p \quad \forall x \in V. \quad (23)$$

Consequently, possibly updating the value of the constant  $M$ , we are left with

$$\begin{aligned} \widehat{B}(J_\lambda^B(u_\lambda(t))) + \int_0^t \|J_\lambda^A(\partial_t u_\lambda^d(s))\|_H^2 ds & \leq M \left( 1 + \int_0^t \widehat{B}(J_\lambda^B(u_\lambda(s))) ds \right) \\ & \quad + \int_0^t (B_\lambda(u_\lambda(s)), G(s, J_\lambda^B(u_\lambda(s))) dW(s))_H \end{aligned} \quad (24)$$

for every  $t \in [0, T]$ ,  $\mathbb{P}$ -almost surely. The stochastic integral is a martingale, so that taking expectations and using the Gronwall Lemma yield

$$\left\| \widehat{B}(J_\lambda^B(u_\lambda)) \right\|_{L^\infty(0, T; L^1(\Omega))} + \|J_\lambda^A(\partial_t u_\lambda^d)\|_{L^2_{\mathcal{D}}(\Omega; L^2(0, T; H))} \leq M. \quad (25)$$

By (23) and the linear growth of  $A$  in assumption **A**, this implies in turn that

$$\|J_\lambda^B(u_\lambda)\|_{L^\infty(0, T; L^p(\Omega; V))} + \|A_\lambda(\partial_t u_\lambda^d)\|_{L^2_{\mathcal{D}}(\Omega; L^2(0, T; H))} \leq M. \quad (26)$$

Consequently, assumptions **F–G** yield directly

$$\|F(\cdot, R_\lambda J_\lambda^B(u_\lambda))\|_{L^2_{\mathcal{F}}(\Omega; L^2(0, T; H))} + \|G(\cdot, J_\lambda^B(u_\lambda))\|_{L^\infty(0, T; L^p(\Omega; \mathcal{L}^2(U, H)))} \leq M, \quad (27)$$

from which we deduce, by comparison in (21), that

$$\|B_\lambda(u_\lambda)\|_{L^2_{\mathcal{F}}(\Omega; L^2(0, T; H))} \leq M. \quad (28)$$

At this point, going back to (24) and using the estimates (27)–(28) together with the Burkholder-Davis-Gundy inequality on the stochastic integral, we get by a standard argument that

$$\|\widehat{B}(J_\lambda^B(u_\lambda))\|_{L^1_{\mathcal{F}}(\Omega; L^\infty(0, T))} + \|J_\lambda^B(u_\lambda)\|_{L^p_{\mathcal{F}}(\Omega; L^\infty(0, T; V))} \leq M, \quad (29)$$

which in turn implies, again by assumptions **F–G**, that, setting  $p_\nu := p/\nu$ ,

$$\|F(\cdot, R_\lambda J_\lambda^B(u_\lambda))\|_{L^2_{\mathcal{F}}(\Omega; L^2(0, T; H))} + \|G(\cdot, J_\lambda^B(u_\lambda))\|_{L^{p_\nu}_{\mathcal{F}}(\Omega; L^\infty(0, T; \mathcal{L}^2(U, H)))} \leq M. \quad (30)$$

It follows in particular by the Hölder inequality that, setting  $q_\nu := 2p_\nu/(p_\nu + 2)$ ,

$$\|B_\lambda(u_\lambda)G(\cdot, J_\lambda^B(u_\lambda))\|_{L^{q_\nu}_{\mathcal{F}}(\Omega; L^2(0, T; \mathcal{L}^2(U, \mathbb{R})))} \leq M,$$

where by assumption on  $\nu$  we always have that  $q_\nu > 1$ . Hence, going back again to (24) the Burkholder-Davis-Gundy inequality and the Gronwall lemma allow to refine the moment estimates as

$$\|\widehat{B}(J_\lambda^B(u_\lambda))\|_{L^{q_\nu}_{\mathcal{F}}(\Omega; L^\infty(0, T))} + \|J_\lambda^B(u_\lambda)\|_{L^{pq_\nu}_{\mathcal{F}}(\Omega; L^\infty(0, T; V))} \leq M \quad (31)$$

and

$$\|J_\lambda^A(\partial_t u_\lambda^d)\|_{L^{2q_\nu}_{\mathcal{F}}(\Omega; L^2(0, T; H))} + \|A_\lambda(\partial_t u_\lambda^d)\|_{L^{2q_\nu}_{\mathcal{F}}(\Omega; L^2(0, T; H))} \leq M, \quad (32)$$

Lastly, the classical result [17, Lem. 2.1] by FLANDOLI & GATAREK ensures that

$$\left\| I_\lambda := \int_0^\cdot G(s, J_\lambda^B(u_\lambda(s))) \right\|_{L^{p_\nu}_{\mathcal{F}}(\Omega; W^{\eta, p_\nu}(0, T; H))} \leq M_\eta \quad \forall \eta \in (0, 1/2), \quad (33)$$

yielding by comparison in (20) that

$$\|u_\lambda\|_{L^2_{\mathcal{F}}(\Omega; H^1(0, T; H)) + L^{p_\nu}_{\mathcal{F}}(\Omega; W^{\eta, p_\nu}(0, T; H))} \leq M_\eta \quad \forall \eta \in (0, 1/2). \quad (34)$$

**4.3. Passage to the limit.** First of all, note that by assumption on  $\nu$  in **G**, we always have that  $p_\nu = p/\nu > 2$  and  $q_\nu > 1$ : hence, we can fix  $\eta \in (\frac{1}{p_\nu}, \frac{1}{2})$ , so that  $\eta p_\nu > 1$ . Since  $V \hookrightarrow H$  compactly, by the classical Aubin-Lions-Simon compactness results [46, Cor. 4–5, p. 85] we have

$$\begin{aligned} W^{\eta, p_\nu}(0, T; H) &\hookrightarrow C^0([0, T]; V^*) \quad \text{compactly,} \\ L^\infty(0, T; V) \cap (H^1(0, T; H) + W^{\eta, p_\nu}(0, T; H)) &\hookrightarrow C^0([0, T]; H) \quad \text{compactly.} \end{aligned}$$

Let us put then

$$\mathcal{X} := C^0([0, T]; H) \times C^0([0, T]; V^*) \times C^0([0, T]; U).$$

By the estimates (29)–(34) and the compactness inclusions above, using the Prokhorov theorem we readily infer that

$$\text{the laws of } \{(J_\lambda^B(u_\lambda), I_\lambda, W)\}_\lambda \text{ are tight on } \mathcal{X}.$$

By the Skorokhod theorem [21, Thm. 2.7], there exist then a probability space  $(\hat{\Omega}, \hat{\mathcal{F}}, \hat{\mathbb{P}})$ , a sequence of measurable random variables

$$\phi_\lambda : (\hat{\Omega}, \hat{\mathcal{F}}) \rightarrow (\Omega, \mathcal{F}), \quad \lambda > 0,$$

with  $\mathbb{P} \circ \phi_\lambda = \hat{\mathbb{P}}$  for all  $\lambda > 0$ , and some measurable random variables

$$(\hat{u}, \hat{I}, \hat{W}) : (\hat{\Omega}, \hat{\mathcal{F}}) \rightarrow \mathcal{X}$$

such that, setting  $\hat{u}_\lambda := u_\lambda \circ \phi_\lambda$ ,

$$J_\lambda^B(\hat{u}_\lambda) \rightarrow \hat{u} \quad \text{in } C^0([0, T]; H), \quad \hat{\mathbb{P}}\text{-a.s.}, \quad (35)$$

$$\hat{I}_\lambda := I_\lambda \circ \phi_\lambda \rightarrow \hat{I} \quad \text{in } C^0([0, T]; V^*), \quad \hat{\mathbb{P}}\text{-a.s.}, \quad (36)$$

$$\hat{W}_\lambda := W \circ \phi_\lambda \rightarrow \hat{W} \quad \text{in } C^0([0, T]; U), \quad \hat{\mathbb{P}}\text{-a.s.} \quad (37)$$

Furthermore, by (25)–(34) and the fact that composition with  $\phi_\lambda$  preserves the laws, we also infer the convergences

$$J_\lambda^B(\hat{u}_\lambda) \rightarrow \hat{u} \quad \text{in } L^\ell(\hat{\Omega}; C^0([0, T]; H)) \quad \forall \ell \in [1, pq_\nu), \quad (38)$$

$$J_\lambda^B(\hat{u}_\lambda) \xrightarrow{*} \hat{u} \quad \text{in } L^{pq_\nu}(\hat{\Omega}; L^\infty(0, T; V)), \quad (39)$$

$$\partial_t \hat{u}_\lambda^d \rightharpoonup \hat{u}' \quad \text{in } L^2(\hat{\Omega}; L^2(0, T; H)), \quad (40)$$

$$B_\lambda(\hat{u}_\lambda) \rightharpoonup \hat{w} \quad \text{in } L^2(\hat{\Omega}; L^2(0, T; H)), \quad (41)$$

$$A_\lambda(J_\lambda^A(\hat{u}_\lambda)) \rightharpoonup \hat{v} \quad \text{in } L^2(\hat{\Omega}; L^2(0, T; H)), \quad (42)$$

for some  $\hat{u}', \hat{w}, \hat{v} \in L^2(\hat{\Omega}; L^2(0, T; H))$ . Now, noting that  $pq_\nu > 2$ , by the strong-weak closure of the maximal monotone operator  $B$  we readily get that

$$\hat{u} \in D(B), \quad \hat{w} = B(\hat{u}) \quad \text{a.e. in } \hat{\Omega} \times (0, T).$$

Consequently, by the strong monotonicity of  $B$  in **B1** we obtain

$$c_B \|J_\lambda^B(\hat{u}_\lambda) - \hat{u}\|_{L^p(\hat{\Omega}; L^p(0, T; V))}^p \leq \mathbb{E} \int_0^T (B_\lambda(\hat{u}_\lambda(s)) - \hat{w}(s), J_\lambda^B(\hat{u}_\lambda(s)) - \hat{u}(s))_H \, ds \rightarrow 0,$$

so that

$$J_\lambda^B(\hat{u}_\lambda) \rightarrow \hat{u} \quad \text{in } L^p(\hat{\Omega}; L^p(0, T; V)). \quad (43)$$

Since  $R_\lambda \rightarrow I$  in  $\mathcal{L}_s(V, V)$ , the family  $(R_\lambda)_\lambda$  is uniformly bounded in  $\mathcal{L}(V, V)$  by the Banach-Steinhaus theorem, hence the strong convergence (43) yields

$$R_\lambda J_\lambda^B(\hat{u}_\lambda) \rightarrow \hat{u} \quad \text{in } L^p(\hat{\Omega}; L^p(0, T; V)).$$

At this point, using **F** it is immediate to infer that

$$F(\cdot, R_\lambda J_\lambda^B(\hat{u}_\lambda)) \rightarrow F(\cdot, \hat{u}) \quad \text{in } L^2(\hat{\Omega}; L^2(0, T; H)), \quad (44)$$

while **G** gives

$$G(\cdot, J_\lambda^B(\hat{u}_\lambda)) \rightarrow G(\cdot, \hat{u}) \quad \text{in } L^\ell(\hat{\Omega}; C^0([0, T]; \mathcal{L}^2(U, H))) \quad \forall \ell \in [1, pq_\nu), \quad (45)$$

$$G(\cdot, J_\lambda^B(\hat{u}_\lambda)) \rightarrow G(\cdot, \hat{u}) \quad \text{in } L^p(\hat{\Omega}; L^p(0, T; \mathcal{L}^2(U, H))). \quad (46)$$

Now, by definition of  $\phi_\lambda$ , from (20)–(21) we have

$$\hat{u}_\lambda(t) = u_0 + \int_0^t \partial_t \hat{u}_\lambda^d(s) \, ds + \hat{I}_\lambda(t) \quad \text{in } H \quad \forall t \in [0, T], \quad \hat{\mathbb{P}}\text{-a.s.}, \quad (47)$$

$$\lambda \partial_t \hat{u}_\lambda^d + A_\lambda(\partial_t \hat{u}_\lambda^d) + B_\lambda(\hat{u}_\lambda) = F(\cdot, R_\lambda J_\lambda^B(\hat{u}_\lambda)) \quad \text{in } H \quad \text{a.e. in } \hat{\Omega} \times (0, T). \quad (48)$$

Let us introduce on the probability space  $(\hat{\Omega}, \hat{\mathcal{F}}, \hat{\mathbb{P}})$  the filtration

$$\hat{\mathcal{F}}_{\lambda, t} := \sigma \left\{ \hat{u}_\lambda(s), \hat{I}_\lambda(s), \hat{W}_\lambda(s) : s \leq t \right\}, \quad t \in [0, T].$$

Using a classical argument (see [15, § 8] and [48]), we have that  $\hat{W}_\lambda$  is an  $(\hat{\mathcal{F}}_{\lambda,t})_t$ -cylindrical Wiener process. Moreover, following the approach in [38, § 4.5] and [41], using the fact that  $\phi_\lambda$  preserves the laws and comparing (20) and (21), we also deduce that, possibly enlarging the filtration  $(\hat{\mathcal{F}}_{\lambda,t})_t$ ,  $\hat{I}_\lambda$  is the square-integrable martingale

$$\hat{I}_\lambda = \int_0^\cdot G(s, J_\lambda^B(\hat{u}_\lambda(s))) d\hat{W}(s).$$

At this point, letting  $\lambda \searrow 0$  in (47)–(48), thanks to the convergences (35)–(45) we get

$$\hat{u}(t) = \hat{u}_0 + \int_0^t \hat{u}'(s) ds + \hat{I}(t) \quad \text{in } V^* \quad \forall t \in [0, T], \quad \hat{\mathbb{P}}\text{-a.s.}, \quad (49)$$

$$\hat{v} + B(\hat{u}) = F(\cdot, \hat{u}) \quad \text{in } H \quad \text{a.e. in } \hat{\Omega} \times (0, T). \quad (50)$$

We introduce the limiting filtration

$$\hat{\mathcal{F}}_t := \sigma \left\{ \hat{u}(s), \hat{I}(s), \hat{W}(s) : s \leq t \right\}, \quad t \in [0, T],$$

as well as the corresponding progressive  $\sigma$  algebra  $\hat{\mathcal{P}}$ . The strong convergence (37) and the representation results [15, § 8] ensure again that  $\hat{W}$  is a  $(\hat{\mathcal{F}}_t)_t$ -cylindrical Wiener process. Furthermore, proceeding as in [38, § 4.5] and [41], possibly enlarging  $(\hat{\mathcal{F}}_t)_t$  we have the representation

$$\hat{I} = \int_0^\cdot G(s, \hat{u}(s)) d\hat{W}(s).$$

In particular, this shows a posteriori that  $\hat{I} \in L^p_{\hat{\mathcal{P}}}(\hat{\Omega}; C^0([0, T]; H))$ . Consequently, equation (49) yields

$$\hat{u}(t) = u_0 + \int_0^t \hat{u}'(s) ds + \int_0^t G(s, \hat{u}(s)) d\hat{W}(s) \quad \forall t \in [0, T], \quad \hat{\mathbb{P}}\text{-a.s.} \quad (51)$$

We deduce that

$$\hat{u} \in \mathcal{I}^{2,p}(H, H), \quad \partial_t \hat{u}^d = \hat{u}', \quad \hat{u}^s = G(\cdot, \hat{u}).$$

**4.4. Identification of the nonlinearity A.** The last thing that we have to show is that  $\hat{v} \in A(\partial_t \hat{u}^d)$  almost everywhere. To this end, from the Itô formula (22), taking expectations and fixing  $t = T$  we immediately deduce

$$\begin{aligned} & \hat{\mathbb{E}} \hat{B}_\lambda(\hat{u}_\lambda(T)) + \lambda \hat{\mathbb{E}} \int_0^T \|\partial_t \hat{u}_\lambda^d(s)\|_H^2 ds + \hat{\mathbb{E}} \int_0^T (A_\lambda(\partial_t \hat{u}_\lambda^d(s)), \partial_t \hat{u}_\lambda^d(s))_H ds \\ &= \hat{B}_\lambda(u_0) + \hat{\mathbb{E}} \int_0^T (F(s, R_\lambda J_\lambda^B(\hat{u}_\lambda(s))), \partial_t \hat{u}_\lambda^d(s))_H ds \\ &+ \frac{1}{2} \hat{\mathbb{E}} \int_0^T \text{Tr} [G(s, J_\lambda^B(\hat{u}_\lambda(s)))^* D_G B_\lambda(\hat{u}_\lambda(s)) G(s, J_\lambda^B(\hat{u}_\lambda(s)))] ds. \end{aligned}$$

Now, by lower semicontinuity we have

$$\liminf_{\lambda \searrow 0} \left( \hat{\mathbb{E}} \hat{B}_\lambda(\hat{u}_\lambda(T)) + \lambda \hat{\mathbb{E}} \int_0^T \|\partial_t \hat{u}_\lambda^d(s)\|_H^2 ds \right) \geq \hat{\mathbb{E}} \hat{B}(\hat{u}(T)),$$

while from the convergences (40) and (44) it follows

$$\lim_{\lambda \searrow 0} \hat{\mathbb{E}} \int_0^T (F(s, R_\lambda J_\lambda^B(\hat{u}_\lambda(s))), \partial_t \hat{u}_\lambda^d(s))_H ds = \hat{\mathbb{E}} \int_0^T (F(s, \hat{u}(s)), \partial_t \hat{u}^d(s))_H ds.$$

As the initial term, we have  $\widehat{B}_\lambda(u_0) \rightarrow \widehat{B}(u_0)$ . Furthermore, for the trace term we note that

$$\begin{aligned} \operatorname{Tr} [G^*(\cdot, J_\lambda^B(\hat{u}_\lambda)) D_G B_\lambda(\hat{u}_\lambda) G(\cdot, J_\lambda^B(\hat{u}_\lambda))] &= \operatorname{Tr} [G(\cdot, J_\lambda^B(\hat{u}_\lambda)) G^*(\cdot, J_\lambda^B(\hat{u}_\lambda)) D_G B_\lambda(\hat{u}_\lambda)] \\ &= \operatorname{Tr} [L(\cdot, J_\lambda^B(\hat{u}_\lambda)) D_G J_\lambda^B(\hat{u}_\lambda)] . \end{aligned}$$

By the strong convergence (43), the continuity of  $L$  in assumption **G**, and Lemma 3.1, we have

$$\begin{aligned} L(\cdot, J_\lambda^B(\hat{u}_\lambda)) &\rightarrow L(\cdot, \hat{u}) \quad \text{in } \mathcal{L}^1(H, H), \quad \text{a.e. in } \hat{\Omega} \times (0, T), \\ D_G J_\lambda^B(\hat{u}_\lambda) &\rightarrow I \quad \text{in } \mathcal{L}_s(H, H), \quad \text{a.e. in } \hat{\Omega} \times (0, T), \end{aligned}$$

so that

$$\operatorname{Tr} [L(\cdot, J_\lambda^B(\hat{u}_\lambda)) D_G J_\lambda^B(\hat{u}_\lambda)] \rightarrow \operatorname{Tr} [L(\cdot, \hat{u})] \quad \text{a.e. in } \hat{\Omega} \times (0, T).$$

Noting also that by assumption **G** we have

$$\begin{aligned} |\operatorname{Tr} [L(\cdot, J_\lambda^B(\hat{u}_\lambda)) D_G J_\lambda^B(\hat{u}_\lambda)]| &\leq \|L(\cdot, J_\lambda^B(\hat{u}_\lambda))\|_{\mathcal{L}^1(H, H)} \|D_G J_\lambda^B(\hat{u}_\lambda)\|_{\mathcal{L}(H, H)} \\ &\leq \|L(\cdot, J_\lambda^B(\hat{u}_\lambda))\|_{\mathcal{L}^1(H, H)} \\ &\leq h_G(\cdot) + C_G \|J_\lambda^B(\hat{u}_\lambda)\|_V^p, \end{aligned}$$

convergence (43) and the Vitali convergence theorem yield

$$\operatorname{Tr} [G^*(\cdot, J_\lambda^B(\hat{u}_\lambda)) D_G B_\lambda(\hat{u}_\lambda) G(\cdot, J_\lambda^B(\hat{u}_\lambda))] \rightarrow \operatorname{Tr} [L(\cdot, \hat{u})] \quad \text{in } L^1_{\mathcal{F}}(\hat{\Omega}; L^1(0, T)).$$

Putting all this information together, we are left with

$$\begin{aligned} \limsup_{\lambda \searrow 0} \hat{\mathbb{E}} \int_0^T (A_\lambda(\partial_t \hat{u}_\lambda^d(s)), \partial_t \hat{u}_\lambda^d(s))_H \, ds \\ \leq \widehat{B}(u_0) - \hat{\mathbb{E}} \widehat{B}(\hat{u}(T)) + \hat{\mathbb{E}} \int_0^T (F(s, \hat{u}(s)), \partial_t \hat{u}^d(s))_H \, ds + \frac{1}{2} \hat{\mathbb{E}} \int_0^T \operatorname{Tr} [L(\cdot, \hat{u}(s))] \, ds. \end{aligned} \quad (52)$$

At this point, from equation (51) and the generalized Itô formula in Proposition 3.3 we infer that

$$\begin{aligned} \hat{\mathbb{E}} \widehat{B}(\hat{u}(T)) + \hat{\mathbb{E}} \int_0^T (\hat{v}(s), \partial_t \hat{u}^d(s))_H \, ds \\ = \widehat{B}(u_0) + \hat{\mathbb{E}} \int_0^T (F(s, \hat{u}(s)), \partial_t \hat{u}^d(s))_H \, ds + \frac{1}{2} \hat{\mathbb{E}} \int_0^T \operatorname{Tr} [L(\cdot, \hat{u}(s))] \, ds. \end{aligned} \quad (53)$$

Hence, comparing (52) and (53) we infer that

$$\limsup_{\lambda \searrow 0} \hat{\mathbb{E}} \int_0^T (A_\lambda(\partial_t \hat{u}_\lambda^d(s)), \partial_t \hat{u}_\lambda^d(s))_H \, ds \leq \hat{\mathbb{E}} \int_0^T (\hat{v}(s), \partial_t \hat{u}^d(s))_H \, ds.$$

Together with the weak convergences (40)–(42) and the maximal monotonicity of  $A$ , this ensures indeed that  $\hat{v} \in A(\hat{u})$  almost everywhere in  $\hat{\Omega} \times (0, T)$ , and concludes the proof of Theorem 2.1.

## 5. UNIQUENESS AND EXISTENCE OF PROBABILISTICALLY STRONG SOLUTIONS

This last section is devoted to the proof of Theorem 2.2, showing that existence and uniqueness of probabilistically strong solutions hold under the additional assumptions that  $B : V \rightarrow V^*$  and  $G(t, \cdot) : H \rightarrow \mathcal{L}^2(U, H)$  are linear and continuous, and  $A : H \rightarrow 2^H$  is strongly monotone. Let us point out that since by **B1** the operator  $B : V \rightarrow V^*$  is a subdifferential, we automatically have that  $B$  is self-adjoint, and necessarily  $p = 2$  and

$$\widehat{B}(x) = \frac{1}{2} \langle B(x), x \rangle_{V^*, V}, \quad D_G B(x) = B \in \mathcal{L}(V, V^*), \quad \forall x \in V.$$



Moreover, the strong monotonicity of  $A$  reads

$$(y_1 - y_2, x_1 - x_2)_H \geq c_A \|x_1 - x_2\|_H^2 \quad \forall x_i \in D(A), \quad \forall y_i \in A(x_i), \quad i = 1, 2.$$

First of all, we show uniqueness of martingale solutions. Let  $(\hat{u}_1, \hat{u}_1^d, \hat{v}_1)$  and  $(\hat{u}_2, \hat{u}_2^d, \hat{v}_2)$  be two martingale solutions in the sense of Theorem 2.1 on the same stochastic basis  $(\hat{\Omega}, \hat{\mathcal{F}}, (\hat{\mathcal{F}}_t)_t, \hat{\mathbb{P}}, \hat{W})$ , with respect to some initial data  $u_{0,1}, u_{0,2} \in V$ . Then, using the fact that  $B$  is linear we get

$$\begin{aligned} \hat{u}_1 - \hat{u}_2 &= u_{0,1} - u_{0,2} + \int_0^\cdot \partial_t(\hat{u}_1^d - \hat{u}_2^d)(s) \, ds + \int_0^\cdot G(s, (\hat{u}_1 - \hat{u}_2)(s)) \, d\hat{W}(s), \\ \hat{v}_1 - \hat{v}_2 + B(\hat{u}_1 - \hat{u}_2) &= F(\cdot, \hat{u}_1) - F(\cdot, \hat{u}_2). \end{aligned}$$

The generalized Itô formula in Proposition 3.3 gives then

$$\begin{aligned} &\hat{B}((\hat{u}_1 - \hat{u}_2)(t)) + \int_0^t ((\hat{v}_1 - \hat{v}_2)(s), \partial_t(\hat{u}_1^d - \hat{u}_2^d)(s))_H \, ds \\ &= \hat{B}(u_{0,1} - u_{0,2}) + \int_0^t (F(s, \hat{u}_1(s)) - F(s, \hat{u}_2(s)), \partial_t(\hat{u}_1^d - \hat{u}_2^d)(s))_H \, ds \\ &\quad + \frac{1}{2} \int_0^t \text{Tr}[L(s, (\hat{u}_1 - \hat{u}_2)(s))] \, ds + \int_0^t (B((\hat{u}_1 - \hat{u}_2)(s)), G(s, (\hat{u}_1 - \hat{u}_2)(s)) \, dW(s))_H \end{aligned}$$

for every  $t \in [0, T]$ ,  $\hat{P}$ -almost surely. On the left-hand side, by the coercivity of  $\hat{B}$  (23) we have

$$\hat{B}((\hat{u}_1 - \hat{u}_2)(t)) \geq \frac{c_B}{2} \|(\hat{u}_1 - \hat{u}_2)(t)\|_V^2,$$

while the strong monotonicity of  $A$  yields

$$\int_0^t ((\hat{v}_1 - \hat{v}_2)(s), \partial_t(\hat{u}_1^d - \hat{u}_2^d)(s))_H \, ds \geq c_A \int_0^t \|\partial_t(\hat{u}_1^d - \hat{u}_2^d)(s)\|_H^2 \, ds.$$

On the right-hand side, the first term gives, by assumption **B1–B2**,

$$\hat{B}(u_{0,1} - u_{0,2}) \leq \frac{1}{2} \|B\|_{\mathcal{L}(V, V^*)} \|u_{0,1} - u_{0,2}\|_V^2,$$

while the second and third ones can be handled using the Young inequality and **F–G** as

$$\begin{aligned} &\int_0^t (F(s, \hat{u}_1(s)) - F(s, \hat{u}_2(s)), \partial_t(\hat{u}_1^d - \hat{u}_2^d)(s))_H \, ds + \frac{1}{2} \int_0^t \text{Tr}[L(s, (\hat{u}_1 - \hat{u}_2)(s))] \, ds \\ &\leq \frac{c_A}{2} \int_0^t \|\partial_t(\hat{u}_1^d - \hat{u}_2^d)(s)\|_H^2 \, ds + \left( \frac{C_F}{2c_A} + \frac{C_G}{2} \right) \int_0^t \|(\hat{u}_1 - \hat{u}_2)(s)\|_V^2 \, ds. \end{aligned}$$

Taking expectations we infer then

$$\begin{aligned} &\frac{c_B}{2} \mathbb{E} \|(\hat{u}_1 - \hat{u}_2)(t)\|_V^2 + \frac{c_A}{2} \mathbb{E} \int_0^t \|\partial_t(\hat{u}_1^d - \hat{u}_2^d)(s)\|_H^2 \, ds \\ &\leq \frac{\|B\|_{\mathcal{L}(V, V^*)}}{2} \|u_{0,1} - u_{0,2}\|_V^2 + \left( \frac{C_F}{2c_A} + \frac{C_G}{2} \right) \mathbb{E} \int_0^t \|(\hat{u}_1 - \hat{u}_2)(s)\|_V^2 \, ds \quad \forall t \in [0, T]. \end{aligned}$$

The Gronwall lemma ensures then the required continuous dependence result.

As far as uniqueness is concerned, the continuous dependence property directly implies that if  $u_{0,1} = u_{0,2}$  then

$$\hat{u}_1 = \hat{u}_2 \quad \text{in } L^\infty(0, T; L^2(\hat{\Omega}; V)), \quad \partial_t \hat{u}_1^d = \partial_t \hat{u}_2^d \quad \text{in } L^2_{\hat{\mathcal{F}}}(\hat{\Omega}; L^2(0, T; H)).$$

The second equality yields straightaway that

$$\hat{u}_1^d = \hat{u}_2^d \quad \text{in } L^2_{\mathcal{F}}(\hat{\Omega}; C^0([0, T]; H)),$$

while the first equality and assumption **G** implies

$$G(\cdot, \hat{u}_1) = G(\cdot, \hat{u}_2) \quad \text{in } L^2_{\mathcal{F}}(\hat{\Omega}; L^2(0, T; \mathcal{L}^2(U, H))),$$

from which also

$$\int_0^\cdot G(s, \hat{u}_1(s)) d\hat{W}(s) = \int_0^\cdot G(s, \hat{u}_2(s)) d\hat{W}(s) \quad \text{in } L^2_{\mathcal{F}}(\hat{\Omega}; C^0([0, T]; H)).$$

Consequently, we deduce that

$$\hat{u}_1 = \hat{u}_2 \quad \text{in } L^2_{\mathcal{F}}(\hat{\Omega}; C^0([0, T]; H)),$$

and pathwise uniqueness of martingale solutions holds.

In order to conclude the proof of Theorem 2.2, we are only left to show existence of probabilistically strong solutions. This can be done in a classical way, using existence and uniqueness of martingale solutions. In particular, we recall the following lemma due to GYÖNGY & KRYLOV [20, Lem. 1.1].

**Lemma 5.1.** *Let  $\mathcal{X}$  be a Polish space and  $(Z_n)_n$  be a sequence of  $\mathcal{X}$ -valued random variables. Then  $(Z_n)_n$  converges in probability if and only if for any pair of subsequences  $(Z_{n_k})_k$  and  $(Z_{n_j})_j$ , there exists a joint sub-subsequence  $(Z_{n_{k_\ell}}, Z_{n_{j_\ell}})_{\ell}$  converging in law to a probability measure  $\nu$  on  $\mathcal{X} \times \mathcal{X}$  such that  $\nu(\{(z_1, z_2) \in \mathcal{X} \times \mathcal{X} : z_1 = z_2\}) = 1$ .*

Under the additional assumptions of Theorem 2.2, recalling the proof of Theorem 2.1, it is readily seen that the Skorokhod Theorem and the just proved pathwise uniqueness of martingale solutions to the limit problem ensure exactly the condition of the lemma above. It follows then that all the arguments for passing to the limit as  $\lambda \searrow 0$  in Subsections 4.3–4.4 can be replicated on the original probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ . This ensures then existence of probabilistically strong solutions, and Theorem 2.2 is proved.

## 6. APPLICATION

We conclude our discussion by presenting a concrete case of an SPDE to which the abstract theory applies. Let us consider the initial and boundary value problem

$$\begin{aligned} \alpha(\partial_t u^d) dt + u^s dW - \operatorname{div}(D\hat{\beta}_1(\nabla u)) dt + \hat{\beta}'_0(u) dt &\ni f(\cdot, u, \nabla u) dt + G(\cdot, u) dW, \\ &\text{in } \Omega \times (0, T) \times \mathcal{O}, \end{aligned} \tag{54}$$

$$u = 0 \quad \text{on } \Omega \times (0, T) \times \partial\mathcal{O}, \tag{55}$$

$$u(0, \cdot) = u_0 \quad \text{on } \Omega \times \mathcal{O}. \tag{56}$$

Here, the solution is a scalar function  $u : \Omega \times (0, T) \times \mathcal{O} \rightarrow \mathbb{R}$  where  $\mathcal{O} \subset \mathbb{R}^d$  is an open, bounded, and Lipschitz domain. The possibly multivalued map  $\alpha : \mathbb{R} \rightarrow 2^{\mathbb{R}}$  is assumed to be strongly maximal monotone and linearly bounded with  $0 \in \alpha(0)$ . The potential  $\hat{\beta}_1 \in C^2(\mathbb{R}^d)$  is asked to be nonnegative, convex, minimized at 0, and with polynomial growth. More precisely, we assume to be given  $p \geq 2$  and  $c_\beta > 0$  such that

$$|D^2\hat{\beta}_1(\xi)| \leq \frac{1}{c_\beta}(1 + |\xi|^{p-2}), \quad (D\hat{\beta}_1(\xi) - D\hat{\beta}_1(\eta)) \cdot (\xi - \eta) \geq c_\beta|\xi - \eta|^p \quad \forall \xi, \eta \in \mathbb{R}^d.$$

The potential  $\widehat{\beta}_0 \in C^2(\mathbb{R})$  is assumed to be nonnegative, convex, minimized at 0, with  $\widehat{\beta}_0''$  bounded by a polynomial of order  $p-2$ . We assume  $f : (0, T) \times \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{R}$  to be Carathéodory and uniformly Lipschitz in the third variable with  $t \mapsto f(t, 0, 0) \in L^1(0, T)$ ,  $G$  to fulfil Assumption **G** with

$$H = L^2(\mathcal{O}), \quad V = W_0^{1,p}(\mathcal{O})$$

and the initial datum  $u_0$  to belong to  $W_0^{1,p}(\mathcal{O})$ . Note that the homogenous Dirichlet boundary conditions are chosen here for the sake of definiteness, other choices also being possible. Note assumption **G** allows for most of the classical choices of the noise coefficient. These include, but are not limited to, the natural case of additive noise  $G \in \mathcal{L}^2(U, V)$  as well as the case of multiplicative noise of superposition type. An explicit form for  $G$  could be

$$G(u) dW = \sum_{j=1}^{\infty} h_j(u) dW_j$$

where  $W_j$  are independent real Brownian motions and  $h_j : H \rightarrow H$  are  $L_j$ -Lipschitz continuous functions with  $\sum_j L_j^2 < \infty$  and  $h_j(V) \subset V$ . As for the map  $h_j$ , in addition to *local* functions one could consider some *nonlocal* operators as well, possibly defined via convolutions.

We introduce a variational formulation of problem (54) in  $(V, H, V^*)$  by letting the operators  $A : H \rightarrow 2^H$  and  $F : (0, T) \times V \rightarrow H$  and the functional  $\widehat{B} : V \rightarrow (-\infty, +\infty]$  be defined as

$$\begin{aligned} \xi \in A(w) &\Leftrightarrow \xi \in \alpha(w) \text{ a.e. in } \mathcal{O}, & \xi, w \in H, \\ F(t, u) &= f(t, u, \nabla u) \text{ a.e. in } \mathcal{O}, & u \in V, \\ \widehat{B}(u) &:= \int_{\mathcal{O}} \left( \widehat{\beta}_1(\nabla u(x)) + \widehat{\beta}_0(u(x)) \right) dx, & u \in V. \end{aligned}$$

The assumptions on  $\alpha$  entail that  $A$  is strongly maximal monotone and linearly bounded with  $0 \in A(0)$ , so that Assumption **A** holds. As  $\widehat{\beta}_1$  and  $\widehat{\beta}_0$  are smooth and  $\widehat{\beta}_1$  has  $p$  growth Assumptions **B1–B2**, follow as well. The uniform Lipschitz continuity of  $f$  entails Assumption **F** with  $h_F(\cdot) = f(\cdot, 0, 0) \in L^1(0, T)$ . Under the above provisions, the abstract relation (4), complemented by the initial condition, is the variational formulation of (54) in  $(V, H, V^*)$ . As the assumptions of Subsection 2.3 are satisfied, Theorem 2.1 entails the existence of solutions. In particular, we have the following.

**Corollary 6.1.** *Under the assumptions of this section, problem (54)–(56) admits an analytically strong martingale solution in the sense of Theorem 2.1.*

#### ACKNOWLEDGEMENT

LS is funded by the Austrian Science Fund (FWF) through the Lise-Meitner project M 2876. US is partially supported by the Austrian Science Fund (FWF) through projects F 65, I 4354, P 32788, and by the Vienna Science and Technology Fund through project MA14-009.

#### REFERENCES

- [1] S. Aizicovici and Q. Yan. Convergence theorems for abstract doubly nonlinear differential equations. *Panamer. Math. J.* 7:1–17, 1997.
- [2] G. Akagi. Doubly nonlinear evolution equations with non-monotone perturbations in reflexive Banach spaces. *J. Evol. Equ.*, 11(1):1–41, 2011.
- [3] G. Akagi. Global attractors for doubly nonlinear evolution equations with non-monotone perturbations, *J. Differential Equations*, 250:1850–1875, 2011.

- [4] G. Akagi and U. Stefanelli. Weighted energy-dissipation functionals for doubly nonlinear evolution, *J. Funct. Anal.*, 260:2541–2578, 2011.
- [5] G. Akagi and U. Stefanelli. A variational principle for doubly nonlinear evolution. *Appl. Math. Lett.*, 23:1120–1124, 2010.
- [6] L. Ambrosio, N. Gigli, and G. Savaré. *Gradient flows in metric spaces and in the space of probability measures*, Birkhäuser Verlag, Basel, second edition, 2008.
- [7] T. Arai. On the existence of the solution for  $\partial\varphi(u \langle t, \cdot \rangle t) + \partial\psi(u(t)) \ni f(t)$ . *J. Fac. Sci. Univ. Tokyo Sect. IA Math.*, 26(1):75–96, 1979.
- [8] V. Barbu. Existence theorems for a class of two point boundary problems. *J. Differential Equations*, 17:236–257, 1975.
- [9] V. Barbu, S. Bonaccorsi, and L. Tubaro. Existence and asymptotic behavior for hereditary stochastic evolution equations. *Appl. Math. Optim.*, 69:273–314, 2014.
- [10] V. Barbu and G. Da Prato. The two phase stochastic Stefan problem, *Probab. Theory Related Fields*. 124:544–560, 2002.
- [11] V. Barbu, G. Da Prato, and M. Röckner. Existence of strong solutions for stochastic porous media equation under general monotonicity conditions. *Ann. Probab.*, 37:428–452, 2009.
- [12] M. A. Biot. Variational principles in irreversible thermodynamics with application to viscoelasticity. *Phys. Rev. (2)*, 97:1463–1469, 1955.
- [13] P. Colli. On some doubly nonlinear evolution equations in Banach spaces. *Japan J. Indust. Appl. Math.*, 9(2):181–203, 1992.
- [14] P. Colli and A. Visintin. On a class of doubly nonlinear evolution equations. *Comm. Partial Differential Equations*, 15(5):737–756, 1990.
- [15] G. Da Prato and J. Zabczyk. *Stochastic equations in infinite dimensions*, volume 152 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, second edition, 2014.
- [16] R. E. Edwards. *Functional analysis. Theory and applications*. Holt, Rinehart and Winston, New York-Toronto-London, 1965.
- [17] F. Flandoli and D. Gatarek. Martingale and stationary solutions for stochastic Navier-Stokes equations. *Probab. Theory Related Fields*, 102(3):367–391, 1995.
- [18] P. Germain. *Cours de mécanique des milieux continus*. Masson et Cie, Éditeurs, Paris, 1973. Tome I: Théorie générale.
- [19] B. Gess. Strong solutions for stochastic partial differential equations of gradient type. *J. Funct. Anal.*, 263:2355–2383, 2012.
- [20] I. Gyöngy and N. Krylov. Existence of strong solutions for Itô’s stochastic equations via approximations. *Probab. Theory Related Fields*, 105(2):143–158, 1996.
- [21] N. Ikeda and S. Watanabe. *Stochastic differential equations and diffusion processes*, volume 24 of *North-Holland Mathematical Library*. North-Holland Publishing Co., Amsterdam; Kodansha, Ltd., Tokyo, second edition, 1989.
- [22] M. Keller-Ressel and M. S. Müller. A Stefan-type stochastic moving boundary problem. *Stoch. Partial Differ. Equ. Anal. Comput.*, 4:746–790, 2016.
- [23] N. V. Krylov and B. L. Rozovskiĭ. Stochastic evolution equations, In: *Current Problems in Mathematics*, vol. 14, Akad. Nauk SSSR, Vsesoyuz. Inst. Nauchn. i Tekhn. Informatsii, Moscow, 1979, pp. 71–147, 256 (in Russian).
- [24] W. Liu and M. Röckner. *Stochastic partial differential equations: an introduction*. Springer, Cham, 2015.
- [25] C. Marinelli and L. Scarpa. Refined existence and regularity results for a class of semilinear dissipative SPDEs, *Infin. Dimens. Anal. Quantum Probab. Relat. Top.*, to appear, 2020.
- [26] C. Marinelli and L. Scarpa. Strong solutions to SPDEs with monotone drift in divergence form. *Stoch. Partial Differ. Equ. Anal. Comput.*, 6:364–396, 2018.
- [27] C. Marinelli and L. Scarpa. A variational approach to dissipative SPDEs with singular drift. *Ann. Probab.*, 46:1455–1497, 2018.
- [28] C. Marinelli and L. Scarpa. Ergodicity and Kolmogorov equations for dissipative spdes with singular drift: a variational approach. *Potential Anal.*, 52(1):69–103, 2020.
- [29] C. Marinelli and L. Scarpa. A note on doubly nonlinear SPDEs with singular drift in divergence form. *Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl.*, 29:619–633, 2018.
- [30] J.-J. Moreau. Sur les lois de frottement, de viscosité et plasticité. *C. R. Acad. Sci. Paris Sér. II Méc. Phys. Chim. Sci. Univers Sci. Terre*, 271:608–611, 1970.
- [31] J.-J. Moreau. Sur l’évolution d’un système élasto-visco-plastique. *C. R. Acad. Sci. Paris Sér. A-B*, 273:A118–A121, 1971.

- [32] A. Mielke and T. Roubíček. *Rate-Independent Systems - Theory and Application*, Appl. Math. Sci. Series, vol. 193. Springer, New York, 2015.
- [33] C. Orrieri and L. Scarpa. Singular stochastic Allen-Cahn equations with dynamic boundary conditions. *J. Differential Equations*, 266:4624–4667, 2019.
- [34] E. Pardoux. *Equations aux dérivées partielles stochastiques nonlinéaires monotones*. PhD thesis, Université Paris XI, 1975.
- [35] É. Pardoux. Sur des équations aux dérivées partielles stochastiques monotones. *C. R. Acad. Sci. Paris Sér. A-B*, 275:A101–A103, 1972.
- [36] C. Prévôt and M. Röckner. *A concise course on stochastic partial differential equations*. Lecture Notes in Mathematics, 1905. Springer, Berlin, 2007.
- [37] T. Roubíček. *Nonlinear partial differential equations with applications*. International Series of Numerical Mathematics, Vol. 153. Birkhäuser Verlag, Basel, 2005.
- [38] N. Sapountzoglou, P. Wittbold, and A. Zimmermann. On a doubly nonlinear PDE with stochastic perturbation. *Stoch. Partial Differ. Equ. Anal. Comput.*, 7(2):297–330, 2019.
- [39] L. Scarpa. Well-posedness for a class of doubly nonlinear stochastic PDEs of divergence type. *J. Differential Equations*, 263:2113–2156, 2017.
- [40] L. Scarpa. On the stochastic Cahn-Hilliard equation with a singular double-well potential. *Nonlinear Anal.*, 171:102–133, 2018.
- [41] L. Scarpa and U. Stefanelli. Doubly nonlinear stochastic evolution equations. *Math. Models Methods Appl. Sci.*, 30(5):991–1031, 2020.
- [42] L. Scarpa and U. Stefanelli. Stochastic PDEs via convex minimization. [arXiv:2004.00337](https://arxiv.org/abs/2004.00337), 2020.
- [43] G. Schimperna, A. Segatti, and U. Stefanelli. Well-posedness and long-time behavior for a class of doubly nonlinear equations, *Discrete Contin. Dyn. Syst.*, 18:15–38, 2007.
- [44] A. Segatti. Global attractor for a class of doubly nonlinear abstract evolution equations. *Discrete Contin. Dyn. Syst.*, 14:801–820, 2006.
- [45] T. Senba. On some nonlinear evolution equation. *Funkcial. Ekv.*, 29:243–257, 1986.
- [46] J. Simon. Compact sets in the space  $L^p(0, T; B)$ . *Ann. Mat. Pura Appl. (4)*, 146:65–96, 1987.
- [47] U. Stefanelli. The Brezis-Ekeland principle for doubly nonlinear equations. *SIAM J. Control Optim.*, 47:1615–1642, 2008.
- [48] G. Vallet and A. Zimmermann. Well-posedness for a pseudomonotone evolution problem with multiplicative noise. *J. Evol. Equ.*, 19(1):153–202, 2019.

(Luca Scarpa) FACULTY OF MATHEMATICS, UNIVERSITY OF VIENNA, OSKAR-MORGENSTERN-PLATZ 1, 1090 VIENNA, AUSTRIA.

*E-mail address:* [luca.scarpa@univie.ac.at](mailto:luca.scarpa@univie.ac.at)

*URL:* <http://www.mat.univie.ac.at/~scarpa>

(Ulisse Stefanelli) FACULTY OF MATHEMATICS, UNIVERSITY OF VIENNA, OSKAR-MORGENSTERN-PLATZ 1, A-1090 VIENNA, AUSTRIA, VIENNA RESEARCH PLATFORM ON ACCELERATING PHOTOREACTION DISCOVERY, UNIVERSITY OF VIENNA, WÄHRINGER STR. 17, 1090 VIENNA, AUSTRIA, & ISTITUTO DI MATEMATICA APPLICATA E TECNOLOGIE INFORMATICHE “E. MAGENES” - CNR, v. FERRATA 1, I-27100 PAVIA, ITALY

*E-mail address:* [ulisse.stefanelli@univie.ac.at](mailto:ulisse.stefanelli@univie.ac.at)

*URL:* <http://www.mat.univie.ac.at/~stefanelli>