

The weak Stokes problem associated with a flow through a profile cascade in L^r –framework

Tomáš Neustupa

Abstract

We study the weak steady Stokes problem, associated with a flow of a Newtonian incompressible fluid through a spatially periodic profile cascade, in the L^r –setup. The used mathematical model is based on the reduction to one spatial period, represented by a bounded 2D domain Ω . The corresponding Stokes problem is formulated by means of three types of boundary conditions: the conditions of periodicity on the “lower” and “upper” parts of the boundary, the Dirichlet boundary conditions on the “inflow” and on the profile and an artificial “do nothing”–type boundary condition on the “outflow”. Under appropriate assumptions on the given data, we prove the existence and uniqueness of a weak solution in $\mathbf{W}^{1,r}(\Omega)$ and its continuous dependence on the data. We explain the sense in which the “do nothing” boundary condition on the “outflow” is satisfied.

AMS math. classification (2000): 35Q30, 76D03, 76D05.

Keywords: The Stokes problem, weak solution, artificial boundary condition.

1 Introduction

One spatial period: domain Ω . Mathematical models of a flow through a three-dimensional turbine wheel often use the reduction to two space dimensions, where the flow is studied as a flow through an infinite planar profile cascade. In an appropriately chosen Cartesian coordinate system, the profiles in the cascade periodically repeat with the period τ in the x_2 –direction. It can be naturally assumed that the flow is τ –periodic in variable x_2 , too. This enables one to study the flow through one spatial period, which contains just one profile – see domain Ω and profile P on Fig. 1. This approach is used e.g. in [7] and [17], where the authors present the numerical analysis of the models or corresponding numerical simulations, and in the papers, [8]–[10] and [26]–[29], devoted to theoretical analysis of the mathematical models.

We assume that the fluid flows into the cascade through the straight line γ_{in} (the x_2 –axis, the inflow) and essentially leaves the cascade through the straight line γ_{out} , whose equation is $x_1 = d$ (the outflow). By “essentially” we mean that we do not exclude possible




Fig. 1: Domain Ω

reverse flows on γ_{out} . The considered spatial period Ω is mainly determined by artificially chosen curves Γ_0 and $\Gamma_1 \equiv \Gamma_0 + \tau \mathbf{e}_2$, which form the “lower” and “upper” parts of $\partial\Omega$ (the boundary of Ω), respectively. (See Fig. 1. We denote by \mathbf{e}_2 the unit vector in the x_2 -direction.) The parts of $\partial\Omega$, lying on the straight lines γ_{in} and γ_{out} , are the line segments $\Gamma_{\text{in}} \equiv A_0A_1$ and $\Gamma_{\text{out}} \equiv B_0B_1$, respectively, of length τ . The last part of $\partial\Omega$, i.e. the boundary of profile the P is denoted by Γ_p . We assume that the domain Ω is Lipschitzian and the curves Γ_0 and Γ_1 are of the class C^∞ .

The Stokes boundary–value problem on one spatial period. The flow of an incompressible Newtonian fluid is described by the Navier-Stokes equations. An important role in theoretical studies of these equations play the properties of solutions of the steady Stokes problem. If one neglects the derivative with respect to time and the nonlinear “convective” term in the momentum equation in the Navier–Stokes system, one obtains

$$-\nu \Delta \mathbf{u} + \nabla p = \mathbf{f}. \quad (1.1)$$

This equation is studied together with the equation of continuity (= condition of incompressibility)

$$\operatorname{div} \mathbf{u} = 0 \quad (1.2)$$

and the equations (1.1), (1.2) represent the so called *steady Stokes system*, or the *steady Stokes equations*. The unknowns are $\mathbf{u} = (u_1, u_2)$ (the velocity) and p (the pressure). The positive constant ν is the kinematic coefficient of viscosity and \mathbf{f} denotes the external body force. The density of the fluid can be without loss of generality supposed to be equal to one. The system (1.1), (1.2) is completed by appropriate boundary conditions on $\partial\Omega$. One can naturally assume that the velocity profile on Γ_{in} is known, which leads to the inhomogeneous Dirichlet boundary condition

$$\mathbf{u} = \mathbf{g} \quad \text{on } \Gamma_{\text{in}}. \quad (1.3)$$

Further, we consider the homogeneous Dirichlet boundary condition

$$\mathbf{u} = \mathbf{0} \quad \text{on } \Gamma_p, \quad (1.4)$$

the condition of periodicity on Γ_0 and Γ_1

$$\mathbf{u}(x_1, x_2 + \tau) = \mathbf{u}(x_1, x_2) \quad \text{for } \mathbf{x} \equiv (x_1, x_2) \in \Gamma_0 \quad (1.5)$$

and the artificial boundary condition

$$-\nu \frac{\partial \mathbf{u}}{\partial \mathbf{n}} + p \mathbf{n} = \mathbf{h} \quad \text{on } \Gamma_{\text{out}}, \quad (1.6)$$

where \mathbf{h} is a given vector–function on Γ_{out} and \mathbf{n} denotes the unit outer normal vector, which is equal to $\mathbf{e}_1 \equiv (1, 0)$ on Γ_{out} . The boundary condition (1.6) (with $\mathbf{h} = \mathbf{0}$) is often called the “do nothing” condition, because it naturally follows from an appropriate weak formulation of the boundary–value problem, see [13] and [15].

On the results of this paper. The main results of the paper are theorems on properties of so called weak Stokes operator (Theorem 1) and on the existence, uniqueness and continuous dependence on the data of a weak solution \mathbf{u} to the Stokes problem (1.1)–(1.6) in the L^r -setting (Theorem 3). The weak solution is defined in Definition 1. Theorem 2 provides the existence of an associated pressure p and explains the sense, in which \mathbf{u} and p satisfy the boundary condition (1.6). These results do not follow from the previous cited papers on the

Stokes problem, mainly because we consider three different types of boundary conditions on $\partial\Omega$, two of which “meet” in the “corner points” A_0, A_1, B_0 and B_1 of domain Ω . Moreover, while the corresponding L^2 -theory is relatively simple (see [29]), the general L^r -case is much more difficult. The key inequality (3.4) is proven in Section 5. The crucial estimate of the $W^{1,r}$ -norm of the velocity in the neighborhood of Γ_{out} is obtained, applying results of S. Agmon, A. Douglis and L. Nirenberg [1] on general elliptic systems. As an auxiliary result, we present Lemma 3 on an appropriate extension of the velocity profile \mathbf{g} from Γ_{in} to Ω .

On some previous related results. The Stokes problem, in various domains and with various boundary conditions, has already been studied in many papers and books. As to weak solutions, the L^2 -existential theory and the proof of uniqueness of an existing weak solution is relatively simple, because one works in a Hilbert space and can apply the Riesz theorem. However, the corresponding L^r -theory for a general $r \in (1, \infty)$ is much more difficult. Nevertheless, also in the L^r -setup, results on the existence and uniqueness of weak solutions of the Stokes problem with Dirichlet’s boundary condition for the velocity can be found e.g. in [6], [30] and [11], with Navier’s boundary condition in [2] and with the Navier-type boundary condition in [3]. Fundamental estimates have been basically obtained by means of the Stokes fundamental solution and the corresponding Green tensor in [6] (in 3D) and [11], respectively also the inf-sup condition in [2] and [3]. The 2D case (with Dirichlet’s boundary condition and in a smooth domain) has also been solved in [30] by expressing the velocity by means of a stream function, application of operator ∇^\perp to the Stokes equation and the results on the biharmonic boundary-value problem.

In studies of the Navier–Stokes equations in channels or profile cascades with artificial boundary conditions on the outflow, many authors use various modifications of condition (1.6). (See e.g. [5]), [8]–[10] and [26]–[28].) The reason is that condition (1.6) does not enable one to control the amount of kinetic energy in Ω in the case of a reverse flow on Γ_{out} . Hence modifications are suggested so that one can derive an energy inequality, and consequently prove the existence of weak solutions. In papers [21] and [22], the authors use the boundary condition on the outflow in connection with a flow in a channel, and they prove the existence of weak solutions of the Navier–Stokes equations for “small data”. Possible reverse flows on the “outflow” of a channel are controlled by means of additional conditions in [18], [19], [20], where the Navier–Stokes equations are replaced by the Navier–Stokes variational inequalities.

The regularity up to the boundary of existing weak solutions (stationary or time-dependent) to the Navier–Stokes equations with the boundary condition (1.6) on a part of the boundary has not been studied in literature yet. This is mainly because one at first needs a deeper information on existence, uniqueness and regularity in the L^r -framework for the corresponding steady Stokes problem. There are, to our knowledge, only two papers which bring an information on regularity of a solution of this Stokes problem: 1) paper [23], where the authors studied a flow in a 2D channel D of a special geometry, considering the homogeneous Dirichlet boundary condition on the walls and condition (1.6) on the outflow, and proved that the velocity is in $\mathbf{W}^{2-\beta,2}(D)$ for certain $\beta \in (0, 1)$, provided that $\mathbf{f} \in \mathbf{L}^2(D)$ (see [23, Theorem 2.1]), and 2) paper [29], where the belonging of the solution (\mathbf{u}, p) of the Stokes problem (1.1)–(1.6) to $\mathbf{W}^{2,2}(\Omega) \times W^{1,2}(\Omega)$ has been recently proven, under natural assumptions on \mathbf{f} , \mathbf{g} and \mathbf{h} .

2 Notation and some preliminary results

Notation. Recall that Ω is a Lipschitzian domain in \mathbb{R}^2 , sketched on Fig. 1. Its boundary consists of the curves Γ_{in} , Γ_{out} , Γ_0 , Γ_1 and Γ_p , described in Section 1. We denote by $\mathbf{n} = (n_1, n_2)$ the outer normal vector field on $\partial\Omega$. We use c as a generic constant, i.e. a constant whose values may change throughout the text.

- Γ_{in}^0 , respectively Γ_{out}^0 , denotes the open line segment without the end points A_0, A_1 , respectively B_0, B_1 . Similarly, Γ_0^0 , respectively Γ_1^0 denotes the curve Γ_0 , respectively Γ_1 , without the end points A_0, B_0 , respectively A_1, B_1 .
- We denote vector functions and spaces of vector functions by boldface letters. If $\mathbf{v} \equiv (v_1, v_2)$ is a vector function then $\nabla \mathbf{v}$ is a tensor function with the entries $\partial v_i / \partial x_j \equiv \partial_j v_i$ on the positions ij (for $i, j = 1, 2$). Spaces of 2nd-order tensor functions are denoted by the superscript 2×2 .
- We denote by $\|\cdot\|_r$ the norm in $L^r(\Omega)$ or in $\mathbf{L}^r(\Omega)$ or in $L^r(\Omega)^{2 \times 2}$. Similarly, $\|\cdot\|_{s,r}$ is the norm in $W^{s,r}(\Omega)$ or in $\mathbf{W}^{s,r}(\Omega)$ or in $W^{s,r}(\Omega)^{2 \times 2}$.
- $\mathcal{C}_\sigma^\infty(\Omega)$ denotes the linear space of all infinitely differentiable divergence-free vector functions in $\bar{\Omega}$, whose support is disjoint with $\Gamma_{\text{in}} \cup \Gamma_p$ and that satisfy, together with all their derivatives (of all orders), the condition of periodicity (1.5). Note that each $\mathbf{w} \in \mathcal{C}_\sigma^\infty(\Omega)$ automatically satisfies the outflow condition $\int_{\Gamma_{\text{out}}} \mathbf{w} \cdot \mathbf{n} \, dl = 0$.
- We denote by $\mathcal{C}_{0,\sigma}^\infty(\Omega)$ the intersection $\mathcal{C}_\sigma^\infty(\Omega) \cap \mathcal{C}_0^\infty(\Omega)$, where $\mathcal{C}_0^\infty(\Omega)$ is the space of all infinitely differentiable vector functions in Ω with a compact support in Ω .
- $\mathbf{V}_\sigma^{1,r}(\Omega)$ (for $r > 1$) is the closure of $\mathcal{C}_\sigma^\infty(\Omega)$ in $\mathbf{W}^{1,r}(\Omega)$. The space $\mathbf{V}_\sigma^{1,r}(\Omega)$ can be characterized as a space of divergence-free vector functions from $\mathbf{W}^{1,r}(\Omega)$, whose traces on $\Gamma_{\text{in}} \cup \Gamma_p$ are equal to zero and the traces on Γ_0 and Γ_1 satisfy the condition of periodicity (1.5). Note that as functions from $\mathbf{V}_\sigma^{1,r}(\Omega)$ are equal to zero on $\Gamma_{\text{in}} \cup \Gamma_p$ (in the sense of traces) and the domain Ω is bounded, the norm in $\mathbf{V}_\sigma^{1,r}(\Omega)$ is equivalent to $\|\nabla \cdot\|_r$.
- We denote by r' the conjugate exponent to r , by $\mathbf{W}^{-1,r}(\Omega)$ the dual space to $\mathbf{W}_0^{1,r'}(\Omega)$ and by $\mathbf{W}_0^{-1,r}(\Omega)$ the dual space to $\mathbf{W}^{1,r'}(\Omega)$. The corresponding norms are denoted by $\|\cdot\|_{\mathbf{W}^{-1,r}}$ and $\|\cdot\|_{\mathbf{W}_0^{-1,r}}$, respectively.
- $\mathbf{V}_\sigma^{-1,r}(\Omega)$ denotes the dual space to $\mathbf{V}_\sigma^{1,r'}(\Omega)$. The duality pairing between $\mathbf{V}_\sigma^{-1,r}(\Omega)$ and $\mathbf{V}_\sigma^{1,r'}(\Omega)$ is denoted by $\langle \cdot, \cdot \rangle_{(\mathbf{V}_\sigma^{-1,r}, \mathbf{V}_\sigma^{1,r'})}$. The norm in $\mathbf{V}_\sigma^{-1,r}(\Omega)$ is denoted by $\|\cdot\|_{\mathbf{V}_\sigma^{-1,r}}$.
- Denote by \mathcal{A}_r the linear mapping of $\mathbf{V}_\sigma^{1,r}(\Omega)$ to $\mathbf{V}_\sigma^{-1,r}(\Omega)$, defined by the equation

$$\langle \mathcal{A}_r \mathbf{v}, \mathbf{w} \rangle_{(\mathbf{V}_\sigma^{-1,r}, \mathbf{V}_\sigma^{1,r'})} := (\nabla \mathbf{v}, \nabla \mathbf{w}) \quad \text{for } \mathbf{v} \in \mathbf{V}_\sigma^{1,r}(\Omega) \text{ and } \mathbf{w} \in \mathbf{V}_\sigma^{1,r'}(\Omega), \quad (2.1)$$

where $(\nabla \mathbf{v}, \nabla \mathbf{w})$ represents the integral $\int_\Omega \nabla \mathbf{v} : \nabla \mathbf{w} \, dx$. We call operator \mathcal{A}_r the *weak Stokes operator*.

Lemma 1. Let $\mathbf{f} \in \mathbf{W}_0^{-1,r}(\Omega)$. Then there exists $\mathbb{F} \in L^r(\Omega)^{2 \times 2}$, satisfying $\text{div } \mathbb{F} = \mathbf{f}$ in the sense of distributions in Ω and

$$\|\mathbb{F}\|_r \leq c \|\mathbf{f}\|_{\mathbf{W}_0^{-1,r}}, \quad (2.2)$$

where c is independent of \mathbf{f} and \mathbb{F} .

Proof. The proof is based on the results from paper [12] by M. Geussert, H. Heck and M. Hieber. If domain $\Omega' \subset \mathbb{R}^3$ is bounded and star-shaped with respect to some ball $K \subset \Omega'$ and ω is a function in $C_0^\infty(K)$, such that $\int_K \omega \, d\mathbf{x} = 1$ then it follows from Proposition 2.1 in [12] that there is a bounded linear operator $\mathfrak{B} : W_0^{-1,r}(\Omega') \rightarrow \mathbf{L}^r(\Omega')$, such that $\operatorname{div} \mathfrak{B}f = f - \omega \int_{\Omega'} f \, d\mathbf{x}$ for $f \in L^r(\Omega')$. Applying an appropriate limit procedure, one can show that if $f \in W_0^{-1,r}(\Omega')$ then

$$\operatorname{div} \mathfrak{B}f = f - \omega \langle f, 1 \rangle_{(W_0^{-1,r}(\Omega'), W^{1,r'}(\Omega'))}$$

in the sense of distributions. Then $\operatorname{div} [\mathfrak{B}f + \mathbf{z}] = f - \omega \langle f, 1 \rangle_{(W_0^{-1,r}(\Omega'), W^{1,r'}(\Omega'))} + \operatorname{div} \mathbf{z}$ for $\mathbf{z} \in \mathbf{L}^r(\Omega')$. Let us choose \mathbf{z} so that $\mathbf{z} = \mathbf{z}_1 + \mathbf{z}_2$, where

$$\operatorname{div} \mathbf{z}_1 = (\omega - \bar{\omega}) \langle f, 1 \rangle_{(W_0^{-1,r}(\Omega'), W^{1,r'}(\Omega'))}, \quad (2.3)$$

and $\mathbf{z}_2 = \frac{1}{3} \mathbf{x} \bar{\omega} \langle f, 1 \rangle_{(W_0^{-1,r}(\Omega'), W^{1,r'}(\Omega'))}$. (We denote by $\bar{\omega}$ the mean value of ω in Ω' .) As the mean value of the right hand side of (2.3) in Ω' is zero, the equation (2.3) is solvable in $W_0^{1,r}(\Omega')$ due to [12, Theorem 2.5]. Moreover,

$$\|\mathbf{z}_1\|_{1,r;\Omega'} \leq c \|\omega - \bar{\omega}\|_{r;\Omega'} |\langle f, 1 \rangle_{(W_0^{-1,r}(\Omega'), W^{1,r'}(\Omega'))}| \leq c \|f\|_{W_0^{-1,r}(\Omega')},$$

where c depends only on Ω' , ω and r . Furthermore,

$$\operatorname{div} \mathbf{z}_2 = \bar{\omega} \langle f, 1 \rangle_{(W_0^{-1,r}(\Omega'), W^{1,r'}(\Omega'))} \quad \text{and} \quad \|\mathbf{z}_2\|_{1,r;\Omega'} \leq c \|f\|_{W_0^{-1,r}(\Omega')},$$

Thus, the function $\mathfrak{B}f + \mathbf{z}$ satisfies the equation $\operatorname{div} [\mathfrak{B}f + \mathbf{z}] = f$ in the sense of distributions in Ω' and $\|\mathfrak{B}f + \mathbf{z}\|_{r;\Omega'} \leq c \|f\|_{W_0^{-1,r}(\Omega')}$.

Domain Ω can be expressed as a finite union of star-shaped domains. This enables us to carry over these results to the whole domain Ω , applying the same arguments as in [12], pp. 116–117. Thus, we can formulate the proposition: *there exists a bounded linear operator $\mathfrak{B} : W_0^{-1,r}(\Omega) \rightarrow \mathbf{L}^r(\Omega)$, such that $\operatorname{div} \mathfrak{B}f = f$ in the sense of distributions in Ω for $f \in W_0^{-1,r}(\Omega)$.* Now, it is just a technical step to extend this proposition from $f \in W_0^{-1,r}(\Omega)$ to $\mathbf{f} \in \mathbf{W}_0^{-1,r}(\Omega)$. This completes the proof. \square

Remark 1. As there is not a complete coincidence on the definition of the divergence of a tensor field in literature, note that if $\mathbb{F} = (F_{ij})$ ($i, j = 1, 2$) then $\operatorname{div} \mathbb{F}$ in Lemma 1 (and also further on throughout the paper) denotes the vector $\partial_j F_{ij}$ ($i = 1, 2$). In accordance with this notation, $\operatorname{div} \nabla \mathbf{v}$ is the vector with the entries $\partial_j (\partial_j v_i) = \Delta v_i$ ($i = 1, 2$).

One can also assume that $\mathbf{f} \in \mathbf{W}^{-1,r}(\Omega)$ (instead of $\mathbf{f} \in \mathbf{W}_0^{-1,r}(\Omega)$) in Lemma 1. However, in this case, one cannot apply [12] in order to obtain the tensorial function $\mathbb{F} \in L^r(\Omega)^{2 \times 2}$ with the properties stated in the lemma. Nevertheless, the existence of \mathbb{F} , satisfying the equation $\operatorname{div} \mathbb{F} = \mathbf{f}$ (in the sense of distributions) and the estimate $\|\mathbb{F}\|_r \leq c \|\mathbf{f}\|_{\mathbf{W}^{-1,r}}$ can be proven in this case, too, just appropriately modifying the proof of Lemma II.1.6.1 in [31], which concerns the case $r = 2$.

3 The weak Stokes operator

The next theorem is an analogue of results, known on the Stokes problem with the homogeneous Dirichlet or Navier or Navier–type boundary conditions on the whole boundary of

domain Ω , see [11], [2] and [3]. Recall that the theorem is non-trivial especially due to the variety of used boundary conditions and the fact that one cannot apply Riesz' theorem in the general L^r -framework in order to establish the existence and uniqueness of a solution of the equation $\mathcal{A}_r \mathbf{v} = \mathbf{f}$ for $\mathbf{f} \in \mathbf{V}_\sigma^{-1,r}(\Omega)$.

Theorem 1 (on the weak Stokes operator \mathcal{A}_r). *The weak Stokes operator \mathcal{A}_r is a bounded, closed and injective operator from $\mathbf{V}_\sigma^{1,r}(\Omega)$ to $\mathbf{V}_\sigma^{-1,r}(\Omega)$ with $D(\mathcal{A}_r) = \mathbf{V}_\sigma^{1,r}(\Omega)$ and $R(\mathcal{A}_r) = \mathbf{V}_\sigma^{-1,r}(\Omega)$. The adjoint operator to \mathcal{A}_r is $\mathcal{A}_r^* = \mathcal{A}_{r'}$.*

Proof. The case $r = 2$ is proven in [29]. Thus, assume that $r \neq 2$. We split the proof to several parts.

(a) Denote by $\|\mathcal{A}_r\|_{\mathbf{V}_\sigma^{1,r} \rightarrow \mathbf{V}_\sigma^{-1,r}}$ the norm of operator \mathcal{A}_r . Since

$$\begin{aligned} \|\mathcal{A}_r\|_{\mathbf{V}_\sigma^{1,r} \rightarrow \mathbf{V}_\sigma^{-1,r}} &= \sup_{\mathbf{v} \in \mathbf{V}_\sigma^{1,r}(\Omega), \mathbf{v} \neq \mathbf{0}} \frac{\|\mathcal{A}_r \mathbf{v}\|_{\mathbf{V}_\sigma^{-1,r}}}{\|\mathbf{v}\|_{1,r}} \leq c \sup_{\mathbf{v} \in \mathbf{V}_\sigma^{1,r}(\Omega), \mathbf{v} \neq \mathbf{0}} \frac{\|\mathcal{A}_r \mathbf{v}\|_{\mathbf{V}_\sigma^{-1,r}}}{\|\nabla \mathbf{v}\|_r} \\ &= c \sup_{\mathbf{v} \in \mathbf{V}_\sigma^{1,r}(\Omega), \mathbf{v} \neq \mathbf{0}} \sup_{\mathbf{w} \in \mathbf{V}_\sigma^{1,r'}(\Omega), \mathbf{w} \neq \mathbf{0}} \frac{|(\nabla \mathbf{v}, \nabla \mathbf{w})|}{\|\nabla \mathbf{v}\|_r \|\nabla \mathbf{w}\|_{r'}} \leq c, \end{aligned}$$

the operator \mathcal{A}_r is bounded. The identity $D(\mathcal{A}_r) = \mathbf{V}_\sigma^{1,r}(\Omega)$ follows from the definition of \mathcal{A}_r . Operator \mathcal{A}_r is closed, as a bounded linear operator, defined on the whole space $\mathbf{V}_\sigma^{1,r}(\Omega)$.

(b) In this part, we consider $\mathbf{F} \in \mathbf{V}_\sigma^{-1,2}(\Omega)$ of a special form and deal with the equation $\mathcal{A}_2 \mathbf{v} = \mathbf{F}$. Concretely, we assume that $\mathbb{F} \in W^{1,2}(\Omega)^{2 \times 2}$ satisfies the condition

$$\mathbb{F}(x_1, x_2 + \tau) = \mathbb{F}(x_1, x_2) \quad \text{for a.a. } (x_1, x_2) \in \Gamma_0 \quad (3.1)$$

and define $\mathbf{F} \in \mathbf{V}_\sigma^{-1,2}(\Omega)$ by the formula

$$\langle \mathbf{F}, \mathbf{w} \rangle_{(\mathbf{V}_\sigma^{-1,2}, \mathbf{V}_\sigma^{1,2})} := - \int_\Omega \mathbb{F} : \nabla \mathbf{w} \, dx \quad (3.2)$$

for all $\mathbf{w} \in \mathbf{V}_\sigma^{1,2}(\Omega)$. Then, due to [29, Lemma 1 and Theorem 2], the equation $\mathcal{A}_2 \mathbf{v} = \mathbf{F}$ has a unique solution $\mathbf{v} \in \mathbf{V}_\sigma^{1,2}(\Omega) \cap \mathbf{W}^{2,2}(\Omega)$, there exists $p \in W^{1,2}(\Omega)$ (an associated pressure), such that $\nabla \mathbf{v} + p\mathbb{I} - \mathbb{F} \in W^{1,2}(\Omega)^{2 \times 2}$, and \mathbf{v}, p satisfy the equation

$$-\Delta \mathbf{v} + \nabla p - \operatorname{div} \mathbb{F} \equiv \operatorname{div} (-\nu \nabla \mathbf{v} + p\mathbb{I} - \mathbb{F}) = \mathbf{0} \quad (3.3)$$

a.e. in Ω . If $r > 2$ then there exists $c_1 > 0$, independent of \mathbf{v} and \mathbb{F} , such that

$$\|\mathbf{v}\|_{1,r} \leq c_1 \|\mathbb{F}\|_r. \quad (3.4)$$

This estimate is the key part of the proof of Theorem 1. As the proof of (3.4) is relatively long, the used technique differs from other sections, and in order not to disrupt the logical sequence of the text, we postpone the derivation of (3.4) to a separate section (Section 5).

(c) In this part, we assume that $r > 2$, $\mathbb{F} \in L^r(\Omega)^{2 \times 2}$ and the functional $\mathbf{F} \in \mathbf{V}_\sigma^{-1,r}(\Omega)$ is defined by the same formula as (3.2) (for all $\mathbf{w} \in \mathbf{V}_\sigma^{1,r'}(\Omega)$). We prove the solvability of the equation $\mathcal{A}_r \mathbf{v} = \mathbf{F}$.

There exists a sequence $\{\mathbb{F}_n\}$ in $W^{1,2}(\Omega)^{2 \times 2}$, satisfying (3.1), such that $\mathbb{F}_n \rightarrow \mathbb{F}$ in $L^r(\Omega)^{2 \times 2}$. Let $\{\mathbf{F}_n\}$ be a sequence of functionals from $\mathbf{V}_\sigma^{-1,r}(\Omega)$, related to \mathbb{F}_n through formula (3.2), which is now valid for all $\mathbf{w} \in \mathbf{V}_\sigma^{1,r'}(\Omega)$. The functionals \mathbf{F} and \mathbf{F}_n satisfy

$$\|\mathbf{F}_n - \mathbf{F}\|_{\mathbf{V}_\sigma^{-1,r}} = \sup_{\mathbf{w} \in \mathcal{C}_\sigma^\infty(\Omega); \mathbf{w} \neq \mathbf{0}} \|\mathbf{w}\|_{1,r'}^{-1} \left| \int_\Omega (\mathbb{F}_n - \mathbb{F}) : \nabla \mathbf{w} \, d\mathbf{x} \right| \leq c \|\mathbb{F}_n - \mathbb{F}\|_r. \quad (3.5)$$

Since $\mathbf{V}_\sigma^{-1,r}(\Omega) \subset \mathbf{V}_\sigma^{-1,2}(\Omega)$, there exists (due to part (b) of this proof) a sequence $\{\mathbf{v}_n\}$ in $\mathbf{V}_\sigma^{1,2}(\Omega) \cap \mathbf{W}^{2,2}(\Omega)$, such that $\mathcal{A}_2 \mathbf{v}_n = \mathbf{F}_n$ and

$$\|\mathbf{v}_n\|_{1,r} \leq c_1 \|\mathbb{F}_n\|_r. \quad (3.6)$$

As the space $\mathbf{V}_\sigma^{1,r}(\Omega)$ is reflexive, there exists a subsequence (which we again denote by $\{\mathbf{v}_n\}$), such that $\mathbf{v}_n \rightharpoonup \mathbf{v}$ in $\mathbf{V}_\sigma^{1,r}(\Omega)$. The functions \mathbf{v}_n satisfy

$$(\nabla \mathbf{v}_n, \nabla \mathbf{w}) = \langle \mathbf{F}_n, \mathbf{w} \rangle_{(\mathbf{V}_\sigma^{-1,2}, \mathbf{V}_\sigma^{1,2})} \quad \text{for all } \mathbf{w} \in \mathbf{V}_\sigma^{1,2}(\Omega).$$

If $\mathbf{w} \in \mathbf{V}_\sigma^{1,r'}(\Omega)$ then the right hand side equals $\langle \mathbf{F}_n, \mathbf{w} \rangle_{(\mathbf{V}_\sigma^{-1,r}, \mathbf{V}_\sigma^{1,r'})}$. Thus, as $\mathbf{V}_\sigma^{1,2}(\Omega)$ is dense in $\mathbf{V}_\sigma^{1,r'}(\Omega)$, we also have

$$(\nabla \mathbf{v}_n, \nabla \mathbf{w}) = \langle \mathbf{F}_n, \mathbf{w} \rangle_{(\mathbf{V}_\sigma^{-1,r}, \mathbf{V}_\sigma^{1,r'})} \quad \text{for all } \mathbf{w} \in \mathbf{V}_\sigma^{1,r'}(\Omega).$$

The limit transition for $n \rightarrow \infty$ yields

$$(\nabla \mathbf{v}, \nabla \mathbf{w}) = \langle \mathbf{F}, \mathbf{w} \rangle_{(\mathbf{V}_\sigma^{-1,r}, \mathbf{V}_\sigma^{1,r'})} \quad \text{for all } \mathbf{w} \in \mathbf{V}_\sigma^{1,r'}(\Omega),$$

which means that $\mathcal{A}_r \mathbf{v} = \mathbf{F}$. It also follows from the limit transition that the function \mathbf{v} satisfies inequality (3.4).

(d) Here, we show that each functional $\mathbf{f} \in \mathbf{V}_\sigma^{-1,r}(\Omega)$ (for any $r \in (1, \infty)$) can be expressed in the form (3.2) for some appropriate $\mathbb{F} \in L^r(\Omega)^{2 \times 2}$. This, together with part (c), shows that if $r > 2$ and $\mathbf{f} \in \mathbf{V}_\sigma^{-1,r}(\Omega)$ then the equation $\mathcal{A}_r \mathbf{v} = \mathbf{f}$ is solvable. In other words, $R(\mathcal{A}_r) = \mathbf{V}_\sigma^{-1,r}(\Omega)$.

Thus, let $\mathbf{f} \in \mathbf{V}_\sigma^{-1,r}(\Omega)$. As $\mathbf{V}_\sigma^{1,r'}(\Omega)$ is a closed subspace of $\mathbf{W}^{1,r'}(\Omega)$, the functional \mathbf{f} can be extended (by the Hahn–Banach theorem) to a bounded linear functional $\mathbf{f}_* \in \mathbf{W}_0^{-1,r}(\Omega)$, such that $\|\mathbf{f}_*\|_{\mathbf{W}_0^{-1,r}} = \|\mathbf{f}\|_{\mathbf{V}_\sigma^{-1,r}}$. There exists (by Lemma 1) $\mathbb{F} \in L^r(\Omega)^{2 \times 2}$, such that $\operatorname{div} \mathbb{F} = \mathbf{f}_*$ in the sense of distributions in Ω and $\|\mathbb{F}\|_r \leq c \|\mathbf{f}_*\|_{\mathbf{W}_0^{-1,r}} \equiv \|\mathbf{f}\|_{\mathbf{V}_\sigma^{-1,r}}$. As $\mathbf{f}_* = \mathbf{f}$ on $\mathbf{V}_\sigma^{1,r'}(\Omega)$, we have

$$\langle \operatorname{div} \mathbb{F}, \mathbf{w} \rangle = \langle \mathbf{f}, \mathbf{w} \rangle \quad \forall \mathbf{w} \in \mathcal{C}_{0,\sigma}^\infty(\Omega),$$

where $\langle \cdot, \cdot \rangle$ denotes the action of a distribution on a function from $\mathcal{C}_0^\infty(\Omega)$. Since the left hand side is equal to

$$-\langle \mathbb{F}, \nabla \mathbf{w} \rangle = - \int_\Omega \mathbb{F} : \nabla \mathbf{w} \, d\mathbf{x},$$

we obtain

$$\langle \mathbf{f}, \mathbf{w} \rangle_{(\mathbf{V}_\sigma^{-1,r}, \mathbf{V}_\sigma^{1,r'})} = \langle \mathbf{f}, \mathbf{w} \rangle = - \int_\Omega \mathbb{F} : \nabla \mathbf{w} \, d\mathbf{x} \quad \text{for all } \mathbf{w} \in \mathcal{C}_{0,\sigma}^\infty(\Omega).$$

Both the left hand– and right hand–sides can be continuously extended so that they equal each other for all $\mathbf{w} \in \mathbf{V}_\sigma^{1,r'}(\Omega)$. This means that $\mathbf{f} = \mathbf{F}$, where \mathbf{F} is related to \mathbb{F} through formula (3.2) (where we consider $\mathbf{w} \in \mathbf{V}_\sigma^{1,r'}(\Omega)$). It follows from part (c) that there exists $\mathbf{v} \in \mathbf{V}_\sigma^{1,r}(\Omega)$, such that $\mathcal{A}_r \mathbf{v} = \mathbf{f}$. There exists $c > 0$, independent of \mathbf{f} and \mathbf{v} , such that the solution satisfies the estimate

$$\|\mathbf{v}\|_{1,r} \leq c \|\mathbf{f}\|_{\mathbf{V}_\sigma^{-1,r}}. \quad (3.7)$$

(e) In this part, we derive information on the adjoint operator \mathcal{A}_r^* to \mathcal{A}_r for any $r \in (1, \infty)$. The adjoint operator acts from $\mathbf{V}_\sigma^{-1,r}(\Omega)^* \equiv \mathbf{V}_\sigma^{1,r'}(\Omega)^{**}$ to $\mathbf{V}_\sigma^{1,r}(\Omega)^* = \mathbf{V}_\sigma^{-1,r'}(\Omega)$. However, as $\mathbf{V}_\sigma^{1,r'}(\Omega)$ is reflexive, $\mathbf{V}_\sigma^{1,r'}(\Omega)^{**}$ can be identified with $\mathbf{V}_\sigma^{1,r'}(\Omega)$. Thus, \mathcal{A}_r^* is an operator from $\mathbf{V}_\sigma^{1,r'}(\Omega)$ to $\mathbf{V}_\sigma^{-1,r}(\Omega)$. The domain of \mathcal{A}_r^* is, by definition, the set of all $\mathbf{w} \in \mathbf{V}_\sigma^{1,r'}(\Omega)$, such that $\langle \mathcal{A}_r \mathbf{z}, \mathbf{w} \rangle_{(\mathbf{V}_\sigma^{-1,r}, \mathbf{V}_\sigma^{1,r'})}$ is, in dependence on \mathbf{z} , a bounded linear functional on $\mathbf{V}_\sigma^{1,r}(\Omega)$. This functional is exactly $\mathcal{A}_r^* \mathbf{w}$ and it satisfies

$$\langle \mathcal{A}_r^* \mathbf{w}, \mathbf{z} \rangle_{(\mathbf{V}_\sigma^{-1,r}, \mathbf{V}_\sigma^{1,r'})} = \langle \mathcal{A}_r \mathbf{z}, \mathbf{w} \rangle_{(\mathbf{V}_\sigma^{-1,r}, \mathbf{V}_\sigma^{1,r'})}.$$

By definition of \mathcal{A}_r , the right hand side equals $(\nabla \mathbf{z}, \nabla \mathbf{w})$. This can be also written as $(\nabla \mathbf{w}, \nabla \mathbf{z})$ and it is, in dependence on \mathbf{z} , a bounded linear functional acting on $\mathbf{V}_\sigma^{1,r}(\Omega)$ for each fixed $\mathbf{w} \in \mathbf{V}_\sigma^{1,r'}(\Omega)$. From this, we deduce that $D(\mathcal{A}_r^*) = \mathbf{V}_\sigma^{1,r'}(\Omega)$ and $\mathcal{A}_r^* = \mathcal{A}_{r'}$.

(f) Here, we assume that $1 < r < \infty$ and prove the uniqueness of the solution \mathbf{v} of the equation $\mathcal{A}_r \mathbf{v} = \mathbf{f}$ for any $\mathbf{f} \in \mathbf{V}_\sigma^{-1,r}(\Omega)$.

Assume at first that $r > 2$. Then, as $\mathbf{V}_\sigma^{1,r}(\Omega) \subset \mathbf{V}_\sigma^{1,r'}(\Omega)$, one can use equation (2.1) with $\mathbf{w} = \mathbf{v}$ and deduce that $\mathcal{A}_r \mathbf{v} = \mathbf{0}$ implies $\mathbf{v} = \mathbf{0}$, which means that the solution of the equation $\mathcal{A}_r \mathbf{v} = \mathbf{f}$ is unique.

Assume now that $1 < r < 2$. The null space of \mathcal{A}_r and range of $\mathcal{A}_{r'}$ satisfy the identity $N(\mathcal{A}_r) = R(\mathcal{A}_{r'})^\perp$, see [16, p. 168]. However, as $R(\mathcal{A}_{r'})$ is the whole space $\mathbf{V}_\sigma^{-1,r'}(\Omega)$ (because $r' > 2$ and due to parts (c) and (d) of this proof), the space of annihilators $R(\mathcal{A}_{r'})^\perp$ is trivial. Thus, we obtain the identity $N(\mathcal{A}_r) = \{\mathbf{0}\}$. This implies the uniqueness of the solution \mathbf{v} and the injectivity of operator \mathcal{A}_r .

(g) Finally, we assume that $1 < r < 2$ and prove the existence of a solution \mathbf{v} of the equation $\mathcal{A}_r \mathbf{v} = \mathbf{f}$ for any $\mathbf{f} \in \mathbf{V}_\sigma^{-1,r}(\Omega)$. Since $R(\mathcal{A}_r)^\perp = N(\mathcal{A}_{r'}) = \{\mathbf{0}\}$, $R(\mathcal{A}_r)$ is dense in $\mathbf{V}_\sigma^{-1,r}(\Omega)$. The operator $\mathcal{A}_{r'}^{-1}$ is bounded from $\mathbf{V}_\sigma^{-1,r'}(\Omega)$ to $\mathbf{V}_\sigma^{1,r'}(\Omega)$, which follows from the inequality $r' > 2$, parts (c) and (d) of this proof and the closed graph theorem. Consequently, due to [16, p. 169], the operator $(\mathcal{A}_{r'})^{-1} \equiv \mathcal{A}_r^{-1}$ is bounded from $\mathbf{V}_\sigma^{-1,r}(\Omega)$ to $\mathbf{V}_\sigma^{1,r}(\Omega)$. Hence \mathcal{A}_r maps a closed set in $\mathbf{V}_\sigma^{1,r}(\Omega)$ onto a closed set in $\mathbf{V}_\sigma^{-1,r}(\Omega)$. It means that $R(\mathcal{A}_r)$ is closed in $\mathbf{V}_\sigma^{-1,r}(\Omega)$. Since it is also dense in $\mathbf{V}_\sigma^{-1,r}(\Omega)$, we have $R(\mathcal{A}_r) = \mathbf{V}_\sigma^{-1,r}(\Omega)$. This completes the proof. \square

4 The weak Stokes problem

The spaces $\mathbf{W}_{\text{per}}^{1-1/r,r}(\Gamma_{\text{in}})$, $\mathbf{W}_{\text{per}}^{1-1/r',r'}(\Gamma_{\text{out}})$ and $\mathbf{W}_{\text{per}}^{-1/r,r}(\Gamma_{\text{out}})$. Recall that the straight lines γ_{in} , γ_{out} and the line segments Γ_{in} , Γ_{out} are sketched on Fig. 1. Let $1 < r < \infty$. We denote by $\mathbf{W}_{\text{per}}^{1-1/r,r}(\gamma_{\text{in}})$ the space of τ -periodic functions in $W_{\text{loc}}^{1-1/r,r}(\gamma_{\text{in}})$ and by $\mathbf{W}_{\text{per}}^{1-1/r,r}(\Gamma_{\text{in}})$ the space of functions from $W^{1-1/r,r}(\Gamma_{\text{in}})$, that can be extended from Γ_{in} to γ_{in} as functions in $\mathbf{W}_{\text{per}}^{1-1/r,r}(\gamma_{\text{in}})$.

The space $\mathbf{W}_{\text{per}}^{1-1/r',r'}(\Gamma_{\text{out}})$ is defined by analogy with $\mathbf{W}_{\text{per}}^{1-1/r,r}(\Gamma_{\text{in}})$. Let us denote by $\mathbf{W}_{\text{per}}^{-1/r,r}(\Gamma_{\text{out}})$ the dual space to $\mathbf{W}_{\text{per}}^{1-1/r',r'}(\Gamma_{\text{out}})$ and by $\langle \cdot, \cdot \rangle_{(\mathbf{W}_{\text{per}}^{-1/r,r}(\Gamma_{\text{out}}), \mathbf{W}_{\text{per}}^{1-1/r',r'}(\Gamma_{\text{out}}))}$ the duality pairing between $\mathbf{W}_{\text{per}}^{-1/r,r}(\Gamma_{\text{out}})$ and $\mathbf{W}_{\text{per}}^{1-1/r',r'}(\Gamma_{\text{out}})$.

An alternative description of $\mathbf{W}_{\text{per}}^{1-1/r,r}(\Gamma_{\text{in}})$ (which also holds for $\mathbf{W}_{\text{per}}^{-1/r,r}(\Gamma_{\text{out}})$) is explained in the following remark.

Remark 2. Assume at first that $r > 2$. In this case, functions from $\mathbf{W}_{\text{per}}^{1-1/r,r}(\Gamma_{\text{in}})$ have traces at the end points A_0 and A_1 of Γ_{in} . Denote $\widetilde{\mathbf{W}}_{\text{per}}^{1-1/r,r}(\Gamma_{\text{in}}) := \{\mathbf{w} \in \mathbf{W}^{1-1/r,r}(\Gamma_{\text{in}}); \mathbf{w}(A_0) = \mathbf{w}(A_1) \text{ in the sense of traces}\}$. Obviously, $\mathbf{W}_{\text{per}}^{1-1/r,r}(\Gamma_{\text{in}}) \subset \widetilde{\mathbf{W}}_{\text{per}}^{1-1/r,r}(\Gamma_{\text{in}})$. Let us show that the opposite inclusion is also true. Thus, let $\mathbf{w} \in \widetilde{\mathbf{W}}_{\text{per}}^{1-1/r,r}(\Gamma_{\text{in}})$. Denote by the same symbol \mathbf{w} the τ -periodic extension from Γ_{in} to γ_{in} . We can assume without loss of generality that $A_0 = (0, 0)$ and $A_1 = (0, \tau)$. Put $A_2 := (0, 2\tau)$. In order to verify that $\mathbf{w} \in \mathbf{W}_{\text{loc}}^{1-1/r,r}(\gamma_{\text{in}})$, it is sufficient to show that \mathbf{w} lies in the space $\mathbf{W}^{1-1/r,r}(A_0, A_2)$. The norm of \mathbf{w} in $\mathbf{W}^{1-1/r,r}(A_0, A_2)$ equals $\|\mathbf{w}\|_{r; A_0, A_2} + \langle\langle \mathbf{w} \rangle\rangle_{1-1/r,r; A_0, A_2}$ (see formulas (2) and (3) in [24], paragraph 8.3.2, p. 386), where

$$\begin{aligned} \langle\langle \mathbf{w} \rangle\rangle_{1-1/r,r; A_0, A_2} &= \int_0^{2\tau} \int_0^{2\tau} \frac{|\mathbf{w}(0, y) - \mathbf{w}(0, z)|^r}{|y - z|^r} dy dz \\ &= \left(\int_0^\tau \int_0^\tau + \int_0^\tau \int_\tau^{2\tau} + \int_\tau^{2\tau} \int_0^\tau + \int_\tau^{2\tau} \int_\tau^{2\tau} \right) \frac{|\mathbf{w}(0, y) - \mathbf{w}(0, z)|^r}{|y - z|^r} dy dz \\ &\leq \langle\langle \mathbf{w} \rangle\rangle_{1-1/r,r; A_0, A_1} + \langle\langle \mathbf{w} \rangle\rangle_{1-1/r,r; A_1, A_2} + 2 \int_0^\tau \int_\tau^{2\tau} \frac{|\mathbf{w}(0, y) - \mathbf{w}(0, z)|^r}{|y - z|^r} dy dz. \end{aligned}$$

The first two terms on the right hand side are equal due to the τ -periodicity of $\mathbf{w}(0, \cdot)$. The integral on the right hand side is less than or equal to

$$\begin{aligned} &c \int_0^\tau \int_\tau^{2\tau} \left(\frac{|\mathbf{w}(0, y) - \mathbf{w}(0, \tau)|^r}{(y - \tau)^r} + \frac{|\mathbf{w}(0, \tau) - \mathbf{w}(0, z)|^r}{(\tau - z)^r} \right) dy dz \\ &\leq c \int_\tau^{2\tau} \frac{|\mathbf{w}(0, y) - \mathbf{w}(0, \tau)|^r}{(y - \tau)^{r-1}} dy + c \int_0^\tau \frac{|\mathbf{w}(0, \tau) - \mathbf{w}(0, z)|^r}{(\tau - z)^{r-1}} dz \\ &\leq c \int_\tau^{2\tau} \int_\tau^{2\tau} \frac{|\mathbf{w}(0, y) - \mathbf{w}(0, x)|^r}{|y - x|^r} dx dy + c \int_0^\tau \int_0^\tau \frac{|\mathbf{w}(0, x) - \mathbf{w}(0, z)|^r}{|x - z|^r} dx dz. \end{aligned}$$

The last estimate holds due to the fractional Hardy inequality, see e.g. [14, Theorem 2.1]. The right hand side is, due to the τ -periodicity of the function $\mathbf{w}(0, \cdot)$, less than or equal to $c \langle\langle \mathbf{w} \rangle\rangle_{1-1/r,r; A_0, A_1}$. This confirms that $\mathbf{w} \in \mathbf{W}^{1-1/r,r}(A_0, A_2)$, indeed.

In the critical case $r = 2$, one cannot characterize $\mathbf{W}_{\text{per}}^{1/2,2}(\Gamma_{\text{in}})$ as in the case $r > 2$, because the traces at the end points A_0 and A_1 generally do not exist. Moreover, although $\mathbf{W}_0^{1/2,2}(\Gamma_{\text{in}}) = \mathbf{W}^{1/2,2}(\Gamma_{\text{in}})$, see [25, Theorem II.11.1]), one cannot identify $\mathbf{W}_{\text{per}}^{1/2,2}(\Gamma_{\text{in}})$ with $\mathbf{W}^{1/2,2}(\Gamma_{\text{in}})$, because e.g. the linear function $g(0, y) := y$ for $(0, y) \in \Gamma_{\text{in}}$ is in $\mathbf{W}^{1/2,2}(\Gamma_{\text{in}})$, but its τ -periodic extension to γ_{in} is not in $\mathbf{W}_{\text{loc}}^{1/2,2}(\gamma_{\text{in}})$. Thus, one only has the inclusion $\mathbf{W}_{\text{per}}^{1/2,2}(\Gamma_{\text{in}}) \subset \mathbf{W}^{1/2,2}(\Gamma_{\text{in}})$.

If $1 < r < 2$ then $\mathbf{W}_0^{1-1/r,r}(\Gamma_{\text{in}}) = \mathbf{W}^{1-1/r,r}(\Gamma_{\text{in}})$ (which follows from the density of $\mathbf{W}^{1-1/r,r}(\Gamma_{\text{in}})$ in $\mathbf{W}^{1/2,2}(\Gamma_{\text{in}})$, the identity $\mathbf{W}^{1/2,2}(\Gamma_{\text{in}}) = \mathbf{W}_0^{1/2,2}(\Gamma_{\text{in}})$ and the density of $\mathbf{W}_0^{1/2,2}(\Gamma_{\text{in}})$ in $\mathbf{C}_0^\infty(\Gamma_{\text{in}}^0)$). This and similar estimates of $\langle\langle \mathbf{w} \rangle\rangle_{1-1/r,r; A_0, A_2}$, as above, enable one to show that every function from $\mathbf{W}^{1-1/r,r}(\Gamma_{\text{in}})$, periodically extended to γ_{in} with the period τ , is in $\mathbf{W}_{\text{loc}}^{1-1/r,r}(\gamma_{\text{in}})$. Thus, one obtains the identity $\mathbf{W}_{\text{per}}^{1-1/r,r}(\Gamma_{\text{in}}) = \mathbf{W}^{1-1/r,r}(\Gamma_{\text{in}})$.

Definition 1 (weak solution of the Stokes problem (1.1)–(1.6)). Let $r \in (1, \infty)$. Let $\mathbf{f} \in \mathbf{W}_0^{-1,r}(\Omega)$, $\mathbf{g} \in \mathbf{W}_{\text{per}}^{1-1/r,r}(\Gamma_{\text{in}})$ and $\mathbf{h} \in \mathbf{W}_{\text{per}}^{-1/r,r}(\Gamma_{\text{out}})$. Let $\mathbb{F} \in L^r(\Omega)^{2 \times 2}$ be a tensor function, satisfying $\text{div } \mathbb{F} = \mathbf{f}$ in the sense of distributions, provided by Lemma 1. A divergence-free

function $\mathbf{u} \in \mathbf{W}^{1,r}(\Omega)$, satisfying the conditions (1.3), (1.4) and (1.5) in the sense of traces on Γ_{in} , Γ_p and Γ_0 , respectively, and the equation

$$\nu \int_{\Omega} \nabla \mathbf{u} : \nabla \mathbf{w} \, d\mathbf{x} = - \int_{\Omega} \mathbb{F} : \nabla \mathbf{w} \, d\mathbf{x} + \langle \mathbf{h}, \mathbf{w} \rangle_{(\mathbf{W}_{\text{per}}^{-1/r,r}(\Gamma_{\text{out}}), \mathbf{W}_{\text{per}}^{1-1/r',r'}(\Gamma_{\text{out}}))} \quad (4.1)$$

for all $\phi \in \mathbf{V}_{\sigma}^{1,r'}(\Omega)$, is said to be a *weak solution* to the Stokes problem (1.1)–(1.6).

We show in Theorem 2 that if \mathbf{u} is a weak solution of the problem (1.1)–(1.6) then there exists an associate pressure p so that \mathbf{u} and p satisfy an analogue of the condition (1.6) in a certain weak sense on Γ_{out} .

Theorem 2 (a posteriori properties of a weak solution). *1) Let r , \mathbf{f} , \mathbb{F} , \mathbf{g} and \mathbf{h} satisfy the assumptions from Definition 1. Let \mathbf{u} be a weak solution of the Stokes problem (1.1)–(1.6). Then there exists an associated pressure $p \in L^r(\Omega)$ such that \mathbf{u} and p satisfy the equations (1.1) and (1.2) in the sense of distributions in Ω and*

$$(-\nu \nabla \mathbf{u} - p \mathbb{I} - \mathbb{F}) \cdot \mathbf{n} = \mathbf{h} \quad (4.2)$$

holds as an equality in $\mathbf{W}_{\text{per}}^{-1/r,r}(\Gamma_{\text{out}})$.

2) If, moreover, $\mathbf{f} \in \mathbf{L}^r(\Omega)$ then the tensor function \mathbb{F} can be constructed so that it lies in $W_{\text{per}}^{1,r}(\Omega)^{2 \times 2}$, satisfies the equation $\text{div } \mathbb{F} = \mathbf{f}$ a.e. in Ω and the condition $\mathbb{F} \cdot \mathbf{n} = \mathbf{0}$ a.e. on Γ_{out} . In this case, (4.2) takes the form

$$(-\nu \nabla \mathbf{u} - p \mathbb{I}) \cdot \mathbf{n} = \mathbf{h}, \quad (4.3)$$

consistent with (1.6).

Proof. Suppose at first that the test function \mathbf{w} , used in (4.1), is in $\mathbf{C}_{0,\sigma}^{\infty}(\Omega)$. Then, since $\mathbf{w} = \mathbf{0}$ on Γ_{out} , (4.1) and the equation $\mathbf{f} = \text{div } \mathbb{F}$ imply that

$$\langle \nu \Delta \mathbf{u} + \text{div } \mathbb{F}, \mathbf{w} \rangle = 0.$$

As this holds for all $\mathbf{w} \in \mathbf{C}_{0,\sigma}^{\infty}(\Omega)$, we can apply De Rham's lemma (see [30, p. 14]) and deduce that there exists a distribution p_0 in Ω , such that

$$\nu \Delta \mathbf{u} + \text{div } \mathbb{F} = \nabla p_0 \quad (4.4)$$

holds in Ω in the sense of distributions. As both $\nu \Delta \mathbf{u}$ and $\text{div } \mathbb{F}$ can also be identified with elements of $\mathbf{W}^{-1,r}(\Omega)$, ∇p_0 belongs to $\mathbf{W}^{-1,r}(\Omega)$, too. It follows from [11, Lemma IV.1.1] that $p_0 \in L^r(\Omega)$ and it can be chosen so that

$$\|p_0\|_r \leq c \|\nu \Delta \mathbf{u} + \text{div } \mathbb{F}\|_{\mathbf{W}^{-1,r}}, \quad (4.5)$$

where c depends only on ν and Ω .

Since $-\nu \nabla \mathbf{u} + p_0 \mathbb{I} - \mathbb{F} \in L^r(\Omega)^{2 \times 2}$ and $\text{div } (-\nu \nabla \mathbf{u} + p_0 \mathbb{I} - \mathbb{F}) = \mathbf{0}$ in the sense of distributions, we can apply Theorem III.2.2 from [11] and deduce that $(-\nu \nabla \mathbf{u} + p_0 \mathbb{I} - \mathbb{F}) \cdot \mathbf{n} \in \mathbf{W}_{\text{per}}^{-1/r,r}(\Gamma_{\text{out}})$ (in the sense of traces).

Let $\mathbf{w} \in \mathbf{C}_{\sigma}^{\infty}(\Omega)$. The equation (4.1) and the generalized Gauss identity (see [11, p. 160]) imply that

$$0 = \langle \text{div } [\nu \nabla \mathbf{u} + \mathbb{F} - p_0 \mathbb{I}], \mathbf{w} \rangle_{(\mathbf{W}_0^{-1,r}(\Omega), \mathbf{W}^{1,r'}(\Omega))}$$

$$\begin{aligned}
&= \langle (\nu \nabla \mathbf{u} + \mathbb{F} - p_0 \mathbb{I}) \cdot \mathbf{n}, \mathbf{w} \rangle_{(\mathbf{W}^{-1/r, r}(\partial\Omega), \mathbf{W}^{1-1/r', r'}(\partial\Omega))} - \int_{\Omega} [\nu \nabla \mathbf{u} + \mathbb{F}] : \nabla \mathbf{w} \, dx \\
&= \langle (\nu \nabla \mathbf{u} + \mathbb{F} - p_0 \mathbb{I}) \cdot \mathbf{n}, \mathbf{w} \rangle_{\mathbf{W}_{\text{per}}^{-1/r, r}(\Gamma_{\text{out}}), \mathbf{W}_{\text{per}}^{1-1/r', r'}(\Gamma_{\text{out}})} - \langle \mathbf{h}, \mathbf{w} \rangle_{(\mathbf{W}_{\text{per}}^{-1, r}(\Gamma_{\text{out}}), \mathbf{W}_{\text{per}}^{1-1/r', r'}(\Gamma_{\text{out}}))} \\
&\quad + \langle \mathbf{h}, \mathbf{w} \rangle_{(\mathbf{W}_{\text{per}}^{-1, r}(\Gamma_{\text{out}}), \mathbf{W}_{\text{per}}^{1-1/r', r'}(\Gamma_{\text{out}}))} - \int_{\Omega} [\nu \nabla \mathbf{u} + \mathbb{F}] : \nabla \mathbf{w} \, dx \\
&= \langle (\nu \nabla \mathbf{u} + \mathbb{F} - p_0 \mathbb{I}) \cdot \mathbf{n} - \mathbf{h}, \mathbf{w} \rangle_{\mathbf{W}_{\text{per}}^{-1/r, r}(\Gamma_{\text{out}}), \mathbf{W}_{\text{per}}^{1-1/r', r'}(\Gamma_{\text{out}})}.
\end{aligned}$$

The set of traces of all functions from $\mathcal{C}_{\sigma}^{\infty}(\Omega)$ on Γ_{out} is dense in the set of all functions $\mathbf{w} \in \mathbf{W}_{\text{per}}^{1-1/r', r'}(\Gamma_{\text{out}})$, such that $\int_{\Gamma_{\text{out}}} \mathbf{w} \cdot \mathbf{n} \, dl = 0$. (This follows from the density of the space of all functions $\mathbf{w} \in \mathcal{C}_{\text{per}}^{\infty}(\Gamma_{\text{out}})$, such that $\int_{\Gamma_{\text{out}}} \mathbf{w} \cdot \mathbf{n} \, dl = 0$, in $\{\mathbf{w} \in \mathbf{W}_{\text{per}}^{1-1/r', r'}(\Gamma_{\text{out}}); \int_{\Gamma_{\text{out}}} \mathbf{w} \cdot \mathbf{n} \, dl = 0\}$, and from the possibility of extension of any function in $\mathcal{C}_{\text{per}}^{\infty}(\Gamma_{\text{out}})$ from Γ_{out} to Ω so that the extended function is in $\mathcal{C}_{\sigma}^{\infty}(\Omega)$.) Hence there exists $c_2 \in \mathbb{R}$ such that \mathbf{u} and p_0 satisfy

$$(-\nu \nabla \mathbf{u} - \mathbb{F} - p_0 \mathbb{I}) \cdot \mathbf{n} - \mathbf{h} = c_2 \mathbf{n}, \quad (4.6)$$

as an equality in $\mathbf{W}_{\text{per}}^{-1/r, r}(\Gamma_{\text{out}})$. Put $p := p_0 - c_2$. Then \mathbf{u} and p satisfy the equation (1.1) in the sense of distributions in Ω and the boundary condition (4.2) as an equality in $\mathbf{W}_{\text{per}}^{-1/r, r}(\Gamma_{\text{out}})$. This completes the proof of part 1). The statements in part 2) follow from the next lemma.

Denote by $W_{\text{per}}^{1, r}(\Omega)$ is the space of functions from $W^{1, r}(\Omega)$, that satisfy in the sense of traces the condition of periodicity on Γ_0 and Γ_1 , analogous to (1.5).

Lemma 2. *Let $\mathbf{f} \in \mathbf{L}^r(\Omega)$ be given. Then there exists $\mathbb{F} \in W_{\text{per}}^{1, r}(\Omega)^{2 \times 2}$, such that $\text{div } \mathbb{F} = \mathbf{f}$ a.e. in Ω , $\mathbb{F} \cdot \mathbf{n} = \mathbf{0}$ a.e. on Γ_{out} in the sense of traces and*

$$\|\mathbb{F}\|_{1, r} \leq c \|\mathbf{f}\|_r, \quad (4.7)$$

where $c = c(\Omega, r)$.

Proof. Denote $\widehat{\Omega} := \Omega \cup P$. Then $|\widehat{\Omega}| = \tau d$. Define $\mathbf{f} := \mathbf{0}$ in P . Thus, $\mathbf{f} \in \mathbf{L}^r(\widehat{\Omega})$. Put $\mathbf{k} := |\widehat{\Omega}|^{-1} \int_{\widehat{\Omega}} \mathbf{f} \, d\mathbf{x}$.

Let $\zeta = \zeta(x_1)$ be a smooth real function in $[0, d]$, such that $\zeta(0) = 1$ and ζ is supported in $[0, \delta]$, where $\delta > 0$ is so small that the profile P (see Fig. 1) lies in the stripe $\delta < x_1 < d$. Since $\int_{\widehat{\Omega}} \zeta'(x_1) \, d\mathbf{x} = -\tau$, we have $\int_{\widehat{\Omega}} (\mathbf{f} + \mathbf{k} d \zeta'(x_1)) \, d\mathbf{x} = \mathbf{0}$. Thus, due to [11, Theorem III.3.3], there exists $\mathbb{F}_0 \in W_0^{1, r}(\widehat{\Omega})^{2 \times 2}$, such that $\text{div } \mathbb{F}_0 = \mathbf{f} + \mathbf{k} d \zeta'$ a.e. in $\widehat{\Omega}$ and

$$\|\mathbb{F}_0\|_{1, r; \widehat{\Omega}} \leq c \|\mathbf{f} + \mathbf{k} d \zeta'\|_{r; \widehat{\Omega}} \leq c \|\mathbf{f}\|_r,$$

where $c = c(\tau, d, \zeta)$. Since $\mathbf{k} d \zeta'(x_1) = \text{div } [\mathbf{k} d \zeta(x_1) \otimes \mathbf{e}_1]$, \mathbb{F}_0 satisfies

$$\text{div } [\mathbb{F}_0 - \mathbf{k} d \zeta(x_1) \otimes \mathbf{e}_1] = \mathbf{f}$$

a.e. in Ω . Put $\mathbb{F} := \mathbb{F}_0 - \mathbf{k} d \zeta(x_1) \otimes \mathbf{e}_1$. The tensor function \mathbb{F} has all the properties, stated in the lemma. \square

The proof of Theorem 2 is completed. \square

Remark 3. The identification of the right hand side of (1.1) with $\text{div } \mathbb{F}$ enables us to deduce that \mathbf{u} and p satisfy (4.2) or (4.3), which are equalities in $\mathbf{W}_{\text{per}}^{-1/r, r}(\Gamma_{\text{out}})$.

If $\mathbf{f} \in \mathbf{W}_0^{-1,2}(\Omega)$ and \mathbb{F} is only in $L^r(\Omega)^{2 \times 2}$, as in part 1) of Theorem 2, then neither $(-\nu \nabla \mathbf{u} - p \mathbb{I}) \cdot \mathbf{n}$, nor $\mathbb{F} \cdot \mathbf{n}$ need not be in $\mathbf{W}_{\text{per}}^{-1/r,r}(\Gamma_{\text{out}})$. Thus, the sole term $\mathbb{F} \cdot \mathbf{n}$ need not have a sense on Γ_{out} and it is therefore generally not possible to require \mathbb{F} to satisfy $\mathbb{F} \cdot \mathbf{n} = \mathbf{0}$ on Γ_{out} . The situation is different if $\mathbf{f} \in \mathbf{L}^r(\Omega)$ and $\mathbb{F} \in W_{\text{per}}^{1,2}(\Omega)$, see part 2) of Theorem 2.

In order to establish the existence of a weak solution of the problem (1.1)–(1.6), we shall need the next lemma. It is a modification of a result, proven in [10, Section 3].

Lemma 3. *Let $1 < r < \infty$ and $\mathbf{g} \in \mathbf{W}^{s,r}(\Gamma_{\text{in}})$, where $s > 1/r$ if $1 < r \leq 2$ and $s = 1 - 1/r$ if $r > 2$. Let \mathbf{g} satisfy the condition $\mathbf{g}(A_0) = \mathbf{g}(A_1)$. There exists a divergence-free extension \mathbf{g}_* of \mathbf{g} from Γ_{in} to Ω and a constant $c_3 > 0$, independent of \mathbf{g} , such that $\mathbf{g}_* \in \mathbf{W}^{1,r}(\Omega)$, $\mathbf{g}_* = \mathbf{0}$ on Γ_p in the sense of traces,*

- a) $\|\mathbf{g}_*\|_{1,r} \leq c_3 \|\mathbf{g}\|_{s,r;\Gamma_{\text{in}}}$,
- b) \mathbf{g}_* satisfies the condition of periodicity (1.5) in the sense of traces on $\Gamma_0 \cup \Gamma_1$,
- c) $\mathbf{g}_* = (\Phi/\tau) \mathbf{e}_1$ in a neighbourhood of Γ_{out} , where $\Phi = -\int_{\Gamma_{\text{in}}} \mathbf{g} \cdot \mathbf{n} \, dl$.

Principles of the proof. The assumptions on number s guarantee that $rs > 1$ and it makes therefore sense to speak about the traces of the function \mathbf{g} at the end points A_0 and A_1 of Γ_{in} . This enables one to extend at first \mathbf{g} from Γ_{in} to $\partial\Omega$ in the way, described in [8, subsection 3.2]. The extended function (which is again denoted by \mathbf{g}) satisfies the condition of periodicity (1.5), the condition $\mathbf{g} = \mathbf{0}$ on Γ_p and the equality $\int_{\partial\Omega} \mathbf{g} \cdot \mathbf{n} \, dl = 0$. The inequality $\|\mathbf{g}\|_{1-1/r,r;\partial\Omega} \leq c \|\mathbf{g}\|_{s,r;\Gamma_{\text{in}}}$ can be proven by analogy with [8, Lemma 1], expressing the norm of \mathbf{g} in $\mathbf{W}^{s,r}(\Gamma_{\text{in}})$ by formulas (2) and (3) in [24], paragraph 8.3.2, p. 386, and the norm of \mathbf{g} in $\mathbf{W}^{1-1/r,r}(\partial\Omega)$ by formulas (II.4.8) and (II.4.9) in [11], p. 64.

The existence of a divergence-free extension $\mathbf{g}_* \in \mathbf{W}^{1,r}(\Omega)$ of function \mathbf{g} from $\partial\Omega$ to Ω follows from [11, Exercise III.3.5]. The extension satisfies the estimate $\|\mathbf{g}_*\|_{1,r} \leq c \|\mathbf{g}\|_{1-1/r,r;\partial\Omega} \leq c \|\mathbf{g}\|_{s,r;\partial\Omega}$.

The function \mathbf{g}_* can be further modified in the way described in [10, Subsection 3.3], so that it finally has the property c), too. Note that $r = 2$ in [10]. However, dealing with general $r \in (1, \infty)$ does not affect the used procedure and the resulting estimates. \square

The next theorem provides an information on the existence of a weak solution to the Stokes problem (1.1)–(1.6).

Theorem 3 (on existence of a weak solution). *Let $1 < r < \infty$, $\mathbf{f} \in \mathbf{W}_0^{-1,r}(\Omega)$, \mathbf{g} be a given velocity profile on Γ_{in} satisfying the assumptions of Lemma 3, and $\mathbf{h} \in \mathbf{W}_{\text{per}}^{-1/r,r}(\Gamma_{\text{out}})$. Then the Stokes problem (1.1)–(1.6) has a unique weak solution \mathbf{u} (in the sense of Definition 1). The functions \mathbf{u} and p (an associated pressure, given by Theorem 2) satisfy*

$$\|\mathbf{u}\|_{1,r} + \|p\|_r \leq c [\|\mathbf{f}\|_{\mathbf{W}_0^{-1,r}} \|\mathbf{g}\|_{s,r;\Gamma_{\text{in}}} + \|\mathbf{h}\|_{\mathbf{W}_{\text{per}}^{-1/r,r}(\Gamma_{\text{out}})}], \quad (4.8)$$

where $c = c(\Omega, \nu)$.

Proof. Note that if \mathbf{g} satisfies the assumptions of Lemma 3 then \mathbf{g} also lies in $\mathbf{W}_{\text{per}}^{1-1/r,r}(\Gamma_{\text{in}})$.

Let \mathbb{F} be a tensor function in $L^r(\Omega)^{2 \times 2}$, provided by Lemma 1 and $\mathbf{F} \in \mathbf{V}_{\sigma}^{-1,r}(\Omega)$ be a functional in $\mathbf{V}_{\sigma}^{-1,r}(\Omega)$, defined by the formula

$$\langle \mathbf{F}, \mathbf{w} \rangle_{(\mathbf{V}_{\sigma}^{-1,r}, \mathbf{V}_{\sigma}^{1,r'})} := - \int_{\Omega} \mathbb{F} : \nabla \mathbf{w} \, dx \quad (4.9)$$

for all $\mathbf{w} \in \mathbf{V}_\sigma^{1,r'}(\Omega)$. The norm of \mathbf{F} in $\mathbf{V}_\sigma^{-1,r}(\Omega)$ satisfies the estimate $\|\mathbf{F}\|_{\mathbf{V}_\sigma^{-1,r}} \leq c \|\mathbb{F}\|_r \leq c \|\mathbf{f}\|_{\mathbf{W}_0^{-1,r}}$. Similarly, let \mathbf{g}_* be the function, given by Lemma 3. Define a functional $\mathbf{G} \in \mathbf{V}_\sigma^{-1,r}(\Omega)$ by the formula

$$\langle \mathbf{G}, \mathbf{w} \rangle_{(\mathbf{V}_\sigma^{-1,r}, \mathbf{V}_\sigma^{1,r'})} := - \int_\Omega \nabla \mathbf{g}_* : \nabla \mathbf{w} \, d\mathbf{x} \quad (4.10)$$

for all $\mathbf{w} \in \mathbf{V}_\sigma^{1,r'}(\Omega)$. Then \mathbf{G} satisfies $\|\mathbf{G}\|_{\mathbf{V}_\sigma^{-1,r}} \leq c \|\nabla \mathbf{g}_*\|_r \leq c \|\mathbf{g}\|_{s,r; \Gamma_{\text{in}}}$. Finally, let \mathbf{H} be a functional in $\mathbf{V}_\sigma^{-1,r}(\Omega)$, defined by the formula

$$\langle \mathbf{H}, \mathbf{w} \rangle_{(\mathbf{V}_\sigma^{-1,r}, \mathbf{V}_\sigma^{1,r'})} := \langle \mathbf{h}, \mathbf{w} \rangle_{(\mathbf{W}_{\text{per}}^{-1/r,r}(\Gamma_{\text{out}}), \mathbf{W}_{\text{per}}^{1-1/r',r'}(\Gamma_{\text{out}}))} \quad (4.11)$$

for all $\mathbf{w} \in \mathbf{V}_\sigma^{1,r'}(\Omega)$. Obviously, \mathbf{H} satisfies the estimate $\|\mathbf{H}\|_{\mathbf{V}_\sigma^{-1,r}} \leq c \|\mathbf{h}\|_{\mathbf{W}^{-1/r,r}(\Gamma_{\text{out}})}$.

Due to Theorem 1, the equation

$$\nu \mathcal{A}_r \mathbf{v} = \mathbf{F} + \nu \mathbf{G} + \mathbf{H} \quad (4.12)$$

has a unique solution $\mathbf{v} \in \mathbf{V}_\sigma^{1,r}(\Omega)$, which satisfies

$$\|\nabla \mathbf{v}\|_r \leq c (\|\mathbf{F}\|_{\mathbf{V}_\sigma^{-1,r}} + \|\mathbf{G}\|_{\mathbf{V}_\sigma^{-1,r}} + \|\mathbf{H}\|_{\mathbf{V}_\sigma^{-1,r}}), \quad (4.13)$$

where $c = c(\nu, \Omega, r)$. The equation (4.12) implies that

$$\nu \int_\Omega \nabla \mathbf{v} : \nabla \mathbf{w} \, d\mathbf{x} = - \int_\Omega \mathbb{F} : \mathbf{w} \, d\mathbf{x} - \nu \int_\Omega \nabla \mathbf{g}_* : \nabla \mathbf{w} \, d\mathbf{x} + \langle \mathbf{h}, \mathbf{w} \rangle_{(\mathbf{W}_{\text{per}}^{-1/r,r}(\Gamma_{\text{out}}), \mathbf{W}_{\text{per}}^{1-1/r',r'}(\Gamma_{\text{out}}))}$$

for all $\mathbf{w} \in \mathbf{V}_\sigma^{1,r'}(\Omega)$. Put $\mathbf{u} := \mathbf{v} + \mathbf{g}_*$. One can now easily verify that \mathbf{u} has all properties, stated in Definition 1, which means that \mathbf{u} is a weak solution to the problem (1.1)–(1.6). The existence of an associated pressure p follows from Theorem 2. The estimate (4.8) follows from (4.13), from the estimates of the norms of the functionals \mathbf{F} , \mathbf{G} and \mathbf{H} in $\mathbf{V}_\sigma^{-1,r}(\Omega)$ and from (4.5) and (4.6). \square

5 Proof of the inequality (3.4)

Recall that in part (b) of the proof of Theorem 1, we assume that $\mathbb{F} \in W^{1,2}(\Omega)^{2 \times 2}$ satisfies (3.1) and the functions $\mathbf{v} \in \mathbf{V}_\sigma^{1,2}(\Omega) \cap \mathbf{W}^{2,2}(\Omega)$ and $p \in W^{1,2}(\Omega)$ satisfy the equation (3.3) a.e. in Ω , the boundary condition

$$\mathbf{v} = \mathbf{0} \quad (5.1)$$

on $\Gamma_{\text{in}} \cup \Gamma_{\text{p}}$ and the condition of periodicity (1.5) on Γ_0 . It follows from [29, Theorem 2] that \mathbf{v} and p also satisfy the conditions of periodicity

$$\frac{\partial \mathbf{v}}{\partial \mathbf{n}}(x_1, x_2 + \tau) = - \frac{\partial \mathbf{v}}{\partial \mathbf{n}}(x_1, x_2) \quad \text{for a.a. } \mathbf{x} \equiv (x_1, x_2) \in \Gamma_0, \quad (5.2)$$

$$p(x_1, x_2 + \tau) = p(x_1, x_2) \quad \text{for a.a. } \mathbf{x} \equiv (x_1, x_2) \in \Gamma_0 \quad (5.3)$$

and the boundary condition

$$(-\nu \nabla \mathbf{v} + p \mathbb{I} - \mathbb{F}) \cdot \mathbf{n} \equiv -\nu \frac{\partial \mathbf{v}}{\partial \mathbf{n}} + p \mathbf{n} - \mathbb{F} \cdot \mathbf{n} = \mathbf{0} \quad \text{a.e. on } \Gamma_{\text{out}}. \quad (5.4)$$

Extending \mathbf{v} , p and \mathbb{F} τ -periodically in the x_2 -direction, we deduce that the extended functions (which we again denote by \mathbf{v} , p and \mathbb{F}) satisfy the equation (3.3) a.e. in

$$\Omega_{\text{ext}} := \Omega \cup \Gamma_0^0 \cup \Gamma_1^0 \cup \{\mathbf{x} = (x_1, x_2) \in \mathbb{R}^2; 0 < x_1 < d, \mathbf{x} \pm \tau \mathbf{e}_2 \in \Omega\}.$$

The extended functions satisfy $\mathbf{v} \in \mathbf{W}^{2,2}(\Omega_{\text{ext}})$, $p \in W^{1,2}(\Omega_{\text{ext}})$ and $\mathbb{F} \in W^{1,2}(\Omega_{\text{ext}})$. Note that

$$\partial\Omega_{\text{ext}} = (\Gamma_{\text{in}})_{\text{ext}} \cup (\Gamma_{\text{out}})_{\text{ext}} \cup (\Gamma_0 - \tau \mathbf{e}_2) \cup (\Gamma_1 + \tau \mathbf{e}_2) \cup (\Gamma_p)_{\text{ext}},$$

where $(\Gamma_{\text{in}})_{\text{ext}}$ is the part of the boundary of Ω_{ext} on γ_{in} , $(\Gamma_{\text{out}})_{\text{ext}}$ is the part of the boundary of Ω_{ext} on γ_{out} and

$$(\Gamma_p)_{\text{ext}} := \Gamma_p \cup (\Gamma_p + \tau \mathbf{e}_2) \cup (\Gamma_p - \tau \mathbf{e}_2).$$

We assume that $r > 2$ throughout this section and we split the derivation of the estimate (3.4) into two parts, in which we obtain the desired estimate in the interior of Ω_{ext} plus a neighborhood of $\Gamma_{\text{in}} \cup \Gamma_p$ (see Lemma 4) and in a neighborhood of Γ_{out} (see Lemma 5).

Lemma 4. *Let Ω' be a sub-domain of Ω_{ext} , such that the distance between $\overline{\Omega'}$ and any of the curves $\Gamma_0 - \tau \mathbf{e}_2$, $\Gamma_1 + \tau \mathbf{e}_2$, $\Gamma_p - \tau \mathbf{e}_2$, $\Gamma_p + \tau \mathbf{e}_2$ and $(\Gamma_{\text{out}})_{\text{ext}}$ is positive. Then \mathbf{v} and p satisfy the estimate*

$$\|\mathbf{v}\|_{1,r;\Omega'} \leq c \|\mathbb{F}\|_r, \quad (5.5)$$

where $c = c(\nu, \Omega, \Omega')$.

Proof. We may assume, without loss of generality, that $\Gamma_p \subset \partial\Omega'$. There exists $\rho > 0$ so small that

$$U_\rho(\Omega') := \{\mathbf{x} = (x_1, x_2) \in \mathbb{R}^2 \setminus P; 0 < x_1 < d, \text{dist}(\mathbf{x}, \Omega') < \rho\}$$

is a subset of Ω_{ext} . (Recall that P is a compact set, see Fig. 1.) There exists an infinitely differentiable function η in Ω_{ext} , such that $\eta = 1$ in Ω' , $\text{supp } \eta \subset U_\rho(\Omega')$ and $\Omega'' := \text{supp } \eta$ is of the class C^2 . Denote $\tilde{\mathbf{v}} := \eta \mathbf{v}$ and $\tilde{p} := \eta p$. The functions $\tilde{\mathbf{v}}$, \tilde{p} represent a strong solution of the problem

$$-\nu \Delta \tilde{\mathbf{v}} + \nabla \tilde{p} = \tilde{\mathbf{f}} \quad \text{in } \Omega'', \quad (5.6)$$

$$\text{div } \tilde{\mathbf{v}} = \tilde{h} \quad \text{in } \Omega'', \quad (5.7)$$

$$\tilde{\mathbf{v}} = \mathbf{0} \quad \text{on } \partial\Omega'', \quad (5.8)$$

where

$$\tilde{\mathbf{f}} := \eta \text{div } \mathbb{F} - 2\nu \nabla \eta \cdot \nabla \mathbf{v} - \nu (\Delta \eta) \mathbf{v} - (\nabla \eta) p \quad \text{and} \quad \tilde{h} := \nabla \eta \cdot \mathbf{v}.$$

Applying Proposition I.2.3 from [30], p. 35, we obtain the estimate

$$\|\tilde{\mathbf{v}}\|_{1,r;\Omega''} \leq c (\|\tilde{\mathbf{f}}\|_{\mathbf{W}^{-1,r}} + \|\tilde{h}\|_r). \quad (5.9)$$

Since $\mathbf{L}^2(\Omega) \hookrightarrow \mathbf{W}^{-1,r}(\Omega)$ and $\mathbf{W}^{1,2}(\Omega) \hookrightarrow \mathbf{L}^r(\Omega)$, we can estimate the terms on the right hand side as follows:

$$\begin{aligned} \|\tilde{\mathbf{f}}\|_{\mathbf{W}^{-1,r}} &\leq c \|\mathbb{F}\|_r + c \|\mathbf{v}\|_r + c \|p\|_{-1,r} \leq \|\mathbb{F}\|_r + c \|\mathbf{v}\|_{1,2} + c \|p\|_2, \\ \|\tilde{h}\|_r &\leq c \|\mathbf{v}\|_r \leq c \|\mathbf{v}\|_{1,2}. \end{aligned}$$

Due to [29, Lemma 1 and estimate (2.5)], we have $\|\mathbf{v}\|_{1,2} + \|p\|_2 \leq c \|\mathbb{F}\|_{\mathbf{V}_\sigma^{-1,2}} \leq c \|\mathbb{F}\|_2 \leq c \|\mathbb{F}\|_r$. Substituting these estimates to (5.9), we obtain: $\|\tilde{\mathbf{v}}\|_{1,r;\Omega''} \leq c \|\mathbb{F}\|_r$. Since $\Omega' \subset \Omega''$ and $\eta = 1$ on Ω' , we obtain (5.5). \square

Lemma 5. *Let Ω' be a sub-domain of Ω_{ext} , such that $\overline{\Omega'}$ has a positive distance from any of the sets $\Gamma_0 - \tau \mathbf{e}_2$, $\Gamma_1 + \tau \mathbf{e}_2$, $(\Gamma_{\text{in}})_{\text{ext}}$ and $(\Gamma_{\text{p}})_{\text{ext}}$. Then \mathbf{v} and p satisfy the estimate*

$$\|\mathbf{v}\|_{1,r;\Omega'} \leq c \|\mathbb{F}\|_r, \quad (5.10)$$

where $c = c(\nu, \Omega, \Omega')$.

Proof. Since $\text{div } \mathbf{v} = 0$, there exists $\varphi \in W^{3,2}(\Omega_{\text{ext}})$, such that $\mathbf{v} \equiv (v_1, v_2) = \nabla^\perp \varphi \equiv (-\partial_2 \varphi, \partial_1 \varphi)$. As $\mathbf{v} = \mathbf{0}$ on $(\Gamma_{\text{in}})_{\text{ext}}$, the function φ satisfies $\nabla^\perp \varphi = \mathbf{0}$ on $(\Gamma_{\text{in}})_{\text{ext}}$. Since φ is determined uniquely up to an additive constant, we can choose φ so that

$$\varphi = 0 \quad \text{and} \quad \nabla \varphi = \mathbf{0} \quad \text{on } (\Gamma_{\text{in}})_{\text{ext}}. \quad (5.11)$$

Put

$$\mathbb{Z} := -\nu \nabla \mathbf{v} + p \mathbb{I} - \mathbb{F}.$$

Thus, if we denote by Z_{ij} ($i, j = 1, 2$) the entries of \mathbb{Z} and by F_{ij} ($i, j = 1, 2$) the entries of \mathbb{F} , we can write this formula in the form

$$\begin{aligned} \begin{pmatrix} Z_{11} & Z_{12} \\ Z_{21} & Z_{22} \end{pmatrix} &= -\nu \begin{pmatrix} \partial_1 v_1 & \partial_2 v_1 \\ \partial_1 v_2 & \partial_2 v_2 \end{pmatrix} + \begin{pmatrix} p & 0 \\ 0 & p \end{pmatrix} - \begin{pmatrix} F_{11} & F_{12} \\ F_{21} & F_{22} \end{pmatrix} \\ &= -\nu \begin{pmatrix} -\partial_{12} \varphi & -\partial_{22} \varphi \\ \partial_{11} \varphi & \partial_{21} \varphi \end{pmatrix} + \begin{pmatrix} p & 0 \\ 0 & p \end{pmatrix} - \begin{pmatrix} F_{11} & F_{12} \\ F_{21} & F_{22} \end{pmatrix}. \end{aligned}$$

Equation (3.3) says that $\text{div } \mathbb{Z} = \mathbf{0}$, which means that $\partial_j Z_{ij} = 0$ for $i = 1, 2$. Hence there exist functions $\psi_1, \psi_2 \in W^{2,2}(\Omega_{\text{ext}})$, such that $Z_{11} = -\partial_2 \psi_1$, $Z_{12} = \partial_1 \psi_1$, $Z_{21} = -\partial_2 \psi_2$, $Z_{22} = \partial_1 \psi_2$. Thus, we obtain the equation

$$\begin{pmatrix} -\partial_2 \psi_1 & \partial_1 \psi_1 \\ -\partial_2 \psi_2 & \partial_1 \psi_2 \end{pmatrix} = -\nu \begin{pmatrix} -\partial_{12} \varphi & -\partial_{22} \varphi \\ \partial_{11} \varphi & \partial_{21} \varphi \end{pmatrix} + \begin{pmatrix} p & 0 \\ 0 & p \end{pmatrix} - \begin{pmatrix} F_{11} & F_{12} \\ F_{21} & F_{22} \end{pmatrix}. \quad (5.12)$$

This tensorial equation can also be considered to be a system of four equations for four unknowns: φ , p , ψ_1 and ψ_2 . Since $\mathbf{n} = \mathbf{e}_1$ on Γ_{out} , the boundary condition (5.4) yields $Z_{11} = Z_{12} = 0$ on Γ_{out} . This means that $\partial_2 \psi_1 = \partial_2 \psi_2 = 0$ on Γ_{out} , which implies that ψ_1 and ψ_2 are constant on Γ_{out} . Let us denote the constants by k_1 and k_2 . Thus, we obtain the boundary conditions

$$\psi_1 = k_1, \quad \psi_2 = k_2 \quad \text{on } (\Gamma_{\text{out}})_{\text{ext}}. \quad (5.13)$$

Let us concretely choose k_1 and k_2 so that

$$\int_{\Omega} \psi_1 \, d\mathbf{x} = \int_{\Omega} \psi_2 \, d\mathbf{x} = 0. \quad (5.14)$$

Later on in this proof, we shall need estimates of $\|\psi_1\|_r$ and $\|\psi_2\|_r$. Let us begin with $\|\psi_1\|_r$:

$$\|\psi_1\|_r = \sup_{w \in L^{r'}(\Omega); w \neq 0} \|w\|_{r'}^{-1} \left| \int_{\Omega} \psi_1 w \, d\mathbf{x} \right|.$$

The function $w \in L^{r'}(\Omega)$ can be written in the form $w = \overline{w} + w'$, where $\overline{w} := \int_{\Omega} w \, d\mathbf{x}$ and $\int_{\Omega} w' \, d\mathbf{x} = 0$. Then, obviously, $\|w'\|_{r'} \leq c \|w\|_{r'}$, where $c = c(\Omega)$. Moreover, there exists $\mathbf{z} \in \mathbf{W}_0^{1,r}(\Omega)$, such that $\text{div } \mathbf{z} = w'$ and $\|\mathbf{z}\|_{1,r'} \leq c \|w'\|_{r'}$, see [11, Theorem III.3.3]. Hence

$$\|\psi_1\|_r \leq c \sup_{w \in L^{r'}(\Omega); w \neq 0} \|w'\|_{r'}^{-1} \left| \overline{w} \int_{\Omega} \psi_1 \, d\mathbf{x} + \int_{\Omega} \psi_1 w' \, d\mathbf{x} \right|$$

$$\begin{aligned}
& \leq c \sup_{\mathbf{z} \in \mathbf{W}_0^{1,r'}(\Omega); \mathbf{z} \neq 0} \|\nabla \mathbf{z}\|_{r'}^{-1} \left| \int_{\Omega} \psi_1 \operatorname{div} \mathbf{z} \, d\mathbf{x} \right| \\
& = c \sup_{\mathbf{z} \in \mathbf{W}_0^{1,r'}(\Omega); \mathbf{z} \neq 0} \|\nabla \mathbf{z}\|_{r'}^{-1} \left| \int_{\Omega} \nabla \psi_1 \cdot \mathbf{z} \, d\mathbf{x} \right| \\
& = c \|\nabla \psi_1\|_{\mathbf{W}^{-1,r}} \leq c \|\mathbb{Z}\|_{\mathbf{W}^{-1,r}} = c \|\nu \nabla \mathbf{v} + p \mathbb{I} - \mathbb{F}\|_{\mathbf{W}^{-1,r}} \\
& \leq c (\|\mathbf{v}\|_r + \|p\|_{\mathbf{W}^{-1,r}} + \|\mathbb{F}\|_r). \tag{5.15}
\end{aligned}$$

As $\mathbf{W}_0^{1,r'}(\Omega) \hookrightarrow \mathbf{L}^2(\Omega)$, we also have $\mathbf{L}^2(\Omega) \hookrightarrow \mathbf{W}^{-1,r}(\Omega)$. Hence $\|p\|_{\mathbf{W}^{-1,r}} \leq c \|p\|_2$. Furthermore, as $\mathbf{W}^{1,2}(\Omega) \hookrightarrow \mathbf{L}^r(\Omega)$, we also have $\|\mathbf{v}\|_r \leq c \|\mathbf{v}\|_{1,2}$. Thus, (5.15) yields

$$\|\psi_1\|_r \leq c (\|\mathbf{v}\|_{1,2} + \|p\|_2 + \|\mathbb{F}\|_r). \tag{5.16}$$

Due to [29, Lemma 1 and estimate (2.5)], the right hand side is less than or equal to $c (\|\mathbb{F}\|_{\mathbf{V}_{\sigma}^{-1,2}} + \|\mathbb{F}\|_r) \leq c (\|\mathbb{F}\|_2 + \|\mathbb{F}\|_r)$, which is less than or equal to $c \|\mathbb{F}\|_r$. Substituting this to (5.16) and taking into account that ψ_2 can be estimated in the same way, we finally obtain

$$\|\psi_1\|_r + \|\psi_2\|_r \leq c \|\mathbb{F}\|_r. \tag{5.17}$$

In order to obtain the desired estimate (3.4), we will apply Theorem 10.2 from the paper [1] by S. Agmon, A. Douglis and L. Nirenberg. This theorem, however, concerns a boundary value problem in a half-space, which means a half-plane in the case of a problem in 2D. In order to transform the problem (5.12), (5.13) to a problem in the half-plane $x_1 < d$, we apply the cut-off function technique; there exists $\rho > 0$ so small that

$$U_{\rho}(\Omega') := \{\mathbf{x} = (x_1, x_2) \in \mathbb{R}^2; x_1 < d, \operatorname{dist}(\mathbf{x}, \Omega') < \rho\}$$

is a subset of Ω_{ext} . Let η be an infinitely differentiable function in Ω_{ext} , such that $\eta = 1$ in Ω' and $\eta = 0$ in $\Omega_{\text{ext}} \setminus U_{\rho}(\Omega')$. Multiplying (5.12) by η , and denoting $\tilde{\varphi} := \eta\varphi$, $\tilde{\psi}_1 := \eta\psi_1$, $\tilde{\psi}_2 := \eta\psi_2$ and $\tilde{p} := \eta p$, we get

$$\begin{aligned}
& \begin{pmatrix} -\partial_2 \tilde{\psi}_1 & \partial_1 \tilde{\psi}_1 \\ -\partial_2 \tilde{\psi}_2 & \partial_1 \tilde{\psi}_2 \end{pmatrix} + \nu \begin{pmatrix} -\partial_{12} \tilde{\varphi} & -\partial_{22} \tilde{\varphi} \\ \partial_{11} \tilde{\varphi} & \partial_{21} \tilde{\varphi} \end{pmatrix} - \begin{pmatrix} \tilde{p} & 0 \\ 0 & \tilde{p} \end{pmatrix} \\
& = - \begin{pmatrix} \eta F_{11} & \eta F_{12} \\ \eta F_{21} & \eta F_{22} \end{pmatrix} + \begin{pmatrix} -(\partial_2 \eta) \psi_1 & (\partial_1 \eta) \psi_1 \\ -(\partial_2 \eta) \psi_2 & (\partial_1 \eta) \psi_2 \end{pmatrix} + \begin{pmatrix} -(\partial_{12} \eta) \varphi & -(\partial_{22} \eta) \varphi \\ (\partial_{11} \eta) \varphi & (\partial_{21} \eta) \varphi \end{pmatrix} \\
& + \begin{pmatrix} -(\partial_1 \eta)(\partial_2 \varphi) - (\partial_2 \eta)(\partial_1 \varphi) & -2(\partial_2 \eta)(\partial_2 \varphi) \\ 2(\partial_1 \eta)(\partial_1 \varphi) & (\partial_1 \eta)(\partial_2 \varphi) + (\partial_2 \eta)(\partial_1 \varphi) \end{pmatrix}. \tag{5.18}
\end{aligned}$$

Since η is supported in the closure of $U_{\rho}(\Omega')$, we can treat (5.18) as a system of four equations in the half-plane $x_1 < d$ for the unknowns $\tilde{\varphi}$, $\tilde{\psi}_1$, $\tilde{\psi}_2$ and \tilde{p} with the boundary conditions

$$\tilde{\psi}_1 = \eta k_1, \quad \tilde{\psi}_2 = \eta k_2 \quad \text{on } \gamma_{\text{out}}. \tag{5.19}$$

In order to simplify the equation (5.18) and to use a notation, consistent with [1], we eliminate \tilde{p} by subtracting the two corresponding equations and denote $u_1 := \tilde{\varphi}$, $u_2 := \tilde{\psi}_1$ and $u_3 := \tilde{\psi}_2$. Then the problem (5.18), (5.19) reduces to the system of three equations

$$2\nu \partial_{12} u_1 + \partial_2 u_2 + \partial_1 u_3 = -\eta(F_{22} - F_{11}) + (\partial_1 \eta) \psi_2 + (\partial_2 \eta) \psi_1 + 2(\partial_{12} \eta) \varphi$$

$$+ 2(\partial_1 \eta)(\partial_2 \varphi) + 2(\partial_2 \eta)(\partial_1 \varphi), \quad (5.20)$$

$$\nu \partial_{11} u_1 - \partial_2 u_3 = -\eta F_{21} - (\partial_2 \eta) \psi_2 + (\partial_{11} \eta) \varphi + 2(\partial_1 \eta)(\partial_1 \varphi), \quad (5.21)$$

$$\nu \partial_{22} u_1 - \partial_1 u_2 = -\eta F_{12} - (\partial_1 \eta) \psi_1 + (\partial_{22} \eta) \varphi + 2(\partial_2 \eta)(\partial_2 \varphi) \quad (5.22)$$

with the boundary conditions

$$u_2 = \eta k_1, \quad u_3 = \eta k_2 \quad \text{on } \gamma_{\text{out}}. \quad (5.23)$$

Obviously, the leading differential operator in (5.20)–(5.22) is

$$\mathcal{L}(\partial_1, \partial_2) := \begin{pmatrix} 2\nu \partial_1 \partial_2, & \partial_2, & \partial_1, \\ \nu \partial_1^2, & 0, & -\partial_2, \\ \nu \partial_2^2, & -\partial_1, & 0 \end{pmatrix}.$$

The determinant of $\mathcal{L}(\xi_1, \xi_2)$ equals $-\nu(\xi_1^2 + \xi_2^2)^2$, which is a polynomial of degree $2m = 4$. With this information, one can verify that the system (5.20)–(5.22) is “uniformly elliptic” of order 4 in the sense of Agmon–Douglis–Nirenberg, which means that it satisfies the conditions (1.1)–(1.7) from [1], pp. 38, 39, and the so called “supplementary condition”, see [1, p. 39]. Moreover, the boundary conditions (5.23), the number of which is $m = 2$, satisfy condition (2.2) (see [1, p. 42]) and have all required properties of the so called “complementing boundary conditions”, formulated in [1, p. 42–43]. The verification is elementary, however technical, hence we do not provide the details here.

Note, that one can also find the description and explanation of the conditions that enable one to call a considered system “uniformly elliptic” and considered boundary conditions to be “complementing” in [4, Appendix D].

Denote by \mathbb{R}_{d-}^2 the half-plane $x_1 < d$.

The inclusions $u_1 \in W^{3,2}(\mathbb{R}_{d-}^2) \hookrightarrow W^{2,r}(\mathbb{R}_{d-}^2)$, $u_2, u_3 \in W^{2,2}(\mathbb{R}_{d-}^2) \hookrightarrow W^{1,r}(\mathbb{R}_{d-}^2)$ and the verification of the aforementioned conditions from [1] enable us to apply Theorem 10.2 from [1], which yields the estimate

$$\begin{aligned} & \|u_1\|_{2,r;\mathbb{R}_{d-}^2} + \|u_2\|_{1,r;\mathbb{R}_{d-}^2} + \|u_3\|_{1,r;\mathbb{R}_{d-}^2} \\ & \leq c \sum_{i=1}^3 \|\tilde{f}_i\|_{r;\mathbb{R}_{d-}^2} + c (\|\eta k_1\|_{1-1/r,r;\gamma_{\text{out}}} + \|\eta k_2\|_{1-1/r,r;\gamma_{\text{out}}}) \\ & \leq c \sum_{i=1}^3 \|\tilde{f}_i\|_{r;\mathbb{R}_{d-}^2} + c (|k_1| + |k_2|), \end{aligned} \quad (5.24)$$

where \tilde{f}_1 , \tilde{f}_2 and \tilde{f}_3 denote the right hand sides of equations (5.20), (5.21) and (5.22), respectively, and c is independent of u_i , \tilde{f}_i ($i = 1, 2, 3$) and k_1, k_2 . The first term on the right hand side of (5.24) can be estimated by means of the inequalities

$$\sum_{i=1}^3 \|\tilde{f}_i\|_{r;\mathbb{R}_{d-}^2} \leq c (\|\mathbb{F}\|_r + \|\psi_1\|_r + \|\psi_2\|_r + \|\varphi\|_{1,r}). \quad (5.25)$$

The norms $\|\psi_1\|_r$ and $\|\psi_2\|_r$ can be estimated by means of (5.17). The norm $\|\varphi\|_{1,r}$ can be estimated as follows:

$$\|\varphi\|_{1,r} \leq c \|\mathbf{v}\|_r \leq c \|\mathbf{v}\|_{1,2} \leq c \|\mathbf{F}\|_{\mathbf{V}_\sigma^{-1,2}} \leq c \|\mathbb{F}\|_2 \leq c \|\mathbb{F}\|_r. \quad (5.26)$$

(Here, the first inequality holds due to the definition of φ and (5.11), the second inequality follows from the continuous imbedding $\mathbf{W}^{1,2}(\Omega) \hookrightarrow \mathbf{L}^r(\Omega)$, the third inequality follows from [29, Lemma 1], the fourth inequality follows from the definition of \mathbf{F} and the last inequality holds, because $r > 2$.) The second term on the right hand side of (5.24) can be estimated by means of these inequalities:

$$\begin{aligned}
|k_1| + |k_2| &= |\Omega|^{-1} \left| \int_{\Omega} (\psi_1 - k_1) \, d\mathbf{x} \right| + |\Omega|^{-1} \left| \int_{\Omega} (\psi_2 - k_2) \, d\mathbf{x} \right| \\
&\leq c \|\nabla(\psi_1 - k_1)\|_2 + c \|\nabla(\psi_2 - k_2)\|_2 \leq c \|\mathbb{Z}\|_2 \leq c (\|\nabla \mathbf{v}\|_2 + \|p\|_2 + \|\mathbb{F}\|_2) \\
&\leq c \|\mathbf{F}\|_{\mathbf{V}_{\sigma}^{-1,2}} + c \|\mathbb{F}\|_2 \leq c \|\mathbb{F}\|_2 \leq c \|\mathbb{F}\|_r.
\end{aligned} \tag{5.27}$$

(The inequalities $\left| \int_{\Omega} (\psi_j - k_j) \, d\mathbf{x} \right| \leq c \|\nabla(\psi_j - k_j)\|_2$ (for $j = 1, 2$) hold due to (5.13). The first inequality on the last line follows from Lemma 1 and estimate (2.5) in [29] and the last estimate follows from the condition $r > 2$.) Estimates (5.24)–(5.27) now yield

$$\|\varphi\|_{2,r;\Omega'} \leq \|\tilde{\varphi}\|_{2,r;\mathbb{R}_{d-}^2} = \|u_1\|_{2,r;\mathbb{R}_{d-}^2} \leq c \|\mathbb{F}\|_r$$

Since $\|\mathbf{v}\|_{1,r;\Omega'} \leq c \|\varphi\|_{2,r;\Omega'}$, we obtain inequality (5.10). \square

Now the estimate (3.4) follows easily from Lemmas 4 and 5.

Acknowledgement. This work was supported by European Regional Development Fund-Project “Center for Advanced Applied Science” No. CZ.02.1.01/0.0/0.0/16_019/0000778.

References

- [1] S. Agmon, A. Douglis, L. Nirenberg: Estimates near the boundary for solutions of elliptic partial differential equations satisfying general boundary conditions II. *Comm. Pure and Appl. Math.* **XVII** (1964), 35–92.
- [2] P. Acevedo, Ch. Amrouche, C. Conca, A. Ghosh: Stokes and Navier–Stokes equations with Navier boundary conditions. *C. R. Acad. Sci. Paris, Ser. I* **357** (2019), 115–119.
- [3] Ch. Amrouche, N. El Houda Seloula: On the Stokes equation with Navier–type boundary conditions. *Diff. Equations & Appl.* **3** (2011), 4, 581–607.
- [4] P. B. Bochev, M. D. Gunzburger: *Least-Squares Finite Element Methods*. Springer 2009.
- [5] C. H. Bruneau, P. Fabrie: New efficient boundary conditions for incompressible Navier–Stokes equations: A well-posedness result. *Math. Modelling and Num. Analysis* **30** (1996), 7, 815–840.
- [6] L. Cattabriga: Su un problema al contorno relativo al sistema il equazioni di Stokes. *Rend. Mat. Sem. Univ. Padova* **31** (1961), 308–340.
- [7] V. Dolejší, M. Feistauer, J. Felcman: Numerical simulation of compressible viscous flow through cascades of profiles. *Z. für Angew. Math. Mech.* **76** (1996), 301–304.
- [8] M. Feistauer, T. Neustupa: On some aspects of analysis of incompressible flow through cascades of profiles. *Operator Theory, Advances and Applications*, Vol. 147, Birkhauser, Basel 2004, 257–276.
- [9] M. Feistauer, T. Neustupa: On non-stationary viscous incompressible flow through a cascade of profiles. *Math. Meth. Appl. Sci.* **29** (2006), No. 16, 1907–1941.

- [10] M. Feistauer, T. Neustupa: On the existence of a weak solution of viscous incompressible flow past a cascade of profiles with an arbitrarily large inflow. *J. Math. Fluid Mech.* **15** (2013), 701–715.
- [11] G. P. Galdi: *An Introduction to the Mathematical Theory of the Navier–Stokes Equations, Steady State Problems*. Springer–Verlag, New York–Berlin–Heidelberg 2011.
- [12] M. Geissert, H. Heck, M. Hieber: On the equation $\operatorname{div} u = g$ and Bogovskii’s operator in Sobolev spaces of negative order. In *Partial differential equations and functional analysis, Oper. Theory Adv. Appl.* **168**, Birkhäuser, Basel 2006, pp. 113–121.
- [13] R. Glowinski: *Numerical Methods for Nonlinear Variational Problems*. Springer–Verlag, New York–Berlin–Heidelberg–Tokyo, 1984.
- [14] H. P. Heinig, A. Kufner, L.-E. Persson: On some fractional order Hardy inequalities. *J. of Inequal. & Appl.* **1** (1997), No. 1, 25–46.
- [15] J. G. Heywood, R. Rannacher and S. Turek: Artificial boundaries and flux and pressure conditions for the incompressible Navier–Stokes equations. *Int. J. for Numerical Methods in Fluids* **22** (1996), 325–352.
- [16] T. Kato: *Perturbation Theory for Linear Operators*. Springer–Verlag, Berlin–Heidelberg–New York, 1966.
- [17] K. Kozel, P. Louda, J. Příhoda: Numerical solution of turbulent flow in a turbine cascade. *Proc. Appl. Math. Mech.* **6** (2006), 743–744.
- [18] S. Kračmar, J. Neustupa: Modelling of flows of a viscous incompressible fluid through a channel by means of variational inequalities. *ZAMM* **74** (1994), No. 6, 637–639.
- [19] S. Kračmar, J. Neustupa: A weak solvability of a steady variational inequality of the Navier–Stokes type with mixed boundary conditions. *Nonlinear Analysis* **47** (2001), No. 6, 4169–4180.
- [20] S. Kračmar, J. Neustupa: Modeling of the unsteady flow through a channel with an artificial outflow condition by the Navier–Stokes variational inequality. *Math. Nachrichten* **291** (2018), Issue 11–12, 1–14.
- [21] P. Kučera, Z. Skalák, Solutions to the Navier–Stokes equations with mixed boundary conditions. *Acta Appl. Math.* **54** (1998), No. 3, 275–288.
- [22] P. Kučera: Basic properties of the non-steady Navier–Stokes equations with mixed boundary conditions in a bounded domain. *Ann. Univ. Ferrara* **55** (2009), 289–308.
- [23] P. Kučera, M. Beneš: Solution to the Navier–Stokes equations with mixed boundary conditions in two-dimensional bounded domains. *Math. Nachr.* **289** (2016), No. 2–3, 194–212.
- [24] A. Kufner, O. John, S. Fučík: *Function Spaces*. Noordhoff International Publishing, Leyden 1977.
- [25] J. L. Lions, E. Magenes: *Problèmes aux limites non homogènes et applications*. Dunod, Paris 1968.
- [26] T. Neustupa: Question of existence and uniqueness of solution for Navier–Stokes Equation with linear “do-nothing” type boundary condition on the outflow. *Lecture Notes in Computer Science* **5434** (2009), 431–438.
- [27] T. Neustupa: The analysis of stationary viscous incompressible flow through a rotating radial blade machine, existence of a weak solution. *Applied Math. and Computation* **219** (2012), 3316–3322.

- [28] T. Neustupa: A steady flow through a plane cascade of profiles with an arbitrarily large inflow: the mathematical model, existence of a weak solution. *Applied Math. and Computation* **272** (2016) 687–691.
- [29] T. Neustupa: The maximum regularity property of the steady Stokes problem associated with a flow through a profile cascade. Re-submitted after revision, see also <https://arxiv.org/abs/2006.15651>
- [30] R. Temam: *Navier–Stokes Equations*. North–Holland, Amsterdam–New York–Oxford 1977.
- [31] H. Sohr: *The Navier–Stokes equations. The Elementary Functional Analytic Approach*. Birkhäuser Verlag, Basel–Boston–Berlin 2001.

Author’s address: Tomáš Neustupa
 Czech Technical University
 Faculty of Mechanical Engineering
 Department of Technical Mathematics
 Karlovo nám. 13, 121 35 Praha 2
 Czech Republic
 e-mail: tomas.neustupa@fs.cvut.cz