

## ON KHOVANOV HOMOLOGY OF QUASI-ALTERNATING LINKS

KHALED QAZAQZEH AND NAFAA CHBILI

**ABSTRACT.** We prove that the length of any gap in the differential grading of the Khovanov homology of any quasi-alternating link is one. As a consequence, we obtain that the length of any gap in the Jones polynomial of any such link is one. This establishes a weaker version of Conjecture 2.3 in [5]. Moreover, we obtain a lower bound for the determinant of any such link in terms of the breadth of its Jones polynomial. This establishes a weaker version of Conjecture 3.8 in [16]. The main tool in obtaining this result is proving the Knight Move Conjecture [2] for the class of quasi-alternating links.

## 1. INTRODUCTION

The class of alternating links has played an important role in the development of knot theory since its early age. In particular, the study of their Jones polynomials led to the solution of long lasting conjectures in knot theory. Thistlethwaite [21] proved that the Jones polynomial of any prime alternating link, which is not a  $(2, p)$ -torus link, is alternating and has no gaps. Moreover, the breadth of the Jones polynomial of any alternating link is equal to its crossing number. Alternating links are also known to have simple Khovanov and link Floer homologies. Indeed, the Khovanov homology of a given alternating link  $L$  is entirely determined by its signature  $\sigma_L$  and its Jones polynomial  $V_L(t)$ . Similarly, the link Floer homology of any alternating link  $L$  is entirely determined by its signature  $\sigma_L$  and its Alexander polynomial  $\Delta_L(t)$ . Furthermore, Ozsváth and Szabó [15] studied the Heegaard Floer Homology of the branched double-cover of alternating links and proved that this homology is determined by the determinant of the link,  $\det(L)$ . This homological property extends to a larger family of links called quasi-alternating. Unlike alternating links which admit a simple diagrammatic definition, quasi-alternating links are defined recursively as follows:

**Definition 1.1.** The set  $\mathcal{Q}$  of quasi-alternating links is the smallest set satisfying the following properties:

- The unknot belongs to  $\mathcal{Q}$ .
- If  $L$  is a link with a diagram  $D$  containing a crossing  $c$  such that
  - (1) both smoothings of the diagram  $D$  at the crossing  $c$ ,  $L_0$  and  $L_1$  as in Figure 1 belong to  $\mathcal{Q}$ , and
  - (2)  $\det(L_0), \det(L_1) \geq 1$ ,
  - (3)  $\det(L) = \det(L_0) + \det(L_1)$ ; then  $L$  is in  $\mathcal{Q}$  and in this case we say  $L$  is quasi-alternating at the crossing  $c$  with quasi-alternating diagram  $D$ .

It is almost impossible to use the definition above to determine whether a given link is not quasi-alternating. On the other hand, several obstructions for a link to be quasi-alternating have been introduced through the past two decades. Many of these have been established for alternating links first, then extended to this new class of links. Some of these main obstructions are listed here.

- (1) the branched double-cover of any quasi-alternating link is an  $L$ -space [15, Proposition. 3.3];

---

*Date:* 18/09/2020.

*Key words and phrases.* quasi-alternating, Khovanov homology, Jones polynomial, breadth.




FIGURE 1. The link diagram  $L$  at the crossing  $c$  and its smoothings  $L_0$  and  $L_1$  respectively.

- (2) the space of branched double-cover of any quasi-alternating link bounds a negative definite 4-manifold  $W$  with  $H_1(W) = 0$  [15, Proof of Lemma. 3.6];
- (3) the  $\mathbb{Z}/2\mathbb{Z}$  knot Floer homology group of any quasi-alternating link is  $\sigma$ -thin [13, Theorem. 2];
- (4) the reduced ordinary Khovanov homology group of any quasi-alternating link is  $\sigma$ -thin [13, Theorem. 1];
- (5) the reduced odd Khovanov homology group of any quasi-alternating link is  $\sigma$ -thin [14, Remark after Proposition. 5.2];
- (6) the determinant of any quasi-alternating link is bigger than the degree of its  $Q$ -polynomial [16, Theorem 2.2]. This inequality was sharpened to the determinant minus one is bigger than or equal to the degree of the  $Q$ -polynomial with equality holds only for  $(2, n)$ -torus links [20, Theorem 1.1].

The main purpose of this paper is to introduce a new and simple obstruction for a link to be quasi-alternating. More precisely, we prove that if the differential grading of the Khovanov homology of any quasi-alternating link has a gap, then the length of this gap is one. As an immediate application, we obtain that any gap in the Jones polynomial of any quasi-alternating link has length one. Moreover, a lower bound for the determinant of any such link is obtained in terms of the breadth of its Jones polynomial. The main result of this paper totally relies on establishing the Knight Move Conjecture for quasi-alternating links.

Consequently, we prove that certain links are not quasi-alternating as the differential grading of their Khovanov homology has a gap of length bigger than one. On the other hand, we show that some subclasses of quasi-alternating links have no gaps in the differential grading of their Khovanov homology. This leads us to suggest Conjecture 4.11 that implies both Conjecture 2.3 in [5] and Conjecture 3.8 in [16].

This paper is organized as follows. In Section 2, we introduce some background and notations needed for the rest of the paper. In Section 3, we give the proof of our main result. Finally, Section 4 will be devoted to give some applications and consequences of our main result.

## 2. BACKGROUND AND NOTATIONS

In this section, we introduce some notations and definitions, and we review some properties of Khovanov homology needed in the rest of this paper. Without loss of generality, we assume that the three unoriented links  $L, L_0$  and  $L_1$  are given according to the scheme in Figure 1. For an oriented link  $L$ , we always assume that we smooth a positive crossing according to Figure 2. All the results of this paper discuss this case. Similar results can be obtained in the case of a negative crossing by taking the mirror image of  $L$ .

**2.1. The Jones polynomial.** The Jones polynomial  $V_L(t)$  is an invariant of oriented links. It is a Laurent polynomial with integral coefficients that can be defined in several ways. In this subsection, we shall briefly recall the definition of this polynomial in terms of the Kauffman bracket and review some of its properties needed in the sequel.

**Definition 2.1.** The Kauffman bracket polynomial is a function from the set of unoriented link diagrams in the oriented plane to the ring of Laurent polynomials with integer coefficients in an indeterminate  $A$ . It maps a link  $L$  to  $\langle L \rangle \in \mathbb{Z}[A^{-1}, A]$  and it is defined by the following relations:

- (1)  $\langle \bigcirc \rangle = 1$ ,
- (2)  $\langle \bigcirc \cup L \rangle = (-A^{-2} - A^2) \langle L \rangle$ ,
- (3)  $\langle L \rangle = A \langle L_0 \rangle + A^{-1} \langle L_1 \rangle$ ,

where  $\bigcirc$ , in relation 2 above, denotes a trivial circle disjoint from the rest of the link, and  $L, L_0$ , and  $L_1$  represent three unoriented links which are identical everywhere except in a small region where they are as indicated in Figure 1.

Given an oriented link diagram  $L$ , we let  $x(L)$  to denote the number of negative crossings and  $y(L)$  to denote the number of positive crossings in  $L$ , see Figure 2. We define the writhe of  $L$  as the integer  $w(L) = y(L) - x(L)$ .

**Definition 2.2.** The Jones polynomial  $V_L(t)$  of an oriented link  $L$  is the Laurent polynomial in  $t^{\pm 1/2}$  with integer coefficients defined by

$$V_L(t) = ((-A)^{-3w(L)} \langle L \rangle)_{t^{1/2}=A^{-2}} \in \mathbb{Z}[t^{-1/2}, t^{1/2}],$$

where  $\langle L \rangle$  is the Kauffman bracket of the unoriented link obtained from  $L$  by ignoring the orientation.




FIGURE 2. Negative and positive crossings respectively.

**2.2. Khovanov Invariant.** This link invariant is a bigraded cohomology theory, with rational coefficients, that was first introduced by Khovanov in [9]. For an oriented link  $L$ , we denote the  $i$ -th homology group of the complex  $\overline{\mathcal{C}}(L)$  and  $\mathcal{C}(L)$  by  $\overline{\mathcal{H}}^i(L)$  and  $\mathcal{H}^i(L)$ , respectively. Also, we denote the  $j$ -th graded component of  $\overline{\mathcal{H}}^i(L)$ , and  $\mathcal{H}^i(L)$  by  $\overline{\mathcal{H}}^{i,j}(L)$ , and  $\mathcal{H}^{i,j}(L)$ , respectively. Therefore, we have

$$\overline{\mathcal{H}}^i(L) = \oplus_{j \in \mathbb{Z}} \overline{\mathcal{H}}^{i,j}(L), \text{ and } \mathcal{H}^i(L) = \oplus_{j \in \mathbb{Z}} \mathcal{H}^{i,j}(L).$$

The homology groups  $\overline{\mathcal{H}}(L)$  and  $\mathcal{H}(L)$  are isomorphic up to some shifts. More precisely

$$(1) \quad \mathcal{H}^{i,j}(L) = \overline{\mathcal{H}}^{i+x(L), j+2x(L)-y(L)}(L),$$

where  $x(L)$  and  $y(L)$  are as defined above. The graded Euler characteristic of this invariant is equal to the normalized Jones polynomial. More precisely,

$$(2) \quad \sum_{i,j \in \mathbb{Z}} (-1)^i q^j \dim \mathcal{H}^{i,j}(L) = (q^{-1} + q) V(L)_{t^{1/2}=-q}.$$

It is worth mentioning here that the graded Euler characteristic is indeed the evaluation, at  $t = -1$ , of the two-variable Khovanov polynomial invariant of the link defined first in [2] as follows:

$$(3) \quad Kh(L)(t, q) = \sum_{i,j \in \mathbb{Z}} t^i q^j \dim \mathcal{H}^{i,j}(L).$$

**2.3. Exact Sequence.** It is clear that  $\overline{\mathcal{C}}(L_0)$  and  $\overline{\mathcal{C}}(L_1)[-1]\{-1\}$  are subcomplexes of  $\overline{\mathcal{C}}(L)$  and form a short exact sequence

$$0 \rightarrow \overline{\mathcal{C}}(L_1)[-1]\{-1\} \rightarrow \overline{\mathcal{C}}(L) \rightarrow \overline{\mathcal{C}}(L_0) \rightarrow 0,$$

with degree preserving maps. This induces the following exact sequences:

$$(4) \quad \cdots \rightarrow \overline{\mathcal{H}}^{i-1,j}(L_0) \xrightarrow{\delta} \overline{\mathcal{H}}^{i,j}(L_1)[-1]\{-1\} \rightarrow \overline{\mathcal{H}}^{i,j}(L) \rightarrow \overline{\mathcal{H}}^{i,j}(L_0) \xrightarrow{\delta} \overline{\mathcal{H}}^{i+1,j}(L_1)[-1]\{-1\} \rightarrow \cdots$$

Consequently, we obtain the following long exact sequences in homology as a result of the facts that  $x(L_0) = x(L)$ ,  $y(L_0) = y(L) - 1$ ,  $x(L_1) = x(L) + e$  and  $y(L_1) = y(L) - e - 1$ :

$$(5) \quad \cdots \rightarrow \mathcal{H}^{i-1,j-1}(L_0) \xrightarrow{\delta} \mathcal{H}^{i-e-1,j-3e-2}(L_1) \rightarrow \mathcal{H}^{i,j}(L) \rightarrow \mathcal{H}^{i,j-1}(L_0) \xrightarrow{\delta} \mathcal{H}^{i-e,j-3e-2}(L_1) \rightarrow \cdots,$$

where  $e$  denotes the difference between the number of negative crossings in  $L_1$  and the number of such crossings in  $L$ .

We recall from [10] that a link  $L$  is said to be *H-thin* if its Khovanov homology  $\mathcal{H}(L)$  is supported on the two diagonals that satisfy  $j - 2i = -\sigma(L) \pm 1$ . It has been proved in [11] that all alternating links are *H-thin*. This fact has been later generalized to quasi-alternating links in [13, Theorem 1].

**Remark 2.3.** The result of [13, Theorem 1] shows that the reduced Khovanov homology of any quasi-alternating link is  $\sigma$ -thin over  $\mathbb{Z}$ . Indeed, this implies that the unreduced rational Khovanov homology that we consider in this paper is *H-thin*.

**2.4. Lee Invariant.** The Lee invariant is a variant of the rational Khovanov homology obtained from the same underlying complex with a differential  $\Phi$  of degree difference  $(1, 4)$ . We let  $H^i(L)$  to denote the  $i$ -th homology group of the complex  $\mathcal{C}(D)$  using Lee's differential. We summarize all results about this invariant needed in the sequel in the following proposition:

**Proposition 2.4.** [11, Section 4] *Let  $L$  be an oriented link with  $k$  components  $L_1, L_2, \dots, L_k$ . Then*

- (1)  $\dim(H(L)) = 2^k$ .
- (2) *For every orientation  $\theta$  of  $L$  there is a generator of homology in degree*

$$2 \times \sum_{l \in E, m \in \bar{E}} \text{lk}(L_l, L_m)$$

*where  $E \subset \{1, 2, \dots, k\}$  indexes the set of components of  $L$  whose original orientation needs to be reversed to get the orientation  $\theta$  and  $\bar{E} = \{1, \dots, k\} \setminus E$ . The linking numbers  $\text{lk}(L_l, L_m)$  are the linking numbers (for the original orientation) between the components  $L_l$  and  $L_m$ .*

- (3) *For any H-thin link  $L$ , we have a spectral sequence converging to  $H(L)$  with  $E_1^{s,t} = \mathcal{H}^{s+t, 4s}(L)$  and  $E_2 = H(L)$ . In other words,*

$$H(L) \cong \frac{\text{Ker}(\Phi : \mathcal{H}(L) \rightarrow \mathcal{H}(L))}{\text{Image}(\Phi : \mathcal{H}(L) \rightarrow \mathcal{H}(L))}.$$

Our main goal in this paper is to prove the following theorem and give some consequences in the last section.

**Theorem 2.5.** *The length of any gap in the differential grading of the Khovanov homology of any quasi-alternating link is one.*

$j \backslash i$	-3	-2	-1	0
-1				1
-3				1
-5		1		
-7				
-9	1			

TABLE 1. The Khovanov homology of the trefoil knot.

## 3. PROOF OF THE MAIN THEOREM

The crucial step of the proof of Theorem 2.5 relies on showing that the Lee homology of any quasi-alternating link is bi-graded and that the Khovanov homology of such link satisfies the Knight Move Conjecture.

A simple way to describe the Khovanov homology of a given link is to locate a non negative integer  $n$  at the point  $(i, j)$ , where  $n$  represents the dimension of  $\mathcal{H}^{i,j}(L)$  over  $\mathbb{Q}$ . An example is illustrated in Table 1. Based on the this description of Khovanov homology, we introduce the following definition.

**Definition 3.1.** For a given link  $L$ ,

- (1) We say that the differential grading of the Khovanov homology of  $L$  has a gap of length  $s$  if there are  $s$  consecutive columns of the Khovanov homology of the link  $L$  with zero entries and these columns correspond to vertical lines of  $x$ -intercepts  $i, i+1, i+2, \dots, i+s-1$  such that  $i_{\min}+1 \leq i < i+1 < i+2 < \dots < i+s-1 \leq i_{\max}-1$ . This is equivalent to say that  $\mathcal{H}^{i-1}(L) \neq 0$  and  $\mathcal{H}^{i+s}(L) \neq 0$  while  $\mathcal{H}^m(L) = 0$  for  $m = i, i+1, \dots, i+s-1$ .
- (2) We say that the quantum grading of the Khovanov homology of  $L$  has a gap of length  $s$  if there are  $s$  consecutive rows of the Khovanov homology of the link  $L$  with zero entries and these rows correspond to horizontal lines of  $y$ -intercepts  $j, j+2, j+4, \dots, j+2s-2$  such that  $j_{\min}+2 \leq j < j+2 < j+4 < \dots < j+2s-2 \leq j_{\max}-2$ .
- (3) We say that the the Jones polynomial of the link  $L$  has a gap of length  $s$  if this polynomial has two monomials  $t^i$  and  $t^{i+s+1}$  of nonzero coefficients and all the monomials  $t^m$  have zero coefficients for  $m = i+1, i+2, \dots, i+s$ . The gap in the Kauffman bracket polynomial of the link  $L$  can be defined in a similar manner.
- (4) We say that there is a gap between the two polynomials  $f(A)$  and  $g(A)$  of length  $s$  if either  $M = \maxdeg(f(A)) < m = \mindeg(g(A))$  or  $M = \maxdeg(g(A)) < m = \mindeg(f(A))$  and all the monomials  $A^{M+1}, A^{M+2}, \dots, A^{M+s} = A^{m-1}$  have zero coefficients in the two polynomials.

To illustrate this definition, we consider the Khovanov homology of the trefoil knot  $3_1$  given in Table 1. This homology has a gap of length one in the differential grading as  $\mathcal{H}^{-1}(3_1) = 0$  while both  $\mathcal{H}^{-2}(3_1)$  and  $\mathcal{H}^0(3_1)$  are nontrivial. Also, the Jone polynomial of the trefoil that is known to be  $-t^{-4} + t^{-3} + t^{-1}$  has a gap of length one as the monomial  $t^{-2}$  has zero coefficient while the monomials  $t^{-1}$  and  $t^{-3}$  have nonzero coefficients.

From now on, we let  $L$  be a quasi-alternating link and  $L_0$  and  $L_1$  be its smoothings at the crossing  $c$  where it is quasi-alternating. The signatures of the skein triple  $(L, L_0, L_1)$  are related as given in the following lemma:

**Lemma 3.2.** [13, Lemma 3] *For the triple  $(L, L_0, L_1)$ , we have*

$$\sigma(L_0) = \sigma(L) + 1$$

and

$$\sigma(L_1) = \sigma(L) + e,$$

where  $e$  is as defined in Section 2.

Unlike Khovanov homology which is bi-graded as explained in Section 2, Lee's homology theory is known to be filtered rather than bi-graded. However, the Lee homology of a quasi-alternating link can be endowed with a bi-grading structure as shown in the following theorem.

**Theorem 3.3.** *If  $L$  is a quasi-alternating link, then the Lee homology  $H(L)$  is bi-graded and  $H^{i,j}(L) \cong H^{i,j+2}(L) \cong \mathbb{Q}^n$  for some non-negative integer  $n$  and for any  $i$  and  $j$  such that  $j - 2i = -\sigma(L) - 1$ .*

*Proof.* The link  $L$  is  $H$ -thin as it is quasi-alternating, [13, Theorem 1]. Consequently, the bi-grading of  $\mathcal{H}(L)$  induces a bi-grading on  $H(L)$  as a result of Part 3 of Proposition 2.4. We use induction on the determinant of the link  $L$  to prove the second claim in the theorem. It is clear that the result holds when  $\det(L) = 1$  since the only quasi-alternating link of determinant 1 is the unknot [7, Prop. 3.2]. Now suppose that the result holds for all links of determinant less than  $m = \det(L)$ . In particular, it holds for the links  $L_0$  and  $L_1$ . It is easy to see that if  $H^i(L) \neq 0$  then  $H^{i\pm 1}(L) = 0$  as a result of Part 2 of Proposition 2.4. If we use this fact in the long exact sequence of the cohomology groups  $H(L), H(L_0)$  and  $H(L_1)$  that is mentioned for the first time in the proof of [11, Theorem 4.2], then we obtain

$$0 \longrightarrow \overline{H}^{i-1}(L_0) \longrightarrow \overline{H}^{i-1}(L_1) \longrightarrow \overline{H}^i(L) \longrightarrow \overline{H}^i(L_0) \longrightarrow \overline{H}^i(L_1) \longrightarrow 0.$$

As a result of the facts that  $x(L_0) = x(L), y(L_0) = y(L) - 1, x(L_1) = x(L) + e$  and  $y(L_1) = y(L) - e - 1$ , we obtain the following exact sequences:

$$(6) \quad 0 \longrightarrow H^{i-1}(L_0) \longrightarrow H^{i-e-1}(L_1) \longrightarrow H^i(L) \longrightarrow H^i(L_0) \longrightarrow H^{i-e}(L_1) \longrightarrow 0.$$

and

$$(7) \quad 0 \longrightarrow H^{i-1,j-1}(L_0) \longrightarrow H^{i-e-1,j-3e-2}(L_1) \longrightarrow H^{i,j}(L) \longrightarrow H^{i,j-1}(L_0) \longrightarrow H^{i-e,j-3e-2}(L_1) \longrightarrow 0$$

The above exact sequence still holds if we replace  $j$  by  $j + 2$  as follows:

$$(8) \quad 0 \longrightarrow H^{i-1,j+1}(L_0) \longrightarrow H^{i-e-1,j-3e}(L_1) \longrightarrow H^{i,j+2}(L) \longrightarrow H^{i,j+1}(L_0) \longrightarrow H^{i-e,j-3e}(L_1) \longrightarrow 0$$

It is easy to see that  $H^{i-1,j+1}(L_0) = 0 = H^{i-e,j-3e-2}(L_1)$  since  $(j+1) - 2(i-1) = j - 2i + 3 = -\sigma(L) - 1 + 3 = -\sigma(L_0) + 3 \neq -\sigma(L_0) \pm 1$  and  $(j - 3e - 2) - 2(i - e) = j - 2i - e - 2 = -\sigma(L) - 1 - e - 2 = -\sigma(L_1) - 3 \neq -\sigma(L_1) \pm 1$ , respectively provided that  $j - 2i = -\sigma(L) - 1$ . Also as a result of the induction hypotheses on  $L_0$  and  $L_1$ , we obtain  $H^{i-e-1,j-3e-2}(L_1) \cong H^{i-e-1,j-3e}(L_1)$  and  $H^{i,j-1}(L_0) \cong H^{i,j+1}(L_0)$  since  $(j - 3e) - 2(i - e - 1) = j - 2i - e + 2 = -\sigma(L) - 1 - e + 2 = -\sigma(L_1) + 1, (j - 3e - 2) - 2(i - e - 1) = j - 2i - e = -\sigma(L) - 1 - e = -\sigma(L_1) - 1$  and  $(j - 1) - 2i = j - 2i - 1 = -\sigma(L) - 1 - 1 = -\sigma(L_0) - 1, (j + 1) - 2i = j - 2i + 1 = -\sigma(L) - 1 + 1 = -\sigma(L_0) + 1$ . Now we have four cases to consider:

- (1) If  $H^{i-e-1,j-3e-2}(L_1) \cong H^{i-e-1,j-3e}(L_1) = 0$  and  $H^{i,j-1}(L_0) \cong H^{i,j+1}(L_0) = 0$ , then  $H^{i,j+2}(L) = 0 = H^{i,j}(L)$ .
- (2) If  $H^{i-e-1,j-3e-2}(L_1) \cong H^{i-e-1,j-3e}(L_1) \neq 0$  and  $H^{i,j-1}(L_0) \cong H^{i,j+1}(L_0) \neq 0$ , then as a result of the fact that if  $H^i(L) \neq 0$  then  $H^{i\pm 1}(L) = 0$  obtained from Part 2 of Proposition 2.4 we conclude that  $H^{i-1,j-1}(L_0) = 0 = H^{i-e,j-3e}(L_1)$ . This implies that  $H^{i,j}(L) \cong H^{i,j-1}(L_0) \oplus H^{i-e-1,j-3e-2}(L_1) \cong H^{i,j+1}(L_0) \oplus H^{i-e-1,j-3e}(L_1) \cong H^{i,j+2}(L)$ .
- (3) If  $H^{i-e-1,j-3e-2}(L_1) \cong H^{i-e-1,j-3e}(L_1) = 0$  and  $H^{i,j-1}(L_0) \cong H^{i,j+1}(L_0) \neq 0$ , then as a result of the fact that if  $H^i(L) \neq 0$  then  $H^{i\pm 1}(L) = 0$  obtained from Part 2 of Proposition 2.4 we conclude that  $H^{i-1,j-1}(L_0) = 0$ . In particular, we obtain  $H^{i,j-1}(L_0) \cong$

$H^{i,j}(L) \oplus H^{i-e,j-3e-2}(L_1) \cong H^{i,j}(L)$  and  $H^{i,j+1}(L_0) \cong H^{i,j+2}(L) \oplus H^{i-e,j-3e}(L_1)$ . To prove the required result it is enough to prove that  $H^{i-e,j-3e}(L_1) = 0$ . We prove this by contradiction. If we assume that  $H^{i-e,j-3e}(L_1) \neq 0$ , then  $H^{i-e}(L_1) \neq 0$  as a result of the fact that  $H^{i-e}(L_1) \cong H^{i-e,j-3e+2}(L_1) \oplus H^{i-e,j-3e}(L_1)$  obtained from Part 3 of Proposition 2.4 because  $L_1$  is  $H$ -thin as it is quasi-alternating. Now according to the long exact sequence 6 and the assumption in this case we conclude that  $H^i(L_0) \cong H^i(L) \oplus H^{i-e}(L_1) \cong (H^{i,j}(L) \oplus H^{i,j+2}(L)) \oplus (H^{i-e,j-3e}(L_1) \oplus H^{i-e,j-3e+2}(L_1))$ . Thus we obtain  $0 = H^{i-e,j-3e+2}(L_1) \cong H^{i-e,j-3e}(L_1)$ , where the isomorphism follows by the induction hypothesis on the link  $L_1$ .

- (4) If  $H^{i-e-1,j-3e-2}(L_1) \cong H^{i-e-1,j-3e}(L_1) \neq 0$  and  $H^{i,j-1}(L_0) \cong H^{i,j+1}(L_0) = 0$ . This case can be discussed using a similar argument as in the previous case.

□

**Remark 3.4.** In fact, if the  $E_2$ - and the  $E_\infty$ -pages of the spectral sequence of the double complex  $(\mathcal{C}(L), d(L), \Phi(L))$  are isomorphic, then the Lee homology  $H(L)$  is bi-graded since these two pages are isomorphic to  $\frac{\text{Ker}(\Phi: \mathcal{H}(L) \rightarrow \mathcal{H}(L))}{\text{Image}(\Phi: \mathcal{H}(L) \rightarrow \mathcal{H}(L))}$  and  $H(L)$ , respectively. As a result of this fact, we can associate a Lee polynomial invariant to any such link as follows:

$$(9) \quad Le(L)(t, q) = \sum_{i,j \in \mathbb{Z}} t^i q^j \dim H^{i,j}(L).$$

The following corollary follows directly from Part 2 of Proposition 2.4 and Remark 3.4.

**Corollary 3.5.** *If  $H(L)$  is bi-graded, then the polynomial  $\sum_{i,j \in \mathbb{Z}} (-1)^i q^j \dim H^{i,j}(L)$  has only monomials of positive coefficients.*

As a consequence of Theorem 3.3 and Part 3 of Proposition 2.4, we establish the Knight Move Conjecture for quasi-alternating links. This generalizes [11, Theorem 4.5] to this class of links.

**Corollary 3.6.** *Let  $L$  be an oriented quasi-alternating link with components  $L_1, L_2, \dots, L_n$ . Let  $l_{jk}$  denote the linking number of the components  $L_j$  and  $L_k$ . Then, we have*

$$Kh(L)(t, q) = q^{-\sigma(L)} \{ (q^{-1} + q) \left( \sum_{E \subset \{2, 3, \dots, n\}} (tq^2)^{\sum_{j \in E, k \notin E} 2l_{jk}} \right) + (q^{-1} + tq^2 \cdot q) Kh'(L)(tq^2) \},$$

for some polynomial  $Kh'(L)$ .

One can define two types of *breadth* for the Khovanov homology of any link  $L$  based on the differential and quantum gradings of Khovanov homology. These two types are denoted hereafter by  $\text{breadth}_i(\mathcal{H}(L))$  and  $\text{breadth}_j(\mathcal{H}(L))$  and are defined to be the difference between the maximum and minimum corresponding gradings. For instance, the  $\text{breadth}_i(\mathcal{H}(3_1)) = 3$  while  $\text{breadth}_j(\mathcal{H}(3_1)) = 8$  as one can see from Table 1. In the case of quasi-alternating links, these two types of breadth are related as follows.

**Corollary 3.7.** *If  $L$  is a quasi-alternating link, then  $\text{breadth}_j(\mathcal{H}(L)) = 2 \text{breadth}_i(\mathcal{H}(L)) + 2$ .*

*Proof.* In the above description of Khovanov homology, we claim that the most left-bottom nonzero entry that comes right after any gap or at the beginning of the Khovanov homology of  $L$  will be on the lower diagonal with  $j - 2i = -\sigma(L) - 1$  and the most right-top nonzero entry right before any gap or at the end of the Khovanov homology of  $L$  will be on the upper diagonal with  $j - 2i = -\sigma(L) + 1$ . Each one of these two entries is the only entry in that row.

We show the first case of the claim and the second case of the claim follows using a similar argument. Suppose that the most left-bottom entry shares the same row with another entry to the right. In this case, the leftmost entry will be on the upper diagonal and therefore it will

be the only entry to survive in  $H^i(L)$  in that column, where  $i$  is the differential grading of the most left-bottom entry. This is impossible according to the result of Theorem 3.3.

Thus the minimum quantum grading  $j_{\min}$  is related to the minimum differential grading  $i_{\min}$  by  $j_{\min} - 2i_{\min} = -\sigma(L) - 1$  and the maximum quantum grading  $j_{\max}$  is related to the maximum differential grading  $i_{\max}$  by  $j_{\max} - 2i_{\max} = -\sigma(L) + 1$ . Now the result follows because  $\text{breadth}_j(\mathcal{H}(L)) = j_{\max} - j_{\min} = (2i_{\max} - \sigma(L) + 1) - (2i_{\min} - \sigma(L) - 1) = 2(i_{\max} - i_{\min}) + 2 = 2 \text{breadth}_i(\mathcal{H}(L)) + 2$ .  $\square$

We can give an explicit formula for the Jones polynomial of any quasi-alternating link in terms of the integers that are given in the above description of its Khovanov homology. Indeed, these integers can be expressed in terms of the dimensions of the Khovanov and Lee homologies of the given link. More precisely, if  $a_l$  is the  $(i_{\min} + l - 1, j_{\min} + 2l - 2)$ -entry of the lower diagonal in the above description of  $\mathcal{H}(L)$  after taking the quotient by  $H(L)$ . Then we have

**Lemma 3.8.** *If  $L$  is a quasi-alternating link, then*

$$a_l = \begin{cases} \dim \mathcal{H}^{i_{\min}}(L) - \dim H^{i_{\min}}(L), & \text{if } l = 1, \\ \dim \mathcal{H}^{i_{\min}+l-1}(L) - \dim H^{i_{\min}+l-1}(L) - a_{l-1}, & \text{if } 1 < l < i_{\max} - i_{\min} \\ \dim \mathcal{H}^{i_{\max}}(L) - \dim H^{i_{\min}}(L), & \text{if } i = i_{\max} - i_{\min}. \end{cases}$$

*Proof.* The result is a straightforward consequence of the fact that  $L$  is  $H$ -thin and Part 3 of Proposition 2.4.  $\square$

**Corollary 3.9.** *If the link  $L$  is quasi-alternating, then its Jones polynomial is given by*

$$(10) \quad V_L(t) = [(-1)^{i_{\min}} a_1 q^{j_{\min}+1} + (-1)^{i_{\max}} a_{(i_{\max}-i_{\min})} q^{j_{\max}-1} \\ + \sum_j n_j q^j + \sum_{l=1}^{\frac{j_{\max}-j_{\min}-4}{2}} (-1)^{l+i_{\min}} (a_l + a_{l+1}) q^{j_{\min}+2l+1}]_{q=-\sqrt{t}},$$

where the first sum is taken over all  $j_{\min} + 1 \leq j \leq j_{\max} - 1$  such that  $H^{i,j+1}(L) \cong H^{i,j-1}(L) \cong \mathbb{Q}^{n_j}$  for some positive integer  $n_j$ .

*Proof.* The entries at each column of  $H(L)$  contribute  $n_j(q^m + q^{m+2}) = n_j q^{m+1}(q^{-1} + q)$  to  $Kh(-1, q)$  for some integer  $m$ . Now the first sum follows after we divide by  $q^{-1} + q$  and go over all the columns of  $H(L)$ .

As a result of Part 3 of Proposition 2.4, we conclude that there is a pairing (knight move) in the entries of  $\mathcal{H}(L)/H(L)$ . In particular, the  $(i_{\min} + l - 1, j_{\min} + 2l - 2)$ -entry of the lower diagonal is equal to the  $(i_{\min} + l, j_{\min} + 2l + 2)$ -entry for  $1 \leq l \leq i_{\max} - i_{\min}$ . It is not too hard to see that the  $(i_{\min} + l - 1, j_{\min} + 2l - 2)$ -entry is simply  $a_l$  as a result of Lemma 3.8. These two entries contribute  $(-1)^{i_{\min}+l-1} a_l (q^{j_{\min}+2l-2} - q^{j_{\min}+2l+2}) = (-1)^{i_{\min}+l-1} a_l (q^{-1} + q)(q^{j_{\min}+2l-1} - q^{j_{\min}+2l+1})$  to the polynomial  $Kh(-1, q)$ . The second sum follows after we divide by  $q^{-1} + q$  and go over all such entries.  $\square$

**Lemma 3.10.** *The coefficient  $b_j$  of the monomial  $q^j$  in the above formula is given by*

$$b_j = \begin{cases} (-1)^{i_{\min}} a_1 + \frac{1}{2} \dim H^{i_{\min}}(L), & \text{if } j = j_{\min} + 1, \\ (-1)^{l+i_{\min}} (a_{i-i_{\min}} + a_{i-i_{\min}+1}) + \frac{1}{2} \dim H^{i_{\min}+l}(L), & \text{if } j_{\min} + 1 < j < j_{\max} - 1 \\ (-1)^{i_{\max}} a_{i_{\max}-i_{\min}} + \frac{1}{2} \dim H^{i_{\max}}(L), & \text{if } j = j_{\max} - 1, \end{cases}$$

where  $l = \frac{j-j_{\min}-1}{2}$ . Moreover,  $b_j$  is zero only if the two terms that define  $b_j$  are equal to zero.

*Proof.* We have three cases to consider according to the value of  $j$ . We discuss the case  $j_{\min} + 1 < j < j_{\max} - 1$  and the other cases follow using a similar argument. If the monomial  $q^j$  appears in the two sums in Equation 10, then we obtain  $n_j = \frac{1}{2} \dim H^i(L)$  with  $j - 2i = -\sigma(L)$  and  $j = j_{\min} + 2l + 1$  for some  $l$ . Now if we use the fact that  $j_{\min} - 2i_{\min} = -\sigma(L) - 1$ , we conclude



that  $i = i_{\min} + l$  and hence the result of the first part follows and the second part follows easily from the fact that  $i$  has to be even if  $\dim H^i(L) \neq 0$ .  $\square$

**Proposition 3.11.** *Let  $L$  be a quasi-alternating link. The differential grading of  $\mathcal{H}(L)$  has a gap of length  $s$  if and only if  $V_L(t)$  has a gap of length  $s$ .*

*Proof.* The result can be proved easily from the following three equivalences which are direct consequences of Lemma 3.10

- (1)  $\dim \mathcal{H}^{i_{\min}}(L) = a_1 + \dim H^{i_{\min}}(L) \neq 0$  is equivalent to  $a_1 + \frac{1}{2} \dim H^{i_{\min}}(L) \neq 0$  which is also equivalent to  $b_{j_{\min}+1} = (-1)^{i_{\min}} a_1 + \frac{1}{2} \dim H^{i_{\min}}(L) \neq 0$ .
- (2)  $\dim \mathcal{H}^i(L) = a_{i-i_{\min}} + a_{i-i_{\min}+1} + \dim H^i(L) \neq 0$  is equivalent to  $a_{i-i_{\min}} + a_{i-i_{\min}+1} + \frac{1}{2} \dim H^i(L) \neq 0$  which is also equivalent to  $b_j = (-1)^{l+i_{\min}} (a_{i-i_{\min}} + a_{i-i_{\min}+1}) + \frac{1}{2} \dim H^{i_{\min}+l}(L) \neq 0$  where  $i_{\min} < i < i_{\max}$  and  $j = 2i - \sigma(L)$ .
- (3)  $\dim \mathcal{H}^{i_{\max}}(L) = a_{i_{\max}-i_{\min}} + \dim H^{i_{\max}}(L) \neq 0$  is equivalent to  $a_{i_{\max}-i_{\min}} + \frac{1}{2} \dim H^{i_{\max}}(L) \neq 0$  which is also equivalent to  $b_{j_{\max}-1} = (-1)^{i_{\max}} a_{i_{\max}-i_{\min}} + \frac{1}{2} \dim H^{i_{\max}}(L) \neq 0$ .  $\square$

According to the long exact sequence in Equation 5, the differential support of  $\mathcal{H}(L)$  is included in the union of the differential support of  $\mathcal{H}(L_0)$  and the differential support of  $\mathcal{H}(L_1)$  after doing the appropriate shift. We denote the differential supports of  $\mathcal{H}(L_0)$  and  $\mathcal{H}(L_1)$  by  $S_0$  and  $S_1$ , respectively. Also, we let  $S\{k\}$  to denote the set  $S$  shifted to the left by  $k$  that is  $i \in S\{k\}$  if  $i + k \in S$ .

**Proposition 3.12.** *If there is a gap between the two polynomials  $A\langle L_0 \rangle$  and  $A^{-1}\langle L_1 \rangle$  of length bigger than three, then  $|i_1 - i_0| > 1$  for any  $i_0 \in S_0$  and  $i_1 \in S_1\{e+1\}$ . In particular, the sets  $S_0$  and  $S_1\{e+1\}$  are disjoint and this implies that*

$$\mathcal{H}^i(L) \cong \begin{cases} \mathcal{H}^i(L_0), & \text{if } i \in S_0, \\ \mathcal{H}^{i-e-1}(L_1), & \text{if } i \in S_1\{e+1\}, \\ 0, & \text{otherwise.} \end{cases}$$

*Proof.* If there is a gap between the two polynomials  $A\langle L_0 \rangle$  and  $A^{-1}\langle L_1 \rangle$  of length bigger than three, then either  $M = \maxdeg(A\langle L_0 \rangle) < \mindeg(A^{-1}\langle L_1 \rangle) = m$  or  $M = \maxdeg(A^{-1}\langle L_1 \rangle) < \mindeg(A\langle L_0 \rangle) = m$  such that the monomials  $A^{M+1}, A^{M+1}, \dots, A^{M+s}$  have zero coefficients in both polynomials with  $s$  at least seven as the powers of the monomials of  $\langle L \rangle$  differ by multiples of four. We discuss the first case and the second case follows using a similar argument. Notice that we have

$$\begin{aligned} \left( (-A)^{-3w(L)} (-A^{-2} - A^2) \langle L \rangle \right) &= (-t^{-1/2} - t^{1/2}) V_L(t)_{t^{1/2}=-q} = (q + q^{-1}) V_L(t)_{t^{1/2}=-q} \\ &= \sum_{i,j \in \mathbb{Z}} (-1)^i q^j \dim \mathcal{H}^{i,j}(L). \end{aligned}$$

In addition, since  $x(L_0) = x(L)$  and  $y(L_0) = y(L) - 1$ , we obtain  $w(L_0) = w(L) - 1$  and

$$\begin{aligned}
\left( A(-A)^{-3w(L)}(-A^{-2} - A^2) \langle L_0 \rangle \right) &= \left( A(-A)^{-3(w(L_0)+1)}(-A^{-2} - A^2) \langle L_0 \rangle \right)_{A^{-2}=t^{1/2}=-q} \\
&= -t^{1/2}(-t^{-1/2} - t^{1/2})V_{L_0}(t)_{t^{1/2}=-q} \\
&= q(q + q^{-1})V_{L_0}(t)_{t^{1/2}=-q} \\
&= \sum_{i,j \in \mathbb{Z}} (-1)^i q^{j+1} \dim \mathcal{H}^{i,j}(L_0) \\
&= \sum_{i,j \in \mathbb{Z}} (-1)^i q^j \dim \mathcal{H}^{i,j-1}(L_0).
\end{aligned}$$

Moreover, from facts that  $x(L_1) = x(L) + e$  and  $y(L_1) = y(L) - e - 1$ , we obtain  $w(L_1) = w(L) - 2e - 1$  where  $e$  is as defined in Lemma 3.2 and

$$\begin{aligned}
\left( A^{-1}(-A)^{-3w(L)}(-A^{-2} - A^2) \langle L_1 \rangle \right) &= \left( A^{-1}(-A)^{-3(w(L_1)+2e+1)}(-A^{-2} - A^2) \langle L_1 \rangle \right)_{A^{-2}=t^{1/2}=-q} \\
&= -t^{\frac{3e+2}{2}}(-t^{-1/2} - t^{1/2})V_{L_1}(t)_{t^{1/2}=-q} \\
&= (-1)^{3e-1} q^{3e+2} (q + q^{-1}) V_{L_1}(t)_{t^{1/2}=-q} \\
&= \sum_{i,j \in \mathbb{Z}} (-1)^{i+3e-1} q^{j+3e+2} \dim \mathcal{H}^{i,j}(L_1) \\
&= \sum_{i,j \in \mathbb{Z}} (-1)^i q^j \dim \mathcal{H}^{i-e-1,j-3e-2}(L_1).
\end{aligned}$$

As a direct consequence of the previous argument and the fact that  $\langle L \rangle = A \langle L_0 \rangle + A^{-1} \langle L_1 \rangle$ , we obtain

$$\sum_{i,j \in \mathbb{Z}} (-1)^i q^j \dim \mathcal{H}^{i,j}(L) = \sum_{i,j \in \mathbb{Z}} (-1)^i q^j \dim \mathcal{H}^{i,j-1}(L_0) + \sum_{i,j \in \mathbb{Z}} (-1)^i q^j \dim \mathcal{H}^{i-e-1,j-3e-2}(L_1),$$

with  
(11)

$$\maxdeg \left( \sum_{i,j \in \mathbb{Z}} (-1)^i q^j \dim \mathcal{H}^{i,j-1}(L_0) \right) < \mindeg \left( \sum_{i,j \in \mathbb{Z}} (-1)^i q^j \dim \mathcal{H}^{i-e-1,j-3e-2}(L_1) \right).$$

Now, if  $(i_{\max}, j_{\max})$  are the coordinates of the most right-top nonzero entry of  $\mathcal{H}(L_0)$  that is part of the upper diagonal as pointed out in the proof of Corollary 3.7, then we obtain  $\mathcal{H}^{i_{\max}, j_{\max}+1}(L) \cong \mathcal{H}^{i_{\max}, j_{\max}}(L_0)$  from the long exact sequence in Equation 5. The last result follows because  $\dim \mathcal{H}^{i_{\max}-e-1, j_{\max}-3e-1}(L_1) = 0 = \dim \mathcal{H}^{i_{\max}-e, j_{\max}-3e-1}(L_1)$  as they are the coefficients of  $q^{j_{\max}+1}$  in  $\sum_{i,j \in \mathbb{Z}} (-1)^i q^j \dim \mathcal{H}^{i-e-1,j-3e-2}(L_1)$  with

$$\begin{aligned}
\mindeg \left( \sum_{i,j \in \mathbb{Z}} (-1)^i q^j \dim \mathcal{H}^{i-e-1,j-3e-2}(L_1) \right) &> \maxdeg \left( \sum_{i,j \in \mathbb{Z}} (-1)^i q^j \dim \mathcal{H}^{i,j-1}(L_0) \right) \\
&= j_{\max} + 1.
\end{aligned}$$

As a result of the fact  $j_{\max} - 2i_{\max} = -\sigma(L_0) + 1$ , we obtain  $j_{\max} + 1 - 2i_{\max} = j_{\max} - 2i_{\max} + 1 = -\sigma(L_0) + 1 + 1 = -(\sigma(L_0) - 1) + 1 = -\sigma(L) + 1$ . In other words, the last statement is equivalent to say that the above nonzero entry of the Khovanov homology of  $L$  with coordinates  $(i_{\max}, j_{\max} + 1)$  will also lie on the upper diagonal of the Khovanov homology of  $L$ .

Similarly, if  $(i_{\min}, j_{\min})$  are the coordinates of the most left-bottom nonzero entry of  $\mathcal{H}(L_1)$  that is part of the lower diagonal as pointed in the proof of Corollary 3.7, then we obtain

$\mathcal{H}^{i_{\min}+e+1, j_{\min}+3e+2}(L) \cong \mathcal{H}^{i_{\min}, j_{\min}}(L_1)$  from the long exact sequence in Equation 5. The last result follows because  $\dim \mathcal{H}^{i_{\min}+e, j_{\min}+3e+1}(L_0) = 0 = \dim \mathcal{H}^{i_{\min}+e+1, j_{\min}+3e+1}(L_0)$  as they are the coefficients of  $q^{j_{\min}+3e+2}$  in  $\sum_{i,j \in \mathbb{Z}} (-1)^i q^j \dim \mathcal{H}^{i,j-1}(L_0)$  with

$$\begin{aligned} j_{\min} + 3e + 2 &= \mindeg \left( \sum_{i,j \in \mathbb{Z}} (-1)^i q^j \dim \mathcal{H}^{i-e-1, j-3e-2}(L_1) \right) \\ &> \maxdeg \left( \sum_{i,j \in \mathbb{Z}} (-1)^i q^j \dim \mathcal{H}^{i,j-1}(L_0) \right). \end{aligned}$$

Since  $j_{\min} - 2i_{\min} = -\sigma(L_1) + 1$ , we obtain  $j_{\min} + 3e + 2 - 2(i_{\min} + e + 1) = j_{\min} - 2i_{\min} + e = -\sigma(L_1) + e - 1 = -(\sigma(L_1) - e) - 1 = -\sigma(L) - 1$ . In other words, the last statement is equivalent to say that the above nonzero entry of the Khovanov homology of  $L$  with coordinates  $(i_{\max} + e + 1, j_{\max} + 3e + 2)$  will also lie on the lower diagonal of the Khovanov homology of  $L$ .

Thus, we conclude that  $2(i_{\min} + e + 1 - i_{\max}) = j_{\min} + 3e + 2 + \sigma(L) + 1 - j_{\max} - \sigma(L) + 1 = j_{\min} + 3e + 2 - j_{\max} + 2 \geq 2 + 2 = 4$ , where the inequality follows from the fact that  $j_{\min} + 3e + 2 - j_{\max} - 1 > 0$  that is obtained from Equation 11. The last inequality implies that  $i_{\min} + e + 1 \geq i_{\max} + 2$  and hence we conclude that  $|i_1 - i_0| \geq |(i_{\min} + e + 1) - i_{\max}| \geq 2$  for any  $i_0 \in S_0$  and  $i_1 \in S_1\{e+1\}$  as a result of  $i_0 \leq i_{\max}$  and  $i_{\min} + e + 1 \leq i_1$ .

To compute the Khovanov homology of the link  $L$ , we need to consider only  $i \in S_0 \cup S_1\{e+1\}$  because otherwise we obtain  $0 = \mathcal{H}^{i-e-1}(L_1) \rightarrow \mathcal{H}^i(L) \rightarrow \mathcal{H}^i(L_0) = 0$  from the long exact sequence and this implies that  $\mathcal{H}^i(L) = 0$ . The other two cases are either  $i \in S_0$  or  $i \in S_1\{e+1\}$  but not both since the two sets  $S_0$  and  $S_1\{e+1\}$  are disjoint. We discuss the first case and the other case follows using a similar argument. If  $i \in S_0$ , then  $i + e + 1, i + e \notin S_1$  as a result of what we just proved. Hence the result follows directly from the long exact sequence  $0 = \mathcal{H}^{i-e-1}(L_1) \rightarrow \mathcal{H}^i(L) \rightarrow \mathcal{H}^i(L_0) \rightarrow \mathcal{H}^{i-e-1}(L_1) = 0$ .

□

**Corollary 3.13.** *If there is a gap between the two polynomials  $A\langle L_0 \rangle$  and  $A^{-1}\langle L_1 \rangle$  of length bigger than three, then we have*

$$H^i(L) \cong \begin{cases} H^i(L_0), & \text{if } i \in S_0, \\ H^{i-e-1}(L_1), & \text{if } i \in S_1\{e+1\}, \\ 0, & \text{otherwise.} \end{cases}$$

*Proof.* The Lee homology of the link  $L$  is isomorphic to the  $E_2$ -page of the spectral sequence of the double complex  $(\bar{\mathcal{C}}(L), d(L), \Phi(L))$  that is simply the disjoint union of the  $E_2$ -pages of the spectral sequences of the double complexes for the links  $L_0$  and  $L_1$  as a consequence of the result of Proposition 3.12. But as a result of Part 3 of Proposition 2.4 and since  $L_0$  and  $L_1$  are quasi-alternating, the  $E_2$ -pages of the spectral sequences for the double complexes for the links  $L_0$  and  $L_1$  are isomorphic to the  $E_\infty$ -pages of the spectral sequences of the double complexes for the links  $L_0$  and  $L_1$ , respectively. Therefore, the  $E_\infty$ -page of the spectral sequence of the double complex for the link  $L$  is simply the disjoint union of the  $E_\infty$ -pages of the spectral sequences for the double complexes for the links  $L_0$  and  $L_1$ . In particular, the Lee homology of the link  $L$  is simply the disjoint union of the Lee homologies of the links  $L_0$  and  $L_1$ . □

**Corollary 3.14.** *If  $l_0 \neq l - 1$  or  $l_1 \neq l - 1$  where  $l, l_0$  and  $l_1$  are the number of components of  $L_0, L_1$  and  $L$  respectively and if there is a gap between the two polynomials  $A\langle L_0 \rangle$  and  $A^{-1}\langle L_1 \rangle$ , then this gap has to be of length three.*


*Proof.* If the gap is of length bigger than three, then the results of Corollary 3.13 and Part 1 of Proposition 2.4 yields  $2^l = \dim H(L) = \dim H(L_0) + \dim H(L_1) = 2^{l_0} + 2^{l_1}$ . Obviously, this holds only if  $l_0 = l - 1 = l_1$ .  $\square$


**Proof of Theorem 2.5.** We use induction on the determinant of the given quasi-alternating link to prove the statement of the theorem. If  $\det(L) = 1$ , then the result holds since the only quasi-alternating link of determinant one is the unknot [7, Prop.3.2]. Now, suppose that the result holds for all quasi-alternating links of determinant less than the determinant of the link  $L$ . In particular, any gap in the differential gradings of  $\mathcal{H}(L_0)$  and  $\mathcal{H}(L_1)$  is of length one.

We have two cases to consider. The first case if there is a gap between the polynomials  $A\langle L_0 \rangle$  and  $A^{-1}\langle L_1 \rangle$  of length seven or less and this includes the case of having no gap. The second case corresponds to a gap of length bigger than seven. In any case, there is no cancellation between these two polynomials as a result of  $L$  being quasi-alternating. Thus any gap in the Jones polynomial of the link  $L$  is induced either by a gap in one of the two polynomials of  $L_0$  and  $L_1$  or by the gap between these two polynomials.

In the first case, it is easy to see that the Jones polynomial of  $L$  has no gap of length bigger than one as a result of the facts that the above polynomials have no gap of length bigger than seven from the induction hypothesis on the links  $L_0$  and  $L_1$  and that the gap between them is of length seven or less. Now the result follows from Proposition 3.11 since having a gap in the differential grading of  $\mathcal{H}(L)$  of length bigger than one induces a gap of length bigger than one in the Jones polynomial of  $L$  and this contradicts with what we know already about the Jones polynomial of  $L$ .

In the second case, the gap between the two polynomials is of length bigger than seven. According to Corollary 3.14, we can assume that  $L$  is a link of more than one component that is quasi-alternating at a crossing consisting of two arcs of two different components with  $l_0 = l - 1 = l_1$ . Without loss of generality, we can assume that the link  $L$  up to isotopy inside

a 3-ball is linked in a clasp as it appears in  and  $c$  is the top crossing in this clasp. This clasp structure can be obtained by sliding, if necessary, the under arc on top of the over arc of

the crossing  $c$  using a second Reidemeister move as shown in . Thus, we can assume that

the two components of the link are linked in a clasp as described above.

As a result of the relations of the Kauffman bracket, we have  $\langle L \rangle = A\langle L_0 \rangle + A^{-1}\langle L_1 \rangle = A(A\langle L_{00} \rangle + A^{-1}\langle L_{01} \rangle) - A^{-4}\langle L_{01} \rangle$ . We write  $A^2\langle L_{00} \rangle = f(A) - \langle L_{01} \rangle$ , where the polynomials  $f(A) = A\langle L_0 \rangle$  and  $\langle L_{01} \rangle = -A^3\langle L_1 \rangle = -A^4(A^{-1}\langle L_1 \rangle)$ . In fact, there is a gap between the two polynomials  $f(A)$  and  $\langle L_{01} \rangle$  of length bigger than or equal to seven as a result of the assumption that there is a gap of length bigger than seven between the polynomials  $A\langle L_0 \rangle$  and  $A^{-1}\langle L_1 \rangle$ . We show that such a case is impossible and thus the result follows.

Using an argument similar to the argument in Proposition 3.12, we obtain  $Kh_{L_0}(-1, q) = Kh_{L_{00}}(-1, q) + Kh_{L_{01}}(-1, q)$  from the equation  $\langle L_0 \rangle = A\langle L_{00} \rangle + A^{-1}\langle L_{01} \rangle$ . We can rewrite this equation as follows:  $Kh_{L_{00}}(-1, q) = Kh_{L_0}(-1, q) - Kh_{L_{01}}(-1, q)$  with a gap of a length bigger than or equal to one between  $Kh_{L_0}(-1, q)$  and  $Kh_{L_{01}}(-1, q)$  as a result of the fact that there is a gap between the polynomials  $f(A)$  and  $\langle L_{01} \rangle$  of length bigger than or equal to seven.

The Khovanov homology of the links  $L_0$  and  $L_{01}$  are supported on two diagonals since both links are quasi-alternating. The homologies of these two links do not overlap since there is a gap between  $Kh_{L_0}(-1, q)$  and  $Kh_{L_{01}}(-1, q)$  of length bigger than or equal to one. In particular, we have two cases to consider either  $M = \maxdeg(Kh_{L_0}(-1, q)) < m = \mindeg(Kh_{L_{01}}(-1, q))$  or  $M = \maxdeg(Kh_{L_{01}}(-1, q)) < m = \mindeg(Kh_{L_0}(-1, q))$ . We discuss the first case and the second case follows using a similar argument.

Now the most right-top nonzero entry in the Khovanov homology of  $L_0$  is on the top diagonal and the most left-bottom nonzero entry in the Khovanov homology of  $L_{01}$  is on the lower diagonal and each one of these two entries is the only entry in that row. Therefore, the  $E_\infty$ -page of the spectral sequence of the double complex  $(\overline{\mathcal{C}}(L_{00}), d(L_{00}), \Phi(L_{00}))$  is isomorphic to the  $E_2$ -page since  $d_2$  must be zero because of their degree. In this case, we obtain that the  $E_2$ -page for the double complex for the link  $L_{00}$  is simply the disjoint union of the  $E_2$ -pages for the double complexes for the links  $L_0$  and  $L_{01}$ . Thus according to Remark 3.4, the Lee polynomial invariant can be defined for this link and it is simply the difference of the Lee polynomial invariants of the links  $L_0$  and  $L_{01}$  that have no monomials in common and this contradicts the result of Corollary 3.5 after substituting  $t = -1$ .  $\square$

#### 4. CONSEQUENCES OF THE MAIN THEOREM

In this section, we discuss some consequences and applications of Theorem 2.5. We start by stating two corollaries. The first of which establishes a weaker version of Conjecture 2.3 in [5] and the second one establishes a weaker version of Conjecture 3.8 in [16].

**Corollary 4.1.** *If  $L$  is a quasi-alternating link, then the length of any gap in the Jones polynomial  $V_L(t)$  is one.*

*Proof.* The result follows easily by combining Theorem 2.5 with Proposition 3.11.  $\square$

**Corollary 4.2.** *If  $L$  is a quasi-alternating link, then  $\left\lceil \frac{\text{breadth } V_L(t)}{2} \right\rceil + 1 \leq \det(L)$ .*

*Proof.* The result follows from the fact that any gap in the Jones polynomial of the link has length one. This implies that the Jones polynomial has at least  $\left(\left\lceil \frac{\text{breadth } V_L(t)}{2} \right\rceil + 1\right)$  monomials each of which contributes an increment of at least one to the value of the determinant due to the fact that the Jones polynomial of a quasi-alternating link is alternating.  $\square$

**Corollary 4.3.** *Let  $L$  be a quasi-alternating link. Then, the length of any gap in the quantum grading of  $\mathcal{H}(L)$  is one.*

*Proof.* Suppose there are two consecutive rows that correspond to horizontal lines of  $y$ -intercepts  $j$  and  $j + 2$  of  $\mathcal{H}(L)$  with zero entries such that  $j_{\min} + 2 \leq j < j + 2 \leq j_{\max} - 2$ . In this case, this implies that  $\mathcal{H}^i(L) = 0$  for some  $i$  with  $j - 2i = -\sigma(L) - 1$ . From the assumption, we have  $\mathcal{H}^{i,j}(L) \cong \mathcal{H}^{i-1,j}(L) \cong \mathcal{H}^{i,j+2}(L) \cong \mathcal{H}^{i+1,j+2}(L) = 0$ . As a result of the knight move, we obtain  $\mathcal{H}^{i-1,j-2}(L) \cong \mathcal{H}^{i,j+2}(L) = 0$  and  $\mathcal{H}^{i+1,j+4}(L) \cong \mathcal{H}^{i,j}(L) = 0$ . This implies that  $\mathcal{H}^{i-1}(L) = \mathcal{H}^i(L) = \mathcal{H}^{i+1}(L) = 0$ . Also, as a result of the knight move we obtain  $j_{\min} + 4 \leq j \leq j_{\max} - 4$ . Therefore, we conclude that  $i_{\min} + 2 \leq i \leq i_{\max} - 2$ . This contradicts the fact that the length of any gap in the differential grading of  $\mathcal{H}(L)$  is one.  $\square$

**Corollary 4.4.** *Let  $L$  be a quasi-alternating link. If there is a gap between the two polynomials  $A\langle L_0 \rangle$  and  $A^{-1}\langle L_1 \rangle$ , then it has to be of length at most seven.*

As another application, we obtain a necessary condition for a Kanenobu knot to be quasi-alternating.

**Corollary 4.5.** *If  $|p + q| > 6$ , then the Kanenobu knot  $K(p, q)$  is not quasi-alternating.*

*Proof.* Any Kanenobu knots  $K(p, q)$  with  $|p + q| > 6$  has gap in the differential grading of  $\mathcal{H}(K(p, q))$  of length bigger than one as it was proved in [19, Theorem 1.3].  $\square$

**Remark 4.6.** The above result can be also obtained in terms of  $V_L(t)$ . In particular, it is proven in [8] that  $V_{K(p,q)}(t) = (-t)^{p+q}(V_{K(0,0)}(t) - 1) + 1 = (-t)^{p+q}((t^{-2} - t^{-1} + 1 - t + t^2)^2 - 1) + 1$ . It is easy to see that if  $|p + q| > 6$ , then the above polynomial has a gap of length bigger than

one. Thus according to Corollary 4.1, the knot  $K(p, q)$  is not quasi-alternating. It is known though that all these knots but finitely many of them are not quasi-alternating as a result of Corollary 3.3 in [16].

Now we investigate the implications of the main result to the class of alternating links being considered as a special class of quasi-alternating links. We know that any alternating link that is not a  $(2, n)$ -torus link has no gap in its Jones polynomial as a result of [21, Theorem 1(iv)]. In addition, it is known that all alternating links are  $H$ -thin [11, Theorem 1.2]. If we combine all these results, we obtain

**Corollary 4.7.** *If  $L$  is an alternating link that is not a  $(2, n)$ -torus link, then it has no gap in the differential grading of  $\mathcal{H}(L)$ .*

*Proof.* If we assume that the differential grading of  $\mathcal{H}(L)$  has a gap, then Proposition 3.11 implies that the Jones polynomial has a gap. This contradicts [21, Theorem 1(iv)].  $\square$

**Remark 4.8.** It is easy to see that if  $L$  is a  $(2, n)$ -torus link, then it has only one gap in the differential grading of  $\mathcal{H}(L)$  according to the computations in [9].

An easy technique to produce new examples of quasi-alternating links from old ones was introduced by Champanerkar and Kofman [3, Page 2452]. It basically replaces the crossing  $c$  where the link  $L$  is quasi-alternating by an alternating rational tangle of the same type. This technique has been later generalized not only to a single rational tangle but also to a product of rational tangles [17, Definition 2.5] and to non-algebraic alternating tangles [4]. The following proposition shows how this twisting technique affects the gaps in the differential grading.

**Proposition 4.9.** *Let  $L$  be a quasi-alternating link at some crossing  $c$  and let  $L^*$  be the link obtained by replacing the crossing  $c$  in  $L$  by a product of rational tangles that extends  $c$ . Then the number of gaps in the differential grading of  $\mathcal{H}(L^*)$  is less than or equal the number of gaps in  $\mathcal{H}(L)$ . In particular, if there are no gaps in differential grading of  $\mathcal{H}(L)$ , then it is also the case for the differential grading of  $\mathcal{H}(L^*)$ .*

*Proof.* For a positive integer  $n$ , we let  $L^n$  denote the link with the crossing  $c$  replaced by an alternating integer tangle of  $n$  vertical or horizontal crossings of the same type. To prove our result, we first show that the result holds for  $L^n$ . By writing the Kauffman bracket skein relations, one can easily prove that:

$$\begin{aligned} \langle L^n \rangle &= A^{n-1} (A \langle L_0 \rangle + A^{-1} \langle L_1 \rangle) + \left( \sum_{i=1}^{n-1} (-1)^i A^{n-4i-2} \right) \langle L_1 \rangle, \text{ and} \\ \langle L^n \rangle &= A^{-n+1} (A \langle L_0 \rangle + A^{-1} \langle L_1 \rangle) + \left( \sum_{i=1}^{n-1} (-1)^i A^{-n+4i+2} \right) \langle L_1 \rangle, \end{aligned}$$

for the case of vertical and horizontal tangles, respectively. Now, we discuss the first case. The second case can be treated in a similar manner. It is not too hard to see that any gap in  $\langle L^n \rangle$  is induced basically by a gap in  $\langle L \rangle$ . Finally the result follows since any product of rational tangles can be obtained by a sequence of integer tangles and the result of Proposition 3.11.  $\square$

Quasi-alternating links of braid index 3 and quasi-alternating Montesinos links have been classified in [1, 6], respectively. Now we apply the above result to these two classes of quasi-alternating links to obtain:

**Corollary 4.10.** *All quasi-alternating Montesinos links and quasi-alternating links of braid index 3 have no gap in the differential grading of their Khovanov homologies.*

*Proof.* The result follows as a result of Corollary 4.9 and the fact that the Jones polynomial of all quasi-alternating Montesinos links and quasi-alternating links of braid index 3 have no gap (Theorem 3.10 and Theorem 4.3 in [5], respectively).  $\square$

Finally, we enclose our discussion with the following conjecture that is supported by the above results and implies both Conjecture 2.3 in [5] and Conjecture 3.8 in [16]. In particular, it implies that the Jones polynomial of any quasi-alternating link that is not a  $(2, n)$ -torus link has no gap and the breadth of the Jones polynomial of such a link is a lower bound of the determinant of this link.

**Conjecture 4.11.** *If  $L$  is prime quasi-alternating that is not a  $(2, n)$ -torus link, then the differential grading of  $\mathcal{H}(L)$  has no gap.*

## REFERENCES

- [1] J. Baldwin, *Heegaard Floer homology and genus one, one boundary component open books*, J. Topol. **1** (4): 963–992, 2008.
- [2] D. Bar-Natan, *On Khovanov’s categorification of the Jones polynomial*, Algebraic and Geometric Topology **2**-16 (2002) 337–370.
- [3] A. Champanerker and I. Kofman, *Twisting Quasi-alternating Links*, Proc. Amer. Math. Soc., **137**(7): 2451–2458, 2009.
- [4] N. Chbili and K. Kaur, *Extending Quasi-alternating Links*, arXiv:2007.07770 [math.GT], 2020.
- [5] N. Chbili and K. Qazaqzeh, *On the Jones polynomial of quasi-alternating links*, Topol. Appl., **264** (2019): 1–11.
- [6] A. Issa, *The classification of quasi-alternating Montesinos links*, Proc. Amer. Math. Soc., **146**(9): 4047–4057, 2018.
- [7] J. Greene, *Homologically thin, non-quasi-alternating links*, Math. Res. Lett., **17**(1): 39–49, 2010.
- [8] T. Kanenobu: *Infinitely Many Knots with the Same Polynomial Invariant*, Proc. Amer. Math. Soc. **97** (1) (1986), 158–162.
- [9] M. Khovanov, *A categorification of the Jones polynomial*, Duke Math. J. **101**(3) (2000) 359–426.
- [10] M. Khovanov, *Patterns in knot cohomology. I*, Experiment. Math. **12**(3) (2003), 365–374.
- [11] E. S. Lee, *An endomorphism of the Khovanov invariant*, Adv. Math. **197**(2) (2005), 554–586.
- [12] C. Manolescu, *An unoriented skein exact triangle for knot Floer homology*, Math. Res. Lett. **14**(5) (2007), 839–852.
- [13] C. Manolescu and P. Ozsváth, *On the Khovanov and knot Floer homologies of quasi-alternating links*. In Proceedings of Gökova Geometry-Topology Conference (2007), 60–81. Gökova Geometry/Topology Conference (GGT), Gökova, 2008.
- [14] P. Ozsváth, J. Rasmussen and Z. Szabó, *Odd Khovanov homology*. arXiv:0710.4300, 2008.
- [15] P. Ozsváth and Z. Szabó, *On the Heegaard Floer homology of branched double-covers*, Adv. Math., **194**(1) (2005), 1–33.
- [16] K. Qazaqzeh and N. Chbili, *A new obstruction of quasi-alternating links*, Algebr. Geom. Topol. **15** (2015) 1847–1862.
- [17] K. Qazaqzeh, N. Chbili, and B. Qublan, *Characterization of quasi-alternating Montesinos links*, J. Knot Theory Ramifications **24** (1):155000 (13 pages), 2015.
- [18] K. Qazaqzeh, B. Qublan, and A. Jaradat, *A remark on the determinant of quasi-alternating links*, J. Knot Theory and its Ram. **22** (6) (2013), 1350031 (13 pages).
- [19] K. Qazaqzeh, and I. Mansour, *Further Study of Kanenobu Knots*, Kobe J. Math., **33** (2016) 3152.
- [20] M. Teragaito, *Quasi-alternating Links and  $Q$ -polynomials*, J. Knot Theory Ram. **23** (12) (2014), 1450068.
- [21] M. Thistlewaite, *A spanning tree expansion of the Jones polynomial*, Topology **26**: 297–309, 1988.

ADDRESS:- DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, KUWAIT UNIVERSITY P. O. BOX 5969 SAFAT-13060, KUWAIT, STATE OF KUWAIT

*E-mail address:* khaled.qazaqzeh@ku.edu.kw

DEPARTMENT OF MATHEMATICAL SCIENCES, COLLEGE OF SCIENCE, UAE UNIVERSITY, 15551 AL AIN, U.A.E.

*E-mail address:* nafaachbili@uaeu.ac.ae

*URL:* <http://faculty.uaeu.ac.ae/nafaachbili>