

CORRECTION TO: A DUAL ITERATIVE SUBSTRUCTURING METHOD WITH A SMALL PENALTY PARAMETER

CHANG-OCK LEE, EUN-HEE PARK, AND JONGHO PARK

ABSTRACT. In this corrigendum, we offer a correction to [J. Korean. Math. Soc., 54 (2017), pp. 461–477]. We construct a counterexample for the strengthened Cauchy–Schwarz inequality used in the original paper. In addition, we provide a new proof for Lemma 5 of the original paper, an estimate for the extremal eigenvalues of the standard unpreconditioned FETI-DP dual operator.

In the first and second authors' previous work [4], the strengthened Cauchy–Schwarz inequality used for [4, Eq. (3.8)] is incorrect and consequently, the statement of [4, Lemma 4] needs to be corrected. We present a new proof for [4, Lemma 5], that does not use [4, Lemma 4]. All notations are adopted from the original paper [4].

In the paragraph containing [4, Eq. (3.8)], it was claimed that by deriving a strengthened Cauchy–Schwarz inequality in a similar way to Lemma 4.3 in [3], it is shown that there exists a constant γ such that

$$2\tilde{a}(v_I + v_\Delta, v_c) \geq -\gamma(\tilde{a}(v_I + v_\Delta, v_I + v_\Delta) + \tilde{a}(v_c, v_c)),$$

where $0 < \gamma < 1$ is independent of H and h . That is, the above inequality is true when there exists a constant γ such that

$$(1) \quad |\tilde{a}(v_I + v_\Delta, v_c)| \leq \gamma(\tilde{a}(v_I + v_\Delta, v_I + v_\Delta))^{1/2} (\tilde{a}(v_c, v_c))^{1/2},$$

where $0 < \gamma < 1$ is independent of h and H .

On the other hand, a specific function $w = w_I + w_c + w_\Delta$ can be constructed, for which γ approaches 1 as H decreases. In fact, it suffices to characterize such w_Δ because w_I and w_c in (1) are determined by w_Δ in terms of the discrete \tilde{a} -harmonic extension $\mathcal{H}^c(w_\Delta)$.

Proposition 1. *There is no γ ($0 < \gamma < 1$), independent of h and H , satisfying (1).*

Proof. Noting that $\mathcal{H}^c(v_\Delta)$ in X_h^c is $\tilde{a}(\cdot, \cdot)$ -orthogonal to all the functions which vanish at the interface nodes except for the subdomain corners, we have that

$$\begin{aligned} \tilde{a}(v_I + v_\Delta, v_c) &= \tilde{a}(\mathcal{H}^c(v_\Delta) - v_c, v_c) \\ &= \tilde{a}(\mathcal{H}^c(v_\Delta), v_c) - \tilde{a}(v_c, v_c) \\ &= -\tilde{a}(v_c, v_c), \end{aligned}$$

which implies that for $\tilde{a}(v_I + v_\Delta, v_I + v_\Delta) \neq 0$, the estimate (1) is equivalent to

$$(2) \quad \frac{\tilde{a}(v_c, v_c)}{\tilde{a}(v_I + v_\Delta, v_I + v_\Delta)} \leq \gamma^2,$$

2020 *Mathematics Subject Classification.* 65F10, 65N30, 65N55.

Key words and phrases. domain decomposition, dual substructuring, FETI-DP.

where $\gamma < 1$ is independent of h and H .

Next, let us divide $\Omega = (0, 1)^2$ into $1/H \times 1/H$ square subdomains with a side length H . Each subdomain is partitioned into $2 \times H/h \times H/h$ uniform right triangles. Associated with such a triangulation, we select the function w in X_h^c such that w is a conforming \mathbb{P}_1 element function in each subdomain, and $w_\Delta = 1$ at all the nodes on the interface except for the subdomain corners. Hence, for w_c and w_I that are computed by the discrete harmonic extension of w_Δ , it is observed that

$$(3a) \quad w_c = 1 \text{ at all the subdomain corners } x_k \text{ that are not on } \partial\Omega,$$

$$(3b) \quad w_I = 1 \text{ in } \Omega_j \text{ for } \partial\Omega_j \cap \partial\Omega = \emptyset,$$

which imply that

$$(4) \quad w \equiv 1 \text{ in all subdomains whose boundary does not touch } \partial\Omega.$$

Let us first estimate $\tilde{a}(w_c, w_c)$ in (2). Using (3a), we have that

$$\tilde{a}(w_c, w_c) = \sum_{k=1}^{(1/H-1)^2} \tilde{a}(\phi_{c,k}, \phi_{c,k}) = 4 \left(\frac{1}{H} - 1 \right)^2,$$

where $\phi_{c,k}$ is the nodal basis function associated with the corner x_k . We next look over $\tilde{a}(w_I + w_\Delta, w_I + w_\Delta)$ based on the fact that, for $\partial\Omega_j \cap \partial\Omega = \emptyset$

$$(5) \quad \tilde{a}_{\Omega_j}(w_I + w_\Delta, w_I + w_\Delta) = \int_{\Omega_j} |\nabla(w_I + w_\Delta)|^2 dx = \int_{\Omega_j} |\nabla w_c|^2 dx = 4,$$

which follows from (4). Hence it suffices to estimate $\tilde{a}_{\Omega_j}(w_I + w_\Delta, w_I + w_\Delta)$ for the following two cases:

- (i) only one of the edges of the subdomain Ω_j is on $\partial\Omega$.
- (ii) two edges of the subdomain Ω_j are on $\partial\Omega$.

Here, the number of subdomains corresponding to the cases (i) and (ii) is $4 \left(\frac{1}{H} - 2 \right)$ and 4, respectively. Let us take $H/h = 3$ to focus only on the dependence of γ on either H or h . By finding the discrete local harmonic extensions for the cases (i) and (ii), it is computed directly that

$$(6) \quad \tilde{a}_{\Omega_j}(w_I + w_\Delta, w_I + w_\Delta) = \begin{cases} \frac{17}{4} & \text{for the case (i),} \\ \frac{14}{4} & \text{for the case (ii).} \end{cases}$$

Then by using (5) and (6), it follows that

$$(7) \quad \begin{aligned} \tilde{a}(w_I + w_\Delta, w_I + w_\Delta) &= \left(\sum_{\substack{j \text{ for} \\ \partial\Omega_j \cap \partial\Omega = \emptyset}} + \sum_{\substack{j \text{ for} \\ \partial\Omega_j \cap \partial\Omega \neq \emptyset}} \right) \tilde{a}_{\Omega_j}(w_I + w_\Delta, w_I + w_\Delta) \\ &= 4 \left(\frac{1}{H} - 2 \right)^2 + 17 \left(\frac{1}{H} - 2 \right) + 14. \end{aligned}$$

Finally, from (3a) and (7), it is confirmed that for a function w given above,

$$\lim_{H \rightarrow 0} \frac{\tilde{a}(w_c, w_c)}{\tilde{a}(w_I + w_\Delta, w_I + w_\Delta)} = 1,$$

which implies that (2) does not hold. \square

In [4, Lemma 5], the extremal eigenvalues of the FETI-DP dual operator $F = B_\Delta S^{-1} B_\Delta^T$ were estimated using [4, Lemma 4], estimates for the extremal eigenvalues of S . Since [4, Lemma 4] is incorrect, we provide a new estimate for F that does not utilize [4, Lemma 4]. We assume that each subdomain Ω_j is the union of elements in a conforming coarse mesh \mathcal{T}_H of Ω . First, we consider the following Poincaré-type inequality that generalizes [4, Proposition 3].

Lemma 2. *For any $v_j \in X_h^j$, let $I_j^H v_j$ be the linear coarse interpolation of v_j such that $I_j^H v_j = v_j$ at vertices of a subdomain $\Omega_j \subset \mathbb{R}^d$. Then we have*

$$|v_j|_{H^1(\Omega_j)}^2 \gtrsim \begin{cases} H^{-1} \left(1 + \ln \frac{H}{h}\right)^{-1} \|v_j - I_j^H v_j\|_{L^2(\partial\Omega_j)}^2 & \text{for } d = 2, \\ h^{-1} \left(\frac{H}{h}\right)^{-2} \|v_j - I_j^H v_j\|_{L^2(\partial\Omega_j)}^2 & \text{for } d = 3. \end{cases}$$

Proof. Since the both sides of the inequality do not change if a constant is added to v_j , we may assume that v_j has the zero average, so that the following Poincaré inequality holds:

$$(8) \quad \|v_j\|_{H^1(\Omega_j)} \lesssim |v_j|_{H^1(\Omega_j)},$$

where $\|\cdot\|_{H^1(\Omega_j)}$ is the weighted H^1 -norm on Ω_j given by

$$\|v_j\|_{H^1(\Omega_j)}^2 = |v_j|_{H^1(\Omega_j)}^2 + \frac{1}{H^2} \|v_j\|_{L^2(\Omega_j)}^2.$$

Since $I_j^H v_j$ attains its extremum at vertices, we have

$$(9) \quad \begin{aligned} \|v_j - I_j^H v_j\|_{L^2(\partial\Omega_j)} &\lesssim H^{\frac{d-1}{2}} \|v_j - I_j^H v_j\|_{L^\infty(\partial\Omega_j)} \\ &\leq H^{\frac{d-1}{2}} (\|v_j\|_{L^\infty(\partial\Omega_j)} + \|I_j^H v_j\|_{L^\infty(\partial\Omega_j)}) \\ &\lesssim H^{\frac{d-1}{2}} \|v_j\|_{L^\infty(\partial\Omega_j)}. \end{aligned}$$

Let $\mathcal{H}_j v_j$ be the generalized harmonic extension of $v_j|_{\partial\Omega_j}$ introduced in [7] such that $\mathcal{H}_j v_j = v_j$ on $\partial\Omega_j$ and

$$(10) \quad \|\mathcal{H}_j v_j\|_{H^1(\Omega_j)} = \min_{\substack{w_j \in H^1(\Omega_j) \\ w_j = v_j \text{ on } \partial\Omega_j}} \|w_j\|_{H^1(\Omega_j)}.$$

Then it follows that

$$\begin{aligned} (11a) \quad H^{d-1} \|v_j\|_{L^\infty(\partial\Omega_j)}^2 &\leq H^{d-1} \|\mathcal{H}_j v_j\|_{L^\infty(\Omega_j)}^2 \\ (11b) \quad &\lesssim C_d(H, h) \|\mathcal{H}_j v_j\|_{H^1(\Omega_j)}^2 \\ (11c) \quad &\leq C_d(H, h) \|v_j\|_{H^1(\Omega_j)}^2 \\ &\lesssim C_d(H, h) |v_j|_{H^1(\Omega_j)}^2, \end{aligned}$$

where

$$C_d(H, h) = \begin{cases} H \left(1 + \ln \frac{H}{h}\right) & \text{for } d = 2, \\ h \left(\frac{H}{h}\right)^2 & \text{for } d = 3, \end{cases}$$

and (11a) is due to the discrete Sobolev inequality [2, Lemma 2.3]. Also (10) and (8) are used in (11b) and (11c), respectively. Combination of (9) and (11) completes the proof. \square

Note that Lemma 2 reduces to [4, Proposition 3] when v_j vanishes at vertices of Ω_j so that $I_j^H v_j = 0$. Using Lemma 2, we obtain the following estimate for F .

Proposition 3. For $F = B_\Delta S^{-1} B_\Delta^T$, we have

$$\underline{C}_F \lambda^T \lambda \lesssim \lambda^T F \lambda \lesssim \overline{C}_F \lambda^T \lambda \quad \forall \lambda$$

where

$$\underline{C}_F = h^{2-d} \text{ for } d = 2, 3,$$

and

$$\overline{C}_F = \begin{cases} \left(\frac{H}{h}\right) \left(1 + \ln \frac{H}{h}\right) & \text{for } d = 2, \\ h^{-1} \left(\frac{H}{h}\right)^2 & \text{for } d = 3. \end{cases}$$

Consequently, the condition number of F satisfies the following bound:

$$\kappa(F) \lesssim \begin{cases} \left(\frac{H}{h}\right) \left(1 + \ln \frac{H}{h}\right) & \text{for } d = 2, \\ \left(\frac{H}{h}\right)^2 & \text{for } d = 3. \end{cases}$$

Proof. As the derivation of the maximum eigenvalue of S in the original paper [4] is correct, the derivation of \underline{C}_F is also correct. Thus, we only estimate \overline{C}_F in the following. We note that our proof closely follows [5, Theorem 4.5].

Similarly to [5, Theorem 4.4], it suffices to prove that

$$(12) \quad (B_\Delta \mathbf{v}_\Delta)^T (B_\Delta \mathbf{v}_\Delta) \lesssim \overline{C}_F \mathbf{v}_\Delta^T S \mathbf{v}_\Delta \quad \forall \mathbf{v}_\Delta.$$

If (12) were true, we get the desired result as follows:

$$\begin{aligned} \lambda^T F \lambda &= \max_{\mathbf{v}_\Delta \neq 0} \frac{((B_\Delta \mathbf{v}_\Delta)^T \lambda)^2}{\mathbf{v}_\Delta^T S \mathbf{v}_\Delta} \\ &\lesssim \overline{C}_F \max_{B_\Delta \mathbf{v}_\Delta \neq 0} \frac{((B_\Delta \mathbf{v}_\Delta)^T \lambda)^2}{(B_\Delta \mathbf{v}_\Delta)^T B_\Delta \mathbf{v}_\Delta} \\ &\leq \overline{C}_F \max_{\mu \neq 0} \frac{(\mu^T \lambda)^2}{\mu^T \mu} \\ &= \overline{C}_F \lambda^T \lambda, \end{aligned}$$

where we used [5, Lemma 4.3] in the first equality.

Take any v_Δ and its discrete \tilde{a} -harmonic extension $v = \mathcal{H}^c(v_\Delta)$. Let $w = v - I^H v$, where $I^H v$ is the linear coarse interpolation of v onto \mathcal{T}^H such that $I^H v = v$ at the subdomain vertices. We write $w = w_I + w_\Delta$. Since $I^H v$ is continuous along Γ , we have $B_\Delta \mathbf{w}_\Delta = B_\Delta \mathbf{v}_\Delta$. Then it follows that

$$\begin{aligned} (B_\Delta \mathbf{v}_\Delta)^T (B_\Delta \mathbf{v}_\Delta) &= (B_\Delta \mathbf{w}_\Delta)^T (B_\Delta \mathbf{w}_\Delta) \\ &= \sum_{j < k} \left(\mathbf{w}_\Delta^{(j)}|_{\Gamma_{jk}} - \mathbf{w}_\Delta^{(k)}|_{\Gamma_{jk}} \right)^T \left(\mathbf{w}_\Delta^{(j)}|_{\Gamma_{jk}} - \mathbf{w}_\Delta^{(k)}|_{\Gamma_{jk}} \right) \\ &\lesssim \sum_{j < k} \left(\left(\mathbf{w}_\Delta^{(j)}|_{\Gamma_{jk}} \right)^T \mathbf{w}_\Delta^{(j)}|_{\Gamma_{jk}} + \left(\mathbf{w}_\Delta^{(k)}|_{\Gamma_{jk}} \right)^T \mathbf{w}_\Delta^{(k)}|_{\Gamma_{jk}} \right) \\ &\lesssim \sum_{j=1}^{N_s} (\mathbf{w}_\Delta^{(j)})^T (\mathbf{w}_\Delta^{(j)}) \\ &\lesssim h^{1-d} \sum_{j=1}^{N_s} \|w\|_{L^2(\partial\Omega_j)}^2 \\ &\lesssim \overline{C}_F \mathbf{v}_\Delta^T S \mathbf{v}_\Delta, \end{aligned}$$

where the last inequality is due to Lemma 2. \square

It must be mentioned that the conclusion of Proposition 3 agrees with Lemma 5 of the original paper [4]. Since the conclusion of [4, Lemma 5] is true, it requires no additional correction in the remaining part of that paper.

For the sake of completeness, we present a correct estimate for the extremal eigenvalues of S that replaces [4, Lemma 4].

Proposition 4. *For $S = A_{\Delta\Delta} - A_{I\Delta}^T A_{II}^{-1} A_{I\Delta}$, we have*

$$\underline{C}_S \mathbf{v}_\Delta^T \mathbf{v}_\Delta \lesssim \mathbf{v}_\Delta^T S \mathbf{v}_\Delta \lesssim \overline{C}_S \mathbf{v}_\Delta^T \mathbf{v}_\Delta \quad \forall \mathbf{v}_\Delta,$$

where

$$\underline{C}_S = \begin{cases} Hh \left(1 + \ln \frac{H}{h}\right)^{-1} & \text{for } d = 2, \\ h^3 & \text{for } d = 3, \end{cases}$$

and

$$\overline{C}_S = h^{d-2} \text{ for } d = 2, 3.$$

Proof. Since the derivation of \overline{C}_S in the original paper [4] is correct, we only consider an estimate for \underline{C}_S . Take any \mathbf{v}_Δ and its corresponding finite element function v_Δ . Let $v = \mathcal{H}^c(v_\Delta)$ be the discrete \tilde{a} -harmonic extension of v_Δ . Proceeding as in [6, Lemma 4.11], we get

$$\begin{aligned} \mathbf{v}_\Delta^T \mathbf{v}_\Delta &\lesssim h^{1-d} \sum_{j=1}^{N_s} \|v_\Delta\|_{L^2(\partial\Omega_j)}^2 \\ &\lesssim Hh^{1-d} \sum_{j=1}^{N_s} \left(|v|_{H^1(\Omega_j)}^2 + H^{-2} \|v\|_{L^2(\Omega_j)}^2 \right) \\ &= Hh^{1-d} \mathbf{v}_\Delta^T S \mathbf{v}_\Delta + H^{-1} h^{1-d} \|v\|_{L^2(\Omega)}^2. \end{aligned}$$

Note that we cannot apply the discrete Poincaré inequality [1, Lemma 5.1] in each subdomain Ω_j since $\mathcal{H}v_\Delta$ does not vanish at the subdomain vertices in general.

It remains to show that

$$(13) \quad \|v\|_{L^2(\Omega)}^2 \lesssim \begin{cases} \left(1 + \ln \frac{H}{h}\right) \mathbf{v}_\Delta^T S \mathbf{v}_\Delta & \text{for } d = 2, \\ \frac{H}{h} \mathbf{v}_\Delta^T S \mathbf{v}_\Delta & \text{for } d = 3. \end{cases}$$

Let $I^H v$ be the linear nodal interpolation of v onto the coarse mesh \mathcal{T}_H . Since $I^H v$ is continuous along the subdomain interfaces Γ , we can apply the Poincaré inequality to obtain

$$\|I^H v\|_{L^2(\Omega)} \lesssim |I^H v|_{H^1(\Omega)}.$$

Then it follows that

$$\begin{aligned} \|v\|_{L^2(\Omega)}^2 &\lesssim \|v - I^H v\|_{L^2(\Omega)}^2 + \|I^H v\|_{L^2(\Omega)}^2 \\ &\lesssim \|v - I^H v\|_{L^2(\Omega)}^2 + |I^H v|_{H^1(\Omega)}^2 \\ &\lesssim \begin{cases} \left(1 + \ln \frac{H}{h}\right) \mathbf{v}_\Delta^T S \mathbf{v}_\Delta & \text{for } d = 2, \\ \frac{H}{h} \mathbf{v}_\Delta^T S \mathbf{v}_\Delta & \text{for } d = 3, \end{cases} \end{aligned}$$

where the last inequality is due to [6, Remark 4.13] for $d = 2$ and [6, Lemma 4.12] for $d = 3$, respectively. \square

REFERENCES

- [1] P. BOCHEV AND R. B. LEHOUCQ, *On the finite element solution of the pure Neumann problem*, SIAM Rev., 47 (2005), pp. 50–66.
- [2] J. H. BRAMBLE AND J. XU, *Some estimates for a weighted L^2 projection*, Math. Comp., 56 (1991), pp. 463–476.
- [3] C.-O. LEE AND E.-H. PARK, *A dual iterative substructuring method with a penalty term in three dimensions*, Comput. Math. Appl., 64 (2012), pp. 2787–2805.
- [4] ———, *A dual iterative substructuring method with a small penalty parameter*, J. Korean Math. Soc., 54 (2017), pp. 461–477.
- [5] J. MANDEL AND R. TEZAUR, *On the convergence of a dual-primal substructuring method*, Numer. Math., 88 (2001), pp. 543–558.
- [6] A. TOSELLI AND O. WIDLUND, *Domain Decomposition Methods—Algorithms and Theory*, Springer, Berlin, 2005.
- [7] J. XU AND J. ZOU, *Some nonoverlapping domain decomposition methods*, SIAM Rev., 40 (1998), pp. 857–914.

(Chang-Ock Lee) DEPARTMENT OF MATHEMATICAL SCIENCES, KAIST, DAEJEON 34141, KOREA
E-mail address: `collee@kaist.edu`

(Eun-Hee Park) DIVISION OF LIBERAL STUDIES, KANGWON NATIONAL UNIVERSITY, SAMCHEOK 25913, KOREA
E-mail address: `eh.park@kangwon.ac.kr`

(Jongho Park) DEPARTMENT OF MATHEMATICAL SCIENCES, KAIST, DAEJEON 34141, KOREA
E-mail address: `jongho.park@kaist.ac.kr`