

ON THE SPECTRAL ν -CONTINUITY

S. SÁNCHEZ-PERALES, S. PALAFOX, AND T. PÉREZ-BECERRA

ABSTRACT. In this paper we study the ν -continuity of the spectrum and some of its parts. We show that the approximate point spectrum σ_{ap} is upper semi- ν -continuous at every Fredholm operator, then we give sufficient conditions to guarantee the ν -continuity of σ_{ap} . Also we show that the restriction of the Weyl spectrum on the class of essentially G_1 operators is ν -continuous. Finally, we investigate the ν -continuity of the spectrum on the class of p -hyponormal operators.

1. INTRODUCTION

The spectral continuity is a relevant subject in Banach theory and operator theory. Several authors have studied this topic using different types of convergence. Regarding norm convergence, Newburgh in [17] was one of the first to do systematically investigated in general Banach algebras. More completely results in the case of the algebra $B(H)$ for H a Hilbert space are given in the series of papers of Conway and Morrel, see [7, 8, 9]. Other articles on this topic are [5, 10, 11, 12, 18, 21]. Another type of convergence for the study of spectral continuity is the ν -convergence given by Ahues in [1]. Also developed by Sánchez et al. [19, 20, 22] and Ammar et al. [3, 4].

In this paper we study the ν -continuity of the spectrum function. In section 2, we review this concept on arbitrary unital Banach algebras. We show that in an unital abelian Banach algebra the spectrum function is ν -continuous. In section 3, we study the ν -continuity of the spectrum and the approximate point spectrum on the algebra $B(X)$, for a Banach space X . We show that for the particular case of a Hilbert space H , the ν -continuity of spectrum is equivalent under certain conditions to the usual continuity of the spectrum. Also, we show that the approximate point spectrum is upper semi- ν -continuous at every Fredholm operator, then we give sufficient conditions to guarantee the ν -continuity of this function. In section 4, we show that the restriction of the Weyl spectrum on the class of essentially G_1 operators is ν -continuous. Finally, we investigate the ν -continuity of the spectrum on the class of p -hyponormal operators.

The next concepts are part of classical point set topology. Let (E_n) be a sequence of arbitrary subsets of \mathbb{C} and define the limits inferior and superior of (E_n) as follows:

1991 *Mathematics Subject Classification.* Primary 47A10; Secondary 47A53.

Key words and phrases. ν -convergence, spectral continuity, Fredholm operators.

$\liminf E_n = \{\lambda \in \mathbb{C} \mid \text{for every } \epsilon > 0, \text{ there exists } N \in \mathbb{N} \text{ such that } B(\lambda, \epsilon) \cap E_n \neq \emptyset \text{ for all } n \geq N\}.$

$\limsup E_n = \{\lambda \in \mathbb{C} \mid \text{for every } \epsilon > 0, \text{ there exists } J \subseteq \mathbb{N} \text{ infinite such that } B(\lambda, \epsilon) \cap E_n \neq \emptyset \text{ for all } n \in J\}.$

Remark 1.1. Let (E_n) be a sequence of non-empty subsets of \mathbb{C} . The following properties are hold:

- (a) $\liminf E_n$ and $\limsup E_n$ are closed subsets of \mathbb{C} .
- (b) $\lambda \in \limsup E_n$ if and only if there exists an increasing sequence of natural numbers $n_1 < n_2 < n_3 < \dots$ and points $\lambda_{n_k} \in E_{n_k}$, for all $k \in \mathbb{N}$, such that $\lim \lambda_{n_k} = \lambda$.
- (c) $\lambda \in \liminf E_n$ if and only if there exists a sequence $\{\lambda_n\}$ such that $\lambda_n \in E_n$ for all $n \in \mathbb{N}$, and $\lim \lambda_n = \lambda$.

Proposition 1.2. Let K, E, E_n be non-empty compact subsets of \mathbb{C} such that $E_n \subseteq K$, for all $n \in \mathbb{N}$. Then $E_n \rightarrow E$ in the Hausdorff metric if and only if $\limsup E_n \subseteq E$ and $E \subseteq \liminf E_n$.

2. SPECTRAL CONTINUITY ON COMPLEX BANACH ALGEBRAS

Let \mathcal{A} be a complex Banach algebra with identity $1_{\mathcal{A}}$. For $x \in \mathcal{A}$ the resolvent of x is defined by $\rho(x) = \{\lambda \in \mathbb{C} : x - \lambda 1_{\mathcal{A}} \text{ is invertible in } \mathcal{A}\}$ and the spectrum of x is given by $\sigma(x) = \mathbb{C} \setminus \rho(x)$. The spectral radius $r(x)$ of x is the number $r(x) = \max\{|\lambda| : \lambda \in \sigma(x)\}$. It is well known that $r(x) = \lim_{n \rightarrow \infty} \|x^n\|^{\frac{1}{n}} = \inf_n \|x^n\|^{\frac{1}{n}}$.

A sequence (x_n) in \mathcal{A} is said to be norm convergent to x (in notation $x_n \xrightarrow{n} x$), if $\|x_n - x\| \rightarrow 0$. M. Ahues in [1] introduces a new mode of convergence on $B(X)$, named ν -convergence. This type of convergence can be generalized in the same way to complex unital Banach algebras: a sequence (x_n) in \mathcal{A} is said to be ν -convergent to x , denoted by $x_n \xrightarrow{\nu} x$, if $(\|x_n\|)$ is bounded, $\|(x_n - x)x\| \rightarrow 0$ and $\|(x_n - x)x_n\| \rightarrow 0$. The ν -convergence is a pseudo-convergence in the sense that it is possible to have $x_n \xrightarrow{\nu} x$ and $x_n \xrightarrow{\nu} y$ but $x \neq y$, see for instance [1, Example 1]. There is a connection between norm convergence and ν -convergence as follows: if $x_n \xrightarrow{n} x$ then $x_n \xrightarrow{\nu} x$, also, if $x_n \xrightarrow{\nu} x$ and x is right invertible then $x_n \xrightarrow{n} x$.

A function τ defined on \mathcal{A} whose values are non-empty compact sets in \mathbb{C} is said to be continuous (ν -continuous) at x , if $\tau(x_n) \rightarrow \tau(x)$ with respect to the Hausdorff metric, for all sequence (x_n) in \mathcal{A} such that $x_n \xrightarrow{n} x$ ($x_n \xrightarrow{\nu} x$). It is clear that if τ is ν -continuous at x , then τ is continuous at x . τ is said to be upper semi-continuous (upper semi- ν -continuous) at x , if $\limsup \tau(x_n) \subseteq \tau(x)$ for all sequence (x_n) in \mathcal{A} such that $x_n \xrightarrow{n} x$ ($x_n \xrightarrow{\nu} x$). Also, τ is said to be lower semi-continuous (lower semi- ν -continuous) at x , if $\tau(x) \subseteq \liminf \tau(x_n)$ for all (x_n) in \mathcal{A} such that $x_n \xrightarrow{n} x$ ($x_n \xrightarrow{\nu} x$). From Proposition 1.2 we have next remark.

Remark 2.1. Let τ be a function defined on \mathcal{A} whose values are non-empty compact sets in \mathbb{C} such that $\tau(y) \subseteq \sigma(y)$ for all $y \in \mathcal{A}$. Then

- (1) τ is continuous at $x \in \mathcal{A}$ if and only if τ is upper and lower semi-continuous at x .
- (2) τ is ν -continuous at $x \in \mathcal{A}$ if and only if τ is upper and lower semi- ν -continuous at x .

Next theorem is proved in [1, Corollary 2.7].

Theorem 2.2. *For each $x \in \mathcal{A}$, σ is upper semi- ν -continuous at x .*

In the following proposition we will use the notation $\sigma_{\mathcal{B}}(x)$ for the spectrum of $x \in \mathcal{B}$ with respect to a given subalgebra \mathcal{B} of \mathcal{A} . A character on an abelian Banach algebra \mathcal{B} is a non-zero homomorphism $\varphi : \mathcal{B} \rightarrow \mathbb{C}$. The set of all characters on \mathcal{B} is denoted by $\mathcal{M}(\mathcal{B})$. It is well known that if \mathcal{B} is an unital abelian Banach algebra then

- (1) $\varphi(x) \in \sigma_{\mathcal{B}}(x)$ for all $x \in \mathcal{B}$ and $\varphi \in \mathcal{M}(\mathcal{B})$;
- (2) for every $\lambda \in \sigma_{\mathcal{B}}(x)$, there exists $\varphi \in \mathcal{M}(\mathcal{B})$ such that $\varphi(x) = \lambda$;
- (3) for each $\varphi \in \mathcal{M}(\mathcal{B})$, $\|\varphi\| = 1$.

A very well known result in spectral continuity is that if the elements of a sequence (x_n) in a Banach algebra \mathcal{A} commute with their limit, i.e. if $x_n \xrightarrow{n} x$ and $x_n x = x x_n$, then $\sigma(x_n) \rightarrow \sigma(x)$. We generalize this result for the ν -convergence, as shown in the following proposition.

Proposition 2.3. *Let $a \in \mathcal{A}$ and (a_n) be a sequence in \mathcal{A} such that $a_n a = a a_n$ and $a_n a_m = a_m a_n$ for all $n, m \in \mathbb{N}$. If $a_n \xrightarrow{\nu} a$ and $0 \in \text{acc } \sigma(a)$, then $\sigma(a_n) \rightarrow \sigma(a)$.*

Proof. By Theorem 2.2, σ is upper semi- ν -continuous at a , thus $\limsup \sigma_{\mathcal{A}}(a_n) \subseteq \sigma_{\mathcal{A}}(a)$. Hence by Proposition 1.2, we only need to prove that $\sigma_{\mathcal{A}}(a) \subseteq \liminf \sigma_{\mathcal{A}}(a_n)$.

Consider \mathcal{B}_0 the subalgebra of \mathcal{A} which consists of all linear combinations of finite products of elements in $\{a_n : n \in \mathbb{N}\} \cup \{a, 1_{\mathcal{A}}\}$. From hypothesis, \mathcal{B}_0 is commutative. Thus by Zorn's lemma, there exists \mathcal{B} the maximal abelian subalgebra of \mathcal{A} such that $\mathcal{B}_0 \subseteq \mathcal{B}$. Therefore by [16, Exercise 8, p. 8], $\sigma_{\mathcal{A}}(a) = \sigma_{\mathcal{B}}(a)$ and $\sigma_{\mathcal{A}}(a_n) = \sigma_{\mathcal{B}}(a_n)$ for all $n \in \mathbb{N}$. Let $\lambda \in \sigma_{\mathcal{A}}(a)$ with $\lambda \neq 0$. Then there exists $\varphi \in \mathcal{M}(\mathcal{B})$ such that $\varphi(a) = \lambda$. Observe that

$$\|(\varphi(a_n) - \varphi(a))\varphi(a)\| = \|\varphi((a_n - a)a)\| \leq \|\varphi\| \|(a_n - a)a\| \rightarrow 0.$$

Thus $\|(\varphi(a_n) - \varphi(a))\lambda\| \rightarrow 0$ which implies that $\varphi(a_n) \rightarrow \lambda$. Now, since $\varphi(a_n) \in \sigma_{\mathcal{B}}(a_n) (= \sigma_{\mathcal{A}}(a_n))$ for all $n \in \mathbb{N}$, it follows from Remark 1.1 that $\lambda \in \liminf \sigma_{\mathcal{A}}(a_n)$. Consequently, since $0 \in \text{acc } \sigma_{\mathcal{A}}(a)$,

$$\sigma_{\mathcal{A}}(a) = \overline{\sigma_{\mathcal{A}}(a) \setminus \{0\}} \subseteq \liminf \sigma_{\mathcal{A}}(a_n).$$

□

An elementary Cauchy domain is a bounded open connected subset of \mathbb{C} whose boundary is the union of a finite number of nonintersecting Jordan

curves. A finite union of elementary Cauchy domains having disjoint closures is called a Cauchy domain. Let D be a Cauchy domain, if each curve involved in the boundary of D is oriented in such a way that points in D lie to the left as the curve is traced out, then the oriented boundary C of D is called a Cauchy contour. The interior of the Cauchy contour C is defined as $\text{int}(C) = D$ and the exterior of C is defined by $\text{ext}(C) = \mathbb{C} \setminus (D \cup C)$.

A set $\Lambda \subseteq \sigma(a)$ is a spectral set for a if Λ is closed as well as open in $\sigma(a)$. We set $\mathcal{C}(a, \Lambda)$ the set of all Cauchy contours C separating Λ from $\sigma(a) \setminus \Lambda$, i.e. $\Lambda \subseteq \text{int}(C)$ and $\sigma(a) \setminus \Lambda \subseteq \text{ext}(C)$. It is clear that if $C \in \mathcal{C}(a, \Lambda)$ then $C \subseteq \rho(a)$. For a spectral set Λ for a and $C \in \mathcal{C}(a, \Lambda)$, define

$$p(a, \Lambda) = -\frac{1}{2\pi i} \int_C (a - z)^{-1} dz.$$

The element $p(a, \Lambda)$ does not depend on the choice of $C \in \mathcal{C}(a, \Lambda)$.

Remark 2.4. Let $a \in \mathcal{A}$, Λ a spectral set for a and $C \in \mathcal{C}(a, \Lambda)$. If $p = p(a, \Lambda)$, then

- (1) $p^2 = p$ and $pa = ap$
- (2) $\Lambda = \emptyset$ if and only if $p(a, \Lambda) = 0$.

Proposition 2.5. Let $p, q \in \mathcal{A}$ be such that $p^2 = p$. If $p \neq 0$ and $r(p - q) < 1$ then $q \neq 0$.

Proof. Since $r(p - q) < 1$ it follows that $(p - q) - 1_{\mathcal{A}}$ is invertible. Suppose that $q = 0$, then $p - 1_{\mathcal{A}}$ is invertible. Thus there exists $z \in \mathcal{A}$ such that $(p - 1_{\mathcal{A}})z = 1_{\mathcal{A}}$. This implies that $p(p - 1_{\mathcal{A}})z = p1_{\mathcal{A}}$ and so $(p^2 - p)z = p$ i.e. $0 = (p - p)z = p$, which is a contradiction. \square

Theorem 2.6. [1, Proposition 2.9] Let $a \in \mathcal{A}$, Λ be a spectral set for a , $C \in \mathcal{C}(a, \Lambda)$ and (a_n) be a sequence in \mathcal{A} such that $a_n \xrightarrow{\nu} a$. Then

- (1) There exists $n_0 \in \mathbb{N}$ such that for every $n \geq n_0$, C lies in $\rho(a_n)$.
- (2) If $\Lambda_n := \sigma(a_n) \cap \text{int}(C)$ for all $n \geq n_0$, then Λ_n is a spectral set for a_n and $C \in \mathcal{C}(a_n, \Lambda_n)$. Further, if $0 \in \text{ext}(C)$ then $p(a_n, \Lambda_n) \xrightarrow{\nu} p(a, \Lambda)$.

Next lemma is a generalization of [7, Lemma 1.5].

Lemma 2.7. Let $a \in \mathcal{A}$ and (a_n) be a sequence in \mathcal{A} such that $a_n \xrightarrow{\nu} a$. If U is an open set for which $0 \notin U$ and U contains a component of $\sigma(a)$, then there exists $n_0 \in \mathbb{N}$ such that U contains a component of $\sigma(a_n)$ for all $n \geq n_0$.

Proof. Let Ω be a component of $\sigma(a)$ and U be an open set of \mathbb{C} such that $0 \notin U$ and $\Omega \subseteq U$. Since $\Omega \cap [\sigma(a) \setminus U] = \emptyset$, $\sigma(a) \setminus U$ is closed and $\sigma(a)$ is compact, it follows that there exists $\Lambda \neq \emptyset$ open and closed set in $\sigma(a)$ such that $\Omega \subseteq \Lambda$ and $\Lambda \cap [\sigma(a) \setminus U] = \emptyset$. This implies that $\Lambda \subseteq U$. From [1, Theorem 1.21], there exists a Cauchy domain D such that

$$\Lambda \subset D \quad \text{and} \quad \overline{D} \subset U \cap [\mathbb{C} \setminus (\sigma(a) \setminus \Lambda)]. \quad (2.1)$$

Let C be the boundary of D oriented in way that C be a Cauchy contour. It is clear by (2.1) that $C \in \mathcal{C}(a, \Lambda)$. Then from Theorem 2.6, there exists $n_1 \in \mathbb{N}$ such that for every $n \geq n_1$, C lies in $\rho(a_n)$. Further, if $\Lambda_n := \sigma(a_n) \cap D$ for all $n \geq n_1$ then since $0 \notin U$ we have that

$$[r(p - p_n)]^2 \leq \|(p - p_n)^2\| \leq \|(p_n - p)p_n\| + \|(p_n - p)p\| \rightarrow 0,$$

where $p_n = p(a_n, \Lambda_n)$ and $p = p(a, \Lambda)$. Thus there exists $n_0 \in \mathbb{N}$ with $n_0 \geq n_1$ such that $r(p - p_n) < 1$ for all $n \geq n_0$. Since $\Lambda \neq \emptyset$ we have from Remark 2.4 that $p \neq 0$. Therefore, by Proposition 2.5, $p_n \neq 0$ for all $n \geq n_0$. Thus by Remark 2.4, $\Lambda_n \neq \emptyset$ for all $n \geq n_0$. This implies, since Λ_n is both open and closed in $\sigma(a_n)$, that there exists Ω_n a component of $\sigma(a_n)$ such that $\Omega_n \subseteq \Lambda_n$. Observe that $\Lambda_n \subseteq D \subseteq U$. Thus $\Omega_n \subseteq U$. Therefore U contains a component of $\sigma(a_n)$ for all $n \geq n_0$. \square

Remark 2.8. From Lemma 2.7 we have that if $\lambda \in \text{iso } \sigma(a)$ with $\lambda \neq 0$, then $\lambda \in \liminf \sigma(a_n)$ for all sequence (a_n) in \mathcal{A} such that $a_n \xrightarrow{\nu} a$.

3. SPECTRAL CONTINUITY ON THE ALGEBRA $B(X)$

Let X be a Banach space and let $B(X)$ be the algebra of all bounded linear operators defined on X . For $T \in B(X)$, let $N(T)$ and $R(T)$ be denote the null space and the range of the mapping T . Let $\alpha(T) = \dim N(T)$ and $\beta(T) = \dim X/R(T)$, if these spaces are finite dimensional, otherwise let $\alpha(T) = \infty$ and $\beta(T) = \infty$. If the range $R(T)$ of $T \in B(X)$ is closed and $\alpha(T) < \infty$ then T is said to be an *upper semi-Fredholm* operator ($T \in \Phi_+(X)$). Similarly, if $\beta(T) < \infty$ then T is said to be a *lower semi-Fredholm* operator ($T \in \Phi_-(X)$). If $T \in \Phi_-(X) \cup \Phi_+(X)$ then T is called a *semi-Fredholm* operator ($T \in \Phi_{\pm}(X)$) and for $T \in \Phi_-(X) \cap \Phi_+(X)$ we say that T is a *Fredholm* operator ($T \in \Phi(X)$). For $T \in \Phi_{\pm}(X)$, the *index* of T is defined by $i(T) = \alpha(T) - \beta(T)$. It is well known that the index is a continuous function on the set of semi-Fredholm operators. This property also holds for the ν -convergence in the following sense:

Theorem 3.1. [19, Theorem 3.4] *Let $T \in \Phi(X)$ and (T_n) be a sequence in $B(X)$ such that $(T_n - T)T \xrightarrow{n} 0$. If λ, λ_n are complex numbers such that $\lambda_n \rightarrow \lambda$ and*

$$T - \lambda, T_n - \lambda_n \in \Phi_{\pm}(X)$$

for all $n \in \mathbb{N}$, then $i(T_n - \lambda_n) \rightarrow i(T - \lambda)$.

For $T \in B(X)$ and $k \in \mathbb{N} \cup \{-\infty, \infty\}$, let $\rho_{sf}^k(T)$ be denote the set of $\lambda \in \mathbb{C}$ for which $T - \lambda \in \Phi_{\pm}(X)$ and $i(T - \lambda) = k$. Put

$$\rho_{sf}^-(T) = \bigcup_{-\infty \leq k \leq -1} \rho_{sf}^k(T), \quad \rho_{sf}^+(T) = \bigcup_{1 \leq k \leq \infty} \rho_{sf}^k(T), \quad \rho_{sf}^{\pm}(T) = \rho_{sf}^-(T) \cup \rho_{sf}^+(T).$$

All these sets are open and are contained in $\sigma(T)$. Let $\sigma_{sf}(T)$ be denote the set of all $\lambda \in \mathbb{C}$ such that $T - \lambda \notin \Phi_{\pm}(X)$. Let $K(X)$ be denote the set of all compact linear operators in $B(X)$. If $\pi : B(X) \rightarrow B(X)/K(X)$ is the canonical homomorphism, then the essential spectrum of an operator $T \in$

$B(X)$, $\sigma_e(T)$, is the spectrum of $\pi(T)$ in the Calkin algebra $B(X)/K(X)$. Also, the left essential spectrum $\sigma_{le}(T)$ (the right essential spectrum $\sigma_{re}(T)$) is the left spectrum (right spectrum) of $\pi(T)$. We set $\sigma_{lre}(T) = \sigma_{le}(T) \cap \sigma_{re}(T)$. It is clear that

$$\sigma_{sf}(T) \subseteq \sigma_{lre}(T) \quad (3.1)$$

but the opposite inclusion is not always satisfied in general Banach spaces. These classes of sets coincide in the case of Hilbert spaces.

The approximate point spectrum, the surjective spectrum, the point spectrum, the Weyl spectrum and the set Riesz points of $T \in B(X)$ are defined by $\sigma_{ap}(T) = \{\lambda \in \mathbb{C} : T - \lambda \text{ is not bounded below}\}$, $\sigma_s(T) = \{\lambda \in \mathbb{C} : T - \lambda \text{ is not surjective}\}$, $\sigma_p(T) = \{\lambda \in \mathbb{C} : \lambda \text{ is an eigenvalue of } T\}$, $\sigma_w(T) = \{\lambda \in \mathbb{C} : T - \lambda \text{ is not a Fredholm operator of index zero}\}$ and $\pi_0(T) = \{\lambda \in \mathbb{C} : \lambda \text{ is an isolated eigenvalue of } T \text{ of finite algebraic multiplicity}\}$.

Remark 3.2. Let $T \in B(X)$ and (T_n) be a sequence in $B(X)$ such that $T_n \xrightarrow{\nu} T$. The following inclusions are hold:

- (1) $\pi_0(T) \subseteq \liminf \pi_0(T_n)$. See, [1, Corollary 2.13].
- (2) $\Delta \subseteq \liminf \sigma(T_n)$, where $\Delta = \{\lambda \in \sigma(T) \setminus (\overline{\rho_{sf}^\pm(T)} \cup \{0\}) : \text{for every } \epsilon > 0, \text{ the ball } B(\lambda, \epsilon) \text{ contains a component of } \sigma_{sf}(T) \cup \pi_0(T) \text{ and } T \text{ satisfies Browder's theorem. Really, take } \lambda \in \Delta \text{ then } \lambda \in \sigma(T) \setminus \overline{\rho_{sf}^\pm(T)} \text{ and } \lambda \neq 0, \text{ so there exists } r > 0 \text{ such that } B(\lambda, r) \cap \overline{\rho_{sf}^\pm(T)} = \emptyset \text{ and } 0 \notin B(\lambda, r). \text{ Let } \epsilon > 0 \text{ with } \epsilon < r, \text{ since } \lambda \in \Delta, \text{ the ball } B(\lambda, \epsilon) \text{ contains a component } \Omega \text{ of } \sigma_{sf}(T) \cup \pi_0(T). \text{ By [21, Lemma 3.6], } \Omega \text{ is a component of } \sigma(T). \text{ Therefore by Lemma 2.7, there exists } n_0 \in \mathbb{N} \text{ such that } B(\lambda, \epsilon) \text{ contains a component } \Omega_n \text{ of } \sigma(T_n) \text{ for all } n \geq n_0. \text{ Thus } B(\lambda, \epsilon) \cap \sigma(T_n) \neq \emptyset \text{ for all } n \geq n_0. \text{ This implies that } \lambda \in \liminf \sigma(T_n).$
- (3) $[\text{iso } \sigma(T)] \setminus \{0\} \subseteq \liminf \sigma(T_n)$.

Moreover, if $T \in \Phi(X)$ then

- (4) $\overline{\rho_{sf}^+(T)} \subseteq \liminf \sigma_a(T_n)$. See, [19, Theorem 3.6].
- (5) $\overline{\rho_{sf}^-(T)} \subseteq \liminf \sigma_s(T_n)$.
- (6) $\overline{\rho_{sf}^\pm(T)} \subseteq \liminf \sigma_w(T_n)$.
- (7) $\Gamma \subseteq \liminf \sigma_{sf}(T_n)$, where $\Gamma = \{\lambda \in \sigma(T) : \text{for every } \epsilon > 0, \text{ there exist points } \mu_1, \mu_2 \in B(\lambda, \epsilon) \text{ such that } T - \mu_1, T - \mu_2 \in \Phi_\pm(X) \text{ and } i(T - \mu_1) \neq i(T - \mu_2)\}$. This result was established in [8, Proposition 2.1] for the norm convergence. The proof for the ν -convergence is in a similar way.

Theorem 3.3. Let $T \in \Phi(X)$ be such that $0 \in \text{acc}\sigma(T)$. If for each $\lambda \in \sigma(T) \setminus \overline{\rho_{sf}^\pm(T)}$ and $\epsilon > 0$, the ball $B(\lambda, \epsilon)$ contains a component of $\sigma_{sf}(T) \cup \pi_0(T)$, then σ is ν -continuous at T .

Proof. First observe that $\sigma(T) \setminus \sigma_w(T) = \pi_0(T) \cup \text{int}[\sigma(T) \setminus \sigma_w(T)]$, see for example [18, Proposition 2.1]. By hypothesis we have that $\text{int}[\sigma(T) \setminus \sigma_w(T)] = \emptyset$, thus T satisfies Browder's theorem.

From Remark 2.1 and Theorem 2.2 it is sufficient to prove that σ is lower semi- ν -continuous at T . Let (T_n) be a sequence in $B(X)$ such that $T_n \xrightarrow{\nu} T$. By Remark 3.2 (6), $\overline{\rho_{sf}^{\pm}(T)} \subseteq \liminf \sigma(T_n)$. Now, from hypothesis, $\sigma(T) \setminus (\overline{\rho_{sf}^{\pm}(T)} \cup \{0\}) \subseteq \Delta$, therefore by Remark 3.2 (2), $\sigma(T) \setminus (\overline{\rho_{sf}^{\pm}(T)} \cup \{0\}) \subseteq \liminf \sigma(T_n)$. Consequently, $\sigma(T) = \overline{\sigma(T) \setminus \{0\}} \subseteq \liminf \sigma(T_n)$. Thus σ is lower semi- ν -continuous at T . \square

Corollary 3.4. *Let H be a Hilbert space and $T \in \Phi(H)$ be such that $0 \in \text{acc}\sigma(T)$. Then, σ is continuous at T if and only if σ is ν -continuous at T .*

Proof. It is clear that the ν -continuity of σ at T implies the continuity of σ at T . Now, if σ is continuous at T , then by [7, Theorem 3.1], for each $\lambda \in \sigma(T) \setminus \overline{\rho_{sf}^{\pm}(T)}$ and $\epsilon > 0$, the ball $B(\lambda, \epsilon)$ contains a component of $\sigma_{sf}(T) \cup \pi_0(T)$. Therefore, by Theorem 3.3, σ is ν -continuous at T . \square

Theorem 3.5. *Let $T \in \Phi(X)$. If (T_n) is a sequence in $B(X)$ such that $T_n \xrightarrow{\nu} T$ then*

$$[\limsup \sigma_{ap}(T_n)] \setminus \{0\} \subseteq \sigma_{ap}(T).$$

Proof. Let D, E be closed subspaces of X with $\dim E < \infty$ such that

$$X = N(T) \oplus D \quad \text{and} \quad X = R(T) \oplus E. \quad (3.2)$$

Consider (T_n) a sequence in $B(X)$ which is ν -convergent to T . Let $\lambda \in \limsup \sigma_{ap}(T_n)$ with $\lambda \neq 0$. By Remark 1.1, there exist an increasing sequence of natural numbers (n_k) and points $\lambda_{n_k} \in \sigma_{ap}(T_{n_k})$ such that $\lambda_{n_k} \rightarrow \lambda$. Suppose that $\lambda \notin \sigma_{ap}(T)$. Then $T - \lambda \in \Phi_+(X)$ and $\alpha(T - \lambda) = 0$. By (3.2), $R(T|_D) = R(T)$ and so $T|_D$ is bounded below, therefore by [23, Theorem 5.26], $(T - \lambda)T|_D \in \Phi_+(D, X)$ and $\alpha((T - \lambda)T|_D) = 0$.

On the other hand, observe that

$$(T_{n_k} - \lambda_{n_k})T|_D = (T - \lambda)T|_D + (T_{n_k} - T)T|_D + (\lambda - \lambda_{n_k})T|_D.$$

From $\|(T_n - T)T\| \rightarrow 0$ we have that $(T_{n_k} - \lambda_{n_k})T|_D$ converges in norm to $(T - \lambda)T|_D$. Consequently by [23, Theorem 5.23], there exists $k_0 \in \mathbb{N}$ such that every $k \geq k_0$, $(T_{n_k} - \lambda_{n_k})T|_D \in \Phi_+(D, X)$ and $\alpha((T_{n_k} - \lambda_{n_k})T|_D) = 0$.

Suppose that for each $k \geq k_0$,

$$N(T_{n_k} - \lambda_{n_k}) \cap E \neq \{0\}.$$

Take $v_k \in N(T_{n_k} - \lambda_{n_k}) \cap E$ with $\|v_k\| = 1$ for all $k \geq k_0$. Since $\dim E < \infty$ it follows that $F := \{e \in E : \|e\| = 1\}$ is a compact set. Therefore we may assume without loss of generality that there exists $v \in F$ such that $v_k \rightarrow v$.

Observe that

$$\begin{aligned} \|(T_{n_k} - T)T_{n_k}\| &\geq \|(T_{n_k} - T)T_{n_k}v_k\| \\ &= \|(T_{n_k} - T)\lambda_{n_k}v_k\| \\ &= |\lambda_{n_k}| \|\lambda_{n_k}v_k - Tv_k\| \end{aligned}$$

for all $k \geq k_0$, and $|\lambda_{n_k}| \|\lambda_{n_k}v_k - Tv_k\| \rightarrow |\lambda| \|\lambda v - Tv\|$. This implies that

$$|\lambda| \|\lambda v - Tv\| = \lim |\lambda_{n_k}| \|\lambda_{n_k}v_k - Tv_k\| \leq \lim \|(T_{n_k} - T)T_{n_k}\| = 0,$$

and so $\|\lambda v - Tv\| = 0$, i.e. $Tv = \lambda v$. Consequently, $\lambda \in \sigma_p(T) (\subseteq \sigma_{ap}(T))$, which is a contradiction. Therefore there exists $k' \geq k_0$ such that $N(T_{n_{k'}} - \lambda_{n_{k'}}) \cap E = \{0\}$. Thus

$$X = R(T) \oplus N(T_{n_{k'}} - \lambda_{n_{k'}}) \oplus E. \quad (3.3)$$

Then by (3.2) and (3.3),

$$\begin{aligned} \dim E &= \dim X/R(T) = \dim[N(T_{n_{k'}} - \lambda_{n_{k'}}) \oplus E] \\ &= \dim N(T_{n_{k'}} - \lambda_{n_{k'}}) + \dim E. \end{aligned}$$

Hence $\dim N(T_{n_{k'}} - \lambda_{n_{k'}}) = 0$ and so $N(T_{n_{k'}} - \lambda_{n_{k'}}) = \{0\}$. From (3.3),

$$R(T_{n_{k'}} - \lambda_{n_{k'}}) = (T_{n_{k'}} - \lambda_{n_{k'}})T(D) + (T_{n_{k'}} - \lambda_{n_{k'}})(E),$$

which implies that $R(T_{n_{k'}} - \lambda_{n_{k'}})$ is closed. Therefore $\lambda_{n_{k'}} \notin \sigma_{ap}(T_{n_{k'}})$, a contradiction. Consequently, $\lambda \in \sigma_{ap}(T)$. \square

Let $\phi_+(T)$ be the set of $\lambda \in \rho_{sf}^+(T)$ such that $N(T - \lambda)$ is complemented in X and $\phi_-(T)$ the set of $\lambda \in \rho_{sf}^-(T)$ such that $R(T - \lambda)$ is complemented in X . We set $\phi_\pm(T) = \phi_+(T) \cup \phi_-(T)$, $\phi_{+\infty}(T) = \{\lambda \in \phi_+(T) \mid i(T - \lambda) = \infty\}$, $\phi_{-\infty}(T) = \{\lambda \in \phi_-(T) \mid i(T - \lambda) = -\infty\}$ and $\phi_{\pm\infty}(T) = \phi_{+\infty}(T) \cup \phi_{-\infty}(T)$. All these sets are open and it is clear that

$$\sigma_e(T) = \sigma_{lre}(T) \cup \phi_{\pm\infty}(T). \quad (3.4)$$

With this equality, [7, Proposition 1.3] can be extended to general Banach spaces:

Proposition 3.6. *If C is a component of $\sigma_{lre}(T)$ and $C \cap \overline{\phi_\pm(T)} = \emptyset$ then C is a component of $\sigma_e(T)$.*

Theorem 3.7. *If the following conditions hold*

- i. $0 \in \text{acc}\sigma_{ap}(T)$
- ii. $T \in \Phi(X)$
- iii. T satisfies Browder's theorem
- iv. $\sigma_p(T) \cap \phi_-(T) = \emptyset$
- v. for every $\epsilon > 0$ and $\lambda \in \sigma_{lre}(T) \setminus \overline{\phi_\pm(T)}$, the ball $B(\lambda, \epsilon)$ contains a component of $\sigma_{lre}(T)$,

then $\sigma_{ap}(T_n) \rightarrow \sigma_{ap}(T)$ for all $T_n \xrightarrow{\nu} T$.

Proof. From conditions (i) and (ii) we have by Theorem 3.5 that

$$\limsup \sigma_{ap}(T_n) \subseteq \sigma_{ap}(T).$$

Now, we show that $\sigma_{ap}(T) \subseteq \liminf \sigma_{ap}(T_n)$. Take $\lambda \in \sigma_{ap}(T)$ with $\lambda \neq 0$. Suppose that $\lambda \in \sigma_{lre}(T) \setminus \overline{\phi_{\pm}(T)}$. Let $\epsilon > 0$, there exists $0 < \epsilon_1 < \epsilon$ such that $B(\lambda, \epsilon_1) \cap \overline{\phi_{\pm}(T)} = \emptyset$ and $0 \notin B(\lambda, \epsilon_1)$. By hypothesis (v), $B(\lambda, \epsilon_1)$ contains a component Ω of $\sigma_{lre}(T)$. Then by Proposition 3.6, Ω is a component of $\sigma_e(T)$. Since $T_n \xrightarrow{\nu} T$ we have that $\pi(T_n) \xrightarrow{\nu} \pi(T)$ in the Calkin algebra $B(X)/K(X)$. Thus by Lemma 2.7, there exists $n_0 \in \mathbb{N}$ such that $B(\lambda, \epsilon_1)$ contains a component Ω_n of $\sigma_e(T_n)$ for all $n \geq n_0$. Consequently

$$\emptyset \neq \partial\Omega_n \subseteq \partial\sigma_e(T_n) \cap B(\lambda, \epsilon) \subseteq \sigma_{sf}(T_n) \cap B(\lambda, \epsilon)$$

for all $n \geq n_0$. Therefore $\lambda \in \liminf \sigma_{sf}(T_n) (\subseteq \liminf \sigma_{ap}(T_n))$.

If $\lambda \notin \sigma_{lre}(T) \setminus \overline{\phi_{\pm}(T)}$, then $\lambda \in \phi_{\pm}(T)$ or $i(T - \lambda) = 0$. From condition (iv), we have that $\lambda \notin \phi_-(T)$. Thus from condition (iii), $\lambda \in \phi_+(T) \cup \pi_0(T)$. Therefore, by Remark 3.2, $\lambda \in \liminf \sigma_{ap}(T_n)$. \square

4. ON CERTAIN CLASS OF OPERATORS

We say that an operator $T \in B(H)$ is essentially G_1 if $\|(\pi(T) - z)^{-1}\| = \frac{1}{d(z, \sigma_e(T))}$ for all $z \notin \sigma_e(T)$. In [15, Theorem 6] it is shown that the restriction of the Wey spectrum on the class of essentially G_1 operators is continuous. This is also true for ν -continuity, as the following theorem states.

Theorem 4.1. *Let $T \in \Phi(H)$ be such that $0 \notin \sigma_w(T)$ or $0 \in \text{acc}\sigma_w(T)$. If (T_n) is a sequence of essentially G_1 operators such that $T_n \xrightarrow{\nu} T$ then $\sigma_w(T_n) \rightarrow \sigma_w(T)$.*

Proof. From [3, Theorem 3.3] we have that $\limsup \sigma_w(T_n) \subseteq \sigma_w(T)$. We show that $\sigma_w(T) \subseteq \liminf \sigma_w(T_n)$. Let $\lambda \in \sigma_w(T) \setminus \{0\}$ and suppose that $\lambda \notin \liminf \sigma_w(T_n)$. Then, there exist $\epsilon > 0$ such that $B_{\epsilon}(\lambda) \cap \sigma_w(T_n) = \emptyset$ for infinite number of n 's. Without loss of generality assume that this holds for all n . This implies that $d(\lambda, \sigma_e(T_n)) \geq d(\lambda, \sigma_w(T_n)) \geq \epsilon > 0$. Thus $\lambda \notin \sigma_e(T_n)$, now from the fact that T_n is essentially G_1 , we have that $\|(\pi(T_n) - \lambda)^{-1}\| = \frac{1}{d(\lambda, \sigma_e(T_n))} \leq \frac{1}{\epsilon}$. This implies that

$$\begin{aligned} & \left\| \left[(\pi(T_n) - \pi(T))(\pi(T_n) - \lambda)^{-1} \right]^2 \right\| \leq \\ & \leq \frac{1}{|\lambda|} \left[\|(\pi(T_n) - \pi(T))\pi(T_n)\| \|(\pi(T_n) - \lambda)^{-1}\| \|\pi(T_n) - \pi(T)\| \right. \\ & \quad \left. + \|(\pi(T_n) - \pi(T))\pi(T)\| + \|(\pi(T_n) - \pi(T))\pi(T_n)\| \right] \|(\pi(T_n) - \lambda)^{-1}\| \\ & \leq \frac{1}{|\lambda|} \left[\|(T - T_n)T_n\| \|T_n - T\| \frac{1}{\epsilon} + \|(T - T_n)T\| + \|(T - T_n)T_n\| \right] \frac{1}{\epsilon}. \quad (4.1) \end{aligned}$$

Now, since $T_n \xrightarrow{\nu} T$, it follows that $\|(T_n - T)T\| \rightarrow 0$, $\|(T_n - T)T_n\| \rightarrow 0$ and $(\|T_n\|)$ is bounded. Therefore the expression (4.1) tends to 0. Thus,

there exists $n^* \in \mathbb{N}$ such that

$$\left\| \left[(\pi(T_{n^*}) - \pi(T))(\pi(T_{n^*}) - \lambda)^{-1} \right]^2 \right\| < 1.$$

Then $r\left((\pi(T_{n^*}) - \pi(T))(\pi(T_{n^*}) - \lambda)^{-1}\right) < 1$, which implies that $(\pi(T_{n^*}) - \pi(T))(\pi(T_{n^*}) - \lambda)^{-1} - 1$ is invertible and so

$$\begin{aligned} \pi(T) - \lambda &= (\pi(T_{n^*}) - \lambda) - (\pi(T_{n^*}) - \pi(T)) \\ &= -\left[(\pi(T_{n^*}) - \pi(T))(\pi(T_{n^*}) - \lambda)^{-1} - 1\right](\pi(T_{n^*}) - \lambda) \end{aligned}$$

is invertible. Consequently, $\lambda \notin \sigma_e(T)$, i.e. $T - \lambda$ is a Fredholm operator. Finally, since $\lambda \notin \sigma_w(T_n)$ for all $n \in \mathbb{N}$, it follows by Theorem 3.1 that $i(T_n - \lambda) \rightarrow i(T - \lambda)$, and so $i(T - \lambda) = 0$, which is a contradiction. \square

Theorem 4.2. *Let $T \in \Phi(H)$ be such that satisfies Browder's theorem. If $0 \in \text{acc}\sigma_w(T)$ and (T_n) is a sequence of essentially G_1 operators such that $T_n \xrightarrow{\nu} T$, then $\sigma(T_n) \rightarrow \sigma(T)$.*

An operator $A \in B(H)$ is called p -hyponormal if $(A^*A)^p - (AA^*)^p \geq 0$. For the case $p = 1$ the operator A is called hyponormal. It is well known that the restriction of the spectrum on the class of p -hyponormal operators is continuous. See, [11] and [14]. In the following theorem we use the idea of [11] and adapt it for the case of ν -convergence. Note first that if $A \in B(H)$ is a p -hyponormal operator and $0 \in \sigma_p(A)$ then $0 \in \sigma_{jp}(A)$ and so $N(A) = N(A^*)$ which implies that $N(A)$ is invariant for both A and A^* . Therefore

$$A = \begin{bmatrix} 0 & 0 \\ 0 & B \end{bmatrix} \quad (4.2)$$

on $N(A) \oplus N(A)^\perp$, where $B = A|_{N(A)^\perp}$ and $0 \notin \sigma_p(B)$. From [24, Lemma 4], B is also p -hyponormal. We claim that $0 \notin \sigma(|B|)$, indeed if $0 \in \sigma(|B|) (= \sigma_{ap}(|B|))$ then there exists a sequence (x_m) of unit vectors such that $|B|x_m \rightarrow 0$. This implies that $Bx_m \rightarrow 0$, thus $0 \in \sigma_{ap}(B)$, but since $R(A)$ is closed i.e. $R(B)$ is closed, it follows that $0 \in \sigma_p(B)$, which is a contradiction.

Theorem 4.3. *If T_n, T are operators in $B(H)$ such that*

- (1) $T_n \xrightarrow{\nu} T$,
- (2) $T \in \Phi(H)$ and T_n is p -hyponormal for all $n \in \mathbb{N}$,
- (3) $0 \in \sigma_p(T_n)$ for all $n \in \mathbb{N}$, and the sequence $(\| |B_n|^{-1} \|)$ is bounded, where the operators $|B_n|$ are as in (4.2),

then $\sigma(T_n) \rightarrow \sigma(T)$.

Proof. First observe that $\|T_n T - T^2\| = \|(T_n - T)T\| \rightarrow 0$, thus $T_n T \xrightarrow{n} T^2$, which implies that $T_n T \in \Phi(H)$ for all n large. Thus we may suppose that $R(T_n)$ is closed for all $n \in \mathbb{N}$. We show that there exist a sequence (S_n) of hyponormal operators and a sequence (X_n) of invertible operators such that $S_n = X_n T_n X_n^{-1}$ for all $n \in \mathbb{N}$, and $(\|X_n\|), (\|X_n^{-1}\|)$ are bounded.

From condition (2), $0 \in \sigma_p(T_n)$ for all $n \in \mathbb{N}$. Then $T_n = \begin{bmatrix} 0 & 0 \\ 0 & B_n \end{bmatrix}$ on $N(T_n) \oplus N(T_n)^\perp$, $0 \notin \sigma_p(B_n)$, B_n is p -hyponormal and $0 \notin \sigma(|B_n|)$. Consider the polar decomposition $B_n = U_n|B_n|$ and define $\widehat{B}_n = |B_n|^{1/2}U_n|B_n|^{1/2}$. Observe that if $x \in N(\widehat{B}_n)$ then $|B_n|^{1/2}U_n|B_n|^{1/2}x = 0$ and so $B_n|B_n|^{-1}|B_n|^{1/2}x = 0$ which implies that $|B_n|^{-1}|B_n|^{1/2}x = 0$ because $0 \notin \sigma_p(B_n)$, hence $x = 0$. Thus $N(\widehat{B}_n) = \{0\}$ i.e. $0 \notin \sigma_p(\widehat{B}_n)$. This implies that $0 \notin \sigma(|\widehat{B}_n|)$. Let \widehat{B}_n have the polar decomposition $\widehat{B}_n = V_n|\widehat{B}_n|$, then by [2, Corollary 3], the operator \widetilde{B}_n , defined by $\widetilde{B}_n = |\widehat{B}_n|^{1/2}V_n|\widehat{B}_n|^{1/2}$, is hyponormal. Define

$$X_n = \begin{bmatrix} 1 & 0 \\ 0 & |\widehat{B}_n|^{1/2}|B_n|^{1/2} \end{bmatrix} \quad \text{and} \quad S_n = \begin{bmatrix} 0 & 0 \\ 0 & \widetilde{B}_n \end{bmatrix}.$$

Then S_n is hyponormal, X_n is invertible and $S_n = X_n T_n X_n^{-1}$. From condition (2) we have that $(\|X_n^{-1}\|)$ is bounded. Also it is clear that $(\|X_n\|)$ is bounded.

We show that $\sigma(T) \subseteq \liminf \sigma(T_n)$. Take $\lambda \in \sigma(T) \setminus \{0\}$ and suppose that $\lambda \notin \liminf \sigma(T_n)$. Then we may assume that there exists $\epsilon > 0$ such that $B(\lambda, \epsilon) \cap \sigma(T_n) = \emptyset$ for all $n \in \mathbb{N}$. This implies that $T_n - \lambda$ is invertible. In a similar way to proof of Theorem 4.1, we have that

$$\begin{aligned} \left\| \left[(T_n - T)(T_n - \lambda)^{-1} \right]^2 \right\| &\leq \frac{1}{|\lambda|} \left[\|(T_n - T)T_n\| \|(T_n - \lambda)^{-1}\| \|T_n - T\| \right. \\ &\quad \left. + \|(T_n - T)T\| + \|(T_n - T)T_n\| \right] \|(T_n - \lambda)^{-1}\|. \end{aligned} \quad (4.3)$$

Since T_n and S_n are similar, it follows that $\sigma(T_n) = \sigma(S_n)$. Therefore, $d(\lambda, \sigma(S_n)) \geq \epsilon$ and $S_n - \lambda$ is invertible. Note that $(T_n - \lambda)^{-1} = X_n^{-1}(S_n - \lambda)^{-1}X_n$. Moreover, since S_n is hyponormal it follows that $\|(S_n - \lambda)^{-1}\| = \frac{1}{d(\lambda, \sigma(S_n))} \leq \frac{1}{\epsilon}$. Thus the right term of (4.3) is bounded by

$$\frac{1}{|\lambda|} \left[\|(T_n - T)T_n\| \frac{M_1 M_2}{\epsilon} \|T_n - T\| + \|(T_n - T)T\| + \|(T_n - T)T_n\| \right] \frac{M_1 M_2}{\epsilon}, \quad (4.4)$$

where M_1, M_2 are constants such that $\|X_n\| \leq M_1$ and $\|X_n^{-1}\| \leq M_2$ for all $n \in \mathbb{N}$. Since $T_n \xrightarrow{\nu} T$ it follows that the expression in (4.4) tends to zero. Proceeding similarly to the final part of the proof of Theorem 4.1, we obtain that $T - \lambda$ is invertible, which is a contradiction. \square

Remark 4.4. The conclusion of Theorem 4.3 holds if we replace the hypothesis by the following conditions:

- (1) $T_n \xrightarrow{\nu} T$, $T^*(T_n - T) \xrightarrow{n} 0$ and $T_n^*(T_n - T) \xrightarrow{n} 0$;
- (2) $T \in \Phi(H)$ and $\{0\} \neq N(T) \subseteq N(T_n)$ for all $n \in \mathbb{N}$;
- (3) T, T_n are p -hyponormal operators.

Indeed, from condition (1), $|T_n|^2 - |T|^2 = T_n^* T_n - T^* T = T_n^*(T_n - T) + [T^*(T_n - T)]^* \xrightarrow{n} 0$, thus $|T_n|^{1/2} \xrightarrow{n} |T|^{1/2}$. Since $0 \in \sigma_p(T)$, it follows that

$T = 0 \oplus B$ on $N(T) \oplus N(T)^\perp$ with $0 \notin \sigma(|B|)$. Then there exists $\alpha > 0$ such that $\alpha\|y\| \leq \| |B|^{1/2}y \|$ for all $y \in N(T)^\perp$. This implies by condition (2) that for $0 < \epsilon < \alpha$, there exists $N \in \mathbb{N}$ such that for every $n \geq N$, $(\alpha - \epsilon)\|y\| \leq \| |B_n|^{1/2}y \|$ for all $y \in N(T_n)^\perp$. Consequently, $\|(|B_n|^{1/2})^{-1}\| \leq \frac{1}{\alpha - \epsilon}$ for all $n \geq N$.

Berberian shows that for every Hilbert space H , there exists a Hilbert space $K \supset H$ and a faithful $*$ -representation $T \rightarrow T^\circ$ from $B(H)$ to $B(K)$: $(S+T)^\circ = S^\circ + T^\circ$, $(\lambda T)^\circ = \lambda T^\circ$, $(ST)^\circ = S^\circ T^\circ$, $(T^*)^\circ = (T^\circ)^*$, $(I_H)^\circ = I_K$ and $\|T^\circ\| = \|T\|$ such that

- (1) $T \geq 0$ if and only if $T^\circ \geq 0$,
- (2) $\sigma_p(T^\circ) = \sigma_a(T^\circ) = \sigma_a(T)$.

Remark 4.5. Observe that in the previous theorem, $\sigma_p(B_n) = \sigma_{ap}(B_n)$ due to $R(T_n)$ is closed. In [11] the authors use the Berberian extension T_n° of a p -hyponormal operator T_n and state that if $0 \in \sigma_p(T_n^\circ)$, then

$$\sigma_p(B_n) = \sigma_{ap}(B_n), \quad (4.5)$$

where $T_n^\circ = 0 \oplus B_n$ on $N(T_n) \oplus N(T_n)^\perp$ and $0 \notin \sigma_p(B_n)$, without the need for $R(T_n)$ to be closed. This fact was also established in [13], page 586, line 20. The authors claim that

$$\sigma_{ap}(B_\lambda) = \sigma_p(B_\lambda) \quad (4.6)$$

for all non-zero $\lambda \in \sigma_p(A^\circ)$, where $A \in \mathcal{C}(i)$, and this collection is defined as the set of all operators $T \in B(H_i)$ for which $\sigma(T) = \{0\}$ implies T is nilpotent and T° satisfies the property:

$$T^\circ = \begin{bmatrix} \lambda & X_\lambda \\ 0 & B_\lambda \end{bmatrix} \quad \text{on } N(T^\circ - \lambda) \oplus N(T^\circ - \lambda)^\perp$$

at every non-zero $\lambda \in \sigma_p(T^\circ)$ for some operators X_λ and B_λ such that $\lambda \notin \sigma_p(B_\lambda)$ and $\sigma(T^\circ) = \{\lambda\} \cup \sigma(B_\lambda)$. However, equalities (4.5) and (4.6) are not necessary hold. We prove only that (4.6) is false. It is clear that $\sigma_a(B_\lambda) \setminus \{\lambda\} = \sigma_p(B_\lambda)$ and $\alpha(B_\lambda - \lambda) = 0$, but $R(B_\lambda - \lambda)$ is not necessarily closed. Indeed, consider a normal operator $A \in B(H_i)$ such that $\sigma(A) = [0, 1]$ (for example, the multiplication operator $A : L^2([0, 1]) \rightarrow L^2([0, 1])$ defined by $A(f)(x) = xf(x)$). Then $A \in \mathcal{C}(i)$. We show that for every $\lambda \in \sigma_p(A^\circ)$, $R(B_\lambda - \lambda)$ is not closed. By contradiction, suppose that there exists $\lambda \in \sigma_p(A^\circ)$ such that $R(B_\lambda - \lambda)$ is closed. Then $B_\lambda - \lambda$ is a semi-Fredholm operator such that $\alpha(B_\lambda - \lambda) = 0$. By [6, Theorem 4.2.1], there exists $\epsilon > 0$ such that if $|\gamma - \lambda| < \epsilon$ then $B_\lambda - \gamma \in \Phi_+(N(A^\circ - \lambda)^\perp)$ and $\alpha(B_\lambda - \gamma) = 0$. This implies that $R(A^\circ - \gamma) = (\lambda - \gamma)N(A^\circ - \lambda) \oplus R(B_\lambda - \gamma)$ is closed and $\alpha(A^\circ - \gamma) = \alpha((\lambda - \gamma)I) + \alpha(B_\lambda - \gamma) = 0$ for all $\gamma \in B(\lambda, \epsilon)$ with $\gamma \neq \lambda$. Therefore, $\lambda \in \text{iso } \sigma_a(A^\circ)$. On the other hand, since A is a normal operator it follows that $\sigma(A) = \sigma_a(A) = \sigma_a(A^\circ)$. Thus, $\lambda \in \text{iso } \sigma(A)$, which is a contradiction, because $\sigma(A) = [0, 1]$. Consequently, the equality (4.6) is not true. This affects the proof of the main result in the paper [13].

REFERENCES

- [1] M. Ahues, A. Largillier, and B. V. Limaye. *Spectral computations for bounded operators*, volume 18 of *Applied Mathematics (Boca Raton)*. Chapman & Hall/CRC, Boca Raton, FL, 2001.
- [2] A. Aluthge. On p -hyponormal operators for $0 < p < 1$. *Integral Equations Operator Theory*, 13(3):307–315, 1990.
- [3] A. Ammar. Some properties of the Wolf and Weyl essential spectra of a sequence of linear operators ν -convergent. *Indag. Math. (N.S.)*, 28(2):424–435, 2017.
- [4] A. Ammar, A. Bouchekoua, and A. Jeribi. Some approximation results in a non-Archimedean Banach space. *Funct. Anal. Approx. Comput.*, 12(1):33–50, 2020.
- [5] L. Burlando. Spectral continuity in some Banach algebras. *Rocky Mountain J. Math.*, 23(1):17–39, 1993.
- [6] S. R. Caradus, W. E. Pfaffenberger, and B. Yood. *Calkin algebras and algebras of operators on Banach spaces*. Marcel Dekker, Inc., New York, 1974. Lecture Notes in Pure and Applied Mathematics, Vol. 9.
- [7] J. B. Conway and B. B. Morrel. Operators that are points of spectral continuity. *Integral Equations Operator Theory*, 2(2):174–198, 1979.
- [8] J. B. Conway and B. B. Morrel. Operators that are points of spectral continuity. II. *Integral Equations Operator Theory*, 4(4):459–503, 1981.
- [9] J. B. Conway and B. B. Morrel. Operators that are points of spectral continuity. III. *Integral Equations Operator Theory*, 6(3):319–344, 1983.
- [10] S. Djordjević. The continuity of the essential approximative point spectrum. *Facta Univ. Ser. Math. Inform.*, (10):97–104, 1995.
- [11] S. V. Djordjević and B. P. Duggal. Weyl’s theorems and continuity of spectra in the class of p -hyponormal operators. *Studia Math.*, 143(1):23–32, 2000.
- [12] S. V. Djordjević and Y. M. Han. Browder’s theorems and spectral continuity. *Glasg. Math. J.*, 42(3):479–486, 2000.
- [13] B. P. Duggal, I. H. Jeon, and I. H. Kim. Continuity of the spectrum on a class of upper triangular operator matrices. *J. Math. Anal. Appl.*, 370(2):584–587, 2010.
- [14] I. S. Hwang and W. Y. Lee. The spectrum is continuous on the set of p -hyponormal operators. *Math. Z.*, 235(1):151–157, 2000.
- [15] G. R. Luecke. A note on spectral continuity and on spectral properties of essentially G_1 operators. *Pacific J. Math.*, 69(1):141–149, 1977.
- [16] G. J. Murphy. *C^* -algebras and operator theory*. Academic Press, Inc., Boston, MA, 1990.
- [17] J. D. Newburgh. The variation of spectra. *Duke Math. J.*, 18:165–176, 1951.
- [18] S. Sánchez-Perales and S. V. Djordjević. Continuity of spectra and compact perturbations. *Bull. Korean Math. Soc.*, 48(6):1261–1270, 2011.
- [19] S. Sánchez-Perales and S. V. Djordjević. Spectral continuity using ν -convergence. *J. Math. Anal. Appl.*, 433(1):405–415, 2016.
- [20] S. Sánchez-Perales and S. V. Djordjević. Spectral continuity relative to invariant subspaces. *Complex Anal. Oper. Theory*, 11(4):927–941, 2017.
- [21] S. Sánchez-Perales, S. V. Djordjević, and S. Palafox. Some results about spectral continuity and compact perturbations. *Filomat*, 2020. Submitted.
- [22] S. Sánchez-Perales, S. Palafox, and S. V. Djordjević. α -Fredholm operators relative to invariant subspaces. *Oper. Matrices*, 13(4):921–936, 2019.
- [23] M. Schechter. *Principles of functional analysis*, volume 36 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, second edition, 2002.
- [24] A. Uchiyama. Berger-Shaw’s theorem for p -hyponormal operators. *Integral Equations Operator Theory*, 33(2):221–230, 1999.

UNIVERSIDAD TECNOLÓGICA DE LA MIXTECA, INSTITUTO DE FÍSICA Y MATEMÁTICAS,
KM. 2.5 CARRETERA A ACATLIMA, 69000 OAXACA, MEXICO.

E-mail address: `es21254@yahoo.com.mx`

UNIVERSIDAD TECNOLÓGICA DE LA MIXTECA, INSTITUTO DE FÍSICA Y MATEMÁTICAS,
KM. 2.5 CARRETERA A ACATLIMA, 69000 OAXACA, MEXICO.

E-mail address: `sergiopalafox@gmail.com`

UNIVERSIDAD TECNOLÓGICA DE LA MIXTECA, DIVISIÓN DE ESTUDIOS DE POSTGRADO,
KM. 2.5 CARRETERA A ACATLIMA, 69000 OAXACA, MEXICO

E-mail address: `tombp55@hotmail.com`