

Note on Sunflowers

Tolson Bell*, Suchakree Chueluecha[†] and Lutz Warnke[‡]

September 17, 2020; revised March 17, 2021

Abstract

A sunflower with p petals consists of p sets whose pairwise intersections are identical. The goal of the sunflower problem is to find the smallest $r = r(p, k)$ such that any family of r^k distinct k -element sets contains a sunflower with p petals. Building upon a breakthrough of Alweiss, Lovett, Wu and Zhang from 2019, Rao proved that $r = O(p \log(pk))$ suffices; this bound was reproved by Tao in 2020. In this short note we record that $r = O(p \log k)$ suffices, by using a minor variant of the probabilistic part of these recent proofs.

1 Introduction

A sunflower with p petals is a family of p sets whose pairwise intersections are identical (the intersections may be empty). Let $\text{Sun}(p, k)$ denote the smallest natural number s with the property that any family of at least s distinct k -element sets contains a sunflower with p petals. In 1960, Erdős and Rado [4] proved that $(p-1)^k < \text{Sun}(p, k) \leq (p-1)^k k! + 1 = O((pk)^k)$, and conjectured that for any $p \geq 2$ there is a constant $C_p > 0$ such that $\text{Sun}(p, k) \leq C_p^k$ for all $k \geq 2$. This famous conjecture in extremal combinatorics was one of Erdős' favorite problems [2], for which he offered a \$1000 reward [3]; it remains open despite considerable attention [7].

In 2019, there was a breakthrough on the sunflower conjecture: using iterative encoding arguments, Alweiss, Lovett, Wu and Zhang [1] proved that $\text{Sun}(p, k) \leq (Cp^3 \log k \log \log k)^k$ for some constant $C > 0$, opening the floodgates for further improvements. Using Shannon's noiseless coding theorem, Rao [8] subsequently simplified their proof and obtained a slightly better bound. Soon thereafter, Frankston, Kahn, Narayanan and Park [5] refined some key counting arguments from [1]. Their ideas were then utilized by Rao [9] to improve the best-known sunflower bound to $\text{Sun}(p, k) \leq (Cp \log(pk))^k$ for some constant $C > 0$, which in 2020 was reproved by Tao in his blog [10] using Shannon entropy arguments.

The aim of this short note is to record, for the convenience of other researchers, that a minor variant of (the probabilistic part of) the arguments from [9, 10] gives $\text{Sun}(p, k) \leq (Cp \log k)^k$ for some constant $C > 0$.

Theorem 1. *There is a constant $C \geq 4$ such that $\text{Sun}(p, k) \leq (Cp \log k)^k$ for all integers $p, k \geq 2$.*

Setting $r(p, k) = Cp \log k + \mathbb{1}_{\{k=1\}}p$, we shall in fact prove $\text{Sun}(p, k) \leq r(p, k)^k$ for all integers $p \geq 2$ and $k \geq 1$. Similarly to the strategy of [1, 9, 10], this upper bound follows easily by induction on $k \geq 1$ from Lemma 2 below, where a family \mathcal{S} of k -element sets is called r -spread if there are at most $r^{k-|T|}$ sets of \mathcal{S} that contain any non-empty set T . (Indeed, the base case $k = 1$ is trivial due to $r(p, 1) = p$, and the induction step $k \geq 2$ uses a simple case distinction: if \mathcal{S} is $r(p, k)$ -spread, then Lemma 2 guarantees a sunflower with p petals; otherwise there is a non-empty set T such that more than $r(p, k)^{k-|T|} \geq r(p, k-|T|)^{k-|T|}$ sets of \mathcal{S} contain T , and among this family of sets we easily find a sunflower with p petals using induction.)

Lemma 2. *There is a constant $C \geq 4$ such that, setting $r(p, k) = Cp \log k$, the following holds for all integers $p, k \geq 2$. If a family \mathcal{S} with $|\mathcal{S}| \geq r(p, k)^k$ sets of size k is $r(p, k)$ -spread, then \mathcal{S} contains p disjoint sets.*

*School of Mathematics, Georgia Institute of Technology, Atlanta, GA 30332, USA. E-mail: tbell137@gatech.edu. Research supported by NSF Grant DMS-1851843.

[†]Department of Mathematics, Lehigh University, Bethlehem, PA 18015, USA. E-mail: suc221@lehigh.edu. Research supported by Georgia Institute of Technology, College of Sciences.

[‡]School of Mathematics, Georgia Institute of Technology, Atlanta, GA 30332, USA. E-mail: warnke@math.gatech.edu. Research supported by NSF Grant DMS-1703516, NSF CAREER grant DMS-1945481 and a Sloan Research Fellowship.

Inspired by [1], in [9, 10] probabilistic arguments are used to deduce Lemma 2 with $r(p, k) = \Theta(p \log(pk))$ from Theorem 3 below, where X_δ denotes the random subset of X in which each element is included independently with probability δ .

Theorem 3 (Main technical estimate of [9, 10]). *There is a constant $B \geq 1$ such that the following holds for any integer $k \geq 2$, any reals $0 < \delta, \epsilon \leq 1/2$, $r \geq B\delta^{-1} \log(k/\epsilon)$, and any family \mathcal{S} of k -element subsets of a finite set X . If \mathcal{S} is r -spread with $|\mathcal{S}| \geq r^k$, then $\mathbb{P}(\exists S \in \mathcal{S} : S \subseteq X_\delta) > 1 - \epsilon$.* \square

The core idea of [1, 9, 10] is to randomly partition the set X into $V_1 \cup \dots \cup V_p$, by independently placing each element $x \in X$ into a randomly chosen V_i . Note that the marginal distribution of each V_i equals the distribution of X_δ with $\delta = 1/p$. Invoking Theorem 3 with $\epsilon = 1/p$ and $r = B\delta^{-1} \log(k/\epsilon)$, a standard union bound argument implies that, with non-zero probability, all of the random partition-classes V_i contain a set from \mathcal{S} . Hence p disjoint sets $S_1, \dots, S_p \in \mathcal{S}$ must exist, which proves Lemma 2 with $r(p, k) = Bp \log(pk)$.

We prove Lemma 2 with $r(p, k) = \Theta(p \log k)$ using a minor twist: by randomly partitioning the vertex-set into more than p classes V_i , and then using linearity of expectation (instead of a union bound).

Proof of Lemma 2. Set $C = 4B$. We randomly partition the set X into $V_1 \cup \dots \cup V_{2p}$, by independently placing each element $x \in X$ into a randomly chosen V_i . Let I_i be the indicator random variable for the event that V_i contains a set from \mathcal{S} . Since V_i has the same distribution as X_δ with $\delta = 1/(2p)$, by invoking Theorem 3 with $\epsilon = 1/2$ and $r = r(p, k) = 2Bp \log(k^2) \geq B\delta^{-1} \log(k/\epsilon)$, we obtain $\mathbb{E} I_i > 1/2$. Using linearity of expectation, the expected number of partition-classes V_i with $I_i = 1$ is thus at least p . Hence there must be a partition where at least p of the V_i contain a set from \mathcal{S} , which gives the desired p disjoint sets $S_1, \dots, S_p \in \mathcal{S}$. \square

Generalizing this idea, Theorem 3 gives $p > \lfloor 1/\delta \rfloor (1 - \epsilon)$ disjoint sets $S_1, \dots, S_p \in \mathcal{S}$, which in the special case $\lfloor 1/\delta \rfloor \epsilon \leq 1$ (used in [1, 9, 10] with $\delta = \epsilon = 1/p$) simplifies to $p \geq \lfloor 1/\delta \rfloor$.

2 Remarks

Our proof of Lemma 2 only invokes Theorem 3 with $\epsilon = 1/2$, i.e., it does not exploit the fact that Theorem 3 has an essentially optimal dependence on ϵ (see Lemma 4 below). In particular, this implies that we could alternatively also prove Lemma 2 and thus the $\text{Sun}(p, k) \leq (Cp \log k)^k$ bound of Theorem 1 using the combinatorial arguments of Frankston, Kahn, Narayanan and Park [5] (we have verified that the proof of [5, Theorem 1.7] can be extended to yield Theorem 3 under the stronger assumption $r \geq B\delta^{-1} \max\{\log k, \log^2(1/\epsilon)\}$, say).

We close by recording that Theorem 3 is essentially best possible with respect to the r -spread assumption, which follows from the construction in [1, Section 2] that in turn builds upon [4, Theorem II].

Lemma 4. *For any reals $0 < \delta, \epsilon \leq 1/2$ and any integers $k \geq 1$, $1 \leq r \leq 0.25\delta^{-1} \log(k/\epsilon)$, there exists an r -spread family \mathcal{S} of k -element subsets of $X = \{1, \dots, rk\}$ with $|\mathcal{S}| = r^k$ and $\mathbb{P}(\exists S \in \mathcal{S} : S \subseteq X_\delta) < 1 - \epsilon$.*

Proof. We fix a partition $V_1 \cup \dots \cup V_k$ of X into sets of equal size $|V_i| = r$, and define \mathcal{S} as the family of all k -element sets containing exactly one element from each V_i . It is easy to check that \mathcal{S} is r -spread, with $|\mathcal{S}| = r^k$. Focusing on the necessary event that X_δ contains at least one element from each V_i , we obtain

$$\mathbb{P}(\exists S \in \mathcal{S} : S \subseteq X_\delta) \leq (1 - (1 - \delta)^r)^k \leq e^{-(1-\delta)r^k} < e^{-e^{-2\delta r^k}} \leq e^{-\sqrt{\epsilon k}} < 1 - \epsilon$$

by elementary considerations (since $e^{-\sqrt{\epsilon}} < 1 - \epsilon$ due to $0 < \epsilon \leq 1/2$). \square

3 Acknowledgements

This research was conducted as part of the 2020 REU program at Georgia Institute of Technology.

References

- [1] R. Alweiss, S. Lovett, K. Wu, and J. Zhang. Improved bounds for the Sunflower lemma. In *Proceedings of the 52nd Annual ACM SIGACT Symposium on Theory of Computing (STOC 2020)*, pp. 624–630, ACM, New York (2020). Extended preprint available at [arXiv:1908.08483v2](https://arxiv.org/abs/1908.08483v2)
- [2] P. Erdős. On the combinatorial problems which I would most like to see solved. *Combinatorica* **1** (1981), 25–42.
- [3] P. Erdős. Some of my old and new combinatorial problems. In *Paths, Flows, and VLSI-layout (Bonn 1988)*, Algorithms & Combinatorics 9, pp. 35–45, Springer, Berlin (1990).
- [4] P. Erdős and R. Rado. Intersection theorems for systems of sets. *Journal of the London Mathematical Society* **35** (1960), 85–90.
- [5] K. Frankston, J. Kahn, B. Narayanan, and J. Park. Thresholds versus fractional Expectation-thresholds. Preprint (2019). [arXiv:1910.13433v2](https://arxiv.org/abs/1910.13433v2)
- [6] S. Janson, T. Luczak, and A. Ruciński. *Random Graphs*. Wiley-Interscience (2000).
- [7] A. Kostochka. Extremal problems on Δ -systems. In *Numbers, Information and Complexity (Bielefeld 1998)*, pp. 143–150, Kluwer Academic Publishers, Boston (2000).
- [8] A. Rao. Coding for Sunflowers. Preprint (2019). [arXiv:1909.04774v1](https://arxiv.org/abs/1909.04774v1)
- [9] A. Rao. Coding for Sunflowers. *Discrete Analysis* **2** (2020), 8 pp. [arXiv:1909.04774v2](https://arxiv.org/abs/1909.04774v2)
- [10] T. Tao. The Sunflower lemma via Shannon entropy. Blog post (2020). Available at <https://terrytao.wordpress.com/2020/07/20/the-sunflower-lemma-via-shannon-entropy/>

Appendix: Theorem 3

Theorem 3 follows from Tao’s proof of Proposition 5 in [10] (noting that any r -spread family \mathcal{S} with $|\mathcal{S}| \geq r^k$ sets of size k is also r -spread in the sense of [10]). We now record that Theorem 3 also follows from Rao’s proof of Lemma 4 in [9] (where the random subset of X is formally chosen in a slightly different way).

Proof of Theorem 3 based on [9]. Set $\gamma = \delta/2$ and $m = \lceil \gamma|X| \rceil$. Let X_i denote a set chosen uniformly at random from all i -element subsets of X . Since X_δ conditioned on containing exactly i elements has the same distribution as X_i , by the law of total probability and monotonicity it routinely follows that $\mathbb{P}(\exists S \in \mathcal{S} : S \subseteq X_\delta)$ is at least $\mathbb{P}(\exists S \in \mathcal{S} : S \subseteq X_m) \cdot \mathbb{P}(|X_\delta| \geq m)$. The proof of Lemma 4 in [9] shows that $\mathbb{P}(\exists S \in \mathcal{S} : S \subseteq X_m) > 1 - \epsilon^2$ whenever $r \geq \alpha \gamma^{-1} \log(k/\epsilon)$, where $\alpha > 0$ is a sufficiently large constant. Noting $|\mathcal{S}| \leq |X|^k$ we see that $|\mathcal{S}| \geq r^k$ enforces $|X| \geq r$, so standard Chernoff bounds (such as [6, Theorem 2.1]) imply that $\mathbb{P}(|X_\delta| < m) \leq \mathbb{P}(|X_\delta| \leq |X|\delta/2)$ is at most $e^{-|X|\delta/8} \leq e^{-r\delta/8} \leq \epsilon^2$ whenever $r \geq 16\delta^{-1} \log(1/\epsilon)$. This completes the proof with $B = \max\{2\alpha, 16\}$, say (since $(1 - \epsilon^2)^2 \geq 1 - \epsilon$ due to $0 < \epsilon \leq 1/2$). \square