

# THE BINARY GOLD FUNCTION AND ITS $c$ -BOOMERANG CONNECTIVITY TABLE

SARTAJ UL HASAN, MOHIT PAL, AND PANTELIMON STĂNICĂ

**ABSTRACT.** Here, we give a complete description of the  $c$ -Boomerang Connectivity Table for the Gold function over finite fields of even characteristic, by using double Weil sums. In the process we generalize a result of Boura and Canteaut (IACR Trans. Symmetric Cryptol. 2018(3) : 290-310, 2018) for the classical boomerang uniformity.

## 1. INTRODUCTION

Let  $\mathbb{F}_q$  be the finite field with  $q = p^n$  elements, where  $p$  is a prime and  $n$  is a positive integer. The multiplicative cyclic group of nonzero elements of the finite field is denoted by  $\mathbb{F}_q^* = \langle g \rangle$ , where  $g$  is a primitive element of  $\mathbb{F}_q$ . A Weil sum is an important character sum defined as follows

$$\sum_{x \in \mathbb{F}_q} \chi(F(x)),$$

where  $\chi$  is an additive character of  $\mathbb{F}_q$  and  $F(x)$  is a polynomial in  $\mathbb{F}_q[x]$ . It is well-known that a polynomial  $F(x)$  over finite field  $\mathbb{F}_q$  is a permutation polynomial (PP) if and only if its Weil sum  $\sum_{x \in \mathbb{F}_q} \chi(F(x)) = 0$  for all nontrivial additive characters  $\chi$  of  $\mathbb{F}_q$ . Permutation polynomials are a very important class of polynomials as they have applications in coding theory and cryptography, especially in the substitution boxes (S-boxes) of the block ciphers. The security of the S-boxes relies on certain properties of the function  $F(x)$ , e.g., its differential uniformity, boomerang uniformity, nonlinearity etc.

Recently, Cid et al. [4] introduced a “new tool” for analyzing the boomerang style attack proposed by Wagner [17]. This new tool is usually referred to as Boomerang Connectivity Table (BCT). Boura and Canteaut [2] further studied BCT and coined the term boomerang uniformity, which is essentially the maximum value in the BCT. Li et al. [9] provided new insights in the study of BCT and presented an equivalent technique to compute BCT, which does not require the compositional inverse of the permutation polynomial  $F(x)$  at all. In fact, Li et al. [9] also gave a characterization of BCT in terms of Walsh transform and gave a class of permutation polynomial with boomerang uniformity 4.

Recently, Stănică [12] extended the notion of BCT and boomerang uniformity. In fact, he defined what he termed as  $c$ -BCT and  $c$ -boomerang uniformity for an arbitrary polynomial function  $F$  over  $\mathbb{F}_q$  and for any  $c \neq 0 \in \mathbb{F}_q$ . Let  $a, b \in \mathbb{F}_q$ , then the entry of the  $c$ -Boomerang Connectivity Table ( $c$ -BCT) at  $(a, b) \in \mathbb{F}_{p^n} \times \mathbb{F}_{p^n}$ , denoted as  ${}_c\mathcal{B}_F(a, b)$ , is the number of solutions in  $\mathbb{F}_{p^n} \times \mathbb{F}_{p^n}$  of the following system

$$(1.1) \quad \begin{cases} F(x) - cF(y) = b \\ F(x + a) - c^{-1}F(y + a) = b. \end{cases}$$

---

2010 *Mathematics Subject Classification.* 12E20, 11T24, 11T06, 94A60.

*Key words and phrases.* Finite fields, double Weil sums, boomerang uniformity,  $c$ -boomerang uniformity.

The  $c$ -boomerang uniformity of  $F$  is defined as

$$\beta_{F,c} = \max_{a,b \in \mathbb{F}_{p^n}^*} {}_c\mathcal{B}_F(a,b).$$

In yet other recent papers, Stănică [13, 14] further studied the  $c$ -BCT for the swapped inverse function and also gave an elegant description of the  $c$ -BCT entries of the power map in terms of double Weil sums. He further simplified his expressions for the Gold function  $x^{p^k+1}$  over  $\mathbb{F}_{p^n}$ , for all  $1 \leq k < n$  and  $p$  odd. In this paper, we shall complement the work of [14] to the finite fields of even characteristic ( $p = 2$ ).

The paper is structured as follows. Section 2 contains some preliminary results that will be used across the sections. Section 3 contains the characterization of  $c$ -BCT entries in terms of double Weil sums. For  $c = 1$ , we further simplify this expression in Section 4. In fact, Theorem 4.1 generalizes previously known results of Boura and Canteaut [2]. In Section 5, we consider the case when  $c \in \mathbb{F}_{2^e} \setminus \{0, 1\}$ , where  $e = \gcd(k, n)$ . In Section 6, we discuss the general case. Finally, in Section 7, we discuss the affine, extended affine and CCZ-equivalence as it relates to  $c$ -boomerang uniformity.

## 2. PRELIMINARIES

We begin this section by first recalling the recent notion of  $c$ -differentials introduced in [7]. We shall assume that  $q = 2^n$  for rest of the paper. For an  $(n, n)$ -function  $F : \mathbb{F}_q \rightarrow \mathbb{F}_q$ , and  $c \in \mathbb{F}_q$ , we define the (*multiplicative*)  $c$ -derivative of  $F$  with respect to  $a \in \mathbb{F}_q$  to be the function

$${}_cD_a F(x) = F(x+a) - cF(x), \text{ for all } x \in \mathbb{F}_q.$$

Further, for  $a, b \in \mathbb{F}_q$ , we let the entries of the  $c$ -Difference Distribution Table ( $c$ -DDT) be defined by  ${}_c\Delta_F(a, b) = \#\{x \in \mathbb{F}_q : F(x+a) - cF(x) = b\}$ . We call the quantity

$$\delta_{F,c} = \max \{ {}_c\Delta_F(a, b) \mid a, b \in \mathbb{F}_q, \text{ and } a \neq 0 \text{ if } c = 1 \},$$

the  $c$ -differential uniformity of  $F$ . Note that the case  $c = 1$  corresponds to the usual notion of differential uniformity. The interested reader may refer to [1, 8, 10, 15, 16, 18] for some recent results concerning  $c$ -differential uniformity.

The following theorem is a “binary” analogue of [14, Theorem 1], which gives a nice connection between  $c$ -BCT and  $c$ -DDT entries of the power map  $x^d$  over  $\mathbb{F}_{2^n}$ .

**Theorem 2.1.** *Let  $F(x) = x^d$  be a power function on  $\mathbb{F}_q$ ,  $q = 2^n$  and  $c \in \mathbb{F}_q^*$ . Then, for fixed  $b \in \mathbb{F}_q^*$ , the  $c$ -Boomerang Connectivity Table entry  ${}_c\mathcal{B}_F(1, b)$  at  $(1, b)$  is given by*

$$\frac{1}{q} \left( \sum_{w \in \mathbb{F}_q} ({}_c\Delta_F(w, b) + {}_{c^{-1}}\Delta_F(w, b)) \right) - 1 + \frac{1}{q^2} \sum_{\alpha, \beta \in \mathbb{F}_q, \alpha\beta \neq 0} \chi_1(b(\alpha + \beta)) S_{\alpha, \beta} S_{\alpha c, \beta c^{-1}},$$

with

$$\begin{aligned} S_{\alpha, \beta} &= \sum_{x \in \mathbb{F}_q} \chi_1(\alpha x^d) \chi_1(\beta(x+1)^d) \\ &= \frac{1}{(q-1)^2} \sum_{j,k=0}^{q-2} G(\bar{\psi}_j, \chi_1) G(\bar{\psi}_k, \chi_1) \sum_{x \in \mathbb{F}_q} \psi_1((\alpha x^d)^j (\beta(x+1)^d)^k), \end{aligned}$$

where  $\chi_1$  is the canonical additive character of the additive group of  $\mathbb{F}_q$ ,  $\psi_k$  is the  $k$ -th multiplicative character of the multiplicative group of  $\mathbb{F}_q$  and  $G(\psi, \chi)$  is the Gauss sum.

We shall now state some lemmas that will be used in the sequel. The following lemma is well-known and has been used in various contexts.

**Lemma 2.2.** *Let  $e = \gcd(k, n)$ . Then*

$$\gcd(2^k + 1, 2^n - 1) = \begin{cases} 1 & \text{if } n/e \text{ is odd,} \\ 2^e + 1 & \text{if } n/e \text{ is even.} \end{cases}$$

We shall also use the following lemma, which appeared in [5], describing the number of roots in  $\mathbb{F}_{2^n}$  of a linearized polynomial  $u^{2^k} x^{2^{2k}} + ux$ , where  $u \in \mathbb{F}_{2^n}^*$ .

**Lemma 2.3.** [5, Theorem 3.1] *Let  $g$  be a primitive element of  $\mathbb{F}_{2^n}$  and let  $e = \gcd(n, k)$ . For any  $u \in \mathbb{F}_{2^n}^*$  consider the linearized polynomial  $L_u(x) = u^{2^k} x^{2^{2k}} + ux$  over  $\mathbb{F}_{2^n}$ . Then for the equation  $L_u(x) = 0$ , the following are true:*

- (1) *If  $n/e$  is odd, then there are  $2^e$  solutions to this equation for any choice of  $u \in \mathbb{F}_{2^n}^*$ ;*
- (2) *If  $n/e$  is even and  $u = g^{t(2^e+1)}$  for some  $t$ , then there are  $2^{2e}$  solutions to the equation;*
- (3) *If  $n/e$  is even and  $u \neq g^{t(2^e+1)}$  for any  $t$ , then  $x = 0$  is the only solution.*

The explicit expression for the Weil sum of the form  $\sum_{x \in \mathbb{F}_{2^n}} \chi_1(ux^{2^k+1} + vx)$ , where  $u, v \in \mathbb{F}_{2^n}$ , is obtained in [5]. In what follows, we shall denote the Weil sum  $\sum_{x \in \mathbb{F}_q} \chi(ux^{2^k+1} + vx)$  by  $\mathfrak{S}(u, v)$ . The following lemma gives the explicit expression for  $\mathfrak{S}(u, 0)$ .

**Lemma 2.4.** [5] *Let  $\chi$  be any nontrivial additive character of  $\mathbb{F}_q$  and  $g$  be the primitive element of the cyclic group  $\mathbb{F}_q^*$ . The following hold:*

- (1) *If  $n/e$  is odd, then*

$$\sum_{x \in \mathbb{F}_q} \chi(ux^{2^k+1}) = \begin{cases} q & \text{if } u = 0, \\ 0 & \text{otherwise.} \end{cases}$$

- (2) *Let  $n/e$  be even so that  $n = 2m$  for some integer  $m$ . Then*

$$\sum_{x \in \mathbb{F}_q} \chi(ux^{2^k+1}) = \begin{cases} (-1)^{m/e} 2^m & \text{if } u \neq g^{t(2^e+1)} \text{ for any integer } t, \\ (-1)^{\frac{m}{e}+1} 2^{m+e} & \text{if } u = g^{t(2^e+1)} \text{ for some integer } t. \end{cases}$$

From Lemma 2.2, it is easy to see that when  $n/e$  is odd, the power map  $x^{2^k+1}$  permutes  $\mathbb{F}_{2^n}$ . Therefore if  $u \neq 0$ , there exists a unique element  $\gamma \in \mathbb{F}_q^*$  such that  $\gamma^{2^k+1} = u$  and hence

$$\begin{aligned} \mathfrak{S}(u, v) &= \sum_{x \in \mathbb{F}_q} \chi(ux^{2^k+1} + vx) \\ &= \sum_{x \in \mathbb{F}_q} \chi(x^{2^k+1} + v\gamma^{-1}x) \\ &= \mathfrak{S}(1, v\gamma^{-1}). \end{aligned}$$

The following lemma gives the expression for the Weil sum  $\mathfrak{S}(1, v)$  for  $v \neq 0$  and  $n/e$  odd.

**Lemma 2.5.** [5, Theorem 4.2] *Let  $v \neq 0$  and  $n/e$  is odd. Then*

$$\mathfrak{S}(1, v) = \begin{cases} 0 & \text{if } \text{Tr}_e(v) \neq 1, \\ \left(\frac{2}{n/e}\right)^e 2^{\frac{n+e}{2}} & \text{if } \text{Tr}_e(v) = 1, \end{cases}$$

where  $\left(\frac{2}{n/e}\right)$  is the Jacobi symbol.

In the case when  $u, v \neq 0$  and  $n/e$  is even, the Weil sum  $\mathfrak{S}(u, v)$  depends whether or not the linearized polynomial  $L_u(x) = u^{2^k} x^{2^{2k}} + ux$  is a permutation of  $\mathbb{F}_{2^n}$ . The following lemma gives the expression for Weil sum  $\mathfrak{S}(u, v)$  for  $u, v \neq 0$  and  $n/e$  even.

**Lemma 2.6.** [5, Theorem 5.3] *Let  $u, v \in \mathbb{F}_q^*$  and  $n/e$  is even so  $n = 2m$  for some integer  $m$ . Then*

- (1) *If  $u \neq g^{t(2^e+1)}$  for any integer  $t$  then  $L_u$  is a PP. Let  $x_u \in \mathbb{F}_q$  be the unique solution of the equation  $L_u(x) = v^{2^k}$ . Then*

$$\mathfrak{S}(u, v) = (-1)^{m/e} 2^m \chi_1(ux_u^{2^k+1}).$$

- (2) *If  $u = g^{t(2^e+1)}$  for some integer  $t$ , then  $\mathfrak{S}(u, v) = 0$  unless the equation  $L_u(x) = v^{2^k}$  is solvable. If the equation  $L_u(x) = v^{2^k}$  is solvable with some solution, say  $x_u$ , then*

$$\mathfrak{S}(u, v) = \begin{cases} (-1)^{m/e} 2^m \chi_1(ux_u^{2^k+1}) & \text{if } \text{Tr}_e(u) \neq 0, \\ (-1)^{\frac{m}{e}+1} 2^{m+e} \chi_1(ux_u^{2^k+1}) & \text{if } \text{Tr}_e(u) = 0. \end{cases}$$

### 3. THE BINARY GOLD FUNCTION

In this section, we shall give the explicit expression for the  $c$ -BCT entries of the Gold function  $x^{2^k+1}$  over  $\mathbb{F}_{2^n}$ , for all  $c \neq 0$ . Recall that the  $c$ -boomerang uniformity of a power function  $F(x) = x^d$  over  $\mathbb{F}_{2^n}$  is given by  $\max_{b \in \mathbb{F}_{2^n}^*} {}_c\mathcal{B}_F(1, b)$ , where  ${}_c\mathcal{B}_F(1, b)$  is the number of solutions in  $\mathbb{F}_q \times \mathbb{F}_q$ ,  $q = 2^n$  of the following system

$$(3.1) \quad \begin{cases} x^d + cy^d = b \\ (x+1)^d + c^{-1}(y+1)^d = b. \end{cases}$$

As done in [14], for  $b \neq 0$  and fixed  $c \neq 0$ , the number of solutions  $(x, y) \in \mathbb{F}_q^2$  of the system (3.1) is given by

$$\begin{aligned} {}_c\mathcal{B}_F(1, b) &= \frac{1}{q^2} \sum_{x, y \in \mathbb{F}_q} \sum_{\alpha \in \mathbb{F}_q} \chi_1(\alpha(x^d + cy^d + b)) \sum_{\beta \in \mathbb{F}_q} \chi_1(\beta((x+1)^d + c^{-1}(y+1)^d + b)) \\ &= \frac{1}{q^2} \sum_{\alpha, \beta \in \mathbb{F}_q} \chi_1(b(\alpha + \beta)) \sum_{x \in \mathbb{F}_q} \chi_1(\alpha x^d + \beta(x+1)^d) \sum_{y \in \mathbb{F}_q} \chi_1(c\alpha y^d + c^{-1}\beta(y+1)^d) \\ &= \frac{1}{q^2} \sum_{\alpha, \beta \in \mathbb{F}_q} \chi_1(b(\alpha + \beta)) S_{\alpha, \beta} S_{c\alpha, c^{-1}\beta}, \end{aligned}$$

where  $S_{\alpha, \beta} = \sum_{x \in \mathbb{F}_q} \chi_1(\alpha x^d + \beta(x+1)^d)$ . Therefore, the problem of computing the  $c$ -BCT entry  ${}_c\mathcal{B}_F(1, b)$  is reduced to the computation of the product of the Weil sums  $S_{\alpha, \beta}$  and  $S_{c\alpha, c^{-1}\beta}$ . Now, in the particular case when  $d = 2^k + 1$ , i.e., for the Gold case, we shall further simplify the expression for  $S_{\alpha, \beta}$  as follows:

$$\begin{aligned} S_{\alpha, \beta} &= \sum_{x \in \mathbb{F}_q} \chi_1(\alpha x^{2^k+1} + \beta(x+1)^{2^k+1}) \\ &= \chi_1(\beta) \sum_{x \in \mathbb{F}_q} \chi_1((\alpha + \beta)x^{2^k+1}) \chi_1(\beta x^{2^k} + \beta x) \\ &= \chi_1(\beta) \sum_{x \in \mathbb{F}_q} \chi_1((\alpha + \beta)x^{2^k+1}) \chi_1((\beta^{2^{n-k}} x)^{2^k} + \beta x) \end{aligned}$$

$$\begin{aligned}
&= \chi_1(\beta) \sum_{x \in \mathbb{F}_q} \chi_1((\alpha + \beta)x^{2^k+1}) \chi_1((\beta^{2^{n-k}} + \beta)x) \\
&= \chi_1(\beta) \sum_{x \in \mathbb{F}_q} \chi_1((\alpha + \beta)x^{2^k+1} + (\beta^{2^{n-k}} + \beta)x) \\
&= \chi_1(\beta) \sum_{x \in \mathbb{F}_q} \chi_1(Ax^{2^k+1} + Bx),
\end{aligned}$$

where  $A = \alpha + \beta$  and  $B = \beta^{2^{n-k}} + \beta$ . Here one may note that  $A = 0$  if and only if  $\alpha = \beta$ . Also,  $B = 0$  if and only if  $\beta \in \mathbb{F}_{2^e}$ , since

$$\begin{aligned}
B = 0 &\Leftrightarrow \beta^{2^{n-k}} = \beta \\
&\Leftrightarrow \beta^{2^{n-k}-1} = 1 \\
&\Leftrightarrow \beta^{2^{\gcd(n-k, n)}-1} = 1 \\
&\Leftrightarrow \beta^{2^e-1} = 1, \text{ (as } \gcd(n-k, n) = e) \\
&\Leftrightarrow \beta \in \mathbb{F}_{2^e}.
\end{aligned}$$

Now we shall calculate  $S_{\alpha, \beta}$  in two cases, namely,  $n/e$  odd and  $n/e$  even, respectively.

**Case 1:**  $n/e$  is odd.

In this case, if  $\alpha = \beta$  and  $\beta \in \mathbb{F}_{2^e}$ , then  $S_{\alpha, \beta} = q\chi_1(\beta)$ . If  $\alpha = \beta$  and  $\beta \in \mathbb{F}_{2^n} \setminus \mathbb{F}_{2^e}$  then  $S_{\alpha, \beta} = 0$ . In the event of  $\alpha \neq \beta$  and  $\beta \in \mathbb{F}_{2^e}$ , again we have  $S_{\alpha, \beta} = 0$ . Finally, if  $\alpha \neq \beta$  and  $\beta \in \mathbb{F}_{2^n} \setminus \mathbb{F}_{2^e}$ , by Lemma 2.5 we have,

$$S_{\alpha, \beta} = \begin{cases} 0 & \text{if } \text{Tr}_e(B\gamma^{-1}) \neq 1, \\ \left(\frac{2}{n/e}\right)^e 2^{\frac{n+e}{2}} \chi_1(\beta) & \text{if } \text{Tr}_e(B\gamma^{-1}) = 1, \end{cases}$$

where  $\gamma \in \mathbb{F}_q$  is the unique element such that  $\gamma^{2^k+1} = A$ .

**Case 2:**  $n/e$  is even.

Let  $n = 2m$ , for some positive integer  $m$  and  $g$  be a primitive element of the finite field  $\mathbb{F}_q$ . When  $\alpha = \beta$  and  $\beta \in \mathbb{F}_{2^e}$  then  $S_{\alpha, \beta} = q\chi_1(\beta)$ . If  $\alpha = \beta$  and  $\beta \in \mathbb{F}_{2^n} \setminus \mathbb{F}_{2^e}$  then again  $S_{\alpha, \beta} = 0$ . In the event of  $\alpha \neq \beta$  and  $\beta \in \mathbb{F}_{2^e}$ , by Lemma 2.4 we have

$$S_{\alpha, \beta} = \begin{cases} (-1)^{m/e} 2^m \chi_1(\beta) & \text{if } A \neq g^{t(2^e+1)} \text{ for any integer } t, \\ (-1)^{\frac{m}{e}+1} 2^{m+e} \chi_1(\beta) & \text{if } A = g^{t(2^e+1)} \text{ for some integer } t. \end{cases}$$

Finally, when  $\alpha \neq \beta$  and  $\beta \in \mathbb{F}_{2^n} \setminus \mathbb{F}_{2^e}$ , we shall consider two cases depending on whether or not the linearized polynomial  $L_A(x) = A^{2^k} x^{2^{2k}} + Ax$  is a permutation polynomial. From Lemma 2.3,  $L_A$  is a permutation polynomial if and only if  $n/e$  is even and  $A \neq g^{t(2^e+1)}$  for any integer  $t$ . Therefore, when  $n/e$  is even and  $A \neq g^{t(2^e+1)}$  for any integer  $t$ , the equation  $L_A(x) = B^{2^k}$  will have a unique solution, say  $x_A$ . Therefore, by Lemma 2.6, we have

$$S_{\alpha, \beta} = (-1)^{m/e} 2^m \chi_1(\beta) \chi_1(Ax_A^{2^k+1}).$$

Now if the linearized polynomial  $L_A$  is not permutation, i.e,  $n/e$  is even and  $A = g^{t(2^e+1)}$  for some integer  $t$ , we again have two cases depending on whether or not the equation  $L_A(x) = B^{2^k}$  is solvable. In the case when equation  $L_A(x) = B^{2^k}$  is solvable, let  $x_A$  be one of its solution. Therefore, by Lemma 2.6 we have,

$$S_{\alpha, \beta} = \begin{cases} (-1)^{\frac{m}{e}+1} 2^{m+e} \chi_1(\beta) \chi_1(Ax_A^{2^k+1}) & \text{if } \text{Tr}_e(A) = 0, \\ (-1)^{\frac{m}{e}} 2^m \chi_1(\beta) \chi_1(Ax_A^{2^k+1}) & \text{if } \text{Tr}_e(A) \neq 0. \end{cases}$$

If  $L_A(x) = B^{2^k}$  is not solvable, again, by Lemma 2.6,  $S_{\alpha,\beta} = 0$ .

Thus we have computed  $S_{\alpha,\beta}$  in all possible cases. Similarly, we can find  $S_{c\alpha, c^{-1}\beta}$  by putting  $c\alpha$  and  $c^{-1}\beta$  in place of  $\alpha$  and  $\beta$ , respectively. We shall now explicitly compute the  $c$ -BCT entry  ${}_cB_F(1, b)$  for  $c = 1$ ,  $c \in \mathbb{F}_{2^e} \setminus \{0, 1\}$  and  $c \in \mathbb{F}_{2^n} \setminus \mathbb{F}_{2^e}$  in the forthcoming sections.

#### 4. THE CASE $c = 1$

When  $c = 1$ ,  $S_{\alpha,\beta}$  and  $S_{c\alpha, c^{-1}\beta}$  coincide, therefore for any fixed  $b \neq 0$ , the  $c$ -BCT entry is given by,

$${}_1B_F(1, b) = \frac{1}{q^2} \sum_{\alpha, \beta \in \mathbb{F}_q} \chi_1(b(\alpha + \beta)) S_{\alpha,\beta}^2.$$

Let us denote  $T_b = S_{\alpha,\beta}^2$ . Now we shall consider two cases, namely,  $n/e$  odd and  $n/e$  even, respectively.

**Case 1:**  $n/e$  is odd. We consider the following subcases.

(1) If  $\alpha = \beta$  and  $\beta \in \mathbb{F}_{2^e}$ , then

$$T_b^{[1]} = q^2 \chi_1(\beta)^2 = q^2.$$

(2) If  $\alpha = \beta$  and  $\beta \in \mathbb{F}_{2^n} \setminus \mathbb{F}_{2^e}$ , then

$$T_b^{[2]} = 0.$$

(3) If  $\alpha \neq \beta$  and  $\beta \in \mathbb{F}_{2^e}$ , then

$$T_b^{[3]} = 0.$$

(4) If  $\alpha \neq \beta$  and  $\beta \in \mathbb{F}_{2^n} \setminus \mathbb{F}_{2^e}$  then

$$T_b^{[4]} = \begin{cases} 0 & \text{if } \text{Tr}_e(B\gamma^{-1}) \neq 1, \\ 2^{n+e} & \text{if } \text{Tr}_e(B\gamma^{-1}) = 1. \end{cases}$$

Nyberg [11, Proposition 3] showed that the differential uniformity of the Gold function  $x \mapsto x^{2^k+1}$  over  $\mathbb{F}_{2^n}$  is  $2^e$ , where  $e = \gcd(k, n)$ . Also, from [4], we know that the boomerang uniformity of the APN function equals 2. Boura and Canteaut [2, Proposition 8] proved that when  $n/e$  is odd and  $n \equiv 2 \pmod{4}$ , then the differential as well as the boomerang uniformity of the Gold function  $x \mapsto x^{2^k+1}$  is 4. Our first theorem in this section generalizes the two previously mentioned results, and gives the boomerang uniformity of the Gold function for any parameters, when  $\frac{n}{e}$  is odd. Note that we would require the notion of Walsh-Hadamard transform in the proof of this theorem, which is defined as follows.

For  $f : \mathbb{F}_{2^n} \rightarrow \mathbb{F}_2$  we define the *Walsh-Hadamard transform* to be the integer-valued function

$$\mathcal{W}_f(u) = \sum_{x \in \mathbb{F}_{2^n}} (-1)^{f(x) + \text{Tr}(ux)}, u \in \mathbb{F}_{2^n}.$$

The Walsh transform  $\mathcal{W}_F(a, b)$  of an  $(n, m)$ -function  $F : \mathbb{F}_{2^n} \rightarrow \mathbb{F}_{2^m}$  at  $a \in \mathbb{F}_{2^n}, b \in \mathbb{F}_{2^m}$  is the Walsh-Hadamard transform of its component function  $\text{Tr}_1^m(bF(x))$  at  $a$ , that is,

$$\mathcal{W}_F(a, b) = \sum_{x \in \mathbb{F}_{2^n}} (-1)^{\text{Tr}_1^m(bF(x)) - \text{Tr}_1^n(ax)}.$$

**Theorem 4.1.** *Let  $F(x) = x^{2^k+1}$ ,  $1 \leq k < n$ , be a function on  $\mathbb{F}_q$ ,  $q = 2^n$ ,  $n \geq 2$ . Let  $c = 1$  and  $n/e$  be odd, where  $e = \gcd(k, n)$ . Then the  $c$ -BCT entry  ${}_1B_F(1, b)$  of  $F$  at  $(1, b)$  is*

$${}_1B_F(1, b) = 0, \text{ or, } 2^e,$$

if  $\text{Tr}_e\left(b^{\frac{1}{2}}\right) = 0$ , respectively,  $\text{Tr}_e\left(b^{\frac{1}{2}}\right) \neq 0$ .

*Proof.* For every  $\alpha, \beta$ , let  $A = \alpha + \beta$ ,  $B = \beta^{2^{-k}} + \beta$ , and  $\gamma \in \mathbb{F}_q$  be the unique element such that  $\gamma^{2^k+1} = A$ . Further, let

$$\begin{aligned}\mathcal{A} &= \{(\alpha, \beta) \in \mathbb{F}_q^2 \mid \alpha = \beta \in \mathbb{F}_{2^e}\}, \\ \mathcal{B} &= \{(\alpha, \beta) \in \mathbb{F}_q^2 \mid \alpha = \beta \in \mathbb{F}_q \setminus \mathbb{F}_{2^e}\}, \\ \mathcal{C} &= \{(\alpha, \beta) \in \mathbb{F}_q^2 \mid \alpha \neq \beta \text{ and } \beta \in \mathbb{F}_{2^e}\}, \\ \mathcal{D} &= \{(\alpha, \beta) \in \mathbb{F}_q^2 \mid \alpha \neq \beta \text{ and } \beta \in \mathbb{F}_q \setminus \mathbb{F}_{2^e}\}, \\ \mathcal{E} &= \{(\alpha, \beta) \in \mathcal{D} \mid \text{Tr}_e(B\gamma^{-1}) \neq 1\}, \\ \mathcal{F} &= \{(\alpha, \beta) \in \mathcal{D} \mid \text{Tr}_e(B\gamma^{-1}) = 1\}.\end{aligned}$$

Then,

$$\begin{aligned}_1\mathcal{B}_F(1, b) &= \frac{1}{q^2} \left( \sum_{(\alpha, \beta) \in \mathcal{A}} \chi_1(b(\alpha + \beta))T_b^{[1]} + \sum_{(\alpha, \beta) \in \mathcal{B}} \chi_1(b(\alpha + \beta))T_b^{[2]} + \sum_{(\alpha, \beta) \in \mathcal{C}} \chi_1(b(\alpha + \beta))T_b^{[3]} + \right. \\ &\quad \left. \sum_{(\alpha, \beta) \in \mathcal{E}} \chi_1(b(\alpha + \beta))T_b^{[4]} + \sum_{\alpha, \beta \in \mathcal{F}} \chi_1(b(\alpha + \beta))T_b^{[4]} \right) \\ &= \frac{1}{q^2} \left( \sum_{(\alpha, \beta) \in \mathcal{A}} q^2 + \sum_{(\alpha, \beta) \in \mathcal{F}} \chi_1(b(\alpha + \beta))2^{n+e} \right) \\ &= 2^e + \frac{2^e}{2^n} \sum_{(\alpha, \beta) \in \mathcal{F}} \chi_1(b(\alpha + \beta)).\end{aligned}$$

As customary,  $t^{-1} = t^{2^n-2}$ , rendering  $0^{-1} = 0$ . For each  $\beta \in \mathbb{F}_{2^n} \setminus \mathbb{F}_{2^e}$ , we let (if  $\beta \in \mathbb{F}_{2^e}$ ,  $Y_\beta = \mathbb{F}_{2^n}$ )

$$Y_\beta = \left\{ \gamma^{-1} \in \mathbb{F}_{2^n} : \text{Tr}_e\left((\beta^{2^{-k}} + \beta)\gamma^{-1}\right) = 1 \right\},$$

and

$$T_\beta = \left\{ d \in \mathbb{F}_{2^n} : \text{Tr}_e((\beta^{2^{-k}} + \beta)d) = 0 \right\} = \langle \beta^{2^{-k}} + \beta \rangle^{\perp_e}.$$

We shall use below that when  $\frac{n}{e}$  is odd, then  $\text{Tr}_e(1) = 1$ . We label by  $\langle S \rangle_e$  the  $\mathbb{F}_{2^e}$ -linear subspace in  $\mathbb{F}_{2^n}$  generate by  $S$  and we write  $S^{\perp_e}$ , for the trace orthogonal (via the relative trace  $\text{Tr}_e$ ) of the subspace  $\langle S \rangle_e$  (if  $e = 1$ , we drop the subscripts). Since  $\text{Tr}_e(1) = 1$ , then,  $(\beta^{2^{-k}} + \beta)^{-1} \in Y_\beta$ . If  $\gamma_1^{-1}, \gamma_2^{-1} \in Y_\beta$ , then  $\gamma_1^{-1} + \gamma_2^{-1} \in T_\beta$ , of cardinality  $|T_\beta| = 2^{n-1}$ . Reciprocally, if  $\gamma^{-1} \in Y_\beta$  and  $d \in T_\beta$ , it is easy to see that  $\gamma^{-1} + d \in Y_\beta$ . Therefore,  $Y_\beta$  is the affine subspace  $Y_\beta = \gamma_\beta + T_\beta$ , where  $\gamma_\beta = (\beta^{2^{-k}} + \beta)^{-1}$ .

Next, we observe that the kernel of  $\phi : \beta \mapsto \beta^{2^{-k}} + \beta$ , say  $\ker(\phi)$ , is an  $\mathbb{F}_2$ -linear space of dimension  $e$  (in fact, it is exactly  $\mathbb{F}_{2^e}$ ) and the image of  $\phi$ , say  $\text{Im}(\phi)$ , is an  $\mathbb{F}_2$ -linear space of dimension  $n - e$ . Further, we show that  $\text{Im}(\phi)^{\perp_e} = \ker(\phi)$ . We use below the fact that  $\text{Tr}_e(x^{2^e}) = \text{Tr}_e(x)$  and  $e \mid k$ . Let  $u \in \text{Im}(\phi)^{\perp_e}$ , that is, for all  $\beta \in \mathbb{F}_{2^n}$ ,

$$0 = \text{Tr}_e(u(\beta^{2^{-k}} + \beta)) = \text{Tr}_e(u\beta^{2^{-k}}) + \text{Tr}_e(u\beta) = \text{Tr}_e(u^{2^k}\beta) + \text{Tr}_e(u\beta) = \text{Tr}_e((u + u^{2^k})\beta),$$

and so,  $u^{2^k} + u = 0$ , which shows the claim. For easy referral, if we speak of the dimension of an  $\mathbb{F}_{2^e}$ -linear space  $S$ , we shall be using the notation  $\dim_e S$  (no subscript if  $e = 1$ ).

We will be using below the Poisson summation formula (see [3, Corollary 8.9] and [6, Theorem 2.15]), which states that if  $f : \mathbb{F}_{2^n} \rightarrow \mathbb{R}$  and  $S$  is a subspace of  $\mathbb{F}_{2^n}$  of dimension  $\dim S$ , then

$$\sum_{u \in \alpha + S} \mathcal{W}_f(u)(-1)^{\text{Tr}(\beta u)} = 2^{\dim S} (-1)^{\text{Tr}(\alpha \beta)} \sum_{u \in \beta + S^\perp} f(u)(-1)^{\text{Tr}(\alpha u)},$$

and in particular,

$$\sum_{u \in S} \mathcal{W}_f(u) = 2^{\dim S} \sum_{u \in S^\perp} f(u).$$

Now, we are able to compute our sum (labelling  $\alpha = \beta + \gamma^{2^k+1}$ , and writing  $\phi^{-1}(t) = \{\beta : \phi(\beta) = t\}$ ; we also note that when  $\frac{n}{e}$  is odd,  $\gcd(2^k + 1, 2^n - 1) = 1$ , and so  $\gamma \mapsto \gamma^{2^k+1}$  is a permutation)

$$\begin{aligned} {}_1\mathcal{B}_F(1, b) &= 2^e + \frac{2^e}{2^n} \sum_{\substack{\beta \in \mathbb{F}_{2^n} \setminus \mathbb{F}_{2^e}, \gamma \in \mathbb{F}_{2^n} \\ \text{Tr}_e((\beta^{2^{-k}} + \beta)\gamma^{-1}) = 1}} \chi_1(b\gamma^{2^k+1}) \\ &= 2^e + \frac{2^e}{2^n} \sum_{\beta \in \mathbb{F}_{2^n} \setminus \mathbb{F}_{2^e}} \sum_{\gamma^{-1} \in Y_\beta} \chi_1(b\gamma^{2^k+1}) \\ &= 2^e + \frac{2^e}{2^n} \sum_{\beta \in \mathbb{F}_{2^n}} \sum_{x \in (\beta^{2^{-k}} + \beta)^{-1} + \langle \beta^{2^{-k}} + \beta \rangle^{\perp_e}} \chi_1(bx^{-2^k-1}) \\ &\quad \text{(we used here that } Y_\beta = (\beta^{2^{-k}} + \beta)^{-1} + T_\beta; \text{ we also added} \\ &\quad \beta \in \mathbb{F}_{2^e}, \text{ as it contributes 0 to the inner sum)} \\ &= 2^e + \frac{2^e}{2^n} \sum_{\beta \in \mathbb{F}_{2^n}} 2^{-\dim S} \sum_{u \in (\langle \beta^{2^{-k}} + \beta \rangle^{\perp_e})^\perp} \mathcal{W}_{g_\beta}(u)(-1)^{\text{Tr}(u(\beta^{2^{-k}} + \beta)^{-1})} \\ &\quad \text{(by Poisson summation with } S^\perp = \langle \beta^{2^{-k}} + \beta \rangle^{\perp_e}, \text{ and } g_\beta(x) = \chi_1(bx^{-2^k-1})). \end{aligned}$$

We now analyze the  $\mathbb{F}_2$ -linear space

$$\left( \langle \beta^{2^{-k}} + \beta \rangle^{\perp_e} \right)^\perp = \{x \in \mathbb{F}_{2^n} : \text{Tr}(dx) = 0, \forall d \text{ with } \text{Tr}_e(d(\beta^{2^{-k}} + \beta)) = 0\}.$$

Further,  $\mathbb{F}_{2^n}$  has dimension  $n/e$  as an  $\mathbb{F}_{2^e}$ -linear space and so,  $\dim_e \langle \beta^{2^{-k}} + \beta \rangle^{\perp_e} = \frac{n}{e} - 1$  as an  $\mathbb{F}_{2^e}$ -linear space, and since  $\mathbb{F}_{2^e}$  has dimension  $e$  as an  $\mathbb{F}_2$ -linear space, then  $\dim \langle \beta^{2^{-k}} + \beta \rangle^{\perp_e} = n - e$  as an  $\mathbb{F}_2$ -linear space. Thus,  $\dim \left( \langle \beta^{2^{-k}} + \beta \rangle^{\perp_e} \right)^\perp = e$ . Moreover,  $\text{Tr}_e(\beta^{2^{-k}} + \beta) = 0$  and if  $u \in \mathbb{F}_{2^e}$  then  $\text{Tr}_e(u(\beta^{2^{-k}} + \beta)) = u \text{Tr}_e(\beta^{2^{-k}} + \beta) = 0$ , and consequently (since the dimensions match and  $(\beta^{2^{-k}} + \beta)\mathbb{F}_{2^e} \subseteq S$ )

$$S = \left( \langle \beta^{2^{-k}} + \beta \rangle^{\perp_e} \right)^\perp = (\beta^{2^{-k}} + \beta)\mathbb{F}_{2^e}.$$

We are now ready to continue the computation, thus,

$$\begin{aligned} {}_1\mathcal{B}_F(1, b) &= 2^e + \frac{2^e}{2^n} 2^{-e} \sum_{\beta \in \mathbb{F}_{2^n}} \sum_{u \in (\beta^{2^{-k}} + \beta)\mathbb{F}_{2^e}} \mathcal{W}_{g_\beta}(u)(-1)^{\text{Tr}(u(\beta^{2^{-k}} + \beta)^{-1})} \\ &= 2^e + \frac{2^e}{2^n} 2^{-e} \sum_{\beta \in \mathbb{F}_{2^n}} \sum_{d' \in \mathbb{F}_{2^e}} \mathcal{W}_{g_\beta}(d'(\beta^{2^{-k}} + \beta))(-1)^{\text{Tr}(d')} \end{aligned}$$



$$\begin{aligned}
&= 2^e + \frac{2^e}{2^n} 2^{-e} \sum_{\beta \in \mathbb{F}_{2^n}} \sum_{d' \in \mathbb{F}_{2^e}} \sum_{x \in \mathbb{F}_{2^n}} \chi_1 \left( bx^{-2^k-1} + d'x(\beta^{2^{-k}} + \beta) + d' \right) \\
&= 2^e + \frac{2^e}{2^n} 2^{-e} \sum_{d' \in \mathbb{F}_{2^e}} \sum_{x \in \mathbb{F}_{2^n}} \chi_1 \left( bx^{-2^k-1} + d' \right) \sum_{\beta \in \mathbb{F}_{2^n}} \chi_1 \left( d'x(\beta^{2^{-k}} + \beta) \right) \\
&= 2^e + \frac{2^e}{2^n} 2^{-e} \sum_{d' \in \mathbb{F}_{2^e}} \sum_{x \in \mathbb{F}_{2^n}} \chi_1 \left( bx^{-2^k-1} + d' \right) \sum_{\beta \in \mathbb{F}_{2^n}} \chi_1 \left( \left( (d'x)^{2^k} + d'x \right) \beta \right) \\
&\quad \left( \text{since } \text{Tr} \left( d'x(\beta^{2^{-k}} + \beta) \right) = \text{Tr} \left( ((d'x)^{2^k} + d'x)\beta \right) = \text{Tr}(d'(x^{2^k} + x)\beta) \right) \\
&= 2^e + \frac{2^e}{2^n} 2^{n-e} \sum_{\substack{d' \in \mathbb{F}_{2^e}, x \in \mathbb{F}_{2^n} \\ d'(x^{2^k} + x) = 0}} \chi_1 \left( bx^{-2^k-1} + d' \right) \\
&= 2^e + \frac{2^e}{2^n} 2^{n-e} \sum_{d' \in \mathbb{F}_{2^e}^*, x \in \mathbb{F}_{2^e}} \chi_1(bx^{-2} + d') + \sum_{x \in \mathbb{F}_{2^n}} \chi_1(bx^{-2^k-1}) \\
&= 2^e + \frac{2^e}{2^n} 2^{n-e} \sum_{d' \in \mathbb{F}_{2^e}^*, x \in \mathbb{F}_{2^e}} \chi_1(bx^{-2} + d') \\
&= 2^e - 2^e \delta_0 \left( \text{Tr}_e \left( b^{\frac{1}{2}} \right) \right),
\end{aligned}$$

where  $\delta_0$  is the Dirac symbol, defined by  $\delta_0(c) = 1$ , if  $c = 0$ , and 0, otherwise. Thus,  ${}_1\mathcal{B}_F(1, b) \in \{0, 2^e\}$ , and the claim of our theorem is shown.  $\square$

**Case 2:**  $n/e$  is even.

(1) If  $\alpha = \beta$  and  $\beta \in \mathbb{F}_{2^e}$ , then

$$T_b^{[1]} = q^2 \chi_1(\beta)^2 = q^2.$$

(2) If  $\alpha = \beta$  and  $\beta \in \mathbb{F}_{2^n} \setminus \mathbb{F}_{2^e}$ , then

$$T_b^{[2]} = 0.$$

(3) If  $\alpha \neq \beta$  and  $\beta \in \mathbb{F}_{2^e}$ , then

$$T_b^{[3]} = \begin{cases} 2^n & \text{if } A \neq g^{t(2^e+1)} \text{ for any integer } t, \\ 2^{n+2e} & \text{if } A = g^{t(2^e+1)} \text{ for some integer } t. \end{cases}$$

(4) If  $\alpha \neq \beta$  and  $\beta \in \mathbb{F}_{2^n} \setminus \mathbb{F}_{2^e}$ , then

(a) If  $A \neq g^{t(2^e+1)}$  for any integer  $t$ , then

$$T_b^{[4(a)]} = 2^n.$$

(b) If  $A = g^{t(2^e+1)}$  for some integer  $t$ , then

(i) If the equation  $L_A(x) = B^{2^k}$  is not solvable, where  $L_A(x) = A^{2^k} x^{2^{2k}} + Ax$ , then

$$T_b^{[4(b)(i)]} = 0.$$

(ii) If the equation  $L_A(x) = B^{2^k}$  is solvable, then

$$T_b^{[4(b)(ii)]} = \begin{cases} 2^n & \text{if } \text{Tr}_e(A) \neq 0, \\ 2^{n+2e} & \text{if } \text{Tr}_e(A) = 0. \end{cases}$$

Now we shall summarize the above discussion in the following theorem.

**Theorem 4.2.** *Let  $F(x) = x^{2^k+1}$ ,  $1 \leq k < n$  be a function on  $\mathbb{F}_{2^n}$ ,  $n \geq 2$ . Let  $c = 1$  and  $n/e$  be even, where  $e = \gcd(k, n)$ . Then the  $c$ -BCT entry  ${}_1\mathcal{B}_F(1, b)$  of  $F$  at  $(1, b)$  is given by*

$$2^e + \frac{1}{2^n} \sum_{(\alpha, \beta) \in \mathcal{G} \cup \mathcal{I} \cup \mathcal{K}} \chi_1(b(\alpha + \beta)) + \frac{2^{2e}}{2^n} \sum_{(\alpha, \beta) \in \mathcal{H} \cup \mathcal{L}} \chi_1(b(\alpha + \beta)),$$

with  $A = \alpha + \beta$ ,  $B = \beta^{2^{n-k}} + \beta$ ,  $L_A(x) = A^{2^k} x^{2^{2k}} + Ax$ , and

$$\mathcal{G} = \{(\alpha, \beta) \in \mathcal{C} \mid A \neq g^{t(2^e+1)} \text{ for any integer } t\},$$

$$\mathcal{H} = \{(\alpha, \beta) \in \mathcal{C} \mid A = g^{t(2^e+1)} \text{ for some integer } t\},$$

$$\mathcal{I} = \{(\alpha, \beta) \in \mathcal{D} \mid A \neq g^{t(2^e+1)} \text{ for any integer } t\},$$

$$\mathcal{K} = \{(\alpha, \beta) \in \mathcal{D} \mid A = g^{t(2^e+1)} \text{ for some integer } t, \text{Tr}_e(A) \neq 0, L_A(x) = B^{2^k} \text{ is solvable}\},$$

$$\mathcal{L} = \{(\alpha, \beta) \in \mathcal{D} \mid A = g^{t(2^e+1)} \text{ for some integer } t, \text{Tr}_e(A) = 0, L_A(x) = B^{2^k} \text{ is solvable}\}.$$

*Proof.* For the proof, we need to define

$$\mathcal{J} = \{(\alpha, \beta) \in \mathcal{D} \mid A = g^{t(2^e+1)} \text{ for an integer } t, L_A(x) = B^{2^k} \text{ is not solvable}\}.$$

Then

$$\begin{aligned} {}_1\mathcal{B}_F(1, b) &= \frac{1}{q^2} \left( \sum_{(\alpha, \beta) \in \mathcal{A}} \chi_1(b(\alpha + \beta)) T_b^{[1]} + \sum_{(\alpha, \beta) \in \mathcal{B}} \chi_1(b(\alpha + \beta)) T_b^{[2]} \right. \\ &\quad + \sum_{(\alpha, \beta) \in \mathcal{G}} \chi_1(b(\alpha + \beta)) T_b^{[3]} + \sum_{(\alpha, \beta) \in \mathcal{H}} \chi_1(b(\alpha + \beta)) T_b^{[3]} \\ &\quad + \sum_{(\alpha, \beta) \in \mathcal{I}} \chi_1(b(\alpha + \beta)) T_b^{[4(a)]} + \sum_{(\alpha, \beta) \in \mathcal{J}} \chi_1(b(\alpha + \beta)) T_b^{[4(b)(i)]} \\ &\quad \left. + \sum_{(\alpha, \beta) \in \mathcal{K}} \chi_1(b(\alpha + \beta)) T_b^{[4(b)(ii)]} + \sum_{(\alpha, \beta) \in \mathcal{L}} \chi_1(b(\alpha + \beta)) T_b^{[4(b)(ii)]} \right) \\ &= \frac{1}{q^2} \left( \sum_{(\alpha, \beta) \in \mathcal{A}} q^2 + 2^n \sum_{(\alpha, \beta) \in \mathcal{G} \cup \mathcal{I} \cup \mathcal{K}} \chi_1(b(\alpha + \beta)) + 2^{n+2e} \sum_{(\alpha, \beta) \in \mathcal{H} \cup \mathcal{L}} \chi_1(b(\alpha + \beta)) \right) \\ &= 2^e + \frac{1}{2^n} \sum_{(\alpha, \beta) \in \mathcal{G} \cup \mathcal{I} \cup \mathcal{K}} \chi_1(b(\alpha + \beta)) + \frac{2^{2e}}{2^n} \sum_{(\alpha, \beta) \in \mathcal{H} \cup \mathcal{L}} \chi_1(b(\alpha + \beta)). \end{aligned}$$

This completes the proof.  $\square$

**Corollary 4.3.** *Let  $F(x) = x^{2^k+1}$ ,  $1 \leq k < n$ , be a function on  $\mathbb{F}_q$ ,  $n \geq 2$ . Let  $c = 1$  and  $n/e$  be even, where  $e = \gcd(k, n)$ . With the notations of the previous theorem, the  $c$ -boomerang uniformity of  $F$  satisfies*

$$\beta_{F,c} \leq 2^e + 2^{-n} |\mathcal{G} \cup \mathcal{I} \cup \mathcal{K}| + 2^{2e-n} |\mathcal{H} \cup \mathcal{L}|.$$

## 5. THE CASE $c \in \mathbb{F}_{2^e} \setminus \{0, 1\}$ .

Since the case  $c = 1$  has already been considered in the previous section, throughout this section we assume that  $c \neq 1$ . Notice that when  $c \in \mathbb{F}_{2^e}^*$ ,  $\beta \in \mathbb{F}_{2^e} \Leftrightarrow \beta c^{-1} \in \mathbb{F}_{2^e}$ . Recall that

for any fixed  $b \neq 0$ , the  $c$ -BCT entry is given by,

$${}_c\mathcal{B}_F(1, b) = \frac{1}{q^2} \sum_{\alpha, \beta \in \mathbb{F}_q} \chi_1(b(\alpha + \beta)) S_{\alpha, \beta} S_{c\alpha, c^{-1}\beta}.$$

Let us denote  $T_b = S_{\alpha, \beta} S_{c\alpha, c^{-1}\beta}$  (we will use superscripts to point out the case we are in, for its value). Recall that  $A = \alpha + \beta$  and  $B = \beta^{2^{n-k}} + \beta$ . Let us denote  $\gamma = A^{\frac{1}{2^k+1}}$ ,  $A' = c\alpha + c^{-1}\beta$  and  $B' = (c^{-1}\beta)^{2^{n-k}} + c^{-1}\beta$ . It is easy to observe that the conditions  $B = 0$  and  $B' = 0$  are equivalent. Now we shall consider two cases namely,  $\frac{n}{e}$  odd and  $\frac{n}{e}$  even, respectively.

**Case 1:**  $\frac{n}{e}$  is odd.

(1) Let  $A = 0, B = 0$ .

(a) If  $A' = 0, B' = 0$ , then

$$T_b^{[1(a)]} = q^2 \chi_1((1 + c^{-1})\beta).$$

(b) If  $A' \neq 0, B' = 0$ , then  $S_{c\alpha, c^{-1}\beta} = 0$  and hence

$$T_b^{[1(b)]} = 0.$$

(2) Let  $A = 0, B \neq 0$ . In this case  $S_{\alpha, \beta} = 0$  and hence

$$T_b^{[2]} = 0.$$

(3) Let  $A \neq 0, B = 0$ . Again  $S_{\alpha, \beta} = 0$  and hence

$$T_b^{[3]} = 0$$

(4) Let  $A \neq 0, B \neq 0$ .

(a) Assume  $A' = 0, B' \neq 0$ , then  $S_{c\alpha, c^{-1}\beta} = 0$  and hence

$$T_b^{[4(a)]} = 0.$$

(b) Assume  $A' \neq 0, B' \neq 0$ . In this case, recall that  $\gamma^{2^k+1} = A$  and let  $\gamma' \in \mathbb{F}_q$  such that  $(\gamma')^{2^k+1} = A'$ .

(i) If  $\text{Tr}_e(B\gamma^{-1}) \neq 1$ , then  $S_{\alpha, \beta} = 0$  and hence

$$T_b^{[4(b)(i)]} = 0.$$

(ii) If  $\text{Tr}_e(B\gamma^{-1}) = 1$  and  $\text{Tr}_e(B'(\gamma')^{-1}) \neq 1$ , then  $S_{c\alpha, c^{-1}\beta} = 0$  and hence

$$T_b^{[4(b)(ii)]} = 0.$$

(iii) If  $\text{Tr}_e(B\gamma^{-1}) = 1$  and  $\text{Tr}_e(B'(\gamma')^{-1}) = 1$ , then

$$T_b^{[4(b)(iii)]} = 2^{n+e} \chi_1((1 + c^{-1})\beta).$$

We now use the above discussion in the following theorem.

**Theorem 5.1.** *Let  $F(x) = x^{2^k+1}$ ,  $1 \leq k < n$  be a function on  $\mathbb{F}_{2^n}$ ,  $n \geq 2$ . Let  $c \in \mathbb{F}_{2^e} \setminus \{0, 1\}$  and  $n/e$  be odd, where  $e = \gcd(k, n)$ . Then the  $c$ -BCT entry  ${}_c\mathcal{B}_F(1, b)$  of  $F$  at  $(1, b)$  is given by*

$$1 + \frac{2^e}{2^n} \sum_{(\alpha, \beta) \in \mathcal{F} \cap \mathcal{F}'} \chi_1(b\alpha + (1 + c^{-1} + b)\beta),$$

where

$$\mathcal{F} = \{(\alpha, \beta) \in \mathbb{F}_q^2 \mid A, B \neq 0 \text{ and } \text{Tr}_e(B\gamma^{-1}) = 1\},$$

$$\mathcal{F}' = \{(\alpha, \beta) \in \mathbb{F}_q^2 \mid A', B' \neq 0 \text{ and } \text{Tr}_e(B'(\gamma')^{-1}) = 1\},$$

and  $A = \alpha + \beta$ ,  $B = \beta^{2^{n-k}} + \beta$ ,  $A' = c\alpha + c^{-1}\beta$  and  $B' = (c^{-1}\beta)^{2^{n-k}} + c^{-1}\beta$ ,  $\gamma = A^{\frac{1}{2^k+1}}$ ,  $\gamma' = A'^{\frac{1}{2^k+1}}$ .

*Proof.* Let

$$\begin{aligned}\mathcal{A}' &= \{(\alpha, \beta) \in \mathbb{F}_q^2 \mid c\alpha = c^{-1}\beta \text{ and } c^{-1}\beta \in \mathbb{F}_{2^e}\}, \\ \mathcal{B}' &= \{(\alpha, \beta) \in \mathbb{F}_q^2 \mid c\alpha = c^{-1}\beta \text{ and } c^{-1}\beta \in \mathbb{F}_q \setminus \mathbb{F}_{2^e}\}, \\ \mathcal{C}' &= \{(\alpha, \beta) \in \mathbb{F}_q^2 \mid c\alpha \neq c^{-1}\beta \text{ and } c^{-1}\beta \in \mathbb{F}_{2^e}\}, \\ \mathcal{D}' &= \{(\alpha, \beta) \in \mathbb{F}_q^2 \mid c\alpha \neq c^{-1}\beta \text{ and } c^{-1}\beta \in \mathbb{F}_q \setminus \mathbb{F}_{2^e}\}, \\ \mathcal{E}' &= \{(\alpha, \beta) \in \mathcal{D}' \mid \text{Tr}_e(B'(\gamma')^{-1}) \neq 1\}.\end{aligned}$$

Then,

$$\begin{aligned}_c\mathcal{B}_F(1, b) &= \frac{1}{q^2} \left( \sum_{(\alpha, \beta) \in \mathcal{A} \cap \mathcal{A}'} \chi_1(b(\alpha + \beta)) T_b^{[1(a)]} + \sum_{(\alpha, \beta) \in \mathcal{A} \cap \mathcal{C}'} \chi_1(b(\alpha + \beta)) T_b^{[1(b)]} \right. \\ &\quad + \sum_{(\alpha, \beta) \in \mathcal{B}} \chi_1(b(\alpha + \beta)) T_b^{[2]} + \sum_{(\alpha, \beta) \in \mathcal{C}} \chi_1(b(\alpha + \beta)) T_b^{[3]} \\ &\quad + \sum_{(\alpha, \beta) \in \mathcal{D} \cap \mathcal{B}'} \chi_1(b(\alpha + \beta)) T_b^{[4(a)]} + \sum_{(\alpha, \beta) \in \mathcal{E}} \chi_1(b(\alpha + \beta)) T_b^{[4(b)(i)]} \\ &\quad \left. + \sum_{(\alpha, \beta) \in \mathcal{F} \cap \mathcal{E}'} \chi_1(b(\alpha + \beta)) T_b^{[4(b)(ii)]} + \sum_{(\alpha, \beta) \in \mathcal{F} \cap \mathcal{F}'} \chi_1(b(\alpha + \beta)) T_b^{[4(b)(iii)]} \right) \\ &= \sum_{(\alpha, \beta) \in \mathcal{A} \cap \mathcal{A}'} \chi_1(b\alpha + (1 + c^{-1} + b)\beta) \\ &\quad + \frac{2^e}{2^n} \sum_{(\alpha, \beta) \in \mathcal{F} \cap \mathcal{F}'} \chi_1(b\alpha + (1 + c^{-1} + b)\beta) \\ &= 1 + \frac{2^e}{2^n} \sum_{(\alpha, \beta) \in \mathcal{F} \cap \mathcal{F}'} \chi_1(b\alpha + (1 + c^{-1} + b)\beta).\end{aligned}$$

This completes the proof.  $\square$

**Corollary 5.2.** Let  $F(x) = x^{2^k+1}$ ,  $1 \leq k < n$ , be a function on  $\mathbb{F}_q$ ,  $n \geq 2$ . Let  $c \in \mathbb{F}_{2^e} \setminus \{0, 1\}$  and  $n/e$  be odd, where  $e = \gcd(k, n)$ . With the notations of the previous theorem, the  $c$ -boomerang uniformity of  $F$  satisfies

$$\beta_{F,c} \leq 1 + 2^{e-n} |\mathcal{F} \cap \mathcal{F}'|.$$

**Case 2:**  $n/e$  is even.

(1) Let  $A = 0, B = 0$ .

(a) If  $A' = 0, B' = 0$ , then

$$T_b^{[1(a)]} = \chi_1((1 + c^{-1})\beta) q^2.$$

(b) If  $A' \neq 0, B' = 0$ , let

$$\mathcal{G}' = \{(\alpha, \beta) \in \mathcal{C}' \mid A' \neq g^{t(2^e+1)} \text{ for any integer } t\},$$

$$\mathcal{H}' = \{(\alpha, \beta) \in \mathcal{C}' \mid A' = g^{t(2^e+1)} \text{ for some integer } t\}.$$

Then,

$$T_b^{[1(b)]} = \begin{cases} (-1)^{\frac{m}{e}} 2^{m+n} \chi_1((1+c^{-1})\beta) & \text{if } (\alpha, \beta) \in \mathcal{A} \cap \mathcal{G}', \\ (-1)^{\frac{m}{e}+1} 2^{m+n+e} \chi_1((1+c^{-1})\beta) & \text{if } (\alpha, \beta) \in \mathcal{A} \cap \mathcal{H}'. \end{cases}$$

(2) Let  $A = 0, B \neq 0$ .

In this case  $S_{\alpha, \beta} = 0$  and hence

$$T_b^{[2]} = 0.$$

(3) Let  $A \neq 0, B = 0$ .

(a) If  $A' = 0, B' = 0$ , then  $T_b^{[3(a)]}$  is given by

$$\begin{cases} (-1)^{\frac{m}{e}} 2^{m+n} \chi_1((1+c^{-1})\beta) & \text{if } (\alpha, \beta) \in \mathcal{A}' \cap \mathcal{G}, \\ (-1)^{\frac{m}{e}+1} 2^{m+n+e} \chi_1((1+c^{-1})\beta) & \text{if } (\alpha, \beta) \in \mathcal{A}' \cap \mathcal{H}. \end{cases}$$

(b) If  $A' \neq 0, B' = 0$ , then

$$T_b^{[3(b)]} = \begin{cases} 2^n \chi_1((1+c^{-1})\beta) & \text{if } (\alpha, \beta) \in \mathcal{G} \cap \mathcal{G}', \\ -2^{n+e} \chi_1((1+c^{-1})\beta) & \text{if } (\alpha, \beta) \in \mathcal{G} \cap \mathcal{H}', \\ -2^{n+e} \chi_1((1+c^{-1})\beta) & \text{if } (\alpha, \beta) \in \mathcal{H} \cap \mathcal{G}', \\ 2^{n+2e} \chi_1((1+c^{-1})\beta) & \text{if } (\alpha, \beta) \in \mathcal{H} \cap \mathcal{H}'. \end{cases}$$

(4) Let  $A \neq 0, B \neq 0$ .

(a) If  $A' = 0, B' \neq 0$ , then  $S_{c\alpha, c^{-1}\beta} = 0$  and hence

$$T_b^{[4(a)]} = 0.$$

(b) If  $A' \neq 0, B' \neq 0$ , let

$$\begin{aligned} \mathcal{I}' &= \{(\alpha, \beta) \in \mathcal{D}' \mid A' \neq g^{t(2^e+1)} \text{ for any integer } t\}, \\ \mathcal{J}' &= \{(\alpha, \beta) \in \mathcal{D}' \mid A' = g^{t(2^e+1)} \text{ for some integer } t, \\ &\quad L_{A'}(x) = (B')^{2^k} \text{ is not solvable}\}, \\ \mathcal{K}' &= \{(\alpha, \beta) \in \mathcal{D}' \mid A' = g^{t(2^e+1)} \text{ for some integer } t, \\ &\quad \text{Tr}_e(A') \neq 0, L_{A'}(x) = (B')^{2^k} \text{ is solvable}\}, \\ \mathcal{L}' &= \{(\alpha, \beta) \in \mathcal{D}' \mid A' = g^{t(2^e+1)} \text{ for some integer } t, \\ &\quad \text{Tr}_e(A') = 0, L_{A'}(x) = (B')^{2^k} \text{ is solvable}\}. \end{aligned}$$

Then,

$$T_b^{[4(b)]} = \begin{cases} 2^n \cdot M & \text{if } (\alpha, \beta) \in (\mathcal{I} \cup \mathcal{K}) \cap (\mathcal{I}' \cup \mathcal{K}'), \\ 0 & \text{if } (\alpha, \beta) \in (\mathcal{I} \cup \mathcal{K} \cup \mathcal{L}) \cap \mathcal{J}', \\ -2^{n+e} \cdot M & \text{if } (\alpha, \beta) \in (\mathcal{I} \cup \mathcal{K}) \cap \mathcal{L}', \\ 0 & \text{if } (\alpha, \beta) \in \mathcal{J} \cap (\mathcal{I}' \cup \mathcal{J}' \cup \mathcal{K}' \cup \mathcal{L}'), \\ -2^{n+e} \cdot M & \text{if } (\alpha, \beta) \in \mathcal{L} \cap (\mathcal{I}' \cup \mathcal{K}'), \\ 2^{n+2e} \cdot M & \text{if } (\alpha, \beta) \in \mathcal{L} \cap \mathcal{L}', \end{cases}$$

where  $M = \chi_1((1+c^{-1})\beta) \chi_1(AA'x_A^{2^k+1}x_{A'}^{2^k+1})$  and  $x_A, x_{A'}$  are the solutions of the equations  $L_A(x) = B^{2^k}$  and  $L_{A'}(x) = (B')^{2^k}$ , respectively.

We now summarize the above discussion in the following theorem.

**Theorem 5.3.** *Let  $F(x) = x^{2^k+1}$ ,  $1 \leq k < n$  be a function on  $\mathbb{F}_{2^n}$ ,  $n \geq 2$ . Let  $c \in \mathbb{F}_{2^e} \setminus \{0, 1\}$  and  $n/e$  be even, where  $e = \gcd(k, n)$ . With the previous notations, the  $c$ -BCT entry  ${}_c\mathcal{B}_F(1, b)$  of  $F$  at  $(1, b)$  is given by*

$$\begin{aligned} \frac{1}{q^2} & \left( \sum_{(\alpha, \beta) \in \mathcal{A} \cap \mathcal{A}'} \chi_1(b(\alpha + \beta)) T_b^{[1(a)]} + \sum_{(\alpha, \beta) \in \mathcal{A} \cap \mathcal{G}'} \chi_1(b(\alpha + \beta)) T_b^{[1(b)]} \right. \\ & + \sum_{(\alpha, \beta) \in \mathcal{A} \cap \mathcal{H}'} \chi_1(b(\alpha + \beta)) T_b^{[1(b)]} + \sum_{(\alpha, \beta) \in \mathcal{A}' \cap \mathcal{G}} \chi_1(b(\alpha + \beta)) T_b^{[3(a)]} \\ & + \sum_{(\alpha, \beta) \in \mathcal{A}' \cap \mathcal{H}} \chi_1(b(\alpha + \beta)) T_b^{[3(a)]} + \sum_{(\alpha, \beta) \in \mathcal{G} \cap \mathcal{G}'} \chi_1(b(\alpha + \beta)) T_b^{[3(b)]} \\ & + \sum_{(\alpha, \beta) \in \mathcal{G} \cap \mathcal{H}'} \chi_1(b(\alpha + \beta)) T_b^{[3(b)]} + \sum_{(\alpha, \beta) \in \mathcal{H} \cap \mathcal{G}'} \chi_1(b(\alpha + \beta)) T_b^{[3(b)]} \\ & + \sum_{(\alpha, \beta) \in \mathcal{H} \cap \mathcal{H}'} \chi_1(b(\alpha + \beta)) T_b^{[3(b)]} + \sum_{(\alpha, \beta) \in (\mathcal{I} \cup \mathcal{K}) \cap (\mathcal{I}' \cup \mathcal{K}')} \chi_1(b(\alpha + \beta)) T_b^{[4(b)]} \\ & + \sum_{(\alpha, \beta) \in (\mathcal{I} \cup \mathcal{K}) \cap \mathcal{L}'} \chi_1(b(\alpha + \beta)) T_b^{[4(b)]} + \sum_{(\alpha, \beta) \in \mathcal{L} \cap (\mathcal{I}' \cup \mathcal{K}')} \chi_1(b(\alpha + \beta)) T_b^{[4(b)]} \\ & \left. + \sum_{(\alpha, \beta) \in \mathcal{L} \cap \mathcal{L}'} \chi_1(b(\alpha + \beta)) T_b^{[4(b)]} \right). \end{aligned}$$

## 6. THE GENERAL CASE

Since the case  $c \in \mathbb{F}_{2^e}$  has already been considered in previous sections, throughout this section we assume that  $c \in \mathbb{F}_{2^n} \setminus \mathbb{F}_{2^e}$ . Recall that for any fixed  $b \neq 0$ , the  $c$ -BCT entry is given by,

$${}_c\mathcal{B}_F(1, b) = \frac{1}{q^2} \sum_{\alpha, \beta \in \mathbb{F}_q} \chi_1(b(\alpha + \beta)) S_{\alpha, \beta} S_{c\alpha, c^{-1}\beta}.$$

Let us denote  $T_b = S_{\alpha, \beta} S_{c\alpha, c^{-1}\beta}$ . Recall that  $A = \alpha + \beta$ ,  $B = \beta^{2^{n-k}} + \beta$ ,  $A' = c\alpha + c^{-1}\beta$  and  $B' = (c^{-1}\beta)^{2^{n-k}} + c^{-1}\beta$ . Notice that, when  $c \in \mathbb{F}_{2^n} \setminus \mathbb{F}_{2^e}$  then  $\beta \in \mathbb{F}_{2^e}^*$ , and so,  $\beta c^{-1} \in \mathbb{F}_{2^n} \setminus \mathbb{F}_{2^e}$ , otherwise  $c \in \mathbb{F}_{2^e}$ . Thus  $B = 0 = B'$  if and only if  $\beta = 0$ . Also, observe that the conditions  $A = 0 = A'$  if and only if  $\alpha = 0 = \beta$ . Now we shall consider two cases namely,  $\frac{n}{e}$  is odd and  $\frac{n}{e}$  is even, respectively.

**Case 1:**  $\frac{n}{e}$  is odd.

(1) Let  $A = 0, B = 0$ .

Notice that the cases  $A' = 0, B' \neq 0$ , and  $A' \neq 0, B' = 0$  would not arise, therefore, we shall calculate  $T_b$  in remaining two cases only.

(a) If  $A' = 0, B' = 0$ , then

$$T_b^{[1(a)]} = \chi_1((1 + c^{-1})\beta) q^2.$$

(b) If  $A' \neq 0, B' \neq 0$ , then

$$T_b^{[1(b)]} = \begin{cases} 0 & \text{if } \text{Tr}_e(B'(\gamma')^{-1}) \neq 1, \\ \left(\frac{2}{n/e}\right)^e 2^{\frac{3n+e}{2}} \chi_1((1 + c^{-1})\beta) & \text{if } \text{Tr}_e(B'(\gamma')^{-1}) = 1. \end{cases}$$

(2) Let  $A = 0, B \neq 0$ . In this case  $S_{\alpha,\beta} = 0$  and hence

$$T_b^{[2]} = 0.$$

(3) Let  $A \neq 0, B = 0$ . Again,  $S_{\alpha,\beta} = 0$  and hence

$$T_b^{[3]} = 0.$$

(4) Let  $A \neq 0, B \neq 0$ .

(a) If  $A' = 0, B' = 0$ , then

$$T_b^{[4(a)]} = \begin{cases} 0 & \text{if } \text{Tr}_e(B\gamma^{-1}) \neq 1, \\ \left(\frac{2}{n/e}\right)^e 2^{\frac{3n+e}{2}} \chi_1((1+c^{-1})\beta) & \text{if } \text{Tr}_e(B\gamma^{-1}) = 1. \end{cases}$$

(b) If  $A' = 0, B' \neq 0$ , then  $S_{c\alpha, c^{-1}\beta} = 0$  and hence

$$T_b^{[4(b)]} = 0.$$

(c) If  $A' \neq 0, B' = 0$ , then again  $S_{c\alpha, c^{-1}\beta} = 0$  and hence

$$T_b^{[4(c)]} = 0.$$

(d) If  $A' \neq 0, B' \neq 0$ , then the only relevant case is and

$$T_b^{[4(d)]} = \begin{cases} 2^{n+e} \chi_1((1+c^{-1})\beta) & \text{if } (\alpha, \beta) \in \mathcal{F} \cap \mathcal{F}', \\ 0 & \text{otherwise.} \end{cases}$$

We now summarize the above discussion in the following theorem.

**Theorem 6.1.** *Let  $F(x) = x^{2^k+1}$ ,  $1 \leq k < n$  be a function on  $\mathbb{F}_{2^n}$ ,  $n \geq 2$ . Let  $c \in \mathbb{F}_{2^n} \setminus \mathbb{F}_{2^e}$  and  $n/e$  be odd, where  $e = \gcd(k, n)$ . Then the  $c$ -BCT entry  ${}_c\mathcal{B}_F(1, b)$  of  $F$  at  $(1, b)$  is given by*

$$\begin{aligned} 1 + \frac{2^{\frac{e}{2}}}{2^n} \sum_{(\alpha, \beta) \in (\mathcal{A} \cap \mathcal{F}') \cup (\mathcal{A}' \cap \mathcal{F})} \chi_1(b\alpha + (1+c^{-1}+b)\beta) \\ + \frac{2^e}{2^n} \sum_{(\alpha, \beta) \in \mathcal{F} \cap \mathcal{F}'} \chi_1(b\alpha + (1+c^{-1}+b)\beta). \end{aligned}$$

*Proof.*

$$\begin{aligned} {}_c\mathcal{B}_F(1, b) &= \frac{1}{q^2} \left( \sum_{(\alpha, \beta) \in \mathcal{A} \cap \mathcal{A}'} \chi_1(b(\alpha + \beta)) T_b^{[1(a)]} + \sum_{(\alpha, \beta) \in \mathcal{A} \cap \mathcal{F}'} \chi_1(b(\alpha + \beta)) T_b^{[1(b)]} \right. \\ &\quad \left. + \sum_{(\alpha, \beta) \in \mathcal{F} \cap \mathcal{A}'} \chi_1(b(\alpha + \beta)) T_b^{[4(a)]} + \sum_{(\alpha, \beta) \in \mathcal{F} \cap \mathcal{F}'} \chi_1(b(\alpha + \beta)) T_b^{[4(d)]} \right) \\ &= 1 + \left(\frac{2}{n/e}\right)^e \cdot 2^{\frac{e-n}{2}} \sum_{(\alpha, \beta) \in (\mathcal{A} \cap \mathcal{F}') \cup (\mathcal{A}' \cap \mathcal{F})} \chi_1(b\alpha + (1+c^{-1}+b)\beta) \\ &\quad + 2^{e-n} \sum_{(\alpha, \beta) \in \mathcal{F} \cap \mathcal{F}'} \chi_1(b\alpha + (1+c^{-1}+b)\beta). \end{aligned}$$

□

**Corollary 6.2.** *Let  $F(x) = x^{2^k+1}$ ,  $1 \leq k < n$ , be a function on  $\mathbb{F}_q$ ,  $n \geq 2$ . Let  $c \in \mathbb{F}_{2^n} \setminus \mathbb{F}_{2^e}$  and  $n/e$  be odd, where  $e = \gcd(k, n)$ . With the notations of the previous theorem, the  $c$ -boomerang uniformity of  $F$  satisfies*

$$\beta_{F,c} \leq 1 + \left( \frac{2}{n/e} \right)^e \cdot 2^{\frac{e-n}{2}} |(\mathcal{A} \cap \mathcal{F}') \cup (\mathcal{A}' \cap \mathcal{F})| + 2^{e-n} |\mathcal{F} \cap \mathcal{F}'|.$$

**Case 2:**  $n/e$  is even.

- (1) Let  $A = 0, B = 0$ . Notice that the cases  $A' = 0, B' \neq 0$ , and  $A' \neq 0, B' = 0$  would not arise, therefore, we shall calculate  $T_b$  in remaining two cases only.

- (a) If  $A' = 0, B' = 0$ , then

$$T_b^{[1(a)]} = \chi_1((1 + c^{-1})\beta) q^2.$$

- (b) If  $A' \neq 0, B' \neq 0$ , then

$$T_b^{[1(b)]} = \begin{cases} (-1)^{\frac{m}{e}} 2^{m+n} M' & \text{if } (\alpha, \beta) \in \mathcal{A} \cap (\mathcal{I}' \cup \mathcal{K}'), \\ 0 & \text{if } (\alpha, \beta) \in \mathcal{A} \cap \mathcal{J}', \\ (-1)^{\frac{m}{e}+1} 2^{m+n+e} M' & \text{if } (\alpha, \beta) \in \mathcal{A} \cap \mathcal{L}', \end{cases}$$

where  $M' = \chi_1((1 + c^{-1})\beta) \chi_1(A' x_{A'}^{2^k+1})$ .

- (2) Let  $A = 0, B \neq 0$ . In this case  $S_{\alpha,\beta} = 0$  and hence

$$T_b^{[2]} = 0$$

- (3) Let  $A \neq 0, B = 0$ . Notice that the case  $A' = 0, B' = 0$  would not arise. Now we shall calculate  $T_b$  in the remaining cases.

- (a) If  $A' = 0, B' \neq 0$ , then  $S_{c\alpha, c^{-1}\beta} = 0$  and hence

$$T_b^{[3(a)]} = 0.$$

- (b) If  $A' \neq 0, B' = 0$ , then

$$T_b^{[3(b)]} = \begin{cases} 2^n \chi_1((1 + c^{-1})\beta) & \text{if } (\alpha, \beta) \in \mathcal{G} \cap \mathcal{G}', \\ -2^{n+e} \chi_1((1 + c^{-1})\beta) & \text{if } (\alpha, \beta) \in \mathcal{G} \cap \mathcal{H}', \\ -2^{n+e} \chi_1((1 + c^{-1})\beta) & \text{if } (\alpha, \beta) \in \mathcal{H} \cap \mathcal{G}', \\ 2^{n+2e} \chi_1((1 + c^{-1})\beta) & \text{if } (\alpha, \beta) \in \mathcal{H} \cap \mathcal{H}'. \end{cases}$$

- (c) If  $A' \neq 0, B' \neq 0$ , then

$$T_b^{[3(c)]} = \begin{cases} 2^n M' & \text{if } (\alpha, \beta) \in \mathcal{G} \cap (\mathcal{I}' \cup \mathcal{K}'), \\ 0 & \text{if } (\alpha, \beta) \in (\mathcal{G} \cup \mathcal{H}) \cap \mathcal{J}', \\ -2^{n+e} M' & \text{if } (\alpha, \beta) \in \mathcal{G} \cap \mathcal{L}', \\ -2^{n+e} M' & \text{if } (\alpha, \beta) \in \mathcal{H} \cap (\mathcal{I}' \cup \mathcal{K}'), \\ 2^{n+2e} M' & \text{if } (\alpha, \beta) \in \mathcal{H} \cap \mathcal{L}'. \end{cases}$$

- (4) Let  $A \neq 0, B \neq 0$ .

- (a) If  $A' = 0, B' = 0$ , then

$$T_b^{[4(a)]} = \begin{cases} (-1)^{\frac{m}{e}} 2^{m+n} M'' & \text{if } (\alpha, \beta) \in \mathcal{A}' \cap (\mathcal{I} \cup \mathcal{K}), \\ 0 & \text{if } (\alpha, \beta) \in \mathcal{A}' \cap \mathcal{J}, \\ (-1)^{\frac{m}{e}+1} 2^{m+n+e} M'' & \text{if } (\alpha, \beta) \in \mathcal{A}' \cap \mathcal{L}, \end{cases}$$

where  $M'' = \chi_1((1 + c^{-1})\beta) \chi_1(A x_A^{2^k+1})$ .



(b) If  $A' = 0, B' \neq 0$ , then  $S_{c\alpha, c^{-1}\beta} = 0$  and hence

$$T_b^{[4(b)]} = 0.$$

(c) If  $A' \neq 0, B' = 0$ , then

$$T_b^{[4(c)]} = \begin{cases} 2^n M'' & \text{if } (\alpha, \beta) \in \mathcal{G}' \cap (\mathcal{I} \cup \mathcal{K}), \\ 0 & \text{if } (\alpha, \beta) \in (\mathcal{G}' \cup \mathcal{H}') \cap \mathcal{J}, \\ -2^{n+e} M'' & \text{if } (\alpha, \beta) \in \mathcal{G}' \cap \mathcal{L}, \\ -2^{n+e} M'' & \text{if } (\alpha, \beta) \in \mathcal{H}' \cap (\mathcal{I} \cup \mathcal{K}), \\ 2^{n+2e} M'' & \text{if } (\alpha, \beta) \in \mathcal{H}' \cap \mathcal{L}. \end{cases}$$

(d) If  $A' \neq 0, B' \neq 0$ , then

$$T_b^{[4(d)]} = \begin{cases} 2^n M''' & \text{if } (\alpha, \beta) \in (\mathcal{I} \cup \mathcal{K}) \cap (\mathcal{I}' \cup \mathcal{K}'), \\ 0 & \text{if } (\alpha, \beta) \in (\mathcal{I} \cup \mathcal{K} \cup \mathcal{L}) \cap \mathcal{J}', \\ -2^{n+e} M''' & \text{if } (\alpha, \beta) \in (\mathcal{I} \cup \mathcal{K}) \cap \mathcal{L}', \\ 0 & \text{if } (\alpha, \beta) \in \mathcal{J} \cap (\mathcal{I}' \cup \mathcal{J}' \cup \mathcal{K}' \cup \mathcal{L}'), \\ -2^{n+e} M''' & \text{if } (\alpha, \beta) \in \mathcal{L} \cap (\mathcal{I}' \cup \mathcal{K}'), \\ 2^{n+2e} M''' & \text{if } (\alpha, \beta) \in \mathcal{L} \cap \mathcal{L}', \end{cases}$$

where  $M''' = \chi_1((1 + c^{-1})\beta)\chi_1(Ax_A^{2^k+1} + A'x_{A'}^{2^k+1})$ .

We now summarize the above discussion in the form of following theorem.

**Theorem 6.3.** *Let  $F(x) = x^{2^k+1}$ ,  $1 \leq k < n$  be a function on  $\mathbb{F}_{2^n}$ ,  $n \geq 2$ . Let  $c \in \mathbb{F}_{2^n} \setminus \mathbb{F}_{2^e}$  and  $n/e$  be even, where  $e = \gcd(k, n)$ . With the prior notations, the  $c$ -BCT entry  ${}_c\mathcal{B}_F(1, b)$  of  $F$  at  $(1, b)$  is given by*

$$\begin{aligned} & \frac{1}{q^2} \left( \sum_{(\alpha, \beta) \in \mathcal{A} \cap \mathcal{A}'} \chi_1(b(\alpha + \beta))T_b^{[1(a)]} + \sum_{(\alpha, \beta) \in \mathcal{A} \cap (\mathcal{I}' \cup \mathcal{K}')} \chi_1(b(\alpha + \beta))T_b^{[1(b)]} \right. \\ & + \sum_{(\alpha, \beta) \in \mathcal{A} \cap \mathcal{L}'} \chi_1(b(\alpha + \beta))T_b^{[1(b)]} + \sum_{(\alpha, \beta) \in \mathcal{G} \cap \mathcal{G}'} \chi_1(b(\alpha + \beta))T_b^{[3(b)]} \\ & + \sum_{(\alpha, \beta) \in \mathcal{G} \cap \mathcal{H}'} \chi_1(b(\alpha + \beta))T_b^{[3(b)]} + \sum_{(\alpha, \beta) \in \mathcal{H} \cap \mathcal{G}'} \chi_1(b(\alpha + \beta))T_b^{[3(b)]} \\ & + \sum_{(\alpha, \beta) \in \mathcal{H} \cap \mathcal{H}'} \chi_1(b(\alpha + \beta))T_b^{[3(b)]} + \sum_{(\alpha, \beta) \in \mathcal{G} \cap (\mathcal{I}' \cup \mathcal{K}')} \chi_1(b(\alpha + \beta))T_b^{[3(c)]} \\ & + \sum_{(\alpha, \beta) \in \mathcal{G} \cap \mathcal{L}'} \chi_1(b(\alpha + \beta))T_b^{[3(c)]} + \sum_{(\alpha, \beta) \in \mathcal{H} \cap (\mathcal{I}' \cup \mathcal{K}')} \chi_1(b(\alpha + \beta))T_b^{[3(c)]} \\ & + \sum_{(\alpha, \beta) \in \mathcal{H} \cap \mathcal{L}'} \chi_1(b(\alpha + \beta))T_b^{[3(c)]} + \sum_{(\alpha, \beta) \in \mathcal{A}' \cap (\mathcal{I} \cup \mathcal{K})} \chi_1(b(\alpha + \beta))T_b^{[4(a)]} \\ & + \sum_{(\alpha, \beta) \in \mathcal{A}' \cap \mathcal{L}} \chi_1(b(\alpha + \beta))T_b^{[4(a)]} + \sum_{(\alpha, \beta) \in \mathcal{G}' \cap (\mathcal{I} \cup \mathcal{K})} \chi_1(b(\alpha + \beta))T_b^{[4(c)]} \\ & + \sum_{(\alpha, \beta) \in \mathcal{G}' \cap \mathcal{L}} \chi_1(b(\alpha + \beta))T_b^{[4(c)]} + \sum_{(\alpha, \beta) \in \mathcal{H}' \cap (\mathcal{I} \cup \mathcal{K})} \chi_1(b(\alpha + \beta))T_b^{[4(c)]} \\ & + \sum_{(\alpha, \beta) \in \mathcal{H}' \cap \mathcal{L}} \chi_1(b(\alpha + \beta))T_b^{[4(c)]} + \sum_{(\alpha, \beta) \in (\mathcal{I} \cup \mathcal{K}) \cap (\mathcal{I}' \cup \mathcal{K}')} \chi_1(b(\alpha + \beta))T_b^{[4(d)]} \end{aligned}$$

$$\begin{aligned}
& + \sum_{(\alpha, \beta) \in (I \cup \mathcal{K}) \cap \mathcal{L}'} \chi_1(b(\alpha + \beta)) T_b^{[4(d)]} + \sum_{(\alpha, \beta) \in (I' \cup \mathcal{K}') \cap \mathcal{L}} \chi_1(b(\alpha + \beta)) T_b^{[4(d)]} \\
& + \sum_{(\alpha, \beta) \in \mathcal{L}' \cap \mathcal{L}} \chi_1(b(\alpha + \beta)) T_b^{[4(d)]} \Bigg).
\end{aligned}$$

## 7. DISCUSSION ON EQUIVALENCE

Boura and Canteaut [2] showed that the BCT table is preserved under the affine equivalence but not under the extended affine equivalence (and consequently under the CCZ-equivalence). It is quite natural to ask a similar question in the context of  $c$ -BCT. It is straightforward to see that in the case of even characteristic,  $c$ -BCT and  $c^{-1}$ -BCT entries of an  $(n, n)$ -function  $F : \mathbb{F}_{2^n} \rightarrow \mathbb{F}_{2^n}$  are the same under the transformations  $x \mapsto x + a$  and  $y \mapsto y + a$ , since the  $c$ -boomerang system

$$\begin{cases} F(x) + cF(y) = b \\ F(x + a) + c^{-1}F(y + a) = b \end{cases}$$

becomes

$$\begin{cases} F(x) + c^{-1}F(y) = b \\ F(x + a) + cF(y + a) = b. \end{cases}$$

We consider the binomial  $G(x) = x^{2^k+1} + ux^{2^{n-k}+1} \in \mathbb{F}_{2^n}[x]$ , which is a PP if and only if  $\frac{n}{e}$  is odd and  $u \neq g^{t(2^e-1)}$ , where  $e = \gcd(n, k) = \gcd(n - k, k)$  and  $g$  is the primitive element of  $\mathbb{F}_{2^n}$ . Notice that  $G(x) = (L \circ F)(x)$  where  $L(x) = x^{2^k} + ux$  and  $F(x) = x^{2^{n-k}+1}$ . When  $n = 6, k = 2$  and  $u = g$ , where  $g$  is a root of the primitive polynomial  $y^6 + y^4 + y^3 + y + 1$  over  $\mathbb{F}_2$ , then  $L(x)$  and  $G(x)$  are PP. It is easy to see from the Table 1 in the Appendix A that the  $c$ -BCT is not preserved under the (output applied) affine equivalence. However, if the affine transformation is applied to the input, that is,  $G(x) = (F \circ L)(x)$ , then the  $c$ -BCT spectrum is preserved, as was the case for the  $c$ -differential uniformity.

## ACKNOWLEDGEMENTS

Sartaj Ul Hasan is partially supported by MATRICS grant MTR/2019/000744 from the Science and Engineering Research Board, Government of India.

## REFERENCES

- [1] D. Bartoli, M. Calderini, *On construction and (non)existence of  $c$ -(almost) perfect nonlinear functions*, Finite Fields Appl., 72 (2021), 101835.
- [2] C. Boura, A. Canteaut, *On the boomerang uniformity of cryptographic Sboxes*, IACR Trans. Symmetric Cryptol., vol. 2018, no. 3, 290–310, 2018.
- [3] C. Carlet, *Boolean functions for cryptography and error correcting codes*, In: Y. Crama, P. Hammer (eds.), Boolean Methods and Models, Cambridge Univ. Press, Cambridge, pp. 257–397, 2010.
- [4] C. Cid, T. Huang, T. Peyrin, Y. Sasaki, and L. Song, *Boomerang connectivity table: a new cryptanalysis tool*. In: Nielsen J., Rijmen V. (eds.), Advances in Cryptology-EUROCRYPT 2018, LNCS 10821, Springer, Cham, pp. 683–714, 2018.
- [5] R. S. Coulter, *On the evaluation of a class of Weil sums in characteristic 2*, New Zealand J. of Math., vol. 28, pp. 171–184, 1999.
- [6] T. W. Cusick, P. Stănică, Cryptographic Boolean Functions and Applications (Ed. 2), Academic Press, San Diego, CA, 2017.
- [7] P. Ellingsen, P. Felke, C. Riera, P. Stănică, A. Tkachenko,  *$C$ -differentials, multiplicative uniformity and (almost) perfect  $c$ -nonlinearity*, IEEE Trans. Inform. Theory 66(9) (2020), 5781–5789.

- [8] S. U. Hasan, M. Pal, C. Riera, P. Stănică, *On the  $c$ -differential uniformity of certain maps over finite fields*, Des. Codes Cryptogr. 89(2) (2021), 221–239.
- [9] K. Li, L. Qu, B. Sun, C. Li, *New results about the boomerang uniformity of permutation polynomials*, IEEE Trans. Inform. Theory 65(11) (2019), 7542–7553.
- [10] S. Mesnager, C. Riera, P. Stănică, H. Yan and Z. Zhou, *Investigations on  $c$ -(almost) perfect nonlinear functions*, IEEE Trans. Inform. Theory (2021), <https://doi.org/10.1109/TIT.2021.3081348>.
- [11] K. Nyberg, *Differentially uniform mappings for cryptography*. In: Helleseeth T. (eds.), Advances in Cryptology–EUROCRYPT 1993, LNCS 765, Springer, Berlin, Heidelberg, pp. 55–64, 1994.
- [12] P. Stănică, *Investigations on  $c$ -boomerang uniformity and perfect nonlinearity*, Discrete Appl. Math., Vol. 304, 297–314, 2021.
- [13] P. Stănică, *Low  $c$ -differential and  $c$ -boomerang uniformity of the swapped inverse function*, Discrete Math. 344(10) (2021), 112543.
- [14] P. Stănică, *Using double Weil sums in finding the  $c$ -boomerang connectivity table for monomial functions on finite fields*, Appl. Algebra Eng. Commun. Comput., (2021). <https://doi.org/10.1007/s00200-021-00520-9>
- [15] P. Stănică, A. Geary, *The  $c$ -differential behaviour of the inverse function under the EA-equivalence*, Cryptogr. Commun. 13 (2021), 295–306.
- [16] P. Stănică, C. Riera, A. Tkachenko, *Characters, Weil sums and  $c$ -differential uniformity with an application to the perturbed Gold function*, Cryptogr. Commun. (2021), <https://doi.org/10.1007/s12095-021-00485-z>.
- [17] D. Wagner, *The boomerang attack*, In: L. R. Knudsen (ed.) Fast Software Encryption-FSE 1999. LNCS 1636, Springer, Berlin, Heidelberg, pp. 156–170, 1999.
- [18] Z. Zha, L. Hu, *Some classes of power functions with low  $c$ -differential uniformity over finite fields*, Des. Codes Cryptogr. (2021), <https://doi.org/10.1007/s10623-021-00866-8>.

## APPENDIX A.

Let  $G(x) = x^5 + gx^{17} = (x^4 + gx) \circ x^{17} \in \mathbb{F}_{2^6}[x]$ , where  $g$  is a root of the primitive polynomial  $y^6 + y^4 + y^3 + y + 1$  over  $\mathbb{F}_2$ . The following Table 1 gives the set of the  $c$ -BCT entries for  $x^{17}$  as well as  $G(x) = x^5 + gx^{17}$  for all  $c \in \mathbb{F}_{2^6} \setminus \mathbb{F}_2$ . In view of the discussion in Section 7, it is sufficient to compute the set of  $c$ -BCT entries for either of  $c$  or  $c^{-1}$  as they are going to be exactly the same.

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY JAMMU, JAMMU 181221, INDIA  
*Email address:* sartaj.hasan@iitjammu.ac.in

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY JAMMU, JAMMU 181221, INDIA  
*Email address:* 2018RMA0021@iitjammu.ac.in

APPLIED MATHEMATICS DEPARTMENT, NAVAL POSTGRADUATE SCHOOL, MONTEREY 93943, USA  
*Email address:* pstanica@nps.edu

$c$	Set of $c$ -BCT entries of $x^{17}$	Set of $c$ -BCT entries of $G(x)$
$g$	$\{0,1,2,3,4\}$	$\{0,1,2,3,4,5\}$
$g^2$	$\{0,1,2,3,4\}$	$\{0,1,2,3,4,5\}$
$g^3$	$\{0,1,2,3,5\}$	$\{0,1,2,3,4,5\}$
$g^4$	$\{0,1,2,3,4\}$	$\{0,1,2,3,4,5\}$
$g^5$	$\{0,1,2,3,4\}$	$\{0,1,2,3,4,5,6\}$
$g^6$	$\{0,1,2,3,5\}$	$\{0,1,2,3,4,5,6\}$
$g^7$	$\{0,1,2,3,4,5\}$	$\{0,1,2,3,4,5,6\}$
$g^8$	$\{0,1,2,3,4\}$	$\{0,1,2,3,4,5\}$
$g^9$	$\{0,1,2,3,4\}$	$\{0,1,2,3,4,5,6\}$
$g^{10}$	$\{0,1,2,3,4\}$	$\{0,1,2,3,4,5\}$
$g^{11}$	$\{0,1,2,3\}$	$\{0,1,2,3,4,5\}$
$g^{12}$	$\{0,1,2,3,5\}$	$\{0,1,2,3,4,5\}$
$g^{13}$	$\{0,1,2,3\}$	$\{0,1,2,3,4,5\}$
$g^{14}$	$\{0,1,2,3,4,5\}$	$\{0,1,2,3,4,5,6\}$
$g^{15}$	$\{0,1,2,3,5\}$	$\{0,1,2,3,4,5\}$
$g^{16}$	$\{0,1,2,3,4\}$	$\{0,1,2,3,4,5\}$
$g^{17}$	$\{0,1,2,3,4\}$	$\{0,1,2,3,4,5,6\}$
$g^{18}$	$\{0,1,2,3,4\}$	$\{0,1,2,3,4,5,6\}$
$g^{19}$	$\{0,1,2,3\}$	$\{0,1,2,3,4,5\}$
$g^{20}$	$\{0,1,2,3,4\}$	$\{0,1,2,3,4,5,6\}$
$g^{21}$	$\{0,1,4\}$	$\{0,1,4\}$
$g^{22}$	$\{0,1,2,3\}$	$\{0,1,2,3,4,5\}$
$g^{23}$	$\{0,1,2,3,4\}$	$\{0,1,2,3,4,5\}$
$g^{24}$	$\{0,1,2,3,5\}$	$\{0,1,2,3,4,5,6\}$
$g^{25}$	$\{0,1,2,3\}$	$\{0,1,2,3,4,5\}$
$g^{26}$	$\{0,1,2,3\}$	$\{0,1,2,3,4,5\}$
$g^{27}$	$\{0,1,2,3,4\}$	$\{0,1,2,3,4,5,6\}$
$g^{28}$	$\{0,1,2,3,4,5\}$	$\{0,1,2,3,4,5,6\}$
$g^{29}$	$\{0,1,2,3,4\}$	$\{0,1,2,3,4,5\}$
$g^{30}$	$\{0,1,2,3,5\}$	$\{0,1,2,3,4,5,6\}$
$g^{31}$	$\{0,1,2,3,4\}$	$\{0,1,2,3,4,5\}$

**Table 1.**  $c$ -BCT entries of  $x^{17}$  and  $x^5 + gx^{17}$