

ON OSCILLATORY INTEGRALS ASSOCIATED TO PHASE FUNCTIONS WITH DEGENERATE SINGULAR POINTS

TOSHIO NAGANO AND NAOYA MIYAZAKI

ABSTRACT. In this note, by using the result in one variable, we obtain asymptotic expansions of oscillatory integrals for certain multivariable phase functions with **degenerate** singular points. Moreover by using this result, we have asymptotic expansions of oscillatory integrals with phase function of type A_k , E_6 , E_8 -function germs.

1. INTRODUCTION

We study asymptotic expansions of oscillatory integrals:

$$Os\text{-}\int_{\mathbb{R}^n} e^{i\lambda\phi(x)} a(x) dx := \lim_{\varepsilon \rightarrow +0} \int_{\mathbb{R}^n} e^{i\lambda\phi(x)} a(x) \chi(\varepsilon x) dx,$$

as a positive real parameter $\lambda \rightarrow \infty$, where $\chi \in \mathcal{S}(\mathbb{R}^n)$ with $\chi(0) = 1$ and $0 < \varepsilon < 1$.

As to a method of asymptotic expansions of oscillatory integrals for phase functions with a **non-degenerate** singular point, the stationary phase method is known ([7], [8], [4], [6]).

We are interested in asymptotic expansions of oscillatory integrals for phase functions with a **degenerate** singular point expressed by real polynomials $\sum_{j=1}^n \pm x^{m_j}$, and for amplitude functions belonging to the class $\mathcal{A}_\delta^r(\mathbb{R}^n)$ (Definition 2.5).

In [10], we obtained the following result in one variable, for a phase function $\phi(x) = \pm x^m$ where $m \in \mathbb{N}$ (Theorem 2.6): for any $N > m$, as $\lambda \rightarrow \infty$,

$$Os\text{-}\int_{-\infty}^{\infty} e^{\pm i\lambda x^m} a(x) dx = \sum_{k=0}^{N-m-1} c_k^\pm \frac{a^{(k)}(0)}{k!} \lambda^{-\frac{k+1}{m}} + O\left(\lambda^{-\frac{N-m+1}{m}}\right),$$

where c_k^\pm is given by (2.5).

In this note, by using this result, we obtained the following result in multivariable, for a phase function $\phi(x) = \sum_{j=1}^n \pm_j x^{m_j}$ where $m_j \in \mathbb{N}$ such that $m_1 \geq \dots \geq m_n \geq 2$ and \pm_j stands for “+” or “−” determined by j (Theorem 4.2): for any $N_1 \in \mathbb{N}$ such that $N_1 > m_1$, as $\lambda \rightarrow \infty$,

$$Os\text{-}\int_{\mathbb{R}^n} e^{i\lambda \sum_{j=1}^n \pm_j x^{m_j}} a(x) dx = \sum_{\alpha \in \Omega} c_\alpha \lambda^{-\sum_{j=1}^n \frac{\alpha_j + 1}{m_j}} + O\left(\lambda^{-\frac{N_1 - m_1 + 1}{m_1}}\right),$$

where Ω and c_α are given by (4.3) and (4.4).

Date: September 20, 2020.

2010 Mathematics Subject Classification. Primary 42B20 ; Secondary 41A60, 58K05.

Key words and phrases. oscillatory integral, asymptotic expansion, stationary phase method.

The first author was supported by Tokyo University of Science Graduate School doctoral program scholarship and an exemption of the cost of equipment from 2016 to 2018 and would like to thank to Professor Minoru Ito for giving me an opportunity of studies and preparing the environment.

Moreover by using our result, in the cases of the phase function ϕ is type A_k , E_6 , E_8 -function germ where $k \in \mathbb{N}$ ([1], [2]):

$$A_k : \pm x_1^{k+1} + x_2^2 + x_3^2, \quad E_6 : \pm x_1^4 + x_2^3 + x_3^2, \quad E_8 : x_1^5 + x_2^3 + x_3^2,$$

we can obtain asymptotic expansions of oscillatory integrals (Corollary 4.4): for any $N \in \mathbb{N}$ such that $N > k + 1$, $N > 4$ and $N > 5$ respectively, as $\lambda \rightarrow \infty$,

$$\begin{aligned} Os\text{-}\int_{\mathbb{R}^3} e^{i\lambda(\pm x_1^{k+1} + x_2^2 + x_3^2)} a(x) dx &= \sum_{\alpha \in \Omega_1} c_\alpha \lambda^{-\frac{\alpha_1+1}{k+1} - \frac{\alpha_2+1}{2} - \frac{\alpha_3+1}{2}} + O\left(\lambda^{-\frac{N-k}{k+1}}\right), \\ Os\text{-}\int_{\mathbb{R}^3} e^{i\lambda(\pm x_1^4 + x_2^3 + x_3^2)} a(x) dx &= \sum_{\alpha \in \Omega_2} c_\alpha \lambda^{-\frac{\alpha_1+1}{4} - \frac{\alpha_2+1}{3} - \frac{\alpha_3+1}{2}} + O\left(\lambda^{-\frac{N-4}{4}}\right), \\ Os\text{-}\int_{\mathbb{R}^3} e^{i\lambda(x_1^5 + x_2^3 + x_3^2)} a(x) dx &= \sum_{\alpha \in \Omega_3} c_\alpha \lambda^{-\frac{\alpha_1+1}{5} - \frac{\alpha_2+1}{3} - \frac{\alpha_3+1}{2}} + O\left(\lambda^{-\frac{N-5}{5}}\right). \end{aligned}$$

As to a study related to this result, there is a [3].

To the end of §1, we remark notation which will be used in this letter:

$\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{Z}_{\geq 0}^n$ is a multi-index with a length $|\alpha| = \alpha_1 + \dots + \alpha_n$, and then, we use $x^\alpha = x_1^{\alpha_1} \dots x_n^{\alpha_n}$, $\alpha! = \alpha_1! \dots \alpha_n!$, $\partial_x^\alpha = \partial_{x_1}^{\alpha_1} \dots \partial_{x_n}^{\alpha_n}$, $D_x^\alpha = D_{x_1}^{\alpha_1} \dots D_{x_n}^{\alpha_n}$ and $\tilde{D}_x^\alpha = \tilde{D}_{x_1}^{\alpha_1} \dots \tilde{D}_{x_n}^{\alpha_n}$, where $\partial_{x_j} = \frac{\partial}{\partial x_j}$, $D_{x_j} = i^{-1} \partial_{x_j}$ and $\tilde{D}_{x_j} = \lambda^{-1} D_{x_j}$ for $x = (x_1, \dots, x_n)$ and $\lambda > 0$.

$C^\infty(\mathbb{R}^n)$ is the set of complex-valued functions of class C^∞ on \mathbb{R}^n . $C_0^\infty(\mathbb{R}^n)$ is the set of all $f \in C^\infty(\mathbb{R}^n)$ with compact support. $\mathcal{S}(\mathbb{R}^n)$ is the Schwartz space of rapidly decreasing functions of class C^∞ on \mathbb{R}^n , that is, the Fréchet space of all $f \in C^\infty(\mathbb{R}^n)$ such that $\max_{k+|\alpha| \leq m} \sup_{x \in \mathbb{R}^n} \langle x \rangle^k |\partial_x^\alpha f(x)| < \infty$ for any $m \in \mathbb{Z}_{\geq 0}$, where $\langle x \rangle := (1 + |x|^2)^{1/2}$.

$[x]$ is the Gauss' symbol for $x \in \mathbb{R}$, that is, $[x] \in \mathbb{Z}$ such that $x - 1 < [x] \leq x$.

O means the Landau's symbol, that is, $f(x) = O(g(x))$ ($x \rightarrow a$) if $|f(x)/g(x)|$ is bounded as $x \rightarrow a$ for functions f and g , where $a \in \mathbb{R} \cup \{\pm\infty\}$.

δ_{ij} is the Kronecker's delta, that is, $\delta_{ii} = 1$, and $\delta_{ij} = 0$ if $i \neq j$.

$\tau^+ := \max\{\tau, 0\}$ for $\tau \in \mathbb{R}$.

2. PRELIMINARIES

In this section, we shall recall asymptotic expansions of oscillatory integrals and the result in one variable. First we recall the oscillatory integrals.

Definition 2.1. Let $\lambda > 0$ and let ϕ be a real-valued function of class C^∞ on \mathbb{R}^n and $a \in C^\infty(\mathbb{R}^n)$. If there exists the following limit of improper integral:

$$\tilde{I}_\phi[a](\lambda) := Os\text{-}\int_{\mathbb{R}^n} e^{i\lambda\phi(x)} a(x) dx := \lim_{\varepsilon \rightarrow +0} \int_{\mathbb{R}^n} e^{i\lambda\phi(x)} a(x) \chi(\varepsilon x) dx$$

independent of $\chi \in \mathcal{S}(\mathbb{R}^n)$ with $\chi(0) = 1$ and $0 < \varepsilon < 1$, then we call $\tilde{I}_\phi[a](\lambda)$ an oscillatory integral where we call ϕ (resp. a) a phase function (resp. an amplitude function).

If we suppose certain suitable conditions for ϕ and a , then we can show $\tilde{I}_\phi[a](\lambda)$ exists independent of χ and ε (Theorem 3.6 (ii)). The fundamental properties are the following (cf. [9] p.47.):

Proposition 2.2. Let $\chi \in \mathcal{S}(\mathbb{R}^n)$ with $\chi(0) = 1$ and $\langle x \rangle := (1 + |x|^2)^{1/2}$. Then

- (i) $\chi(\varepsilon x) \rightarrow 1$ uniformly on any compact set in \mathbb{R}^n as $\varepsilon \rightarrow +0$.
- (ii) For each multi-index $\alpha \in \mathbb{Z}_{\geq 0}^n$, there exists a positive constant C_α independent of $0 < \varepsilon < 1$ such that for any $x \in \mathbb{R}^n$

$$|\partial_x^\alpha(\chi(\varepsilon x))| \leq C_\alpha \langle x \rangle^{-|\alpha|}.$$

- (iii) For any multi-index $\alpha \in \mathbb{Z}_{\geq 0}^n$ with $\alpha \neq 0$, $\partial_x^\alpha \chi(\varepsilon x) \rightarrow 0$ uniformly in \mathbb{R}^n as $\varepsilon \rightarrow +0$.

Proof. (i) Let K be an any compact set in \mathbb{R}^n . Since $\chi \in \mathcal{S}(\mathbb{R}^n)$, put $C_K := n \sup_{x \in K} |x| \cdot \sup_{x \in \mathbb{R}^n} |\partial_x \chi(x)| > 0$, then by Taylor expansion of $\chi(x)$ at $x = 0$, for each $\varepsilon' > 0$, there exists $\delta := \varepsilon'/C_K > 0$, for any $\varepsilon > 0$ such that $0 < \varepsilon < \delta$ and for any $x \in K$,

$$|\chi(\varepsilon x) - 1| \leq \sum_{k=1}^n \varepsilon |x| \int_0^1 |\partial_{x_k} \chi(\theta \varepsilon x)| d\theta \leq C_K \varepsilon < C_K \delta = C_K \cdot \frac{\varepsilon'}{C_K} = \varepsilon'.$$

Therefore $\chi(\varepsilon x) \rightarrow 1$ uniformly on any compact set K in \mathbb{R}^n as $\varepsilon \rightarrow +0$.

- (ii) For any multi-index $\alpha \in \mathbb{Z}_{\geq 0}^n$,

$$(2.1) \quad \partial_x^\alpha(\chi(\varepsilon x)) = (\partial_x^\alpha \chi)(y) \Big|_{y=\varepsilon x} \cdot \varepsilon^{|\alpha|}$$

$$(2.2) \quad = \{ |y|^{|\alpha|} (\partial_x^\alpha \chi)(y) \} \Big|_{y=\varepsilon x} \cdot |x|^{-|\alpha|}.$$

Here since $\chi \in \mathcal{S}(\mathbb{R}^n)$, put $C'_{\alpha,k} := \sup_{x \in \mathbb{R}^n} \langle x \rangle^k |\partial_x^\alpha \chi(x)| > 0$ for any $k \in \mathbb{Z}_{\geq 0}$.

If $|x| \leq 1$, since $\langle x \rangle \leq \sqrt{2}$, then by (2.1), there exists a positive constant $C_\alpha := 2^{|\alpha|/2} C'_{\alpha,0}$ independent of $0 < \varepsilon < 1$ such that

$$|\partial_x^\alpha(\chi(\varepsilon x))| \leq C'_{\alpha,0} \varepsilon^{|\alpha|} \leq C'_{\alpha,0} \{ \sqrt{2} \langle x \rangle^{-1} \}^{|\alpha|} = C_\alpha \langle x \rangle^{-|\alpha|}.$$

If $|x| > 1$, since $\langle x \rangle \leq \sqrt{2}|x|$, then by (2.2), there exists a positive constant $C_\alpha := 2^{|\alpha|/2} C'_{\alpha,|\alpha|}$ independent of $0 < \varepsilon < 1$ such that

$$|\partial_x^\alpha(\chi(\varepsilon x))| \leq C'_{\alpha,|\alpha|} |x|^{-|\alpha|} \leq C'_{\alpha,|\alpha|} \{ \sqrt{2} \langle x \rangle^{-1} \}^{|\alpha|} = C_\alpha \langle x \rangle^{-|\alpha|}.$$

- (iii) If $\alpha \neq 0$, since $|\alpha| \geq 1$, by (2.1), then for each $\varepsilon' > 0$, there exists $\delta := (\varepsilon'/C'_{\alpha,0})^{1/|\alpha|} > 0$, for any $\varepsilon > 0$ such that $\varepsilon < \delta$ and for any $x \in \mathbb{R}^n$,

$$|\partial_x^\alpha(\chi(\varepsilon x)) - 0| \leq C'_{\alpha,0} \varepsilon^{|\alpha|} < C'_{\alpha,0} \delta^{|\alpha|} = C'_{\alpha,0} (\varepsilon'/C'_{\alpha,0}) = \varepsilon'.$$

Therefore $\partial_x^\alpha \chi(\varepsilon x) \rightarrow 0$ uniformly in \mathbb{R}^n as $\varepsilon \rightarrow +0$. □

In the stationary phase method, the typical case is the following:

Theorem 2.3. *Let $a \in \mathcal{S}(\mathbb{R}^n)$. Then for any $N \in \mathbb{N}$, as $\lambda \rightarrow \infty$,*

$$\begin{aligned} & \int_{\mathbb{R}^n} e^{i\lambda(\sum_{j=1}^p x_j^2 - \sum_{j=p+1}^n x_j^2)} a(x) dx \\ &= \pi^{\frac{n}{2}} e^{i\frac{\pi}{4}\{p-(n-p)\}} \sum_{|\beta|=0}^{N-1} \frac{(-1)^{\sum_{j=p+1}^n \beta_j} i^{|\beta|} \partial_x^{2\beta} a(0)}{4^{|\beta|} \beta!} \lambda^{-k-\frac{n}{2}} + O(\lambda^{-N-\frac{n}{2}}). \end{aligned}$$

First we assume as the phase function the following:

Definition 2.4. Let $m_j \in \mathbb{N}$ such that $\mu_n := \min_{j=1,\dots,n} m_j \geq 2$. Then we define the phase function ϕ_n as follows: for any $x = (x_1, \dots, x_n) \in \mathbb{R}^n$,

$$(2.3) \quad \phi_n(x) := \sum_{j=1}^n \pm_j x_j^{m_j}.$$

Next according to the phase function, we define the class of amplitude functions as follows (cf. [9] p.46.):

Definition 2.5. Assume that $\mu_n = \min_{j=1,\dots,n} m_j$ as above. Let $\tau \in \mathbb{R}$ and $-1 \leq \delta < \mu_n - 1$. Then the class of amplitude functions $\mathcal{A}_\delta^\tau(\mathbb{R}^n)$ is the set of all $a \in C^\infty(\mathbb{R}^n)$ such that for $l \in \mathbb{Z}_{\geq 0}$,

$$(2.4) \quad |a|_l^{(\tau)} := \max_{k \leq l} \sup_{x \in \mathbb{R}^n} |x|^{-\tau - \delta|\alpha|} |\partial_x^\alpha a(x)| < \infty.$$

In [10], we obtained the following result in one variable.

Theorem 2.6. Assume that $\lambda > 0$, $m \in \mathbb{N}$ and $a \in \mathcal{A}_\delta^\tau(\mathbb{R})$. Then for any $N \in \mathbb{N}$ such that $N > m$,

$$Os- \int_{-\infty}^{\infty} e^{\pm i\lambda x^m} a(x) dx = \sum_{k=0}^{N-m-1} c_k^\pm \frac{a^{(k)}(0)}{k!} \lambda^{-\frac{k+1}{m}} + R_N^\pm(\lambda)$$

and

$$\begin{aligned} R_N^\pm(\lambda) &:= \sum_{k=N-m}^{N-1} c_k^\pm \frac{a^{(k)}(0)}{k!} \lambda^{-\frac{k+1}{m}} + \frac{1}{N!} Os- \int_{-\infty}^{\infty} e^{\pm i\lambda x^m} x^N a^{(N)}(\theta x) dx \\ &= O\left(\lambda^{-\frac{N-m+1}{m}}\right) \quad (\lambda \rightarrow \infty), \end{aligned}$$

where

$$(2.5) \quad c_k^\pm := m^{-1} \left\{ e^{\pm i\frac{\pi}{2}\frac{k+1}{m}} + (-1)^k e^{\pm(-1)^m i\frac{\pi}{2}\frac{k+1}{m}} \right\} \Gamma\left(\frac{k+1}{m}\right),$$

$0 < \theta < 1$ and double signs \pm are in the same order.

Proof. We obtained this theorem in [10] (Theorem 5.2 (ii)). However, we recall the proof for the readers.

First if $p > q > 0$, then applying Cauchy's integral theorem to a homeomorphic function $e^{iz^p} z^{q-1}$ on the domain with the anticlockwise oriented boundary $\sum_{j=1}^4 C_j$ defined by $C_1 := \{z = r \in \mathbb{C} | 0 < \varepsilon \leq r \leq R\}$, $C_2 := \{z = Re^{i\theta} \in \mathbb{C} | 0 \leq \theta \leq \pi/2p\}$, $C_3 := \{z = -se^{i(\pi/2p)} \in \mathbb{C} | -R \leq s \leq -\varepsilon\}$ and $C_4 := \{z = \varepsilon e^{-i\tau} \in \mathbb{C} | -\pi/2p \leq \tau \leq 0\}$, and using $\varepsilon \rightarrow +0$ and $R \rightarrow \infty$ give

$$(2.6) \quad \int_0^\infty e^{\pm ix^p} x^{q-1} dx = p^{-1} e^{\pm i\frac{\pi}{2}\frac{q}{p}} \Gamma\left(\frac{q}{p}\right).$$

Secondly if $p > 0$ and $q > 0$, then dividing $\int_0^\infty e^{\pm i\lambda x^p} x^{q-1} a(x) \chi(\varepsilon x) dx$ where $\chi \in \mathcal{S}(\mathbb{R})$ with $\chi(0) = 1$ and $0 < \varepsilon < 1$ by cutoff function φ , repeating integration by parts by $L := \frac{1}{px^{p-1}} \frac{1}{i\lambda} \frac{d}{dx}$ on the support of $1 - \varphi$, making the order of integrand descend to be integrable in the sense of Lebesgue by $L^* := -\frac{1}{i\lambda} \frac{d}{dx} \frac{1}{px^{p-1}}$, and applying Lebsgue's convergence theorem with Proposition 2.2 (i) and (iii) give the existence

of oscillatory integrals $\tilde{I}_{p,q}^\pm[a](\lambda) := Os-\int_0^\infty e^{\pm i\lambda x^p} x^{q-1} a(x) dx$. In particular, if $\lambda = 1$ and $a \equiv 1$, then by (2.6),

$$(2.7) \quad \tilde{I}_{p,q}^\pm := Os-\int_0^\infty e^{\pm i x^p} x^{q-1} dx = p^{-1} e^{\pm i \frac{\pi}{2} \frac{q}{p}} \Gamma\left(\frac{q}{p}\right),$$

and if $q > p$, then

$$(2.8) \quad |\tilde{I}_{p,q}^\pm[a](\lambda)| = O\left(\lambda^{-\frac{q-p}{p}}\right) \quad (\lambda \rightarrow \infty).$$

Thirdly dividing $\tilde{I}_{p,1}^\pm[a](\lambda) := Os-\int_0^\infty e^{\pm i\lambda x^p} a(x) dx$ into the principal part and the remainder term by Taylor expansion of $a(x)$ at $x = 0$. And then applying (2.7) after using change of variable $y = \lambda^{1/p} x$ in the former and applying (2.8) in the latter give for any $N \in \mathbb{N}$ such that $N \geq p + 1$,

$$(2.9) \quad Os-\int_0^\infty e^{\pm i\lambda x^p} a(x) dx = \sum_{k=0}^{N-[p]-1} \tilde{I}_{p,k+1}^\pm \frac{a^{(k)}(0)}{k!} \lambda^{-\frac{k+1}{p}} + \tilde{R}_N^\pm(\lambda),$$

where

$$(2.10) \quad \begin{aligned} \tilde{R}_N^\pm(\lambda) &:= \sum_{k=N-[p]}^{N-1} \tilde{I}_{p,k+1}^\pm \frac{a^{(k)}(0)}{k!} \lambda^{-\frac{k+1}{p}} + \frac{1}{N!} Os-\int_0^\infty e^{\pm i\lambda x^p} x^N a^{(N)}(\theta x) dx \\ &= O\left(\lambda^{-\frac{N-m+1}{m}}\right) \quad (\lambda \rightarrow \infty). \end{aligned}$$

Finally set $p = m \in \mathbb{N}$ on (2.9) and (2.10), and if $x < 0$, then changing a variable $x = -y$ completes the proof. \square

3. EXISTENCE AND ESTIMATES OF OSCILLATORY INTEGRALS

In this section, we shall show existence and estimates of oscillatory integrals used in the next section. First as to the phase function, we have the following:

Proposition 3.1. *Assume that ϕ_n is defined by (2.3). Then there exists a positive constant C such that for any $|x_j| \geq 1$ for $j = 1, \dots, n$,*

$$C|x|^{\mu_n-1} \leq |\nabla \phi_n(x)|.$$

Proof. By Hölder's inequality: $\sum_{j=1}^n x_j^2 \cdot 1 < (\sum_{j=1}^n (x_j^2)^{\mu_n-1})^{1/(\mu_n-1)} (\sum_{j=1}^n 1^q)^{1/q}$ where $q > 1$ and $\{1/(\mu_n-1)\} + (1/q) = 1$ for $\mu_n > 2$. \square

Proposition 3.2. *Assume that $\lambda > 0$ and ϕ_n is defined by (2.3). Let*

$$(3.1) \quad L_n := \sum_{j=1}^n b_j(x) \tilde{D}_{x_j}, \quad b_j(x) := \frac{\partial_{x_j} \phi_n(x)}{|\nabla \phi_n(x)|^2},$$

Then $L_n e^{i\lambda \phi_n(x)} = e^{i\lambda \phi_n(x)}$ and $b_j \in C^\infty(\mathbb{R}^n \setminus \{0\})$, and then for each multi-index $\alpha \in \mathbb{Z}_{\geq 0}^n$, there exists a positive constant $C_{j,\alpha}$ such that for any $|x_k| \geq 1$ for $k = 1, \dots, n$,

$$(3.2) \quad |\partial_x^\alpha b_j(x)| \leq C_{j,\alpha} |x|^{-(\mu_n-1)}.$$

Proof. By induction on $|\alpha|$. If $|\alpha| = 0$, by Proposition 3.1, then (3.2) holds. If (3.2) holds for $|\alpha| - 1$, then applying ∂_x^α to $|\nabla\phi_n(x)|^2 b_j(x) = \partial_{x_j}\phi_n(x)$ and using Leibniz's formula give

$$|\nabla\phi_n(x)|^2 \partial_x^\alpha b_j(x) = - \sum_{0 < \beta \leq \alpha} \binom{\alpha}{\beta} \partial_x^\beta |\nabla\phi_n(x)|^2 \partial_x^{\alpha-\beta} b_j(x) + \partial_x^\alpha \partial_{x_j} \phi_n(x).$$

Moreover using $|\partial_x^\beta |\nabla\phi_n(x)|^2| \leq C_\beta |\nabla\phi_n(x)|^2$, the hypothesis, $|\partial_x^\alpha \partial_{x_j} \phi_n(x)| \leq C_\alpha |\nabla\phi_n(x)|$ and Proposition 3.1 show (3.2) for $|\alpha|$. \square

By Proposition 3.2, we obtain the following:

Proposition 3.3. *Let*

$$(3.3) \quad L_n^* := - \sum_{j=1}^n \tilde{D}_{x_j} b_j(x)$$

be a formal adjoint operator of L_n , and denote

$$(3.4) \quad L_n^* = \sum_{j=1}^n d_j(x) \tilde{D}_{x_j} + d_0(x), \quad d_j(x) := -b_j(x), \quad d_0(x) := - \sum_{j=1}^n \tilde{D}_{x_j} (b_j(x)).$$

Then $d_j \in C^\infty(\mathbb{R}^n \setminus \{0\})$ for $j = 0, \dots, n$, and for each multi-index $\alpha \in \mathbb{Z}_{\geq 0}^n$, there exist positive constants $C_{j,\alpha}$ such that for any $|x_k| \geq 1$ for $k = 1, \dots, n$,

$$(3.5) \quad |\partial_x^\alpha d_j(x)| \leq C_{j,\alpha} |x|^{-(\mu_n-1)} \quad \text{for } j = 1, \dots, n,$$

$$(3.6) \quad |\partial_x^\alpha d_0(x)| \leq C_{0,\alpha} \lambda^{-1} |x|^{-(\mu_n-1)}.$$

Proof. The estimates (3.2) and (3.4) above give (3.5) and (3.6). \square

By Proposition 3.3, we obtain the following:

Proposition 3.4. *For any $l \in \mathbb{Z}_{\geq 0}$, denote*

$$(3.7) \quad L_n^{*l} f = \sum_{|\alpha| \leq l} d_{l,\alpha}(x) \tilde{D}_x^\alpha f$$

*for $f \in C^\infty(\mathbb{R}^n \setminus \{0\})$, where L_n^{*0} is an identity operator. Then $d_{l,\alpha} \in C^\infty(\mathbb{R}^n \setminus \{0\})$ and for each multi-index $\beta \in \mathbb{Z}_{\geq 0}^n$, there exists a positive constant $C_{l,\beta}$ such that for any $|x_k| \geq 1$ for $k = 1, \dots, n$,*

$$(3.8) \quad |\partial_x^\beta d_{l,\alpha}(x)| \leq C_{l,\beta} \lambda^{-(l-|\alpha|)} |x|^{-(\mu_n-1)l}.$$

Proof. By induction on $l \in \mathbb{Z}_{\geq 0}$. If $l = 0$, since $d_{0,0} \equiv 1$, then (3.8) holds. If (3.8) holds for $l-1$ with $l \geq 1$, then for any $l \in \mathbb{N}$ and for any multi-index $\alpha \in \mathbb{Z}_{\geq 0}^n$ with $|\alpha| \leq l$, for any $f \in C^\infty(\mathbb{R}^n \setminus \{0\})$,

$$L_n^{*l} f = \sum_{|\alpha| \leq l-1} \left\{ \left(\sum_{j=1}^n d_j \tilde{D}_{x_j} d_{l-1,\alpha} + d_0 d_{l-1,\alpha} \right) \tilde{D}_x^\alpha f + \sum_{j=1}^n d_j d_{l-1,\alpha} \tilde{D}_{x_j} \tilde{D}_x^\alpha f \right\}$$

Applying ∂_x^β to the coefficients and using Leibniz's formula gives

$$\begin{aligned} |\partial_x^\beta d_{l,\alpha}(x)| &\leq \sum_{|\alpha| \leq l-1} \sum_{\beta' \leq \beta} \binom{\beta}{\beta'} \\ &\times \left\{ \left| \partial_x^{\beta'} \left(\sum_{j=1}^n d_j \tilde{D}_{x_j} d_{l-1,\alpha} + d_0 d_{l-1,\alpha} \right) \right| + \left| \partial_x^{\beta'} \left(\sum_{j=1}^n d_j d_{l-1,\alpha} \lambda^{-1} \right) \right| \right\}. \end{aligned}$$

Moreover using (3.5), (3.6) and the induction hypothesis shows (3.8) for l . \square

We also have

Proposition 3.5. *Assume that $\lambda > 0$. For any $s = 1, \dots, n$, let $\phi_s(x') := \sum_{j=1}^s \pm_j x_j^{m_j}$, $L_s := \sum_{j=1}^s b_j(x') \tilde{D}_{x_j}$ and $L_s^* := -\sum_{j=1}^s \tilde{D}_{x_j} b_j(x')$ be defined by (2.3), (3.1) and (3.3) for $x = (x', x'') \in \mathbb{R}^s \times \mathbb{R}^{n-s}$ respectively. And let $a \in \mathcal{A}_\delta^\tau(\mathbb{R}^n)$, $\varphi_k \in C_0^\infty(\mathbb{R})$ be a cutoff function such that $\varphi_k \equiv 1$ on $|x_k| \leq 1$ and $\varphi_k \equiv 0$ on $|x_k| \geq r > 1$, and $\psi_k := 1 - \varphi_k$ for $k = 1, \dots, n$. And for any $s = 0, \dots, n$, let*

$$(3.9) \quad a_s(x) := a(x) \prod_{k=s+1}^n \varphi_k(x_k) \prod_{k=1}^s \psi_k(x_k),$$

where $\prod_{k=n+1}^n \varphi_k := 1$ and $\prod_{k=1}^0 \psi_k := 1$. And let $\chi \in \mathcal{S}(\mathbb{R}^n)$ with $\chi(0) = 1$ and $0 < \varepsilon < 1$. Then for any $s = 1, \dots, n$, the following hold:

(i) For any $j = 1, \dots, s$, for any $l \in \mathbb{Z}_{\geq 0}$ and for any $h = 0, 1$, let

$$f_{j,l,h}(x) := (b_j(x') \tilde{D}_{x_j})^{1-h} (e^{i\lambda\phi_s(x')}) (-\tilde{D}_{x_j})^h (b_j(x')^h L_s^{*l}(a_s(x)\chi(\varepsilon x))),$$

Then for any $0 < \varepsilon < 1$ and for any $l \in \mathbb{Z}_{\geq 0}$, the improper integrals

$$\int_{\mathbb{R}^n} e^{i\lambda\phi_n(x)} L_s^{*l}(a_s(x)\chi(\varepsilon x)) dx \text{ and } \int_{\mathbb{R}^n} e^{i\lambda(\phi_n(x) - \phi_s(x'))} f_{j,l,h}(x) dx$$

are absolutely convergent.

(ii) For any $j = 1, \dots, s$, for any $(x_1, \dots, \hat{x}_j, \dots, x_n) \in \mathbb{R}^{n-1}$, for any $l \in \mathbb{N}$ and for any $0 < \varepsilon < 1$, as $|x_j| \rightarrow \infty$,

$$|b_j(x')(i\lambda)^{-1} e^{i\lambda\phi_s(x')} L_s^{*(l-1)}(a_s(x)\chi(\varepsilon x))| \rightarrow 0.$$

(iii) For any $l \in \mathbb{N}$ and for any $0 < \varepsilon < 1$,

$$\int_{\mathbb{R}^n} e^{i\lambda\phi_n(x)} a_s(x)\chi(\varepsilon x) dx = \int_{\mathbb{R}^n} e^{i\lambda\phi_n(x)} L_s^{*l}(a_s(x)\chi(\varepsilon x)) dx.$$

(iv) For each $l \in \mathbb{Z}_{\geq 0}$, there exists a positive constant C_l independent of $0 < \varepsilon < 1$ such that for any $x \in \mathbb{R}^n$,

$$|L_s^{*l}(a_s(x)\chi(\varepsilon x))| \leq C_l \lambda^{-l} \prod_{k=s+1}^n \|\varphi_k\| \|a\|_l^{(\tau)} |\psi_{1,s}|_r^{(l)} \langle x \rangle^t,$$

where $t := \tau^+ - (\mu_n - 1 - \delta^+)l$, $\|\varphi_k\| := \sup_{x_k \in \mathbb{R}} |\varphi_k(x_k)|$ and

$$(3.10) \quad |\psi_{1,s}|_r^{(l)} := \max_{|\gamma| \leq l} \sup_{\substack{|x_k| < r, \\ k=1, \dots, n}} \langle x \rangle^{|\gamma|} \prod_{k=1}^s |\partial_{x_k}^{\gamma_k} \psi_k(x_k)| + 1.$$

Proof. First we show (i). A function $\chi \in \mathcal{S}(\mathbb{R}^n)$ gives $L_s^{*l}(a_s(x)\chi(\varepsilon x)) = O(|x|^t)$ ($|x| \rightarrow \infty$) when $t < -n$ and $e^{i\lambda(\phi_n(x) - \phi_s(x'))} f_{j,l,h}(x) = O(|x|^t)$ ($|x| \rightarrow \infty$) when $t < -n$. This shows (i). Secondly we show (ii). A function $\chi \in \mathcal{S}(\mathbb{R}^n)$ also gives $b_j(x')(i\lambda)^{-1} e^{i\lambda\phi_s(x')} L_s^{*(l-1)}(a_s(x)\chi(\varepsilon x)) = O(|x|^t)$ ($|x| \rightarrow \infty$) when $t < 0$. This shows (ii). Next we show (iii). By induction on $l \in \mathbb{N}$. Applying integration

by part with (i), (ii) and Fubini's theorem show (iii). Finally we show (iv). If $x \in \text{supp} L_s^{*l}(a_s(x)\chi(\varepsilon x))$, since

$$\text{supp} L_s^{*l}(a_s(x)\chi(\varepsilon x)) \subset \text{supp} a_s \subset \prod_{k=1}^s (\mathbb{R} \setminus (-1, 1)) \times \prod_{k=s+1}^n (-r, r),$$

then $|x_k| \geq 1$ for $k = 1, \dots, s$ and $|x_k| \leq r$ for $k = s+1, \dots, n$. And then since $|x'| \geq 1$ and $|x''| \leq (n-s)^{1/2}r$, then $|x'|^{-(\mu_s-1)l} \leq C_{n,l} \langle x \rangle^{-(\mu_n-1)l}$ where $C_{n,l} := \max_{s=0, \dots, n} \{2 + 2(n-s)r^2\}^{(\mu_s-1)l/2}$. Hence (3.8), (2.4), (3.10) and Proposition 2.2 (ii) give

$$\begin{aligned} |L_s^{*l}(a_s(x)\chi(\varepsilon x))| &= \left| \prod_{k=s+1}^n \varphi_k(x_k) \sum_{|\alpha| \leq l} d_{l,\alpha}(x') \tilde{D}_{x'}^\alpha \left(a(x) \prod_{k=1}^s \psi_k(x_k) \chi(\varepsilon x) \right) \right| \\ &\leq \prod_{k=s+1}^n |\varphi_k(x_k)| \sum_{|\alpha| \leq l} |d_{l,\alpha}(x')| \lambda^{-|\alpha|} \\ &\quad \times \sum_{\beta+\gamma+\zeta=\alpha} \frac{\alpha!}{\beta! \gamma! \zeta!} |\partial_{x'}^\beta a(x)| \left| \partial_{x'}^\gamma \prod_{k=1}^s \psi_k(x_k) \right| |\partial_{x'}^\zeta (\chi(\varepsilon x))| \\ &\leq \prod_{k=s+1}^n \|\varphi_k\| \sum_{|\alpha| \leq l} C_{l,0} \lambda^{-(l-|\alpha|)} |x'|^{-(\mu_s-1)l} \lambda^{-|\alpha|} \\ &\quad \times \sum_{\beta+\gamma+\zeta=\alpha} \frac{\alpha!}{\beta! \gamma! \zeta!} |a|_l^{(\tau)} \langle x \rangle^{\tau+\delta|\beta|} \langle x \rangle^{-|\gamma|} \langle x \rangle^{|\gamma|} \prod_{k=1}^s |\partial_{x_k}^{\gamma_k} \psi_k(x_k)| C_\zeta \langle x \rangle^{\delta|\zeta|} \\ &\leq \lambda^{-l} \prod_{k=s+1}^n \|\varphi_k\| \sum_{|\alpha| \leq l} C_{l,0} C_{n,l} \langle x \rangle^{-(\mu_n-1)l} \\ &\quad \times \sum_{\beta+\gamma+\zeta=\alpha} \frac{\alpha!}{\beta! \gamma! \zeta!} |a|_l^{(\tau)} \langle x \rangle^{\tau+\delta|\alpha|} |\psi_{1,s}|_r^{(l)} C_\zeta. \end{aligned}$$

This shows (iv). \square

By Proposition 3.5, we obtain the following theorem:

Theorem 3.6. *Assume that $\lambda > 0$ and $s = 0, \dots, n$. Let ϕ_n , a_s and L_s^* be defined by (2.3), (3.9) and (3.3) respectively. Then the following hold:*

(i) *There exists the following oscillatory integral, and the following holds:*

$$\tilde{I}_{\phi_n}[a_0](\lambda) := Os- \int_{\mathbb{R}^n} e^{i\lambda\phi_n(x)} a_0(x) dx = \int_{\mathbb{R}^n} e^{i\lambda\phi_n(x)} a_0(x) dx.$$

(ii) *If $s \neq 0$, then there exists the following oscillatory integral, and for any $l \in \mathbb{N}$ such that $l \geq l_n$, the following holds:*

$$\tilde{I}_{\phi_n}[a_s](\lambda) := Os- \int_{\mathbb{R}^n} e^{i\lambda\phi_n(x)} a_s(x) dx = \int_{\mathbb{R}^n} e^{i\lambda\phi_n(x)} L_s^{*l}(a_s(x)) dx,$$

where $l_n := [(\tau^+ + n)/(\mu_n - 1 - \delta^+)] + 1$. And then for each $l \in \mathbb{N}$ such that $l \geq l_n$, there exists a positive constant $\tilde{C}_{s,l}$ such that for any $\lambda > 0$,

$$|\tilde{I}_{\phi_n}[a_s](\lambda)| \leq \tilde{C}_{s,l} \prod_{k=s+1}^n \|\varphi_k\| |a|_{l_n}^{(\tau)} |\psi_{1,s}|_r^{(l)} \lambda^{-l},$$

where $\|\varphi_k\| := \sup_{x_k \in \mathbb{R}} |\varphi_k(x_k)|$ and $|\psi_{1,s}|_r^{(l)}$ is defined by (3.10).

Proof. First we prove (i). Applying Lebesgue's convergence theorem to $e^{i\lambda\phi_n(x)} a_0(x)\chi(\varepsilon x) \in C_0^\infty(\mathbb{R}^n)$ with Proposition 2.2 shows (i). Next we prove (ii). Proposition 3.5 (iv) makes the order of integrand descend to be integrable in the sense of Lebesgue: For any $l \in \mathbb{N}$ such that $l \geq l_n := [(\tau^+ + n)/(\mu_n - 1 - \delta^+)] + 1$,

$$|e^{i\lambda\phi_n(x)} L_s^{*l}(a_s(x)\chi(\varepsilon x))| \leq C_l \lambda^{-l} \prod_{k=s+1}^n \|\varphi_k\| |a|_l^{(\tau)} |\psi_{1,s}|_r^{(l)} \langle x \rangle^t =: M(x),$$

where $t := \tau^+ - (\mu_n - 1 - \delta^+)l < -n$. Since $\int_{\mathbb{R}^n} M(x)dx$ is absolutely convergent independent of χ and ε , applying Lebesgue's convergence theorem with Proposition 2.2 on Proposition 3.5 (iii) as $\varepsilon \rightarrow +0$, completes the proof. \square

4. ASYMPTOTIC EXPANSIONS OF OSCILLATORY INTEGRALS

In this section, we show asymptotic expansions of oscillatory integrals in multi-variable. First the following holds.

Lemma 4.1. *Let φ_j and ψ_j be functions for $j = 1, \dots, n$. Then*

$$(4.1) \quad \prod_{j=1}^n (\varphi_j + \psi_j) = \sum_{(j_1, \dots, j_n) \in S_n} \sum_{s=0}^n \frac{1}{(n-s)!s!} \prod_{k=s+1}^n \varphi_{j_k} \prod_{k=1}^s \psi_{j_k},$$

where S_n is a symmetric group, $\prod_{k=n+1}^n \varphi_{j_k} := 1$ and $\prod_{k=1}^0 \psi_{j_k} := 1$.

Proof. By induction on $n \in \mathbb{N}$. \square

Next by Lemma 4.1, Theorem 2.6 in §2 and Theorem 3.6 in §3, we obtain the following theorem:

Theorem 4.2. *Assume that $\lambda > 0$, $\phi(x) := \sum_{j=1}^n \pm_j x_j^{m_j}$ where $m_j \in \mathbb{N}$ such that $m_1 \geq \dots \geq m_n \geq 2$ and \pm_j stands for “+” or “−” determined by j , and $a \in \mathcal{A}_\delta^r(\mathbb{R}^n)$. Then for any $N_1 \in \mathbb{N}$ such that $N_1 > m_1$, as $\lambda \rightarrow \infty$,*

$$(4.2) \quad Os\text{-} \int_{\mathbb{R}^n} e^{i\lambda\phi(x)} a(x) dx = \sum_{\alpha \in \Omega} c_\alpha \lambda^{-\sum_{j=1}^n \frac{\alpha_j + 1}{m_j}} + O\left(\lambda^{-\frac{N_1 - m_1 + 1}{m_1}}\right)$$

where

$$(4.3) \quad \Omega := \left\{ (\alpha_j) \in \mathbb{Z}_{\geq 0}^n \left| \sum_{j=1}^n \frac{\alpha_j + (1 - \delta_{1j})}{m_j} < \frac{N_1}{m_1} - 1 \right. \right\}$$

and

$$(4.4) \quad c_\alpha := \prod_{j=1}^n c_{j, \alpha_j}^{\pm_j} \frac{\partial_x^\alpha a(0)}{\alpha!}$$

with

$$\begin{aligned}
 c_{j,\alpha_j}^{\pm_j} &= m_j^{-1} \left\{ e^{\pm_j i \frac{\pi}{2} \frac{\alpha_j+1}{m_j}} + (-1)^{\alpha_j} e^{\pm_j (-1)^{m_j} i \frac{\pi}{2} \frac{\alpha_j+1}{m_j}} \right\} \Gamma\left(\frac{\alpha_j+1}{m_j}\right) \\
 (4.5) \quad &= \begin{cases} \frac{1}{k_j} e^{\pm_j i \frac{\pi}{2} \frac{2\beta_j+1}{2k_j}} \Gamma\left(\frac{2\beta_j+1}{2k_j}\right) & \text{for } m_j = 2k_j, \alpha_j = 2\beta_j, \\ 0 & \text{for } m_j = 2k_j, \alpha_j = 2\beta_j + 1, \\ \frac{2}{2k_j+1} \cos \frac{\pi(2\beta_j+1)}{2(2k_j+1)} \Gamma\left(\frac{2\beta_j+1}{2k_j+1}\right) & \text{for } m_j = 2k_j+1, \alpha_j = 2\beta_j, \\ \frac{\pm_j 2i}{2k_j+1} \sin \frac{\pi(\beta_j+1)}{2k_j+1} \Gamma\left(\frac{2\beta_j+2}{2k_j+1}\right) & \text{for } m_j = 2k_j+1, \alpha_j = 2\beta_j+1, \end{cases}
 \end{aligned}$$

where $k_j \in \mathbb{N}$ and $\beta_j \in \mathbb{Z}_{\geq 0}$ for $j = 1, \dots, n$.

Proof. First we divide $\int_{\mathbb{R}^n} e^{i\lambda\phi(x)} a(x) \chi(\varepsilon x) dx$ into an integral on compact support around the singular point of ϕ and integrals on non-compact support: After using cutoff function φ_j for $j = 1, \dots, n$ with Lemma 4.1, by proper permutations of variables in each term, without lost of generality, we can assume that $\sum_{s=0}^n \frac{n!}{(n-s)!s!} \int_{\mathbb{R}^n} e^{i\lambda\tilde{\phi}(x)} a_s(x) \chi(\varepsilon x) dx$ with (3.9). If $s = 0$, since $a_0 \in C_0^\infty(\mathbb{R}^n)$, then by Theorem 3.6 (i) and Theorem 2.6, for any $N_1 \in \mathbb{N}$ such that $N_1 > m_1$, as $\lambda \rightarrow \infty$,

$$\begin{aligned}
 I_0 &:= O_s \int_{\mathbb{R}^n} e^{i\lambda\phi(x)} a_0(x) dx = \int_{\mathbb{R}^n} e^{i\lambda\phi(x)} a_0(x) = \prod_{j=1}^n \int_{-\infty}^{\infty} dx_j e^{\pm_j i \lambda x_j^{m_j}} a_0(x) \\
 &= \sum_{\alpha_1=0}^{N_1-m_1-1} \prod_{j=2}^n \int_{-\infty}^{\infty} dx_j e^{\pm_j i \lambda x_j^{m_j}} c_{1,\alpha_1}^{\pm_1} \lambda^{-\frac{\alpha_1+1}{m_1}} \frac{\partial x_1^{\alpha_1}}{\alpha_1!} \Big|_{x_1=0} a_0(x) + O\left(\lambda^{-\frac{N_1-m_1+1}{m_1}}\right),
 \end{aligned}$$

where $c_{1,\alpha_1}^{\pm_1}$ is given by (4.5). Here for any $j = 2, \dots, n$, let $N_j \in \mathbb{N}$ such that $N_j \geq N_1$, then $N_j > m_j$, and for any $\lambda \geq 1$,

$$(4.6) \quad \frac{N_j}{m_j} > \frac{N_1}{m_1} - \sum_{k=2}^j \frac{1}{m_k} \iff \lambda^{-\sum_{k=1}^{j-1} \frac{1}{m_k} - \frac{N_j-m_j+1}{m_j}} \leq \lambda^{-\frac{N_1-m_1+1}{m_1}}.$$

Repeating to apply Theorem 2.6 with (4.6) gives inductively

$$I_0 = \sum_{\alpha_1=0}^{N_1-m_1-1} \cdots \sum_{\alpha_n=0}^{N_n-m_n-1} c_\alpha \lambda^{-\sum_{j=1}^n \frac{\alpha_j+1}{m_j}} + O\left(\lambda^{-\frac{N_1-m_1+1}{m_1}}\right) \quad (\lambda \rightarrow \infty),$$

where c_α is given by (4.4). Let Ω be defined by (4.3). Then

$$\begin{aligned}
 \frac{\alpha_j + (1 - \delta_{1j})}{m_j} &< \sum_{j=1}^n \frac{\alpha_j + (1 - \delta_{1j})}{m_j} < \frac{N_1}{m_1} - 1 \leq \frac{N_j}{m_j} - 1 \\
 &\implies \alpha_j \leq N_j - m_j - 1,
 \end{aligned}$$

for $j = 1, \dots, n$, and for any $\lambda \geq 1$,

$$\sum_{j=1}^n \frac{\alpha_j + (1 - \delta_{1j})}{m_j} < \frac{N_1}{m_1} - 1 \iff \lambda^{-\sum_{j=1}^n \frac{\alpha_j+1}{m_j}} > \lambda^{-\frac{N_1-m_1+1}{m_1}}.$$

Hence $\lambda \rightarrow \infty$,

$$I_0 := Os- \int_{\mathbb{R}^n} e^{i\lambda\phi(x)} a_0(x) dx = \sum_{\alpha \in \Omega} c_\alpha \lambda^{-\sum_{j=1}^n \frac{\alpha_j+1}{m_j}} + O\left(\lambda^{-\frac{N_1-m_1+1}{m_1}}\right).$$

If $s \neq 0$, since a_s has a non-compact support, then Theorem 3.6 (ii) gives

$$\sum_{s=1}^n \frac{n!}{(n-s)!s!} Os- \int_{\mathbb{R}^n} e^{i\lambda\phi(x)} a_s(x) dx = O\left(\lambda^{-\frac{N_1-m_1+1}{m_1}}\right) \quad (\lambda \rightarrow \infty).$$

□

By Theorem 4.2, we obtain the following immediately.

Corollary 4.3. *Assume that $\lambda > 0$ and $a \in \mathcal{A}_\delta^\tau(\mathbb{R}^n)$. Then the following hold:*

- (i) *Let $m \in \mathbb{N}$ such that $m \geq 2$. Then for any $N \in \mathbb{N}$ such that $N > m$, as $\lambda \rightarrow \infty$,*

$$Os- \int_{\mathbb{R}^n} e^{i\lambda \sum_{j=1}^n \pm_j x_j^m} a(x) dx = \sum_{\alpha \in \Omega} c_\alpha \lambda^{-\frac{|\alpha|+n}{m}} + O\left(\lambda^{-\frac{N-m+1}{m}}\right),$$

where Ω and c_α are defined by (4.3) when $N_1 = N$, and (4.4) with

$$c_{j,\alpha_j}^{\pm_j} = m^{-1} \left\{ e^{\pm_j i \frac{\pi}{2} \frac{\alpha_j+1}{m}} + (-1)^{\alpha_j} e^{\pm_j (-1)^m i \frac{\pi}{2} \frac{\alpha_j+1}{m}} \right\} \Gamma\left(\frac{\alpha_j+1}{m}\right).$$

- (ii) *For any $N \in \mathbb{N}$ such that $N > 2$, as $\lambda \rightarrow \infty$,*

$$\begin{aligned} & \int_{\mathbb{R}^n} e^{i\lambda(\sum_{j=1}^p x_j^2 - \sum_{j=p+1}^n x_j^2)} a(x) dx \\ &= \pi^{\frac{n}{2}} e^{i\frac{\pi}{4}\{p-(n-p)\}} \sum_{\beta \in \Omega'} \frac{(-1)^{\sum_{j=p+1}^n i|\beta|} \partial_x^{2\beta} a(0)}{4^{|\beta|} \beta!} \lambda^{-|\beta| - \frac{n}{2}} + O\left(\lambda^{-\frac{N}{2} + \frac{1}{2}}\right), \end{aligned}$$

where $\Omega' = \{(\beta_j) \in \mathbb{Z}_{\geq 0}^n \mid |\beta| < (N - n - 1)/2\}$.

Proof. (i) Set $m_j = m \geq 2$ for $j = 1, \dots, n$ on Theorem 4.2. (ii) Set $m = 2$, $\pm_j = +$ for $j = 1, \dots, p$ and $\pm_j = -$ for $j = p+1, \dots, n$ on (i), and apply (4.5). □

By Theorem 2.3 in §2 and Corollary 4.3, we can consider that Theorem 4.2 is an extension of the stationary phase method.

Corollary 4.4. *Assume that $\lambda > 0$ and $a \in \mathcal{A}_\delta^\tau(\mathbb{R}^3)$. Then the following hold:*

- (i) *For any $N \in \mathbb{N}$ such that $N > k+1$, as $\lambda \rightarrow \infty$,*

$$Os- \int_{\mathbb{R}^3} e^{i\lambda(\pm x_1^{k+1} + x_2^2 + x_3^2)} a(x) dx = \sum_{\alpha \in \Omega_1} c_\alpha \lambda^{-\frac{\alpha_1+1}{k+1} - \frac{\alpha_2+1}{2} - \frac{\alpha_3+1}{2}} + O\left(\lambda^{-\frac{N-k}{k+1}}\right),$$

- (ii) *For any $N \in \mathbb{N}$ such that $N > 4$, as $\lambda \rightarrow \infty$,*

$$Os- \int_{\mathbb{R}^3} e^{i\lambda(\pm x_1^4 + x_2^3 + x_3^2)} a(x) dx = \sum_{\alpha \in \Omega_2} c_\alpha \lambda^{-\frac{\alpha_1+1}{4} - \frac{\alpha_2+1}{3} - \frac{\alpha_3+1}{2}} + O\left(\lambda^{-\frac{N-3}{4}}\right),$$

- (iii) *For any $N \in \mathbb{N}$ such that $N > 5$, as $\lambda \rightarrow \infty$,*

$$Os- \int_{\mathbb{R}^3} e^{i\lambda(x_1^5 + x_2^3 + x_3^2)} a(x) dx = \sum_{\alpha \in \Omega_3} c_\alpha \lambda^{-\frac{\alpha_1+1}{5} - \frac{\alpha_2+1}{3} - \frac{\alpha_3+1}{2}} + O\left(\lambda^{-\frac{N-4}{5}}\right),$$

where Ω_j for $j = 1, 2, 3$, and c_α with c_{j,α_j}^\pm are defined by (4.3) when $N_1 = N$, and (4.4) with (4.5).

Proof. Set $n = 3$ and $(m_1, m_2, m_3; \pm_1, \pm_2, \pm_3) = (k + 1, 2, 2; \pm, +, +)$
 $= (4, 3, 2; \pm, +, +) = (5, 3, 2; +, +, +)$ on Theorem 4.2 respectively. \square

REFERENCES

- [1] V.I. ARNOLD, *Normal forms of functions with simple singular points, the Weyl groups A_k , D_k , E_k , and Lagrange immersions*, Functional Anal. Appl. 254-272, (Russian original, Funkc. anal. i prilozh. **6**-4 (1972), 3–25).
- [2] V.I. ARNOLD, S.M. GUSEIN-ZADE AND A.N. VARČENKO, *Singularities of differentiable maps volume I, II*, Birkhäuser, I, 1985, II, 1988.
- [3] J.J. DUISTERMAAT, *Oscillatory integrals, Lagrange immersions and unfoldings of singularities*, Comm. Pure Appl. Math., **27** (1974), 207–281.
- [4] J.J. DUISTERMAAT, *Fourier integral operators*, Birkhäuser, Boston. 1996.
- [5] D. FUJIWARA, *Rigorous time slicing approach to Feynman path integrals*, Springer, 2017.
- [6] D. FUJIWARA, *Asymptotic method in linear partial differential equations*, Iwanami Shoten, 2019 (in Japanese).
- [7] L. HÖRMANDER, *Fourier integral operators I*, Acta Math., **127** (1971), 79–183.
- [8] L. HÖRMANDER, *The analysis of linear partial differential operators I*, Springer, 1983.
- [9] H. KUMANO-GO, *Pseudo-differential operators*, MIT Press, 1981.
- [10] T. NAGANO AND N. MIYAZAKI, *Generalized Fresnel integrals as oscillatory integrals with positive real power phase functions and applications to asymptotic expansions*, arXiv:2005.12754v1 [math.CA], 24 May 2020.
- [11] T. NAGANO AND N. MIYAZAKI, *Asymptotic expansions of oscillatory integrals for certain phase functions with a degenerate singular point including type A_k , E_6 , E_8 -function germs*, in preparation.

DEPARTMENT OF LIBERAL ARTS, FACULTY OF SCIENCE AND TECHNOLOGY, TOKYO UNIVERSITY OF SCIENCE, 2641, YAMAZAKI, NODA, CHIBA 278-8510, JAPAN
E-mail address: `tonagan@rs.tus.ac.jp`

DEPARTMENT OF MATHEMATICS, FACULTY OF ECONOMICS, KEIO UNIVERSITY, YOKOHAMA, 223-8521, JAPAN
E-mail address: `miyazaki@a6.keio.jp`