

Multiple solutions of Kazdan-Warner equation on graphs in the negative case

Shuang Liu, Yunyan Yang¹

Department of Mathematics, Renmin University of China, Beijing 100872, P. R. China

Abstract

Let $G = (V, E)$ be a finite connected graph, and let $\kappa : V \rightarrow \mathbb{R}$ be a function such that $\int_V \kappa d\mu < 0$. We consider the following Kazdan-Warner equation on G :

$$\Delta u + \kappa - K_\lambda e^{2u} = 0,$$

where $K_\lambda = K + \lambda$ and $K : V \rightarrow \mathbb{R}$ is a non-constant function satisfying $\max_{x \in V} K(x) = 0$ and $\lambda \in \mathbb{R}$. By a variational method, we prove that there exists a $\lambda^* > 0$ such that when $\lambda \in (-\infty, \lambda^*]$ the above equation has solutions, and has no solution when $\lambda \geq \lambda^*$. In particular, it has only one solution if $\lambda \leq 0$; at least two distinct solutions if $0 < \lambda < \lambda^*$; at least one solution if $\lambda = \lambda^*$. This result complements earlier work of Grigor'yan-Lin-Yang [7], and is viewed as a discrete analog of that of Ding-Liu [4] and Yang-Zhu [17] on manifolds.

Key words: Kazdan-Warner problem on graph, variation problem on graph
2010 MSC: 35R02, 34B45

1. Introduction

Variational method is always a powerful tool in partial differential equations and geometric analysis. Recently, using this tool, Grigor'yan-Lin-Yang [7, 8, 9] obtained existence results for solutions to various partial differential equations on graphs. In particular, Kazdan-warner equation was proposed on graphs in [7]. The Kazdan-Warner equation arises from the basic geometric problem on prescribing Gaussian curvature of Riemann surface, which systematically studied by Kazdan-Warner [12, 13]. On a closed Riemann surface (Σ, g) with the Gaussian curvature κ , let $\tilde{g} = e^{2u}g$ be a smooth metric conformal to g and K be the Gaussian curvature with respect to \tilde{g} . Then u satisfies the equation

$$\Delta_g u + \kappa - K e^{2u} = 0, \tag{1}$$

where Δ_g denotes the Laplace-Beltrami operator with respect to the metric g . Let v be a solution to $\Delta_g v = \bar{\kappa} - \kappa$ and $f = 2(u - v)$, where $\bar{\kappa}$ is the averaged integral of κ . Then the above equation is transformed to

$$\Delta_g f + 2\bar{\kappa} - (2K e^{2v})e^f = 0.$$

¹Corresponding author

Email addresses: shuangliu@ruc.edu.cn (Shuang Liu), yunyan yang@ruc.edu.cn (Yunyan Yang)

*Preprint submitted to ****

November 25, 2021

Hence, one can free (1) from the geometric situation, and just studies the equation

$$\Delta_g f + c - h e^f = 0, \quad (2)$$

where c is a constant and h is a function. On graphs, it seems to be out of reach to resemble this topic in terms of Gaussian curvature. Therefore, in [7], the authors focused on the equation similar to the form of (2), namely the Kazdan-Warner equation on graph, and obtained the following: when $c = 0$, it has a solution if and only if h changes sign and the integral of h is negative; when $c > 0$, it has a solution if and only if h is positive somewhere; when $c < 0$, there is a threshold $c_h < 0$ such that it has a solution if $c \in (c_h, 0)$, but it has no solution for any $c < c_h$. Later, Ge [5] found a solution in the critical case $c = c_h$. More recently Ge-Jiang [6] studied the Kazdan-Warner equation on infinite graphs and Keller-Schwarz [11] on canonically compactifiable graphs; Camilli-Marchi [3] extended the Kazdan-Warner equation on network; for other related works, we refer the readers to [10, 14].

Let us come back to a closed Riemann surface (Σ, g) , whose Euler characteristic is negative, or equivalently $\int_{\Sigma} K dv_g < 0$. Replacing K by $K + \lambda$ in (1) with $K \leq 0$, $K \not\equiv 0$, and $\lambda \in \mathbb{R}$, Ding-Liu [4] obtained the following conclusion by using a method of upper and lower solutions and a variational method: there exists a $\lambda^* > 0$ such that if $\lambda \leq 0$, then (1) has a unique solution; if $0 < \lambda < \lambda^*$, then (1) has at least two distinct solutions; if $\lambda = \lambda^*$, then (1) has at least one solution; if $\lambda > \lambda^*$, then (1) has no solution. Recently, this result was partly reproved by Borer-Galimberti-Struwe [2] via a monotonicity technique due to Struwe [15, 16], and was extended to the case of conical metrics by Yang-Zhu [17].

Our aim is to extend results of Ding-Liu [4] to graphs. Let us recall some notations from graph theory. Throughout this paper, $G = (V, E)$ is assumed to be a finite connected graph. The edges on the graph are allowed to be weighted. Weights are given by a function $\omega : V \times V \rightarrow [0, \infty)$, the edge xy from x to y has weight $\omega_{xy} > 0$. We assume this weight function is symmetric, $\omega_{xy} = \omega_{yx}$. Let $\mu : V \rightarrow \mathbb{R}^+$ be a positive measure on the vertices of the G . Denote by $V^{\mathbb{R}}$ the space of real functions on V . and by $\ell_{\mu}^p = \{f \in V^{\mathbb{R}} : \sum_{x \in V} \mu(x) |f(x)|^p < \infty\}$, for any $1 \leq p < \infty$, the space of ℓ^p integrable functions on V with respect to the measure μ . For $p = \infty$, let $\ell^{\infty} = \{f \in V^{\mathbb{R}} : \sup_{x \in V} |f(x)| < \infty\}$ be the set of all bounded functions. As usual, we define the ℓ_{μ}^p norm of $f \in \ell_{\mu}^p$, $1 \leq p \leq \infty$, by

$$\|f\|_p = \left(\sum_{x \in V} \mu(x) |f(x)|^p \right)^{1/p}, \quad 1 \leq p < \infty, \quad \|f\|_{\infty} = \sup_{x \in V} |f(x)|.$$

We define the Laplacian $\Delta : V^{\mathbb{R}} \rightarrow V^{\mathbb{R}}$ on G by

$$\Delta f(x) = \frac{1}{\mu(x)} \sum_{y \sim x} \omega_{xy} (f(x) - f(y)). \quad (3)$$

Given the weight ω on E , there are two typical choices of Laplacian as follows:

- $\mu(x) = \deg(x) := \sum_{y \sim x} \omega_{xy}$ for all $x \in V$, which is called the normalized graph Laplacian;
- $\mu(x) \equiv 1$ for all $x \in V$, which is the combinatorial graph Laplacian.

In this paper, we do not restrict $\mu(x)$ to the above two forms, but only require $\mu(x) > 0$ for all $x \in V$. Note that the Laplace operator defined in (3) is the negative usual Laplace operator. The

gradient form is defined by

$$\begin{aligned} 2\Gamma(f, g)(x) &= (f \cdot \Delta g + g \cdot \Delta f - \Delta(f \cdot g))(x) \\ &= \frac{1}{\mu(x)} \sum_{y \sim x} \omega_{xy} (f(x) - f(y))(g(x) - g(y)). \end{aligned}$$

For the sake of simplicity, we write $\Gamma(f, f) = \Gamma(f)$. Sometimes we use the notation $\nabla f \nabla g = \Gamma(f, g)$. The length of the gradient is denoted by

$$|\nabla f|(x) = \sqrt{\Gamma(f)(x)}.$$

From now on, we write $\int_V u d\mu = \sum_{x \in V} \mu(x)u(x)$. Define a Sobolev space with a norm on the graph G by

$$W^{1,2}(V) = \left\{ u \in V^{\mathbb{R}} : \int_V (|\nabla u|^2 + u^2) d\mu < +\infty \right\},$$

and

$$\|u\|_{W^{1,2}(V)} = \left(\int_V (|\nabla u|^2 + u^2) d\mu \right)^{1/2}$$

respectively. Since G is a finite graph, we have that $W^{1,2}(V)$ is exactly $V^{\mathbb{R}}$, a finite dimensional linear space. This implies the following Sobolev embedding:

Lemma 1 ([7], Lemma 5). *If G is a finite graph, then the Sobolev space $W^{1,2}(V)$ is precompact. Namely, if $\{u_j\}$ is bounded in $W^{1,2}(V)$, then there exists some $u \in W^{1,2}(V)$ such that up to a subsequence, $u_j \rightarrow u$ in $W^{1,2}(V)$.*

The Kazdan-Warner equation we are interested in this paper reads as

$$\Delta u + \kappa - K_\lambda e^{2u} = 0 \quad \text{on } V, \tag{4}$$

where $\kappa \in V^{\mathbb{R}}$ is a function, and $K_\lambda = K + \lambda$, $\lambda \in \mathbb{R}$, $K \in V^{\mathbb{R}}$ is a function. Now we are ready to state our main results.

Theorem 2. *Let $G = (V, E)$ be a finite graph, κ and K_λ be given as in (4) such that $\int_V \kappa d\mu < 0$, $K \leq \max_V K = 0$, and $K \not\equiv 0$. Then there exists a $\lambda^* \in (0, -\min_V K)$ satisfying*

1. *if $\lambda \leq 0$, then (4) has a unique solution;*
2. *if $0 < \lambda < \lambda^*$, then (4) has at least two distinct solutions;*
3. *if $\lambda = \lambda^*$, then (4) has at least one solution;*
4. *if $\lambda > \lambda^*$, then (4) has no solution;*

Remark 1. *The assertion of $\lambda^* < -\min_V K$ comes from the conclusion of Step 2 in Subsection 3.3.*

Remark 2. *Compared to the existence of solutions in the literature (see for example [7, 5]), the above results firstly reveal the multiple solution problem of Kazdan-Warner equation on graphs in the negative case.*

The proof of Theorem 2 is based on the method of variation. It can be viewed as a discrete analog of the result of Ding-Liu [4]. The remaining part of this paper will be organized as follows: In Section 2, we give several preliminary lemmas for our use later; In Section 3, we finish the proof of Theorem 2.

2. Preliminaries

In this section, we provide discrete versions of the maximum principle, the Palais-Smale condition and the upper and lower solution principle. Note that $G = (V, E)$ is a finite connected graph.

2.1. maximum principle

To proceed, we need the following maximum principles, which are known for experts (see for examples [7, 8]). For readers' convenience, we include the detailed proofs here.

Lemma 3 (Weak maximum principle). *For any constant $c > 0$, if u satisfies $\Delta u + cu \geq 0$, then $u \geq 0$ on V .*

Proof. Let $u^- = \min\{u, 0\}$. For any $x \in V$, we claim that

$$\Delta u^-(x) + cu^-(x) \geq 0, \quad (5)$$

from which, one has

$$\int_V \Gamma(u^-) d\mu + c \|u^-\|_{\ell_\mu^2}^2 = \langle u^-, \Delta u^- + cu^- \rangle \leq 0.$$

This leads to $u^- \equiv 0$ on V .

To prove this claim, we first consider the case $u(x) \geq 0$. Therefore, $cu^-(x) = 0$ and

$$\Delta u^-(x) = \frac{1}{\mu(x)} \sum_{y \sim x} \omega_{xy}(u^-(x) - u^-(y)) = -\frac{1}{\mu(x)} \sum_{y \sim x} \omega_{xy} u^-(y) \geq 0,$$

due to $u^-(z) \leq 0$ for any $z \in V$. In the case $u(x) < 0$, one has $cu^-(x) = cu(x)$ and thus

$$\begin{aligned} \Delta u^-(x) &= \frac{1}{\mu(x)} \sum_{y \sim x} \omega_{xy}(u^-(x) - u^-(y)) = \frac{1}{\mu(x)} \sum_{y \sim x} \omega_{xy}(u(x) - u^-(y)) \\ &\geq \frac{1}{\mu(x)} \sum_{y \sim x} \omega_{xy}(u(x) - u(y)) = \Delta u(x). \end{aligned}$$

It follows that $\Delta u^-(x) + cu^-(x) \geq \Delta u(x) + cu(x) \geq 0$, which confirms (5) and ends the proof of the lemma. \square

Lemma 4 (Strong maximum principle). *Suppose that $u \geq 0$, and that $\Delta u + cu \geq 0$ for some constant $c > 0$. If there exists $x_0 \in V$ such that $u(x_0) = 0$, then $u \equiv 0$ on V .*

Proof. Let $x = x_0$, we have

$$\frac{1}{\mu(x_0)} \sum_{y \sim x_0} \omega_{yx_0}(u(x_0) - u(y)) + cu(x_0) \geq 0,$$

which implies

$$\frac{1}{\mu(x_0)} \sum_{y \sim x_0} \omega_{yx_0} u(y) \leq 0.$$

Since $u \geq 0$ and $\omega_{yx_0} > 0$ for all $y \sim x_0$, we obtain

$$u(y) = 0, \quad \text{for all } y \sim x_0.$$

Therefore, $u \equiv 0$ on V by the connectedness of G . \square

2.2. Palais-Smale condition

We define a functional $E_\lambda : W^{1,2}(V) \rightarrow \mathbb{R}$ by

$$E_\lambda(u) = \int_V (|\nabla u|^2 + 2\kappa u - K_\lambda e^{2u}) d\mu,$$

where κ and K_λ are given as in the assumptions of Theorem 2, in particular $\int_V \kappa d\mu < 0$. For any $\phi \in W^{1,2}(V)$, denote by $dE_\lambda(u)(\phi)$ the Frechet derivative of the functional, by $d^k E_\lambda(u)(\phi, \dots, \phi)$ the Frechet derivative of order $k \geq 2$.

Lemma 5 (Palais-Smale condition). *Suppose that $V_\lambda^- = \{x \in V : K_\lambda(x) < 0\}$ is nonempty for some $\lambda \in \mathbb{R}$. Then E_λ satisfies the $(PS)_c$ condition for all $c \in \mathbb{R}$, i.e. if (u_j) is a sequence of functions in $W^{1,2}(V)$ such that $E_\lambda(u_j) \rightarrow c$ and $dE_\lambda(u_j)(\phi) \rightarrow 0$ for all $\phi \in W^{1,2}(V)$ as $j \rightarrow \infty$, then there exists some $u_0 \in W^{1,2}(V)$ satisfying $u_j \rightarrow u_0$ in $W^{1,2}(V)$.*

Proof. Let (u_j) be a function sequence such that $E_\lambda(u_j) \rightarrow c$ and $dE_\lambda(u_j)(\phi) \rightarrow 0$, or equivalently

$$\int_V (|\nabla u_j|^2 + 2\kappa u_j - K_\lambda e^{2u_j}) d\mu = c + o_j(1), \quad (6)$$

$$\int_V (\nabla u_j \nabla \phi + \kappa \phi - K_\lambda e^{2u_j} \phi) d\mu = o_j(1) \|\phi\|_{W^{1,2}(V)}, \quad \forall \phi \in W^{1,2}(V), \quad (7)$$

where $o_j(1) \rightarrow 0$ as $j \rightarrow \infty$.

Let $\phi \equiv 1$ in (7), one has

$$\int_V (\kappa - K_\lambda e^{2u_j}) d\mu = o_j(1) \mu(V)^{1/2},$$

which implies

$$\int_V K_\lambda e^{2u_j} d\mu = \int_V \kappa d\mu + o_j(1). \quad (8)$$

Inserting (8) into (6), we obtain

$$\int_V (|\nabla u_j|^2 + 2\kappa u_j) d\mu = \int_V \kappa d\mu + c + o_j(1). \quad (9)$$

We now *claim* that u_j is bounded in ℓ_μ^2 . Suppose not, there holds $\|u_j\|_{\ell_\mu^2} \rightarrow \infty$. We set $v_j = \frac{u_j}{\|u_j\|_{\ell_\mu^2}}$. By the Cauchy-Schwarz inequality, one has

$$\int_V \kappa \frac{u_j}{\|u_j\|_{\ell_\mu^2}^2} d\mu = o_j(1).$$

This together with (9) leads to

$$\int_V |\nabla v_j|^2 d\mu = o_j(1). \quad (10)$$

Hence, v_j is bounded in $W^{1,2}(V)$. In view of Lemma 1 and (10), $v_j \rightarrow \gamma$ in $W^{1,2}(V)$ for some constant γ . Here and in the sequel, we do not distinguish sequence and subsequence. Since $\|v_j\|_{\ell_\mu^2} = 1$, we have $\gamma \neq 0$. It follows from (9) that

$$\int_V \kappa v_j d\mu \leq o_j(1).$$

Passing to the limit $j \rightarrow \infty$ in the above inequality, we conclude that $\gamma \geq 0$ since $\int_V \kappa d\mu < 0$. Therefore $\gamma > 0$.

On the other hand, for any $x \in V_\lambda^-$, if there exists $N \in \mathbb{N}$, if $j > N$ such that $u_j(x) \leq 0$, then $\lim_{j \rightarrow \infty} v_j(x) \leq 0$, which contradicts $\gamma > 0$ and confirms our claim. If not, let $x_* \in V_\lambda^-$, due to the finiteness of V , we can choose a subsequence $\{j_k\}_{k=0}^\infty$ such that $u_{j_k}(x_*) > 0$. Set

$$\phi(x) = \begin{cases} u_{j_k}(x_*), & x = x_* \\ 0, & x \neq x_*. \end{cases}$$

Then

$$\|\phi\|_{W^{1,2}(V)}^2 = 2 \sum_{y \sim x_*} \omega_{x_* y} u_{j_k}^2(x_*) + \mu(x_*) u_{j_k}^2(x_*) = (2 \deg(x_*) + \mu(x_*)) u_{j_k}^2(x_*).$$

Substituting it into (7), we have

$$\Delta u_{j_k}(x_*) + \kappa(x_*) - K_\lambda(x_*) e^{2u_{j_k}(x_*)} \leq C'. \quad (11)$$

Since $v_j \rightarrow \gamma$, $\|u_j\|_{\ell_\mu^2} \rightarrow +\infty$ and V has finite points, we conclude

$$u_j = (\gamma + o_j(1)) \|u_j\|_{\ell_\mu^2} \quad \text{uniformly on } V.$$

This together with (11) leads to

$$\|u_{j_k}\|_{\ell_\mu^2}^2 O_{j_k}(1) + \kappa(x_*) - K_\lambda(x_*) e^{2(\gamma + o_{j_k}(1)) \|u_{j_k}\|_{\ell_\mu^2}^2} \leq C',$$

which is impossible since $K_\lambda(x_*) < 0$. Then our claim follows immediately.

Since u_j is bounded in ℓ_μ^2 , we have u_j is bounded in $W^{1,2}(V)$ due to the finiteness of V . Therefore, by Lemma 1, there exists some $u_0 \in W^{1,2}(V)$ such that up to subsequence, $u_j \rightarrow u_0$ in $W^{1,2}(V)$. \square

2.3. Upper and lower solutions principle

Let $f : V \times \mathbb{R} \rightarrow \mathbb{R}$ be a function, and f is smooth with respect to the second variable. We say that $u \in V^\mathbb{R}$ is an upper (lower) solution to the following equation

$$\Delta u(x) + f(x, u(x)) = 0, \quad x \in V, \quad (12)$$

if u satisfies $\Delta u(x) + f(x, u(x)) \geq (\leq) 0$ for any $x \in V$. We generalize ([7], Lemma 8) to the following:

Lemma 6. Suppose that φ, ψ are lower and upper solution to (12) respectively with $\varphi \leq \psi$ on V . Then (12) has a solution u with $\varphi \leq u \leq \psi$ on V .

Proof. This is a discrete version of the argument of Kazdan-Warner ([12], Lemma 9.3), and the method of proof carries over to the setting of graphs.

Since the graph is finite, there exists a constant A such that $-A \leq \varphi \leq \psi \leq A$. One can find a sufficient large constant c such that $F(x, t) = ct - f(x, t)$ is increasing with respect to $t \in [-A, A]$ for any fixed $x \in V$. We define an operator $Lu = \Delta u + cu$, and L is a compact operator and $\text{Ker}(L) = \text{span}\{1\}$ due to the finiteness of the graph. Hence, we can define $\varphi_{j+1}, \psi_{j+1}$ inductively as the unique solution to

$$\varphi_0 = \varphi, \quad L\varphi_{j+1}(x) = c\varphi_j(x) - f(x, \varphi_j(x)), \quad \forall j \geq 0, x \in V,$$

$$\psi_0 = \psi, \quad L\psi_{j+1}(x) = c\psi_j(x) - f(x, \psi_j(x)), \quad \forall j \geq 0, x \in V$$

respectively. Combining with the definition of upper (lower) solution and the monotonicity of $F(x, t)$ with respect to t , we obtain

$$L\varphi_0(x) \leq L\varphi_1(x) = F(x, \varphi(x)) \leq F(x, \psi(x)) = L\psi_1(x) \leq L\psi(x), \quad x \in V.$$

Then the weak maximum principle (see Lemma 3) yields that

$$\varphi \leq \varphi_1 \leq \psi_1 \leq \psi.$$

Moreover, it turns out that φ_1 and ψ_1 are lower and upper solution to (12) respectively. By induction, we have

$$\varphi \leq \varphi_j \leq \varphi_{j+1} \leq \psi_{j+1} \leq \psi_j \leq \psi, \quad j = 1, 2, \dots$$

Since V is finite, it is easy to see that up to a subsequence, $\varphi_j \rightarrow u_1, \psi_j \rightarrow u_2$ uniformly on V , and $u = u_1$ or u_2 is a solution to (12) with $\varphi \leq u \leq \psi$ on V . \square

3. Proof of Theorem 2

3.1. *Unique solution in the case $\lambda \leq 0$.*

Claim 1. E_λ is strictly convex on $W^{1,2}(V)$.

Proof. We only need to show that there exists some constant $C > 0$ such that

$$d^2 E_\lambda(u)(h, h) \geq C \|h\|_{W^{1,2}(V)}^2, \quad \forall u, h \in W^{1,2}(V). \quad (13)$$

Suppose not, there would be a function $u \in W^{1,2}(V)$ and a function sequence $h_j \in W^{1,2}(V)$ such that $\|h_j\|_{W^{1,2}(V)} = 1$ for all j and $d^2 E_\lambda(u)(h_j, h_j) \rightarrow 0$ as $j \rightarrow \infty$. From Lemma 1, there exists $h_\infty \in W^{1,2}(V)$, such that up to a subsequence, $h_j \rightarrow h_\infty$ as $j \rightarrow \infty$ in $W^{1,2}(V)$. Since

$$d^2 E_\lambda(u)(h_j, h_j) = 2 \int_V (|\nabla h_j|^2 - 2K_\lambda e^{2u} h_j^2) d\mu,$$

and $K_\lambda \leq 0$, it follows that $\int_V |\nabla h_j|^2 d\mu \rightarrow 0$ and $\int_V K_\lambda e^{2u} h_j^2 d\mu \rightarrow 0$, which lead to $h_\infty \equiv c$ for some constant c , and moreover

$$c^2 \int_V K_\lambda e^{2u} d\mu = \lim_{j \rightarrow \infty} \int_V K_\lambda e^{2u} h_j^2 d\mu = 0.$$

It is easily seen that $\int_V K_\lambda e^{2u} d\mu < 0$ by $K \not\equiv 0$, thus $c = 0$. This contradicts

$$\|h_\infty\|_{W^{1,2}(V)} = \lim_{j \rightarrow \infty} \|h_j\|_{W^{1,2}(V)} = 1.$$

Hence (13) holds. \square

Claim 2. For any $\varepsilon > 0$, there exist constants $C, C(\varepsilon) > 0$ such that

$$E_\lambda(u) \geq (C - 2\varepsilon) \|u\|_{W^{1,2}(V)}^2 - 2C(\varepsilon).$$

Proof. By Young's inequality, for any $\varepsilon > 0$, there exists a constant $C(\varepsilon) > 0$ such that

$$\left| \int_V \kappa u d\mu \right| \leq \varepsilon \|u\|_{W^{1,2}(V)}^2 + C(\varepsilon).$$

Thus, it is sufficient to find some constant $C > 0$ such that for all $u \in W^{1,2}(V)$

$$\int_V (|\nabla u|^2 - K_\lambda e^{2u}) d\mu \geq C \|u\|_{W^{1,2}(V)}^2.$$

Suppose not, there would exist a sequence of functions u_j satisfying

$$\int_V (|\nabla u_j|^2 + u_j^2) d\mu = 1, \quad \int_V (|\nabla u_j|^2 - K_\lambda e^{2u_j}) d\mu = o_j(1).$$

Clearly, u_j is bounded in $W^{1,2}(V)$, it follows from Lemma 1 that there exists some function $u \in W^{1,2}(V)$ such that up to a subsequence, $u_j \rightarrow u$ in $W^{1,2}(V)$ as $j \rightarrow \infty$. Due to $K_\lambda \not\equiv 0$ and $K_\lambda \leq 0$, we have

$$0 < \int_V (|\nabla u|^2 - K_\lambda e^{2u}) d\mu = \lim_{j \rightarrow \infty} \int_V (|\nabla u_j|^2 - K_\lambda e^{2u_j}) d\mu = 0,$$

which gets a contradiction. \square

Proof of (1) in Theorem 2. It is a consequence of Claim 1 and Claim 2. Precisely we denote $\Lambda = \inf_{u \in W^{1,2}(V)} E_\lambda(u)$. By Claim 2, we see that Λ is a definite real number. Take a function sequence $u_j \in W^{1,2}(V)$ such that $E_\lambda(u_j) \rightarrow \Lambda$ as $j \rightarrow \infty$. Applying Claim 2, we have that u_j is bounded in $W^{1,2}(V)$. Then, in view of Lemma 1, there exists a subsequence of u_j (still denoted by u_j) and a function $u_0 \in W^{1,2}(V)$ such that $u_j \rightarrow u_0$ in $W^{1,2}(V)$. Obviously $E_\lambda(u_0) = \Lambda$, and thus u_0 is a critical point of E_λ . We also need to explain why E_λ has only one critical point. For otherwise, we assume u^* is another critical point of E_λ . Note that $dE_\lambda(u_0) = dE_\lambda(u^*) = 0$. It follows from Claim 1, particularly from (13), that

$$\begin{aligned} E_\lambda(u_0) &= E_\lambda(u^*) + dE_\lambda(u^*)(u_0 - u^*) + \frac{1}{2} d^2 E_\lambda(\xi)(u_0 - u^*, u_0 - u^*) \\ &\geq E_\lambda(u^*) + C \|u_0 - u^*\|_{W^{1,2}(V)}^2 \end{aligned}$$

for some positive constant C , where ξ is a function lies between u^* and u_0 . Hence we have $E_\lambda(u_0) > E_\lambda(u^*)$, contradicting the fact that $E_\lambda(u_0) = \Lambda$. This implies the uniqueness of the critical point of E_λ . \square

3.2. Multiplicity of solutions for $0 < \lambda < \lambda^*$.

Fixing $\lambda \in (0, \lambda^*)$, we will seek two different solutions of (4). One is a strict local minimum of the functional E_λ , and the other is from mountain-pass theorem. We firstly prove the existence of λ^* . Consider the case $\lambda = 0$ in the equation (4) as follows

$$\Delta u + \kappa - K e^{2u} = 0. \tag{14}$$

For the solution u_0 of (14), the linearized equation of (14) at u_0

$$\Delta v - 2K e^{2u_0} v = 0$$

has only a trivial solution $v \equiv 0$, since $K \leq 0$ and $K \not\equiv 0$. Indeed,

$$0 \leq \int_V |\nabla v|^2 d\mu = \int_V v \Delta v d\mu = 2 \int_V K e^{2u_0} v^2 d\mu \leq 0,$$

which implies $v(x) = 0$ when $K(x) < 0$. In the case $K(x) = 0$, we have $\Delta v(x) = 0$. Thus

$$\int_V |\nabla v|^2 d\mu = \int_V v \Delta v d\mu = 0.$$

It follows that v is a constant function and hence $v \equiv 0$. By the implicit function theorem, there exists a small enough $s > 0$ such that the equation (4) has a solution for any $\lambda \in (0, s)$. Indeed, let $u = tv + u_0$, $v \not\equiv 0$ on V , we consider $G(\lambda, t) = \Delta u + \kappa - K_\lambda e^u$. It is easy to see that $G(0, 0) = 0$, $G(\lambda, t)$ and $\partial_t G(\lambda, t) = \Delta v - 2v(K + \lambda)e^{2(u_0 + tv)}$ are continuous on any domain $D \subset \mathbb{R}^2$. Furthermore, $\partial_t G(0, 0) = \Delta v - 2Ke^{2u_0}v \not\equiv 0$ unless $v \equiv 0$ on V . Therefore, by the implicit function theorem, there exists $s > 0$ such that $t = g(\lambda)$ and $G(\lambda, g(\lambda)) = 0$ for any $\lambda \in (0, s)$. In other words, $u_\lambda = g(\lambda)v + u_0$, $\forall \lambda \in (0, s)$ is the solution of (4). Define

$$\lambda^* = \sup\{s : \text{the equation (4) has a solution for any } \lambda \in (0, s)\}.$$

One can see that $\lambda^* \leq -\min_V K$. For otherwise, $K_\lambda = K + \lambda > 0$ for some $\lambda < \lambda^*$. Adding up the equation (4) for all $x \in V$, we have

$$0 > \int_V \kappa d\mu = \int_V K_\lambda e^{2u} d\mu > 0,$$

which is impossible. In conclusion, we have $0 < \lambda^* \leq -\min_V K$.

Proof of (2) in Theorem 2. We separate the proof into the following three steps.

Step 1. *The existence of the upper and lower solution of (4).*

Take λ_1 with $\lambda < \lambda_1 < \lambda^*$, let u_{λ_1} be a solution of (4) at λ_1 . It is easily seen that $\psi = u_{\lambda_1}$ is a strict upper solution of (4) at λ , namely

$$\Delta \psi + \kappa - K_\lambda e^{2\psi} > 0.$$

Let v be the solution to the following equation

$$\Delta v = -\kappa + \frac{1}{\mu(V)} \int_V \kappa d\mu. \quad (15)$$

The existence of solution to (15) was proved in [7]. Set $\varphi = v - s$, where s is a sufficiently large constant such that $\varphi < \psi$ on V and

$$\Delta \varphi + \kappa - K_\lambda e^{2\varphi} = \frac{1}{\mu(V)} \int_V \kappa d\mu - K_\lambda e^{2v-2s} < 0$$

since $\int_V \kappa d\mu < 0$. Therefore, φ is a strict lower solution of (4). Let $[\varphi, \psi]$ be the order interval defined by

$$[\varphi, \psi] = \{u \in V^{\mathbb{R}} : \varphi \leq u \leq \psi \text{ on } V\}.$$

The upper and lower-solution method (Lemma 6) asserts that (4) has a solution $u_\lambda \in [\varphi, \psi]$ on V .

Step 2. u_λ can be chosen as a strict local minimum of E_λ .

Let $f_\lambda(x, t) = ct - \kappa(x) + K_\lambda(x)e^{2t}$, where c is sufficiently large such that $f_\lambda(x, t)$ is increasing in $t \in [-A, A]$, A is a constant such that $-A \leq \varphi < \psi \leq A$ on V . Let $F_\lambda(x, u(x)) = \int_0^{u(x)} f_\lambda(x, t)dt$. It is easy to rewrite $E_\lambda(u)$ as

$$E_\lambda(u) = \int_V (|\nabla u|^2 + cu^2)d\mu - 2 \sum_{x \in V} \mu(x)F_\lambda(x, u(x)) - \int_V K_\lambda d\mu.$$

It is obvious that E_λ is bounded from below on $[\varphi, \psi]$. Therefore, we denote

$$a := \inf_{u \in [\varphi, \psi]} E_\lambda(u).$$

Taking a function sequence $u_j \subset [\varphi, \psi]$ such that $E_\lambda(u_j) \rightarrow a$ as $j \rightarrow \infty$. From it, we can get that u_j is bounded in $W^{1,2}(V)$, and thus up to subsequence, u_j converges to some u_λ in $W^{1,2}(V)$ and ℓ_μ^q for any $q \geq 1$, and e^{2u_j} converges to e^{2u_λ} in ℓ_μ^1 . Hence

$$E_\lambda(u_\lambda) = \inf_{u \in [\varphi, \psi]} E_\lambda(u).$$

As a consequence, u_λ satisfies the Euler-Lagrange equation

$$\Delta u_\lambda(x) + cu_\lambda(x) = f_\lambda(x, u_\lambda(x)).$$

From it, one can conclude that

$$\varphi(x) < u_\lambda(x) < \psi(x), \forall x \in V. \quad (16)$$

Indeed, noting that $f_\lambda(x, t)$ is increasing with respect to $t \in [-A, A]$, we have

$$\Delta\varphi(x) + c\varphi(x) \leq f_\lambda(x, \varphi(x)) \leq f_\lambda(x, \psi(x)) \leq \Delta\psi(x) + c\psi(x).$$

One can conclude (16) by the strong maximum principle (Lemma 4), and the fact $\varphi < \psi$. For any $h \in W^{1,2}(V)$, we define a function $\eta(t) = E_\lambda(u_\lambda + th)$, $t \in \mathbb{R}$. There holds $\varphi \leq u_\lambda + th \leq \psi$ for sufficiently small $|t|$. Since u_λ is a minimum of E_λ on (φ, ψ) , we have $\eta'(0) = dE_\lambda(u_\lambda)(h) = 0$ and $\eta''(0) = d^2E_\lambda(u_\lambda)(h, h) \geq 0$. Furthermore, there exists a constant $C > 0$ such that

$$d^2E_\lambda(u_\lambda)(h, h) \geq C\|h\|_{W^{1,2}(V)}, \quad \forall h \in W^{1,2}(V), \quad (17)$$

which implies u_λ is strict local minimum of E_λ on $W^{1,2}(V)$. It remains to prove (17). We first denote

$$\theta := \inf_{\|h\|_{W^{1,2}(V)}=1} d^2E_\lambda(u_\lambda)(h, h),$$

which is nonnegative. It is sufficient to prove $\theta > 0$, (17) follows. Suppose $\theta = 0$, we claim that there exists some \tilde{h} with $\|\tilde{h}\|_{W^{1,2}(V)} = 1$ such that $d^2E_\lambda(u_\lambda)(\tilde{h}, \tilde{h}) = 0$. To see this, let h_j be a function sequence satisfying $\|h_j\|_{W^{1,2}(V)} = 1$ for all j and $d^2E_\lambda(u_\lambda)(h_j, h_j) \rightarrow 0$ as $j \rightarrow \infty$. Up to subsequence, $h_j \rightarrow \tilde{h}$ in $W^{1,2}(V)$ from Lemma 1, and confirms our claim. To put it another way, the functional $v \mapsto d^2E_\lambda(u_\lambda)(v, v)$ attains its minimum at $v = \tilde{h}$, it follows that $d^2E_\lambda(u_\lambda)(\tilde{h}, v) = 0$ for all $v \in W^{1,2}(V)$. Hence, \tilde{h} is a solution of the following equation

$$\Delta h = 2K_\lambda e^{2u_\lambda} h, \quad \lambda \in (0, \lambda^*). \quad (18)$$

It is easy to see that \tilde{h} is not a constant. For otherwise (18) yields

$$0 > \int_V \kappa d\mu = \int_V K_\lambda e^{2u_\lambda} d\mu = 0,$$

which is impossible. Multiplying (18) by \tilde{h}^3 , we obtain

$$\begin{aligned} d^4 E_\lambda(u_\lambda)(\tilde{h}, \tilde{h}, \tilde{h}, \tilde{h}) &= -16 \int_V K_\lambda e^{2u_\lambda} \tilde{h}^4 d\mu \\ &= -8 \int_V \tilde{h}^3 \Delta \tilde{h} d\mu \\ &= -8 \sum_{x,y \in V} \omega_{xy} (\tilde{h}^3(x) - \tilde{h}^3(y)) (\tilde{h}(x) - \tilde{h}(y)) \\ &= -8 \sum_{x,y \in V} \omega_{xy} (\tilde{h}^2(x) + \tilde{h}(x)\tilde{h}(y) + \tilde{h}^2(y)) (\tilde{h}(x) - \tilde{h}(y))^2 < 0. \end{aligned}$$

The last inequality is due to the fact $a^2 + ab + b^2 > 0$ for any $a, b \in \mathbb{R}$ satisfying $ab \neq 0$. Since $d^2 E_\lambda(u_\lambda + t\tilde{h})(\tilde{h}, \tilde{h})$ attains its minimum at $t = 0$, we have $d^3 E_\lambda(u_\lambda)(\tilde{h}, \tilde{h}, \tilde{h}) = 0$, which together with $dE_\lambda(u_\lambda)(\tilde{h}) = 0$ and $d^2 E_\lambda(u_\lambda)(\tilde{h}, \tilde{h}) = 0$ leads to

$$E_\lambda(u_\lambda + \epsilon\tilde{h}) = E_\lambda(u_\lambda) + \frac{\epsilon^4}{24} d^4 E_\lambda(u_\lambda)(\tilde{h}, \tilde{h}, \tilde{h}, \tilde{h}) + 0(\epsilon^5) < E_\lambda(u_\lambda) \quad (19)$$

for small $\epsilon > 0$. Let ϵ small enough such that $\varphi \leq u_\lambda + \epsilon\tilde{h} \leq \psi$, thus by (19),

$$E_\lambda(u_\lambda + \epsilon\tilde{h}) < E_\lambda(u_\lambda),$$

which contradicts the fact that u_λ is the minimum of E_λ on $[\varphi, \psi]$. Therefore $\theta > 0$, which concludes (17).

Step 3. *The second solution of (4) is given by the mountain-pass theorem.*

We shall use the mountain-pass theorem due to Ambrosetti and Rabinowitz [1], which reads as follows: *Let $(X, \|\cdot\|)$ be a Banach space, $J \in C^1(X, \mathbb{R})$, $e_0, e \in X$ and $r > 0$ be such that $\|e - e_0\| > r$ and*

$$b := \inf_{\|u - e_0\| = r} J(u) > J(e_0) \geq J(e).$$

If J satisfies the $(PS)_c$ condition with $c := \inf_{\gamma \in \Gamma} \max_{t \in [0,1]} J(\gamma(t))$, where

$$\Gamma := \{\gamma \in C([0, 1], X) : \gamma(0) = e_0, \gamma(1) = e\},$$

then c is a critical value of J . In our case, $W^{1,2}(V)$ is a Banach space, and $E_\lambda : W^{1,2}(V) \rightarrow \mathbb{R}$ is a smooth functional.

Since u_λ is a strict local minimum of E_λ on $W^{1,2}(V)$, there exists a small enough number $r > 0$ such that

$$\inf_{\|u - u_\lambda\|_{W^{1,2}(V)} = r} E_\lambda(u) > E_\lambda(u_\lambda). \quad (20)$$

Moreover, for any $\lambda > 0$, E_λ has no lower bound on $W^{1,2}(V)$, namely, there exists $v \in W^{1,2}(V)$ such that

$$E_\lambda(v) < E_\lambda(u_\lambda), \quad \|v - u_\lambda\|_{W^{1,2}(V)} > r. \quad (21)$$

To see this, we set $V_\varepsilon = \{x \in V : K_\lambda(x) > \varepsilon\}$ for small $\varepsilon > 0$. Note that V_ε is nonempty since $\max_V K = 0$ and $\lambda > 0$. Let $f \in W^{1,2}(V)$ be a function which equals to 1 in V_ε and vanishes on V/V_ε , then

$$\begin{aligned} E_\lambda(tf) &= t^2 \int_V |\nabla f|^2 d\mu + t \int_{V_\varepsilon} \kappa d\mu - \int_{V_\varepsilon} K_\lambda e^{2t} d\mu - \int_{V/V_\varepsilon} K_\lambda d\mu \\ &\leq At^2 + Bt + C - \varepsilon \mu(V_\varepsilon) e^{2t} \rightarrow -\infty, \quad t \rightarrow +\infty. \end{aligned}$$

In view of (20), (21) and Lemma 5, the mountain pass theorem of Ambrosetti and Rabinowitz gives another critical point u^λ of E_λ other than u_λ . In particular,

$$E_\lambda(u^\lambda) = \min_{\gamma \in \Gamma} \max_{u \in \gamma} E_\lambda(u),$$

where $\Gamma = \{\gamma \in C([0, 1], W^{1,2}(V)) : \gamma(0) = u_\lambda, \gamma(1) = v\}$, and $dE_\lambda(u^\lambda)(h) = 0$ for any $h \in W^{1,2}(V)$. \square

3.3. Solvability at λ^* .

For any $\lambda, 0 < \lambda < \lambda^*$, let u_λ be the local minimum of E_λ obtained in the previous subsection. That is, u_λ is the solution of (4), and

$$d^2 E_\lambda(u_\lambda)(h, h) = 2 \int_V (|\nabla h|^2 - 2K_\lambda e^{2u_\lambda} h^2) d\mu \geq 0, \quad \forall h \in W^{1,2}(V). \quad (22)$$

Proof of (3) in Theorem 2. The crucial point in this proof is to show that u_λ is uniformly bounded in $W^{1,2}(V)$ as $\lambda \rightarrow \lambda^*$. If it is true, then up to subsequence, u_λ converges to some u in $W^{1,2}(V)$, and u is the solution of

$$\Delta u + \kappa - K_{\lambda^*} e^{2u} = 0.$$

Hence, we aim to prove the $W^{1,2}(V)$ boundedness of u_λ . To this end, we divide the proof into the following three steps.

Step 1. *There exists a constant $C > 0$ such that*

$$u_\lambda \geq -C \quad \text{on } V \quad (23)$$

uniformly for any $0 < \lambda < \lambda^$.*

Let v satisfy (15), and $\varphi_s = v - s$ for $s > 0$. Then for sufficient large s , say $s \geq s_0$, φ_s is a continuous family with respect to s of strict lower solution of (4) at $\lambda = 0$, i.e.

$$\Delta \varphi_s + \kappa - K e^{2\varphi_s} < 0.$$

It is clear that φ_s is also a strict lower solution of (4) at $\lambda \in (0, \lambda^*)$ for any $s \in [s_0, \infty)$. Now, we prove that $u_\lambda \geq \varphi_{s_0}$, and thus (23) holds. For otherwise, we can find for some $s \in (s_0, \infty)$

$$u_\lambda \geq \varphi_s \quad \text{on } V, \quad \text{and } u_\lambda(\tilde{x}) = \varphi_s(\tilde{x}) \quad \text{for some } \tilde{x} \in V.$$

It follows from the strong maximum principle (Lemma 4) that $u_\lambda \equiv \varphi_s$, which contradicts that φ_s is strict low solution of (4) at $\lambda \in [0, \lambda^*)$.

Step 2. *The set $V_{\lambda^*}^- = \{x \in V : K_{\lambda^*}(x) < 0\}$ is not empty.*

From the case (1) in Theorem 2, there exists unique solution w_0 of the equation

$$\Delta w + \kappa + e^{2w} = 0.$$

Together with the solution u_λ to the equation (4) at λ , and let $v_\lambda = u_\lambda - w_0$, we have

$$\Delta v_\lambda - K_\lambda e^{2u_\lambda} - e^{2w_0} = 0.$$

Multiplying the above equation by e^{-2v_λ} and intergrading by parts, one has

$$\int_V K_\lambda e^{w_0} d\mu = \int_V e^{-2v_\lambda} \Delta v_\lambda d\mu - \int_V e^{-2(u_\lambda - 2w_0)} d\mu \leq 0.$$

Indeed, by Green formula

$$\int_V e^{-2v_\lambda} \Delta v_\lambda d\mu = \sum_{x,y \in V} \omega_{xy} (v_\lambda(x) - v_\lambda(y)) (e^{-2v_\lambda(x)} - e^{-2v_\lambda(y)}) \leq 0$$

since $(v_\lambda(x) - v_\lambda(y))(e^{-2v_\lambda(x)} - e^{-2v_\lambda(y)}) \leq 0$ for any $x, y \in V$. Therefore

$$\int_V K_{\lambda^*} e^{w_0} d\mu = \lim_{\lambda \rightarrow \lambda^*} \int_V K_\lambda e^{w_0} d\mu \leq 0.$$

Suppose that $K_{\lambda^*} \geq 0$, thus $K_{\lambda^*} \equiv 0$. This contradicts the assumption that K_{λ^*} is not a constant.

Step 3. u_λ is uniformly bounded in $W^{1,2}(V)$ as $\lambda \rightarrow \lambda^*$.

Fixing $x_0 \in V_{\lambda^*}^-$, we set $\rho \in V^{\mathbb{R}}$ to be a function which vanishes besides x_0 and $\rho(x_0) < 0$. Consider the equation

$$\Delta w + \kappa - \rho e^{2w} = 0,$$

which always has the unique solution from the case (1) in Theorem 2, set w_1 . As above, the function $v_\lambda = u_\lambda - w_1$ satisfies the equation

$$\Delta v_\lambda + \rho e^{2w_1} - K_\lambda e^{2(v_\lambda + w_1)} = 0. \quad (24)$$

Multiplying (24) by e^{2v_λ} and integrating by parts over V gives

$$\int_V e^{2v_\lambda} \Delta v_\lambda d\mu + \int_V \rho e^{2(v_\lambda + w_1)} d\mu - \int_V K_\lambda e^{2(2v_\lambda + w_1)} d\mu = 0. \quad (25)$$

Utilizing the fact that

$$(e^{2a} - e^{2b})(a - b) \geq (e^a - e^b)^2, \quad a, b \in \mathbb{R},$$

one can estimate the first term in (25) as follows.

$$\begin{aligned} \int_V e^{2v_\lambda} \Delta v_\lambda d\mu &= \frac{1}{2} \sum_{x,y \in V} \omega_{xy} (e^{2v_\lambda(x)} - e^{2v_\lambda(y)}) (v_\lambda(x) - v_\lambda(y)) \\ &\geq \frac{1}{2} \sum_{x,y \in V} \omega_{xy} (e^{v_\lambda(x)} - e^{v_\lambda(y)})^2 \\ &= \int_V |\nabla e^{v_\lambda}|^2 d\mu. \end{aligned}$$

Inserting this estimate into (25), we obtain

$$\int_V |\nabla e^{v_\lambda}|^2 d\mu + \int_V \rho e^{2(v_\lambda+w_1)} d\mu - \int_V K_\lambda e^{2(2v_\lambda+w_1)} d\mu \leq 0. \quad (26)$$

On the other hand, let $h = e^{v_\lambda}$ in (22), yields

$$\int_V |\nabla e^{v_\lambda}|^2 d\mu - 2 \int_V K_\lambda e^{2(2v_\lambda+w_1)} d\mu \geq 0.$$

Together with (26), we have

$$\int_V |\nabla e^{v_\lambda}|^2 d\mu \leq -2 \int_V \rho e^{2(v_\lambda+w_1)} d\mu = -2\rho(x_0) e^{2u_\lambda(x_0)}. \quad (27)$$

One may derive that $u_\lambda(x_0)$ is the uniform bound, i.e. there exists a constant $C > 0$ such that

$$e^{2u_\lambda(x_0)} \leq C.$$

Indeed, if $u_\lambda(x_0) \leq 0$, the above inequality is obvious; if $u_\lambda(x_0) > 0$, as in the proof of Lemma 5, see (11), one can get the boundedness of $u_\lambda(x_0)$ as well. Together with (27), yields

$$\int_V |\nabla e^{v_\lambda}|^2 d\mu \leq C'. \quad (28)$$

Next, we *claim* that e^{v_λ} and thus e^{u_λ} is uniformly bounded in ℓ_μ^2 . Suppose not, we may assume that $\|e^{v_\lambda}\|_{\ell_\mu^2} \rightarrow \infty$ as $\lambda \rightarrow \lambda^*$. Let

$$w_\lambda = \frac{e^{v_\lambda}}{\|e^{v_\lambda}\|_{\ell_\mu^2}},$$

then $\|w_\lambda\|_{\ell_\mu^2} = 1$, and $\|\nabla w_\lambda\|_{\ell_\mu^2} \rightarrow 0$ from (28). It follows that w_λ converges to a constant in $W^{1,2}(V)$. From (3.3), we have $w_\lambda(x_0) \rightarrow 0$, and hence $w_\lambda \equiv 0$ on V , which contradicts $\|w_\lambda\|_{\ell_\mu^2} = 1$ and then confirms our claim. Since u_λ is bounded below by (23) in Step 1, one has the ℓ_μ^2 -boundedness of u_λ , and thus the $W^{1,2}(V)$ -boundedness of u_λ . \square

3.4. No solution when $\lambda > \lambda^*$.

Proof of this case is a consequence of the upper and lower solutions principle, as follows.

Proof of (4) in Theorem 2. Let u_{λ_1} be the solution of (4) at some $\lambda_1 > \lambda^*$. For any $0 < \lambda < \lambda_1$, u_{λ_1} is an upper solution of (4) at λ . Indeed,

$$\Delta u_{\lambda_1} + \kappa - (K + \lambda) e^{2u_{\lambda_1}} = (\lambda_1 - \lambda) e^{2u_{\lambda_1}} > 0.$$

From (15), it is easy to get a lower bound solution φ of (4) at λ such that $\varphi \leq u_{\lambda_1}$. By the upper and lower solutions principle, there exists a solution of (4) at λ , which contradicts the definition of λ^* . \square

References

References

- [1] A. Ambrosetti, P. Rabinowitz, Dual variational methods in critical point theory and applications, *J. Funct. Anal.* 14 (1973) 349-381.
- [2] F. Borer, L. Galimberti, M. Struwe, “Large” conformal metrics of prescribing Gauss curvature on surfaces of high genus, *Comment. Math. Helv.* 90 (2015) 407-428.
- [3] F. Camilli, C. Marchi, A note on Kazdan-Warner equation on networks, arXiv:1909.08472.
- [4] W. Ding, J. Liu, A note on the problem of prescribing Gaussian curvature on surfaces, *Trans. Amer. Math. Soc.* 347 (1995) 1059-1066.
- [5] H. Ge, Kazdan-Warner equation on graph in the negative case, *J. Math. Anal. Appl.* 453 (2017) 1022-1027.
- [6] H. Ge, W. Jiang, Kazdan-warner equation on infinite graphs, *J. Korean Math. Soc.* 55 (2018) 1091-1101.
- [7] A. Grigor’yan, Y. Lin, Y. Yang, Kazdan-Warner equation on graph, *Calc. Var. Partial Differential Equations* 55 (2016), no. 4, Art. 92, 13 pp.
- [8] A. Grigor’yan, Y. Lin, Y. Yang, Yamabe type equations on graphs. *J. Differential Equations* 261 (2016) 4924-4943.
- [9] A. Grigor’yan, Y. Lin, Y. Yang, Existence of positive solutions to some nonlinear equations on locally finite graphs, *Sci. China Math.* 60 (2017) 1311-1324.
- [10] X. Han, M. Sha, L. Zhao, Existence and convergence of solutions for nonlinear biharmonic equations on graphs, to appear in *J. Differential Equations*, <https://doi.org/10.1016/j.jde.2019.10.007>
- [11] M. Keller, M. Schwarz, The Kazdan-Warner equation on canonically compactifiable graphs, *Calc. Var. Partial Differential Equations* 57 (2018), no. 2, Art. 70, 18 pp.
- [12] J. Kazdan, F. Warner, Curvature functions for compact 2-manifolds, *Ann. of Math.* 99 (1974) 14-47.
- [13] J. Kazdan, F. Warner, Curvature functions for open 2-manifolds, *Ann. of Math.* 99 (1974) 203-219.
- [14] Y. Lin, Y. Wu, The existence and nonexistence of global solutions for a semilinear heat equation on graphs, *Calc. Var. Partial Differential Equations* 56 (2017), no. 4, Art. 102, 22 pp.
- [15] M. Struwe, Critical points of embeddings of $H_0^{1,n}$ into Orlicz spaces, *Ann. Inst. H. Poincaré Anal. Non Linéaire* 5 (1988) 425-464.
- [16] M. Struwe, The existence of surfaces of constant mean curvature with free boundaries, *Acta Math.* 160 (1988) 19-64.
- [17] Y. Yang, X. Zhu, Prescribing Gaussian curvature on closed Riemann surface with conical singularity in the negative case, *Ann. Acad. Sci. Fenn. Math.* 44 (2019) 167-181.