

A NOTE ON UNCONDITIONALLY CONVERGENT SERIES IN A COMPLETE TOPOLOGICAL RING

ALEX RAVSKY

ABSTRACT. We answer a question concerning classes of complete topological rings where unconditionally convergent series have a special property.

In this note we answer Question 1 posed by a user dch at Mathematics Stack Exchange a few months ago [3]. In order to formulate it, we introduce the following notion. A topological ring R is a *dch ring*, if for any unconditionally convergent series $\sum_{i=1}^{\infty} a_i$ in R and any neighborhood S of the additive identity 0 of R there exists a neighborhood S' of 0 such that for any finite subset F of S' and any sequence (f_i) of elements of F , we have that $\sum_{i=1}^{\infty} f_i a_i$ converges to a point in S .

Question 1. *Are there well-known classes (e.g. locally compact, locally connected, regular) of complete Hausdorff topological rings which are dch rings?*

We show that a topological ring which is a Banach space (and so a connected complete metric space) can fail to be a dch ring, see Proposition 4. On the other hand, a topological ring R is a dch ring provided R is locally compact Hausdorff, see Proposition 7 or R has a base at the zero consisting of open ideals and the additive topological group of R is sequentially complete, see Proposition 3.

All groups explicitly considered in this paper are supposed to be Abelian, add all topological groups (including additive groups of topological rings) are Hausdorff. An endomorphism of a group G is a homomorphism from G to itself. A series $\sum_{i=1}^{\infty} a_i$ in a topological group, is *unconditionally convergent*, if there exists an element $a \in G$ such that for each permutation σ of \mathbb{N} , a series $\sum_{i=1}^{\infty} a_{\sigma(i)}$ converges to a .

A non-negative function $\|\cdot\|$ defined in a group G is called a *norm*, provided for each $g, h \in G$, $\|g\| = 0$ iff $g = 0$, $\|-g\| = \|g\|$, and $\|g+h\| \leq \|g\| + \|h\|$. It is well known and easy to show that in this case a function d such that $d(g, h) = \|g - h\|$ for each $g, h \in G$, is a metric on G and the group G endowed with the metric d is a topological group. A series $\sum_{i=1}^{\infty} a_i$ on this group (G, d) is *absolutely convergent*, if a series $\sum_{i=1}^{\infty} \|a_i\|$ converges in \mathbb{R} . It is well known and easy to show (see, for instance, [2, Exercise 1.36]) that each absolutely convergent series on (G, d) is unconditionally convergent iff (G, d) is complete.

Whether unconditional convergence implies absolute convergence is a more delicate issue. If $G = \mathbb{R}^n$ with the standard Euclidean norm, then by the Riemann Rearrangement Theorem unconditional convergence implies absolute convergence. On the other hand, by Dvoretzky-Rogers theorem [4], in any infinite dimensional real Banach space there exists a series which is unconditionally convergent but not absolutely convergent.

1991 *Mathematics Subject Classification.* 13J10, 22A10, 54H11.

Key words and phrases. topological ring, unconditional convergence, locally compact topological ring, locally compact Abelian topological group.

By Proposition 1.27 from [2], convergence in a complete, non-Archimedean normed group G is much simpler. If (a_i) is a sequence in G then the series $\sum_{i=1}^{\infty} a_i$ is unconditionally convergent iff it is convergent iff (a_i) converges to the identity of G .

A sequence (x_n) on a topological group G is *Cauchy* if for every neighborhood U of the identity of G there is natural n such that $x_i - x_j \in U$, for each $i, j \geq n$. A topological group G is *sequentially complete*, if each Cauchy sequence on G converges.

Lemma 2. *Let G be a sequentially complete topological group, $\sum_{i=1}^{\infty} a_i$ be an unconditionally convergent in G series, and $(f_i)_{i=1}^{\infty}$ be a sequence of continuous endomorphisms of G such that a set F of values of (f_i) is finite. Then a series $\sum_{i=1}^{\infty} f_i(a_i)$ converges.*

Proof. Let $f \in F$ be any element. For each natural i put $x_i^f = \sum_{k \leq i, f_k = f} a_i$ (we put that the sum over an empty set equals the identity of G). It is easy to show that the unconditional convergence of the series $\sum_{i=1}^{\infty} a_i$, implies that a sequence (x_i^f) is Cauchy. Since the group G is sequentially complete, the sequence (x_i^f) converges to a point $x^f \in G$. Let n be any natural number. Since each map $f \in F$ is a homomorphism, $\sum_{i=1}^n f_i(a_i) = \sum_{f \in F} f(x_n^f)$. Since each map $f \in F$ is continuous and the sequence (x_i^f) converges to x^f , the sequence $\left(\sum_{f \in F} f(x_n^f) \right)_{n \in \mathbb{N}}$ converges to $\sum_{f \in F} f(x^f)$. \square

Proposition 3. *Each topological ring R which has a base at the zero consisting of open ideals, and whose additive topological group is sequentially complete, is a dch ring.*

Proof. Let $\sum_{i=1}^{\infty} a_i$ be any convergent series in R and S be any neighborhood of the zero 0 of R . Let S' be an arbitrary open ideal contained in S . Since each open subgroup of a topological group is closed in it, S' is closed in R . Let (f_i) be any sequence of elements of S' and T be any open ideal in R . Since the sequence (a_i) converges to 0, there exists N such that $a_n \in T$ for each $n \geq N$. Since T is ideal, it follows that $\sum_{i=m}^n f_i a_i \in T$ for each $m, n \geq N$. Since R has a base at the zero consisting of open ideals, $(\sum_{i=1}^n f_i a_i)$ is a Cauchy sequence. Since the additive group of R is sequentially complete, the series $\sum_{i=1}^{\infty} f_i a_i$ converges to a point $a \in R$. Since S' is an ideal, for any natural n , $\sum_{i=1}^n f_i a_i \in S'$, so we have $a \in \overline{S'} = S' \subset S$. \square

Let $p \in [1, \infty]$ be any number and R_p be a ring of bounded linear operators on a normed space $(\ell_p, \|\cdot\|)$ of sequences $x = (x_n)_{n \in \mathbb{N}}$ of real numbers such that $\|x\|^p = \sum |x_n|^p < \infty$ for $p < \infty$ and $\|x\| = \sup |x_n| < \infty$ for $p = \infty$, such that $(ab)(x) = b(a(x))$ for each $a, b \in R_p$ and $x \in \ell_p$. It is easy to check that R_p endowed with a topology generated by an operator norm $\|a\| = \sup\{\|a(x)\| : x \in \ell_p \text{ with } \|x\| \leq 1\}$ is a topological ring and a Banach space.

Proposition 4. *For any $p \in [1, \infty]$, the ring R_p is non-dch.*

Proof. For each i and each $x = (x_n) \in \ell_p$ let $a_i(x)$ has the only non-zero i -th coordinate, which equals x_i/i . Then a series $\sum_{i=1}^{\infty} a_i$ unconditionally converges. Now let S be a unit ball centered at the zero 0 of R_p and S' be any open neighborhood of 0. There exists $\varepsilon > 0$ such that S' contains a closed ball of radius ε centered at 0. For each i and each $x = (x_n) \in \ell_p$ let $f'_i(x)$ has the only non-zero n -th coordinate, which equals εx_1 . Since the harmonic series $\sum_{n=1}^{\infty} 1/n$ diverges, there exists N such that $\varepsilon \sum_{n=1}^N 1/n > 1$. Put $f_i = f'_i$, if $i \leq N$ and $f_i = 0$, otherwise. Then $f'_i \in S'$ for each i . Let $x_0 = (1, 0, 0, \dots) \in \ell_p$. Then

$\|x_0\|_p = 1$, but

$$\left\| \left(\sum_{i=0}^{\infty} f_i a_i \right) (x_0) \right\| = \varepsilon \sum_{n=1}^N 1/n > 1.$$

So $\sum_{i=0}^{\infty} f_i a_i \notin S$. \square

Let X be a non-empty set and $\ell_{\infty}(X)$ be a linear space of bounded real-valued functions on X endowed with the supremum norm $\|a\| = \sup_{x \in X} |a(x)|$ for each $a \in \ell_{\infty}(X)$. Then X is a Banach space and a topological ring with respect to the coordinatewise addition and multiplication.

Proposition 5. *For each non-empty set X , $R = \ell_{\infty}(X)$ is a dch ring.*

Proof. Let $\sum_{i=1}^{\infty} a_i$ be any unconditionally convergent series in R and S be any neighborhood of 0 of R . There exists $\varepsilon > 0$ such that $0 \in B_{\varepsilon} = \{b \in R : \|b\| \leq \varepsilon\} \subset S$. Since for each $x \in X$ a series $\sum_{i=1}^{\infty} a_i(x)$ is unconditionally convergent, it is absolutely convergent, that is $a(x) = \sum_{i=1}^{\infty} |a_i(x)| < \infty$. If $A = \sup_{x \in X} a(x) = \infty$ then we can inductively construct a sequence (x_n) of elements of X and an increasing sequence (m_n) of natural numbers and a permutation σ of \mathbb{N} such that a sequence $(|\sum_{i=1}^{m_n} a_{\sigma(i)}(x_n)|)$ is unbounded, which contradicts the unconditional convergence of $\sum_{i=1}^{\infty} a_i$. Pick $\delta > 0$ such that $\delta A \leq \varepsilon$. Then $S' = B_{\delta} = \{b \in R : \|b\| \leq \delta\}$ is a neighborhood of 0 of R . Let (f_i) be any sequence of elements of S' such that a set of values of (f_i) is finite. By Lemma 2, a series $\sum_{i=1}^{\infty} f_i a_i$ converges to some point $a \in R$. Since for each $x \in X$ and each natural n we have

$$\left\| \sum_{i=1}^n f_i(x) a_i(x) \right\| \leq \sum_{i=1}^n \|f_i(x)\| \|a_i(x)\| \leq \sum_{i=1}^n \delta \|a_i(x)\| \leq \delta \sum_{i=1}^{\infty} \|a_i(x)\| \leq \delta A \leq \varepsilon,$$

we see that $\sum_{i=1}^{\infty} f_i a_i \in B_{\varepsilon}$ and thus $\in \overline{B_{\varepsilon}} = B_{\varepsilon} \subset S$. \square

Recall that a *character* of a topological group G is a continuous homomorphism from G to a group $\mathbb{T} = \{z \in \mathbb{C} : |z| = 1\}$, endowed with the multiplication and topology inherited from \mathbb{C} .

Lemma 6. *Let G be a locally compact topological group. Let E be a set of endomorphisms of G containing the zero endomorphism 0_E and endowed with a topology such that the evaluation map from $E \times G$ to G , $(f, x) \mapsto f(x)$ for each $f \in E$ and $x \in G$, is continuous. Let a series $\sum_{i=1}^{\infty} a_i$ of elements a_i of G unconditionally converges. Then for any neighborhood S of the identity of G there exists an open neighborhood S' of 0_E such that for any sequence (f_i) of elements of S' and any natural n , $\sum_{i=1}^n f_i(a_i) \in S$.*

Proof. Shrinking the neighborhood S , if needed, we can assume that its closure \overline{S} is compact. Since the series $\sum_{i=1}^{\infty} a_i$ converges, all but finitely many a_i 's belong to S . Thus a set $\overline{S} \cup \{a_i\}$ is compact. Let G' be the smallest subgroup of G containing $\overline{S} \cup \{a_i\}$. Since $G' \supset S$, G' is an open subgroup of G . By Example 69 from [5], the group G' is topologically isomorphic to a product $H \times \mathbb{Z}^p \times \mathbb{R}^q$, where H is a compact group, \mathbb{Z} and \mathbb{R} are the groups of integer and real numbers, respectively, endowed with the usual topologies, and p, q are non-negative integers. By [5, §34.C], for each element x of H there exists a character χ from H to \mathbb{T} such that $\chi(x) \neq 0$. It follows that H can be topologically isomorphically embedded into a power \mathbb{T}^{κ} for some cardinal κ . So there exists an isomorphic topological embedding $i : G' \rightarrow \mathbb{T}^{\kappa} \times \mathbb{R}^s$ for a non-negative integer $s = p + q$.

Thus there exist $0 < \varepsilon \leq \pi$ and a finite set Γ of continuous homomorphisms (which are compositions of i and projections on the factors) from G' to a group \mathbb{K} (which is \mathbb{R} or \mathbb{T}) such that $S \supset S_0 = \{x \in G' : \|\gamma(x)\| \leq \varepsilon \text{ for each } \gamma \in \Gamma\}$, where $\|g\| = |g|$, if $g \in \mathbb{R}$ and $\|g\| = |\arg g| \leq \pi$, if $g \in \mathbb{T}$. Remark that for each $g, h \in \mathbb{K}$ we have $\|g + h\| \leq \|g\| + \|h\|$ and if $\|g\| \leq \pi/2$ then $\|g + g\| = 2\|g\|$. For each $x \in G'$ put $\|x\|_\Gamma = \max_{\gamma \in \Gamma} \|\gamma(x)\|$.

Let γ be any element of Γ . Since the series $\sum_{i=1}^\infty a_i$ unconditionally converges and γ is continuous, a series $\sum_{i=1}^\infty \gamma(a_i)$ unconditionally converges too. This easily follows that a series $\sum_{i=1}^\infty \|\gamma(a_i)\|$ converges. Since the set Γ is finite,

$$M = 1 + \frac{1}{\varepsilon} \sum_{i=1}^\infty \|a_i\|_\Gamma \leq 1 + \frac{1}{\varepsilon} \sum_{i=1}^\infty \sum_{\gamma \in \Gamma} \|\gamma(a_i)\| < \infty.$$

Pick $0 < \delta \leq \varepsilon/(2M)$.

Since S_0 is a closed subset of a compact space \bar{S} , it is compact. Let $x \in S_0$ be any element. Since $0 = \|0_E(x)\|_\Gamma < \delta$ and both the evaluation map from $E \times G$ to G and $\|\cdot\|_\Gamma$ are continuous, there exist a neighborhood S''_x of 0_E and an open neighborhood O_x of x in S_0 such that $\|g\|_\Gamma < \delta$ for each $g \in S''_x(O_x)$, where $S''_x(O_x) = \{f(y) : f \in S''_x, y \in O_x\}$. Since the set S_0 is compact, there exists a finite subset F_0 of S_0 such that $S_0 \subset \bigcup \{O_x : x \in F_0\}$. Put $S'' = \bigcap \{S''_x : x \in F_0\}$. It follows $\|f(x)\|_\Gamma < \delta$ for each $f \in S''$ and $x \in S_0$.

Suppose to the contrary that there exist $f \in S''$ and $x \in S_0$ such that $M\|f(x)\|_\Gamma > \|x\|_\Gamma$. Since $0 < \|f(x)\|_\Gamma < \delta$, there exists the largest natural k such that $2^k\|f(x)\|_\Gamma < 2\delta \leq \varepsilon \leq \pi$. Then $2^k\|f(x)\|_\Gamma \geq \delta$. Let $2^k x$ be a sum of 2^k instances of x . It is easy to see that

$$\varepsilon \geq 2M\delta > M\|f(2^k x)\|_\Gamma = 2^k M\|f(x)\|_\Gamma > 2^k \|x\|_\Gamma.$$

It follows $2^k x \in S_0$ and so $\|f(2^k x)\|_\Gamma < \delta$, a contradiction.

Since the series $\sum_{i=1}^\infty a_i$ converges, a set $\{a_i\} \setminus S_0$ is finite. Since $\|0_E(x)\|_\Gamma = 0$ for each $x \in \{a_i\} \setminus S_0$ and both the evaluation map from $E \times G$ to G and $\|\cdot\|_\Gamma$ are continuous, there exists an open neighborhood $S' \subset S''$ of 0_E such that $\|f(x)\|_\Gamma < 2\delta < \|x\|_\Gamma/M$ for each $f \in S'$ and each $x \in \{a_i\} \setminus S_0$. Thus we have $\|f(a_i)\|_\Gamma \leq \|a_i\|_\Gamma/M$ for each $f \in S'$ each a_i .

Let (f_i) be any sequence of elements of S' , $x = \sum_{i=1}^n f_i(a_i)$. Then $f_i(a_i) \in G'$ for each i , so $x \in G'$ and

$$\|x\|_\Gamma \leq \sum_{i=1}^n \|f_i(a_i)\|_\Gamma \leq \sum_{i=1}^n \|a_i\|_\Gamma/M \leq \varepsilon.$$

Thus $x \in S_0 \subset S$. □

Proposition 7. *Each locally compact Hausdorff topological ring R is dch.*

Proof. Let $\sum_{i=1}^\infty a_i$ be any unconditionally convergent series in R and S be any neighborhood of the zero 0 of R . It is well known that each topological group is regular, so it suffices to show that there exists a neighborhood S' of 0 such that for any finite subset F of S' and any sequence (f_i) of elements of F , we have that $\sum_{i=1}^\infty f_i a_i$ converges to a point in \bar{S} . Since for each $f \in R$ a map $R \rightarrow R$, $x \mapsto fx$ for each $x \in R$ is a continuous homomorphism of the additive group of R , Lemma 6 implies that there exists a neighborhood S' of 0 such that for any sequence (f_i) of elements of S' and any natural n , $\sum_{i=1}^n f_i a_i \in S$. Since the group G is locally compact, it is sequentially complete, for instance, by Theorem 3.6.24 from [1], by Lemma 2, the series $\sum_{i=1}^\infty f_i a_i$ converges to a point $a \in R$. Since for any natural n , $\sum_{i=1}^n f_i a_i \in S$, we have $a \in \bar{S}$. □

Acknowledgment. The author thanks to Taras Banakh for the careful reading of the paper manuscript and its helpful discussion.

REFERENCES

- [1] Alexander Arhangel'skiĭ, Mikhail Tkachenko, *Topological Groups and Related Structures*, Atlantis Press, 2008.
- [2] Pete L. Clark, *Algebraic Number Theory II: Valuations, Local Fields and Adeles*, math.uga.edu/~pete/8410FULL.pdf.
- [3] dch, *Unconditional convergence of a sum of elements in a complete Hausdorff topological ring*, math.stackexchange.com/q/3702472.
- [4] A. Dvoretzky, C.A. Rogers, *Absolute and unconditional convergence in normed linear spaces*, Proc. Nat. Acad. Sci. USA **36** (1950), 192–197.
- [5] L.S. Pontrjagin, *Continuous groups*, 2nd ed., M., 1954 (in Russian).

ALEX RAVSKY: PIDSTRYHACH INSTITUTE FOR APPLIED PROBLEMS OF MECHANICS AND MATHEMATICS NATIONAL ACADEMY OF SCIENCES OF UKRAINE

E-mail address: alexander.ravsky@uni-wuerzburg.de