

EQUIPARTITION OF A SEGMENT

SERGEY AVVAKUMOV[♠] AND ROMAN KARASEV[♣]

ABSTRACT. We prove that, for any positive integer m , a segment may be partitioned into m possibly degenerate or empty segments with equal values of a continuous function f evaluated on segments, assuming that f may take positive and negative values, but its value on degenerate or empty segments is zero.

1. INTRODUCTION

The mathematical theory of fair division develops along two main lines of research. On the one hand, it looks for partitions of a body or measure into m pieces (the positive integer m is fixed) of a certain shape and equal in some sense, e.g. convex polygons of identical area. An early example is the ham sandwich theorem [17, 18] about equipartitioning several measures by hyperplanes. More recent examples are the Nandakumar and Ramana Rao conjecture [13] that every convex planar polygon can be partitioned into m convex polygons of equal area and perimeter, solved in [1], and higher-dimensional analogues of the Nandakumar and Ramana Rao problem that were solved in [4, 8] under the assumption that m is a prime power. See also [16] for a result in between the ham sandwich theorem and the Nandakumar and Ramana Rao problem.

On the other hand, the theory of fair division contributes key existence results to the concept of fairness favored by economists and many social scientists, known as Envy Freeness, that is, each one of m agents compares pieces of the partition in her own way. We look for an m -partition where each agent gets, in her own view, one of the best pieces. In this paper all the agents use the same utility function. For further references on it, see the foundational work of Gale [6] and popular reviews [3, 19].

Here we prove a topological property for partitions of a segment that we believe to be useful for both types of results just mentioned. There is a single agent who evaluates each subsegment $[a, b]$ of $[0, 1]$ by a continuous utility function $f(a, b)$ such that $f(a, a) = 0$ for all a . Apart from the continuity requirement (understood as continuity of a function in two real variables), f is very general, in particular it can take both positive and negative values. We show the existence of an m -partition of $[0, 1]$ into subsegments all of equal utility. This is a key ingredient used in the companion paper [5], where it implies the existence of a universal Fair Guarantee, which is a utility level that can be achieved simultaneously by any m agents, each with her own utility function.

Theorem 1.1. *Let \mathcal{I} be the space of possibly degenerate subsegments*

$$[a, b] \subseteq [0, 1], \quad 0 \leq a \leq b \leq 1.$$

Assume we have a continuous function $f : \mathcal{I} \rightarrow \mathbb{R}$ such that for degenerate segments we have $f([a, a]) \equiv 0$ for all $a \in [0, 1]$. Then for any positive integer m it is possible to partition the segment $[0, 1]$ into m possibly degenerate segments

$$[0, 1] = [0, x_1] \cup [x_1, x_2] \cup \cdots \cup [x_{m-1}, 1], \quad 0 \leq x_1 \leq x_2 \leq \cdots \leq x_{m-1} \leq 1,$$

2010 *Mathematics Subject Classification.* 91B32, 55M20, 55M35.

Key words and phrases. Fair partition, Configuration space.

[♠] Supported by the European Research Council under the European Union's Seventh Framework Programme ERC Grant agreement ERC StG 716424 – CAsE.

[♣] Supported by the Russian Foundation for Basic Research grant 19-01-00169.

so that

$$f([0, x_1]) = f([x_1, x_2]) = \cdots = f([x_{m-1}, 1]).$$

Let us first comment on the novelty of this result. This theorem might look like a particular case $d = 1$ of [1, Theorem 6.1], but it is not. The difference is that in Theorem 1.1 we additionally require a certain behavior of f on degenerate segments, while in [1, Theorem 6.1] the assumption $d \geq 2$ eliminates the necessity to consider degenerate parts. The function of a convex body f in the proof of [1, Theorem 6.1] is only applied to convex bodies of a certain positive volume (measure), thus excluding the need to consider degenerate parts and extend the function to them.

On the level of proofs, we use the same approach of using “nice multivalued functions”, adapting [1, Lemma 4.2] to our problem in the form of Lemma 2.3 below. The proof of this lemma is also very similar to the proof of [1, Lemma 4.2], but in this paper the proof is made self-contained and independent of the very technical results of [4], which are used in [1], with the help of a simpler configuration space.

Let us also comment on the previously known particular cases of Theorem 1.1. The case of non-negative f in Theorem 1.1 follows from the Knaster–Kuratowski–Mazurkiewicz theorem [10] in a standard way. The case of m a prime power and f of varying sign follows from the more general result of [2], the case of prime m following from [11]. The case of f additive on segments is an elementary exercise. Hence *the new case here is when the sign of f varies, f is not additive, and m is not a prime power*. Of course, our proof is also an essentially new proof for those known particular cases.

A generalization of Theorem 1.1 for envy-free divisions, when m players divide a segment into m possibly empty parts and each of the players wants to receive one of the best parts according to his/her individual function $f_i : \mathcal{I} \rightarrow \mathbb{R}$, is open for m not a prime power. See further explanations and definitions on envy-free division of the segment in [15, 11, 2].

Acknowledgments. We thank Pablo Soberón, Peter Landweber, and Hervé Moulin for numerous remarks that helped improve the exposition. We also thank the anonymous referees for convincing us to rewrite the paper in a more elementary and self-contained way.

2. REDUCTION TO THE LEMMA ON MULTIVALUED FUNCTIONS

First, we pass from single-valued functions to multi-valued functions. After rescaling we may assume that f in the statement of the theorem takes values in $(-1, 1)$. Let the *cylinder* be the set $\mathcal{I} \times [-1, 1]$.

Definition 2.1. A *nice* multi-valued function $\mathcal{I} \rightarrow [-1, 1]$ is a compact subset $Z \subset \mathcal{I} \times (-1, 1)$ (also called the *graph of the multi-valued function*) that *separates the top from the bottom*, that is, the sets $\mathcal{I} \times \{-1\}$ and $\mathcal{I} \times \{1\}$ belong to different connected components of $\mathcal{I} \times [-1, 1] \setminus Z$.

The following lemma allows us to represent nice multi-valued functions by single-valued continuous functions on the whole cylinder. This will be needed to build maps out of single-valued functions and apply Borsuk–Ulam-type arguments to the maps, see the proof of Claim 4.1 below.

Lemma 2.2. (a) *For any nice multi-valued function of \mathcal{I} , its graph is the zero set of an ordinary continuous function $\varphi : \mathcal{I} \times [-1, 1] \rightarrow \mathbb{R}$ such that $\varphi(\mathcal{I} \times \{-1\}) < 0$ and $\varphi(\mathcal{I} \times \{1\}) > 0$.*

(b) *For any ordinary continuous function $\varphi : \mathcal{I} \times [-1, 1] \rightarrow \mathbb{R}$ such that $\varphi(\mathcal{I} \times \{-1\}) < 0$ and $\varphi(\mathcal{I} \times \{1\}) > 0$, its zero set is a graph of a nice multi-valued function.*

Proof. Claim (b) is trivial, so we prove (a). For a graph $Z \subset \mathcal{I} \times (-1, 1)$ of a nice multi-valued function let φ be the distance to Z with a sign. It is possible to choose the sign arbitrarily for each connected component of $\mathcal{I} \times [-1, 1] \setminus Z$; any such signed distance function is continuous. The requirement for the sign of φ is achieved if we choose the sign of φ positive on the top,

negative on the bottom, and arbitrarily on other connected components. Note that the “nice” property allows choosing the sign on the top and on the bottom independently. \square

In what follows we pass back and forth between the two points of view on multi-valued functions using Lemma 2.2. The function f from the statement of Theorem 1.1 can be considered as a nice multi-valued function with $\varphi([a, b], y) = y - f([a, b])$. The boundary assumption $f([a, a]) \equiv 0$ for all a means that $\varphi([a, a], y) \equiv y$ for all a .

Lemma 2.3 (A modification of Lemma 4.2 from [1]). *Assume a continuous $\varphi : \mathcal{I} \times [-1, 1] \rightarrow \mathbb{R}$ corresponds to a nice multi-valued function and $\varphi([a, a], y) \equiv y$ for all y . Let p be a prime. Then there exists another nice multi-valued function, represented by a continuous $\psi : \mathcal{I} \times [-1, 1] \rightarrow \mathbb{R}$ such that $\psi([a, a], y) \equiv y$ for all y and, whenever $I \in \mathcal{I}$ and $y \in (-1, 1)$ satisfy*

$$\psi(I, y) = 0$$

then there exists a partition $I = I_1 \cup \dots \cup I_p$ into possibly degenerate segments such that

$$(2.1) \quad \varphi(I_1, y) = \dots = \varphi(I_p, y) = 0.$$

Proof of Theorem 1.1 assuming Lemma 2.3. Let $m = p_1 p_2 \dots p_n$ be a decomposition into primes. Let φ_1 be the initial single-valued function f . Apply the lemma to φ_1 and p_1 to obtain a nice multi-valued function φ_2 . Then apply the lemma to φ_2 and p_2 and so on. The final function φ_{n+1} will be a nice multi-valued function of a segment.

From the definition it follows that a nice multi-valued function assigns at least one value to any segment. Hence there exists $y \in (-1, 1)$ such that

$$\varphi_{n+1}([0, 1], y) = 0.$$

It means that $[0, 1]$ may be partitioned into p_n possibly degenerate segments I_1, \dots, I_{p_n} of the same value y of the multi-valued function φ_n ,

$$\varphi_n(I_1, y) = \dots = \varphi_n(I_{p_n}, y) = 0.$$

Each of these segments may in turn be partitioned into p_{n-1} segments of the same value y of the multi-valued function φ_{n-1} , and so on. Eventually, we obtain a partition of $[0, 1]$ into $m = p_1 \dots p_n$ parts of the same value y of the multi-valued function φ_1 , which is in fact the single-valued function f we have started from. \square

3. PARAMETRIZING PARTITIONS OF A SEGMENT INTO A PRIME NUMBER OF PARTS

Our proof of Lemma 2.3 will use equivariant maps and certain claims of Borsuk–Ulam type. We generally follow [1], but instead of using the configuration space of [4], we make a simplification. In this case the partitions of a segment into segments are easier to parametrize more directly, using the idea of [14] with a slightly different construction.

In order to present a proof of Lemma 2.3, we need to parametrize all possibly degenerate partitions of the segment $[a, b]$ into p parts (p is a prime number). A direct parametrization by the relative lengths of the segments,

$$t_1, \dots, t_p \geq 0, \quad t_1 + \dots + t_p = 1,$$

produces the simplex Δ^{p-1} . If the segment $[a, b]$ is degenerate, $a = b$, then we think of all such partitions as the partition of the degenerate segment into degenerate segments. But we still distinguish which of the degenerate segments has its corresponding $t_i = 0$ and which has $t_i > 0$.

Let us additionally label the parts of the partition by numbers from 1 to p , where part i is labeled by $\sigma(i)$. Thus we obtain the space $\Delta^{p-1} \times \mathfrak{S}_p$, where \mathfrak{S}_p is the group of permutations of $\{1, \dots, p\}$. So far our configuration space is just a disjoint union of $p!$ simplices.

After that we identify the pairs $(t, \sigma) \sim (t', \sigma')$ if the labelings σ and σ' become the same sequence of integers after erasing the labels corresponding to degenerate segments with $t_i = 0$ and $t'_j = 0$. This results in identifying some faces of $\Delta^{p-1} \times \{\sigma\}$ and $\Delta^{p-1} \times \{\sigma'\}$.

The obtained space Q_p is a $(p-1)$ -dimensional cell complex. More precisely, this is a Δ -complex in the sense of [7, Section 2.1]. This complex becomes a *simplicial complex*, that is, a union of faces of a simplex spanned by its vertices, after taking the barycentric subdivision of every simplex in its construction.

One may also think of Q_p as partitions of $[a, b]$ into at most p non-degenerate parts with the parts carrying distinct labels from $\{1, \dots, p\}$. This space Q_p has the natural action of the cyclic group $G_p \subseteq \mathfrak{S}_p$ by cyclic permutations of the labels. Since a partition uses at least one label, it is clear that this action of G_p is *free*, that is, for $g \in G_p$, $g \neq e$, and any $\xi \in Q_p$, we have $g\xi \neq \xi$.

An *orientation* of a simplex is an order of its vertices up to even permutations. If the order of vertices of a simplex F is (v_0, \dots, v_k) then its boundary is oriented so that a boundary face $(v_0, \dots, \widehat{v_i}, \dots, v_k)$ (omitted v_i) has the given orientation for even i and the opposite orientation for odd i .

Lemma 3.1. *Consider the orientation of all $(p-1)$ -dimensional cells of Q_p by the order of the labels corresponding to the cell. This orientation makes Q_p a pseudomanifold modulo p , that is, for every $(p-2)$ -dimensional cell F' with any orientation one has the sum over cells containing F'*

$$\sum_{F \supset F'} [F : F'] \equiv 0 \pmod{p},$$

where $[F : F']$ is the sign by which the orientation of F' differs from the orientation of ∂F .

Proof. The generic points of the cell F' correspond to partitions of $[a, b]$ into $p-1$ non-degenerate parts and a certain fixed order of assigned labels from $\{1, \dots, p\}$, one label $k \in \{1, \dots, p\}$ is not assigned. Each cell $F \supset F'$ has the property that generic points of F are partitions of $[a, b]$ into p non-degenerate parts and a certain fixed order of assigned labels from $\{1, \dots, p\}$, all labels assigned.

Since we orient F and F' by the order of the labels, the sign $[F : F']$ equals $(-1)^{k-1}$ by the standard convention on the orientation of the boundary of a simplex. Hence this sign does not depend on F and the sum in the statement of the lemma equals $(-1)^{k-1}p$, since there are precisely p such F containing F' , corresponding to inserting the label k into any of the p places of the sequence labeling F' . \square

Lemma 3.2. *The orientation of all $(p-1)$ -dimensional cells of Q_p by the order of the labels corresponding to the cell is invariant with respect to the action of G_p when p is an odd prime, and changes under the action of the nontrivial element of G_2 when $p = 2$. The collection of $(p-1)$ -faces of Q_p with given orientations is then a G_p -equivariant $(p-1)$ -dimensional cellular cycle modulo p .*

Proof. The proof follows from the fact that G_p , viewed as a subgroup of the permutations, consists of even permutations when p is odd.

For $p = 2$ there is no need in choosing orientations since $+1 \equiv -1 \pmod{2}$, every vertex of Q_2 has two edges connected to it, and equivariance modulo 2 is trivial. \square

Example 3.3. Q_2 is constructed from two segments with endpoints labeled by $\{1, 2\}$, glued according to their labels. Topologically Q_2 is a circle, though our choice of orientations on the segments according to their labels does not produce an orientation of the circle, since the orientations do not match where the segments are glued. We may only say that with such a “wrong” orientation this circle becomes a pseudomanifold modulo 2, see Figure 1.

Example 3.4. Q_3 is constructed from six triangles, each having labels $\{1, 2, 3\}$ on its vertices and each having an order of vertices corresponding to the order on the segment that we partition. We may index those triangles as

$$(1, 2, 3), (1, 3, 2), (2, 1, 3), (2, 3, 1), (3, 1, 2), (3, 2, 1).$$




FIGURE 1. Above: Labeled partitions of the segment into two segments and corresponding points in the 1-dimensional simplices of Q_2 . Below: The complex Q_2 with orientation of its simplices.

The gluing rules mean that Q_3 has three vertices with the labels 1, 2, 3 and six edges labeled

$$(1, 2), (2, 1), (2, 3), (3, 2), (1, 3), (3, 1),$$

attached to the vertices accordingly. In particular, the 1-skeleton of Q_3 is not a simple graph. The six triangles are glued to the edges so that, for example, triangle with labels (3, 2, 1) is glued to the edges (3, 2), (2, 1), (3, 1) according to the labels, see Figure 2.




FIGURE 2. The 1-dimensional skeleton of Q_3 and its six triangular faces shown separately.

We continue to investigate the properties of Q_p that are needed in the proof of Lemma 2.3. Let $W_p \subset \mathbb{R}^p$ be the linear subspace of vectors with the sum of coordinates equal to zero, which is $(p - 1)$ -dimensional. Let the cyclic group G_p act on W_p by cyclic permutations of the coordinates. This action flips the orientation of W_p only when $p = 2$, when we do not care about orientations.

The following construction together with the pseudomanifold property from Lemma 3.1 ensures a Borsuk–Ulam-type property for maps from Q_p to W_p . Define the map $\Psi_0 : Q_p \rightarrow W_p$

as taking a labeled partition (t, σ) to the sequence

$$(3.1) \quad \left(t_{\sigma^{-1}(1)} - \frac{1}{p}, \dots, t_{\sigma^{-1}(p)} - \frac{1}{p} \right).$$

This map may be viewed as sending a configuration $\xi \in Q_p$ to the lengths of the segments of the corresponding partition, listed in the order of their labels, and then normalized by subtracting the vector $(1/p, \dots, 1/p)$. It clearly agrees with the identifications made in the construction of Q_p and is *equivariant*, that is, commuting with the action of G_p on Q_p and W_p .

We need the standard notion of transversality:

Definition 3.5. For a simplicial complex Q , a continuous map $f : Q \rightarrow \mathbb{R}^n$ is *transverse to zero* if for every face $F \subseteq Q$ and every point $x \in F$ such that $f(x) = 0$ the map f is linear in a neighborhood of x in F and this linear map is surjective.

In particular, the images (under a transverse to zero $f : Q \rightarrow \mathbb{R}^n$) of faces $F \subset Q$ of dimension strictly less than n do not contain 0. When a continuous map does not have 0 in its image then it is transverse to zero by definition. In what follows we simply write *transverse* instead of *transverse to zero*; this should not lead to a confusion.

Claim 3.6. *Suppose that a finite group G acts on a simplicial complex Q freely, and acts on \mathbb{R}^n linearly. Then any equivariant continuous map $f : Q \rightarrow \mathbb{R}^n$ can be approximated by a transverse equivariant PL map $g : Q \rightarrow \mathbb{R}^n$.*

Proof. Consider an iterated barycentric subdivision of Q . Perturb the f map on the vertices of this iterated barycentric subdivision generically and equivariantly; the latter can be achieved because the action is free. Then extend the map linearly to the faces of the subdivision, obtaining $g : Q \rightarrow \mathbb{R}^n$.

Since the action of G on Q is free, the vertices of any face F of the barycentric subdivision belong to different G -orbits, and hence generically the restriction $g|_F$ is transverse, and the whole g is transverse.

If the number of iterations of barycentric subdivision is sufficiently large, g will approximate f arbitrarily close. \square

See also [2, Lemma 3.4] for a generalization of Claim 3.6 that does not assume that the action is free.

When $\dim Q = n$ and $f : Q \rightarrow \mathbb{R}^n$ is transverse, at any point $\xi \in f^{-1}(0)$ the derivative Df is a well-defined linear map from the n -dimensional tangent space of the n -face of Q containing ξ to \mathbb{R}^n . Having an orientation on n -faces of Q (as we have for Q_p) we define the *local degree* of f at ξ as the sign of the determinant of Df (compare to the more general definition in [7, Section 2.2]).

Lemma 3.7. *The test map $\Psi_0 : Q_p \rightarrow W_p$ is transverse. The local degrees of Ψ_0 at every point of $\Psi_0^{-1}(0)$ are the same. There are in total $p!$ points and $(p-1)!$ of G_p -orbits in the set $\Psi_0^{-1}(0)$.*

Proof. Evidently, $\Psi_0^{-1}(0)$ corresponds to the partition of $[a, b]$ into parts of equal length labeled in each of $p!$ possible ways. Since the orientation on every $(p-1)$ -dimensional face F is given by the order of labels and the map (3.1) is linear on F , one easily sees the map Ψ_0 around any point $\xi \in \Psi_0^{-1}(0)$ does not flip the given orientation of Q_p compared to the orientation of W_p and all the signs are $+1$. \square

4. PROOF OF LEMMA 2.3

We are going to parametrize partitions of a segment $[a, b]$ into p possibly degenerate segments by the above described Q_p . The proof below, except for the ending, almost literally follows the proof of Lemma 4.2 in [1], but we choose to present the self-contained argument here.

For any $\xi \in Q_p$ and a label $i \in \{1, \dots, p\}$ we denote by I_i the subsegment of the segment $[a, b]$ having label i in ξ . If the label goes to a degenerate segment then this segment is not well-defined, though it is correct to insert such $I_i(\xi)$ into the function φ from the statement of the lemma, since φ is the same for all degenerate segments by the assumptions of the lemma. In this notation, the equations on the varying segment $[a, b]$, $\xi \in Q_p$, and $t \in [-1, 1]$

$$(4.1) \quad \varphi(I_1(\xi), y) = \dots = \varphi(I_p(\xi), y) = 0,$$

define a closed subset $S \subset \mathcal{I} \times Q_p \times [-1, 1]$.

The set S is G_p -invariant, where G_p acts on Q_p as described and acts trivially on \mathcal{I} and $[-1, 1]$. In other words, the set S is the preimage of zero under the G_p -equivariant continuous map

$$\Phi : \mathcal{I} \times Q_p \times [-1, 1] \rightarrow \mathbb{R}^p, \quad \Phi(I, \xi, y) = (\varphi(I_1(\xi), y), \varphi(I_2(\xi), y), \dots, \varphi(I_p(\xi), y)).$$

As the first step in understanding S , we fix a segment $I \in \mathcal{I}$ and study the structure of the fiber set

$$S_I = S \cap (\{I\} \times Q_p \times [-1, 1]).$$

Put $\Phi_I = \Phi|_{\{I\} \times Q_p \times [-1, 1]}$.

Claim 4.1. *For a transverse Φ_I , the set S_I is a finite point sets representing a nontrivial G_p -equivariant 0-cycle modulo p , that is, the local degrees of Φ_I at the orbits of S_I sum up to a number not divisible by p . The projection of the quotient S_I/G_p to the segment $[-1, 1]$ is a nontrivial 0-cycle modulo p .*

Proof. When Φ_I is transverse, the solution set S_I is a finite number of points by dimensional considerations. If we deform Φ_I by a G_p -equivariant homotopy keeping the boundary conditions on its components $\varphi(I_i, y)$ then the solution set S_I changes, but it changes in a definite way. If the homotopy $H : Q_p \times [-1, 1] \times [0, 1] \rightarrow \mathbb{R}^p$ is transverse (this can be achieved by a small perturbation from Claim 3.6) then $H^{-1}(0)$ represents a G_p -equivariant 1-dimensional cycle modulo p relative to $Q_p \times [-1, 1] \times \{0, 1\} \subset Q_p \times [-1, 1] \times [0, 1]$.

The domain of the homotopy $Q_p \times [-1, 1] \times [0, 1]$ is a product of a pseudomanifold modulo p (by Lemma 3.1) and two segments, and therefore it satisfies the pseudomanifold property for faces not contained in its “boundary modulo p ” $Q_p \times \partial([s_0, s_1] \times [-1, 1])$. The mentioned “1-dimensional cycle modulo p ” property means that $H^{-1}(0)$ is a graph (with possible loops), whose edges have orientations (induced by the orientation of $Q_p \times [-1, 1] \times [0, 1]$ and \mathbb{R}^p) and whose every vertex, except for those in $Q_p \times [-1, 1] \times \{0, 1\}$, has algebraically 0 mod p edges connected to it. Note that no vertex of $H^{-1}(0)$ is contained in $Q_p \times \{-1, 1\} \times [0, 1]$ by construction.

The “1-dimensional cycle modulo p ” property holds because under the transversality assumption every vertex $v \in H^{-1}(0)$ is a point of intersection of $H^{-1}(0)$ with a codimension 1 face $F' \subset Q_p \times [-1, 1] \times [0, 1]$. If this vertex is not in $Q_p \times [-1, 1] \times \{0, 1\}$ then there are 0 mod p full-dimensional faces F containing such F' of codimension 1, and each F' contributes the same sign to the orientation of F' . Under the transversality assumption each $F \supset F'$ contributes an edge of $H^{-1}(0)$ attached to v , all such edges contribute the same sign and their total number is 0 mod p . The vertices of the graph $H^{-1}(0)$ contained in $Q_p \times [-1, 1] \times \{0, 1\}$ correspond to the start and the end of the homotopy.

G_p -equivariance of $H^{-1}(0)$ as a 1-dimensional cycle modulo p follows from Lemma 3.2 and the equivariance of the map H . Again, for $p = 2$ the orientation is actually not needed.

Summarizing, the zero set of a transverse Φ_I changes equivariantly homologically modulo p to itself under G_p -equivariant homotopies of the map Φ_I .

Let us present an instance of a transverse map $\Phi_0 : Q_p \times [-1, 1] \rightarrow \mathbb{R}^p$ (a test map), which is G_p -equivariant, satisfies the boundary conditions that we impose on Φ_I , and for which the set $\Phi_0^{-1}(0)$ is homologically nontrivial. By the above homotopy consideration (connecting Φ_0 to Φ_I by a convex interpolation of their coordinates), the existence of such a test map implies

the homological nontriviality of S_I for any transverse map Φ_I . In order to produce the needed test map, we take the G_p -equivariant test map

$$\Psi_0 : Q_p \rightarrow W_p \subset \mathbb{R}^p$$

from Lemma 3.7. The transverse preimage of zero $\Psi_0^{-1}(0)$ consists of $(p-1)! \not\equiv 0 \pmod p$ G_p -orbits, and all orientations (signs) of those points are equal. This verifies the homological nontriviality of $\Psi_0^{-1}(0)$ as a 0-dimensional G_p -equivariant cycle (the number of signed orbits is not divisible by p).

We augment Ψ_0 to the map (assuming the coordinates of Ψ_0 are in the interval $(-1, 1)$)

$$\Phi_0(\xi, y) = \Psi_0(\xi) + (y, \dots, y) \in \mathbb{R}^p.$$

Then $\Phi_0^{-1}(0) = \Psi_0^{-1}(0) \times \{0\}$ and this preimage is still a nontrivial G_p -equivariant 0-cycle modulo p . This finishes the proof of the first part of the claim.

The second part of the claim means that we consider the set of orbits S_I/G_p with coefficients equal to the local degrees of Φ_I . Every orbit projects to a single point $y \in [-1, 1]$, and we assign to such y the coefficient equal to the sum of all coefficients of the orbits of S_I/G_p mapped to y . Thus we evidently obtain a non-trivial modulo p cycle in the segment $[-1, 1]$. \square

Remark 4.2. Note that our homotopy observation is an almost direct generalization of Imre Bárány's geometric proof of the classical Borsuk–Ulam theorem to the case of a pseudomanifold modulo p as a domain. See, for example, [12, 9] where this technique is explained for honest manifolds as domains.

Now we understand that the set S_I is always non-empty, since were it empty, the map Φ_I would be transverse by definition and S_I would have to be non-empty by the claim.

As the second step in our understanding of S , we change the segment I in a continuous one-parameteric family $\{I(s) \mid s \in [s_0, s_1]\}$ and obtain a G_p -equivariant map with one more parameter

$$\tilde{\Phi} : Q_p \times [s_0, s_1] \times [-1, 1] \rightarrow \mathbb{R}^p, \quad \tilde{\Phi}(\xi, s, y) = (\varphi(I_1(\xi, s), y), \varphi(I_2(\xi, s), y), \dots, \varphi(I_p(\xi, s), y)),$$

where $I_i(\xi, s)$ is the i th part of the partition of $I(s)$ corresponding to Q_p .

Claim 4.3. *For a family of segments $I(s)$, the set*

$$S_{I(s)} = S \cap (\{I(s)\}_{s \in [s_0, s_1]} \times Q_p \times [-1, 1])$$

separates the top $[s_0, s_1] \times \{1\}$ from the bottom $[s_0, s_1] \times \{-1\}$ when projected to the rectangle $[s_0, s_1] \times [-1, 1]$.

Proof. Assume first that $\tilde{\Phi}$ is transverse. The solution set $\tilde{\Phi}^{-1}(0)$ then represents a G_p -equivariant 1-dimensional cycle modulo p relative to $Q_p \times [s_0, s_1] \times [-1, 1]$. As in the beginning of the proof of Claim 4.1, under the transversality assumption $\tilde{\Phi}^{-1}(0)$ is a graph formed by oriented paths or loops in the top-dimensional faces of the domain $Q_p \times [s_0, s_1] \times [-1, 1]$ whose ends are isolated points of intersection of $\tilde{\Phi}^{-1}(0)$ with the 1-codimensional skeleton of the domain.

The domain $Q_p \times [s_0, s_1] \times [-1, 1]$ is a product of a pseudomanifold modulo p (by Lemma 3.1) and two segments, and therefore it satisfies the pseudomanifold property for faces not contained in its “boundary modulo p ” $Q_p \times \partial([s_0, s_1] \times [-1, 1])$. Since $\tilde{\Phi}^{-1}(0)$ does not intersect $Q_p \times [s_0, s_1] \times \{-1, 1\}$ by construction, the edges of $\tilde{\Phi}^{-1}(0)$ are attached to every its vertex of 0 modulo p times, except for the vertices with $s = s_0$ or $s = s_1$, corresponding to the start and the end of the homotopy.

Projecting the 1-cycle $\tilde{\Phi}^{-1}(0)$ to the rectangle $[s_0, s_1] \times [-1, 1]$ and noting that every G_p -orbit goes to a single point under this projection, we get a 1-dimensional cycle S' modulo p relative to $[s_0, s_1] \times [-1, 1]$, intersecting a generic line $s = \text{const}$ nontrivially modulo p by Claim 4.1, since this is the solution set of a generic problem with a fixed segment $I(s)$. More generally, any curve connecting the bottom $[s_0, s_1] \times \{-1\}$ and the top $[s_0, s_1] \times \{1\}$ of the rectangle is

homologous to such a line, and it must intersect this cycle by the homological invariance of the intersection number modulo p . The homological invariance holds because the rectangle $[s_0, s_1] \times [0, 1]$ is a PL manifold, S' has boundary on the left and the right sides of the rectangle, and we only consider the curves connecting the top and the bottom of the rectangle and not touching the sides.

We have proved the claim for a transverse $\tilde{\Phi}$, and now consider the general case. Assume that we have a curve Γ from $[s_0, s_1] \times \{-1\}$ to $[s_0, s_1] \times \{1\}$ not touching the projection of the solution set for a not necessarily transverse G_p -equivariant $\tilde{\Phi}$, satisfying the boundary conditions. From the compactness considerations, the minimum of $|\tilde{\Phi}|$ on the preimage $\tilde{\Gamma}$ of Γ in $Q_p \times [s_0, s_1] \times [-1, 1]$ is some $\varepsilon > 0$. Hence, if the transverse equivariant perturbation $\tilde{\Phi}_\varepsilon$ provided by Claim 3.6 is less than ε close to $\tilde{\Phi}$ then the solution set $\tilde{\Phi}_\varepsilon^{-1}(0)$ will still be disjoint from $\tilde{\Gamma}$ and its projection to the rectangle will still be disjoint from Γ . But for a transverse $\tilde{\Phi}_\varepsilon$ the existence of such a curve Γ is already shown to be impossible. \square

Now perform the third step in our understanding of S , working in the full cylinder of parameters, $\mathcal{I} \times [-1, 1]$.

Claim 4.4. *The projection Z of S to $\mathcal{I} \times [-1, 1]$ separates the top $\mathcal{I} \times \{1\}$ from the bottom $\mathcal{I} \times \{-1\}$.*

Proof. Assume that a continuous curve

$$\Gamma : [s_0, s_1] \rightarrow \mathcal{I} \times [-1, 1]$$

passes from the bottom $\mathcal{I} \times \{-1\}$ to the top $\mathcal{I} \times \{1\}$ in the cylinder. Its first coordinate may be considered as a one-parametric family of segments $I(s)$, to which we apply Claim 4.3 and conclude that Γ must meet Z . \square

We have the crucial separation property of $Z \subset \mathcal{I} \times [-1, 1]$, considered as a graph of a multi-valued function. The separation property implies that this multi-valued function is nice; the first approximation to its corresponding

$$\psi : \mathcal{I} \times [-1, 1] \rightarrow \mathbb{R}$$

could be obtained as a signed distance function, as in the proof of Lemma 2.2. Whenever $\psi(I, y) = 0$, the pair (I, y) is in Z and corresponds to some $(I, \xi, y) \in S$. By the definition of S , ξ provides a labeled partition of I into p segments I_1, \dots, I_p satisfying

$$\varphi(I_1, y) = \dots = \varphi(I_p, y) = 0.$$

But we also need to ensure that $\psi([a, a], y) \equiv y$ for all y , which may be not true for the signed distance function obtained from the proof of Lemma 2.2. We had no such difficulty in the proof of [1, Lemma 4.2] and the need to overcome it here is essentially what makes this proof different.

Put for brevity $\mathcal{I}' = \{[a, a] \mid a \in [0, 1]\} \subset \mathcal{I}$. Examining our construction of S and Z in case of a degenerate segment (which may only be partitioned into degenerate segments) and using the fact that $\varphi([a, a], y) \equiv y$ for all y , we see that

$$(4.2) \quad Z \cap (\mathcal{I}' \times \mathbb{R}) = \mathcal{I}' \times \{0\}.$$

In order to obtain the property $\psi([a, a], y) \equiv y$, we use a modification of the argument in the proof of Lemma 2.2 to build the function $\psi : \mathcal{I} \times [-1, 1] \rightarrow \mathbb{R}$ with zero set Z . Define

$$\psi(Z) = 0, \quad \psi(\mathcal{I}', y) \equiv y, \quad \psi(I, 1) \equiv 1, \quad \psi(I, -1) \equiv -1.$$

After this and because of (4.2) ψ is continuously defined on the closed set $Y = Z \cup (\mathcal{I}' \times [-1, 1]) \cup (\mathcal{I} \times \{-1, 1\}) \supseteq Z$.

Then we extend ψ by the Tietze extension theorem to the connected components of $\mathcal{I} \times (-1, 1) \setminus Y$ so that on the connected components touching the top it remains non-negative. By adding to ψ the distance function to the set Y in such components (and still denoting the

resulting function by ψ) we make ψ strictly positive in top components of the complement of Y . We do the same on the components of the complement of Y touching the bottom with the minus sign, thus extending ψ to a negative function there. In effect, we obtain ψ with zero set Z satisfying $\psi(\mathcal{I}', y) \equiv y$, $\psi(\mathcal{I}, 1) > 0$, and $\psi(\mathcal{I}, -1) < 0$.

REFERENCES

- [1] A. Akopyan, S. Avvakumov, and R. Karasev. Convex fair partitions into an arbitrary number of pieces. 2018. [arXiv:1804.03057](#), version 7 and higher.
- [2] S. Avvakumov and R. Karasev. Envy-free division using mapping degree. *Mathematika*, 67(1):36–53, 2021. [arXiv:1907.11183](#).
- [3] J. B. Barbanel, S. J. Brams, and W. Stromquist. Cutting a pie is not a piece of cake. *American Mathematical Monthly*, 116(6):496–514, 2009.
- [4] P. Blagojević and G. Ziegler. Convex equipartitions via equivariant obstruction theory. *Israel Journal of Mathematics*, 200(1):49–77, 2014. [arXiv:1202.5504](#).
- [5] A. Bogomolnaia and H. Moulin. Guarantees in fair division: general or monotone preferences. 2019. [arXiv:1911.10009](#), version 3 and higher.
- [6] D. Gale. Equilibrium in a discrete exchange economy with money. *International Journal of Game Theory*, 13(1):61–64, 1984.
- [7] A. Hatcher. *Algebraic Topology*. Cambridge University Press, 2001. pi.math.cornell.edu/~hatcher/AT/AT.pdf.
- [8] R. Karasev, A. Hubard, and B. Aronov. Convex equipartitions: the spicy chicken theorem. *Geometriae Dedicata*, 170(1):263–279, 2014. [arXiv:1306.2741](#).
- [9] B. Klartag. Convex geometry and waist inequalities. *Geometric and Functional Analysis*, 27(1):130–164, 2017. [arXiv:1608.04121](#).
- [10] B. Knaster, C. Kuratowski, and S. Mazurkiewicz. Ein Beweis des Fixpunktsatzes für n -dimensionale Simplexe. *Fundamenta Mathematicae*, 14(1):132–137, 1929.
- [11] F. Meunier and S. Zerbib. Envy-free cake division without assuming the players prefer nonempty pieces. *Israel Journal of Mathematics*, 234:907–925, 2019. [arXiv:1804.00449](#).
- [12] O. R. Musin. Borsuk–Ulam type theorems for manifolds. *Proceedings of the American Mathematical Society*, 140:2551–2560, 2012.
- [13] R. Nandakumar and N. Ramana Rao. ‘Fair’ partitions of polygons – an introduction. 2008. [arXiv:0812.2241](#).
- [14] G. Panina and R. Živaljević. Envy-free division via configuration spaces. 2021. [arXiv:2102.06886](#).
- [15] E. Segal-Halevi. Fairly dividing a cake after some parts were burnt in the oven. *Proceedings of the 17th International Conference on Autonomous Agents and MultiAgent Systems (AAMAS 2018)*, pages 1276–1284, 2018.
- [16] P. Soberón. Balanced convex partitions of measures in \mathbb{R}^d . *Mathematika*, 58(1):71–76, 2012.
- [17] H. Steinhaus. Sur la division des ensembles de l’espace par les plans et des ensembles plans par les cercles. *Fund. Math.*, 33:245–263, 1945.
- [18] A. Stone and J. Tukey. Generalized “sandwich” theorems. *Duke Mathematical Journal*, 9:356–359, 1942.
- [19] F. E. Su. Rental harmony: Sperner’s lemma in fair division. *American Mathematical Monthly*, 106(10):930–942, 1999.

SERGEY AVVAKUMOV, DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF COPENHAGEN, UNIVERSITETSPARK 5, 2100 COPENHAGEN, DENMARK

Email address: savvakumov@gmail.com

ROMAN KARASEV, INSTITUTE FOR INFORMATION TRANSMISSION PROBLEMS RAS, BOLSHOY KARETNY PER. 19, MOSCOW, RUSSIA 127994

MOSCOW INSTITUTE OF PHYSICS AND TECHNOLOGY, INSTITUTSKIY PER. 9, DOLGOPRUDNY, RUSSIA 141700

Email address: r.n.karasev@mail.ru

URL: <http://www.rkarasev.ru/en/>