

THE CAUCHY PROBLEM FOR 3-EVOLUTION EQUATIONS WITH DATA IN GELFAND-SHILOV SPACES

ALEXANDRE ARIAS JUNIOR, ALESSIA ASCANELLI, AND MARCO CAPPIELLO

ABSTRACT. We consider the Cauchy problem for a third order evolution operator P with (t, x) -depending coefficients and complex valued lower order terms. We assume the initial data to be Gevrey regular with an exponential decay at infinity, that is, the data belong to some Gelfand-Shilov space of type \mathcal{S} . Under suitable assumptions on the decay at infinity of the imaginary parts of the coefficients of P we prove the existence of a solution with the same Gevrey regularity of the data and we describe its behavior for $|x| \rightarrow \infty$.

2010 Mathematics Subject Classification: 35Q41, 35S05, 46F05

Keywords and phrases: p -evolution equations, Gelfand-Shilov spaces, infinite order pseudodifferential operators

1. Introduction and main result

Let us consider for $(t, x) \in [0, T] \times \mathbb{R}$ the Cauchy problem in the unknown $u = u(t, x)$:

$$(1.1) \quad \begin{cases} P(t, x, D_t, D_x)u(t, x) = f(t, x) \\ u(0, x) = g(x) \end{cases},$$

where

$$(1.2) \quad P(t, x, D_t, D_x) = D_t + a_p(t)D_x^p + \sum_{j=0}^{p-1} a_j(t, x)D_x^j,$$

with $D = \frac{1}{i}\partial$, $p \geq 2$, $a_p \in C([0, T], \mathbb{R})$, $a_p(t) \neq 0$ for $t \in [0, T]$, and $a_j \in C([0, T], C^\infty(\mathbb{R}; \mathbb{C}))$, $j = 0, \dots, p-1$. The operator P is known in literature as p -evolution operator, cf. [29], and p is the evolution degree. The well posedness of (1.1), (1.2) has been investigated in various functional settings for arbitrary p , cf. [3–5, 10]. Further results concern special values of p which correspond to classes of operators of particular interest in Mathematical Physics, cf. [6, 8, 9, 11, 22, 24] for the case $p = 2$ and [1] in the case $p = 3$. The condition that a_p is real valued means that the principal symbol of P (in the sense of Petrowski) has the real characteristic $\tau = -a_p(t)\xi^p$; this guarantees that the operator (1.2) satisfies the assumptions of Lax-Mizohata theorem. The presence of complex valued coefficients in the lower order terms of (1.2) plays a crucial role in the analysis of the problem (1.1) in all the above mentioned papers. In fact, when the coefficients $a_j(t, x)$, $j = 0, \dots, p-1$, are real valued and of class \mathcal{B}^∞ with respect to x (uniformly bounded together with all their x -derivatives), it is well known that the problem (1.1) is well-posed in $L^2(\mathbb{R})$ (and in L^2 -based Sobolev spaces H^m , $m \in \mathbb{R}$). On the contrary if any of the coefficients $a_j(t, x)$ are complex valued, then in order to obtain

The first author would like to thank Fundação Araucária for the financial support during the development of this paper. The second author has been supported by the Italian national research funds FFABR2017 and GNAMPA 2020. The third author is supported by the fund GNAMPA 2020.

well-posedness either in $L^2(\mathbb{R})$, or in $H^\infty(\mathbb{R}) = \bigcap_{m \in \mathbb{R}} H^m(\mathbb{R})$, some decay conditions at infinity on the imaginary part of the coefficients a_j are needed (see [4], [22]).

Sufficient conditions for well-posedness in L^2 and H^∞ have been given in [9] and [24] for the case $p = 2$, in [3] for larger p . Considering the Cauchy problem (1.1) in the framework of weighted Sobolev-Kato spaces $H^m = H^{(m_1, m_2)}$, with $m = (m_1, m_2) \in \mathbb{R}^2$, defined as

$$(1.3) \quad H^m(\mathbb{R}) = \{u \in \mathcal{S}'(\mathbb{R}) : \langle x \rangle^{m_2} \langle D_x \rangle^{m_1} u \in L^2(\mathbb{R})\},$$

where the analysis of regularity and decay/growth of the solution can be performed simultaneously, and assuming the coefficients a_j to be polynomially bounded, the second and the third author obtained in [5] well-posedness also in the Schwartz space $\mathcal{S}(\mathbb{R})$ of smooth and rapidly decreasing functions and in the dual space $\mathcal{S}'(\mathbb{R})$. We recall that $\mathcal{S}(\mathbb{R}) = \bigcap_{m \in \mathbb{R}^2} H^m(\mathbb{R})$ and $\mathcal{S}'(\mathbb{R}) = \bigcup_{m \in \mathbb{R}^2} H^m(\mathbb{R})$. In short, the above mentioned results can be summarized as follows: if

$$(1.4) \quad |\partial_x^\beta \operatorname{Im} a_j(t, x)| \leq C_\beta \langle x \rangle^{-\frac{j}{p-1}\sigma - |\beta|}, \quad (t, x) \in [0, T] \times \mathbb{R}, \beta \in \mathbb{N}_0, C_\beta > 0, j = 0, \dots, p-1,$$

the problem (1.1) is well-posed in:

- $L^2(\mathbb{R})$, $H^m(\mathbb{R})$ for every $m \in \mathbb{R}^2$ when $\sigma > 1$;
- $H^\infty(\mathbb{R})$, $\mathcal{S}(\mathbb{R})$ when $\sigma = 1$. In general, a finite loss of regularity of the solution with respect to the initial data is observed in the case $\sigma = 1$.

Now we want to consider the case when an estimate of the form (1.4) for $j = p-1$ holds for some $\sigma \in (0, 1)$. In this situation, there are no results in the literature for p -evolution operators of arbitrary order. In [11, 24] the case $p = 2$, which corresponds to Schrödinger-type equations, is considered assuming $0 < \sigma < 1$ and $C_\beta = C^{|\beta|+1} \beta!^{s_0}$ for some $s_0 \in (1, 1/(1-\sigma))$ in (1.4). The authors find well posedness results in certain Gevrey spaces of order θ with $s_0 \leq \theta < 1/(1-\sigma)$, namely in the class

$$\mathcal{H}_\theta^\infty = \bigcup_{\rho > 0} H_{\rho; \theta}^m, \quad H_{\rho; \theta}^m := \{u \in L^2 \mid e^{\rho \langle D \rangle^{1/\theta}} u \in H^m\}, \quad m \in \mathbb{R}.$$

In both papers, starting from data f, g in $H_{\rho; \theta}^m$ for some $\rho > 0$ the authors obtain a solution in $H_{\rho-\delta; \theta}^m$ for some $\delta > 0$ such that $\rho - \delta > 0$. This means a sort of loss of regularity in the constant ρ which rules the Gevrey behavior. We also notice that the condition $s_0 \leq \theta < 1/(1-\sigma)$ means that the rate of decay of the coefficients of P imposes a restriction on the spaces $\mathcal{H}_\theta^\infty$ in which the problem (1.1) is well posed. Finally, the case $\theta > s_0 = 1/(1-\sigma)$ is investigated in [8] where the authors prove that a decay condition as $|x| \rightarrow \infty$ on a datum in H^m , $m \geq 0$, produces a solution with (at least locally) the same regularity as the data, but with a different behavior at infinity. In the recent paper [6] the role of data with exponential decay on the regularity of the solution has been also analyzed for 2-evolution equations in arbitrary space dimension, in the frame of Gelfand-Shilov type spaces which can be seen as the global counterpart of classical Gevrey spaces, cf. Subsection 2.1. In particular, it is proved that starting from data with an exponential decay at infinity, we can find a solution with the same Gevrey regularity of the data but with a possible exponential growth at infinity in x . Moreover, this holds for every $\theta > s_0$. Finally, the result in [6] is proved under the more general assumption with respect to (1.4) that the coefficients a_1, a_0 may admit an algebraic growth at infinity, namely

$$(1.5) \quad |\partial_x^\beta \operatorname{Im} a_1(t, x)| \leq C^{|\beta|+1} \beta!^{s_0} \langle x \rangle^{-\sigma - |\beta|},$$

$$(1.6) \quad |\partial_x^\beta \operatorname{Re} a_1(t, x)| + |\partial_x^\beta a_0(t, x)| \leq C^{|\beta|+1} \beta!^{s_0} \langle x \rangle^{1-\sigma - |\beta|}.$$

Recently, we started to consider the case $p = 3$ in a Gevrey setting in one space dimension under assumption (1.4) with $j = p-1$ taking $\sigma \in (0, 1)$. This case is of particular interest because linear 3-evolution equations can be regarded as linearizations of relevant physical

semilinear models like KdV and KdV-Burgers equation and their generalizations, see for instance [23, 26–28, 32]. There are some results concerning KdV-type equations with coefficients not depending on (t, x) in the Gevrey setting, see [18–20]. Our aim is to treat the more general case of variable coefficients. The present paper and [1] are devoted to establish the linear theory which is a preliminary step to treat the semilinear case. In a future paper we shall consider the case when the coefficients a_j may depend also on u following the approach developed in [2] in the H^∞ setting.

Also for the case $p = 3$, assuming a condition of the form (1.4) with $j = 2$ on the term a_2 for some $\sigma \in (0, 1)$, namely

$$(1.7) \quad |\partial_x^\beta \operatorname{Im} a_2(t, x)| \leq C_\beta \langle x \rangle^{-\sigma-\beta},$$

is enough to lose in general well posedness both in H^∞ and in \mathcal{S} , since the necessary condition for H^∞ well-posedness

$$(1.8) \quad \sup_{x \in \mathbb{R}} \min_{0 \leq \tau \leq t \leq T} \int_{-\varrho}^{\varrho} \operatorname{Im} a_2(t, x + 3a_3(\tau)\theta) d\theta \leq M \log(1 + \varrho) + N, \quad \forall \varrho > 0, \text{ for some } M, N > 0$$

proved in [4] is no more satisfied. Namely, well posedness in H^∞ or in \mathcal{S} may fail due to an infinite loss of regularity or of decay. To give an idea of the latter phenomenon, consider the following initial value problem

$$(1.9) \quad \begin{cases} D_t u + D_x^3 u + a_2(t, x) D_x^2 u + a_1(t, x) D_x u + a_0(t, x) u = 0 \\ u(0, x) = e^{-\langle x \rangle^{1-\sigma}} \end{cases}, \quad (t, x) \in [0, T] \times \mathbb{R},$$

where

$$\begin{aligned} a_2(t, x) &= i(t-1)(1-\sigma)x\langle x \rangle^{-\sigma-1}, \\ a_1(t, x) &= 2(t-1)(1-\sigma)[\langle x \rangle^{-\sigma-1} - (\sigma+1)x^2\langle x \rangle^{-\sigma-3}], \\ a_0(t, x) &= i\langle x \rangle^{1-\sigma} + i(t-1)(1-\sigma^2)[3x\langle x \rangle^{-\sigma-3} - (\sigma+3)x^3\langle x \rangle^{-\sigma-5}]. \end{aligned}$$

Notice that the coefficients a_j are analytic and satisfy conditions (1.4) for $j = 2$ and (1.5), (1.6) for $j = 0, 1$. Moreover the initial datum belongs to $\mathcal{S}(\mathbb{R})$ since $\sigma \in (0, 1)$. It is easy to verify that the problem (1.9) admits the solution

$$u(t, x) = e^{(t-1)\langle x \rangle^{1-\sigma}} \notin C([0, T], \mathcal{S}(\mathbb{R})),$$

if $T \geq 1$. Analogously, $u \notin C([0, T], H^\infty(\mathbb{R}))$. More precisely, we notice that the solution has the same regularity as the initial data but it grows exponentially for $|x| \rightarrow \infty$ when $t \geq 1$. This motivates us to study the effect of an exponential decay of the data on the solution of (1.1).

In the recent paper [1] we proved a result of well posedness in the space $\mathcal{H}_\theta^\infty$ for the problem (1.1) which extends to the case $p = 3$ the results obtained in [11, 24] for the case $p = 2$ (at least in one space dimension). As in the latter case, also in [1] a loss of regularity in the index ρ appears. However, the previous example suggests that this loss can be avoided assuming the initial data to admit a suitable exponential decay. The price to pay is a considerable loss of decay which may produce solutions admitting an exponential growth. In view of the considerations above, it is quite natural to analyze the problem (1.1) when the initial data belong to Gelfand-Shilov spaces, cf. Subsection 2.1 for the definition.

In order to state our main result we need to recall the definition of Gevrey type **SG**-symbol classes and of Gelfand-Shilov Sobolev spaces.

Given $\mu, \nu \geq 1$, $m = (m_1, m_2) \in \mathbb{R}^2$, we denote by $\mathbf{SG}_{\mu, \nu}^{m_1, m_2}(\mathbb{R}^2)$ (or by $\mathbf{SG}_{\mu, \nu}^m(\mathbb{R}^2)$) the space of all functions $p \in C^\infty(\mathbb{R}^2)$ for which there exist $C, C_1 > 0$ such that

$$|\partial_\xi^\alpha \partial_x^\beta p(x, \xi)| \leq C_1 C^{|\alpha+\beta|} \alpha!^\mu \beta!^\nu \langle \xi \rangle^{m_1-|\alpha|} \langle x \rangle^{m_2-|\beta|}, \quad x, \xi \in \mathbb{R}, \alpha, \beta \in \mathbb{N},$$

see also Definition 2. In the case $\mu = \nu$ we write $\mathbf{SG}_\mu^{m_1, m_2}(\mathbb{R}^2)$ instead of $\mathbf{SG}_{\mu, \mu}^{m_1, m_2}(\mathbb{R}^2)$. In the following we shall obtain our results via energy estimates, hence we need to introduce the Gelfand-Shilov-Sobolev spaces $H_{\rho; s, \theta}^m(\mathbb{R})$ defined, for $m = (m_1, m_2)$, $\rho = (\rho_1, \rho_2)$ in \mathbb{R}^2 and $\theta, s > 1$, by

$$H_{\rho; s, \theta}^m(\mathbb{R}) = \{u \in \mathcal{S}'(\mathbb{R}) : \langle x \rangle^{m_2} \langle D \rangle^{m_1} e^{\rho_2 \langle x \rangle^{\frac{1}{s}}} e^{\rho_1 \langle D \rangle^{\frac{1}{\theta}}} u \in L^2(\mathbb{R})\},$$

where $e^{\rho_1 \langle D \rangle^{\frac{1}{\theta}}}$ is the Fourier multiplier with symbol $e^{\rho_1 \langle \xi \rangle^{\frac{1}{\theta}}}$. When $\rho = (0, 0)$ we recover the usual notion of weighted Sobolev spaces (1.3).

Our pseudodifferential approach allows to consider more general 3-evolution operators of the form

$$(1.10) \quad P(t, D_t, x, D_x) = D_t + a_3(t, D_x) + a_2(t, x, D_x) + a_1(t, x, D_x) + a_0(t, x, D_x),$$

$t \in [0, T]$, $x \in \mathbb{R}$, where $a_3(t, D_x)$ is a pseudodifferential operator with symbol $a_3(t, \xi) \in \mathbb{R}$ while, for $j = 0, 1, 2$, $a_j(t, x, D_x)$ are pseudodifferential operators with symbols $a_j(t, x, \xi) \in \mathbb{C}$. Notice that (1.2) in the case $p = 3$ is a particular case of (1.10). Our main result reads as follows.

Theorem 1. *Let $P(t, x, D_t, D_x)$ be an operator as in (1.10) and assume the following:*

- (i) $a_3 \in C([0, T], \mathbf{SG}_1^{3,0}(\mathbb{R}^2))$, a_3 is real valued and there are $C_{a_3}, R_{a_3} > 0$ and $\sigma \in (0, 1)$ such that

$$|\partial_\xi a_3(t, \xi)| \geq C_{a_3} |\xi|^2, \quad \forall |\xi| \geq R_{a_3}, \quad \forall t \in [0, T];$$

- (ii) $\operatorname{Re} a_2 \in C([0, T], \mathbf{SG}_{1, s_0}^{2,0}(\mathbb{R}^2))$, $\operatorname{Im} a_2 \in C([0, T], \mathbf{SG}_{1, s_0}^{2, -\sigma}(\mathbb{R}^2))$;

- (iii) $\operatorname{Re} a_1 \in C([0, T], \mathbf{SG}_{1, s_0}^{1, 1-\sigma}(\mathbb{R}^2))$, $\operatorname{Im} a_1 \in C([0, T], \mathbf{SG}_{1, s_0}^{1, -\frac{\sigma}{2}}(\mathbb{R}^2))$;

- (iv) $a_0 \in C([0, T], \mathbf{SG}_{1, s_0}^{0, 1-\sigma}(\mathbb{R}^2))$.

Let $s, \theta > 1$ such that $s_0 \leq s < \frac{1}{1-\sigma}$ and $\theta > s_0$. Let moreover $f \in C([0, T]; H_{\rho; s, \theta}^m(\mathbb{R}))$ and $g \in H_{\rho; s, \theta}^m(\mathbb{R})$, where $m = (m_1, m_2)$, $\rho = (\rho_1, \rho_2) \in \mathbb{R}^2$ and $\rho_2 > 0$. Then the Cauchy problem (1.1) admits a solution $u \in C([0, T]; H_{(\rho_1, -\tilde{\delta}); s, \theta}^m(\mathbb{R}))$ for every $\tilde{\delta} > 0$, which satisfies the following energy estimate

$$(1.11) \quad \|u(t)\|_{H_{(\rho_1, -\tilde{\delta}); s, \theta}^m}^2 \leq C \left(\|g\|_{H_{\rho; s, \theta}^m}^2 + \int_0^t \|f(\tau)\|_{H_{\rho; s, \theta}^m}^2 d\tau \right),$$

for all $t \in [0, T]$ and for some $C > 0$.

Remark 1. We notice that the solution obtained in Theorem 1 has the same Gevrey regularity as the initial data but it may lose the decay exhibited at $t = 0$ and admit an exponential growth for $|x| \rightarrow \infty$ when $t > 0$. Moreover, the loss $\rho_2 + \tilde{\delta}$ for an arbitrary $\tilde{\delta} > 0$ in the behavior at infinity is independent of θ, s and ρ_1 . Both these phenomena had been already observed in the case $p = 2$, see [6].

Remark 2. Let us compare Theorem 1 with the recent result obtained in [1]. In the latter paper, taking a_0 uniformly bounded, $a_1 \sim \langle x \rangle^{-\sigma/2}$ and $a_2 \sim \langle x \rangle^{-\sigma}$ for some $\sigma \in (1/2, 1)$, and the Cauchy data $f(t), g \in H_{(\rho_1, 0); s, \theta}^{(m_1, 0)}$ with $s_0 < \theta < 1/(2(1-\sigma))$ we prove the existence of a unique solution $u \in C([0, T], H_{(\rho'_1, 0); s, \theta}^{(m_1, 0)})$, for some $\rho'_1 \in (0, \rho_1)$ i.e. a solution less regular than the data. Theorem 1 in the present paper shows that if the data $f(t), g \in H_{(\rho_1, \rho_2); s, \theta}^{(m_1, m_2)}$ with $\rho_2 > 0, \theta > s_0$ and $s_0 \leq s < 1/(1-\sigma)$, then there exists a solution $u \in C([0, T], H_{(\rho_1, -\tilde{\delta}); s, \theta}^{(m, 0)})$, $\forall \tilde{\delta} > 0$, i.e. a solution with the same index ρ_1 as the data, but with a possible worse behavior at infinity: in particular, this solution may grow exponentially for $|x| \rightarrow \infty$. Concerning the

assumptions, with respect to the existing literature, in particular [1, 24], in our result we allow:

- a polynomial growth of exponent $1 - \sigma \in (0, 1)$ for the coefficients $\operatorname{Re} a_1$ and a_0 ;
- an arbitrary Gevrey regularity index $\theta > s_0$ both for the data and for the solution, without any upper bound: namely there is no relation between the rate of decay of the data and the Gevrey regularity of the solution.

Remark 3. Part of the recent literature on p -evolution equations is focused on the research of necessary conditions for the well posedness of the problem (1.1) in various functional settings, see [4, 14, 22]. Necessary conditions are usually expressed in an integral form as in (1.8) instead than via pointwise decay estimates as in (1.7). As far as we know the only result of this type in the Gevrey setting concerns the case $p = 2$, see [14]. Our purpose is to investigate this problem in the next future for generic p .

In order to help the reading of the next sections we briefly outline the strategy of the proof of Theorem 1. Let

$$iP = \partial_t + ia_3(t, D) + \sum_{j=0}^2 ia_j(t, x, D) = \partial_t + ia_3(t, D) + A(t, x, D).$$

Noticing that $a_3(t, \xi)$ is real valued, we have

$$\begin{aligned} \frac{d}{dt} \|u(t)\|_{L^2}^2 &= 2\operatorname{Re} (\partial_t u(t), u(t))_{L^2} \\ &= 2\operatorname{Re} (iPu(t), u(t))_{L^2} - 2\operatorname{Re} (ia_3(t, D)u(t), u(t))_{L^2} - 2\operatorname{Re} (Au(t), u(t))_{L^2} \\ &\leq \|Pu(t)\|_{L^2}^2 + \|u(t)\|_{L^2}^2 - ((A + A^*)u(t), u(t))_{L^2}. \end{aligned}$$

Since $(A + A^*)(t) \in \mathbf{SG}^{2,1-\sigma}(\mathbb{R}^2)$ we cannot derive an energy inequality in L^2 from the estimate above. The idea is then to conjugate the operator iP by a suitable pseudodifferential operator $e^\Lambda(t, x, D)$ in order to obtain

$$(iP)_\Lambda := e^\Lambda(iP)\{e^\Lambda\}^{-1} = \partial_t + ia_3(t, D) + A_\Lambda(t, x, D),$$

where A_Λ still has symbol $A_\Lambda(t, x, \xi) \in \mathbf{SG}^{2,1-\sigma}(\mathbb{R}^2)$ but with $\operatorname{Re} A_\Lambda \geq 0$. In this way, with the aid of Fefferman-Phong (see [15]) and sharp Gårding (see Theorem 1.7.15 of [30]) inequalities, we obtain the estimate from below

$$\operatorname{Re} (A_\Lambda v(t), v(t))_{L^2} \geq -c\|v(t)\|_{L^2}^2,$$

and therefore for the solution v of the Cauchy problem associated to the operator P_Λ we get

$$\frac{d}{dt} \|v(t)\|_{L^2}^2 \leq C(\|(iP)_\Lambda v(t)\|_{L^2}^2 + \|v(t)\|_{L^2}^2).$$

Gronwall inequality then gives the desired energy estimate for the conjugated operator $(iP)_\Lambda$. By standard arguments in the energy method we then obtain that the Cauchy problem associated with P_Λ

$$(1.12) \quad \begin{cases} P_\Lambda(t, D_t, x, D_x)v(t, x) = e^\Lambda(t, x, D_x)f(t, x), & (t, x) \in [0, T] \times \mathbb{R} \\ v(0, x) = e^\Lambda(0, x, D_x)g(x), & x \in \mathbb{R} \end{cases},$$

is well-posed in the weighted Sobolev spaces $H^m(\mathbb{R})$ in (1.3). Finally, we derive the existence of a solution of (1.1) from the well posedness of (1.12). In fact, if u solves (1.1) then $v = e^\Lambda u$ solves (1.12), and if v solves (1.12) then $u = \{e^\Lambda\}^{-1}v$ solves (1.1). In this step the continuous mapping properties of e^Λ and $\{e^\Lambda\}^{-1}$ will play an important role.

The construction of the operator e^Λ will be the core of the proof. The function $\Lambda(t, x, \xi)$ will be of the form

$$\Lambda(t, x, \xi) = k(t)\langle x \rangle_h^{1-\sigma} + \lambda_2(x, \xi) + \lambda_1(x, \xi), \quad t \in [0, T], \quad x, \xi \in \mathbb{R},$$

where $\lambda_1, \lambda_2 \in \mathbf{SG}_\mu^{0,1-\sigma}(\mathbb{R}^2)$, $k \in C^1([0, T]; \mathbb{R})$ is a non increasing function to be chosen later on and $\langle x \rangle_h = \sqrt{h^2 + x^2}$ for some $h \geq 1$ to be chosen later on. The role of the terms λ_1, λ_2, k will be the following:

- the transformation with λ_2 will change the terms of order 2 into the sum of a positive operator of the same order plus a remainder of order 1;
- the transformation with λ_1 will not change the terms of order 2, but it will turn the terms of order 1 into the sum of a positive operator of order 1 plus a remainder of order 0, with some growth with respect to x ;
- the transformation with $k(t)\langle x \rangle_h^{1-\sigma}$ will correct this remainder term, making it positive.

The precise definitions of λ_2 and λ_1 will be given in Section 4. Since Λ admits an algebraic growth on the x variable, then e^Λ presents an exponential growth; this is the reason why we need to work with pseudodifferential operators of infinite order.

The paper is organized as follows. In Section 2 we recall some basic definitions and properties of Gelfand-Shilov spaces and the calculus for pseudodifferential operators of infinite order that we will use in the next sections. Section 3 is devoted to prove a result of spectral invariance for pseudodifferential operators with Gevrey regular symbols which is new in the literature and interesting per se. In this paper the spectral invariance will be used to prove the continuity properties of the inverse $\{e^\Lambda(t, x, D)\}^{-1}$. In Section 4 we introduce the functions λ_1, λ_2 mentioned above and prove the invertibility of the operator $e^{\tilde{\Lambda}}(x, D)$, $\tilde{\Lambda} = \lambda_1 + \lambda_2$. In Section 5 we perform the change of variable and the conjugation of the operator P . Section 6 concerns the choice of the parameters appearing in the definition of Λ in order to obtain a positive operator on $L^2(\mathbb{R})$. Finally, in Section 7, we give the proof of Theorem 1.

2. Gelfand-Shilov spaces and pseudodifferential operators of infinite order on \mathbb{R}^n

2.1. Gelfand-Shilov spaces. Given $s, \theta \geq 1$ and $A, B > 0$ we say that a smooth function f belongs to $\mathcal{S}_{s,B}^{\theta,A}(\mathbb{R}^n)$ if there is a constant $C > 0$ such that

$$|x^\beta \partial_x^\alpha f(x)| \leq C A^{|\alpha|} B^{|\beta|} \alpha!^\theta \beta!^s,$$

for every $\alpha, \beta \in \mathbb{N}_0^n$ and $x \in \mathbb{R}^n$. The norm

$$\|f\|_{\theta,s,A,B} = \sup_{\substack{x \in \mathbb{R}^n \\ \alpha, \beta \in \mathbb{N}_0^n}} |x^\beta \partial_x^\alpha f(x)| A^{-|\alpha|} B^{-|\beta|} \alpha!^{-\theta} \beta!^{-s}$$

turns $\mathcal{S}_{s,B}^{\theta,A}(\mathbb{R}^n)$ into a Banach space. We define

$$\mathcal{S}_s^\theta(\mathbb{R}^n) = \bigcup_{A,B>0} \mathcal{S}_{s,B}^{\theta,A}(\mathbb{R}^n)$$

and we can equip it with the inductive limit topology of the Banach spaces $\mathcal{S}_{s,B}^{\theta,A}(\mathbb{R}^n)$. The spaces $\mathcal{S}_s^\theta(\mathbb{R}^n)$ have been originally introduced in the book [17], see [31]. We also consider the projective version, that is

$$\Sigma_s^\theta(\mathbb{R}^n) = \bigcap_{A,B>0} \mathcal{S}_{s,B}^{\theta,A}(\mathbb{R}^n)$$

equipped with the projective limit topology. When $\theta = s$ we simply write $\mathcal{S}_\theta, \Sigma_\theta$ instead of $\mathcal{S}_\theta^\theta, \Sigma_\theta^\theta$. We can also define, for $C, \varepsilon > 0$, the normed space $\mathcal{S}_{s,C}^{\theta,\varepsilon}(\mathbb{R}^n)$ given by the functions $f \in C^\infty(\mathbb{R}^n)$ such that there is $C > 0$ satisfying

$$\|f\|_{s,\theta}^{\varepsilon,C} := \sup_{\substack{x \in \mathbb{R}^n \\ \alpha \in \mathbb{N}_0^n}} C^{-|\alpha|} \alpha!^{-\theta} e^{\varepsilon|x|^{\frac{1}{s}}} |\partial_x^\alpha f(x)| < \infty,$$

and we have (with equivalent topologies)

$$\mathcal{S}_s^\theta(\mathbb{R}^n) = \bigcup_{C,\varepsilon>0} \mathcal{S}_{s,C}^{\theta,\varepsilon}(\mathbb{R}^n) \quad \text{and} \quad \Sigma_s^\theta(\mathbb{R}^n) = \bigcap_{C,\varepsilon>0} \mathcal{S}_{s,C}^{\theta,\varepsilon}(\mathbb{R}^n).$$

The following inclusions are continuous (for every $\varepsilon > 0$)

$$\Sigma_s^\theta(\mathbb{R}^n) \subset \mathcal{S}_s^\theta(\mathbb{R}^n) \subset \Sigma_{s+\varepsilon}^{\theta+\varepsilon}(\mathbb{R}^n).$$

All the previous spaces can be written in terms of the Gelfand-Shilov Sobolev spaces $H_{\rho;s,\theta}^m$, with $\rho, m \in \mathbb{R}^2$ defined in the Introduction. Namely, we have

$$\mathcal{S}_s^\theta(\mathbb{R}^n) = \bigcup_{\substack{\rho \in \mathbb{R}^2 \\ \rho_j > 0, j=1,2}} H_{\rho;s,\theta}^m(\mathbb{R}^n), \quad \Sigma_s^\theta(\mathbb{R}^n) = \bigcap_{\substack{\rho \in \mathbb{R}^2 \\ \rho_j > 0, j=1,2}} H_{\rho;s,\theta}^m(\mathbb{R}^n).$$

From now and on we shall denote by $(\mathcal{S}_s^\theta)'(\mathbb{R}^n), (\Sigma_s^\theta)'(\mathbb{R}^n)$ the respective dual spaces. Concerning the action of the Fourier transform we have the following isomorphisms

$$\begin{aligned} \mathcal{F} : \Sigma_s^\theta(\mathbb{R}^n) &\rightarrow \Sigma_\theta^s(\mathbb{R}^n), \quad \mathcal{F} : \mathcal{S}_s^\theta(\mathbb{R}^n) \rightarrow \mathcal{S}_\theta^s(\mathbb{R}^n), \\ \mathcal{F} : H_{(\rho_1,\rho_2);s,\theta}^{(m_1,m_2)}(\mathbb{R}^n) &\rightarrow H_{(\rho_2,\rho_1);\theta,s}^{(m_2,m_1)}(\mathbb{R}^n). \end{aligned}$$

2.2. Pseudodifferential operators of infinite order. We start defining the symbol classes of infinite order.

Definition 1. Let $\tau \in \mathbb{R}, \kappa, \theta, \mu, \nu > 1$ and $C, c > 0$.

- (i) We denote by $\mathbf{SG}_{\mu,\nu;\kappa}^{\tau,\infty}(\mathbb{R}^{2n}; C, c)$ the Banach space of all functions $p \in C^\infty(\mathbb{R}^{2n})$ satisfying the following condition:

$$\|p\|_{C,c} := \sup_{\alpha,\beta \in \mathbb{N}_0^n} C^{-|\alpha+\beta|} \alpha!^{-\mu} \beta!^{-\nu} \sup_{x,\xi \in \mathbb{R}^n} \langle \xi \rangle^{-\tau+|\alpha|} \langle x \rangle^{|\beta|} e^{-c|x|^{\frac{1}{\kappa}}} |\partial_\xi^\alpha \partial_x^\beta p(x, \xi)| < \infty.$$

We set $\mathbf{SG}_{\mu,\nu;\kappa}^{\tau,\infty}(\mathbb{R}^{2n}) := \bigcup_{C,c>0} \mathbf{SG}_{\mu,\nu;\kappa}^{\tau,\infty}(\mathbb{R}^{2n}; C, c)$ with the topology of inductive limit of the Banach spaces $\mathbf{SG}_{\mu,\nu;\kappa}^{\tau,\infty}(\mathbb{R}^{2n}; C, c)$.

- (ii) We denote by $\mathbf{SG}_{\mu,\nu;\theta}^{\infty,\tau}(\mathbb{R}^{2n}; C, c)$ the Banach space of all functions $p \in C^\infty(\mathbb{R}^{2n})$ satisfying the following condition:

$$\|p\|^{C,c} := \sup_{\alpha,\beta \in \mathbb{N}_0^n} C^{-|\alpha+\beta|} \alpha!^{-\mu} \beta!^{-\nu} \sup_{x,\xi \in \mathbb{R}^n} \langle \xi \rangle^{|\alpha|} \langle x \rangle^{-\tau+|\beta|} e^{-c|\xi|^{\frac{1}{\theta}}} |\partial_\xi^\alpha \partial_x^\beta p(x, \xi)| < \infty.$$

We set $\mathbf{SG}_{\mu,\nu;\theta}^{\infty,\tau}(\mathbb{R}^{2n}) := \bigcup_{C,c>0} \mathbf{SG}_{\mu,\nu;\theta}^{\infty,\tau}(\mathbb{R}^{2n}; C, c)$ with the topology of inductive limit of the spaces $\mathbf{SG}_{\mu,\nu;\theta}^{\infty,\tau}(\mathbb{R}^{2n}; C, c)$.

We also need the following symbol classes of finite order.

Definition 2. Let $\mu, \nu \geq 1, m = (m_1, m_2) \in \mathbb{R}^2$ and $C > 0$. We denote by $\mathbf{SG}_{\mu,\nu}^m(\mathbb{R}^{2n}; C)$ the Banach space of all functions $p \in C^\infty(\mathbb{R}^{2n})$ satisfying the following condition:

$$\sup_{\alpha,\beta \in \mathbb{N}_0^n} C^{-|\alpha+\beta|} \alpha!^{-\mu} \beta!^{-\nu} \sup_{x,\xi \in \mathbb{R}^n} \langle \xi \rangle^{-m_1+|\alpha|} \langle x \rangle^{-m_2+|\beta|} |\partial_\xi^\alpha \partial_x^\beta p(x, \xi)| < \infty.$$

We set $\mathbf{SG}_{\mu,\nu}^m(\mathbb{R}^{2n}) := \bigcup_{C>0} \mathbf{SG}_{\mu,\nu}^m(\mathbb{R}^{2n}; C)$.

Finally we say that $p \in \mathbf{SG}^m(\mathbb{R}^{2n})$ if for any $\alpha, \beta \in \mathbb{N}_0^n$ there is $C_{\alpha, \beta} > 0$ satisfying

$$|\partial_\xi^\alpha \partial_x^\beta p(x, \xi)| \leq C_{\alpha, \beta} \langle \xi \rangle^{m_1 - |\alpha|} \langle x \rangle^{m_2 - |\beta|}, \quad x, \xi \in \mathbb{R}^n.$$

When $\mu = \nu$ we write $\mathbf{SG}_\mu^m(\mathbb{R}^{2n})$, $\mathbf{SG}_{\mu, \kappa}^{\tau, \infty}(\mathbb{R}^{2n})$, $\mathbf{SG}_{\mu, \theta}^{\infty, \tau}(\mathbb{R}^{2n})$ instead of $\mathbf{SG}_{\mu, \mu}^m(\mathbb{R}^{2n})$, $\mathbf{SG}_{\mu, \mu; \kappa}^{\tau, \infty}(\mathbb{R}^{2n})$, $\mathbf{SG}_{\mu, \mu; \theta}^{\infty, \tau}(\mathbb{R}^{2n})$.

As usual, given a symbol $p(x, \xi)$ we shall denote by $p(x, D)$ or by $\text{op}(p)$ the pseudodifferential operator defined as standard by

$$p(x, D)u(x) = \int e^{ix \cdot \xi} p(x, \xi) \widehat{u}(\xi) d\xi, \quad x \in \mathbb{R}^n,$$

where u belongs to some suitable function space depending on the assumptions on p , and $d\xi$ stands for $(2\pi)^{-n} d\xi$. We have the following continuity results.

Proposition 1. *Let $\tau \in \mathbb{R}$, $s > \mu \geq 1$, $\nu \geq 1$ and $p \in \mathbf{SG}_{\mu, \nu; s}^{\tau, \infty}(\mathbb{R}^{2n})$. Then for every $\theta > \nu$ the pseudodifferential operator $p(x, D)$ with symbol $p(x, \xi)$ is continuous on $\Sigma_s^\theta(\mathbb{R}^n)$ and it extends to a continuous map on $(\Sigma_s^\theta)'(\mathbb{R}^n)$.*

Proposition 2. *Let $\tau \in \mathbb{R}$, $\theta > \nu \geq 1$, $\mu \geq 1$ and $p \in \mathbf{SG}_{\mu, \nu; \theta}^{\infty, \tau}(\mathbb{R}^{2n})$. Then for every $s > \mu$ the operator $p(x, D)$ is continuous on $\Sigma_s^\theta(\mathbb{R}^n)$ and it extends to a continuous map on $(\Sigma_s^\theta)'(\mathbb{R}^n)$.*

The proof of Propositions 1 and 2 can be derived following the argument in the proof of [6, Proposition 2.3] and [1, Proposition 1]. We leave details to the reader.

Now we define the notion of asymptotic expansion and recall some fundamental results, which can be found in the Appendix A of [6]. For $t_1, t_2 \geq 0$ set

$$Q_{t_1, t_2} = \{(x, \xi) \in \mathbb{R}^{2n} : \langle x \rangle < t_1 \text{ and } \langle \xi \rangle < t_2\}$$

and $Q_{t_1, t_2}^e = \mathbb{R}^{2n} \setminus Q_{t_1, t_2}$. When $t_1 = t_2 = t$ we simply write Q_t and Q_t^e .

Definition 3. *We say that:*

- (i) $\sum_{j \geq 0} a_j \in FSG_{\mu, \nu; \kappa}^{\tau, \infty}$ if $a_j(x, \xi) \in C^\infty(\mathbb{R}^{2n})$ and there are $C, c, B > 0$ satisfying

$$|\partial_\xi^\alpha \partial_x^\beta a_j(x, \xi)| \leq C^{|\alpha| + |\beta| + 2j + 1} \alpha!^\mu \beta!^\nu j!^{\mu + \nu - 1} \langle \xi \rangle^{\tau - |\alpha| - j} \langle x \rangle^{-|\beta| - j} e^{c|x|^\frac{1}{\kappa}},$$

for every $\alpha, \beta \in \mathbb{N}_0^n$, $j \geq 0$ and $(x, \xi) \in Q_{B(j)}^e$, where $B(j) := Bj^{\mu + \nu - 1}$;

- (ii) $\sum_{j \geq 0} a_j \in FSG_{\mu, \nu; \theta}^{\infty, \tau}$ if $a_j(x, \xi) \in C^\infty(\mathbb{R}^{2n})$ and there are $C, c, B > 0$ satisfying

$$|\partial_\xi^\alpha \partial_x^\beta a_j(x, \xi)| \leq C^{|\alpha| + |\beta| + 2j + 1} \alpha!^\mu \beta!^\nu j!^{\mu + \nu - 1} \langle \xi \rangle^{-|\alpha| - j} e^{c|\xi|^\frac{1}{\theta}} \langle x \rangle^{\tau - |\beta| - j},$$

for every $\alpha, \beta \in \mathbb{N}_0^n$, $j \geq 0$ and $(x, \xi) \in Q_{B(j)}^e$;

- (iii) $\sum_{j \geq 0} a_j \in FSG_{\mu, \nu}^m$ if $a_j(x, \xi) \in C^\infty(\mathbb{R}^{2n})$ and there are $C, B > 0$ satisfying

$$|\partial_\xi^\alpha \partial_x^\beta a_j(x, \xi)| \leq C^{|\alpha| + |\beta| + 2j + 1} \alpha!^\mu \beta!^\nu j!^{\mu + \nu - 1} \langle \xi \rangle^{m_1 - |\alpha| - j} \langle x \rangle^{m_2 - |\beta| - j},$$

for every $\alpha, \beta \in \mathbb{N}_0^n$, $j \geq 0$ and $(x, \xi) \in Q_{B(j)}^e$.

Definition 4. *Let $\sum_{j \geq 0} a_j$, $\sum_{j \geq 0} b_j$ in $FSG_{\mu, \nu; \kappa}^{\tau, \infty}$. We say that $\sum_{j \geq 0} a_j \sim \sum_{j \geq 0} b_j$ in $FSG_{\mu, \nu; \kappa}^{\tau, \infty}$ if there are $C, c, B > 0$ satisfying*

$$|\partial_\xi^\alpha \partial_x^\beta \sum_{j < N} (a_j - b_j)(x, \xi)| \leq C^{|\alpha| + |\beta| + 2N + 1} \alpha!^\mu \beta!^\nu N!^{\mu + \nu - 1} \langle \xi \rangle^{\tau - |\alpha| - N} \langle x \rangle^{-|\beta| - N} e^{c|x|^\frac{1}{\kappa}},$$

for every $\alpha, \beta \in \mathbb{N}_0^n$, $N \geq 1$ and $(x, \xi) \in Q_{B(N)}^e$. Analogous definitions for the classes $FSG_{\mu, \nu; \theta}^{\infty, \tau}$, $FSG_{\mu, \nu}^m$.

Remark 4. If $\sum_{j \geq 0} a_j \in FSG_{\mu,\nu;\kappa}^{\tau,\infty}$, then $a_0 \in SG_{\mu,\nu;\kappa}^{\tau,\infty}$. Given $a \in SG_{\mu,\nu;\kappa}^{\tau,\infty}$ and setting $b_0 = a$, $b_j = 0$, $j \geq 1$, we have $a = \sum_{j \geq 0} b_j$. Hence we can consider $SG_{\mu,\nu;\kappa}^{\tau,\infty}$ as a subset of $FSG_{\mu,\nu;\kappa}^{\tau,\infty}$.

Proposition 3. Given $\sum_{j \geq 0} a_j \in FSG_{\mu,\nu;\kappa}^{\tau,\infty}$, there exists $a \in SG_{\mu,\nu;\kappa}^{\tau,\infty}$ such that $a \sim \sum_{j \geq 0} a_j$ in $FSG_{\mu,\nu;\kappa}^{\tau,\infty}$. Analogous results for the classes $FSG_{\mu,\nu;\theta}^{\infty,\tau}$ and $FSG_{\mu,\nu}^m$.

Proposition 4. Let $a \in SG_{\mu,\nu;\kappa}^{0,\infty}$ such that $a \sim 0$ in $FSG_{\mu,\nu;\kappa}^{0,\infty}$. If $\kappa > \mu + \nu - 1$, then $a \in \mathcal{S}_\delta(\mathbb{R}^{2n})$ for every $\delta \geq \mu + \nu - 1$. Analogous results for the classes $FSG_{\mu,\nu;\theta}^{\infty,\tau}$ and $FSG_{\mu,\nu}^m$.

Concerning the symbolic calculus and the continuous mapping properties on the Gelfand-Shilov Sobolev spaces we have the following results, cf. [6, Propositions A.12 and A.13].

Theorem 2. Let $p \in SG_{\mu,\nu;\kappa}^{\tau,\infty}(\mathbb{R}^{2n})$, $q \in SG_{\mu,\nu;\kappa}^{\tau',\infty}(\mathbb{R}^{2n})$ with $\kappa > \mu + \nu - 1$. Then the L^2 adjoint p^* and the composition $p \circ q$ have the following structure:

$$p^*(x, D) = a(x, D) + r(x, D) \text{ where } r \in \mathcal{S}_{\mu+\nu-1}(\mathbb{R}^{2n}), a \in SG_{\mu,\nu;\kappa}^{\tau,\infty}(\mathbb{R}^{2n}), \text{ and}$$

$$a(x, \xi) \sim \sum_{\alpha} \frac{1}{\alpha!} \partial_{\xi}^{\alpha} D_x^{\alpha} \overline{p(x, \xi)} \text{ in } FSG_{\mu,\nu;\kappa}^{\tau,\infty}(\mathbb{R}^{2n});$$

$$p(x, D) \circ q(x, D) = b(x, D) + s(x, D), \text{ where } s \in \mathcal{S}_{\mu+\nu-1}(\mathbb{R}^{2n}), b \in SG_{\mu,\nu;\kappa}^{\tau+\tau',\infty}(\mathbb{R}^{2n}) \text{ and}$$

$$b(x, \xi) \sim \sum_{\alpha} \frac{1}{\alpha!} \partial_{\xi}^{\alpha} p(x, \xi) D_x^{\alpha} q(x, \xi) \text{ in } FSG_{\mu,\nu;\kappa}^{\tau+\tau',\infty}(\mathbb{R}^{2n}).$$

Analogous results for the classes $SG_{\mu,\nu;\theta}^{\infty,\tau}(\mathbb{R}^{2n})$ and $SG_{\mu,\nu}^m(\mathbb{R}^{2n})$.

Theorem 3. Let $p \in SG_{\mu,\nu}^{m'}(\mathbb{R}^{2n})$ for some $m' \in \mathbb{R}^2$. Then for every $m, \rho \in \mathbb{R}^2$ and s, θ such that $\min\{s, \theta\} > \mu + \nu - 1$ the operator $p(x, D)$ maps $H_{\rho;s,\theta}^m(\mathbb{R}^n)$ into $H_{\rho;s,\theta}^{m-m'}(\mathbb{R}^n)$ continuously.

A simple application of the Faà formula gives us the following result.

Proposition 5. If $\lambda \in SG_{\mu}^{0,\frac{1}{\kappa}}(\mathbb{R}^{2n})$, then $e^{\lambda} \in SG_{\mu;\kappa}^{0,\infty}(\mathbb{R}^{2n})$. If $\lambda \in SG_{\mu}^{\frac{1}{\theta},0}(\mathbb{R}^{2n})$, then $e^{\lambda} \in SG_{\mu;\theta}^{\infty,0}(\mathbb{R}^{2n})$.

We conclude this section proving the following theorem.

Theorem 4. Let $\rho, m \in \mathbb{R}^2$ and $s, \theta, \mu > 1$ with $\min\{s, \theta\} > 2\mu - 1$. Let $\lambda \in SG_{\mu}^{0,\frac{1}{\kappa}}(\mathbb{R}^{2n})$. Then:

- i) if $\kappa > s$, the operator $e^{\lambda}(x, D) : H_{\rho;s,\theta}^m(\mathbb{R}^n) \longrightarrow H_{\rho-\delta e_2;s,\theta}^m(\mathbb{R}^n)$ is continuous for every $\delta > 0$, where $e_2 = (0, 1)$;
- ii) if $\kappa = s$, the operator $e^{\lambda}(x, D) : H_{\rho;s,\theta}^m(\mathbb{R}^n) \longrightarrow H_{\rho-\delta e_2;s,\theta}^m(\mathbb{R}^n)$ is continuous for every

$$\delta > C(\lambda) := \sup_{(x,\xi) \in \mathbb{R}^{2n}} \frac{\lambda(x, \xi)}{\langle x \rangle^{1/s}}.$$

Proof. (i) Let $\phi(x) \in \gamma^{\mu}(\mathbb{R}^n)$, that is, a uniform Gevrey function of index μ , be a cut-off function such that, for a positive constant K , $\phi(x) = 1$ for $|x| < K/2$, $\phi(x) = 0$ for $|x| > K$ and $0 \leq \phi(x) \leq 1$ for every $x \in \mathbb{R}^n$. We split the symbol $e^{\lambda(x,\xi)}$ as

$$(2.1) \quad e^{\lambda(x,\xi)} = \phi(x) e^{\lambda(x,\xi)} + (1 - \phi(x)) e^{\lambda(x,\xi)} = a_1(x, \xi) + a_2(x, \xi).$$

Since ϕ has compact support and λ has order zero with respect to ξ , we have $a_1 \in \mathbf{SG}_{\mu}^{0,0}$. On the other hand, given any $\delta > 0$ and choosing K large enough, since $\kappa > s$ we may write $|\lambda(x, \xi)|\langle x \rangle^{-1/s} < \delta$ on the support of $a_2(x, \xi)$. Hence we obtain

$$a_2(x, \xi) = e^{\delta\langle x \rangle^{1/s}} (1 - \phi(x)) e^{\lambda(x, \xi) - \delta\langle x \rangle^{1/s}},$$

with $(1 - \phi(x))e^{\lambda(x, \xi) - \delta\langle x \rangle^{1/s}}$ of order $(0, 0)$ because $\lambda(x, \xi) - \delta\langle x \rangle^{1/s} < 0$ on the support of $(1 - \phi(x))$. Thus, (2.1) becomes

$$e^{\lambda(x, \xi)} = a_1(x, \xi) + e^{\delta\langle x \rangle^{1/s}} \tilde{a}_2(x, \xi),$$

a_1 and \tilde{a}_2 of order $(0, 0)$. Since by Theorem 3 the operators $a_1(x, D)$ and $\tilde{a}_2(x, D)$ map continuously $H_{\rho, s, \theta}^m$ into itself, then we obtain i). The proof of (ii) follows a similar argument and can be found in [6, Theorem 2.4]. \square

3. Spectral invariance for SG - Ψ DO with Gevrey estimates

Let $p \in \mathbf{SG}^{0,0}(\mathbb{R}^{2n})$, then $p(x, D)$ extends to a continuous operator on $L^2(\mathbb{R}^n)$. Suppose that $p(x, D) : L^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n)$ is bijective. The question is to determine whether or not the inverse p^{-1} is also an \mathbf{SG} operator of order $(0, 0)$. This is known as the spectral invariance problem and it has an affirmative answer, see [12].

Following the ideas presented in [12, pp. 51-57], we will prove that the symbol of p^{-1} satisfies Gevrey estimates, whenever the symbol $p \in \mathbf{SG}_{\mu, \nu}^{0,0}(\mathbb{R}^{2n})$. This is an important step in the study of the continuous mapping properties of $\{e^\Lambda(x, D)\}^{-1}$ on Gelfand-Shilov Sobolev spaces $H_{\rho, s, \theta}^m$.

Theorems 5, 6, 7 here below can be found in [33, Chapters 20, 21].

Theorem 5. *Let X, Y separable Hilbert spaces. Then a bounded operator $A : X \rightarrow Y$ is Fredholm if and only if there are $B : Y \rightarrow X$ bounded, $K_1 : X \rightarrow Y$ and $K_2 : Y \rightarrow X$ compact operators such that*

$$BA = I_X - K_1, \quad AB = I_Y - K_2.$$

Theorem 6. *Let X, Y, Z be separable Hilbert spaces and let $A : X \rightarrow Y$, $B : Y \rightarrow Z$ be Fredholm operators. Then:*

- $B \circ A : X \rightarrow Z$ is Fredholm and $i(BA) = i(B) + i(A)$, where $i(\cdot)$ stands for the index of a Fredholm operator;
- $Y = N(A^t) \oplus R(A)$.

Remark 5. *Let X be a Hilbert space and $K : X \rightarrow X$ a compact operator. Then $I - K$ is Fredholm and $i(I - K) = 0$.*

Theorem 7. *Let $p \in \mathbf{SG}^{m_1, m_2}(\mathbb{R}^{2n})$ such that $p(x, D) : H^{s_1 + m_1, s_2 + m_2}(\mathbb{R}^n) \rightarrow H^{m_1, m_2}(\mathbb{R}^n)$ is Fredholm for some $s_1, s_2 \in \mathbb{R}$. Then p is \mathbf{SG} -elliptic, that is there exist $C, R > 0$ such that*

$$|p(x, \xi)| \geq C \langle \xi \rangle^{m_1} \langle x \rangle^{m_2} \quad \text{for } (x, \xi) \in Q_R^c.$$

Theorem 8. *Let $p \in \mathbf{SG}_{\mu, \nu}^{m_1, m_2}(\mathbb{R}^{2n})$ be \mathbf{SG} -elliptic. Then there is $q \in \mathbf{SG}_{\mu, \nu}^{-m_1, -m_2}(\mathbb{R}^{2n})$ such that*

$$p(x, D) \circ q(x, D) = I + r_1(x, D), \quad q(x, D) \circ p(x, D) = I + r_2(x, D),$$

where $r_1, r_2 \in S_{\mu + \nu - 1}(\mathbb{R}^{2n})$.

Proof. See [30, Theorem 6.3.16]. \square

In order to prove the main result of this section, we need the following technical lemma.

Lemma 1. *Let $A : L^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n)$ be a bounded operator such that A and A^* map $L^2(\mathbb{R}^n)$ into $\Sigma_r(\mathbb{R}^n)$ continuously. Then the Schwartz kernel of A belongs to $\Sigma_r(\mathbb{R}^{2n})$.*

Proof. Since $\Sigma_r(\mathbb{R}^n) \subset L^2(\mathbb{R}^n)$ is a nuclear Fréchet space, (cf. [13]), by [21, Propositions 2.1.7 and 2.1.8], we have that A is defined by a kernel $H(x, y)$ and we can write

$$H(x, y) = \sum_{j \in \mathbb{N}_0} a_j f_j(x) g_j(y) = \sum_{j \in \mathbb{N}_0} \tilde{a}_j \tilde{f}_j(x) \tilde{g}_j(y),$$

where $a_j, \tilde{a}_j \in \mathbb{C}$, $\tilde{f}_j(x), g_j(y) \in \Sigma_r(\mathbb{R}^n)$, $f_j(x), \tilde{g}_j(y) \in L^2(\mathbb{R}^n)$, $\sum_j |a_j| < \infty$, $\sum_j |\tilde{a}_j| < \infty$, $\tilde{f}_j(x), g_j(y)$ converge to zero in $\Sigma_r(\mathbb{R}^n)$ and $f_j(x), \tilde{g}_j(y)$ converge to zero in $L^2(\mathbb{R}^n)$.

We now use the following characterization: $H \in \Sigma_r(\mathbb{R}^{2n})$ if and only if

$$\sup_{\alpha, \beta \in \mathbb{N}_0^n} \left\| \frac{x^\alpha y^\beta H(x, y)}{C^{|\alpha|+|\beta|} \alpha!^r \beta!^r} \right\|_{L^2} < \infty \quad \text{and} \quad \sup_{\alpha, \beta \in \mathbb{N}_0^n} \left\| \frac{\xi^\alpha \eta^\beta \widehat{H}(\xi, \eta)}{C^{|\alpha|+|\beta|} \alpha!^r \beta!^r} \right\|_{L^2} < \infty$$

for every $C > 0$, and prove that both the latter conditions hold. Note that

$$\begin{aligned} \|y^\beta H(x, y)\|_{L^2}^2 &= \iint \left| \sum_{j \in \mathbb{N}_0} a_j f_j(x) y^\beta g_j(y) \right|^2 dx dy \\ &= \iint \left| \sum_{j \in \mathbb{N}_0} a_j^{\frac{1}{2}} f_j(x) a_j^{\frac{1}{2}} y^\beta g_j(y) \right|^2 dx dy \\ &\leq \int \sum_{j \in \mathbb{N}_0} |a_j^{\frac{1}{2}} f_j(x)|^2 dx \int \sum_{j \in \mathbb{N}_0} |a_j^{\frac{1}{2}} y^\beta g_j(y)|^2 dy \\ &= \sum_{j \in \mathbb{N}_0} |a_j| \|f_j\|_{L^2}^2 \sum_{j \in \mathbb{N}_0} |a_j| \|y^\beta g_j\|_{L^2}^2. \end{aligned}$$

Since g_j converges to zero in $\Sigma_r(\mathbb{R}^n)$, we have

$$\|y^\beta g_j(y)\|_{L^2} = \left\| \frac{y^\beta g_j(y)}{C^{|\beta|} \beta!^r} \right\|_{L^2} C^{|\beta|} \beta!^r \leq C^{|\beta|} \beta!^r \sup_{j \in \mathbb{N}_0} \left\| \frac{y^\beta g_j(y)}{C^{|\beta|} \beta!^r} \right\|_{L^2},$$

for every $C > 0$, and therefore

$$\|y^\beta H(x, y)\|_{L^2} \leq \left(\sum_{j \in \mathbb{N}_0} |a_j| \right) \sup_{j \in \mathbb{N}_0} \|f_j\|_{L^2} \sup_{j \in \mathbb{N}_0} \left\| \frac{y^\beta g_j(y)}{C^{|\beta|} \beta!^r} \right\|_{L^2} C^{|\beta|} \beta!^r.$$

Hence

$$\sup_{\beta \in \mathbb{N}_0^n} \left\| \frac{y^\beta H(x, y)}{C^{|\beta|} \beta!^r} \right\|_{L^2} < \infty \iff \sup_{N \in \mathbb{N}_0} \left\| \frac{\langle y \rangle^N H(x, y)}{C^N N!^r} \right\|_{L^2} < \infty,$$

for every $C > 0$. Using the representation $\sum \tilde{a}_j \tilde{f}_j(x) \tilde{g}_j(y)$, analogously we can obtain

$$\sup_{\alpha \in \mathbb{N}_0^n} \left\| \frac{x^\alpha H(x, y)}{C^{|\alpha|} \alpha!^r} \right\|_{L^2} < \infty \iff \sup_{N \in \mathbb{N}_0} \left\| \frac{\langle x \rangle^N H(x, y)}{C^N N!^r} \right\|_{L^2} < \infty,$$

for every $C > 0$.

Now note that, for every $N \in \mathbb{N}_0$, $x, y \in \mathbb{R}^n$,

$$\langle x, y \rangle^N = (\langle x \rangle^2 + |y|^2)^{\frac{N}{2}} \leq (\langle x \rangle + \langle y \rangle)^N \leq 2^{N-1} (\langle x \rangle^N + \langle y \rangle^N).$$

Therefore, for every $C > 0$,

$$\left\| \frac{\langle x, y \rangle^N H(x, y)}{C^N N!^r} \right\|_{L^2} \leq \left\| \frac{\langle x \rangle^N H(x, y)}{C_1^N N!^r} \right\|_{L^2} + \left\| \frac{\langle y \rangle^N H(x, y)}{C_1^N N!^r} \right\|_{L^2},$$

where $C_1 = (2^{-1}C)^N$. Hence, for every $C > 0$,

$$\sup_{N \in \mathbb{N}_0} \left\| \frac{\langle x, y \rangle^N H(x, y)}{C^N N!^r} \right\|_{L^2} < \infty.$$

Since the Fourier transformation is an isomorphism on L^2 and on Σ_r , we have

$$\widehat{H}(\xi, \eta) = \sum_{j \in \mathbb{N}_0} a_j \widehat{f}_j(\xi) \widehat{g}_j(\eta) = \sum_{j \in \mathbb{N}_0} \widetilde{a}_j \widehat{\widetilde{f}}_j(\xi) \widehat{\widetilde{g}}_j(\eta),$$

where $a_j, \widetilde{a}_j \in \mathbb{C}$, $\widehat{f}_j(\xi), \widehat{g}_j(\eta) \in \Sigma_r(\mathbb{R}^n)$, $\widehat{\widetilde{f}}_j(\xi), \widehat{\widetilde{g}}_j(\eta) \in L^2(\mathbb{R}^n)$, $\sum_j |a_j| < \infty$, $\sum_j |\widetilde{a}_j| < \infty$, $\widehat{\widetilde{f}}_j(\xi), \widehat{\widetilde{g}}_j(\eta)$ converge to zero in $\Sigma_r(\mathbb{R}^n)$ and $\widehat{f}_j(\xi), \widehat{g}_j(\eta)$ converge to zero in $L^2(\mathbb{R}^n)$. In an analogous way as before we get, for every $C > 0$,

$$\sup_{N \in \mathbb{N}_0} \left\| \frac{\langle \xi, \eta \rangle^N \widehat{H}(\xi, \eta)}{C^N N!^r} \right\|_{L^2} < \infty.$$

Hence $H \in \Sigma_r(\mathbb{R}^{2n})$. □

Theorem 9. *Let $p \in \mathbf{SG}_{\mu, \nu}^{0,0}(\mathbb{R}^{2n})$ such that $p(x, D) : L^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n)$ is bijective. Then $\{p(x, D)\}^{-1} : L^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n)$ is a pseudodifferential operator given by a symbol $\tilde{p} = q + \tilde{k}$ where $q \in \mathbf{SG}_{\mu, \nu}^{0,0}(\mathbb{R}^{2n})$ and $\tilde{k} \in \Sigma_r(\mathbb{R}^{2n})$ for every $r > \mu + \nu - 1$.*

Proof. Since $p(x, D) : L^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n)$ is bijective, then $p(x, D)$ is Fredholm and

$$i(p(x, D)) = \dim N(p(x, D)) - \dim N(p^t(x, D)) = 0.$$

Therefore by Theorem 7 p is **SG**-elliptic and by Theorem 8 there is $q \in \mathbf{SG}_{\mu, \nu}^{0,0}(\mathbb{R}^{2n})$ such that

$$q(x, D) \circ p(x, D) = I + r(x, D), \quad p(x, D) \circ q(x, D) = I + s(x, D),$$

for some $r, s \in \mathcal{S}_{\mu+\nu-1}(\mathbb{R}^{2n})$. In particular $r(x, D), s(x, D)$ are compact operators on $L^2(\mathbb{R}^n)$. By Theorem 5 $q(x, D)$ is a Fredholm operator and we have

$$i(q(x, D)) = i(q(x, D)) + i(p(x, D)) = i(q(x, D) \circ p(x, D)) = i(I + r(x, D)) = 0.$$

Note that $N(q(x, D))$ and $N(q^t(x, D))$ are subspaces of $\mathcal{S}_{\mu+\nu-1}(\mathbb{R}^n)$. Indeed, let $f \in N(q)$ and $g \in N(q^t)$, then

$$0 = p(x, D) \circ q(x, D)f = (I + s(x, D))f \implies f = -s(x, D)f,$$

$$0 = p^t(x, D) \circ q^t(x, D)g = (q(x, D) \circ p(x, D))^t g = (I + r(x, D))^t g \implies g = -r^t(x, D)g.$$

Since $L^2(\mathbb{R}^n)$ is a separable Hilbert space and $N(q(x, D))$ is closed, we have the following decompositions

$$L^2 = N(q) \oplus N(q)^\perp, \quad L^2 = N(q^t) \oplus R_{L^2}(q),$$

where $R_{L^2}(q)$ denotes the range of $q(x, D)$ as an operator on $L^2(\mathbb{R}^n)$.

Let $\pi : L^2 \rightarrow N(q)$ the projection of L^2 onto $N(q)$ with null space $N(q)^\perp$, $F : N(q) \rightarrow N(q^t)$ an isomorphism and $i : N(q^t) \rightarrow L^2$ the inclusion. Set $Q = i \circ F \circ \pi$. Then $Q : L^2 \rightarrow L^2$ is bounded and its image is contained in $N(q^t) \subset \mathcal{S}_{\mu+\nu-1}$. It is not difficult to see that $Q^* = \tilde{i} \circ F^* \circ \pi_{N(q^t)}$, where \tilde{i} is the inclusion of $N(q)$ into L^2 and $\pi_{N(q^t)}$ is the orthogonal projection of L^2 onto $N(q)$. Since $\mathcal{S}_{\mu+\nu-1} \subset \Sigma_r$, then by Lemma 1, Q is given by a kernel in Σ_r .

We will now show that $q + Q$ is a bijective parametrix of p . Indeed, let $u = u_1 + u_2 \in N(q) \oplus N(q^t)$ such that $(q + Q)u = 0$. Then $0 = qu_2 + (i \circ F)u_1 \in R_{L^2}(q) \oplus N(q^t)$. Hence $qu_2 = 0$ and $i \circ Fu_1 = 0$ which implies that $u = 0$. In order to prove that Q is surjective, consider $f = f_1 + f_2 \in R_{L^2}(q) \oplus N(q^t)$. There exist $u_1 \in L^2$ and $u_2 \in N(q)$ such that $qu_1 = f_1$ and $Fu_2 = f_2$. Now write $u_1 = v_1 + v_2 \in N(q) \oplus N(q)^\perp$. Then $q(u_1) = q(v_2)$ and therefore $(q + Q)(v_2 + u_2) = f_1 + f_2 = f$. Finally notice that

$$p(x, D) \circ (q(x, D) + Q) = I + s(x, D) + p(x, D) \circ Q = I + s'(x, D),$$

$$(q(x, D) + Q) \circ p(x, D) = I + r(x, D) + Q \circ p(x, D) = I + r'(x, D),$$

where $r', s' \in \Sigma_r(\mathbb{R}^{2n})$.

Now set $\tilde{q} = q(x, D) + Q$. Therefore $\tilde{q} \circ p(x, D) = I + r'(x, D) : L^2 \rightarrow L^2$ is bijective. Set $k = -(I + r')^{-1} \circ r'$. Then $(I + r')(I + k) = I$ and $k = -r' - r'k$. Observe that

$$k^t = -\{r'\}^t - k^t\{r'\}^t = -\{r'\}^t + \{r'\}^t\{(I + r')^{-1}\}^t\{r'\}^t.$$

Hence k, k^t map L^2 into Σ_r and by Lemma 1 we have that k is given by a kernel in $\Sigma_r(\mathbb{R}^{2n})$.

To finish the proof, it is enough to notice that

$$p^{-1} \circ \tilde{q}^{-1} = (\tilde{q} \circ p)^{-1} = (I + r')^{-1} = I + k \implies p^{-1} = (I + k) \circ \tilde{q}.$$

□

4. Change of variables

4.1. Definition and properties of λ_2 and λ_1 . Let $M_2, M_1 > 0$ and $h \geq 1$ to be chosen later on. We define

$$(4.1) \quad \lambda_2(x, \xi) = M_2 w\left(\frac{\xi}{h}\right) \int_0^x \langle y \rangle^{-\sigma} dy, \quad (x, \xi) \in \mathbb{R}^2,$$

$$(4.2) \quad \lambda_1(x, \xi) = M_1 w\left(\frac{\xi}{h}\right) \langle \xi \rangle_h^{-1} \int_0^x \langle y \rangle^{-\frac{\sigma}{2}} \psi\left(\frac{\langle y \rangle^\sigma}{\langle \xi \rangle_h^2}\right) dy, \quad (x, \xi) \in \mathbb{R}^2,$$

where

$$w(\xi) = \begin{cases} 0, & |\xi| \leq 1 \\ -\text{sgn}(\partial_\xi a_3(t, \xi)), & |\xi| > R_{a_3} \end{cases}, \quad \psi(y) = \begin{cases} 1, & |y| \leq \frac{1}{2} \\ 0, & |y| \geq 1 \end{cases},$$

$|\partial^\alpha w(\xi)| \leq C_w^{\alpha+1} \alpha!^\mu$, $|\partial^\beta \psi(y)| \leq C_\psi^{\beta+1} \beta!^\mu$ for some $\mu > 1$ which will be chosen later. Notice that by the assumption (i) of Theorem 1 the function $w(\xi)$ is constant for $\xi \geq R_{a_3}$ and for $\xi \leq -R_{a_3}$.

Lemma 2. *Let $\lambda_2(x, \xi)$ as in (4.1). Then there exists $C > 0$ such that for all $\alpha, \beta \in \mathbb{N}$ and $(x, \xi) \in \mathbb{R}^2$:*

- (i) $|\lambda_2(x, \xi)| \leq \frac{M_2}{1-\sigma} \langle x \rangle^{1-\sigma}$;
- (ii) $|\partial_x^\beta \lambda_2(x, \xi)| \leq M_2 C^\beta \beta! \langle x \rangle^{1-\sigma-\beta}$, for $\beta \geq 1$;
- (iii) $|\partial_\xi^\alpha \partial_x^\beta \lambda_2(x, \xi)| \leq M_2 C^{\alpha+\beta+1} \alpha!^\mu \beta!^\mu \chi_{E_{h, R_{a_3}}}(\xi) \langle \xi \rangle_h^{-\alpha} \langle x \rangle^{1-\sigma-\beta}$, for $\alpha \geq 1, \beta \geq 0$,

where $E_{h, R_{a_3}} = \{\xi \in \mathbb{R} : h \leq \xi \leq R_{a_3} h\}$. In particular $\lambda_2 \in \mathbf{SG}_\mu^{0, 1-\sigma}(\mathbb{R}^2)$.

Proof. First note that

$$|\lambda_2(x, \xi)| = M_2 \left| w\left(\frac{\xi}{h}\right) \right| \int_0^{|x|} \langle y \rangle^{-\sigma} dy \leq M_2 \int_0^{\langle x \rangle} y^{-\sigma} dy = \frac{M_2}{1-\sigma} \langle x \rangle^{1-\sigma}.$$

For $\beta \geq 1$

$$|\partial_x^\beta \lambda_2(x, \xi)| \leq M_2 \left| w \left(\frac{\xi}{h} \right) \right| |\partial_x^{\beta-1} \langle x \rangle^{-\sigma}| \leq M_2 C^{\beta-1} (\beta-1)! \langle x \rangle^{1-\sigma-\beta}.$$

For $\alpha \geq 1$

$$\begin{aligned} |\partial_\xi^\alpha \lambda_2(x, \xi)| &\leq M_2 h^{-\alpha} \left| w^{(\alpha)} \left(\frac{\xi}{h} \right) \right| \int_0^{\langle x \rangle} y^{-\sigma} dy \\ &\leq \frac{M_2}{1-\sigma} C_w^{\alpha+1} \langle R_{a_3} \rangle^\alpha \alpha!^\mu \chi_{E_{h,R_{a_3}}}(\xi) \langle \xi \rangle_h^{-\alpha} \langle x \rangle^{1-\sigma}. \end{aligned}$$

Finally, for $\alpha, \beta \geq 1$

$$\begin{aligned} |\partial_\xi^\alpha \partial_x^\beta \lambda_2(x, \xi)| &\leq M_2 h^{-\alpha} \left| w^{(\alpha)} \left(\frac{\xi}{h} \right) \right| \partial_x^{\beta-1} \langle x \rangle^{-\sigma} \\ &\leq M_2 C_w^{\alpha+1} \langle R_{a_3} \rangle^\alpha C^{\beta-1} \alpha!^\mu (\beta-1)! \chi_{E_{h,R_{a_3}}}(\xi) \langle \xi \rangle_h^{-\alpha} \langle x \rangle^{1-\sigma-\beta}. \end{aligned}$$

□

For the function λ_1 we can prove the following alternative estimates.

Lemma 3. *Let $\lambda_1(x, \xi)$ as in (4.2). Then there exists $C > 0$ such that for all $\alpha, \beta \geq 0$ and $(x, \xi) \in \mathbb{R}^2$:*

- (i) $|\partial_\xi^\alpha \partial_x^\beta \lambda_1(x, \xi)| \leq M_1 C^{\alpha+\beta+1} (\alpha! \beta!)^\mu \langle \xi \rangle_h^{-1-\alpha} \langle x \rangle^{1-\frac{\sigma}{2}-\beta};$
- (ii) $|\partial_\xi^\alpha \partial_x^\beta \lambda_1(x, \xi)| \leq M_1 C^{\alpha+\beta+1} (\alpha! \beta!)^\mu \langle \xi \rangle_h^{-\alpha} \langle x \rangle^{1-\sigma-\beta}.$

In particular $\lambda_1 \in \mathcal{SG}_\mu^{0,1-\sigma}(\mathbb{R}^2)$.

Proof. Denote by $\chi_\xi(x)$ the characteristic function of the set $\{x \in \mathbb{R} : \langle x \rangle^\sigma \leq \langle \xi \rangle_h^2\}$. For $\alpha = \beta = 0$ we have

$$|\lambda_1(x, \xi)| \leq M_1 \left| w \left(\frac{\xi}{h} \right) \right| \langle \xi \rangle_h^{-1} \int_0^{\langle x \rangle} y^{-\frac{\sigma}{2}} dy \leq \frac{2}{2-\sigma} M_1 \langle \xi \rangle_h^{-1} \langle x \rangle^{1-\frac{\sigma}{2}},$$

and

$$|\lambda_1(x, \xi)| \leq M_1 \left| w \left(\frac{\xi}{h} \right) \right| \int_0^{\langle x \rangle} \langle \xi \rangle_h^{-1} \langle y \rangle^{-\frac{\sigma}{2}} \chi_\xi(y) dy \leq \frac{M_1}{1-\sigma} \langle x \rangle^{1-\sigma}.$$

For $\alpha \geq 1$, with the aid of Faà di Bruno formula, we have

$$\begin{aligned}
|\partial_\xi^\alpha \lambda_1(x, \xi)| &\leq M_1 \sum_{\alpha_1+\alpha_2+\alpha_3=\alpha} \frac{\alpha!}{\alpha_1!\alpha_2!\alpha_3!} h^{-\alpha_1} \left| w^{(\alpha_1)} \left(\frac{\xi}{h} \right) \right| \partial_\xi^{\alpha_2} \langle \xi \rangle_h^{-1} \left| \int_0^x \langle y \rangle^{-\frac{\sigma}{2}} \partial_\xi^{\alpha_3} \psi \left(\frac{\langle y \rangle^\sigma}{\langle \xi \rangle_h^2} \right) dy \right| \\
&\leq M_1 \sum_{\alpha_1+\alpha_2+\alpha_3=\alpha} \frac{\alpha!}{\alpha_1!\alpha_2!\alpha_3!} C_w^{\alpha_1+1} \langle R_{a_3} \rangle^{\alpha_3} \alpha_1! \langle \xi \rangle_h^{-\alpha_1} C^{\alpha_2} \alpha_2! \langle \xi \rangle_h^{-1-\alpha_2} \\
&\quad \times \int_0^{\langle x \rangle} \langle y \rangle^{-\frac{\sigma}{2}} \chi_\xi(y) \sum_{j=1}^{\alpha_3} \frac{|\psi^{(j)} \left(\frac{\langle y \rangle^\sigma}{\langle \xi \rangle_h^2} \right)|}{j!} \sum_{\gamma_1+\dots+\gamma_j=\alpha_3} \frac{\alpha_3!}{\gamma_1! \dots \gamma_j!} \prod_{\ell=1}^j \partial_\xi^{\gamma_\ell} \langle \xi \rangle_h^{-2} \langle y \rangle^\sigma dy \\
&\leq M_1 \sum_{\alpha_1+\alpha_2+\alpha_3=\alpha} \frac{\alpha!}{\alpha_1!\alpha_2!\alpha_3!} C_w^{\alpha_1+1} \langle R_{a_3} \rangle^{\alpha_3} \alpha_1! \langle \xi \rangle_h^{-\alpha_1} C^{\alpha_2} \alpha_2! \langle \xi \rangle_h^{-1-\alpha_2} \\
&\quad \times \int_0^{\langle x \rangle} \langle y \rangle^{-\frac{\sigma}{2}} \chi_\xi(y) \sum_{j=1}^{\alpha_3} C_\psi^{j+1} j!^{\mu-1} \sum_{\gamma_1+\dots+\gamma_j=\alpha_3} \frac{\alpha_3!}{\gamma_1! \dots \gamma_j!} \prod_{\ell=1}^j C^{\gamma_\ell+1} \gamma_\ell! \langle \xi \rangle_h^{-\gamma_\ell} dy \\
&\leq M_1 C_{\{w, \psi, \sigma, R_{a_3}\}}^{\alpha+1} \alpha!^\mu \langle \xi \rangle_h^{-1-\alpha} \langle y \rangle^{1-\frac{\sigma}{2}},
\end{aligned}$$

and

$$\begin{aligned}
|\partial_\xi^\alpha \lambda_1(x, \xi)| &\leq M_1 \sum_{\alpha_1+\alpha_2+\alpha_3=\alpha} \frac{\alpha!}{\alpha_1!\alpha_2!\alpha_3!} h^{-\alpha_1} \left| w^{(\alpha_1)} \left(\frac{\xi}{h} \right) \right| \partial_\xi^{\alpha_2} \langle \xi \rangle_h^{-1} \left| \int_0^x \langle y \rangle^{-\frac{\sigma}{2}} \partial_\xi^{\alpha_3} \psi \left(\frac{\langle y \rangle^\sigma}{\langle \xi \rangle_h^2} \right) dy \right| \\
&\leq M_1 \sum_{\alpha_1+\alpha_2+\alpha_3=\alpha} \frac{\alpha!}{\alpha_1!\alpha_2!\alpha_3!} C_w^{\alpha_1+1} \langle R_{a_3} \rangle^{\alpha_3} \alpha_1! \langle \xi \rangle_h^{-\alpha_1} C^{\alpha_2} \alpha_2! \langle \xi \rangle_h^{-\alpha_2} \\
&\quad \times \int_0^{\langle x \rangle} \langle \xi \rangle_h^{-1} \langle y \rangle^{-\frac{\sigma}{2}} \chi_\xi(y) \sum_{j=1}^{\alpha_3} \frac{|\psi^{(j)} \left(\frac{\langle y \rangle^\sigma}{\langle \xi \rangle_h^2} \right)|}{j!} \sum_{\gamma_1+\dots+\gamma_j=\alpha_3} \frac{\alpha_3!}{\gamma_1! \dots \gamma_j!} \prod_{\ell=1}^j \partial_\xi^{\gamma_\ell} \langle \xi \rangle_h^{-2} \langle y \rangle^\sigma dy \\
&\leq M_1 \sum_{\alpha_1+\alpha_2+\alpha_3=\alpha} \frac{\alpha!}{\alpha_1!\alpha_2!\alpha_3!} C_w^{\alpha_1+1} \langle R_{a_3} \rangle^{\alpha_3} \alpha_1! \langle \xi \rangle_h^{-\alpha_1} C^{\alpha_2} \alpha_2! \langle \xi \rangle_h^{-\alpha_2} \\
&\quad \times \int_0^{\langle x \rangle} \langle y \rangle^{-\sigma} \chi_\xi(y) \sum_{j=1}^{\alpha_3} C_\psi^{j+1} j!^{\mu-1} \sum_{\gamma_1+\dots+\gamma_j=\alpha_3} \frac{\alpha_3!}{\gamma_1! \dots \gamma_j!} \prod_{\ell=1}^j C^{\gamma_\ell+1} \gamma_\ell! \langle \xi \rangle_h^{-\gamma_\ell} dy \\
&\leq M_1 C_{\{w, \psi, \sigma, R_{a_3}\}}^{\alpha+1} \alpha!^\mu \langle \xi \rangle_h^{-\alpha} \langle y \rangle^{1-\sigma}.
\end{aligned}$$

For $\beta \geq 1$ we have

$$\begin{aligned}
|\partial_x^\beta \lambda_1(x, \xi)| &\leq M_1 \langle \xi \rangle_h^{-1} \chi_\xi(x) \sum_{\beta_1 + \beta_2 = \beta - 1} \frac{(\beta - 1)!}{\beta_1! \beta_2!} \partial_x^{\beta_1} \langle x \rangle^{-\frac{\sigma}{2}} \sum_{j=1}^{\beta_2} \frac{|\psi^{(j)}(\frac{\langle x \rangle^\sigma}{\langle \xi \rangle_h^2})|}{j!} \\
&\times \sum_{\delta_1 + \dots + \delta_j = \beta_2} \frac{\beta_2!}{\delta_1! \dots \delta_j!} \prod_{\ell=1}^j \langle \xi \rangle_h^{-2} \partial_x^{\delta_\ell} \langle x \rangle^\sigma \\
&\leq M_1 \langle \xi \rangle_h^{-1} \chi_\xi(x) \sum_{\beta_1 + \beta_2 = \beta - 1} \frac{(\beta - 1)!}{\beta_1! \beta_2!} C^{\beta_1+1} \beta_1!^\mu \langle x \rangle^{-\frac{\sigma}{2} - \beta_1} \sum_{j=1}^{\beta_2} C_\psi^{j+1} j!^{\mu-1} \\
&\times \sum_{\delta_1 + \dots + \delta_j = \beta_2} \frac{\beta_2!}{\delta_1! \dots \delta_j!} \prod_{\ell=1}^j C^{\delta_\ell+1} \delta_\ell! \langle x \rangle^{-\delta_\ell} \\
&\leq M_1 C_\psi^{\alpha+\beta+1} (\beta - 1)^\mu \langle \xi \rangle_h^{-1} \chi_\xi(x) \langle x \rangle^{1-\sigma-\beta} \\
&\leq M_1 C_\psi^{\alpha+\beta+1} (\beta - 1)^\mu \langle x \rangle^{1-\sigma-\beta}.
\end{aligned}$$

Finally, for $\alpha, \beta \geq 1$ we have

$$\begin{aligned}
|\partial_\xi^\alpha \partial_x^\beta \lambda_1(x, \xi)| &\leq M_1 \sum_{\alpha_1 + \alpha_2 + \alpha_3 = \alpha} \frac{\alpha!}{\alpha_1! \alpha_2! \alpha_3!} h^{-\alpha_1} \left| w^{(\alpha_1)} \left(\frac{\xi}{h} \right) \right| \partial_\xi^{\alpha_2} \langle \xi \rangle_h^{-1} \sum_{\beta_1 + \beta_2 = \beta - 1} \frac{(\beta - 1)!}{\beta_1! \beta_2!} \\
&\times \partial_x^{\beta_1} \langle x \rangle^{\frac{\sigma}{2}} \left| \partial_\xi^{\alpha_3} \partial_x^{\beta_2} \psi \left(\frac{\langle x \rangle^\sigma}{\langle \xi \rangle_h^2} \right) \right| \\
&\leq M_1 \chi_\xi(x) \sum_{\alpha_1 + \alpha_2 + \alpha_3 = \alpha} \frac{\alpha!}{\alpha_1! \alpha_2! \alpha_3!} h^{-\alpha_1} \left| w^{(\alpha_1)} \left(\frac{\xi}{h} \right) \right| \partial_\xi^{\alpha_2} \langle \xi \rangle_h^{-1} \sum_{\beta_1 + \beta_2 = \beta - 1} \frac{(\beta - 1)!}{\beta_1! \beta_2!} \partial_x^{\beta_1} \langle x \rangle^{-\frac{\sigma}{2}} \\
&\times \sum_{j=1}^{\alpha_3 + \beta_2} \frac{|\psi^{(j)}(\frac{\langle x \rangle^\sigma}{\langle \xi \rangle_h^2})|}{j!} \sum_{\gamma_1 + \dots + \gamma_j = \alpha_3} \sum_{\delta_1 + \dots + \delta_j = \beta_2} \frac{\alpha_3!}{\gamma_1! \dots \gamma_j!} \frac{\beta_2!}{\delta_1! \dots \delta_j!} \prod_{\ell=1}^j \partial_\xi^{\gamma_\ell} \langle \xi \rangle_h^{-2} \partial_x^{\delta_\ell} \langle x \rangle^\sigma \\
&\leq M_1 \chi_\xi(x) \sum_{\alpha_1 + \alpha_2 + \alpha_3 = \alpha} \frac{\alpha!}{\alpha_1! \alpha_2! \alpha_3!} C_w^{\alpha_1+1} \alpha_1!^\mu \langle R_{a_3} \rangle^{\alpha_1} \langle \xi \rangle_h^{-\alpha_1} C^{\alpha_2+1} \alpha_2! \langle \xi \rangle_h^{-1-\alpha_2} \\
&\times \sum_{\beta_1 + \beta_2 = \beta - 1} \frac{(\beta - 1)!}{\beta_1! \beta_2!} C^{\beta_1+1} \beta_1! \langle x \rangle^{-\frac{1}{2}(1-\frac{1}{s})-\beta_1} \sum_{j=1}^{\alpha_3 + \beta_2} C_\psi^{j+1} j!^{\mu-1} \\
&\times \sum_{\gamma_1 + \dots + \gamma_j = \alpha_3} \sum_{\delta_1 + \dots + \delta_j = \beta_2} \frac{\alpha_3!}{\gamma_1! \dots \gamma_j!} \frac{\beta_2!}{\delta_1! \dots \delta_j!} \prod_{\ell=1}^j C^{\gamma_\ell+1} \gamma_\ell! \langle \xi \rangle_h^{-2-\gamma_\ell} C^{\delta_\ell+1} \delta_\ell! \langle x \rangle^{\sigma-\delta_\ell} \\
&\leq M_1 \chi_\xi(x) C_{\{w, \sigma, \psi, R_{a_3}\}}^{\alpha+\beta+1} \alpha!^\mu (\beta - 1)^\mu \langle \xi \rangle_h^{-1-\alpha} \langle x \rangle^{1-\frac{\sigma}{2}-\beta} \\
&\leq M_1 C_{\{w, \sigma, \psi, R_{a_3}\}}^{\alpha+\beta+1} \alpha!^\mu (\beta - 1)^\mu \langle \xi \rangle_h^{-\alpha} \langle x \rangle^{1-\sigma-\beta}.
\end{aligned}$$

□

4.2. Invertibility of $e^{\tilde{\Lambda}}$, $\tilde{\Lambda} = \lambda_2 + \lambda_1$. In this section we construct an inverse for the operator $e^{\tilde{\Lambda}}(x, D)$ with $\tilde{\Lambda}(x, \xi) = \lambda_2(x, \xi) + \lambda_1(x, \xi)$ and we prove that the inverse acts continuously on Gelfand Shilov-Sobolev spaces. By Lemmas 2 and 3 we have $\tilde{\Lambda} \in \mathbf{SG}_\mu^{0,1-\sigma}(\mathbb{R}^2)$. Therefore, by Proposition 5, $e^{\tilde{\Lambda}} \in \mathbf{SG}_{\mu;1/(1-\sigma)}^{0,\infty}(\mathbb{R}^2)$. To construct the inverse of $e^{\tilde{\Lambda}}(x, D)$ we need to use the

L^2 adjoint of $e^{-\tilde{\Lambda}}(x, D)$, denoted in the sequel by ${}^R e^{-\tilde{\Lambda}}$ and defined as an oscillatory integral by

$${}^R e^{-\tilde{\Lambda}} u(x) = \iint e^{i(x-y)\xi} e^{-\tilde{\Lambda}(y, \xi)} u(y) dy d\xi.$$

Assuming $\mu > 1$ such that $1/(1-\sigma) > 2\mu - 1$, by results from calculus, we may write

$${}^R e^{-\tilde{\Lambda}} = a_1(x, D) + r_1(x, D),$$

where $a_1 \sim \sum_{\alpha} \frac{1}{\alpha!} \partial_{\xi}^{\alpha} D_x^{\alpha} e^{-\tilde{\Lambda}}$ in $FSG_{\mu; 1/(1-\sigma)}^{0, \infty}(\mathbb{R}^2)$, $r_1 \in \mathcal{S}_{2\mu-1}(\mathbb{R}^2)$, and

$$e^{\tilde{\Lambda}} \circ {}^R e^{-\tilde{\Lambda}} = e^{\tilde{\Lambda}} \circ a_1(x, D) + e^{\tilde{\Lambda}} \circ r_1(x, D) = a_2(x, D) + r_2(x, D) + e^{\tilde{\Lambda}} \circ r_1(x, D),$$

where

$$a_2 \sim \sum_{\alpha, \beta} \frac{1}{\alpha! \beta!} \partial_{\xi}^{\alpha} e^{\tilde{\Lambda}} \partial_{\xi}^{\beta} D_x^{\alpha+\beta} e^{-\tilde{\Lambda}} = \sum_{\gamma} \frac{1}{\gamma!} \partial_{\xi}^{\gamma} (e^{\tilde{\Lambda}} D_x^{\gamma} e^{-\tilde{\Lambda}}) \text{ in } FSG_{\mu; 1/(1-\sigma)}^{0, \infty}(\mathbb{R}^2)$$

and $r_2 \in \mathcal{S}_{2\mu-1}(\mathbb{R}^2)$. Therefore

$$e^{\tilde{\Lambda}} \circ {}^R e^{-\tilde{\Lambda}} = a(x, D) + r(x, D),$$

where $a \sim \sum_{\gamma} \frac{1}{\gamma!} \partial_{\xi}^{\gamma} (e^{\tilde{\Lambda}} D_x^{\gamma} e^{-\tilde{\Lambda}})$ in $FSG_{\mu; 1/(1-\sigma)}^{0, \infty}(\mathbb{R}^2)$ and $r \in \mathcal{S}_{2\mu-1}(\mathbb{R}^2)$.

Now let us study more carefully the asymptotic expansion

$$\sum_{\gamma \geq 0} \frac{1}{\gamma!} \partial_{\xi}^{\gamma} (e^{\tilde{\Lambda}} D_x^{\gamma} e^{-\tilde{\Lambda}}) = \sum_{\gamma \geq 0} r_{1, \gamma}.$$

Note that

$$e^{\tilde{\Lambda}(x, \xi)} D_x^{\gamma} e^{-\tilde{\Lambda}(x, \xi)} = \sum_{j=1}^{\gamma} \frac{(-1)^j}{j!} \sum_{\gamma_1 + \dots + \gamma_j = \gamma} \frac{\gamma!}{\gamma_1! \dots \gamma_j!} \prod_{\ell=1}^j D_x^{\gamma_{\ell}} \tilde{\Lambda}(x, \xi),$$

hence, for $\alpha, \beta \geq 0$,

$$\begin{aligned} |\partial_{\xi}^{\alpha} \partial_x^{\beta} r_{1, \gamma}| &\leq \frac{1}{\gamma!} \sum_{j=1}^{\gamma} \frac{1}{j!} \sum_{\gamma_1 + \dots + \gamma_j = \gamma} \frac{\gamma!}{\gamma_1! \dots \gamma_j!} \sum_{\alpha_1 + \dots + \alpha_j = \alpha + \gamma} \sum_{\beta_1 + \dots + \beta_j = \beta} \frac{(\alpha + \gamma)!}{\alpha_1! \dots \alpha_j!} \frac{\beta!}{\beta_1! \dots \beta_j!} \\ &\quad \times \prod_{\ell=1}^j |\partial_{\xi}^{\alpha_{\ell}} \partial_x^{\beta_{\ell} + \gamma_{\ell}} \tilde{\Lambda}(x, \xi)| \\ &\leq \frac{1}{\gamma!} \sum_{j=1}^{\gamma} \frac{1}{j!} \sum_{\gamma_1 + \dots + \gamma_j = \gamma} \frac{\gamma!}{\gamma_1! \dots \gamma_j!} \sum_{\alpha_1 + \dots + \alpha_j = \alpha + \gamma} \sum_{\beta_1 + \dots + \beta_j = \beta} \frac{(\alpha + \gamma)!}{\alpha_1! \dots \alpha_j!} \frac{\beta!}{\beta_1! \dots \beta_j!} \\ &\quad \times \prod_{\ell=1}^j C_{\tilde{\Lambda}}^{\alpha_{\ell} + \beta_{\ell} + \gamma_{\ell} + 1} \alpha_{\ell}!^{\mu} (\beta_{\ell} + \gamma_{\ell})!^{\mu} \langle \xi \rangle_h^{-\alpha_{\ell}} \langle x \rangle^{1-\sigma-\beta_{\ell}-\gamma_{\ell}} \\ &\leq C^{\alpha+\beta+2\gamma+1} \alpha!^{\mu} \beta!^{\mu} \gamma!^{2\mu-1} \langle \xi \rangle_h^{-\alpha-\gamma} \sum_{j=1}^{\gamma} \frac{\langle x \rangle^{(1-\sigma)j-\beta-\gamma}}{j!}. \end{aligned}$$

In the following we shall consider the sets

$$Q_{t_1, t_2; h} = \{(x, \xi) \in \mathbb{R}^2 : \langle x \rangle < t_1 \text{ and } \langle \xi \rangle_h < t_2\}$$

and $Q_{t_1, t_2; h}^e = \mathbb{R}^2 \setminus Q_{t_1, t_2; h}$. When $t_1 = t_2 = t$ we simply write $Q_{t; h}$ and $Q_{t; h}^e$.

Let $\psi(x, \xi) \in C^{\infty}(\mathbb{R}^2)$ such that $\psi \equiv 0$ on $Q_{2; h}$, $\psi \equiv 1$ on $Q_{3; h}^e$, $0 \leq \psi \leq 1$ and

$$|\partial_{\xi}^{\alpha} \partial_x^{\beta} \psi(x, \xi)| \leq C_{\psi}^{\alpha+\beta+1} \alpha!^{\mu} \beta!^{\mu},$$

for every $x, \xi \in \mathbb{R}$ and $\alpha, \beta \in \mathbb{N}_0$. Now set $\psi_0 \equiv 1$ and, for $j \geq 1$,

$$\psi_j(x, \xi) := \psi\left(\frac{x}{R(j)}, \frac{\xi}{R(j)}\right),$$

where $R(j) := Rj^{2\mu-1}$ and $R > 0$ is a large constant. Let us recall that

- $(x, \xi) \in Q_{3R(j)}^e \implies \left(\frac{x}{R(j)}, \frac{\xi}{R(j)}\right) \in Q_3^e \implies \psi_i(x, \xi) = 1$, for $i \leq j$;
- $(x, \xi) \in Q_{R(j)} \implies \left(\frac{x}{R(j)}, \frac{\xi}{R(j)}\right) \in Q_2 \implies \psi_i(x, \xi) = 0$, for $i \geq j$.

Defining $b(x, \xi) = \sum_{j \geq 0} \psi_j(x, \xi) r_{1,j}(x, \xi)$ we have that $b \in \mathbf{SG}_{\mu; \frac{1}{1-\sigma}}^{0, \infty}(\mathbb{R}^2)$ and

$$b(x, \xi) \sim \sum_{j \geq 0} r_{1,j}(x, \xi) \text{ in } FSG_{\mu; \frac{1}{1-\sigma}}^{0, \infty}(\mathbb{R}^2).$$

We will show that $b \in SG_\mu^{0,0}(\mathbb{R}^{2n})$. Indeed, first we write

$$b(x, \xi) = 1 + \sum_{j \geq 1} \psi_j(x, \xi) r_{1,j}(x, \xi) = 1 + \sum_{j \geq 0} \psi_{j+1}(x, \xi) r_{1,j+1}(x, \xi).$$

On the support of $\partial_\xi^{\alpha_1} \partial_x^{\beta_1} \psi_{j+1}$ we have

$$\langle x \rangle \leq 3R(j+1) \quad \text{and} \quad \langle \xi \rangle_h \leq 3R(j+1),$$

whenever $\alpha_1 + \beta_1 \geq 1$. Hence

$$\begin{aligned} |\partial_\xi^\alpha \partial_x^\beta \sum_{j \geq 0} \psi_{j+1} r_{1,j+1}(x, \xi)| &\leq \sum_{\substack{\alpha_1 + \alpha_2 = \alpha \\ \beta_1 + \beta_2 = \beta}} \frac{\alpha!}{\alpha_1! \alpha_2!} \frac{\beta!}{\beta_1! \beta_2!} |\partial_\xi^{\alpha_1} \partial_x^{\beta_1} \psi_{j+1}(x, \xi)| |\partial_\xi^{\alpha_2} \partial_x^{\beta_2} r_{1,j+1}(x, \xi)| \\ &\leq \sum_{j \geq 0} \sum_{\substack{\alpha_1 + \alpha_2 = \alpha \\ \beta_1 + \beta_2 = \beta}} \frac{\alpha!}{\alpha_1! \alpha_2!} \frac{\beta!}{\beta_1! \beta_2!} \frac{1}{R(j+1)^{(\alpha_1 + \beta_1)}} C_\psi^{\alpha_1 + \beta_1 + 1} \alpha_1!^\mu \beta_1!^\mu \\ &\quad \times C^{\alpha_2 + \beta_2 + 2(j+1)+1} \alpha_2!^\mu \beta_2!^\mu (j+1)!^{2\mu-1} \langle \xi \rangle_h^{-\alpha_2 - (j+1)} \langle x \rangle^{-\beta_1 - (j+1)} \sum_{\ell=1}^{j+1} \frac{\langle x \rangle^{(1-\sigma)\ell}}{\ell!} \\ &\leq \sum_{j \geq 0} \sum_{\substack{\alpha_1 + \alpha_2 = \alpha \\ \beta_1 + \beta_2 = \beta}} \frac{\alpha!}{\alpha_1! \alpha_2!} \frac{\beta!}{\beta_1! \beta_2!} \frac{1}{R(j+1)^{(\alpha_1 + \beta_1)}} C_\psi^{\alpha_1 + \beta_1 + 1} \alpha_1!^\mu \beta_1!^\mu \\ &\quad \times C^{\alpha_2 + \beta_2 + 2(j+1)+1} \alpha_2!^\mu \beta_2!^\mu (j+1)!^{2\mu-1} \langle \xi \rangle_h^{-\alpha_2 - (j+1)} \langle x \rangle^{-\sigma - \beta_1} \sum_{\ell=1}^{j+1} \frac{\langle x \rangle^{(1-\sigma)(\ell-1)-j}}{\ell!} \\ &\leq \tilde{C}^{\alpha + \beta + 1} (\alpha! \beta!)^\mu \langle \xi \rangle_h^{-1-\alpha} \langle x \rangle^{-\sigma - \beta} \sum_{j \geq 0} C^{2j} (j+1)!^{2\mu-1} \langle \xi \rangle_h^{-j} \sum_{\ell=0}^{j+1} \frac{\langle x \rangle^{(1-\sigma)(\ell-1)-j}}{\ell!}. \end{aligned}$$

We also have that

$$\langle x \rangle \geq R(j+1) \quad \text{or} \quad \langle \xi \rangle_h \geq R(j+1)$$

holds true on the support of $\partial_\xi^{\alpha_1} \partial_x^{\beta_1} \psi_{j+1}$. If $\langle \xi \rangle_h \geq R(j+1)$, then

$$\langle \xi \rangle_h^{-j} \leq R^{-j} (j+1)^{-j(2\mu-1)} \leq R^{-j} (j+1)!^{-(2\mu-1)}.$$

On the other hand, since we are assuming $\mu > 1$ such that $2\mu - 1 < \frac{1}{1-\sigma}$, if $\langle x \rangle \leq R(j+1)$ we obtain

$$\begin{aligned} \langle x \rangle^{(1-\sigma)(\ell-1)-j} &\leq R^{(1-\sigma)(\ell-1)-j} \{(j+1)^{2\mu-1}\}^{(1-\sigma)(\ell-1)-j} \\ &\leq R^{-\sigma j} (j+1)^{\ell-1-j(2\mu-1)} \\ &= R^{-\sigma j} (j+1)^{\ell-1} (j+1)!^{-(2\mu-1)}. \end{aligned}$$

Enlarging $R > 0$ if necessary, we can infer that $\sum_{j \geq 1} r_{1,j} \in \mathbf{SG}_\mu^{-1,-\sigma}(\mathbb{R}^2)$.

In analogous way it is possible to prove that $\sum_{j \geq k} r_{1,j} \in \mathbf{SG}_\mu^{-k,-\sigma k}(\mathbb{R}^2)$. Hence, we may conclude

$$b(x, \xi) - \sum_{j < k} r_{1,j}(x, \xi) \in \mathbf{SG}_\mu^{-k,-\sigma k}(\mathbb{R}^2), \quad k \in \mathbb{N},$$

that is, $b \sim \sum_j r_{1,j}$ in $\mathbf{SG}_\mu^{0,0}(\mathbb{R}^2)$.

Since $a \sim \sum_j r_{1,j}$ in $FSG_{\mu;1/(1-\sigma)}^{0,\infty}(\mathbb{R}^2)$, $b \sim \sum_j r_{1,j}$ in $FSG_{\mu;1/(1-\sigma)}^{0,\infty}(\mathbb{R}^2)$ we have $a - b \in \mathcal{S}_{2\mu-1}(\mathbb{R}^2)$. Thus we may write

$$e^{\tilde{\Lambda}}(x, D) \circ^R e^{-\tilde{\Lambda}} = I + \tilde{r}(x, D) + \bar{r}(x, D) = I + r(x, D),$$

where $\tilde{r} \in \mathbf{SG}_\mu^{-1,-\sigma}(\mathbb{R}^2)$, $\tilde{r} \sim \sum_{\gamma \geq 1} r_{1,\gamma}$ in $\mathbf{SG}_\mu^{-1,-\sigma}(\mathbb{R}^2)$ and $\bar{r} \in \mathcal{S}_{2\mu-1}(\mathbb{R}^2)$. In particular $r \in \mathbf{SG}_{2\mu-1}^{-1,-\sigma}(\mathbb{R}^2)$, therefore we obtain

$$\begin{aligned} |\partial_\xi^\alpha \partial_x^\beta r(x, \xi)| &\leq C_{\alpha,\beta} \langle \xi \rangle_h^{-1-\alpha} \langle x \rangle^{-\sigma-\beta} \\ &\leq C_{\alpha,\beta} h^{-1} \langle \xi \rangle_h^{-\alpha} \langle x \rangle^{-\sigma-\beta}. \end{aligned}$$

This implies that the $(0,0)$ -seminorms of r are bounded by h^{-1} . Choosing h large enough, we obtain that $I + r(x, D)$ is invertible on $L^2(\mathbb{R})$ and its inverse $(I + r(x, D))^{-1}$ is given by the Neumann series $\sum_{j \geq 0} (-r(x, D))^j$.

By Theorem 9 we have

$$(I + r(x, D))^{-1} = q(x, D) + k(x, D),$$

where $q \in \mathbf{SG}_{2\mu-1}^{0,0}(\mathbb{R}^2)$, $k \in \Sigma_\delta(\mathbb{R}^2)$ for every $\delta > 2(2\mu - 1) - 1 = 4\mu - 3$. Choosing $\mu > 1$ close enough to 1, we have that δ can be chosen arbitrarily close to 1. Hence, by Theorem 3, for every fixed $s > 1, \theta > 1$, we can find $\mu > 1$ such that

$$(I + r(x, D))^{-1} : H_{\rho';s,\theta}^{m'}(\mathbb{R}) \rightarrow H_{\rho';s,\theta}^{m'}(\mathbb{R})$$

is continuous for every $m', \rho' \in \mathbb{R}^2$. Analogously one can show the existence of a left inverse of $e^{\tilde{\Lambda}}$ with the same properties. Summing up, we obtain the following result.

Lemma 4. *Let $s, \theta > 1$ and take $\mu > 1$ such that $\min\{s, \theta\} > 4\mu - 3$. For $h > 0$ large enough, the operator $e^{\tilde{\Lambda}}(x, D)$ is invertible on $L^2(\mathbb{R})$ and on $\Sigma_{\min\{s,\theta\}}(\mathbb{R})$ and its inverse is given by*

$$\{e^{\tilde{\Lambda}}(x, D)\}^{-1} =^R e^{-\tilde{\Lambda}(x,D)} \circ (I + r(x, D))^{-1} =^R e^{-\tilde{\Lambda}(x,D)} \circ \sum_{j \geq 0} (-r(x, D))^j,$$

where $r \in \mathbf{SG}_{2\mu-1}^{-1,-\sigma}(\mathbb{R}^2)$ and $r \sim \sum_{\gamma \geq 1} \frac{1}{\gamma!} \partial_\xi^\gamma (e^{\tilde{\Lambda}} D_x^\gamma e^{-\tilde{\Lambda}})$ in $\mathbf{SG}_{2\mu-1}^{-1,-\sigma}(\mathbb{R}^2)$. Moreover, the symbol of $(I + r(x, D))^{-1}$ belongs to $\mathbf{SG}_\delta^{0,0}(\mathbb{R}^2)$ for every $\delta > 4\mu - 3$ and it maps continuously $H_{\rho';s,\theta}^{m'}(\mathbb{R})$ into itself for any $\rho', m' \in \mathbb{R}^2$.

We conclude this section writing $\{e^{\tilde{\Lambda}}(x, D)\}^{-1}$ in a more precise way. From the asymptotic expansion of the symbol $r(x, \xi)$ we may write

$$\{e^{\tilde{\Lambda}}(x, D)\}^{-1} =^R e^{-\tilde{\Lambda}} \circ (I - r(x, D) + (r(x, D))^2 + q_{-3}(x, D)),$$

where q_{-3} denotes an operator with symbol in $\mathbf{SG}_\delta^{-3,-3\sigma}(\mathbb{R}^2)$ for every $\delta > 4\mu - 3$. Now note that

$$r = i\partial_\xi \partial_x \tilde{\Lambda} + \frac{1}{2} \partial_\xi^2 (\partial_x^2 \tilde{\Lambda} - [\partial_x \tilde{\Lambda}]^2) + q_{-3} = q_{-1} + q_{-2} + q_{-3}.$$

Hence

$$\begin{aligned} (r(x, D))^2 &= (q_{-1} + q_{-2} + q_{-3})(x, D) \circ (q_{-1} + q_{-2} + q_{-3})(x, D) \\ &= q_{-1}(x, D) \circ q_{-1}(x, D) + q_{-3}(x, D) \\ &= \text{op} \left\{ -[\partial_\xi \partial_x \tilde{\Lambda}]^2 + q_{-3} \right\} \end{aligned}$$

for a new element q_{-3} in the same space. We finally obtain:

$$(4.3) \quad \{e^{\tilde{\Lambda}}(x, D)\}^{-1} = {}^R e^{-\tilde{\Lambda}} \circ \left[I + \text{op} \left(-i\partial_\xi \partial_x \tilde{\Lambda} - \frac{1}{2} \partial_\xi^2 (\partial_x^2 \tilde{\Lambda} - [\partial_x \tilde{\Lambda}]^2) - [\partial_\xi \partial_x \tilde{\Lambda}]^2 + q_{-3} \right) \right],$$

where $q_{-3} \in \mathbf{SG}_\delta^{-3,-3\sigma}(\mathbb{R}^2)$. Since we deal with operators whose order does not exceed 3, in the next sections we are going to use frequently formula (4.3) for the inverse of $e^{\tilde{\Lambda}}(x, D)$.

5. Conjugation of iP

In this section we will perform the conjugation of iP by the operator $e^{\rho_1 \langle D \rangle^{\frac{1}{\theta}}} \circ e^{\Lambda(t, x, D)}$ and its inverse, where $\Lambda(t, x, \xi) = k(t) \langle x \rangle_h^{1-\sigma} + \tilde{\Lambda}(x, \xi)$ and $k \in C^1([0, T]; \mathbb{R})$ is a non increasing function such that $k(T) \geq 0$. Since the arguments in the following involve also derivatives with respect to t these derivatives will be denoted by D_t , whereas the symbol D in the notation for pseudodifferential operators will always correspond to derivatives with respect to x .

More precisely, we will compute

$$e^{\rho_1 \langle D \rangle^{\frac{1}{\theta}}} \circ e^{k(t) \langle x \rangle_h^{1-\sigma}} \circ e^{\tilde{\Lambda}}(x, D) \circ (iP)(t, x, D_t, D_x) \circ \{e^{\tilde{\Lambda}}(x, D)\}^{-1} \circ e^{-k(t) \langle x \rangle_h^{1-\sigma}} \circ e^{-\rho_1 \langle D \rangle^{\frac{1}{\theta}}},$$

where $\rho_1 \in \mathbb{R}$ and $P(t, x, D_t, D_x)$ is given by (1.10). As we discussed before, the role of this conjugation is to make positive the lower order terms of the conjugated operator.

Since the operator ${}^R e^{-\tilde{\Lambda}}$ appears in the inverse $\{e^{\tilde{\Lambda}}(x, D)\}^{-1}$, we need the following technical lemma.

Lemma 5. *Let $\tilde{\Lambda} \in \mathbf{SG}_\mu^{0,1-\sigma}(\mathbb{R}^2)$ and $a \in \mathbf{SG}_{1,s_0}^{m_1,m_2}(\mathbb{R}^2)$, with $\mu > 1$ such that $1/(1-\sigma) > \mu + s_0 - 1$ and $s_0 > \mu$. Then, for $M \in \mathbb{N}$,*

$$e^{\tilde{\Lambda}}(x, D) \circ a(x, D) \circ {}^R e^{-\tilde{\Lambda}} = a(x, D) + \text{op} \left(\sum_{1 \leq \alpha + \beta < M} \frac{1}{\alpha! \beta!} \partial_\xi^\alpha \{ \partial_\xi^\beta e^{\tilde{\Lambda}} D_x^\beta a D_x^\alpha e^{-\tilde{\Lambda}} \} + q_M \right) + r(x, D),$$

where $q_M \in \mathbf{SG}_{\mu,s_0}^{m_1-M, m_2-M\sigma}(\mathbb{R}^2)$ and $r \in \mathcal{S}_{\mu+s_0-1}(\mathbb{R}^2)$.

Proof. Since $e^{\pm \tilde{\Lambda}} \in \mathbf{SG}_{\mu, \frac{1}{1-\sigma}}^{0,\infty}(\mathbb{R}^2)$ and $a \in \mathbf{SG}_{1,s_0}^{m_1,m_2}(\mathbb{R}^2)$, we have $e^{\pm \tilde{\Lambda}}, a \in \mathbf{SG}_{\mu,s_0; \frac{1}{1-\sigma}}^{0,\infty}(\mathbb{R}^2)$. Therefore, by results from calculus, we obtain

$${}^R e^{-\tilde{\Lambda}} = a_1(x, D) + r_1(x, D) \quad \text{and} \quad e^{\tilde{\Lambda}}(x, D) \circ a(x, D) = a_2(x, D) + r_2(x, D),$$

where $a_1 \in \mathbf{SG}_{\mu,s_0; \frac{1}{1-\sigma}}^{0,\infty}(\mathbb{R}^2)$, $a_2 \in \mathbf{SG}_{\mu,s_0; \frac{1}{1-\sigma}}^{m_1,\infty}(\mathbb{R}^2)$, $r_1, r_2 \in \mathcal{S}_{\mu+s_0-1}(\mathbb{R}^2)$ and

$$a_1 \sim \sum_{\alpha} \frac{1}{\alpha!} \partial_\xi^\alpha D_x^\alpha e^{-\tilde{\Lambda}} \text{ in } FSG_{\mu,s_0; \frac{1}{1-\sigma}}^{0,\infty}(\mathbb{R}^2),$$

$$a_2 \sim \sum_{\beta} \frac{1}{\beta!} \partial_{\xi}^{\beta} e^{\tilde{\Lambda}} D_x^{\beta} a \text{ in } FSG_{\mu, s_0; \frac{1}{1-\sigma}}^{m_1, \infty}(\mathbb{R}^2).$$

Hence

$$\begin{aligned} e^{\tilde{\Lambda}} \circ a(x, D) \circ^R e^{-\tilde{\Lambda}} &= a_2(x, D) \circ a_1(x, D) + a_2(x, D) \circ r_1(x, D) \\ &+ r_2(x, D) \circ a_1(x, D) + r_2(x, D) \circ r_1(x, D) \\ &= a_3(x, D) + r_3(x, D) + a_2(x, D) \circ r_1(x, D) \\ &+ r_2(x, D) \circ a_1(x, D) + r_2(x, D) \circ r_1(x, D), \end{aligned}$$

with $a_2(x, D) \circ a_1(x, D) = a_3(x, D) + r_3(x, D)$, where $a_3 \in \mathbf{SG}_{\mu, s_0; \frac{1}{1-\sigma}}^{m_1, \infty}(\mathbb{R}^2)$, $r_3 \in \mathcal{S}_{\mu+s_0-1}(\mathbb{R}^2)$ and

$$\begin{aligned} a_3 &\sim \sum_{\gamma, \alpha, \beta} \frac{1}{\alpha! \beta! \gamma!} \partial_{\xi}^{\gamma} \{ \partial_{\xi}^{\beta} e^{\tilde{\Lambda}} D_x^{\beta} a \} \partial_{\xi}^{\alpha} D_x^{\alpha+\gamma} e^{-\tilde{\Lambda}} = \sum_{\alpha, \beta} \frac{1}{\alpha! \beta!} \partial_{\xi}^{\alpha} \{ \partial_{\xi}^{\beta} e^{\tilde{\Lambda}} D_x^{\beta} a D_x^{\alpha} e^{-\tilde{\Lambda}} \} \\ &= \sum_{j \geq 0} \sum_{\alpha+\beta=j} \frac{1}{\alpha! \beta!} \partial_{\xi}^{\alpha} \{ \partial_{\xi}^{\beta} e^{\tilde{\Lambda}} D_x^{\beta} a D_x^{\alpha} e^{-\tilde{\Lambda}} \} =: \sum_{j \geq 0} c_j \text{ in } FSG_{\mu, s_0; \frac{1}{1-\sigma}}^{m_1, \infty}. \end{aligned}$$

Thus

$$e^{\tilde{\Lambda}}(x, D) \circ a(x, D) \circ^R e^{-\tilde{\Lambda}} = a_3(x, D) + r(x, D),$$

for some $r \in \mathcal{S}_{\mu+s_0-1}(\mathbb{R}^2)$.

Now let us study the asymptotic expansion of a_3 . For $\alpha, \beta \in \mathbb{N}_0$ we have (omitting the dependence (x, ξ)):

$$\begin{aligned} \partial_{\xi}^{\beta} e^{\tilde{\Lambda}} \partial_x^{\beta} a \partial_x^{\alpha} e^{-\tilde{\Lambda}} &= \partial_x^{\beta} a \sum_{h=1}^{\beta} \frac{1}{h!} \sum_{\beta_1+\dots+\beta_h=\beta} \frac{\beta!}{\beta_1! \dots \beta_h!} \prod_{\ell=1}^h \partial_{\xi}^{\beta_{\ell}} \tilde{\Lambda} \\ &\times \sum_{k=1}^{\alpha} \frac{1}{k!} \sum_{\alpha_1+\dots+\alpha_k=\alpha} \frac{\alpha!}{\alpha_1! \dots \alpha_k!} \prod_{\ell=1}^k \partial_x^{\alpha_{\ell}} (-\tilde{\Lambda}). \end{aligned}$$

Therefore, by Faà di Bruno formula, for $\gamma, \delta \in \mathbb{N}_0$, we have

$$\begin{aligned}
\partial_\xi^{\gamma+\alpha} \partial_x^\delta \{ \partial_\xi^\beta e^{\tilde{\Lambda}} \partial_x^\beta a \partial_x^\alpha e^{-\tilde{\Lambda}} \} &= \sum_{\gamma_1+\gamma_2+\gamma_3=\gamma+\alpha} \sum_{\delta_1+\delta_2+\delta_3=\delta} \frac{(\gamma+\alpha)!}{\gamma_1!\gamma_2!\gamma_3!} \frac{\delta!}{\delta_1!\delta_2!\delta_3!} \partial_\xi^{\gamma_1} \partial_x^{\beta+\delta_1} a \\
&\times \partial_\xi^{\gamma_2} \partial_x^{\delta_2} \left(\sum_{h=1}^{\beta} \frac{1}{h!} \sum_{\beta_1+\dots+\beta_h=\beta} \frac{\beta!}{\beta_1!\dots\beta_h!} \prod_{\ell=1}^h \partial_\xi^{\beta_\ell} \tilde{\Lambda} \right) \\
&\times \partial_\xi^{\gamma_3} \partial_x^{\delta_3} \left(\sum_{k=1}^{\alpha} \frac{1}{k!} \sum_{\alpha_1+\dots+\alpha_k=\alpha} \frac{\alpha!}{\alpha_1!\dots\alpha_k!} \prod_{\ell=1}^k \partial_x^{\alpha_\ell} (-\tilde{\Lambda}) \right) \\
&= \sum_{\gamma_1+\gamma_2+\gamma_3=\gamma+\alpha} \sum_{\delta_1+\delta_2+\delta_3=\delta} \frac{(\gamma+\alpha)!}{\gamma_1!\gamma_2!\gamma_3!} \frac{\delta!}{\delta_1!\delta_2!\delta_3!} \partial_\xi^{\gamma_1} \partial_x^{\beta+\delta_1} a \\
&\times \sum_{h=1}^{\beta} \frac{1}{h!} \sum_{\beta_1+\dots+\beta_h=\beta} \frac{\beta!}{\beta_1!\dots\beta_h!} \sum_{\theta_1+\dots+\theta_h=\gamma_2} \sum_{\sigma_1+\dots+\sigma_h=\delta_2} \frac{\gamma_2!}{\theta_1!\dots\theta_h!} \frac{\delta_2!}{\sigma_1!\dots\sigma_h!} \\
&\times \prod_{\ell=1}^h \partial_\xi^{\theta_\ell+\beta_\ell} \partial_x^{\sigma_\ell} \tilde{\Lambda} \\
&\times \sum_{k=1}^{\alpha} \frac{1}{k!} \sum_{\alpha_1+\dots+\alpha_k=\alpha} \frac{\alpha!}{\alpha_1!\dots\alpha_k!} \sum_{\theta_1+\dots+\theta_k=\gamma_3} \sum_{\sigma_1+\dots+\sigma_k=\delta_3} \frac{\gamma_3!}{\theta_1!\dots\theta_k!} \frac{\delta_3!}{\sigma_1!\dots\sigma_k!} \\
&\times \prod_{\ell=1}^k \partial_\xi^{\theta_\ell} \partial_x^{\alpha_\ell+\sigma_\ell} (-\tilde{\Lambda}),
\end{aligned}$$

hence

$$\begin{aligned}
|\partial_\xi^{\gamma+\alpha} \partial_x^\delta (\partial_\xi^\beta e^{\tilde{\Lambda}} D_x^\beta a D_x^\alpha e^{-\tilde{\Lambda}})| &\leq \sum_{\substack{\gamma_1+\gamma_2+\gamma_3=\gamma+\alpha \\ \delta_1+\delta_2+\delta_3=\delta}} \frac{(\gamma+\alpha)!}{\gamma_1!\gamma_2!\gamma_3!} \frac{\delta!}{\delta_1!\delta_2!\delta_3!} C_a^{\gamma_1+\beta+\delta_1+1} \gamma_1!^\mu (\beta+\gamma_1)!^{s_0} \\
&\times \langle \xi \rangle^{m_1-\gamma_1} \langle x \rangle^{m_2-\beta-\delta_1} \\
&\times \sum_{h=1}^\beta \frac{1}{h!} \sum_{\beta_1+\dots+\beta_h=\beta} \frac{\beta!}{\beta_1!\dots\beta_h!} \sum_{\theta_1+\dots+\theta_h=\gamma_2} \sum_{\sigma_1+\dots+\sigma_h=\delta_2} \frac{\gamma_2!}{\theta_1!\dots\theta_h!} \frac{\delta_2!}{\sigma_1!\dots\sigma_h!} \\
&\times \prod_{\ell=1}^h C_{\tilde{\Lambda}}^{\theta_\ell+\beta_\ell+\sigma_\ell+1} (\theta_\ell+\beta_\ell)!^\mu \sigma_\ell!^\mu \langle \xi \rangle^{-\theta_\ell-\beta_\ell} \langle x \rangle^{1-\sigma-\sigma_\ell} \\
&\times \sum_{k=1}^\alpha \frac{1}{k!} \sum_{\alpha_1+\dots+\alpha_k=\alpha} \frac{\alpha!}{\alpha_1!\dots\alpha_k!} \sum_{\theta_1+\dots+\theta_k=\gamma_3} \sum_{\sigma_1+\dots+\sigma_k=\delta_3} \frac{\gamma_3!}{\theta_1!\dots\theta_k!} \frac{\delta_3!}{\sigma_1!\dots\sigma_k!} \\
&\times \prod_{\ell=1}^k C_{\tilde{\Lambda}}^{\theta_\ell+\beta_\ell+\sigma_\ell+1} (\theta_\ell)!^\mu (\alpha_\ell+\sigma_\ell)!^\mu \langle \xi \rangle^{-\theta_\ell} \langle x \rangle^{1-\sigma-\alpha_\ell-\sigma_\ell} \\
&\leq C_1^{\gamma+\delta+2(\alpha+\beta)+1} \gamma!^\mu \delta!^{s_0} (\alpha+\beta)!^{\mu+s_0} \langle \xi \rangle^{m_1-\gamma-(\alpha+\beta)} \langle x \rangle^{m_2-\delta-(\alpha+\beta)} \\
&\times \sum_{k=1}^\alpha \frac{\langle x \rangle^{k(1-\sigma)}}{k!} \sum_{h=1}^\beta \frac{\langle x \rangle^{h(1-\sigma)}}{h!} \\
&\leq C_1^{\gamma+\delta+2(\alpha+\beta)+1} \gamma!^\mu \delta!^{s_0} (\alpha+\beta)!^{\mu+s_0} \langle \xi \rangle^{m_1-\gamma-(\alpha+\beta)} \langle x \rangle^{m_2-\delta-(\alpha+\beta)} \\
&\times 2^{\alpha+\beta} \sum_{k=1}^{\alpha+\beta} \frac{\langle x \rangle^{k(1-\sigma)}}{k!}.
\end{aligned}$$

The above estimate implies

$$|\partial_\xi^\alpha \partial_x^\beta c_j(x, \xi)| \leq C^{\alpha+\beta+2j+1} \alpha!^\mu \beta!^{s_0} (j)!^{\mu+s_0-1} \langle \xi \rangle^{m_1-\alpha-j} \langle x \rangle^{m_2-\beta-j} \sum_{k=1}^j \frac{\langle x \rangle^{k(1-\sigma)}}{k!},$$

for every $j \geq 0$, $\alpha, \beta \in \mathbb{N}_0$ and $x, \xi \in \mathbb{R}$.

Let $\psi(x, \xi) \in C^\infty(\mathbb{R}^2)$ such that $\psi \equiv 0$ on Q_2 , $\psi \equiv 1$ on Q_3^e , $0 \leq \psi \leq 1$ and

$$|\partial_\xi^\alpha \partial_x^\beta \psi(x, \xi)| \leq C^{\alpha+\beta+1} \alpha!^\mu \beta!^{s_0},$$

for every $x, \xi \in \mathbb{R}$ and $\alpha, \beta \in \mathbb{N}_0$. Now set $\psi_0 \equiv 1$ and, for $j \geq 1$,

$$\psi_j(x, \xi) := \psi\left(\frac{x}{R(j)}, \frac{\xi}{R(j)}\right),$$

where $R(j) = Rj^{s_0+\mu-1}$, for a large constant $R > 0$.

Setting $b(x, \xi) = \sum_{j \geq 0} \psi_j(x, \xi) c_j(x, \xi)$ we have $b \in \mathbf{SG}_{\mu, s_0}^{m_1, \infty}(\mathbb{R}^2)$ and

$$b(x, \xi) \sim \sum_{j \geq 0} c_j(x, \xi) \text{ in } FSG_{\mu, s_0}^{m_1, \infty}(\mathbb{R}^2).$$

By similar arguments as the ones used in Section 4.2 we can prove that

$$\sum_{j \geq k} \psi_j(x, \xi) c_j(x, \xi) \in \mathbf{SG}_{\mu, s_0}^{m_1-k, m_2-\sigma k}(\mathbb{R}^2), \quad k \in \mathbb{N}_0.$$

Hence

$$b(x, \xi) - \sum_{j < k} c_j(x, \xi) \in \mathbf{SG}_{\mu, s_0}^{m_1 - k, m_2 - \sigma k}, \quad k \in \mathbb{N}.$$

Since $1/(1 - \sigma) > \mu + s_0 - 1$ we can conclude that $b - a_3 \in \mathcal{S}_{\mu + s_0 - 1}(\mathbb{R}^2)$ and we obtain

$$e^{\tilde{\Lambda}} \circ a \circ^R e^{-\tilde{\Lambda}} = b(x, D) + \tilde{r}(x, D),$$

where $\tilde{r} \in \mathcal{S}_{\mu + s_0 - 1}(\mathbb{R}^2)$. This concludes the proof. \square

5.1. Conjugation by $e^{\tilde{\Lambda}}$. We start noting that $e^{\tilde{\Lambda}} \partial_t \{e^{\tilde{\Lambda}}\}^{-1} = \partial_t$ since $\tilde{\Lambda}$ does not depend on t .

- Conjugation of $ia_3(t, D)$.

Since a_3 does not depend of x , applying Lemma 5, we have

$$e^{\tilde{\Lambda}}(x, D)(ia_3)(t, D)^R(e^{-\tilde{\Lambda}}) = ia_3(t, D) + s(t, x, D) + r_3(t, x, D),$$

with

$$s \sim \sum_{j \geq 1} \frac{1}{j!} \partial_\xi^j \{e^{\tilde{\Lambda}} ia_3 D_x^j e^{-\tilde{\Lambda}}\} \in FSG_{\mu, s_0}^{2, -\sigma}(\mathbb{R}^2), \quad r_3 \in C([0, T], \mathcal{S}_{\mu + s_0 - 1}(\mathbb{R}^2)).$$

Hence, using (4.3) we can write more explicitly $s(t, x, D)$ and we obtain

$$\begin{aligned} e^{\tilde{\Lambda}}(ia_3)\{e^{\tilde{\Lambda}}\}^{-1} &= \text{op} \left(ia_3 - \partial_\xi(a_3 \partial_x \tilde{\Lambda}) + \frac{i}{2} \partial_\xi^2 [a_3(\partial_x^2 \tilde{\Lambda} - (\partial_x \tilde{\Lambda})^2)] + a_3^{(0)} + r_3 \right) \\ &\quad \circ \left[I + \text{op} \left(-i \partial_\xi \partial_x \tilde{\Lambda} - \frac{1}{2} \partial_\xi^2 (\partial_x^2 \tilde{\Lambda} - [\partial_x \tilde{\Lambda}]^2) - [\partial_\xi \partial_x \tilde{\Lambda}]^2 + q_{-3} \right) \right] \\ &= ia_3 - \text{op} \left(\partial_\xi(a_3 \partial_x \tilde{\Lambda}) + \frac{i}{2} \partial_\xi^2 \{a_3(\partial_x^2 \tilde{\Lambda} - \{\partial_x \tilde{\Lambda}\}^2)\} - a_3 \partial_\xi \partial_x \tilde{\Lambda} + i \partial_\xi a_3 \partial_\xi \partial_x^2 \tilde{\Lambda} \right) \\ &\quad + \text{op} \left(i \partial_\xi(a_3 \partial_x \tilde{\Lambda}) \partial_\xi \partial_x \tilde{\Lambda} - \frac{i}{2} a_3 \{\partial_\xi^2 (\partial_x^2 \tilde{\Lambda} + [\partial_x \tilde{\Lambda}]^2) + 2[\partial_\xi \partial_x \tilde{\Lambda}]^2 + r_0 + \bar{r} \} \right) \\ &= ia_3 - \text{op} \left(\partial_\xi a_3 \partial_x \tilde{\Lambda} + \frac{i}{2} \partial_\xi^2 \{a_3[\partial_x^2 \tilde{\Lambda} - (\partial_x \tilde{\Lambda})^2]\} + i \partial_\xi a_3 \partial_\xi \partial_x^2 \tilde{\Lambda} \right) \\ &\quad + \text{op} \left(i \partial_\xi(a_3 \partial_x \tilde{\Lambda}) \partial_\xi \partial_x \tilde{\Lambda} - \frac{i}{2} a_3 \{\partial_\xi^2 (\partial_x^2 \tilde{\Lambda} + [\partial_x \tilde{\Lambda}]^2) + 2(\partial_\xi \partial_x \tilde{\Lambda})^2\} + r_0 + \bar{r} \right) \\ &= ia_3 - \text{op} \left(\partial_\xi a_3 \partial_x \lambda_2 + \partial_\xi a_3 \partial_x \lambda_1 + \frac{i}{2} \partial_\xi^2 \{a_3(\partial_x^2 \lambda_2 - \{\partial_x \lambda_2\}^2)\} + i \partial_\xi a_3 \partial_\xi \partial_x^2 \lambda_2 \right) \\ &\quad + \text{op} \left(i \partial_\xi(a_3 \partial_x \lambda_2) \partial_\xi \partial_x \lambda_2 - \frac{i}{2} a_3 \{\partial_\xi^2 (\partial_x^2 \lambda_2 + [\partial_x \lambda_2]^2) + 2[\partial_\xi \partial_x \lambda_2]^2\} + r_0 + \bar{r} \right), \end{aligned}$$

where $a_3^{(0)} \in C([0, T]; \mathbf{SG}_{\mu, s_0}^{0,0}(\mathbb{R}^2))$, $r_0 \in C([0, T]; \mathbf{SG}_\delta^{0,0}(\mathbb{R}^2))$ and, since we may assume $\delta < \mu + s_0 - 1$, $\bar{r} \in C([0, T]; \mathcal{S}_{\mu + s_0 - 1}(\mathbb{R}^2))$.

- Conjugation of $ia_2(t, x, D)$.

By Lemma 5 with $M = 2$ and using (4.3) we get

$$\begin{aligned}
e^{\tilde{\Lambda}}(x, D)(ia_2)(t, x, D)\{e^{\tilde{\Lambda}}(x, D)\}^{-1} &= \text{op} \left(ia_2 - \partial_{\xi} \{a_2 \partial_x \tilde{\Lambda}\} + \partial_{\xi} \tilde{\Lambda} \partial_x a_2 + a_2^{(0)} + r_2 \right) \\
&\quad \circ \left[I - \text{op} \left(i \partial_{\xi} \partial_x \tilde{\Lambda} + \frac{1}{2} \partial_{\xi}^2 (\partial_x^2 \tilde{\Lambda} - [\partial_x \tilde{\Lambda}]^2) + [\partial_{\xi} \partial_x \tilde{\Lambda}]^2 + q_{-3} \right) \right] \\
&= ia_2(t, x, D) + \text{op}(-\partial_{\xi} \{a_2 \partial_x \tilde{\Lambda}\} + \partial_{\xi} \tilde{\Lambda} \partial_x a_2 + a_2 \partial_{\xi} \partial_x \tilde{\Lambda} + r_0 + \bar{r}) \\
&= ia_2(t, x, D) + \text{op}(-\partial_{\xi} a_2 \partial_x \tilde{\Lambda} + \partial_{\xi} \tilde{\Lambda} \partial_x a_2 + r_0 + \bar{r}) \\
&= ia_2(t, x, D) + \text{op}(-\partial_{\xi} a_2 \partial_x \lambda_2 + \partial_{\xi} \lambda_2 \partial_x a_2 + r_0 + \bar{r}),
\end{aligned}$$

where $a_2^{(0)} \in C([0, T]; \mathbf{SG}_{\mu, s_0}^{0,0})$, $r_0 \in C([0, T]; \mathbf{SG}_{\delta}^{0,0}(\mathbb{R}^2))$ and $\bar{r} \in C([0, T]; \Sigma_{\mu+s_0-1}(\mathbb{R}^2))$.

- Conjugation of $ia_1(t, x, D)$:

$$e^{\tilde{\Lambda}}(x, D)ia_1(t, x, D)\{e^{\tilde{\Lambda}}(x, D)\}^{-1} = \text{op}(ia_1 + a_1^{(0)} + r_1) \circ \sum_{j \geq 0} (-r(t, x, D))^j = \text{op}(ia_1 + \tilde{r}_0 + \tilde{r}),$$

where $a_1^{(0)} \in C([0, T]; \mathbf{SG}_{\mu, s_0}^{0,1-2\sigma}(\mathbb{R}^2))$, $\tilde{r}_0 \in C([0, T]; \mathbf{SG}_{\delta}^{0,1-2\sigma}(\mathbb{R}^2))$, $\tilde{r} \in C([0, T]; \Sigma_{\mu+s_0-1}(\mathbb{R}^2))$.

- Conjugation of $ia_0(t, x, D)$:

$$e^{\tilde{\Lambda}}(x, D)ia_0(t, x, D)\{e^{\tilde{\Lambda}}(x, D)\}^{-1} = \text{op}(ia_0 + a_0^{(0)} + r_0) \sum_{j \geq 0} (-r(t, x, D))^j = \text{op}(ia_0 + \tilde{\tilde{r}}_0 + \tilde{r}_1),$$

where $a_0^{(0)} \in C([0, T]; \mathbf{SG}_{\mu, s_0}^{0,1-2\sigma})$, $\tilde{\tilde{r}}_0 \in C([0, T]; \mathbf{SG}_{\delta}^{-1,1-2\sigma}(\mathbb{R}^2))$ and $\tilde{r}_1 \in C([0, T]; \Sigma_{\mu+s_0-1}(\mathbb{R}^2))$.

Summing up we obtain

$$\begin{aligned}
(5.1) \quad e^{\tilde{\Lambda}}(x, D)(iP(t, x, D_t, D_x))\{e^{\tilde{\Lambda}}(x, D)\}^{-1} &= \partial_t + ia_3(t, D) + ia_2(t, x, D) \\
&\quad - \text{op}(\partial_{\xi} a_3 \partial_x \lambda_2) + ia_1(t, x, D) - \text{op}(\partial_{\xi} a_3 \partial_x \lambda_1 + \partial_{\xi} a_2 \partial_x \lambda_2 - \partial_{\xi} \lambda_2 \partial_x a_2 + ib_1) \\
&\quad + ia_0(t, x, D) + r_{\sigma}(t, x, D) + r_0(t, x, D) + \bar{r}(t, x, D),
\end{aligned}$$

where

$$(5.2) \quad b_1 \in C([0, T]; \mathbf{SG}_{\mu, s_0}^{1,-2\sigma}(\mathbb{R}^2)), \quad b_1(t, x, \xi) \in \mathbb{R}, \quad b_1 \text{ does not depend on } \lambda_1,$$

and

$$r_0 \in C([0, T]; \mathbf{SG}_{\delta}^{0,0}(\mathbb{R}^2)), \quad r_{\sigma} \in C([0, T]; \mathbf{SG}_{\delta}^{0,1-2\sigma}(\mathbb{R}^2)), \quad \bar{r} \in C([0, T]; \Sigma_{\mu+s_0-1}(\mathbb{R}^2)).$$

5.2. Conjugation by $e^{k(t)\langle x \rangle_h^{1-\sigma}}$. Let us recall that we are assuming that $k \in C^1([0, T]; \mathbb{R})$, $k'(t) \leq 0$ and $k(t) \geq 0$ for every $t \in [0, T]$. Following the same ideas of the proof of Lemma 5 one can prove the following result.

Lemma 6. *Let $a \in C([0, T], \mathbf{SG}_{\mu, s_0}^{m_1, m_2}(\mathbb{R}^2))$, where $1 < \mu < s_0$ and $\mu + s_0 - 1 < \frac{1}{1-\sigma}$. Then*

$$e^{k(t)\langle x \rangle_h^{1-\sigma}} a(t, x, D) e^{-k(t)\langle x \rangle_h^{1-\sigma}} = a(t, x, D) + b(t, x, D) + \bar{r}(t, x, D),$$

where $b \sim \sum_{j \geq 1} \frac{1}{j!} e^{k(t)\langle x \rangle_h^{1-\sigma}} \partial_{\xi}^j a D_x^j e^{-k(t)\langle x \rangle_h^{1-\sigma}}$ in $\mathbf{SG}_{\mu, s_0}^{m_1-1, m_2-\sigma}(\mathbb{R}^2)$ and $\bar{r} \in C([0, T], \mathcal{S}_{\mu+s_0-1}(\mathbb{R}^2))$.

Let us perform the conjugation by $e^{k(t)\langle x \rangle_h^{1-\sigma}}$ of the operator $e^{\tilde{\Lambda}}(iP)\{e^{\tilde{\Lambda}}\}^{-1}$ in (5.1) with the aid of Lemma 6.

- Conjugation of ∂_t : $e^{k(t)\langle x \rangle_h^{1-\sigma}} \partial_t e^{-k(t)\langle x \rangle_h^{1-\sigma}} = \partial_t - k'(t)\langle x \rangle_h^{1-\sigma}$.

- Conjugation of $ia_3(t, D)$:

$$e^{k(t)\langle x \rangle_h^{1-\sigma}} ia_3(t, D) e^{-k(t)\langle x \rangle_h^{1-\sigma}} = ia_3 + \text{op}(-k(t)\partial_\xi a_3 \partial_x \langle x \rangle_h^{1-\sigma}) \\ + \text{op}\left(\frac{i}{2}\partial_\xi^2 a_3 \{k(t)\partial_x^2 \langle x \rangle_h^{1-\sigma} - k^2(t)[\partial_x \langle x \rangle_h^{1-\sigma}]^2\}\right) + a_3^{(0)} + r_3,$$

where $a_3^{(0)} \in C([0, T]; \mathbf{SG}_{\mu, s_0}^{0, -3\sigma}(\mathbb{R}^2))$ and $r_3 \in C([0, T]; \Sigma_{\mu+s_0-1}(\mathbb{R}^2))$.

- Conjugation of $\text{op}(ia_2 - \partial_\xi a_3 \partial_x \lambda_2)$:

$$e^{k(t)\langle x \rangle_h^{1-\sigma}} \text{op}(ia_2 - \partial_\xi a_3 \partial_x \lambda_2) e^{-k(t)\langle x \rangle_h^{1-\sigma}} \\ = ia_2 + \text{op}(-\partial_\xi a_3 \partial_x \lambda_2) + \text{op}(-k(t)\partial_\xi a_2 \partial_x \langle x \rangle_h^{1-\sigma} - ik(t)\partial_\xi \{\partial_\xi a_3 \partial_x \lambda_2\} \partial_x \langle x \rangle_h^{1-\sigma} + a_2^{(0)} + r_2),$$

where $a_2^{(0)} \in C([0, T]; \mathbf{SG}_{\mu, s_0}^{0, -2\sigma}(\mathbb{R}^2))$ and $r_2 \in C([0, T]; \Sigma_{\mu+s_0-1}(\mathbb{R}^2))$.

- Conjugation of $i(a_1 + a_0)(t, x, D)$:

$$e^{k(t)\langle x \rangle_h^{1-\sigma}} i(a_1 + a_0)(t, x, D) e^{-k(t)\langle x \rangle_h^{1-\sigma}} = \text{op}(ia_1 + ia_0 + a_{1,0} + r_1),$$

where $r_1 \in C([0, T]; \Sigma_{\mu+s_0-1}(\mathbb{R}^2))$ and

$$a_{1,0} \sim \sum_{j \geq 1} e^{k(t)\langle x \rangle_h^{1-\sigma}} \frac{1}{j!} \partial_\xi^j i(a_1 + a_0) D_x^j e^{-k(t)\langle x \rangle_h^{1-\sigma}} \text{ in } \mathbf{SG}_{\mu, s_0}^{0, 1-2\sigma}(\mathbb{R}^2).$$

It is not difficult to verify that the following estimate holds

$$(5.3) \quad |a_{1,0}(t, x, \xi)| \leq \max\{1, k(t)\} C_T \langle x \rangle_h^{1-2\sigma},$$

where C_T depends on a_1 and does not depend on $k(t)$.

- Conjugation of $\text{op}(-\partial_\xi a_3 \partial_x \lambda_1 - \partial_\xi a_2 \partial_x \lambda_2 + \partial_\xi \lambda_2 \partial_x a_2 + ib_1)$: taking into account i) of Lemma 3

$$e^{k(t)\langle x \rangle_h^{1-\sigma}} \text{op}(-\partial_\xi a_3 \partial_x \lambda_1 - \partial_\xi a_2 \partial_x \lambda_2 + \partial_\xi \lambda_2 \partial_x a_2 + ib_1) e^{-k(t)\langle x \rangle_h^{1-\sigma}} \\ = \text{op}(-\partial_\xi a_3 \partial_x \lambda_1 - \partial_\xi a_2 \partial_x \lambda_2 + \partial_\xi \lambda_2 \partial_x a_2 + ib_1 + r_0 + \bar{r}),$$

where $r_0 \in C([0, T]; \mathbf{SG}_{\mu, s_0}^{0, 0}(\mathbb{R}^2))$ and $\bar{r} \in C([0, T]; \Sigma_{\mu+s_0-1}(\mathbb{R}^2))$.

- Conjugation of $r_\sigma(t, x, D)$:

$$e^{k(t)\langle x \rangle_h^{1-\sigma}} r_\sigma(t, x, D) e^{-k(t)\langle x \rangle_h^{1-\sigma}} = r_{\sigma,1}(t, x, D) + \bar{r}(t, x, D),$$

where $r \in C([0, T]; \Sigma_{\mu+s_0-1}(\mathbb{R}^2))$, $r_{\sigma,1} \in C([0, T]; \mathbf{SG}_\delta^{0, 1-2\sigma}(\mathbb{R}^2))$ and the estimate

$$(5.4) \quad |r_{\sigma,1}(t, x, \xi)| \leq C_{T, \tilde{\Lambda}} \langle x \rangle_h^{1-2\sigma}$$

holds with $C_{T, \tilde{\Lambda}}$ independent of $k(t)$.

Gathering all the previous computations we may write

$$(5.5) \quad e^{k(t)\langle x \rangle_h^{1-\sigma}} e^\Lambda(iP) \{e^\Lambda\}^{-1} e^{-k(t)\langle x \rangle_h^{1-\sigma}} = \partial_t + \text{op}(ia_3 - \partial_\xi a_3 \partial_x \lambda_2 + ia_2 - k(t)\partial_\xi a_3 \partial_x \langle x \rangle_h^{1-\sigma}) \\ + \text{op}(-\partial_\xi a_3 \partial_x \lambda_1 + ia_1 - \partial_\xi a_2 \partial_x \lambda_2 + \partial_\xi \lambda_2 \partial_x a_2 - k(t)\partial_\xi a_2 \partial_x \langle x \rangle_h^{1-\sigma}) \\ + \text{op}(ib_1 + ic_1 + ia_0 - k'(t)\langle x \rangle_h^{1-\sigma} + a_{1,0} + r_{\sigma,1}) + r_0(t, x, D) + \bar{r}(t, x, D),$$

where b_1 satisfies (5.2),

$$(5.6) \quad c_1 \in C([0, T]; \mathbf{SG}_{\mu, s_0}^{1, -2\sigma}(\mathbb{R}^2)), \quad c_1(t, x, \xi) \in \mathbb{R}, \quad c_1 \text{ does not depend on } \lambda_1,$$

(but c_1 depends of $\lambda_2, k(t)$), $a_{1,0}$ as in (5.3), $r_{\sigma,1}$ as in (5.4), and for some new operators

$$r_0 \in C([0, T]; \mathbf{SG}_\delta^{0, 0}(\mathbb{R}^2)), \quad \bar{r} \in C([0, T]; \mathcal{S}_{\mu+s_0-1}(\mathbb{R}^2)).$$

5.3. Conjugation by $e^{\rho_1 \langle D \rangle^{\frac{1}{\theta}}}$. Since we are considering $\theta > s_0$ and $\mu > 1$ arbitrarily close to 1, we may assume that all the previous smoothing remainder terms have symbols in $\Sigma_\theta(\mathbb{R}^2)$.

Lemma 7. *Let $a \in \mathbf{SG}_{\mu, s_0}^{m_1, m_2}$, where $1 < \mu < s_0$ and $\mu + s_0 - 1 < \theta$. Then*

$$e^{\rho_1 \langle D \rangle^{\frac{1}{\theta}}} a(x, D) e^{-\rho_1 \langle D \rangle^{\frac{1}{\theta}}} = a(x, D) + b(x, D) + r(x, D),$$

where $b \sim \sum_{j \geq 1} \frac{1}{j!} \partial_\xi^j e^{\rho_1 \langle \xi \rangle^{\frac{1}{\theta}}} D_x^j a e^{-\rho_1 \langle \xi \rangle^{\frac{1}{\theta}}}$ in $\mathbf{SG}_{\mu, s_0}^{m_1 - (1 - \frac{1}{\theta}), m_2 - 1}(\mathbb{R}^2)$ and $r \in \mathcal{S}_{\mu + s_0 - 1}(\mathbb{R}^2)$.

Let us now conjugate by $e^{\rho_1 \langle D \rangle^{\frac{1}{\theta}}}$ the operator $e^{k(t) \langle x \rangle_h^{1-\sigma}} e^\Lambda(iP) \{e^\Lambda\}^{-1} e^{-k(t) \langle x \rangle_h^{1-\sigma}}$ in (5.5). First of all we observe that since a_3 does not depend of x , we simply have

$$e^{\rho_1 \langle D \rangle^{\frac{1}{\theta}}} i a_3(t, D) e^{-\rho_1 \langle D \rangle^{\frac{1}{\theta}}} = i a_3(t, D).$$

- Conjugation of $\text{op}(-\partial_\xi a_3 \partial_x \lambda_2 + i a_2 - k(t) \partial_\xi a_3 \partial_x \langle x \rangle_h^{1-\sigma})$:

$$\begin{aligned} & e^{\rho_1 \langle D \rangle^{\frac{1}{\theta}}} \text{op}(-\partial_\xi a_3 \partial_x \lambda_2 + i a_2 - k(t) \partial_\xi a_3 \partial_x \langle x \rangle_h^{1-\sigma}) e^{-\rho_1 \langle D \rangle^{\frac{1}{\theta}}} \\ &= \text{op}(-\partial_\xi a_3 \partial_x \lambda_2 + i a_2 - k(t) \partial_\xi a_3 \partial_x \langle x \rangle_h^{1-\sigma}) + (a_{2, \rho_1} + \bar{r})(t, x, D), \end{aligned}$$

where $a_{2, \rho_1} \in C([0, T], \mathbf{SG}_{\mu, s_0}^{1 + \frac{1}{\theta}, -1}(\mathbb{R}^2))$, $\bar{r} \in C([0, T], \Sigma_\theta(\mathbb{R}^2))$ and the following estimate holds

$$(5.7) \quad |\partial_\xi^\alpha \partial_x^\beta a_{2, \rho_1}(t, x, \xi)| \leq \max\{1, k(t)\} |\rho_1| C_{\lambda_2, T}^{\alpha + \beta + 1} \alpha!^\mu \beta!^{s_0} \langle \xi \rangle^{1 + \frac{1}{\theta} - \alpha} \langle x \rangle^{-\sigma - \beta}.$$

In particular

$$|a_{2, \rho_1}(t, x, \xi)| \leq \max\{1, k(t)\} |\rho_1| C_{\lambda_2, T} \langle \xi \rangle_h^{1 + \frac{1}{\theta}} \langle x \rangle^{-\sigma}.$$

- Conjugation of $\text{op}(-\partial_\xi a_3 \partial_x \lambda_1 + i a_1 - \partial_\xi a_2 \partial_x \lambda_2 + \partial_\xi \lambda_2 \partial_x a_2 - k(t) \partial_\xi a_2 \partial_x \langle x \rangle_h^{1-\sigma} + i b_1 + i c_1)$: the conjugation of this term is given by

$$\begin{aligned} & \text{op}(-\partial_\xi a_3 \partial_x \lambda_1 + i a_1 - \partial_\xi a_2 \partial_x \lambda_2 + \partial_\xi \lambda_2 \partial_x a_2 - k(t) \partial_\xi a_2 \partial_x \langle x \rangle_h^{1-\sigma} + i b_1 + i c_1) \\ &+ a_{1, \rho_1}(t, x, D) + \bar{r}(t, x, D), \end{aligned}$$

where $\bar{r} \in C([0, T], \Sigma_\theta(\mathbb{R}^2))$ and a_{1, ρ_1} satisfies the following estimate

$$(5.8) \quad |\partial_\xi^\alpha \partial_x^\beta a_{1, \rho_1}(t, x, \xi)| \leq \max\{1, k(t)\} |\rho_1| C_{\tilde{\lambda}, T}^{\alpha + \beta + 1} \alpha!^\mu \beta!^{s_0} \langle \xi \rangle^{\frac{1}{\theta} - \alpha} \langle x \rangle^{-\sigma - \beta}.$$

In particular

$$|a_{1, \rho_1}(t, x, \xi)| \leq \max\{1, k(t)\} |\rho_1| C_{\tilde{\lambda}, T} \langle \xi \rangle_h^{\frac{1}{\theta}} \langle x \rangle^{-\sigma}.$$

- Conjugation of $\text{op}(i a_0 - k'(t) \langle x \rangle_h^{1-\sigma} + a_{1,0} + r_{\sigma,1})$:

$$\begin{aligned} & e^{\rho_1 \langle D \rangle^{\frac{1}{\theta}}} \text{op}(i a_0 - k'(t) \langle x \rangle_h^{1-\sigma} + a_{1,0} + r_{\sigma,1}) e^{-\rho_1 \langle D \rangle^{\frac{1}{\theta}}} \\ &= \text{op}(i a_0 - k'(t) \langle x \rangle_h^{1-\sigma} + a_{1,0} + r_{\sigma,1}) + r_0(t, x, D) + \bar{r}(t, x, D), \end{aligned}$$

where $r_0 \in C([0, T], \mathbf{SG}_\delta^{(0,0)}(\mathbb{R}^2))$ and $\bar{r} \in C([0, T], \Sigma_\theta(\mathbb{R}^2))$.

Summing up, from the previous computations we obtain

$$\begin{aligned} & e^{\rho_1 \langle D \rangle^{\frac{1}{\theta}}} e^{k(t) \langle x \rangle_h^{1-\sigma}} e^{\tilde{\Lambda}}(iP) \{e^{\rho_1 \langle D \rangle^{\frac{1}{\theta}}} e^{k(t) \langle x \rangle_h^{1-\sigma}} e^{\tilde{\Lambda}}\}^{-1} = \partial_t + i a_3(t, D) \\ &+ \text{op}(-\partial_\xi a_3 \partial_x \lambda_2 + i a_2 - k(t) \partial_\xi a_3 \partial_x \langle x \rangle_h^{1-\sigma} + a_{2, \rho_1}) \\ &+ \text{op}(-\partial_\xi a_3 \partial_x \lambda_1 + i a_1 - \partial_\xi a_2 \partial_x \lambda_2 + \partial_\xi \lambda_2 \partial_x a_2 - k(t) \partial_\xi a_2 \partial_x \langle x \rangle_h^{1-\sigma} + i b_1 + i c_1 + a_{1, \rho_1}) \\ &+ \text{op}(i a_0 - k'(t) \langle x \rangle_h^{1-\sigma} + a_{1,0} + r_{\sigma,1}) + r_0(t, x, D) + \bar{r}(t, x, D), \end{aligned}$$

with a_{2,ρ_1} as in (5.7), b_1 as in (5.2), c_1 as in (5.6), a_{1,ρ_1} as in (5.8), $a_{1,0}$ as in (5.3), $r_{\sigma,1}$ as in (5.4), and for some operators

$$r_0 \in C([0, T]; \mathbf{SG}_\delta^{(0,0)}(\mathbb{R}^2)), \quad \bar{r} \in C([0, T]; \Sigma_\theta(\mathbb{R}^2)).$$

6. Estimates from below

In this section we will choose M_2, M_1 and $k(t)$ in order to apply Fefferman-Phong and sharp Gårding inequalities to get the desired energy estimate for the conjugated problem. By the computations of the previous section we have

$$\begin{aligned} & e^{\rho_1 \langle D \rangle^{\frac{1}{\theta}}} e^{k(t) \langle x \rangle_h^{1-\sigma}} e^{\tilde{\Lambda}}(x, D) (iP) \{ e^{\rho_1 \langle D \rangle^{\frac{1}{\theta}}} e^{k(t) \langle x \rangle_h^{1-\sigma}} e^{\tilde{\Lambda}}(x, D) \}^{-1} \\ &= \partial_t + ia_3(t, D) + \sum_{j=0}^2 \tilde{a}_j(t, x, D) + r_0(t, x, D) + \bar{r}(t, x, D), \end{aligned}$$

where \tilde{a}_2, \tilde{a}_1 represent the part with ξ -order 2, 1 respectively and \tilde{a}_0 represents the part with ξ -order 0, but with a positive order (less than or equal to $1 - \sigma$) with respect to x . The real parts are given by

$$\begin{aligned} \operatorname{Re} \tilde{a}_2 &= -\partial_\xi a_3 \partial_x \lambda_2 - \operatorname{Im} a_2 - k(t) \partial_\xi a_3 \partial_x \langle x \rangle_h^{1-\sigma} + \operatorname{Re} a_{2,\rho_1}, \\ \operatorname{Im} \tilde{a}_2 &= \operatorname{Re} a_2 + \operatorname{Im} a_{2,\rho_1}, \\ \operatorname{Re} \tilde{a}_1 &= -\partial_\xi a_3 \partial_x \lambda_1 - \operatorname{Im} a_1 - \partial_\xi \operatorname{Re} a_2 \partial_x \lambda_2 + \partial_\xi \lambda_2 \partial_x \operatorname{Re} a_2 \\ &\quad - k(t) \partial_\xi \operatorname{Re} a_2 \partial_x \langle x \rangle_h^{1-\sigma} + \operatorname{Re} a_{1,\rho_1}, \\ \operatorname{Re} \tilde{a}_0 &= -\operatorname{Im} a_0 - k'(t) \langle x \rangle_h^{1-\sigma} + \operatorname{Re} a_{1,0} + \operatorname{Re} r_{\sigma,1}. \end{aligned}$$

Since the Fefferman-Phong inequality holds true only for scalar symbols, we need to decompose $\operatorname{Im} \tilde{a}_2$ into its Hermitian and anti-Hermitian part:

$$i \operatorname{Im} \tilde{a}_2 = \frac{i \operatorname{Im} \tilde{a}_2 + (i \operatorname{Im} \tilde{a}_2)^*}{2} + \frac{i \operatorname{Im} \tilde{a}_2 - (i \operatorname{Im} \tilde{a}_2)^*}{2} = t_1 + t_2,$$

where $2 \operatorname{Re} \langle t_2(t, x, D) u, u \rangle = 0$ and $t_1(t, x, \xi) = - \sum_{\alpha \geq 1} \frac{i}{2\alpha!} \partial_\xi^\alpha D_x^\alpha \operatorname{Im} \tilde{a}_2(t, x, \xi)$. Observe that, using (5.7),

$$\begin{aligned} |t_1(t, x, \xi)| &\leq C_{a_2} \langle \xi \rangle \langle x \rangle^{-1} + \max\{1, k(t)\} |\rho_1| C_{\lambda_2} \langle \xi \rangle^{\frac{1}{\theta}} \langle x \rangle^{-\sigma-1} \\ (6.1) \quad &\leq \{C_{a_2} + h^{-(1-\frac{1}{\theta})} \max\{1, k(0)\} |\rho_1| C_{\lambda_2}\} \langle \xi \rangle_h \langle x \rangle^{-\frac{\sigma}{2}}. \end{aligned}$$

In this way we may write

$$\begin{aligned} (6.2) \quad & e^{\rho_1 \langle D \rangle^{\frac{1}{\theta}}} e^{k(t) \langle x \rangle_h^{1-\sigma}} e^{\tilde{\Lambda}}(x, D) (iP) \{ e^{\rho_1 \langle D \rangle^{\frac{1}{\theta}}} e^{k(t) \langle x \rangle_h^{1-\sigma}} e^{\tilde{\Lambda}}(x, D) \}^{-1} \\ &= \partial_t + ia_3(t, D) + (\operatorname{Re} \tilde{a}_2 + t_2 + t_1 + \tilde{a}_1 + \tilde{a}_0)(t, x, D) + \tilde{r}_0(t, x, D), \end{aligned}$$

where \tilde{r}_0 has symbol of order $(0, 0)$.

Now we are ready to choose M_2, M_1 and $k(t)$. The function $k(t)$ will be of the form $k(t) = K(T - t)$, $t \in [0, T]$, for a positive constant K to be chosen. In the following computations we

shall consider $|\xi| > hR_{a_3}$. Observe that $2|\xi|^2 \geq \langle \xi \rangle_h^2$ whenever $|\xi| \geq h > 1$. For $Re \tilde{a}_2$ we have:

$$\begin{aligned}
Re \tilde{a}_2 &= M_2 |\partial_\xi a_3| \langle x \rangle^{-\sigma} - Im a_2 - k(t) \partial_\xi a_3 \partial_x \langle x \rangle_h^{1-\sigma} + Re a_{2,\rho_1} \\
&\geq M_2 C_{a_3} |\xi|^2 \langle x \rangle^{-\sigma} - C_{a_2} \langle \xi \rangle_h^2 \langle x \rangle^{-\sigma} \\
&\quad - \tilde{C}_{a_3} k(0) (1 - \sigma) \langle \xi \rangle_h^2 \langle x \rangle_h^{-\sigma} - \max\{1, k(0)\} C_{\lambda_2, \rho_1} \langle \xi \rangle_h^{1+\frac{1}{\theta}} \langle x \rangle^{-\sigma} \\
&\geq (M_2 \frac{C_{a_3}}{2} - C_{a_2} - \tilde{C}_{a_3} k(0) (1 - \sigma) - \max\{1, k(0)\} C_{\lambda_2, \rho_1} \langle \xi \rangle_h^{-(1-\frac{1}{\theta})}) \langle \xi \rangle_h^2 \langle x \rangle^{-\sigma} \\
&\geq (M_2 \frac{C_{a_3}}{2} - C_{a_2} - \tilde{C}_{a_3} k(0) (1 - \sigma) - \max\{1, k(0)\} C_{\lambda_2, \rho_1} h^{-(1-\frac{1}{\theta})}) \langle \xi \rangle_h^2 \langle x \rangle^{-\sigma} \\
&= (M_2 \frac{C_{a_3}}{2} - C_{a_2} - \tilde{C}_{a_3} KT (1 - \sigma) - \max\{1, KT\} C_{\lambda_2, \rho_1} h^{-(1-\frac{1}{\theta})}) \langle \xi \rangle_h^2 \langle x \rangle^{-\sigma}
\end{aligned}$$

For $Re \tilde{a}_1$, we have:

$$\begin{aligned}
Re \tilde{a}_1 &= M_1 |\partial_\xi a_3| \langle \xi \rangle_h^{-1} \langle x \rangle^{-\frac{\sigma}{2}} \psi \left(\frac{\langle x \rangle^\sigma}{\langle \xi \rangle_h^2} \right) - Im a_1 - \partial_\xi Re a_2 \partial_x \lambda_2 + \partial_\xi \lambda_2 \partial_x Re a_2 \\
&\quad - k(t) \partial_\xi Re a_2 \partial_x \langle x \rangle_h^{1-\sigma} + Re a_{1,\rho_1} \\
&\geq M_1 C_{a_3} |\xi|^2 \langle \xi \rangle_h^{-1} \langle x \rangle^{-\frac{\sigma}{2}} \psi \left(\frac{\langle x \rangle^\sigma}{\langle \xi \rangle_h^2} \right) - C_{a_1} \langle \xi \rangle_h \langle x \rangle^{-\frac{\sigma}{2}} - \tilde{C}_{a_2, \lambda_2} \langle \xi \rangle_h \langle x \rangle^{-\sigma} \\
&\quad - C k(0) (1 - \sigma) \langle \xi \rangle_h \langle x \rangle_h^{-\sigma} - \max\{1, k(0)\} C_{\tilde{\Lambda}, \rho_1} \langle \xi \rangle_h^{\frac{1}{\theta}} \langle x \rangle_h^{-\sigma} \\
&\geq M_1 \frac{C_{a_3}}{2} \langle \xi \rangle_h \langle x \rangle^{-\frac{\sigma}{2}} \psi \left(\frac{\langle x \rangle^\sigma}{\langle \xi \rangle_h^2} \right) - C_{a_1} \langle \xi \rangle_h \langle x \rangle^{-\frac{\sigma}{2}} - \tilde{C}_{a_2, \lambda_2} \langle \xi \rangle_h \langle x \rangle^{-\frac{\sigma}{2}} \\
&\quad - C k(0) (1 - \sigma) \langle \xi \rangle_h \langle x \rangle_h^{-\frac{\sigma}{2}} \langle x \rangle^{-\frac{\sigma}{2}} - \max\{1, k(0)\} C_{\tilde{\Lambda}, \rho_1} \langle \xi \rangle_h^{-(1-\frac{1}{\theta})} \langle \xi \rangle_h \langle x \rangle^{-\frac{\sigma}{2}} \\
&= (M_1 \frac{C_{a_3}}{2} - C_{a_1} - \tilde{C}_{a_2, \lambda_2} - C K T (1 - \sigma) \langle x \rangle_h^{-\frac{\sigma}{2}}) \langle \xi \rangle_h \langle x \rangle^{-\frac{\sigma}{2}} \\
&\quad - \max\{1, K T\} C_{\tilde{\Lambda}, \rho_1} \langle \xi \rangle_h^{-(1-\frac{1}{\theta})} \langle \xi \rangle_h \langle x \rangle^{-\frac{\sigma}{2}} - M_1 \frac{C_{a_3}}{2} \langle \xi \rangle_h \langle x \rangle^{-\frac{\sigma}{2}} \left(1 - \psi \left(\frac{\langle x \rangle^\sigma}{\langle \xi \rangle_h^2} \right) \right).
\end{aligned}$$

Since $\langle \xi \rangle_h \langle x \rangle^{-\frac{\sigma}{2}} \leq \sqrt{2}$ on the support of $1 - \psi \left(\frac{\langle x \rangle^\sigma}{\langle \xi \rangle_h^2} \right)$, we may conclude

$$\begin{aligned}
Re \tilde{a}_1 &\geq (M_1 \frac{C_{a_3}}{2} - C_{a_1} - \tilde{C}_{a_2, \lambda_2} - C K T (1 - \sigma) h^{-\frac{\sigma}{2}}) \langle \xi \rangle_h \langle x \rangle^{-\frac{\sigma}{2}} \\
&\quad - \max\{1, K T\} C_{\tilde{\Lambda}, \rho_1} h^{-(1-\frac{1}{\theta})} \langle \xi \rangle_h \langle x \rangle^{-\frac{\sigma}{2}} - M_1 \frac{C_{a_3}}{2} \sqrt{2}.
\end{aligned}$$

Taking (6.1) into account we obtain

$$\begin{aligned}
Re (\tilde{a}_1 + t_1) &\geq (M_1 \frac{C_{a_3}}{2} - C_{a_2} - C_{a_1} - \tilde{C}_{a_2, \lambda_2} - C K T (1 - \sigma) h^{-\frac{\sigma}{2}}) \langle \xi \rangle_h \langle x \rangle^{-\frac{\sigma}{2}} \\
&\quad - \max\{1, K T\} (C_{\tilde{\Lambda}, \rho_1} + |\rho_1| C_{\lambda_2}) h^{-(1-\frac{1}{\theta})} \langle \xi \rangle_h \langle x \rangle^{-\frac{\sigma}{2}} - M_1 \frac{C_{a_3}}{2} \sqrt{2}.
\end{aligned}$$

For $Re \tilde{a}_0$, we have:

$$\begin{aligned}
Re \tilde{a}_0 &= -Im a_0 - k'(t) \langle x \rangle_h^{1-\sigma} + Re a_{1,0} + Re r_{\sigma,1} \\
&= -Im a_0 + (-k'(t) - \max\{1, k(0)\} C_T \langle x \rangle_h^{-\sigma} - C_{T, \tilde{\Lambda}} \langle x \rangle_h^{-\sigma}) \langle x \rangle_h^{1-\sigma} \\
&\geq (-C_{a_0} + K - \max\{1, K T\} C_T h^{-\sigma} - C_{T, \tilde{\Lambda}} h^{-\sigma}) \langle x \rangle_h^{1-\sigma}.
\end{aligned}$$

Finally, let us proceed with the choices of M_1, M_2 and K . First we choose K larger than $\max\{C_{a_0}, 1/T\}$, then we set M_2 large in order to obtain $M_2 \frac{C_{a_3}}{2} - C_{a_2} - \tilde{C}_{a_3} K T (1 - \sigma) > 0$ and

after that we take M_1 such that $M_1 \frac{C_{a_3}}{2} - C_{a_2} - C_{a_1} - \tilde{C}_{a_2, \lambda_2} > 0$ (choosing M_2 , M_1 we determine $\tilde{\Lambda}$). Enlarging the parameter h we may assume

$$\begin{aligned} KTC_{\lambda_2, \rho_1} h^{-(1-\frac{1}{\theta})} &< \frac{1}{4} (M_2 \frac{C_{a_3}}{2} - C_{a_2} - \tilde{C}_{a_3} KT(1-\sigma)), \\ CKT(1-\sigma)h^{-\frac{\sigma}{2}} + KT(C_{\tilde{\Lambda}, \rho_1} + |\rho_1|C_{\lambda_2})h^{-(1-\frac{1}{\theta})} &< \frac{1}{4} (M_1 \frac{C_{a_3}}{2} - C_{a_2} - C_{a_1} - \tilde{C}_{a_2, \lambda_2}), \\ KTC_T h^{-\sigma} + C_{T, \tilde{\Lambda}} h^{-\sigma} &< \frac{K - C_{a_0}}{4}. \end{aligned}$$

With these choices we obtain that $Re \tilde{a}_2 \geq 0$, $Re(\tilde{a}_1 + t_1) + M_1 \frac{C_{a_3}}{2} \sqrt{2} \geq 0$ and $Re \tilde{a}_0 \geq 0$. Let us also remark that the choices of M_2 , M_1 and $k(t)$ do not depend of ρ_1 and θ .

7. Proof of Theorem 1

Let us denote

$$\tilde{P}_\Lambda := e^{\rho_1 \langle D \rangle^{\frac{1}{\theta}}} e^\Lambda(t, x, D) iP \{e^{\rho_1 \langle D \rangle^{\frac{1}{\theta}}} e^\Lambda(t, x, D)\}^{-1}.$$

By (6.2), with the choices of M_2 , M_1 , $k(t)$ in the previous section, we get

$$i\tilde{P}_\Lambda = \partial_t + ia_3(t, D) + (Re \tilde{a}_2 + t_2)(t, x, D) + (\tilde{a}_1 + t_1)(t, x, D) + \tilde{a}_0(t, x, D) + \tilde{r}_0(t, x, D),$$

with

$$(7.1) \quad Re \tilde{a}_2 \geq 0, \quad Re(\tilde{a}_1 + t_1) + M_1 \frac{C_{a_3}}{2} \sqrt{2} \geq 0, \quad Re \tilde{a}_0 \geq 0.$$

Fefferman-Phong inequality applied to $Re \tilde{a}_2$ and sharp Gårding inequality applied to $\tilde{a}_1 + t_1 + M_1 \frac{C_{a_3}}{2} \sqrt{2}$ and \tilde{a}_0 give

$$\begin{aligned} Re \langle Re \tilde{a}_2(t, x, D)v, v \rangle &\geq -c \|v\|_{L^2}^2, \\ Re \langle (\tilde{a}_1 + t_1)(t, x, D)v, v \rangle &\geq - \left(c + M_1 \frac{C_{a_3}}{2} \sqrt{2} \right) \|v\|_{L^2}^2, \\ Re \langle \tilde{a}_0(t, x, D)v, v \rangle &\geq -c \|v\|_{L^2}^2 \end{aligned}$$

for a positive constant c . Now applying Gronwall inequality we come to the following energy estimate:

$$\|v(t)\|_{L^2}^2 \leq C \left(\|v(0)\|_{L^2}^2 + \int_0^t \|(i\tilde{P}_\Lambda v(\tau))\|_{L^2}^2 d\tau \right), \quad t \in [0, T],$$

for every $v(t, x) \in C^1([0, T]; \mathcal{S}(\mathbb{R}))$. By usual computations, this estimate provides well-posedness of the Cauchy problem associated with \tilde{P}_Λ in $H^m(\mathbb{R})$ for every $m = (m_1, m_2) \in \mathbb{R}^2$: for every $\tilde{f} \in C([0, T]; H^m(\mathbb{R}))$ and $\tilde{g} \in H^m(\mathbb{R})$, there exists a unique $v \in C([0, T]; H^m(\mathbb{R}))$ such that $\tilde{P}_\Lambda v = \tilde{f}$, $v(0) = \tilde{g}$ and

$$(7.2) \quad \|v(t)\|_{H^m}^2 \leq C \left(\|\tilde{g}\|_{H^m}^2 + \int_0^t \|\tilde{f}(\tau)\|_{H^m}^2 d\tau \right), \quad t \in [0, T].$$

Let us now turn back to our original Cauchy problem (1.10), (1.1). Fixing initial data $f \in C([0, T], H_{\rho; s, \theta}^m(\mathbb{R}))$ and $g \in H_{\rho; s, \theta}^m(\mathbb{R})$ for some $m, \rho \in \mathbb{R}^2$ with $\rho_2 > 0$ and positive s, θ such that $\theta > s_0$, we can define Λ as at the beginning of Section 6 with $\mu > 1$ such that $s_0 > 2\mu - 1$ and $M_1, M_2, k(0)$ such that (7.1) holds. Then by Theorem 4 we get

$$f_{\rho_1, \Lambda} := e^{\rho_1 \langle D \rangle^{\frac{1}{\theta}}} e^\Lambda(t, x, D) f \in C([0, T], H_{(0, \rho_2 - \delta); s, \theta}^m(\mathbb{R}))$$

and

$$g_{\rho_1, \Lambda} := e^{\rho_1 \langle D \rangle^{\frac{1}{\theta}}} e^{\Lambda}(0, x, D)g \in H_{(0, \rho_2 - \bar{\delta}); s, \theta}^m(\mathbb{R})$$

for every $\bar{\delta} > 0$, because $1/(1 - \sigma) > s$. Since $\bar{\delta}$ can be taken arbitrarily small, we have that $f_{\rho_1, \Lambda} \in C([0, T], H^m)$ and $g_{\rho_1, \Lambda} \in H^m$. Hence the Cauchy problem

$$\begin{cases} \tilde{P}_\Lambda v = f_{\rho_1, \Lambda} \\ v(0) = g_{\rho_1, \Lambda} \end{cases}$$

admits a unique solution $v \in C([0, T], H^m) \cap C^1([0, T], H^{m_1-3, m_2-1+1/\sigma})$ satisfying the energy estimate (7.2). Taking now $u = (e^{\Lambda(t, x, D)})^{-1} e^{-\rho_1 \langle D \rangle^{1/\theta}} v$, we easily see that u solves the Cauchy problem (1.1), it satisfies

$$e^{\rho_1 \langle D \rangle^{1/\theta}} e^{K(T-t) \langle x \rangle_h^{1-\sigma}} e^{\tilde{\Lambda}(x, D)} u \in H^m(\mathbb{R})$$

and it is the unique solution with this property. Namely, $u \in C([0, T], H_{(\rho_1, -\bar{\delta}); s, \theta}^m(\mathbb{R})) \cap C^1([0, T], H_{(\rho_1, -\bar{\delta}); s, \theta}^{(m_1-3, m_2)}(\mathbb{R}))$ for every $\bar{\delta} > 0$. Moreover, from (7.2) we get

$$\begin{aligned} \|u\|_{H_{(\rho_1, -\bar{\delta}); s, \theta}^m} &\leq C \|v\|_{H^m} \leq C \left(\|g_{\rho_1, \Lambda}\|_{H^m}^2 + \int_0^t \|f_{\rho_1, \Lambda}(\tau)\|_{H^m}^2 d\tau \right) \\ &\leq C \left(\|g\|_{H_{\rho; s, \theta}^m}^2 + \int_0^t \|f(\tau)\|_{H_{\rho; s, \theta}^m}^2 d\tau \right). \end{aligned}$$

which gives (1.11) This concludes the proof. \square

Remark 6. Notice that the argument of the proof of Theorem 1 and in particular the energy estimate (1.11) implies that the solution of the problem (1.1) is unique in the space of all functions u such that

$$e^{\rho_1 \langle D \rangle^{1/\theta}} e^{K(T-t) \langle x \rangle_h^{1-\sigma}} e^{\tilde{\Lambda}(x, D)} u \in H^m(\mathbb{R}).$$

In general we cannot conclude that it is unique in $C([0, T], H_{(\rho_1, -\bar{\delta}); s, \theta}^m(\mathbb{R}))$.

Remark 7. In our main result we assume that the symbol of the leading term $a_3(t, D)$ is independent of x . This assumption is crucial in the argument of the proof. As a matter of fact, we observe that if a_3 depends on x , even allowing it to decay like $\langle x \rangle^{-m}$ for $m \gg 0$, the conjugation of this term with the operator $e^{\rho_1 \langle D \rangle^{1/\theta}}$ would give

$$e^{\rho_1 \langle D \rangle^{1/\theta}} (ia_3(t, x, D)) e^{-\rho_1 \langle D \rangle^{1/\theta}} = ia_3(t, x, D_x) + op \left(\rho_1 \partial_\xi \langle \xi \rangle^{\frac{1}{\theta}} \cdot \partial_x a_3 \right) + l.o.t.$$

We observe that

$$\rho_1 \partial_\xi \langle \xi \rangle^{\frac{1}{\theta}} \cdot \partial_x a_3(t, x, \xi) \sim \langle \xi \rangle^{2+\frac{1}{\theta}} \langle x \rangle^{m-1}.$$

Hence since $2 + \frac{1}{\theta} > 2$, this remainder term could not be controlled by other lower order terms whose order in ξ does not exceed 2. Notice that this represents a difference in comparison with the H^∞ frame where a dependence on x in the leading term can be allowed by assuming a suitable decay assumption with respect to x , cf. [2, Section 4].

Remark 8. With a major technical effort one can consider 3-evolution equations in higher space dimension, that is for $x \in \mathbb{R}^n, n > 1$. At this moment, results of this type exist only for the case $p = 2$, cf. [6, 11, 24]. When passing to higher dimension, the main difficulty is the choice of the functions λ_1, λ_2 defining the change of variable, which must be chosen in order to satisfy certain partial differential inequalities, see also [2, Section 4]. These may be non trivial for $p > 2$. In this paper we prefer to restrict to the one space dimensional case both since the content is already quite technical and since the main physical models to which our results could be applied are included in this case.

Acknowledgements. The authors are grateful to the referees for their valuable and stimulating suggestions which helped them to improve the presentation of their results.

References

- [1] A. Arias Junior, A. Ascanelli, M. Capiello, *Gevrey well posedness for 3-evolution equations with variable coefficients*. Preprint (2021), <https://arxiv.org/abs/2106.09511>
- [2] A. Ascanelli, C. Boiti, *Semilinear p -evolution equations in Sobolev spaces*, J. Differential Equations **260** (2016), 7563-7605.
- [3] A. Ascanelli, C. Boiti, L. Zanghirati, *Well-posedness of the Cauchy problem for p -evolution equations*. J. Differential Equations **253** (10) (2012), 2765-2795.
- [4] A. Ascanelli, C. Boiti, L. Zanghirati, *A Necessary condition for H^∞ well-posedness of p -evolution equations*. Adv. Differential Equations **21** (2016), 1165-1196.
- [5] A. Ascanelli, M. Capiello, *Weighted energy estimates for p -evolution equations in SG classes*. J. Evol. Eqs, **15** (3) (2015), 583-607.
- [6] A. Ascanelli, M. Capiello, *Schrödinger-type equations in Gelfand-Shilov spaces*. J. Math. Pures Appl. **132** (2019), 207-250.
- [7] A. Ascanelli, M. Cicognani, F. Colombini, *The global Cauchy problem for a vibrating beam equation*, J. Differential Equations **247** (2009), 1440-1451.
- [8] A. Ascanelli, M. Cicognani, M. Reissig, *The interplay between decay of the data and regularity of the solution in Schrödinger equations*. Ann. Mat. Pura Appl. (1923-) **199** (4) (2020), 1649-1671.
- [9] A. Baba, *The H_∞ -wellposed Cauchy problem for Schrödinger type equations*. Tsukuba J. Math. **18** (1) (1994), 101-117.
- [10] M. Cicognani, F. Colombini, *The Cauchy problem for p -evolution equations*. Trans. Amer. Math. Soc. **362** (9) (2010), 4853-4869.
- [11] M. Cicognani, M. Reissig, *Well-posedness for degenerate Schrödinger equations*. Evol. Equ. Control Theory **3** (1) (2014), 15-33.
- [12] A. Dasgupta, M.W. Wong, *Spectral Invariance of SG Pseudo-Differential Operators on $L^p(\mathbb{R}^n)$* . Pseudo-Differential Operators: Complex Analysis and Partial Differential Equations, Springer (2009), 51-57.
- [13] A. Debrouwere, L. Neyt, J. Vindas, *The nuclearity of Gelfand-Shilov spaces and kernel theorems*. Collect. Math. **72** (1) (2021), 203-227.
- [14] M. Dreher, *Necessary conditions for the well-posedness of Schrödinger type equations in Gevrey spaces*, Bull. Sci. Mat. **127** (2003), 485-503.
- [15] C. Fefferman, D.H. Phong, *On positivity of pseudo-differential operators*. Proc. Nat. Acad. Sci. U.S.A. **75** (1978) (10), 4673-4674.
- [16] R. Figueroa, A.A. Himonas, F. Yan, *A higher dispersion KdV equation on the line*. Nonlinear Anal. **199** (2020), 112055, 38 pp.
- [17] I.M. Gelfand, G.E. Shilov, *Generalized functions, Vol. 2*, Academic Press, New York-London, 1967.
- [18] J. Gorsky, A.A. Himonas, C. Holliman, G. Petronilho, *The Cauchy problem of a periodic KdV equation in analytic Gevrey spaces*, J. Math. Anal. Appl. **405** (2) (2013), 349-361.
- [19] O. Goubet, *Gevrey regularity of the global attractor for damped forced KdV equation on the real line*, J. Math. Study **51** (3) (2018), 294-308.
- [20] H. Hannah, A.A. Himonas, G. Petronilho, *Anisotropic Gevrey regularity for m KdV on the circle*. Discrete Contin. Dyn. Syst. 2011, Dynamical systems, differential equations and applications. 8th AIMS Conference. Suppl. Vol. I, 634-642.
- [21] G. Harutyunyan, B.-W. Schulze, *Elliptic mixed, transmission and singular crack problems*. EMS Tracts in Mathematics, 4. European Mathematical Society (EMS), Zürich (2008), xii+765 pp.
- [22] W. Ichinose, *Some remarks on the Cauchy problem for Schrödinger type equations*. Osaka J. Math. **21** (3) (1984), 565-581.
- [23] A. Jeffrey, M.N.B. Mohamad, *Exact solutions to the KdV-Burgers' equation*, Wave Motion **14** (4) (1991), 369-375.
- [24] K. Kajitani, A. Baba, *The Cauchy problem for Schrödinger type equations*. Bull. Sci. Math. **119** (5) (1995), 459-473.
- [25] K. Kajitani, T. Nishitani, *The hyperbolic Cauchy problem*. Lecture Notes in Mathematics **1505** Springer-Verlag, Berlin, 1991.

- [26] T. Kato, *On the Cauchy problem for the (generalized) Korteweg-de Vries equation*, Studies in Applied Mathematics, 93-128, Adv. Math. Suppl. Stud. **8** Academic Press, New York, 1983.
- [27] K. Kato, M. Kawamoto, K. Nambu, *Singularity for solutions of linearized KdV equations*, J. Math. Phys. **61** (5) (2020), 051502, 11 pp.
- [28] D.J. Korteweg, G. De Vries, *On the change of form of long waves advancing in a rectangular canal, and on a new type of long stationary waves*, Philos. Mag. (5) **39** (240) (1895), 422-443.
- [29] S. Mizohata, *Some remarks on the Cauchy problem*, J. Math. Kyoto Univ. **1** (1) (1961), 109-127.
- [30] F. Nicola, L. Rodino, *Global pseudo-differential calculus on Euclidean spaces*. Series Pseudo-Differential Operators **4**, Springer, 2010.
- [31] S. Pilipović, *Tempered ultradistributions*, Boll. Unione Mat. Ital., VII. Ser., B, **2** (2) (1988), 235-251.
- [32] M. Tsutsumi, T. Musaka, I. Yoshi, *On the generalized Korteweg-de Vries equation*, Proc. Japan Acad. **46** (1970), 921-925.
- [33] M. W. Wong, *An introduction to pseudo-differential operators*. World Scientific Publishing Company (2014) **6**.

DEPARTMENT OF MATHEMATICS, FEDERAL UNIVERSITY OF PARANÁ, CEP 81531-980, CURITIBA, BRAZIL
Email address: arias@ufpr.br

DIPARTIMENTO DI MATEMATICA ED INFORMATICA, UNIVERSITÀ DI FERRARA, VIA MACHIAVELLI 30,
 44121 FERRARA, ITALY
Email address: alessia.ascanelli@unife.it

DIPARTIMENTO DI MATEMATICA "G. PEANO", UNIVERSITÀ DI TORINO, VIA CARLO ALBERTO 10, 10123
 TORINO, ITALY
Email address: marco.cappiello@unito.it