ON DISPERSION MANAGED NONLINEAR SCHRÖDINGER EQUATIONS WITH LUMPED AMPLIFICATION

MI-RAN CHOI[†], YOUNGHOON KANG[‡], YOUNG-RAN LEE[‡]

ABSTRACT. We show the global well–posedness of the nonlinear Schrödinger equation with periodically varying coefficients and a small parameter $\varepsilon > 0$, which is used in optical–fiber communications. We also prove that the solutions converge to the solution for the Gabitov–Turitsyn or averaged equation as ε tends to zero.

1. INTRODUCTION

We consider the nonlinear Schrödinger equation (NLS) with periodically varying coefficients

$$i\partial_t u + d(t)\partial_x^2 u + c(t)|u|^2 u = 0$$

$$\tag{1.1}$$

which describes the behavior of a signal transmitted on an optical-fiber cable. Here, x denotes the (retarded) time, t the position along the cable, and periodic functions $d(\cdot)$ and $c(\cdot)$ the dispersion and the fiber loss/amplification along the cable respectively.

The original evolution of optical pulses in a dispersion managed system with lumped amplification is described by the nonlinear Schrödinger equation

$$i\partial_t E + d(t)\partial_x^2 E + |E|^2 E = ig(t)E$$

The fiber loss and amplification coefficient along the cable is given by

$$g(t) = -\frac{\Gamma}{2} + \Gamma \sum_{j=1}^{\infty} \delta(t - t_j),$$

where $\Gamma > 0$ is the fiber loss, t_j corresponds to the location of amplifiers, and $\delta(\cdot)$ is the Dirac delta function. For more information on this equation, see, e.g., [2]. Taking

$$E(x,t) = u(x,t) \exp\left(\int_0^t g(t')dt'\right),$$

we obtain equation (1.1) with $c(t) = \exp\left(2\int_0^t g(t')dt'\right)$ provided that g is a periodic function with mean zero.

The dispersion management with alternating sections of positive and negative dispersion in fibers was introduced in 1980, see [23]. It was successful to transfer the data at ultra-high speed over long distances, see, e.g., [1, 12, 14, 15, 22, 27]. For more information on the dispersion management, see [26] and references therein.

In the strong dispersion management regime, the dispersion is given by

$$d(t) = d_{\mathrm{av}} + \frac{1}{\varepsilon} d_0 \left(\frac{t}{\varepsilon}\right),$$

Key words and phrases. nonlinear Schrödinger equation, dispersion management, well-posedness, averaging .

MSC2020 classification.35Q55, 35Q60, 35A01.

^{©2021} by the authors. Faithful reproduction of this article, in its entirety, by any means is permitted for non-commercial purposes.

where $d_0(\cdot)$ is the mean zero part of the dispersion which is a 2-periodic function satisfying

$$d_0(t) = \begin{cases} 1, & 0 \le t < 1, \\ -1, & 1 \le t < 2, \end{cases}$$

 $d_{\rm av} \in \mathbb{R}$ the average dispersion over one period, and $\varepsilon > 0$ a small parameter. The fiber loss and amplification is defined to be

$$c(t) = G\left(\frac{t}{\varepsilon}\right),$$

where G is also a 2-periodic function given by

$$G(t) = \exp\left(2\int_0^t \left(-\frac{\Gamma}{2} + \Gamma \sum_{j \in \mathbb{Z}} \delta(t' - 2j)\right) dt'\right).$$

The first main result of this paper is the well–posedness of the Cauchy problem

$$\begin{cases} i\partial_t u + \left(d_{\mathrm{av}} + \frac{1}{\varepsilon}d_0\left(\frac{t}{\varepsilon}\right)\right)\partial_x^2 u + G\left(\frac{t}{\varepsilon}\right)|u|^2 u = 0,\\ u(x,0) = u_0(x). \end{cases}$$
(1.2)

Theorem 1.1 (Global well–posedness). Let $d_{av} \in \mathbb{R}$. For every $u_0 \in H^1(\mathbb{R})$, there exists a unique solution $u \in \mathcal{C}(\mathbb{R}, H^1(\mathbb{R}))$ of (1.2). Moreover, u depends continuously on the initial datum in the following sense. For every M > 0, the map $u_0 \mapsto u(t)$ from $H^1(\mathbb{R})$ to $\mathcal{C}([-M, M], H^1(\mathbb{R}))$ is locally Lipschitz continuous.

Now we change the variables $u = T_{D(t/\varepsilon)}v$ in (1.2) to obtain

$$\begin{cases} i\partial_t v + d_{\rm av}\partial_x^2 v + G\left(\frac{t}{\varepsilon}\right)T_{D(t/\varepsilon)}^{-1}\left(|T_{D(t/\varepsilon)}v|^2 T_{D(t/\varepsilon)}v\right) = 0,\\ v(x,0) = u_0(x), \end{cases}$$
(1.3)

where $D(t) = \int_0^t d_0(t')dt'$ and T_t is the solution operator for the free Schrödinger equation in dimension one. Note that since d_0 is a 2-periodic function with mean zero, D is also 2-periodic and, therefore, the map $t \mapsto T_{D(t/\varepsilon)}$ is 2ε -periodic.

For small $\varepsilon > 0$, that is, in the regime of strong dispersion management, equation (1.3) contains the fast oscillating terms $T_{D(t/\varepsilon)}$ and $G(t/\varepsilon)$ in the nonlinearity and hence Gabitov and Turitsyn suggested averaging the equation over one period, see [14, 15]. This yields the following "averaged" equation

$$i\partial_t v + d_{\rm av}\partial_x^2 v + \frac{1}{2}\int_0^2 G(\tau)T_{D(\tau)}^{-1}\Big(|T_{D(\tau)}v|^2 T_{D(\tau)}v\Big)d\tau = 0.$$

We make the change of variables $D(\tau) = r$, then we have

$$i\partial_t v + d_{\rm av}\partial_x^2 v + \int_0^1 T_r^{-1} \Big(|T_r v|^2 T_r v \Big) \psi(r) dr = 0, \tag{1.4}$$

where

$$\psi(r) = e^{-\Gamma} \cosh \Gamma(r-1).$$

The well-posedness of the averaged equation (1.4) in $H^s(\mathbb{R})$ for all $s \geq 0$ is proved in [3] for a general dispersion profile. For more general nonlinearities including saturated nonlinearities, see [11].

The averaging procedure is rigorously justified in [28] when the fiber loss and amplification are not present. More precisely, it is shown that for $\varepsilon > 0$, the solutions of (1.3) and (1.4) with the same initial datum in $H^s(\mathbb{R})$ stay ε -close in $H^{s-3}(\mathbb{R})$ for a long time in $O(\varepsilon^{-1})$ when s is sufficiently large. Note that the convergence is not shown in $H^s(\mathbb{R})$ where the solutions exist. However, we prove that the solutions for (1.3) converge to the solution for (1.4) in $H^1(\mathbb{R})$, where the solutions exist, as $\varepsilon \to 0$.

Theorem 1.2 (Averaging Theorem). Let $d_{av} \in \mathbb{R}$, M > 0 and $v \in C(\mathbb{R}, H^1(\mathbb{R}))$ be the solution of the averaged equation (1.4) with the initial datum $v_0 \in H^1(\mathbb{R})$. Then there exist C > 0 and $\varepsilon_0 > 0$ such that if $0 < \varepsilon \le \varepsilon_0$ and $||u_0 - v_0||_{H^1(\mathbb{R})} \le \varepsilon$, then

$$\|v_{\varepsilon} - v\|_{\mathcal{C}([-M,M],H^1(\mathbb{R}))} \le C\varepsilon, \tag{1.5}$$

where v_{ε} is the solution of (1.3) with the initial datum $u_0 \in H^1(\mathbb{R})$

In our main theorems, Theorems 1.1 and 1.2, we prove the well-posedness of the Cauchy problem (1.2) and the validity of the averaging process in the strong dispersion, $\frac{1}{\varepsilon}d_0(\frac{t}{\varepsilon})$, while the well-posedness of the averaged equation is already proved in [3]. There are analogous results for the fast dispersion, $d_0(\frac{t}{\varepsilon})$, and the random dispersion, $\frac{1}{\varepsilon}d_0(\frac{t}{\varepsilon^2})$, with some centered stationary random process d_0 , in [5] and [6], respectively.

We here remark some provable facts which are not dealt with in this paper. Related to standing wave solutions $v(x,t) = e^{i\omega t}f(x)$, $\omega \in \mathbb{R}$, of the averaged equation (1.4), a constrained minimization problem is well studied. The existence of minimizers can be found in [10, 19] when $d_{av} \geq 0$. One can easily show that every minimizer is a weak solution of the corresponding Euler–Lagrange equation. Each weak solution and its Fourier transform decay exponentially, which can be proven by modifying the proofs in [13, 17] a little. Particularly, every minimizer is smooth. Moreover, the set of ground states is orbitally stable, see [11, 18] as well as [8]. In the case $d_{av} > 0$, using the averaging theorem and the orbital stability, it is possible to obtain the stable soliton–like solution for (1.2).

The paper is organized as follows. In Section 2, we prove Theorem 1.1, the global well– posedness result in $H^1(\mathbb{R})$. We start by showing the local well–posedness in $H^1(\mathbb{R})$ and the global existence in $L^2(\mathbb{R})$. Although we do not have the energy conservation, we prove the existence of a global solution in $H^1(\mathbb{R})$ based on the mass conservation and the boundedness of the mixed norm of the solution for Strichartz admissible pairs. In Section 3, we prove Theorem 1.2, the averaging theorem. In Appendix A, we gather basic properties of the free Schrödinger time evolution to prove Theorem 1.1.

2. Well-posedness

To begin with, let us introduce some notations. For $1 \leq p < \infty$, we use $L^p(\mathbb{R})$ to denote the Banach space of functions f whose norm

$$||f||_{L^p} := \left(\int_{\mathbb{R}} |f(x)|^p dx\right)^{\frac{1}{p}}$$

is finite with the essential supremum instead when $p = \infty$. The space $L^2(\mathbb{R})$ is a Hilbert space with scalar product given by $\langle f, g \rangle = \int_{\mathbb{R}} f(x)\overline{g(x)}dx$. We use $L^q_t(J, L^p_x(I))$ to denote, for $1 \leq p, q < \infty$ and intervals $I, J \subset \mathbb{R}$, the Banach space of all functions u with the mixed norm

$$||u||_{L^{q}_{t}(J,L^{p}_{x}(I))} := \left(\int_{J} \left(\int_{I} |u(x,t)|^{p} dx \right)^{\frac{q}{p}} dt \right)^{\frac{1}{q}}$$

If $p = \infty$ or $q = \infty$, use the usual modification. For notational simplicity, we use $L^q(J, L^p)$ for $L^q_t(J, L^p_x(\mathbb{R}))$. We say that $u \in L^q_{loc}(J, L^p)$ when $u \in L^q(\widetilde{J}, L^p)$ for every bounded interval $\widetilde{J} \subset J$.

The Fourier transform on \mathbb{R} is defined by

$$\widehat{f}(\xi) := \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{-ix\xi} f(x) \, dx$$

for $f \in \mathcal{S}(\mathbb{R})$, the Schwartz space of infinitely smooth, rapidly decreasing functions. For $s \in \mathbb{R}$, the Sobolev space $H^s(\mathbb{R})$ is defined as the space of all tempered distributions $f \in S'(\mathbb{R})$ for which

$$||f||_{H^s} := \left(\int_{\mathbb{R}} (1+\xi^2)^s |\widehat{f}(\xi)|^2 d\xi\right)^{1/2} < \infty.$$

For a Banach space X with norm $\|\cdot\|_X$ and an interval J, C(J,X) is the space of all continuous functions $u: J \to X$. When J is compact, it is a Banach space with norm

$$||u||_{C(J,X)} = \sup_{t \in J} ||u(t)||_X$$

and $C^1(J, X)$ is the Banach space of all continuously differentiable functions $u: J \to X$.

Let T_t denote the solution operator for the free Schrödinger equation in spatial dimension one. In terms of the Fourier transform, this is given by

$$\widehat{T_t f}(\xi) = \widehat{e^{it\partial_x^2} f}(\xi) = e^{-it\xi^2} \widehat{f}(\xi)$$

for $f \in \mathcal{S}(\mathbb{R})$, thus, one can express

$$T_t f(x) = e^{it\partial_x^2} f(x) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{ix\xi} e^{-it\xi^2} \widehat{f}(\xi) d\xi.$$

It is a unitary operator on $L^2(\mathbb{R})$ and also on $H^1(\mathbb{R})$. Therefore, for every $t \in \mathbb{R}$,

 $||T_t f||_{L^2} = ||f||_{L^2}$ and $||T_t f||_{H^1} = ||f||_{H^1}.$

We use the notation $f \leq g$ when there exists a positive constant C such that $f \leq Cg$.

Now we prove the well–posedness of the Cauchy problem (1.2). Since the proof does not rely on the factor ε in (1.2), we only consider the case $\varepsilon = 1$,

$$\begin{cases} i\partial_t u + (d_{av} + d_0(t))\partial_x^2 u + G(t)|u|^2 u = 0, \\ u(x,0) = u_0(x), \end{cases}$$
(2.1)

or, equivalently,

$$\begin{cases} i\partial_t v + d_{\rm av}\partial_x^2 v + G(t)T_{D(t)}^{-1} \left(|T_{D(t)}v|^2 T_{D(t)}v \right) = 0, \\ v(x,0) = u_0(x), \end{cases}$$
(2.2)

where $u = T_{D(t)}v$. We first prove the local existence of a unique solution for the integral equation of (2.2),

$$v(t) = e^{itd_{av}\partial_x^2} u_0 + i \int_0^t e^{i(t-t')d_{av}\partial_x^2} Q(v(t'))dt',$$
(2.3)

with $u_0 \in H^1(\mathbb{R})$, where

$$Q(v)(t) = G(t)T_{D(t)}^{-1} \Big(|T_{D(t)}v(t)|^2 T_{D(t)}v(t) \Big)$$

Here and below, we use C to denote various constants.

Lemma 2.1. For every $f, g \in H^1(\mathbb{R})$, we have

$$\|Q(f)(t)\|_{H^1} \lesssim \|G\|_{L^{\infty}} \|f\|_{H^1}^3 \tag{2.4}$$

and

$$\|Q(f)(t) - Q(g)(t)\|_{H^1} \lesssim \|G\|_{L^{\infty}} (\|f\|_{H^1}^2 + \|g\|_{H^1}^2) \|f - g\|_{H^1}.$$
(2.5)

Proof. We define, for every $f_1, f_2, f_3 \in H^1(\mathbb{R})$,

$$Q(f_1, f_2, f_3)(t) := G(t)T_{D(t)}^{-1} \Big(T_{D(t)}f_1 \overline{T_{D(t)}f_2} T_{D(t)}f_3 \Big),$$

which is multi-linear. Note that Q(f)(t) = Q(f, f, f)(t) for any $f \in H^1(\mathbb{R})$. Since

 $\|fg\|_{H^1} \le C \|f\|_{H^1} \|g\|_{H^1}$

and $T_{D(s)}$ is unitary on $H^1(\mathbb{R})$, we obtain

$$\|Q(f_1, f_2, f_3)(t)\|_{H^1} \lesssim \|G\|_{L^{\infty}} \|f_1\|_{H^1} \|f_2\|_{H^1} \|f_3\|_{H^1}$$
(2.6)

which proves (2.4). Observing

$$Q(f)(t) - Q(g)(t) = Q(f - g, f, f)(t) + Q(g, f - g, f)(t) + Q(g, g, f - g)(t),$$

one can easily obtain (2.5) from (2.6).

Proposition 2.2. Let $d_{av} \in \mathbb{R}$. For every K > 0, there exist $M_{\pm} > 0$ such that for every initial datum $u_0 \in H^1(\mathbb{R})$ with $||u_0||_{H^1} \leq K$, there is a unique solution $v \in \mathcal{C}([-M_-, M_+], H^1)$ of (2.3). Moreover,

$$||v(t)||_{H^1} \le 2K$$
 for all $t \in [-M_-, M_+]$.

Corollary 2.3. Let $d_{av} \in \mathbb{R}$. For any initial datum $u_0 \in H^1(\mathbb{R})$, there exist maximal life times $T_{\pm} \in (0, \infty]$ such that there is a unique solution $v \in \mathcal{C}((-T_-, T_+), H^1)$ of (2.3). Moreover, the blow-up alternative for solutions holds: if T_+ is finite, then

$$\lim_{t \to T_+} \|v(t)\|_{H^1} = \infty$$

and if T_{-} is finite, then

$$\lim_{t \to -T_{-}} \|v(t)\|_{H^{1}} = \infty.$$

Proof of Proposition 2.2. Without loss of generality, we assume that t > 0.

To prove the existence of a solution, we use a fixed point argument. For each M > 0 and a > 0, let

$$B_{M,a} = \{ v \in \mathcal{C}([0, M], H^1) : \|v\|_{\mathcal{C}([0, M], H^1)} \le a \}$$

be equipped with the distance

$$d(v,w) = \|v - w\|_{\mathcal{C}([0,M],H^1)}.$$

Let K > 0 and $u_0 \in H^1(\mathbb{R})$ with $||u_0||_{H^1} \leq K$ be fixed. Define the map Φ on $B_{M,a}$ by

$$\Phi(v)(t) = e^{itd_{\mathrm{av}}\partial_x^2}u_0 + i\int_0^t e^{i(t-t')d_{\mathrm{av}}\partial_x^2}Q(v)(t')dt'.$$

It follows from Lemma 2.1 that if $v(t), w(t) \in H^1(\mathbb{R})$, then

$$\|\Phi(v)(t)\|_{H^1} \le \|u_0\|_{H^1} + \int_0^t \|Q(v)(t')\|_{H^1} dt' \le \|u_0\|_{H^1} + C \int_0^M \|v(t')\|_{H^1}^3 dt'$$

and

$$\begin{split} \|\Phi(v)(t) - \Phi(w)(t)\|_{H^{1}} &\leq \int_{0}^{t} \|Q(v)(t') - Q(w)(t')\|_{H^{1}} dt' \\ &\leq C \int_{0}^{M} \left(\|v(t')\|_{H^{1}}^{2} + \|w(t')\|_{H^{1}}^{2} \right) \|v(t') - w(t')\|_{H^{1}} dt' \end{split}$$

$$(2.7)$$

for every $0 \le t \le M$. Therefore, there is a positive constant C such that for all $v, w \in B_{M,a}$,

 $\|\Phi(v)\|_{\mathcal{C}([0,M],H^1)} \le K + CMa^3$

and

$$d(\Phi(v), \Phi(w)) \le CMa^2 d(v, w).$$

Now set a = 2K and choose $M_+ > 0$ satisfying

$$CM_+(2K)^2 < \frac{1}{2},$$

then we obtain that Φ is a contraction from $B_{M_+,2K}$ into itself. Thus, Banach's contraction mapping theorem shows that there exists a unique solution v of (2.3) in $B_{M_+,2K}$ and, moreover,

$$||v||_{\mathcal{C}([0,M_+],H^1)} \le 2K$$

To show the uniqueness of a solution, let $v_1, v_2 \in \mathcal{C}([0, M_+], H^1)$ be solutions of (2.3). Then it follows from (2.5) that for every $t \in [0, M_+]$

$$\|v_1(t) - v_2(t)\|_{H^1} \le C \left(\|v_1\|_{\mathcal{C}([0,M_+],H^1)}^2 + \|v_2\|_{\mathcal{C}([0,M_+],H^1)}^2 \right) \int_0^t \|v_1(t') - v_2(t')\|_{H^1} dt' \quad (2.8)$$

which implies $||v_1 - v_2||_{\mathcal{C}([0, M_+], H^1)} = 0.$

Proof of Corollary 2.3. Given the initial datum $0 \neq u_0 \in H^1(\mathbb{R})$, let us define the maximal life time T_+ by

 $T_+ = \sup \{M > 0 : \text{a unique solution of } (2.3) \text{ exists in } \mathcal{C}([0, M], H^1) \}.$

Then it immediately follows from Proposition 2.2 that $T_+ \in (0, \infty]$.

Note that if a solution exists in $\mathcal{C}([0, M], H^1)$ for any M > 0, then it is a unique solution in $\mathcal{C}([0, M], H^1)$, by the same argument in the proof of the uniqueness in Proposition 2.2. Thus, there exists a unique solution $v \in \mathcal{C}([0, T_+), H^1)$ of (2.3). To prove the blow-up alternative, let $T_+ < \infty$. Suppose to the contrary that there exist a positive number K and a sequence $\{t_j\}$ in $(0, T_+)$ such that $\|v(t_j)\|_{H^1} \leq K$ and $t_j \to T_+$ as $j \to \infty$. Then, by Proposition 2.2, there exists M > 0 such that a unique solution of (2.3) with the initial datum $v(t_j)$ exists in $\mathcal{C}([t_j, t_j + M], H^1)$ for all j. Since we can choose j^* such that $t_{j^*} + M > T_+$, this contradicts the definition of T_+ . The case of T_- can be done similarly.

Next, we show the continuous dependence of the solutions for (2.3) on the initial data to finish the local well–posedness. Indeed, the map $u_0 \mapsto v(t)$ is locally Lipschitz continuous on $H^1(\mathbb{R})$.

Proposition 2.4. Let $d_{av} \in \mathbb{R}$. For every K > 0, there exists a positive constant C such that for all initial data $v_0, w_0 \in H^1(\mathbb{R})$ with $\|v_0\|_{H^1}, \|w_0\|_{H^1} \leq K$, we have

$$||v - w||_{\mathcal{C}([-M_{-},M_{+}],H^{1})} \le e^{C \max(M_{-},M_{+})} ||v_{0} - w_{0}||_{H^{1}},$$

where v and w are the corresponding local solutions of (2.3) with initial data v_0 , w_0 on the time interval $[-M_-, M_+]$ of existence, guaranteed by Proposition 2.2.

Proof. We consider positive t only and fix $t \in (0, M_+]$. From Proposition 2.2, we know that $\|v(t)\|_{H^1} \leq 2K$ and $\|w(t)\|_{H^1} \leq 2K$.

By a similar argument of (2.7), we obtain

$$\begin{aligned} \|v(t) - w(t)\|_{H^{1}} &\leq \|v_{0} - w_{0}\|_{H^{1}} + \int_{0}^{t} \|Q(v)(t') - Q(w)(t')\|_{H^{1}} dt' \\ &\leq \|v_{0} - w_{0}\|_{H^{1}} + C \int_{0}^{t} \left(\|v(t')\|_{H^{1}}^{2} + \|w(t')\|_{H^{1}}^{2}\right) \|v(t') - w(t')\|_{H^{1}} dt' \\ &\leq \|v_{0} - w_{0}\|_{H^{1}} + CK^{2} \int_{0}^{t} \|v(t') - w(t')\|_{H^{1}} dt'. \end{aligned}$$

Thus, it follows from Gronwall's inequality that

$$\|v(t) - w(t)\|_{H^1} \le e^{CK^2 t} \|v_0 - w_0\|_{H^1} \le e^{CK^2 M_+} \|v_0 - w_0\|_{H^1},$$

which completes the proof.

Remark 2.5. If we define the energy E(v(t)) of the solution v for (2.2) by

$$E(v(t)) = \frac{d_{\mathrm{av}}}{2} \|\partial_x v(t)\|_{L^2}^2 - \frac{G(t)}{4} \int_{\mathbb{R}} |T_{D(t)}v(t)|^4 dx,$$

then, however, the energy is neither conserved nor decreasing. Indeed, its derivative is given by

$$\frac{dE(v(t))}{dt} = -\frac{1}{4}G'(t)\int_{\mathbb{R}}|T_{D(t)}v(t)|^4dx$$
(2.9)

for all $t \in \mathbb{R} \setminus 2\mathbb{Z}$. Note that E(v(t)) is not differentiable nor continuous at $t \in 2\mathbb{Z}$. If there is no fiber loss nor amplification, i.e., $G \equiv 1$, then $T_{\pm} = \infty$ by the conservation of energy and the blow-up alternative, which immediately gives the global well-posedness. However, as you see in (2.9), the energy is no longer conserved in our case.

Now we consider the integral form of the Cauchy problem (2.1)

$$u(t) = U(0,t)u_0 + i \int_0^t U(t',t) \Big(G(t')|u(t')|^2 u(t') \Big) dt'.$$
(2.10)

Here and below, U(0,t) is the solution operator for the linear Schrödinger equation associated with (2.1), i.e., for every $f \in L^2(\mathbb{R})$, U(0,t)f solves the initial value problem

$$\begin{cases} i\partial_t w + (d_{\rm av} + d_0(t))\partial_x^2 w = 0, \\ w(x,0) = f(x). \end{cases}$$

Next we define, for all $t_0, t \in \mathbb{R}$,

$$U(t_0, t) := U(0, t) (U(0, t_0))^{-1}$$
(2.11)

on $L^2(\mathbb{R})$. Then $U(t_0, t)$ is unitary on $L^2(\mathbb{R})$ and also on $H^1(\mathbb{R})$, and therefore, for every $t, t_0 \in \mathbb{R}$,

$$||U(t_0,t)f||_{L^2} = ||f||_{L^2}$$
 and $||U(t_0,t)f||_{H^1} = ||f||_{H^1}$.

For more properties of $U(t_0, t)$, see Appendix A.

Note that, given $u_0 \in H^1(\mathbb{R})$, there exists a unique solution $v \in \mathcal{C}((-T_-, T_+), H^1)$ for equation (2.3) by Corollary 2.3. If we let $u = T_{D(t)}v$, then $u \in \mathcal{C}((-T_-, T_+), H^1)$ solves equation (2.10) and the blow-up alternative holds since $\{T_{D(t)} : t \in \mathbb{R}\}$ is a strongly continuous group of unitary operators on $H^1(\mathbb{R})$. Moreover, since $H^1(\mathbb{R}) \hookrightarrow L^{\infty}(\mathbb{R})$, we have $u \in L^{\infty}((-T_{-}, T_{+}), L^{2}) \cap L^{4}_{loc}((-T_{-}, T_{+}), L^{\infty})$ and, therefore, by the Riesz-Thorin interpolation Theorem,

$$u \in L^q_{\text{loc}}((-T_-, T_+), L^p)$$
 (2.12)

for every admissible pair (p, q). Before we prove the global existence of a solution, we show the existence of a unique global solution of (2.10) with the initial datum $u_0 \in L^2(\mathbb{R})$ when $d_{av} \neq \pm 1$. As usual, we use the Strichartz estimates (Lemma A.2) to prove the existence of a local solution for (2.10), see [7, 20] for example.

Proposition 2.6. Let $d_{av} \neq \pm 1$. For any $u_0 \in L^2(\mathbb{R})$, there exists a unique global solution $u \in \mathcal{C}(\mathbb{R}, L^2) \cap L^6_{loc}(\mathbb{R}, L^6)$ of (2.10). Moreover, the solution u satisfies

$$||u(t)||_{L^2} = ||u_0||_{L^2}$$
 for all $t \in \mathbb{R}$.

Furthermore, for every M > 0 and admissible pair (p,q), there exists a positive constant C depending on d_{av} and $||u_0||_{L^2}$ such that

$$\|u\|_{L^q([-M,M],L^p)} \le C.$$
(2.13)

Proof. Let $0 \neq u_0 \in L^2(\mathbb{R})$ be fixed. Without loss of generality, we consider positive t only. First, to prove the existence of a unique solution in $\mathcal{C}([0,1], L^2) \cap L^6([0,1], L^6)$ of (2.10), let us define the closed ball

$$B_{M,a} := \{ u \in L^{\infty}([0,M], L^2) \cap L^6([0,M], L^6) : \|u\|_{L^{\infty}([0,M], L^2)} + \|u\|_{L^6([0,M], L^6)} \le a \}$$

equipped with the distance

$$d(u,v) = \|u - v\|_{L^{\infty}([0,M],L^2)} + \|u - v\|_{L^6([0,M],L^6)}$$

for each $0 < M \leq 1$ and a > 0. Define the map Φ on $B_{M,a}$ by

$$\Phi(u)(t) = U(0,t)u_0 + i \int_0^t U(t',t) \Big(G(t') |u(t')|^2 u(t') \Big) dt'.$$

For appropriate values of M and a, the map Φ is a contraction on $(B_{M,a}, d)$. Indeed, it follows from the Stricharz estimates (Lemma A.2) and the Cauchy–Schwarz inequality that

$$\begin{split} \|\Phi(u)\|_{L^{6}([0,M],L^{6})} &\leq C \|u_{0}\|_{L^{2}} + C \|G(\cdot)|u|^{2}u\|_{L^{1}([0,M],L^{2})} \\ &\leq C \|u_{0}\|_{L^{2}} + CM^{1/2} \|u\|_{L^{6}([0,M],L^{6})}^{3} . \end{split}$$

$$(2.14)$$

On the other hand, using the unitarity of U(0,t) on $L^2(\mathbb{R})$ and the argument used in (2.14), we obtain

$$\begin{split} \|\Phi(u)\|_{L^{\infty}([0,M],L^{2})} &\leq \|u_{0}\|_{L^{2}} + C\|G(\cdot)|u|^{2}u\|_{L^{1}([0,M],L^{2})} \\ &\leq \|u_{0}\|_{L^{2}} + CM^{1/2}\|u\|_{L^{6}([0,M],L^{6})}^{3}. \end{split}$$

Next, noting

$$||z_1|^2 z_1 - |z_2|^2 z_2| \le C(|z_1|^2 + |z_2|^2)|z_1 - z_2| \quad \text{for all } z_1, z_2 \in \mathbb{C},$$
(2.15)

by the same arguments above, we get

$$\begin{split} \|\Phi(u) - \Phi(v)\|_{L^{6}([0,M],L^{6})} &\leq C \|G(\cdot)(|u|^{2}u - |v|^{2}v)\|_{L^{1}([0,M],L^{2})} \\ &\leq C \int_{0}^{M} (\|u(t)\|_{L^{6}}^{2} + \|v(t)\|_{L^{6}}^{2})\|u(t) - v(t)\|_{L^{6}} dt \qquad (2.16) \\ &\leq C M^{1/2} (\|u\|_{L^{6}([0,M],L^{6})}^{2} + \|v\|_{L^{6}([0,M],L^{6})}^{2})\|u - v\|_{L^{6}([0,M],L^{6})} \end{split}$$

and

$$\begin{split} \|\Phi(u) - \Phi(v)\|_{L^{\infty}([0,M],L^{2})} &\leq C \|G(\cdot)(|u|^{2}u - |v|^{2}v)\|_{L^{1}([0,M],L^{2})} \\ &\leq CM^{1/2} \left(\|u\|_{L^{6}([0,M],L^{6})}^{2} + \|v\|_{L^{6}([0,M],L^{6})}^{2}\right) \|u - v\|_{L^{6}([0,M],L^{6})}. \end{split}$$

Therefore, we have a positive constant C such that for all $u, v \in B_{M,a}$,

$$\|\Phi(u)\|_{L^{\infty}([0,M],L^{2})} + \|\Phi(u)\|_{L^{6}([0,M],L^{6})} \le C \|u_{0}\|_{L^{2}} + CM^{1/2}a^{3}$$

and

$$d(\Phi(u), \Phi(v)) \le CM^{1/2}a^2d(u, v).$$

Now set $a = 2C ||u_0||_{L^2}$ and choose $0 < M \le 1$ satisfying

$$CM^{1/2}(2C||u_0||_{L^2})^2 < \frac{1}{2},$$

then we obtain that Φ is a contraction from $B_{M,2C||u_0||_{L^2}}$ into itself. Thus, Φ has a unique fixed point $u \in B_{M,2C||u_0||_{L^2}}$ and

$$\|u\|_{L^{\infty}([0,M],L^{2})} + \|u\|_{L^{6}([0,M],L^{6})} \le 2C\|u_{0}\|_{L^{2}}.$$
(2.17)

Moreover, the Stricharz estimates guarantee that u is even in $\mathcal{C}([0, M], L^2) \cap L^6([0, M], L^6)$.

To prove the uniqueness in $\mathcal{C}([0, M], L^2) \cap L^6([0, M], L^6)$, it is enough to find a small $\delta > 0$ so that the uniqueness is guaranteed in $\mathcal{C}([0, \delta], L^2) \cap L^6([0, \delta], L^6)$. Suppose that $u_1, u_2 \in \mathcal{C}([0, \delta], L^2) \cap L^6([0, \delta], L^6)$ solve (2.10), for $0 < \delta \leq M$. It follows from (2.16) that

$$\|u_1 - u_2\|_{L^6([0,\delta],L^6)} \le C\delta^{1/2} \left(\|u_1\|_{L^6([0,\delta],L^6)}^2 + \|u_2\|_{L^6([0,\delta],L^6)}^2 \right) \|u_1 - u_2\|_{L^6([0,\delta],L^6)}.$$

Choosing $\delta > 0$ small enough, we obtain

$$||u_1 - u_2||_{L^6([0,\delta],L^6)} \le \frac{1}{2} ||u_1 - u_2||_{L^6([0,\delta],L^6)},$$

which implies that $u_1 = u_2$ on $\mathcal{C}([0, \delta], L^2) \cap L^6([0, \delta], L^6)$.

Let $u \in \mathcal{C}([0, M], L^2) \cap L^6([0, M], L^6)$ be a unique solution of (2.10). We now show that $u \in L^q([0, M], L^p)$ for every admissible pair (p, q). Applying the Strichartz estimates to (2.10) and the Hölder inequality with exponents $\frac{5}{2}$ and $\frac{5}{3}$ in x-integral and t-integral, we get

$$\begin{aligned} \|u\|_{L^{q}([0,M],L^{p})} &\leq C \|u_{0}\|_{L^{2}} + C \|G(\cdot)|u|^{2}u\|_{L^{6/5}([0,M],L^{6/5})} \\ &\leq C \|u_{0}\|_{L^{2}} + C \left(\int_{0}^{M} \|u(t)\|_{L^{2}}^{6/5} \|u(t)\|_{L^{6}}^{12/5} dt\right)^{5/6} \\ &\leq C \|u_{0}\|_{L^{2}} + CM^{1/2} \|u\|_{L^{\infty}([0,M],L^{2})} \|u\|_{L^{6}([0,M],L^{6})}^{2} \\ &\leq C \|u_{0}\|_{L^{2}} + CM^{1/2} \|u_{0}\|_{L^{2}}^{3}, \end{aligned}$$

where (2.17) is used. Moreover, since u satisfies the mass conservation, that is,

$$||u(t)||_{L^2} = ||u_0||_{L^2}$$
 for all $t \in [0, M]$,

one can iterate this argument to obtain a unique solution in $C([0,1], L^2) \cap L^6([0,1], L^6)$ of (2.10). Iterating this process, we obtain a unique global solution and it satisfies (2.13).

Now we give the proof of Theorem 1.1.

Proof of Theorem 1.1. Let $u_0 \in H^1(\mathbb{R})$ be fixed and let us consider positive times only. We first consider the case $d_{av} \neq \pm 1$. Let $u \in \mathcal{C}([0,\infty), L^2) \cap L^6_{loc}([0,\infty), L^6)$ be the solution of (2.10), obtained in Proposition 2.6. On the other hand, Corollary 2.3 gives us the solution $\tilde{u} \in \mathcal{C}([0, T_+), H^1)$ for (2.10), where $T_+ > 0$ is the maximal life time. Moreover, we have $\tilde{u} \in \mathcal{C}([0, T_+), L^2) \cap L^6_{loc}([0, T_+), L^6)$ from (2.12). By the uniqueness of a solution in $\mathcal{C}([0, T_+), L^2) \cap L^6_{loc}([0, T_+), L^6)$, $u = \tilde{u}$ and, therefore, u is also in $\mathcal{C}([0, T_+), H^1)$ and it remains to show $T_+ = \infty$.

Suppose to the contrary that $T_+ < \infty$ and choose $n \in \mathbb{N}_0$ such that $n < T_+ \le n + 1$. It follows from (2.13) that

$$|u||_{L^q([n,T_+],L^p)} < \infty \tag{2.18}$$

for every admissible pair (p, q).

For now, we assume to have

$$\frac{d}{dt} \|\partial_x u(t)\|_{L^2}^2 = -2G(t) \operatorname{Im} \int_{\mathbb{R}} \left(u(t) \overline{\partial_x u(t)} \right)^2 dx \tag{2.19}$$

for all $t \in (n, T_+)$. Then, for every $t \in (n, T_+)$,

$$\frac{d}{dt} \|\partial_x u(t)\|_{L^2}^2 \le 2\|u(t)\|_{L^\infty}^2 \|\partial_x u(t)\|_{L^2}^2$$

and, therefore, by Gronwall's inequality and the Cauchy-Schwarz inequality, we have

$$\begin{aligned} \|\partial_x u(t)\|_{L^2}^2 &\leq \|\partial_x u(n)\|_{L^2}^2 \exp\left(2\int_n^t \|u(t')\|_{L^\infty}^2 dt'\right) \\ &\leq \|\partial_x u(n)\|_{L^2}^2 \exp\left(2\|u\|_{L^4([n,T_+),L^\infty)}^2\right). \end{aligned}$$

Thus, we use (2.18) and the mass conservation to obtain

$$\sup_{n < t < T_{+}} \|u(t)\|_{H^{1}}^{2} < \infty$$
(2.20)

which contradicts the blow-up alternative.

To finish this case, it remains to show (2.19). We use the twisted solution

$$w(t) = (U(n,t))^{-1}u(t).$$

Since u solves (2.10), w solves

$$w(t) = u(n) + i \int_{n}^{t} (U(n, t'))^{-1} G(t') |u(t')|^{2} u(t') dt'$$

and, therefore, w is differentiable on (n, T_+) and

$$\dot{w}(t) = \partial_t w(t) = i(U(n,t))^{-1} G(t) |u(t)|^2 u(t)$$

which is in $H^1(\mathbb{R})$. Using this, one sees that

$$\frac{d}{dt} \|\partial_x u(t)\|_{L^2}^2 = \frac{d}{dt} \|\partial_x w(t)\|_{L^2}^2 = 2\operatorname{Re}\langle\partial_x w(t), \partial_x \dot{w}(t)\rangle$$
$$= -2G(t)\operatorname{Im}\langle\partial_x u(t), \partial_x (|u(t)|^2 u(t))\rangle$$

which yields (2.19).

Next, we assume that $d_{av} = 1$ and solve (2.1) recursively. First, we find a solution in $\mathcal{C}([0,1], H^1)$ of (2.1), i.e.,

$$\begin{cases} i\partial_t u + 2\partial_x^2 u + G(t)|u|^2 u = 0 & \text{for } t \in (0,1), \\ u = u_0 & \text{for } t = 0. \end{cases}$$

It is well-known that there exists a unique solution $u \in \mathcal{C}([0, 1], H^1)$, for example, see [9]. Denote u(1) by u_1 and solve the ordinary differential equation with the initial datum u_1

$$\begin{cases} i\partial_t u + G(t)|u|^2 u = 0 & \text{for } t \in (1,2) \\ u = u_1 & \text{for } t = 1 \end{cases}$$
(2.21)

of which solution is given by

$$u(t) = u_1 \exp\left(i|u_1| \int_1^t G(t')dt'\right).$$
 (2.22)

Here, we used that $|u(t)| = |u_1|$ for all $t \in (1, 2)$. Indeed, multiplying the ordinary equation in (2.21) by \overline{u} and taking the imaginary part of the resulting identity, we obtain $\partial_t |u(t)|^2 = 0$ for all $t \in (1, 2)$. Noting that (2.22) has the limit at t = 2, we continuously extend u to [1, 2]. Repeating this process completes the proof. The case $d_{av} = -1$ is done similarly.

Remark 2.7. To get (2.19), one can use a formal calculation as follows:

$$\frac{d}{dt}\|\partial_x u(t)\|_{L^2}^2 = -2\operatorname{Re}\langle\partial_x u, \partial_x \partial_t u\rangle = 2\operatorname{Im}\left((d_{\mathrm{av}} + d_0(t))\langle\partial_x u, \partial_x \partial_x^2 u\rangle + G(t)\langle\partial_x u, \partial_x |u|^2 u\rangle\right).$$

However, the scalar product $\langle \partial_x u, \partial_x \partial_x^2 u \rangle$ in $L^2(\mathbb{R})$ may not be defined for $u \in H^1(\mathbb{R})$ since $\partial_x u \in L^2(\mathbb{R})$ and $\partial_x \partial_x^2 u = \partial_x^3 u \in H^{-2}(\mathbb{R})$. Thus, as in [11], we used the twisting argument, see, also, [4, 24].

3. Averaging theorem

In this section, to prove the averaging theorem (Theorem 1.2), we compare the solutions of equations (1.3) and (1.4)

$$i\partial_t v + d_{\rm av}\partial_x^2 v + Q_\varepsilon(v) = 0, \qquad (3.1)$$

and

$$i\partial_t v + d_{\rm av}\partial_x^2 v + \langle Q \rangle(v) = 0 \tag{3.2}$$

with close initial data $u_0, v_0 \in H^1(\mathbb{R})$, where the nonlinearities are given by

$$Q_{\varepsilon}(v) := G\left(\frac{t}{\varepsilon}\right) T_{D(t/\varepsilon)}^{-1} \left(|T_{D(t/\varepsilon)}v|^2 T_{D(t/\varepsilon)}v \right)$$

and

$$\langle Q \rangle(v) := \int_0^1 T_r^{-1} \Big(|T_r v|^2 T_r v \Big) \psi(r) dr,$$

respectively. Recall that

$$\psi(r) = e^{-\Gamma} \cosh \Gamma(r-1).$$

We, first, show a lemma which is inspired by the proof of Theorem 4.1 in [28].

Lemma 3.1. For every M > 0, if $v \in \mathcal{C}([-M, M], H^1)$, then

$$\int_{0}^{t} e^{id_{\mathrm{av}}(t-t')\partial_{x}^{2}} Q_{\varepsilon}(v(t'))dt' \to \int_{0}^{t} e^{id_{\mathrm{av}}(t-t')\partial_{x}^{2}} \langle Q \rangle(v(t'))dt'$$
(3.3)

in $\mathcal{C}([-M,M],H^1)$ as $\varepsilon \to 0$.

Proof. We consider positive times only. Let $R_{\varepsilon}(v) := Q_{\varepsilon}(v) - \langle Q \rangle(v)$. Then, by the Plancherel identity, its Fourier transform in x can be expressed by

$$\widehat{R_{\varepsilon}(v)}(\xi,t) = \frac{1}{2\pi} \int_{\mathbb{R}^3} \delta(\xi_1 - \xi_2 + \xi_3 - \xi) A_{\varepsilon}(\xi_1, \xi_2, \xi_3, \xi, t) \hat{v}(\xi_1, t) \overline{\hat{v}(\xi_2, t)} \hat{v}(\xi_3, t) d\xi_1 d\xi_2 d\xi_3$$

for all $\xi \in \mathbb{R}$ and $t \in [0, M]$, where

$$A_{\varepsilon}(\xi_1,\xi_2,\xi_3,\xi,t) := G(t/\varepsilon)e^{-iD(t/\varepsilon)(\xi_1^2 - \xi_2^2 + \xi_3^2 - \xi^2)} - \int_0^1 e^{-ir(\xi_1^2 - \xi_2^2 + \xi_3^2 - \xi^2)}\psi(r)dr.$$

Now, we define $B_{\varepsilon} : \mathbb{R}^4 \times [0, M] \to \mathbb{C}$ by

$$B_{\varepsilon}(\xi_1,\xi_2,\xi_3,\xi,t) := \int_0^t A_{\varepsilon}(\xi_1,\xi_2,\xi_3,\xi,t')dt',$$

then

$$B_{\varepsilon}(\xi_1,\xi_2,\xi_3,\xi,t) = \int_0^t A_1(\xi_1,\xi_2,\xi_3,\xi,t'/\varepsilon)dt' = \varepsilon \int_0^{t/\varepsilon} A_1(\xi_1,\xi_2,\xi_3,\xi,t')dt',$$

and therefore,

$$|B_{\varepsilon}(\xi_1,\xi_2,\xi_3,\xi,t)| \le \varepsilon \int_0^2 |A_1(\xi_1,\xi_2,\xi_3,\xi,t')| dt' \le 4\varepsilon$$

since A_1 is a 2-periodic function in t with mean zero and bounded by 2. Thus,

$$\|B_{\varepsilon}\|_{L^{\infty}(\mathbb{R}^{4}\times[0,M])} \le 4\varepsilon.$$
(3.4)

Using the same argument as in (2.7), for every $v_1, v_2 \in \mathcal{C}([0, M], H^1(\mathbb{R}))$, we see

$$\left\| \int_{0}^{\cdot} e^{id_{\mathrm{av}}(\cdot - t')\partial_{x}^{2}} \left(Q_{\varepsilon}(v_{1})(t') - Q_{\varepsilon}(v_{2})(t') \right) dt' \right\|_{L^{\infty}([0,M],H^{1})} \leq CM \|v_{1} - v_{2}\|_{L^{\infty}([0,M],H^{1})}.$$
(3.5)

The estimate (3.5) also holds when Q_{ε} is replaced by $\langle Q \rangle$. Therefore, by a density argument, it suffices to prove (3.3) for $v \in \mathcal{C}^1([0, M], \mathcal{S}(\mathbb{R}))$ only. Since

$$\frac{\partial}{\partial t}B_{\varepsilon}(\xi_1,\xi_2,\xi_3,\xi,t) = A_{\varepsilon}(\xi_1,\xi_2,\xi_3,\xi,t)$$

for almost every $t \in [0, M]$, by the integration by parts, we obtain

$$\int_0^t e^{id_{\mathrm{av}}(t-t')\xi^2} \widehat{R_{\varepsilon}(v)}(\xi,t') dt' = \widehat{\mathcal{I}_1(v)}(\xi,t) - \int_0^t e^{id_{\mathrm{av}}(t-t')\xi^2} \left(\widehat{\mathcal{I}_2(v)}(\xi,t') - id_{\mathrm{av}}\widehat{\mathcal{I}_3(v)}(\xi,t')\right) dt',$$

where

where

$$\widehat{\mathcal{I}_{1}(v)}(\xi,t) = \int_{\mathbb{R}^{3}} \delta(\xi_{1} - \xi_{2} + \xi_{3} - \xi) B_{\varepsilon}(\xi_{1},\xi_{2},\xi_{3},\xi,t) \hat{v}(\xi_{1},t) \overline{\hat{v}(\xi_{2},t)} \hat{v}(\xi_{3},t) d\xi_{1} d\xi_{2} d\xi_{3},$$

$$\widehat{\mathcal{I}_{2}(v)}(\xi,t) = \int_{\mathbb{R}^{3}} \delta(\xi_{1} - \xi_{2} + \xi_{3} - \xi) B_{\varepsilon}(\xi_{1},\xi_{2},\xi_{3},\xi,t) \partial_{t} \left(\hat{v}(\xi_{1},t) \overline{\hat{v}(\xi_{2},t)} \hat{v}(\xi_{3},t) \right) d\xi_{1} d\xi_{2} d\xi_{3},$$
and

and

$$\widehat{\mathcal{I}_{3}(v)}(\xi,t) = \int_{\mathbb{R}^{3}} \delta(\xi_{1} - \xi_{2} + \xi_{3} - \xi) B_{\varepsilon}(\xi_{1},\xi_{2},\xi_{3},\xi,t) \xi^{2} \hat{v}(\xi_{1},t) \overline{\hat{v}(\xi_{2},t)} \hat{v}(\xi_{3},t) d\xi_{1} d\xi_{2} d\xi_{3}.$$

Then, fix $t \in [0, M]$, we have

$$\left\| \int_{0}^{t} e^{id_{\mathrm{av}}(t-t')\partial_{x}^{2}} R_{\varepsilon}(v)(\cdot,t')dt' \right\|_{H^{1}} = \left(\int_{\mathbb{R}} (1+\xi^{2}) \left| \int_{0}^{t} e^{id_{\mathrm{av}}(t-t')\xi^{2}} \widehat{R_{\varepsilon}(v)}(\xi,t')dt' \right|^{2} d\xi \right)^{1/2} \\ \lesssim \|\mathcal{I}_{1}(v)(\cdot,t)\|_{H^{1}} + \int_{0}^{t} \left(\|\mathcal{I}_{2}(v)(\cdot,t')\|_{H^{1}} + \|\mathcal{I}_{3}(v)(\cdot,t')\|_{H^{1}} \right) dt',$$
(3.6)

where we use Minkowski's inequality. First, to get a bound of $\mathcal{I}_1(v)$, note

$$|\widehat{\mathcal{I}_{1}(v)}(\xi,t)| \leq ||B_{\varepsilon}||_{L^{\infty}(\mathbb{R}^{4} \times [0,M])} \left| \int_{\mathbb{R}^{2}} \hat{v}(\xi_{1},t) \overline{\hat{v}(\xi_{2},t)} \hat{v}(\xi-\xi_{1}+\xi_{2},t) d\xi_{1} d\xi_{2} \right|^{2} d\xi_{1} d\xi_{2}$$

for all ξ and t. If we define $I(f_1, f_2, f_3)$ by its Fourier transform

$$\widehat{I}(f_1, f_2, f_3)(\xi) = \int_{\mathbb{R}^2} \widehat{f}_1(\xi_1) \overline{\widehat{f}_2(\xi_2)} \widehat{f}_3(\xi - \xi_1 + \xi_2) d\xi_1 d\xi_2$$

for all $f_1, f_2, f_3 \in \mathcal{S}(\mathbb{R})$, then by a straightforward calculation, we obtain

$$||I(f_1, f_2, f_3)||_{H^1} \le C ||f_1||_{H^1} ||f_2||_{H^1} ||f_3||_{H^1}.$$

This together with (3.4), we have

$$\|\mathcal{I}_{1}(v)(\cdot,t)\|_{H^{1}} \leq \|B_{\varepsilon}\|_{L^{\infty}(\mathbb{R}^{4}\times[0,M])}\|I(v(t),v(t),v(t))\|_{H^{1}} \lesssim \varepsilon \|v(t)\|_{H^{1}}^{3}$$

for all t. By a similar argument, we have

$$\|\mathcal{I}_{2}(v)(\cdot,t)\|_{H^{1}} \lesssim \varepsilon \|\partial_{t}v(t)\|_{H^{1}} \|v(t)\|_{H^{1}}^{2}$$

and

$$\|\mathcal{I}_{3}(v)(\cdot,t)\|_{H^{1}} \lesssim \varepsilon \Big(\|\partial_{x}^{2}v(t)\|_{H^{1}} \|v(t)\|_{H^{1}}^{2} + \|\partial_{x}v(t)\|_{H^{1}}^{2} \|v(t)\|_{H^{1}} \Big)$$

for all t. Substituting the last three inequalities into (3.6) completes the proof.

Now we are ready to give

Proof of Theorem 1.2. Fix M > 0 and consider positive times only. Let

$$K = 2 \sup_{t \in [0,M]} \|v(t)\|_{H^1}$$

and let $0 < \varepsilon \leq \frac{K}{2}$ for now. Then we have $||u_0||_{H^1} \leq K$ since $||u_0 - v_0||_{H^1} \leq \varepsilon$ and $||v_0||_{H^1} \leq \frac{K}{2}$. Then, it follows from Proposition 2.2 that there exists $M_+ = M_+(K)$, independent of ε , such that $v_{\varepsilon} \in \mathcal{C}([0, M_+], H^1(\mathbb{R}))$ and

$$\sup_{0 < \varepsilon \le \frac{K}{2}} \sup_{t \in [0, M_+]} \| v_{\varepsilon}(t) \|_{H^1} \le 2K.$$
(3.7)

We now prove that (1.5) holds on $[0, M_+]$, i.e., there exists C > 0 such that

 $\|v_{\varepsilon} - v\|_{\mathcal{C}([0,M+],H^1(\mathbb{R}))} \le C\varepsilon.$

By Duhamel's formula, we have

$$v_{\varepsilon}(t) - v(t) = e^{id_{\mathrm{av}}t\partial_x^2}(u_0 - v_0) + i\mathcal{I}_1(t) + i\mathcal{I}_2(t)$$

for all $0 \leq t \leq M_+$, where

$$\mathcal{I}_{1}(t) = \int_{0}^{t} e^{id_{\mathrm{av}}(t-t')\partial_{x}^{2}} \left[Q_{\varepsilon}(v_{\varepsilon})(t') - Q_{\varepsilon}(v)(t') \right] dt'$$

and

$$\mathcal{I}_2(t) = \int_0^t e^{id_{\mathrm{av}}(t-t')\partial_x^2} \left[Q_\varepsilon(v)(t') - \langle Q \rangle(v)(t') \right] dt'.$$

It follows from Lemma 3.1 that there exists a constant C > 0 such that

$$\sup_{t \in [0, M_+]} \| \mathcal{I}_2(t) \|_{H^1} \le C\varepsilon.$$
(3.8)

To bound \mathcal{I}_1 , we use Minkowski's inequality and Lemma 2.1, then we obtain

$$\begin{aligned} \|\mathcal{I}_{1}(t)\|_{H^{1}} &\leq \int_{0}^{t} \|Q_{\varepsilon}(v_{\varepsilon})(t') - Q_{\varepsilon}(v)(t')\|_{H^{1}} dt' \\ &\lesssim \int_{0}^{t} \left(\|v_{\varepsilon}(t')\|_{H^{1}}^{2} + \|v(t')\|_{H^{1}}^{2} \right) \|v_{\varepsilon}(t') - v(t')\|_{H^{1}} dt' \end{aligned}$$
(3.9)

for all $0 \le t \le M_+$. Since $||u_0 - v_0||_{H^1} \le \varepsilon$, it follows from (3.8) and (3.9) that, for all $0 \le t \le M_+$, there exist positive constants C_1 , depending only on K, and C_2 such that

$$\begin{aligned} \|v_{\varepsilon}(t) - v(t)\|_{H^{1}} &\leq \|u_{0} - v_{0}\|_{H^{1}} + \|\mathcal{I}_{1}(t)\|_{H^{1}} + \|\mathcal{I}_{2}(t)\|_{H^{1}} \\ &\leq C_{2}\varepsilon + C_{1} \int_{0}^{t} \|v_{\varepsilon}(t') - v(t')\|_{H^{1}} dt'. \end{aligned}$$

Thus, by Gronwall's inequality, we obtain

$$\sup_{t \in [0, M_+]} \| v_{\varepsilon}(t) - v(t) \|_{H^1} \le C_2 \varepsilon e^{C_1 M_+}.$$
(3.10)

If $M_+ \geq M$, the proof is complete and now we assume that $M_+ < M$. It follows from (3.10) and $v_{\varepsilon} - v \in \mathcal{C}([0, M_+], H^1(\mathbb{R}))$ that

$$\|v_{\varepsilon}(M_{+}) - v(M_{+})\|_{H^{1}} \le C_{2}\varepsilon e^{C_{1}M_{+}}.$$

Now choose $\varepsilon_1 > 0$ such that $C_2 \varepsilon_1 e^{C_1 M_+} \leq \frac{K}{2}$ and $\varepsilon_1 \leq \frac{K}{2}$. Let $0 < \varepsilon \leq \varepsilon_1$. Then

$$||v_{\varepsilon}(M_{+})||_{H^{1}} \le \frac{K}{2} + C_{2}\varepsilon e^{C_{1}M_{+}} \le K.$$

Applying Proposition 2.2 to (2.3) with the initial datum $v_{\varepsilon}(M_{+})$, which satisfies

$$\sup_{0<\varepsilon\leq\varepsilon_1}\|v_\varepsilon(M_+)\|_{H^1}\leq K$$

we have

$$\sup_{0<\varepsilon\leq\varepsilon_1}\sup_{t\in[M_+,2M_+]}\|v_\varepsilon(t)\|_{H^1}\leq 2K$$

Combining this and (3.7), we have

$$\sup_{0<\varepsilon\leq\varepsilon_1}\sup_{t\in[0,2M_+]}\|v_\varepsilon(t)\|_{H^1}\leq 2K.$$

Repeat the above argument to obtain

$$\sup_{t\in[0,2M_+]} \|v_{\varepsilon}(t) - v(t)\|_{H^1} \le C_2 \varepsilon e^{2C_1 M_+}.$$

Iterating this argument, we can choose $\varepsilon_0 > 0$ such that $C_2 \varepsilon_0 \exp\left(\left(\lfloor \frac{M}{M_+} \rfloor + 1\right) C_1 M_+\right) \le \frac{K}{2}$ and $\varepsilon_0 \le \frac{K}{2}$ to get (1.5) with $C = C_2 \exp\left(\left(\lfloor \frac{M}{M_+} \rfloor + 2\right) C_1 M_+\right)$, where $\lfloor a \rfloor = \max\{n \in \mathbb{Z} : n \le a\}$. This completes the proof.

APPENDIX A. LINEAR PROPAGATOR

For the reader's convenience, the properties of the linear propagator $U(t_0, t)$ defined in (2.11) are referred in this section, which can be also found in [3, 5].

Lemma A.1. Let $d_{av} \neq \pm 1$. Then there exists a constant C depending only on d_{av} such that

$$||U(t_0,t)f||_{L^{\infty}} \le C|t-t_0|^{-1/2}||f||_{L^1}$$

for all $f \in L^1(\mathbb{R})$ and all distinct t, t_0 with $\lfloor t \rfloor = \lfloor t_0 \rfloor$.

Proof. Using the kernel of the solution operator for the free Schrödinger equation

$$(T_t f)(x) = \frac{1}{\sqrt{4\pi i t}} \int_{\mathbb{R}} e^{i\frac{|x-y|^2}{4t}} f(y) dy, \quad t \neq 0,$$

14

for Schwartz functions f, we obtain that

$$||U(t_0,t)f||_{L^{\infty}} \le \frac{||f||_{L^1}}{(4\pi |\int_{t_0}^t d(t')dt'|)^{1/2}}, \quad t \ne t_0.$$

Using this and the fact that for all t and t_0 with $\lfloor t \rfloor = \lfloor t_0 \rfloor$,

$$\left| \int_{t_0}^t d(t')dt' \right| = \begin{cases} |d_{\mathrm{av}} + 1||t - t_0|, \quad \lfloor t_0 \rfloor \text{ even} \\ |d_{\mathrm{av}} - 1||t - t_0|, \quad \lfloor t_0 \rfloor \text{ odd}, \end{cases}$$

we complete the proof.

Using Lemma A.1 and the unitarity of $U(t_0, t)$ on $L^2(\mathbb{R})$, via the well-known arguments, we obtain the one-dimensional Strichartz estimates on each time interval [n, n + 1], for every $n \in \mathbb{Z}$. For the classical Strichartz estimates, see, e.g., [25, 16, 21]. As usual, we say that a pair of exponents (p, q) is admissible if $2 \leq p \leq \infty$ and

$$\frac{1}{p} + \frac{2}{q} = \frac{1}{2}$$

and, also, for every $p \ge 1$, denote by p' the Hölder conjugate, i.e.,

$$\frac{1}{p} + \frac{1}{p'} = 1.$$

Lemma A.2. Assume $d_{av} \neq \pm 1$. Let (p,q) and (p_0,q_0) be admissible pairs and $t_0 \in [n, n+1]$ for some $n \in \mathbb{Z}$.

(i) If $f \in L^2(\mathbb{R})$, then the function $t \mapsto U(t_0, t)f$ on [n, n+1] belongs to $L^q([n, n+1], L^p) \cap C([n, n+1], L^2)$. Moreover, there exists a constant C depending only on p and d_{av} such that for all $f \in L^2(\mathbb{R})$

$$||U(t_0,t)f||_{L^q([n,n+1],L^p)} \le C||f||_{L^2}.$$

(ii) Let I be an interval contained in [n, n+1] and $t_0 \in \overline{I}$. If $F \in L^{q'_0}(I, L^{p'_0})$, then the function

$$t\mapsto \int_{t_0}^t U(t',t)F(t')dt'$$

on I belongs to $L^q(I, L^p) \cap C(\overline{I}, L^2)$. Moreover, there exists a constant C depending only on p, p_0 , and d_{av} such that for all $F \in L^{q'_0}(I, L^{p'_0})$

$$\left(\int_{I} \left\| \int_{t_{0}}^{t} U(t',t)F(\cdot,t')dt' \right\|_{L^{p}}^{q} dt \right)^{1/q} \leq C \left(\int_{I} \left\| F(\cdot,t) \right\|_{L^{p_{0}'}}^{q_{0}'} dt \right)^{1/q_{0}'}.$$

Acknowledgements:

Young–Ran Lee and Mi–Ran Choi are supported by the National Research Foundation of Korea(NRF) grants funded by the Korean government(MSIT–2020R1A2C1A01010735 and MOE–2019R111A1A01058151).

References

- M. J. Ablowitz and G. Biondini, Multiscale pulse dynamics in communication systems with strong dispersion management. Opt. Lett. 23 (1998), 1668–1670.
- [2] G. P. Agrawal, Nonlinear Fiber Optics (Fifth Edition). Academic Press, San Diego, (2012).
- [3] J. Albert and E. Kahlil, On the well-posedness of the Cauchy problem for some nonlocal nonlinear Schrödinger equations. Nonlinearity 30 (2017), 2308–2333.

- [4] I. Anapolitanos, M. Hott, and D. Hundertmark, Derivation of the Hartree equation for compound Bose gases in the mean field limit. Rev. Math. Phys. 29 (2017), no. 7, 1750022 (28 pages).
- [5] P. Antonelli, J. C. Saut, and C. Sparber, Well-posedness and averaging of NLS with timeperiodic dispersion management. Adv. Differential Equations 18 (2013), 49–68.
- [6] A. de Bouard and A. Debussche, The nonlinear Schrödinger equation with white noise dispersion. J. Funct. Anal. 259 (2010), no. 5, 1300–1321.
- [7] T. Cazenave, Semilinear Schrödinger Equations. Courant Lecture Notes in Mathematics, vol. 10. American Mathematical Society, Providence (2003).
- [8] T. Cazenave and P. L. Lions Orbital stability if standing waves for some nonlinear Schrödinger Equations. Comm. Math. Phys. 85 (1982), 549–561.
- T. Cazenave and M. Scialom, A Schrödinger equation with time-oscillating nonlinearity. Rev. Mat. Complut. 23 (2010), no. 2, 321–339.
- [10] M.-R. Choi, D. Hundertmark, and Y.-R. Lee, Thresholds for existence of dispersion management solitons for general nonlinearities. SIAM J. Math. Anal. 49 (2017), no. 2, 1519–1569.
- M.-R. Choi, D. Hundertmark, and Y.-R. Lee, Well-posedness of dispersion managed nonlinear Schrödinger equations. arXiv:2003.09076
- [12] A. R. Chraplyvy, A. H. Gnauck, R. W. Tkach, and R. M. Derosier, 8 × 10 Gb/s transmission through 280 km of dispersion-managed fiber. IEEE Photon. Technol. Lett. 5 (1993), 1233–1235.
- [13] M. B. Erdoğan, D. Hundertmark, Y.-R. Lee, Exponential decay of dispersion managed solitons for vanishing average dispersion. Math. Res. Lett. 18(1) (2011), 13–26.
- [14] I. Gabitov and S. K. Turitsyn, Averaged pulse dynamics in a cascaded transmission system with passive dispersion compensation. Opt. Lett. 21 (1996), 327–329.
- [15] I. Gabitov and S. K. Turitsyn, Breathing solitons in optical fiber links. JETP Lett. 63 (1996), 861–866.
- [16] J. Ginibre and G. Velo, The global Cauchy problem for the nonlinear Schrödinger equation revisited. Ann. Inst. H. Poincaré Anal. Non Linéaire 2 (1985), 309–327.
- [17] W. Green and D. Hundertmark, Exponential decay of dispersion managed solitons for general dispersion profiles. Lett. Math. Phys. 106 (2016), no. 2, 221–249.
- [18] D. Hundertmark, P. Kunstmann, and R. Schnaubelt, Stability of dispersion managed solitons for vanishing average dispersion. Arch. Math. 104 (2015), no. 3, 283–288.
- [19] D. Hundertmark and Y.-R. Lee, On non-local variational problems with lack of compactness related to non-linear optics. J. Nonlinear Sci. 22 (2012), 1–38.
- [20] T. Kato, On nonlinear Schrödinger equatuions. Ann. Inst. H. Poincaré Phys. Théor. 46 (1987), 113–129.
- [21] M. Keel and T. Tao, Endpoint Strichartz estimates. Am. J. Math. 120 (1998), 955–980.
- [22] S. Kumar and A. Hasegawa, Quasi-soliton propagation in dispersion-managed optical fibers. Opt. Lett. 22 (1997), 372–374.
- [23] C. Lin, H. Kogelnik, and L. G. Cohen, Optical pulse equalization and low dispersion transmission in singlemode fibers in the 1.3–1.7 μm spectral region. Opt. Lett. 5 (1980), 476–478.
- [24] T. Ozawa, Remarks on proofs of conservation laws for nonlinear Schrödinger equations Calc. Var. Partial Differential Equations 25 (2006), 403–408.
- [25] R. S. Strichartz, Restrictions of Fourier transforms to quadratic surfaces and decay of solutions of wave equations. Duke Math. J. 44 (1977), 705–714.
- [26] S. K. Turitsyn, B. Bale, and M. P. Fedoruk, Dispersion-managed solitons in fibre systems and lasers, Phys. Rep. 521 (2012), no. 4, 135–203.
- [27] S. K. Turitsyn, E. G. Shapiro, S. B. Medvedev, M. P. Fedoruk, and V. K. Mezentsev, *Physics and mathematics of dispersion-managed optical solitons*. C. R. Phys. 4 (2003), 145–161.
- [28] V. Zharnitsky, E. Grenier, C. K. R. T. Jones, and S. K. Turitsyn, Stabilizing effects of dispersion management. Physica D 152-153 (2001), 794–817.

 † Research Institute for Basic Science, Sogang University, 35 Baekbeom–ro (Sinsu–dong), Mapo-gu, Seoul 04107, South Korea.

Email address: rani9030@sogang.ac.kr

[‡] Department of Mathematics, Sogang University, 35 Baekbeom–ro (Sinsu–dong), Mapo–gu, Seoul 04107, South Korea.

Email address: kkyh0409@sogang.ac.kr, younglee@sogang.ac.kr