

# Could higher dimensional rotating black holes be formed by gravitational collapse?

Naresh Dadhich<sup>1,\*</sup> and Sanjar Shaymatov<sup>2,3,4,5,†</sup>

<sup>1</sup>*Inter University Centre for Astronomy & Astrophysics, Post Bag 4, Pune 411007, India*

<sup>2</sup>*Institute for Theoretical Physics and Cosmology,*

*Zhejiang University of Technology, Hangzhou 310023, China*

<sup>3</sup>*Ulugh Beg Astronomical Institute, Astronomicheskaya 33, Tashkent 100052, Uzbekistan*

<sup>4</sup>*National University of Uzbekistan, Tashkent 100174, Uzbekistan*

<sup>5</sup>*Tashkent Institute of Irrigation and Agricultural Mechanization Engineers, Kori Niyoziy 39, Tashkent 100000, Uzbekistan*

Black holes are generally formed by collapse of a dust cloud under its own gravity or accretion of matter onto a gravitating center. For rotating black hole, cloud has to have some rotation which also contributes to gravitational potential producing repulsive component that counters attraction due to mass. In higher dimensions greater than four, non-zero angular momentum particles would always encounter potential barrier which is greater than particle's energy at infinity. Thus arises the critical question how does then a rotating black hole form in higher dimensions?

PACS numbers:

## I. INTRODUCTION

The primary requirement for formation of astrophysical black hole is accretion of matter under its own gravity onto a gravitating centre. In contrast to Newtonian gravity in general relativity rotation also contributes to gravitational potential but repulsively. Attractive contribution due to mass goes as  $1/r^{D-3}$  while repulsive component due to rotation would always go as  $1/r^2$  and higher. Hence in dimensions greater than five<sup>1</sup>, the latter would though override the former yet the overall effective potential for zero angular momentum accretion onto the Myers-Perry rotating black hole [1] would remain less than particle's energy at infinity. Zero angular momentum particles would therefore encounter no potential barrier to fall onto black hole. This however gets critically offset even in five dimension when centrifugal component of non-zero angular momentum accreting matter is included. Thus matter with non-zero angular momentum would always encounter potential barrier, and hence cannot accrete to a center under its own gravity in higher dimensions greater than four. It therefore raises the critical question how do rotating black holes form in higher dimensions?

There does however exist higher dimensional analogue of the four dimensional rotating Kerr black hole in the Myers-Perry solution [1] of Einstein's vacuum equation describing a rotating black hole. This is all very fine, the question is, how does it get formed in astrophysical setting? As we argued above that in higher dimensions dust cloud with rotation cannot collapse under its own gravity. Could it however be possible to impart rota-

tion to already existing non-rotating black hole? So long as black hole overall remains non-rotating, some rotation could be imparted to it by bombardment of particles with angular momentum. The moment rotation becomes significant enough so as to require the Myers-Perry metric for its description, the situation turns to what is alluded above — no further accretion could occur because of potential barrier it would encounter in all higher dimensions greater than four. Thus gravitational collapse cannot occur in all higher dimensions greater than the usual four.

Following [2], the phenomenon of overspinning a near extremal black hole has been pursued quite intensely as evidenced by large number of papers (see some representative examples [3–8]). It turns out that it is in general possible to overspin a black hole under linear test particle accretion<sup>2</sup>. Recently in a breakthrough paper, Sorce and Wald [11] have shown that this result is always overturned when non-linear perturbations are included. Thus weak cosmic censorship conjecture (WCCC) which may be violated at linear order is always restored at non-linear order.

Higher dimensional Myers-Perry black hole can have more than one rotation parameters, given by  $n = [(D - 1)/2]$ . If number of rotations are less than  $n$ ; i.e. one or more parameters are switched off, black hole has only one horizon, hence has no extremal limit and consequently no overspinning [12]. It is a pertinent question to ask how does occurrence of potential barrier in higher dimensions impact on process of overspinning? Clearly particles with suitable parameters that could cause overspinning would find it difficult to reach horizon. It has been shown [13] with an explicit calculation that a six dimensional black hole cannot be overspun even under linear order ac-

\*Electronic address: [nkd@iucaa.in](mailto:nkd@iucaa.in)

†Electronic address: [sanjar@astrin.uz](mailto:sanjar@astrin.uz)

<sup>1</sup> In five dimensions, both the components have the same fall off  $1/r^2$ , but since mass would override rotation parameter for a black hole, overall it would be attractive.

<sup>2</sup> It is however not possible to convert a non-extremal into extremal black hole [9], and nor an extremal into over-extremal state [10] by geodesic accretion. What is envisaged here is that extremality cannot be reached but it could perhaps be jumped over by a discrete non-geodesic but linear order perturbative process.

tion. In all dimensions greater than six, the same situation of potential barrier would prevail, hence a black hole cannot be overspun in  $D \geq 6$  [12]. In five dimension it could however be overspun at linear order which gets overturned when second order perturbations are included [14]. Thus higher dimensions favour WCCC for rotating black holes.

We have also investigated [15] overspinning of pure Lovelock<sup>3</sup> rotating black hole and have shown that it cannot be overspun in dimension  $> 4N + 1$ . For  $N = 1$ , Einstein gravity, it is in  $D > 5$  rotating black hole cannot be overspun.

The principal aim of this letter is to raise the question of formation of rotating black holes in higher dimensions. It is clear that it cannot be formed by gravitational collapse. This is what we wish to demonstrate in the following by studying effective potential for zero and non-zero angular momentum motion of test particles in higher dimensional Myers-Perry rotating black hole metric. We shall consider only one rotation parameter to make the analysis simple and transparent. This is by no means any restriction on generality of consideration because the main argument hinges on occurrence of repulsive potential barrier which would get further strengthened when more than one rotation parameters are involved. We shall first recall the Myers-Perry metric for higher dimensional rotating black hole and its some salient features.

## II. MYERS-PERRY ROTATING BLACK HOLE

The metric describing the higher dimensional Myers-Perry rotating black hole [1] is given by

$$ds^2 = -dt^2 + (r^2 + a_i^2) (d\mu_i^2 + \mu_i^2 d\phi_i^2) + \frac{\mu r^2}{\Pi F} (dt + a_i \mu_i^2 d\phi_i)^2 + \frac{\Pi F}{\Delta} dr^2, \quad (1)$$

where

$$F = 1 - \frac{a_i^2 \mu_i^2}{r^2 + a_i^2}, \quad \Pi = (r^2 + a_1^2) \dots (r^2 + a_n^2), \quad (2)$$

and  $i = 1, 2, \dots, (D-1)/2$ . In odd  $D = 2n+1$  dimensions,  $\Delta = \Pi - \mu r^2$  and  $\Sigma \mu_i^2 = 1$  while for even  $D = 2n+2$ , there is an additional term  $r^2 d\alpha^2$  in the above metric,  $\Delta = \Pi - \mu r$  and  $\Sigma \mu_i^2 + \alpha^2 = 1$  where  $\mu$  and  $a$  are respectively

mass and rotation parameters and  $n = [(D-1)/2]$  is number of rotation parameters a black hole can have.

The black hole horizons are located at positive roots of  $\Delta = 0$ ; i.e.  $(r^2 + a_1^2) \dots (r^2 + a_n^2) - \mu r^2 = 0$  and  $(r^2 + a_1^2) \dots (r^2 + a_n^2) - \mu r = 0$  respectively for odd and even dimensions.

It should however be noted whenever one or more of rotation parameters are switched off,  $\Delta = 0$  has only one positive root indicating occurrence of only one horizon [12]. As argued earlier for our purpose it would suffice to consider only one rotation parameter for black hole. In particular we would examine the case of 5 and 6 dimensions and show how effective potential barrier occurs? Following the standard procedure for geodesic motion in a rotating black hole spacetime, we would write effective potential for zero and non-zero angular momentum motion.

Horizon for five dimension is given by  $r_+ = \sqrt{\mu - a^2}$  while for six dimension it reads as

$$r_+ = (\mu A)^{1/3} - \frac{2^{1/3} a^2 / \mu^{1/3}}{3A^{1/3}}, \quad (3)$$

where  $A = 1 + \sqrt{1 + 4a^6/27\mu^2}$ .

## III. EFFECTIVE POTENTIAL

Effective potential for non-zero angular momentum motion in equatorial plane for a rotating black hole with a single rotation is generically given by

$$V_{eff} = \Omega \mathcal{L} + \sqrt{\Delta/g_{\phi\phi}(\mathcal{L}^2/g_{\phi\phi} + 1)}, \quad (4)$$

where  $\mathcal{L}$  is angular momentum of particle and  $\Omega = -g_{t\phi}/g_{\phi\phi}$  is the frame dragging angular velocity. This would in 5 and 6 dimensions read as

$$V_{eff}^{5D} = \frac{r \left[ r^4 + (r^2 + \mu) a^2 + r^2 \mathcal{L}^2 \right]^{1/2}}{r^4 + (r^2 + \mu) a^2} (r^2 - \mu + a^2)^{1/2} + \frac{a\mu\mathcal{L}}{r^4 + (r^2 + \mu) a^2}, \quad (5)$$

$$V_{eff}^{6D} = \frac{r \left[ r^5 + (r^3 + \mu) a^2 + r^3 \mathcal{L}^2 \right]^{1/2}}{r^5 + (r^3 + \mu) a^2} (r^3 - \mu + a^2 r)^{1/2} + \frac{a\mu\mathcal{L}}{r^5 + (r^3 + \mu) a^2}, \quad (6)$$

and for  $\mathcal{L} = 0$  to

$$(V_{eff}^{5D})^2 = \left( 1 - \frac{\mu}{r^2} + \frac{a^2}{r^2} \right) \left( 1 + \frac{a^2}{r^2} + \frac{\mu a^2}{r^4} \right)^{-1}, \quad (7)$$

$$(V_{eff}^{6D})^2 = \left( 1 - \frac{\mu}{r^3} + \frac{a^2}{r^2} \right) \left( 1 + \frac{a^2}{r^2} + \frac{\mu a^2}{r^5} \right)^{-1}. \quad (8)$$

<sup>3</sup> Pure Lovelock means the Lovelock Lagrangian and the following equation of motion has only one  $N$ th order term without sum over lower orders. Here  $N$  is degree of homogeneous Riemann curvature polynomial. It should however be noted that the metric we employ for rotating black hole is extrapolated [16] from the corresponding Myers-Perry metric but it is not an exact solution of pure Lovelock vacuum equation. It however has all the desirable and expected features.




FIG. 1: Effective potential plots: Upper and lower panels respectively refer to  $\mathcal{L} = 0$  and  $\mathcal{L} = 2$  while left/right panels to  $D = 5/6$ .

Note that for zero angular momentum potential even though repulsive component due to rotation dominates over attractive due to mass in the numerator yet it would not turn overall repulsive because of the denominator. This becomes clear as we expand it for large  $r$ ,  $V_{eff}^{6D} \sim 1 - \mu/2r^3$ . It is interesting that at large  $r$  contribution due to rotation gets completely annulled out. That is why potential in the upper panel in Fig. 1 is similar to that of non rotating black hole.

The situation however gets radically altered for non-zero angular momentum case where additional centrifugal component,  $\mathcal{L}^2/r^2$  comes in (Fig. 1, lower panel). Note that  $V_{eff} = 1$  at infinity, and then it rises as  $r$  decreases and reaches maximum before coming down at horizon. It is interesting that  $V_{eff} \geq 1$  all through except very close to horizon. For a single rotation, there occurs only one horizon and hence there is no upper limit on rotation parameter  $a$  except for five dimension where it has to respect  $a^2 \leq \mu$  lest it turns into a naked singularity<sup>4</sup>. This is an interesting case of a rotating black hole with

one horizon yet having an extremal limit for its rotation parameter. However it cannot be overspun [14] even for linear order accretion. It is also interesting to note that in  $D = 6$ , as  $a \rightarrow \infty$ ,  $V_{eff} \rightarrow 1$  at all  $r$ . This is why initially maximum of curve rises with increasing  $a$  until  $a \gtrsim 1.3$ , then it starts coming down, see (Fig. 1 lower right panel).

#### IV. DISCUSSION

We have seen in the above that particles with angular momentum would always encounter potential barrier exceeding their rest mass energy at infinity and hence they would not be able to reach rotating black hole horizon unless they are somehow energised to a value overriding maximum of potential. Such an energising avenue is not known in realistic astrophysical scenario except of course of merger and collision of black holes.

Also notice that there exists only one maximum and no minimum in the lower panel of Fig. 1 indicating non-existence of bound orbits. This was however known for non rotating black holes in higher dimensions [17]. This feature is also carried over to rotating black holes. This also raises question about accretion process for non rotating black hole. Accretion is mediated through accretion disk which cannot occur in higher dimensions because

<sup>4</sup> This happens only in the special case of five dimension and not in general for  $D = 2n + 1$ , because in this case contribution to potential for both mass and rotation fall as  $1/r^2$ .

there exist no bound orbits. Of course a gas cloud free of angular momentum could collapse under its own gravity — purely radial accretion, to form a non rotating black hole.

It thus appears that in Einstein gravity rotating black holes may not be able to form in higher dimensions greater than usual four by gravitational collapse or accretion process. This was what we had set out to demonstrate.

How about taking the question to generalized theories of gravity? The most natural generalization of general relativity in higher dimensions is the Lovelock theory which is quintessentially higher dimensional. It is only the pure Lovelock theory [18], which has only one  $N$ th order term without sum over lower orders in the Lovelock Lagrangian, that admits bound orbits in the dimension range  $2N + 1 < D < 4N + 1$  [17]. For  $N = 1$ , Einstein gravity bound orbits exist only in four dimension while for  $N = 2$  pure Gauss-Bonnet gravity, they exist for  $D = 6, 7, 8$ . However non rotating black hole [19] is stable [20] only in dimensions  $D \geq 3N + 1$ ; i.e. for pure Gauss-Bonnet in seven and eight dimensions. That is, in these dimensions rotating black holes could,

in principle like Kerr black hole in four dimension, be formed in pure Gauss-Bonnet gravity by collapse and accretion process. In general for the dimension window,  $2N + 1 < D < 4N + 1$ , pure Lovelock rotating black holes could be formed.

Finally higher dimensional rotating black holes cannot be formed in Einstein gravity by collapse and accretion process, they could however be formed only in pure Lovelock gravity, in particular pure Gauss-Bonnet gravity in dimensions  $D = 6, 7, 8$ . Besides kinematicity of gravity in critical odd dimensions  $D = 2N + 1$  and existence of bound orbits [18], formation of rotating black holes in higher dimensions is yet another property that singles out pure Lovelock gravity.

### Acknowledgments

S.S. acknowledges the support of Uzbekistan Ministry for Innovative Development Grants No. VA-FA-F-2-008 and No. MRB-AN- 2019-29.

- 
- [1] R. C. Myers and M. J. Perry, *Ann. Phys. (N.Y.)* **172**, 304 (1986).
- [2] T. Jacobson and T. P. Sotiriou, *Phys. Rev. Lett.* **103**, 141101 (2009), [arXiv:0907.4146 \[gr-qc\]](#) .
- [3] E. Barausse, V. Cardoso, and G. Khanna, *Phys. Rev. Lett.* **105**, 261102 (2010), [arXiv:1008.5159 \[gr-qc\]](#) .
- [4] A. Saa and R. Santarelli, *Phys. Rev. D* **84**, 027501 (2011), [arXiv:1105.3950 \[gr-qc\]](#) .
- [5] Z. Li and C. Bambi, *Phys. Rev. D* **87**, 124022 (2013), [arXiv:1304.6592 \[gr-qc\]](#) .
- [6] J. V. Rocha and R. Santarelli, *Phys. Rev. D* **89**, 064065 (2014), [arXiv:1402.4840 \[gr-qc\]](#) .
- [7] M. Colleoni, L. Barack, A. G. Shah, and M. van de Meent, *Phys. Rev. D* **92**, 084044 (2015), [arXiv:1508.04031 \[gr-qc\]](#) .
- [8] S. Shaymatov, M. Patil, B. Ahmedov, and P. S. Joshi, *Phys. Rev. D* **91**, 064025 (2015), [arXiv:1409.3018 \[gr-qc\]](#) .
- [9] N. Dadhich and K. Narayan, *Phys. Lett. A* **231**, 335 (1997).
- [10] R. Wald, *Ann. Phys. (N.Y.)* **82**, 548 (1974).
- [11] J. Sorce and R. M. Wald, *Phys. Rev. D* **96**, 104014 (2017), [arXiv:1707.05862 \[gr-qc\]](#) .
- [12] S. Shaymatov and N. Dadhich, arXiv e-prints (2020), [arXiv:2004.09242 \[gr-qc\]](#) .
- [13] S. Shaymatov, N. Dadhich, and B. Ahmedov, *Phys. Rev. D* **101**, 044028 (2020), [arXiv:1908.07799 \[gr-qc\]](#) .
- [14] S. Shaymatov, N. Dadhich, and B. Ahmedov, *Eur. Phys. J. C* **79**, 585 (2019), [arXiv:1809.10457 \[gr-qc\]](#) .
- [15] S. Shaymatov and N. Dadhich, arXiv e-prints (2020), [arXiv:2008.04092 \[gr-qc\]](#) .
- [16] N. Dadhich and S. G. Ghosh, arXiv e-prints (2013), [arXiv:1307.6166 \[gr-qc\]](#) .
- [17] N. Dadhich, S. G. Ghosh, and S. Jhingan, *Phys. Rev. D* **88**, 124040 (2013), [arXiv:1308.4770 \[gr-qc\]](#) .
- [18] N. Dadhich, *Eur. Phys. J. C* **76**, 104 (2016), [arXiv:1506.08764 \[gr-qc\]](#) .
- [19] N. Dadhich, J. M. Pons, and K. Prabhu, *Gen. Relativ. Gravit.* **45**, 1131 (2013), [arXiv:1201.4994 \[gr-qc\]](#) .
- [20] R. Gannouji, Y. Rodríguez Baez, and N. Dadhich, *Phys. Rev. D* **100**, 084011 (2019), [arXiv:1907.09503 \[gr-qc\]](#) .