

# On finite groups with all simple modules of low dimension in characteristic $p$

Geoffrey R. Robinson

November 10, 2021

## Abstract

We offer a short and reasonably elementary proof that if  $G$  is a finite group,  $F$  is an algebraically closed field of prime characteristic  $p$ , and all simple  $FG$ -modules have dimension less than  $p$ , then  $G$  has a normal Sylow  $p$ -subgroup. The algebraic closure of  $F$  is not really necessary, but simplifies the exposition in the essential case.

## 1 Introduction

Let  $G$  be a finite group,  $p$  be a prime, and let  $F$  be an algebraically closed field of characteristic  $p$ . We recall that  $O_p(G)$  acts trivially on each simple  $FG$ -module. We will prove that if each simple  $FG$ -module has dimension less than  $p$ , then  $G$  has a normal Sylow  $p$ -subgroup. This may be compared with an analogous result for ordinary irreducible representations proved in [2] by Isaacs and Passman, where the resulting normal Sylow  $p$ -subgroup is necessarily Abelian, which need not be the case in the present modular context.

We will make use of the following reciprocity theorem, which appears in [3] with a proof using properties of the Reynolds ideal of  $Z(FG)$ , but which may also be proved using projective homomorphisms (see, for example, Lemma 5.2 of Grodal [1]): if  $H$  is a subgroup of  $G$  and  $S$  is simple  $FG$ -module,  $T$  is a simple  $FH$ -module, then the multiplicity of the projective cover of  $S$  as a summand of  $\text{Ind}_H^G(T)$  is equal to the multiplicity of the projective cover of  $T$  as a summand of  $\text{Res}_H^G(S)$ .

Now we prove our main:

**Theorem:** *Let  $G$  be a finite group and  $F$  be an algebraically closed field of prime characteristic  $p$ . Suppose that every simple  $FG$ -module has dimension less than  $p$ . Then  $G$  has a normal Sylow  $p$ -subgroup.*

**Proof :** We proceed by induction on  $|G|$ . Since  $O_p(G)$  acts trivially on every simple  $FG$ -module, we may suppose that  $O_p(G) = 1$ , but that  $p$  divides  $|G|$ .

Since the hypotheses imply (on consideration of the composition factors of the respective regular modules) that for any section  $K$  of  $G$ , each simple  $FK$ -module has dimension less than  $p$ , we may suppose by induction that every proper section  $K$  of  $G$  has a normal Sylow  $p$ -subgroup. Let  $P$  be a Sylow  $p$ -subgroup of  $G$ .

We claim that  $P \cap P^g = 1$  for all  $g \in G \setminus N_G(P)$ . It suffices to prove that  $N_G(V) \leq N_G(P)$  whenever  $1 \neq V \leq P$ . Since  $O_p(G) = 1$ , we have  $N_G(V) < G$  for all such  $V$ . Hence, by induction, we know that  $N_G(V)$  has a normal (and hence unique) Sylow  $p$ -subgroup for each such  $V$ . It follows that  $N_G(V)$  is contained in the normalizer of its unique Sylow  $p$ -subgroup, so, using induction on  $[P : V]$  (and the fact that  $N_P(V) > V$  whenever  $V < P$ ), the claim follows.

Now  $N_G(P) < G$  and we have  $\text{Ind}_{N_G(P)}^G(F) = F \oplus M$  where  $M$  is a (non-zero) projective  $FG$ -module, using Mackey decomposition, for example. Let  $Q$  be a projective indecomposable summand of  $M$ , and let  $S$  be the socle of  $Q$  (which is also isomorphic to the head of  $Q$ ). Then the multiplicity of the projective cover of the trivial  $FN_G(P)$ -module  $F$  as a summand of  $\text{Res}_{N_G(P)}^G(S)$  is strictly positive by the above reciprocity theorem (applied with the trivial  $FN_G(P)$ -module  $F$  in place of  $T$ ). Hence  $\dim_F(S) \geq |P|$ , contrary to our assumption that each simple  $FG$ -module has dimension less than  $p$ . This contradiction shows that  $|G|$  is not divisible by  $p$ , and so completes the proof of the Theorem.

## References

- [1] Grodal, J. ; *Higher limits via subgroup complexes*, Annals of Math. (2), **155** (2), 2002, 405-457.
- [2] Isaacs, I.M and Passman, D.S. ; *A characterization of groups in terms of the degree of their characters*, Pacific J. Math., **15**, 3, (1965), 877-903.
- [3] Robinson, G.R.; *On projective summands of induced modules*, Journal of Algebra, **122**, (1989), 106-111.