

Non-asymptotic oracle inequalities for the Lasso in high-dimensional mixture of experts

TrungTin Nguyen

TRUNG-TIN.NGUYEN@INRIA.FR

Univ. Grenoble Alpes, Inria, CNRS, Grenoble INP, LJK, Inria Grenoble Rhone-Alpes, 655 av. de l'Europe, 38335 Montbonnot, France.

Hien D Nguyen

H.NGUYEN7@UQ.EDU.AU

School of Mathematics and Physics, University of Queensland, St. Lucia, Brisbane, Australia.

Faïcel Chamroukhi

FAICEL.CHAMROUKHI@IRT-SYSTEMX.FR

IRT SystemX, Palaiseau, France.

Geoffrey J McLachlan

G.MCLACHLAN@UQ.EDU.AU

School of Mathematics and Physics, University of Queensland, St. Lucia, Brisbane, Australia.

Abstract

Mixture of experts (MoE) has a well-principled finite mixture model construction for prediction, allowing the gating network (mixture weights) to learn from the predictors (explanatory variables) together with the experts' network (mixture component densities). We investigate the estimation properties of MoEs in a high-dimensional setting, where the number of predictors is much larger than the sample size, for which the literature lacks computational and especially theoretical results. We consider the class of finite MoE models with softmax gating functions and Gaussian regression experts, and focus on the theoretical properties of their l_1 -regularized estimation via the Lasso. We provide a lower bound on the regularization parameter of the Lasso penalty that ensures an l_1 -oracle inequality is satisfied by the Lasso estimator according to the Kullback–Leibler loss. We further state an l_1 -ball oracle inequality for the l_1 -penalized maximum likelihood estimator from the model selection.

Keywords: Mixture of experts, mixture of regressions, penalized maximum likelihood, l_1 -oracle inequality, high-dimensional statistics, Lasso.

1. Introduction

1.1. High-dimensional mixture of experts

Mixture of experts (MoE), introduced by [Jacobs et al. \(1991\)](#); [Jordan and Jacobs \(1994\)](#), is a flexible finite mixture model (MM) construction for conditional density estimation and prediction, and is widely used in statistics and machine learning due to its flexibility and wealth of available statistical estimation and model selection tools. The MoE model construction allows the mixture weights (or gating functions) to depend on the explanatory variables (or predictors) together with the experts (or mixture component densities). This permits the modeling of data arising from more complex data generating processes than those that can be analyzed using the classical MM and mixture of regressions (MoR) models, whose mixing parameters are independent of the covariates. Finite mixture type models have also been popularized thanks to their universal approximation properties, which have been extensively studied in [Genovese and Wasserman \(2000\)](#); [Rakhlin et al. \(2005\)](#); [Nguyen \(2013\)](#); [Ho and Nguyen \(2016a,b\)](#); [Nguyen et al. \(2020b, 2022b\)](#). Analogously, the approximation properties of MoEs for conditional density estimation and mean function estimation

have been established in [Jiang and Tanner \(1999\)](#); [Norets \(2010\)](#); [Nguyen et al. \(2016\)](#); [Ho et al. \(2022\)](#); [Nguyen et al. \(2019, 2021a\)](#).

In particular, for high-dimensional, multimodal, heterogeneous, streaming, lifelong and large-scale data, it is worth noting that MoE and its generic, deep neural network-MoE (DNN-MoE), are extremely well suited. In fact, the MoEs is a specific instantiation of conditional computation [Bengio \(2013\)](#), where different model “experts” are responsible for different regions of the input space. Thus, by applying only a subset of parameters to each example, MoE can increase model capacity while keeping training and inference costs roughly constant. For instance, sparse-MoEs (see, *e.g.*, recent review in [Fedus et al. \(2022a\)](#) and its references) have been successfully used in a variety of domains including natural language processing (*e.g.*, scaling neural machine translation models [Shazeer et al. \(2017\)](#); [Lepikhin et al. \(2021\)](#); [Ruiz et al. \(2021\)](#); [Nguyen et al. \(2022a\)](#), GPT-3 [Brown et al. \(2020\)](#); [Du et al. \(2022\)](#)), speech recognition [Yang et al. \(2021\)](#); [You et al. \(2022\)](#), multi-modal learning [Mustafa et al. \(2022\)](#) and computer vision [Puigcerver et al. \(2021\)](#); [Ruiz et al. \(2021\)](#). In particular, when combined with transformers [Vaswani et al. \(2017\)](#); [Fedus et al. \(2022b\)](#), such models allow us to perform training and inference with fewer resources, while at the same time keeping the computational time of such large networks practical. Recently, [Ye and Bors \(2023\)](#) proposed an end-to-end lifelong learning MoEs, where each expert is implemented by a variational autoencoder ([Kingma and Welling, 2014](#), VAE). This mechanism is advantageous both in terms of storage efficiency and computational cost. Only one expert is used in the inference process. In [Kopf et al. \(2021\)](#), the MoE similarity VAE was proposed, which can learn multi-modal distributions of high-dimensional data and use them to generate realistic data with a high degree of effectiveness and efficiency.

However, there is only a small amount of work aimed at understanding the mathematical and statistical properties of such sparse MoEs in the context of DDNs, or even in the simpler context of just MoEs. To my knowledge, [Chen et al. \(2022\)](#) provided the first result towards a formal understanding of the mechanism of the MoE layer for deep learning. More specifically, they showed that the cluster structure of the data plays an important role in the success of the MoE layer. Furthermore, [Zoph et al. \(2022\)](#) only provided practical guidance to partially answer important and difficult questions about the number of experts for each MoE. In the same spirit, [Ye and Bors \(2022\)](#) used the idea of probabilistic dropout to empirically select the appropriate number of VAE components during the training of their mixture VAE. However, while their method provides an efficient end-to-end training mechanism, it has no theoretical underpinning.

This provides the motivation to investigate more mathematical and statistical properties of MoEs, one of the core components of the sparse DNN-MoEs framework, its regularized estimations for sparse solution in heterogeneous and multimodal high-dimensional data. In the context of regression, softmax-gated MoE (SGaME) models, defined as MoEs with Gaussian experts and softmax gating functions, are a standard choice and a powerful tool for modeling more complex nonlinear relationships between response and predictor, arising from different subpopulations, compared to the MoR models. The reader is referred to [Yuksel et al. \(2012\)](#); [Masoudnia and Ebrahimpour \(2014\)](#); [Nguyen and Chamroukhi \(2018\)](#); [Nguyen \(2021\)](#) for recent reviews on this topic. Statistical estimation and model selection MoEs in the high-dimensional regression setting, where the number of explanatory variables can be much larger than the sample size, remain challenging, particularly from a theoretical point of view, where there is still a lack of results. In such situations, we need to reduce the dimension of the problem by looking for the most relevant relationships to avoid numerical problems while ensuring identifiability.

1.2. Related literature

We focus on the use of the Lasso, originally introduced by Tibshirani (1996), also known as an l_1 penalised maximum likelihood estimator (PMLE). Using l_1 -PMLE tends to produce sparse solutions and can be viewed as a convex surrogate for the non-convex l_0 -penalization problem. These methods have attractive computational and theoretical properties (cf. Fan and Li (2001)). First introduced for the linear regression model, the Lasso estimator has since been studied and extended to many statistical problems. The Lasso has been studied in Chen and Derezhinski (2021, for analysing Lasso’s query complexity via robust uniform convergence), Varre and Flammarion (2022, explaining the success of overparameterized neural networks via Lasso), Kifer et al. (2012); Thakurta and Smith (2013, dealing with homogeneous high-dimensional data using linear sparse regression via Lasso). In particular, to deal with heterogeneous high-dimensional data, several researchers have studied the Lasso for variable selection in the context of mixture of regression (MoR) models, see, *e.g.*, Khalili and Chen (2007); Stadler et al. (2010); Meynet (2013); Devijver (2015a); Lloyd-Jones et al. (2018) suggested using mixture of regression (MoR) models with Lasso for variable selection. More precisely, Stadler et al. (2010) provided an l_0 -oracle inequality, satisfied by the Lasso estimators, conditional on the restricted eigenvalue condition, namely the Fisher information matrix is positive definite. Furthermore, they have to introduce some margin conditions so as to link the Kullback–Leibler (KL) loss function to the l_2 -norm of the parameters. Another direction of studying this problem is to look at its l_1 regularisation properties; see *e.g.*, Massart and Meynet (2011); Meynet (2013); Devijver (2015a). As indicated by Devijver (2015a), in contrast to results for the l_0 penalty, some results for the l_1 penalty are valid without any assumptions, either on the Gram matrix or on the bound. However, such results can only produce error bounds that converge at a rate of $1/n$ instead of $1/\sqrt{n}$, where n is the sample size.

In this article, we extend such results for MoR models to the more general SGaME models as defined in (1). Since each mixture weight is modeled by a softmax function of the covariates, the dependence on each feature appears both in the experts and in the mixing proportion functions (gating functions), which allows one to capture more complex nonlinear relationships between the response and predictors arising from different subpopulations, compared to MoR models. This is demonstrated via numerical experiments in several articles such as Chamroukhi and Huynh (2018, 2019); Montuelle et al. (2014); Nguyen and Chamroukhi (2018).

In the context of studying the statistical properties of the PMLE for SGaME, we may consider the prior works of Khalili (2010) and Montuelle et al. (2014). More precisely, by extending the theoretical developments for MoR models in Khalili and Chen (2007), standard asymptotic theorems for SGaME are established in Khalili (2010). More precisely, under several strict regularity conditions on the true joint density function and the choice of tuning parameter, the PMLE using the Scad penalty function from Fan and Li (2001), instead of Lasso, is proved to be both consistent in feature selection and maintains root- n consistency. Therefore, our non-asymptotic result in Theorem 1 can be considered as a complement to such asymptotics for SGaME.

Other related results to our work are the weak oracle inequalities for MoEs from Montuelle et al. (2014); Devijver (2015b, 2017); Nguyen et al. (2022c, 2021b). In particular, regarding SGaME models, Montuelle et al. (2014, Theorem 1) focused on the model selection procedure instead of investigating the l_1 -regularization properties for the Lasso estimator. A detailed comparison between our work and these related results can be found in Section 3.2.

1.3. Main contributions

To the best of our knowledge, we are the first to study the l_1 -regularization properties of the MoE models. Our main contribution, Theorem 1, provides the lower bound on the regularization parameter of the Lasso penalty that ensures non-asymptotic theoretical control of the KL loss of the Lasso estimator for SGaME models. To be more precise, from here on we consider $q, p, n \in \mathbb{N}^*$, $[n] := \{1, \dots, n\}$, where $\mathbb{N}^*$ stands for the positive integers, and $\mathbb{R}^p$ denotes the real number of dimension p . We assume that we have n observations $((x_i, y_i))_{i \in [n]} \in ([0, 1]^p \times \mathbb{R}^q)$, where typically $p \gg n$, coming from the unknown conditional PDF $s_0 \equiv s_{\psi_0} \in S$, which will be formally defined in (5) as the SGaME models with K mixture components. We shall also introduce a KL loss function in (6) between s_0 and a Lasso estimator $\hat{s}^L(\lambda)$ in (7) defined by an l_1 -norm $\|\cdot\|_1$, where $\lambda \geq 0$ is a regularization parameter to be tuned. Then, if $\lambda \geq \kappa \frac{K}{\sqrt{n}} C_{0n}$ for some absolute constant $\kappa \geq 148$, the Lasso estimator $\hat{s}^L(\lambda)$ satisfies the following l_1 -oracle inequality:

$$\mathbb{E} [\text{KL}_n(s_0, \hat{s}^L(\lambda))] \leq (1 + \kappa^{-1}) \inf_{s_\psi \in S} \left(\text{KL}_n(s_0, s_\psi) + \lambda \left\| \psi^{[1,2]} \right\|_1 \right) + \lambda + \sqrt{\frac{K}{n}} C_{1n}.$$

Here, we suppress the dependence of positive deterministic constants C_{0n} and C_{1n} on (p, q, K, n, C_0) and (q, K, n, C_0) , respectively, for ease of interpretation. Here, $C_0 \equiv (A_\gamma, A_\beta, a_\Sigma, A_\Sigma)$ are boundedness positive constants, defining our conditional class S in Section 2.1.2. For the explicit dependences, we refer the reader to (13) and (15). In particular, it should be emphasized that both C_{0n} and C_{1n} depend on the sample size n only via the logarithmic term $\ln n$. Note that this result is non-asymptotic. More specifically, the number of observations n is fixed, while the number of predictors p can grow, with respect to n , and can be much larger than n . Good discussions of non-asymptotic statistics can be found in Massart (2007) and Wainwright (2019).

In this paper, we focus on a simplified but a standard setting in which the means of the experts are linear functions, with respect to explanatory variables. Although simplified, this model captures the core of the MoE regression problem, which concerns the interactions among the different mixture components. We believe that the general techniques that we develop here can be extended to more general experts, such as Gaussian experts with polynomial means (*e.g.*, Mendes and Jiang, 2012) or hierarchical MoE for exponential family regression models, as in Jiang and Tanner (1999). We leave such nontrivial developments for future work.

The rest of this article proceeds as follows. In Section 2, we discuss the construction and framework of SGaME models. In Section 3, we present our main results, an l_1 -oracle inequality satisfied by the Lasso estimators and an l_1 -ball oracle inequality for the l_1 -PMLE in SGaME models. Some conclusions and perspectives of this work are given in Section 3.2. Appendix A is devoted to the proof of these main results. The proof of technical lemmas can be found in Appendix B.

2. Notation and framework

2.1. High-dimensional SGaME models

In high-dimensional regression frameworks, we observe n couples $(x_{[n]}, y_{[n]}) \equiv ((x_i, y_i))_{i \in [n]} \in (\mathcal{X} \times \mathcal{Y})^n \subset (\mathbb{R}^p \times \mathbb{R}^q)^n$, where typically $p \gg n$, x_i is fixed and y_i is a realization of the random variable Y_i , for all $i \in [n]$. We assume that, conditional on $\{x_i\}_{i \in [n]}$, $\{Y_i\}_{i \in [n]}$ are independent and identically distributed (IID) with conditional PDF $s_0(\cdot | x_i)$. Our goal is to estimate s_0 from

the observations using K -component SGaME models (cf., [Jacobs et al. \(1991\)](#); [Jordan and Jacobs \(1994\)](#); [Nguyen \(2021\)](#)) defined as

$$s_\psi(y|x) = \sum_{k=1}^K g_k(x; \gamma) \phi(y; v_k(x), \Sigma_k). \quad (1)$$

Here, the softmax gating function $g_k(x; \gamma)$ is described as

$$g_k(x; \gamma) = \frac{\exp(w_k(x))}{\sum_{l=1}^K \exp(w_l(x))}, w_k(x) = \gamma_{k0} + \gamma_k^\top x, \gamma = (\gamma_{k0}, \gamma_k^\top)_{k \in [K]} \in \Gamma = \mathbb{R}^{(p+1)K}. \quad (2)$$

Next, $\phi(\cdot; v_k, \Sigma_k)$ is the multivariate Gaussian density with mean v_k , which is a function of x that specifies the mean of the k th component, and with covariance matrix Σ_k . In particular, $(v, \Sigma) \equiv ((v_1, \dots, v_K), (\Sigma_1, \dots, \Sigma_K)) \in (\Upsilon \times V)$, where Υ is a set of K -tuples of mean functions from $\mathcal{X}$ to $\mathbb{R}^q$ and V is a sets of K -tuples of symmetric positive definite matrices on $\mathbb{R}^q$. We shall define the parameter vector ψ in the sequel.

2.1.1. FIXED PREDICTORS AND NUMBER OF COMPONENTS WITH LINEAR MEAN FUNCTIONS

The explanatory variables x_i and the number of components $K \in \mathbb{N}^*$ are both fixed. We assume that $\mathcal{X}$ is a compact subset of $\mathbb{R}^p$ and the observations $x_{[n]}$ are finite. Without loss of generality, we choose to rescale x , so that $\|x\|_\infty \leq 1$. Therefore, we can assume that the explanatory variables $x_{[n]} \in \mathcal{X} = [0, 1]^p$.

For simplicity, we consider the case where the means of Gaussian experts are linear functions of the explanatory variables. That is,

$$\Upsilon = \left\{ v : \mathcal{X} \mapsto v_\beta(x) := (\beta_{k0} + \beta_k x)_{k \in [K]} \in (\mathbb{R}^q)^K \mid \beta = (\beta_{k0}, \beta_k)_{k \in [K]} \in \mathcal{B} = \left(\mathbb{R}^{q \times (p+1)} \right)^K \right\},$$

where β_{k0} and β_k are respectively the $q \times 1$ vector of bias and the $q \times p$ regression coefficients matrix for the k th expert. In summary, we wish to estimate s_0 via conditional PDFs belonging to the class:

$$\{(x, y) \mapsto s_\psi(y|x) \mid \psi = (\gamma, \beta, \Sigma) \in \Psi\}, \text{ where } \Psi = \Gamma \times \Xi, \text{ and } \Xi = \mathcal{B} \times V. \quad (3)$$

2.1.2. BOUNDEDNESS ASSUMPTION ON THE SOFTMAX GATING AND GAUSSIAN PARAMETERS

For a matrix A , let $m(A)$ be the modulus of the smallest eigenvalue, and $M(A)$ the modulus of the largest eigenvalue. We assume that there exists deterministic constants $A_\gamma, A_\beta, a_\Sigma, A_\Sigma > 0$, such that $\psi \in \tilde{\Psi}$, where

$$\begin{aligned} \tilde{\Psi} &= \tilde{\Gamma} \times \tilde{\Xi}, \tilde{\Gamma} = \left\{ \gamma \in \Gamma \mid \forall k \in [K], \sup_{x \in \mathcal{X}} (|\gamma_{k0}| + |\gamma_k^\top x|) \leq A_\gamma \right\}, \\ \tilde{\Xi} &= \left\{ \xi \in \Xi \mid \forall k \in [K], \max_{z \in [q]} \sup_{x \in \mathcal{X}} (|[\beta_{k0}]_z| + |[\beta_k x]_z|) \leq A_\beta, a_\Sigma \leq m(\Sigma_k^{-1}) \leq M(\Sigma_k^{-1}) \leq A_\Sigma \right\}. \end{aligned} \quad (4)$$

Finally, we wish to estimate s_0 via the following conditional PDFs

$$S = \left\{ (x, y) \mapsto s_\psi(y|x) \mid \psi = (\gamma, \beta, \Sigma) \in \tilde{\Psi} \right\}. \quad (5)$$

To simplify the proofs, we shall assume that the true density s_0 belongs to S . That is to say, there exists $\psi_0 = (\gamma_0, \beta_0, \Sigma_0) \in \tilde{\Psi}$, such that $s_0 = s_{\psi_0}$. Under the assumption of bounded parameters only, (4), we are able to establish a powerful oracle inequality from Theorem 1.

2.2. Loss functions, Lasso estimators and l_1 -ball PMLEs

In MLE, the KL divergence is the most natural loss function, defined for densities s and t by

$$\text{KL}(s, t) = \begin{cases} \int_{\mathbb{R}^q} \ln \left(\frac{s(y)}{t(y)} \right) s(y) dy, & \text{if } s dy \text{ is absolutely continuous with respect to } t dy, \\ +\infty, & \text{otherwise.} \end{cases}$$

However, to take into account the structure of conditional PDFs, for fixed explanatory variables $(x_i)_{1 \leq i \leq n}$, we consider the following average KL (AKL) loss function:

$$\text{KL}_n(s, t) = \frac{1}{n} \sum_{i=1}^n \text{KL}(s(\cdot|x_i), t(\cdot|x_i)) = \frac{1}{n} \sum_{i=1}^n \int_{\mathbb{R}^q} \ln \left(\frac{s(y|x_i)}{t(y|x_i)} \right) s(y|x_i) dy. \quad (6)$$

Conditioned on $(x_i)_{1 \leq i \leq n}$, the MLE approach suggests estimating s_0 by the conditional PDF s_ψ that minimizes the negative log-likelihood (NLL):

$$-\frac{1}{n} \sum_{i=1}^n \ln(s_\psi(y_i|x_i)).$$

However, since we are dealing with high-dimensional data, we need to regularize the MLE in order to obtain reasonable estimates. Here we first consider the l_1 -PMLE, also known as the Lasso estimator:

$$\hat{s}^L(\lambda) := \underset{s_\psi \in S}{\operatorname{argmin}} \left\{ -\frac{1}{n} \sum_{i=1}^n \ln(s_\psi(y_i|x_i)) + \text{pen}_\lambda(\psi) \right\}, \quad (7)$$

where $\lambda \geq 0$ is a regularization parameter to be tuned, $\psi = (\gamma, \beta, \Sigma)$ and

$$\text{pen}_\lambda(\psi) = \lambda \left\| \psi^{[1,2]} \right\|_1 := \lambda \left(\left\| \psi^{[1]} \right\|_1 + \left\| \psi^{[2]} \right\|_1 \right), \quad (8)$$

$$\left\| \psi^{[1]} \right\|_1 = \|\gamma\|_1 = \sum_{k=1}^K \sum_{j=1}^p |\gamma_{kj}|, \quad \left\| \psi^{[2]} \right\|_1 = \|\text{vec}(\beta)\|_1 = \sum_{k=1}^K \sum_{j=1}^p \sum_{z=1}^q \left| [\beta_k]_{z,j} \right|. \quad (9)$$

Here, $\text{vec}(\cdot)$ is the vectorization operator that stacks the columns of a matrix into a vector. From now on, we denote $\|\beta\|_p$ ($p \in \{1, 2, \infty\}$) by the induced p -norm of a matrix; see Definition 18, which differs from $\|\text{vec}(\beta)\|_p$. Note that $\text{pen}_\lambda(\psi)$ is a Lasso regularization term encouraging sparsity for both the gating and expert parameters. Note that without considering the l_2 -penalty function considered in Khalili (2010), Chamroukhi and Huynh (2018, 2019), their penalty functions belong to our framework and the l_1 -oracle inequality from Theorem 1 can be obtained for it.

Next, in a more general l_1 -ball PMLE, where for some class of model S_m characterized by index $m \in \mathbb{N}^*$, we consider an l_1 -ball

$$S_m = \left\{ s_\psi \in S, \left\| \psi^{[1,2]} \right\|_1 \leq m \right\}. \quad (10)$$

Then let $\hat{s}_m$ be a η_m -log-likelihood estimator (LLE) in S_m , that is, for some $\eta_m \geq 0$, it holds that

$$-\frac{1}{n} \sum_{i=1}^n \ln(\hat{s}_m(y_i|x_i)) \leq \inf_{s_m \in S_m} \left(-\frac{1}{n} \sum_{i=1}^n \ln(s_m(y_i|x_i)) \right) + \eta_m. \quad (11)$$

As is always the case, it is not sufficient to use the LLE of the estimate in each model as a criterion. It underestimates the risk of the estimate and the result is a choice of models that are too complex. In the context of the PMLE, by adding an appropriate penalty $\text{pen}(m)$, one hopes to create a trade-off between good data fit and model complexity. Suppose that for all $m \in \mathbb{N}^*$, the penalty function satisfies $\text{pen}(m) = \lambda m$, where λ will be determined later in (13). Then, for some $\eta \geq 0$, an l_1 -ball PMLE is defined as $\hat{s}_{\hat{m}}$, where $\hat{m}$ satisfies

$$-\frac{1}{n} \sum_{i=1}^n \ln(\hat{s}_{\hat{m}}(y_i|x_i)) + \text{pen}(\hat{m}) \leq \inf_{m \in \mathbb{N}^*} \left(-\frac{1}{n} \sum_{i=1}^n \ln(\hat{s}_m(y_i|x_i)) + \text{pen}(m) \right) + \eta. \quad (12)$$

Note that the error terms η_m and η are necessary to avoid any existence problems, *e.g.*, the infimum may not be reached. Roughly speaking, the Ekeland variational principle states that for any extended-valued lower semicontinuous function which is bounded below, one can add a small perturbation to ensure the existence of the minimum, see *e.g.*, [Borwein and Zhu \(2004\)](#). This framework is also used in [Montuelle et al. \(2014\)](#); [Nguyen et al. \(2022c\)](#).

3. Main results

In this section, we state our main contribution, Theorem 1, an l_1 -oracle inequality for the Lasso estimator for SGaME models, which is proved in Appendix A.2. To simplify the statement of Theorem 1, given some constants $\kappa \geq 148$, first we define the following different condition for λ :

$$\lambda \geq \kappa \frac{K}{\sqrt{n}} C_{0n}, \quad C_{0n} = B_{0n} \left(q \ln n \sqrt{\ln(2p+1)} + 1 \right), \quad (13)$$

$$B_{0n} = \max(A_\Sigma, 1 + K A_G) \left[1 + 2q\sqrt{q} A_\Sigma (5A_\beta^2 + 4A_\Sigma \ln n) \right], \quad (14)$$

$$C_{1n} = \sqrt{2q} A_\gamma \left(\frac{e^{q/2-1} \pi^{q/2}}{A_\Sigma^{q/2}} + H_{s_0} \right) + B_{0n} C_{2n}, \quad (15)$$

$$H_{s_0} = \max \left\{ 0, \ln \left[(4\pi)^{-q/2} A_\Sigma^{q/2} \right] \right\}, \quad C_{2n} = 302qK \left[1 + \left(A_\gamma + qA_\beta + \frac{q\sqrt{q}}{a_\Sigma} \right)^2 \right] \quad (16)$$

$$\lambda \geq \kappa \frac{K}{\sqrt{n}} C_{3n}, \quad C_{3n} = B_n \left(q \ln n \sqrt{\ln(2p+1)} + 1 \right), \quad (17)$$

$$B_n = \max(A_\Sigma, 1 + K A_G) \left[1 + q\sqrt{q} (M_n + A_\beta)^2 A_\Sigma \right], \text{ for some constant } M_n > 0, \quad (18)$$

Note that both C_{0n} and C_{1n} depend on the sample size n only via the logarithm $\ln n$ term. Furthermore, H_{s_0} is related to the negative of the differential entropy of s_0 , see more details via Lemma 6.

Theorem 1 (l_1 -oracle inequality) *Assume that we observe $(x_{[n]}, y_{[n]}) \in ([0, 1]^p \times \mathbb{R}^q)^n$, coming from an unknown conditional PDF $s_0 \equiv s_{\psi_0} \in S$, defined in (5). Given C_{1n} in (15), if λ satisfies*

(13), the Lasso estimator $\hat{s}^L(\lambda)$, defined in (7), satisfies the following l_1 -oracle inequality

$$\mathbb{E} [\text{KL}_n(s_0, \hat{s}^L(\lambda))] \leq (1 + \kappa^{-1}) \inf_{s_\psi \in S} \left(\text{KL}_n(s_0, s_\psi) + \lambda \left\| \psi^{[1,2]} \right\|_1 \right) + \lambda + \sqrt{\frac{K}{n}} C_{1n}. \quad (19)$$

Next, we state an l_1 -ball model selection in Theorem 2, which is proved in Appendix A.3. Note that Theorem 1 is an immediate consequence of Theorem 2.

Theorem 2 (l_1 -ball model selection) Assume that we observe $(x_{[n]}, y_{[n]}) \in ([0, 1]^p \times \mathbb{R}^q)^n$, coming from an unknown conditional PDF $s_0 \equiv s_{\psi_0} \in S$ defined in (5). Given C_{1n} in (15), if λ satisfies (13), the l_1 -ball PMLE $\hat{s}_{\hat{m}}$, defined in (12), satisfies the following l_1 -oracle inequality

$$\mathbb{E} [\text{KL}_n(s_0, \hat{s}_{\hat{m}})] \leq (1 + \kappa^{-1}) \inf_{m \in \mathbb{N}^*} \left(\inf_{s_m \in S_m} \text{KL}_n(s_0, s_m) + \text{pen}(m) + \eta_m \right) + \eta + \sqrt{\frac{K}{n}} C_{1n}. \quad (20)$$

3.1. Sketch of the proof of Theorem 1

At a high level, the main idea here is to study the Lasso estimator, Theorem 1, as a solution of the l_1 -ball PMLE, Theorem 2, from the model selection framework. Then the proof of Theorem 2 can be deduced from Propositions 3 and 4, which deal with the cases for small and large values of Y and are proved in the Appendices A.4 and A.5, respectively.

Given any $M_n > 0$, this can be done by introducing an event $\mathcal{T}$ and a space F_m as follows:

$$\mathcal{T} = \left\{ \max_{i \in [n]} \|Y_i\|_\infty = \max_{i \in [n]} \max_{z \in [q]} |[Y_i]_z| \leq M_n \right\}, \quad (21)$$

$$\mathcal{T}^C = \left\{ \max_{i \in [n]} \|Y_i\|_\infty = \max_{i \in [n]} \max_{z \in [q]} |[Y_i]_z| > M_n \right\}, \quad (22)$$

$$F_m = \left\{ f_m = -\ln \left(\frac{s_m}{s_0} \right) = \ln(s_0) - \ln(s_m), s_m \in S_m \right\}. \quad (23)$$

Conditional on $\{x_i\}_{i \in [n]}$, let $Y'_{[n]}|x_{[n]} \equiv (Y'_i|x_i)_{i \in [n]}$ be IID random samples from Y arising from the conditional PDF $s_0(\cdot|x_i)$, $i \in [n]$. They are independence copies of the sample $Y_{[n]}|x_{[n]}$. By taking into account the definition of AKL loss function from (6), and the conditional expectation property, we obtain

$$\begin{aligned} \text{KL}_n(s_0, \hat{s}_{\hat{m}}) &= \frac{1}{n} \sum_{i=1}^n \int_{\mathbb{R}^q} \ln \left(\frac{s_0(y|x_i)}{\hat{s}_{\hat{m}}(y|x_i)} \right) s_0(y|x_i) dy = \frac{1}{n} \sum_{i=1}^n \int_{\mathbb{R}^q} \hat{f}_{\hat{m}}(y|x_i) s_0(y|x_i) dy \\ &= \frac{1}{n} \sum_{i=1}^n \mathbb{E}_{Y'_i|x_i} [\hat{f}_{\hat{m}}(Y_i|x_i)] = \mathbb{E}_{Y'_{[n]}|x_{[n]}} \left[\frac{1}{n} \sum_{i=1}^n \hat{f}_{\hat{m}}(Y_i|x_i) \right] \\ &= \mathbb{E}_{Y'_{[n]}|x_{[n]}} \left[\frac{1}{n} \sum_{i=1}^n \hat{f}_{\hat{m}}(Y_i|x_i) | \mathcal{T} \right] \mathbb{1}_{\mathcal{T}} + \mathbb{E}_{Y'_{[n]}|x_{[n]}} \left[\frac{1}{n} \sum_{i=1}^n \hat{f}_{\hat{m}}(Y_i|x_i) | \mathcal{T}^C \right] \mathbb{1}_{\mathcal{T}^C} \\ &\equiv (\text{KL}_n(s_0, \hat{s}_{\hat{m}}) | \mathcal{T}) \mathbb{1}_{\mathcal{T}} + (\text{KL}_n(s_0, \hat{s}_{\hat{m}}) | \mathcal{T}^C) \mathbb{1}_{\mathcal{T}^C}. \end{aligned} \quad (24)$$

From now on, when there is no confusion, for the sake of simpler notation, the expectation of (24) can be written as follows

$$\begin{aligned}\mathbb{E} [\text{KL}_n (s_0, \widehat{s}_{\widehat{m}})] &\equiv \mathbb{E}_{Y_{[n]}} [\text{KL}_n (s_0, \widehat{s}_{\widehat{m}})] \\ &= \mathbb{E}_{Y_{[n]}} [(\text{KL}_n (s_0, \widehat{s}_{\widehat{m}}) | \mathcal{T}) \mathbb{1}_{\mathcal{T}}] + \mathbb{E}_{Y_{[n]}} [(\text{KL}_n (s_0, \widehat{s}_{\widehat{m}}) | \mathcal{T}^C) \mathbb{1}_{\mathcal{T}^C}] \\ &\equiv \mathbb{E} [\text{KL}_n (s_0, \widehat{s}_{\widehat{m}}) \mathbb{1}_{\mathcal{T}}] + \mathbb{E} [\text{KL}_n (s_0, \widehat{s}_{\widehat{m}}) \mathbb{1}_{\mathcal{T}^C}].\end{aligned}\quad (25)$$

Therefore, on the basis of the above remark (25), the Theorem 2 is proved by obtaining the upper bound for each of the following terms using Propositions 3 and 4:

$$\mathbb{E} [\text{KL}_n (s_0, \widehat{s}_{\widehat{m}})] = \mathbb{E} [\text{KL}_n (s_0, \widehat{s}_{\widehat{m}}) \mathbb{1}_{\mathcal{T}}] + \mathbb{E} [\text{KL}_n (s_0, \widehat{s}_{\widehat{m}}) \mathbb{1}_{\mathcal{T}^C}]. \quad (26)$$

Proposition 3 (Small values of Y) Assume that we observe $(x_{[n]}, y_{[n]}) \in ([0, 1]^p \times \mathbb{R}^q)^n$, coming from an unknown conditional PDF $s_0 \equiv s_{\psi_0} \in S$ defined in (5). Given C_{2n} and B_n defined as in (16) and (18), respectively, if λ satisfies (17), the l_1 -ball PMLE $\widehat{s}_{\widehat{m}}$, defined in (12), satisfies:

$$\mathbb{E} [\text{KL}_n (s_0, \widehat{s}_{\widehat{m}}) \mathbb{1}_{\mathcal{T}}] \leq (1 + \kappa^{-1}) \inf_{m \in \mathbb{N}^*} \left(\inf_{s_m \in S_m} \text{KL}_n (s_0, s_m) + \text{pen}(m) + \eta_m \right) + \eta + \sqrt{\frac{K}{n}} B_n C_{2n}.$$

Proposition 4 (Large values of Y) Consider $s_0, \mathcal{T}$, and $\widehat{s}_{\widehat{m}}$ as defined in Proposition 3. Then,

$$\mathbb{E} [\text{KL}_n (s_0, \widehat{s}_{\widehat{m}}) \mathbb{1}_{\mathcal{T}^C}] \leq \left(\frac{e^{q/2-1} \pi^{q/2}}{A_{\Sigma}^{q/2}} + H_{s_0} \right) \sqrt{2KnqA_{\gamma}} e^{-\frac{M_n^2 - 2M_n A_{\beta}}{4A_{\Sigma}}},$$

where H_{s_0} is as defined in (16).

Proposition 3 constitutes our important technical contribution. Via Lemma 5, the main idea to prove Proposition 3 is to control the following deviation on the event $\mathcal{T}$:

$$\sup_{f_m \in F_m} |\nu_n (-f_m)| \equiv \sup_{f_m \in F_m} \left| \frac{1}{n} \sum_{i=1}^n \{f_m (Y_i | x_i) - \mathbb{E} [f_m (Y_i | x_i)]\} \right|, \quad (27)$$

$$F_m = \left\{ f_m = -\ln \left(\frac{s_m}{s_0} \right) = \ln(s_0) - \ln(s_m), s_m \in S_m \right\}. \quad (28)$$

Lemma 5 (Control of deviation) For each $m' \in \mathbb{N}^*$, let

$$\Delta_{m'} = m' \sqrt{\ln(2p+1) \ln n} + 2\sqrt{K} \left(A_{\gamma} + qA_{\beta} + \frac{q\sqrt{q}}{a_{\Sigma}} \right). \quad (29)$$

Then, on the event $\mathcal{T}$, for all $m' \in \mathbb{N}^*$, and for all $t > 0$, with probability greater than $1 - e^{-t}$,

$$\sup_{f_{m'} \in F_{m'}} |\nu_n (-f_{m'})| \leq \frac{4KB_n}{\sqrt{n}} \left[37q\Delta_{m'} + \sqrt{2} \left(A_{\gamma} + qA_{\beta} + \frac{q\sqrt{q}}{a_{\Sigma}} \right) \sqrt{t} \right]. \quad (30)$$

Its proof appears in Appendix B.1 and follows the arguments developed in the proof of a more general model selection theorem for MLEs of Massart (2007, Theorem 7.11). More precisely, the proof of Proposition 3 is in the spirit of Vapnik's method of structural risk minimization, first

established in [Vapnik \(1982\)](#) and briefly summarized in Section 8.2 of [Massart \(2007\)](#). In particular, we will use concentration inequalities combined with symmetrization arguments to obtain an upper bound on the empirical process in expectation from (27). More specifically, our technique combines Vapnik’s structural risk minimization paradigm (*e.g.*, [Vapnik, 1982](#)) and model selection theory for conditional density estimation (*e.g.*, [Cohen and Le Pennec, 2011](#)), which is an extended version of the density estimation results of [Massart \(2007\)](#).

3.2. Discussion of our contributions

Theorem 1 provides information on the performance of the Lasso as the l_1 PMLEs for SGaME models. If the regularization parameter λ is properly chosen, the Lasso estimator, which is the solution to the l_1 penalized empirical risk minimization problem, behaves in a manner comparable to the deterministic Lasso (or so-called *oracle*). This oracle is the solution to the l_1 penalized true risk minimization problem, up to an error term of order λ . Note that the best model is defined by

$$\inf_{s_\psi \in S} \left(\text{KL}_n(s_0, s_\psi) + \lambda \left\| \psi^{[1,2]} \right\|_1 \right) \quad (31)$$

is the one with the smallest l_1 penalized true risk. However, since we do not have access to the true density s_0 , we cannot select this best model, which we call the *oracle*. The oracle is by definition the model in the collection that minimizes the l_1 -penalized risk in (31), which is generally assumed to be unknown.

Our result is non-asymptotic: the number n of observations is fixed, while the number of co-variables p can grow with respect to n , and in fact can be much larger than n . Note that, as in [Khalili \(2010\)](#), the true number of mixtures K of the SGaME model is assumed to be known. From a pragmatic perspective, it can be estimated using the AIC of [Akaike \(1974\)](#), the BIC of [Schwarz \(1978\)](#), or the slope heuristic of [Birgé and Massart \(2007\)](#).

Much attention has been paid to obtaining a lower bound of λ in (13), for the oracle inequality, with optimal dependence on p and q , which are the only parameters not to be fixed and which can increase the possibility that $p \gg n$. In fact, the condition that $\kappa \geq 148$ is implied by the Theorem 2. for some absolute constants $\kappa \geq 148$. Such conditions are obtained from the proof of Theorem 2, which requires the results of Proposition 3. In the proof of Proposition 3, the original condition for κ is as follows: let $\kappa \geq 1$ and assume that $\text{pen}(m) = \lambda m$, for all $m \in \mathbb{N}^*$ with

$$\lambda \geq \kappa \frac{4KB_n}{\sqrt{n}} \left(37q \ln n \sqrt{\ln(2p+1)} + 1 \right).$$

Then, to simplify the constant on the lower bound for λ , we replaced $4 \times 37\kappa, \kappa \geq 1$, by $\kappa \equiv \kappa' = 4 \times 37\kappa \geq 148$.

Note that we recover the same dependence of the form $\sqrt{\ln(2p+1)}$ as for the homogeneous linear regression in [Stadler et al. \(2010\)](#) and of the form $\sqrt{\ln(2p+1)} (\ln n)^2 / \sqrt{n}$ for the MoR models in [Meynet \(2013\)](#). On the contrary, the dependence on q for the mixture of multivariate Gaussian regression models in [Devijver \(2015a\)](#) was of the form $q^2 + q$, while here we get the form $q^2 \sqrt{q}$. The main reason is that we need to control the larger class S of the SGaME model, and we use a different technique to evaluate the upper bound of the uniform norm of the gradient for each element in S . Furthermore, the dependence on n for the homogeneous linear regression in [Stadler et al. \(2010\)](#) was of the order of $1/\sqrt{n}$, while here we have an additional $(\ln n)^2$ factor. In

fact, the same situation can be found in the l_1 -oracle inequalities of [Meynet \(2013\)](#), and [Devijver \(2015a\)](#). As explained in [Meynet \(2013\)](#), the use of nonlinear KL information leads to a scenario where the linearity arguments developed in [Stadler et al. \(2010\)](#) with the quadratic loss function cannot be exploited. Instead, we need to use the entropy arguments to handle our model, which leads to an additional $(\ln n)^2$ factor. Motivated by the frameworks of [Meynet \(2013\)](#), [Devijver \(2015a\)](#), we have taken care to give an explicit dependence not only on n , p and q , but also on the number of mixture components K , as well as on the regressors and A_β, A_Σ, A_G —all the quantities that constrain the parameters of the model. However, we should be aware that these dependencies may not be optimal. In our lower bound we get the factor K^2 dependence instead of K as in [Meynet \(2013\)](#), [Devijver \(2015a\)](#). This can be explained by the fact that we used a different technique to handle the more complex model when dealing with the upper bound on the uniform norm of the gradient of $\ln s_\psi$, for $s_\psi \in S$, in Lemma 14. We refer to [Meynet \(2013, Remark 5.8\)](#) for some data sets where the dependence on K can be reduced to the order of $\sqrt{K}$ for the mixture Gaussian regression models. The determination of optimal rates for such problems is still open.

Finally, it is worth noting that the constant $1 + \kappa^{-1}$ from the upper bound in Theorem 2 and C_1 from [Montuelle et al. \(2014, Theorem 1\)](#) cannot be taken to be equal to 1. This fact follows if s_0 does not belong to the approximation class, i.e. if the model is misspecified. This problem also occurred in the l_1 -oracle inequalities of [Meynet \(2013\)](#); [Devijver \(2015a\)](#). Deriving an oracle inequality such that $1 + \kappa^{-1}$ can be replaced by 1 for the KL loss is still an open problem. However, one way to deal with this difficulty is to use the universal approximation of the class SGaME models (see, e.g., [Nguyen et al. \(2021a, 2019, 2020b,a\)](#)). In fact, if we take a large enough number of mixture components, K , we could well approximate a wide class of densities, then the term on the first right hand side, $(1 + \kappa^{-1}) \inf_{s_\psi \in S} (\text{KL}_n(s_0, s_\psi) + \lambda \|\psi^{[1,2]}\|_1)$, where S depends on K , is small for well chosen K .

3.3. Comparison of our contributions

Note that Theorem 1 is also complementary to [Montuelle et al. \(2014, Theorem 1\)](#), where [Montuelle et al. \(2014\)](#) also considered SGaME models for low-dimensional data. Note that they focused on model selection and obtained a *weak oracle inequality* for the PMLE, while we aimed to study the l_1 regularization properties of the Lasso estimators. However, we can compare their procedure with Theorem 2. The main reason why their result is considered a *weak oracle inequality* is that we can see that Theorem 1 of [Montuelle et al. \(2014\)](#) uses a difference divergence on the left (the $\text{JKL}_\rho^{\otimes n}$, tensorised Jensen KL divergence) to that on the right (the $\text{KL}^{\otimes n}$, tensorised KL divergence). However, under a strong assumption, the two divergences are *equivalent* for the conditional PDFs considered. This strong assumption is, however, satisfied if we assume that $\mathcal{X}$ is compact, as in the case of $\mathcal{X} = [0, 1]^p$ in Theorem 2, s_0 is compactly supported, the regression functions are uniformly bounded, and there is a uniform lower bound on the eigenvalues of the covariance matrices.

To illustrate the strictness of the compactness assumption for s_0 , we need only consider s_0 as a univariate Gaussian PDF, which obviously does not satisfy such a hypothesis. Therefore, in such a case, Theorem 1 in [Montuelle et al. \(2014\)](#) is actually weaker than Theorem 2 with respect to the compact support assumption on the true conditional PDF s_0 . On the contrary, the only assumption used to establish Theorem 2 is the boundedness of the parameters of the mixtures, which is also assumed in [Montuelle et al. \(2014, Theorem 1\)](#).

Furthermore, these boundedness assumptions also appeared in [Stadler et al. \(2010\)](#); [Meynet \(2013\)](#); [Devijver \(2015a\)](#). They are quite natural when working with MLE ([Baudry, 2009](#); [Maugis and Michel, 2011](#)), at least when considering the problem of the unboundedness of the likelihood at the boundary of the parameter space ([Redner and Walker, 1984](#); [McLachlan and Peel, 2000](#)), and to prevent the likelihood from diverging. However, by using the smaller divergence $\text{JKL}_\rho^{\otimes n}$ (or stricter assumptions on s_0 and s_m , so that the same divergence $\text{KL}^{\otimes n}$ appears on both sides of the oracle inequality in Theorem 2), [Montuelle et al. \(2014, Theorem 1\)](#) obtained the faster convergence rate of order $1/n$ of the error upper bound, while in Theorem 2 we only seek a convergence rate of order $1/\sqrt{n}$. Therefore, in cases where there are no guarantees on the strict conditions such as the compactness of the support of s_0 and the uniform boundedness of the regression functions, Theorem 2 provides a theoretical basis for the Lasso estimators with an order of convergence of $1/\sqrt{n}$ with only a boundedness assumption on the parameter space.

4. Conclusions and perspectives

We have studied an l_1 PMLEs for SGaME models. For the first time in the literature, we have established the l_1 -oracle inequality, which provides the lower bound for the regularization of the Lasso that ensures non-asymptotic theoretical control of the KL loss of the estimator. Here we assume for simplicity that the regressors belong to $\mathcal{X} = [0, 1]^p$. However, the arguments in our proof are valid for covariates of any scale. To the best of our knowledge, Theorem 2.8 of [Maugis-Rabusseau and Michel \(2013\)](#) is the only result in the literature that investigates the minimax estimator for Gaussian MM models. We believe that establishing such an extension for SGaME models is not trivial. We hope to take up such a challenge in the future.

We also note that our theorem ensures that there exists a sufficiently large λ for which the estimate has good properties, but does not give an explicit value for λ . However, in the Theorem 1 we at least give the lower bound on the value of λ via the bound $\lambda \geq \kappa C(p, q, n, K)$, where $\kappa \geq 148$, although this value is obviously too pessimistic. Possible solutions for calibrating the penalties from the data are the AIC and BIC approaches, motivated by asymptotic arguments. Another method is the slope heuristic, first introduced by [Birgé and Massart \(2007\)](#) and further discussed in [Baudry et al. \(2012\)](#), which is a non-asymptotic criterion for selecting a model from a collection of models. However, such a strategy will not be considered further here, as the implementations would lead to an over-extension of the length and scope of the manuscript. Furthermore, the technical developments required for such methods are non-trivial and represent an important line of research that we hope to pursue in the future. Nevertheless, for a first look at joint rank and variable selection in high-dimensional SGaME models via Lasso using the slope heuristic based non-asymptotic approach, we refer to the work of [Nguyen \(2021, Chapter 4\)](#). For numerical simulations, the reader is referred to the work of [Montuelle et al. \(2014, using the slope heuristic for low-dimensional data\)](#) and [Chamroukhi and Huynh \(2018, 2019, using BIC\)](#) for the practical implementation of penalisation methods in the context of SGaME models.

Apart from some remaining questions regarding the tightness of the bounds and the form of the penalization functions, we believe that our contribution contributes to the further popularization of SGaME models by providing a theoretical basis for their application to high-dimensional problems.

Acknowledgments

TTN is supported by a “Contrat doctoral” from the French Ministry of Higher Education and Research and FC is granted by the French National Research Agency (ANR) grant SMILES ANR-18-CE40-0014. HDN and GJM are funded by Australian Research Council grant number DP180101192.

References

H Akaike. A new look at the statistical model identification. *IEEE Transactions on Automatic Control*, 19(6):716–723, 1974. doi: 10.1109/TAC.1974.1100705.

Jean-Patrick Baudry. *Sélection de modèle pour la classification non supervisée. Choix du nombre de classes*. PhD thesis, Université Paris-Sud XI, 2009.

Jean-Patrick Baudry, Cathy Maugis, and Bertrand Michel. Slope heuristics: overview and implementation. *Statistics and Computing*, 22(2):455–470, 2012. ISSN 1573-1375. doi: 10.1007/s11222-011-9236-1. URL <https://doi.org/10.1007/s11222-011-9236-1>.

Yoshua Bengio. Deep Learning of Representations: Looking Forward. In Adrian-Horia Dediu, Carlos Martín-Vide, Ruslan Mitkov, and Bianca Truthe, editors, *Statistical Language and Speech Processing*, pages 1–37, Berlin, Heidelberg, 2013. Springer Berlin Heidelberg. ISBN 978-3-642-39593-2.

Lucien Birgé and Pascal Massart. Minimal penalties for Gaussian model selection. *Probability Theory and Related Fields*, 138(1):33–73, 2007. doi: 10.1007/s00440-006-0011-8. URL <https://doi.org/10.1007/s00440-006-0011-8>.

Jonathan M Borwein and Qiji J Zhu. *Techniques of Variational Analysis*. Springer, 2004.

Tom Brown, Benjamin Mann, Nick Ryder, Melanie Subbiah, Jared D Kaplan, Prafulla Dhariwal, Arvind Neelakantan, Pranav Shyam, Girish Sastry, Amanda Askell, Sandhini Agarwal, Ariel Herbert-Voss, Gretchen Krueger, Tom Henighan, Rewon Child, Aditya Ramesh, Daniel Ziegler, Jeffrey Wu, Clemens Winter, Chris Hesse, Mark Chen, Eric Sigler, Mateusz Litwin, Scott Gray, Benjamin Chess, Jack Clark, Christopher Berner, Sam McCandlish, Alec Radford, Ilya Sutskever, and Dario Amodei. Language Models are Few-Shot Learners. In H. Larochelle, M. Ranzato, R. Hadsell, M. F. Balcan, and H. Lin, editors, *Advances in Neural Information Processing Systems*, volume 33, pages 1877–1901. Curran Associates, Inc., 2020. URL <https://proceedings.neurips.cc/paper/2020/file/1457c0d6bfc4967418bfb8ac142f64>

Faïcel Chamroukhi and Bao Tuyen Huynh. Regularized maximum-likelihood estimation of mixture-of-experts for regression and clustering. In *2018 International Joint Conference on Neural Networks (IJCNN)*, pages 1–8, 2018.

Faïcel Chamroukhi and Bao Tuyen Huynh. Regularized maximum likelihood estimation and feature selection in mixtures-of-experts models. *Journal de la Société Française de Statistique*, 160(1): 57–85, 2019.

- Xue Chen and Michal Dereziński. Query complexity of least absolute deviation regression via robust uniform convergence. In Mikhail Belkin and Samory Kpotufe, editors, *Proceedings of Thirty Fourth Conference on Learning Theory*, volume 134 of *Proceedings of Machine Learning Research*, pages 1144–1179. PMLR, August 2021. URL <https://proceedings.mlr.press/v134/chen21d.html>.
- Zixiang Chen, Yihe Deng, Yue Wu, Quanquan Gu, and Yuanzhi Li. Towards understanding mixture of experts in deep learning. *arXiv preprint arXiv:2208.02813*, 2022.
- S X Cohen and Erwan Le Pennec. Conditional density estimation by penalized likelihood model selection and applications. *Technical report, INRIA*, 2011.
- Thomas M Cover. *Elements of information theory*. John Wiley & Sons, 1999.
- E Devijver. An l_1 -oracle inequality for the Lasso in finite mixture of multivariate Gaussian regression. *ESAIM: Probability and Statistics*, 19:649–670, 2015a.
- Emilie Devijver. Finite mixture regression: a sparse variable selection by model selection for clustering. *Electronic journal of statistics*, 9(2):2642–2674, 2015b.
- Emilie Devijver. Joint rank and variable selection for parsimonious estimation in a high-dimensional finite mixture regression model. *Journal of Multivariate Analysis*, 157:1–13, 2017.
- Nan Du, Yanping Huang, Andrew M Dai, Simon Tong, Dmitry Lepikhin, Yuanzhong Xu, Maxim Krikun, Yanqi Zhou, Adams Wei Yu, Orhan Firat, Barret Zoph, Liam Fedus, Maarten P Bosma, Zongwei Zhou, Tao Wang, Emma Wang, Kellie Webster, Marie Pellat, Kevin Robinson, Kathleen Meier-Hellstern, Toju Duke, Lucas Dixon, Kun Zhang, Quoc Le, Yonghui Wu, Zhifeng Chen, and Claire Cui. GLaM: Efficient Scaling of Language Models with Mixture-of-Experts. In Kamalika Chaudhuri, Stefanie Jegelka, Le Song, Csaba Szepesvari, Gang Niu, and Sivan Sabato, editors, *Proceedings of the 39th International Conference on Machine Learning*, volume 162 of *Proceedings of Machine Learning Research*, pages 5547–5569. PMLR, July 2022. URL <https://proceedings.mlr.press/v162/du22c.html>.
- Johannes Jisse Duistermaat and Johan AC Kolk. *Multidimensional real analysis I: differentiation*, volume 86. Cambridge University Press, 2004.
- Jianqing Fan and Runze Li. Variable selection via nonconcave penalized likelihood and its oracle properties. *Journal of the American statistical Association*, 96(456):1348–1360, 2001.
- William Fedus, Jeff Dean, and Barret Zoph. A review of sparse expert models in deep learning. *arXiv preprint arXiv:2209.01667*, 2022a.
- William Fedus, Barret Zoph, and Noam Shazeer. Switch Transformers: Scaling to Trillion Parameter Models with Simple and Efficient Sparsity. *Journal of Machine Learning Research*, 23(120): 1–39, 2022b. URL <http://jmlr.org/papers/v23/21-0998.html>.
- Christopher R Genovese and Larry Wasserman. Rates of convergence for the Gaussian mixture sieve. *The Annals of Statistics*, 28(4):1105–1127, aug 2000. doi: 10.1214/aos/1015956709. URL <https://doi.org/10.1214/aos/1015956709>.

- Gene H Golub and Charles F Van Loan. *Matrix computations*, volume 3. JHU press, 2012.
- Nhat Ho and XuanLong Nguyen. Convergence rates of parameter estimation for some weakly identifiable finite mixtures. *The Annals of Statistics*, 44(6):2726 – 2755, 2016a. doi: 10.1214/16-AOS1444. URL <https://doi.org/10.1214/16-AOS1444>. Publisher: Institute of Mathematical Statistics and Bernoulli Society.
- Nhat Ho and XuanLong Nguyen. On strong identifiability and convergence rates of parameter estimation in finite mixtures. *Electronic Journal of Statistics*, 10(1):271–307, 2016b. Publisher: The Institute of Mathematical Statistics and the Bernoulli Society.
- Nhat Ho, Chiao-Yu Yang, and Michael I Jordan. Convergence Rates for Gaussian Mixtures of Experts. *Journal of Machine Learning Research*, 2022.
- Roger A Horn and Charles R Johnson. *Matrix analysis*. Cambridge University Press, 2012.
- Robert A Jacobs, Michael I Jordan, Steven J Nowlan, and Geoffrey E Hinton. Adaptive mixtures of local experts. *Neural computation*, 3(1):79–87, 1991.
- J L W V Jensen. Sur les fonctions convexes et les inégalités entre les valeurs moyennes. *Acta Mathematica*, 30(1):175–193, 1906. ISSN 1871-2509. doi: 10.1007/BF02418571. URL <https://doi.org/10.1007/BF02418571>.
- Wenxin Jiang and Martin A Tanner. Hierarchical mixtures-of-experts for exponential family regression models: approximation and maximum likelihood estimation. *Annals of Statistics*, pages 987–1011, 1999.
- Michael I Jordan and Robert A Jacobs. Hierarchical mixtures of experts and the EM algorithm. *Neural computation*, 6(2):181–214, 1994.
- Abbas Khalili. New estimation and feature selection methods in mixture-of-experts models. *Canadian Journal of Statistics*, 38(4):519–539, 2010.
- Abbas Khalili and Jiahua Chen. Variable selection in finite mixture of regression models. *Journal of the american Statistical association*, 102(479):1025–1038, 2007.
- Daniel Kifer, Adam Smith, and Abhradeep Thakurta. Private Convex Empirical Risk Minimization and High-dimensional Regression. In Shie Mannor, Nathan Srebro, and Robert C. Williamson, editors, *Proceedings of the 25th Annual Conference on Learning Theory*, volume 23 of *Proceedings of Machine Learning Research*, pages 25.1–25.40, Edinburgh, Scotland, June 2012. PMLR. URL <https://proceedings.mlr.press/v23/kifer12.html>.
- Diederik P. Kingma and Max Welling. Auto-Encoding Variational Bayes. In Yoshua Bengio and Yann LeCun, editors, *2nd International Conference on Learning Representations, ICLR 2014, Banff, AB, Canada, April 14-16, 2014, Conference Track Proceedings*, 2014. URL <http://arxiv.org/abs/1312.6114>.
- Andreas Kopf, Vincent Fortuin, Vignesh Ram Somnath, and Manfred Claassen. Mixture-of-Experts Variational Autoencoder for clustering and generating from similarity-based representations on single cell data. *PLOS Computational Biology*, 17(6):1–17, June 2021. doi: 10.1371/journal.pcbi.

1009086. URL <https://doi.org/10.1371/journal.pcbi.1009086>. Publisher: Public Library of Science.
- Dmitry Lepikhin, HyoukJoong Lee, Yuanzhong Xu, Dehao Chen, Orhan Firat, Yanping Huang, Maxim Krikun, Noam Shazeer, and Zhifeng Chen. {GS}hard: Scaling Giant Models with Conditional Computation and Automatic Sharding. In *International Conference on Learning Representations*, 2021. URL <https://openreview.net/forum?id=qrwe7XHTmYb>.
- Luke R Lloyd-Jones, Hien D Nguyen, and Geoffrey J McLachlan. A globally convergent algorithm for lasso-penalized mixture of linear regression models. *Computational Statistics & Data Analysis*, 119:19–38, 2018.
- Jan R Magnus and Heinz Neudecker. *Matrix differential calculus with applications in statistics and econometrics*. John Wiley & Sons, 2019.
- Masud Mansuripur. *Introduction to information theory*. Prentice-Hall, Inc., 1987.
- Saeed Masoudnia and Reza Ebrahimpour. Mixture of experts: a literature survey. *Artificial Intelligence Review*, 42(2):275–293, 2014. ISSN 1573-7462. doi: 10.1007/s10462-012-9338-y. URL <https://doi.org/10.1007/s10462-012-9338-y>.
- Pascal Massart. *Concentration Inequalities and Model Selection: Ecole d’Eté de Probabilités de Saint-Flour XXXIII-2003*. Springer, 2007.
- Pascal Massart and C Meynet. The Lasso as an l_1 -ball model selection procedure. *Electronic Journal of Statistics*, 5:669–687, 2011.
- Cathy Maugis and Bertrand Michel. A non asymptotic penalized criterion for gaussian mixture model selection. *ESAIM: Probability and Statistics*, 15:41–68, 2011.
- Cathy Maugis-Rabusseau and Bertrand Michel. Adaptive density estimation for clustering with Gaussian mixtures. *ESAIM: Probability and Statistics*, 17:698–724, 2013.
- G J McLachlan and D Peel. *Finite Mixture Models*. John Wiley & Sons, 2000.
- Eduardo F Mendes and Wenxin Jiang. On convergence rates of mixtures of polynomial experts. *Neural computation*, 24(11):3025–3051, 2012.
- C Meynet. An l_1 -oracle inequality for the Lasso in finite mixture Gaussian regression models. *ESAIM: Probability and Statistics*, 17:650–671, 2013.
- Lucie Montuelle, Erwan Le Pennec, et al. Mixture of gaussian regressions model with logistic weights, a penalized maximum likelihood approach. *Electronic Journal of Statistics*, 8(1):1661–1695, 2014.
- Basil Mustafa, Carlos Riquelme Ruiz, Joan Puigcerver, Rodolphe Jenatton, and Neil Houlsby. Multimodal contrastive learning with LIMoe: the language-image mixture of experts. In Alice H. Oh, Alekh Agarwal, Danielle Belgrave, and Kyunghyun Cho, editors, *Advances in Neural Information Processing Systems*, 2022. URL <https://openreview.net/forum?id=Qy1D9JyMBg0>.

- Hien D Nguyen and Faicel Chamroukhi. Practical and theoretical aspects of mixture-of-experts modeling: An overview. *Wiley Interdisciplinary Reviews: Data Mining and Knowledge Discovery*, 8(4):e1246, 2018.
- Hien D Nguyen, Luke R Lloyd-Jones, and Geoffrey J McLachlan. A universal approximation theorem for mixture-of-experts models. *Neural computation*, 28(12):2585–2593, 2016.
- Hien D Nguyen, Faicel Chamroukhi, and Florence Forbes. Approximation results regarding the multiple-output Gaussian gated mixture of linear experts model. *Neurocomputing*, 366: 208–214, 2019. ISSN 0925-2312. doi: <https://doi.org/10.1016/j.neucom.2019.08.014>. URL <https://www.sciencedirect.com/science/article/pii/S0925231219311336>.
- Hien D Nguyen, TrungTin Nguyen, Faicel Chamroukhi, and Geoffrey John McLachlan. Approximations of conditional probability density functions in Lebesgue spaces via mixture of experts models. *Journal of Statistical Distributions and Applications*, 8(1):13, 2021a. ISSN 2195-5832. doi: 10.1186/s40488-021-00125-0. URL <https://doi.org/10.1186/s40488-021-00125-0>.
- Tan Minh Nguyen, Tam Minh Nguyen, Hai Ngoc Do, Khai Nguyen, Vishwanath Saragadam, Minh Pham, Nguyen Duy Khuong, Nhat Ho, and Stanley Osher. Improving Transformer with an Admixture of Attention Heads. In Alice H. Oh, Alekh Agarwal, Danielle Belgrave, and Kyunghyun Cho, editors, *Advances in Neural Information Processing Systems*, 2022a. URL <https://openreview.net/forum?id=0VFQhPGF1M3>.
- Trung Tin Nguyen, Faicel Chamroukhi, Hien Duy Nguyen, and Florence Forbes. Non-asymptotic model selection in block-diagonal mixture of polynomial experts models. *arXiv preprint arXiv:2104.08959*, 2021b. URL <https://arxiv.org/abs/2104.08959>.
- TrungTin Nguyen. *Model Selection and Approximation in High-dimensional Mixtures of Experts Models: from Theory to Practice*. PhD Thesis, Normandie Université, December 2021. URL <https://tel.archives-ouvertes.fr/tel-03524749>.
- TrungTin Nguyen, Faicel Chamroukhi, Hien D Nguyen, and Geoffrey J McLachlan. Approximation of probability density functions via location-scale finite mixtures in Lebesgue spaces. *arXiv preprint arXiv:2008.09787*. *Communications in Statistics - Theory and Methods*, 2020a.
- TrungTin Nguyen, Hien D Nguyen, Faicel Chamroukhi, and Geoffrey J McLachlan. Approximation by finite mixtures of continuous density functions that vanish at infinity. *Cogent Mathematics & Statistics*, 7(1):1750861, 2020b.
- TrungTin Nguyen, Faicel Chamroukhi, Hien D. Nguyen, and Geoffrey J. McLachlan. Approximation of probability density functions via location-scale finite mixtures in Lebesgue spaces. *Communications in Statistics - Theory and Methods*, pages 1–12, May 2022b. ISSN 0361-0926. doi: 10.1080/03610926.2021.2002360. URL <https://doi.org/10.1080/03610926.2021.2002360>. Publisher: Taylor & Francis.
- TrungTin Nguyen, Hien Duy Nguyen, Faicel Chamroukhi, and Florence Forbes. A non-asymptotic approach for model selection via penalization in high-dimensional mixture of experts models.

- Electronic Journal of Statistics*, 16(2):4742 – 4822, 2022c. doi: 10.1214/22-EJS2057. URL <https://doi.org/10.1214/22-EJS2057>. Publisher: Institute of Mathematical Statistics and Bernoulli Society.
- XuanLong Nguyen. Convergence of latent mixing measures in finite and infinite mixture models. *The Annals of Statistics*, 41(1):370–400, 2013. doi: 10.1214/12-AOS1065. URL <https://doi.org/10.1214/12-AOS1065>.
- Andriy Norets. Approximation of conditional densities by smooth mixtures of regressions. *The Annals of Statistics*, 38(3):1733 – 1766, 2010. doi: 10.1214/09-AOS765. URL <https://doi.org/10.1214/09-AOS765>. Publisher: Institute of Mathematical Statistics.
- Joan Puigcerver, Carlos Riquelme Ruiz, Basil Mustafa, Cedric Renggli, André Susano Pinto, Sylvain Gelly, Daniel Keysers, and Neil Houlsby. Scalable Transfer Learning with Expert Models. In *International Conference on Learning Representations*, 2021. URL <https://openreview.net/forum?id=23ZjUGpjcc>.
- Alexander Rakhlin, Dmitry Panchenko, and Sayan Mukherjee. Risk bounds for mixture density estimation. *ESAIM: PS*, 9:220–229, 2005. doi: 10.1051/ps:2005011. URL <https://doi.org/10.1051/ps:2005011>.
- Richard A Redner and Homer F Walker. Mixture densities, maximum likelihood and the EM algorithm. *SIAM review*, 26(2):195–239, 1984. Publisher: SIAM.
- Carlos Riquelme Ruiz, Joan Puigcerver, Basil Mustafa, Maxim Neumann, Rodolphe Jenatton, André Susano Pinto, Daniel Keysers, and Neil Houlsby. Scaling Vision with Sparse Mixture of Experts. In A. Beygelzimer, Y. Dauphin, P. Liang, and J. Wortman Vaughan, editors, *Advances in Neural Information Processing Systems*, 2021. URL https://openreview.net/forum?id=NGPmH3vbAA_.
- Gideon Schwarz. Estimating the dimension of a model. *The annals of statistics*, 6(2):461–464, 1978. Publisher: Institute of Mathematical Statistics.
- Noam Shazeer, *Azalia Mirhoseini, *Krzysztof Maziarsz, Andy Davis, Quoc Le, Geoffrey Hinton, and Jeff Dean. Outrageously Large Neural Networks: The Sparsely-Gated Mixture-of-Experts Layer. In *International Conference on Learning Representations*, 2017. URL <https://openreview.net/forum?id=BlckMDqlg>.
- N Stadler, P Buhlmann, and S van de Geer. l_1 -penalization for mixture regression models. *TEST*, 19:209–256, 2010.
- Abhradeep Guha Thakurta and Adam Smith. Differentially Private Feature Selection via Stability Arguments, and the Robustness of the Lasso. In Shai Shalev-Shwartz and Ingo Steinwart, editors, *Proceedings of the 26th Annual Conference on Learning Theory*, volume 30 of *Proceedings of Machine Learning Research*, pages 819–850, Princeton, NJ, USA, June 2013. PMLR. URL <https://proceedings.mlr.press/v30/Guha13.html>.
- Robert Tibshirani. Regression shrinkage and selection via the Lasso. *Journal of the Royal Statistical Society: Series B (Methodological)*, 58(1):267–288, 1996. Publisher: Wiley Online Library.

AW Van Der Vaart and JA Wellner. Weak convergence and empirical processes: With applications to statistics springer series in statistics. *Springer*, 58:59, 1996.

Vladimir Vapnik. *Estimation of Dependences Based on Empirical Data (Springer Series in Statistics)*. Springer-Verlag, 1982.

Aditya Varre and Nicolas Flammarion. Accelerated SGD for Non-Strongly-Convex Least Squares. In Po-Ling Loh and Maxim Raginsky, editors, *Proceedings of Thirty Fifth Conference on Learning Theory*, volume 178 of *Proceedings of Machine Learning Research*, pages 2062–2126. PMLR, July 2022. URL <https://proceedings.mlr.press/v178/varre22a.html>.

Ashish Vaswani, Noam Shazeer, Niki Parmar, Jakob Uszkoreit, Llion Jones, Aidan N Gomez, Lukasz Kaiser, and Illia Polosukhin. Attention is All you Need. In I. Guyon, U. Von Luxburg, S. Bengio, H. Wallach, R. Fergus, S. Vishwanathan, and R. Garnett, editors, *Advances in Neural Information Processing Systems*, volume 30. Curran Associates, Inc., 2017. URL <https://proceedings.neurips.cc/paper/2017/file/3f5ee243547dee91fbd053c1c4a845a>

Martin J Wainwright. *High-dimensional statistics: A non-asymptotic viewpoint*, volume 48. Cambridge University Press, 2019.

Christopher K Williams and Carl Edward Rasmussen. *Gaussian processes for machine learning*, volume 2. MIT press Cambridge, MA, 2006.

An Yang, Junyang Lin, Rui Men, Chang Zhou, Le Jiang, Xianyan Jia, Ang Wang, Jie Zhang, Jiamang Wang, and Yong Li. M6-t: Exploring sparse expert models and beyond. *arXiv preprint arXiv:2105.15082*, 2021.

Fei Ye and Adrian G. Bors. Deep Mixture Generative Autoencoders. *IEEE Transactions on Neural Networks and Learning Systems*, 33(10):5789–5803, 2022. doi: 10.1109/TNNLS.2021.3071401.

Fei Ye and Adrian G. Bors. Lifelong Mixture of Variational Autoencoders. *IEEE Transactions on Neural Networks and Learning Systems*, 34(1):461–474, 2023. doi: 10.1109/TNNLS.2021.3096457.

Zhao You, Shulin Feng, Dan Su, and Dong Yu. Speechmoe2: Mixture-of-Experts Model with Improved Routing. In *ICASSP 2022 - 2022 IEEE International Conference on Acoustics, Speech and Signal Processing (ICASSP)*, pages 7217–7221, 2022. doi: 10.1109/ICASSP43922.2022.9747065.

S E Yuksel, J N Wilson, and P D Gader. Twenty Years of Mixture of Experts. *IEEE Transactions on Neural Networks and Learning Systems*, 23(8):1177–1193, 2012. ISSN 2162-2388 VO - 23. doi: 10.1109/TNNLS.2012.2200299.

Barret Zoph, Irwan Bello, Sameer Kumar, Nan Du, Yanping Huang, Jeff Dean, Noam Shazeer, and William Fedus. Designing effective sparse expert models. *arXiv preprint arXiv:2202.08906*, 2022.

Appendix A. Proof of main results

First, it is important to note that the negative of the differential entropy (see, *e.g.*, [Mansuripur \(1987, Chapter 9\)](#)) of the true unknown conditional density $s_0 \in S$, defined in (5), is finite, see more in Lemma 6, which is proved in Appendix B.3.

Lemma 6 *There exists a constant $H_{s_0} = \max \{0, \ln C_{s_0}\}$, where $C_{s_0} = (4\pi)^{-q/2} A_{\Sigma}^{q/2}$, s.t.*

$$\max \left\{ 0, \sup_{x \in \mathcal{X}} \int_{\mathbb{R}^q} \ln(s_0(y|x)) s_0(y|x) dy \right\} \leq H_{s_0} < \infty. \quad (32)$$

We then introduce some definitions and notations that we shall use in the proofs.

A.1. Additional notation

For any measurable function $f : \mathbb{R}^q \rightarrow \mathbb{R}$, consider its empirical norm

$$\|f\|_n := \sqrt{\frac{1}{n} \sum_{i=1}^n f^2(Y_i|x_i)},$$

and its conditional expectation

$$\mathbb{E}_{Y|X=x}[f] := \mathbb{E}[f(Y|X) | X = x] = \int_{\mathbb{R}^q} f(y|x) s_0(y|x) dy.$$

Furthermore, we also define its empirical process

$$P_n(f) := \frac{1}{n} \sum_{i=1}^n f(Y_i|x_i), \quad (33)$$

with expectation

$$P(f) = \frac{1}{n} \sum_{i=1}^n \mathbb{E}_{Y_i|X_i=x_i}[f(Y_i|X_i) | X_i = x_i] = \frac{1}{n} \sum_{i=1}^n \int_{\mathbb{R}^q} f(y|x_i) s_0(y|x_i) dy, \quad (34)$$

and the recentered process

$$\nu_n(f) := P_n(f) - P(f) = \frac{1}{n} \sum_{i=1}^n \left[f(Y_i|x_i) - \int_{\mathbb{R}^q} f(y|x_i) s_0(y|x_i) dy \right]. \quad (35)$$

For all $m \in \mathbb{N}^*$, recall that we consider the model

$$\begin{aligned} S_m &= \left\{ s_\psi \in S, \left\| \psi^{[1,2]} \right\|_1 \leq m \right\}, \\ F_m &= \left\{ f_m = -\ln \left(\frac{s_m}{s_0} \right) = \ln(s_0) - \ln(s_m), s_m \in S_m \right\}. \end{aligned}$$

By using the basic properties of the infimum: for every $\epsilon > 0$, there exists $x_\epsilon \in A$, such that $x_\epsilon < \inf A + \epsilon$. Then let $\delta_{\text{KL}} > 0$ for all $m \in \mathbb{N}^*$, and let $\eta_m \geq 0$. It holds that there exist two functions $\widehat{s}_m$ and $\overline{s}_m$ in S_m , such that

$$P_n(-\ln \widehat{s}_m) \leq \inf_{s_m \in S_m} P_n(-\ln s_m) + \eta_m, \text{ and} \quad (36)$$

$$\text{KL}_n(s_0, \overline{s}_m) \leq \inf_{s_m \in S_m} \text{KL}_n(s_0, s_m) + \delta_{\text{KL}}. \quad (37)$$

Define

$$\widehat{f}_m := -\ln \left(\frac{\widehat{s}_m}{s_0} \right), \text{ and } \overline{f}_m := -\ln \left(\frac{\overline{s}_m}{s_0} \right). \quad (38)$$

Let $\eta \geq 0$ and fix $m \in \mathbb{N}^*$. Further, define

$$\widehat{\mathcal{M}}(m) = \{m' \in \mathbb{N}^* | P_n(-\ln \widehat{s}_{m'}) + \text{pen}(m') \leq P_n(-\ln \widehat{s}_m) + \text{pen}(m) + \eta\}. \quad (39)$$

A.2. Proof of Theorem 1

Let $\lambda > 0$ and define $\widehat{m}$ to be the smallest integer such that $\widehat{s}^{\text{L}}(\lambda)$ belongs to $S_{\widehat{m}}$, i.e., $\widehat{m} := \lceil \|\psi^{[1,2]}\|_1 \rceil \leq \|\psi^{[1,2]}\|_1 + 1$. Then using the definition of $\widehat{m}$, (7), (10), and $S = \bigcup_{m \in \mathbb{N}^*} S_m$, we get

$$\begin{aligned} & -\frac{1}{n} \sum_{i=1}^n \ln(\widehat{s}^{\text{L}}(\lambda)(y_i|x_i)) + \lambda \widehat{m} \\ & \leq -\frac{1}{n} \sum_{i=1}^n \ln(\widehat{s}^{\text{L}}(\lambda)(y_i|x_i)) + \lambda \left(\|\psi^{[1,2]}\|_1 + 1 \right) \\ & = \inf_{s_\psi \in S} \left(-\frac{1}{n} \sum_{i=1}^n \ln(s_\psi(y_i|x_i)) + \lambda \|\psi^{[1,2]}\|_1 \right) + \lambda \\ & = \inf_{m \in \mathbb{N}^*} \left(\inf_{s_\psi \in S_m} \left(-\frac{1}{n} \sum_{i=1}^n \ln(s_\psi(y_i|x_i)) + \lambda \|\psi^{[1,2]}\|_1 \right) \right) + \lambda \\ & = \inf_{m \in \mathbb{N}^*} \left(\inf_{s_\psi \in S, \|\psi^{[1,2]}\|_1 \leq m} \left(-\frac{1}{n} \sum_{i=1}^n \ln(s_\psi(y_i|x_i)) + \lambda \|\psi^{[1,2]}\|_1 \right) \right) + \lambda \\ & \leq \inf_{m \in \mathbb{N}^*} \left(\inf_{s_m \in S_m} \left(-\frac{1}{n} \sum_{i=1}^n \ln(s_m(y_i|x_i)) + \lambda m \right) \right) + \lambda, \end{aligned}$$

which implies

$$-\frac{1}{n} \sum_{i=1}^n \ln(\widehat{s}^{\text{L}}(\lambda)(y_i|x_i)) + \text{pen}(\widehat{m}) \leq \inf_{m \in \mathbb{N}^*} \left(-\frac{1}{n} \sum_{i=1}^n \ln(\widehat{s}_m(y_i|x_i)) + \text{pen}(m) \right) + \eta$$

with $\text{pen}(m) = \lambda m$, $\eta = \lambda$, and $\widehat{s}_m$ is a η_m -log-likelihood minimizer in S_m , with $\eta_m \geq 0$ defined by (11). Thus, $\widehat{s}^{\text{L}}(\lambda)$ satisfies (12) with $\widehat{s}^{\text{L}}(\lambda) \equiv \widehat{s}_{\widehat{m}}$, i.e.,

$$-\frac{1}{n} \sum_{i=1}^n \ln(\widehat{s}_{\widehat{m}}(y_i|x_i)) + \text{pen}(\widehat{m}) \leq \inf_{m \in \mathbb{N}^*} \left(-\frac{1}{n} \sum_{i=1}^n \ln(\widehat{s}_m(y_i|x_i)) + \text{pen}(m) \right) + \eta. \quad (40)$$

Then, Theorem 2 implies that if

$$\lambda \geq \kappa \frac{KB_{0n}}{\sqrt{n}} \left(q \ln n \sqrt{\ln(2p+1)} + 1 \right), B_{0n} = \max(A_\Sigma, 1 + KA_G) [1 + 2q\sqrt{q}A_\Sigma (5A_\beta^2 + 4A_\Sigma \ln n)],$$

for some absolute constants $\kappa \geq 148$, it holds that, as required

$$\begin{aligned} \mathbb{E} [\text{KL}_n(s_0, \hat{s}^L(\lambda))] &\leq (1 + \kappa^{-1}) \inf_{s_\psi \in S} \left(\text{KL}_n(s_0, s_\psi) + \lambda \left\| \psi^{[1,2]} \right\|_1 \right) + \lambda \\ &\quad + \sqrt{\frac{K}{n}} \left(\frac{e^{q/2-1} \pi^{q/2}}{A_\Sigma^{q/2}} + H_{s_0} \right) \sqrt{2qA_\gamma} \\ &\quad + 302q \sqrt{\frac{K}{n}} \max(A_\Sigma, 1 + KA_G) (1 + 2q\sqrt{q}A_\Sigma (5A_\beta^2 + 4A_\Sigma \ln n)) \\ &\quad \times K \left(1 + \left(A_\gamma + qA_\beta + \frac{q\sqrt{q}}{a_\Sigma} \right)^2 \right). \end{aligned}$$

A.3. Proof of Theorem 2

Let $M_n > 0$ and $\kappa \geq 148$. Assume that, for all $m \in \mathbb{N}^*$, the penalty function satisfies $\text{pen}(m) = \lambda m$, with

$$\lambda \geq \kappa \frac{KB_n}{\sqrt{n}} \left(q \ln n \sqrt{\ln(2p+1)} + 1 \right). \quad (41)$$

We derive, from Propositions 3 and 4, that any penalized likelihood estimator $\hat{s}_{\hat{m}}$ with $\hat{m}$, satisfying

$$-\frac{1}{n} \sum_{i=1}^n \ln(\hat{s}_{\hat{m}}(y_i|x_i)) + \text{pen}(\hat{m}) \leq \inf_{m \in \mathbb{N}^*} \left(-\frac{1}{n} \sum_{i=1}^n \ln(\hat{s}_m(y_i|x_i)) + \text{pen}(m) \right) + \eta,$$

for some $\eta \geq 0$, yields

$$\begin{aligned} \mathbb{E} [\text{KL}_n(s_0, \hat{s}_{\hat{m}})] &= \mathbb{E} [\text{KL}_n(s_0, \hat{s}_{\hat{m}}) \mathbb{1}_{\mathcal{T}}] + \mathbb{E} [\text{KL}_n(s_0, \hat{s}_{\hat{m}}) \mathbb{1}_{\mathcal{T}^c}] \\ &\leq (1 + \kappa^{-1}) \inf_{m \in \mathbb{N}^*} \left(\inf_{s_m \in S_m} \text{KL}_n(s_0, s_m) + \text{pen}(m) + \eta_m \right) \\ &\quad + \frac{302K^{3/2}qB_n}{\sqrt{n}} \left(1 + \left(A_\gamma + qA_\beta + \frac{q\sqrt{q}}{a_\Sigma} \right)^2 \right) + \eta \\ &\quad + \left(\frac{e^{q/2-1} \pi^{q/2}}{A_\Sigma^{q/2}} + H_{s_0} \right) \sqrt{2KnqA_\gamma} e^{-\frac{M_n^2 - 2M_n A_\beta}{4A_\Sigma}}. \end{aligned} \quad (42)$$

To obtain inequality (20), it only remains to optimize the inequality (42), with respect M_n . Since the two terms depending on M_n , in (42), have opposite monotonicity with respect to M_n , we are looking for a value of M_n such that these two terms are the same order with respect to n . Consider the positive solution $M_n = A_\beta + \sqrt{A_\beta^2 + 4A_\Sigma \ln n}$ of the equation $\frac{X(X-2A_\beta)}{4A_\Sigma} - \ln n = 0$. Then, on the one hand,

$$e^{-\frac{M_n^2 - 2M_n A_\beta}{4A_\Sigma}} \sqrt{n} = e^{-\ln n} \sqrt{n} = \frac{1}{\sqrt{n}}.$$

On the other hand, using the inequality $(a + b)^2 \leq 2(a^2 + b^2)$, we have

$$\begin{aligned} B_n &= \max(A_\Sigma, 1 + KA_G) \left(1 + q\sqrt{q} (M_n + A_\beta)^2 A_\Sigma \right) \\ &= \max(A_\Sigma, 1 + KA_G) \left(1 + q\sqrt{q} A_\Sigma \left(2A_\beta + \sqrt{A_\beta^2 + 4A_\Sigma \ln n} \right)^2 \right) \\ &\leq \max(A_\Sigma, 1 + KA_G) \left(1 + 2q\sqrt{q} A_\Sigma (5A_\beta^2 + 4A_\Sigma \ln n) \right), \end{aligned}$$

hence (42) implies (20).

A.4. Proof of Proposition 3

For every $m' \in \widehat{\mathcal{M}}(m)$, from (39), (38), and (36), we obtain

$$\begin{aligned} P_n(\widehat{f}_{m'}) + \text{pen}(m') &= P_n(\ln(s_0) - \ln(\widehat{s}_{m'})) + \text{pen}(m') \text{ (using (38))} \\ &\leq P_n(\ln(s_0) - \ln(\widehat{s}_m)) + \text{pen}(m) + \eta \text{ (using (39))} \\ &\leq P_n(\ln(s_0) - \ln(\overline{s}_m)) + \eta_m + \text{pen}(m) + \eta \\ &\quad \text{(using (36) with } \overline{s}_m \text{ in } S_m \text{ and linearity of } P_n) \\ &= P_n(\overline{f}_m) + \text{pen}(m) + \eta_m + \eta \text{ (using (38))}. \end{aligned}$$

By the definition of recentered process, $\nu_n(\cdot)$, in (35), it holds that

$$P(\widehat{f}_{m'}) + \text{pen}(m') \leq P(\overline{f}_m) + \text{pen}(m) + \nu_n(\overline{f}_m) - \nu_n(\widehat{f}_{m'}) + \eta + \eta_m.$$

Taking into account (6) and (33), we obtain

$$\begin{aligned} \text{KL}_n(s_0, \widehat{s}_{m'}) &= \frac{1}{n} \sum_{i=1}^n \int_{\mathbb{R}^q} \ln \left(\frac{s_0(y|x_i)}{\widehat{s}_{m'}(y|x_i)} \right) s_0(y|x_i) dy \\ &= \frac{1}{n} \sum_{i=1}^n \int_{\mathbb{R}^q} \widehat{f}_{m'}(y|x_i) s_0(y|x_i) dy \text{ (using (38))} \\ &= P(\widehat{f}_{m'}) \text{ (using (34))}. \end{aligned}$$

Similarly, we also obtain $\text{KL}_n(s_0, \overline{s}_m) = P(\overline{f}_m)$. Hence, (37) implies that

$$\begin{aligned} \text{KL}_n(s_0, \widehat{s}_{m'}) + \text{pen}(m') &\leq \text{KL}_n(s_0, \overline{s}_m) + \text{pen}(m) + \nu_n(\overline{f}_m) - \nu_n(\widehat{f}_{m'}) + \eta + \eta_m \\ &\leq \inf_{s_m \in S_m} \text{KL}_n(s_0, s_m) + \text{pen}(m) + \nu_n(\overline{f}_m) \\ &\quad - \nu_n(\widehat{f}_{m'}) + \eta_m + \delta_{\text{KL}} + \eta. \end{aligned} \tag{43}$$

All that remains is to control the deviation of $-\nu_n(\widehat{f}_{m'}) = \nu_n(-\widehat{f}_{m'})$. To handle the randomness of $\widehat{f}_{m'}$, we shall control the deviation of $\sup_{f_{m'} \in F_{m'}} \nu_n(-f_{m'})$, since $\widehat{f}_{m'} \in F_{m'}$. Such control is

provided by Lemma 5. From (43) and (30), we derive that on the event $\mathcal{T}$, for all $m \in \mathbb{N}^*$, and $t > 0$, with probability larger than $1 - e^{-t}$,

$$\begin{aligned}
 \text{KL}_n(s_0, \widehat{s}_{m'}) + \text{pen}(m') &\leq \inf_{s_m \in S_m} \text{KL}_n(s_0, s_m) + \text{pen}(m) + \nu_n(\overline{f}_m) - \nu_n(\widehat{f}_{m'}) + \eta_m + \delta_{\text{KL}} + \eta \\
 &\leq \inf_{s_m \in S_m} \text{KL}_n(s_0, s_m) + \text{pen}(m) + \nu_n(\overline{f}_m) + \eta_m + \delta_{\text{KL}} + \eta \\
 &\quad + \frac{4KB_n}{\sqrt{n}} \left[37q\Delta_{m'} + \sqrt{2} \left(A_\gamma + qA_\beta + \frac{q\sqrt{q}}{a_\Sigma} \right) \sqrt{t} \right] \\
 &\leq \inf_{s_m \in S_m} \text{KL}_n(s_0, s_m) + \text{pen}(m) + \nu_n(\overline{f}_m) + \eta_m + \delta_{\text{KL}} + \eta \\
 &\quad + \frac{4KB_n}{\sqrt{n}} \left[37q\Delta_{m'} + \frac{1}{2} \left(A_\gamma + qA_\beta + \frac{q\sqrt{q}}{a_\Sigma} \right)^2 + t \right], \text{ if } m' \in \widehat{\mathcal{M}}(m).
 \end{aligned} \tag{44}$$

Here, we use the fact that $\widehat{f}_{m'} \in F_{m'}$ and get the last inequality using the fact that $2ab \leq a^2 + b^2$ for $b = \sqrt{t}$, and $a = \left(A_\gamma + qA_\beta + \frac{q\sqrt{q}}{a_\Sigma} \right) / \sqrt{2}$.

It remains to sum up the tail bounds (44) over all possible values of $m \in \mathbb{N}^*$ and $m' \in \widehat{\mathcal{M}}(m)$. To get an inequality valid on a set of high probability, we need to adequately choose the value of the parameter t , depending on $m \in \mathbb{N}^*$ and $m' \in \widehat{\mathcal{M}}(m)$. Let $z > 0$, for all $m \in \mathbb{N}^*$ and $m' \in \widehat{\mathcal{M}}(m)$, and apply (44) to obtain $t = z + m + m'$. Then, on the event $\mathcal{T}$, for all $m \in \mathbb{N}^*$, with probability larger than $1 - e^{-(z+m+m')}$,

$$\begin{aligned}
 \text{KL}_n(s_0, \widehat{s}_{m'}) + \text{pen}(m') &\leq \inf_{s_m \in S_m} \text{KL}_n(s_0, s_m) + \text{pen}(m) + \nu_n(\overline{f}_m) + \eta_m + \delta_{\text{KL}} + \eta \\
 &\quad + \frac{4KB_n}{\sqrt{n}} \left[37q\Delta_{m'} + \frac{1}{2} \left(A_\gamma + qA_\beta + \frac{q\sqrt{q}}{a_\Sigma} \right)^2 + (z + m + m') \right], \\
 \text{if } m' \in \widehat{\mathcal{M}}(m).
 \end{aligned} \tag{45}$$

Here, (45) is equivalent to

$$\begin{aligned}
 \text{KL}_n(s_0, \widehat{s}_{m'}) - \nu_n(\overline{f}_m) &\leq \inf_{s_m \in S_m} \text{KL}_n(s_0, s_m) + \left[\text{pen}(m) + \frac{4KB_n}{\sqrt{n}} m \right] + \eta_m + \delta_{\text{KL}} + \eta \\
 &\quad + \left[\frac{4KB_n}{\sqrt{n}} (37q\Delta_{m'} + m') - \text{pen}(m') \right] \\
 &\quad + \frac{4KB_n}{\sqrt{n}} \left[\frac{1}{2} \left(A_\gamma + qA_\beta + \frac{q\sqrt{q}}{a_\Sigma} \right)^2 + z \right].
 \end{aligned} \tag{46}$$

Note that with probability larger than $1 - e^{-z}$, (45) holds simultaneously for all $m \in \mathbb{N}^*$ and $m' \in \widehat{\mathcal{M}}(m)$. Indeed, by defining the event

$$\cap_{(m,m') \in \mathbb{N}^* \times \widehat{\mathcal{M}}(m)} \Omega_{m,m'} = \{w : w \in \Omega \text{ such that the event in (45) holds}\},$$

it holds that, on the event $\mathcal{T}$,

$$\begin{aligned}
 \mathbb{P} \left(\bigcap_{(m,m') \in \mathbb{N}^* \times \widehat{\mathcal{M}}(m)} \Omega_{m,m'} \right) &= 1 - \mathbb{P} \left(\bigcup_{(m,m') \in \mathbb{N}^* \times \widehat{\mathcal{M}}(m)} \Omega_{m,m'}^C \right) \\
 &\geq 1 - \sum_{(m,m') \in \mathbb{N}^* \times \widehat{\mathcal{M}}(m)} \mathbb{P}(\Omega_{m,m'}^C) \\
 &\geq 1 - \sum_{(m,m') \in \mathbb{N}^* \times \widehat{\mathcal{M}}(m)} e^{-(z+m+m')} \\
 &\geq 1 - \sum_{(m,m') \in \mathbb{N}^* \times \mathbb{N}^*} e^{-(z+m+m')} \\
 &= 1 - e^{-z} \left(\sum_{m \in \mathbb{N}^*} e^{-m} \right)^2 \\
 &\geq 1 - e^{-z},
 \end{aligned}$$

where we get the last inequality by using the the geometric series

$$\sum_{m=1}^{\infty} (e^{-1})^m = \sum_{m=0}^{\infty} (e^{-1})^m - 1 = \frac{1}{1-e^{-1}} - 1 = \frac{e}{e-1} - 1 = \frac{1}{e-1} < 1.$$

Taking into account (29), we get

$$\begin{aligned}
 \text{KL}_n(s_0, \widehat{s}_{m'}) - \nu_n(\bar{f}_m) &\leq \inf_{s_m \in S_m} \text{KL}_n(s_0, s_m) + \left[\text{pen}(m) + \frac{4KB_n}{\sqrt{n}}m \right] + \eta_m + \delta_{\text{KL}} + \eta \\
 &\quad + \left[\frac{4KB_n}{\sqrt{n}} \left(37q \ln n \sqrt{\ln(2p+1)} + 1 \right) m' - \text{pen}(m') \right] \\
 &\quad + \frac{4KB_n}{\sqrt{n}} \left[\frac{1}{2} \left(A_\gamma + qA_\beta + \frac{q\sqrt{q}}{a_\Sigma} \right)^2 + 74q\sqrt{K} \left(A_\gamma + qA_\beta + \frac{q\sqrt{q}}{a_\Sigma} \right) + z \right].
 \end{aligned} \tag{47}$$

Now, let $\kappa \geq 1$ and assume that $\text{pen}(m) = \lambda m$, for all $m \in \mathbb{N}^*$ with

$$\lambda \geq \kappa \frac{4KB_n}{\sqrt{n}} \left(37q \ln n \sqrt{\ln(2p+1)} + 1 \right). \tag{48}$$

Then, (47) implies

$$\begin{aligned}
 \text{KL}_n(s_0, \widehat{s}_{m'}) - \nu_n(\bar{f}_m) &\leq \inf_{s_m \in S_m} \text{KL}_n(s_0, s_m) + \left[\lambda m + \frac{4KB_n}{\sqrt{n}} m \right] + \eta_m + \delta_{\text{KL}} + \eta \\
 &\quad + \left[\underbrace{\frac{4KB_n}{\sqrt{n}} \left(37q \ln n \sqrt{\ln(2p+1)} + 1 \right)}_{\leq \lambda \kappa^{-1}} m' - \lambda m' \right] \\
 &\quad + \frac{4KB_n}{\sqrt{n}} \left[\frac{1}{2} \left(A_\gamma + qA_\beta + \frac{q\sqrt{q}}{a_\Sigma} \right)^2 + 74q\sqrt{K} \left(A_\gamma + qA_\beta + \frac{q\sqrt{q}}{a_\Sigma} \right) + z \right] \\
 &\leq \inf_{s_m \in S_m} \text{KL}_n(s_0, s_m) + \left[\text{pen}(m) + \underbrace{\frac{4KB_n}{\sqrt{n}} m}_{\leq \kappa^{-1} \text{pen}(m)} \right] + \eta_m + \delta_{\text{KL}} + \eta \\
 &\quad + \underbrace{[\lambda \kappa^{-1} m' - \lambda m']}_{\leq 0} \\
 &\quad + \frac{4KB_n}{\sqrt{n}} \left[\frac{1}{2} \left(A_\gamma + qA_\beta + \frac{q\sqrt{q}}{a_\Sigma} \right)^2 + 74q\sqrt{K} \left(A_\gamma + qA_\beta + \frac{q\sqrt{q}}{a_\Sigma} \right) + z \right] \\
 &\leq \inf_{s_m \in S_m} \text{KL}_n(s_0, s_m) + (1 + \kappa^{-1}) \text{pen}(m) + \eta_m + \delta_{\text{KL}} + \eta \\
 &\quad + \frac{4KB_n}{\sqrt{n}} \left[\frac{1}{2} \left(A_\gamma + qA_\beta + \frac{q\sqrt{q}}{a_\Sigma} \right)^2 + 74q\sqrt{K} \left(A_\gamma + qA_\beta + \frac{q\sqrt{q}}{a_\Sigma} \right) + z \right].
 \end{aligned}$$

Next, using the inequality $2ab \leq \beta^{-1}a^2 + \beta^{-1}b^2$ for $a = \sqrt{K}$, $b = K \left(A_\gamma + qA_\beta + \frac{q\sqrt{q}}{a_\Sigma} \right)$, and $\beta = \sqrt{K}$, and the fact that $K \leq K^{3/2}$, for all $K \in \mathbb{N}^*$, it follows that

$$\begin{aligned}
 &\text{KL}_n(s_0, \widehat{s}_{m'}) - \nu_n(\bar{f}_m) \\
 &\leq \inf_{s_m \in S_m} \text{KL}_n(s_0, s_m) + (1 + \kappa^{-1}) \text{pen}(m) + \eta_m + \delta_{\text{KL}} + \eta \\
 &\quad + \frac{4B_n}{\sqrt{n}} \left[\frac{qK^{3/2}}{2} \left(A_\gamma + qA_\beta + \frac{q\sqrt{q}}{a_\Sigma} \right)^2 + \underbrace{74q\sqrt{K}K \left(A_\gamma + qA_\beta + \frac{q\sqrt{q}}{a_\Sigma} \right)}_{37q \times 2ab} + Kz \right] \\
 &\leq \inf_{s_m \in S_m} \text{KL}_n(s_0, s_m) + (1 + \kappa^{-1}) \text{pen}(m) + \eta_m + \delta_{\text{KL}} + \eta \\
 &\quad + \frac{4B_n}{\sqrt{n}} \left[37qK^{1/2} + \frac{75qK^{3/2}}{2} \left(A_\gamma + qA_\beta + \frac{q\sqrt{q}}{a_\Sigma} \right)^2 + Kz \right]. \tag{49}
 \end{aligned}$$

By (12) and (39), $\widehat{m}$ belongs to $\widehat{\mathcal{M}}(m)$, for all $m \in \mathbb{N}^*$, so we deduce from (49) that on the event $\mathcal{T}$, for all $z > 0$, with probability greater than $1 - e^{-z}$,

$$\begin{aligned} \text{KL}_n(s_0, \widehat{s}_{\widehat{m}}) - \nu_n(\overline{f}_m) &\leq \inf_{m \in \mathbb{N}^*} \left(\inf_{s_m \in S_m} \text{KL}_n(s_0, s_m) + (1 + \kappa^{-1}) \text{pen}(m) + \eta_m \right) + \eta + \delta_{\text{KL}} \\ &\quad + \frac{4B_n}{\sqrt{n}} \left[37qK^{1/2} + \frac{75qK^{3/2}}{2} \left(A_\gamma + qA_\beta + \frac{q\sqrt{q}}{a_\Sigma} \right)^2 + Kz \right]. \end{aligned} \quad (50)$$

Note that for any non-negative random variable Z and any $a > 0$, $\mathbb{E}[Z] = a \int_{z \geq 0} \mathbb{P}(Z > az) dz$. Indeed, if we let $t = az$, then $dz = a dt$ and

$$\begin{aligned} a \int_{z \geq 0} \mathbb{P}(Z > az) dz &= a \int_0^\infty \int_{az}^\infty f_Z(u) du dz = \int_0^\infty \int_t^\infty f_Z(u) du dt = \int_0^\infty \int_0^u f_Z(u) dt du \\ &= \int_0^\infty f_Z(u) \int_0^u dt du = \int_0^\infty f_Z(u) u du = \mathbb{E}[Z]. \end{aligned} \quad (51)$$

Then, we define the following random variable w.r.t. the random response $Y_{[n]} := (Y_i)_{i \in [n]}$:

$$\begin{aligned} Z &:= \text{KL}_n(s_0, \widehat{s}_{\widehat{m}}) - \nu_n(\overline{f}_m) - \left[\inf_{m \in \mathbb{N}^*} \left(\inf_{s_m \in S_m} \text{KL}_n(s_0, s_m) + (1 + \kappa^{-1}) \text{pen}(m) + \eta_m \right) + \eta + \delta_{\text{KL}} \right] \\ &\quad - \frac{4B_n}{\sqrt{n}} \left[37qK^{1/2} + \frac{75qK^{3/2}}{2} \left(A_\gamma + qA_\beta + \frac{q\sqrt{q}}{a_\Sigma} \right)^2 \right]. \end{aligned}$$

Then by (50), on the even $\mathcal{T}$, it holds that $P(Z \leq az) \geq 1 - e^{-z}$ and if $Z \leq 0$ then $P(Z < az) = 1 \geq 1 - e^{-z}$, for all $z > 0$, where $a = \frac{4B_n K}{\sqrt{n}} > 0$. Therefore, it is sufficient to consider $Z \geq 0$ and it holds that $P(Z > az | \mathcal{T}) \leq e^{-z}$. In this case, by (51) and the fact that $\mathbb{P}(\mathcal{T}) \leq 1$, it holds that

$$\begin{aligned} \mathbb{E}_{Y_{[n]}}[Z \mathbb{1}_{\mathcal{T}}] &= \mathbb{P}(\mathcal{T}) \mathbb{E}_{Y_{[n]}}[Z | \mathcal{T}] \leq \mathbb{E}_{Y_{[n]}}[Z | \mathcal{T}] \\ &= a \int_{z \geq 0} \mathbb{P}(Z > az | \mathcal{T}) dz \leq a \int_{z \geq 0} e^{-z} dz = a. \end{aligned} \quad (52)$$

Then, by integrating (50) over $z > 0$ using (52), the fact that

$$\mathbb{E}_{Y_{[n]}}[\nu_n(\overline{f}_m)] = \mathbb{E}_{Y_{[n]}}[P_n(\overline{f}_m)] - \mathbb{E}_{Y_{[n]}}[P(\overline{f}_m)] = 0, \quad (53)$$

$\delta_{\text{KL}} > 0$ can be chosen arbitrary small, and $\mathbb{E}_{Y_{[n]}} [\mathbb{1}_{\mathcal{T}}] = \mathbb{P}(\mathcal{T}) \leq 1$, we obtain that

$$\begin{aligned}
 \mathbb{E} [\text{KL}_n(s_0, \widehat{s}_{\widehat{m}}) \mathbb{1}_{\mathcal{T}}] &\leq \left[\inf_{m \in \mathbb{N}^*} \left(\inf_{s_m \in S_m} \text{KL}_n(s_0, s_m) + (1 + \kappa^{-1}) \text{pen}(m) + \eta_m \right) + \eta \right] \mathbb{E}_{Y_{[n]}} [\mathbb{1}_{\mathcal{T}}] \\
 &\quad + \frac{4B_n}{\sqrt{n}} \left[37qK^{1/2} + \frac{75qK^{3/2}}{2} \left(A_\gamma + qA_\beta + \frac{q\sqrt{q}}{a_\Sigma} \right)^2 + K \right] \mathbb{E}_{Y_{[n]}} [\mathbb{1}_{\mathcal{T}}] \\
 &\leq \inf_{m \in \mathbb{N}^*} \left(\inf_{s_m \in S_m} \text{KL}_n(s_0, s_m) + (1 + \kappa^{-1}) \text{pen}(m) + \eta_m \right) + \eta \\
 &\quad + \frac{4B_n}{\sqrt{n}} \left[37qK^{3/2} + \frac{75qK^{3/2}}{2} \left(A_\gamma + qA_\beta + \frac{q\sqrt{q}}{a_\Sigma} \right)^2 + qK^{3/2} \right] \\
 &\leq \inf_{m \in \mathbb{N}^*} \left(\inf_{s_m \in S_m} \text{KL}_n(s_0, s_m) + (1 + \kappa^{-1}) \text{pen}(m) + \eta_m \right) + \eta \\
 &\quad + \frac{302K^{3/2}qB_n}{\sqrt{n}} \left(1 + \left(A_\gamma + qA_\beta + \frac{q\sqrt{q}}{a_\Sigma} \right)^2 \right). \tag{54}
 \end{aligned}$$

A.5. Proof of Proposition 4

By the Cauchy-Schwarz inequality,

$$\mathbb{E} [\text{KL}_n(s_0, \widehat{s}_{\widehat{m}}) \mathbb{1}_{\mathcal{T}^C}] \leq \sqrt{\mathbb{E} [\text{KL}_n^2(s_0, \widehat{s}_{\widehat{m}})]} \sqrt{\mathbb{E} [\mathbb{1}_{\mathcal{T}^C}^2]} = \sqrt{\mathbb{E} [\text{KL}_n^2(s_0, \widehat{s}_{\widehat{m}})]} \sqrt{\mathbb{P}(\mathcal{T}^C)}. \tag{55}$$

We seek to bound the two terms of the right-hand side of (55).

For the first term, let us bound $\text{KL}(s_0(\cdot|x), s_\psi(\cdot|x))$, for all $s_\psi \in S$ and $x \in \mathcal{X}$. Let $s_\psi \in S$ and $x \in \mathcal{X}$. Then, we obtain

$$\begin{aligned}
 \text{KL}(s_0(\cdot|x), s_\psi(\cdot|x)) &= \int_{\mathbb{R}^q} \ln \left(\frac{s_0(y|x)}{s_\psi(y|x)} \right) s_0(y|x) dy \\
 &= \int_{\mathbb{R}^q} \ln(s_0(y|x)) s_0(y|x) dy - \int_{\mathbb{R}^q} \ln(s_\psi(y|x)) s_0(y|x) dy \\
 &\leq - \int_{\mathbb{R}^q} \ln(s_\psi(y|x)) s_0(y|x) dy + H_{s_0}, \forall x \in \mathcal{X} \text{ (using (32))}. \tag{56}
 \end{aligned}$$

Since

$$a_G := \frac{\exp(-A_\gamma)}{\sum_{l=1}^K \exp(A_\gamma)} \leq \sup_{x \in \mathcal{X}, \gamma \in \tilde{\Gamma}} \frac{\exp(\gamma_{k0} + \gamma_k^\top x)}{\sum_{l=1}^K \exp(\gamma_{l0} + \gamma_l^\top x)} \leq \frac{\exp(A_\gamma)}{\sum_{l=1}^K \exp(-A_\gamma)} =: A_G,$$

there exists deterministic positive constants a_G, A_G , such that

$$a_G \leq \sup_{x \in \mathcal{X}, \gamma \in \tilde{\Gamma}} g_k(x; \gamma) \leq A_G. \tag{57}$$

Thus, for all $y \in \mathbb{R}^q$,

$$\begin{aligned}
 & \ln(s_\psi(y|x)) s_0(y|x) \\
 & \geq \ln \left[\sum_{k=1}^K \frac{a_G \det(\Sigma_k^{-1})^{1/2}}{(2\pi)^{q/2}} \exp \left(- \left(y^\top \Sigma_k^{-1} y + (\beta_{k0} + \beta_k x)^\top \Sigma_k^{-1} (\beta_{k0} + \beta_k x) \right) \right) \right] \\
 & \quad \times \sum_{k=1}^K \frac{a_G \det(\Sigma_{0,k}^{-1})^{1/2}}{(2\pi)^{q/2}} \exp \left(- \left(y^\top \Sigma_{0,k}^{-1} y + (\beta_{0,k0} + \beta_{0,k} x)^\top \Sigma_{0,k}^{-1} (\beta_{0,k0} + \beta_{0,k} x) \right) \right) \\
 & \quad \left(\text{using (57) and } -(a-b)^\top A(a-b)/2 \geq -(a^\top Aa + b^\top Ab), \text{ e.g., } a = y, b = \beta_{k0} + \beta_k x, A = \Sigma_k \right) \\
 & \geq \ln \left[\sum_{k=1}^K \frac{a_G a_\Sigma^{q/2}}{(2\pi)^{q/2}} \exp \left(- \left(y^\top \Sigma_k^{-1} y + (\beta_{k0} + \beta_k x)^\top \Sigma_k^{-1} (\beta_{k0} + \beta_k x) \right) \right) \right] \\
 & \quad \times \sum_{k=1}^K \frac{a_G a_\Sigma^{q/2}}{(2\pi)^{q/2}} \exp \left(- \left(y^\top \Sigma_{0,k}^{-1} y + (\beta_{0,k0} + \beta_{0,k} x)^\top \Sigma_{0,k}^{-1} (\beta_{0,k0} + \beta_{0,k} x) \right) \right) \quad (\text{using (4)}) \\
 & \geq \ln \left[K \frac{a_G a_\Sigma^{q/2}}{(2\pi)^{q/2}} \exp \left(- \left(y^\top y + q A_\beta^2 \right) A_\Sigma \right) \right] \times K \frac{a_G a_\Sigma^{q/2}}{(2\pi)^{q/2}} \exp \left(- \left(y^\top y + q A_\beta^2 \right) A_\Sigma \right) \quad (\text{using (4)}), \\
 & \tag{58}
 \end{aligned}$$

where, in the last inequality, we use the fact that for all $u \in \mathbb{R}^q$. By using the eigenvalue decomposition of $\Sigma_1 = P^\top D P$,

$$\left| u^\top \Sigma_1 u \right| = \left| u^\top P^\top D P u \right| \leq \|P u\|_2 \leq M(D) \|P u\|_2^2 \leq A_\Sigma \|u\|_2^2 \leq A_\Sigma q \|u\|_\infty^2,$$

where in the last inequality, we used the fact that (102). Therefore, setting $u = \sqrt{2A_\Sigma} y$ and $h(t) = t \ln t$, for all $t \in \mathbb{R}$, and noticing that $h(t) \geq h(e^{-1}) = -e^{-1}$, for all $t \in \mathbb{R}$, and from (56) and (58), we get that

$$\begin{aligned}
 & \text{KL}(s_0(\cdot|x), s_\psi(\cdot|x)) - H_{s_0} \\
 & \leq - \int_{\mathbb{R}^q} \left[\ln \left[K \frac{a_\gamma a_\Sigma^{q/2}}{(2\pi)^{q/2}} \exp \left(- \left(y^\top y + q A_\beta^2 \right) A_\Sigma \right) \right] K \frac{a_\gamma a_\Sigma^{q/2}}{(2\pi)^{q/2}} \exp \left(- \left(y^\top y + q A_\beta^2 \right) A_\Sigma \right) \right] dy \\
 & = - \frac{K a_\gamma a_\Sigma^{q/2} e^{-q A_\beta^2 A_\Sigma}}{(2A_\Sigma)^{q/2}} \int_{\mathbb{R}^q} \left[\ln \left(K \frac{a_\gamma a_\Sigma^{q/2}}{(2\pi)^{q/2}} \right) - q A_\beta^2 A_\Sigma - \frac{u^\top u}{2} \right] \frac{e^{-\frac{u^\top u}{2}}}{(2\pi)^{q/2}} du \\
 & = - \frac{K a_\gamma a_\Sigma^{q/2} e^{-q A_\beta^2 A_\Sigma}}{(2A_\Sigma)^{q/2}} \mathbb{E}_U \left[\left[\ln \left(K \frac{a_\gamma a_\Sigma^{q/2}}{(2\pi)^{q/2}} \right) - q A_\beta^2 A_\Sigma - \frac{U^\top U}{2} \right] \right] \quad (\text{with } U \sim \mathcal{N}_q(0, I_q)) \\
 & = - \frac{K a_\gamma a_\Sigma^{q/2} e^{-q A_\beta^2 A_\Sigma}}{(2A_\Sigma)^{q/2}} \left[\ln \left(K \frac{a_\gamma a_\Sigma^{q/2}}{(2\pi)^{q/2}} \right) - q A_\beta^2 A_\Sigma - \frac{q}{2} \right] \\
 & = - \frac{K a_\gamma a_\Sigma^{q/2} e^{-q A_\beta^2 A_\Sigma - \frac{q}{2}}}{(2\pi)^{q/2} (A_\Sigma)^{q/2}} e^{q/2 \pi^{q/2} \ln \left(\frac{K a_\gamma a_\Sigma^{q/2} e^{-q A_\beta^2 A_\Sigma - \frac{q}{2}}}{(2\pi)^{q/2}} \right)} \leq \frac{e^{q/2 - 1} \pi^{q/2}}{A_\Sigma^{q/2}}, \\
 & \tag{59}
 \end{aligned}$$

where we used the fact that $t \ln(t) \geq -e^{-1}$, for all $t \in \mathbb{R}$.

Then, for all $s_\psi \in S$,

$$\text{KL}_n(s_0, s_\psi) = \frac{1}{n} \sum_{i=1}^n \text{KL}(s_0(\cdot|x_i), s_\psi(\cdot|x_i)) \leq \frac{e^{q/2-1} \pi^{q/2}}{A_\Sigma^{q/2}} + H_{s_0},$$

and note that $\widehat{s}_{\widehat{m}} \in S$, thus

$$\sqrt{\mathbb{E} [\text{KL}_n^2(s_0, \widehat{s}_{\widehat{m}})]} \leq \frac{e^{q/2-1} \pi^{q/2}}{A_\Sigma^{q/2}} + H_{s_0}. \quad (60)$$

We now provide an upper bound for $\mathbb{P}(\mathcal{T}^C)$:

$$\mathbb{P}(\mathcal{T}^C) \leq \sum_{i=1}^n \mathbb{P}(\|Y_i\|_\infty > M_n). \quad (61)$$

For all $i \in [n]$,

$$Y_i|x_i \sim \sum_{k=1}^K g_k(x_i; \gamma) \mathcal{N}_q(\beta_{k0} + \beta_k x_i, \Sigma_k),$$

so we see from (61) that we need to provide an upper bound on $\mathbb{P}(|Y_x| > M_n)$, with

$$Y_x \sim \sum_{k=1}^K g_k(x; \gamma) \mathcal{N}_q(\beta_{k0} + \beta_k x, \Sigma_k), x \in \mathcal{X}.$$

First, using Chernoff's inequality for a centered Gaussian variable (see Lemma 23), and the fact that ψ belongs to the bounded space $\tilde{\Psi}$ (defined by (4)), and that $\sum_{k=1}^K g_k(x; \gamma) = 1$, we get

$$\begin{aligned}
 & \mathbb{P}(\|Y_x\|_\infty > M_n) \\
 &= \sum_{k=1}^K \frac{g_k(x; \gamma)}{(2\pi)^{q/2} \det(\Sigma_k)^{1/2}} \int_{\{\|y\|_\infty > M_n\}} \exp\left(-\frac{(y - (\beta_{k0} + \beta_k x))^\top \Sigma_k^{-1} (y - (\beta_{k0} + \beta_k x))}{2}\right) dy \\
 &= \sum_{k=1}^K g_k(x; \gamma) \mathbb{P}(\|Y_{x,k}\|_\infty > M_n) \leq \sum_{k=1}^K g_k(x; \gamma) \sum_{z=1}^q \mathbb{P}(|[Y_{x,k}]_z| > M_n) \\
 &= \sum_{k=1}^K g_k(x; \gamma) \sum_{z=1}^q (\mathbb{P}([Y_{x,k}]_z < -M_n) + \mathbb{P}([Y_{x,k}]_z > M_n)) \\
 &= \sum_{k=1}^K g_k(x; \gamma) \sum_{z=1}^q \left(\mathbb{P}\left(U > \frac{M_n - [\beta_{k0} + \beta_k x]_z}{[\Sigma_k]_{z,z}^{1/2}}\right) + \mathbb{P}\left(U < \frac{-M_n - [\beta_{k0} + \beta_k x]_z}{[\Sigma_k]_{z,z}^{1/2}}\right) \right) \\
 &= \sum_{k=1}^K g_k(x; \gamma) \sum_{z=1}^q \left(\mathbb{P}\left(U > \frac{M_n - [\beta_{k0} + \beta_k x]_z}{[\Sigma_k]_{z,z}^{1/2}}\right) + \mathbb{P}\left(U > \frac{M_n + [\beta_{k0} + \beta_k x]_z}{[\Sigma_k]_{z,z}^{1/2}}\right) \right) \\
 &\leq \sum_{k=1}^K g_k(x; \gamma) \sum_{z=1}^q \left[e^{-\frac{1}{2} \left(\frac{M_n - [\beta_{k0} + \beta_k x]_z}{[\Sigma_k]_{z,z}^{1/2}} \right)^2} + e^{-\frac{1}{2} \left(\frac{M_n + [\beta_{k0} + \beta_k x]_z}{[\Sigma_k]_{z,z}^{1/2}} \right)^2} \right] \text{ (using Lemma 23, (113))} \\
 &\leq 2 \sum_{k=1}^K g_k(x; \gamma) \sum_{z=1}^q e^{-\frac{1}{2} \left(\frac{M_n - |[\beta_{k0} + \beta_k x]_z|}{[\Sigma_k]_{z,z}^{1/2}} \right)^2} \leq 2 \sum_{k=1}^K g_k(x; \gamma) \sum_{z=1}^q e^{-\frac{1}{2} \frac{M_n^2 - 2M_n |[\beta_{k0} + \beta_k x]_z| + |[\beta_{k0} + \beta_k x]_z|^2}{[\Sigma_k]_{z,z}}} \\
 &\leq 2KA_\gamma q e^{-\frac{M_n^2 - 2M_n A_\beta}{2A_\Sigma}}, \tag{62}
 \end{aligned}$$

where

$$Y_{x,k} \sim \mathcal{N}_q(\beta_{k0} + \beta_k x, \Sigma_k), [Y_{x,k}]_z \sim \mathcal{N}([\beta_{k0} + \beta_k x]_z, [\Sigma_k]_{z,z}), \text{ and } U = \frac{[Y_{x,k}]_z - [\beta_k x]_z}{[\Sigma_k]_{z,z}^{1/2}} \sim \mathcal{N}(0, 1),$$

and using the facts that $e^{-\frac{1}{2} \frac{|[\beta_{k0} + \beta_k x]_z|^2}{A_\Sigma}} \leq 1$ and $\max_{1 \leq z \leq q} |[\Sigma_k]_{z,z}| \leq \|\Sigma_k\|_2 = M(\Sigma_k) = m(\Sigma_k^{-1}) \leq A_\Sigma$. We derive from (61) and (62) that

$$\mathbb{P}(\mathcal{T}^c) \leq 2KnqA_\gamma e^{-\frac{M_n^2 - 2M_n A_\beta}{2A_\Sigma}}, \tag{63}$$

and finally from (55), (60), and (63), we obtain

$$\mathbb{E}[\text{KL}_n(s_0, \hat{s}_{\hat{m}}) \mathbb{1}_{\mathcal{T}^c}] \leq \left(\frac{e^{q/2-1} \pi^{q/2}}{A_\Sigma^{q/2}} + H_{s_0} \right) \sqrt{2KnqA_\gamma} e^{-\frac{M_n^2 - 2M_n A_\beta}{4A_\Sigma}}. \tag{64}$$

Appendix B. Proofs of technical lemmas

B.1. Proof of Lemma 5

Let $m \in \mathbb{N}^*$, on the event $\mathcal{T}$, to control the deviation

$$\sup_{f_m \in F_m} |\nu_n(-f_m)| = \sup_{f_m \in F_m} \left| \frac{1}{n} \sum_{i=1}^n \{f_m(Y_i|x_i) - \mathbb{E}[f_m(Y_i|x_i)]\} \right|, \quad (65)$$

we shall use concentration and symmetrization arguments. We shall first use the following concentration inequality, Lemma 7, which is an adaption of [Wainwright \(2019, Theorem 4.10\)](#).

Lemma 7 (Theorem 4.10 from [Wainwright \(2019\)](#)) *Let $Z_1, \dots, Z_n$ be independent random variables with values in some space $\mathcal{Z}$ and let $\mathcal{F}$ be a class of integrable real-valued functions with domain on $\mathcal{Z}$. Assume that*

$$\sup_{f \in \mathcal{F}} \|f\|_\infty \leq R_n \quad \text{for some non-random constants } R_n < \infty. \quad (66)$$

Then, for all $t > 0$,

$$\mathbb{P} \left(\sup_{f \in \mathcal{F}} \left| \frac{1}{n} \sum_{i=1}^n [f(Z_i) - \mathbb{E}[f(Z_i)]] \right| > \mathbb{E} \left[\sup_{f \in \mathcal{F}} \left| \frac{1}{n} \sum_{i=1}^n [f(Z_i) - \mathbb{E}[f(Z_i)]] \right| \right] + 2\sqrt{2}R_n \sqrt{\frac{t}{n}} \right) \leq e^{-t}. \quad (67)$$

That is, with probability greater than $1 - e^{-t}$,

$$\sup_{f \in \mathcal{F}} \left| \frac{1}{n} \sum_{i=1}^n [f(Z_i) - \mathbb{E}[f(Z_i)]] \right| \leq \mathbb{E} \left[\sup_{f \in \mathcal{F}} \left| \frac{1}{n} \sum_{i=1}^n [f(Z_i) - \mathbb{E}[f(Z_i)]] \right| \right] + 2\sqrt{2}R_n \sqrt{\frac{t}{n}} \quad (68)$$

Then, we propose to bound $\mathbb{E} \left[\sup_{f \in \mathcal{F}} \left| \frac{1}{n} \sum_{i=1}^n [f(Z_i) - \mathbb{E}[f(Z_i)]] \right| \right]$ due to the following symmetrization argument. The proof of this result can be found in [Van Der Vaart and Wellner \(1996\)](#).

Lemma 8 (See Lemma 2.3.6 in [Van Der Vaart and Wellner \(1996\)](#)) *Let $Z_1, \dots, Z_n$ be independent random variables with values in some space $\mathcal{Z}$ and let $\mathcal{F}$ be a class of real-valued functions on $\mathcal{Z}$. Let $(\epsilon_1, \dots, \epsilon_n)$ be a Rademacher sequence independent of $(Z_1, \dots, Z_n)$. Then,*

$$\mathbb{E} \left[\sup_{f \in \mathcal{F}} \left| \frac{1}{n} \sum_{i=1}^n [f(Z_i) - \mathbb{E}[f(Z_i)]] \right| \right] \leq 2\mathbb{E} \left[\sup_{f \in \mathcal{F}} \left| \frac{1}{n} \sum_{i=1}^n \epsilon_i f(Z_i) \right| \right]. \quad (69)$$

From (69), the problem is to provide an upper bound on

$$\mathbb{E} \left[\sup_{f \in \mathcal{F}} \left| \frac{1}{n} \sum_{i=1}^n \epsilon_i f(Z_i) \right| \right].$$

To do so, we shall apply the following lemma, which is adapted from Lemma 6.1 in [Massart \(2007\)](#).

Lemma 9 (See Lemma 6.1 in Massart (2007)) *Let $Z_1, \dots, Z_n$ be independent random variables with values in some space $\mathcal{Z}$ and let $\mathcal{F}$ be a class of real-valued functions on $\mathcal{Z}$. Let $(\epsilon_1, \dots, \epsilon_n)$ be a Rademacher sequence, independent of $(Z_1, \dots, Z_n)$. Define R_n , a non-random constant, such that*

$$\sup_{f \in \mathcal{F}} \|f\|_n \leq R_n. \quad (70)$$

Then, for all $S \in \mathbb{N}^*$,

$$\mathbb{E} \left[\sup_{f \in \mathcal{F}} \left| \frac{1}{n} \sum_{i=1}^n \epsilon_i f(Z_i) \right| \right] \leq R_n \left(\frac{6}{\sqrt{n}} \sum_{s=1}^S 2^{-s} \sqrt{\ln [1 + M(2^{-s} R_n, \mathcal{F}, \|\cdot\|_n)]} + 2^{-S} \right), \quad (71)$$

where $M(\delta, \mathcal{F}, \|\cdot\|_n)$ stands for the δ -packing number (see Definition 20) of the set of functions $\mathcal{F}$, equipped with the metric induced by the norm $\|\cdot\|_n$.

We are now able to prove Lemma 5. Indeed, given any fixed values $x_1, \dots, x_n \in \mathcal{X}$, in order to control $\sup_{f_m \in F_m} |\nu_n(-f_m)| \mathbb{1}_{\mathcal{T}}$ from the (65), we would like to apply Lemmas 7–9. On the one hand, we see from (70) that we need an upper bound of $\sup_{f_m \in F_m} \|f_m\|_\infty \mathbb{1}_{\mathcal{T}}$. On the other hand, we see from (71) that on the event $\mathcal{T}$, we need to bound the entropy of the set of functions F_m , equipped with the metric induced by the norm $\|\cdot\|_n$. Such bounds are provided by the two following lemmas.

Recall that given $M_n > 0$, we considered the event

$$\mathcal{T} = \left\{ \max_{i \in [n]} \|Y_i\|_\infty = \max_{i \in [n]} \max_{z \in [q]} |[Y_i]_z| \leq M_n \right\}, \quad (72)$$

$$\text{let } B_n = \max(A_\Sigma, 1 + K A_G) \left(1 + q\sqrt{q} (M_n + A_\beta)^2 A_\Sigma \right).$$

Lemma 10 *On the event $\mathcal{T}$, for all $m \in \mathbb{N}^*$,*

$$\sup_{f_m \in F_m} \|f_m\|_\infty \leq 2K B_n \left(A_\gamma + q A_\beta + \frac{q\sqrt{q}}{a_\Sigma} \right) =: R_n. \quad (73)$$

Proof See Appendix B.2.1. ■

Lemma 11 *Let $\delta > 0$ and $m \in \mathbb{N}^*$. On the event $\mathcal{T}$, we have the following upper bound of the δ -packing number of the set of functions F_m , equipped with the metric induced by the norm $\|\cdot\|_n$:*

$$M(\delta, F_m, \|\cdot\|_n) \leq (2p+1)^{\frac{72B_n^2 q^2 K^2 m^2}{\delta^2}} \left(1 + \frac{18B_n K q A_\beta}{\delta} \right)^K \left(1 + \frac{18B_n K A_\gamma}{\delta} \right)^K \left(1 + \frac{18B_n K q \sqrt{q}}{a_\Sigma \delta} \right)^K.$$

Proof See Appendix B.2.2. ■

Lemma 12 (Lemma 5.9 from [Meynet \(2013\)](#)) *Let $\delta > 0$ and $(x_{ij})_{i \in [n]; j=1, \dots, p} \in \mathbb{R}^{np}$. There exists a family $\mathcal{B}$ of $(2p+1)^{\|x\|_{\max, n}^2 / \delta^2}$ vectors in $\mathbb{R}^p$, such that for all $\beta \in \mathbb{R}^p$, with $\|\beta\|_1 \leq 1$, where $\|x\|_{\max, n}^2 = \frac{1}{n} \sum_{i=1}^n \max_{j \in \{1, \dots, p\}} x_{ij}^2$, there exists $\beta' \in \mathcal{B}$, such that*

$$\frac{1}{n} \sum_{i=1}^n \left(\sum_{j=1}^p (\beta_j - \beta'_j) x_{ij} \right)^2 \leq \delta^2.$$

Proof See in the proof of [Meynet \(2013, Lemma 5.9\)](#). ■

Via the upper bounds provided in Lemmas [10](#) and [11](#), we can apply Lemma [9](#) to get an upper bound of

$$\mathbb{E} \left[\sup_{f_m \in \mathcal{F}_m} \left| \frac{1}{n} \sum_{i=1}^n \epsilon_i f_m(Y_i | x_i) \right| \right] \text{ on the event } \mathcal{T}. \quad (74)$$

In order to provide such an upper bound, Lemmas [7](#) and [9](#) can be utilized via defining a suitable class of integrable real-valued functions as follows:

$$\mathcal{F} := \{f := f_m \mathbb{1}_{|f_m| \leq R_n} : f_m \in F_m, \}, \quad Z_i := Y_i | x_i, \forall i \in [n]. \quad (75)$$

Indeed, by definition, it holds that

$$\sup_{f \in \mathcal{F}} \|f\|_n \leq \sup_{f \in \mathcal{F}} \|f\|_\infty = \sup_{f \in \mathcal{F}} \sup_{z \in \mathcal{Z}} |f(z)| = \sup_{f_m \in F_m} \sup_{z \in \mathcal{Z}} |f_m(z) \mathbb{1}_{|f_m(z)| \leq R_n}| \leq R_n. \quad (76)$$

Note that the last inequality is valid since if $|f_m(z)| \leq R_n$ then $|f_m(z) \mathbb{1}_{|f_m(z)| \leq R_n}| = |f_m(z)| \leq R_n$. Otherwise, if $|f_m(z)| > R_n$, then $|f_m(z) \mathbb{1}_{|f_m(z)| \leq R_n}| = |f_m(z) \times 0| = 0 \leq R_n$. We thus obtain the following results.

Lemma 13 *Let $m \in \mathbb{N}^*$, consider $(\epsilon_1, \dots, \epsilon_n)$, a Rademacher sequence independent of $(Y_1, \dots, Y_n)$. Then, on the event $\mathcal{T}$, it holds that*

$$\mathbb{E} \left[\sup_{f_m \in \mathcal{F}_m} \left| \frac{1}{n} \sum_{i=1}^n \epsilon_i f_m(Y_i | x_i) \right| \right] \leq \frac{74KB_n q}{\sqrt{n}} \Delta_m, \text{ where} \\ \Delta_m := m \sqrt{\ln(2p+1) \ln n} + 2\sqrt{K} \left(A_\gamma + qA_\beta + \frac{q\sqrt{q}}{a_\Sigma} \right). \quad (77)$$

Proof See Appendix [B.2.3](#). ■

We now return to the proof of the Lemma [5](#).

Finally, on the event $\mathcal{T}$, using (75) and Lemma 10, for all $m \in \mathbb{N}^*$ and $t > 0$, with probability greater than $1 - e^{-t}$, we obtain

$$\sup_{f_m \in F_m} |\nu_n(-f_m)| = \sup_{f \in \mathcal{F}} \left| \frac{1}{n} \sum_{i=1}^n \{f(Z_i) - \mathbb{E}[f(Z_i)]\} \right| \quad (78)$$

$$\leq \mathbb{E} \left[\sup_{f \in \mathcal{F}} \left| \frac{1}{n} \sum_{i=1}^n [f(Z_i) - \mathbb{E}[f(Z_i)]] \right| \right] + 2\sqrt{2}R_n \sqrt{\frac{t}{n}} \text{ (using Lemma 7)} \quad (79)$$

$$\leq \mathbb{E} \left[\sup_{f \in \mathcal{F}} \left| \frac{1}{n} \sum_{i=1}^n [f(Z_i) - \mathbb{E}[f(Z_i)]] \right| \right] + 2\sqrt{2}R_n \sqrt{\frac{t}{n}} \text{ (since } \leq 1) \quad (80)$$

$$\leq 2\mathbb{E} \left[\sup_{f \in \mathcal{F}} \left| \frac{1}{n} \sum_{i=1}^n \epsilon_i f(Z_i) \right| \right] + 2\sqrt{2}R_n \sqrt{\frac{t}{n}} \text{ (Lemma 8)} \quad (81)$$

$$= 2\mathbb{E} \left[\sup_{f_m \in \mathcal{F}_m} \left| \frac{1}{n} \sum_{i=1}^n \epsilon_i f_m(Y_i|x_i) \right| \right] + 2\sqrt{2}R_n \sqrt{\frac{t}{n}} \quad (82)$$

$$\leq \frac{148KB_nq}{\sqrt{n}} \Delta_m + 4\sqrt{2}KB_n \left(A_\gamma + qA_\beta + \frac{q\sqrt{q}}{a_\Sigma} \right) \sqrt{\frac{t}{n}} \quad (83)$$

$$\left(\text{using Lemma 13 and } R_n = 2KB_n \left(A_\gamma + qA_\beta + \frac{q\sqrt{q}}{a_\Sigma} \right) \right) \quad (84)$$

$$\leq \frac{4KB_n}{\sqrt{n}} \left[37q\Delta_m + \sqrt{2} \left(A_\gamma + qA_\beta + \frac{q\sqrt{q}}{a_\Sigma} \right) \sqrt{t} \right]. \quad (85)$$

B.2. Proofs of Lemmas 10–13

The proofs of Lemmas 10–11 require an upper bound on the uniform norm of the gradient of $\ln s_\psi$, for $s_\psi \in S$. We begin by providing such an upper bound.

Lemma 14 *Given s_ψ , as described in (5), it holds that*

$$\sup_{x \in \mathcal{X}} \sup_{\psi \in \tilde{\Psi}} \left\| \frac{\partial \ln(s_\psi(\cdot|x))}{\partial \psi} \right\|_\infty \leq G(\cdot),$$

$$G : \mathbb{R}^q \ni y \mapsto G(y) = \max(A_\Sigma, 1 + KA_G) \left(1 + q\sqrt{q}(\|y\|_\infty + A_\beta)^2 A_\Sigma \right). \quad (86)$$

Proof Let $s_\psi \in S$, with $\psi = (\gamma, \beta, \Sigma)$. From now on, we consider any $x \in \mathcal{X}$, any $y \in \mathbb{R}^q$, and any $k \in [K]$. We can write

$$\begin{aligned} \ln(s_\psi(y|x)) &= \ln\left(\sum_{k=1}^K g_k(x; \gamma) \phi(y; \beta_{k0} + \beta_k x, \Sigma_k)\right) = \ln\left(\sum_{k=1}^K f_k(x, y)\right), \\ g_k(x; \gamma) &= \frac{\exp(w_k(x))}{\sum_{l=1}^K \exp(w_l(x))}, w_k(x) = \gamma_{k0} + \gamma_k^\top x, \\ \phi(y; \beta_{k0} + \beta_k x, \Sigma_k) &= \frac{1}{(2\pi)^{q/2} \det(\Sigma_k)^{1/2}} \exp\left(-\frac{(y - (\beta_{k0} + \beta_k x))^\top \Sigma_k^{-1} (y - (\beta_{k0} + \beta_k x))}{2}\right), \\ f_k(x, y) &= g_k(x; \gamma) \phi(y; \beta_{k0} + \beta_k x, \Sigma_k) \\ &= \frac{g_k(x; \gamma)}{(2\pi)^{q/2} \det(\Sigma_k)^{1/2}} \exp\left[-\frac{1}{2} (y - (\beta_{k0} + \beta_k x))^\top \Sigma_k^{-1} (y - (\beta_{k0} + \beta_k x))\right]. \end{aligned}$$

By using the chain rule, for all $l \in [K]$,

$$\begin{aligned} \frac{\partial \ln(s_\psi(y|x))}{\partial \gamma_{l0}} &= \sum_{k=1}^K \frac{f_k(x, y)}{g_k(x; \gamma) \sum_{k=1}^K f_k(x, y)} \frac{\partial g_k(x; \gamma)}{\partial w_l(x)} \underbrace{\frac{\partial w_l(x)}{\partial \gamma_{l0}}}_{=1}, \text{ and} \\ \frac{\partial \ln(s_\psi(y|x))}{\partial (\gamma_l^\top x)} &= \sum_{k=1}^K \frac{f_k(x, y)}{g_k(x; \gamma) \sum_{k=1}^K f_k(x, y)} \frac{\partial g_k(x; \gamma)}{\partial w_l(x)} \underbrace{\frac{\partial w_l(x)}{\partial (\gamma_l^\top x)}}_{=1}. \end{aligned}$$

Furthermore,

$$\begin{aligned} \frac{\partial g_k(x; \gamma)}{\partial w_l(x)} &= \frac{\partial}{\partial w_l(x)} \left(\frac{\exp(w_k(x))}{\sum_{l=1}^K \exp(w_l(x))} \right) \\ &= \frac{\delta_{lk} \exp(w_k(x))}{\sum_{l=1}^K \exp(w_l(x))} - \frac{\exp(w_k(x))}{\sum_{l=1}^K \exp(w_l(x))} \frac{\exp(w_l(x))}{\sum_{l=1}^K \exp(w_l(x))} = g_k(x; \gamma) (\delta_{lk} - g_l(x; \gamma)), \\ \text{where } \delta_{lk} &= \begin{cases} 1 & \text{if } l = k, \\ 0 & \text{if } l \neq k. \end{cases} \end{aligned}$$

Therefore, we obtain

$$\begin{aligned} \left| \frac{\partial \ln(s_\psi(y|x))}{\partial (\gamma_l^\top x)} \right| &= \left| \frac{\partial \ln(s_\psi(y|x))}{\partial \gamma_{l0}} \right| = \left| \sum_{k=1}^K \frac{f_k(x, y)}{g_k(x; \gamma) \sum_{k=1}^K f_k(x, y)} g_k(x; \gamma) (\delta_{lk} - g_l(x; \gamma)) \right| \\ &= \left| \sum_{k=1}^K \frac{f_k(x, y)}{\sum_{k=1}^K f_k(x, y)} (\delta_{lk} - g_l(x; \gamma)) \right| \leq \left| \sum_{k=1}^K (\delta_{lk} - g_l(x; \gamma)) \right| \\ &= \left| 1 - \sum_{k=1}^K g_l(x; \gamma) \right| = |1 - K g_l(x; \gamma)| \leq 1 + K g_l(x; \gamma) \leq 1 + K A_G \text{ (using (57))}. \end{aligned}$$

Similarly, by using the fact that ψ belongs to the bounded space $\tilde{\Psi}$, $f_l(x, y) / \sum_{k=1}^K f_k(x, y) \leq 1$,

$$\begin{aligned}
 & \left\| \frac{\partial \ln(s_\psi(y|x))}{\partial \beta_{l0}} \right\|_\infty = \left\| \frac{\partial \ln(s_\psi(y|x))}{\partial (\beta_l x)} \right\|_\infty \\
 &= \left\| \frac{f_l(x, y)}{\sum_{k=1}^K f_k(x, y)} \frac{\partial}{\partial (\beta_{l0} + \beta_l x)} \left[-\frac{1}{2} (y - (\beta_{l0} + \beta_l x))^\top \Sigma_l^{-1} (y - (\beta_{l0} + \beta_l x)) \right] \right\|_\infty \\
 &\leq \left\| \frac{\partial}{\partial (\beta_{l0} + \beta_l x)} \left[-\frac{1}{2} (y - (\beta_{l0} + \beta_l x))^\top \Sigma_l^{-1} (y - (\beta_{l0} + \beta_l x)) \right] \right\|_\infty \\
 &= \left\| \Sigma_l^{-1} (y - (\beta_{l0} + \beta_l x)) \right\|_\infty \leq \left\| \Sigma_l^{-1} \right\|_\infty \|y - (\beta_{l0} + \beta_l x)\|_\infty \text{ (using (103))} \\
 &\leq \sqrt{q} \left\| \Sigma_l^{-1} \right\|_2 (\|y\|_\infty + \|\beta_{l0} + \beta_l x\|_\infty) \text{ (using (108))} \\
 &\leq \sqrt{q} M (\Sigma_l^{-1}) (\|y\|_\infty + \|\beta_{l0} + \beta_l x\|_\infty) \text{ (using (107))} \\
 &\leq \sqrt{q} A_\Sigma (\|y\|_\infty + A_\beta) \text{ (using (4))}.
 \end{aligned}$$

Now, we need to calculate the gradient w.r.t. to the covariance matrices of the Gaussian experts. To do this, we need the following result: given any $l \in [K]$, $v_l = \beta_{l0} + \beta_l x$, it holds that

$$\frac{\partial}{\partial \Sigma_l} \phi(x; v_l, \Sigma_l) = \phi(x; v_l, \Sigma_l) \underbrace{\frac{1}{2} \left[\Sigma_l^{-1} (x - v_l) (x - v_l)^\top \Sigma_l^{-1} - (\Sigma_l^{-1})^\top \right]}_{T(x, v_l, \Sigma_l)}, \quad (87)$$

noting that

$$\frac{\partial}{\partial \Sigma_l} \left((x - v_l)^\top \Sigma_l^{-1} (x - v_l) \right) = -\Sigma_l^{-1} (x - v_l) (x - v_l)^\top \Sigma_l^{-1} \text{ (using Lemma 16)}, \quad (88)$$

$$\frac{\partial}{\partial \Sigma_l} (\det(\Sigma_l)) = \det(\Sigma_l) (\Sigma_l^{-1})^\top \text{ (using Jacobi formula, Lemma 17)}. \quad (89)$$

For any $l \in [K]$,

$$\begin{aligned}
 & \left| \frac{\partial \ln(s_\psi(y|x))}{\partial ([\Sigma_l]_{z_1, z_2})} \right| \leq \left\| \frac{\partial \ln(s_\psi(y|x))}{\partial \Sigma_l} \right\|_2 \text{ (using (107))} \\
 &= \left\| \frac{f_l(x, y)}{\sum_{k=1}^K f_k(x, y)} \left\| \frac{\partial}{\partial \Sigma_l} \left[-\frac{1}{2} (y - (\beta_{l0} + \beta_l x))^\top \Sigma_l^{-1} (y - (\beta_{l0} + \beta_l x)) \right] \right\|_2 \right\|_2 \\
 &\leq \left\| \frac{\partial}{\partial \Sigma_l} \left[-\frac{1}{2} (y - (\beta_{l0} + \beta_l x))^\top \Sigma_l^{-1} (y - (\beta_{l0} + \beta_l x)) \right] \right\|_2 \\
 &= \frac{1}{2} \left\| \Sigma_l^{-1} (y - (\beta_{l0} + \beta_l x)) (y - (\beta_{l0} + \beta_l x))^\top \Sigma_l^{-1} - (\Sigma_l^{-1})^\top \right\|_2 \text{ (using (87))} \\
 &\leq \frac{1}{2} \left[A_\Sigma + \sqrt{q} \left\| (y - (\beta_{l0} + \beta_l x)) (y - (\beta_{l0} + \beta_l x))^\top \right\|_\infty A_\Sigma^2 \right] \text{ (using (108))} \\
 &\leq \frac{1}{2} \left[A_\Sigma + q\sqrt{q} (\|y\|_\infty + A_\beta)^2 A_\Sigma^2 \right] \text{ (using (4))},
 \end{aligned}$$

where, in the last inequality given $a = y - (\beta_{l0} + \beta_l x)$, we use the fact that

$$\left\| aa^\top \right\|_\infty = \max_{1 \leq i \leq q} \sum_{j=1}^q |a_i a_j| = \max_{1 \leq i \leq q} \sum_{j=1}^q |a_i a_j| = \max_{1 \leq i \leq q} |a_i| \sum_{j=1}^q |a_j| \leq q \|a\|_\infty^2.$$

Thus,

$$\begin{aligned}
 & \sup_{x \in \mathcal{X}} \sup_{\psi \in \tilde{\Psi}} \left\| \frac{\partial \ln(s_\psi(y|x))}{\partial \psi} \right\|_\infty \\
 & \leq \max \left[1 + K A_G, \sqrt{q} (\|y\|_\infty + A_\beta) A_\Sigma, \frac{1}{2} \left[A_\Sigma + q \sqrt{q} (\|y\|_\infty + A_\beta)^2 A_\Sigma^2 \right] \right] \\
 & \leq \max \left[1 + K A_G, \max(A_\Sigma, 1) \left(1 + q \sqrt{q} (\|y\|_\infty + A_\beta)^2 A_\Sigma \right) \right] \\
 & \leq \max(A_\Sigma, 1 + K A_G) \left(1 + q \sqrt{q} (\|y\|_\infty + A_\beta)^2 A_\Sigma \right) =: G(y),
 \end{aligned}$$

where we use the fact that

$$\begin{aligned}
 \sqrt{q} (\|y\|_\infty + A_\beta) A_\Sigma & =: \theta \leq 1 + \theta^2 = 1 + q (\|y\|_\infty + A_\beta)^2 A_\Sigma^2 \\
 & \leq \max(A_\Sigma, 1) \left(1 + q \sqrt{q} (\|y\|_\infty + A_\beta)^2 A_\Sigma \right).
 \end{aligned}$$

■

B.2.1. PROOF OF LEMMA 10

Let $m \in \mathbb{N}^*$ and $f_m \in F_m$. By (28), there exists $s_m \in S_m$, such that $f_m = -\ln(s_m/s_0)$. For all $x \in \mathcal{X}$, let $\psi(x) = (\gamma_{k0}, \gamma_k^\top x, \beta_{k0}, \beta_k x, \Sigma_k)_{k \in [K]}$ be the parameters of $s_m(\cdot|x)$. In our case, we approximate $f(\psi) = \ln(s_\psi(y_i|x_i))$ around $\psi_0(x_i)$ by the $n = 0^{th}$ degree Taylor polynomial of $f(\psi)$. That is,

$$\begin{aligned}
 |\ln(s_m(y_i|x_i)) - \ln(s_0(y_i|x_i))| & =: |f(\psi) - f(\psi_0)| = |R_0(\psi)| \text{ (defined in Lemma 24)} \\
 & \leq \sup_{x \in \mathcal{X}} \sup_{\psi \in \tilde{\Psi}} \left\| \frac{\partial \ln(s_\psi(y_i|x))}{\partial \psi} \right\|_\infty \|\psi(x_i) - \psi_0(x_i)\|_1.
 \end{aligned}$$

First applying Taylor's inequality and then Lemma 14 on the event $\mathcal{T}$. For all $i \in [n]$, it holds that

$$\begin{aligned}
 |f_m(y_i|x_i)| \mathbb{1}_{\mathcal{T}} & = |\ln(s_m(y_i|x_i)) - \ln(s_0(y_i|x_i))| \mathbb{1}_{\mathcal{T}} \leq \sup_{x \in \mathcal{X}} \sup_{\psi \in \tilde{\Psi}} \left\| \frac{\partial \ln(s_\psi(y_i|x))}{\partial \psi} \right\|_\infty \|\psi(x_i) - \psi_0(x_i)\|_1 \mathbb{1}_{\mathcal{T}} \\
 & \leq \underbrace{\max(A_\Sigma, 1 + K A_G) \left(1 + q \sqrt{q} (M_n + A_\beta)^2 A_\Sigma \right)}_{=: B_n} \|\psi(x_i) - \psi_0(x_i)\|_1 \text{ (using Lemma 14)} \\
 & \leq B_n \sum_{k=1}^K \left(|\gamma_{k0} - \gamma_{0,k0}| + \left| \gamma_k^\top x_i - \gamma_{0,k}^\top x_i \right| + \|\beta_{k0} - \beta_{0,k0}\|_1 + \|\beta_k x_i - \beta_{0,k} x_i\|_1 + \|\text{vec}(\Sigma_k - \Sigma_{0,k})\|_1 \right) \\
 & \leq 2B_n \sum_{k=1}^K \left(|\gamma_{k0}| + \left| \gamma_k^\top x_i \right| + \|\beta_{k0}\|_1 + \|\beta_k x_i\|_1 + q \|\Sigma_k\|_1 \right) \text{ (using (105))} \\
 & \leq 2KB_n (A_\gamma + q \|\beta_{k0}\|_\infty + q \|\beta_k x_i\|_\infty + q \sqrt{q} \|\Sigma_k\|_2) \text{ (using (4), (100), (101), (109))} \\
 & \leq 2KB_n \left(A_\gamma + q A_\beta + \frac{q \sqrt{q}}{a_\Sigma} \right) \text{ (using (4))}.
 \end{aligned}$$

Therefore, on the event $\mathcal{T}$,

$$\sup_{f_m \in F_m} \|f_m\|_\infty \leq 2KB_n \left(A_\gamma + qA_\beta + \frac{q\sqrt{q}}{a_\Sigma} \right) =: R_n.$$

B.2.2. PROOF OF LEMMA 11

Let $m \in \mathbb{N}^*$, $f_m^{[1]} \in F_m$, and $x \in [0, 1]^p$. By (28), there exists $s_m^{[1]} \in S_m$, such that $f_m^{[1]} = -\ln(s_m^{[1]}/s_0)$. Introduce the notation $s_m^{[2]} \in S$ and $f_m^{[2]} = -\ln(s_m^{[2]}/s_0)$. Let

$$\psi^{[1]}(x) = \left(\gamma_{k0}^{[1]}, \gamma_k^{[1]} x, \beta_{k0}^{[1]}, \beta_k^{[1]} x, \Sigma_k^{[1]} \right)_{k \in [K]}, \text{ and } \psi^{[2]}(x) = \left(\gamma_{k0}^{[2]}, \gamma_k^{[2]} x, \beta_{k0}^{[2]}, \beta_k^{[2]} x, \Sigma_k^{[2]} \right)_{k \in [K]},$$

be the parameters of the PDFs $s_m^{[1]}(\cdot|x)$ and $s_m^{[2]}(\cdot|x)$, respectively. By applying Taylor's inequality and then Lemma 14 on the event $\mathcal{T}$, for all $i \in [n]$, it holds that

$$\begin{aligned} & \left| f_m^{[1]}(y_i|x_i) - f_m^{[2]}(y_i|x_i) \right| = \left| \ln(s_m^{[1]}(y_i|x_i)) - \ln(s_m^{[2]}(y_i|x_i)) \right| \\ & \leq \sup_{x \in \mathcal{X}} \sup_{\psi \in \tilde{\Psi}} \left| \frac{\partial \ln(s_\psi(y_i|x))}{\partial \psi} \right| \left\| \psi^{[1]}(x_i) - \psi^{[2]}(x_i) \right\|_1 \text{ (using Taylor's inequality in Lemma 24)} \\ & \leq \underbrace{\max(A_\Sigma, C(p, K)) \left(1 + q\sqrt{q}(M_n + A_\beta)^2 A_\Sigma \right)}_{B_n} \left\| \psi^{[1]}(x_i) - \psi^{[2]}(x_i) \right\|_1 \text{ (using Lemma 14)} \\ & \leq B_n \sum_{k=1}^K \left(\left| \gamma_{k0}^{[1]} - \gamma_{k0}^{[2]} \right| + \left| \gamma_k^{[1]\top} x_i - \gamma_k^{[2]\top} x_i \right| \right. \\ & \quad \left. + \left\| \beta_{k0}^{[1]} - \beta_{k0}^{[2]} \right\|_1 + \left\| \beta_k^{[1]} x_i - \beta_k^{[2]} x_i \right\|_1 + \left\| \text{vec}(\Sigma_k^{[1]} - \Sigma_k^{[2]}) \right\|_1 \right). \end{aligned}$$

By the Cauchy-Schwarz inequality, $(\sum_{i=1}^m a_i)^2 \leq m \sum_{i=1}^m a_i^2$ ($m \in \mathbb{N}^*$), we get

$$\begin{aligned} & \left| f_m^{[1]}(y_i|x_i) - f_m^{[2]}(y_i|x_i) \right|^2 \\ & \leq 3B_n^2 \left[\left(\sum_{k=1}^K \left| \gamma_k^{[1]\top} x_i - \gamma_k^{[2]\top} x_i \right| \right)^2 + \left(\sum_{k=1}^K \sum_{z=1}^q \left| [\beta_k^{[1]} x_i]_z - [\beta_k^{[2]} x_i]_z \right| \right)^2 \right] \\ & \quad + 3B_n^2 \left(\left\| \beta_0^{[1]} - \beta_0^{[2]} \right\|_1 + \left\| \gamma_0^{[1]} - \gamma_0^{[2]} \right\|_1 + \left\| \text{vec}(\Sigma^{[1]} - \Sigma^{[2]}) \right\|_1 \right)^2 \\ & \leq 3B_n^2 \left[K \sum_{k=1}^K \left(\sum_{j=1}^p \gamma_{kj}^{[1]\top} x_{ij} - \sum_{j=1}^p \gamma_{kj}^{[2]\top} x_{ij} \right)^2 + Kq \sum_{k=1}^K \sum_{z=1}^q \left(\sum_{j=1}^p [\beta_k^{[1]}]_{z,j} x_{ij} - \sum_{j=1}^p [\beta_k^{[2]}]_{z,j} x_{ij} \right)^2 \right] \\ & \quad + 3B_n^2 \left(\left\| \beta_0^{[1]} - \beta_0^{[2]} \right\|_1 + \left\| \gamma_0^{[1]} - \gamma_0^{[2]} \right\|_1 + \left\| \text{vec}(\Sigma^{[1]} - \Sigma^{[2]}) \right\|_1 \right)^2, \end{aligned}$$

and

$$\begin{aligned}
 \|f_m^{[1]} - f_m^{[2]}\|_n^2 &= \frac{1}{n} \sum_{i=1}^n \left| f_m^{[1]}(y_i|x_i) - f_m^{[2]}(y_i|x_i) \right|^2 \\
 &\leq \underbrace{3B_n^2 K \sum_{k=1}^K \frac{1}{n} \sum_{i=1}^n \left(\sum_{j=1}^p \gamma_{kj}^{[1]} x_{ij} - \sum_{j=1}^p \gamma_{kj}^{[2]} x_{ij} \right)^2}_{=:a} \\
 &\quad + \underbrace{3B_n^2 K q \sum_{k=1}^K \sum_{z=1}^q \frac{1}{n} \sum_{i=1}^n \left(\sum_{j=1}^p [\beta_k^{[1]}]_{z,j} x_{ij} - \sum_{j=1}^p [\beta_k^{[2]}]_{z,j} x_{ij} \right)^2}_{=:b} \\
 &\quad + 3B_n^2 \left(\|\beta_0^{[1]} - \beta_0^{[2]}\|_1 + \|\gamma_0^{[1]} - \gamma_0^{[2]}\|_1 + \|\text{vec}(\Sigma^{[1]} - \Sigma^{[2]})\|_1 \right)^2.
 \end{aligned}$$

So, for all $\delta > 0$, if

$$\begin{aligned}
 a &\leq \delta^2 / (36B_n^2), \quad b \leq \delta^2 / (36B_n^2), \quad \|\beta_0^{[1]} - \beta_0^{[2]}\|_1 \leq \delta / (18B_n), \\
 \|\gamma_0^{[1]} - \gamma_0^{[2]}\|_1 &\leq \delta / (18B_n), \quad \text{and} \quad \|\text{vec}(\Sigma^{[1]} - \Sigma^{[2]})\|_1 \leq \delta / (18B_n),
 \end{aligned}$$

then $\|f_m^{[1]} - f_m^{[2]}\|_n^2 \leq \delta^2/4$. To bound a and b , we can write

$$\begin{aligned}
 a &= Km^2 \sum_{k=1}^K \frac{1}{n} \sum_{i=1}^n \left(\sum_{j=1}^p \frac{\gamma_{kj}^{[1]}}{m} x_{ij} - \sum_{j=1}^p \frac{\gamma_{kj}^{[2]}}{m} x_{ij} \right)^2, \quad \text{and} \\
 b &= Kqm^2 \sum_{k=1}^K \sum_{z=1}^q \frac{1}{n} \sum_{i=1}^n \left(\sum_{j=1}^p \frac{[\beta_k^{[1]}]_{z,j}}{m} x_{ij} - \sum_{j=1}^p \frac{[\beta_k^{[2]}]_{z,j}}{m} x_{ij} \right)^2.
 \end{aligned}$$

Then, we apply Lemma 12 to obtain $\frac{\gamma_{k,\cdot}^{[1]}}{m} = \left(\frac{\gamma_{kj}^{[1]}}{m} \right)_{j \in [q]}$ and $\frac{[\beta_k^{[1]}]_{z,\cdot}}{m} = \left(\frac{[\beta_k^{[1]}]_{z,j}}{m} \right)_{j \in [q]}$, for all $k \in [K], z \in [q]$. Since $s_m^{[1]} \in S_m$, and using (10), we have $\|\gamma_k^{[1]}\| \leq m$ and $\|\text{vec}(\beta_k^{[1]})\|_1 \leq m$, which leads to $\sum_{j=1}^p \left| \frac{\gamma_{kj}^{[1]}}{m} \right| \leq 1$ and $\sum_{z=1}^q \sum_{j=1}^p \left| \frac{[\beta_k^{[1]}]_{z,j}}{m} \right| \leq 1$, respectively. Furthermore, given $x \in \mathcal{X} = [0, 1]^p$, we have $\|x\|_{\max, n}^2 = 1$. Thus, there exist families $\mathcal{A}$ of $(2p+1)^{36B_n^2 K^2 m^2 / \delta^2}$ vectors and $\mathcal{B}$ of $(2p+1)^{16B_n^2 q^2 K^2 m^2 / \delta^2}$ vectors of $\mathbb{R}^p$, such that for all $k \in [K], z \in [q], \gamma_{k,\cdot}^{[1]}$,

and $\left[\beta_k^{[1]}\right]_{z,\cdot}$, there exist $\gamma_{k,\cdot}^{[1]} \in \mathcal{A}$ and $\left[\beta_k^{[2]}\right]_{z,\cdot} \in \mathcal{B}$, such that

$$\begin{aligned} \frac{1}{n} \sum_{i=1}^n \left(\sum_{j=1}^p \frac{\gamma_{kj}^{[1]}}{m} x_{ij} - \sum_{j=1}^p \frac{\gamma_{kj}^{[2]}}{m} x_{ij} \right)^2 &\leq \frac{\delta^2}{36B_n^2 K^2 m^2}, \text{ and} \\ \frac{1}{n} \sum_{i=1}^n \left(\sum_{j=1}^p \frac{\left[\beta_k^{[1]}\right]_{z,j}}{m} x_{ij} - \sum_{j=1}^p \frac{\left[\beta_k^{[2]}\right]_{z,j}}{m} x_{ij} \right)^2 &\leq \frac{\delta^2}{36B_n^2 q^2 K^2 m^2}, \end{aligned}$$

which leads to $a \leq \delta^2/36B_n^2$ and $b \leq \delta^2/36B_n^2$. Moreover, (4) leads to

$$\begin{aligned} \left\| \beta_0^{[1]} \right\|_1 &= \sum_{k=1}^K \left\| \beta_{0k}^{[1]} \right\|_1 \leq Kq \left\| \beta_{0k}^{[1]} \right\|_\infty \leq KqA_\beta \text{ (using (100))}, \\ \left\| \gamma_0^{[1]} \right\|_1 &= \sum_{k=1}^K \left| \gamma_{0k}^{[1]} \right| \leq KA_\gamma, \left\| \text{vec} \left(\Sigma^{[1]} \right) \right\|_1 \leq \frac{Kq\sqrt{q}}{a_\Sigma}. \end{aligned}$$

Therefore, on the event $\mathcal{T}$,

$$\begin{aligned} M(\delta, F_m, \|\cdot\|_n) &\leq N(\delta/2, F_m, \|\cdot\|_n) \text{ (using Lemma 22)} \\ &\leq \text{card}(\mathcal{A}) \text{card}(\mathcal{B}) N\left(\frac{\delta}{18B_n}, B_1^K(KqA_\beta), \|\cdot\|_1\right) \\ &\quad \times N\left(\frac{\delta}{18B_n}, B_1^K(KA_\gamma), \|\cdot\|_1\right) N\left(\frac{\delta}{18B_n}, B_1^K\left(\frac{Kq\sqrt{q}}{a_\Sigma}\right), \|\cdot\|_1\right) \\ &\leq (2p+1)^{\frac{72B_n^2 q^2 K^2 m^2}{\delta^2}} \left(1 + \frac{18B_n KqA_\beta}{\delta}\right)^K \left(1 + \frac{18B_n KA_\gamma}{\delta}\right)^K \left(1 + \frac{18B_n Kq\sqrt{q}}{a_\Sigma \delta}\right)^K. \end{aligned}$$

B.2.3. PROOF OF LEMMA 13

Let $m \in \mathbb{N}^*$. From Lemma 10, on the event $\mathcal{T}$, it holds that

$$\sup_{f_m \in F_m} \|f_m\|_n \leq 2KB_n \left(A_\gamma + qA_\beta + \frac{q\sqrt{q}}{a_\Sigma} \right) =: R_n. \quad (90)$$

From Lemma 11, on the event $\mathcal{T}$ for all $S \in \mathbb{N}^*$, with $\delta = 2^{-s} R_n$,

$$\begin{aligned}
 & \sum_{s=1}^S 2^{-s} \sqrt{\ln [1 + M (2^{-s} R_n, F_m, \|\cdot\|_n)]} \leq \sum_{s=1}^S 2^{-s} \sqrt{\ln [2M (\delta, F_m, \|\cdot\|_n)]} \\
 & \leq \sum_{s=1}^S 2^{-s} \left[\sqrt{\ln 2} + \frac{6\sqrt{2}B_n q K m}{\delta} \sqrt{\ln (2p+1)} \right. \\
 & \quad \left. + \sqrt{K \ln \left[\left(1 + \frac{18B_n K q A_\beta}{\delta}\right) \left(1 + \frac{18B_n K A_\gamma}{\delta}\right) \left(1 + \frac{18B_n K q \sqrt{q}}{a_\Sigma \delta}\right) \right]} \right] \\
 & \leq \sum_{s=1}^S 2^{-s} \left[\sqrt{\ln 2} + \frac{2^s 6\sqrt{2}B_n q K m}{R_n} \sqrt{\ln (2p+1)} \right. \\
 & \quad \left. + \sqrt{K \ln \left[\left(1 + \frac{2^s 18B_n K q A_\beta}{R_n}\right) \left(1 + \frac{2^s 18B_n K A_\gamma}{R_n}\right) \left(1 + \frac{2^s 18B_n K q \sqrt{q}}{a_\Sigma R_n}\right) \right]} \right]. \quad (91)
 \end{aligned}$$

Notice from (90), that $R_n \geq 2K B_n \max \left(A_\gamma, q A_\beta, \frac{q\sqrt{q}}{a_\Sigma} \right)$. Moreover, it holds that $1 \leq 2^{s+3}$, and $\sum_{s=1}^S 2^{-s} = 1 - 2^{-S} \leq 1$, $\sum_{s=1}^S (\sqrt{e}/2)^s \leq \sqrt{e}/(2 - \sqrt{e})$, and since for all $s \in \mathbb{N}^*$, $e^s \geq s$, and thus $2^{-s} \sqrt{s} \leq (\sqrt{e}/2)^s$. Therefore, from (91):

$$\begin{aligned}
 & \sum_{s=1}^S 2^{-s} \sqrt{\ln [1 + M (2^{-s} R_n, F_m, \|\cdot\|_n)]} \\
 & \leq \sum_{s=1}^S 2^{-s} \left[\sqrt{\ln 2} + \frac{2^s 6\sqrt{2}B_n q K m}{R_n} \sqrt{\ln (2p+1)} + \sqrt{K \ln [(2^{s+1} 3^2) (2^{s+1} 3^2) (2^{s+1} 3^2)]} \right] \\
 & = \sum_{s=1}^S 2^{-s} \left[\sqrt{\ln 2} + \frac{2^s 6\sqrt{2}B_n q K m}{R_n} \sqrt{\ln (2p+1)} + \sqrt{K} \sqrt{3((s+1) \ln 2 + 2 \ln 3)} \right] \\
 & \leq \frac{6\sqrt{2}B_n K q m}{R_n} \sqrt{\ln (2p+1)} S + \sqrt{K} \sqrt{3 \ln 2} \sum_{s=1}^S 2^{-s} \sqrt{s} + \sqrt{\ln 2} (1 + \sqrt{3K}) + \sqrt{6 \ln 3 K} \\
 & \leq \frac{6\sqrt{2}B_n K q m}{R_n} \sqrt{\ln (2p+1)} S + \sqrt{K} \sqrt{3 \ln 2} \sum_{s=1}^S \left(\frac{\sqrt{e}}{2} \right)^s + \sqrt{\ln 2} (1 + \sqrt{3K}) + \sqrt{6 \ln 3 K} \\
 & \leq \frac{6\sqrt{2}B_n q K m}{R_n} \sqrt{\ln (2p+1)} S + \sqrt{K \ln 2} \underbrace{\left(\frac{\sqrt{3e}}{2 - \sqrt{e}} + 1 + \sqrt{3} + \sqrt{\frac{6 \ln 3}{\ln 2}} \right)}_{=: C_1}. \quad (92)
 \end{aligned}$$

Then, for all $S \in \mathbb{N}^*$, on the event $\mathcal{T}$:

$$\begin{aligned}
 & \mathbb{E} \left[\sup_{f_m \in \mathcal{F}_m} \left| \frac{1}{n} \sum_{i=1}^n \epsilon_i f_m(Z_i) \right| \right] \\
 &= \mathbb{E} \left[\sup_{f_m \in \mathcal{F}_m} \left| \frac{1}{n} \sum_{i=1}^n \epsilon_i f_m(Z_i) \mathbb{1}_{|f_m(Z_i)| \leq R_n} \right| \right] \quad (\text{using Lemma 10}) \\
 &= \mathbb{E} \left[\sup_{f \in \mathcal{F}} \left| \frac{1}{n} \sum_{i=1}^n \epsilon_i f(Z_i) \right| \right] \\
 &\leq R_n \left(\frac{6}{\sqrt{n}} \sum_{s=1}^S 2^{-s} \sqrt{\ln[1 + M(2^{-s} R_n, \mathcal{F}, \|\cdot\|_n)]} + 2^{-S} \right) \quad (\text{using Lemma 9}) \\
 &\leq R_n \left[\frac{6}{\sqrt{n}} \left(\frac{6\sqrt{2} B_n K m q}{R_n} \sqrt{\ln(2p+1)} S + \sqrt{K \ln 2} C_1 \right) + 2^{-S} \right] \quad (\text{using Lemma 92}). \quad (93)
 \end{aligned}$$

We choose $S = \ln n / \ln 2$ so that the two terms depending on S in (93) are of the same order. In particular, for this value of S , $2^{-S} \leq 1/n$, and we deduce from (93) and (90) that on the event $\mathcal{T}$,

$$\begin{aligned}
 & \mathbb{E} \left[\sup_{f_m \in \mathcal{F}_m} \left| \frac{1}{n} \sum_{i=1}^n \epsilon_i f_m(Z_i) \right| \right] \\
 &\leq \frac{36\sqrt{2} B_n K m q}{\sqrt{n}} \sqrt{\ln(2p+1)} \frac{\ln n}{\ln 2} + 2K B_n \left(A_\gamma + q A_\beta + \frac{q\sqrt{q}}{a_\Sigma} \right) \left(6\sqrt{\ln 2} C_1 \frac{\sqrt{K}}{\sqrt{n}} + \frac{1}{n} \right) \\
 &\leq \frac{B_n K m q}{\sqrt{n}} \sqrt{\ln(2p+1)} \ln n \underbrace{\frac{36\sqrt{2}}{\ln 2}}_{\approx 73.45} + \frac{K\sqrt{K}}{\sqrt{n}} B_n \left(A_\gamma + q A_\beta + \frac{q\sqrt{q}}{a_\Sigma} \right) \underbrace{2(6\sqrt{\ln 2} C_1 + 1)}_{\approx 141.32} \\
 &< \frac{74K B_n}{\sqrt{n}} \left[m q \sqrt{\ln(2p+1)} \ln n + 2\sqrt{K} \left(A_\gamma + q A_\beta + \frac{q\sqrt{q}}{a_\Sigma} \right) \right].
 \end{aligned}$$

B.3. Proof of Lemma 6

Since $\ln(z)$ is concave in z , Jensen's inequality (see e.g. Jensen (1906); Cover (1999)) implies that $\ln(\mathbb{E}_Z[Z]) \geq \mathbb{E}_Z[\ln(Z)]$, where Z is a random variable. Thus, for all $x \in \mathcal{X}$, Jensen's inequality

and Lemma 15 lead us the following upper bound

$$\begin{aligned}
 & \int_{\mathbb{R}^q} \ln(s_0(y|x)) s_0(y|x) dy \\
 &= \int_{\mathbb{R}^q} \ln[s_0(y|x)] \sum_{k=1}^K [g_k(x; \gamma_0) \phi(y; v_{0k}(x), \Sigma_{0k})] dy \\
 &= \sum_{k=1}^K g_k(x; \gamma_0) \int_{\mathbb{R}^q} \ln[s_0(y|x)] \phi(y; v_{0k}(x), \Sigma_{0k}) dy \\
 &\leq \sum_{k=1}^K g_k(x; \gamma_0) \ln \left[\int_{\mathbb{R}^q} s_0(y|x) \phi(y; v_{0k}(x), \Sigma_{0k}) dy \right] \\
 &= \sum_{k=1}^K g_k(x; \gamma_0) \ln \left[\int_{\mathbb{R}^q} \sum_{l=1}^K g_l(x; \gamma_0) \phi(y; v_{0l}(x), \Sigma_{0l}) \phi(y; v_{0k}(x), \Sigma_{0k}) dy \right] \\
 &= \sum_{k=1}^K g_k(x; \gamma_0) \ln \left[\sum_{l=1}^K g_l(x; \gamma_0) \int_{\mathbb{R}^q} \phi(y; v_{0l}(x), \Sigma_{0l}) \phi(y; v_{0k}(x), \Sigma_{0k}) dy \right] \\
 &\leq \sum_{k=1}^K g_k(x; \gamma_0) \ln \left[\sum_{l=1}^K g_l(x; \gamma_0) C_{s_0} \right], C_{s_0} = (4\pi)^{-q/2} A_{\Sigma}^{q/2}, (\text{using Lemma 15}) \\
 &= \ln C_{s_0} < \infty.
 \end{aligned} \tag{94}$$

Therefore, we obtain

$$\max \left\{ 0, \sup_{x \in \mathcal{X}} \int_{\mathbb{R}^q} \ln(s_0(y|x)) s_0(y|x) dy \right\} \leq \max \{0, \ln C_{s_0}\} =: H_{s_0} < \infty.$$

Next, we state the following important Lemma 15, which is used in the proof of Lemma 6.

Lemma 15 *There exists a positive constant $C_{s_0} := (4\pi)^{-q/2} A_{\Sigma}^{q/2}$, $0 < C_{s_0} < \infty$, such that for all $k \in [K], l \in [L]$,*

$$\int_{\mathbb{R}^q} \phi(y; v_{0l}(x), \Sigma_{0l}) \phi(y; v_{0k}(x), \Sigma_{0k}) dy < C_{s_0}, \quad \forall x \in \mathcal{X}. \tag{95}$$

Proof [Proof of Lemma 15] Firstly, for all $k \in [K], l \in [L]$, given

$$c_{lk}(x) = C_{lk} [\Sigma_{0l}^{-1} v_{0l}(x) + \Sigma_{0k}^{-1} v_{0k}(x)], \quad C_{lk} = (\Sigma_{0l}^{-1} + \Sigma_{0k}^{-1})^{-1},$$

Lemma 25 leads to

$$\begin{aligned}
 & \int_{\mathbb{R}^q} [\phi(y; v_{0l}(x), \Sigma_{0l}) \phi(y; v_{0k}(x), \Sigma_{0k})] dy \\
 &= Z_{lk}^{-1} \underbrace{\int_{\mathbb{R}^q} \phi(y; c_{lk}(x), C_{lk}) dy}_{=1}, \text{ where} \\
 &= (2\pi)^{-q/2} \det(\Sigma_{0l} + \Sigma_{0k})^{-1/2} \exp \left(-\frac{1}{2} (v_{0l}(x) - v_{0k}(x))^{\top} (\Sigma_{0l} + \Sigma_{0k})^{-1} (v_{0l}(x) - v_{0k}(x)) \right)
 \end{aligned} \tag{96}$$

Next, since the determinant is the product of the eigenvalues, counted with multiplicity, and Weyl's inequality, see *e.g.*, Lemma 26, for all $k \in [K], l \in [L]$, we have

$$\begin{aligned} \det(\Sigma_{0l} + \Sigma_{0k}) &\geq [m(\Sigma_{0l} + \Sigma_{0k})]^q \\ &\geq [m(\Sigma_{0l}) + m(\Sigma_{0k})]^q \text{ (using (116) from Lemma 26)} \\ &= \left[M(\Sigma_{0l}^{-1})^{-1} + M(\Sigma_{0k}^{-1})^{-1} \right]^q \\ &\geq (2A_\Sigma^{-1})^q \text{ (using boundedness assumptions in (4))}. \end{aligned}$$

Therefore, for all $k \in [K], l \in [L]$, it holds that

$$\det(\Sigma_{0l} + \Sigma_{0k})^{-1/2} \leq 2^{-q/2} (A_\Sigma)^{q/2} \text{ (using boundedness assumptions in (4))}. \quad (97)$$

Since $(\Sigma_{0l} + \Sigma_{0k})^{-1}$ is a positive definite matrix, it holds that

$$(v_{0l}(x) - v_{0k}(x))^\top (\Sigma_{0l} + \Sigma_{0k})^{-1} (v_{0l}(x) - v_{0k}(x)) \geq 0, \quad \forall x \in \mathcal{X}, l \in [L], k \in [K].$$

Then, since the exponential function is increasing, $\forall x \in \mathcal{X}, l \in [L], k \in [K]$, we have

$$\exp\left(-\frac{1}{2} (v_{0l}(x) - v_{0k}(x))^\top (\Sigma_{0l} + \Sigma_{0k})^{-1} (v_{0l}(x) - v_{0k}(x))\right) \leq \exp(0) = 1. \quad (98)$$

Finally, from (96), (97) and (98), we obtain

$$\int_{\mathbb{R}^q} [\phi(y; v_{0l}(x), \Sigma_{0l}) \phi(y; v_{0k}(x), \Sigma_{0k})] dy \leq (2\pi)^{-q/2} 2^{-q/2} A_\Sigma^{q/2} = (4\pi)^{-q/2} A_\Sigma^{q/2} =: C_{s_0} < \infty.$$

■

Appendix C. Technical results

We denote the vector space of all q -by- q real matrices by $\mathbb{R}^{q \times q}$ ($q \in \mathbb{N}^*$):

$$A \in \mathbb{R}^{q \times q} \iff A = (a_{i,j}) = \begin{bmatrix} a_{1,1} & \cdots & a_{1,q} \\ \vdots & & \vdots \\ a_{q,1} & \cdots & a_{q,q} \end{bmatrix}, a_{i,j} \in \mathbb{R}.$$

If a capital letter is used to denote a matrix (*e.g.*, A, B), then the corresponding lower-case letter with subscript i, j refers to the (i, j) th entry (*e.g.*, $a_{i,j}, b_{i,j}$). When required, we also designate the elements of a matrix with the notation $[A]_{i,j}$ or $A(i, j)$. Denote the q -by- q identity and zero matrices by I_q and 0_q , respectively.

Lemma 16 (Derivative of quadratic form, see Magnus and Neudecker (2019)) *Assume that X and a are non-singular matrix in $\mathbb{R}^{q \times q}$ and vector in $\mathbb{R}^{q \times 1}$, respectively. Then*

$$\frac{\partial a^\top X^{-1} a}{\partial X} = -X^{-1} a a^\top X^{-1}.$$

Lemma 17 (Jacobi's formula, Theorem 8.1 from Magnus and Neudecker (2019)) *If X is a differentiable map from the real numbers to q -by- q matrices,*

$$\frac{d}{dt} \det(X(t)) = \text{tr} \left(\text{Adj}(X(t)) \frac{dX(t)}{dt} \right), \quad \frac{\partial \det(X)}{\partial X} = (\text{Adj}(X))^\top = \det(X) (X^{-1})^\top.$$

Lemma 18 (Operator (induced) p -norm) *We recall an operator (induced) p -norms of a matrix $A \in \mathbb{R}^{q \times q}$ ($q \in \mathbb{N}^*$, $p \in \{1, 2, \infty\}$),*

$$\|A\|_p = \max_{x \neq 0} \frac{\|Ax\|_p}{\|x\|_p} = \max_{x \neq 0} \left\| A \left(\frac{x}{\|x\|_p} \right) \right\|_p = \max_{\|x\|_p=1} \|Ax\|_p, \quad (99)$$

where for all $x \in \mathbb{R}^q$,

$$\|x\|_\infty \leq \|x\|_1 = \sum_{i=1}^q |x_i| \leq q \|x\|_\infty, \quad (100)$$

$$\|x\|_2 = \left(\sum_{i=1}^q |x_i|^2 \right)^{\frac{1}{2}} = \left(x^\top x \right)^{\frac{1}{2}} \leq \|x\|_1 \leq \sqrt{q} \|x\|_2, \text{ and} \quad (101)$$

$$\|x\|_\infty = \max_{1 \leq i \leq q} |x_i| \leq \|x\|_2 \leq \sqrt{q} \|x\|_\infty. \quad (102)$$

Lemma 19 (Some matrix p -norm properties, Golub and Van Loan (2012)) *By definition, we always have the important property that for every $A \in \mathbb{R}^{q \times q}$ and $x \in \mathbb{R}^q$,*

$$\|Ax\|_p \leq \|A\|_p \|x\|_p, \quad (103)$$

and every induced p -norm is submultiplicative, i.e., for every $A \in \mathbb{R}^{q \times q}$ and $B \in \mathbb{R}^{q \times q}$,

$$\|AB\|_p \leq \|A\|_p \|B\|_p. \quad (104)$$

In particular, it holds that

$$\|A\|_1 = \max_{1 \leq j \leq q} \sum_{i=1}^q |a_{ij}| \leq \sum_{j=1}^q \sum_{i=1}^q |a_{ij}| := \|\text{vec}(A)\|_1 \leq q \|A\|_1, \quad (105)$$

$$\|\text{vec}(A)\|_\infty := \max_{1 \leq i \leq q, 1 \leq j \leq q} |a_{ij}| \leq \|A\|_\infty = \max_{1 \leq j \leq q} \sum_{i=1}^q |a_{ij}| \leq q \|\text{vec}(A)\|_\infty, \quad (106)$$

$$\|\text{vec}(A)\|_\infty \leq \|A\|_2 = \lambda_{\max}(A) \leq q \|\text{vec}(A)\|_\infty, \quad (107)$$

where $\lambda_{\max}$ is the largest eigenvalue of a positive definite symmetric matrix A . The p -norms, when $p \in \{1, 2, \infty\}$, satisfy

$$\frac{1}{\sqrt{q}} \|A\|_\infty \leq \|A\|_2 \leq \sqrt{q} \|A\|_\infty, \quad (108)$$

$$\frac{1}{\sqrt{q}} \|A\|_1 \leq \|A\|_2 \leq \sqrt{q} \|A\|_1. \quad (109)$$

Given $\delta > 0$, we need to define the δ -packing number and δ -covering number.

Definition 20 (δ -packing number, e.g., Definition 5.4 from Wainwright (2019)) Let $(\mathcal{F}, \|\cdot\|)$ be a normed space and let $\mathcal{G} \subset \mathcal{F}$. With $(g_i)_{i=1,\dots,m} \in \mathcal{G}$, $\{g_1, \dots, g_m\}$ is an δ -packing of $\mathcal{G}$ of size $m \in \mathbb{N}^*$, if $\|g_i - g_j\| > \delta, \forall i \neq j, i, j \in \{1, \dots, m\}$, or equivalently, $\bigcap_{i=1}^m B(g_i, \delta/2) = \emptyset$. Upon defining δ -packing, we can measure the maximal number of disjoint closed balls with radius $\delta/2$ that can be “packed” into $\mathcal{G}$. This number is called the δ -packing number and is defined as

$$M(\delta, \mathcal{G}, \|\cdot\|) := \max \{m \in \mathbb{N}^* : \exists \delta\text{-packing of } \mathcal{G} \text{ of size } m\}. \quad (110)$$

Definition 21 (δ -covering number, Definition 5.1 from Wainwright (2019)) Let $(\mathcal{F}, \|\cdot\|)$ be a normed space and let $\mathcal{G} \subset \mathcal{F}$. With $(g_i)_{i \in [n]} \in \mathcal{G}$, $\{g_1, \dots, g_n\}$ is an δ -covering of $\mathcal{G}$ of size n if $\mathcal{G} \subset \bigcup_{i=1}^n B(g_i, \delta)$, or equivalently, $\forall g \in \mathcal{G}, \exists i$ such that $\|g - g_i\| \leq \delta$. Upon defining the δ -covering, we can measure the minimal number of closed balls with radius δ , which is necessary to cover $\mathcal{G}$. This number is called the δ -covering number and is defined as

$$N(\delta, \mathcal{G}, \|\cdot\|) := \min \{n \in \mathbb{N}^* : \exists \delta\text{-covering of } \mathcal{G} \text{ of size } n\}. \quad (111)$$

The covering entropy (metric entropy) is defined as follows $H_{\|\cdot\|}(\delta, \mathcal{G}) = \ln(N(\delta, \mathcal{G}, \|\cdot\|))$.

The relation between the packing number and the covering number is described in the following lemma.

Lemma 22 (Lemma 5.5 from Wainwright (2019)) Let $(\mathcal{F}, \|\cdot\|)$ be a normed space and let $\mathcal{G} \subset \mathcal{F}$. Then

$$M(2\delta, \mathcal{G}, \|\cdot\|) \leq N(\delta, \mathcal{G}, \|\cdot\|) \leq M(\delta, \mathcal{G}, \|\cdot\|).$$

Lemma 23 (Chernoff’s inequality, e.g., Chapter 2 in Wainwright (2019)) Assume that the random variable has a moment generating function in a neighborhood of zero, meaning that there is some constant $b > 0$ such that the function $\varphi(\lambda) = \mathbb{E}[e^{\lambda(U-\mu)}]$ exists for all $\lambda \leq |b|$. In such a case, we may apply Markov’s inequality to the random variable $Y = e^{\lambda(U-\mu)}$, thereby obtaining the upper bound

$$\mathbb{P}(U - \mu \geq a) = \mathbb{P}(e^{\lambda(U-\mu)} \geq e^{\lambda a}) \leq \frac{\mathbb{E}[e^{\lambda(U-\mu)}]}{e^{\lambda a}}.$$

Optimizing our choice of λ so as to obtain the tightest result yields the Chernoff bound

$$\ln(\mathbb{P}(U - \mu \geq a)) \leq \sup_{\lambda \in [0, b]} \left\{ \lambda a - \ln \left(\mathbb{E}[e^{\lambda(U-\mu)}] \right) \right\}. \quad (112)$$

In particular, if $U \sim \mathcal{N}(\mu, \sigma)$ is a Gaussian random variable with mean μ and variance σ^2 . By a straightforward calculation, we find that U has the moment generating function

$$\mathbb{E}[e^{\lambda U}] = e^{\mu\lambda + \frac{\sigma^2\lambda^2}{2}}, \text{ valid for all } \lambda \in \mathbb{R}.$$

Substituting this expression into the optimization problem defining the optimized Chernoff bound (112), we obtain

$$\sup_{\lambda \geq 0} \left\{ \lambda t - \ln \left(\mathbb{E} \left[e^{\lambda(U-\mu)} \right] \right) \right\} = \sup_{\lambda \geq 0} \left\{ \lambda t - \frac{\sigma^2 \lambda^2}{2} \right\} = -\frac{t^2}{2\sigma^2},$$

where we have taken derivatives in order to find the optimum of this quadratic function. So, (112) leads to

$$\mathbb{P}(X \geq \mu + t) \leq e^{-\frac{t^2}{2\sigma^2}}, \text{ for all } t \geq 0. \quad (113)$$

Recall that a multi-index $\alpha = (\alpha_1, \dots, \alpha_p)$, $\alpha_i \in \mathbb{N}^*$, $\forall i \in \{1, \dots, p\}$ is a p -tuple of non-negative integers. Let

$$|\alpha| = \sum_{i=1}^p \alpha_i, \quad \alpha! = \prod_{i=1}^p \alpha_i!, \quad x^\alpha = \prod_{i=1}^p x_i^{\alpha_i}, \quad x \in \mathbb{R}^p, \quad \partial^\alpha f = \partial_1^{\alpha_1} \partial_2^{\alpha_2} \dots \partial_p^{\alpha_p} = \frac{\partial^{|\alpha|} f}{\partial x_1^{\alpha_1} \partial x_2^{\alpha_2} \dots \partial x_p^{\alpha_p}}.$$

The number $|\alpha|$ is called the *order* or *degree* of α . Thus, the order of α is the same as the order of x^α as a monomial or the order of ∂^α as a partial derivative.

Lemma 24 (Taylor's Theorem in Several Variables from Duistermaat and Kolk (2004)) Suppose $f : \mathbb{R}^p \mapsto \mathbb{R}$ is in the class C^{k+1} , of continuously differentiable functions, on an open convex set S . If $a \in S$ and $a + h \in S$, then

$$f(a + h) = \sum_{|\alpha| \leq k} \frac{\partial^\alpha f(a)}{\alpha!} h^\alpha + R_{a,k}(h),$$

where the remainder is given in Lagrange's form by

$$R_{a,k}(h) = \sum_{|\alpha|=k+1} \partial^\alpha f(a + ch) \frac{h^\alpha}{\alpha!} \text{ for some } c \in (0, 1),$$

or in integral form by

$$R_{a,k}(h) = (k+1) \sum_{|\alpha|=k+1} \frac{h^\alpha}{\alpha!} \int_0^1 (1-t)^k \partial^\alpha f(a + th) dt.$$

In particular, we can estimate the remainder term if $|\partial^\alpha f(x)| \leq M$ for $x \in S$ and $|\alpha| = k+1$, then

$$|R_{a,k}(h)| \leq \frac{M}{(k+1)!} \|h\|_1^{k+1}, \quad \|h\|_1 = \sum_{i=1}^p |h_i|.$$

Recall that the multivariate Gaussian (or Normal) distribution has a joint density given by

$$\phi(y; \mu; \Sigma) = (2\pi)^{-q/2} \det(\Sigma)^{-1/2} \exp \left(-\frac{1}{2} (y - \mu)^\top \Sigma^{-1} (y - \mu) \right), \quad (114)$$

where μ is the mean vector (of length q) and Σ is the symmetric, positive definite covariance matrix (of size $q \times q$). Then, we have the following well-known Gaussian identity, see more in Lemma 25.

Lemma 25 (Product of two Gaussians, see e.g., Equation (A.7) in Williams and Rasmussen (2006))
The product of two Gaussians gives another (un-normalized) Gaussian

$$\begin{aligned} \phi(y; a, A) \phi(y; b, B) &= Z^{-1} \phi(y; c, C), \text{ where,} \\ c &= C (A^{-1}a + B^{-1}b) \text{ and } C = (A^{-1} + B^{-1})^{-1}, \\ Z^{-1} &= (2\pi)^{-q/2} \det(A + B)^{-1/2} \exp\left(-\frac{1}{2}(a - b)^\top (A + B)^{-1} (a - b)\right). \end{aligned} \quad (115)$$

We recall the following inequality of Hermann Weyl, see e.g., Horn and Johnson (2012, Theorem 4.3.1)

Lemma 26 (Weyl's inequality, e.g., Theorem 4.3.1 from Horn and Johnson (2012)) *Let $A, B \in \mathbb{R}^{q \times q}$ be Hermitian and let the respective eigenvalues of A, B , and $A+B$ be $\{\lambda_i(A)\}_{i \in [q]}$, $\{\lambda_i(B)\}_{i \in [q]}$, and $\{\lambda_i(A+B)\}_{i \in [q]}$, each algebraically nondecreasing order as follows:*

$$m(A) = \lambda_1(A) \leq \lambda_2(A) \leq \dots \leq \lambda_q(A) = M(A).$$

Then, for each $i \in [q]$,

$$\begin{aligned} \lambda_i(A+B) &\leq \lambda_{i+j}(A) + \lambda_{q-j}(B), \quad j \in \{0\} \cup [q-i], \\ \lambda_{i-j+1}(A) + \lambda_j(B) &\leq \lambda_i(A+B), \quad j \in [i]. \end{aligned}$$

In particular, we have

$$\begin{aligned} M(A+B) &\leq M(A) + M(B), \\ m(A+B) &\geq m(A) + m(B). \end{aligned} \quad (116)$$