

# Weak gravitational lensing by Einstein-non-linear-Maxwell-Yukawa black hole

Wajiha Javed,<sup>1,\*</sup> Muhammad Bilal Khadim,<sup>1,†</sup> and Ali Övgün<sup>2,‡</sup>

<sup>1</sup>*Division of Science and Technology, University of Education, Township-Lahore, Pakistan*

<sup>2</sup>*Physics Department, Eastern Mediterranean University,  
99628, Famagusta, North Cyprus via Mersin 10, Turkey.*

(Dated: January 17, 2022)

In this article, we analyze the weak gravitational lensing in the context of Einstein-non-linear Maxwell-Yukawa black hole. To this desire, we derive the deflection angle of light by Einstein-non-linear Maxwell-Yukawa black hole using the Gibbons and Werner method. For this purpose, we obtain the Gaussian curvature and apply the Gauss-Bonnet theorem to find the deflection angle of Einstein-non-linear Maxwell-Yukawa black hole in weak field limits. Moreover, we derive the deflection angle of light in the influence of plasma medium. We also analyze the graphical behavior of deflection angle by Einstein-non-linear Maxwell-Yukawa black hole in the presence of plasma as well as non-plasma medium.

PACS numbers: 95.30.Sf, 98.62.Sb, 97.60.Lf

Keywords: Weak gravitational lensing; Einstein-non-linear Maxwell-Yukawa black holes; Deflection angle; Gauss-Bonnet theorem.

## I. INTRODUCTION

In our universe the black holes (BHs) are essential components and when stars are died, they become very dense objects which is the most important discoveries of astrophysics. To test the fundamental laws of universe black holes provide a golden opportunity. For example neutron star merges and the gravitational waves from black holes has been discover recently. Therefore, the gravitational lensing of black holes has attain incredible attention in the last few decade, essentially because of the solid proof of supermassive black holes at the focal point of galaxies [1, 2]. By utilizing the gravitational lensing we can simplified the study of black holes, which is a general investigative method for acquiring the time delays of the images, magnification and positions. Darwin [3] was the first who analyses the Schwarzschild geometry. After that in 1985 Herschel [4] published his similar article and after that many authors [5, 6], extended this geometry to general spherically symmetric black holes and Reissner-Nordstrom geometry. Kerr black holes [7]-[10] was also discussed for acquiring the time delays of the images, magnification and positions by using gravitational lensing. Many modification has been done through deflection of light and modification in the reference of non-linear electrodynamics (NLE) [11] has been studied through the various alternative gravity theories [12]. Classically, a black hole contains singularity and also horizon because in general theory of relativity the spacetime singularities create a dozen of problems which are physical as well as mathematical. Therefore many people use various methods to remove these singularities from the black holes like modified gravity, effect of quantum gravity and NLE. Bozza [13] discussed these topics in his recent article in details which also include observational prospectus and additional references [14]-[20].

The main focus of this article is to calculate the deflection angle of photon by using GBT. The aspect of deflection of light has been widely studied in different astrophysical system in the influence of strong as well as weak gravitational lensing [18]-[21]. Gibbons and Werner [22] made a very important role who contended about the significance of topological effect on the deflection of angle by utilizing the GBT and optical geometry. Additionally, they have calculated deflection angle for the schwarzschild black holes which is different from the other method by supposing light ray by taking domain outside, where the mass is closed in the given area on space is strongly related to the lensing effect. Recently, Gibbons and Werner (GW) [22] computed the deflection angle by applying GBT as follows

$$\sigma = - \int \int_{D_\infty} \mathcal{K} dS.$$

\*Electronic address: [wajiha.javed@ue.edu.pk](mailto:wajiha.javed@ue.edu.pk); [wajihajaved84@yahoo.com](mailto:wajihajaved84@yahoo.com)

†Electronic address: [blaljutt723@gmail.com](mailto:blaljutt723@gmail.com)

‡Electronic address: [ali.ovgun@emu.edu.tr](mailto:ali.ovgun@emu.edu.tr); URL: <https://aovgun.weebly.com>

Here,  $\mathcal{K}$  is Gaussian curvature and  $dS$  represents the surface element. Afterwards, Werner [23] extend this method and obtained the deflection of light by Kerr-Newman BHs by applying the Nazims's method for Rander-Finsler metric. Recently, deflection angle for a asymptotically flat and spherically symmetric spacetime by using the finite distance from an observer to a light source has been calculated by Ishihara et al. [24]-[26]. Moreover, Asada et al. [27] in stationary axisymmetric spacetime have calculated the weak gravitational lensing. In all of these methods by using GBT they have calculated the weak gravitational lensing which shows the global properties. The study of gravitational lensing in the presence of plasma medium discussed in number of cases. Initially, Bisnovatyi-kogan [28]-[29] show that the gravitational deflection in plasma is different from vacuum deflection angle due to presive properties of plasma. Recently, Gallo and Crisnejo [30] discussed the deflection angle of photon in a plasma medium.

In Einstein-Maxwell theory (EMT) for hairy BH in the context of Weak gravitational lensing with a non-minimally coupled dilaton has been examined by Javed et al. [31]. After that, there is a lot of literature [32]-[75] related to the investigation of weak gravitational lensing through the method of GBT on various black holes, cosmic strings and wormholes. In this paper, we study the weak gravitational lensing by Einstein-non-linear Maxwell-Yukawa black holes.

This paper is composed as; In Section 2, we concisely review about Einstein-non-linear Maxwell-Yukawa black hole. In Section 3, by using the Gauss-Bonnet theorem we compute the deflection angle of Einstein-non-linear Maxwell-Yukawa black hole. In Section 4, we work to investigate the deflection angle in the influence of plasma medium. We also demonstrate the graphical effect of deflection angle in the context of Einstein-non-linear Maxwell-Yukawa black hole for plasma and also for non-plasma medium. In Section 5, we present our result.

## II. COMPUTATION OF WEAK LENSING BY EINSTEIN-NON-LINEAR MAXWELL-YUKAWA BLACK HOLE AND GAUSS-BONNET THEOREM

The Einstein-non-linear Maxwell-Yukawa BH in a static and spherically symmetric form is given as [76]

$$ds^2 = -f(r)dt^2 + \frac{dr^2}{f(r)} + r^2(d\theta^2 + \sin^2\theta d\phi^2), \quad (1)$$

The metric function  $f(r)$  yields

$$f(r) \simeq 1 + \frac{2M}{r} - \frac{4qC_0}{r^2} + \frac{4qC_0\alpha}{3r} - qC_0\alpha^2 + \mathcal{O}(\alpha^3). \quad (2)$$

Here,  $M$  rendered as mass of BH,  $C_0$  is an integration constant [77],  $q$  represents charge of BH that is located at the origin and  $\alpha$  is a positive constant and it can be chosen as  $\alpha = 1$ .

The optical spacetime simply written as

$$dt^2 = \frac{dr^2}{f(r)^2} + \frac{r^2 d\phi^2}{f(r)} \quad (3)$$

From Eq. 1 we obtain non-zero Christopher symbols given below

$$\Gamma_{00}^0 = \frac{-f'(r)}{f(r)}, \quad \Gamma_{11}^0 = \frac{2f(r) - rf'(r)}{2rf(r)}, \quad \Gamma_{01}^1 = \frac{r^2 f'(r) - f(r)2r}{2}. \quad (4)$$

Now, we calculate the Ricci Scalar corresponds to the optical metric by using the non-zero Christopher symbols that are stated as:

$$\mathcal{R} = \frac{-f'(r)2}{2} + f''(r)f(r). \quad (5)$$

The Gaussian curvature that is computed as follows:

$$\mathcal{K} = \frac{RicciScalar}{2} \quad (6)$$

After simplifying, Gaussian optical curvature for Einstein-non-linear Maxwell-Yukawa black hole is stated as:

$$\mathcal{K} = \frac{-f'(r)^2}{4} + \frac{f(r)f''(r)}{2}, \quad (7)$$

where  $f(r)$  is given in Eq. 2 so that Gaussian optical curvature for Einstein-non-linear Maxwell-Yukawa black hole in leading order term is obtained as;

$$\mathcal{K} \approx -12 \frac{C_0 q}{r^4} + 4 \frac{C_0 q \alpha}{3r^3} + \left( 2r^{-3} - 24 \frac{C_0 q}{r^5} + 4 \frac{C_0 q \alpha}{r^4} \right) M. \quad (8)$$

### III. DEFLECTION ANGLE OF EINSTEIN-NON-LINEAR MAXWELL-YUKAWA BLACK HOLES AND GAUSS-BONNET THEOREM

Now, by utilizing GBT, we will calculate the deflection angle of photon by Einstein-non-linear Maxwell-Yukawa BH. By using GBT in the region  $\mathcal{G}_R$ , given as

$$\int_{\mathcal{G}_R} \mathcal{K} dS + \oint_{\partial \mathcal{G}_R} k dt + \sum_t \hat{\alpha} = 2\pi \mathcal{X}(\mathcal{G}_R), \quad (9)$$

where,  $k$  represent the geodesic curvature,  $\mathcal{K}$  denotes the Gaussian curvature and one can define  $k$  as  $k = \bar{g}(\nabla_{\hat{\alpha}} \hat{\alpha}, \check{\alpha})$  in that way  $\bar{g}(\hat{\alpha}, \hat{\alpha}) = 1$ , where  $\hat{\alpha}$  represent the unit acceleration vector and  $\alpha_t$  denotes the exterior angle at  $t^{th}$  vertex respectively. As  $R \rightarrow \infty$ , we obtain the jump angles approximate to  $\pi/2$ , Thus  $\alpha_O + \alpha_S \rightarrow \pi$ . Here,  $\mathcal{X}(\mathcal{G}_R) = 1$  is a Euler characteristic number and  $\mathcal{G}_R$  denotes the non-singular domain. Therefore, we obtain

$$\int \int_{\mathcal{G}_R} \mathcal{K} dS + \oint_{\partial \mathcal{G}_R} k dt + \hat{\alpha} = 2\pi \mathcal{X}(\mathcal{G}_R). \quad (10)$$

where, the total jump angle is  $\hat{\alpha} = \pi$ , When  $R$  approaching to infinity, then the remaining part is  $k(D_R) = |\nabla_{\dot{D}_R} \dot{D}_R|$ . For geodesic curvature the radial component is described as:

$$(\nabla_{\dot{D}_R} \dot{D}_R)^r = \dot{D}_R^\phi \partial_\phi \dot{D}_R^r + \Gamma_{\phi\phi}^r (\dot{D}_R^\phi)^2. \quad (11)$$

At  $R$  very high,  $D_R := r(\phi) = R = \text{const.}$  Thus, the first term of Eq. 11 vanishes and  $(\dot{D}_R^\phi)^2 = \frac{1}{f^2(r^*)}$ . Recalling  $\Gamma_{\phi\phi}^r = \frac{2f(r)-rf'(r)}{2rf(r)}$ , we get

$$(\nabla_{\dot{D}_R} \dot{D}_R)^r \rightarrow \frac{-1}{R}, \quad (12)$$

and  $k(D_R) \rightarrow R^{-1}$ , so we write  $dt = R d\phi$ . Thus;

$$k(D_R) dt = d\phi. \quad (13)$$

From the pervious results, we get

$$\int \int_{\mathcal{G}_R} \mathcal{K} ds + \oint_{\partial \mathcal{G}_R} k dt \stackrel{R \rightarrow \infty}{=} \int \int_{S_\infty} \mathcal{K} dS + \int_0^{\pi+\sigma} d\phi. \quad (14)$$

At  $0^{th}$  order weak field deflection limit of the light is defined as  $r(t) = b/\sin \phi$ . Hence, the deflection angle given as:

$$\sigma = - \int_0^\pi \int_{b/\sin \phi}^\infty \mathcal{K} \sqrt{\det g} du d\phi, \quad (15)$$

here we put the first terms of Eq. 8 into above Eq. 15, so we get the deflection angle upto leading order term is computed as:

$$\sigma \simeq \frac{4M}{b} + \frac{C_0 M \alpha q \pi}{b^2} + 8 \frac{C_0 q \alpha}{3b} - 3 \frac{C_0 q \pi}{b^2} - \frac{32 C_0 M q}{3 b^3}. \quad (16)$$

Note that the solution (16) with  $\alpha = q = 0$  reduces to deflection angle of Schwarzschild BH in the leading order terms. Moreover, it can be seen that the  $\alpha$  parameter increases the deflection angle.

#### IV. WEAK LENSING BY EINSTEIN-NON-LINEAR MAXWELL-YUKAWA BLACK HOLE IN A PLASMA MEDIUM

This section is based on the calculation of weak gravitational lensing by Einstein-non-linear Maxwell-Yukawa black hole in plasma medium. For Einstein non-linear Maxwell-Yukawa black hole the refractive index  $n(r)$ , is obtain as,

$$n(r) = \sqrt{1 - \frac{\omega_e^2}{\omega_\infty^2} f(r)}, \quad (17)$$

where electron plasma frequency is  $\omega_e$  and photon frequency measured by an observer at infinity is  $\omega_\infty$  are and respectively, then the corresponding optical metric illustrated as

$$dt^2 = g_{ij}^{opt} dx^i dx^j = n^2(r) \left[ -f(r) dt^2 + \frac{dr^2}{f(r)} + r^2 (d\theta^2) \right]. \quad (18)$$

The metric function  $f(r)$  is defined in Eq. 2.4 and the non-zero christopher symbols corresponding to the optical metric are computed as:

$$\begin{aligned} \Gamma_{00}^0 &= \left( 1 + \frac{\omega_e^2 f(r)}{\omega_\infty^2} \right) \left[ \frac{-\omega_e^2 f'(r)}{2\omega_\infty^2} \right] - \frac{-f'(r)}{f(r)}, \\ \Gamma_{01}^1 &= \left( 1 + \frac{\omega_e^2 f(r)}{\omega_\infty^2} \right) \left( \frac{-\omega_e^2 f'(r)}{2\omega_\infty^2} \right) + \frac{2f(r) - r f'(r)}{2r f(r)}, \\ \Gamma_{11}^0 &= \left( 1 + \frac{\omega_e^2 f(r)}{\omega_\infty^2} \right) \left( \frac{\omega_e^2 f(r) f'(r) r^2}{2\omega_\infty^2} \right) + \frac{r^2 f'(r) - f(r) 2r}{2}. \end{aligned} \quad (19)$$

Now, by using the above non-zero christopher symbols the value of Gaussian optical curvature is found as:

$$\begin{aligned} \mathcal{K} \approx 3 \frac{\omega_e^2 M}{r^3 \omega_\infty^2} + 2 \frac{M}{r^3} - 104 \frac{C_0 q M \omega_e^2}{\omega_\infty^2 r^5} - 24 \frac{C_0 q M}{r^5} - 20 \frac{C_0 q \omega_e^2}{\omega_\infty^2 r^4} - 12 \frac{C_0 q}{r^4} + 16 \frac{q \alpha C_0 \omega_e^2 M}{\omega_\infty^2 r^4} + 4 \frac{q \alpha C_0 M}{r^4} \\ + 2 \frac{q \alpha C_0 \omega_e^2}{r^3 \omega_\infty^2} + 4/3 \frac{q \alpha C_0}{r^3}. \end{aligned} \quad (20)$$

For this, we use GBT to compute the deflection angle in order to relate it with non-plasma. As follows, for calculating angle in the weak field limits, as the light beams become a straight line. Therefore, the condition at zero order is  $r = \frac{b}{\sin \phi}$ . Then the GBT reduces to this simple form for calculating the deflection angle  $\sigma$ :

$$\sigma = - \lim_{R \rightarrow 0} \int_0^\pi \int_{\frac{b}{\sin \phi}}^R \mathcal{K} dS \quad (21)$$


So, the deflection angle of Einstein-non-linear Maxwell-Yukawa BH in the presence of plasma medium is obtained as follows:

$$\begin{aligned} \sigma \simeq \frac{4M}{b} + 4 \frac{q \alpha C_0 M \omega_e^2 \pi}{b^2 \omega_\infty^2} + \frac{C_0 q M \alpha \pi}{b^2} + 4 \frac{C_0 q \alpha \omega_e^2}{b \omega_\infty^2} + 8/3 \frac{q C_0 \alpha}{b} - 5 \frac{C_0 q \omega_e^2 \pi}{b^2 \omega_\infty^2} - 3 \frac{q C_0 \pi}{b^2} - \frac{416 q C_0 M \omega_e^2}{9 b^3 \omega_\infty^2} \\ - \frac{32 C_0 q M}{3 b^3} + 6 \frac{M \omega_e^2}{b \omega_\infty^2}. \end{aligned} \quad (22)$$


The above result tells us that photon rays are moving into medium of homogeneous plasma. One can see that the plasma effect can be removed by neglecting  $\frac{\omega_e}{\omega_\infty}$  term from Eq. 22 and it reduced into Eq. 16.

#### V. GRAPHICAL INFLUENCE OF DEFLECTION ANGLE ON EINSTEIN-NON-LINEAR MAXWELL-YUKAWA BLACK HOLE

This section of the paper comprises the graphical influence of deflection angle of Einstein-non-linear Maxwell-Yukawa BH. We examine the impact of different parameters on deflection angle. Here for simplicity we take  $C_0 = c$

Figure 1:  $\sigma$  versus  $b$ .


- **Figure 1 Left Plot** indicates the behavior of deflection angle w.r.t  $b$  by fixing the value of  $M, c, \alpha$  and varying  $q$ . It is to be observed that for small constant value of  $M$  and  $q \geq 0$  the behavior of deflection angle gradually decreasing with respect to impact parameter  $b$ . For increasing value of  $q$ , deflection angle increases.
- **Figure 1 Right Plot** represents the behavior of deflection angle w.r.t  $b$  by varying the mass  $M$  and taking  $q, c, \alpha$  fixed. We noticed that for values of  $M \geq 0$  the behavior of deflection angle gradually positively decreasing with respect to  $b$ . But for the increasing values of  $M$ , the behavior of the deflection angle is increasing.

Figure 1(a):  $\sigma$  versus  $b$ .


- **Figure 1(a) Left Plot** indicates the behavior of deflection angle w.r.t  $b$  by fixing the value of  $\alpha$  and varying  $q, c, M$ . It is to be observed that for decreasing the value of  $\alpha < 0$  the deflection angle gradually decreasing. On the other hand, the deflection angle is increasing gradually with respect to  $b$  for fixed  $\alpha$ .
- **Figure 1(a) Right Plot** represents the behavior of deflection angle w.r.t  $b$  by varying the  $c$  and taking  $q, \alpha, M$  fixed. We noticed that for values of  $b > 5$  and  $c \geq 0$  the behavior of deflection angle gradually positively decreasing. For the increasing the value of  $c$ , the behavior of deflection angle is negatively decreasing.

## VI. GRAPHICAL ANALYSIS FOR PLASMA MEDIUM

This section gives us graphical analysis of deflection angle  $\alpha$  in the presence of plasma medium. For simplicity we take  $C_0 = c$  in the Eq. 22.

Figure 2:  $\sigma$  versus  $b$ .

- **Figure 2 Left Plot**, shows the behavior of deflection angle w.r.t  $b$  by taking the fixed values of  $M, q, c, \alpha$  and by varying  $\beta$ . It is to analyzed that initially the behavior of deflection angle is positively decreasing for  $\beta \geq 0$ . Furthermore, for the increasing value of  $\beta$  the deflection angle positively increases for fixed  $b$ .
- **Figure 2 Right Plot**, illustrate the behavior of deflection angle w.r.t  $b$  by varying BH charge  $q$  and for fixed  $M, \alpha, c$  and  $\beta$ . We examined that for positive values of  $q$  the deflection angle gradually positively decreasing. While, for increasing values of  $q$ , the behavior of deflection angle is gradually increasing for fixed  $b$ .

Figure 2(a):  $\tilde{\alpha}$  versus  $b$ .

- **Figure 2(a) Left Plot**, examines the behavior of deflection angle w.r.t  $b$  by taking the fixed values of  $\beta, q, c, M$  and by varying  $\alpha$ . We noticed that for small values of  $\alpha$ , and  $\beta = 1, q = 1$  then the deflection angle cannot define the behavior. While, for large values of  $\alpha$  the behavior of deflection angle is positively decreasing. For values of  $\alpha < 0$  the behavior of deflection angle gradually increasing.
- **Figure 2(a) Right Plot**, we analyzed the behavior of deflection angle w.r.t  $b$  by varying  $c$  and for fixed  $M, \alpha, q$  and  $\beta$ . we see that for  $c \geq 0$  the deflection angle positively decreasing. For increasing values of  $c$ , deflection angle is increasing for fixed  $b$ .




Figure 3:  $\sigma$  versus  $b$ .

- **Figure 3**, manifest the influence of deflection angle w.r.t  $b$  for constant values of  $q, \beta, c, \alpha$  and varying  $M$ . It has been noted that for small range of  $b < 15$  the behavior of deflection angle not defined. On the other hand for large  $b \geq 15$  and for  $M \geq 0$  the deflection angle positively decreasing and the behavior of deflection angle is increasing for increasing values of  $M$  at fixed  $b$ .

## VII. CONCLUSION

The present paper is about the investigation of deflection angle by Einstein-non-linear Maxwell-Yukawa BH in non-plasma as well as plasma medium. In this regard, we study the weak gravitational lensing by using GBT and obtain the deflection angle of photon for Einstein-non-linear Maxwell-Yukawa BH. The obtained deflection angle is given in Eq. 16 as follows:

$$\sigma \simeq \frac{4M}{b} + \frac{C_0 M \alpha q \pi}{b^2} + 8 \frac{C_0 q \alpha}{3b} - 3 \frac{C_0 q \pi}{b^2} - \frac{32 C_0 M q}{3 b^3} + \mathcal{O}(q^2, M^2, C_0^2).$$

We examine that by the reduction of some parameters the obtained deflection angle converted into the Schwarzschild deflection angle up to the first order terms. We also discuss the graphical effect of different parameters on deflection angle by Einstein-non-linear Maxwell-Yukawa BH. We also observed that deflection angle in the presence of plasma medium given by Eq. 22 which is;

$$\begin{aligned} \sigma \simeq \frac{4M}{b} + 4 \frac{q \alpha C_0 M \omega_e^2 \pi}{b^2 \omega_\infty^2} + \frac{C_0 q M \alpha \pi}{b^2} + 4 \frac{C_0 q \alpha \omega_e^2}{b \omega_\infty^2} + 8/3 \frac{q C_0 \alpha}{b} - 5 \frac{C_0 q \omega_e^2 \pi}{b^2 \omega_\infty^2} - 3 \frac{q C_0 \pi}{b^2} - \frac{416 q C_0 M \omega_e^2}{9 b^3 \omega_\infty^2} \\ - \frac{32 C_0 q M}{3 b^3} + 6 \frac{M \omega_e^2}{b \omega_\infty^2} + \mathcal{O}(q^2, M^2, C_0^2, \omega_e^3) \end{aligned} \quad (23)$$

When the  $\frac{\omega_e}{\omega_\infty}$  approaches to zero, the plasma effect are removed. Furthermore, we scrutinized the graphical impact of deflection angle on Einstein-non-linear Maxwell-Yukawa BH in plasma medium versus some parameters. One can conclude that the deflection of photon is found outside of the lensing area which points that the gravitational lensing effect is a global and even topological effect. Hence, it is our belief that studying on the Virbhadrha-Ellis lens equation, can provide us to improve the computation of the image positions, Einstein ring radii, magnification factors and the magnification ratio. This is going to be our next problem in the near future. Moreover, it will be interesting to investigate the deflection angle of black holes in MAXWELL  $f(R)$  gravity theories and in fourth order gravity theories using the GBT in future [78–80].

- 
- [1] S. Gillessen, F. Eisenhauer, S. Trippe, T. Alexander, R. Genzel, F. Martins and T. Ott, “Monitoring stellar orbits around the Massive Black Hole in the Galactic Center,” *Astrophys. J.* **692**, 1075-1109 (2009).
  - [2] B. P. Abbott *et al.* [LIGO Scientific and Virgo], “GW150914: The Advanced LIGO Detectors in the Era of First Discoveries,” *Phys. Rev. Lett.* **116**, no.13, 131103 (2016).
  - [3] C. G. Darwin, “The gravity field of a particle,” *Proc. Roy. Soc London*, **249**, 180 (1959).

- [4] J. P. Luminet, "Image of a spherical black hole with thin accretion disk," *Astron. Astrophys.* **75**, 228-235 (1979).
- [5] E. F. Eiroa, G. E. Romero and D. F. Torres, "Reissner-Nordstrom black hole lensing," *Phys. Rev. D* **66**, 024010 (2002).
- [6] R. D. Viollier, F. R. Leimgruber and D. Trautmann, "Halos of heavy neutrinos around baryonic stars," *Phys. Lett. B* **297**, 132-137 (1992).
- [7] N. Bilic and R. D. Viollier, "Gravitational phase transition of heavy neutrino matter," *Phys. Lett. B* **408**, 75-80 (1997).
- [8] E. T. Akhmedov, V. Akhmedova and D. Singleton, "Hawking temperature in the tunneling picture," *Phys. Lett. B* **642**, 124 (2006).
- [9] S. H. Rhie and D. P. Bennett, "Global monopoles do not 'collapse'," *Phys. Rev. Lett.* **67**, 1173 (1991).
- [10] P. Jetzer, "Boson stars," *Phys. Rept.* **220**, 163-227 (1992).
- [11] W. Javed, J. Abbas, and A. Övgün, "Deflection angle of photon from magnetized black hole and effect of nonlinear electrodynamics," *Eur. Phys. J. C* **79**, no. 8, 694 (2019).
- [12] S. E. Vazquez and E. P. Esteban, "Strong field gravitational lensing by a Kerr black hole," *Nuovo Cim. B* **119**, 489-519 (2004).
- [13] V. Bozza, "Gravitational lensing in the strong field limit," *Phys. Rev. D* **66**, 103001 (2002).
- [14] H. S. Vieira, V. B. Bezerra and C. R. Muniz, "Exact solutions of the Klein-Gordon equation in the Kerr-Newman background and Hawking radiation," *Annals Phys.* **350**, 14 (2014).
- [15] M. Barriola and A. Vilenkin, "Gravitational Field of a Global Monopole," *Phys. Rev. Lett.* **63**, 341 (1989).
- [16] J. Badia and E. F. Eiroa, "Gravitational lensing by a Horndeski black hole," *Eur. Phys. J. C* **77**, no.11, 779 (2017).
- [17] D. Valls-Gabaud, "The conceptual origins of gravitational lensing," *AIP Conf. Proc.* **861**, no.1, 1163-1171 (2006).
- [18] K. Jusufi, A. Övgün, J. Saavedra, P. A. Gonzalez and Y. Vasquez, "Deflection of light by rotating regular black holes using the Gauss-Bonnet theorem," *Phys. Rev. D* **97**, 124024 (2018).
- [19] V. Akhmedova, T. Pilling, A. de Gill and D. Singleton, "Temporal contribution to gravitational WKB-like calculations," *Phys. Lett. B* **666**, 269 (2008).
- [20] M. K. Parikh and F. Wilczek, "Hawking radiation as tunneling," *Phys. Rev. Lett.* **85**, 5042 (2000).
- [21] A. Einstein, "Lens-Like Action of a Star by the Deviation of Light in the Gravitational Field," *Science* **84**, 506-507 (1936).
- [22] G. W. Gibbons and M. C. Werner, "Applications of the Gauss-Bonnet theorem to gravitational lensing," *Class. Quant. Grav.* **25**, 235009 (2008).
- [23] M. C. Werner, "Gravitational lensing in the Kerr-Randers optical geometry," *Gen. Rel. Grav.* **44**, 3047 (2012).
- [24] A. Ishihara, Y. Suzuki, T. Ono and H. Asada, "Finite distance corrections to the gravitational bending angle of light in the strong deflection limit," *Phys. Rev. D* **95**, no. 4, 044017 (2017).
- [25] A. Ishihara, Y. Suzuki, T. Ono, T. Kitamura and H. Asada, "Gravitational bending angle of light for finite distance and the Gauss-Bonnet theorem," *Phys. Rev. D* **94**, no.8, 084015 (2016).
- [26] T. Ono, A. Ishihara and H. Asada, "Deflection angle of light for an observer and source at finite distance from a rotating global monopole," *Phys. Rev. D* **99**, no. 12, 124030 (2019).
- [27] T. Ono, A. Ishihara and H. Asada, "Deflection angle of light for an observer and source at finite distance from a rotating wormhole," *Phys. Rev. D* **98**, no. 4, 044047 (2018).
- [28] O. Y. Tsupko and G. S. Bisnovatyi-Kogan, "Gravitational lensing in the presence of plasmas and strong gravitational fields," *Grav. Cosmol.* **20**, no.3, 220-225 (2014).
- [29] O. Y. Tsupko and G. S. Bisnovatyi-Kogan, "Gravitational lensing in plasma: Relativistic images at homogeneous plasma," *Phys. Rev. D* **87**, no. 12, 124009 (2013).
- [30] G. Crisnejo and E. Gallo, "Weak lensing in a plasma medium and gravitational deflection of massive particles using the Gauss-Bonnet theorem. A unified treatment," *Phys. Rev. D* **97**, no.12, 124016 (2018).
- [31] W. Javed, J. Abbas and A. Övgün, "Effect of the Hair on Deflection Angle by Asymptotically Flat Black Holes in Einstein-Maxwell-Dilaton Theory," *Phys. Rev. D* **100**, no. 4, 044052 (2019).
- [32] K. Nozari and S. Hamid Mehdipour, "Quantum Gravity and Recovery of Information in Black Hole Evaporation," *EPL* **84**, no. 2, 20008 (2008).
- [33] M. Sharif and W. Javed, "Fermions Tunneling from Charged Accelerating and Rotating Black Holes with NUT Parameter," *Eur. Phys. J. C* **72**, 1997 (2012).
- [34] K. Jusufi and A. Övgün, "Gravitational Lensing by Rotating Wormholes," *Phys. Rev. D* **97**, no. 2, 024042 (2018).
- [35] G. W. Gibbons and C. M. Warnick, "Universal properties of the near-horizon optical geometry," *Phys. Rev. D* **79**, 064031 (2009).
- [36] G. W. Gibbons, C. A. R. Herdeiro, C. M. Warnick and M. C. Werner, "Stationary Metrics and Optical Zermelo-Randers-Finsler Geometry," *Phys. Rev. D* **79**, 044022 (2009).
- [37] G. W. Gibbons and M. Vyska, "The Application of Weierstrass elliptic functions to Schwarzschild Null Geodesics," *Class. Quant. Grav.* **29**, 065016 (2012).
- [38] C. Bloomer, "Optical Geometry of the Kerr Space-time," [arXiv:1111.4998 [math-ph]].
- [39] G. W. Gibbons, "The Jacobi-metric for timelike geodesics in static spacetimes," *Class. Quant. Grav.* **33**, no.2, 025004 (2016).
- [40] P. Das, R. Sk and S. Ghosh, "Motion of charged particle in Reissner-Nordstrom spacetime: a Jacobi-metric approach," *Eur. Phys. J. C* **77**, no.11, 735 (2017).
- [41] I. Sakalli and A. Övgün, "Hawking Radiation and Deflection of Light from Rindler Modified Schwarzschild Black Hole," *EPL* **118**, no.6, 60006 (2017).
- [42] K. Jusufi, M. C. Werner, A. Banerjee and A. Övgün, "Light Deflection by a Rotating Global Monopole Spacetime," *Phys. Rev. D* **95**, no.10, 104012 (2017).
- [43] T. Ono, A. Ishihara and H. Asada, "Gravitomagnetic bending angle of light with finite-distance corrections in stationary axisymmetric spacetimes," *Phys. Rev. D* **96**, no. 10, 104037 (2017).



- [44] K. Jusufi, I. Sakalli and A. Övgün, "Effect of Lorentz Symmetry Breaking on the Deflection of Light in a Cosmic String Spacetime," *Phys. Rev. D* **96**, no. 2, 024040 (2017).
- [45] H. Arakida, "Light deflection and Gauss-Bonnet theorem: definition of total deflection angle and its applications," *Gen. Rel. Grav.* **50**, no.5, 48 (2018).
- [46] A. Övgün, I. Sakalli and J. Saavedra, "Shadow cast and Deflection angle of Kerr-Newman-Kasuya spacetime," *JCAP* **1810**, no. 10, 041 (2018)
- [47] A. Övgün, I. Sakalli and J. Saavedra, "Weak gravitational lensing by Kerr-MOG Black Hole and Gauss-Bonnet theorem," *Annals Phys.* **411**, 167978 (2019).
- [48] A. Övgün, G. Gyulchev and K. Jusufi, "Weak Gravitational lensing by phantom black holes and phantom wormholes using the Gauss-Bonnet theorem," *Annals Phys.* **406**, 152 (2019).
- [49] A. Övgün, K. Jusufi and I. Sakalli, "Gravitational lensing under the effect of Weyl and bumblebee gravities: Applications of Gauss-Bonnet theorem," *Annals Phys.* **399**, 193 (2018).
- [50] K. Jusufi and A. Övgün, "Effect of the cosmological constant on the deflection angle by a rotating cosmic string," *Phys. Rev. D* **97**, no. 6, 064030 (2018).
- [51] A. Övgün, "Light deflection by Damour-Solodukhin wormholes and Gauss-Bonnet theorem," *Phys. Rev. D* **98**, no. 4, 044033 (2018)
- [52] T. Ono, A. Ishihara and H. Asada, "Deflection angle of light for an observer and source at finite distance from a rotating global monopole," *Phys. Rev. D* **99**, no. 12, 124030 (2019).
- [53] G. Crisnejo, E. Gallo and A. Rogers, "Finite distance corrections to the light deflection in a gravitational field with a plasma medium," *Phys. Rev. D* **99**, no. 12, 124001 (2019).
- [54] Z. Li and T. Zhou, "Equivalence of Gibbons-Werner method to geodesics method in the study of gravitational lensing," *Phys. Rev. D* **101**, no.4, 044043 (2020).
- [55] Z. Li and J. Jia, "The finite-distance gravitational deflection of massive particles in stationary spacetime: a Jacobi metric approach," *Eur. Phys. J. C* **80**, no. 2, 157 (2020).
- [56] Z. Li, G. He and T. Zhou, "Gravitational deflection of relativistic massive particles by wormholes," *Phys. Rev. D* **101**, no.4, 044001 (2020).
- [57] A. Övgün, "Deflection Angle of Photons through Dark Matter by Black Holes and Wormholes Using Gauss-Bonnet Theorem," *Universe* **5**, no. 5, 115 (2019).
- [58] W. Javed, R. Babar, A. Övgün, "The effect of the Brane-Dicke coupling parameter on weak gravitational lensing by wormholes and naked singularities," *Phys. Rev. D* **99**, no. 8, 084012 (2019).
- [59] W. Javed, A. Hazma and A. Övgün, "Effect of Non-Linear Electrodynamics on Weak Field Deflection Angle by Black Hole," *Phys. Rev. D* **101**, no.10, 103521 (2020).
- [60] A. Övgün, "Weak field deflection angle by regular black holes with cosmic strings using the Gauss-Bonnet theorem," *Phys. Rev. D* **99**, no. 10, 104075 (2019).
- [61] K. Jusufi and A. Övgün, "Light Deflection by a Quantum Improved Kerr Black Hole Pierced by a Cosmic String," *Int. J. Geom. Meth. Mod. Phys.* **16**, no. 08, 1950116 (2019).
- [62] G. Crisnejo, E. Gallo and J. R. Villanueva, "Gravitational lensing in dispersive media and deflection angle of charged massive particles in terms of curvature scalars and energy-momentum tensor," *Phys. Rev. D* **100**, no.4, 044006 (2019)
- [63] W. Javed, R. Babar, and A. Övgün, "Effect of the dilaton field and plasma medium on deflection angle by black holes in Einstein-Maxwell-dilaton-axion theory," *Phys. Rev. D* **100**, no. 10, 104032 (2019)
- [64] K. de Leon and I. Vega, "Weak gravitational deflection by two-power-law densities using the Gauss-Bonnet theorem," *Phys. Rev. D* **99**, no.12, 124007 (2019).
- [65] Y. Kumaran and A. Övgün, "Weak Deflection Angle of Extended Uncertainty Principle Black Holes," *Chin. Phys. C* **44**, no.2, 025101 (2020).
- [66] W. Javed, J. Abbas, Y. Kumaran and A. Övgün, Preprints 2019, 2019110210 (doi: 10.20944/preprints201911.0210.v1).
- [67] A. Övgün, I. Sakalli and J. Saavedra, "Effect of aether field on weak deflection angle of black holes," [arXiv:1908.04261 [gr-qc]], accepted for publication in Chinese Physics C, 10.1088/1674-1137/abb532.
- [68] K. Jusufi, A. Övgün, A. Banerjee, and I. Sakalli, "Gravitational lensing by wormholes supported by electromagnetic, scalar, and quantum effects," *Eur. Phys. J. Plus* **134**, no. 9, 428 (2019).
- [69] K. Jusufi, A. Övgün, and A. Banerjee, "Light deflection by charged wormholes in Einstein-Maxwell-dilaton theory," *Phys. Rev. D* **96**, no. 8, 084036 (2017).
- [70] Z. Li and A. Övgün, "Finite-Distance Gravitational Deflection of Massive Particles by the Kerr-like Black Hole in the Bumblebee Gravity Model," *Phys. Rev. D* **101**, no. 2, 024040 (2020).
- [71] A. Övgün and I. Sakalli, "Testing Generalized Einstein-Cartan-Kibble-Sciama Gravity Using Weak Deflection Angle and Shadow Cast," doi:10.1088/1361-6382/abb579 [arXiv:2005.00982 [gr-qc]].
- [72] H. El Mounni, K. Masmar and A. Övgün, "Weak Deflection angle of some classes of non-linear electrodynamics black holes via Gauss-Bonnet Theorem," [arXiv:2008.06711 [gr-qc]].
- [73] W. Javed, M. B. Khadim and A. Övgün, "Weak gravitational lensing by BocharovaBronnikovMelnikovBekenstein black holes using Gauss-Bonnet theorem," *Eur. Phys. J. Plus* **135**, no.7, 595 (2020).
- [74] Z. Li, G. Zhang and A. Övgün, "Circular Orbit of a Particle and Weak Gravitational Lensing," *Phys. Rev. D* **101**, no.12, 124058 (2020).
- [75] W. Javed, M. B. Khadim, A. Övgün and J. Abbas, "Weak gravitational lensing by stringy black holes," *Eur. Phys. J. Plus* **135**, no.3, 314 (2020).
- [76] S. H. Mazharimousavi and M. Halilsoy, "Einstein-nonlinear Maxwell-Yukawa black hole," *Int. J. Mod. Phys. D* **28**, no.09,

1950120 (2019).

- [77] M. Abramowitz and Stegun, Handbook of Mathematical Functions, (United States Department of Commerce, National Institute of Standards and Technology (NBS), 1967).
- [78] G. G. L. Nashed and S. Capozziello, "Charged spherically symmetric black holes in  $f(R)$  gravity and their stability analysis," Phys. Rev. D **99**, no.10, 104018 (2019).
- [79] Z. Horvath, L. A. Gergely, D. Hobill, S. Capozziello and M. De Laurentis, "Weak gravitational lensing by compact objects in fourth order gravity," Phys. Rev. D **88**, no.6, 063009 (2013).
- [80] S. Capozziello, V. F. Cardone and A. Troisi, "Gravitational lensing in fourth order gravity," Phys. Rev. D **73**, 104019 (2006).