

# COHOMOLOGY THEORY OF AVERAGING ALGEBRAS, $L_\infty$ -STRUCTURES AND HOMOTOPY AVERAGING ALGEBRAS

KAI WANG AND GUODONG ZHOU

**ABSTRACT.** This paper studies averaging algebras, say, associative algebras endowed with averaging operators. We develop a cohomology theory for averaging algebras and justify it by interpreting lower degree cohomology groups as formal deformations and abelian extensions of averaging algebras. We make explicit the  $L_\infty$ -algebra structure over the cochain complex defining cohomology groups and introduce the notion of homotopy averaging algebras as Maurer-Cartan elements of this  $L_\infty$ -algebra.

## CONTENTS

|   |    |
|---|----|
| 1. Introduction                                 | 1  |
| 2. Averaging algebras and their bimodule        | 2  |
| 3. Cohomology theory of averaging algebras      | 5  |
| 3.1. Cohomology of averaging operators          | 5  |
| 3.2. Cohomology of averaging algebras           | 7  |
| 3.3. Relationship among the cohomologies.       | 8  |
| 4. Formal deformation of averaging algebras     | 9  |
| 5. Abelian extensions of averaging algebras     | 11 |
| 6. $L_\infty$ -structure on the cochain complex | 14 |
| 7. Homotopy averaging algebras                  | 19 |
| Appendix: Proof of Theorem 6.5                  | 20 |
| References                                      | 26 |

## 1. INTRODUCTION

Throughout this paper,  $\mathbf{k}$  denotes a field. All the vector spaces and algebras are over  $\mathbf{k}$  and all tensor products are also taking over  $\mathbf{k}$ .

Let  $R$  be an associative algebra over field  $\mathbf{k}$ . An averaging operator over  $R$  is a  $\mathbf{k}$ -linear map  $A : R \rightarrow R$  such that

$$A(x)A(y) = A(A(x)y) = A(xA(y))$$

for all  $x, y \in R$ .

The research on averaging operators has a long history and it appears in various mathematical branches from turbulence theory to functional analysis, probability theory etc. When investigating turbulence theory, Reynolds already studied implicitly the averaging operator in a famous

---

*Date:* October 19, 2021.

*1991 Mathematics Subject Classification.* 16E40 16S80 16S70 17B38 .

*Key words and phrases.* averaging algebra, averaging operator, cohomology, extension, deformation, homotopy averaging algebra,  $L_\infty$ -algebra.

paper [20] published in 1895. In a series of papers of 1930s, Kolmogoroff and Kampé de Fériet introduced explicitly the averaging operator in the context of turbulence theory and functional analysis [12][17]. Birkhoff continued the line of research in [3]. Moy investigated averaging operators from the viewpoint of conditional expectation in probability theory [18]. Kelley [13] and Rota [22] studied the role of averaging operators in Banach algebras.

Contrary to the above studies of analytic nature, the algebraic study on averaging operators began with the Ph.D thesis of Cao [5]. He constructed explicitly free unitary commutative averaging algebras and discovered the Lie algebra structures induced naturally from averaging operators. Inspired by the work of Loday [16] who defined a dialgebra as the enveloping algebra of a Leibniz algebra, Aguiar associated a dissociative algebra to an averaging associative algebra [1]. In a paper [19] Guo and Pei studied averaging operators from an algebraic and combinatorial point of view. They first construct free nonunital averaging algebras in terms of a class of bracketed words called averaging words, related them to large Schröder numbers and unreduced trees from an operadic point of view. Another paper by Gao and Zhang [9] contains an explicit construction of free unital averaging algebras in terms of bracketed polynomials and the main tools were rewriting systems and Gröbner-Shirshov bases.

The averaging operators attract much attention also partly because of their closely relationship to other operators such as Reynolds operators, symmetric operators and Rota-Baxter operators [4][8][25], the latter having many applications in many other fields of mathematics [2][11].

In the paper we study averaging operators from the viewpoint of deformation theory. We construct a cohomology theory which controls simultaneous deformations of associative multiplication and the averaging operator. Because of the complexity of the problem, the cochain complex is no longer a differential graded Lie algebra (DGLA) but an  $L_\infty$ -algebra. We give explicitly the higher Lie brackets over the cochain complex. As a byproduct, we define homotopy averaging algebras as Maurer-Cartan elements of this  $L_\infty$ -algebra.

We would like to mention several papers which are somehow parallel to this paper. Motivated from geometry and physics, Sheng, Tang and Zhu [23] studies embedding tensors (another name of averaging operators in physics) for Lie algebras and they construct a cohomology theory for such operators on Lie algebras using derived brackets as a main tool. Another work of Chen, Ge and Xiang [6] studies embedding tensors using operadic tool and they compute explicitly the Boardmann-Vogt resolution of the colored operad governing embedding tensors. We warmly recommend the comparative reading of their papers.

The layout of this paper is as follows: The first section contains basic definitions and facts about averaging algebras and bimodules over them. We define the cohomology theory of averaging operators and averaging algebras in Section 2. Deformation theory of averaging algebras is developed in Section 4 and abelian extensions of averaging algebras are interpreted as the second cohomology group in Section 5. We construct explicitly the  $L_\infty$ -algebra structure over the cochain complex computing cohomology theory of averaging algebras in Section 6 and in the last section we introduce the notion of homotopy averaging algebras as Maurer-Cartan elements of this  $L_\infty$ -algebra.

## 2. AVERAGING ALGEBRAS AND THEIR BIMODULE

In this section, we introduce averaging algebras and bimodules over them and present some basic observations.

**Definition 2.1.** Let  $(R, \cdot)$  be an associative algebra over field  $\mathbf{k}$ . If is endowed with a linear operator  $A : R \rightarrow R$  such that

$$(1) \quad A(x)A(y) = A(xA(y)) = A(A(x)y), \forall x, y \in R,$$

then we call  $(R, \cdot, A)$  an averaging algebra.

Given two averaging algebras  $(R, \cdot, A)$  and  $(R', \cdot', A')$ , a morphism of averaging algebras from  $(R, \cdot, A)$  to  $(R', \cdot', A')$  is a homomorphism of algebras  $\phi : (R, \cdot) \rightarrow (R', \cdot')$  satisfying  $\phi \circ A = A' \circ \phi$ .

**Definition 2.2.** Let  $(R, \cdot, A)$  be an averaging algebra.

(i) A bimodule over the averaging algebra  $(R, \cdot, A)$  is a bimodule  $M$  over associative algebra  $(R, \cdot)$  endowed with an operator  $A_M : M \rightarrow M$ , such that for any  $r \in R, m \in M$ , the following equalities hold:

$$(2) \quad A(r)A_M(m) = A_M(A(r)m) = A_M(rA_M(m)),$$

$$(3) \quad A_M(m)A(r) = A_M(A_M(m)r) = A_M(mA(r)).$$

(ii) Given two bimodules  $(M, A_M)$  and  $(N, A_N)$  over averaging algebra  $(R, \cdot, A)$ , a morphism from  $(M, A_M)$  to  $(N, A_N)$  is a bimodule morphism  $f : M \rightarrow N$  over  $(R, \cdot)$  such that :

$$f \circ A_M = A_N \circ f.$$

The algebra  $R$  itself is naturally a bimodule over  $(R, \cdot, A)$  with  $A_R = A$ , called the regular bimodule.

The following easy result is left as an exercise.

**Proposition 2.3.** Let  $(M, A_M)$  be a bimodule over averaging algebra  $(R, \cdot, A)$ . Then  $R \oplus M$  is an averaging algebra, where the averaging operator is  $A \oplus A_M$  and the multiplication is:

$$(a, m) \cdot (b, n) = (ab, an + mb),$$

for all  $a, b \in R, m, n \in M$ .

**Proposition 2.4.** Let  $(R, \cdot, A)$  be an averaging algebra. Then the following operations:

$$a \star b := aA(b),$$

$$a \diamond b := A(a)b$$

are both associative.

Conversely, let  $(R, \cdot)$  be a unital algebra over  $\mathbf{k}$  endowed with a linear operator  $A$  such that operations  $\star, \diamond$  are associative. Then  $(R, \cdot, A)$  is an averaging algebra.

*Proof.* By definition, we have identities for any  $a, b, c \in R$ :

$$\begin{aligned} (a \star b) \star c &= (a \cdot A(b)) \star c \\ &= a \cdot A(b)A(c) \\ &= a \cdot (A(b \cdot Ac)) \\ &= a \star (b \star c). \end{aligned}$$

Thus the operation  $\star$  is associative. Analogously, one can check the associativity of operation  $\diamond$ .

Conversely, assume that unital algebra  $(R, \cdot)$  is endowed with a linear map  $A : R \rightarrow R$  such that operations  $\star, \diamond$  is associative. Then

$$\begin{aligned}
 A(a)A(b) &= (1_R \cdot A(a))A(b) \\
 &= (1_R \star a) \star b \\
 &= 1_R \star (a \star b) \\
 &= 1_R \star (a \cdot A(b)) \\
 &= 1_R \cdot A(a \cdot A(b)) \\
 &= A(a \cdot (A(b))).
 \end{aligned}$$

hold for any  $a, b \in R$ . Similarly, equality  $A(a)A(b) = A(A(a)b)$  also holds. Thus  $(R, \cdot, A)$  is an averaging algebra.  $\square$

**Proposition 2.5.** *Let  $(R, \cdot, A)$  be an averaging algebra. Then  $(R, \star, A), (R, \diamond, A)$  are also averaging algebras.*

*Proof.* For any  $a, b \in R$ , we have

$$\begin{aligned}
 A(a) \star A(b) &= A(a) \cdot (A(A(b))) \\
 &= A(a \cdot A^2(b)) \\
 &= A(A(a) \cdot A(b)),
 \end{aligned}$$

and

$$\begin{aligned}
 A(a \star A(b)) &= A(a \cdot A^2(b)), \\
 A(A(a) \star b) &= A(A(a) \cdot A(b)).
 \end{aligned}$$

Thus the relations  $A(a) \star A(b) = A(a \star A(b)) = A(A(a) \star b)$  hold. So  $(R, \star, A)$  is an averaging algebra. It is analogous to prove that  $(R, \diamond, A)$  is an averaging algebra.  $\square$

Let  $(R, \cdot, A)$  be an averaging algebra and  $(M, A_M)$  be a bimodule over  $(R, \cdot, A)$ . Then we can make  $M$  into a bimodule over  $(R, \star)$  and  $(R, \diamond)$ . For any  $a \in R$  and  $m \in M$ , we define the following actions:

$$a \vdash m := A(a)m - A_M(am), \quad m \dashv a := mA(a)$$

and

$$a \triangleright m := A(a)m, \quad m \triangleleft a := mA(a) - A_M(mb).$$

**Proposition 2.6.** *The action  $(\vdash, \dashv)$  (resp.  $(\triangleright, \triangleleft)$ ) makes  $M$  become a bimodule over associative algebra  $(R, \star)$  (resp.  $(R, \diamond)$ ).*

*Proof.* For any  $a, b \in R$  and  $m \in M$ , we have

$$\begin{aligned}
 a \vdash (b \vdash m) &= a \vdash (A(b)m - A_M(bm)) \\
 &= A(a) \cdot (A(b)m - A_M(bm)) - A_M(a \cdot A(b) \cdot m - a \cdot A_M(b \cdot m)) \\
 &= A(a) \cdot A(b) \cdot m - A_M(a \cdot A(b) \cdot m),
 \end{aligned}$$

$$\begin{aligned}
(a \star b) \vdash m & \\
&= (a \cdot A(b)) \vdash m \\
&= A(a \cdot A(b)) \cdot m - A_M(a \cdot A(b)) \cdot m \\
&= a \vdash (b \vdash m).
\end{aligned}$$

Thus operation  $\vdash$  makes  $M$  into a left module over  $(R, \star)$ . Analogously, operation  $\dashv$  makes  $M$  into a right module over  $(R, \star)$ .

Moreover, we have :

$$\begin{aligned}
a \vdash (m \dashv b) & \\
&= a \vdash (m \cdot A(b)) \\
&= A(a) \cdot m \cdot A(b) - A_M(a \cdot m \cdot A(b)) \\
&= A(a) \cdot m \cdot A(b) - A_M(a \cdot m) \cdot A(b) \\
&= (A(a) \cdot m - A_M(a \cdot m)) \cdot A(b) \\
&= (a \vdash m) \cdot A(b) \\
&= (a \vdash m) \dashv b
\end{aligned}$$

Thus operations  $(\vdash, \dashv)$  makes  $M$  into a bimodule over associative algebra  $(R, \star)$ .

Similarly, one can check that operations  $(\triangleright, \triangleleft)$  makes  $M$  into a bimodule over  $(R, \diamond)$ .

□

### 3. COHOMOLOGY THEORY OF AVERAGING ALGEBRAS

Let  $M$  be a bimodule over an associative algebra  $R$ . Recall that the Hochschild cohomology of  $R$  with coefficients in  $M$ :

$$(C_{\text{Alg}}^\bullet(A, M) = \bigoplus_{n=0}^{\infty} C_{\text{Alg}}^n(R, M), \delta),$$

where  $C_{\text{Alg}}^n(R, M) = (\text{Hom}(R^{\otimes n}, M))$  and the differential  $\delta : C_{\text{Alg}}^n(R, M) \rightarrow C_{\text{Alg}}^{n+1}(R, M)$  is given by

$$\begin{aligned}
\delta(f)(x_1 \otimes \dots \otimes x_{n+1}) &= x_1 f(x_2 \otimes \dots \otimes x_n) + \sum_{i=1}^n (-1)^i f(x_1 \otimes \dots \otimes x_i x_{i+1} \otimes \dots \otimes x_{n+1}) \\
&\quad + (-1)^{n+1} f(x_1 \otimes \dots \otimes x_n) x_{n+1}
\end{aligned}$$

for all  $f \in C^n(R, M)$ ,  $x_1, \dots, x_{n+1} \in R$ . The corresponding Hochschild cohomology is denoted  $\text{HH}^\bullet(R, M)$ . When  $M = R$ , just denote the Hochschild cochain complex with coefficients in  $R$  by  $C_{\text{Alg}}^\bullet(R)$  and denote the Hochschild cohomology by  $\text{HH}^\bullet(R)$ .

**3.1. Cohomology of averaging operators.** Let  $(R, \cdot, A)$  be an averaging algebra and  $(M, A_M)$  be a bimodule over  $(R, \cdot, A)$ . In this subsection, we define the cohomology of averaging operators. By Proposition 2.4, the averaging operator  $A$  induces two new multiplications  $\star$  and  $\diamond$  on  $R$ , and by Proposition 2.6, operations  $(\vdash, \dashv)$  (resp.  $(\triangleright, \triangleleft)$ ) makes  $M$  into a bimodule over  $(R, \star)$  (resp.  $(R, \diamond)$ ).

Consider the Hochschild cochain complex of  $(R, \star)$  with the coefficient in bimodule  $(M, \vdash, \dashv)$ . Denote this cochain complex by

$$C_r^\bullet(R, M) = \bigoplus_{i=0}^{\infty} C_r^n(R, M),$$

where  $C_r^n(R, M) = \text{Hom}(R^{\otimes n}, M)$  and its differential  $\partial_r : C_r^n(R, M) \rightarrow C_r^{n+1}(R, M)$  is given by :

$$\begin{aligned} \partial_r(f)(x_1 \otimes \dots \otimes x_{n+1}) &= x_1 \vdash f(x_2 \otimes \dots \otimes x_{n+1}) + \sum_{i=1}^n (-1)^i f(x_1 \otimes \dots \otimes x_i \star x_{i+1} \otimes \dots \otimes x_{n+1}) \\ &\quad + (-1)^{n+1} f(x_1 \otimes \dots \otimes x_n) \dashv x_{n+1} \\ &= A(x_1)f(x_2 \otimes \dots \otimes x_{n+1}) - A_M(x_1 f(x_2 \otimes \dots \otimes x_{n+1})) \\ &\quad + \sum_{i=1}^n (-1)^i f(x_1 \otimes \dots \otimes x_i A(x_{i+1}) \otimes \dots \otimes x_{n+1}) \\ &\quad + (-1)^{n+1} f(x_1 \otimes \dots \otimes x_n) A(x_{n+1}). \end{aligned}$$

for all  $f \in C_r^n(R, M)$ ,  $x_1, \dots, x_{n+1} \in R$ .

For algebra  $(R, \diamond)$ , we denote its Hochschild cochain complex with coefficients in bimodule  $(M, \triangleright, \triangleleft)$  by

$$C_l^\bullet(R, M) = \bigoplus_{i=0}^{\infty} C_l^n(R, M),$$

where  $C_l^n(R, M) = \text{Hom}(R^{\otimes n}, M)$  and by definition, its differential  $\partial_l : C_l^n(R, M) \rightarrow C_l^{n+1}(R, M)$  is given by:

$$\begin{aligned} \partial_l(g)(x_1 \otimes \dots \otimes x_{n+1}) &= x_1 \triangleright g(x_2 \otimes \dots \otimes x_{n+1}) + \sum_{i=1}^n (-1)^i g(x_1 \otimes \dots \otimes x_i \diamond x_{i+1} \otimes \dots \otimes x_{n+1}) \\ &\quad + (-1)^{n+1} g(x_1 \otimes \dots \otimes x_n) \triangleleft x_{n+1} \\ &= A(x_1)g(x_2 \otimes \dots \otimes x_{n+1}) + \sum_{i=1}^n (-1)^i g(x_1 \otimes \dots \otimes A(x_i)x_{i+1} \otimes \dots \otimes x_{n+1}) \\ &\quad + (-1)^{n+1} g(x_1 \otimes \dots \otimes x_n) A(x_{n+1}) + (-1)^n A_M(g(x_1 \otimes \dots \otimes x_n)x_{n+1}) \end{aligned}$$

for all  $g \in C_l^n(R, M)$ ,  $x_1, \dots, x_n \in R$ . When  $M = R$ , we just denote  $C_l^\bullet(R, R)$  by  $C_l^\bullet(R)$ , and denote  $C_r^\bullet(R, R)$  by  $C_r^\bullet(R)$ .

Identifying  $C_r^0(R, M)$  with  $C_l^0(R, M)$ , identifying  $C_l^1(R, M)$  with  $C_r^1(R, M)$  and taking the direct sum of  $C_l^n(R, M)$  and  $C_r^n(R, M)$  for  $n \geq 2$ , we get the following complex:

$$\text{Hom}(k, M) \xrightarrow{\partial_0} \text{Hom}(R, M) \xrightarrow{\begin{pmatrix} \partial_r \\ \partial_l \end{pmatrix}} \begin{pmatrix} C_r^2(R, M) \\ \bigoplus \\ C_l^2(R, M) \end{pmatrix} \xrightarrow{\begin{pmatrix} \partial_r & 0 \\ 0 & \partial_l \end{pmatrix}} \dots \dots \begin{pmatrix} C_r^n(R, M) \\ \bigoplus \\ C_l^n(R, M) \end{pmatrix} \xrightarrow{\begin{pmatrix} \partial_r & 0 \\ 0 & \partial_l \end{pmatrix}} \begin{pmatrix} C_r^{n+1}(R, M) \\ \bigoplus \\ C_l^{n+1}(R, M) \end{pmatrix} \dots \dots,$$

where  $\partial_0$  is defined as: for  $f \in \text{Hom}(k, M)$ , assume that  $f(1) = m$ , then

$$\partial_0(f)(r) = A_M(mr) - A_M(rm) - mA(r) + A(r)m$$

for any  $r \in R$ . One can check that

$$\begin{pmatrix} \partial_r \\ \partial_l \end{pmatrix} \circ \partial_0 = 0.$$

We denote the above complex by  $C_{\text{AvO}}^\bullet(R, M)$ , called the *cochain complex of the averaging operator  $A$  with coefficients in bimodule  $(M, A_M)$* . Its cohomology is denoted by  $H_{\text{AvO}}^\bullet(R, M)$ , called the *cohomology of the averaging operator  $A$  of averaging algebra  $(R, \cdot, A)$  with coefficients in  $(M, A_M)$* . When  $(M, A_M) = (R, A)$ , we just denote  $C_{\text{AvO}}^\bullet(R, R)$  by  $C_{\text{AvO}}^\bullet(R)$  and call it the cochain complex of averaging operator. We denote  $H_{\text{AvO}}^\bullet(R, M)$  by  $H_{\text{AvO}}^\bullet(R)$ , called the cohomology of averaging operator  $A$ .

**3.2. Cohomology of averaging algebras.** In this subsection, we'll define the cohomology of averaging algebras and such a cohomology theory involves both the multiplication and the averaging operator in an averaging algebra.

Firstly, let's build a morphism of complexes:  $\Phi : C_{\text{Alg}}^\bullet(R, M) \rightarrow C_{\text{AvO}}^\bullet(R, M)$ ,

$$\begin{array}{ccccccc} C_{\text{Alg}}^0(R, M) & \xrightarrow{\delta^0} & C_{\text{Alg}}^1(R, M) & \xrightarrow{\delta^1} & C_{\text{Alg}}^2(R, M) & \cdots & C_{\text{Alg}}^n(R, M) \xrightarrow{\delta^n} C_{\text{Alg}}^{n+1}(R, M) \cdots \\ \downarrow \Phi^0 & & \downarrow \Phi^1 & & \downarrow \Phi^2 & & \downarrow \Phi^n \\ \text{Hom}(k, M) & \xrightarrow{\partial^0} & \text{Hom}(R, M) & \xrightarrow{\begin{pmatrix} \partial_r^1 \\ \partial_l^1 \end{pmatrix}} & C_r^2(R, M) \oplus C_l^2(R, M) & \cdots & C_r^n(R, M) \oplus C_l^n(R, M) \xrightarrow{\begin{pmatrix} \partial_r^n & 0 \\ 0 & \partial_l^n \end{pmatrix}} C_r^{n+1}(R, M) \oplus C_l^{n+1}(R, M) \cdots \\ & & & & \downarrow \Phi^{n+1} & & \downarrow \Phi^{n+1} \end{array}$$

Here we define

$$\Phi^0 = \text{id}, \Phi^1(f) = f \circ A - A_M \circ f$$

for any  $f \in \text{Hom}(R, M)$ , and when  $n \geq 2$ ,  $\Phi^n = \begin{pmatrix} \Phi_r^n \\ \Phi_l^n \end{pmatrix}$ , where  $\Phi_r^n : C^n(R)$

$$\Phi_r^n(f) = f \circ A^{\otimes n} - A_M \circ f \circ (\text{id} \otimes A^{\otimes n-1}),$$

$$\Phi_l^n(f) = f \circ A^{\otimes n} - A_M \circ f \circ (A^{\otimes n-1} \otimes \text{id})$$

for any  $f \in C^n(R, M)$ .

**Proposition 3.1.** *The morphism  $\Phi : C^\bullet(R, M) \rightarrow C_{\text{AO}}^\bullet(R, M)$  is compatible with differentials.*

*Proof.* The equalities

$$\Phi^1 \circ \delta^0 = \partial^0 \circ \Phi^0$$

and

$$\Phi^2 \circ \delta^1 = \begin{pmatrix} \partial_r^1 \\ \partial_l^1 \end{pmatrix} \circ \Phi^1$$

are easy to verify.

For  $n \geq 2$ , we need to show

$$\begin{pmatrix} \Phi_r^{n+1} \\ \Phi_l^{n+1} \end{pmatrix} \circ \delta^n = \begin{pmatrix} \partial_r^n & 0 \\ 0 & \partial_l^n \end{pmatrix} \circ \begin{pmatrix} \Phi_r^n \\ \Phi_l^n \end{pmatrix},$$

that is,

$$\Phi_r^{n+1} \circ \delta^n = \partial_r^n \circ \Phi_r^n$$

and

$$\Phi_l^{n+1} \circ \delta^n = \partial_r^n \circ \Phi_l^n.$$

We only prove the equality the first one, the second being similar.

In fact, for  $f \in \text{Hom}(R^{\otimes n+1}, M)$  and  $x_1, \dots, x_{n+1} \in R$ , we have

$$\begin{aligned} \Phi_r^{n+1}(\delta^n f)(x_{1,n+1}) &= \delta^n(f)(A(x)_{1,n+1}) - A_M \circ \delta^n(f)(x_1 \otimes A(x)_{2,n+1}) \\ &= A(x_1)f(A(x)_{2,n+1}) + \sum_{i=1}^n (-1)^i f(A(x)_{1,i-1} \otimes A(x_i)A(x_{i+1}) \otimes A(x)_{i+2,n+1}) \\ &\quad + (-1)^{n+1} f(A(x)_{1,n})A(x_{n+1}) \\ &\quad - A_M(x_1 f(A(x)_{2,n+1})) + A_M \circ f(x_1 A(x_2) \otimes A(x)_{3,n+1}) \\ &\quad - \sum_{i=2}^n (-1)^i A_M(f(x_1 \otimes A(x)_{2,i-1} \otimes A(x_i)A(x_{i+1}) \otimes A(x)_{i+2,n+1})) \\ &\quad - (-1)^{n+1} A_M \circ f(x_1 \otimes A(x)_{2,n})A(x_{n+1}), \end{aligned}$$

where we use the notations: for  $i \leq j$ ,  $x_{i,j} = x_i \otimes \dots \otimes x_j$  and  $A(x)_{i,j} = A(x_i) \otimes \dots \otimes A(x_j)$ ; for  $i > j$ , they are  $1 \in \mathbf{k}$ . On the other hand,

$$\begin{aligned} \partial_r^n \circ \Phi_l^n(x_{1,n+1}) &= A(a_1)\Phi_r^n(f)(x_{2,n+1}) - A_M(x_1 \Phi_r^n(f)(x_{2,n+1})) \\ &\quad + \sum_{i=1}^n (-1)^i \Phi_r^n(f)(x_{1,i-1} \otimes x_i A(x_{i+1}) \otimes x_{i+2,n+1}) \\ &\quad + (-1)^{n+1} \Phi_r^n(f)(x_{1,n})A(x_{n+1}) \\ &= A(x_1)f(A(x)_{2,n+1}) - A(a_1)A_M \circ f(a_2 \otimes A(a)_{3,n+1}) \\ &\quad - A_M(x_1 f(A(x)_{2,n+1})) + A_M(x_1 f(x_2 \otimes A(a)_{3,n+1})) \\ &\quad + \sum_{i=1}^n f(A(a)_{1,i-1} \otimes A(x_i)A(x_{i+1})) \otimes A(x)_{i+2,n+1} \\ &\quad - \sum_{i=2}^n (-1)^i A_M \circ f(x_1 \otimes A(x)_{2,i-1} \otimes A(x_i)A(x_{i+1})) \otimes A(x)_{i+2,n+1} \\ &\quad + A_M(f(x_1 A(x_2) \otimes A(x)_{3,n+1}) + (-1)^{n+1} f(A(x)_{1,n})A(x_{n+1})) \\ &\quad - (-1)^{n+1} A_M \circ f(x_1 \otimes A(x)_{2,n})A(x_{n+1}). \end{aligned}$$

We obtain  $\Phi_r^{n+1}(\delta^n f)(x_{1,n+1}) = \partial_r^n \circ \Phi_l^n(x_{1,n+1})$ , because

$$A(a_1)A_M \circ f(a_2 \otimes A(a)_{3,n+1}) = A_M(x_1 f(x_2 \otimes A(a)_{3,n+1}))$$

and  $A(x_i A(x_{i+1})) = A(x_i)A(x_{i+1})$ .

The proof is done.  $\square$

Multiplying  $\Phi^n$  by  $(-1)^n$ , then the above commutative diagram becomes a bicomplex. Taking its totalization, we obtain a cochain complex, and denote it by  $C_{\text{AvA}}^\bullet(R, M)$ .

**Definition 3.2.** The cohomology of the cochain complex  $C_{\text{AvA}}^\bullet(R)$ , denoted by  $H_{\text{AvA}}^\bullet(R, M)$  is called the cohomology of the averaging algebra  $(R, \cdot, A)$  with coefficients in bimodule  $(M, A_M)$ . When  $(M, A_M) = (R, A)$ ,  $H_{\text{AvA}}^\bullet(R, R)$  is called the cohomology of averaging algebra  $(R, \cdot, A)$  and denoted by  $H_{\text{AvA}}^\bullet(R)$ .

**3.3. Relationship among the cohomologies.** By the commutative diagram in the last subsection, we have a canonical short exact sequences of complexes:

$$0 \rightarrow C_{\text{AO}}^{\bullet-1}(R, M) \rightarrow C_{\text{AvA}}^\bullet(R, M) \rightarrow C^\bullet(R, M) \rightarrow 0.$$

Then it is trivial to obtain the following result:

**Theorem 3.3.** *We have the following long exact sequence of cohomology groups,*

$$\dots \rightarrow H_{\text{AvO}}^{n-1}(R, M) \rightarrow H_{\text{AvA}}^n(R, M) \rightarrow \text{HH}^n(R, M) \rightarrow H_{\text{AvO}}^n(R, M) \rightarrow \dots$$



## 4. FORMAL DEFORMATION OF AVERAGING ALGEBRAS

Let  $(R, \bullet, A)$  be an averaging algebra. Denote by  $\mu_A$  the multiplication  $\bullet$  of  $R$ . Consider the 1-parameterized family

$$\mu_t = \sum_{i=0}^{\infty} \mu_i t^i, \mu_i \in C_{\text{Alg}}^2(R), A_t = \sum_{i=0}^{\infty} A_i t^i, A_i \in C_{\text{AvO}}^1(R).$$

**Definition 4.1.** A **1-parameter formal deformation** of an averaging algebra  $(R, \mu, A)$  is a pair  $(\mu_t, A_t)$  which endows the  $\mathbf{k}[[t]]$ -module  $(R[[t]], \mu_t, A_t)$  with the averaging algebra structure over  $\mathbf{k}[[t]]$  such  $(\mu_0, A_0) = (\mu, A)$ .

Power series  $\mu_t$  and  $A_t$  determines a 1-parameter formal deformation of the averaging algebra  $(R, \mu, A)$  if and only if for all  $x, y, z \in R$ , the following equations hold:

$$\begin{aligned} \mu_t(\mu_t(x \otimes y) \otimes z) &= \mu_t(x \otimes \mu_t(y \otimes z)), \\ \mu_t(A_t(x) \otimes A_t(y)) &= A_t(\mu_t(A_t(x) \otimes y)), \\ \mu_t(A_t(x) \otimes A_t(y)) &= A_t(\mu_t(x \otimes A_t(y))). \end{aligned}$$

Expand these equations and compare the coefficients of  $t^n$ , we get the conditions that  $\{\mu_i\}_{i \in \mathbb{N}}$  and  $\{A_i\}_{i \in \mathbb{N}}$  should satisfy:

$$(4) \quad \sum_{i=0}^n \mu_i \circ (\mu_{n-i} \otimes \text{id}) = \sum_{i=0}^n \mu_i \circ (\text{id} \otimes \mu_{n-i}),$$

$$(5) \quad \sum_{i+j=0}^n \mu_{n-i-j} \circ (A_i \otimes A_j) = \sum_{i+j=0}^n A_{n-i-j} \circ \mu_j \circ (A_i \otimes \text{id}),$$

$$(6) \quad \sum_{i+j=0}^n \mu_{n-i-j} \circ (A_i \otimes A_j) = \sum_{i+j=0}^n A_{n-i-j} \circ \mu_j \circ (\text{id} \otimes A_i).$$

**Proposition 4.2.** Let  $(R[[t]], \mu_t, A_t)$  be a 1-parameter formal deformation of an averaging algebra  $(R, \mu, A)$ . Then  $(\mu_1, A_1)$  is a 2-cocycle in the cochain complex  $C_{\text{AvA}}^\bullet(R)$ .

*Proof.* Compute the equations (4) (5) (6) for  $n = 1$ , we have :

$$\begin{aligned} \mu_1 \circ (\mu \otimes \text{id}) + \mu \circ (\mu_1 \otimes \text{id}) &= \mu_1 \circ (\text{id} \otimes \mu) + \mu \circ (\text{id} \otimes \mu_1), \\ \mu_1 \circ (A \otimes A) + \mu \circ (A \otimes A_1) + \mu \circ (A_1 \otimes A) &= A_1 \circ \mu \circ (A \otimes \text{id}) + A \circ \mu \circ (A_1 \otimes \text{id}) + A \circ \mu_1 \circ (A \otimes \text{id}), \\ \mu_1 \circ (A \otimes A) + \mu \circ (A \otimes A_1) + \mu \circ (A_1 \otimes A) &= A_1 \circ \mu \circ (\text{id} \otimes A) + A \circ \mu \circ (\text{id} \otimes A_1) + A \circ \mu_1 \circ (\text{id} \otimes A). \end{aligned}$$

They are equivalent to :

$$\begin{aligned} \delta(\mu_1) &= 0, \\ \partial_r(A_1) + \Phi_r^2(\mu_1) &= 0, \\ \partial_l(A_1) + \Phi_l^2(\mu_1) &= 0. \end{aligned}$$

That is,  $(\mu_1, A_1)$  is a 2-cocycle in  $C_{\text{AvA}}^\bullet(R)$ . □

If  $\mu_t = \mu_A$  in the above 1-parameter formal deformation of the averaging algebra  $(R, \mu, A)$ , we get a 1-parameter formal deformation of the averaging operator  $A$ . Consequently, we have:

**Corollary 4.3.** Let  $A_t$  be a 1-parameter formal deformation of the averaging operator  $A$ . Then  $A_1$  is a 1-cocycle in the cochain complex  $C_{\text{AvO}}^\bullet(R)$ .

**Definition 4.4.** The 2-cocycle  $(\mu_1, A_1)$  is called the infinitesimal of the 1-parameter formal deformation  $(R[[t]], \mu_t, A_t)$  of averaging  $(R, \mu, A)$ .

**Definition 4.5.** Let  $(R[[t]], \mu_t, A_t)$  and  $(R[[t]], \mu'_t, A'_t)$  be two 1-parameter formal deformations of averaging algebra  $(R, \mu, A)$ . A formal isomorphism from  $(R[[t]], \mu'_t, A'_t)$  to  $(R[[t]], \mu_t, A_t)$  is a power series  $\phi_t = \sum_{i \geq 0} \phi_i t^i : R[[t]] \rightarrow R[[t]]$ , where  $\phi_i : R \rightarrow R, i \in \mathbb{N}$  are linear maps with  $\phi_0 = \text{id}$ , such that

$$(7) \quad \phi_t \circ \mu'_t = \mu_t \circ (\phi_t \otimes \phi_t),$$

$$(8) \quad \phi_t \circ A'_t = A_t \circ \phi_t.$$

Two 1-parameter formal deformations  $(R[[t]], \mu_t, A_t)$  and  $(R[[t]], \mu'_t, A'_t)$  are said to be equivalent if there exists a formal isomorphism  $\phi_t : (R[[t]], \mu'_t, A'_t) \rightarrow (R[[t]], \mu_t, A_t)$ .

**Theorem 4.6.** *The infinitesimals of two equivalent 1-parameter formal deformations of  $(R, \mu, A)$  are in the same cohomology class in  $H^2_{\text{AvA}}(R)$ . Conversely, if the infinitesimals of two 1-parameter formal deformations of  $(R, \mu, A)$  fall into the same cohomology class, they are equivalent.*

*Proof.* Let  $\phi_t : (R[[t]], \mu'_t, A'_t) \rightarrow (R[[t]], \mu_t, A_t)$  be a formal isomorphism. For all  $x, y \in R$ , we have

$$(9) \quad \phi_t \circ \mu'_t(x \otimes y) = \mu_t \circ (\phi_t \otimes \phi_t)(x \otimes y),$$

$$(10) \quad \phi_t \circ A'_t(x) = A_t \circ \phi_t(x).$$

Expanding the identities comparing the coefficients of  $t$ , we get:

$$(11) \quad \mu'_1(x \otimes y) = \mu_1(x \otimes y) + x\phi_1(y) - \phi_1(xy) + \phi_1(x)y,$$

$$(12) \quad A'_1(x) = A_1(x) + A(\phi_1(x)) - \phi_1(A(x)).$$

So we have,

$$(\mu'_1, A'_1) - (\mu_1, A_1) = d(\phi_1) \in C^{\bullet}_{\text{AvA}}(R).$$

That is,  $[(\mu'_1, A'_1)] = [(\mu_1, A_1)] \in H^2_{\text{AvA}}(R)$ .

Conversely, given two formal deformations  $(R[[t]], \mu'_t, A'_t)$  and  $(R[[t]], \mu_t, A_t)$  of an averaging algebra  $(R, \mu, A)$ , suppose that  $(\mu_1, A_1)$  and  $(\mu'_1, A'_1)$  are in the same cohomology class of  $H^2_{\text{AvA}}(R)$ .  $\square$

**Definition 4.7.** A 1-parameter formal deformation  $(R[[t]], \mu_t, A_t)$  of  $(R, \mu, A)$  is said to be trivial if it is equivalent to be deformation  $(R[[t]], \mu, A)$ , that is, there exists  $\phi_t = \sum_{i=0}^{\infty} \phi_i t^i : R[[t]] \rightarrow R[[t]]$ , where  $\phi : A \rightarrow A$  are linear maps with  $\phi = \text{id}$ , such that

$$(13) \quad \phi_t \circ \mu_t = \mu_A \circ (\phi_t \otimes \phi_t),$$

$$(14) \quad \phi_t \circ A_t = A \circ \phi_t.$$

**Definition 4.8.** An averaging algebra  $(R, \mu, A)$  is said to be rigid if every 1-parameter formal deformation is trivial.

**Proposition 4.9.** *Let  $(R, \mu, A)$  be an averaging algebra. If  $H^2_{\text{AvA}}(R) = 0$ ,  $(R, \mu, A)$  is rigid.*

*Proof.* Let  $(R[[t]], \mu_t, A_t)$  be a 1-parameter formal deformation of  $(R, \mu, A)$ . By Proposition ,  $(\mu_1, A_1)$  is a 2-cocycle. By  $H^2_{\text{AvA}}(R) = 0$ , there exists a 1-cochain  $(\phi'_1, x) \in C^1_{\text{Alg}}(R, R) \oplus \text{Hom}(k, M)$  such that  $(\mu_1, A_1) = -d_{\text{AvA}}(\phi'_1, x)$ , that is,  $\mu_1 = -\delta^1(\phi'_1)$  and  $A_1 = -\partial(x) - \Phi^1(\phi'_1)$ . Since  $\Phi^0 = \text{Id}$ , let  $\phi_1 = \phi'_1 + \delta^0(x)$ . Then  $\mu_1 = -\delta^1(\phi_1)$  and  $A_1 = -\Phi^1(\phi_1)$ .

Setting  $\phi_t = Id_R + \phi_1 t$ , we have a deformation  $(R[[t]], \bar{\mu}_t, \bar{A}_t)$ , where

$$\bar{\mu}_t = \phi_t^{-1} \circ \mu_t \circ (\phi_t \times \phi_t)$$

and

$$\bar{A}_t = \phi_t^{-1} \circ A_t \circ \phi_t.$$

It is not difficult to see that

$$\begin{aligned} \bar{\mu}_t &= \mu + \bar{\mu}_2 t^2 + \dots, \\ \bar{A}_t &= A + \bar{A}_2 t^2 + \dots. \end{aligned}$$

Then by repeating the argument, we can show that  $(R[[t]], \mu_t, A_t)$  is equivalent to the trivial extension  $(R[[t]], \mu, A)$ . Thus,  $(R, \mu, A)$  is rigid.  $\square$

## 5. ABELIAN EXTENSIONS OF AVERAGING ALGEBRAS

In this section, we study abelian extensions of averaging algebras and show that they are classified by the second cohomology, as one would expect of a good cohomology theory.

**Definition 5.1.** An **abelian extension** of averaging algebras is a short exact sequence of homomorphisms of averaging algebras

$$\begin{array}{ccccccc} 0 & \longrightarrow & M & \xrightarrow{i} & \hat{R} & \xrightarrow{p} & R \longrightarrow 0 \\ & & \downarrow A_M & & \downarrow \hat{A} & & \downarrow A \\ 0 & \longrightarrow & M & \xrightarrow{i} & \hat{R} & \xrightarrow{p} & R \longrightarrow 0 \end{array}$$

such that  $uv = 0$  for all  $u, v \in M$ .

We will call  $(\hat{R}, \hat{A})$  an abelian extension of  $(R, A)$  by  $(M, A_M)$ .

**Definition 5.2.** Let  $(\hat{R}_1, \hat{A}_1)$  and  $(\hat{R}_2, \hat{A}_2)$  be two abelian extensions of  $(R, A)$  by  $(M, A_M)$ . They are said to be **isomorphic** if there exists an isomorphism of averaging algebras  $\zeta : (\hat{R}_1, \hat{A}_1) \rightarrow (\hat{R}_2, \hat{A}_2)$  such that the following commutative diagram holds:

$$\begin{array}{ccccccc} 0 & \longrightarrow & (M, A_M) & \xrightarrow{i} & (\hat{R}_1, \hat{A}_1) & \xrightarrow{p} & (R, A) \longrightarrow 0 \\ & & \parallel & & \downarrow \zeta & & \parallel \\ 0 & \longrightarrow & (M, A_M) & \xrightarrow{i} & (\hat{R}_2, \hat{A}_2) & \xrightarrow{p} & (R, A) \longrightarrow 0. \end{array}$$

A **section** of an abelian extension  $(\hat{R}, \hat{A})$  of  $(R, A)$  by  $(M, A_M)$  is a linear map  $s : R \rightarrow \hat{R}$  such that  $p \circ s = Id_R$ .

Now for an abelian extension  $(\hat{R}, \hat{A})$  of  $(R, A)$  by  $(M, A_M)$  with a section  $s : R \rightarrow \hat{R}$ , we define linear maps  $\rho_l : R \rightarrow \text{End}_{\mathbf{k}}(M)$ ,  $r \mapsto (m \mapsto rm)$  and  $\rho_r : R \rightarrow \text{End}_{\mathbf{k}}(M)$ ,  $r \mapsto (m \mapsto mr)$  respectively by

$$rm := s(r)m, \quad mr := ms(r), \quad \forall r \in R, m \in M.$$

**Proposition 5.3.** With the above notations,  $(M, \rho_l, \rho_r, A_M)$  is a bimodule over the averaging algebra  $(R, A)$ .

*Proof.* For any  $x, y \in R$ ,  $v \in M$ , since  $s(xy) - s(x)s(y) \in M$  implies  $s(xy)m = s(x)s(y)m$ , we have

$$\rho_l(xy)(m) = s(xy)m = s(x)s(y)m = \rho_l(x) \circ \rho_l(y)(m).$$

Hence,  $\rho_l$  is an algebra homomorphism. Similarly,  $\rho_r$  is an algebra anti-homomorphism. Moreover,  $\hat{A}(s(r)) - s(A(r)) \in M$  means that  $\hat{A}(s(r))m = s(A(r))m$ . Thus we have

$$\begin{aligned} A(r)A_M(m) &= s(A(r))A_M(m) \\ &= \hat{A}(s(r))A_M(m) \\ &= \hat{A}(s(r)A_M(m)) = \hat{A}(\hat{A}(sr)m) \\ &= A_M(rA(m)) = A_M(A(r)m). \end{aligned}$$

It is similar to see  $A_M(m)A(r) = A_M(A_M(m)r) = A_M(mA(r))$ . Hence,  $(M, \rho_l, \rho_r, A_M)$  is a bimodule over  $(R, A)$ .  $\square$

We further define linear maps  $\psi : R \otimes R \rightarrow M$  and  $\chi : R \rightarrow M$  respectively by

$$\begin{aligned} \psi(x \otimes y) &= s(x)s(y) - s(xy), \quad \forall x, y \in R, \\ \chi(x) &= \hat{A}(s(x)) - s(A(x)), \quad \forall x \in R. \end{aligned}$$

We transfer the averaging algebra structure on  $\hat{R}$  to  $R \oplus M$  by endowing  $R \oplus M$  with a multiplication  $\cdot_\psi$  and an averaging operator  $A_\chi$  defined by

$$(15) \quad (x, m) \cdot_\psi (y, n) = (xy, xn + my + \psi(x, y)), \quad \forall x, y \in R, m, n \in M,$$

$$(16) \quad A_\chi(x, m) = (A(x), \chi(x) + A_M(m)), \quad \forall x \in R, m \in M.$$

**Proposition 5.4.** *The triple  $(R \oplus M, \cdot_\psi, A_\chi)$  is a averaging algebra if and only if  $(\psi, \chi)$  is a 2-cocycle of the averaging algebra  $(R, A)$  with the coefficient in  $(M, A_M)$ .*

*Proof.* If  $(A \oplus M, \cdot_\psi, A_\chi)$  is a averaging algebra, then the associativity of  $\cdot_\psi$  implies

$$(17) \quad x\psi(y \otimes z) - \psi(xy \otimes z) + \psi(x \otimes yz) - \psi(x \otimes y)z = 0,$$

which means  $\delta(\phi) = 0$  in  $C^\bullet(A, M)$ . Since  $A_\chi$  is an averaging operator, for any  $x, y \in R, m, n \in M$ , we have

$$A_\chi((x, m)) \cdot_\psi A_\chi((y, n)) = A_\chi(A_\chi(x, m) \cdot_\psi (y, n)) = A_\chi((x, m) \cdot_\psi A_\chi(y, n))$$

Then  $\chi, \psi$  satisfy the following equations:

$$[\psi(A(x) \otimes A(y)) - A_M(\psi(A(x) \otimes y))] + [A(x)\chi(y) - A_M(\chi(x)y) - \chi(A(x)y) + \chi(x)A(y)] = 0$$

$$[\psi(A(x) \otimes A(y)) - A_M(\psi(x \otimes A(y)))] + [A(x)\chi(y) - A_M(x\chi(y)) - \chi(xA(y)) + \chi(x)A(y)] = 0$$

That is,

$$\partial_r^1(\chi) + \Phi_r^2(\psi) = 0, \partial_l^1(\chi) + \Phi_l^2(\psi) = 0.$$

Hence,  $(\psi, \chi)$  is a 2-cocycle.

Conversely, if  $(\psi, \chi)$  is a 2-cocycle, one can easily check that  $(R \oplus M, \cdot_\psi, A_\chi)$  is an averaging algebra.  $\square$

Now we are ready to classify abelian extensions of an averaging algebra.

**Theorem 5.5.** *Let  $M$  be a vector space and  $A_M \in \text{End}_{\mathbf{k}}(M)$ . Then abelian extensions of an averaging algebra  $(R, A)$  by  $(M, A_M)$  are classified by the second cohomology group  $H_{\text{AvA}}^2(R, M)$  of  $(R, A)$  with coefficients in the bimodule  $(M, A_V)$ .*

*Proof.* Let  $(\hat{R}, \hat{A})$  be an abelian extension of  $(R, A)$  by  $(M, A_M)$ . We choose a section  $s : R \rightarrow \hat{R}$  to obtain a 2-cocycle  $(\psi, \chi)$  by Proposition 5.4. We first show that the cohomological class of  $(\psi, \chi)$  does not depend on the choice of sections. Indeed, let  $s_1$  and  $s_2$  be two distinct sections providing 2-cocycles  $(\psi_1, \chi_1)$  and  $(\psi_2, \chi_2)$  respectively. We define  $\phi : R \rightarrow M$  by  $\gamma(r) = s_1(r) - s_2(r)$ . Then

$$\begin{aligned} \psi_1(x, y) &= s_1(x)s_1(y) - s_1(xy) \\ &= (s_2(x) + \gamma(x))(s_2(y) + \gamma(y)) - (s_2(xy) + \gamma(xy)) \\ &= (s_2(x)s_2(y) - s_2(xy)) + s_2(x)\gamma(y) + \gamma(x)s_2(y) - \gamma(xy) \\ &= (s_2(x)s_2(y) - s_2(xy)) + x\gamma(y) + \gamma(x)y - \gamma(xy) \\ &= \psi_2(x, y) + \delta(\gamma)(x, y) \end{aligned}$$

and

$$\begin{aligned} \chi_1(x) &= \hat{A}(s_1(x)) - s_1(A(x)) \\ &= \hat{A}(s_2(x) + \gamma(x)) - (s_2(A(x)) + \gamma(A(x))) \\ &= (\hat{A}(s_2(x)) - s_2(A(x))) + \hat{A}(\gamma(x)) - \gamma(A(x)) \\ &= \chi_2(x) + d_V(\gamma(x)) - \gamma(d_A(x)) \\ &= \chi_2(x) - \Phi^1(\gamma)(x). \end{aligned}$$

That is,  $(\psi_1, \chi_1) = (\psi_2, \chi_2) + d(\gamma)$ . Thus  $(\psi_1, \chi_1)$  and  $(\psi_2, \chi_2)$  are in the same cohomological class in  $H_{\text{AvA}}^2(R, M)$ .

Next we prove that isomorphic abelian extensions give rise to the same element in  $H_{\text{AvA}}^2(R, M)$ . Assume that  $(\hat{R}_1, \hat{A}_1)$  and  $(\hat{R}_2, \hat{A}_2)$  are two isomorphic abelian extensions of  $(R, A)$  by  $(M, A_M)$  with the associated homomorphism  $\zeta : (\hat{R}_1, \hat{A}_1) \rightarrow (\hat{R}_2, \hat{A}_2)$ . Let  $s_1$  be a section of  $(\hat{R}_1, \hat{A}_1)$ . As  $p_2 \circ \zeta = p_1$ , we have

$$p_2 \circ (\zeta \circ s_1) = p_1 \circ s_1 = \text{Id}_R.$$

Therefore,  $\zeta \circ s_1$  is a section of  $(\hat{R}_2, \hat{A}_2)$ . Denote  $s_2 := \zeta \circ s_1$ . Since  $\zeta$  is a homomorphism of differential algebras such that  $\zeta|_M = \text{Id}_M$ , we have

$$\begin{aligned} \psi_2(x \otimes y) &= s_2(x)s_2(y) - s_2(xy) = \zeta(s_1(x))\zeta(s_1(y)) - \zeta(s_1(xy)) \\ &= \zeta(s_1(x)s_1(y) - s_1(xy)) = \zeta(\psi_1(x, y)) \\ &= \psi_1(x, y) \end{aligned}$$

and

$$\begin{aligned} \chi_2(x) &= \hat{A}_2(s_2(x)) - s_2(A(x)) = \hat{A}_2(\zeta(s_1(x))) - \zeta(s_1(A(x))) \\ &= \zeta(\hat{A}_1(s_1(x)) - s_1(A(x))) = \zeta(\chi_1(x)) \\ &= \chi_1(x). \end{aligned}$$

Consequently, all isomorphic abelian extensions give rise to the same element in  $H_{\text{AvA}}^2(R, M)$ .

Conversely, given two 2-cocycles  $(\psi_1, \chi_1)$  and  $(\psi_2, \chi_2)$ , we can construct two abelian extensions  $R \oplus M, \cdot_{\psi_1}, A_{\chi_1}$  and  $(R \oplus M, \cdot_{\psi_2}, A_{\chi_2})$  via equalities (15) and (16). If they represent the same cohomological class in  $H_{\text{AvA}}^2(R, M)$ , then there exists a linear map  $\gamma : R \rightarrow M$  such that

$$(\psi_1, \chi_1) = (\psi_2, \chi_2) + (\delta(\gamma), \Phi^1(\gamma)).$$

Define  $\zeta : R \oplus M \rightarrow R \oplus M$  by

$$\zeta(r, m) := (r, \gamma(r) + m).$$

Then  $\zeta$  is an isomorphism of these two abelian extensions.  $\square$

## 6. $L_\infty$ -STRUCTURE ON THE COCHAIN COMPLEX

Let's recall the definition of  $L_\infty$ -algebras.

For graded indeterminates  $t_1, \dots, t_n$  and  $\sigma \in S_n$ , the Koszul sign  $\epsilon(\sigma, t_1, \dots, t_n)$  is defined by

$$t_1 \cdot t_2 \cdots t_n = \epsilon(\sigma, t_1, \dots, t_n) t_{\sigma(1)} \cdot t_{\sigma(2)} \cdots t_{\sigma(n)},$$

where “ $\cdot$ ” is the multiplication in the free graded commutative algebra  $k\langle t_1, \dots, t_n \rangle / (t_i t_j - (-1)^{|t_i||t_j|} t_j t_i)$  generated by  $t_1, \dots, t_n$ . Define also  $\chi(\sigma, t_1, \dots, t_n) = \text{sgn}(\sigma) \epsilon(\sigma, t_1, \dots, t_n)$ .

**Definition 6.1.** ([24, 15, 14]) Let  $L = \bigoplus_{i \in \mathbb{Z}} L_i$  be a graded space over  $\mathbf{k}$ . Assume that  $L$  is endowed with a family of linear operators  $\{l_n : L^{\otimes n} \rightarrow L\}_{n \geq 1}$  with  $|l_n| = n - 2$  satisfying the following conditions:

- (i)  $l_n(x_{\sigma(1)} \otimes \dots \otimes x_{\sigma(n)}) = \chi(\sigma, x_1, \dots, x_n) l_n(x_1 \otimes \dots \otimes x_n)$ ,  $\forall \sigma \in S_n, x_1, \dots, x_n \in L$ ,
- (ii)  $\sum_{i=1}^n \sum_{\sigma \in S(i, n-i)} \chi(\sigma, x_1, \dots, x_n) (-1)^{i(n-i)} l_{n-i+1}(l_i(x_{\sigma(1)} \otimes \dots \otimes x_{\sigma(i)}) \otimes x_{\sigma(i+1)} \otimes \dots \otimes x_{\sigma(n)}) = 0$ ,  
where  $S(i, n-i)$  is the set of all  $(i, n-i)$ -shuffles, i.e.,  $S(i, n-i) = \{\sigma \in S_n, |\sigma(1) < \sigma(2) < \dots < \sigma(i), \sigma(i+1) < \sigma(i+2) < \dots < \sigma(n)\}$ ,

Then  $(L, \{l_n\}_{n \geq 1})$  is called a  $L_\infty$ -algebra.

Let  $sL$  be the suspension of  $L$ , i.e.,  $(sL)_n = L_{n-1}$ . Let  $S^\bullet(sL)$  be the cofree cocommutative coalgebra generated by  $sL$ . Then  $\{l_n\}_{n \geq 1}$  will determine a family of operators  $\{d_n\}_{n \geq 1}$ , where  $d_n : (sL)^{\otimes n} \rightarrow sL$  is defined as  $d_n = (-1)^{\frac{n(n-1)}{2}} s \circ l_n \circ (s^{-1})^{\otimes n}$ . Then  $\{d_n\}_{n \geq 1}$  satisfies the equation

$$(18) \quad \sum_{\sigma \in S(i, n-i)} \epsilon(\sigma, sx_1, \dots, sx_n) d_{n-i+1}(d_i(sx_{\sigma(1)}, \dots, sx_{\sigma(i)}), sx_{\sigma(i+1)}, \dots, sx_{\sigma(n)}) = 0.$$

Denote  $\sum_{i=1}^{\infty} d_n : S^\bullet(sL) \rightarrow sL$  by  $d'$ . Then  $d'$  can induce a coderivation  $d$  on  $S^\bullet(sL)$ . Equation (18) ensures that the coderivation  $d$  is a differential, i.e.,  $d^2 = 0$ . This can be considered as an equivalent definition of  $L_\infty$ -algebra.

**Definition 6.2.** Let  $(L, \{l_n\}_{n \geq 1})$  be a  $L_\infty$ -algebra and  $\alpha \in L_{-1}$ . We call  $\alpha$  a Maurer-Cartan element if it satisfies the following equation:

$$\sum_{n=1}^{\infty} \frac{1}{n!} (-1)^{\frac{n(n-1)}{2}} l_n(\alpha^{\otimes n}) = 0.$$

**Proposition 6.3.** Given a Maurer-Cartan element  $\alpha$  in  $L_\infty$ -algebra  $L$ , we can define a new  $L_\infty$ -structure  $\{l_n^\alpha\}_{n \geq 1}$  on  $L$ , where  $l_n^\alpha : L^{\otimes n} \rightarrow L$  is defined as:

$$l_n^\alpha(x_1 \otimes \dots \otimes x_n) = \sum_{i=0}^{\infty} \frac{1}{i!} (-1)^{in + \frac{i(i+1)}{2} + \sum_{k=1}^{n-1} \sum_{j=1}^k |x_j|} l_{n+i}(x_1 \otimes \dots \otimes x_i \otimes \alpha^{\otimes i} \otimes x_1 \otimes \dots \otimes x_n).$$

Let  $\alpha$  be a Maurer Cartan element in  $L_\infty$  algebra  $L$ . Use the equivalent definition of  $L_\infty$ -algebra, we can see  $s\alpha \in sL$  satisfies the following equation on  $S^\bullet(sL)$ :

$$\sum_{n=1}^{\infty} \frac{1}{n!} d_n((s\alpha)^{\otimes n}) = 0.$$

Define

$$d_n^\alpha(sx_1 \otimes \dots \otimes sx_n) = \sum_{i=0}^{\infty} \frac{1}{i!} d_{n+1}((s\alpha)^{\otimes i} \otimes sx_1 \otimes \dots \otimes sx_n).$$

Then the family  $\{d_n^\alpha\}_{n \geq 1}$  can also induce a differential on  $S^\bullet(sL)$ .

Let  $W$  be a graded space and  $T^c(W)$  be the cofree coalgebra generated by  $W$ . For  $f \in \text{Hom}(W^{\otimes n}, W)$ ,  $g_i \in \text{Hom}(W^{\otimes i}, W)$ ,  $1 \geq i \geq m$ ,  $f \circ (g_1, \dots, g_m) \in \text{Hom}(W^{\otimes l_1 + \dots + l_m + n - m}, W)$  is defined as follows:

$$(f \circ (g_1, \dots, g_m))(w_1 \otimes \dots \otimes w_{l_1 + \dots + l_m + n - m}) = \sum_{0 \leq i_1 \leq i_2 \leq \dots \leq i_m} (-1)^\eta f(w_1 \otimes \dots \otimes w_{i_1} \otimes g_1(w_{i_1+1} \otimes \dots) \otimes \dots \otimes g_k(w_{i_k+1} \otimes \dots) \otimes \dots \otimes g_m(w_{i_m+1} \otimes \dots) \otimes \dots)$$

$$\text{where } \eta = \sum_{k=1}^m |g_k| \left( \sum_{j=1}^{i_k} |w_j| \right).$$

For any  $f, g \in \text{Hom}(T^c(W), W)$ , their Gerstenhaber bracket  $[f, g]_G \in \text{Hom}(T^c(W), W)$  is defined as :

$$[f, g]_G = f \circ g - (-1)^{|f||g|} g \circ f.$$

**Lemma 6.4.** [10] *Let  $W$  be a graded space. Then  $(\text{Hom}(T^c(W), W), [-, -]_G)$  forms a graded Lie-algebra.*

Now, given a graded space  $V = \bigoplus_{i \in \mathbb{Z}} V_i$ , we will define a  $L_\infty$ -algebra  $C_{AvA}(V)$ . We will see that when  $V$  is concentrated in degree 0, an averaging algebra structure on  $V$  is equivalent to a Maurer-Cartan element in this  $L_\infty$ -algebra.

Firstly, the underlying graded space of  $C_{AvA}(V)$  is

$$C_A(V) \oplus \text{Hom}(k, V) \oplus \text{Hom}(sV, V) \oplus C_{AvO}(V)_l^{\geq 2} \oplus C_{AvO}(V)_r^{\geq 2}.$$

where

$$\begin{aligned} C_A(V) &= \text{Hom}(T^c(sV), sV), \\ C_{AvO}(V)_l^{\geq 2} &= \text{Hom}\left(\bigoplus_{n=2}^{\infty} (sV)^{\otimes n}, V\right), \\ C_{AvO}(V)_r^{\geq 2} &= \text{Hom}\left(\bigoplus_{n=2}^{\infty} (sV)^{\otimes n}, V\right). \end{aligned}$$

And we define

$$\begin{aligned} C_{AvO}(V)_r &= \text{Hom}(k, V) \oplus \text{Hom}(sV, V) \oplus C_{AvO}(V)_r^{\geq 2} \\ C_{AvO}(V)_l &= \text{Hom}(k, V) \oplus \text{Hom}(sV, V) \oplus C_{AvO}(V)_l^{\geq 2} \end{aligned}$$

Now, let's give a  $L_\infty$ -algebra structure on  $C_{AvA}(V)$ , i.e., we need to give a family of operators  $\{l_n\}_{n \geq 1}$  to satisfy the conditions in Definition 6.1. Here, we identify  $\text{Hom}(T^c(sV), sV)$  with  $s\text{Hom}(T^c(sV), V)$ .

The family  $\{l_i\}_{i \geq 1}$  are defined by the following processes:

(I) For  $sh \in \text{Hom}(k, sV) \subset C_A(V)$ ,  $l_1(sh) = h$ . And  $l_1$  vanishes elsewhere.

(II) For  $sg, sh \in C_A(V)$ ,  $l_2(sg \otimes sh) = [sg, sh]_G$ .

(III) For homogeneous elements  $sh \in \text{Hom}((sV)^{\otimes n}, sV)$  in  $C_A(V)$  and  $g_1, \dots, g_n \in C_{AvO}(V)_r$ , define  $l'_{n+1}(sh \otimes g_1 \otimes \dots \otimes g_n) \in C_{AvO}(V)_r$  in the following ways:

(i) If  $g_1, \dots, g_n$  are all contained in  $\text{Hom}(sV, V) \oplus C_{AvO}(V)_r^{\geq 2}$ , then

$$l'_{n+1}(sh \otimes g_1 \otimes \dots \otimes g_n) = \sum_{\sigma \in S_n} (-1)^\varepsilon \left( h \circ (sg_{\sigma(1)} \otimes \dots \otimes sg_{\sigma(n)}) - (-1)^{(|g_{\sigma(1)}|+1)(|h|+1)} g_{\sigma(1)} \bar{\circ} (sh \circ (\text{id}_{sV} \otimes sg_{\sigma(2)} \otimes \dots \otimes sg_{\sigma(n)})) \right)$$

(ii) If there exists some  $g_i$  coming from  $\text{Hom}(k, V)$ , then

$$l'_{n+1}(sh \otimes g_1 \otimes \dots \otimes g_n) = \sum_{\sigma \in S_n} (-1)^\varepsilon \left( h \circ (sg_{\sigma(1)} \otimes \dots \otimes sg_{\sigma(n)}) - (-1)^{(|g_{\sigma(p)}|+1)(|h|+1+\sum_{k=1}^{p-1}(|g_{\sigma(k)}|+1))} g_{\sigma(p)} \bar{\circ} (sh \circ (g_{\sigma(1)} \otimes sg_{\sigma(2)} \otimes \dots \otimes g_{\sigma(p-1)} \otimes \text{id}_{sV} \otimes sg_{\sigma(p+1)} \otimes \dots \otimes sg_{\sigma(n)})) \right)$$

where  $(-1)^\varepsilon = \chi(\sigma; g_1, \dots, g_n) \cdot (-1)^{n(|h|+1)+\sum_{k=1}^{n-1} \sum_{j=1}^k |g_{\sigma(j)}|}$ , and  $p$  is the integer such that  $g_{\sigma(1)}, \dots, g_{\sigma(p-1)} \in \text{Hom}(k, V)$  and  $g_{\sigma(p)} \in \text{Hom}(sV, V) \oplus C_{AvO}(V)_r^{\geq 2}$ .

(IV) Let  $sh \in \text{Hom}((sV)^{\otimes n}, sV)$  in  $C_A(V)$ ,  $g_1, \dots, g_n \in C_{AvO}(V)_l$  be homogeneous elements. Define  $l^l_{n+1}(sh \otimes g_1 \otimes \dots \otimes g_n) \in C_{AvO}(V)_l$  in the following ways:

(i) If  $g_1, \dots, g_n$  all belong to  $\text{Hom}(sV, V) \oplus C_{AvO}(V)_l^{\geq 2}$ , then define

$$l^l_{n+1}(sh \otimes g_1 \otimes \dots \otimes g_n) = \sum_{\sigma \in S_n} (-1)^\varepsilon \left( h \circ (sg_{\sigma(1)} \otimes \dots \otimes sg_{\sigma(n)}) - (-1)^{(|g_{\sigma(n)}|+1)(|h|+1+\sum_{k=1}^{n-1}(|g_{\sigma(k)}|+1))} g_{\sigma(n)} \bar{\circ} (sh \circ (sg_{\sigma(1)} \otimes \dots \otimes sg_{\sigma(n-1)} \otimes \text{id}_{sV})) \right),$$

(ii) If there exists some  $g_i \in \text{Hom}(k, V)$ , then define:

$$l^l_{n+1}(sh \otimes g_1 \otimes \dots \otimes g_n) = \sum_{\sigma \in S_n} (-1)^\varepsilon \left( h \circ (sg_{\sigma(1)} \otimes \dots \otimes sg_{\sigma(n)}) - (-1)^{(|g_{\sigma(q)}|+1)(|h|+1+\sum_{k=1}^{q-1}(|g_{\sigma(k)}|+1))} g_{\sigma(q)} \bar{\circ} (sh \circ (sg_{\sigma(1)} \otimes \dots \otimes sg_{\sigma(q-1)} \otimes \text{id}_{sV} \otimes sg_{\sigma(q+1)} \otimes \dots \otimes sg_{\sigma(n)})) \right).$$

Where  $q$  is the integer such that  $g_{\sigma(q+1)}, \dots, g_{\sigma(n)} \in \text{Hom}(k, V)$  and  $g_{\sigma(q)} \in \text{Hom}(sV, V) \oplus C_{AvO}(V)_l^{\geq 2}$ .

(V) (i) If  $g_1, \dots, g_n \in \text{Hom}(k, V) \oplus \text{Hom}(sV, V) = C_{AvO}(V)_l \cap C_{AvO}(V)_r$  with at most one  $g_i \in \text{Hom}(sV, V)$ , then the definitions of  $l'_{n+1}, l^l_{n+1}$  coincide. Then we define

$$l_{n+1}(sh, g_1, \dots, g_n) := l'_{n+1}(sh, g_1, \dots, g_n) = l^l_{n+1}(sh, g_1, \dots, g_n).$$

(ii) If  $g_1, \dots, g_n \in \text{Hom}(k, V) \oplus \text{Hom}(sV, V) = C_{AvO}(V)_l \cap C_{AvO}(V)_r$  with at least two  $g_i \in \text{Hom}(sV, V)$ , then  $l^l_{n+1}(sh, g_1, \dots, g_n) \in C_{AvO}(V)_l^{\geq 2}$  and  $l'_{n+1}(sh, g_1, \dots, g_n) \in C_{AvO}(V)_r^{\geq 2}$ . Then we define

$$l_{n+1}(sh, g_1, \dots, g_n) = (l'_{n+1}(sh, g_1, \dots, g_n), l^l_{n+1}(sh, g_1, \dots, g_n)) \in C_{AvO}(V)_r^{\geq 2} \oplus C_{AvO}(V)_l^{\geq 2}.$$



(iii) For  $g_1, \dots, g_n \in C_{AvO}(V)_r$  with some  $g_i \in C_{AvO}(V)_r^{\geq 2}$ , define

$$l_{n+1}(sh, g_1, \dots, g_n) := l_{n+1}^r(sh, g_1, \dots, g_n).$$

For  $g_1, \dots, g_n \in C_{AvO}(V)_l$  with some  $g_i \in C_{AvO}(V)_l^{\geq 2}$ , define

$$l_{n+1}(sh, g_1, \dots, g_n) := l_{n+1}^r(sh, g_1, \dots, g_n).$$

(VI) At last we define

$$l_{n+1}(g_1 \otimes \dots \otimes g_i \otimes sh \otimes g_{i+1} \otimes \dots \otimes g_n) = (-1)^{(|h|+1)(\sum_{k=1}^i |g_k|)+i} l_{n+1}(sh \otimes g_1 \otimes \dots \otimes g_n).$$

And  $l_n$  vanishes elsewhere.

**Theorem 6.5.** *Let  $V$  be a graded space. Then  $C_{AvA}(V)$  endowed with operations  $\{l_n\}_{n \geq 1}$  defined above forms a  $L_\infty$ -algebra.*

Now, let's realize averaging algebra structures as Maurer-Cartan elements in this  $L_\infty$ -algebra.

**Theorem 6.6.** *Let  $R$  be a vector space. Then an averaging structure on  $R$  is equivalent to a Maurer-Cartan elements in  $C_{AvA}(R)$ .*

*Proof.* Consider  $R$  as a graded vector space concentrated in degree 0. Then the degree -1 part of  $C_{AvA}(R)$  is

$$C_{AvA}(R)_{-1} = \text{Hom}((sV)^{\otimes 2}, sV) \bigoplus \text{Hom}(sV, V).$$

Assume that  $\alpha = (m, \tau)$  be a Maurer-Cartan element in  $C_{AvA}(R)$ , that is,  $\alpha$  satisfy the equation:

$$(19) \quad \sum_{k=1}^{\infty} \frac{1}{k!} (-1)^{\frac{k(k-1)}{2}} l_k(\alpha^{\otimes k}) = 0.$$

By the definition of  $\{l_n\}_{n \geq 1}$ ,

$$\begin{aligned} & \frac{1}{2!} (-1) l_2(\alpha \otimes \alpha) = -\frac{1}{2} [m, m]_G, \\ & \frac{1}{3!} (-1) l_3(\alpha^{\otimes 3}) \\ &= -\frac{1}{3!} (l_3^r(m \otimes \tau \otimes \tau) + l_3^r(\tau \otimes m \otimes \tau) + l_3^r(\tau \otimes \tau \otimes m), l_3^l(m \otimes \tau \otimes \tau) + l_3^l(\tau \otimes m \otimes \tau) + l_3^l(\tau \otimes \tau \otimes m)) \\ &= -\frac{1}{3!} (3l_3^r(m \otimes \tau \otimes \tau), 3l_3^l(m \otimes \tau \otimes \tau)) \\ &= -\left(s^{-1}m \circ (s\tau \otimes s\tau) - \tau \circ (sm \circ (\text{id} \otimes s\tau)), s^{-1}m \circ (s\tau \otimes s\tau) - \tau \circ (sm \circ (s\tau \otimes \text{id}))\right) \end{aligned}$$

Then Equation (19) implies that

$$(20) \quad [m, m]_G = 0$$

$$(21) \quad s^{-1}m \circ ((s\tau)^{\otimes 2}) - \tau \circ (sm \circ (\text{id} \otimes s\tau)) = 0$$

$$(22) \quad s^{-1}m \circ ((s\tau)^{\otimes 2}) - \tau \circ (sm \circ (s\tau \otimes \text{id})) = 0.$$

Define  $\mu = s^{-1} \circ m \circ s^{\otimes 2} : A^{\otimes 2} \rightarrow A, A = \tau \circ s : A \rightarrow A$ . Then Equation (20) is equivalent to

$$\mu \circ (\text{id} \otimes \mu) = \mu \circ (\mu \otimes \text{id}).$$

That is,  $\mu$  is associative. Equations (21) (22) imply

$$\mu \circ (A \otimes A) = A \circ (\mu \circ (\text{id} \otimes A)), \mu \circ (A \otimes A) = A \circ (\mu \circ (A \otimes \text{id})),$$

which means that  $A$  is an averaging operator on associative algebra  $(R, \mu)$ . Conversely, let  $(R, \mu, A)$  be an averaging algebra. Define  $m : (sR)^{\otimes 2} \rightarrow sR$ ,  $\tau : sR \rightarrow R$  to be  $m(sa \otimes sb) = s\mu(a \otimes b)$ ,  $\tau(sa) = A(a)$ . Then  $(m, \tau)$  is a Maurer-Cartan element in  $C_{AvA}(R)$ .  $\square$

**Proposition 6.7.** *Let  $(R, \mu, A)$  be an averaging algebra and  $\alpha = (m, \tau)$  be the corresponding Maurer-Cartan element in  $C_{AvA}(R)$ . Then the underground complex of  $L_\infty$ -algebra  $(C_{AvA}(R), \{l^\alpha\})$  is exactly the cochain complex  $sC_{AvA}^\bullet(R)$  defined in Section 3.*

*Proof.* Identify  $C^\bullet(R)$  with  $C_A(R)$  and identify  $C_{AvO}^\bullet$  with  $\text{Hom}(k, R) \oplus \text{Hom}(sR, R) \oplus C_{AvO}(R)_r^{\geq 2} \oplus C_{AvO}(R)_l^{\geq 2}$ . Let  $sf \in \text{Hom}((sR)^{\otimes n}, sR)$ . Then

$$l_1^\alpha(sf) = \sum_{k=0}^{\infty} \frac{1}{k!} (-1)^{\frac{k(k+1)}{2}+k} l_{k+1}(\alpha^{\otimes k} \otimes sf) = l_2(m \otimes sf) + \frac{1}{n!} (-1)^{\frac{n(n-1)}{2}} l_{n+1}(\tau^{\otimes n} \otimes sf),$$

where  $l_2(m \otimes sf) = [m, sf]_G$ , it's just the differential  $\delta$  of Hochschild cochain complex. We have

$$\frac{1}{n!} (-1)^{\frac{n(n-1)}{2}} l_{n+1}(\tau^{\otimes n} \otimes sf) = \frac{1}{n!} (-1)^{\frac{n(n-1)}{2}+n} (l_{n+2}^\tau(sf \otimes \tau^{\otimes n}), l_{n+1}^\tau(sf \otimes \tau^{\otimes n})).$$

Then by definition,

$$\frac{1}{n!} (-1)^{\frac{n(n-1)}{2}+n} l_{n+2}^\tau(sf \otimes \tau^{\otimes n}) = (-1)^n (f \circ ((s\tau)^{\otimes n}) - \tau \circ (sf \circ (\text{id} \otimes (s\tau)^{\otimes n-1}))),$$

which is exactly the same as  $(-1)^n \Phi_r^n$  defined in Section 3.2. Similarly, we can see  $l_{n+1}^\tau(sf \otimes \tau^{\otimes n})$  is just  $(-1)^n \Phi_l^n$ . In particular, for  $sf \in \text{Hom}(sR, sR)$ ,  $l_2(sf \otimes \tau) = l_2^\tau(sf \otimes \tau) = l_2^\tau(sf \otimes \tau) = -(f \circ s\tau - \tau \circ (sf))$ , it is the same as  $-\Phi^1$ .

For  $g \in C_{AvO}(R)^r$ , we have

$$\begin{aligned} \sum_{k=0}^{\infty} \frac{1}{k!} (-1)^{\frac{k(k+1)}{2}+k} l_{k+1}(\alpha^{\otimes k} \otimes g) &= -\frac{1}{2} (l_3(m \otimes \tau \otimes g) + l_3(\tau \otimes m \otimes g)) \\ &= -l_3(m \otimes \tau \otimes g) \\ &= -((-s^{-1}m_2 \circ (s\tau \otimes sg) + \tau \circ (m \circ (\text{id} \otimes sg))) \\ &\quad - (s^{-1}m \circ (sg \otimes s\tau) - g \circ (m \circ (\text{id} \otimes s\tau)))) \\ &= s^{-1}m_2 \circ (s\tau \otimes sg) - \tau \circ (m \circ (\text{id} \otimes sg)) \\ &\quad - g \circ (m \circ (\text{id} \otimes \tau)) + s^{-1}m \circ (sg \otimes s\tau), \end{aligned}$$

and it is the same as  $\partial_r$  defined in Section 3.1. For  $g \in C_{AvO}(R)^l$ , it is all the same. Especially, when  $g \in \text{Hom}(k, R)$ , then by definition, we have

$$\begin{aligned} \sum_{k=0}^{\infty} \frac{1}{k!} (-1)^{\frac{k(k+1)}{2}+k} l_{k+1}(\alpha^{\otimes k} \otimes g) &= -\frac{1}{2} (l_3(m \otimes \tau \otimes g) + l_3(\tau \otimes m \otimes g)) \\ &= -l_3(m \otimes \tau \otimes g) \\ &= -((-s^{-1}m \circ (s\tau \otimes sg) + \tau \circ (m \circ (\text{id} \otimes sg))) \\ &\quad - (s^{-1}m \circ (sg \otimes s\tau) - \tau \circ (m \circ (sg \otimes \text{id})))) \end{aligned}$$

And it is the same as the operator  $\partial_0$  defined in Section 3.1.

□

**Proposition 6.8.** *Let  $V$  be a graded space. Then a Maurer-Cartan element  $\alpha$  in  $C_{AvA}(V)$  will induce a  $L_\infty$ -algebra structure on  $C_{AvO}(V)$ . In particular, for an averaging algebra  $(R, \mu, A)$ , the cochain complex  $C_{AvO}(R)$  is a differential graded Lie-algebra.*

*Proof.* Given a Maurer-Cartan element  $\alpha$ ,  $\alpha$  will induce  $L_\infty$ -algebra structure  $\{l_n^\alpha\}_{n \geq 1}$  on  $C_{AvA}(V)$ . And notice that all operations  $l_n^\alpha$  can be restricted to  $C_{AvO}(V)$ . Thus  $C_{AvO}(V)$  forms a  $L_\infty$ -subalgebra of  $(C_{AvA}(V), \{l_n^\alpha\}_{n \geq 1})$ .

Let  $(R, \mu, A)$  be an averaging algebra and  $(m, \tau)$  be the corresponding Maurer-Cartan element in  $C_{AvA}(R)$ . Identify  $C_{AvO}(R)$  with  $C_{AvO}(R)$ . Since  $m = -s \circ \mu \circ (s^{-1})^{\otimes 2} \in \text{Hom}((sR)^{\otimes 2}, sR)$  in  $C_A(R)$ , the restriction of  $l_n^\alpha$  on  $C_{AvO}(R)$  is zero for  $n \geq 3$ . Thus  $C_{AvO}(R)$  is just a differential graded Lie-algebra. □

## 7. HOMOTOPY AVERAGING ALGEBRAS

In this subsection, we'll define homotopy averaging algebras.

Recall that an  $A_\infty$ -algebra structure is equivalent to a Maurer-Cartan element in the graded Lie-algebra  $\overline{C_A}(V) := (\text{Hom}(\overline{T^c(sV)}, sV), [-, -]_G)$  where  $\overline{T^c(sV)} = \bigoplus_{n=1}^{\infty} (sV)^{\otimes n}$ . We can define homotopy averaging algebra structure in similar way. For a graded vector space  $V$ , consider the following subspace of  $C_{AvA}(V)$ :

$$\overline{C_{AvA}}(V) := \overline{C_A}(V) \bigoplus \text{Hom}(sV, V) \bigoplus C_{AvO}(V)_r^{\geq 2} \bigoplus C_{AvO}(V)_l^{\geq 2}.$$

Obviously,  $\overline{C_{AvA}}(V)$  is a  $L_\infty$ -subalgebra of  $\overline{C_{AvA}}(V)$  with the operations  $\{l_n\}_{n \geq 1}$  restricted on  $\overline{C_{AvA}}(V)$ .

Then we have the following definition:

**Definition 7.1.** Let  $V$  be a graded space. A homotopy averaging algebra( $Av_\infty$ ) structure on  $V$  is a Maurer-Cartan element in the  $L_\infty$ -algebra  $\overline{C_{AvA}}(V)$ .

Let's describe this structure explicitly.

A Maurer-Cartan element  $\alpha$  in  $\overline{C_{AvA}}(V)$  corresponds to a family of operators:

$$\{b_n\}_{n \geq 1} \cup \{c\} \cup \{c_n^r\}_{n \geq 2} \cup \{c_n^l\}_{n \geq 2}$$

where  $b_n : (sV)^{\otimes n} \rightarrow sV$ ,  $c : sV \rightarrow V$  belongs  $\overline{C_{AvA}}(V)$ ,  $\text{Hom}(sV, V)$  respectively, and  $c_n^r : (sV)^{\otimes n} \rightarrow V$ ,  $c_n^l : (sV)^{\otimes n} \rightarrow V$  belongs to  $C_{AvO}(V)(V)_r^{\geq 2}$ ,  $C_{AvO}(V)_l^{\geq 2}$  respectively. All these operators are of degree  $-1$ . Then  $\alpha$  satisfying the Maurer-Cartan equation implies that the family of operators satisfies the following equations:

$$\begin{aligned} \sum_{i+j=n} b_{n-j+1} \circ (\text{id}^{\otimes i} \otimes b_j \otimes \text{id}^{\otimes n-i-j}) &= 0, \\ \sum_{m=1}^n \sum_{l_1+\dots+l_m=n} \left( s^{-1} b_m \circ (sc_{l_1}^r \otimes \dots \otimes sc_{l_m}^r) - c_{l_1}^r \circ (b_m \circ (\text{id} \otimes sc_{l_2}^r \otimes \dots \otimes sc_{l_m}^r)) \right) &= 0, \\ \sum_{m=1}^n \sum_{l_1+\dots+l_m=n} \left( s^{-1} b_m \circ (sc_{l_1}^l \otimes \dots \otimes sc_{l_m}^l) - c_{l_m}^l \circ (b_m \circ (sc_{l_1}^l \otimes \dots \otimes sc_{l_{m-1}}^l \otimes \text{id})) \right) &= 0. \end{aligned}$$

for any  $n \geq 1$ , where  $c_1^r = c_1^l = c$ .

Then we define  $m_n = s^{-1} \circ b_n \circ s^{\otimes n} : V^{\otimes n} \rightarrow V$ ,  $A = c \circ s : V \rightarrow V$ ,  $A_n^r = c_n^r \circ s^{\otimes n} : V^{\otimes n} \rightarrow V$  and  $A_n^l = c_n^l \circ s^{\otimes n} : V^{\otimes n} \rightarrow V$  for all  $n \geq 1$ , where  $|m_n| = n - 2$ ,  $|A| = 0$ ,  $|A_n^r| = |A_n^l| = n - 1$ .

These operators  $\{m_n\}_{n \geq 1} \cup \{A\} \cup \{A_n^r\}_{n \geq 2} \cup \{A_n^l\}_{n \geq 2}$  satisfies the following identities:

(i)

$$\sum_{i+j=n} (-1)^{i+jk} m_{n-j+1} \circ (\text{id}^{\otimes i} \otimes m_j \otimes \text{id}^{\otimes n-i-j}) = 0,$$

(ii)

$$\begin{aligned} & \sum_{k=1}^n \sum_{l_1+\dots+l_k=n} (-1)^{\varepsilon} \{m_k \circ (A_{l_1}^r \otimes \dots \otimes A_{l_k}^r) \\ & - \sum_{p+q+1=l_1} (-1)^{q \cdot \sum_{j=2}^k (l_j-1) + kp+q} A_{l_1}^r \circ (\text{id}^{\otimes p} \otimes m_k \circ (\text{id} \otimes A_{l_2}^r \otimes \dots \otimes A_{l_k}^r) \otimes \text{id}^{\otimes q})\} = 0, \end{aligned}$$

(iii)

$$\begin{aligned} & \sum_{k=1}^n \sum_{l_1+\dots+l_k=n} (-1)^{\varepsilon} \{m_k \circ (A_{l_1}^l \otimes \dots \otimes A_{l_k}^l) \\ & - \sum_{p+q+1=l_k} (-1)^{p \cdot \sum_{j=1}^{k-1} (l_j-1) + (l_k-1) \cdot \sum_{j=1}^{k-1} l_j + kp+q} A_{l_k}^l \circ (\text{id}^{\otimes p} \otimes m_k \circ (A_{l_1}^l \otimes \dots \otimes A_{l_{k-1}}^l \otimes \text{id}) \otimes \text{id}^{\otimes q})\} = 0 \end{aligned}$$

where  $A_1^l = A_1^r = A$ , and  $\varepsilon = \sum_{j=1}^k \frac{l_j(l_j-1)}{2} + \frac{k(k-1)}{2} + \sum_{j=1}^k (l_j-1)(l_1+\dots+l_{j-1}) = \frac{n(n-1)}{2} + \frac{k(k-1)}{2} + \sum_{j=1}^k (k-j+1)l_j$ .

The equation (i) is just the definition of  $A_{\infty}$ -algebras.

Let's compute the equation (ii)(iii) for  $n = 1, 2$ :

$$(1) \ n = 1, -m_1 \circ A + A \circ m_1 = 0$$

$$\begin{aligned} (2) \ n = 2, m_2 \circ (A \otimes A) - A \circ (m_2 \circ (\text{id} \otimes A)) &= -m_1 \circ A_2^r + A_2^r \circ (m_1 \otimes \text{id}) + A_2^r \circ (\text{id} \otimes m_1), \\ m_2 \circ (A \otimes A) - A \circ (m_2 \circ (A \otimes \text{id})) &= -m_1 \circ A_2^l + A_2^l \circ (m_1 \otimes \text{id}) + A_2^l \circ (\text{id} \otimes m_1). \end{aligned}$$

The equation (1) above implies that  $A$  is compatible with the differential, so operator  $A$  can be induced on the homology of the complex  $(V, m_1)$ . The equation (2) says that  $A$  is an averaging operator up to homotopy with respect to  $m_2$  on  $V$  and the obstructions are  $A_2^r$  and  $A_2^l$ .

So we get another definition of homotopy averaging algebra.

**Definition 7.2.** Let  $V$  be a graded space. Assume that  $V$  is endowed with a family of operators  $\{m_n : V^{\otimes n} \rightarrow V\}_{n \geq 1} \cup \{A : V \rightarrow V\} \cup \{A_n^r : V^{\otimes n} \rightarrow V\}_{n \geq 2} \cup \{A_n^l : V^{\otimes n} \rightarrow V\}_{n \geq 2}$  with  $|m_n| = n - 2$ ,  $|A| = 0$ ,  $|A_n^r| = |A_n^l| = n - 1$ . If these operators satisfy the equations (i) (ii) (iii) above, we call  $V$  a homotopy averaging algebra.

#### APPENDIX: PROOF OF THEOREM 6.5

Firstly, let's display some facts we need to prove Theorem 6.5.

**Lemma 7.3.** Let  $n \geq 1$ ,  $1 \leq i \leq n - 1$ . Then for any  $\pi \in S_n$ , there exists a unique triple  $(\delta, \sigma, \tau)$  with  $\sigma \in S(i, n-i)$ ,  $\delta \in S_i$ ,  $\tau \in S_{n-i}$  such that  $\pi(l) = \sigma\delta(l)$  for  $1 \leq l \leq i$ , and  $\pi(i+m) = \sigma(i+\tau(m))$  for  $1 \leq m \leq n-i$ .

Let  $V$  be a graded space. The operation “ $\bar{\circ}$ ” on  $\text{Hom}(T^c(V), V)$  satisfying the Pre-Jacobi identity:

**Lemma 7.4.** *For any homogeneous elements  $f; g_1, \dots, g_m; h_1, \dots, h_n$  in  $\text{Hom}(T^c(V), V)$ , the following identity holds:*

$$(f \bar{\circ}(g_1, \dots, g_m)) \bar{\circ}(h_1, \dots, h_n) = \sum_{0 \leq i_1 \leq i_2 \leq \dots \leq i_m \leq n} (-1)^{\sum_{k=1}^m |g_k|(\sum_{j=1}^{i_k} |h_j|)} f \bar{\circ}(h_1, \dots, h_{i_1}, g_1 \bar{\circ}(h_{i_1+1}, \dots), \dots, h_{i_m}, g_m \bar{\circ}(h_{i_m+1}, \dots) \dots)$$

Now, let's prove the theorem 6.5.

**Theorem 6.5.** *Let  $V$  be a graded space. Then  $C_{\text{AvA}}(V)$  endowed with operations  $\{l_n\}_{n \geq 1}$  defined above forms a  $L_\infty$ -algebra.*

*Proof.* By the definition of  $L_\infty$ -algebras, we need to check the equation in  $C_{\text{AvA}}(V)$  :

$$(6.1) \sum_{i+j=n} \sum_{\sigma \in S(i, n-i)} (-1)^{i(n-i)} \chi(\sigma, x_1, \dots, x_n) l_{n-i+1}(l_i(x_{\sigma(1)}, \dots, x_{\sigma(i)}), x_{\sigma(i+1)}, \dots, x_{\sigma(n)}) = 0.$$

If all  $x_i$  are contained in  $C_A(V) = \text{Hom}(T^c(sV), sV)$ , then equation 6.1 is just the Jacobi identity on the graded Lie algebra  $C_A(A)$ . Apart from this, by the definition of  $l_n$ , the term  $l_{n-i+1}(l_i(x_{\sigma(1)}, \dots, x_{\sigma(i)}), x_{\sigma(i+1)}, \dots, x_{\sigma(n)})$  is trivial unless there are exactly two elements in  $\{x_1, \dots, x_n\}$  coming from  $C_A(V)$  and all other  $n-2$  elements are contained in  $C_{\text{AvO}}(V)_l$  or  $C_{\text{AvO}}(V)_r$ . We assume that  $x_1 = sh_1 \in \text{Hom}((sV)^{\otimes n_1}, sV)$  and  $x_2 = sh_2 \in \text{Hom}((sV)^{\otimes n_2}, sV)$ . Then  $n$  must be  $n_1 + n_2 + 1$ . And we assume  $x_i = g_{i-2} \in C_{\text{AvO}}(V)_r$ . When  $g_i \in \text{Hom}(k, V) \oplus \text{Hom}(sV, V) \oplus \text{Hom}(T^c(sV)_{\geq 2}, V)_l$ , the proof is all the same.

For simplicity, we assume that all  $g_i$  are not in  $\text{Hom}(s, V)$ . If there is some  $g_i \in \text{Hom}(k, V)$ , the calculation is similar.

So now, we are going to check the equation 6.1 for  $x_1 = sh_1, x_2 = sh_2, x_3 = g_1, \dots, x_{n_1+n_2+1} = g_{n_1+n_2-1}$  with  $g_i \in \text{Hom}(sV, V) \oplus \text{Hom}(k, V) \oplus \text{Hom}(sV, V) \oplus \text{Hom}(T^c(sV)_{\geq 2}, V)_r$ . Then the term  $l_{n-i+1}(l_i(x_{\sigma(1)}, \dots, x_{\sigma(i)}), x_{\sigma(i+1)})$  vanishes unless  $i = 2, n_1 + 1, n_2 + 1$ . Let's compute them one by one.

(A) When  $i = 2$ , the  $(2, n-2)$ -shuffle is identity map, we have

$$\begin{aligned} & l_{n_1+n_2-1}(l_2(sh_1, sh_2), g_1, \dots, g_{n_1+n_2-1}) \\ &= l_{n_1+n_2-1}([sh_1, sh_2]_G, g_1, \dots, g_{n_1+n_2-1}) \\ &= \sum_{\pi \in S_{n_1+n_2-1}} (-1)^\varepsilon \left\{ s^{-1}(sh_1 \bar{\circ} sh_2) \circ (sg_{\pi(1)}, \dots, sg_{\pi(n_1+n_2-1)}) \right. \\ & \quad \left. - (-1)^{(|g_{\pi(1)}|+1)(|h_1|+|h_2|)} g_{\pi(1)} \bar{\circ} ((sh_1 \bar{\circ} sh_2) \circ (\text{id}_{sV} \otimes sg_{\pi(2)} \otimes \dots \otimes sg_{\pi(n)})) \right\} \\ & \quad + \sum_{\pi \in S_{n_1+n_2-1}} (-1)^\varepsilon (-1)^{(|h_1|+1)(|h_2|+1)+1} \left\{ s^{-1}(sh_2 \bar{\circ} sh_1) \circ (sg_{\pi(1)}, \dots, sg_{\pi(n_1+n_2-1)}) \right. \\ & \quad \left. - (-1)^{(|g_{\pi(1)}|+1)(|h_1|+|h_2|)} g_{\pi(1)} \bar{\circ} ((sh_2 \bar{\circ} sh_1) \circ (\text{id}_{sV} \otimes sg_{\pi(2)} \otimes \dots \otimes sg_{\pi(n_1+n_2-1)})) \right\} \\ &= \sum_{\pi \in S_{n_1+n_2-1}} (-1)^\varepsilon \left\{ \sum_{i=0}^{n_1-1} (-1)^{(|h_2|+1)(\sum_{j=1}^i (|g_{\pi(j)}|+1))} h_1 \circ (sg_{\pi(1)}, \dots, sg_{\pi(i)}, sh_2 \circ (sg_{\pi(i+1)}, \dots, sg_{\pi(i+n_2)}), \dots, sg_{\pi(n_1+n_2-1)}) \right\} \end{aligned}$$

$$\begin{aligned}
& + \sum_{\pi \in \mathcal{S}_{n_1+n_2-1}} (-1)^\varepsilon (-1)^{1+(\lg_{\pi(1)}|+1)(|h_1|+|h_2|)} g_{\pi(1)} \bar{\circ} \left\{ sh_1 \circ \left( sh_2 \circ (\text{id}_{sV} \otimes sg_{\pi(2)} \otimes \dots \otimes sg_{\pi(n_2)}) \otimes \dots \otimes sg_{\pi(n_1+n_2-1)} \right) \right. \\
& + \sum_{i=1}^{n_1-1} (-1)^{(|h_2|+1)(\sum_{j=2}^i (\lg_{\pi(j)}|+1))} sh_1 \circ \left( \text{id}_{sV} \otimes sg_{\pi(2)} \otimes \dots \otimes sg_{\pi(i)} \otimes sh_2 \circ (sg_{\pi(i+1)}, \dots, sg_{\pi(i+n_2)}) \otimes \dots \otimes sg_{\pi(n_1+n_2-1)} \right) \Big\} \\
& + \sum_{\pi \in \mathcal{S}_{n_1+n_2-1}} (-1)^\varepsilon (-1)^{(|h_1|+1)(|h_2|+1)+1} \sum_{i=0}^{n_2-1} (-1)^{(|h_1|+1)(\sum_{j=1}^i (\lg_{\pi(j)}|+1))} \bullet \\
& \quad \left\{ h_2 \circ \left( sg_{\pi(1)} \otimes \dots \otimes sg_{\pi(i)} \otimes sh_1 \circ (sg_{\pi(i+1)} \otimes \dots \otimes sg_{\pi(i+n_1)}) \otimes \dots \otimes sg_{\pi(n_1+n_2-1)} \right) \right\} \\
& + \sum_{\pi \in \mathcal{S}_{n_1+n_2-1}} (-1)^\varepsilon (-1)^{(|h_1|+1)(|h_2|+1)+(\lg_{\pi(1)}|+1)(|h_1|+|h_2|)} \bullet \\
& \quad g_{\pi(1)} \bar{\circ} \left\{ sh_2 \circ \left( sh_1 \circ (\text{id}_{sV} \otimes sg_{\pi(2)} \otimes \dots \otimes sg_{\pi(n_2)}) \otimes \dots \otimes sg_{\pi(n_1+n_2-1)} \right) \right. \\
& + \sum_{i=1}^{n_2-1} (-1)^{(|h_1|+1)(\sum_{j=2}^i (\lg_{\pi(j)}|+1))} sh_2 \circ \left( \text{id}_{sV}, sg_{\pi(2)}, \dots, sg_{\pi(i)}, sh_1 \circ (sg_{\pi(i+1)}, \dots, sg_{\pi(i+n_2)}), \dots, sg_{\pi(n_1+n_2-1)} \right) \Big\}.
\end{aligned}$$

$$\text{where } (-1)^\varepsilon = \chi(\pi, g_1, \dots, g_{n_1+n_2-1}) (-1)^{(n_1+n_2-1)(h_1+h_2)+\sum_{k=1}^{n_1+n_2-2} \sum_{j=1}^k |\lg_{\pi(j)}|}.$$

(B) When  $i = n_2 + 1$ ,

$$\begin{aligned}
& \sum_{\rho \in \mathcal{S}(n_2+1, n_1)} (-1)^{(n_2+1)n_1} \chi(\rho, x_1, \dots, x_{n_1+n_2+1}) l_{n_1+1}(l_{n_2+1}(x_{\rho(1)}, \dots, x_{\rho(n_2+1)}), \dots, x_{\rho(n_1+n_2+1)}) \\
& = \sum_{\sigma \in \mathcal{S}(n_2, n_1-1)} (-1)^{(n_2+1)n_1} \chi(\sigma, g_1, \dots, g_{n_1+n_2-1}) (-1)^{(|h_1|+1)(|h_2|+1+\sum_{j=1}^{n_2} |\lg_{\sigma(j)}|)+n_2+1} \bullet \\
& \quad l_{n_1+1}(l_{n_2+1}(sh_2, g_{\sigma(1)}, \dots, g_{\sigma(n_2)}), sh_1, g_{\sigma(n_2+1)}, \dots, g_{\sigma(n_1+n_2-1)}) \\
& = \sum_{\sigma \in \mathcal{S}(n_2, n_1-1)} (-1)^{(n_2+1)n_1} \chi(\sigma, g_1, \dots, g_{n_1+n_2-1}) (-1)^{(|h_1|+1)(|h_2|+1+\sum_{j=1}^{n_2} |\lg_{\sigma(j)}|)+n_2+1} \bullet \\
& \quad (-1)^{(|h_1|+1)(|h_2|+1+\sum_{j=1}^{n_2} |\lg_{\sigma(j)}|+n_2-1)+1} l_{n_1+1} \left( sh_1, \underbrace{l_{n_2+1}(sh_2, g_{\sigma(1)}, \dots, g_{\sigma(n_2)})}_{:=\widehat{h_2}}, g_{\sigma(n_2+1)}, \dots, g_{\sigma(n_1+n_2-1)} \right) \\
& = \sum_{\sigma \in \mathcal{S}(n_2, n_1-1)} \chi(\sigma, g_1, \dots, g_{n_1+n_2-1}) (-1)^{(|h_1|+1)(n_2-1)+(n_2+1)n_1+n_2} l_{n_1+1} \left( sh_1, \widehat{h_2}, g_{\sigma(n_2+1)}, \dots, g_{\sigma(n_1+n_2-1)} \right) \\
& = \sum_{\sigma \in \mathcal{S}(n_2, n_1-1)} \sum_{\tau \in \mathcal{S}_{n_1-1}} (-1)^{\theta_1} \left\{ \underbrace{h_1 \circ (s\widehat{h_2} \otimes sg_{\sigma(n_2+\tau(1))} \otimes \dots \otimes sg_{\sigma(n_2+\tau(n_1-1))})}_{B_1} \right. \\
& \quad \left. - (-1)^{(|\widehat{h_2}|+1)(|h_1|+1)} \underbrace{\widehat{h_2} \bar{\circ} (sh_1 \circ (\text{id}_{sV} \otimes sg_{\sigma(n_2+\tau(1))} \otimes \dots \otimes sg_{\sigma(n_2+\tau(n_1-1))})}_{B_2} \right\}
\end{aligned}$$

$$\begin{aligned}
& + \sum_{\sigma \in S(n_2, n_1-1)} \sum_{\tau \in S(n_1-1)} \sum_{i=1}^{n_1-1} (-1)^{\theta_2} \left\{ \underbrace{h_1 \circ (sg_{\sigma(n_2+\tau(1))} \otimes \dots \otimes sg_{\sigma(n_2+\tau(i))} \otimes \widehat{sh_2} \otimes \dots \otimes sg_{\sigma(n_2+\tau(n_1-1))})}_{B_3} \right. \\
& \left. - (-1)^{(|h_1|+1)(|g_{\sigma(n_2+\tau(1))}|+1)} \underbrace{g_{\sigma(n_2+\tau(1))} \bar{\circ} (sh_1 \circ (\text{id}_{sV} \otimes sg_{\sigma(n_2+\tau(2))} \otimes \dots \otimes sg_{\sigma(n_2+\tau(i))} \otimes \widehat{sh_2} \otimes \dots \otimes sg_{\sigma(n_2+\tau(n_1-1))}))}_{B_4} \right\}
\end{aligned}$$

where

$$\begin{aligned}
(-1)^{\theta_1} &= \chi(\sigma, g_1, \dots, g_{n_1+n_2-1}) \chi(\tau, g_{\sigma(n_2+1)}, \dots, g_{\sigma(n_2+n_1-1)}) \\
&\cdot (-1)^{(|h_1|+1)(n_2-1)+(n_2+1)n_1+n_2} (-1)^{n_1(|h_1|+1)+(n_1-1)(|\widehat{h_2}|) + \sum_{k=1}^{n_1-2} \sum_{j=1}^k |g_{\sigma(n_2+\tau(j))}|} \\
(-1)^{\theta_2} &= \chi(\sigma, g_1, \dots, g_{n_1+n_2-1}) \chi(\tau, g_{\sigma(n_2+1)}, \dots, g_{\sigma(n_2+n_1-1)}) \cdot (-1)^{(|h_1|+1)(n_2-1)+(n_2+1)n_1+n_2} \\
&\cdot (-1)^{n_1(|h_1|+1) + \sum_{k=1}^{n_1-2} \sum_{j=1}^k |g_{\sigma(n_2+\tau(j))}| + \sum_{j=1}^i |g_{\sigma(n_2+\tau(j))}| + (n_1-i-1)|\widehat{h_2}| + (|\widehat{h_2}|)(\sum_{j=1}^i |g_{\sigma(n_2+\tau(j))}|) + i}
\end{aligned}$$

Let's compute the terms  $B_1, B_2, B_3, B_4$  together with their coefficients.

$$\begin{aligned}
B_1 &= \sum_{\sigma \in S(n_2, n_1-1)} \sum_{\tau \in S_{n_1-1}} (-1)^{\theta_1} h_1 \circ (\widehat{sh_2} \otimes \dots \otimes sg_{\sigma(n_2+\tau(1))} \otimes \dots \otimes sg_{\sigma(n_2+\tau(n_1-1))}) \\
&= \sum_{\sigma \in S(n_2, n_1-1)} \sum_{\tau \in S_{n_1-1}} (-1)^{\theta_1} h_1 \circ (sl_{n_2+1}(sh_2, g_{\sigma(1)}, \dots, g_{\sigma(n_2)}), sg_{\sigma(n_2+\tau(1))} \otimes \dots \otimes sg_{\sigma(n_2+\tau(n_1-1))}) \\
&= \sum_{\sigma \in S(n_2, n_1-1)} \sum_{\tau \in S_{n_1-1}} (-1)^{\theta_1} \sum_{\delta \in S_{n_2}} \chi(\delta, g_{\sigma(1)}, \dots, g_{\sigma(n_2)}) (-1)^{n_2(|h_2|+1) + \sum_{k=1}^{n_2-1} \sum_{j=1}^k |g_{\sigma(\delta(j))}|} \bullet \\
&\quad h_1 \circ \left\{ (sh_2 \circ (sg_{\sigma\delta(1)}, \dots, sg_{\sigma\delta(n_2)}) \otimes sg_{\sigma(n_2+\tau(1))} \otimes \dots \otimes sg_{\sigma(n_2+\tau(n_1-1))}) - (-1)^{(|g_{\sigma\delta(1)}|+1)(|h_2|+1)} \bullet \right. \\
&\quad \left. (sg_{\sigma\delta(1)} \bar{\circ} (sh_2 \circ (\text{id}_{sV} \otimes sg_{\sigma\delta(2)} \otimes \dots \otimes sg_{\sigma\delta(n_2)})) \otimes sg_{\sigma(n_2+\tau(1))} \otimes \dots \otimes sg_{\sigma(n_2+\tau(n_1-1))}) \right\} \\
&= \sum_{\pi \in S(n_1+n_2-1)} (-1)^{\eta_1} h_1 \circ \left\{ (sh_2 \circ (sg_{\pi(1)} \otimes \dots \otimes sg_{\pi(n_2)}) \otimes sg_{\pi(n_2+1)} \otimes \dots \otimes sg_{\pi(n_2+n_1-1)}) \right. \\
&\quad \left. - (-1)^{(|g_{\pi(1)}|+1)(|h_2|+1)} (sg_{\pi(1)} \bar{\circ} (sh_2 \circ (\text{id}_{sV} \otimes sg_{\pi(2)} \otimes \dots \otimes sg_{\pi(n_2)})) \otimes sg_{\pi(n_2+1)} \otimes \dots \otimes sg_{\pi(n_2+n_1-1)}) \right\}
\end{aligned}$$

where

$$\begin{aligned}
(-1)^{\eta_1} &= (-1)^{\theta_1} \chi(\delta, g_{\sigma(1)}, \dots, g_{\sigma(n_2)}) (-1)^{n_2(|h_2|+1) + \sum_{k=1}^{n_2-1} \sum_{j=1}^k |g_{\sigma(\delta(j))}|} \\
&= \chi(\pi, g_1, \dots, g_{n_1+n_2-1}) (-1)^{(|h_1|+|h_2|)(n_1+n_2-1)+1 + \sum_{k=1}^{n_1+n_2-2} \sum_{j=1}^k |g_{\pi(j)}|}
\end{aligned}$$

Notice that, in the last equality above, we use lemma 7.3 to replace the triple  $(\delta, \sigma, \tau)$  by its corresponding permutation  $\pi \in S_{n_1+n_2-1}$  and we use the fact

$$\chi(\pi, g_1, \dots, g_{n_1+n_2-1}) = \chi(\sigma, g_1, \dots, g_{n_1+n_2-1}) \chi(\tau, g_{\sigma(n_2+1)}, \dots, g_{\sigma(n_2+n_1-1)}) \chi(\delta, g_{\sigma(1)}, \dots, g_{\sigma(n_2)}).$$

Similarly, we have

$$\begin{aligned}
\mathbf{B}_2 &= \sum_{\sigma \in S(n_2, n_1-1)} \sum_{\tau \in S_{n_1-1}} (-1)^{\theta_1} (-1)^{1+(\widehat{h_2}+1)(|h_1|+1)} \widehat{h_2} \bar{\circ} (sh_1 \circ (\text{id}_{sV} \otimes sg_{\sigma(n_2+\tau(1))} \otimes \dots \otimes sg_{\sigma(n_2+\tau(n_1-1))})) \\
&= \sum_{\pi \in S(n_1+n_2-1)} (-1)^{\eta_2} \left\{ h_2 \circ (sg_{\pi(1)} \otimes \dots \otimes sg_{\pi(n_2)}) - (-1)^{(|h_2|+1)(|g_{\pi(1)}|+1)} \bullet \right. \\
&\quad \left. (g_{\pi(1)} \bar{\circ} (sh_2 \circ (\text{id}_{sV} \otimes sg_{\pi(2)} \otimes \dots \otimes sg_{\pi(n_2)}))) \right\} \underbrace{\bar{\circ} (sh_1 \circ (\text{id}_{sV} \otimes sg_{\pi(n_2+1)} \otimes \dots \otimes sg_{\pi(n_2+n_1-1)}))}_{:= \widetilde{h_1}} \\
&= \sum_{\pi \in S(n_1+n_2-1)} (-1)^{\eta_2} \left\{ \sum_{k=1}^{n_2} (-1)^{\left(\sum_{j=k+1}^{n_2} (|g_{\pi(j)}|+1)\right)(|\widetilde{h_1}|)} h_2 \circ (sg_{\pi(1)} \otimes \dots \otimes sg_{\pi(k)} \bar{\circ} \widetilde{h_1} \otimes sg_{\pi(k+1)} \otimes \dots \otimes sg_{\pi(n_2)}) \right\} \\
&\quad + (-1)^{\eta_2} (-1)^{1+(|h_2|+1)(|g_{\pi(1)}|+1)} \left\{ g_{\pi(1)} \bar{\circ} (sh_2 \circ (\text{id}_{sV} \otimes sg_{\pi(2)} \otimes \dots \otimes sg_{\pi(n_2)}), \widetilde{h_1}) \right\} \\
&\quad + (-1)^{|\widetilde{h_1}| \left(\sum_{j=2}^{n_2} (|g_{\pi(j)}|+1)\right)} g_{\pi(1)} \bar{\circ} (sh_2 \circ (\widetilde{h_1} \otimes sg_{\pi(2)} \otimes \dots \otimes sg_{\pi(n_2)})) \\
&\quad + \sum_{k=2}^{n_2} (-1)^{|\widetilde{h_1}| \left(\sum_{j=k+1}^{n_2} (|g_{\pi(j)}|+1)\right)} g_{\pi(1)} \bar{\circ} (sh_2 \circ (\text{id}_{sV} \otimes sg_{\pi(2)} \otimes \dots \otimes sg_{\pi(k)} \bar{\circ} \widetilde{h_1} \otimes sg_{\pi(k+1)} \otimes \dots \\
&\quad \dots \otimes sg_{\pi(n_2)})) + (-1)^{|\widetilde{h_1}| \left(\sum_{j=2}^{n_2} (|g_{\pi(j)}|+1)+|h_2|+1\right)} g_{\pi(1)} \bar{\circ} (\widetilde{h_1}, sh_2 \circ (\text{id}_{sV} \otimes sg_{\pi(2)} \otimes \dots \otimes sg_{\pi(n_2)})) \Big\}
\end{aligned}$$

where

$$\begin{aligned}
(-1)^{\eta_2} &= (-1)^{\eta_1} (-1)^{1+(\widehat{h_2}+1)(|h_1|+1)} \\
&= \chi(\pi, g_1, \dots, g_{n_1+n_2-1}) (-1)^{(|h_1|+|h_2|)(n_1+n_2-1)+1+\sum_{k=1}^{n_1+n_2-2} \sum_{j=1}^k |g_{\pi(j)}|+1+(\widehat{h_2}+1)(|h_1|+1)} \\
&= \chi(\pi, g_1, \dots, g_{n_1+n_2-1}) (-1)^{(|h_1|+|h_2|)(n_1+n_2-1)+\sum_{k=1}^{n_1+n_2-2} \sum_{j=1}^k |g_{\pi(j)}|+(|h_2|+n_2-1+\sum_{j=1}^{n_2} |g_{\pi(j)}|)(|h_1|+1)}
\end{aligned}$$

$$\begin{aligned}
\mathbf{B}_3 &= \sum_{\sigma \in S(n_2, n_1-1)} \sum_{\tau \in S(n_1-1)} \sum_{i=1}^{n_1-1} (-1)^{\theta_2} h_1 \circ \left\{ sg_{\sigma(n_2+\tau(1))} \otimes \dots \otimes sg_{\sigma(n_2+\tau(i))} \otimes \widehat{sh_2} \otimes \dots \otimes sg_{\sigma(n_2+\tau(n_1-1))} \right\} \\
&= \sum_{\pi \in S(n_2+n_1-1)} \sum_{i=1}^{n_1-1} (-1)^{\eta_3} h_1 \circ \left\{ sg_{\pi(n_2+1)} \otimes \dots \otimes sg_{\pi(n_2+i)} \otimes sh_2 \circ (sg_{\pi(1)} \otimes \dots \otimes sg_{\pi(n_2)}) \otimes \dots \otimes sg_{\pi(n_1+n_2-1)} \right. \\
&\quad \left. - (-1)^{(|h_2|+1)(|g_{\pi(1)}|+1)} sg_{\pi(n_2+1)} \otimes \dots \otimes sg_{\pi(n_2+i)} \otimes sg_{\pi(1)} \bar{\circ} (sh_2 \circ (\text{id}_{sV} \otimes sg_{\pi(2)} \otimes \dots \otimes sg_{\pi(n_2)})) \otimes \dots \right. \\
&\quad \left. \dots \otimes sg_{\pi(n_1+n_2-1)} \right\}
\end{aligned}$$

where

$$\begin{aligned}
(-1)^{\eta_3} &= (-1)^{\theta_2} \chi(\delta, g_{\sigma(1)}, \dots, g_{\sigma(n_2)}) (-1)^{n_2(|h_2|+1)+\sum_{k=1}^{n_2-1} |g_{\pi(j)}|} \\
&= \chi(\sigma, g_1, \dots, g_{n_1+n_2-1}) \chi(\tau, g_{\sigma(n_2+1)}, \dots, g_{\sigma(n_2+n_1-1)}) \cdot (-1)^{(|h_1|+1)(n_2-1)+(n_2+1)n_1+n_2}
\end{aligned}$$



$$\begin{aligned}
& \cdot (-1)^{n_1(|h_1|+1)+\sum_{k=1}^{n_1-2}\sum_{j=1}^k|g_{\sigma(\tau(j))|+\sum_{j=1}^i|g_{\sigma(n_2+\tau(j))|+(n_1-i-1)|\widehat{h_2}|+|\widehat{h_2}|(\sum_{j=n_2+1}^{n_2+i}|g_{\pi(j)}|)+i} \\
& \cdot (-1)^{n_2(|h_2|+1)+\sum_{k=1}^{n_2-1}|g_{\pi(j)}|}\chi(\delta, g_{\sigma(1)}, \dots, g_{\sigma(n_2)}) \\
& = \chi(\pi, g_1, \dots, g_{n_1+n_2-1})(-1)^{(|h_1|+|h_2|)(n_1+n_2-1)+1+\sum_{k=1}^{n_1+n_2-2}\sum_{j=1}^k|g_{\pi(j)}|+(1+|h_2|+n_2+\sum_{j=1}^{n_2}|g_{\pi(j)}|)(i+\sum_{j=1}^i|g_{\pi(n_2+j)}|)}
\end{aligned}$$

$$\begin{aligned}
B_4 = & \sum_{\pi \in S(n_2+n_1-1)} \sum_{i=1}^{n_1-1} (-1)^{\eta_4} g_{\pi(n_2+1)} \bar{\circ} \left\{ sh_1 \circ (\text{id}_{sV} \otimes sg_{\pi(n_2+2)} \otimes \dots \otimes sg_{\pi(n_2+i)} \otimes sh_2 \circ (sg_{\pi(1)} \otimes \dots \otimes sg_{\pi(n_2)})) \otimes \dots \right. \\
& \otimes sg_{\pi(n_2+n_1-1)}) - (-1)^{(|h_2|+1)(|g_{\pi(1)}|+1)} sh_1 \circ (\text{id}_{sV} \otimes sg_{\pi(n_2+2)} \otimes \dots \otimes sg_{\pi(n_2+i)} \otimes sg_{\pi(1)} \bar{\circ} (sh_2 \circ (1_{sV} \otimes sg_{\pi(2)} \otimes \dots \\
& \dots \otimes sg_{\pi(n_2)})) \otimes \dots \otimes sg_{\pi(n_2+n_1-1)}) \left. \right\}
\end{aligned}$$

where

$$(-1)^{\eta_4} = (-1)^{\eta_3} (-1)^{1+(|h_1|+1)(|g_{\pi(n_2+1)}|+1)}.$$

(C) When  $i = n_1 + 1$ ,

$$\begin{aligned}
& \sum_{\rho \in S(n_1+1, n_2)} (-1)^{(n_1+1)n_2} \chi(\rho, x_1, \dots, x_{n_1+n_2+1}) l_{n_2+1}(l_{n_1+1}(x_{\rho(1)}, \dots, x_{\rho(n_2+1)}), \dots, x_{\rho(n_1+n_2+1)}) \\
& = \sum_{\sigma \in S(n_1, n_2-1)} (-1)^{(n_1+1)n_2} (-1)^{n_1+(|h_2|+1)(\sum_{j=1}^{n_1} g_{\sigma(j)})} \chi(\sigma, g_1, g_2, \dots, g_{n_1+n_2-1}) \bullet \\
& \quad l_{n_2+1}(l_1(sh_1, g_{\sigma(1)}, \dots, g_{\sigma(n_1)}), sh_2, g_{\sigma(n_1+1)}, \dots, g_{\sigma(n_1+n_2-1)}) \\
& = \sum_{\sigma \in S(n_1, n_2-1)} (-1)^{(n_1+1)n_2} (-1)^{n_1+(|h_2|+1)(\sum_{j=1}^{n_1} g_{\sigma(j)})} (-1)^{(|h_2|+1)(\sum_{j=1}^{n_1} g_{\sigma(j)}+|h_1|+n_1)+1} \chi(\sigma, x_1, \dots, x_{n_1+n_2+1}) \bullet \\
& \quad l_{n_2+1}(sh_2, l_{n_1+1}(sh_1, g_{\sigma(1)}, \dots, g_{\sigma(n_1)}), g_{\sigma(n_1+1)}, \dots, g_{\sigma(n_2+n_1-1)}) \\
& = \sum_{\sigma \in S(n_1, n_2-1)} (-1)^{(n_1+1)(n_2+1)+(|h_2|+1)(|h_1|+n_1)} \chi(\sigma, g_1, \dots, g_{n_1}) \bullet \\
& \quad l_{n_2+1}(sh_2, l_{n_1+1}(sh_1, g_{\sigma(1)}, \dots, g_{\sigma(n_1)}), g_{\sigma(n_1+1)}, \dots, g_{\sigma(n_2+n_1-1)})
\end{aligned}$$

Switch the roles of  $h_1, n_1$  and  $h_2, n_2$  in the expansion of

$$l_{n_1+1}(sh_1, l_{n_2+1}(sh_2, g_{\sigma(1)}, \dots, g_{\sigma(n_2)}), g_{\sigma(n_2+1)}, \dots, g_{\sigma(n_1+n_2-1)}),$$

which has been computed in case (B), then we will get the expansion of

$$l_{n_2+1}(sh_2, l_{n_1+1}(sh_1, g_{\sigma(1)}, \dots, g_{\sigma(n_1)}), g_{\sigma(n_1+1)}, \dots, g_{\sigma(n_2+n_1-1)}).$$

Then the expansion for case (C) comes out. Then, we can found that the terms of the same form appear exactly twice in the expansion of (A)+(B)+(C), and they have opposite signs. Thus (A) + (B) + (C) = 0 holds. So  $C_{AvA}(V)$  forms a  $L_\infty$ -algebra.  $\square$

## REFERENCES

- [1] M. Aguiar, *Pre-Poisson algebras*, Lett. Math. Phys. **54** (2000) 263-277. 2
- [2] G. Baxter, *An analytic problem whose solution follows from a simple algebraic identity*, Pacific J. Math. **10** (1960) 731-742. 2
- [3] G. Birkhoff, *Moyennes de fonctions bornées*, Coil. Internat. Centre Nat. Recherche Sci. (Paris), Algèbre Théorie Nombres **24** (1949) 149-153. 2
- [4] N. H. Bong, *Some apparent connection between baxter and averaging operators*, J. Math. Anal. Appl. **56** (1976) 330-345. 2
- [5] W. Cao, *An algebraic study of averaging operators*, Ph.D. thesis, Rutgers University at Newark (2000). 2
- [6] Z. Chen, H. Ge and M. Xiang, *Talk at The Third Conference of Operad Theory and Related Topics*, September 2020. 2
- [7] J. Kampé de Fériet, *Introduction to the statistical theory of turbulence, correlation and spectrum*, in The Institute of Fluid Dynamics and Applied Mathematics, Lecture Series No. **8**, prepared by S. I. Pai (University of Maryland, 1950-51). 2
- [8] J. L. B. Gamlen and J. B. Miller, *Averaging and Reynolds operators on Banach algebras II. spectral properties of averaging operators*, J. Math. Anal. Appl. **23** (1968) 183-197. 2
- [9] X. Gao and T. Zhang, *Averaging algebras, rewriting systems and GröbnerCSchirshov bases*, J. Algebra Appl. **17** (2018), no. 7, 1850130, 26 pp. 2
- [10] M. Gerstenhaber, *The cohomology structure of an associative ring*. Ann. of Math. (2) **78** 1963 267-288. 15
- [11] L. Guo, *An Introduction to RotaCBaxter Algebra* (International Press (US) and Higher Education Press (China), 2012). 2
- [12] J. Kampé de Fériet, *L'état actuel du problème de la turbulence (I and II)*, La Sci. Aérienne **3** (1934) 9-34, **4** (1935) 12-52. 2
- [13] J. L. Kelley, *Averaging operators on  $C^\infty(X)$* , Illinois J. Math. **2** (1958) 214-223. 2
- [14] T. Lada and M. Markl, *Strongly homotopy Lie algebras*, Communications in algebra, **23** (1995), no. 6, 2147-2161. 14
- [15] T. Lada and J. Stasheff, *Introduction to sh Lie algebras for physicists*, Internat. J. Theoret. Phys. **32** (1993), 1087-1103. 14
- [16] J. L. Loday, *Dialgebras*, in Dialgebras and related operads, Lecture Notes in Math. 1763 (2002) 7-66. 2
- [17] J. B. Miller, *Averaging and Reynolds operators on Banach algebra I, Representation by derivation and antiderivations*, J. Math. Anal. Appl. **14** (1966) 527-548. 2
- [18] S. T. C. Moy, *Characterizations of conditional expectation as a transformation on function spaces*, Pacific J. Math. **4** (1954) 47-63. 2
- [19] J. Pei and L. Guo, *Averaging algebras, Schröder numbers, rooted trees and operads*, J. Algebra Comb. **42** (2015) 73-109. 2
- [20] O. Reynolds, *On the dynamic theory of incompressible viscous fluids and the determination of the criterion*, Phil. Trans. Roy. Soc. A **136** (1895) 123-164. 2
- [21] G. C. Rota, *Reynolds operators*, Proceedings of Symposia in Applied Mathematics, Vol. XVI (American Mathematical Society, Providence, R.I., 1964), pp. 70-83. 2
- [22] G. C. Rota, *Baxter algebras and combinatorial identities I, II*, Bull. Amer. Math. Soc. **75** (1969) 325-329, 330-334. 2
- [23] Y. Sheng, R. Tang and C. Zhu, *The controlling  $L_\infty$ -algebras, cohomology and homotopy of embedding tensors and Lie-Leibniz triples*, arXiv:2009.11096. 2
- [24] J. Stasheff, *Differential graded Lie algebras, quasi-Hopf algebras and higher homotopy algebras*, Quantum groups (Leningrad, 1990), 120-137, Lecture Notes in Math 1510, Springer, Berlin, 1992. 14
- [25] A. Triki, *Extensions of positive projections and averaging operators*, J. Math. Anal. **153** (1990) 486C496. 2

SCHOOL OF MATHEMATICAL SCIENCES, SHANGHAI KEY LABORATORY OF PMMP, EAST CHINA NORMAL UNIVERSITY, SHANGHAI 200241, CHINA

E-mail address: wangkai@math.ecnu.edu.cn

SCHOOL OF MATHEMATICAL SCIENCES, SHANGHAI KEY LABORATORY OF PMMP, EAST CHINA NORMAL UNIVERSITY, SHANGHAI 200241, CHINA

E-mail address: gdzhou@math.ecnu.edu.cn