

Impulsively Excited Gravitational Quantum States: Echoes and Time-resolved Spectroscopy

I. Tutunnikov,¹ K. V. Rajitha,¹ A. Yu. Voronin,^{2,*} V. V. Nesvizhevsky,^{3,†} and I. Sh. Averbukh^{1,‡}

¹*AMOS and Department of Chemical and Biological Physics,
The Weizmann Institute of Science, Rehovot 7610001, Israel*

²*Lebedev Institute, 53 Leninsky pr., Moscow, Russia, Ru-119333*

³*Institut Max von Laue-Paul Langevin (ILL), 71 avenue des Martyrs, Grenoble, France, F-38042*

We theoretically study an *impulsively* excited quantum bouncer (QB) - a particle bouncing off a surface in the presence of gravity. A pair of time-delayed pulsed excitations is shown to induce a wave-packet echo effect - a partial rephasing of the QB wave function appearing at twice the delay between pulses. In addition, an appropriately chosen observable [here, the population of the ground gravitational quantum state (GQS)] recorded as a function of the delay is shown to contain the transition frequencies between the GQSs, their populations, and partial phase information about the wave packet quantum amplitudes. The wave-packet echo effect is a promising candidate method for precision studies of GQSs of ultra-cold neutrons, atoms, and anti-atoms confined in closed gravitational traps.

Introduction—In last decades, massive quantum particle bouncing off a surface under the influence of gravity turned from being an issue of textbooks and pedagogical essays [1–4] into a subject of precision experiments on atom-optics gravitational cavities [5, 6] and physics of ultra-cold neutrons (UCNs) [7]. The observation of GQSs [8–11] and whispering gallery states [12, 13] of neutrons (n) fueled a vast research in this area, which among other goals, aims to the search for new fundamental short-range interactions and physics beyond Standard Model, as well as verification of weak equivalence principle in the quantum regime (see e.g. the introduction of [14], and references therein).

Cold atoms and anti-atoms can also bounce on surfaces and form GQSs [15] due to the quantum reflection from a rapidly changing attractive van der Waals/Casimir-Polder surface potential (see, e.g. [16] and references therein). In contrast to the extremely precise measurements of gravitational properties of matter [17–19], the best constraint [20] for the gravitational mass (acceleration) of antimatter does not allow even to define the sign of acceleration. Several collaborations perform experiments at CERN [21–23] aiming to improve the accuracy. The GQS method seems to promise the best accuracy for anti-hydrogen atoms ($\bar{H}$) [24].

Resonant spectroscopy of neutron GQSs was proposed in [25], measured using periodic excitation of QBs by mechanical vibrations of the surface [26–30], and being implemented using a periodically changing magnetic field gradient [31, 32]. Spatial distribution of GQSs of n was measured with micron resolution [33]. For bouncing $\bar{H}$ atoms, a resonant spectroscopy [34, 35] and interferometry [24, 36, 37] approaches have been developed.

Here, we study the physics of *impulsively* excited QBs, and consider two example excitations: (i) by applying a pulsed magnetic field gradient interacting with the QB's magnetic dipole moment, and (ii) by a jolt caused by an impulsive shake of the surface. Short laser pulses

have been widely used for time-resolved molecular spectroscopy, however the related aspects of the GQS spectroscopy are unexplored yet. A spectacular effect in the dynamics of kick-excited nonlinear systems is the echo phenomenon first discovered by E. Hahn in spin systems [38, 39] (spin echo). Since then, various types of echoes have been observed, including photon echoes [40, 41], cyclotron echoes [42], plasma-wave echoes [43], neutron spin echo [44], cold atom echoes in optical traps [45–47], echoes in particle accelerators [48–52], and more recently—alignment and orientation echoes in molecular gases [53–61]. In these examples, echo appears in inhomogeneous ensembles of many particles evolving at different frequencies. Echoes were also observed in single quantum objects: in a single mode of quantized electromagnetic field interacting with atoms passing through a cavity [62, 63], and in single vibrationally excited molecules [64].

In the first part of this Letter, we demonstrate, for the first time, that highly nonlinear dynamics of quantum gravitational wave packets favors observation of the echo in a *single* QB. Then, we explore a response of QB to a pair of time-delayed kicks, and analyze its dependence on the delay between kicks. The population of ground GQS as a function of the delay is shown to contain the transition frequencies between the populated GQSs, as well as partial phase information about the QB wave packet. This paves the way to a new kind of time-resolved GQSs spectroscopy which has a number of advantages. It doesn't require fine tuning of the excitation frequency to a specific resonance between the GQSs, and eliminates some frequency shifts characteristic to the resonant GQSs spectroscopy [32].

Free quantum bouncer—The vertical motion of the QB (along Z axis) is quantized and decoupled from the motion along the X, Y axes. The eigenfunctions ψ_i and




Figure 1. Phase space analysis. $N = 2 \times 10^4$ particles bounce on a surface, and kicked at $t = t_k = 60$. Kick parameters: $a_k = 0.5$ (e.g. for neutrons: $|\mu| = 60.3$ neV/T, $\hat{\beta} \approx 0.8$ T/m), $\sigma_k = 0.5$. Initial distribution [light blue, (a)] parameters: $\mu_z = 20.0$, $\mu_v = 0.0$, $\sigma_z = 4$, $\sigma_v = 1/8$. (a) In blue - filamented phase space before the kick. (b) Shortly after the kick. (c) Close to echo event, at $t \approx 2t_k$. Arrows point at the tips (see the text). (d) Average position.

energies E_i of the QB of mass m are found from

$$H_g \psi_i = -\frac{\hbar^2}{2m} \frac{\partial^2 \psi_i}{\partial z^2} + mgz \psi_i = E_i \psi_i, \quad (1)$$

where g is the gravitational acceleration, and z is the vertical position. Inertial and gravitational masses are taken as equal. The perfect reflection off the surface is accounted for by the boundary condition $\psi_i(z = 0) = 0$. The second boundary condition is $\psi_i(z \rightarrow \infty) = 0$. Position, time, and energy are measured in units of [4]: $z_g = (\hbar^2/2m^2g)^{1/3}$, $t_g = \hbar/E_g$, and $E_g = mgz_g$ (e.g. for neutron: $z_g = 5.87 \mu\text{m}$, $t_g = 1.094$ ms, $E_g = 0.60$ peV). The solutions of Eq. (1) are shifted Airy functions [4]

$$\psi_i(z) = N_i \text{Ai}(z - z_i) = \frac{\text{Ai}(z - z_i)}{|\text{Ai}'(-z_i)|}, \quad (2)$$

where $-z_i$ are the zeroes of $\text{Ai}(z)$, and $N_i = |\text{Ai}'(-z_i)|^{-1}$ are the normalization constants [65]. The (positive) energies are $E_i = z_i$ [4].

Echo in a classical ensemble of gravitational bouncers—It is instructive to start from considering the dynamics of $N \gg 1$ classical bouncing particles subject to a pair of delayed pulsed excitations (“kicks”). The first kick initiates nonequilibrium dynamics in the phase space. Here, for clarity of presentation, we model the resulting phase space distribution by a displaced Gaussian with means $\mu_{z,v}$ and variances $\sigma_{z,v}$, for vertical position and velocity [see the bright blue spot in Fig. 1(a)].

Due to the energy dependence of the bouncing frequency, the initial smooth phase space distribution evolves into a spiral-like structure [see the blue filaments

in Fig. 1(a)]. The number of spiral turns increases with time, and they become thinner to conserve the phase-space volume. Such “filamentation” is characteristic of non-linear systems [51, 66, 67]. The spiral in the phase space exhibits itself via multiple sharp peaks (“density waves” [68]) in the QB’s spatial distribution.

The filamented phase space serves as a basis for the echo formation induced by the second kick applied at $t = t_k$. Depending on QB type and specific experimental implementation, various kicking mechanisms can be utilized. As a first example here, we consider particles with nonzero magnetic moment μ , and kick them using pulsed inhomogeneous magnetic field, $\mathbf{B}$. For simplicity, we assume $\mathbf{B}$ has a uniform gradient near the surface [31, 32], and fix μ along/against $\mathbf{B}$. Then, the dimensionless interaction potential is $V_B(z, t) = -s\beta(t)z$ ($s = \pm 1$), $\beta(t) = a_k \exp[-(t - t_k)^2/\sigma_k^2]$, $a_k = |\mu|\hat{\beta}/(mg)$, and $\hat{\beta}$ is the magnitude of the gradient. Figure 1(b) shows the phase space distribution shortly after the kick, leading to particles bunching, and formation of localized tips on each branch of the spiral. The filamented structure provides a quasi-discrete set of oscillation frequencies for the tips [55, 56], which continue evolving freely and, with time, get out of phase. However, due to their quasi-discrete frequencies, the tips synchronize at twice the delay, at $t \approx 2t_k$ [see Fig. 1(c)], resulting in the echo response [48, 51, 55, 56]. Echo manifests in various physical observables. Here, we choose to focus on the average position $\langle z \rangle(t)$ (also averaged over $s = \pm 1$). Figure 1(d) clearly shows the echo response emerging at twice the kick delay, at $t \approx 2t_k$. Although the tips fade with time, they synchronize quasi-periodically producing higher order echoes [51, 54, 55] visible at $3t_k, 4t_k \dots$

Gravitational wave packet echo—Initially, the QB is assumed to be in a pure quantum state, e.g. a wave packet of GQSSs. Pure GQS has not been selected experimentally yet, due to tunneling of particles through a gravitational barrier [10], but we count on the major reduction of contamination of neighboring GQSSs in the future [69]. The QB may be set into motion either by kicking it, or by dropping it on the surface from a step [70]/ion trap [24]. We start from the latter, and model the initial state by a displaced Gaussian

$$\Psi(z, t = 0) = \left(\frac{2}{\pi \sigma_z^2} \right)^{1/4} \exp \left[-\frac{(z - \mu_z)^2}{\sigma_z^2} \right]. \quad (3)$$

This state is similar to the initial phase space distribution used in the classical analysis. In the quantum case, the observable is the expectation value, $\langle z \rangle(t) = \int_0^\infty \Psi^*(z, t) z \Psi(z, t) dz$. In principle, the echo effect can be observed in a variety of experimentally accessible observables, e.g. a flux through the surface [71].

Figure 2 shows that after several bounces, the wave packet collapses [$\langle z \rangle(t)$ oscillations decay] because of the




Figure 2. Echo induced by pulsed inhomogeneous magnetic field kick. The kick is applied at $t = t_k = 60$. Initial state parameters: $\mu_z = 20$, $\sigma_z = 8$ [see Eq. (3)]. Excitation parameters: $a_k = \mu\hat{\beta}/mg = 0.5$ (e.g., for neutrons: $\hat{\beta} \approx 0.8$ T/m), and $\sigma_k = 0.5$. The classical result [see Fig. 1(d)] is added for comparison.

differences in the transition frequencies of GQSs forming the wave packet (a direct consequence of the anharmonicity of the potential). The echo is induced by a kick applied after a delay t_k . Following the example considered classically, we assume that the QB (an atom, anti-atom, or neutron) has spin 1/2, and kick it by a pulsed inhomogeneous magnetic field, $\mathbf{B}$. The Hamiltonian is $H = H_g - s\beta(t)z$, where H_g is defined in Eq. (1), and $s = \pm 1$ corresponds to the spin states oriented along/against the field (see for details the Supplemental Material). The echo response is clearly visible at twice the kick delay, at $t \approx 2t_k$. The result is the average of $\langle z \rangle(t)$ obtained for $s = \pm 1$.

GQSs echo is conceptually different from classical echo emerging in ensemble of many non-identical bouncers. The former can be observed in single bouncers by repeating the experiment many times starting from the same initial state. The interference pattern developing after many measurements is a time-domain analogue of the spatial interference fringes formed in the double slit experiment with single electrons (the famous Feynman gedanken experiment, see [72] and references therein). Related echoes have been observed in single atoms interacting with a single mode of cavity [62, 63], and in single vibrationally excited molecules [64]. The GQSs echo also differs from quantum revivals, which happen in wave packets containing many states without additional kicks. The periodicity of revivals depends only on the energy spectrum [73–76], while the echo period is controlled by the kick delay.

Time-resolved GQSs spectroscopy—An appropriately chosen observable measured as a function of the kick delay contains spectroscopic information about the QB. Here, for example, we choose to follow the population of the ground GQS. The suggested measurement can be realized in the typical flow-through configuration (see Fig. 3) [8, 27, 33, 77], or using closed traps for QBs [14, 77]. The experiment includes three stages: preparation, interaction, and detection. Initially, particles pass through a narrow slit [(a) in Fig. 3], whose top surface




Figure 3. Schematic of a flow-through experimental setup. Bouncing (along Z axis) particles propagate in X direction. (a) First slit with a rough top surface used for preparation. (b) Interaction region. (c) Second slit used for detection.

is rough leading to the loss of highly excited particles. A sufficiently long and properly sized slit allows preparing ground GQS, ψ_1 [78, 79]. Then, the QB enters the interaction region [(b) in Fig. 3] where it is subject to two kicks. In the detection stage, the particles pass through the second slit [(c) in Fig. 3] allowing only the population trapped in the ground state to reach the detector (not shown). The delay, τ between the kicks is varied and the population of the ground state is recorded as a function of τ .

For impulsive (and identical) kicks, the Hamiltonian during the excitations is $H \approx V(z)f(t)$. The wave function after the first kick is given by $\Psi_+ = \mathbf{P}\Psi_-$, where Ψ_- is the wave function before the kick, $\mathbf{P} = \exp[-i\alpha V(z)]$, and $\alpha = \int_{-\infty}^{\infty} f(t)dt$. For the initial ground GQS, ψ_1 , $\Psi_+ = \sum_{i=1}^{\infty} P_{i1}\psi_i$, where P_{ij} is the matrix representation of $\mathbf{P}$ in the basis of ψ_i s. After a delay τ (just before the second excitation), the wave function is $\Psi_-(\tau) = \sum_{i=1}^{\infty} P_{i1}\psi_i e^{-iz_i\tau}$. The delay-dependent amplitude of the ground state after the second kick is given by $c_1(\tau) = \sum_{i=1}^{\infty} P_{1i}^2 \exp[-iz_i\tau]$, while the population reads $|c_1|^2(\tau) = \sum_{i,j=1}^{\infty} (P_{1i}P_{1j}^*)^2 \exp[-i(z_i - z_j)\tau]$. This signal oscillates at transition frequencies between the GQSs populated by the first kick. The second kick affects the amplitudes of the GQSs, but not their transition frequencies. In the limit of weak kicks (keeping only terms with $i = 1, j \geq 1$ and $i \geq 1, j = 1$) $|c_1|^2(\tau)$ reads

$$|c_1|^2(\tau) \approx \sum_{i=1}^{\infty} (P_{11}^* P_{1i})^2 e^{-i(z_i - z_1)\tau} + \text{c.c.}, \quad (4)$$

where “c.c.” stands for complex conjugate. The function in Eq. (4) contains the transition frequencies between the excited states ψ_i and the ground state ψ_1 . Notice that the signal contains *phase information* allowing, in principle, to retrieve the complex-valued wave function expansion coefficients P_{1i} (up to a π phase). This is analogous to the “quantum holography” procedure [80–83]. The access to phase information may open new possibilities for constraining the parameters of extra interactions [84–87].

Figure 4(a) shows the numerically calculated $|c_1|^2(\tau)$ for the case of two delayed kicks by pulsed inhomogeneous magnetic field. Here, the time dependence of the field is defined by




Figure 4. Time-resolved QGS spectroscopy: kicks by pulsed inhomogeneous magnetic field. (a) $|c_1|^2(\tau)$, kicks' parameters: $a_{k1} = 2$, $a_{k2} = 1$ (e.g. for neutron: $|\mu| = 60.3$ neV/T, $\hat{\beta}_1 \approx 2.4$ T/m, $\hat{\beta}_2 \approx 1.2$ T/m), $\sigma_{k1} = \sigma_{k2} = 0.2$. (b) Spectrum of $|c_1|^2(\tau)$. Peaks correspond to energy differences $\mathcal{E}_{i1}$ ($i = 2, \dots, 6$). Theoretical energy differences [see Eq. (2)]: $z_{21} = 1.750$, $z_{31} = 3.182$, $z_{41} = 4.449$, $z_{51} = 5.606$, $z_{61} = 6.684$.

$\beta(t) = a_{k1} \exp[-t^2/\sigma_{k1}^2] + a_{k2} \exp[-(t - \tau)^2/\sigma_{k2}^2]$. The maximal delay is close to the typical time-of-flight through the interaction region [(b) in Fig. 3] in experiments with UCNs (see Ref. [31, 32] for details). Figure 4(b) shows the spectrum of the signal in Fig. 4(a), which contains mainly the energy differences $\mathcal{E}_{ij} = \mathcal{E}_i - \mathcal{E}_j$ between the low-lying excited states ψ_i ($i = 2, \dots, 6$) and the ground state ψ_1 [see Eq. (4)]. The relative errors defined by $100\% \times (\mathcal{E}_{i1} - z_{i1})/z_{i1}$, where $z_{i1} = z_i - z_1$ [see Eq. (2)], are 0.52%, -1.27%, 1.69%, 0.87%, 1.52% for $i = 2, \dots, 6$, typical for flow-through experiments. The precision is determined by the maximal delay, and may be increased in closed traps [14].

Kick by a jolt from the surface—Both the wave packet echoes and the QGSs spectroscopy approach discussed above are general and do not depend on the specific type of kicks, as long as they are short. Here we consider an additional kind of kicks caused by a sudden displacement of the reflecting boundary. The corresponding Schrödinger equation has a time-dependent boundary condition $\Psi[z = h(t)] = 0$, where $h(t)$ is the surface height (see for details the Supplemental Material). Such a model can, in principle, describe several experimental scenarios in which the kicks are induced by shaking the surface as a whole, or by existence of protrusions, grooves, or steps on the surface. Such inhomogeneities appear as a time-dependent boundary in the reference frame co-propagating transversally with the QB moving along the surface. Here, $h(t) = a_{k1} \exp[-t^2/\sigma_{k1}^2] + a_{k2} \exp[-(t - \tau)^2/\sigma_{k2}^2]$, where a_{k1}, a_{k2} are the amplitudes of the kicks, and σ_{k1}, σ_{k2} define their widths.




Figure 5. Echo induced by surface shake. The first kick is applied at $t = 0$, the delay of the second kick is $t_k = 150$. The echo emerges at $t \approx 2t_k, 3t_k$. Kicks' parameters: $a_{k1} = 1.5$, $\sigma_{k1} = 1.0$, $a_{k2} = 0.10$, $\sigma_{k2} = 0.16$.

Figure 5 shows the echo response of $\langle z \rangle(t)$. Here, the QB is initially in the ground state ψ_1 . A single kick at $t = 0$ excites a wave packet which collapses after several oscillations (dashed blue). However, when a second kick is applied at $t = t_k$, echo responses emerge at $t \approx 2t_k, 3t_k$ (solid orange).

Figure 6(a) shows $|c_1|^2(\tau)$ in this case, while the corresponding spectrum is shown in Fig. 6(b). The maximal delay is close to the typical time-of-flight through the interaction region in experiments with UCNs [8, 27, 33, 77]. The relative errors of the extracted energy differences are 0.52%, -1.27%, 0.28%, 0.87%, -0.37% for $i = 2, \dots, 6$. In the limit of weak kicks ($a_{k1}, a_{k2} \ll 1$), $|c_1|^2(\tau)$ can be obtained using time-dependent perturbation theory [see Eq. (9) in the Supplemental Material]. In agreement with Eq. (4), the signal contains the transition frequencies between the excited states ψ_i and the ground state ψ_1 , and the Fourier amplitudes are proportional to the squared expansion coefficients of the wave packet after the first excitation.

Conclusions.—Echo effect in impulsively excited QBs is considered and the formation mechanism is discussed using the auxiliary classical model. Echoes may be used for probing decoherence effects originating from interactions with the environment or other particles. The population of the ground state recorded as a function of the delay is shown to contain the transition frequencies between QGSs excited by the first kick, populations, and partial phases information. The retrieved phases may open opportunities for constraining the parameters of extra fundamental interactions [84–87]. Various initial states, detection schemes, probe particles, and kicking mechanisms can be envisioned for both inducing the echo effect and QGSs spectroscopy. This method can be used by the current collaborations working with QGSs of UCNs (Tokyo, qBounce, Los Alamos, GRANIT),




Figure 6. Time-resolved GQS spectroscopy: kicks by surface shake. (a) $|c_1|^2(\tau)$, kicks' parameters: $a_{k1} = 0.6z_g$, $a_{k2} = 0.1z_g$, and $\sigma_{k1} = \sigma_{k2} = 0.2t_g$. (b) Spectrum of $|c_1|^2(\tau)$ shown in panel (a). Peaks correspond to energy differences $\mathcal{E}_{i1}$ ($i = 2, \dots, 6$). Theoretical differences [see Eq. (2)]: $z_{21} = 1.750$, $z_{31} = 3.182$, $z_{41} = 4.449$, $z_{51} = 5.606$, $z_{61} = 6.684$, where $z_{i1} = z_i - z_1$.

with $\bar{H}$ (GBAR), with hydrogen atoms (GRASIAN), with whispering gallery states of neutrons, atoms, and anti-atoms.

This work was supported in part by the Israel Science Foundation (Grant No. 746/15). I.A. acknowledges support as the Patricia Elman Bildner Professorial Chair. This research was made possible in part by the historic generosity of the Harold Perlman Family.

* dr.a.voronin@gmail.com

† nesvizhevsky@ill.eu

‡ ilya.averbukh@weizmann.ac.il

- [1] I. I. Gol'dman, V. D. Krivchenkov, V. I. Kogan, and V. M. Galitskii, *Selected Problems in Quantum Mechanics*, 2nd ed., edited by D. ter Haar (Infosearch Ltd, London, 1964).
- [2] P. W. Langhoff, "Schrödinger Particle in a Gravitational Well," *Am. J. Phys.* **39**, 954 (1971).
- [3] R. L. Gibbs, "The quantum bouncer," *Am. J. Phys.* **43**, 25 (1975).
- [4] J. Gea-Banacloche, "A quantum bouncing ball," *Am. J. Phys.* **67**, 776 (1999).
- [5] H. Wallis, J. Dalibard, and C. Cohen-Tannoudji, "Trapping atoms in a gravitational cavity," *Appl. Phys. B* **54**, 407 (1992).
- [6] M. F. Andersen, A. Kaplan, N. Friedman, and N. Davidson, "Stable islands in chaotic atom-optics billiards, caused by curved trajectories," *J. Phys. B* **35**, 2183 (2002).
- [7] V. V. Nesvizhevsky and A. Voronin, *Surprising Quantum Bounces* (Imperial College Press, 2015).
- [8] V. V. Nesvizhevsky, H. G. Börner, A. K. Petukhov, H. Abele, S. Baeßler, F. J. Rueß, T. Stöferle, A. Westphal, A. M. Gagarski, G. A. Petrov, and A. V. Strelkov,

- "Quantum states of neutrons in the Earth's gravitational field," *Nature* **415**, 297 (2002).
- [9] V. V. Nesvizhevsky, H. G. Börner, A. M. Gagarski, A. K. Petukhov, G. A. Petrov, H. Abele, S. Baeßler, G. Divkovic, F. J. Rueß, T. Stöferle, A. Westphal, A. V. Strelkov, K. V. Protasov, and A. Yu. Voronin, "Measurement of quantum states of neutrons in the Earth's gravitational field," *Phys. Rev. D* **67**, 102002 (2003).
- [10] V. V. Nesvizhevsky, A. K. Petukhov, H. G. Börner, T. A. Baranova, A. M. Gagarski, G. A. Petrov, K. V. Protasov, A. Yu. Voronin, S. Baeßler, H. Abele, A. Westphal, and L. Lucovac, "Study of the neutron quantum states in the gravity field," *Eur. Phys. J. C* **40**, 479 (2005).
- [11] A. Westphal, H. Abele, S. Baeßler, V. V. Nesvizhevsky, K. V. Protasov, and A. Yu. Voronin, "A quantum mechanical description of the experiment on the observation of gravitationally bound states," *Eur. Phys. J. C* **51**, 367 (2007).
- [12] V. V. Nesvizhevsky, A. K. Petukhov, K. V. Protasov, and A. Yu. Voronin, "Centrifugal quantum states of neutrons," *Phys. Rev. A* **78**, 033616 (2008).
- [13] V. V. Nesvizhevsky, A. Yu. Voronin, R. Cubitt, and K. V. Protasov, "Neutron whispering gallery," *Nat. Phys.* **6**, 114 (2010).
- [14] V. V. Nesvizhevsky, F. Nez, S. A. Vasiliev, E. Widmann, P. Crivelli, S. Reynaud, and A. Yu. Voronin, "A magneto-gravitational trap for studies of gravitational quantum states," *Eur. Phys. J. C* **80**, 520 (2020).
- [15] A. Yu. Voronin, P. Froelich, and V. V. Nesvizhevsky, "Gravitational quantum states of Antihydrogen," *Phys. Rev. A* **83**, 032903 (2011).
- [16] P.-P. Crépin, *Quantum reflection of a cold antihydrogen wave packet*, *Theses*, Sorbonne Universités, UPMC University of Paris 6 (2019).
- [17] E. G. Adelberger, B. R. Heckel, C. W. Stubbs, and Y. Su, "Does antimatter fall with the same acceleration as ordinary matter?" *Phys. Rev. Lett.* **66**, 850 (1991).
- [18] T. W. Darling, F. Rossi, G. I. Opat, and G. F. Moorhead, "The fall of charged particles under gravity: A study of experimental problems," *Rev. Mod. Phys.* **64**, 237 (1992).
- [19] F. M. Huber, R. A. Lewis, E. W. Messerschmid, and G. A. Smith, "Precision tests of Einstein's Weak Equivalence Principle for antimatter," *Adv. Space Res.* **25**, 1245 (2000).
- [20] C. Amole, M. D. Ashkezari, M. Baquero-Ruiz, W. Bertsche, E. Butler, A. Capra, C. L. Cesar, M. Charlton, S. Eriksson, J. Fajans, T. Friesen, M. C. Fujiwara, D. R. Gill, A. Gutierrez, J. S. Hangst, W. N. Hardy, M. E. Hayden, C. A. Isaac, S. Jonsell, L. Kurchaninov, A. Little, N. Madsen, J. T. K. McKenna, S. Menary, S. C. Napoli, P. Nolan, A. Olin, P. Pusa, C. Ø. Rasmussen, F. Robicheaux, E. Sarid, D. M. Silveira, C. So, R. I. Thompson, D. P. van der Werf, J. S. Wurtele, A. I. Zhmoginov, A. E. Charman, and The A. L. P. H. A Collaboration, "Description and first application of a new technique to measure the gravitational mass of antihydrogen," *Nat. Commun.* **4**, 1785 (2013).
- [21] A. Kellerbauer, M. Amoretti, A. S. Belov, G. Bonomi, I. Boscolo, R. S. Brusa, M. Büchner, V. M. Byakov, L. Cabaret, C. Canali, C. Carraro, F. Castelli, S. Cialdi, M. de Combarieu, D. Comparat, G. Consolati, N. Djourellov, M. Doser, G. Drobychev, A. Dupasquier, G. Ferrari, P. Forget, L. Formaro, A. Gervasini, M. G. Giammarchi, S. N. Gninenko, G. Gribakin, S. D. Hogan,

- M. Jacquey, V. Lagomarsino, G. Manuzio, S. Mariazzi, V. A. Matveev, J. O. Meier, F. Merkt, P. Nedelec, M. K. Oberthaler, P. Pari, M. Prevedelli, F. Quasso, A. Rondini, D. Sillou, S. V. Stepanov, H. H. Stroke, G. Testera, G. M. Tino, G. Tréneç, A. Vairo, J. Vigué, H. Walters, U. Warring, S. Zavatarelli, and D. S. Zvezhinskij, “Proposed antimatter gravity measurement with an antihydrogen beam,” *Nucl. Instrum. Methods Phys. Res. B* **266**, 351 (2008).
- [22] P. Indelicato, G. Chardin, P. Grandemange, D. Lunney, V. Manea, A. Badertscher, P. Crivelli, A. Curioni, A. Marchionni, B. Rossi, A. Rubbia, V. Nesvizhevsky, D. Brook-Roberge, P. Comini, P. Debu, P. Dupré, L. Liskay, B. Mansoulié, P. Pérez, J.-M. Rey, B. Reymond, N. Ruiz, Y. Sacquin, B. Vallage, F. Biraben, P. Cladé, A. Douillet, G. Dufour, S. Guellati, L. Hilico, A. Lambrecht, R. Guérout, J.-P. Karr, F. Nez, S. Reynaud, C. I. Szabo, V.-Q. Tran, J. Trapateau, A. Mohri, Y. Yamazaki, M. Charlton, S. Eriksson, N. Madsen, D. P. van der Werf, N. Kuroda, H. Torii, Y. Nagashima, F. Schmidt-Kaler, J. Walz, S. Wolf, P.-A. Hervieux, G. Manfredi, A. Voronin, P. Froelich, S. Wronka, and M. Staszczak, “The GBAR project, or how does antimatter fall?” *Hyperfine Interact.* **228**, 141 (2014).
- [23] W. A. Bertsch, “Prospects for comparison of matter and antimatter gravitation with ALPHA-g,” *Philos. Trans. R. Soc. A* **376**, 20170265 (2018).
- [24] P.-P. Crépin, C. Christen, R. Guérout, V. V. Nesvizhevsky, A. Yu. Voronin, and S. Reynaud, “Quantum interference test of the equivalence principle on antihydrogen,” *Phys. Rev. A* **99**, 042119 (2019).
- [25] V. V. Nesvizhevsky and K. V. Protasov, “Trends in Quantum Gravity Research,” (Nova Science Publications, New York, 2006) Chap. Quantum states of neutrons in the Earth’s gravitational field: state of the art, applications, perspectives, pp. 65–107.
- [26] H. Abele, T. Jenke, H. Leeb, and J. Schmiedmayer, “Ramsey’s method of separated oscillating fields and its application to gravitationally induced quantum phase shifts,” *Phys. Rev. D* **81**, 065019 (2010).
- [27] T. Jenke, P. Geltenbort, H. Lemmel, and H. Abele, “Realization of a gravity-resonance-spectroscopy technique,” *Nat. Phys.* **7**, 468 (2011).
- [28] T. Jenke, G. Cronenberg, J. Burgdörfer, L. A. Chizhova, P. Geltenbort, A. N. Ivanov, T. Lauer, T. Lins, S. Rotter, H. Saul, U. Schmidt, and H. Abele, “Gravity Resonance Spectroscopy Constrains Dark Energy and Dark Matter Scenarios,” *Phys. Rev. Lett.* **112**, 151105 (2014).
- [29] G. Cronenberg, P. Brax, H. Filter, P. Geltenbort, T. Jenke, G. Pignol, M. Pitschmann, M. Thalhammer, and H. Abele, “Acoustic Rabi oscillations between gravitational quantum states and impact on symmetron dark energy,” *Nat. Phys.* **14**, 1022 (2018).
- [30] G. Manfredi, O. Morandi, L. Friedland, T. Jenke, and H. Abele, “Chirped-frequency excitation of gravitationally bound ultracold neutrons,” *Phys. Rev. D* **95**, 025016 (2017).
- [31] G. Pignol, S. Baeßler, V. V. Nesvizhevsky, K. Protasov, D. Rebreyend, and A. Voronin, “Gravitational Resonance Spectroscopy with an Oscillating Magnetic Field Gradient in the GRANIT Flow through Arrangement,” *Adv. High Energy Phys.* **2014**, 628125 (2014).
- [32] S. Baeßler, V. V. Nesvizhevsky, G. Pignol, K. V. Protasov, D. Rebreyend, E. A. Kupriyanova, and A. Yu. Voronin, “Frequency shifts in gravitational resonance spectroscopy,” *Phys. Rev. D* **91**, 042006 (2015).
- [33] G. Ichikawa, S. Komamiya, Y. Kamiya, Y. Minami, M. Tani, P. Geltenbort, K. Yamamura, M. Nagano, T. Sanuki, S. Kawasaki, M. Hino, and M. Kitaguchi, “Observation of the Spatial Distribution of Gravitationally Bound Quantum States of Ultracold Neutrons and Its Derivation Using the Wigner Function,” *Phys. Rev. Lett.* **112**, 071101 (2014).
- [34] A. Yu. Voronin, V. V. Nesvizhevsky, O. D. Dalkarov, E. A. Kupriyanova, and P. Froelich, “Resonance spectroscopy of gravitational states of antihydrogen,” *Hyperfine Interact.* **228**, 133 (2014).
- [35] P.-P. Crépin, G. Dufour, R. Guérout, A. Lambrecht, and S. Reynaud, “Casimir-Polder shifts on quantum levitation states,” *Phys. Rev. A* **95**, 032501 (2017).
- [36] A. Yu. Voronin, V. V. Nesvizhevsky, G. Dufour, and S. Reynaud, “Quantum ballistic experiment on antihydrogen fall,” *J. Phys. B* **49**, 054001 (2016).
- [37] V. V. Nesvizhevsky, A. Yu. Voronin, P.-P. Crépin, and S. Reynaud, “Interference of several gravitational quantum states of antihydrogen in GBAR experiment,” *Hyperfine Interact.* **240**, 32 (2019).
- [38] E. L. Hahn, “Spin Echoes,” *Phys. Rev.* **80**, 580 (1950).
- [39] E. L. Hahn, “Free nuclear induction,” *Phys. Today* **6**, 4 (1953).
- [40] N. A. Kurnit, I. D. Abella, and S. R. Hartmann, “Observation of a Photon Echo,” *Phys. Rev. Lett.* **13**, 567 (1964).
- [41] S. Mukamel, *Principles of Nonlinear Optical Spectroscopy* (Oxford Univ. Press, 1995).
- [42] R. M. Hill and D. E. Kaplan, “Cyclotron Resonance Echo,” *Phys. Rev. Lett.* **14**, 1062 (1965).
- [43] R. W. Gould, T. M. O’Neil, and J. H. Malmberg, “Plasma Wave Echo,” *Phys. Rev. Lett.* **19**, 219 (1967).
- [44] F. Mezei, “Neutron spin echo: A new concept in polarized thermal neutron techniques,” *Z. Phys. A* **255**, 146 (1972).
- [45] A. Bulatov, A. Kuklov, B. E. Vugmeister, and H. Rabitz, “Echo in optical lattices: Stimulated revival of breathing oscillations,” *Phys. Rev. A* **57**, 3788 (1998).
- [46] F. B. J. Buchkremer, R. Dumke, H. Levsen, G. Birkel, and W. Ertmer, “Wave Packet Echoes in the Motion of Trapped Atoms,” *Phys. Rev. Lett.* **85**, 3121 (2000).
- [47] M. Herrera, T. M. Antonsen, E. Ott, and S. Fishman, “Echoes and revival echoes in systems of anharmonically confined atoms,” *Phys. Rev. A* **86**, 023613 (2012).
- [48] G. Stupakov, *Echo Effect in Hadron Colliders*, Tech Report 579 (SSC Report SSCL-579, SSCL, 1992).
- [49] G. V. Stupakov and S. K. Kauffmann, “Echo effect in accelerators,” *Proc. Par. Accel. Conf.*, **1**, 197 (1993).
- [50] L. K. Spentzouris, J.-F. Ostiguy, and P. L. Colestock, “Direct Measurement of Diffusion Rates in High Energy Synchrotrons Using Longitudinal Beam Echoes,” *Phys. Rev. Lett.* **76**, 620 (1996).
- [51] G. Stupakov, “Handbook of Accelerator Physics and Engineering,” (World Scientific, 2013) Chap. 2.3.13, pp. 121–123, 2nd ed.
- [52] T. Sen and Y. S. Li, “Nonlinear theory of transverse beam echoes,” *Phys. Rev. Accel. Beams* **21**, 021002 (2018).
- [53] G. Karras, E. Hertz, F. Billard, B. Lavorel, J.-M. Hartmann, O. Faucher, E. Gershnabel, and Y. and I. Sh. Averbukh, “Orientation and Alignment Echoes,” *Phys. Rev. Lett.* **114**, 153601 (2015).
- [54] G. Karras, E. Hertz, F. Billard, B. Lavorel, G. Siour, J.-

- M. Hartmann, O. Faucher, E. Gershnabel, and Y. and I. Sh. Averbukh, “*Experimental observation of fractional echoes*,” *Phys. Rev. A* **94**, 033404 (2016).
- [55] K. Lin, P. Lu, J. Ma, X. Gong, Q. Song, Q. Ji, W. Zhang, H. Zeng, J. Wu, G. Karras, G. Siour, J.-M. Hartmann, O. Faucher, E. Gershnabel, Y. Prior, and I. Sh. Averbukh, “*Echoes in Space and Time*,” *Phys. Rev. X* **6**, 041056 (2016).
- [56] K. Lin, I. Tutunnikov, J. Ma, J. Qiang, L. Zhou, O. Faucher, Y. Prior, I. Sh. Averbukh, and J. Wu, “*Spatiotemporal rotational dynamics of laser-driven molecules*,” *Adv. Photon.* **2**, 1 (2020).
- [57] H. Zhang, B. Lavorel, F. Billard, J.-M. Hartmann, E. Hertz, O. Faucher, J. Ma, J. Wu, E. Gershnabel, Y. Prior, and I. Sh. Averbukh, “*Rotational Echoes as a Tool for Investigating Ultrafast Collisional Dynamics of Molecules*,” *Phys. Rev. Lett.* **122**, 193401 (2019).
- [58] J. Ma, H. Zhang, B. Lavorel, F. Billard, E. Hertz, J. Wu, C. Boulet, J.-M. Hartmann, and O. Faucher, “*Observing collisions beyond the secular approximation limit*,” *Nat. Comm.* **10**, 5780 (2019).
- [59] D. Rosenberg, R. Damari, and S. Fleischer, “*Echo Spectroscopy in Multilevel Quantum-Mechanical Rotors*,” *Phys. Rev. Lett.* **121**, 234101 (2018).
- [60] D. Rosenberg and S. Fleischer, “*Intrinsic calibration of laser-induced molecular alignment using rotational echoes*,” *Phys. Rev. Research* **2**, 023351 (2020).
- [61] J.-M. Hartmann, J. Ma, T. Delahaye, F. Billard, E. Hertz, J. Wu, B. Lavorel, C. Boulet, and O. Faucher, “*Molecular alignment echoes probing collision-induced rotational-speed changes*,” *Phys. Rev. Research* **2**, 023247 (2020).
- [62] G. Morigi, E. Solano, B.-G. Englert, and H. Walther, “*Measuring irreversible dynamics of a quantum harmonic oscillator*,” *Phys. Rev. A* **65**, 040102 (2002).
- [63] T. Meunier, S. Gleyzes, P. Maioli, A. Auffeves, G. Nogues, M. Brune, J. Raimond, and S. Haroche, “*Rabi Oscillations Revival Induced by Time Reversal: A Test of Mesoscopic Quantum Coherence*,” *Phys. Rev. Lett.* **94**, 010401 (2005).
- [64] J. Qiang, I. Tutunnikov, P. Lu, K. Lin, W. Zhang, F. Sun, Y. Silberberg, Y. Prior, I. Sh. Averbukh, and J. Wu, “*Echo in a single vibrationally excited molecule*,” *Nat. Phys.* **16**, 328 (2020).
- [65] J. R. Albright, “*Integrals of products of Airy functions*,” *J. Phys. A* **10**, 485 (1977).
- [66] D. Lynden-Bell, “*Statistical Mechanics of Violent Relaxation in Stellar Systems*,” *Mon. Not. R. Astron. Soc.* **136**, 101 (1967).
- [67] G. Guignard, in *Frontiers of Particle Beams*, edited by M. Month and S. Turner (Springer Berlin Heidelberg, Berlin, Heidelberg, 1988) pp. 1–50.
- [68] E. J. Kolmes, V. I. Geyko, and N. J. Fisch, “*Density waves in a system of non-interacting particles*,” *Phys. Lett. A* **380**, 3061 (2016).
- [69] M. Escobar, F. Lamy, A. E. Meyerovich, and V. V. Nesvizhevsky, “*Rough Mirror as a Quantum State Selector: Analysis and Design*,” *Adv. High Energy Phys.* **2014**, 764182 (2014).
- [70] V. V. Nesvizhevskii, “*Investigation of quantum neutron states in the terrestrial gravitational field above a mirror*,” *Physics-Uspekhi* **47**, 515 (2004).
- [71] V. V. Nesvizhevsky, R. Cubitt, K. V. Protasov, and A. Yu. Voronin, “*The whispering gallery effect in neutron scattering*,” *New J. Phys.* **12**, 113050 (2010).
- [72] R. Bach, D. Pope, S.-H. Liou, and H. Batelaan, “*Controlled double-slit electron diffraction*,” *New J. Phys.* **15** (2013).
- [73] J. H. Eberly, N. B. Narozhny, and J. J. Sanchez-Mondragon, “*Periodic Spontaneous Collapse and Revival in a Simple Quantum Model*,” *Phys. Rev. Lett.* **44**, 1323 (1980).
- [74] J. Parker and C. R. Stroud, “*Coherence and decay of Rydberg wave Packets*,” *Phys. Rev. Lett.* **56**, 716 (1986).
- [75] I. Sh. Averbukh and N. F. Perelman, “*Fractional revivals: Universality in the long-term evolution of quantum wave packets beyond the correspondence principle dynamics*,” *Phys. Lett. A* **139**, 449 (1989).
- [76] R. Robinett, “*Quantum wave packet revivals*,” *Physics Reports* **392**, 1 (2004).
- [77] D. Roulier, F. Vezzu, S. Baeßler, B. Clément, D. Morton, V. V. Nesvizhevsky, G. Pignol, and D. Rebreyend, “*Status of the GRANIT Facility*,” *Adv. High Energy Phys.* **2015**, 730437 (2015).
- [78] V. I. Luschikov and A. I. Frank, “*Quantum effects occurring when ultracold neutrons are stored on a plane*,” *JETP Letters* **28**, 559 (1978).
- [79] V. V. Nesvizhevsky, H. Boerner, A. M. Gagarski, G. A. Petrov, A. K. Petukhov, H. Abele, S. Baessler, T. Stoeferle, and S. M. Soloviev, “*Search for quantum states of the neutron in a gravitational field: gravitational levels*,” *Nucl. Instrum. Methods Phys. Res.* **440**, 754 (2000).
- [80] C. Leichtle, W. P. Schleich, I. Sh. Averbukh, and M. Shapiro, “*Quantum State Holography*,” *Phys. Rev. Lett.* **80**, 1418 (1998).
- [81] I. Sh. Averbukh, M. Shapiro, C. Leichtle, and W. P. Schleich, “*Reconstructing wave packets by quantum-state holography*,” *Phys. Rev. A* **59**, 2163 (1999).
- [82] T. C. Weinacht, J. Ahn, and P. H. Bucksbaum, “*Measurement of the amplitude and phase of a sculpted Rydberg wave packet*,” *Phys. Rev. Lett.* **80**, 5508 (1998).
- [83] T. C. Weinacht, J. Ahn, and P. H. Bucksbaum, “*Controlling the shape of a quantum wavefunction*,” *Nature* **397**, 233 (1999).
- [84] H. Abele, S. Baeßler, and A. Westphal, “*Quantum States of Neutrons in the Gravitational Field and Limits for Non-Newtonian Interaction in the Range between 1 μm and 10 μm* ,” in *Quantum Gravity: From Theory to Experimental Search*, edited by D. J. W. Giulini, C. Kiefer, and C. Lämmerzahl (Springer Berlin Heidelberg, Berlin, Heidelberg, 2003) pp. 355–366.
- [85] S. Baeßler, V. V. Nesvizhevsky, K. V. Protasov, and A. Yu. Voronin, “*Constraint on the coupling of axionlike particles to matter via an ultracold neutron gravitational experiment*,” *Phys. Rev. D* **75**, 075006 (2007).
- [86] I. Antoniadis, S. Baessler, M. Büchner, V. V. Fedorov, S. Hoedl, A. Lambrecht, V. V. Nesvizhevsky, G. Pignol, K. V. Protasov, S. Reynaud, and Y. Sobolev, “*Short-range fundamental forces*,” *C. R. Phys.* **12**, 755 (2011).
- [87] P. Brax and G. Pignol, “*Strongly Coupled Chameleons and the Neutronic Quantum Bouncer*,” *Phys. Rev. Lett.* **107**, 111301 (2011).