

A note on higher order fractional Hardy-Sobolev inequalities

Roberta Musina* and Alexander I. Nazarov**

Abstract

We establish some qualitative properties of minimizers in the fractional Hardy-Sobolev inequalities of arbitrary order.

Assume $n \geq 1$ is a given integer, and take exponents $s \in (0, \frac{n}{2})$, q, β satisfying

$$2 < q < 2_s^* := \frac{2n}{n-2s}, \quad \frac{n}{q} - \beta = \frac{n}{2} - s.$$

By Hölder interpolation between the Hardy and Sobolev inequalities one obtains the existence of a positive best constant S_q such that

$$S_q \cdot \| |x|^{-\beta} u \|_q^2 \leq \| (-\Delta)^{\frac{s}{2}} u \|_2^2, \quad u \in \mathcal{D}^s(\mathbb{R}^n), \quad (1)$$

where

$$\mathcal{D}^s(\mathbb{R}^n) = \{ u \in L^{2_s^*}(\mathbb{R}^n) \mid (-\Delta)^{\frac{s}{2}} u \in L^2(\mathbb{R}^n) \}, \quad \mathcal{F}[(-\Delta)^{\frac{s}{2}} u] = |\xi|^s \mathcal{F}[u]$$

and \mathcal{F} is the Fourier transform in \mathbb{R}^n .

*Dipartimento di Scienze Matematiche, Informatiche e Fisiche, Università di Udine, via delle Scienze, 206 – 33100 Udine, Italy. Email: roberta.musina@uniud.it. Partially supported by PRID project *VARPROGE*.

**St.Petersburg Dept of Steklov Institute, Fontanka 27, St.Petersburg, 191023, Russia, and St.Petersburg State University, Universitetskii pr. 28, St.Petersburg, 198504, Russia. E-mail: al.il.nazarov@gmail.com. Partially supported by RFBR grant 20-01-00630.

The existence of a minimizer for S_q was proved in [12]. Also, it was claimed in [12, Theorem 1.2], that any minimizer of (1) has constant sign, and is radially symmetric and strictly decreasing. However, the proof of this statement contains serious gaps. Namely, author of [12] refers to the results of [11] and [7], which were formulated and proved only for $s < 1$ and which are in general false for $s > 1$, compare with [9]. Moreover, even in the case $s < 1$ the argument in [12] does not provide the *strict* monotonicity of the minimizer¹.

In this note we prove the following statement.

Theorem 1 *Let u be a minimizer for S_q . Up to a change of sign, u is everywhere positive, radially symmetric and strictly monotone decreasing with respect to $|x|$.*

Before proving Theorem 1 we recall some notation. For $f \geq 0$ measurable and vanishing at infinity, f^* stands for the symmetric-decreasing rearrangement of f , see [8, Section 3.2].

Accordingly with [4], [6], for nonnegative measurable functions f, g vanishing at infinity, we write $f \prec g$ if

$$\int_{B_r(0)} f^*(x) dx \leq \int_{B_r(0)} g^*(x) dx \quad \text{for any } r > 0.$$

Trivially, the pointwise inequality $f \leq g$ implies $f \prec g$, while the inverse implication is not true, in general. Finally, we recall the next result.

Proposition 1 (see, e.g., [6, Lemma 2.1]). *The relation $f \prec g$ is equivalent to any of the following statements:*

1. *For any non-negative convex functions ϕ such that $\phi(0) = 0$, we have*

$$\int_{\mathbb{R}^n} \phi(f^*(x)) dx \leq \int_{\mathbb{R}^n} \phi(g^*(x)) dx. \quad (2)$$

2. *For any non-negative symmetric-decreasing function φ we have*

$$\int_{\mathbb{R}^n} f^*(x) \varphi(x) dx \leq \int_{\mathbb{R}^n} g^*(x) \varphi(x) dx; \quad (3)$$

¹In fact, for $s < 1$ this can be proved by adapting the moving plane argument in [3] or [5], as pointed out in [10].

Proof of Theorem 1. We introduce the functions $U, V \in \mathcal{D}^s(\mathbb{R}^n)$ as (unique) weak solutions to

$$(-\Delta)^{\frac{s}{2}}U = f := |(-\Delta)^{\frac{s}{2}}u| \in L^2(\mathbb{R}^n), \quad (-\Delta)^{\frac{s}{2}}V = f^* \in L^2(\mathbb{R}^n), \quad (4)$$

respectively. It is well known that U and V are explicitly given by convolutions

$$U(x) = C(n, s) \int_{\mathbb{R}^n} \frac{f(y)}{|x-y|^{n-s}} dy, \quad V(x) = C(n, s) \int_{\mathbb{R}^n} \frac{f^*(y)}{|x-y|^{n-s}} dy, \quad (5)$$

where $C(n, s)$ is an explicitly known constant. In particular, $U, V > 0$.

The Riesz rearrangement inequality, see, e.g., [8, Sec. 3.7], immediately implies (3), and thus $U \prec V$, or, equivalently, $U^* \prec V^* = V$.

Further, we proceed in 3 steps.

Step 1. We claim that u strictly preserves the sign. Indeed, (4) and (5) give

$$(-\Delta)^{\frac{s}{2}}(U \pm u) = |(-\Delta)^{\frac{s}{2}}u| \pm (-\Delta)^{\frac{s}{2}}u \geq 0 \quad \Rightarrow \quad U \pm u \geq 0 \quad \Rightarrow \quad U \geq |u|.$$

Therefore,

$$\| |x|^{-\beta} U \|_q^2 \geq \| |x|^{-\beta} u \|_q^2 \quad \text{and} \quad \| (-\Delta)^{\frac{s}{2}} U \|_2^2 = \| (-\Delta)^{\frac{s}{2}} u \|_2^2.$$

Thus U achieves S_q as well. But then

$$\| |x|^{-\beta} U \|_q^2 = \| |x|^{-\beta} u \|_q^2 \quad \Rightarrow \quad |u| \equiv U > 0,$$

and the claim follows. From now on, we assume that $u > 0$.

Step 2. We claim that $u = u^*$. Indeed, by Step 1 we have $u^* = U^* \prec V$. If we assume that $u \neq u^*$, then a basic rearrangement inequality, see, e.g., [8, Sec. 3.4] together with (2), (3) gives

$$\int_{\mathbb{R}^n} |x|^{-\beta q} u^q dx < \int_{\mathbb{R}^n} |x|^{-\beta q} (u^*)^q dx \leq \int_{\mathbb{R}^n} |x|^{-\beta q} V^q dx.$$

Since evidently

$$\| (-\Delta)^{\frac{s}{2}} V \|_2 = \| ((-\Delta)^{\frac{s}{2}} u)^* \|_2 = \| (-\Delta)^{\frac{s}{2}} u \|_2,$$

we can conclude that

$$S_q = \frac{\|(-\Delta)^{\frac{s}{2}}u\|_2^2}{\||x|^{-\beta}u\|_q^2} > \frac{\|(-\Delta)^{\frac{s}{2}}V\|_2^2}{\||x|^{-\beta}V\|_q^2} \geq S_q,$$

and the claim follows via a contradiction argument.

Step 3. It remains to show that u is strictly decreasing. We recall that u is a minimizer in (1) and therefore satisfies the integral equation

$$u(x) = \int_{\mathbb{R}^n} \frac{|y|^{-\beta q} u^{q-1}(y)}{|x-y|^{n-2s}} dy.$$

up to a multiplicative constant. The right-hand side here is the convolution of two symmetric-decreasing functions, and one of them is strictly symmetric-decreasing. This completes the proof. \square

Remark 1 *In contrast to the critical case $q = 2_s^*$, the value of the infimum S_q and its minimizers are not explicitly known even for $s < 1$. The regularity, uniqueness and nondegeneracy of the minimizer in this case has been recently proved in [10], see also [2, 1]. For fractional $s > 1$, these properties have never been investigated.*

References

- [1] S. Alarcén, B. Barrios and A. Quaas, Linear non-degeneracy and uniqueness of the bubble solution for the critical fractional Hénon equation in R^N , arXiv:2009.09481 (2020), 26 pp.
- [2] B. Barrios and A. Quaas, The sharp exponent in the study of the nonlocal Hénon equation in \mathbb{R}^N : a Liouville theorem and an existence result, Calc. Var. Part. Diff. Eqs. **59** (2020), no. 4, Art. 114, 22 pp.
- [3] W. Chen, C. Li and B. Ou, Classification of solutions for an integral equation, Comm. Pure Appl. Math. **59** (2006), no. 3, 330–343.
- [4] K.M. Chong, Some extensions of a theorem of Hardy, Littlewood and Plya and their applications, Canad. J. Math., **26** (1974), 1321–1340.

- [5] S. Dipierro, L. Montoro, I. Peral and B. Sciunzi, Qualitative properties of positive solutions to nonlocal critical problems involving the Hardy-Leray potential, *Calc. Var. Part. Diff. Eqs.* **55** (2016), no. 4, Art. 99, 29 pp.
- [6] V. Ferone and B. Volzone, Symmetrization for fractional elliptic problems: a direct approach, Preprint arXiv:2007.13195 (2020), 40 pp.
- [7] R.L. Frank and R. Seiringer, Non-linear ground state representations and sharp Hardy inequalities, *J. Funct. Anal.* **255** (2008), 3407–3430.
- [8] E. Lieb and M. Loss, *Analysis*. 2nd.ed. Graduate studies in mathematics, **14**. AMS, Providence, RI, 2001.
- [9] R. Musina and A. I. Nazarov, A note on truncations in fractional Sobolev spaces, *Bull. Math. Sci.* **9** (2019), no. 1, 1950001, 7 pp.
- [10] R. Musina and A.I. Nazarov, Complete classification and nondegeneracy of minimizers for the fractional Hardy-Sobolev inequality, and applications, Preprint arXiv:2008.11186 (2020). 28 pp.
- [11] R. Servadei and E. Valdinoci, Variational methods for non-local operators of elliptic type, *Discrete Contin. Dyn. Syst.* **33** (2013), 2105–2137.
- [12] J. Yang, Fractional Sobolev-Hardy inequality in \mathbb{R}^N , *Nonlinear Anal.* **119** (2015), 179–185.