

Emergence and Breaking of Duality Symmetry in Thermodynamic Behavior: Repeated Measurements and Macroscopic Limit

Zhiyue Lu*

Department of Chemistry, University of North Carolina, Chapel Hill, NC 27599-3290, U.S.A.

Hong Qian†

Department of Applied Mathematics, University of Washington, Seattle, WA 98195-3925, U.S.A.

Thermodynamic laws are limiting behavior of the statistics of repeated measurements of an arbitrary system with *a priori* probability distribution. A duality symmetry arises, between Massieu-Guggenheim entropy and Gibbs entropy, in the limit of large number of measurements. This yields the fundamental thermodynamic relation and Hill-Gibbs-Duhem (HGD) equation as a dual pair. We show if the system itself has a second macroscopic limit that satisfies Callen's postulate that entropy being an Eulerian homogeneous function of all extensive variables, the symmetry is lost: the HGD equation reduces to the Gibbs-Duhem equation. This theory provides better logic to textbook thermodynamics, a clarification on nanothermodynamics, as well as novel ideas for a thermodynamic-like framework for single-cell biology.

Thermodynamic descriptions which are organized around the concept of *entropy* and its variants are asymptotic limiting behavior of statistics [1, 2]. The mathematical theory of large deviations in probability provides not only powerful tools for statistical mechanics [3–7], but more importantly a novel organizational principle for the theory of thermodynamics [8]. For a stochastic system with state space \mathfrak{S} and an *a priori* probability density function $f_{\mathbf{x}}(\mathbf{x})$, $\mathbf{x} \in \mathfrak{S}$, the large deviations theory (LDT) states that for a large number M repeated measurements of an array of K real observables $\mathbf{g}(\mathbf{x}) = (g_1, \dots, g_K)(\mathbf{x})$, one can construct the mean value

$$\bar{\mathbf{g}}_M = \frac{\mathbf{g}^{(1)} + \dots + \mathbf{g}^{(M)}}{M}, \quad (1)$$

in which $\mathbf{g}^{(m)}$ is the m^{th} measurement of the \mathbf{g} ; they are treated as independent and identically distributed (i.i.d.) random variables. As $M \rightarrow \infty$, one expects $\bar{\mathbf{g}}_M \rightarrow \mathbb{E}[\mathbf{g}]$, the expected value of $\mathbf{g}(\mathbf{x})$:

$$\mathbb{E}[\mathbf{g}] = \int_{\mathfrak{S}} f_{\mathbf{x}}(\mathbf{x}) \mathbf{g}(\mathbf{x}) d\mathbf{x}.$$

The LDT then predicts that the probability distribution for $\bar{\mathbf{g}}_M$ has an asymptotic expression as $M \rightarrow \infty$,

$$\ln \Pr \{ \mathbf{y} < \bar{\mathbf{g}}_M \leq \mathbf{y} + d\mathbf{y} \} = M\eta(\mathbf{y}) + o(M), \quad (2)$$

$$\eta(\mathbf{y}) = \min_{\boldsymbol{\beta}} \{ \boldsymbol{\beta} \cdot \mathbf{y} + \psi(\boldsymbol{\beta}) \}, \quad (3)$$

$$\psi(\boldsymbol{\beta}) = \ln \int_{\mathfrak{S}} f_{\mathbf{x}}(\mathbf{x}) \exp(-\boldsymbol{\beta} \cdot \mathbf{g}(\mathbf{x})) d\mathbf{x}, \quad (4)$$

in which $\mathbf{y} = (y_1, \dots, y_K)$, $\boldsymbol{\beta} = (\beta_1, \dots, \beta_K)$, and the notation $o(M)$ indicates that $o(M)/M \rightarrow 0$ as $M \rightarrow \infty$. See [9] for a highly condensed but self-contained account of the mathematics. It is known as Cramér's theorem [3, 7].

There is a unmistakable resemblance between these equations and those in statistical mechanics. We emphasize that the K -vector \mathbf{g} and $\boldsymbol{\beta}$ are treated as a single variable in the formalism. Partial transformations with a subset of the components of \mathbf{g} and $\boldsymbol{\beta}$ in (3) and (4), which generates a multitude of different ensembles in the classical theory, will be discussed elsewhere. Letting $K = 1$ and identifying $g(\mathbf{x})$ as mechanical energy, \mathbf{x} being phase space point, the integral in (4) is the canonical partition function with $\beta = (k_B T)^{-1}$. Then $k_B \psi(\beta)$ is known as Massieu-Planck free entropy, and $-\beta^{-1} \psi(\beta)$ is free energy [10]. More interestingly, noting that $\psi(\beta)$ is a convex function, the Legendre-Fenchel transformation in (3) can be carried out with differentiation:

$$\eta(y) = \left[-\beta \left(\frac{\partial \psi}{\partial \beta} \right) + \psi(\beta) \right]_{\beta: \partial \psi / \partial \beta = -y} = \frac{\partial [\beta^{-1} \psi(\beta)]}{\partial (1/\beta)}. \quad (5)$$

This shows that $k_B \eta(y)$ corresponds to the entropy of Gibbs' canonical ensemble.

If one lets $K = 3$ and identifies the $\mathbf{g} = (U, V, N)$, representing energy U , volume V and number of particles N of a mechanical system, then the integral in (4), with conjugate variables $\boldsymbol{\beta} = (k_B T)^{-1}(1, P, -\mu)$, was systematically discussed in [11].¹ The partition function $\exp(\psi(\boldsymbol{\beta}))$ is widely known in the literature as Guggenheim's generalized partition function; the corresponding thermodynamic potential is zero on the order of extensive quantities. T. L. Hill named $-k_B T \psi$ *subdivision potential* [12].

Outside mechanical systems, \mathfrak{S} could be the state space of a living cell and $\mathbf{g}(\mathbf{x})$ be a set of K biomarkers. The same

¹ We discover that the spirit of Guggenheim's setup was similar to J. G. Kirkwood's introduction of the *potential of mean force*, in *J. Chem. Phys.* **3**, 305–313, (1935).

mathematics applies [9]. Not to be confused with the concepts in statistical mechanics, in the present work, we shall fix our terminology by borrowing from the above discussion, calling ψ Massieu-Guggenheim entropy and correspondingly η Gibbs entropy.

Massieu-Guggenheim entropy and Gibbs entropy duality.

The mathematics in Eqs. 2-5 reproduces some of the key steps in statistical mechanics of canonical ensemble and ensemble change. More importantly, it also shows that $\eta(\mathbf{y})$ is itself a convex function since it is the Legendre-Fenchel transformation of the convex function $\psi(\beta)$, and (3) has an inverse relation [13]:

$$\psi(\beta) = \max_{\mathbf{y}} \left\{ -\beta \cdot \mathbf{y} + \eta(\mathbf{y}) \right\}. \quad (6)$$

Even though we first obtain $\psi(\beta)$ from $f_{\mathbf{x}}(\mathbf{x})$ and $\mathbf{g}(\mathbf{x})$ through the partition function, some detailed information concerning the original system is lost in the statistics of the mean value ². Transcending the original stochastic system, the Massieu-Guggenheim entropy $\psi(\beta)$ and Gibbs entropy $\eta(\mathbf{y})$ in (3) and (6) now form a dual: A duality symmetry emerges. This symmetry further gives two differential equations in parallel:

$$d\eta(\mathbf{y}) = \beta \cdot d\mathbf{y}, \quad (7)$$

$$d\psi(\beta) = -\mathbf{y} \cdot d\beta. \quad (8)$$

If we again identify \mathbf{y} with (U, V, N) in statistical mechanics, $k_B\eta$ with entropy S , and correspondingly $\beta = (k_B T)^{-1}(1, P, -\mu)$, we see that (7) is actually the fundamental thermodynamic relation

$$dS = T^{-1} \left(dU + PdV - \mu dN \right). \quad (9a)$$

Eq. 8 is a key result in Hills' thermodynamics of small systems [12], dubbed in recent years nanothermodynamics [14, 15], in which subdivision potential $\mathcal{E} = -k_B T \psi$:

$$\begin{aligned} d\mathcal{E} &= -k_B \psi dT + T \left[U d \left(\frac{1}{T} \right) + V d \left(\frac{P}{T} \right) - N d \left(\frac{\mu}{T} \right) \right] \\ &= \frac{(\mathcal{E} - U - PV + \mu N) dT}{T} + V dP - N d\mu \\ &= -S dT + V dP - N d\mu. \end{aligned} \quad (9b)$$

Bedeaux et. al. termed relation in (9b) Hill-Gibbs-Duhem equation [16]. We shall follow their terminology to call our Eq. 8. All the mathematical equations so far presented are

generally true for any stochastic systems. For a macroscopic system, one additional supposition for $\eta(\mathbf{y})$ being an extensive quantity will reduce the l.h.s. of Eq. 8 to sub-extensive in the thermodynamic limit.

Large system thermodynamic limit. According to classical thermodynamics, a macroscopic system in the *thermodynamic limit* is represented by a set of *extensive variables*, which we take as the vector \mathbf{g} . We adopt Callen's postulate on classical thermodynamics which states that entropy being an Eulerian first-order homogeneous function of all the extensive variables [17]. Thus for a large system approaching the macroscopic limit, we have $\mathbf{y} \rightarrow \infty$ and

$$\eta(\mathbf{y}) = y_1 \frac{\partial \eta}{\partial y_1} + \dots + y_K \frac{\partial \eta}{\partial y_K} + o(\mathbf{y}) = \mathbf{y} \cdot \beta + o(\mathbf{y}), \quad (10)$$

in which the notion $o(\mathbf{y})$ represents a sub-extensive term that scales sub-linear with respect to the system's size: $o(\mathbf{y})/\mathbf{y} \rightarrow 0$ as $\mathbf{y} \rightarrow \infty$. Substituting (10) into (6), one can show that $\psi(\beta)$ is purely sub-extensive, and thus Eq. (8) becomes $\mathbf{y} \cdot d\beta = 0$ in the (macroscopic) thermodynamic limit. Correspondingly Eq. 9b becomes:

$$-S dT + V dP - N d\mu = 0. \quad (11)$$

This is the Gibbs-Duhem equation in classical thermodynamics.

In the thermodynamic limit for extensive large systems, the duality symmetry between (7) and (8) is lost.

Legendre-Fenchel transformation of a homogeneous function. An Eulerian homogeneous function of order 1 cannot be a strictly convex function: If $\eta(\alpha \mathbf{y}) = \alpha \eta(\mathbf{y})$ where α is any real number, then $\eta(\mathbf{y}) = \mathbf{y} \cdot \nabla_{\mathbf{y}} \eta$, and $\nabla_{\mathbf{y}} \eta = \nabla_{\mathbf{y}} \eta + \mathbf{y} \cdot \nabla_{\mathbf{y}} \nabla_{\mathbf{y}} \eta$. This implies

$$\mathbf{y} \cdot \nabla_{\mathbf{y}} \nabla_{\mathbf{y}} \eta = 0. \quad (12)$$

So $\eta(\mathbf{y})$ loses strict convexity along the constant \mathbf{y} direction in which the Hessian matrix $\nabla_{\mathbf{y}} \nabla_{\mathbf{y}} \eta$ is singular. The Legendre-Fenchel transformation (LFT) of such an $\eta(\mathbf{y})$ can exist only if the \mathbf{y} is restricted to a compact and convex domain, and the LFT is non-differentiable on the sub-manifold defined by (12).³ When the \mathbf{y} is extended to the entire \mathbb{R}^K , the domain of its LFT is contracted to a $K - 1$ sub-manifold

² One of course could construct mean square value and many other nonlinear statistics to gain additional information. The relationship between such an approach and traditional thermodynamics remains to be elucidated.

³ An illustrative example is $\eta(y_1, y_2) = y_1^2/y_2$ with $y_2 \in [b_1, b_2]$. The corresponding LFT $\psi(\beta_1, \beta_2) = b_1 z$ if $z(\beta_1, \beta_2) \equiv (\beta_2 - \beta_1^2/4) > 0$, and $= b_2 z$ if $z < 0$. The sub-manifold is $z(\beta_1, \beta_2) = 0$. When $[b_1, b_2]$ extends to the entire \mathbb{R} , $\psi(\beta_1, \beta_2)$ diverges almost everywhere; whose support contracts to the sub-manifold.

which defines the equation of state in classical thermodynamics. Since thermodynamic limit dictates $\eta(\mathbf{y})$ being a homogeneous function of \mathbf{y} with order 1 [17], this is the mathematical origin of duality symmetry breaking in thermodynamic limit.

Integral and differential β 's. Consider again $K = 1$ with both \mathbf{y} and β being scalars. Then Eq. 6 immediately suggests that

$$-\frac{d\eta(y)}{dy} + \frac{\eta(y)}{y} = -\beta + \frac{\eta(y)}{y} = \frac{\psi}{y}. \quad (13)$$

If both $\eta(y)$ and y are extensive quantities, as both tending to ∞ if their ratio $\eta(y)/y$ exists, then according to l'Hôpital's rule the $d\eta(y)/dy$ will have the same limit as $y \rightarrow \infty$. Then the right-hand-side of (13) vanishes, i.e., ψ is sub-extensive. In [12] Hill has introduced a notation $\eta(y)/y = \hat{\beta}$, called integral form of β in contrast to the differential $\beta = \partial\eta(y)/\partial y$. Then

$$-\beta + \hat{\beta} = \frac{\psi}{y} = \frac{\eta(y) - y\eta'(y)}{y} = -\frac{d\hat{\beta}}{d \ln y}. \quad (14)$$

This is another key result in Hills' thermodynamics of small systems [12, 16]. The right-hand-side of (14) has a very clear meaning. To illustrate we take y as the number of particles N for example, then

$$\frac{\psi}{N} = -N \frac{d}{dN} \left(\frac{\eta}{N} \right),$$

in which η/N is the entropy per particle. Accordingly $d(\eta/N)/dN$ is its change due to introducing one additional particle, and $Nd(\eta/N)/dN$ is the change in the entire system, of all N particles, due to one additional particle. In other words, when introducing one additional particle into a system of N particles, the effect is "subdivided" into all N particles.

The meaning of β . A thorough discussion of the meaning of β 's is outside the scope of the present Letter. In the context of a large number, M i.i.d. measurements, it suffices to say that when conditioned on an observation that $\bar{g}_M = y$, there is a posterior distribution $f_{\mathbf{x}}(\mathbf{x})e^{-\beta(y)g(\mathbf{x})}$ among the M samples [18]. If one has two sets of samples with $\bar{g}^{(1)} = y_1$ and $\bar{g}^{(2)} = y_2$, then the joint samples that pooled two together will have $\min\{y_1, y_2\} \leq \bar{g}^{(1 \cup 2)} \leq \max\{y_1, y_2\}$. Since $\beta(y)$ is a monotonic function of y , this also implies a new β that is between the β_1 and β_2 . It has been shown recently that canonical distribution can also be understood as a limit theorem of an infinitesimal portion of a large system [19], which gives the canonical distribution and its β an alternative interpretation.

Three entropies and two limits. In contrast to the classical thermodynamics with one entropy and one limit, the present theory is about three entropies and two limiting processes.

In addition to Gibbs entropy η and Massieu-Guggenheim entropy ψ , there is a third $\ln \Omega(\mathbf{y})$ corresponding to the probability density function for the observable $\mathbf{g}(\mathbf{x})$,

$$\begin{aligned} \Omega(\mathbf{y})d\mathbf{y} &= \Pr \{ \mathbf{y} < \mathbf{g}(\mathbf{x}) \leq \mathbf{y} + d\mathbf{y} \} \\ &= \int_{\mathbf{y} < \mathbf{g}(\mathbf{x}) \leq \mathbf{y} + d\mathbf{y}} f_{\mathbf{x}}(\mathbf{x})d\mathbf{x}, \end{aligned} \quad (15)$$

which completely determines the partition function and $\psi(\beta)$,

$$\psi(\beta) = \ln \int_{\mathbb{R}^K} \Omega(\mathbf{y}) \exp(-\beta \cdot \mathbf{y}) d\mathbf{y}. \quad (16)$$

$\Omega(\mathbf{y})$ is the *density of state* in terms of the observable $\mathbf{g}(\mathbf{x})$. $\ln \Omega$ can be identified as Boltzmann's microcanonical entropy if $\mathbf{g} = (U, V, N)$. In an essence our theory has a logic flow captured by the following scheme:

$$\ln \Omega(\mathbf{y}) \longrightarrow \left\{ \psi(\beta) \longleftrightarrow \eta(\mathbf{y}) \right\} \quad (17a)$$

$$\longrightarrow \left\{ \frac{\psi(\beta)}{\mathbf{y}} = 0, \frac{\eta(\mathbf{y})}{y_i} = \frac{d\eta}{dy_i} = \beta_i \right\}. \quad (17b)$$

If $\ln \Omega(\mathbf{y})$ is convex, then $\ln \Omega(\mathbf{y}) = \eta(\mathbf{y})$. In this case, the microcanonical description in terms of Boltzmann's entropy with \mathbf{y} as independent variables is equivalent to the canonical description in terms of Massieu-Guggenheim entropy with β as independent variables [20]. The y_i in (17b) can be any one component of the extensive \mathbf{y} . The derivative is understood as expressing all other components of \mathbf{y} normalized by the y_i . The " \rightarrow " in Eq. 17a represents the repeated-measurement limit; in general there is a loss of information on the small system. The " \leftrightarrow " in Eq. 17a indicates the emergent duality symmetry. This is the domain of Hill's nanothermodynamics [12]. The " \rightarrow " in Eq. 17b then represents the large system, macroscopic thermodynamics limit. It results in a breaking of the duality symmetry.

The thermodynamic structure presented in the present work, while assumes a probability distribution *a priori*, does not require the concept of *equilibrium* in connection to detailed balance in stochastic dynamics, nor *ergodicity*. Therefore, it is applicable to measurements on biomarkers from isogenic single living cells [9]. Of course, if a large system consists of many statistically identical but independent smaller parts, then the entire argument based on i.i.d. measurements can be applied to a single measurement of extensive variables of the large system as a whole. This mathematics was precisely in Boltzmann's theory of 1877, with the probabilistic concept of "conditioning on the value y " being replaced by Newtonian "conservation with the value y ".

The present result augments the current understanding of the nature of thermodynamic behavior, which so far has been focused on large systems limit and phase transition through

symmetry breaking as a key route for emergent phenomena [21, 22]. We now see there is actually a *large measurements limit* that generates a different kind of emergent order, a duality symmetry, for any small stochastic systems. This symmetry is lost, however, in the large systems limit.

Massieu-Guggenheim entropy of a fluctuating entropy?

The new theory might also help in further investigating some of the standing issues in the field of statistical thermodynamics. Take a discrete system with probability mass function $\Pr\{X = k\} = p(k)$ for example, it has been shown recently that $\ln p(X)$, as a function of the random variable X , is the *fluctuating entropy* in stochastic thermodynamics [2]. Then if one constructs the Massieu-Guggenheim entropy $\psi(\beta)$ corresponding to the random variable $\ln p(X)$:

$$\psi(\beta) = \ln \sum_k p(k) e^{-\beta \ln p(k)} = \ln \sum_k p^{1-\beta}(k), \quad (18)$$

which looks like the Tsallis entropy. In fact, if $\beta = 1 - q \ll 1$, one has the corresponding negative free energy

$$\frac{\psi(\beta)}{\beta} = \frac{1}{1-q} \left(\sum_k p^q(k) - 1 \right). \quad (19)$$

Acknowledgements. H. Q. is grateful to Prof. Lu Yu (ITP, CAS) for advice and encouragement. We thank the members of Summer 2020 Online Club Nanothermodynamica, especially Quanhui Liu and Hong Zhao, for extensive discussions.

* zhiyuelu@unc.edu

† hqian@u.washington.edu

- [1] Chibbaro, S., Rondoni, L. and Vulpiani, A. (2014) *Reductionism, Emergence and Levels of Reality*. Springer, New York.
- [2] Qian, H., Cheng, Y.-C. and Thompson, L. F. (2019) Ternary representation of stochastic change and the origin of entropy and its fluctuations. arXiv:1902.09536.
- [3] Ellis, R. S. (1985) *Entropy, Large Deviations, and Statistical Mechanics*, Springer, Berlin.
- [4] Oono, Y. (1989) Large deviation and statistical physics. *Progr. Theoret. Phys. Supp.* **99**, 165–205.
- [5] Touchette, H. (2009) The large deviation approach to statistical mechanics. *Phys. Rep.* **478**, 1–69.
- [6] Ge, H. and Qian, H. (2012) Analytical mechanics in stochastic dynamics: Most probable path, large-deviation rate function and Hamilton-Jacobi equation. *Intern. J. Mod. Phys. B* **26**, 1230012.
- [7] Dembo, A. and Zeitouni, O. (2014) *Large Deviations Techniques and Applications*, 2nd ed. Springer, New York.
- [8] Smith, E. (2011) Large-deviation principles, stochastic effective actions, path entropies, and the structure and meaning of thermodynamic description. *Rep. Prog. Phys.* **74**, 046601.
- [9] Qian, H. and Cheng, Y.-C. (2020) Counting single cells and computing their heterogeneity: from phenotypic frequencies to mean value of a quantitative biomarker. *Quant. Biol.* **8**, 172–176.
- [10] Planck, M. (2010) *Treatise on Thermodynamics*, Dover, New York.
- [11] Guggenheim, E. A. (1939) Grand partition functions and the so-called “thermodynamic probability”. *J. Chem. Phys.* **7**, 103–107.
- [12] Hill, T. L. (1963) *Thermodynamics of Small Systems*, Dover, New York.
- [13] Rockafellar, R. T. (1996) *Convex Analysis*. Princeton Univ. Press, NJ.
- [14] Hill, T. L. (2001) A different approach to nanothermodynamics. *Nano Lett.* **1**, 273–275.
- [15] Chamberlin, R. V., Clark, M. R., Mujica, V. and Wolf, G. H. (2020) An introduction to nanothermodynamics: Thermal equilibrium for heterogeneous and finite-sized systems. arXiv:2007.13031.
- [16] Bedeaux, D., Kjelstrup, S. and Schnell, S. K. (2020) *Nanothermodynamics: General Theory*. NTNU, Trondheim, Norway.
- [17] Callen, H. B. (1985) *Thermodynamics and an Introduction to Thermostatistics*, 2nd ed., Wiley, New York.
- [18] van Campenhout, J. M. and Cover, T. M. (1981) Maximum entropy and conditional probability. *IEEE Tran. Info. Th.* **IT-27**, 483–489.
- [19] Cheng, Y.-C., Qian, H. and Zhu, Y. (2019) Asymptotic behavior of a sequence of conditional probability distributions and the canonical ensemble. arXiv:1912.11137.
- [20] Touchette, H. (2015) Equivalence and nonequivalence of ensembles: Thermodynamic, macrostate, and measure levels. *J. Stat. Phys.* **159**, 987–1016.
- [21] Anderson, P. W. (1972) More is different. *Science*, **177**, 393–396.
- [22] Qian, H., Cheng, Y.-C. and Yang, Y.-J. (2020) Kinematic basis of emergent energetics of complex dynamics. *EPL*, **131**, 50002.