The smoothed complexity of Frank-Wolfe methods via conditioning of random matrices and polytopes

Luis Rademacher University of California, Davis lrademac@ucdavis.edu Chang Shu University of California, Davis ccshu@ucdavis.edu

Abstract


Frank-Wolfe methods are popular for optimization over a polytope. One of the reasons is because they do not need projection onto the polytope but only linear optimization over it. To understand its complexity, a fruitful approach in many works has been the use of condition measures of polytopes. Lacoste-Julien and Jaggi introduced a condition number for polytopes and showed linear convergence for several variations of the method. The actual running time can still be exponential in the worst case (when the condition number is exponential). We study the smoothed complexity of the condition number, namely the condition number of small random perturbations of the input polytope and show that it is polynomial for any simplex and exponential for general polytopes. Our results also apply to other condition measures of polytopes that have been proposed for the analysis of Frank-Wolfe methods: vertex-facet distance (Beck and Shtern) and facial distance (Peña and Rodríguez).

Our argument for polytopes is a refinement of an argument that we develop to study the conditioning of random matrices. The basic argument shows that for c > 1 a d-by-n random Gaussian matrix with $n \ge cd$ has a d-by-d submatrix with minimum singular value that is exponentially small with high probability. This also has consequences on known results about the robust uniqueness of tensor decompositions, the complexity of the simplex method and the diameter of polytopes.

1 Introduction

Frank-Wolfe methods (FWMs) [FW56] are a family of algorithms that attempt to minimize a differentiable function over a convex set. For concreteness we start by describing the basic Frank-Wolfe method to minimize a differentiable function $f: C \mapsto \mathbb{R}$ where $C \subseteq \mathbb{R}^d$ is a compact convex set. It is an iterative method and proceeds as follows:

Let
$$x_0 \in C$$
.
for $k = 0, ..., K$ do
Compute $y \in \operatorname{argmin}_{x \in C}(\nabla f(x_k))^T x$.
Let $x_{k+1} = x_k + \alpha^*(y - x_k)$, where α^* is a suitable step size.
end for



Some of our results are about Wolfe's method [Wol76], which is a variation of Frank-Wolfe methods specialized to the minimum norm point problem in a polytope (that is, a bounded convex polyhedron).

1.1 Our contributions and related work

In this paper we are interested in the complexity of FWMs. The time complexity of Wolfe's method is know to be exponential in the worst case (by an upper bound in [Wol76] and a lower bound in [LHR20]). There is a large body of work proving linear convergence of several variations of FWMs [GM86, GH13, LJJ13, LJJ15, BS17, PnRS16, PnR19, PNAJ20]. We are particularly interested in [LJJ13, LJJ15, BS17, PnRS16, PnR19] which prove global linear convergence of certain variations of FWMs: F-W with away steps, pairwise F-W and Wolfe's method when the feasible region is a polytope C = conv(A) for finite $A \subseteq \mathbb{R}^d$. In these results the upper bound on the running time (actual speed of linear convergence) depends on a condition number of C. Informally speaking, the dependence is of the following kind: if x_t is the current point after t iterations, then the function value satisfies $f(x_t) - f^* \le (1 - \kappa)^t (f(x_0) - f^*)$ where f^* is the optimal value, x_0 is the initial point and $0 \le \kappa \le 1$ is a measure of conditioning. If κ is small, then convergence is slow. In the previously mentioned papers, κ is of the form "something" / diam(C), where "something" can be:

- [LJJ15] minimum width, minwidth(A) = $\min_{S\subseteq A}$ width(S) (width is standard, see Section 2.8.1);
- [LJJ15] pyramidal width, PWidth(A);
- [BS17] vertex-facet distance, vf(C) = $\min_{F \in \text{facets}(C)} d(\text{aff } F, \text{vertices}(C) \setminus F)$; or
- [PnR19] facial distance, $\Phi(C) = \min_{\substack{F \in \text{faces}(C) \\ \emptyset \subsetneq F \subsetneq C}} d(F, \text{conv}(\text{vertices}(C) \setminus F)).$

We do not provide a definition of pyramidal width at this point as it is complicated and it was shown in [PnR19] that PWidth(A) = $\Phi(C)$ (Theorem 2.24 here). It is also known that minwidth(A) \leq PWidth(A) [LJJ15, Section 3.1]. We start with the observation that $\Phi(C) \leq \text{vf}(C)$ (Theorem 2.25). (Note that the reverse inequality was claimed in [PnR19], but the cube $[0,1]^d$ is a counterexample: $\Phi([0,1]^d) = 1/\sqrt{d}$ while $\text{vf}([0,1]^d) = 1$.) This implies that all four quantities lie between minwidth(A) and vf(C) (Theorem 2.25). It follows from [LHR20] that all of them can be exponentially small as a function of the bit-length of A. In fact, a stronger result follows from the work of Alon and Vu [AV97] combined with the stated inequalities. Alon and Vu showed that there is a 0–1 simplex S such that vf(S) is sub-exponentially small in the dimension (Corollary 3.3). The connection between polytope conditioning for FWMs and the Alon and Vu result was observed in [LJJ15].

The main contributions of this paper are about the smoothed analysis of FWMs and the condition numbers of matrices and polytopes. Smoothed analysis [ST01] is an approach to understand the behavior of algorithms that are efficient in practice but are inefficient in the worst case. The main idea is to study small random perturbations of any given instance of a problem. Suppose that the instance is described by a vector $x \in \mathbb{R}^n$. Then one aims to understand T(x+g), where $g \in \mathbb{R}^n$ is a random vector with distribution $N(0, \sigma^2 \mathbf{I}_n)$ and T is a measure of complexity (for example, T(x) could be the running time of a particular algorithm on input x). We adopt a definition that first appeared in [BV04, BV06].

Definition 1.1 ([RV05] [RV07, Section 1.1]). We say T has (probabilistic) polynomial smoothed complexity if there is a polynomial p such that

$$\max_{x \in \mathbb{R}^n, ||x|| \le 1} \mathbb{P}_g \big(T(x+g) \ge p(n, 1/\sigma, 1/\delta) \big) \le \delta$$

Our first smoothed analysis result concerns FWMs minimizing a convex function on a simplex (Section 3). We show that minwidth has good smoothed complexity (Lemma 3.6). This implies the following result on polytope conditioning that can be combined with results in [LJJ15] to show polynomial smoothed time complexity of several FWMs for the minimization of a convex function in any simplex:

Theorem 1.2. Let $A = \{A_1, \ldots, A_{d+1}\}$ be a set of independent Gaussian random vectors with means μ_i , $\|\mu_i\| \leq 1, i \in [d+1]$, and covariance matrix $\sigma^2 \mathbf{I}_d$. Then for $\delta > 0$, with probability at least $1 - \delta$, the measure of conditioning $\kappa = \frac{\mathrm{PWidth}(A)}{\mathrm{diam}(A)}$ of A is at least some inverse polynomial in d, $1/\sigma$ and $1/\delta$.

Note that even the problem of finding the minimum norm point in a simplex is not known to have a simple polynomial time algorithm. All polynomial time algorithms we know for such a special case are general purpose convex programming algorithms such as the ellipsoid method. Moreover, [LHR20] shows that the linear programming problem reduces in strongly polynomial time to the minimum norm point in a simplex problem. This suggests that to find a simple polynomial time algorithm for the minimum norm point in a simplex is hard and, in particular, to find a strongly polynomial time algorithm would imply the existence of a strongly polynomial time algorithm for linear programming, which would solve a major open problem.

Our second smoothed analysis result concerns condition measures of general polytopes (Section 7). We show that the standard global linear convergence results for FWMs mentioned above based on polytope conditioning cannot guarantee polynomial complexity for general polytopes in the average or smoothed sense. More specifically, for V-polytopes conv(A) with |A| and d large and comparable, $d \approx \delta |A|$, $\delta \in (0,1)$, we show that vertex-facet distance does not have polynomial smoothed complexity. Given that the complexity here increases as vf(A) gets smaller, in the context of Definition 1.1 one sets T = 1/vf. It is enough to take x = 0 there and we show:

Theorem 1.3. Let $\delta \in (0,1)$. Suppose $A = \{A_1, \ldots, A_{n+1}\}$ is a set of iid. standard Gaussian random vectors in \mathbb{R}^d and $d = \lfloor \delta n \rfloor$. Let $P_{n+1} = \text{conv}(A_1, \ldots, A_{n+1})$. Then

$$\mathbb{P}(\operatorname{diam}(P_{n+1}) \ge \sqrt{d}) \ge 1 - e^{-\frac{nd}{32}},$$

and there exists constants 0 < c, c' < 1 (that depends only δ) such that,

$$\lim_{n \to \infty} \mathbb{P}(\operatorname{vf}(P_{n+1}) \le c^d) \ge c'.$$

Hence the measure of conditioning $\kappa = \frac{\operatorname{vf}(P_{n+1})}{\operatorname{diam}(P_{n+1})}$ of A is exponentially small in d with constant probability.

Theorem 1.3 combined with Theorem 2.25 implies that none of the four measures of polytope conditioning (minwidth, PWidth, Φ , vf) has polynomial smoothed complexity.

A way of interpreting Theorem 1.3 is that the standard conditioning measures of polytopes for FWMs are somewhat pessimistic and can appear ill-conditioned even then polytope is bad only

locally. For example, vertex-facet distance can be small even if one vertex and one facet are bad while the rest of the polytope is good. In other words, it may still be possible to show smoothed polynomial complexity of FWMs in a different way.

Theorem 1.3 is a statement about the minimum distance between the affine hull of d points that form a facet and a vertex not on that facet. In order to understand this problem we study first a simplified version where we replace affine hull by span and we remove the restriction that the d-1 points form a facet. Namely, we study the following question: given n standard Gaussian random points in \mathbb{R}^d , how close can one of the points be to the span of some d-1 others when n is somewhat larger than d, say, n=2d? This question is easier to understand than the polytope version and it relates to conditioning of random matrices and the restricted isometry property in compressive sensing. The relation starts from the known observation (Lemma 2.8) that the minimum point-hyperplane distance is, up to polynomial factors, the same as the smallest singular value of a matrix. Given this, our question is essentially equivalent to: given an d-by-n random matrix with iid. standard Gaussian entries, what is the minimum of the smallest singular values over d-by-d submatrices? We answer this question by showing that when $n/d \ge c > 1$ the minimum smallest singular value above (and, equivalently, minimum point-hyperplane distance) is exponentially small:

Theorem 1.4. Let A be an d-by-n random matrix with iid. standard Gaussian entries with $d \ge 2$ and $\frac{n}{d} \ge c_0 > 1$. Then, there exist constants $c_2, c_4 > 1$, $0 < c_6 < 1$ (that depend only on c_0) such that with probability at least $1 - 2c_4c_6^d$,

$$\min_{S\subseteq[n],|S|=d} \sigma_d(A_S) \le \frac{1}{c_4 c_2^{d-1}}.$$

Theorem 1.5. Let A be an d-by-n random matrix with iid. standard Gaussian entries with $d \ge 2$ and $1 < \frac{n}{d-1} \le C_0$. Then, there exist constants $C_1 > 1$, $0 < C_2 < 1$ (that depend only on C_0) such that with probability at least $1 - nC_2^{d-1}$,

$$\min_{S\subseteq[n],|S|=d}\sigma_d(A_S)\geq \frac{1}{C_1^{d-1}}.$$

While Theorems 1.4 and 1.5 are new as far as we know, there is a large body of work, partly motivated by compressive sensing, that studies questions related to them. In that area one is generally interested in showing that all d-by-k submatrices of A are well-conditioned, say, σ_1/σ_k is no more than a constant (the restricted isometry property of Candès and Tao [CT05, CT06]). This can only happen when k is much smaller than d, a regime very different from our case k = d. The standard analyses in compressive sensing as well as recent results such as [CJL19] do not seem to be able to clarify the behavior in our regime.

The idea of the proof of Theorem 1.4 (Section 4) is the following: Consider the case n=2d for concreteness and aim to show that with constant probability one point is exponentially close to the span of d-1 others. Let S be the family of sets of d-1 columns of A. For $S \in S$, let \mathcal{B}_S be the set of points in \mathbb{R}^d within distance ϵ of span S. Let $V_{\epsilon} = \bigcup_{S \in S} \mathcal{B}_S$. It is enough to show that for $\epsilon = 1/c^d$, c > 1, the Gaussian volume $\mathcal{G}(V_{\epsilon})$ is at least a constant. We do this by lower bounding it using the first two terms of the inclusion-exclusion principle (Bonferroni inequality):

$$\mathcal{G}(V_{\epsilon}) \geq \sum_{S} \mathcal{G}(\mathcal{B}_{S}) - \frac{1}{2} \sum_{S,T:S \neq T} \mathcal{G}(\mathcal{B}_{S} \cap \mathcal{B}_{T}).$$

Note that $\mathcal{B}_S \cap \mathcal{B}_T$ can be large if S and T share many columns. To deal with this difficulty, replace S above with a large subfamily $T \subseteq S$ of subsets of columns where each pair of subsets has few columns in common by picking separated subsets *greedily (Gilbert-Varshamov bound)*. See [Raz88], [Juk11, Lemma 19.3] for another instance of Bonferroni's inequality with almost pairwise independence.

While Theorems 1.4 and 1.5 are results about random matrices, they have direct implications in the analysis of algorithms: In Section 5 we discuss how Theorem 1.4 conditions the applicability of the robustness of tensor decomposition result by Bhaskara, Charikar and Vijayaraghavan [BCV14]. In Section 6 we discuss how Theorem 1.4 conditions the applicability of results about the complexity of the simplex method and the diameter of polytopes in [BR13, BGR15, DH16, EV17].

2 Preliminaries

2.1 Notations

For $v \in \mathbb{R}^d$ and $i \in [d]$, let v_{-i} denote vector v with coordinate v_i removed, that is $v_{-i} := (v_1, \ldots, v_{i-1}, v_{i+1}, \ldots, v_d)$. If $v \neq 0$, let $\hat{v} := v/\|v\|_2$. Let $B(x, \epsilon) := \{y \in \mathbb{R}^d : \|y - x\|_2 \leq \epsilon\}$. Let \mathcal{S}^{d-1} denote the (d-1)-dimensional unit sphere in \mathbb{R}^d . For $v \in \mathcal{S}^{d-1}$, denote the spherical cap centered at v with angle α as $\mathcal{C}_{\alpha}(v) := \{x \in \mathcal{S}^{d-1} : v \cdot x \geq \cos \alpha\}$. For $A \subseteq \mathbb{R}^d$, let $A_{\epsilon} := \{x \in \mathbb{R}^d : \operatorname{dist}(x, A) \leq \epsilon\}$, $A_{-\epsilon} := \{x \in \mathbb{R}^d : B(x, \epsilon) \subset A\}$. Let \mathcal{G} denote the standard multivariate Gaussian probability measure. For random variables or distributions X, Y, notation $X \stackrel{d}{=} Y$ states that X and Y have the same distribution.

2.2 Noncentral Chi-Square Distribution

We first recall the definition of noncentral chi-square distribution.

Definition 2.1. Let Y_1, Y_2, \ldots, Y_k be independent Gaussian random variables with means μ_i and unit variance. Then the random variable $\sum_{i=1}^k Y_i^2$ is distributed according to the noncentral chisquare distribution with k degrees of freedom and noncentrality parameter $\lambda = \sum_{i=1}^k \mu_i^2$. The probability density function of noncentral chi-square distribution is given by

$$f_Y(x; k, \lambda) = \sum_{i=0}^{\infty} \frac{e^{-\lambda/2} (\lambda/2)^i}{i!} f_{Z_{k+2i}}(x),$$

where Z_q is distributed as (central) chi-square with q degrees of freedom, denoted as χ_q^2 .

We show an anti-concentration inequality of noncentral chi-square distribution by comparing to (central) chi-square distribution.

Lemma 2.2. Let $\mu \in \mathbb{R}$. Let $X_i \sim \mathcal{N}(0, \sigma^2)$, $Y_i \sim \mathcal{N}(\mu, \sigma^2)$, $i \in [k]$, be independent. Then

$$\mathbb{P}\left(\sum_{i=1}^k X_i^2 \ge t^2\right) \le \mathbb{P}\left(\sum_{i=1}^k Y_i^2 \ge t^2\right).$$

Proof. The proof follows directly from the density function of noncentral chi-square distribution and some basic facts about the chi-square distribution. We consider random variables $Y_i/\sigma \sim$

 $\mathcal{N}(\mu/\sigma, 1)$. Then $\sum_{i=1}^k Y_i^2/\sigma^2$ is distributed as noncentral chi-square with k degrees of freedom and noncentrality parameter $k\mu^2/\sigma^2$. From Definition 2.1,

$$\mathbb{P}\left(\sum_{i=1}^{k} \frac{Y_i^2}{\sigma^2} \ge \frac{t^2}{\sigma^2}\right) = \sum_{i=0}^{\infty} \frac{e^{-k\mu^2/2\sigma^2} (k\mu^2/2\sigma^2)^i}{i!} \mathbb{P}\left(Z_{k+2i} \ge \frac{t^2}{\sigma^2}\right)$$

$$\ge \sum_{i=0}^{\infty} \frac{e^{-k\mu^2/2\sigma^2} (k\mu^2/2\sigma^2)^i}{i!} \mathbb{P}\left(Z_k \ge \frac{t^2}{\sigma^2}\right)$$

$$= \mathbb{P}\left(Z_k \ge \frac{t^2}{\sigma^2}\right)$$

$$= \mathbb{P}\left(\sum_{i=1}^{k} X_i^2 \ge t^2\right),$$

where $Z_k \sim \chi_k^2$. The inequality comes from the fact that chi-square random variable with (k+i) degrees of freedom is equal in distribution to the sum of a chi-square random variable with k degrees of freedom and squares of i independent standard Gaussian random variables, so that Z_{k+2i} has larger tail than Z_k .

The following lemma provides a comparison inequality between the ratio of noncentral chi-square random variables and chi-square random variables, which is used in the proof of Lemma 7.4.

Lemma 2.3. Let $\mu \in \mathbb{R}$. Let $X_0, X_i, Y_0 \sim \mathcal{N}(0, \sigma^2)$, $Y_i \sim \mathcal{N}(\mu, \sigma^2)$, $i \in [k]$ and be independent. Then for any $t \in (0, 1)$,

$$\mathbb{P}\left(\frac{Y_0^2}{Y_0^2 + \sum_{i=1}^n Y_i^2} \ge t^2\right) \le \mathbb{P}\left(\frac{X_0^2}{X_0^2 + \sum_{i=1}^n X_i^2} \ge t^2\right).$$

Proof. Let f denote the probability density function of X_0^2 and Y_0^2 . By the law of total expectation,

$$\begin{split} \mathbb{P}\left(\frac{Y_0^2}{Y_0^2 + \sum_{i=1}^n Y_i^2} \geq t^2\right) &= \int_0^\infty \mathbb{P}\left(\frac{y}{y + \sum_{i=1}^n Y_i^2} \geq t^2\right) f(y) dy \\ &= \int_0^\infty \mathbb{P}\left(\sum_{i=1}^n Y_i^2 \leq (1/t^2 - 1)y\right) f(y) dy \\ &\leq \int_0^\infty \mathbb{P}\left(\sum_{i=1}^n X_i^2 \leq (1/t^2 - 1)x\right) f(x) dx \quad \text{(Lemma 2.2)} \\ &= \int_0^\infty \mathbb{P}\left(\frac{x}{x + \sum_{i=1}^n X_i^2} \geq t^2\right) f(x) dx \\ &= \mathbb{P}\left(\frac{X_0^2}{X_0^2 + \sum_{i=1}^n X_i^2} \geq t^2\right). \end{split}$$

2.3 Concentration and tail inequalities

Lemma 2.4. Let $X \sim \mathcal{N}(0, \sigma^2)$, $Y \sim \mathcal{N}(c, \sigma^2)$, $c \in \mathbb{R}$, then for any t > 0,

$$\mathbb{P}(\|Y\|_2 < t) \leq \mathbb{P}(\|X\|_2 < t).$$

Proof. Without loss of generality, we may assume c > 0. Then

$$\begin{split} \mathbb{P}(\|Y\|_2 < t) &= \int_{-t}^t e^{\frac{-(x-c)^2}{2\sigma^2}} dx \\ &= \int_{-t-c}^{t-c} e^{\frac{-x^2}{2\sigma^2}} dx \\ &= \int_{-t-c}^{-t} e^{\frac{-x^2}{2\sigma^2}} dx + \int_{-t}^t e^{\frac{-x^2}{2\sigma^2}} dx - \int_{-t-c}^t e^{\frac{-x^2}{2\sigma^2}} dx \\ &= \mathbb{P}\big(\|X\|_2 < t\big) + \left(\int_t^{t+c} e^{\frac{-x^2}{2\sigma^2}} dx - \int_{t-c}^t e^{\frac{-x^2}{2\sigma^2}} dx\right) \\ &\leq \mathbb{P}\big(\|X\|_2 < t\big). \end{split}$$

Lemma 2.5 ([LM00]). Let (X_1, \ldots, X_n) be iid. standard Gaussian variables. Let $\alpha_1, \ldots, \alpha_n$ be nonnegative. Let $Z = \sum_{i=1}^n \alpha_i(X_i^2 - 1)$. Then, the following inequalities hold for any positive t:

$$\mathbb{P}(Z \ge 2\|\alpha\|_2 \sqrt{t} + 2\|\alpha\|_{\infty} t) \le \exp(-t),$$

$$\mathbb{P}(Z \le -2\|\alpha\|_2 \sqrt{t}) \le \exp(-t).$$

2.4 Gilbert-Varshamov bound

Lemma 2.6. Let A(n,t,w) be the maximum number of binary n-vectors with exactly w ones and pairwise Hamming distance greater than or equal to t. Then for any $c_0 > 1$, there exist constants $c_1 > 0$ and $c_2 > 1$ (that depend only on c_0) such that for all $d \ge 1$ and $n/d \ge c_0$ we have $A(n,c_1d,d) \ge c_2^d$.

Proof. Pick vectors greedily (Gilbert-Varshamov bound) to get, for integral t:

$$A(n,t,w) \ge \frac{\binom{n}{w}}{B(n,t)},$$

where B(n,t) is the number of binary vectors at Hamming distance less than or equal to t from the zero vector. We have $B(n,t) = \sum_{k=0}^{t} \binom{n}{k} \leq \left(\frac{ne}{t}\right)^t$ (see footnote¹). Note that this last inequality is valid also for fractional $0 < t \leq n$ using the fact that $(a/b)^b$ is increasing in b for a > 0, $0 < b \leq a/e$. Thus, for any 0 < c < 1 and $d \geq 1$ we have

$$A(n, cd, d) \ge \frac{\binom{n}{d}}{\left(\frac{ne}{cd}\right)^{cd}} \ge \frac{(n/d)^d}{\left(\frac{ne}{cd}\right)^{cd}} = \frac{(n/d)^{d(1-c)}}{(e/c)^{cd}} \ge \frac{c_0^{d(1-c)}}{(e/c)^{cd}} = \left(\frac{c_0}{(c_0e/c)^c}\right)^d \tag{1}$$

We have $\lim_{c\to 0^+} (a/c)^c = 1$ and $(a/c)^c$ is increasing again for $a>0,\ 0\le c\le a/e$. Given this, choose $c_1\in (0,1)$ such that $(c_0e/c_1)^{c_1}< c_0$. Let $c_2:=\frac{c_0}{(c_0e/c_1)^{c_1}}>1$. We have $A(n,c_1d,d)\ge c_2^d$ for $d\ge 1$. The claim follows.

$$\frac{1}{\sum_{k=0}^{t} \binom{n}{k}} \le \sum_{k=0}^{t} \frac{n^k}{k!} = \sum_{k=0}^{t} \frac{t^k}{k!} (n/t)^k \le e^t (n/t)^t.$$

2.5 Generalization of Archimedes' formula

Lemma 2.7. Let $d \ge 3$. Let U be a uniformly random d-dimensional unit vector. Then (U_1, \ldots, U_{d-2}) is uniform in B^{d-2} and $\Pr(\|(U_1, \ldots, U_{d-2})\| \le t) = t^{d-2}$.

Proof. The first part is well-known, a proof can be found in [BGMN05, Corollary 4]. The second part follows immediately from the first part. \Box

2.6 One-off-distance vs sigma min

Lemma 2.8 (see e.g. [BCMV13, Lemma 3.5] for a proof). If $A \in \mathbb{R}^{m \times n}$ has columns a_1, \ldots, a_n and $m \ge n$, then denoting $a_{-i} = \operatorname{span}(a_i : j \ne i)$, we have

$$\frac{1}{\sqrt{n}} \min_{i \in [n]} \operatorname{dist}(a_i, a_{-i}) \le \sigma_n(A) \le \min_{i \in [n]} \operatorname{dist}(a_i, a_{-i}).$$

2.7 Facts about Gaussian random polytopes

2.7.1 Gaussian ϵ -neighborhood

Corollary 2.9. Let Q be a convex set in \mathbb{R}^d . Then there exists an absolute constant c > 0 such that $\mathcal{G}(Q \setminus Q_{-\epsilon}) \leq c\epsilon d^{1/4}$.

Proof. Follows immediately from [CCK17, Lemma A.2] and the fact $||I||_{HS} = \sqrt{d}$ (Hilbert-Schmidt norm). Their proof is based on [Bal93, Naz03].

2.7.2 Distances of facets

Lemma 2.10 ([Ver18, Theorem 4.4.5]). Let X be an $m \times n$ random matrix whose entries are iid. standard Gaussian random variables. Then for t > 0,

$$\mathbb{P}\left(\sigma_{\max}(X) > c(\sqrt{m} + \sqrt{n} + t)\right) \le 2e^{-t^2},$$

where c is some absolute positive constant.

Lemma 2.11. Given linearly independent vectors $p_1, p_2, \dots, p_d \in \mathbb{R}^d$, the shortest vector in their affine hull is $v = P^{-1} \mathbb{1}/\|P^{-1}\mathbb{1}\|^2$, where $P = (p_1 \cdots p_d)^{\top}$. In particular, $\|v\| = 1/\|P^{-1}\mathbb{1}\|$.

Proof. From [LHR20, Lemma 1.2], the shortest vector in the affine hull, v, satisfies $Pv = ||v||^2 \mathbb{1}$. Since P has independent columns, $v = ||v||^2 P^{-1} \mathbb{1}$. Compute norms of vectors in the above equation to get $||v|| = 1/||P^{-1}\mathbb{1}||$. The claim follows.

The following lemma can directly generalize to Gaussian random vectors with mean zero and covariance matrix $\sigma^2 \mathbf{I}_d$ by scaling by σ .

Lemma 2.12. Let X_1, \ldots, X_n be iid. standard Gaussian random vectors in \mathbb{R}^d . For $S \subseteq [n]$, |S| = d, define V_S as the shortest vector in $\operatorname{aff}(X_S)$. Then there exists a constant c > 0 such that

$$\mathbb{P}\left(\max_{S\subseteq[n],|S|=d} \|V_S\| \le c(2+\sqrt{n/d})\right) \ge 1 - 2e^{-d}.$$

Proof. Let X be the matrix whose column vectors are X_1, \ldots, X_n . For any $S \subseteq [n], |S| = d, X_S$ is linearly independent with probability 1. By Lemma 2.11,

$$||V_S|| = \frac{1}{||X_S^{-1}\mathbb{1}||} \le \frac{1}{\sqrt{d} \ \sigma_{\min}(X_S^{-1})} = \frac{\sigma_{\max}(A_S)}{\sqrt{d}} \le \frac{\sigma_{\max}(A)}{\sqrt{d}}.$$
 (2)

From Lemma 2.10 we know $\mathbb{P}\left(\sigma_{\max}(A) > c(\sqrt{d} + \sqrt{n} + t)\right) \leq 2e^{-t^2}$. The claim follows by letting $t = \sqrt{d}$ and applying (2).

2.7.3 Number of facets

We will need the fact that the number of facets of the convex hull of n Gaussian random points in \mathbb{R}^d is exponential in d with high probability when $n=cd,\,c>1$. We could not find such a result in the literature and we do not see how to deduce it from results on the asymptotic number of facets in stochastic geometry [Ray70, AW91, HMR04, HR05, BLR18] (the difficulties are: either they only determine the expectation or variance of the number of facets, or the bounds are as n goes to infinity for fixed d). Nevertheless, it is easy to deduce what we want from the work of Donoho and Tanner on compressive sensing and the neighborliness of random polytopes. We build on top of basic polytope theory from [Zie93].

Definition 2.13 (Neighborliness). A polytope P is k-neighborly if every subset of k vertices forms a (k-1)-face.

Let $f_l(P)$ denote the number of *l*-faces of polytope P.

Theorem 2.14 ([DT05], Corollary 1.1, Lemma 3.2). There exists a function (threshold) $\rho(\delta)$: $(0,1) \to \mathbb{R}$, $\rho(\delta) > 0$ with the following property: Let $\delta \in (0,1)$. Let $d = \lfloor \delta n \rfloor$. Let $\rho < \rho(\delta)$. Let X_1, \ldots, X_n be iid. samples from a Gaussian distribution in \mathbb{R}^d with non-singular covariance. Let $P = \text{conv}\{X_1, \ldots, X_n\}$. Then $\lim_{n \to \infty} \mathbb{P}(f_1(P) = n \text{ and } P \text{ is } \lfloor \rho d \rfloor \text{-neighborly}) = 1$.

The above theorem demonstrates, given its assumptions, that when n is large enough, P has $\binom{n}{\lfloor \rho d \rfloor}$ many $\lfloor \rho d \rfloor$ -faces with high probability. Note also that P is simplicial (every facet is a simplex) a.s. Thus, a.s. each facet of P provides at most $\binom{d}{\lfloor \rho d \rfloor}$ many $\lfloor \rho d \rfloor$ -faces, and the number of facets is at least

$$\frac{\binom{n}{\lfloor \rho d \rfloor}}{\binom{d}{\lfloor \rho d \rfloor}} \ge \left(\frac{n}{d}\right)^{\lfloor \rho d \rfloor} \ge \left(\frac{1}{\delta}\right)^{\lfloor \rho d \rfloor} \ge c^d,$$

for some c > 1 (and d large enough). We conclude:

Corollary 2.15. Let $\delta \in (0,1)$. Let P be the convex hull of n iid. standard Gaussian random points in \mathbb{R}^d , $d = \lfloor \delta n \rfloor$. Then there exists a constant c > 1 (that depends only on δ) such that $\lim_{n\to\infty} \mathbb{P}(f_d(P) \geq c^d) = 1$.

Corollary 2.15 can probably also be proven directly from different but related neighborliness results by Vershik and Sporyshev [VS92], [DT05, Theorem 2].

2.8 Condition measures of polytopes

2.8.1 Width and minwidth

Definition 2.16 (Directional width and width). The directional width of a set $A \subseteq \mathbb{R}^d$ with respect to a direction $r \in \mathbb{R}^d$ is defined as $\operatorname{dirW}(A,r) := \sup_{s,v \in \mathcal{A}} \langle \frac{r}{\|r\|}, s-v \rangle$. The width of A, denoted width (A) is the infimum of the directional width over all directions on its affine hull.

Definition 2.17 (Minwidth, [LJJ15, Section 3.1]). The minwidth of a finite set $A \subseteq \mathbb{R}^d$, denoted minwidth(A), is the minimum width over all subsets of A.

2.8.2 Pyramidal width

Definition 2.18 (Pyramidal directional width, [LJJ15]). We define the pyramidal directional width of a finite set $A \subseteq \mathbb{R}^d$ with respect to a direction $r \in \mathbb{R}^d$ and a base point $x \in \text{conv}(A)$ to be

$$PDirW(A, r, x) := \min_{S \in S_x} dirW(S \cup \{s(A, r)\}, r) = \min_{S \in S_x} \max_{s \in A, v \in S} \left\langle \frac{r}{\|r\|}, s - v \right\rangle,$$

where $S_x := \{T \subseteq A : x \text{ is a proper convex combination of all the elements in } T\}$ and $s(A, r) := \operatorname{argmax}_{v \in A} \langle r, v \rangle$.

Definition 2.19 (Feasible direction, [LJJ15]). A direction r is feasible for A from x if it points inwards conv(A), i.e. $r \in cone(A-x)$. A direction r is feasible for A if it is feasible for A from some $x \in A$.

Definition 2.20 (Pyramidal width, [LJJ15]). We define the pyramidal width of a finite set $A \subseteq \mathbb{R}^d$ to be the smallest pyramidal directional width of all its faces,

$$\operatorname{PWidth}(A) := \min_{\substack{K \in \operatorname{faces}(\operatorname{conv}(A)) \\ x \in K \\ r \in \operatorname{cone}(K-x) \backslash \{0\}}} \operatorname{PDirW}(K \cap A, r, x).$$

2.8.3 Vertex-facet distance

The vertex-facet distance polytope conditioning parameter for the analysis of FWMs was introduced in [BS17]. We adopt here the slightly specialized definition in [PnR19], which is defined as a property of a polytope independent of the representation, while the original version in [BS17] can depend on the numbers used to represent a polytope.

Definition 2.21 (vertex-facet distance [BS17, PnR19]). Let $P \subseteq \mathbb{R}^d$ be a polytope with dim(aff(P)) \geq 1. The vertex-facet distance of P is

$$\operatorname{vf}(P) := \min_{F \in \operatorname{facets}(P)} \operatorname{dist}(\operatorname{aff}(F), \operatorname{vertices}(P) \setminus F).$$

2.8.4 Relation between vertex-facet distance and pyramidal width

We show $\operatorname{vf}(\operatorname{conv}(A)) \geq \operatorname{PWidth}(A)$. It seems that this result may have already been know to [PnR19, comment before Theorem 1, combined with Theorem 2], but it is claimed there in the wrong direction. That direction is impossible as the example of a unit cube shows: $\operatorname{PWidth}([0,1]^d) = 1/\sqrt{d}$ [LJJ15, Lemma 4], but $\operatorname{vf}([0,1]^d) = 1$.

Proposition 2.22. Let $A \subseteq \mathbb{R}^d$ be a finite set with at least two points. Then $\operatorname{vf}(\operatorname{conv}(A)) \geq \operatorname{PWidth}(A)$.




Figure 1: Proof of Proposition 2.22

Proof. Let $P = \operatorname{conv}(A)$. Let F be a facet of P and pick $v \in \operatorname{vertices}(P) \setminus F$ so that $\operatorname{dist}(v, \operatorname{aff}(F)) = \epsilon := \operatorname{vf}(P)$. Pick $x \in \operatorname{relint}(\operatorname{conv}(F \cup \{v\}))$ and let r be the unit outer normal vector to F (in $\operatorname{aff}(P)$ if P is not full-dimensional). We set K = P in Definition 2.20 so that $r \in \operatorname{cone}(K - x) = \operatorname{aff}(P)$ and $\operatorname{PWidth}(A) \leq \operatorname{PDirW}(K \cap A, r, x) = \operatorname{PDirW}(A, r, x)$. Now, set $S = A \cap (F \cup \{v\})$ in Definition 2.18 so that, with these choices, $\operatorname{PDirW}(A, r, x) \leq \operatorname{dirW}(S, r) \leq \epsilon$. The claim follows. \square

2.8.5 Facial distance

Definition 2.23 ([PnR19]). Let $C \subseteq \mathbb{R}^d$ be a polytope with $\dim(\operatorname{aff}(C)) \geq 1$. The facial distance of C is

$$\Phi(C) := \min_{\substack{F \in \text{faces}(C) \\ \emptyset \subseteq F \subseteq C}} d(F, \text{conv}(\text{vertices}(C) \setminus F)).$$

2.8.6 Relation between facial distance and pyramidal width

One of the motivations of [PnR19] to introduce parameter Φ is that it is the same as PWidth (except in degenerate cases) while the definition of Φ is simpler to use in many cases. We quote their result next.

Theorem 2.24 ([PnR19, Theorem 2]). Let $A \subseteq \mathbb{R}^d$ be a finite set with at least two points. Then $\Phi(\text{conv}(A)) = \text{PWidth}(A)$.

2.8.7 Summary result

Theorem 2.25. Let $A \subseteq \mathbb{R}^d$ be a finite set with at least two points. Then

$$minwidth(A) < \Phi(conv(A)) = PWidth(A) < vf(conv(A)).$$

Proof. Immediate from [LJJ15, Section 3.1], Theorem 2.24 and Proposition 2.22. \Box

3 Conditioning of simplices

In this section we show that the smoothed conditioning of any simplex is polynomial. This implies that several FWMs have smoothed polynomial complexity on the minimum norm point in a simplex problem and the minimization of many convex functions on a simplex. To put this result in context, we first argue (based on known results) that even a simplex with vertices having 0–1 coordinates can have bad conditioning. Another relevant context to keep in mind is the fact that linear programming reduces in strongly polynomial time to the minimum norm point in a simplex [LHR20].

3.1 Equality of width and minwidth of a simplex

We start with the observation that the minwidth of a simplex is the same as its width.

Lemma 3.1. Let A be the vertex set of a simplex in \mathbb{R}^d and $A_0 \subset A$ which includes more than one vertex. Then width(A) \leq width(A₀). In particular, minwidth(A) = width(A).

Proof. We prove by induction in d. The width of a polytope is the minimum distance between parallel supporting hyperplanes in its affine hull. Width of a 2-simplex is the minimum height of triangle, which is smaller than the length of any edge. For a k-simplex A, suppose the width of one of its facet is given by the distance between two parallel (k-2)-dimensional planes, p_1^{k-2} and p_2^{k-2} . One can extend p_1^{k-2} and p_2^{k-2} to parallel hyperplanes in \mathbb{R}^k that enclose A. Suppose extensions p_1^{k-1} and p_2^{k-1} give the minimum distance. Then,

$$\operatorname{dist} \left(p_1^{k-1}, p_2^{k-1} \right) = \min_{a \in p_1^{k-1}, b \in p_2^{k-1}} \|a - b\| \leq \min_{a \in p_1^{k-2}, b \in p_2^{k-2}} \|a - b\| = \operatorname{dist} \left(p_1^{k-2}, p_2^{k-2} \right)$$

which shows that the width of a k-simplex is less than the width of any of its facets. The claim then follows by induction.

3.2 Bad worst case conditioning of a 0–1 simplex

Lacoste-Julien and Jaggi [LJJ15] observed that the minwidth of the unit cube in \mathbb{R}^d is exponentially small in d. This example was one of their motivations for introducing PWidth, which is $1/\sqrt{d}$ for the cube. Their observation is based on the following result by Alon and Vu:

Theorem 3.2 ([AV97, Theorem 3.2.2], [Zie00, Corollary 27]). There are d + 1 vectors in $\{0, 1\}^d$ that form the vertices of a d-dimensional simplex S so that

$$\frac{2^{d-1}}{d^{d/2}} \le \operatorname{vf}(S) \le \frac{2^{d(2+o(1))}}{d^{d/2}}.$$

[LHR20] observed that PWidth can be exponentially small in the size (bitlength) of a set of points with integer coordinates. Using Theorem 3.2 and the relationships between polytope condition measures, we can immediately strengthen this result and show that this is not just a "large numbers" phenomenon, namely, all condition measures are exponentially small even for a 0–1 simplex:

Corollary 3.3. There are d+1 vectors in $\{0,1\}^d$ that form the vertices of a d-dimensional simplex S so that width(vertices(S)) = minwidth(vertices(S)) \leq PWidth(vertices(S)) = $\Phi(S) \leq$ vf $(S) \leq \frac{2^{d(2+o(1))}}{d^{d/2}}$.

Proof. Let S be the d-dimensional simplex given by Theorem 3.2. Lemma 3.1 gives the leftmost equality. The rightmost inequality is one of the conclusions of Theorem 3.2. The other relations follow from Theorem 2.25.

3.3 Polynomial smoothed complexity of FWMs on a simplex

Now we start analyzing smoothed complexity of FWMs on the minimization of a strongly convex function with Lipschitz gradient on a simplex.

Definition 3.4. A differentiable function f is said to have L-Lipschitz gradient if for some L > 0 and for all x, y in its domain we have $\|\nabla f(x) - \nabla f(y)\| \le L\|x - y\|$.

Definition 3.5. A differentiable function f is μ -strongly convex if for some $\mu > 0$ and for all x, y in its domain, we have

$$f(y) \ge f(x) + \nabla f(x)^T (y - x) + \frac{\mu}{2} ||y - x||^2.$$

In [LJJ15, Theorem 1], Lacoste-Julien and Jaggi proved the global linear convergence of FWMs on the minimization of a strongly convex function with Lipschitz gradient: suppose u_t is the current point after t good iterations², $f(u_t)$ satisfies

$$f(u_t) - f^* \le \left(1 - \frac{\mu}{4L} \left(\frac{\text{PWidth}(A)}{\text{diam}(A)}\right)^2\right)^t (f(u_0) - f^*),\tag{3}$$

where f^* is the optimal value and u_0 is the initial point. To show polynomial smoothed complexity, we need to prove that the measure of conditioning $\kappa = \frac{\text{PWidth}(A)}{\text{diam}(A)}$ is at least inverse polynomial in $d, 1/\sigma, 1/\delta$. We are going to get this by giving a polynomial lower bound on PWidth(A) and a polynomial upper bound on diam(A).

3.3.1 Inverse polynomial smoothed minwidth

We know from Theorem 2.25 that minwidth \leq PWidth, and from Lemma 3.1 that minwidth = width for any simplex. Thus, we instead find a lower bound on width, namely the diameter of a ball contained in the simplex, which is also a lower bound on PWidth. In the next lemma, we prove that a random simplex contains a ball of radius $\Omega(d^{-2})$ with probability close to 1.

Lemma 3.6. Let $A = \{A_1, \ldots, A_{d+1}\}$ be a set of independent Gaussian random vectors with means μ_i , $\|\mu_i\| \le 1$, $i \in [d+1]$, and covariance matrix $\sigma^2 \mathbf{I}_d$. Then for $\delta > 0$,

$$\mathbb{P}\left(\operatorname{minwidth}(\operatorname{conv}(A)) \ge \sqrt{2\pi}\sigma\delta(d+1)^{-2}\right) \ge 1 - \delta.$$

Moreover,

$$\mathbb{P}\left(\operatorname{PWidth}(\operatorname{conv}(A)) \ge \sqrt{2\pi}\sigma\delta(d+1)^{-2}\right) \ge 1 - \delta.$$

²The number of good iterations depends on variants of FWMs being used. It is always lower bounded by some linear function of the actual number of iterations. See details in [LJJ15, Theorem 1].

Proof. It is easy to see that A forms a simplex with probability 1. From Lemma 3.1, we know the minwidth of a simplex is its width. Let D_i be the distance from A_i to the affine hull of its opposite facet, aff $\{A_j: j \neq i\}$. Conditioning on aff $\{A_j: j \neq i\}$, by the rotational invariance of Gaussian distribution, D_i is equal in distribution to the absolute value of a Gaussian random variable with mean $\mu \in \mathbb{R}$ (not necessarily be zero) and variance σ^2 . Let $X \sim \mathcal{N}(0, \sigma^2)$. By Lemma 2.4, we have $\mathbb{P}(D_i < t) \leq \mathbb{P}(\|X\| < t)$ for all t. The right hand side is upper bounded by $2t/\sqrt{2\pi}\sigma$, which is the product of maximal Gaussian density and length of interval. Apply union bound to get

$$\mathbb{P}\left(\bigcap_{i=1}^{d+1} \{D_i \ge t\}\right) \ge 1 - \frac{2t(d+1)}{\sqrt{2\pi}\sigma}.$$

Let C_i be the distance between the center of mass of conv(A) and $aff(A_j: j \neq i)$. Note that $C_i = D_i/(d+1)$. Then

$$\mathbb{P}\left(\bigcap_{i=1}^{d+1} \{C_i \ge \frac{t}{d+1}\}\right) \ge 1 - \frac{2t(d+1)}{\sqrt{2\pi}\sigma}.$$

The above expression states that with some probability the ball centered at the center of mass and of radius t/(d+1) lies inside $\operatorname{conv}(A)$. Setting $t = \frac{\delta\sigma\sqrt{\pi}}{\sqrt{2}(d+1)}$ and using the fact that the width of the simplex is at least the diameter of the inscribed ball, we get

$$\mathbb{P}\left(\operatorname{width}\left(\operatorname{conv}(A)\right) \ge \sqrt{2\pi}\sigma\delta(d+1)^{-2}\right) \ge 1 - \delta.$$

The claim follows immediately from Lemma 3.1 and Theorem 2.25.

3.3.2 Smoothed diameter

Lemma 3.7. Let $A = \{A_1, \ldots, A_{d+1}\}$ be a set of independent Gaussian random vectors with means μ_i , $\|\mu_i\| \le 1$, $i \in [d+1]$, and covariance matrix $\sigma^2 \mathbf{I}_d$. Then for $\delta > 0$,

$$\mathbb{P}\left(\operatorname{diam}(A) \le 2\left(\sigma\sqrt{2d+3\ln\left(\frac{d+1}{\delta}\right)}+1\right)\right) \ge 1-\delta.$$

Proof. Let $A_i = \mu_i + X_i$, where $X_i \sim \mathcal{N}(0, \sigma^2 \mathbf{I}_d)$. Let t > 0. Triangle inequality gives that $\mathbb{P}(\|A_i\| > t+1) = \mathbb{P}(\|X_i + \mu_i\| > t+1) \leq \mathbb{P}(\|X_i\| > t)$. Apply Lemma 2.5 with $\alpha = (\sigma^2, \dots, \sigma^2)$, we have

$$\mathbb{P}\left(\|A_i\| > \sigma\sqrt{d+2\sqrt{dt}+2t}+1\right) \le \mathbb{P}\left(\|X_i\| \ge \sigma\sqrt{d+2\sqrt{dt}+2t}\right) \le e^{-t},$$

which shows that every A_i is contained in a ball of radius $\sigma \sqrt{d+2\sqrt{dt}+2t}+1 \le \sigma \sqrt{2d+3t}+1$ with high probability. With union bound, we see the diameter of the ball is an upper bound of the diameter of convex hull of A:

$$\mathbb{P}\Big(\operatorname{diam}(\operatorname{conv}(A)) \le 2(\sigma\sqrt{2d+3t}+1)\Big) \ge 1 - (d+1)e^{-t}.$$

The claim then follows by setting $t = \ln((d+1)/\delta)$.

Next we restate and prove our main theorem for this section:

Theorem 1.2. Let $A = \{A_1, \ldots, A_{d+1}\}$ be a set of independent Gaussian random vectors with means μ_i , $\|\mu_i\| \leq 1, i \in [d+1]$, and covariance matrix $\sigma^2 \mathbf{I}_d$. Then for $\delta > 0$, with probability at least $1 - \delta$, the measure of conditioning $\kappa = \frac{\mathrm{PWidth}(A)}{\mathrm{diam}(A)}$ of A is at least some inverse polynomial in d, $1/\sigma$ and $1/\delta$.

Proof. We proved in Lemma 3.6 and Lemma 3.7 that,

$$\mathbb{P}\left(\operatorname{PWidth}(\operatorname{conv}(A)) \ge \sqrt{2\pi}\sigma\delta(d+1)^{-2}\right) \ge 1 - \delta.$$

and

$$\mathbb{P}\left(\operatorname{diam}(A) \le 2\left(\sigma\sqrt{2d+3\ln\left(\frac{d+1}{\delta}\right)}+1\right)\right) \ge 1-\delta.$$

Thus with probability at least $1-2\delta$, we have

$$\frac{\text{PWidth}(A)}{\text{diam}(A)} \ge \frac{\sqrt{2\pi}\sigma\delta(d+1)^{-2}}{2\left(\sigma\sqrt{2d+3\ln\left(\frac{d+1}{\delta}\right)}+1\right)}$$
$$= \frac{\delta\sqrt{\pi/2}}{(d+1)^2\left(\sqrt{2d+3\ln\left(\frac{d+1}{\delta}\right)}+\frac{1}{\sigma}\right)}$$
$$\ge 1/\rho(d,1/\sigma,1/\delta)$$

where ρ is a polynomial function of $d, 1/\sigma, 1/\delta$.

Going back to (3), let $h_t = f(u_t) - f^*$. We have

$$h_t \le \left(1 - \frac{\mu}{4L} \left(\frac{\operatorname{PWidth}(A)}{\operatorname{diam}(A)}\right)^2\right)^t h_0.$$

Based on our smoothed analysis on the measure of conditioning in Theorem 1.2, with probability at least $1-2\delta$

$$h_t \le \left(1 - \frac{\mu}{4L\rho^2}\right)^t h_0 \le e^{-\frac{\mu t}{4L\rho^2}} h_0.$$

Hence one needs at most $\frac{4L\rho^2\ln(\frac{1}{\epsilon})}{\mu}$ good iterations to get a solution whose value is within distance $\epsilon(f_0-f^*)$ of f^* . Let T denote the number of good iterations, we have (using the notation from Definition 1.1)

$$\max_{\substack{A \subseteq B(0,1) \subseteq \mathbb{R}^d \\ |A| = d+1}} \mathbb{P}_g \left(T(A+g) \ge \frac{4L\rho(d, \frac{1}{\sigma}, \frac{1}{\delta})^2 \ln(\frac{1}{\epsilon})}{\mu} \right) \le 2\delta.$$

4 Conditioning of random matrices

In this section we prove that the smallest singular value of some square submatrix of a d-by-n Gaussian random matrix is exponentially small with probability exponentially close to 1 when $n/d \ge c > 1$. From Lemma 2.8, we know that the smallest singular value of a square matrix is comparable to the minimum distance between one column vector and the span of the other column vectors (one-off-distance). If we consider exponentially narrow bands around each span of d-1 column vectors of a rectangular matrix, the matrix will have exponentially small minimum singular value if some other column vector falls in one of those bands. We lower bound the Gaussian measure of the union of bands by a constant using the first two terms of the inclusion-exclusion principle (Bonferroni inequality). See Section 1 for a high level overview of the proof.

We start by giving an upper bound of the intersection of two bands in Gaussian measure, which appears in the second term of the inclusion-exclusion principle. The following lemma shows that the Gaussian measure of the intersection depends on the width of bands and the angle between two bands

Lemma 4.1. Let $u, v \in \mathbb{R}^d$ be unit length vectors, let $\epsilon > 0$, and let $c_S, c_T \in \mathbb{R}$. Let

$$\mathcal{B}_S = \{ x \in \mathbb{R}^d : c_S \le x \cdot u \le c_S + \epsilon \},$$

$$\mathcal{B}_T = \{ x \in \mathbb{R}^d : c_T \le x \cdot v \le c_T + \epsilon \}.$$

Then

$$\mathcal{G}(\mathcal{B}_S \cap \mathcal{B}_T) \le \frac{\epsilon^2}{\sqrt{2\pi(1-(u\cdot v)^2)}}.$$

Proof. If u and v are parallel then the claim holds. If they are not parallel, then by the structure of the Gaussian measure \mathcal{G} this is a two-dimensional problem in the plane spanned by u,v. Identify this plane with \mathbb{R}^2 . $\mathcal{G}(\mathcal{B}_S \cap \mathcal{B}_T)$ is at most the maximum density $1/\sqrt{2\pi}$ multiplied by the area of the parallelogram $P' := \{x \in \mathbb{R}^2 : c_S \leq x \cdot u \leq c_S + \epsilon, c_T \leq x \cdot u \leq c_T + \epsilon\}$. One can see that P' has the same area as $P := \{x \in \mathbb{R}^2 : |x \cdot u| \leq \epsilon/2, |x \cdot v| \leq \epsilon/2\}$. Defining A to be the matrix with rows u, v, we have $P = \{x : ||Ax||_{\infty} \leq \epsilon/2\}$. This implies $\operatorname{area}(P) = \epsilon^2 |\det A^{-1}| = \epsilon^2/|\det A| = \epsilon^2/\sqrt{\det AA^T} = \epsilon^2/\sqrt{1 - (u \cdot v)^2}$. The claim follows.

We now switch our focus to the random regime. The following lemma gives a probabilistic upper bound of the intersection of two bands around the spans of two (possibly not disjoint) subsets of random vectors in high dimensional space. The bound is good when not too many points are shared by the subsets (so that the behavior is not very different from two independent bands).

Lemma 4.2. Let $d \ge 1$. Let $0 \le k \le d-1$. Let $A_1, \ldots, A_k, S_1, \ldots, S_{d-k-1}, T_1, \ldots, T_{d-k-1}$ be d-dimensional iid. standard Gaussian random vectors. Let

$$\mathcal{B}_S = (\text{span}\{A_1, \dots, A_k, S_1, \dots, S_{d-k-1}\})_{\epsilon/2},$$

$$\mathcal{B}_T = (\text{span}\{A_1, \dots, A_k, T_1, \dots, T_{d-k-1}\})_{\epsilon/2}.$$

Then for any $t \geq 1$,

$$\mathbb{P}\left(\mathcal{G}(\mathcal{B}_S \cap \mathcal{B}_T) \ge \frac{\epsilon^2 t}{\sqrt{2\pi}}\right) \le \frac{1}{t^{d-k-2}}.$$

Proof. If $d \leq 2$ or $k \geq d-2$, then the claim is immediate. Otherwise, $0 \leq k \leq d-3$ and we argue in the following way: By the structure of the Gaussian measure \mathcal{G} this is a (d-k)-dimensional problem in $\{A_1,\ldots,A_k\}^{\perp}$. More precisely, let U,V be two (d-k)-dimensional iid. uniformly random unitlength vectors and define $\mathcal{B}'_S = \{x \in \mathbb{R}^{d-k} : |x \cdot U| \leq \epsilon/2\}$ and $\mathcal{B}'_T = \{x \in \mathbb{R}^{d-k} : |x \cdot V| \leq \epsilon/2\}$. Then $\mathcal{G}(\mathcal{B}_S \cap \mathcal{B}_T)$ has the same distribution as $\mathcal{G}(\mathcal{B}'_S \cap \mathcal{B}'_T)$. From Lemma 4.1 we have $\mathcal{G}(\mathcal{B}'_S \cap \mathcal{B}'_T) \leq \frac{\epsilon^2}{\sqrt{2\pi(1-(U \cdot V)^2)}}$.

Using the rotational symmetry of the distribution of U and V and then Lemma 2.7 we get

$$\mathbb{P}(\sqrt{(1 - (U \cdot V)^2)} \le 1/t) = \mathbb{P}\left(\sqrt{U_1^2 + \dots + U_{d-k-1}^2} \le 1/t\right)$$
$$\le \mathbb{P}\left(\sqrt{U_1^2 + \dots + U_{d-k-2}^2} \le 1/t\right)$$
$$= 1/t^{d-k-2}.$$

The claim follows. \Box

The main technical content of our singular value bound is the following lower bound on the Gaussian volume of the union of bands around any d-1 columns of a d-by-n Gaussian random matrix. We also include an upper bound on the volume.

Lemma 4.3. Let $\epsilon \geq 0$, $d \geq 2$. For $\{A_1, \ldots, A_n\} \subseteq \mathbb{R}^d$, define

$$V = \mathcal{G}\left(\left(\bigcup_{S\subseteq[n],|S|=d-1}\operatorname{span} A_S\right)_{\epsilon}\right).$$

- 1. $V \leq \frac{2\epsilon}{\sqrt{2\pi}} \binom{n}{d-1}$.
- 2. Suppose A_1, \ldots, A_n are d-dimensional iid. standard Gaussian random vectors with $\frac{n}{d-1} \geq c_0 > 1$. Then there exist constants $c_2, c_4 > 1$ (that depend only on c_0) such that when $\epsilon \leq 1/(c_4c_2^{d-1})$ and with probability at least $1 c_4e^{-d}$ we have $V \geq \frac{c_2^{d-1}}{\sqrt{2\pi}}\epsilon$.

Proof of part 1. The upper bound follows from the union bound and the fact that the 1-dimensional Gaussian density is upper bounded by $1/\sqrt{2\pi}$.

Proof of part 2. Let $S = \{S \subseteq [n], |S| = d - 1\}$. Use Lemma 2.6 to get the bound $A(n, c_1(d - 1), d - 1) \ge c_2^{d-1}$. We get a subfamily $\mathcal{T} \subseteq S$ such that for all $S, T \in \mathcal{T}$ with $S \ne T$ we have $|S \cap T| \le (1 - \frac{c_1}{2})(d - 1)$ and $|\mathcal{T}| = c_2^{d-1}$ for some constants $0 < c_1 < 1, c_2 > 1$ (that depend only on c_0), and any $d \ge 2$. Let $N = |\mathcal{T}|$.

Let $\mathcal{B}_S = (\operatorname{span} A_S)_{\epsilon}$. Use the first two terms of the inclusion-exclusion principle (Bonferroni inequality) and use Lemma 4.2 in a union bound applied to all pairs of sets in \mathcal{T} to get $\mathcal{G}(\mathcal{B}_S \cap \mathcal{B}_T) \leq \frac{4\epsilon^2 t}{\sqrt{2\pi}}$ for all $S, T \in \mathcal{T}, S \neq T$. We get a bound on V that holds with probability at least $1 - \frac{\binom{N}{2}}{t^{d-1-(1-c_1/2)(d-1)-2}} = 1 - \frac{\binom{N}{2}}{t^{c_1(d-1)/2-1}} \geq 1 - \frac{N^2}{t^{c_1(d-1)/2-1}} = 1 - t \left(\frac{c_2^2}{t^{c_1/2}}\right)^{d-1} = 1 - c_3 e^{-d}$ (choosing a constant t > 1 that depends on $c_1(c_0)$ and $c_2(c_0)$ such that $c_2^2/t^{c_1/2} = 1/e$ and then setting $c_3 = t/e$,

which ultimately depends only on c_0). The bound on V is

$$V \ge \mathcal{G}\left(\left(\bigcup_{S \in \mathcal{T}} \operatorname{span} A_S\right)_{\epsilon}\right)$$

$$\ge \sum_{S \in \mathcal{T}} \mathcal{G}(\mathcal{B}_S) - \frac{1}{2} \sum_{S,T \in \mathcal{T},S \ne T} \mathcal{G}(\mathcal{B}_S \cap \mathcal{B}_T)$$

$$\ge \frac{2N\epsilon}{\sqrt{2\pi}} e^{-\epsilon^2/2} - \binom{N}{2} \frac{4\epsilon^2 t}{\sqrt{2\pi}}$$

$$\ge \frac{2N\epsilon}{\sqrt{2\pi}} (e^{-\epsilon^2/2} - 2tN\epsilon)$$

$$\ge \frac{2N\epsilon}{\sqrt{2\pi}} (1 - \epsilon^2/2 - 2tN\epsilon)$$

$$\ge \frac{N\epsilon}{\sqrt{2\pi}} \qquad \text{(for } \epsilon \le 1/(8tN)).$$

In other words, $V \geq \frac{c_2^{d-1}\epsilon}{\sqrt{2\pi}}$ for $\epsilon \leq 1/(8ec_3c_2^{d-1})$. We finish our proof by taking $c_4 = 8ec_3$.

We are ready now to restate and prove the main results of the section.

Theorem 1.4. Let A be an d-by-n random matrix with iid. standard Gaussian entries with $d \ge 2$ and $\frac{n}{d} \ge c_0 > 1$. Then, there exist constants $c_2, c_4 > 1$, $0 < c_6 < 1$ (that depend only on c_0) such that with probability at least $1 - 2c_4c_6^d$,

$$\min_{S\subseteq[n],|S|=d} \sigma_d(A_S) \le \frac{1}{c_4 c_2^{d-1}}.$$

Proof. Pick $c_1 \in (1, c_0)$. Let $m = \lfloor c_1 d \rfloor$. Note that $m \geq c_1 d - 1 \geq c_1 d - c_1 \geq c_1 (d - 1)$, so that we can apply Lemma 4.3 to columns A_1, \ldots, A_m with $\epsilon = 1/c_4 c_2^{d-1}$. Then we get $V \geq \frac{1}{\sqrt{2\pi}c_4}$ with probability greater than $1 - c_4 e^{-d}$. This implies that with probability greater than

$$(1 - c_4 e^{-d}) \left(1 - \left(1 - \frac{1}{\sqrt{2\pi}c_4} \right)^{n-m} \right) \ge \left(1 - c_4 e^{-d} \right) \left(1 - \left(1 - \frac{1}{\sqrt{2\pi}c_4} \right)^{(c_0 - c_1)d} \right)$$

$$\ge 1 - c_4 e^{-d} - \left(1 - \frac{1}{\sqrt{2\pi}c_4} \right)^{(c_0 - c_1)d}$$

$$\ge 1 - 2c_4 c_6^d$$

where $c_6 = \max\{1/e, (1-\frac{1}{\sqrt{2\pi}c_4})^{(c_0-c_1)}\}$, at least one of A_{m+1}, \ldots, A_n , say A_* , falls in V, that is, falls within distance $\epsilon = 1/c_4c_2^{d-1}$ of $\mathrm{span}(A_S)$ for some $S \subseteq [m], |S| = d-1$. Lemma 2.8 gives $\sigma_d(A_S, A_*) \leq 1/c_4c_2^{d-1}$.

Theorem 1.5. Let A be an d-by-n random matrix with iid. standard Gaussian entries with $d \ge 2$ and $1 < \frac{n}{d-1} \le C_0$. Then, there exist constants $C_1 > 1$, $0 < C_2 < 1$ (that depend only on C_0) such that with probability at least $1 - nC_2^{d-1}$,

$$\min_{S\subseteq[n],|S|=d}\sigma_d(A_S)\geq \frac{1}{C_1^{d-1}}.$$

Proof. Apply Lemma 4.3 to columns A_1, \ldots, A_{n-1} to get

$$V \le \frac{2\epsilon}{\sqrt{2\pi}} \binom{n}{d-1} \le \frac{2\epsilon}{\sqrt{2\pi}} \left(\frac{en}{d-1}\right)^{d-1} \le \frac{2\epsilon}{\sqrt{2\pi}} (eC_0)^{d-1}.$$

By picking $\epsilon = 1/C_1^{d-1}$ where $C_1 > eC_0$, there exists a constant $\frac{eC_0}{C_1} < C_2 < 1$ such that $V \le C_2^{d-1}$. This implies that, with probability at most C_2^{d-1} , column A_n is within distance $1/C_1^{d-1}$ of span A_S for some $S \subseteq [n-1], |S| = d-1$. A similar claim holds for columns A_1, \ldots, A_{n-1} as well. Applying the union bound, we get that no A_i falls within distance $1/C_1^{d-1}$ of span A_S for any $S \subseteq [n-1], |S| = d-1$ with probability at least $1 - nC_2^{d-1}$. Lemma 2.8 gives $\sigma_d(A_S, A_n) \ge 1/C_1^{d-1}$ with probability at least $1 - nC_2^{d-1}$.

5 On the stability of tensor decomposition

Kruskal [Kru77] showed a sufficient condition under which the component vectors $a_i, b_i, c_i, i = 1, \ldots, n$ of an order-3 tensor $T = \sum_{i=1}^n a_i \otimes b_i \otimes c_i$ are uniquely determined by the tensor (up to inherent ambiguities). The condition depends on a parameter now known as the Kruskal rank of a matrix: For a d-by-n matrix A, the Kruskal rank of A, denoted K-rank(A), is the maximum $r \in [n]$ such that any r columns of A are linearly independent. The condition is K-rank(A) + K-rank(B) + K-rank $(C) \geq 2n + 2$, where A, B, C are the matrices with columns (a_i) , (b_i) , (c_i) , respectively. For concreteness, it is helpful to consider the symmetric case $A = B = C \in \mathbb{R}^{d \times n}$. Kruskal's condition becomes 3 K-rank $(A) \geq 2n + 2$. Informally, for a generic matrix A we have K-rank(A) = d and so Kruskal's result guarantees uniqueness for generic A when $n \leq 3d/2 - 1$.

Bhaskara, Charikar and Vijayaraghavan [BCV14, Theorem 5] extended Kruskal's uniqueness to a result that guarantees robust decomposition. That is, when the observed tensor is a small perturbation of the original tensor, the components of the perturbed tensor are uniquely determined and close to the components of the original tensor. Their condition for robust unique decomposition is a refinement of Kruskal's condition: Let $\tau > 0$. The robust Kruskal rank (with threshold τ) of A, denoted K-rank $_{\tau}(A)$, is the maximum $k \in [n]$ such that for any subset $S \subseteq [n]$ of size k we have $\sigma_k(A_S) \geq 1/\tau$ (σ_k denotes the kth largest singular value). The condition is K-rank $_{\tau}(A)$ + K-rank $_{\tau}(B)$ +K-rank $_{\tau}(C) \geq 2n+2$ and the error in the recovered components depends polynomially on τ .

In this context, Theorem 1.4 can be stated in the following equivalent way:

Theorem 5.1. Let A be an d-by-n random matrix with iid. standard Gaussian entries with $d \ge 2$ and $n/d \ge c_0 > 1$. Then, there exist constants $c_4, c_5 > 1$, $0 < c_6 < 1$ (that depend only on c_0) such that with probability at least $1 - 2c_4c_6^d$,

$$K$$
-rank $_{\tau}(A) = d \Rightarrow \tau \geq c_4 c_5^{d-1}$.

This has the following implication for Bhaskara, Charikar and Vijayaraghavan's result: Even though Kruskal's result guarantees uniqueness for generic A when n = 3d/2-1 (say, with probability 1 for a random Gaussian matrix we have K-rank(A) = d), Bhaskara, Charikar and Vijayaraghavan's robust uniqueness can give a polynomial bound on the reconstruction error on no more than an exponentially small fraction of matrices A when the fraction is measured by the Gaussian measure. This rarity of sufficiently well-conditioned matrices A is somewhat surprising.

6 On the complexity of the simplex method and the diameter of polytopes

In [BR13], Brunsch and Röglin introduced the following property of a matrix:

Definition 6.1 (δ -distance property, [BGR15]). Let $A = (a_1, \ldots, a_m)^{\top}$ be an m-by-n matrix with unit rows. We say that A satisfies the δ -distance property if: for any $I \subseteq [m]$ and any $j \in [m]$ whenever $a_i \notin \operatorname{span}\{a_i : i \in I\}$ we have $d(a_i, \operatorname{span}\{a_i : i \in I\}) \geq \delta$.

This property has been used in several papers [BR13, BGR15, DH16, EV17] to study polytopes of the form $\{x \in \mathbb{R}^n : Ax \leq b\}$ to provide upper bounds of the form $\mathrm{poly}(m,n,1/\delta)$ on their diameter and the number of pivot steps of the simplex method. Our Theorem 1.4 combined with Lemma 2.8 and concentration of the length of a Gaussian random vector implies that, for $m/n \geq c' > 1$, matrices A with the δ -distance property for $\delta \geq c^n$, 0 < c < 1, are "rare": they are exponentially unlikely when the rows are iid. random unit vectors. As in Section 5, this rarity of well-conditioned matrices A is somewhat surprising.

7 On the smoothed analysis of polytope conditioning

In this section we prove that the vertex-facet distance of the convex hull of a linear number of d-dimensional iid. Gaussian points can be exponentially small with probability at least some constant. The argument is a more elaborate version of the argument for the minimum singular value in Section 4 and works in the following way. Figure 2 shows a polytope, the convex hull of a partial sequence of random points, and ϵ -innner bands at all facets. If a new point falls into the blue region, then the new polytope, which is the convex hull of the old polytope plus the new point, will have vertex-facet distance no larger than ϵ : the new point is a vertex and its distance to the affine hull of the facet associated to the band where the point lies in is less than ϵ .

To get a lower bound on the Gaussian measure of the blue region (Lemma 7.5), we add the measures of the bands and then we subtract the measures of pairwise intersections of bands and ϵ -inner neighbourhood (grey region). Lemma 7.4 gives a bound on the measure of a pairwise intersection. Its proof is divided into two cases: Lemma 7.1 for the case where the two facets do not share vertices and Lemma 7.2 for the case where they do share vertices. This argument is a refinement of the proof of Lemma 4.2.




Figure 2: A polytope (triangle) and the region (blue) where a new point would create a small vertex-facet distance.

Lemma 7.1. Let $S_1, \ldots, S_d, T_1, \ldots, T_d$ be iid. standard Gaussian random vectors in \mathbb{R}^d . Let

$$\mathcal{B}_S = \left(\operatorname{aff} \{ S_1, \dots, S_d \} \right)_{\epsilon/2},$$

$$\mathcal{B}_T = \left(\operatorname{aff} \{ T_1, \dots, T_d \} \right)_{\epsilon/2}.$$

Then for $t \geq 1$

$$\mathbb{P}\left(\mathcal{G}(\mathcal{B}_S \cap \mathcal{B}_T) \ge \frac{\epsilon^2 t}{\sqrt{2\pi}}\right) \le \frac{1}{t^{d-2}}.$$

Proof. By the rotational invariance of the Gaussian distribution, unit normal vectors U, V to $\mathcal{B}_S, \mathcal{B}_T$ are independent and are uniformly distributed on \mathcal{S}^{d-1} . Define

$$\mathcal{B}'_S = \{ x \in \mathbb{R}^d : |x \cdot U| \le \epsilon/2 \},$$

$$\mathcal{B}'_T = \{ x \in \mathbb{R}^d : |x \cdot V| \le \epsilon/2 \}.$$

By a standard argument (say, using logconcavity) we have $\mathbb{P}(\mathcal{G}(\mathcal{B}_S \cap \mathcal{B}_T) \geq t) \leq \mathbb{P}(\mathcal{G}(\mathcal{B}_S' \cap \mathcal{B}_T') \geq t)$. Then, by the argument in the proof of Lemma 4.2 we get that for any $t \geq 1$, $\mathbb{P}\left(\mathcal{G}(\mathcal{B}_S' \cap \mathcal{B}_T') \geq \frac{\epsilon^2 t}{\sqrt{2\pi}}\right) \leq \frac{1}{t^{d-2}}$. The claim follows.

Lemma 7.2. Let $A_1, \ldots, A_k, S_1, \ldots, S_{d-k}, T_1, \ldots, T_{d-k}$ be iid. standard Gaussian random vectors in \mathbb{R}^d , and $1 \leq k \leq d$. Let

$$\mathcal{B}_S = \left(\operatorname{aff} \{ A_1, \dots, A_k, S_1, \dots, S_{d-k} \} \right)_{\epsilon/2},$$

$$\mathcal{B}_T = \left(\operatorname{aff} \{ A_1, \dots, A_k, T_1, \dots, T_{d-k} \} \right)_{\epsilon/2}.$$

Then for $0 < 2\alpha \le \beta < \pi/2$,

$$\mathbb{P}\left(\mathcal{G}(\mathcal{B}_S \cap \mathcal{B}_T) \ge \frac{\epsilon^2}{\sqrt{2\pi}\sin\alpha}\right) \le (\sin\beta)^{d-k-1} + 2\left(\frac{\sin\alpha}{\sin(\beta - \alpha)}\right)^{d-k-2}.$$
 (4)

In particular, for $t > 2\pi$ we have

$$\mathbb{P}\left(\mathcal{G}(\mathcal{B}_S \cap \mathcal{B}_T) \ge \frac{\epsilon^2 t}{\sqrt{2\pi}}\right) \le 3\left(\frac{\pi^{3/2}}{\sqrt{2t}}\right)^{d-k-2}.$$

Proof. If $d-k \leq 2$, then the bound holds immediately. Otherwise, d-k > 2 and we argue in the following way. By the structure of the Gaussian measure, this reduces to a (d-k+1)-dimensional problem: Conditioning on $A_i = a_i$, i = 1, ..., k, we project onto the orthogonal complement of the linear subspace parallel to aff $\{a_1, ..., a_k\}$. We will then prove the bound claimed in (4) conditioning on $A_1, ..., A_k$, which implies the claimed bound by total probability.

With a slight abuse of notation, we denote the projection of $aff\{a_1,\ldots,a_k\}$ as a_1 and the projections of S_i, T_i as S_i, T_i , $i = 1, \ldots, d - k$. Using the fact that the Gaussian distribution is rotationally invariant, we may assume without loss of generality that $a_1 = \mu e_1$ for some $\mu \geq 0$. A

normal vector to aff $\{a_1, S_1, \dots, S_{d-k}\}$ is³

$$U = \det \begin{pmatrix} e_1 & e_2 & \cdots & e_{d-k+1} \\ & & P_1^T & \\ & & \vdots & \\ & & P_{d-k}^T & \end{pmatrix}, \tag{5}$$

where $P_i := S_i - a_1$, $i \in [d - k]$. Define the matrix $P = (P_1 \cdots P_{d-k})$. Let V be a normal vector to aff $\{a_1, T_1, \ldots, T_{d-k}\}$, defined similarly.

Set $\binom{H_i}{P_i'} = P_i$, where $H_i \sim \mathcal{N}(\mu, 1)$ and $P_i' \sim \mathcal{N}(0, \mathbf{I}_{d-k})$. Denote $H^T = (H_1 \cdots H_{d-k})$ as the first row of matrix P and $P' = (P_1' \cdots P_{d-k}')$ as the rest. H and P' are independent.

Note that $||U||^2 = \det(P^T P)$ (follows from (5) and the Cauchy-Binet formula). Also, $U_1 = U \cdot e_1 = \det(P')$. We now compute the distribution of the first coordinate of unit normal vector \hat{U} (using the *matrix determinant lemma* to compute the determinant of a rank-1 update).

$$\begin{split} \hat{U}_{1}^{2} &= \frac{\det(P'^{T}P')}{\det(P^{T}P)} \\ &= \frac{\det(P'^{T}P')}{\det(P'^{T}P' + HH^{T})} \\ &= \frac{\det(P'^{T}P' + HH^{T})}{(1 + H^{T}P'^{-1}P'^{-T}H)\det(P'^{T}P')} \\ &= \frac{1}{1 + H^{T}P'^{-1}P'^{-T}H}. \end{split}$$

Claim 7.3. We have $H^T P'^{-1} P'^{-T} H \stackrel{d}{=} \frac{\sum_{i=1}^{d-k} Y_i^2}{Y_0^2}$, where $Y_0 \sim \mathcal{N}(0,1)$, $Y_i \sim \mathcal{N}(\mu,1)$, $i \in [d-k]$ and Y_0, Y_1, \dots, Y_{d-k} are independent.

Proof of claim. Random variables P' and H are independent. Moreover, P' is a Gaussian matrix and therefore the distribution of P'^{-1} is invariant under any orthogonal transformation applied to rows or columns. Thus, it is enough to consider the case $H = \|H\|e_1$. Note that $\|H\|^2 \stackrel{d}{=} \sum_{i=1}^{d-k} Y_i^2$, and $e_1^T P'^{-1} P'^{-T} e_1 = \|\text{first row of } P'^{-1}\|^2 \stackrel{d}{=} \frac{1}{Y_0^2}$. The claim follows.

Recall that \hat{U}, \hat{V} are unit normal vectors to $\mathcal{B}_S, \mathcal{B}_T$, respectively. We aim to show that $\mathbb{P}(\hat{V} \in \mathcal{C}_{\alpha}(\hat{U}) \cup \mathcal{C}_{\alpha}(-\hat{U}))$, i.e. $\mathbb{P}(|\hat{U} \cdot \hat{V}| \geq \cos \alpha)$, is upper bounded by an expression of the form $c(\alpha)^d$ with $c(\alpha) \to 0$ as $\alpha \to 0$ (where $\mathcal{C}_{\alpha}(\hat{U})$ denotes the spherical cap centered at \hat{U} with angle α). To see this, we divide the analysis into two cases, depending on whether the cap is close to e_1 . The case analysis depends on a parameter β that will need to satisfy the constraint $\beta \geq 2\alpha$.

Case 1:
$$C_{\alpha}(\hat{U}) \subseteq C_{\beta}(e_1) \cup C_{\beta}(-e_1)$$
 (equivalently, $|\hat{U}_1| \ge \cos(\beta - \alpha)$).

 $^{^{3}}$ In the formula for U, the determinant should be interpreted as a formal cofactor expansion along the first row; the entries in the first row are the canonical vectors and the expansion gives the coefficients of these vectors (as subdeterminants).

In this case, the α -cap around \hat{U} is contained in a larger cap centered at e_1 .

$$\mathbb{P}\Big(\big\{\hat{V}\in\mathcal{C}_{\alpha}(\hat{U})\cup\mathcal{C}_{\alpha}(-\hat{U})\big\}\cap\big\{\mathcal{C}_{\alpha}(\hat{U})\subseteq\mathcal{C}_{\beta}(e_{1})\cup\mathcal{C}_{\beta}(-e_{1})\big\}\Big)\leq\mathbb{P}\big(\hat{V}\in\mathcal{C}_{\beta}(e_{1})\cup\mathcal{C}_{\beta}(-e_{1})\big)
(\text{using }\beta\leq\pi/2) = \mathbb{P}(\hat{V}_{1}^{2}\geq\cos^{2}\beta).$$
(6)

From claim 7.3 we get

$$\hat{V}_1^2 \stackrel{d}{=} \frac{Y_0^2}{Y_0^2 + \sum_{i=1}^n Y_i^2}.$$

To upper bound (6), we get from Lemma 2.3 that making $a_1 = 0$ (equivalently, $\mu = 0$) only makes the rhs larger and we then bound the case $a_1 = 0$ explicitly. More precisely, let W be a normal vector to span $\{T_1, \ldots, T_{d-k}\}$ defined similarly to U and V:

$$W = \det \begin{pmatrix} e_1 & e_2 & \cdots & e_{d-k+1} \\ & & T_1^T & \\ & & \vdots & \\ & & T_{d-k}^T & \end{pmatrix}.$$

Note that \hat{W} is a uniformly random unit vector. Following the same computation as for V, one can derive

$$\hat{W}_1^2 \stackrel{d}{=} \frac{X_0^2}{X_0^2 + \sum_{i=1}^n X_i^2},$$

where $X_0, X_i \sim \mathcal{N}(0,1), i \in [d-k]$. Then, by Lemma 2.3, $\mathbb{P}(\hat{V}_1^2 \geq \cos^2 \beta) \leq \mathbb{P}(\hat{W}_1^2 \geq \cos^2 \beta)$. Hence,

$$\mathbb{P}\Big(\big\{\hat{V} \in \mathcal{C}_{\alpha}(\hat{U}) \cup \mathcal{C}_{\alpha}(-\hat{U})\big\} \cap \big\{\mathcal{C}_{\alpha}(\hat{U}) \subseteq \mathcal{C}_{\beta}(e_{1}) \cup \mathcal{C}_{\beta}(-e_{1})\big\}\Big) \leq \mathbb{P}(\hat{W}_{1}^{2} \geq \cos^{2}\beta)$$

$$= \mathbb{P}\left(\sum_{i=2}^{d-k+1} \hat{W}_{i}^{2} \leq \sin^{2}\beta\right)$$

$$\leq \mathbb{P}\left(\sum_{i=2}^{d-k} \hat{W}_{i}^{2} \leq \sin^{2}\beta\right)$$

$$\leq (\sin\beta)^{d-k-1} \quad \text{(Lemma 2.7)}.$$

Case 2: $\mathcal{C}_{\alpha}(\hat{U}) \not\subseteq \mathcal{C}_{\beta}(e_1) \cup \mathcal{C}_{\beta}(-e_1)$.

If $C_{\alpha}(\hat{U})$ is not contained in $C_{\beta}(e_1) \cup C_{\beta}(-e_1)$, then \hat{U} makes an angle at least $\beta - \alpha$ with e_1 and $-e_1$, that is

$$|\hat{U}_1| < \cos(\beta - \alpha). \tag{7}$$

Our goal here is to bound

$$\mathbb{P}\Big(\big\{\hat{V}\in\mathcal{C}_{\alpha}(\hat{U})\cup\mathcal{C}_{\alpha}(-\hat{U})\big\}\cap\big\{\mathcal{C}_{\alpha}(\hat{U})\not\subseteq\mathcal{C}_{\beta}(e_{1})\cup\mathcal{C}_{\beta}(-e_{1})\big\}\Big) \\
=\mathbb{P}\Big(\big\{\hat{V}\in\mathcal{C}_{\alpha}(\hat{U})\big\}\cap\big\{\mathcal{C}_{\alpha}(\hat{U})\not\subseteq\mathcal{C}_{\beta}(e_{1})\cup\mathcal{C}_{\beta}(-e_{1})\big\}\Big) \\
+\mathbb{P}\Big(\big\{\hat{V}\in\mathcal{C}_{\alpha}(-\hat{U})\big\}\cap\big\{\mathcal{C}_{\alpha}(\hat{U})\not\subseteq\mathcal{C}_{\beta}(e_{1})\cup\mathcal{C}_{\beta}(-e_{1})\big\}\Big) \\
=2\cdot\mathbb{P}\Big(\big\{\hat{V}\in\mathcal{C}_{\alpha}(\hat{U})\big\}\cap\big\{\mathcal{C}_{\alpha}(\hat{U})\not\subseteq\mathcal{C}_{\beta}(e_{1})\cup\mathcal{C}_{\beta}(-e_{1})\big\}\Big). \tag{8}$$




Figure 3: Case 2 of proof of Lemma 7.4

Observe that the distribution of \hat{U} and the distribution of \hat{V} are invariant under rotations orthogonal to e_1 . Thus, if we let $\hat{U}_{-1}, \hat{V}_{-1}$ be the projections of \hat{U}, \hat{V} orthogonal to e_1 and $\widehat{U}_{-1}, \widehat{V}_{-1}$ be their normalizations, respectively, then $\widehat{U}_{-1}, \widehat{V}_{-1} \sim \text{Unif}(\mathcal{S}^{d-k})$.

This observation motivates us to use the corresponding probability of projections to bound (8). We will show that under condition (7) of case 2, $\hat{V} \in \mathcal{C}_{\alpha}(\hat{U})$ implies that $\widehat{V}_{-1} \in \mathcal{C}_{f(\alpha)}(\widehat{U}_{-1})$, where $f(\alpha)$ is a bound (to be understood) on the angle that depends only on α . As events,

$$\{\hat{V} \in \mathcal{C}_{\alpha}(\hat{U})\} \subseteq \{\hat{V}_{-1} \in \operatorname{Proj}_{e_{1}^{\perp}} \mathcal{C}_{\alpha}(\hat{U})\}$$

$$\subseteq \{\widehat{V}_{-1} \in \mathcal{C}_{f(\alpha)}(\widehat{U}_{-1})\}. \tag{9}$$

Bounding $f(\alpha)$ is a three-dimensional problem since $\widehat{U_{-1}}, \widehat{V_{-1}}$ are in span $\{e_1, \hat{U}, \hat{V}\}$. From now on, the analysis lives in the above three-dimensional space to get an upper bound on $f(\alpha)$. Let $\tilde{e}_2 = (\hat{U} - \hat{U} \cdot e_1)/\|\hat{U} - \hat{U} \cdot e_1\|$ (so that $\{e_1, \tilde{e}_2\}$ is an orthonormal basis of span $\{e_1, \hat{U}\}$). Let $\{e_1, \tilde{e}_2, \tilde{e}_3\}$ be an orthonormal basis of span $\{e_1, \hat{U}, \hat{V}\}$, and let $\hat{U} = (\hat{U}_1, \hat{U}_2, 0)$ be the coordinate tuple of \hat{U} relative to $\{e_1, \tilde{e}_2, \tilde{e}_3\}$. Consider $x \in \mathcal{C}_{\alpha}(\hat{U})$ such that $x \cdot \hat{U} = \cos \gamma$. Note that its coordinates (x_1, x_2, x_3) in our chosen basis satisfy the following system of equations:

$$x_1^2 + x_2^2 + x_3^2 = 1$$

$$x_1\hat{U}_1 + x_2\hat{U}_2 = \cos\gamma.$$

The projections of all such x (for fixed γ) onto span $\{\tilde{e}_2, \tilde{e}_3\}$ form the ellipse:

$$(x_2 - \hat{U}_2 \cos \gamma)^2 + x_3^2 \hat{U}_1^2 = \hat{U}_1^2 \sin^2 \gamma.$$

If $\hat{U}_1 = 0$, then $\hat{U}_2 = 1$, and the projection is the line segment inside unit circle at $x_2 = \cos \gamma$. The angle between x_{-1} and \widehat{U}_{-1} is upper bounded by γ . As γ ranges from 0 to α , \widehat{U}_{-1} and \widehat{V}_{-1} form an angle at most α when $\widehat{U}_1 = 0$.

If $\hat{U}_1 \neq 0$, the projection is an ellipse inside the unit circle. As shown in Fig. 3, angle between x_{-1} and \widehat{U}_{-1} can be upper bounded by angle formed by \widehat{U}_{-1} and tangent line $x_2 = \frac{\sqrt{\cos^2 \gamma - \hat{U}_1^2}}{\sin \gamma} x_3$. Note that from (7) we know $\hat{U}_1^2 < \cos^2(\beta - \alpha) \leq \cos^2 \alpha \leq \cos^2 \gamma$ (here we use $\beta \geq 2\alpha$ explicitly), so the tangent line always exists.

Hence the angle between x_{-1} and \widehat{U}_{-1} is at most $\arctan\left(\frac{\sin\gamma}{\sqrt{\cos^2\gamma-\hat{U}_1^2}}\right)$. Furthermore, since $\arctan\left(\frac{\sin\gamma}{\sqrt{\cos^2\gamma-\hat{U}_1^2}}\right)$ is increasing in γ , we can conclude that for any $\hat{V}\in\mathcal{C}_{\alpha}(\hat{U})$, its normalized projection orthogonal to e_1,\widehat{V}_{-1} , is contained in the spherical cap centered at \widehat{U}_{-1} with polar angle at most $\arctan\left(\frac{\sin\alpha}{\sqrt{\cos^2\alpha-\hat{U}_1^2}}\right)$ when $\hat{U}_1\neq 0$.

Therefore with (7), we can take

$$f(\alpha) = \max\{\arctan\left(\frac{\sin\alpha}{\sqrt{\cos^2\alpha - \cos^2(\beta - \alpha)}}\right), \alpha\}.$$

Combine with (8) and (9),

$$\mathbb{P}(\{\hat{V} \in \mathcal{C}_{\alpha}(\hat{U}) \cup \mathcal{C}_{\alpha}(-\hat{U})\} \cap \{\mathcal{C}_{\alpha}(\hat{U}) \not\subseteq \mathcal{C}_{\beta}(e_{1}) \cup \mathcal{C}_{\beta}(-e_{1})\}) \\
\leq 2 \cdot \mathbb{P}(\{\widehat{V}_{-1} \in \mathcal{C}_{f(\alpha)}(\widehat{U}_{-1})\} \cap \{\mathcal{C}_{\alpha}(\hat{U}) \not\subseteq \mathcal{C}_{\beta}(e_{1}) \cup \mathcal{C}_{\beta}(-e_{1})\}) \\
\leq 2 \cdot \mathbb{P}\left(|\widehat{U}_{-1} \cdot \widehat{V}_{-1}| \geq \cos(f(\alpha))\right) \\
= 2 \cdot \mathbb{P}\left(\sqrt{1 - (\widehat{U}_{-1} \cdot \widehat{V}_{-1})^{2}} \leq \sin f(\alpha)\right) \\
\leq 2 (\sin f(\alpha))^{d-k-2} \quad \text{(Lemma 2.7)} \\
= 2 \left(\max\{\frac{\sin \alpha}{\sqrt{\cos^{2} \alpha - \cos^{2}(\beta - \alpha) + \sin^{2} \alpha}}, \sin \alpha\}\right)^{d-k-2} \\
= 2 \left(\max\{\frac{\sin \alpha}{\sin(\beta - \alpha)}, \sin \alpha\}\right)^{d-k-2} \\
= 2 \left(\frac{\sin \alpha}{\sin(\beta - \alpha)}\right)^{d-k-2}.$$

Therefore,

$$\mathbb{P}\left(|\hat{U}\cdot\hat{V}| \ge \cos\alpha\right) \le (\sin\beta)^{d-k-1} + 2\left(\frac{\sin\alpha}{\sin(\beta-\alpha)}\right)^{d-k-2}.$$
 (10)

Note that we proved bound (10) conditioning on A_i 's, hence it is also a valid bound for random A_i 's (unconditionally). By Lemma 4.1, (10) implies

$$\mathbb{P}\left(\mathcal{G}(\mathcal{B}_S \cap \mathcal{B}_T) \ge \frac{\epsilon^2}{\sqrt{2\pi} \sin \alpha}\right) \le (\sin \beta)^{d-k-1} + 2\left(\frac{\sin \alpha}{\sin(\beta - \alpha)}\right)^{d-k-2}.$$
 (11)

Use inequalities $(2/\pi)x \le \sin x \le x$ for $0 \le x \le \pi/2$ to get

$$\mathbb{P}\left(\mathcal{G}(\mathcal{B}_S \cap \mathcal{B}_T) \ge \frac{\epsilon^2 \sqrt{\pi}}{\sqrt{8}\alpha}\right) \le \beta^{d-k-1} + 2\left(\frac{\pi\alpha}{2(\beta-\alpha)}\right)^{d-k-2}.$$

Set $\beta = \sqrt{\alpha}$ and restrict $0 < \alpha < 1/4$ so that $\sqrt{\alpha} \le 1/2$. The above probabilistic bound simplifies to

$$\mathbb{P}\left(\mathcal{G}(\mathcal{B}_S \cap \mathcal{B}_T) \ge \frac{\epsilon^2 \sqrt{\pi}}{\sqrt{8}\alpha}\right) \le \alpha^{(d-k-1)/2} + 2\left(\frac{\pi\alpha}{2(\sqrt{\alpha} - \alpha)}\right)^{d-k-2} \\
= \alpha^{(d-k-1)/2} + 2\left(\frac{\pi\sqrt{\alpha}}{2(1 - \sqrt{\alpha})}\right)^{d-k-2} \\
\le \alpha^{(d-k-1)/2} + 2\left(\pi\sqrt{\alpha}\right)^{d-k-2} \quad (\text{use } 1 - \sqrt{\alpha} > 1/2) \\
\le 3\left(\pi\sqrt{\alpha}\right)^{d-k-2}$$

The claim follows by setting $\alpha = \frac{\pi}{2t}$.

Combining Lemmas 7.1 and 7.2 we get

Lemma 7.4. Let $A_1, \ldots, A_k, S_1, \ldots, S_{d-k}, T_1, \ldots, T_{d-k}$ be iid. standard Gaussian random vectors in \mathbb{R}^d and $0 \le k \le d$. Let

$$\mathcal{B}_S = (\text{aff}\{A_1, \dots, A_k, S_1, \dots, S_{d-k}\})_{\epsilon/2}, \mathcal{B}_T = (\text{aff}\{A_1, \dots, A_k, T_1, \dots, T_{d-k}\})_{\epsilon/2}.$$

Then for $t > 2\pi$ we have

$$\mathbb{P}\left(\mathcal{G}(\mathcal{B}_S \cap \mathcal{B}_T) \ge \frac{\epsilon^2 t}{\sqrt{2\pi}}\right) \le 3\left(\frac{\pi^{3/2}}{\sqrt{2t}}\right)^{d-k-2}.$$

Suppose $P_n = \text{conv}(A_1, \ldots, A_n)$ is a full-dimensional simplicial polytope in \mathbb{R}^d and \mathcal{F}_n is its set of facets. For $S \in \mathcal{F}_n$, we abuse notation so that S also denotes the index set of vertices of S. Let U_S be an unit inner normal vector of $\text{aff}(A_S)$ to P_n . Define $(\text{aff }A_S)_{\epsilon^-} := \{x \in \mathbb{R}^d : 0 < d(x, \text{aff }A_S) \le \epsilon, U_S \cdot (x - A_s) \ge 0, s \in S\}$.

Lemma 7.5. Let $\delta \in (0,1)$. Suppose A_1, \ldots, A_n are d-dimensional iid. standard Gaussian random vectors with $d = \lfloor \delta n \rfloor$. Let $P_n = \text{conv}(A_1, \ldots, A_n)$, which is full-dimensional simplicial a.s. For $\epsilon > 0$, define a.s.

$$V_n = \mathcal{G}\left(\bigcup_{S \in \mathcal{F}_n} (\operatorname{aff} A_S)_{\epsilon^-} \setminus P_n\right).$$

- 1. $V_n \leq \frac{\epsilon}{\sqrt{2\pi}} \binom{n}{d}$.
- 2. There exist $c_2, c_7, c_8 > 1$ (that depend only on δ) such that when $\epsilon = \epsilon(d) \le 1/(c_8 c_2^d)$ we have $\lim_{n\to\infty} \mathbb{P}\left(V_n \ge (c_2^d/c_7)\epsilon\right) = 1$.

Proof of part 1. The upper bound follows from the union bound of at most $\binom{n}{d}$ facets and the fact that the 1-dimensional Gaussian density is upper bounded by $1/\sqrt{2\pi}$.

Proof of part 2. From Corollary 2.15, there exists a constant $c_{\mathcal{F}} > 1$ (that depends only on δ) such that $\mathbb{P}\left(|\mathcal{F}_n| \geq c_{\mathcal{F}}^d\right) \to 1$ as $n \to \infty$. Since P_n is simplicial a.s., we may present \mathcal{F}_n as a set of binary n-vectors with exactly d ones. Let $A_{\mathcal{F}_n}(t)$ be the maximum number of vectors in \mathcal{F}_n with pairwise

Hamming distance greater than or equal to t. Similarly as in the proof of Lemma 2.6, one can pick vectors greedily (Gilbert-Varshamov bound) so that when $|\mathcal{F}_n| \geq c_{\mathcal{F}}^d$ and $c \in (0,1)$, and using $n/d < 2/\delta$ when $d \geq 2$,

$$A_{\mathcal{F}_n}(cd) \ge \frac{c_{\mathcal{F}}^d}{(ne/cd)^{cd}} \ge \frac{c_{\mathcal{F}}^d}{(2e/c\delta)^{cd}}.$$

Since $\lim_{c\to 0^+} (2e/c\delta)^c = 1$ and $(2e/c\delta)^c$ is increasing for $0 \le c \le 2/\delta$, we can pick $c_1 \in (0,1)$ such that $(2e/c_1\delta)^{c_1} < c_{\mathcal{F}}$. Let $c_2 = \frac{c_{\mathcal{F}}}{(2e/c_1\delta)^{c_1}} > 1$. Then we have,

$$\lim_{n \to \infty} \mathbb{P}(A_{\mathcal{F}_n}(c_1 d) \ge c_2^d) = 1. \tag{12}$$

Here we get a subset of facets $\mathcal{T} \subseteq \mathcal{F}_n$ such that any two different facets in \mathcal{T} share no more than $(1-\frac{c_1}{2})d$ vertices, and $|\mathcal{T}| = c_2^d$ for some constants $0 < c_1 < 1$, $c_2 > 1$ (that depend only on δ). Let $N = |\mathcal{T}|$. Let $\mathcal{B}_S = (\text{aff } A_S)_{\epsilon^-}$, $S \in \mathcal{F}_n$. Using an argument similar to the proof of Lemma 4.3, we get

$$V_{n} = \mathcal{G}\left(\bigcup_{S \in \mathcal{F}_{n}} (\operatorname{aff} A_{S})_{\epsilon^{-}} \setminus P_{n}\right)$$

$$= \mathcal{G}\left(\bigcup_{S \in \mathcal{F}_{n}} (\operatorname{aff} A_{S})_{\epsilon^{-}}\right) - \mathcal{G}(P_{n} \setminus (P_{n})_{-\epsilon})$$

$$\geq \mathcal{G}\left(\bigcup_{S \in \mathcal{T}} (\operatorname{aff} A_{S})_{\epsilon^{-}}\right) - \mathcal{G}(P_{n} \setminus (P_{n})_{-\epsilon})$$

$$\geq \sum_{S \in \mathcal{T}} \mathcal{G}(\mathcal{B}_{S}) - \frac{1}{2} \sum_{S,T \in \mathcal{T},S \neq T} \mathcal{G}(\mathcal{B}_{S} \cap \mathcal{B}_{T}) - \mathcal{G}(P_{n} \setminus (P_{n})_{-\epsilon}).$$

We are going to bound each of the three terms in the last expression.

First term: $\sum_{S \in \mathcal{T}} \mathcal{G}(\mathcal{B}_S)$. From Lemma 2.12, there exists a constant $c_3 > 0$ (that depends only on δ) such that $\mathbb{P}\left(\max_{S \subseteq [n], |S| = d} \operatorname{dist}(\operatorname{aff} A_S, 0) \le c_3\right) \ge 1 - 2e^{-d}$. Moreover, we increase c_3 so that $c_3 > 1$, which ensures that $c_3 \ge \epsilon$. Recall that $\mathcal{B}_S = (\operatorname{aff} A_S)_{\epsilon^-}$. We get

$$\mathbb{P}\left(\sum_{S\in\mathcal{T}}\mathcal{G}(\mathcal{B}_S) \ge \frac{N\epsilon}{\sqrt{2\pi}}e^{-2c_3^2}\right) \ge 1 - 2e^{-d}.$$
 (13)

Second term: $\frac{1}{2} \sum_{S,T \in \mathcal{T}, S \neq T} \mathcal{G}(\mathcal{B}_S \cap \mathcal{B}_T)$. Use Lemma 7.4 in a union bound applied to all pairs of sets in \mathcal{T} . For $t > 2\pi$ we have,

$$\frac{1}{2} \sum_{S,T \in \mathcal{T}, S \neq T} \mathcal{G}(\mathcal{B}_S \cap \mathcal{B}_T) \le \binom{N}{2} \frac{\epsilon^2 t}{\sqrt{2\pi}}$$

holds with probability at least

$$1 - 3 \binom{N}{2} \left(\frac{\pi^{3/2}}{\sqrt{2t}}\right)^{d - (1 - c_1/2)d - 2} \ge 1 - \frac{3N^2 t^2}{\pi^3} \left(\frac{\pi^{3/2}}{\sqrt{2t}}\right)^{c_1 d/2}$$

$$= 1 - \frac{3t^2}{\pi^3} \left(c_2^2 \left(\frac{\pi^{3/2}}{\sqrt{2t}}\right)^{c_1/2}\right)^d.$$

Choose $t = c_4 := \frac{1}{2}\pi^3 (ec_2^2)^{\frac{4}{c_1}}$ to get

$$\mathbb{P}\left(\frac{1}{2}\sum_{S,T\in\mathcal{T},S\neq T}\mathcal{G}(\mathcal{B}_S\cap\mathcal{B}_T)\leq \binom{N}{2}\frac{\epsilon^2c_4}{\sqrt{2\pi}}\right)\geq 1-\frac{3c_4^2}{\pi^3}e^{-d}.$$
 (14)

Third term: $\mathcal{G}(P_n \setminus (P_n)_{-\epsilon})$. From Corollary 2.9, we know $\mathcal{G}(P_n \setminus (P_n)_{-\epsilon}) \leq c_5 \epsilon d^{1/4}$ for some absolute constant c_5 .

Combining (12), (13) and (14) we conclude that with probability 1 - o(1) as $d \to \infty$:

$$V_n \ge \sum_{S \in \mathcal{T}} \mathcal{G}(B_S) - \frac{1}{2} \sum_{S,T \in \mathcal{T},S \ne T} \mathcal{G}(B_S \cap B_T) - \mathcal{G}(P_n \setminus (P_n)_{-\epsilon})$$

$$\ge \frac{N\epsilon}{\sqrt{2\pi}} e^{-2c_3^2} - \binom{N}{2} \frac{\epsilon^2 c_4}{\sqrt{2\pi}} - c_5 \epsilon d^{1/4}$$

$$\ge \frac{N\epsilon}{\sqrt{2\pi}} \left(e^{-2c_3^2} - \frac{N\epsilon c_4}{2} - \frac{\sqrt{2\pi}c_5 d^{1/4}}{N} \right).$$

Note that $\sqrt{2\pi}c_5d^{1/4}/N$ decays exponentially in d. Therefore, when $\epsilon \leq 1/e^{2c_3^2}c_4N$,

$$\lim_{n \to \infty} \mathbb{P}\left(V_n \ge \frac{N\epsilon}{3\sqrt{2\pi}e^{2c_3^2}}\right) = 1.$$

The proof is finished by setting $c_7 = 3\sqrt{2\pi}e^{2c_3^2}$ and $c_8 = e^{2c_3^2}c_4$.

We are ready now to restate and prove the main result of the section.

Theorem 1.3. Let $\delta \in (0,1)$. Suppose $A = \{A_1, \ldots, A_{n+1}\}$ is a set of iid. standard Gaussian random vectors in \mathbb{R}^d and $d = \lfloor \delta n \rfloor$. Let $P_{n+1} = \text{conv}(A_1, \ldots, A_{n+1})$. Then

$$\mathbb{P}(\operatorname{diam}(P_{n+1}) \ge \sqrt{d}) \ge 1 - e^{-\frac{nd}{32}},$$

and there exists constants 0 < c, c' < 1 (that depends only δ) such that,

$$\lim_{n \to \infty} \mathbb{P}(\operatorname{vf}(P_{n+1}) \le c^d) \ge c'.$$

Hence the measure of conditioning $\kappa = \frac{\text{vf}(P_{n+1})}{\text{diam}(P_{n+1})}$ of A is exponentially small in d with constant probability.

Proof. For diam (P_{n+1}) , by Lemma 2.5 we have

$$\mathbb{P}\left(\operatorname{diam}(P_{n+1})^{2} \leq 2d - 4\sqrt{dt}\right) = \mathbb{P}\left(\|A_{i} - A_{j}\|^{2} \leq 2d - 4\sqrt{dt}, \ \forall i \neq j \in [n+1]\right) \\
\leq \mathbb{P}\left(\bigcap_{i=1}^{\lfloor (n+1)/2 \rfloor} \|A_{2i-1} - A_{2i}\|^{2} \leq 2d - 4\sqrt{dt}\right) \\
\leq \left(e^{-t}\right)^{n/2}.$$

We get the claimed bound by setting t = d/16.

Apply Lemma 7.5 to $P_n = \text{conv}(A_1, \dots, A_n)$ with $\epsilon = 1/(c_8 c_2^d)$, we have

$$\lim_{n \to \infty} \mathbb{P}\left(V_n \ge \frac{1}{c_7 c_8}\right) = 1.$$

Since $\operatorname{vf}(P_{n+1}) \leq \epsilon$ when $A_{n+1} \in V_n$, $\lim_{n \to \infty} \mathbb{P}\left(\operatorname{vf}(P_{n+1}) \leq 1/(c_8 c_2^d)\right) \geq \frac{1}{c_7 c_8}$. The claim follows by picking $c = 1/c_2$ and $c' = 1/c_7 c_8$.

Acknowledgments. We would like to thank Nina Amenta, Jesús De Loera, Miles Lopes, Javier Peña, Thomas Strohmer, Roman Vershynin and Van Vu for helpful discussions. This material is based upon work supported by the National Science Foundation under Grants CCF-1657939, CCF-1422830, CCF-2006994 and CCF-1934568.

References

- [AV97] Noga Alon and Van Vu. Anti-hadamard matrices, coin weighing, threshold gates, and indecomposable hypergraphs. *Journal of Combinatorial Theory, Series A*, 79(1):133–160, 1997.
- [AW91] Fernando Affentranger and John A. Wieacker. On the convex hull of uniform random points in a simple d-polytope. *Discret. Comput. Geom.*, 6:291–305, 1991.
- [Bal93] Keith Ball. The reverse isoperimetric problem for Gaussian measure. Discrete & Computational Geometry, 10(4):411–420, 1993.
- [BCMV13] Aditya Bhaskara, Moses Charikar, Ankur Moitra, and Aravindan Vijayaraghavan. Smoothed analysis of tensor decompositions. *CoRR*, abs/1311.3651, 2013.
- [BCV14] Aditya Bhaskara, Moses Charikar, and Aravindan Vijayaraghavan. Uniqueness of tensor decompositions with applications to polynomial identifiability. In *Conference on Learning Theory*, pages 742–778, 2014.
- [BGMN05] Franck Barthe, Olivier Guédon, Shahar Mendelson, and Assaf Naor. A probabilistic approach to the geometry of the ℓ_p^n -ball. The Annals of Probability, 33(2):480–513, 2005.
- [BGR15] Tobias Brunsch, Anna Großwendt, and Heiko Röglin. Solving totally unimodular LPs with the shadow vertex algorithm. In 32nd International Symposium on Theoretical Aspects of Computer Science, volume 30 of LIPIcs, pages 171–183, 2015.

- [BLR18] Karoly J Boroczky, Gabor Lugosi, and Matthias Reitzner. Facets of high-dimensional Gaussian polytopes. arXiv preprint arXiv:1808.01431, 2018.
- [BR13] Tobias Brunsch and Heiko Röglin. Finding short paths on polytopes by the shadow vertex algorithm. In *International Colloquium on Automata*, *Languages*, and *Programming*, pages 279–290. Springer, 2013.
- [BS17] Amir Beck and Shimrit Shtern. Linearly convergent away-step conditional gradient for non-strongly convex functions. *Mathematical Programming*, 164(1):1–27, Jul 2017.
- [BV04] René Beier and Berthold Vöcking. Typical properties of winners and losers in discrete optimization. In *Proceedings of the 36th Annual ACM Symposium on Theory of Computing*, pages 343–352, 2004.
- [BV06] René Beier and Berthold Vöcking. Typical properties of winners and losers in discrete optimization. SIAM J. Comput., 35(4):855–881, 2006.
- [CCK17] Victor Chernozhukov, Denis Chetverikov, and Kengo Kato. Central limit theorems and bootstrap in high dimensions. *The Annals of Probability*, 45(4):2309–2352, 2017.
- [CJL19] T Tony Cai, Tiefeng Jiang, and Xiaoou Li. Asymptotic analysis for extreme eigenvalues of principal minors of random matrices. arXiv preprint arXiv:1905.08757, 2019.
- [CT05] Emmanuel J. Candès and Terence Tao. Decoding by linear programming. *IEEE Trans.* Inf. Theory, 51(12):4203–4215, 2005.
- [CT06] Emmanuel J. Candès and Terence Tao. Near-optimal signal recovery from random projections: Universal encoding strategies? *IEEE Trans. Inf. Theory*, 52(12):5406– 5425, 2006.
- [DH16] Daniel Dadush and Nicolai Hähnle. On the shadow simplex method for curved polyhedra. *Discret. Comput. Geom.*, 56(4):882–909, 2016.
- [DT05] David L. Donoho and Jared Tanner. Neighborliness of randomly projected simplices in high dimensions. *Proceedings of the National Academy of Sciences*, 102(27):9452–9457, 2005.
- [EV17] Friedrich Eisenbrand and Santosh S. Vempala. Geometric random edge. *Math. Pro-gram.*, 164(1-2):325–339, 2017.
- [FW56] Marguerite Frank and Philip Wolfe. An algorithm for quadratic programming. *Naval Research Logistics Quarterly*, 3(1-2):95–110, 1956.
- [GH13] Dan Garber and Elad Hazan. A polynomial time conditional gradient algorithm with applications to online and stochastic optimization. *CoRR*, abs/1301.4666, 2013.
- [GM86] Jacques Guélat and Patrice Marcotte. Some comments on Wolfe's 'away step'. *Mathematical Programming*, 35(1):110–119, 1986.
- [HMR04] Daniel Hug, Götz Olaf Munsonius, and Matthias Reitzner. Asymptotic mean values of Gaussian polytopes. *Beiträge Algebra Geom.*, 45(2):531–548, 2004.

- [HR05] Daniel Hug and Matthias Reitzner. Gaussian polytopes: variances and limit theorems. *Adv. in Appl. Probab.*, 37(2):297–320, 2005.
- [Juk11] Stasys Jukna. Extremal combinatorics: with applications in computer science. Springer Science & Business Media, 2011.
- [Kru77] Joseph B. Kruskal. Three-way arrays: rank and uniqueness of trilinear decompositions, with application to arithmetic complexity and statistics. *Linear Algebra Appl.*, 18(2):95–138, 1977.
- [LHR20] Jesús A. De Loera, Jamie Haddock, and Luis Rademacher. The minimum euclideannorm point in a convex polytope: Wolfe's combinatorial algorithm is exponential. *SIAM J. Comput.*, 49(1):138–169, 2020.
- [LJJ13] Simon Lacoste-Julien and Martin Jaggi. An affine invariant linear convergence analysis for Frank-Wolfe algorithms. 2013.
- [LJJ15] Simon Lacoste-Julien and Martin Jaggi. On the global linear convergence of Frank-Wolfe optimization variants. In *Advances in Neural Information Processing Systems* 28, pages 496–504. 2015.
- [LM00] B. Laurent and P. Massart. Adaptive estimation of a quadratic functional by model selection. *Ann. Statist.*, 28(5):1302–1338, 10 2000.
- [Naz03] Fedor Nazarov. On the maximal perimeter of a convex set in \mathbb{R}^n with respect to a Gaussian measure. In *Geometric aspects of functional analysis*, pages 169–187. Springer, 2003.
- [PNAJ20] Fabian Pedregosa, Geoffrey Negiar, Armin Askari, and Martin Jaggi. Linearly convergent Frank-Wolfe with backtracking line-search. In *International Conference on Artificial Intelligence and Statistics*, pages 1–10. PMLR, 2020.
- [PnR19] Javier Peña and Daniel Rodríguez. Polytope conditioning and linear convergence of the Frank-Wolfe algorithm. *Math. Oper. Res.*, 44(1):1–18, 2019.
- [PnRS16] Javier Peña, Daniel Rodríguez, and Negar Soheili. On the von Neumann and Frank-Wolfe algorithms with away steps. SIAM Journal on Optimization, 26(1):499–512, 2016.
- [Ray70] H. Raynaud. Sur l'enveloppe convexe des nuages de points aléatoires dans \mathbb{R}^n . I. J. Appl. Probability, 7:35–48, 1970.
- [Raz88] Alexander A Razborov. Bounded-depth formulae over $\{\&, \oplus\}$ and some combinatorial problems. Problems of Cybernetics. Complexity Theory and Applied Mathematical Logic, pages 149–166, 1988.
- [RV05] Heiko Röglin and Berthold Vöcking. Smoothed analysis of integer programming. In *International Conference on Integer Programming and Combinatorial Optimization*, pages 276–290. Springer, 2005.

- [RV07] Heiko Röglin and Berthold Vöcking. Smoothed analysis of integer programming. Math. Program., 110(1):21-56, 2007.
- [ST01] Daniel Spielman and Shang-Hua Teng. Smoothed analysis of algorithms: Why the simplex algorithm usually takes polynomial time. In *Proceedings of the Thirty-Third Annual ACM Symposium on Theory of Computing*, page 296–305, 2001.
- [Ver18] Roman Vershynin. High-dimensional probability: An introduction with applications in data science. 47, 2018.
- [VS92] A. M. Vershik and P. V. Sporyshev. Asymptotic behavior of the number of faces of random polyhedra and the neighborliness problem. volume 11, pages 181–201. 1992.
- [Wol76] Philip Wolfe. Finding the nearest point in a polytope. *Mathematical Programming*, 11(1):128–149, Dec 1976.
- [Zie93] Günter M Ziegler. Lectures on Polytopes. 1993.
- [Zie00] Günter M Ziegler. Lectures on 0/1-polytopes. In *Polytopes—combinatorics and computation*, pages 1–41. 2000.