

THE SEMICLASSICAL RESOLVENT ON CONIC MANIFOLDS AND APPLICATION TO SCHRÖDINGER EQUATIONS

XI CHEN

ABSTRACT. In this article, we shall construct the resolvent of the Laplacian at high energies near the spectrum on non-product conic manifolds with a single cone tip. Microlocally, the resolvent kernel is the sum of b -pseudodifferential operators, scattering pseudodifferential operators and intersecting Legendrian distributions. As an application, we shall establish Strichartz estimates for Schrödinger equations on non-compact manifolds with multiple non-product conic singularities.

CONTENTS

| | |
|---|----|
| 1. Introduction | 2 |
| 1.1. The semiclassical resolvent | 3 |
| 1.2. The Schrödinger equation | 6 |
| Outline | 9 |
| Acknowledgement | 9 |
| 2. The stretched spaces | 9 |
| 2.1. The real blow-up | 9 |
| 2.2. The stretched single space | 10 |
| 2.3. The multi-stretched double space | 13 |
| 2.4. An atlas of $M_{b,0}$ | 15 |
| 2.5. The compressed cotangent bundle and density of $M_{b,0}$ | 16 |
| 3. The abstract b -calculus | 17 |
| 3.1. The small b -calculus | 17 |
| 3.2. The full b -calculus | 18 |
| 4. The b -calculus on $M_{b,0}$ | 20 |
| 4.1. The b -pseudodifferential operator on $M_{b,0}$ | 20 |
| 4.2. The indicial operator at bf | 20 |
| 5. The abstract sc -calculus | 21 |
| 5.1. The sc -pseudodifferential calculus | 21 |
| 5.2. The Legendrian distributions | 23 |
| 6. The fibred sc -calculus on $M_{b,0}$ | 26 |
| 6.1. The sc -pseudodifferential operators on $M_{sc,b,0}$ | 26 |
| 6.2. The fibred sc -bundles on $M_{b,0}$ | 27 |
| 6.3. The contact structures on fibred sc -bundles | 29 |
| 6.4. The Legendrian distributions on $M_{b,0}$ | 30 |
| 7. The normal operator at the transitional face | 33 |

| | | |
|-------|--|----|
| 8. | The b -regime parametrix | 36 |
| 8.1. | Construction near the diagonal | 36 |
| 8.2. | Error at lb | 36 |
| 8.3. | Error at bf | 37 |
| 8.4. | Error at bf_0 | 37 |
| 9. | The (b, sc) -regime parametrix | 38 |
| 9.1. | Boundary error removal | 39 |
| 9.2. | Vanishing at lb_0 | 39 |
| 10. | The sc -regime parametrix | 40 |
| 10.1. | Construction near the diagonal | 41 |
| 10.2. | Error near the boundary of the diagonal | 41 |
| 10.3. | The bicharacteristic flow | 43 |
| 10.4. | Error at the bicharacteristic | 44 |
| 10.5. | Error at L^\sharp | 47 |
| 10.6. | Error at bf_0 | 47 |
| 11. | The (sc, b) -regime parametrix | 48 |
| 11.1. | Boundary error removal | 48 |
| 11.2. | Vanishing order at rb | 49 |
| 12. | The true resolvent | 49 |
| 13. | Application to Schrödinger equations | 52 |
| 13.1. | The Schrödinger propagator on \mathcal{C} | 52 |
| 13.2. | Microlocalization and the microsupport | 53 |
| 13.3. | The conjugated spectral measure | 54 |
| 13.4. | Local Strichartz estimates on \mathcal{C} | 59 |
| 13.5. | Global Strichartz estimates on \mathcal{X} | 60 |
| | References | 62 |

1. INTRODUCTION

Let $\mathcal{C} = [0, 1) \times \mathcal{Y}$ be a conic n -manifold with a single cone tip and endowed with a non-product conic metric g , where

- $\dim \mathcal{C} = n \geq 3$,
- the submanifold \mathcal{Y} is a compact boundaryless $(n - 1)$ -manifold,
- x is the defining function of the cone tip, i.e. $\partial \mathcal{C} = \{x = 0\} = 0 \times \mathcal{Y}$,
- g is a smooth Riemannian metric outside the vicinity of the cone tip,
- g , in a compact neighbourhood \mathcal{K} of $\partial \mathcal{C}$, has the expression,

$$g = dx^2 + x^2 h(x, y, dy),^1$$

where $h(x, y, dy) = h_{ij}(x, y) dy^i dy^j$ is a family of smooth metrics on \mathcal{Y} with a smooth parameter x ,

¹This can be slightly more general, i.e. we can assume h is also dependent of dx . In fact, one can reduce the case $h(x, y, dx, dy)$ to the case $h(x, y, dy)$, by choosing appropriate coordinates in a collar neighbourhood of $\partial \mathcal{C}$. The latter metric is called the normal form of the former metric. See [31, Theorem 1.2].

- \mathcal{K} is taken to be a sufficiently small neighbourhood of $\partial\mathcal{C}$ such that h is stable in the sense

$$(1) \quad \mathbf{e} = \sup_{(x,y) \in \mathcal{K}} \frac{x \partial_x \det(h_{ij}(x,y))_{ij}}{2 \det(h_{ij}(x,y))_{ij}} \ll n-1,$$

- the non-productness of g refers to the dependence of h in x ,
- all geodesics in \mathcal{C} are simple, i.e. there are no conjugate points. ²

The Friedrichs extension of the Laplacian on \mathcal{C} , near the cone tip (i.e. in \mathcal{K}), reads

$$\Delta_{\mathcal{C}} = -\partial_x^2 - \frac{(n-1) + xe}{x} \partial_x + \frac{\Delta_h}{x^2},$$

where Δ_h is the Laplacian with respect to (\mathcal{Y}, h) with a parameter x . The non-productness of the metric in x creates the error function e ,

$$(2) \quad e(x, y) = \frac{1}{2} \frac{\partial \log \det(h_{ij}(x, y))_{ij}}{\partial x} = \frac{1}{2} \text{tr} \left(h^{ik}(x, y) \frac{\partial h_{kj}(x, y)}{\partial x} \right)_{ij},$$

where we denote the inverse matrix $(h_{ij})_{ij}^{-1} = (h^{ij})_{ij}$. We view $\Delta_{\mathcal{C}}$ as an operator acting on smooth sections of the Riemannian half density bundle of the conic metric g , which takes the form,

$$f|dg|^{1/2} = f x^{(n-1)/2} |dxdy|^{1/2} \in C^\infty(\mathcal{C}; |dg|^{1/2}).$$

1.1. The semiclassical resolvent. The main problem we want to solve in this paper is to describe the asymptotic microlocal structure of the resolvent kernel of

$$(\Delta_{\mathcal{C}} - (\lambda \pm i0)^2)^{-1},$$

at high energies near the spectrum, i.e. $|\lambda| \rightarrow \infty$.

In the product case i.e. $\partial_x h \equiv 0$, the resolvent kernel can be computed by the method of separation of variables and expressed explicitly in terms of Bessel functions. See for example [39, (8.30)]. However, this strategy can not be extended to the non-product case, as the inhomogeneity of $\Delta_{\mathcal{C}}$ in x (the presence of the error term $e(x, y)$) makes an essential challenge. To overcome this, we shall employ microlocal analysis to symbolically construct the resolvent kernel.

A natural microlocal approach to tackle this type of high energy problem is to convert the Helmholtz operator into a semiclassical operator by denoting $\hbar = |\lambda|^{-1}$. This conversion changes the high energy asymptotic of the resolvent of $\Delta_{\mathcal{C}}$ to the semiclassical resolvent of

$$(3) \quad P_{\hbar}(\pm i0) = -\hbar^2 \partial_x^2 - \frac{(n-1) + xe}{x} \hbar^2 \partial_x + \hbar^2 \frac{\Delta_h}{x^2} - (1 \pm i0).$$

To compare the conic singularity when $x \rightarrow 0$ and the semiclassical singularity when $\hbar \rightarrow 0$, we further introduce $r = x/\hbar$ and adopt coordinates $\{(r, \hbar)\}$ instead of $\{(x, \hbar)\}$. The semiclassical operator $P_{\hbar}(\pm i0)$ will lift to

$$\tilde{P}_{\hbar}(\pm i0) = -\partial_r^2 - \frac{(n-1) + \tilde{e}r}{r} \partial_r + \frac{\Delta_h}{r^2} - (1 \pm i0),$$

where the lifted error term is

$$\tilde{e}(\hbar, r, y) = \hbar e(\hbar r, y).$$

²This assumption is virtually unnecessary to understand the effect of conic singularities. But conjugate points cause extra technicalities away from the cone tip, despite being tractable, which are beyond the scope of the present paper.

We consider the incoming operator

$$(4) \quad \tilde{P}_h = -\partial_r^2 - \frac{(n-1) + \tilde{e}r}{r} \partial_r + \frac{\Delta_h}{r^2} - (1 + \imath 0).$$

The operator \tilde{P}_h now lives in a space with an artificial infinity and exhibits distinct behaviours around the two endpoints $\{r = 0\}$ and $\{r = \infty\}$:

- For small r , \tilde{P}_h reduces to a b -operator, $P_b = r\tilde{P}_hr$, where

$$(5) \quad P_b = -(r\partial_r)^2 - r\tilde{e}(r\partial_r) + \Delta_h - r\tilde{e}(1 - n/2) - r^2(1 \pm \imath 0) + (1 - n/2)^2.$$

To see this, one has to note that \tilde{P}_h acts on the conic half densities, whilst the b -operator P_b acts on the b -half densities $f|drdy/r|^{1/2} \in C^\infty({}^b\Omega^{1/2})$. By changing half densities, we have

$$\begin{aligned} P_b \left(f \left| \frac{drdy}{r} \right|^{1/2} \right) &= r^{-n/2} \left(r^{1+n/2} \left(-\partial_r^2 - \frac{(n-1) + \tilde{e}r}{r} \partial_r + \frac{\Delta_h}{r^2} - (1 + \imath 0) \right) r^{1-n/2} \right) (f|dg|^{1/2}) \\ &= \left(\left(-(r\partial_r)^2 - r\tilde{e}(r\partial_r) + \Delta_h - r\tilde{e}(1 - n/2) - r^2(1 + \imath 0) + (1 - n/2)^2 \right) f \right) \left| \frac{drdy}{r} \right|^{1/2}. \end{aligned}$$

- For large r , \tilde{P}_h turns out to be a (scattering) sc -operator. To see this, we change coordinates by making $\rho = r^{-1}$. Since ∂_r lifts to $-\rho^2\partial_\rho$, this transformation yields

$$(6) \quad P_{sc} = -(\rho^2\partial_\rho)^2 + \left((n-1)\rho - \tilde{e}(x, \rho, y) \right) \rho^2\partial_\rho + \rho^2\Delta_h - (1 + \imath 0), \quad \text{for } \rho \rightarrow 0,$$

where the error term reads

$$\tilde{e}(\rho, x, y) = \tilde{e}(\hbar, r, y)|_{\hbar=\rho x, r=1/\rho} = \rho x e(x, y).$$

This operator acts on the sc -half densities $f|d\rho dy/\rho^{n+1}|^{1/2} \in C^\infty({}^{sc}\Omega^{1/2})$.

The operators P_b and P_{sc} fit into Melrose's framework of differential analysis on manifolds with corners. Specifically, the resolvent of b -operators is a full b -pseudodifferential operator [28, Chapter 4-6], whilst the resolvent of sc -operators is a sc -pseudodifferential operator [29] plus Legendrian distributions [32, 17, 18].

These observations suggest that the resolvent \tilde{P}_h^{-1} can be constructed by the following scheme:

- use the full b -calculus to construct a local parametrix G_b for $\tilde{P}_h = r^{-1}P_br^{-1}$ with a compact remainder E_b for small r ;
- use the sc -calculus and Legendrian distributions to construct a local parametrix G_{sc} for P_{sc} with a compact remainder E_{sc} for large r ;
- improve the parametrices by understanding the transition between the b -regime (small r region) and the sc -regime (large r region);
- use Fredholm theory to get the true resolvent from these local parametrices.

Geometrically, the resolvent kernel \tilde{P}_h^{-1} has to live in a multi-stretched version of $\mathcal{C}^2 \times [0, 1)$ to incorporate these distinct microlocal structures. It is well-known that Melrose's framework of differential analysis on manifolds with corners, including b -calculus and sc -calculus, rests on some stretched spaces to desingularize the corners and express the resolvent kernel smoothly. However a simple combination of the sc -stretched and b -stretched spaces is insufficient to address the transition between the two regimes. Motivated by an unpublished idea of Melrose-Sá Barreto, previously used in [12, 13, 14, 22], we perform additional blow-ups on $\mathcal{C}^2 \times [0, 1)$ and end up with the multi-stretched spaces $M_{b,0}$ and $M_{sc,b,0}$ defined in (13) and (14) respectively. We divide such spaces into

- the b -regime $\{r < 1\} \cap \{r' < 1\}$,
- the sc -regime $\{\rho < 1\} \cap \{\rho' < 1\}$,
- the (b, sc) -regime $\{r < 1\} \cap \{\rho' < 1\}$,
- the (sc, b) -regime $\{\rho < 1\} \cap \{r' < 1\}$,

where we adopt the following projective coordinates on \mathcal{C} , $M_{b,0}$ and $M_{sc,b,0}$,

$$r = x/\hbar, \quad r' = x'/\hbar, \quad \rho = \hbar/x, \quad \rho' = \hbar/x',$$

and use \cdot' to denote the coordinates for the right subspace of the double space throughout the paper. The regime borders at the boundaries are the dashed curves in Figure 6.

We shall construct the semiclassical resolvent as follows.

Theorem 1. *Let χ_b and χ_{sc} be smooth bump functions on \mathcal{C} supported near $\{r < 1\}$ and $\{\rho < 1\}$ respectively, but also a partition of unity $\chi_b + \chi_{sc} = 1$. The resolvent kernel R of P_\hbar is a distribution on $M_{b,0}$ consisting of*

- $R_b = \chi_b R \chi_b$ is a distribution supported near the b -regime such that
 - R_b is a b -pseudodifferential operator near the diagonal,
 - R_b is polyhomogeneous conormal to lb and rb ,
 - R_b vanishes to order 2 at bf , to order $n/2$ at lb and rb ;
- $R_{(b,sc)} = \chi_b R \chi_{sc}$ is a distribution supported near the (b, sc) -regime such that
 - $R_{(b,sc)}$ vanishes to order $n/2$ at lb and to order $(n-1)/2$ at lb_0 ;
- $R_{sc} = \chi_{sc} R \chi_{sc}$ is a distribution supported near the sc -regime such that
 - R_{sc} is a sc -pseudodifferential operator near the diagonal of $M_{sc,b,0}$,³
 - R_{sc} is intersecting Legendrian distributions associated with geometric Legendrians L and diffractive Legendrians L^\sharp over $mf \cup lf \cup rf$,
 - the principal symbol of R_{sc} , in the non-product case i.e. $e \neq 0$, exhibits additional oscillations near lf , rf , and L^\sharp respectively, determined by the error function e ,
 - R_{sc} vanishes to order $(n-1-e)/2$ at both lf and rf ;
- $R_{(sc,b)} = \chi_{sc} R \chi_b$ is a distribution supported near the (sc, b) -regime such that
 - $R_{(sc,b)}$ vanishes to order $n/2$ at rb and to order $(n-1-e)/2$ at rb_0 .

The precise statement of the resolvent kernel will be given in Theorem 26. The geometry of $M_{b,0}$ and $M_{sc,b,0}$ will be described in Section 2. The involved microlocal machinery will all be defined in Section 3-6. It is also worth pointing out that the stability condition (1), i.e. $\sup_{\mathcal{K}} xe(x, y) = e \ll n-1$, is essential to dominate the additional oscillations near lf , rf , and L^\sharp . Otherwise it will go out of control and produce excess singularities.

Such a resolvent is inspired by a model problem, the resolvent construction at finite energies on non-compact metric cones, due to Guillarmou-Hassell-Sikora [14]. The non-compact metric cone $\mathfrak{C} = [0, \infty] \times \mathcal{D}$ features a global metric

$$\mathfrak{g} = d\mathfrak{x}^2 + \mathfrak{x}^2 \mathfrak{h}(y, dy)$$

with \mathfrak{h} independent of $\mathfrak{x} \in [0, \infty)$ (in contrast with our local non-product case without infinity). Then the Friedrichs extension of the Laplacian on \mathfrak{C} reads

$$\Delta_{\mathfrak{C}} = -\partial_{\mathfrak{x}}^2 - \frac{(n-1)}{\mathfrak{x}} \partial_{\mathfrak{x}} + \frac{\Delta_{\mathfrak{h}}}{\mathfrak{x}^2}.$$

The inhomogeneous e term vanishes in $\Delta_{\mathfrak{C}}$ due to the productness of $(\mathfrak{C}, \mathfrak{g})$. The resolvent kernel $(\Delta_{\mathfrak{C}} - 1 \pm 0i)^{-1}$ is a full b -pseudodifferential operator near the small end $\{\mathfrak{x} = 0, \mathfrak{x}' = 0\}$, and a sc -pseudodifferential operator plus intersecting Legendrian distributions

³The space $M_{sc,b,0}$ is only needed for this part of the resolvent.

associated with geometric Legendrians L and diffractive Legendrians L^\sharp near the big end $\{\mathfrak{x} = \infty, \mathfrak{x}' = \infty\}$. See Theorem 20 for a full description.

The Legendrian structure at the big end of \mathfrak{C}^2 is developed from the geodesic flow at infinity on asymptotically conic manifolds, introduced by Melrose-Zworski [32] for the scattering matrix, and further formulated by Hassell-Vasy [17, 18] and Hassell-Wunsch [20] for the resolvent and spectral measure. (Intersecting) Legendrian distributions on the cotangent bundle restricted to the boundary are the contact analogue of (intersecting) Lagrangian distributions on the cotangent bundle restricted in the interior of the space, pioneered by Hörmander [24], Duistermaat-Hörmander [10] and Melrose-Uhlmann [30].


It is useful to note that the productness of \mathfrak{h} grants the Laplacian $\Delta_{\mathfrak{C}}$ homogeneity in x . Taking advantage of this, one can get the microlocal structure of the semiclassical resolvent $(\hbar^2 \Delta_{\mathfrak{C}} - 1 \pm \imath 0)^{-1}$ on metric cones, which, at the small end, agrees with our result. This indeed implies that the Legendrian structure of the semiclassical resolvent over the cone tip, without the additional oscillations in the product case, is related to the conic singularities over infinity of non-compact metric cones. This observation is essential in the present paper to understand the semiclassical conic singularities over the cone tip on non-product conic manifolds.

In addition, the Legendrian structure in the sc -regime characterises the singularities of the conic geometry and diffractions at the cone tip. The Legendrian pair (L, L^\sharp) corresponds to the geometric propagation and diffractive propagation, in terms of the language of Melrose-Wunsch [31]. Microlocally, 'diffraction' refers to the splitting of singularities (waves). In the conic geometry, it is well-known that the incoming waves split, at the cone tip, into geometric propagating waves and diffractive propagating waves. The wavefront sets of such two beams of waves are indeed inherited from the geometric Legendrian L and the diffractive Legendrian L^\sharp respectively.

The resolvent on this class of local non-product conic manifolds, but away from the spectrum, has been considered by Loya [26] and lately by Hintz [22, 23]. In the classical case, Loya introduced the cone calculus to construct the resolvent $(\Delta_{\mathfrak{C}} - \lambda^2)^{-1}$ for $|\arg \lambda^2| > \theta$ and $|\lambda| < C$. The cone calculus plays the role of b -calculus and sc -calculus discussed above and establishes a 'global' elliptic pseudodifferential calculus for the cone Laplacian away from the spectrum. This idea has been further developed by Hintz for the complex power of the semiclassical resolvent $(\hbar^2 \Delta_{\mathfrak{C}} + 1)^{-1}$, which were used to investigate the propagation of semiclassical regularity and high energy estimates for potential scattering. The semiclassical resolvent, living also in the stretched double space of Melrose-Sá Barreto type, exhibits only polyhomogeneous conormal singularities due to the elliptic nature (away from the spectrum), in contrast with the Legendrian propagation in light of the vanishing boundary principal symbols (near the spectrum). Combining Hintz's work and the present paper together will give a complete description of the semiclassical resolvent on conic manifolds.

Beyond Melrose's framework, Schulze's cone calculus [35, 36, 37] also deals with the differential analysis on manifolds with conic singularities. Schrohe-Seiler [33, 34], utilizing this machinery, constructed a resolvent of the cone Laplacian in the off-spectrum case, which was applied to the study of the heat equation and the porous medium equation on conic manifolds.

1.2. The Schrödinger equation. Consider the Schrödinger equation on a non-compact manifold \mathcal{X} with multiple conic singularities as in Figure 1. We assume that \mathcal{X} is an n -manifold satisfying

FIGURE 1. The non-compact manifold \mathcal{X} with multiple conic singularities

- the boundary $\partial\mathcal{X}$ has a finite number of components $\{\partial_1\mathcal{X}, \dots, \partial_m\mathcal{X}\}$,
- for each $i = 1, \dots, m$, there exists an open neighbourhood \mathcal{C}_i of $\partial_i\mathcal{X}$ which is a conic manifold with a single cone tip $\partial_i\mathcal{X}$ and endowed with a non-product conic metric $g_i = dx_i^2 + x_i^2 h_i(x_i, y, dy)$,
- $\{\mathcal{C}_1, \dots, \mathcal{C}_m\}$ are pairwise disjoint,
- for each $i = 1, \dots, m$, there exists a compact subset $\mathcal{K}_i \subset \mathcal{C}_i$ as well as a compact ball $\mathbb{B}_i \subset \mathbb{R}^n$ such that $\mathcal{X} \setminus \bigcup_{i=1}^m \mathcal{K}_i$ is isometric to $\mathbb{R}^n \setminus \bigcup_{i=1}^m \mathbb{B}_i$ and (1) holds for the metric h_i in \mathcal{K}_i ,
- no geometric geodesic passes through three cone tips, where geometric geodesics are those missing the cone tips and their local limits.

The Schrödinger equation on \mathcal{X} with a Cauchy data $f \in L^2(\mathcal{X})$ reads

$$(7) \quad \begin{cases} i\partial_t u(t, z) + \Delta_{\mathcal{X}} u(t, z) = 0, & t > 0 \\ u(t, z) = f(z), & t = 0 \end{cases},$$

where $\Delta_{\mathcal{X}}$ is the Friedrichs extension of the Laplacian on \mathcal{X} . Using the precise semiclassical resolvent in Theorem 26, we can show

Theorem 2. *For any Schrödinger admissible pair (q, r) , i.e. (q, r) satisfies*

$$r \geq 2, q \geq 2, (q, r) \neq (2, \infty), \quad \frac{2}{q} + \frac{n}{r} = \frac{n}{2},$$

the solution to (7) on \mathcal{X} satisfies the short time Strichartz estimates, i.e.

$$(8) \quad \|u\|_{L^q((0,1]; L^r(\mathcal{X}))} \lesssim \|f\|_{H^{\mathbf{k}}(\mathcal{X})}$$

where we use the usual notations of L^2 -Sobolev spaces and space-time norms

$$H^k(\mathcal{X}) = \left\{ f \in L^2(\mathcal{X}) : \sum_{l=0}^{[k]} \|D^l f\|_{L^2(\mathcal{X})} + \left(\int_{\mathcal{X}^2} \frac{|f(z) - f(z')|^2}{|z - z'|^{2[k]+n}} dg_{\mathcal{X}^2}(z, z') \right)^{1/2} < \infty \right\},$$

$$\|u(t, z)\|_{L^q(\mathcal{T}; L^r(\mathcal{X}))} = \left(\int_{\mathcal{T}} \left(\int_{\mathcal{X}} |u(t, z)|^r dg_{\mathcal{X}}(z) \right)^{q/r} dt \right)^{1/q},$$

and the constant \mathbf{k} is defined by

$$(9) \quad \mathbf{k} = \frac{\mathbf{e}}{q(n + \mathbf{e}/2)},$$

$$(10) \quad \mathbf{e} = \max_{i=1, \dots, m} \left\{ \sup_{(x_i, y) \in \mathcal{K}_i} \left(x_i e_i(x_i, y) \right) \right\},$$

where $e_i(x, y)$ is defined by $h_i(x_i, y, dy)$ in \mathcal{K}_i as in (2) and the same constant \mathbf{e} is defined in (1) for the case of single cone tip.

Strichartz estimates have been proved in a variety of product type conic manifolds in [11, 5, 3, 41]. Nevertheless, their approaches all rely on the explicit expression of the Schrödinger propagator in terms of Bessel and Hankel functions. In the meantime, it is also possible to take advantage of the homogeneity in \mathfrak{x} of $\Delta_{\mathfrak{C}}$ on product metric cone \mathfrak{C} to prove Strichartz estimates by using Guillarmou-Hassell-Sikora's resolvent construction at finite energies. However, these methods all break down on non-product cones due to the presence of the error term e in $\Delta_{\mathfrak{C}}$. Hence Strichartz estimates on non-product type conic manifolds were still unknown. Theorem 2 indeed solves this open problem by utilizing the b -calculus and sc -calculus.

The proof of Theorem 2 integrates several existing techniques, which appeared for example in [25, 38, 15, 3, 21]. The strategy involves the TT^* argument and Keel-Tao endpoint method for (8) locally on each \mathcal{C}_i as well as (dual) local smoothing for (8) globally on \mathcal{X} . Specifically, we break Theorem 2 into the following steps:

- obtain the structure of the spectral measure $dE_{\sqrt{\Delta_{\mathcal{C}_i}}}(\lambda)$ on each \mathcal{C}_i by applying Stone's formula to the semiclassical resolvent,
- (micro)localize the spectral measure around the diagonal of \mathcal{C}_i^2 by conjugation of (microlocal) smooth cut-offs on $T^*\mathcal{C}_i$,
- prove dispersive and energy estimates on \mathcal{C}_i for the cut-off of the Schrödinger propagator $e^{it\Delta_{\mathcal{C}_i}}$, which, through the endpoint method, leads to the local Strichartz estimates on each $\mathcal{C}_i \subset \mathcal{X}$,
- prove Strichartz estimates on \mathcal{X} by combining the local estimates on \mathcal{C}_i and the local smoothing estimates on \mathcal{X} .

The key ingredient in this program, carrying the information from the conic geometry, is the microlocal structure of the resolvent/spectral measure on \mathcal{C} for large spectral parameters.

We also remark that (8) is indeed lossless when $\mathbf{e} = 0$ (i.e. $e(x, y) \leq 0$). Otherwise, the loss \mathbf{k} appears to be inevitable. This is because the resolvent \tilde{P}_h^{-1} exhibits extra growth near $\text{lf} \cup \text{rf} \subset \{\hbar = 0\}$ as pointed out in the main theorem. Consequently, this produces an excess energy growth $\lambda^{\mathbf{e}/2}$ as $\lambda \rightarrow \infty$ for the spectral measure $dE_{\sqrt{\Delta_{\mathcal{C}_i}}}(\lambda)$ in Proposition 27, and subsequently an excess time growth $t^{-\mathbf{e}/4}$ as $t \rightarrow 0$ for $e^{it\Delta_{\mathcal{C}_i}}$ in Proposition 29. Then the power of time in dispersive estimates is no longer the usual $t^{-n/2}$ but $t^{-n/2-\mathbf{e}/4}$. To compensate this larger bound, the Cauchy data thus must bring more regularities.

Alternatively, one can adapt the framework in exterior domains formulated by Burq-Gérard-Tzvetkov [8] and Burq [6, 7] to show Strichartz estimates on \mathcal{X} as follows:

- establish the high energy cut-off resolvent estimates,

$$(11) \quad \|\chi(\Delta_{\mathcal{X}} - (\lambda \pm i0)^2)^{-1}\chi\|_{L^2(\mathcal{X}) \rightarrow L^2(\mathcal{X})} \lesssim |\lambda|^{-1}, \quad |\lambda| > 1,$$

for any $\chi \in C_c^\infty(\mathcal{X})$,

- make use of the resolvent estimates (11) to show the local smoothing estimates for the solution to Schrödinger equations,

$$(12) \quad \int_0^1 \|\chi u(t, z)\|_{\mathcal{D}^{1/2}(\mathcal{X})}^2 dt \lesssim \|f\|_{L^2(\mathcal{X})},$$

where $\mathcal{D}^{s/2}(\mathcal{X})$ is the Friedrichs domain of $\Delta_{\mathcal{X}}^{s/2}$, which is the Sobolev space $H^s(\mathcal{X})$ away from the boundary and locally a weighted b -Sobolev space $x^{-n/2+s}H_b^s(\mathcal{X})$ near the boundary,⁴

- prove the Strichartz estimates by utilizing (12) and interpolation.

On \mathcal{X} , Baskin-Wunsch [4] established (11) as well as (12). However, this program only yields Strichartz estimates with a $1/q$ loss, even if $\mathbf{e} = 0$. This is far less accurate than (8), since the resolvent estimates (11) do not contain the complete information of the semiclassical resolvent on each \mathcal{C}_i . For example the Legendrian structure described in Theorem 26 is not addressed by the resolvent estimates. It is the propagation along the geometric Legendrian that prompts the various oscillations of the resolvent / spectral measure, and then leads to the dispersive estimates (51).

Outline. To begin with, the multi-stretched spaces $M_{b,0}$ and $M_{sc,b,0}$ to accommodate the resolvent kernel will be defined in Section 2. The abstract microlocal machinery, including the b -calculus, the sc -calculus, and Legendrian distributions, will be reviewed Section 3 and Section 5 and then applied to the multi-stretched spaces in Section 4 and Section 6 respectively. To alleviate technicality, we only present what will be needed in the resolvent construction. The reader is referred to [28, 29, 32, 17, 18, 20] for a complete and detailed presentation. With these microlocal tools, the local parametrices in the four regimes will be constructed in Section 8-11 respectively. Subsequently, Section 12 will obtain the true resolvent. In the end, we establish Strichartz estimates.

Acknowledgement. The author is supported by NSFC grant 11701094 and EPSRC grant EP/R001898/1. The author would like to extend his gratitude to the following people. Nicolas Burq suggested the author prove Strichartz estimates without loss by using semiclassical estimates for the resolvent, which was one of the original motivations to construct the semiclassical resolvent in this paper. Andrew Hassell taught the author the general philosophy of Melrosian calculus extensively used in this project, referred the author to [14] for the model problem on non-compact metric cones, and explained to the author a few details in [17, 18]. Peter Hintz shared the author privately with his article [22]. Jared Wunsch suggested the study of Schrödinger equations on conic manifolds and had numerous illuminative discussions with the author over this project. Part of the work was done during the author's visits to Université Paris-Sud and Northwestern University. The author would like to thank their hospitality and support. The author is also indebted to anonymous referees for their careful reading and insightful comments.



2. THE STRETCHED SPACES

2.1. The real blow-up. We wish to understand the microlocal structure of the resolvent of P_h , expressed in (3), with a parameter $\hbar \in [0, 1)$. The resolvent kernel P_h^{-1} is normally viewed as living in the double space $M = \mathcal{C}^2 \times [0, 1)$, with local coordinates (x, y, x', y', \hbar) near the boundaries. See Figure 2.

However, the resolvent kernel is singular near the boundaries of M , as the b -calculus and sc -calculus suggests, even if $\mathcal{C} = \mathbb{R}^n$. For example, the resolvent kernel on \mathbb{R}^3 , given explicitly by

$$(\Delta_{\mathbb{R}^3} - \lambda^2)^{-1}(z, z', \pm\lambda) = \frac{e^{\pm i\lambda|z-z'|}}{4\pi|z-z'|},$$

⁴Here x is a local boundary defining function and the b -Sobolev space $H_b^s(\mathcal{X})$ will be defined in (15). See [31, Section 3] for details.

FIGURE 2. The space $M = \mathcal{C}^2 \times [0, 1)$ FIGURE 3. The stretched single space \mathcal{C}_b

could be singular when $|z - z'| \rightarrow 0$ and $|\lambda| \rightarrow \infty$ simultaneously.

To remedy this and express the asymptotic limit of the kernel explicitly, Melrose pioneered the use of real blow-up with respect to a p -submanifold. Given a manifold X with corners, we say Y is a (product type) p -submanifold if for every point $y \in Y$, there exist local coordinates $\{(x_1, \dots, x_r, y', y'')\}$ of X such that

- $x_i \geq 0$ are boundary defining functions,
- $y' \in \mathbb{R}^{s'}$ and $y'' \in \mathbb{R}^{s''}$,
- $\dim X = r + s' + s''$,
- Y is locally parametrized by $\{x_1 = \dots = x_q = 0, y' = 0\}$ where $q \leq r$.

The real blow-up of X with respect to Y is a new manifold with an extra boundary hypersurface,

$$[X; Y] = (X \setminus Y) \cup SN_+ Y,$$

where Y is replaced by the inward pointing spherical normal bundle $SN_+ Y$. The manifold $[X; Y]$ is a desingularized version of X , in the sense that the fractions, which appear in the resolvent kernel, such as x_i/ϱ for $1 \leq i \leq q$ and y'/ϱ where $\varrho = (\sum_{i=1}^q x_i^2 + |y'|^2)^{1/2}$, lift to smooth functions in $[X; Y]$.

2.2. The stretched single space. The differential operator P_h in (3) is defined on the single space $\mathcal{C} \times [0, 1)$. The motivation computation suggests one can convert P_h to a

b -operator P_b in (5) and a sc -operator P_{sc} in (6) by changing variable x to $r = x/\hbar$ and $\rho = \hbar/x$ respectively.

In terms of the real blow-up, this variable changing amounts to blowing up the corner of $\mathcal{C} \times [0, 1)$. In other words, the operator P_b and the operator P_{sc} are defined on the stretched single space

$$\mathcal{C}_b = [\mathcal{C} \times [0, 1); \partial\mathcal{C} \times \{0\}].$$

We denote by β_\hbar the blow-down map corresponding to such real blow-up. Then the boundary of \mathcal{C}_b consists of $\text{lb}_0, \text{lb}, \text{mf}$, where

$$\begin{aligned} \text{lb}_0 &= \overline{\beta_\hbar^{-1}(\partial\mathcal{C} \times \{0\})} \\ \text{lb} &= \overline{\beta_\hbar^{-1}(\partial\mathcal{C} \times [0, 1))} \\ \text{mf} &= \overline{\beta_\hbar^{-1}(\mathcal{C} \times \{0\})}. \end{aligned}$$

There is a natural atlas for \mathcal{C}_b .

- In an open neighbourhood of $\text{mf} \cap \text{lb}_0$ away from lb , we have local coordinates (ρ, y, \hbar) , where \hbar is the total defining function of $\text{lb}_0 \cup \text{mf}$ and $\hbar/\rho, \rho$ locally define lb_0, mf respectively.
- In an open neighbourhood of lb away from mf , we have local coordinates (r, y, \hbar) , where $\hbar r$ is the total defining function of $\text{lb}_0 \cup \text{lb}$ and \hbar, r locally define lb_0, lb respectively.
- In an open neighbourhood away from $\text{lb} \cup \text{lb}_0$, we have local coordinates (z, \hbar) , where z is in the interior of \mathcal{C} .

To see the b -structure⁵ of P_b and the sc -structure⁶ of P_{sc} , we consider the space $\mathcal{V}_b(\mathcal{C})$ of b -vector fields on \mathcal{C} (which are tangent to $\partial\mathcal{C}$)

$$\mathcal{V}_b(\mathcal{C}) = \text{span}\{x\partial_x, \partial_y\} \quad \text{near } \partial\mathcal{C}$$

and the vector fields on $\mathcal{C} \times [0, 1)$, which are tangent to $\partial\mathcal{C}$ and vanish at the high energy face $\mathcal{C} \times \{0\}$,

$$\hbar\mathcal{V}_b(\mathcal{C}) = \text{span}\{\hbar x\partial_x, \hbar\partial_y\}, \quad \mathcal{V}_b([0, 1)) = \text{span}\{\hbar\partial_\hbar\} \quad \text{near } \partial\mathcal{C} \times [0, 1).$$

We define the space $\mathcal{V}_{b,sc}(\mathcal{C}_b)$ of vector fields on \mathcal{C}_b as

$$\mathcal{V}_{b,sc}(\mathcal{C}_b) = \rho_{\text{lb}_0}^{-1}(\mathcal{P}_{\mathcal{C} \times [0, 1) \rightarrow \mathcal{C}} \circ \beta_\hbar)^*(\hbar\mathcal{V}_b(\mathcal{C})) + \mathcal{V}_b([0, 1)),$$

where ρ_\bullet denotes the boundary defining function of the boundary hypersurface \bullet and $\mathcal{P}_{\mathcal{C} \times [0, 1) \rightarrow \mathcal{C}}$ is the canonical projection from $\mathcal{C} \times [0, 1)$ to \mathcal{C} . With such a space of vector fields on \mathcal{C}_b , we further introduce the compressed tangent bundle ${}^{b,sc}T\mathcal{C}_b$ as a bundle over \mathcal{C}_b the smooth sections of which are $\mathcal{V}_{b,sc}(\mathcal{C}_b)$, the compressed cotangent bundle ${}^{b,sc}T^*\mathcal{C}_b$ as the dual bundle of ${}^{b,sc}T\mathcal{C}_b$, the compressed density bundle ${}^{b,sc}\Omega(\mathcal{C}_b)$ as the line bundle the smooth sections of which are the wedge products of a basis of sections for ${}^{b,sc}T^*\mathcal{C}_b$.

We see the following local properties:


- near $\text{lb} \subset \mathcal{C}_b$ away from mf ,
 - we have the local coordinates (r, y, \hbar) and $\rho_{\text{lb}_0} = \hbar$,
 - $\mathcal{V}_{b,sc}(\mathcal{C}_b)$ is locally spanned by

$$r\partial_r, \quad \partial_y, \quad \hbar\partial_\hbar,$$

- the operator P_b is then an elliptic combination of the sections of ${}^{b,sc}T\mathcal{C}_b$,

⁵The abstract b -calculus will be reviewed in Section 3.

⁶The abstract sc -calculus will be reviewed in Section 5.

FIGURE 4. The sc -stretched and b -stretched double spaces

- a basis of sections of ${}^{b,sc}T^*\mathcal{C}_b$ is locally given by the following singular 1-forms,

$$\frac{dr}{r}, \quad dy, \quad \frac{d\hbar}{\hbar},$$

- the compressed density takes the form

$$\left| \frac{drdyd\hbar}{r\hbar} \right|;$$

- near $mf \subset \mathcal{C}_b$ away from lb ,
 - we have the local coordinates (ρ, y, \hbar) and $\rho|_{b_0} = \hbar/\rho$
 - $\mathcal{V}_{b,sc}(\mathcal{C}_b)$ is locally spanned by

$$\rho^2 \partial_\rho, \quad \rho \partial_y, \quad \hbar \partial_\hbar,$$

- the operator P_{sc} is then an elliptic combination of the sections of ${}^{b,sc}T\mathcal{C}_b$,
- a basis of sections of ${}^{b,sc}T^*\mathcal{C}_b$ is locally given by the following singular 1-forms,

$$\frac{d\rho}{\rho^2}, \quad \frac{dy}{\rho}, \quad \frac{d\hbar}{\hbar},$$

- the compressed density takes the form

$$\left| \frac{d\rho dy d\hbar}{\rho^{n+1} \hbar} \right|;$$

- away from $lb \cup lb_0$, i.e. a neighbourhood of $(\mathcal{C} \setminus \mathcal{K}) \times [0, 1)$,
 - we have the local coordinates (z, \hbar)
 - $\mathcal{V}_{b,sc}(\mathcal{C}_b)$ is locally spanned by

$$\partial_z, \quad \hbar \partial_\hbar,$$

- a basis of sections of ${}^{b,sc}T^*\mathcal{C}_b$ is locally given by the following singular 1-forms,

$$dz, \quad \frac{d\hbar}{\hbar},$$

- the compressed density takes the form

$$\left| \frac{dz d\hbar}{\hbar} \right|,$$

- this region is viewed as a part of $\{\rho < 1\} \subset \mathcal{C}_b$ though there is no $\rho = \hbar/x$ coordinate.

2.3. The multi-stretched double space. However, the stretched single space \mathcal{C}_b is insufficient to address all singularities of the resolvent, e.g. the diagonal singularity as in the expression of $(\Delta_{\mathbb{R}^3} - \lambda^2)^{-1}$, as well as its interactions with boundary singularities generally in Melrosian calculus on manifolds with ends.

Via the canonical 'left' ('right') projection $\mathcal{P}_L(\mathcal{P}_R) : M \rightarrow \mathcal{C}$, we can pull back P_h to the double space M and view it as a distribution acting on the 'left half' ('right half') of M when we construct the 'right-sided' ('left-sided') parametrix. To avoid confusion, we shall use the following notations for a differential operator P on the single space and distributions G on the double space

- by $P(f)$ we mean P acting on a function f on the single space,
- by $P \circ G$ we mean P acting on the left half of G on the double space,
- the usual composition of G_1 and G_2 and the power of G on the double space is denoted by $G_1 G_2$ and $GG = G^2$.

As we have seen on the single space, the operator P_h turns out to be a b -operator P_b and a sc -operator P_{sc} near the two ends $\{r < 1\}$ and $\{\rho < 1\}$ respectively. This implies we need to perform blow-ups to accommodate both the b -structure and the sc -structure.

In an abstract setting, a b -differential operator \mathcal{O}_b and a sc -differential operator \mathcal{O}_{sc} individually act as a distribution on a manifold X with local coordinates (x, y) near the boundary where x is a boundary defining function. We shall follow the convention that the coordinates for the right copy of the double space are primed. Then the kernels of \mathcal{O}_{sc} and \mathcal{O}_b are defined in X^2 with two boundary faces $B_l = \{(0, y, z') \in \partial X \times X\}$ and $B_r = \{(z, 0, y') \in X \times \partial X\}$ as well as a corner $C_X = \{(0, y, 0, y') \in \partial X \times \partial X\}$. The resolvent kernels of \mathcal{O}_{sc} and \mathcal{O}_b , in terms of the sc -calculus and b -calculus, live in $X_b^2 = [X^2; C_X]$ and $X_{sc,b}^2 = [X_b^2; \{x/x' = 1, y = y'\}]$ respectively. See Figure 4.

Hence we aim to describe the resolvent kernel of P_h by consolidating such two distinct microlocal structures in M . The resolvent exhibits separate singularities as we approach the left face, the right face, and the high energy face respectively. We thus need to perform a series of blow-ups to compare these singularities.

The order of these blow-ups matters on M . First, the reduction to (4) by coordinate changing $r = x/\hbar$ ($r' = x'/\hbar$) means we blow up the corner between the left (right) face and the high energy face. Second, as P_b for small $r = x/\hbar$ is a b -differential operator, we need to blow up the intersection of the left and right faces to employ the b -calculus. Third, a blow-up of the diagonal at the high energy faces is required to apply the sc -calculus for P_{sc} . These blown up faces will affect one another and meet at the codimension three corner $\{x = x' = \hbar = 0\} \subset M$.

It is natural to firstly blow up the corner with the highest codimension,


$$\beta_0 : M_0 = [M; \{x = x' = \hbar = 0\}] \longrightarrow M,$$

and then perform blow-ups,

$$(13) \quad \beta_b : M_{b,0} = [M_0; \{x = x' = 0\}, \{x = \hbar = 0\}, \{x' = \hbar = 0\}] \longrightarrow M_0.$$

This two-fold blow-up $\beta_{b,0} = \beta_0 \circ \beta_b$ is illustrated in Figure 5. We introduce the following notations for the boundary hypersurfaces of $M_{b,0}$,

$$\begin{aligned} \text{bf}_0 &= \overline{\beta_{b,0}^{-1}(\partial\mathcal{C} \times \partial\mathcal{C} \times \{0\})} \\ \text{bf} &= \overline{\beta_{b,0}^{-1}(\partial\mathcal{C} \times \partial\mathcal{C} \times [0, 1] \setminus \partial\mathcal{C} \times \partial\mathcal{C} \times \{0\})} \\ \text{lb}_0 &= \overline{\beta_{b,0}^{-1}(\partial\mathcal{C} \times \mathcal{C} \times \{0\} \setminus \partial\mathcal{C} \times \partial\mathcal{C} \times \{0\})} \\ \text{rb}_0 &= \overline{\beta_{b,0}^{-1}(\mathcal{C} \times \partial\mathcal{C} \times \{0\} \setminus \partial\mathcal{C} \times \partial\mathcal{C} \times \{0\})} \end{aligned}$$

FIGURE 5. The space $M_{b,0} = \beta_b^{-1} \beta_0^{-1} M$

$$\begin{aligned} \text{mf} &= \overline{\beta_{b,0}^{-1}(\mathcal{C} \times \mathcal{C} \times \{0\})} \\ \text{lb} &= \overline{\beta_{b,0}^{-1}(\partial\mathcal{C} \times \mathcal{C} \times [0,1])} \\ \text{rb} &= \overline{\beta_{b,0}^{-1}(\mathcal{C} \times \partial\mathcal{C} \times [0,1])}. \end{aligned}$$

Let diag_\bullet be the diagonal of a double space \bullet . The diagonal of the interior of M is lifted via $\beta_{b,0}^{-1}$ to the diagonal of $M_{b,0}$

$$\text{diag}_{M_{b,0}} = \overline{\beta_{b,0}^{-1}(\{(z, z, \hbar) : z \in \mathcal{C} \setminus \partial\mathcal{C}, \hbar \in [0,1]\})}.$$

We denote by $\partial_b \text{diag}_{M_{b,0}}$ and $\partial_{sc} \text{diag}_{M_{b,0}}$ the intersections of $\text{diag}_{M_{b,0}}$ with bf and mf respectively. The dashed line in mf in Figure 5 stands for $\partial_{sc} \text{diag}_{M_{b,0}}$.

The multi-stretched double space $M_{b,0}$ is the suitable double space for the resolvent kernel of the operator \tilde{P}_\hbar defined on the stretched single space. On the one hand, the above blow-up scheme $(\beta_0 \circ \beta_b)^{-1}$ to create the multi-stretched spaces $M_{b,0}$ from $\mathcal{C}^2 \times [0,1]$ is designed in accordance with the variable changing performed from P_\hbar to \tilde{P}_\hbar . On the other hand, the original differential operator P_\hbar , defined on the single space $\mathcal{C} \times [0,1]$ with a parameter \hbar , lifts to P_b and P_{sc} locally defined on the stretched single space \mathcal{C}_b . Although there is no canonical projection from the multi-stretched double space $M_{b,0}$ to the stretched single space \mathcal{C}_b , one can prove there exists a left (right) stretched projection $\tilde{\mathcal{P}}_L(\tilde{\mathcal{P}}_R) : M_{b,0} \rightarrow \mathcal{C}_b$ by invoking [16, Lemma 2.7] and the fact

$$M_{b,0} = [[\mathcal{C}_b \times \mathcal{C}; \text{lb}_0 \times \partial\mathcal{C}]; \text{lb} \times \partial\mathcal{C}, \text{mf} \times \partial\mathcal{C}].$$


In summary, the operator P_\hbar defined on M is lifted, via the left (right) stretched projection $\tilde{\mathcal{P}}_L(\tilde{\mathcal{P}}_R)$, to \tilde{P}_\hbar defined on $M_{b,0}$, which acts on the left (right) 'half' of $M_{b,0}$. The operator \tilde{P}_\hbar is further converted to a b -operator P_b in (5) near bf for small r (r'), whilst it is a sc -operator P_{sc} in (6) near mf for small ρ (ρ').

In addition, the sc -calculus requires a further blow-up on $M_{b,0}$ as we shall see. Then we create the face sc by blowing up $\partial_{sc} \text{diag}_{M_{b,0}}$

$$(14) \quad \beta_{sc} : M_{sc,b,0} = [M_{b,0}; \partial_{sc} \text{diag}_{M_{b,0}}] \longrightarrow M_{b,0},$$

as in Figure 6.

We shall mostly live in $M_{b,0}$ and only visit $M_{sc,b,0}$ briefly while constructing the sc -pseudodifferential parametrix of P_{sc} .

FIGURE 6. The space $M_{sc,b,0} = \beta_{sc}^{-1} \beta_b^{-1} \beta_0^{-1} M$

2.4. An atlas of $M_{b,0}$. To separate the distinct behaviours of \tilde{P}_h as the left (right) operator on $M_{b,0}$, we decompose the stretched space $M_{b,0}$ (and $M_{sc,b,0}$ similarly), by the defining functions

$$r = x/\hbar, \quad r' = x'/\hbar, \quad \rho = \hbar/x, \quad \rho' = \hbar/x',$$

into the following regimes. The two intersecting dashed curves in Figure 5 and Figure 6 stand for the borders of the regimes over the boundary faces.

- The b -regime is defined by $\{r < 1\} \cap \{r' < 1\} \subset M_{b,0}$, furnished with local coordinates (r, y, r', y', \hbar) , and has boundary hypersurfaces:
 - $lb = \overline{\{r = 0, 0 < r' < 1\}}$ is the left boundary in the b -calculus for P_b ,
 - $rb = \overline{\{0 < r < 1, r' = 0\}}$ is the right boundary in the b -calculus for P_b ,
 - $bf = \overline{\{r = 0, r' = 0\}}$ is the front face in the b -calculus for P_b .
- The (b, sc) -regime is defined by $\{r < 1\} \cap \{\rho' < 1\} \subset M_{b,0}$, furnished with local coordinates (r, y, ρ', y', \hbar) , and has boundary hypersurfaces:
 - $lb_0 = \overline{\{0 < r < 1, \rho' = 0\}}$,
 - $lb = \overline{\{0 < \rho' < 1, r = 0\}}$.
- The sc -regime is defined by $\{\rho < 1\} \cap \{\rho' < 1\} \subset M_{b,0}$, furnished with local coordinates $(\rho, y, \rho', y', \hbar)$, and has boundary hypersurfaces:
 - $lf = \{\rho = 0, 0 < \rho' < 1\} \subset rb_0$, defined by ρ , acts as the left face in the sc -calculus for P_{sc} , hence the name lf ,
 - $rf = \{\rho' = 0, 0 < \rho < 1\} \subset lb_0$, defined by ρ' , acts as the right face in the sc -calculus for P_{sc} , hence the name rf ,
 - $mf = \{\rho = 0, \rho' = 0\}$ is the main face, which plays a central role in the discussion of Legendrian structures in the sc -calculus.
- The (sc, b) -regime is defined by $\{\rho < 1\} \cap \{r' < 1\} \subset M_{b,0}$, furnished with local coordinates (ρ, y, r', y', \hbar) , and has boundary hypersurfaces:
 - $rb_0 = \overline{\{0 < r' < 1, \rho = 0\}}$,
 - $rb = \overline{\{0 < \rho < 1, r' = 0\}}$.
- The transitional face bf_0 is defined by $\overline{\{x^2 + x'^2 + \hbar^2 = 0\}} \subset M_{b,0}$ and likewise decomposed into four pieces by the defining functions r, r', ρ, ρ' . Restricted to bf_0 , the operator \tilde{P}_h reduces to $\Delta_{\mathfrak{C}} - (1 + \imath 0)$. Therefore, bf_0 can be viewed as a stretched double metric cone \mathfrak{C}^2 , whilst the resolvent kernel $(\Delta_{\mathfrak{C}} - (1 + \imath 0))^{-1}$ on a

stretched version of \mathfrak{C}^2 is equivalent to the resolvent kernel \tilde{P}_h^{-1} on bf_0 . This will be discussed in detail in Section 7.

- The interior of $M_{b,0}$ is defined by a neighbourhood of $(\mathcal{C} \setminus \mathcal{K})^2 \times [0, 1)$ and furnished with local coordinates (z, z', \hbar) . We remark that the interior does have a boundary face in $\{\hbar = 0\}$ but it has nothing to do with the conic singularities of \mathcal{C} . We view the interior as a subset of the sc -regime as we did for the stretched single space.

2.5. The compressed cotangent bundle and density of $M_{b,0}$. Understanding the microlocal structure of the resolvent kernel on $M_{b,0}$, especially the singularities near the boundary, requires a suitable compressed cotangent bundle.

By the left and right stretched projections $\tilde{\mathcal{P}}_L$ and $\tilde{\mathcal{P}}_R$, we pull back the space $\mathcal{V}_{b,sc}(\mathcal{C}_b)$ of vector fields on the stretched single space \mathcal{C}_b to the space $\mathcal{V}_{b,sc}(M_{b,0})$ of vector fields on the stretched double space $M_{b,0}$,

$$\mathcal{V}_{b,sc}(M_{b,0}) = \tilde{\mathcal{P}}_L^*(\mathcal{V}_{b,sc}(\mathcal{C}_b)) + \tilde{\mathcal{P}}_R^*(\mathcal{V}_{b,sc}(\mathcal{C}_b)).$$

This Lie algebra determines the smooth sections of the compressed tangent bundle ${}^{b,sc}TM_{b,0}$ and then defines the desired compressed cotangent bundle ${}^{b,sc}T^*M_{b,0}$ as well as the compressed density bundle ${}^{b,sc}\Omega(M_{b,0})$.

With the atlas of $M_{b,0}$, we have the following local expressions.

- In the b -regime $\{r < 1\} \cup \{r' < 1\}$, under coordinates (r, y, r', y', \hbar) ,
 - the compressed cotangent bundle ${}^{b,sc}T^*M_{b,0}$ is spanned by singular 1-forms

$$\frac{dr}{r}, \quad \frac{dr'}{r'}, \quad dy, \quad dy', \quad \frac{d\hbar}{\hbar},$$

and essentially a b -cotangent bundle;

- the compressed density ${}^{b,sc}\Omega(M_{b,0})$ thus takes the form,

$$\left| \frac{dr dy dr' dy' d\hbar}{rr' \hbar} \right|,$$

and is essentially a b -density.

- In the sc -regime $\{\rho < 1\} \cap \{\rho' < 1\}$, under coordinates $(\rho, y, \rho', y', \hbar)$,
 - the compressed cotangent bundle ${}^{b,sc}T^*M_{b,0}$ is spanned by singular 1-forms

$$\frac{d\rho}{\rho^2}, \quad \frac{d\rho'}{\rho'^2}, \quad \frac{dy}{\rho}, \quad \frac{dy'}{\rho'}, \quad \frac{d\hbar}{\hbar},$$

and essentially a sc -cotangent bundle;

- the compressed density ${}^{b,sc}\Omega(M_{b,0})$ thus takes the form,

$$\left| \frac{d\rho dy d\rho' dy' d\hbar}{\rho^{n+1} \rho'^{n+1} \hbar} \right|,$$

and is essentially a sc -density.

- In the (b, sc) -regime $\{r < 1\} \cap \{\rho' < 1\}$, under coordinates (r, y, ρ', y', \hbar) ,
 - the compressed cotangent bundle ${}^{b,sc}T^*M_{b,0}$ is spanned by singular 1-forms

$$\frac{dr}{r}, \quad \frac{d\rho'}{\rho'^2}, \quad dy, \quad \frac{dy'}{\rho'}, \quad \frac{d\hbar}{\hbar},$$

- the compressed density ${}^{b,sc}\Omega(M_{b,0})$ thus takes the form,

$$\left| \frac{dr dy d\rho' dy' d\hbar}{r \rho'^{n+1} \hbar} \right|.$$

- In the (sc, b) -regime $\{\rho < 1\} \cap \{r' < 1\}$, under coordinates (ρ, y, r', y', \hbar) ,

- the compressed cotangent bundle ${}^{b,sc}T^*M_{b,0}$ is spanned by singular 1-forms

$$\frac{d\rho}{\rho^2}, \quad \frac{dr'}{r'}, \quad \frac{dy}{\rho}, \quad dy', \quad \frac{d\hbar}{\hbar},$$

- the compressed density ${}^{b,sc}\Omega(M_{b,0})$ thus takes the form,

$$\left| \frac{d\rho dy dr' dy' d\hbar}{\rho^{n+1} r' \hbar} \right|.$$

- The transitional region, bf_0 , can be viewed as the space where the resolvent kernel of $\Delta_{\mathcal{C}}$ lives, which will be discussed in detail in Section 7.

3. THE ABSTRACT b -CALCULUS

In this section, we shall review the abstract b -calculus. The notations in this section will be self-contained and independent of the conic manifold \mathcal{C} .

3.1. The small b -calculus. Suppose X is an n -manifold which near any boundary point has a coordinate chart $\{(x, y) : x \in [0, 1), y \in \mathbb{R}^{n-1}\}$. The space $\mathcal{V}_b(X)$ of b -vector fields is the collection of smooth sections of TX tangent to the boundary ∂X . Then the b -vector fields are smooth sections of the b -tangent bundle bTX , which has a local frame $\{x\partial_x, \partial_y\}$ near the boundary. The b -cotangent bundle ${}^bT^*X$ is the dual bundle of bTX .

The space $\text{Diff}_b^m(X)$ of m th order b -differential operators consists of finite sums of m -fold products of b -vector fields. Near ∂X , a b -differential operator $A \in \text{Diff}_b^m(X)$ acts on b -half densities

$$f \left| \frac{dx dy}{x} \right|^{1/2} \in C^\infty(X; {}^b\Omega^{1/2}),$$

and takes the explicit form,

$$A = \sum_{\alpha+|\beta|\leq m} a_{\alpha\beta}(x, y) (x\partial_x)^\alpha \partial_y^\beta, \quad \alpha \in \mathbb{N}_0 \text{ and } \beta \in \mathbb{N}_0^{n-1}.$$

The b -differential operators can be generalized to b -pseudodifferential operators. The Schwartz kernel of a b -pseudodifferential operator is a distribution living in the b -stretched space

$$X_b^2 = \beta_b^{-1}(X^2) = [X^2; \partial X \times \partial X]$$

with three boundary faces

$$\begin{aligned} \text{bf} &= \overline{\beta_b^{-1}(\partial X \times \partial X)} \\ \text{lb} &= \overline{\beta_b^{-1}(\partial X \times X^\circ)} \\ \text{rb} &= \overline{\beta_b^{-1}(X^\circ \times \partial X)}. \end{aligned}$$

and conormal to the b -diagonal $\text{diag}_b = \overline{\beta_b^{-1}(\text{diag}^\circ)}$. To be concrete,

Definition 3. The space $\Psi_b^m(X; {}^b\Omega^{1/2})$ is the space of operators with a kernel $\kappa |d\mu_b|^{1/2}$ living in X_b^2 such that

- κ is a distribution, conormal of degree m to diag_b , smooth up to bf , and rapidly vanishing at lb and rb .
- $|d\mu_b|^{1/2}$ is the lift of $\pi_l^* {}^b\Omega^{1/2} \otimes \pi_r^* {}^b\Omega^{1/2}$ to X_b^2 , where π_l and π_r are the projections from X_b^2 down to a single space X .

In analogy with classical pseudodifferential operators, b -pseudodifferential operators have a well-defined symbol map

$$\Psi_b^m(X; {}^b\Omega^{1/2}) \longrightarrow S^m({}^bT^*X)/S^{m-1}({}^bT^*X),$$

where $S^m(\cdot)$ is the classical symbol space of degree m . This is extended to a short exact sequence

$$0 \rightarrow \Psi_b^{m-1}(X; {}^b\Omega^{1/2}) \hookrightarrow \Psi_b^m(X; {}^b\Omega^{1/2}) \rightarrow S^m({}^bT^*X) \rightarrow 0.$$

3.2. The full b -calculus. Since the small b -calculus only addresses the singularities of the resolvent near diag_b , the full b -calculus is introduced to describe the boundary behaviours of the true resolvent.

First of all, the boundary singularities are measured in terms of b -Sobolev spaces and polyhomogeneous conormal distributions to boundaries. The b -Sobolev space of order m is defined by b -differential operators as

$$(15) \quad H_b^m(X; {}^b\Omega^{1/2}) = \{u \in L^2(X; {}^b\Omega^{1/2}) : \text{Diff}_b^m(X; {}^b\Omega^{1/2})u \subset L^2(X; {}^b\Omega^{1/2})\}.$$

The space of conormal distributions of degree $(\alpha_{\text{bf}}, \alpha_{\text{lb}}, \alpha_{\text{rb}})$ to $(\text{bf}, \text{lb}, \text{rb})$ on X_b^2 , denoted by $\mathcal{A}^{(\alpha_{\text{bf}}, \alpha_{\text{lb}}, \alpha_{\text{rb}})}(X_b^2; {}^b\Omega^{1/2})$, consists of distributions such that

$$\{u : \mathcal{V}_b u \subset \rho_{\text{bf}}^{\alpha_{\text{bf}}} \rho_{\text{lb}}^{\alpha_{\text{lb}}} \rho_{\text{rb}}^{\alpha_{\text{rb}}} L^\infty(X_b^2; {}^b\Omega^{1/2})\} \quad \text{with } (\alpha_{\text{bf}}, \alpha_{\text{lb}}, \alpha_{\text{rb}}) \in \mathbb{R}^3,$$

where ρ_{bf} , ρ_{lb} and ρ_{rb} are defining functions for bf , lb and rb respectively.

The polyhomogeneity for a boundary \bullet is measured by the index set \mathcal{E}_\bullet , which is a discrete subset of $\mathbb{C} \times \mathbb{N}_0$ such that

- $(z_j, k_j) \in \mathcal{E}_\bullet, |(z_j, k_j)| \rightarrow \infty \Rightarrow \text{Re } z_j \rightarrow \infty$,
- $(z, k) \in \mathcal{E}_\bullet \Rightarrow (z, l) \in \mathcal{E}_\bullet, l \in \mathbb{N}_0, 0 \leq l \leq k$,
- $(z, k) \in \mathcal{E}_\bullet \Rightarrow (z + j, k) \in \mathcal{E}_\bullet, j \in \mathbb{N}_0$.

We denote $\text{Re } \mathcal{E}_\bullet = \min\{\text{Re } z : \exists (z, k) \in \mathcal{E}_\bullet\}$ for \mathcal{E}_\bullet .

A collection \mathcal{E} of index sets for each boundary hypersurface is called the index family. The space $\mathcal{A}_{phg}^{(\mathcal{E}_{\text{bf}}, \mathcal{E}_{\text{lb}}, \mathcal{E}_{\text{rb}})}(X_b^2; {}^b\Omega^{1/2})$ of polyhomogeneous conormal distributions to bf , lb and rb consists of distributions having the following local expressions

$$\begin{aligned} u &\sim \sum_{(z_{\text{bf}}, k_{\text{bf}}) \in \mathcal{E}_{\text{bf}}} \rho_{\text{bf}}^{z_{\text{bf}}} (\log \rho_{\text{bf}})^{k_{\text{bf}}} a_{(z_{\text{bf}}, k_{\text{bf}})} \quad \text{with } a_{(z_{\text{bf}}, k_{\text{bf}})} \in C^\infty(X_b^2; {}^b\Omega^{1/2}), \quad \text{near bf;} \\ u &\sim \sum_{(z_{\text{lb}}, k_{\text{lb}}) \in \mathcal{E}_{\text{lb}}} \rho_{\text{lb}}^{z_{\text{lb}}} (\log \rho_{\text{lb}})^{k_{\text{lb}}} a_{(z_{\text{lb}}, k_{\text{lb}})} \quad \text{with } a_{(z_{\text{lb}}, k_{\text{lb}})} \in C^\infty(X_b^2; {}^b\Omega^{1/2}), \quad \text{near lb;} \\ u &\sim \sum_{(z_{\text{rb}}, k_{\text{rb}}) \in \mathcal{E}_{\text{rb}}} \rho_{\text{rb}}^{z_{\text{rb}}} (\log \rho_{\text{rb}})^{k_{\text{rb}}} a_{(z_{\text{rb}}, k_{\text{rb}})} \quad \text{with } a_{(z_{\text{rb}}, k_{\text{rb}})} \in C^\infty(X_b^2; {}^b\Omega^{1/2}), \quad \text{near rb.} \end{aligned}$$

To measure the polyhomogeneity of the product and composition of two conormal distributions, the operations of addition and extended union for two index sets \mathcal{E}_1 and \mathcal{E}_2 are defined to be

$$\begin{aligned} \mathcal{E}_1 + \mathcal{E}_2 &= \{(z_1 + z_2, k_1 + k_2) | (z_1, k_1) \in \mathcal{E}_1, (z_2, k_2) \in \mathcal{E}_2\} \\ \mathcal{E}_1 \sqcup \mathcal{E}_2 &= \mathcal{E}_1 \cup \mathcal{E}_2 \cup \{(z, k) : \exists (z, k_1) \in \mathcal{E}_1 \text{ and } (z, k_2) \in \mathcal{E}_2 \text{ with } k = k_1 + k_2 + 1\}. \end{aligned}$$

To measure the least power which can occur, the infimum of an index set \mathcal{E} is defined by

$$\inf \mathcal{E} = \min\{\text{Re } z : (z, 0) \in \mathcal{E}\}.$$

The space $\Psi_b^{m,\mathcal{E}}(X_b^2; {}^b\Omega^{1/2})$ of full b -pseudodifferential operators with $\mathcal{E} = (\mathcal{E}_{\text{lb}}, \mathcal{E}_{\text{rb}}, \mathcal{E}_{\text{bf}})$ is the collection of the sums of small b -pseudodifferential operators $\Psi_b^m(X_b^2; {}^b\Omega^{1/2})$ and the operators with a polyhomogeneous conormal kernel in $\mathcal{A}_{phg}^{(\mathcal{E}_{\text{lb}}, \mathcal{E}_{\text{rb}}, \mathcal{E}_{\text{bf}})}(X_b^2; {}^b\Omega^{1/2})$.

Second, the boundary behaviour of the resolvent of A involves the indicial operator $\mathcal{I}(A)$,⁷ which is the restriction of A to ∂X . It has a local expression,

$$\mathcal{I}(A) = \sum_{\alpha+|\beta|=m} a_{\alpha\beta}(0, y) (x\partial_x)^\alpha \partial_y^\beta.$$

In the same spirit with the Fourier transform approach to evolution equations, the inversion of $\mathcal{I}(A)$, via the Mellin transform \mathcal{M} ,

$$\mathcal{M}(u)(\lambda) = \int_0^\infty x^{-i\lambda} u(x, y) \frac{dx}{x},$$

reduces to the indicial family $\mathcal{I}(A, \lambda)$,

$$\mathcal{I}(A, \lambda) = \mathcal{M}(\mathcal{I}(A))(\lambda) = \sum_{\alpha+|\beta|=m} a_{\alpha\beta}(0, y) (\iota\lambda)^\alpha \partial_y^\beta.$$

The Mellin transform gives an isomorphism

$$x^\alpha L^2(\mathbb{R}_+, \frac{dx}{x}) \longrightarrow L^2(\{\text{Im } \lambda = \alpha\}, d\lambda),$$

for any $\alpha \in \mathbb{R}$. The inverse Mellin transform corresponds to integration along $\{\text{Im } \lambda = \alpha\}$,

$$\mathcal{M}^{-1}(v)(x) = \frac{1}{2\pi} \int_{\{\text{Im } \lambda = \alpha\}} x^{i\lambda} v(\lambda, y) d\lambda.$$

To address the invertibility of the indicial operator, we define the boundary spectrum,

$$\begin{aligned} \text{spec}_b(\mathcal{I}(A)) &= \left\{ \lambda \in \mathbb{C} : \mathcal{I}(A, \lambda) \text{ is not invertible on } C^\infty(\partial X) \right\}, \\ \text{Spec}_b(\mathcal{I}(A)) &= \left\{ (\lambda, k) : \text{the inverse of } \mathcal{I}(A, \cdot) \text{ has a pole of order } k+1 \text{ at } \lambda \right\}. \end{aligned}$$

Through the Mellin transform, the invertibility of $\mathcal{I}(A, \lambda)$ amounts to the invertibility of $\mathcal{I}(A)$ on weighted b -Sobolev spaces.

Proposition 4. *The indicial operator $\mathcal{I}(A)$ of $A \in \text{Diff}_b^m(X)$ is an isomorphism between $x^\alpha H_b^j(X; {}^b\Omega^{1/2})$ and $x^\alpha H_b^{j-m}(X; {}^b\Omega^{1/2})$ for all j if and only if $\alpha \in \mathbb{R}$, $\alpha \neq -\text{Im } \lambda$ for any $\lambda \in \text{spec}_b(\mathcal{I}(A))$.*

Finally, the composition of the full b -pseudodifferential operators will be useful in the construction of the parametrix. See [2, Theorem 4.20].

Proposition 5. *For two full b -pseudodifferential operators on X_b^2 ,*

$$\begin{aligned} A &\in \Psi_b^{m_1, \mathcal{E}}(X_b^2, {}^b\Omega^{1/2}) && \text{with } \mathcal{E} = (\mathcal{E}_{\text{lb}}, \mathcal{E}_{\text{rb}}, \mathcal{E}_{\text{bf}}), \\ B &\in \Psi_b^{m_2, \mathcal{F}}(X_b^2, {}^b\Omega^{1/2}) && \text{with } \mathcal{F} = (\mathcal{F}_{\text{lb}}, \mathcal{F}_{\text{rb}}, \mathcal{F}_{\text{bf}}), \end{aligned}$$

satisfying $\text{Re}(\mathcal{E}_{\text{rb}} + \mathcal{F}_{\text{lb}}) > 0$, the composition operator $A \circ B$ lies in the space

$$\Psi_b^{m_1+m_2, \mathcal{G}}(X_b^2, {}^b\Omega^{1/2}) \quad \text{with } \mathcal{G} = (\mathcal{G}_{\text{lb}}, \mathcal{G}_{\text{rb}}, \mathcal{G}_{\text{bf}})$$

⁷It is also called normal operator. In general, the normal operator $\mathcal{N}_\bullet(\cdot)$ is the restriction of \cdot to the face \bullet . When the indicial family is further involved, we call it indicial operator, to be consistent with the convention in [28]. Otherwise, it is named normal operator.

where the index family \mathcal{G} is given by

$$\begin{aligned}\mathcal{G}_{\text{lb}} &= (\mathcal{E}_{\text{bf}} + \mathcal{F}_{\text{lb}}) \sqcup \mathcal{E}_{\text{lb}}, \\ \mathcal{G}_{\text{rb}} &= (\mathcal{E}_{\text{rb}} + \mathcal{F}_{\text{bf}}) \sqcup \mathcal{F}_{\text{rb}}, \\ \mathcal{G}_{\text{bf}} &= (\mathcal{E}_{\text{bf}} + \mathcal{F}_{\text{bf}}) \sqcup (\mathcal{E}_{\text{lb}} + \mathcal{F}_{\text{rb}}).\end{aligned}$$

4. THE b -CALCULUS ON $M_{b,0}$

4.1. The b -pseudodifferential operator on $M_{b,0}$. We now apply the abstract b -calculus to the b -regime of $M_{b,0}$, where $r < 1$ and $r' < 1$. The relevant boundary hypersurfaces here are bf, lb, rb and bf₀.

Suppose $m \in \mathbb{R}$ and $\mathcal{E} = (\mathcal{E}_{\text{lb}}, \mathcal{E}_{\text{rb}}, \mathcal{E}_{\text{bf}}, \mathcal{E}_{\text{bf}_0})$ is the index family for the boundary. The space $\Psi_b^{m, \mathcal{E}}(M_{b,0}; {}^{b,sc}\Omega^{1/2})$ of b -pseudodifferential operators in the b -regime is defined to be the collection of the sums of small b -pseudodifferential operators in $\Psi_b^m(M_{b,0}; {}^{b,sc}\Omega^{1/2})$ and operators with a polyhomogeneous conormal kernel in $\mathcal{A}_{phg}^{\mathcal{E}}(M_{b,0}; {}^{b,sc}\Omega^{1/2})$ supported in the b -regime, where

- $\Psi_b^m(M_{b,0}; {}^{b,sc}\Omega^{1/2})$ is the space of ${}^{b,sc}\Omega^{1/2}$ -valued distributions conormal of degree m to the diagonal $\text{diag}_{M_{b,0}}|_{b\text{-regime}}$, uniformly up to bf and bf₀;
- $\mathcal{A}_{phg}^{\mathcal{E}}(M_{b,0}; {}^{b,sc}\Omega^{1/2})$ is the space of ${}^{b,sc}\Omega^{1/2}$ -valued polyhomogeneous conormal distributions with respect to the index family \mathcal{E} for lb, rb, bf and bf₀.

Adapting the abstract b -calculus, we have analogous composition formulas.

Proposition 6. *For two full b -pseudodifferential operators in the b -regime,*

$$\begin{aligned}A &\in \Psi_b^{m_1, \mathcal{E}}(M_{b,0}, {}^{b,sc}\Omega^{1/2}) \quad \text{with } \mathcal{E} = (\mathcal{E}_{\text{lb}}, \mathcal{E}_{\text{rb}}, \mathcal{E}_{\text{bf}}, \mathcal{E}_{\text{bf}_0}), \\ B &\in \Psi_b^{m_2, \mathcal{F}}(M_{b,0}, {}^{b,sc}\Omega^{1/2}) \quad \text{with } \mathcal{F} = (\mathcal{F}_{\text{lb}}, \mathcal{F}_{\text{rb}}, \mathcal{F}_{\text{bf}}, \mathcal{F}_{\text{bf}_0}),\end{aligned}$$

satisfying $\text{Re}(\mathcal{E}_{\text{rb}} + \mathcal{F}_{\text{lb}}) > 0$, the composition operator AB lies in the space

$$\Psi_b^{m_1+m_2, \mathcal{G}}(M_{b,0}, {}^{b,sc}\Omega^{1/2}) \quad \text{with } \mathcal{G} = (\mathcal{G}_{\text{bf}}, \mathcal{G}_{\text{lb}}, \mathcal{G}_{\text{rb}}, \mathcal{G}_{\text{bf}_0})$$

where the index family \mathcal{G} is given by

$$\begin{aligned}\mathcal{G}_{\text{lb}} &= (\mathcal{E}_{\text{bf}} + \mathcal{F}_{\text{lb}}) \sqcup \mathcal{E}_{\text{lb}}, \\ \mathcal{G}_{\text{rb}} &= (\mathcal{E}_{\text{rb}} + \mathcal{F}_{\text{bf}}) \sqcup \mathcal{F}_{\text{rb}}, \\ \mathcal{G}_{\text{bf}} &= (\mathcal{E}_{\text{bf}} + \mathcal{F}_{\text{bf}}) \sqcup (\mathcal{E}_{\text{lb}} + \mathcal{F}_{\text{rb}}), \\ \mathcal{G}_{\text{bf}_0} &= \mathcal{E}_{\text{bf}_0} + \mathcal{F}_{\text{bf}_0}.\end{aligned}$$

4.2. The indicial operator at bf. We next consider the invertibility of the indicial operator of P_b at bf.

Recall from (5) that P_b is a second order b -differential operator and takes the form

$$-(r\partial_r)^2 - r\tilde{e}(r\partial_r) + \Delta_h - r\tilde{e}(1 - n/2) - r^2(1 + \imath 0) + (1 - n/2)^2.$$

Near $\text{lb} \cap \text{bf}$, we adopt coordinates

$$\{(\tilde{r}, r', y, y', \hbar) : \tilde{r} = r/r'\}.$$

Here \tilde{r} , r' and \hbar are the boundary defining functions for lb, bf and bf₀ respectively. Then the blowup β_b^{-1} lifts P_b to

$$-(\tilde{r}\partial_{\tilde{r}})^2 - \tilde{r}r'\tilde{e}(\tilde{r}\partial_{\tilde{r}}) + \Delta_h - \tilde{r}r'\tilde{e}(1 - n/2) - (\tilde{r}r')^2(1 + \imath 0) + (1 - n/2)^2.$$

The indicial operator $\mathcal{I}(P_b)$ and the indicial family $\mathcal{I}(P_b, \lambda)$ read

$$\mathcal{I}(P_b) = -(\tilde{r}\partial_{\tilde{r}})^2 + \Delta_{h_0} + (1 - n/2)^2.$$

$$\mathcal{I}(P_b, \lambda) = \mathcal{M}(\mathcal{I}(P_b)) = \lambda^2 + \Delta_{h_0} + (1 - n/2)^2,$$

where h_0 denotes $h(0, y, dy)$. Moreover, these operators lift, through the stretched projection $\tilde{\mathcal{P}}_L$, to the b -regime of $M_{b,0}$. They are \hbar -independent so that the face bf_0 is irrelevant for the inverse of $\mathcal{I}(P_b)$ and $\mathcal{I}(P_b, \lambda)$.

Since Δ_{h_0} is the Laplacian on the compact manifold (\mathcal{Y}, h_0) , the resolvent family $(\Delta_{h_0} - \tau)^{-1}$ only has simple poles

$$\sigma_0^2, \sigma_1^2, \sigma_2^2, \dots \quad \text{with} \quad 0 = \sigma_0^2 < \sigma_1^2 < \sigma_2^2 < \dots.$$

Then the indicial family $\mathcal{I}(P_b, \lambda)$ is invertible on $C^\infty(\partial\mathcal{C})$ if and only if $-\lambda^2 - (1 - n/2)^2$ is NOT an eigenvalue of Δ_{h_0} . Hence, we define the indicial roots

$$\lambda_j = \iota((1 - n/2)^2 + \sigma_j^2)^{1/2},$$

and the boundary spectrum

$$\text{spec}_b(\mathcal{I}(P_b)) = \{\pm\lambda_j \in \mathbb{C}\}, \quad \text{Spec}_b(\mathcal{I}(P_b)) = \{(\pm\lambda_j, 0)\}.$$

By adapting the proof from [28, (5.84)], we obtain the inverse of the indicial operator $\mathcal{I}(P_b)$, which is the inverse Mellin transform of the indicial family at bf .

Proposition 7. *The kernel of $\mathcal{I}(P_b)^{-1}$, modulo $\Psi_b^{-2}(M_{b,0}; {}^{b,sc}\Omega^{1/2})$, lies in the space,*

$$(16) \quad \mathcal{A}_{phg}^{(\mathcal{E}_{\text{lb}}, \mathcal{E}_{\text{rb}}, \mathcal{E}_{\text{bf}}, \mathcal{E}_{\text{bf}_0})}(M_{b,0}; {}^{b,sc}\Omega^{1/2}),$$

with $\mathcal{E}_{\text{lb}} = \mathcal{E}_{\text{rb}} = \{(\text{Im } \lambda_j, 0)\}$ and $\mathcal{E}_{\text{bf}} = \mathcal{E}_{\text{bf}_0} = \mathbb{N}_0$.

5. THE ABSTRACT sc -CALCULUS

In this section, we shall review the abstract scattering calculus and Legendrian distributions. The notations in this section will be self-contained and independent of the conic manifold \mathcal{C} .

5.1. The sc -pseudodifferential calculus. Let X be an n -manifold with boundary. It is furnished with local coordinates z in the interior and (x, y) near the boundary with a boundary defining function x .

Consider the Lie algebra,

$$\mathcal{V}_{sc}(X) = \{V \mid V = xW, W \text{ is a } C^\infty \text{ vector field tangent to } \partial X\}.$$

Near ∂X , this Lie algebra, smoothly spanned by the vector fields $x^2\partial_x$ and $x\partial_y$, gives smooth sections of the sc -tangent bundle ${}^{sc}TX$. Corresponding to this, the sc -cotangent bundle ${}^{sc}T^*X$, near ∂X , is generated by dx/x^2 and dy/x . In other words, a sc -cotangent vector, near ∂X , takes the form

$$d(f/x) = \tau dx/x^2 + \mu dy/x$$

for some smooth function f .

The sc -differential operators of order k are the sums of products of at most k sc -vector fields. They act on sc -half densities, which, near the boundary, take the form

$$f \left| \frac{dx dy}{x^{n+1}} \right|^{1/2} \in C^\infty(X; {}^{sc}\Omega^{1/2}).$$

The space of such operators is denoted by $\text{Diff}_{sc}^k(X; {}^{sc}\Omega^{1/2})$.

We extend this to sc -pseudodifferential operators by viewing their Schwartz kernels as conormal distributions on certain double spaces. The appropriate double space to accommodate the sc -pseudodifferential operators is the two-fold blown-up space

$$X_{sc,b} = [X_b^2; \partial \text{diag}_{X_b^2}] = [[X^2; (\partial X)^2]; \partial \text{diag}_{X_b^2}],$$

where $\text{diag}_{X_b^2}$ is the lift of the diagonal of X^2 to X_b^2 . The boundary hypersurfaces and the diagonal diag_{X^2} of X^2 lift via the two-fold blow-up as follows,

$$\begin{aligned} \partial X \times X &\longrightarrow \text{lf}, \\ X \times \partial X &\longrightarrow \text{rf}, \\ (\partial X \times \partial X) \setminus \partial \text{diag}_{X^2} &\longrightarrow \text{mf}, \\ \partial \text{diag}_{X^2} &\longrightarrow \text{sc}, \\ \text{diag}_{X^2} &\longrightarrow \text{diag}_{X_{sc,b}^2}, \end{aligned}$$

where $\text{diag}_{X_{sc,b}^2}$ denotes the diagonal of $X_{sc,b}^2$.

If we view the kernel of a sc -differential operator on X as a distribution on $X_{sc,b}^2$, then it is microlocally supported on and conormal to $\text{diag}_{X_{sc,b}^2}$. In the same spirit,

Definition 8. *The space $\Psi_{sc}^{k,0}(X; {}^{sc}\Omega^{1/2})$ of sc -pseudodifferential operators of degree k consists of all operators with a kernel $\kappa|d\mu_{sc}|^{1/2}$, such that*

- κ defined on $X_{sc,b}^2$ is a distribution, conormal of degree k to $\text{diag}_{X_{sc,b}^2}$, smooth up to sc , rapidly decreasing at lf , rf , mf ,
- $|d\mu_{sc}|^{1/2}$ is the lift of $\pi_l^{*sc}\Omega^{1/2}(X) \otimes \pi_r^{*sc}\Omega^{1/2}(X)$ to X_b^2 , where π_l is the projection from X_b^2 to the left space X and π_r is the projection from X_b^2 to the right space X .

Furthermore, we define the space $\Psi_{sc}^{k,l}(X; {}^{sc}\Omega^{1/2}) = x^l \Psi_{sc}^k(X; {}^{sc}\Omega^{1/2})$.

There are two symbol maps defined for $\Psi_{sc}^{k,0}(X; {}^{sc}\Omega^{1/2})$. One is the usual symbol in the interior of X with respect to ${}^{sc}T^*X$,

$$\sigma_{int}^k : \Psi_{sc}^{k,0}(X, {}^{sc}\Omega^{1/2}) \longrightarrow S^k({}^{sc}T^*X)/S^{k-1}({}^{sc}T^*X),$$

where $S^k(\cdot)$ is the classical symbol space of degree k . This symbol map yields a short exact sequence,

$$0 \longrightarrow \Psi_{sc}^{k-1,0}(X; {}^{sc}\Omega^{1/2}) \longrightarrow \Psi_{sc}^{k,0}(X; {}^{sc}\Omega^{1/2}) \longrightarrow S^k({}^{sc}T^*X)/S^{k-1}({}^{sc}T^*X) \longrightarrow 0.$$

In the meantime, the other symbol map,

$$\sigma_{\partial} : \Psi_{sc}^{k,0}(X, {}^{sc}\Omega^{1/2}) \longrightarrow S^k({}^{sc}T_{\partial X}^*X),$$

is defined as the restriction of the full symbol to $\partial X = \{x = 0\}$. This is well-defined due to the fact $[\mathcal{V}_{sc}(X), \mathcal{V}_{sc}(X)] \subset x\mathcal{V}_{sc}(X)$. It likewise yields a short exact sequence,

$$0 \longrightarrow \Psi_{sc}^{k,1}(X, {}^{sc}\Omega^{1/2}) \longrightarrow \Psi_{sc}^{k,0}(X; {}^{sc}\Omega^{1/2}) \longrightarrow S^k({}^{sc}T_{\partial X}^*X)/S^{k-1}({}^{sc}T_{\partial X}^*X) \longrightarrow 0.$$

If one radially compactifies the fibre of ${}^{sc}T^*X$, then the two symbol maps combined give a joint symbol of $P \in \text{Diff}_{sc}^k(X; {}^{sc}\Omega^{1/2})/\text{Diff}_{sc}^{k-1}(X; {}^{sc}\Omega^{1/2})$,

$$j_{sc}^k(P) = \left(\frac{\sigma_{int}^k(P)}{|\xi|^k} \Big|_{{}^{sc}S^*X}, \sigma_{\partial}(P) \right) \in S^k(C_{sc}(X)),$$

where $C_{sc}(X) = {}^{sc}S^*X \cup {}^{sc}T_{\partial X}^*X$. The joint symbol map j_{sc}^k extends from $\text{Diff}_{sc}^k(X; {}^{sc}\Omega^{1/2})$ to $\Psi_{sc}^{k,0}(X; {}^{sc}\Omega^{1/2})$ multiplicatively,

$$j_{sc}^{k_1}(P_1) \cdot j_{sc}^{k_2}(P_2) = j_{sc}^{k_1+k_2}(P_1 P_2),$$

but also admits a short exact sequence,

$$0 \longrightarrow \Psi_{sc}^{k-1,1}(X, {}^{sc}\Omega^{1/2}) \longrightarrow \Psi_{sc}^{k,0}(X; {}^{sc}\Omega^{1/2}) \longrightarrow S^k(C_{sc}(X))/S^{k-1}(C_{sc}(X)) \longrightarrow 0.$$

As in the classical theory, the ellipticity at a point in $C_{sc}(X)$ of a sc -pseudodifferential operator A means its joint symbol $j_{sc}^k(A)$ does not vanish, whilst the characteristic variety $\Sigma(A)$ of A is the zero set of $j_{sc}^k(A)$. For example, the interior symbol of the sc -differential operator

$$-(x^2\partial_x)^2 + (n-1)x^3\partial_x + x^2\Delta_h - 1$$

is obviously elliptic but the boundary symbol vanishes in a subset of ${}^{sc}T_{\partial X}^*X$. To understand the resolvent of such an operator, we will need the notion of the sc -wavefront set of a tempered distribution u on X ,

$${}^{sc}\text{WF}(u) = \{q \in C_{sc}(X) \mid \chi u \notin \dot{C}^\infty(X) \text{ for any } \chi \in \Psi_{sc}^{0,0}(X) \text{ elliptic at } q\}.$$

In the classical propagation of singularities of hyperbolic equations, the wavefront set of the fundamental solution consists of the bicharacteristic flow curves, which comprise a Lagrangian of the cotangent bundle. If we write the smooth sections of ${}^{sc}T^*X$ as

$$\tau \frac{dx}{x^2} + \mu \frac{dy}{x},$$

then ${}^{sc}T^*X$ is a symplectic manifold with a non-degenerate 2-form,

$$\omega = d\tau \wedge \frac{dx}{x^2} + d\mu \wedge \frac{dy}{x} - \mu \cdot dy \wedge \frac{dx}{x^2}.$$

The symplectic form ω on ${}^{sc}T^*X$ induces a contact structure on ${}^{sc}T_{\partial X}^*X$ with a non-degenerate 1-form

$$\chi = \iota_{x^2\partial_x}\omega = -d\tau + \mu \cdot dy.$$

The integral curves of the Hamilton vector field determined by the contact form are called the bicharacteristics as well. For sc -pseudodifferential operators, we have a variant of the theorem of microlocalization and propagation of singularities.

Proposition 9. *Suppose $A \in \Psi_{sc}^{k,0}(X, {}^{sc}\Omega^{1/2})$ has a real boundary symbol. For a tempered distribution u on X but smooth in the interior,*

$$\text{WF}_{sc}(Au) \subset \text{WF}_{sc}(u)$$

and $\text{WF}_{sc}(u) \setminus \text{WF}_{sc}(Au)$ is a union of maximally extended bicharacteristics of A inside $\Sigma(A) \setminus \text{WF}_{sc}(Au)$.

5.2. The Legendrian distributions. Due to the non-ellipticity of the sc -Laplacian in a subset of $T_{\partial X}^*X$, it is necessary to formulate a theory of Lagrangian distributions over the boundary, i.e. Legendrian distributions. We review some details from [17, Section 2] and [18, Section 2.2].

In contact geometry, an isotropic submanifold is a submanifold of a contact manifold, on which the contact form vanishes, and a Legendrian is a maximal isotropic submanifold. Suppose \mathcal{X} is a suitable double space on which the resolvent kernel is defined, F is an isotropic submanifold of ${}^{sc}T_{\partial\mathcal{X}}^*\mathcal{X}$ such that the Hamiltonian vector field is nowhere tangent to F , the union of bicharacteristics passing through F forms a Legendrian L . The classical theorem of propagation of singularities tells us that the resolvent has to be microlocally supported on the Legendrian L , i.e. it is a Legendrian distribution.

Definition 10. On an n -manifold X furnished with local coordinates (x, y) near ∂X , a Legendrian distribution $u \in I^m(X; L; {}^{sc}\Omega^{1/2})$ locally takes the form

$$u = x^{m+n/4-k/2} (2\pi)^{-k/2-n/4} \int_{\mathbb{R}^k} e^{i\phi(y,v)/x} a(x, y, v) dv,$$

where

- the phase function ϕ is non-degenerate, i.e.
 $d(\partial_{v_i}\phi)$ are linearly independent at $\{(y, v) : d_v\phi = 0\}$ for $i = 1, \dots, k$,
- ϕ locally parametrizes the Legendrian L ,
 $L = \{(y, d_{(x,y)}(\phi(y, v)/x)) : d_v\phi(y, v) = 0\}$,
- the amplitude a is a compactly supported smooth section of the sc -half density bundle.

Suppose λ is a set of functions in the (y, v) -space such that $(d_v\phi, \lambda)$ form local coordinates on $\{(y, v) : d_v\phi = 0\}$. The symbol of u is defined by

$$\sigma_{(x,y),\phi}^m(u) = a(0, y, v)|_{\{d_v\phi=0\}} \left| \frac{\partial(d_v\phi, \lambda)}{\partial(y, v)} \right|^{-1/2} |d\lambda|^{1/2}.$$

The symbol is neither coordinate invariant nor phase function invariant, but changes by the following transition functions

$$(17) \quad \begin{aligned} \sigma_{(x',y'),\phi}^m(u) &= \sigma_{(x,y),\phi}^m(u) \left(\frac{x'}{x}\right)^{n/4-m} e^{-i((x'/x)\mu \cdot \partial_{x'}y - \tau \partial_{x'}(x'/x))|_{x=0}}, \\ \sigma_{(x,y),\psi}^m(u) &= \sigma_{(x,y),\phi}^m(u) e^{i\pi(\text{sign} d_{vv}^2\psi - \text{sign} d_{vv}^2\phi)/4}. \end{aligned}$$

These two transition functions define a line bundle $E \otimes M$. The symbol map of degree m is invariantly defined

$$\sigma^m : I^m(X, L; {}^{sc}\Omega^{1/2}) \longrightarrow C^\infty(L; \Omega^{1/2}(L) \otimes S^{[m]}),$$

where $S^{[m]} = |N^*\partial X|^{m-n/4} \otimes E \otimes M$ and the bundle $N^*\partial X$ is given by differentials of functions vanishing at (each) boundary hypersurface. See [32, Section 12-13] and [20, Section 4.6] for more information of this symbol space and generalizations to manifolds with corners.

This symbol map gives an exact sequence,

$$0 \longrightarrow I^{m+1}(X, L; {}^{sc}\Omega^{1/2}) \longrightarrow I^m(X, L; {}^{sc}\Omega^{1/2}) \longrightarrow C^\infty(L; {}^{sc}\Omega^{1/2}(L) \otimes S^{[m]}) \longrightarrow 0,$$

and the symbol calculus of $u \in I^m(X; L; {}^{sc}\Omega^{1/2})$ with a sc -pseudodifferential operator $P \in \Psi_{sc}(X; {}^{sc}\Omega^{1/2})$ with a real principal symbol vanishing on L ,

$$(18) \quad (-i\mathcal{L}_{H_p} - i(1/2 + m - n/4)\partial_\tau p + p_{\text{sub}})\sigma^m(u),$$

where H_p denotes the Hamilton vector field of the principal symbol of P and p_{sub} is the boundary subprincipal symbol of P . Explicitly, if the full left symbol of a sc -pseudodifferential operator P takes the form

$$p(y, \tau, \mu) + xp_1(y, \tau, \mu) + O(x^2)$$

in the scattering cotangent bundle with local coordinates $\{(x, y, \tau, \mu)\}$, its boundary subprincipal symbol is written as the following expansion in x ,

$$p_{\text{sub}} = p_1 + \frac{i}{2} \left(\frac{\partial^2 p}{\partial y_i \partial \mu_i} - (n-1) \frac{\partial p}{\partial \tau} + \mu_i \frac{\partial^2 p}{\partial \mu_i \partial \tau} \right).$$

For instance, the full left symbol of P_{sc} defined in (6) reads

$$\tau^2 + h^{ij} \mu_i \mu_j + \rho(n-1 - xe(x, y))i\tau - 1,$$

where this x is the defining function in \mathcal{C} . If we adopt coordinates $\{(\rho, y, \tau, \mu)\}$, it follows that the boundary subprincipal symbol is equal to

$$(19) \quad -ixe(x, y)\tau + i\left(\frac{\partial h^{ij}}{\partial y_i}\mu_j\right).$$

When $\mu = 0$, the second term vanishes and then the subprincipal symbol is just equal to the first term $-ixe(x, y)\tau$.

Intersecting Legendrian distributions are used to describe the resolvent near the intersection of the propagating Legendrian with other Legendrians which appear in the microlocal structure of the resolvent. The Legendrian pair accommodates the sc -wavefront set of the resolvent. Assume $\tilde{L} = (L_0, L_1)$ is a pair of Legendrians intersecting cleanly at ∂L_1 . Then near a point $q \in \partial L_1$ there is a unique local phase function $\phi(y, v, s) = \phi_0(y, v) + s\phi_1(y, v, s)$ (up to a diffeomorphism) parameterizing the pair

$$\begin{aligned} L_0 &= \{(y, d_{(x,y)}(\phi/x)) | s = 0, d_v\phi = 0\} \\ L_1 &= \{(y, d_{(x,y)}(\phi/x)) | s \geq 0, d_s\phi = 0, d_v\phi = 0\}. \end{aligned}$$

Definition 11. The space $I^m(X; \tilde{L}; {}^{sc}\Omega^{1/2})$ consists of the sums of elements in

$$I^m(X; L_1 \setminus \partial L_1; {}^{sc}\Omega^{1/2}) + I^{m+1/2}(X; L_0; {}^{sc}\Omega^{1/2}),$$

and distributions in a coordinate patch $\{(x, y)\}$ near ∂X with an expression

$$x^{m+n/4-(k+1)/2} \int_0^\infty ds \int_{\mathbb{R}^k} e^{i\phi(y,v,s)} a(x, y, v, s) dv,$$

where a is a compactly supported smooth sc -half density.

The symbol of $u \in I^m(X; \tilde{L}; {}^{sc}\Omega^{1/2})$ is a smooth section of a bundle over $L_0 \cup L_1$. Let ρ_1 be a boundary defining function for $\partial L_1 = L_0 \cap L_1$ within L_0 , and ρ_0 a boundary defining function for ∂L_1 within L_1 . On L_0 but away from L_1 , the symbol of u lies in the space

$$\rho_1^{-1} C^\infty(\Omega^{1/2}(L_0) \otimes S^{[m+1/2]}(L_0)) = \rho_1^{-1/2} C^\infty(\Omega_b^{1/2}(L_0 \setminus \partial L_1) \otimes S^{[m+1/2]}(L_0)),$$

and the symbol of u on L_1 takes values in the space

$$C^\infty(\Omega^{1/2}(L_1) \otimes S^{[m]}(L_1)) = \rho_0^{1/2} C^\infty(\Omega_b^{1/2}(L_1) \otimes S^{[m]}(L_1)).$$

Here $\Omega^{1/2}(L)$ is the half density bundle over the Legendrian L , and $\Omega_b^{1/2}(L)$ is the b -half density bundle near intersections of the Legendrian pairs. Then the symbol space $C^\infty(\tilde{L}; {}^b\Omega^{1/2}(\tilde{L}) \otimes S^{[m]}(\tilde{L}))$ for $I^m(X; \tilde{L}; {}^{sc}\Omega^{1/2})$ consists of smooth section pairs (a, b) such that

$$(20) \quad \begin{aligned} a &\in \rho_1^{-1/2} C^\infty(\Omega_b^{1/2}(L_0 \setminus \partial L_1) \otimes S^{[m+1/2]}(L_0)) && \text{on } L_0 \setminus \partial L_1 \\ b &\in \rho_0^{1/2} C^\infty(\Omega_b^{1/2}(L_1) \otimes S^{[m]}(L_1)) && \text{on } L_1 \\ \rho_1^{1/2} b &= e^{i\pi/4} (2\pi)^{1/4} \rho_0^{-1/2} a && \text{at } L_0 \cap L_1 \end{aligned}.$$

This symbol map likewise yields an exact sequence,

$$(21) \quad 0 \longrightarrow I^{m+1}(X; \tilde{L}; {}^{sc}\Omega^{1/2}) \longrightarrow I^m(X; \tilde{L}; {}^{sc}\Omega^{1/2}) \longrightarrow C^\infty(\tilde{L}; \Omega_b^{1/2}(\tilde{L}) \otimes S^{[m]}) \longrightarrow 0.$$

More care is needed to deal with a special type of Legendrian pair, which has a conic singularity at the intersection. We say that $\tilde{L} = (L, L^\sharp)$ is a conic Legendrian pair, if

- L^\sharp is a projectable Legendrian, which means that the projection map ${}^{sc}T^*X|_{L^\sharp} \rightarrow \partial X$ is a diffeomorphism,
- $\tau \neq 0$ on L^\sharp , which implies that L^\sharp can be parametrized by the phase function 1,
- L is an open Legendrian such that $\tilde{L} \setminus L$ is contained in L^\sharp ,

- \bar{L} has at most a conic singularity at L^\sharp , which means that $\hat{L} = \beta_\sharp^{-1}\bar{L}$ is transverse to $\beta_\sharp^{-1}\{x = 0, \mu = 0\}$, where the blowdown map β_\sharp defines the blown-up space $[^{sc}T^*X; \{x = 0, \mu = 0\}]$.

A local parametrization of \tilde{L} near $q \in \bar{L} \cap L^\sharp$ is a function $\phi(y, v, s) = 1 + s\psi(y, v, s)$ defined in a neighbourhood of $q' = (y_0, v_0, 0)$ in $\partial X \times \mathbb{R}^k \times [0, \infty)$ satisfying

- 1 parametrizes L^\sharp near q ,
- $d_v\phi = 0$ at q' ,
- $q = (y, d_{(x,y)}(\phi/x))(q')$,
- ϕ satisfies the non-degeneracy hypothesis

$ds, d\psi$, and $d\left(\frac{\partial\psi}{\partial v_i}\right)$ are linearly independent at q' , $1 \leq i \leq k$,

- \hat{L} has the expression near q in terms of coordinates $\{(x/|\mu|, y, \tau, |\mu|, \hat{\mu})\}$,

$$\hat{L} = \{(0, y, -\phi, sd_y\psi, \widehat{d_y\psi}) \mid d_s\phi = 0, d_v\psi = 0, s \geq 0\}.$$

Then we define Legendrians associated with this local parametrization.

Definition 12. A Legendrian distribution of order (m, p) associated to (L, L^\sharp) is a sc-half density of the form $u = u_0 + (\sum_{i=1}^N u_i)\nu$, where

- ν is a smooth sc-half density,
- u_0 is a Legendrian distribution lying in

$$I_c^m(X; L; {}^{sc}\Omega^{1/2}) + I^p(X; L^\sharp; {}^{sc}\Omega^{1/2}),$$

where the subscript c indicates that the wavefront set is contained in the interior of the Legendrian L ,

- u_j , near the boundary, has an expression

$$\int_0^\infty ds \int e^{i\phi_j(y,v,s)/x} a_j(y, v, x/s, s) \left(\frac{x}{s}\right)^{m+n/4-(k+1)/2} s^{p+n/4-1} dv,$$

where ϕ_j locally parametrizes (L, L^\sharp) and $a_j \in C^\infty(X \times \mathbb{R}^k \times [0, \infty) \times [0, \infty))$, with compact support in $v, x/s$ and s .

The space of such Legendrian distributions is denoted by $I^{m,p}(X; \tilde{L}; {}^{sc}\Omega^{1/2})$.

The symbol map is defined on \hat{L} and yields an exact sequence

$$\begin{aligned} 0 \longrightarrow I^{m+1,p}(X; \tilde{L}; {}^{sc}\Omega^{1/2}) &\longrightarrow I^m(X; \tilde{L}; {}^{sc}\Omega^{1/2}) \\ &\longrightarrow s^{p-m}C^\infty(\hat{L}, \Omega_b^{1/2}(\hat{L}) \otimes S^{[m]}(\hat{L})) \longrightarrow 0. \end{aligned}$$

6. THE FIBRED sc -CALCULUS ON $M_{b,0}$

We now come back to the sc -regime of the space $M_{b,0}$ where the resolvent kernel of P_{sc} lives. Similar to the resolvent of the sc -Laplacian, the resolvent kernel of P_{sc} consists of sc -pseudodifferential operators and Legendrian distributions. To understand this microlocal structure, we apply to $M_{b,0}$ the sc -calculus and Legendrian machinery.

6.1. The sc -pseudodifferential operators on $M_{sc,b,0}$. The appropriate stretched double space for the sc -pseudodifferential calculus is $M_{sc,b,0}$ instead of $M_{b,0}$ as required in the abstract sc -calculus.

The space $\Psi_{sc}^k(M_{sc,b,0}; {}^{b,sc}\Omega^{1/2})$ is defined to be the collection of ${}^{b,sc}\Omega^{1/2}$ -valued distributions in the sc -regime of $M_{sc,b,0}$, which are conormal of degree k to $\text{diag}_{M_{sc,b,0}}$, smooth up to sc and bf_0 , rapidly decreasing at lf, rf, mf .

6.2. The fibred sc -bundles on $M_{b,0}$. Dealing with the Legendrian structure near the corners in the sc -regime requires working on the finer structure of boundary fibrations. We shall see that these fibrations are trivial at bf_0 but non-trivial at mf , lf and rf such that the regions near the corners $\text{mf} \cup \text{lf}$ and $\text{mf} \cup \text{rf}$ are of interest. In the meantime, since lf and rf are away from the diagonal conormal bundle and the fibration is trivial at bf_0 , we can blow down the sc face and return to $M_{b,0}$.

Now we are concerned about four boundary faces $\text{mf}, \text{lf}, \text{rf}, \text{bf}_0$ of $M_{b,0}$. But it suffices to discuss the boundary fibration and Legendrian structures away from rf as the structure away from lf is analogous. In addition, we shall denote by ϕ_\bullet the fibration at the face \bullet , by Z_\bullet the base of ϕ_\bullet , by F_\bullet the fibre of \bullet , i.e.

$$\phi_\bullet : \bullet \longrightarrow Z_\bullet.$$

Recall the local expressions of $\mathcal{V}_{b,sc}(M_{b,0})$ in the sc -regime away from rf .

- The region near $\text{lf} \cap \text{mf} \cap \text{bf}_0$ admits local coordinates

$$(\tilde{\rho}, \rho', y, y', x),$$

where $x, \rho', \tilde{\rho} = \rho/\rho'$ are the boundary defining functions of $\text{bf}_0, \text{mf}, \text{lf}$ respectively. The Lie algebra $\mathcal{V}_{b,sc}(M_{b,0})$ thus lifts to

$$\text{span}\{\tilde{\rho}^2 \rho' \partial_{\tilde{\rho}}, \rho'^2 \partial_{\rho'}, \rho \partial_y, \rho' \partial_{y'}, x \partial_x\} \quad \text{near } \text{lf} \cap \text{mf} \cap \text{bf}_0.$$

- The region near the interior of lf has local coordinates

$$(\rho, y, z', x) \quad \text{where } z' \text{ is in the interior of } \mathcal{C}.$$

The Lie algebra $\mathcal{V}_{b,sc}(M_{b,0})$ is then locally expressed by

$$\text{span}\{\rho^2 \partial_\rho, \rho \partial_y, \partial_{z'}, x \partial_x\} \quad \text{near the interior of } \text{lf}.$$

We are interested in the restrictions of $\mathcal{V}_{b,sc}(M_{b,0})$ to the boundary hypersurfaces and the induced fibrations.

- The Lie algebra $\mathcal{V}_{b,sc}(M_{b,0})$ restricted to mf is simply spanned by $x \partial_x$. This implies the fibration over mf reads

$$\phi_{\text{mf}} : \text{mf} \longrightarrow Z_{\text{mf}}, \quad (\tilde{\rho}, y, y', x) \longmapsto (\tilde{\rho}, y, y'),$$

and the base Z_{mf} can be identified with $\text{mf} \cap \text{bf}_0$.

- The Lie algebra $\mathcal{V}_{b,sc}(M_{b,0})$ restricted to lf takes the form

$$\begin{cases} \text{span}\{\rho'^2 \partial_{\rho'}, \rho' \partial_{y'}, x \partial_x\} & \text{near } \text{lf} \cap \text{mf} \cap \text{bf}_0 \\ \text{span}\{\partial_{z'}, x \partial_x\} & \text{near the interior of } \text{lf} \end{cases}.$$

This yields the fibration at lf , which

$$\phi_{\text{lf}} : \text{lf} \longrightarrow Z_{\text{lf}}, \quad \begin{cases} (\rho', y, y', x) \longmapsto y & \text{near } \text{lf} \cap \text{mf} \cap \text{bf}_0 \\ (y, z', x) \longmapsto y & \text{near the interior of } \text{lf} \end{cases}.$$

The base Z_{lf} of the fibration ϕ_{lf} is a compact manifold without boundary, and the fibres of ϕ_{lf} intersect mf transversally.

- The Lie algebra $\mathcal{V}_{b,sc}(M_{b,0})$ restricted to bf_0 reduces to

$$\text{span}\{\tilde{\rho}^2 \rho' \partial_{\tilde{\rho}}, \rho'^2 \partial_{\rho'}, \rho \partial_y, \rho' \partial_{y'}\} \quad \text{near } \text{lf} \cap \text{mf} \cap \text{bf}_0.$$

This gives a trivial fibration at bf_0 , which does not really matter in the fibred sc -calculus.

Denote $\Phi = \{\phi_{\text{mf}}, \phi_{\text{lf}}, \phi_{\text{rf}}\}$ and $\tilde{M}_{b,0} = M_{b,0} \setminus (\text{lf} \cup \text{rf})$. The Lie algebra $\mathcal{V}_{s\Phi}(M_{b,0})$ of the fibred sc -vector fields, with respect to the stratification Φ and total boundary defining function ρ of $\text{lf} \cup \text{mf}$, consists of smooth vector fields V such that

- $V \in \mathcal{V}_b(M_{b,0})$,
- $V\rho \in \rho^2 C^\infty(M_{b,0})$,
- V is tangent to the fibres of Φ .

The Lie algebra $\mathcal{V}_{s\Phi}(M_{b,0})$ generates the differential operator algebra $\text{Diff}_{s\Phi}(M_{b,0})$, which are smooth sections of the fibred sc -bundle ${}^{s\Phi}TM_{b,0}$. Its dual bundle is denoted by ${}^{s\Phi}T^*M_{b,0}$. The wedge product of the basis of ${}^{s\Phi}T^*M_{b,0}$ determines the fibred sc -density bundle ${}^{s\Phi}\Omega(M_{b,0})$. One can write down the expression of ${}^{s\Phi}T^*M_{b,0}$ and ${}^{s\Phi}\Omega(M_{b,0})$ in local coordinates.

- Near $\text{mf} \cap \text{lf} \cap \text{bf}_0$, we have local coordinates $(\rho', \tilde{\rho}, y, y', x)$. The basis of ${}^{s\Phi}T^*M_{b,0}$ is given by

$$\left\{ \frac{d\rho}{\rho^2}, \frac{d\tilde{\rho}}{\rho}, \frac{dy}{\rho}, \frac{dy'}{\rho'}, \frac{dx}{x} \right\} \quad \text{or} \quad \left\{ \frac{d\rho}{\rho^2}, \frac{d\rho'}{\rho'^2}, \frac{dy}{\rho}, \frac{dy'}{\rho'}, \frac{dx}{x} \right\}.$$

For $q \in {}^{s\Phi}T^*M_{b,0}$, we write

$$q = \tau \frac{d\rho}{\rho^2} + \eta \frac{d\tilde{\rho}}{\rho} + \mu \cdot \frac{dy}{\rho} + \mu' \cdot \frac{dy'}{\rho'} + \xi \frac{dx}{x}$$

$$\quad \text{or} \quad q = \tilde{\tau} \frac{d\rho}{\rho^2} + \tau' \frac{d\rho'}{\rho'^2} + \mu \cdot \frac{dy}{\rho} + \mu' \cdot \frac{dy'}{\rho'} + \xi \frac{dx}{x},$$

where $\tau = \tilde{\tau} + \tilde{\rho}\tau'$ and $\eta = -\tau'$. The fibred sc -density ${}^{s\Phi}\Omega(M_{b,0})$ locally reads

$$\left| \frac{d\rho d\tilde{\rho} dy dy' dx}{\rho^{n+2} \rho'^{n-1} x} \right| \quad \text{or} \quad \left| \frac{d\rho dy d\rho' dy' dx}{\rho^{n+1} \rho'^{n+1} x} \right|.$$

- Near the interior of lf , we have local coordinates $\{(\rho, y, z')\}$. The basis of ${}^{s\Phi}T^*M_{b,0}$ is given by

$$\left\{ \frac{d\rho}{\rho^2}, \frac{dy}{\rho}, dz', \frac{dx}{x} \right\}.$$

For $q \in {}^{s\Phi}T^*M_{b,0}$, we write

$$q = \tau \frac{d\rho}{\rho^2} + \mu \cdot \frac{dy}{\rho} + \zeta' \cdot dz' + \xi \frac{dx}{x}.$$

The fibred sc -density ${}^{s\Phi}\Omega(M_{b,0})$ locally reads

$$\left| \frac{d\rho dy dz' dx}{\rho^{n+1} x} \right|.$$

We are interested in two natural subbundles of ${}^{s\Phi}T_{\text{lf}}^*M_{b,0}$.

- The restriction of ${}^{s\Phi}T_{\text{lf}}^*M_{b,0}$ to each fibre F_{lf} of lf is isomorphic to the sc -cotangent bundle of F_{lf} , denoted by ${}^{b,sc}T^*(F_{\text{lf}}; \text{lf})$. Recall that the fibre F_{lf} is furnished with local coordinates

$$\begin{cases} (\rho', y', x) & \text{near } \text{lf} \cap \text{mf} \cap \text{bf}_0 \\ (z', x) & \text{near the interior of } \text{lf} \end{cases}.$$

The basis of ${}^{b,sc}T^*(F_{\text{lf}}; \text{lf})$ is given by

$$\begin{cases} \text{span} \left\{ \frac{d\rho'}{\rho'^2}, \frac{dy'}{\rho'}, \frac{dx}{x} \right\} & \text{near } \text{lf} \cap \text{mf} \cap \text{bf}_0 \\ \text{span} \left\{ dz', \frac{dx}{x} \right\} & \text{near the interior of } \text{lf} \end{cases}.$$

- The quotient bundle, ${}^{s\Phi}T_{\text{lf}}^*M_{b,0}/{}^{b,sc}T^*(F_{\text{lf}}; \text{lf})$, is the pull back of a bundle ${}^{s\Phi}N^*Z_{\text{lf}}$ from the base Z_{lf} of the fibration ϕ_{lf} , and

$${}^{s\Phi}N^*Z_{\text{lf}} = {}^{b,sc}T_{Z_{\text{lf}} \times \{0\}}^*((Z_{\text{lf}})_y \times [0, \epsilon)_\rho).$$

The induced map,

$$\begin{aligned} \tilde{\phi}_{\text{lf}} : {}^{s\Phi}T_{\text{lf}}^*M_{b,0} &\rightarrow {}^{s\Phi}N^*Z_{\text{lf}}, \\ \left\{ \begin{array}{ll} (\rho', y, y', x, \tau, \eta, \mu, \mu', \xi) \mapsto (y, \tau, \mu) & \text{near } \text{lf} \cap \text{mf} \cap \text{bf}_0 \\ (z', y, x, \tau, \mu, \zeta', \xi) \mapsto (y, \tau, \mu) & \text{near the interior of lf} \end{array} \right. \end{aligned}$$

has fibres isomorphic to ${}^{b,sc}T^*F_{\text{lf}}$.

6.3. The contact structures on fibred sc -bundles. We next introduce the contact structures and Legendrians in ${}^{s\Phi}T^*M_{b,0}$. The symplectic form of ${}^{s\Phi}T^*M_{b,0}$, denoted by ω , induces the following contact forms on ${}^{s\Phi}T_{\text{mf}}^*M_{b,0}$, ${}^{s\Phi}N^*Z_{\text{lf}}$ and ${}^{b,sc}T_{\partial F_{\text{lf}}}^*(F_{\text{lf}}; \text{lf})$ respectively.

- At mf , we use the following local expression for ${}^{s\Phi}T^*M_{b,0}$

$$\tau \frac{d\rho}{\rho^2} + \eta \frac{d\tilde{\rho}}{\rho} + \mu \cdot \frac{dy}{\rho} + \mu' \cdot \frac{dy'}{\rho'} + \xi \frac{dx}{x}$$

It follows that the symplectic form ω takes the form

$$\begin{aligned} d\tau \wedge \frac{d\rho}{\rho^2} + d\eta \wedge \frac{d\tilde{\rho}}{\rho} + d\mu \wedge \frac{dy}{\rho} + d\mu' \wedge \frac{dy'}{\rho'} + d\xi \wedge \frac{dx}{x} \\ - \eta d\tilde{\rho} \wedge \frac{d\rho}{\rho^2} - \mu \cdot dy \wedge \frac{d\rho}{\rho^2} + \mu' \cdot dy' \wedge \frac{d\tilde{\rho}}{\rho} - \frac{\tilde{\rho}\mu' \cdot dy' \wedge d\rho}{\rho^2}. \end{aligned}$$

Then ω induces a contact form χ on ${}^{s\Phi}T_{\text{mf}}^*M_{b,0}$,

$$\chi = \iota_{\rho^2 \partial_\rho}(\omega) = -d\tau + \eta d\tilde{\rho} + \mu \cdot dy + \tilde{\rho}\mu' \cdot dy'.$$

- At the corner $\text{mf} \cap \text{lf}$, the one-form χ vanishes identically on ${}^{b,sc}T^*(F_{\text{lf}}; \text{lf})$ but we have a contact form on ${}^{s\Phi}N^*Z_{\text{lf}}$. In local coordinates, this is

$$-d\tau + \mu \cdot dy.$$

- At ∂F_{lf} , we use the following local expression for ${}^{s\Phi}T^*M_{b,0}$

$$\tilde{\tau} \frac{d\rho}{\rho^2} + \tau' \frac{d\rho'}{\rho'^2} + \mu \cdot \frac{dy}{\rho} + \mu' \cdot \frac{dy'}{\rho'} + \xi \frac{dx}{x}.$$

Then ${}^{b,sc}T_{\partial F_{\text{lf}}}^*(F_{\text{lf}}; \text{lf})$ admits a contact form of the form

$$-d\tau' + \mu' \cdot dy'.$$

By incorporating these contact structures, we define Legendrians in ${}^{s\Phi}T^*M_{b,0}$.

Definition 13. A Legendrian L of ${}^{s\Phi}T^*M_{b,0}$ is a Legendrian of ${}^{s\Phi}T_{\text{mf}}^*M_{b,0}$ such that

- L is transversal to ${}^{s\Phi}T_{\text{lf} \cap \text{mf}}^*M_{b,0}$ and ${}^{s\Phi}T_{\text{rf} \cap \text{mf}}^*M_{b,0}$;
- the maps $\tilde{\phi}_{\text{lf}}$ and $\tilde{\phi}_{\text{rf}}$ induce fibrations

$$\begin{aligned} \check{\phi}_{\text{lf}} : L \cap {}^{s\Phi}T_{\text{lf} \cap \text{mf}}^*M_{b,0} &\rightarrow L_1 \\ \check{\phi}_{\text{rf}} : L \cap {}^{s\Phi}T_{\text{rf} \cap \text{mf}}^*M_{b,0} &\rightarrow L'_1, \end{aligned}$$

where L_1 and L'_1 are Legendrians of ${}^{s\Phi}N^*Z_{\text{lf}}$ and ${}^{s\Phi}N^*Z_{\text{rf}}$;

- the fibres of $\check{\phi}_{\text{lf}}$ and $\check{\phi}_{\text{rf}}$ are Legendrians of ${}^{b,sc}T_{\partial F_{\text{lf}}}^*F_{\text{lf}}$ and ${}^{b,sc}T_{\partial F_{\text{rf}}}^*F_{\text{rf}}$.

6.4. The Legendrian distributions on $M_{b,0}$. We shall apply to $M_{b,0}$ the parametrization of the Legendrians and local expressions of Legendrian distributions. We only focus on the part away from rf, as the part away from lf is defined analogously.

We will say that a function $\phi(\tilde{\rho}, y, y', v, w)/\rho$ with $v \in \mathbb{R}^k$ and $w \in \mathbb{R}^l$ is a non-degenerate parametrization of L locally near $q \in {}^{s\Phi}T_{\text{mf} \cap \text{lf}}^* M_{b,0} \cap L$ which is given in local coordinates as $(\tilde{\rho}_0, y_0, y'_0, x_0, \tau_0, \eta_0, \mu_0, \mu', \xi)$ with $\tilde{\rho}_0 = 0$, if

- ϕ has the form

$$\phi(\tilde{\rho}, y, y', v, w) = \phi_1(y, v) + \tilde{\rho}\phi_2(\tilde{\rho}, y, y', v, w),$$

such that ϕ_1, ϕ_2 are smooth in neighbourhoods of $(y_0, v_0) \in \partial\mathcal{C} \times \mathbb{R}^k$ and $q' = (0, y_0, y'_0, v_0, w_0) \in ([0, 1]_{\tilde{\rho}} \times (\partial\mathcal{C})^2) \times \mathbb{R}^{k+k'}$ respectively with

$$(0, y_0, y'_0, x_0, d_{(\rho, \tilde{\rho}, y, y')}(\phi/\rho)(q'), 0) = q, \quad d_v\phi(q') = 0, \quad d_w\phi_2(q') = 0;$$

- ϕ is non-degenerate in the sense that

$$d_{(y,v)} \frac{\partial \phi_1}{\partial v}, \quad d_{(y',w)} \frac{\partial \phi_2}{\partial w}$$

are independent at (y'_0, v_0) and q' respectively;

- locally near q , L is given by

$$L = \{(\tilde{\rho}, y, y', x, d_{(\rho, \tilde{\rho}, y, y')}(\phi/\rho), 0) : d_v\phi = 0, \quad d_w\phi_2 = 0\}.$$

Such a phase function does parametrize the three Legendrians affiliated with L ,

- ϕ/ρ is a non-degenerate phase function for the Legendrian L in ${}^{s\Phi}T_{\text{int}(\text{mf})}^* M_{b,0}$;
- ϕ_1 parametrizes L_1 ;
- $\phi_2(0, y, \cdot, v, \cdot)$, for fixed $(y, v) \in \{(y, v) : d_v\phi_1 = 0\}$, parametrizes the fibre of $\tilde{\phi}_{\text{lf}}$.

Remark 14. The phase function ϕ is indeed independent of x (the boundary defining function of bf_0) as well as its dual cotangent variable ξ , since the three contact forms introduced above are all independent of x and ξ . Consequently, we see x freely varies in $[0, 1)$ and $\xi \equiv 0$ on the Legendrian L , i.e. L extends in x smoothly to $\text{bf}_0 \cap (\text{mf} \cup \text{lf})$.

Away from $\text{lf} \cup \text{rf}$, we only have the contact structure in ${}^{b,sc}T_{\text{mf}}^* \tilde{M}_{b,0}$. The compressed bundle ${}^{b,sc}T_{\text{mf}}^* \tilde{M}_{b,0}$ is a sc -cotangent bundle with respect to mf which is irrelevant to the fibred sc -bundle. The Legendrian distributions are thus defined as in the abstract calculus. Specifically,

Definition 15. The space $I_{sc,c}^m(\tilde{M}_{b,0}; L; {}^{b,sc}\Omega^{1/2})$ of Legendrian distribution associated with a Legendrian $L \subset {}^{b,sc}T_{\text{mf}}^* \tilde{M}_{b,0}$ consists of distributions the form

$$u = \rho'^{m+n/2-k/2} (2\pi)^{-k/2-n/2} \int_{\mathbb{R}^k} e^{i\phi(\tilde{\rho}, y, y', v)/\rho'} a(\tilde{\rho}, \rho', y, y', x, v) dv,$$

where the phase function ϕ is non-degenerate and locally parametrizes the Legendrian L and $a \in C_c^\infty(\tilde{M}_{b,0} \times \mathbb{R}^k; {}^{b,sc}\Omega^{1/2})$.

Remark 16. Despite $\dim M_{b,0} = 2n+1$, the contact structures are independent of the extra variable x such that the powers in the oscillatory integral don't take the extra dimension into account, hence the exponent $n/2$ as on $2n$ -manifolds.⁸ But the amplitude a does depend on x .

The Legendrian distributions associated with the Legendrian L near lf are defined on the fibred sc -bundle ${}^{s\Phi}T^* \tilde{M}_{b,0}$ as follows.

⁸In the abstract setting, we live in n -manifolds, hence the exponent $n/4$.

Definition 17. A Legendrian distribution $u \in I_{s\Phi}^{m; (r_{\text{lf}}, \infty, r_{\text{bf}_0})}(M_{b,0}; L; {}^{s\Phi}\Omega^{1/2})$ associated to a Legendrian L of ${}^{s\Phi}T^*M_{b,0}$ is a distribution in $C^{-\infty}(M_{b,0}; {}^{s\Phi}\Omega^{1/2})$ taking the form

$$u = x^{r_{\text{bf}_0}} u_0 + x^{r_{\text{bf}_0}} u_1 + \sum_i x^{r_{\text{bf}_0}} \omega_i \cdot \nu + \sum_i x^{r_{\text{bf}_0}} \omega'_i \cdot \nu,$$

away from rf , where

- $u_0 \in C^\infty(M_{b,0}; {}^{b,sc}\Omega^{1/2})$ is a ${}^{b,sc}\Omega^{1/2}$ -valued smooth function,
- $u_1 \in I_{sc,c}^m(\tilde{M}_{b,0}, L; {}^{b,sc}\Omega^{1/2})$ is a Legendrian distribution associated with L and supported in $\tilde{M}_{b,0}$,
- $\nu \in C^\infty(M_{b,0}; {}^{s\Phi}\Omega^{1/2})$ is a smooth section of the $s\Phi$ -half density bundle,
- ω_i is an oscillatory integral supported near the corner $\text{lf} \cap \text{mf}$ and takes the form

$$\omega_i(\rho', \tilde{\rho}, y, y', x) = \rho^{m-(k+k')/2+n/2} \tilde{\rho}^{r_{\text{lf}}-k/2} \int_{\mathbb{R}^{k+k'}} e^{i\phi_i(\tilde{\rho}, y, y', v, w)/\rho} a_i(\rho', \tilde{\rho}, y, y', x, v, w) dv dw,$$

where the amplitude $a_i \in C_c^\infty([0, \epsilon) \times U \times \mathbb{R}^{k+k'})$ with open $U \subset \text{lf}$ and the phase function ϕ_i parametrizes the Legendrian L on U ,

- ω'_i is an oscillatory integral supported in the interior of lf and takes the form

$$\omega'_i(\rho, y, z', x) = \rho^{r_{\text{lf}}-k/2} \int_{\mathbb{R}^k} e^{i\psi_i(y, w)/\rho} a_i(\rho, y, z', x, w) dw,$$

where the amplitude $a_i \in C_c^\infty([0, \epsilon) \times U \times \mathbb{R}^k)$ with open $U \subset \text{lf}$ and the phase function $\psi_i = \phi_i|_{\tilde{\rho}=0}$ parametrizes the Legendrian L_1 on $\phi_{\text{lf}}(U)$.

The space $I_{s\Phi}^{m; (\infty, r_{\text{rf}}, r_{\text{bf}_0})}(M_{b,0}; L; {}^{s\Phi}\Omega^{1/2})$ away from lf is defined in the same manner.

A Legendrian pair with conic points, (L, L^\sharp) , in ${}^{s\Phi}T^*M_{b,0}$ consists of two Legendrians L and L^\sharp of ${}^{s\Phi}T^*M_{b,0}$ which form an intersecting pair with conic points in ${}^{b,sc}T_{\tilde{M}_{b,0}}^* \tilde{M}_{b,0}$ such that the fibrations $\check{\phi}_{\text{lf}}$ and $\check{\phi}_{\text{lf}}^\sharp$ of L and L^\sharp have the same Legendrian L_1 of ${}^{s\Phi}N^*Z_{\text{lf}}$ as base and the fibres of $\check{\phi}_{\text{lf}}$ and $\check{\phi}_{\text{lf}}^\sharp$ are intersecting pairs of Legendrians with conic points of ${}^{b,sc}T_{\partial F}^*F$.

One can introduce a parametrization for (L, L^\sharp) . First of all, we note that L^\sharp is parametrized by a function on mf of the form

$$\phi_0/\rho = (\phi_1(y) + \tilde{\rho}\phi_2(\tilde{\rho}, y, y'))/\rho$$

near the boundary with lf , where $\phi_1 = 1$ on L^\sharp near $L \cap L^\sharp$. Then L_1^\sharp is parametrized by $\phi_1(y)/\rho$. The fibres of $\check{\phi}_{\text{lf}}^\sharp$ are pairs of Legendrians with conic points $(L(y), L^\sharp(y))$. For fixed y , $L^\sharp(y)$ is parametrized by $\phi_2(0, y, y')/x'$. Second, we perform the blow-up

$$[{}^{s\Phi}T_{\text{mf}}^*(M_{b,0}); \{\mu' = 0, \tau' = 0, \mu = 0\}],$$

to desingularize the conic singularity of (L, L^\sharp) . Then we can introduce a non-degenerate parametrization ϕ/ρ of (L, L^\sharp) , near a point $q \in \partial\hat{L} \cap {}^{s\Phi}T_{\text{mf} \cap \text{lf}}^*M_{b,0}$, where

$$\phi(\tilde{\rho}, y, y', s, w) = \phi_1(y) + \tilde{\rho}\phi_2(\tilde{\rho}, y, y') + s\tilde{\rho}\psi(\tilde{\rho}, y, y', s, w)$$

is a smooth function defined on a neighbourhood of $q' = (0, y_0, y'_0, 0, w_0)$ with

$$q = (0, y_0, y'_0, x, -\phi_1, -\frac{\psi}{|d_{y'}\psi|}, \frac{d_y\psi}{|d_{y'}\psi|}, 0, \widehat{d_{y'}\psi}(q'), 0), \quad d_s\phi(q') = 0, \quad d_w\psi(q') = 0,$$

and the lift \hat{L} of L under the blow-up is given by

$$\hat{L} = \{(\tilde{\rho}, y, y', x, -\phi_1, -\frac{\psi}{|d_{y'}\psi|}, \frac{d_y\psi}{|d_{y'}\psi|}, s|d_{y'}\psi|, \widehat{d_{y'}\psi}, 0) : d_s\phi = 0, d_w\psi = 0, s \geq 0\}.$$

Note that the phase function for (L, L^\sharp) is again independent of x and ξ due to the contact forms being independent of these variables. Therefore, (L, L^\sharp) extends in x smoothly to $\text{bf}_0 \cap (\text{mf} \cup \text{lf})$.

Away from $\text{lf} \cup \text{rf}$, there are no conic points. The intersecting Legendrian distributions are defined on ${}^{b,sc}T^*\tilde{M}_{b,0}$ as in the abstract settings. Specifically,

Definition 18. *The space $I_{sc,c}^{m,p_\sharp}(\tilde{M}_{b,0}; \tilde{L}; {}^{b,sc}\Omega^{1/2})$ of intersecting Legendrian distributions, associated with $\tilde{L} = (L, L^\sharp)$ in ${}^{b,sc}T_{\text{mf}}^*\tilde{M}_{b,0}$, consists of distributions of the form*

$$u = u_0 + \sum_{i=1}^N u_i \nu_i,$$

where

- $\nu_i \in C^\infty(\tilde{M}_{b,0}; {}^{b,sc}\Omega^{1/2})$ is a ${}^{b,sc}\Omega^{1/2}$ -valued smooth function on $\tilde{M}_{b,0}$,
- u_0 is a Legendrian distribution lying in

$$I_{sc,c}^m(\tilde{M}_{b,0}; L; {}^{b,sc}\Omega^{1/2}) + I_{sc,c}^{p_\sharp}(\tilde{M}_{b,0}; L^\sharp; {}^{b,sc}\Omega^{1/2}),$$

- u_i has a local expression

$$\int_0^\infty ds \int_{\mathbb{R}^k} e^{i\phi_i(\tilde{\rho}, y, y', v, s)/\rho'} a_i(\tilde{\rho}, y, y', x, v, \rho'/s, s) \left(\frac{\rho'}{s}\right)^{m+n/2-(k+1)/2} s^{p_\sharp+n/2-1} dv,$$

where $a_i \in C_c^\infty(\tilde{M}_{b,0} \times \mathbb{R}^k \times [0, \infty) \times [0, \infty))$, ϕ_i is non-degenerate and locally parametrizes (L, L^\sharp) .

The intersecting Legendrian distributions with conic points are defined on ${}^{s\Phi}T^*M_{b,0}$ as follows.

Definition 19. *The space $I_{s\Phi}^{m,p_\sharp;(r_{\text{lf}}, \infty, r_{\text{bf}_0})}(M_{b,0}, \tilde{L}; {}^{s\Phi}\Omega^{1/2})$ associated to a Legendrian pair with conic points $\tilde{L} = (L, L^\sharp)$ of ${}^{s\Phi}T^*M_{b,0}$ consists of distributions $u \in C^{-\infty}(M_{b,0}; {}^{s\Phi}\Omega^{1/2})$ that takes the form*

$$u = u_0 + u_1 + u_2 + \sum_j w_j \cdot \nu,$$


away from rf , where

- $u_0 \in I_{s\Phi}^{p_\sharp;(r_{\text{lf}}, \infty, r_{\text{bf}_0})}(M_{b,0}, L^\sharp; {}^{s\Phi}\Omega^{1/2})$ is a Legendrian distribution associated with L^\sharp ,
- $u_1 \in I_{s\Phi}^{m;(r_{\text{lf}}, \infty, r_{\text{bf}_0})}(M_{b,0}, L; {}^{s\Phi}\Omega^{1/2})$ is a Legendrian distribution associated with L ,
- $u_2 \in x^{r_{\text{bf}_0}} I_{sc,c}^{m,p_\sharp}(\tilde{M}_{b,0}, \tilde{L}; {}^{b,sc}\Omega^{1/2})$ is an intersecting Legendrian distribution associated with \tilde{L} and compactly supported in $\tilde{M}_{b,0}$,
- $\nu \in C^\infty(M_{b,0}; {}^{s\Phi}\Omega^{1/2})$ is a smooth section of the $s\Phi$ -half density bundle,
- ω_j is an oscillatory integral of the form,

$$\omega_j(\rho', \tilde{\rho}, y, y', x) = \int_{[0, \infty) \times \mathbb{R}^{k'}} e^{i\phi_j(\tilde{\rho}, y, y', s, w)/\rho} a_j(\rho', \tilde{\rho}, y, y', x, \rho'/s, s, w) \left(\frac{\rho'}{s}\right)^{m-(k'+1)/2+n/2} s^{p_\sharp-1+n/2} \tilde{\rho}^{r_{\text{lf}}} x^{r_{\text{bf}_0}} ds dw,$$

where the amplitude $a_j \in C_c^\infty([0, \epsilon) \times U \times [0, \infty) \times [0, \infty) \times \mathbb{R}^{k'})$ with U open in mf and the phase function ϕ_j parametrizes the Legendrian pair (L, L^\sharp) on U .

The space $I_{s\Phi}^{m,p_\sharp;(\infty, r_{\text{rf}}, r_{\text{bf}_0})}(M_{b,0}, \tilde{L}; {}^{s\Phi}\Omega^{1/2})$ away from lf is defined in the same manner.

FIGURE 7. The transitional face bf_0 and the stretched double metric cones $\mathfrak{C}_{sc,b}^2$

In the end, we define the 'global' Legendrian distributions, to incorporate the boundary behaviours with respect to both lf and rf , by

$$\begin{aligned}
 I_{s\Phi}^{m;(r_{\text{lf}}, r_{\text{rf}}, r_{\text{bf}_0})}(M_{b,0}; L; s\Phi\Omega^{1/2}) \\
 &= I_{s\Phi}^{m;(r_{\text{lf}}, \infty, r_{\text{bf}_0})}(M_{b,0}; L; s\Phi\Omega^{1/2}) + I_{s\Phi}^{m;(\infty, r_{\text{rf}}, r_{\text{bf}_0})}(M_{b,0}; L; s\Phi\Omega^{1/2}) \\
 I_{s\Phi}^{m,p_{\sharp};(r_{\text{lf}}, r_{\text{rf}}, r_{\text{bf}_0})}(M_{b,0}; \tilde{L}; s\Phi\Omega^{1/2}) \\
 &= I_{s\Phi}^{m,p_{\sharp};(r_{\text{lf}}, \infty, r_{\text{bf}_0})}(M_{b,0}; \tilde{L}; s\Phi\Omega^{1/2}) + I_{s\Phi}^{m,p_{\sharp};(\infty, r_{\text{rf}}, r_{\text{bf}_0})}(M_{b,0}; \tilde{L}; s\Phi\Omega^{1/2}).
 \end{aligned}$$

The space of Legendrian distributions on $M_{b,0}$ admits a short exact sequence

$$\begin{aligned}
 (22) \quad 0 \longrightarrow I_{s\Phi}^{m+1,p_{\sharp};(r_{\text{lf}}, r_{\text{rf}}, r_{\text{bf}_0})}(M_{b,0}; \tilde{L}; s\Phi\Omega^{1/2}) &\longrightarrow I_{s\Phi}^{m,p_{\sharp};(r_{\text{lf}}, r_{\text{rf}}, r_{\text{bf}_0})}(M_{b,0}; \tilde{L}; s\Phi\Omega^{1/2}) \\
 &\longrightarrow \rho_{\text{lf}}^{r_{\text{lf}}-m} \rho_{\text{rf}}^{r_{\text{rf}}-m} s^{p_{\sharp}-m} \rho_{\text{bf}_0}^{r_{\text{bf}_0}} C^\infty(\hat{L}, \Omega_b^{1/2}(\hat{L}) \otimes S^{[m]}(\hat{L})) \longrightarrow 0.
 \end{aligned}$$

7. THE NORMAL OPERATOR AT THE TRANSITIONAL FACE

If we pull back the differential operators \tilde{P}_h , P_b and P_{sc} on \mathcal{C} in (4), (5) and (6) to the stretched double spaces $M_{b,0}$ and $M_{sc,b,0}$ through the canonical left (right) projection, the normal operator $\mathcal{N}_{\text{bf}_0}(\tilde{P}_h)$ of \tilde{P}_h at bf_0 turns out to be $\Delta_{\mathfrak{C}} - (1 + \imath 0)$ where $\Delta_{\mathfrak{C}}$ is the Laplacian on the metric cone $(\mathfrak{C}, \mathfrak{g})$.

In fact, $\Delta_{\mathfrak{C}}$ on $(\mathfrak{C}, \mathfrak{g})$ has the global expression on \mathfrak{C} ,

$$-\partial_{\mathfrak{r}}^2 - \frac{(n-1)}{\mathfrak{r}} \partial_{\mathfrak{r}} + \frac{\Delta_{\mathfrak{h}}}{\mathfrak{r}^2}, \quad \text{for } 0 < \mathfrak{r} < \infty.$$

By making $\mathfrak{h} = h_0 = h(0, y, dy)$, $r = \mathfrak{r}$ near the small end $\{\mathfrak{r} = 0\}$ and $\rho = 1/\mathfrak{r}$ near the big end $\{\mathfrak{r} = \infty\}$, we find

$$(23) \quad \begin{aligned} \Delta_{\mathfrak{C}} - (1 + \imath 0) &= r^{-1} \mathcal{N}_{\text{bf}_0}(P_b) r^{-1} && \text{near } \{r = 0\} \\ \Delta_{\mathfrak{C}} - (1 + \imath 0) &= \mathcal{N}_{\text{bf}_0}(P_{sc}) && \text{near } \{\rho = 0\} \end{aligned} .$$

Therefore, we want to understand the resolvent kernel on the metric cone $(\mathfrak{C}, \mathfrak{g})$ to see the transitional behaviours of the resolvent kernel R at bf_0 between the regimes of $M_{b,0}$ and $M_{sc,b,0}$.

First of all, we need to work out the appropriate double space to accommodate the resolvent kernel of $\Delta_{\mathfrak{C}} - (1 + \iota 0)$. The equations (23) suggest $\Delta_{\mathfrak{C}}$ is a b -differential operator near the small end $\{r = 0\}$ and a sc -differential operator near the big end $\{r = \infty\}$. Hence the resolvent kernel lives in a stretched version of \mathfrak{C}^2 to accommodate these features. For this, we simultaneously blow up the two separate corners $C_{\mathfrak{C}^2,0} = \{r = 0, r' = 0\}$ and $C_{\mathfrak{C}^2,\infty} = \{\rho = \rho' = 0\}$,

$$\mathfrak{C}_b^2 = [\mathfrak{C}^2; C_{\mathfrak{C}^2,\infty}, C_{\mathfrak{C}^2,0}].$$

Subsequently, we perform the blow-up at the boundary at ∞ of the lifted diagonal $\text{diag}_{\mathfrak{C}_b^2}$ of \mathfrak{C}_b^2 ,

$$\mathfrak{C}_{sc,b}^2 = [\mathfrak{C}_b^2; \partial_{\infty} \text{diag}_{\mathfrak{C}_b^2}].$$

Figure 7 depicts the space $\mathfrak{C}_{sc,b}^2$. The dashed line is the lifted diagonal $\text{diag}_{\mathfrak{C}_{sc,b}^2}$ of $\mathfrak{C}_{sc,b}^2$. The boundary at ∞ of the diagonal $\text{diag}_{\mathfrak{C}^2}$ lifts to the face sc .

We assert that the space $\mathfrak{C}_{sc,b}^2$ (\mathfrak{C}_b^2) is isometric to the transitional face bf_0 in $M_{sc,b,0}$ ($M_{b,0}$). On the one hand, we may view $\mathfrak{C}_{sc,b}^2$ (\mathfrak{C}_b^2) as the stretched version of the double space of the front face of the stretched single space \mathcal{C}_b . Recall that $\mathcal{C}_b = [\mathcal{C} \times [0, 1); \partial\mathcal{C} \times \{0\}]$. The lift of $\partial\mathcal{C} \times \{0\}$, i.e. lb_0 or rb_0 , is isometric to \mathfrak{C} . This face can be parametrized by

$$\overline{\{(r, y) : 0 < r = x/\hbar \leq \infty\}}$$

and the constraint $x = 0$ reduces the metric $g = dx^2 + x^2 h(x, y, dy)$ multiplied by the rescaling factor $1/\hbar^2$ to a product type metric $\mathfrak{g} = dr^2 + r^2 \mathfrak{h}(y, dy)$. It follows $\text{lb}_0 \times \text{rb}_0 \cong \mathfrak{C}^2$. On the other hand, the face bf_0 in $M_{sc,b,0}$ ($M_{b,0}$) can be obtained by blowing up $\text{lb}_0 \times \text{rb}_0$, where the blow-down map is just given by the left and right stretched projections $\tilde{\mathcal{P}}_L$ and $\tilde{\mathcal{P}}_R$ in Section 2.

The three faces created by the two fold blow-ups are viewed as the three boundaries of bf_0 with bf , mf , sc respectively, whilst the four side edges of the square \mathfrak{C}^2 are linked to the four boundaries of bf_0 with lb , rb , lb_0 , rb_0 respectively. The dashed curves separate the b -regime, the sc -regime, the (b, sc) -regime and the (sc, b) -regime. We name these boundaries of $\mathfrak{C}_{sc,b}^2$ (\mathfrak{C}_b^2)

$$\{\text{lb}, \text{rb}, \text{lb}_0, \text{rb}_0, \text{bf}, \text{mf}, \text{sc}\},$$

which are isometric to the corners of $M_{sc,b,0}$ ($M_{b,0}$),

$$\{\text{lb} \cap \text{bf}_0, \text{rb} \cap \text{bf}_0, \text{lb}_0 \cap \text{bf}_0, \text{rb}_0 \cap \text{bf}_0, \text{bf} \cap \text{bf}_0, \text{mf} \cap \text{bf}_0, \text{sc} \cap \text{bf}_0\}.$$

In addition, we denote, only in the sc -regime,

$$\begin{aligned} \text{lf} &= \text{rb}_0 \cap \text{sc-regime} = \overline{\{\rho/\rho' = 0, 1 > \rho' > 0\}} \\ \text{rf} &= \text{lb}_0 \cap \text{sc-regime} = \overline{\{\rho'/\rho = 0, 1 > \rho > 0\}} \end{aligned}$$

to be consistent with the stretched double spaces $M_{sc,b,0}$ and $M_{b,0}$.

Through this isomorphism, the compressed cotangent bundles ${}^{b,sc}T^*\mathfrak{C}_b^2$ and ${}^{b,sc}T^*\mathfrak{C}_{sc,b}^2$ are induced from ${}^{b,sc}T^*(\text{bf}_0)$. These bundles further generate the compressed densities ${}^{b,sc}\Omega(\mathfrak{C}_b^2)$ and ${}^{b,sc}\Omega(\mathfrak{C}_{sc,b}^2)$.

Moreover, there exist induced contact structures and Legendrian submanifolds \mathfrak{L}_{\pm} and \mathfrak{L}^{\sharp} over $\text{lb}_0 \cup \text{rb}_0 \cup \text{mf} \subset \mathfrak{C}_b^2$. This is inherited through the isometry from L_{\pm} and L^{\sharp} over $\text{lb}_0 \cup \text{rb}_0 \cup \text{mf} \subset M_{b,0}$, since the contact structure on $M_{b,0}$ is independent of the boundary

defining function of bf_0 and extends smoothly to bf_0 . Then Legendrian distributions are analogously defined over $\text{lb}_0 \cup \text{rb}_0 \cup \text{mf} \subset \mathfrak{C}_b^2$ as we present in Section 6.

Guillarmou-Hassell-Sikora [14, Section 5] proved that the resolvent kernel $(\Delta_{\mathfrak{C}} - (1 + \imath 0))^{-1}$ is the sum of full b -pseudodifferential operators, sc -pseudodifferential operators, and Legendrian distributions. Specifically, the resolvent, in terms of our notations, reads

Theorem 20. *Let χ_b and χ_{sc} be smooth bump functions on \mathfrak{C} in a neighbourhood of $\{r < 1\}$ and $\{\rho < 1\}$ respectively, and also a partition of identity $\chi_b + \chi_{sc} = 1$. The resolvent kernel $R_{\mathfrak{C}}$ of $(\Delta_{\mathfrak{C}} - (1 + \imath 0))^{-1}$ is a distribution on \mathfrak{C}_b^2 consisting of,*

- $R_{\mathfrak{C},b} = \chi_b R_{\mathfrak{C}} \chi_b$ is a distribution supported in the b -regime of \mathfrak{C}_b^2 such that
 - $R_{\mathfrak{C},b}$ is a $^{b,sc}\Omega^{1/2}(\mathfrak{C}_b^2)$ -valued full b -pseudodifferential operator conormal at order -2 to the diagonal $\text{diag}_{\mathfrak{C}_b^2}$ and polyhomogeneous conormal to lb and rb ,
 - $R_{\mathfrak{C},b}$ vanishes to order 2 at bf , $n/2$ at lb and rb ;
- $R_{\mathfrak{C},sc} = \chi_{sc} R_{\mathfrak{C}} \chi_{sc}$ is a distribution supported in the sc -regime of \mathfrak{C}_b^2 such that
 - $R_{\mathfrak{C},sc}$ is conormal to order -2 to the diagonal $\text{diag}_{\mathfrak{C}_{sc,b}^2}$ (with the diagonal at infinity blown up),
 - $R_{\mathfrak{C},sc}$, near the big end of $\text{diag}_{\mathfrak{C}_b^2}$ in \mathfrak{C}_b^2 , is an intersecting Legendrian distribution lying in

$$I^{-1/2}(\mathfrak{C}_b^2; (N^* \text{diag}_{\mathfrak{C}_b^2}, \mathfrak{L}^\circ); ^{b,sc}\Omega^{1/2}(\mathfrak{C}_b^2)),$$

where \cdot° denotes the interior of \cdot ,

- $R_{\mathfrak{C},sc}$, near the intersection of $\text{mf} \cap \text{lf}$ and $\text{mf} \cap \text{rf}$, is intersecting Legendrian distributions lying in

$$I^{-1/2,(n-2)/2;(n-1)/2,(n-1)/2}(\mathfrak{C}_b^2; (\mathfrak{L}, \mathfrak{L}^\#); ^{s\Phi}\Omega^{1/2}),$$

- $R_{\mathfrak{C},sc}$ vanishes to order $(n-1)/2$ at both lf and rf .
- $R_{\mathfrak{C},(b,sc)} = \chi_b R_{\mathfrak{C}} \chi_{sc}$ is a distribution supported in the (b, sc) -regime of $\mathfrak{C}_{sc,b}^2$ such that
 - $R_{\mathfrak{C},(b,sc)}$ is given explicitly by

$$(24) \quad \frac{\pi \imath r}{2\rho} \sum_{j=0}^{\infty} \Pi_{E_j}(y, y') J_{\nu_j}(r) \text{Ha}_{\nu_j}^{(1)}(1/\rho') \left| \frac{dr dy d\rho' dy'}{r \rho'} \right|^{1/2},$$

where Π_{E_j} is the projection on the j th eigenspace E_j of the operator $\Delta_b + (n/2 - 1)^2$, ν_j^2 is the corresponding eigenvalue, and $J_\nu(\cdot)$, $\text{Ha}_\nu^{(1)}(\cdot)$ are standard Bessel and Hankel functions.

- $R_{\mathfrak{C},(b,sc)}$ vanishes to order $n/2$ at lb , $(n-1)/2$ at lb_0 ;
- $R_{\mathfrak{C},(sc,b)} = \chi_{sc} R_{\mathfrak{C}} \chi_b$ is a distribution supported in the (sc, b) -regime of $\mathfrak{C}_{sc,b}^2$ such that
 - $R_{\mathfrak{C},(sc,b)}$ is given explicitly by

$$(25) \quad \frac{\pi \imath r'}{2\rho} \sum_{j=0}^{\infty} \Pi_{E_j}(y, y') J_{\nu_j}(r') \text{Ha}_{\nu_j}^{(1)}(1/\rho) \left| \frac{dp dy dr' dy'}{\rho r'} \right|^{1/2},$$

- $R_{\mathfrak{C},(sc,b)}$ vanishes to order $n/2$ at rb , $(n-1)/2$ at rb_0 .

Consequently, the resolvent kernel of $\mathcal{N}_{\text{bf}_0}(\tilde{P}_h)$ is described as above, if we replace $R_{\mathfrak{C}}$ by $\mathcal{N}_{\text{bf}_0}(\tilde{P}_h)$, $\mathfrak{C}_{sc,b}^2$ by $\text{bf}_0 \subset M_{sc,b,0}$, and \mathfrak{C}_b^2 by $\text{bf}_0 \subset M_{b,0}$.

8. THE b -REGIME PARAMETRIX

In this section, we shall construct a parametrix G_b for P_b with a compact remainder E_b in the b -regime of $M_{b,0}$, where the relevant boundary faces include lb, rb, bf, bf₀. More precisely,

Proposition 21. *There exists a full b -pseudodifferential operator*

$$G_b \in \Psi_b^{-2,(\hat{\mathcal{E}}_b+1,\check{\mathcal{E}}_b+1,\tilde{\mathcal{E}}_b+2,\mathbb{N}_0)}(M_{b,0}; {}^b\Omega^{1/2}),$$

in the b -regime such that

$$r^{-1}P_b r^{-1} \circ G_b - \chi_b \text{Id} \chi_b = -E_b \in \Psi_b^{-\infty,(\infty,\hat{\mathcal{E}}_b+1,\infty,\infty)}(M_{b,0}; {}^b\Omega^{1/2}),$$

where the index sets are defined to be

$$(26) \quad \hat{\mathcal{E}}_b = \overline{\bigcup_{i \in \mathbb{N}_0} (\mathcal{E}_b + i)}$$

$$(27) \quad \check{\mathcal{E}}_b = \hat{\mathcal{E}}_b \sqcup \hat{\mathcal{E}}_b,$$

$$(28) \quad \tilde{\mathcal{E}}_b = \{(0,0)\} \cup \left((\mathbb{N}_0 + 1) \sqcup (\hat{\mathcal{E}}_b + \check{\mathcal{E}}_b) \right),$$

provided the index set $\mathcal{E}_b = \{(\text{Im } \lambda_j, 0)\}$ as in (16).

Proof. The operator P_b is lifted through the stretched left projection $\tilde{\mathcal{P}}_L$ to the b -regime in the stretched double $M_{b,0}$. The construction away from bf₀ is a direct application of the abstract full b -calculus. We follow the b -parametrix construction in [28, Chapter 4-6]. See also Albin's lecture notes [2, Section 4-5] for more details on the index set of the face bf. Moreover, we need additional correction terms near bf₀.

8.1. Construction near the diagonal. First of all, we invoke the small b -calculus to remove the diagonal singularity. Since $P_b \in \text{Diff}_b^2(M_{b,0}; {}^{b,sc}\Omega^{1/2})$ is an elliptic b -differential operator in the b -regime, the routine elliptic construction shows that there exists a small b -pseudodifferential operator $G_{b,1} \in \Psi_b^{-2}(M_{b,0}; {}^{b,sc}\Omega^{1/2})$ in the b -regime such that

$$P_b \circ G_{b,1} - \chi_b \text{Id} \chi_b = -E_{b,1} \in \Psi_b^{-\infty}(M_{b,0}; {}^{b,sc}\Omega^{1/2}).$$

8.2. Error at lb. We next wish to find a $G_{b,2} \in \Psi_b^{-\infty,(\hat{\mathcal{E}}_b,\infty,\mathbb{N}_0,\mathbb{N}_0)}(M_{b,0}; {}^{b,sc}\Omega^{1/2})$ such that

$$P_b \circ G_{b,2} - E_{b,1} = -E_{b,2} \in \Psi_b^{-\infty,(\infty,\hat{\mathcal{E}}_b,\mathbb{N}_0+1,\mathbb{N}_0)}(M_{b,0}; {}^{b,sc}\Omega^{1/2}).$$

Through this step, we reduce the remainder to a distribution vanishing at bf and to infinite order at lb.

Having the remainder vanishing at bf amounts to solving the indicial equation,

$$\mathcal{I}(E_{b,1}) = \mathcal{I}(P_b) \circ \mathcal{I}(G_{b,2}).$$

We solve this equation through the inverse Mellin transform of the inverse of the indicial family $\mathcal{I}(P_b, \lambda)$, which is known to be a polyhomogeneous conormal distribution in Proposition 7. Then part of G_2 is taken to be

$$G_{b,2}^{(1)} = -\frac{1}{2\pi} \int_{-\infty}^{\infty} s^{i\lambda} \mathcal{I}^{-1}(P_b, \lambda) \circ \mathcal{M}(E_{b,1})(\lambda) d\lambda,$$

which satisfies

$$P_b \circ G_{b,2}^{(1)} - E_{b,1} = -E_{b,2}^{(1)} \in \Psi_b^{-\infty,(\mathcal{E}_b,\mathcal{E}_b,\mathbb{N}_0+1,\mathbb{N}_0)}(M_{b,0}; {}^{b,sc}\Omega^{1/2}).$$

In addition, the error at lb can be removed completely by [28, Lemma 5.44]. In fact, this lemma implies that there exists $G_{b,2}^{(2)} \in \mathcal{A}_{phg}^{(\hat{\mathcal{E}}_b, \infty, \mathbb{N}_0+1, \mathbb{N}_0)}(M_{b,0}; {}^{b,sc}\Omega^{1/2})$ such that

$$P \circ G_{b,2}^{(2)} - E_{b,2}^{(1)} = -E_{b,2} \in \mathcal{A}_{phg}^{(\infty, \mathcal{E}_b, \mathbb{N}_0+1, \mathbb{N}_0)}(M_{b,0}; {}^{b,sc}\Omega^{1/2}).$$

If we write $G_{b,2} = G_{b,2}^{(1)} + G_{b,2}^{(2)}$, the parametrix $G_{b,1} + G_{b,2}$ obeys that

$$P_b \circ (G_{b,1} + G_{b,2}) - \chi_b \text{Id} \chi_b = -E_{b,2} \in \Psi_b^{-\infty, (\infty, \mathcal{E}_b, \mathbb{N}_0+1, \mathbb{N}_0)}(M_{b,0}; {}^{b,sc}\Omega^{1/2}).$$

8.3. Error at bf. We can iterate this construction to remove the error at bf. By the calculus of full b -pseudodifferential operators, the j th-power of the remainder is

$$E_{b,2}^j \in \Psi_b^{-\infty, (\infty, \mathcal{E}_{b,j}, \mathbb{N}_0+j, \mathbb{N}_0)}(M_{b,0}; {}^{b,sc}\Omega^{1/2}),$$

where the index sets obey

$$\begin{aligned} \mathcal{E}_{b,1} &= \mathcal{E}_b, \\ \mathcal{E}_{b,j+1} - 1 &= \mathcal{E}_{b,j} \cup (\mathcal{E}_b - 1), \\ \lim_{j \rightarrow \infty} \mathcal{E}_{b,j} &= \hat{\mathcal{E}}_b. \end{aligned}$$

If it is sufficiently close to bf, say $\chi_b = 1$, we have

$$\chi_b (\text{Id} - E_{b,2}) (\text{Id} + \sum_{j=1}^{\infty} E_{b,2}^j) \chi_b - \chi_b \text{Id} \chi_b \in \bigcap_{j>0} \Psi_b^{-\infty, (\infty, \hat{\mathcal{E}}_b, \mathbb{N}_0+j, \mathbb{N}_0)}(M_{b,0}; {}^{b,sc}\Omega^{1/2}).$$

By the composition of index sets given Proposition 6, the improved parametrix

$$G_{b,1,2} = (G_{b,1} + G_{b,2}) (\chi_b \text{Id} \chi_b + \sum_{j=1}^{\infty} E_{b,2}^j)$$

lies in a slightly larger space $\Psi_b^{-2, (\hat{\mathcal{E}}_b, \hat{\mathcal{E}}_b, \hat{\mathcal{E}}_b, \mathbb{N}_0)}(M_{b,0}; {}^{b,sc}\Omega^{1/2})$, whilst the new remainder $E_{b,1,2} \in \Psi_b^{-\infty, (\infty, \hat{\mathcal{E}}_b, \infty, \mathbb{N}_0)}(M_{b,0}; {}^{b,sc}\Omega^{1/2})$ vanishes to infinite order at bf.

8.4. Error at bf_0 . We next construct a correction term $G_{b,3}$ supported around bf_0 in the b -regime and lying in

$$G_{b,3} \in \Psi_b^{-2, (\hat{\mathcal{E}}_b, \hat{\mathcal{E}}_b, \hat{\mathcal{E}}_b, \mathbb{N}_0)}(M_{b,0}; {}^{b,sc}\Omega),$$

which leads to an improved remainder vanishing at bf_0 ,

$$P_b \circ G_{b,3} - E_{b,1,2} = -E_{b,3} \in \Psi_b^{-\infty, (\infty, \hat{\mathcal{E}}_b, \infty, \infty)}(M_{b,0}; {}^b\Omega^{1/2}).$$

This again amounts to solving the normal operator equation

$$\mathcal{N}_{\text{bf}_0}(P_b) \circ \mathcal{N}_{\text{bf}_0}(G_{b,3}) = \mathcal{N}_{\text{bf}_0}(E_{b,1,2}).$$

Due to

$$\mathcal{N}_{\text{bf}_0}(P_b) = \mathcal{N}_{\text{bf}_0}(\chi_b r \tilde{P}_{\hbar} r \chi_b) = \chi_b r (\Delta_{\mathfrak{C}} - 1 \pm i0) r \chi_b,$$

this can be done by using Theorem 20.

If we take

$$G_{b,3}^{(1)} = (r^{-1} R_{\mathfrak{C}, b} r'^{-1}) \mathcal{N}_{\text{bf}_0}(E_{b,1,2}),$$

it results in a better remainder $E_{b,3}^{(1)}$, which vanishes at bf_0 as well as to infinite order at lb and bf and lies in

$$E_{b,3}^{(1)} \in \Psi_b^{-\infty, (\infty, \hat{\mathcal{E}}_b, \infty, \mathbb{N}_0+1)}(M_{b,0}; {}^{b,sc}\Omega^{1/2}).$$

The calculus of full b -pseudodifferential operators, Proposition 6, yields that $\mathcal{N}_{\text{bf}_0}(G_{b,3}^{(1)})$ is polyhomogeneous conormal to lb, rb, bf with respect to index sets $(\hat{\mathcal{E}}_b, \check{\mathcal{E}}_b, \tilde{\mathcal{E}}_b)$ as desired. Repeating this procedure j times yields parametrices

$$G_{b,3}^{(j)} \in \Psi_b^{-2,(\hat{\mathcal{E}}_b, \check{\mathcal{E}}_b, \tilde{\mathcal{E}}_b, \mathbb{N}_0+j-1)}(M_{b,0}; {}^{b,sc}\Omega^{1/2})$$

and remainders

$$E_{b,3}^{(j)} \in \Psi_b^{-\infty,(\infty, \hat{\mathcal{E}}_b, \infty, \mathbb{N}_0+j)}(M_{b,0}; {}^{b,sc}\Omega^{1/2}).$$

Then the desired parametrix $G_{b,3}$ and remainder $E_{b,3}$ are constructed by

$$G_{b,3} = \sum_{j=1}^{\infty} G_{b,3}^{(j)} \quad \text{and} \quad E_{b,3} = \lim_{j \rightarrow \infty} E_{b,3}^{(j)}.$$

In the end, we obtain a parametrix G_b with a remainder E_b such that

$$\begin{aligned} r^{-1}G_b r'^{-1} &= G_{b,1,2} + G_{b,3}, & r^{-1}G_b r'^{-1} &\in \Psi_b^{-2,(\hat{\mathcal{E}}_b, \check{\mathcal{E}}_b, \tilde{\mathcal{E}}_b, \mathbb{N}_0)}(M_{b,0}; {}^{b,sc}\Omega^{1/2}), \\ r^{-1}E_b r'^{-1} &= E_{b,3}, & r^{-1}E_b r'^{-1} &\in \Psi_b^{-\infty,(\infty, \hat{\mathcal{E}}_b, \infty, \infty)}(M_{b,0}; {}^{b,sc}\Omega^{1/2}). \end{aligned}$$

Then G_b and E_b satisfy the conditions in Proposition 21, since rr' lifts to $\rho_{\text{bf}}^2 \rho_{\text{lb}} \rho_{\text{rb}}$ in the b -regime of $M_{b,0}$. \square

9. THE (b, sc) -REGIME PARAMETRIX

Recall that the (b, sc) -regime is a neighbourhood of $\{r < 1\} \cup \{\rho' < 1\}$ in $M_{b,0}$ and furnished with local coordinates (r, y, ρ', y', \hbar) . It has three boundary hypersurfaces lb, lb₀, bf₀ with boundary defining functions $r, \rho', \hbar/\rho'$ respectively. The compressed density ${}^{b,sc}\Omega(M_{b,0})$ locally takes the form

$$\left| \frac{dr dy d\rho dy' d\hbar}{r \rho'^{n+1} \hbar} \right|.$$

Since \tilde{P}_{\hbar} is viewed as a left operator on $M_{b,0}$ via the stretched left projection \tilde{P}_L , we can convert it, as in the b -regime, to a b -operator P_b in the (b, sc) -regime. The b -calculus in the b -regime extends trivially to the (b, sc) -regime, though only the polyhomogeneous conormal distributions in the full calculus are relevant. We shall construct the following parametrix for \tilde{P}_{\hbar} in this regime.

Proposition 22. *In the (b, sc) -regime of $M_{b,0}$, there is a parametrix $G_{(b,sc)}$ with a kernel*

$$G_{(b,sc)} \in \rho_{\text{lb}_0}^{(n-1)/2} \sum_{(z,k) \in \hat{\mathcal{E}}_b + 1} \rho_{\text{lb}}^z (\log \rho_{\text{lb}})^k C^{\infty}(M_{b,0}; {}^{b,sc}\Omega^{1/2}),$$

provided the index set $\hat{\mathcal{E}}_b$ in (26), such that

$$\tilde{P}_{\hbar} \circ G_{(b,sc)} - \chi_b \text{Id} \chi_{sc} = -E_{(b,sc)} \in \rho_{\text{lb}}^{\infty} \rho_{\text{lb}_0}^{\infty} \rho_{\text{bf}_0}^{\infty} C^{\infty}(M_{b,0}; {}^{b,sc}\Omega^{1/2}).$$

Proof. Constructing the parametrix in the (b, sc) -regime involves removing errors at lb, lb₀ and bf₀ and obtaining the polyhomogeneity or vanishing order at lb and lb₀.

9.1. Boundary error removal. The parametrices near lb and lb₀ are similar to the b -regime. Since $r < 1$ in this regime, $P_b = r\tilde{P}_h r$ is still a b -differential operator but supported away from the diagonal $\text{diag}_{M_{b,0}}$ and the face bf.

Since the operator P_b in the (b, sc) -regime takes the same form with the b -regime. The error at lb can be removed through the indicial family which is similar to Section 8.2 and Section 8.3. But the difference is that the indicial operator/family at bf is replaced by that at lb₀ ∪ bf₀ = $\{\bar{h} = 0\}$, since lb₀ plays the same role as bf in the b -regime. Specifically, we find a parametrix $G_{(b,sc),1}$ in the (b, sc) -regime which is polyhomogeneous conormal to lb with an index set $\hat{\mathcal{E}}_b$ such that

$$P_b \circ G_{(b,sc),1} - \chi_b \text{Id} \chi_{sc} = -E_{(b,sc),1} \in \rho_{\text{lb}}^\infty \hbar C^\infty(M_{b,0}; {}^{b,sc}\Omega).$$

Then the standard power series argument removes the error at lb₀ ∪ bf₀ completely. It yields that there a parametrix $G_{(b,sc),2}$ polyhomogeneous conormal to lb₀ ∪ bf₀ with respect to $\tilde{\mathcal{E}}_b$ and to lb with respect to $\hat{\mathcal{E}}_b$ such that

$$P_b \circ G_{(b,sc),2} - E_{(b,sc),1} = -E_{(b,sc),2} \in \rho_{\text{lb}}^\infty \hbar^\infty C^\infty(M_{b,0}; {}^{b,sc}\Omega).$$

From this, the corresponding parametrix for \tilde{P}_h is

$$G_{(b,sc)} = r(G_{(b,sc),1} + G_{(b,sc),2})\rho'^{-1}.$$

9.2. Vanishing at lb₀. The given polyhomogeneity of the parametrix $G_{b,sc}$ in \hbar is crude, since \hbar is the total defining function for lb₀ ∪ bf₀. The index set $\tilde{\mathcal{E}}_b$ in (28) for \hbar , by definition, does not address any vanishing, though $\inf \hat{\mathcal{E}}_b = n/2 - 1$ does give an $n/2$ -vanishing of $G_{b,sc}$ at lb. It remains to work out the exact vanishing order of $G_{(b,sc)}$ at lb₀. This is determined by the resolvent of the normal operator.⁹

Since $x = \hbar = 0$ at lb₀ ∪ bf₀, we see the normal operator of $P_{\tilde{h}}$ at lb₀ ∪ bf₀ is

$$\mathcal{N}_{\text{lb}_0 \cup \text{bf}_0}(\tilde{P}_{\tilde{h}}) = -\partial_r^2 - \frac{n-1}{r} \partial_r + \frac{\Delta_{h_0}}{r^2} - (1 + i0),$$

where $h_0 = h(0, y, dy)$ is a metric on \mathcal{Y} and thus the Laplacian Δ_{h_0} on (\mathcal{Y}, h_0) is independent of r . This is equal to $\Delta_{\mathfrak{C}} - (1 + i0)$ and then converted to a b -operator

$$P_{(b,sc)} = r\mathcal{N}_{\text{lb}_0 \cup \text{bf}_0}(\tilde{P}_{\tilde{h}})r = -(r\partial_r)^2 + \Delta_{h_0} + (n/2 - 1)^2 - r^2(1 \pm i0).$$

One can apply separation of variables to compute $(P_{(b,sc)})^{-1}$. This is done in [14, (5.5) (5.6)] for the resolvent of $\Delta_{\mathfrak{C}} - (1 + i0)$ given in Theorem 20. But the resolvent kernel of $P_{(b,sc)}$ lives in a $(2n + 1)$ -manifold. Using this result, we have

$$(\mathcal{N}_{\text{lb}_0 \cup \text{bf}_0}(\tilde{P}_{\tilde{h}}))^{-1} = \frac{\pi i r}{2\rho'} \sum_{j=0}^{\infty} \Pi_{E_j}(y, y') J_{\nu_j}(r) \text{Ha}_{\nu_j}^{(1)}(1/\rho') \left| \frac{dr dy d\rho' dy' d\hbar}{r \rho' \hbar} \right|^{1/2},$$

where Π_{E_j} is the projection on the j th eigenspace E_j of the operator $\Delta_{h_0} + (n/2 - 1)^2$, ν_j^2 is the corresponding eigenvalue, and $J_\nu(\cdot)$, $\text{Ha}_\nu^{(1)}(\cdot)$ are standard Bessel and Hankel functions as in (24) and (25).

Invoking the asymptotic behaviours of Bessel and Hankel functions,

$$(29) \quad J_\nu(r) \sim \frac{1}{\Gamma(\nu + 1)} \left(\frac{r}{2}\right)^\nu + O(r^{\nu+1}), \quad \text{for } r \rightarrow 0,$$

⁹If $\tilde{P}_{\tilde{h}}$ is written near lb₀ in the form of an asymptotic expansion in ρ_{lb_0} , the normal operator is just the leading terms of such expansion. On the other hand, the resolvent of $\tilde{P}_{\tilde{h}}$ also has an asymptotic expansion in ρ_{lb_0} , whilst the resolvent of the normal operator is just the leading terms of the resolvent of $\tilde{P}_{\tilde{h}}$.

$$(30) \quad \text{Ha}_\nu^{(1)}(r') \sim r'^{-1/2} e^{ir' - \nu\pi/2 + i\pi/4} \sum_{k=0}^{\infty} a_k(\nu) r'^{-k}, \quad \text{for } r' \rightarrow \infty,$$

where the coefficients are given by

$$a_0(\nu) = 1, \quad a_k(\nu) = \frac{\prod_{j=1}^k (4\nu^2 - (2j-1)^2)}{k! 8^k} \quad \text{for } k \in \mathbb{Z}_+,$$

we see that

$$(\mathcal{N}_{\text{lb}_0 \cup \text{bf}_0}(\tilde{P}_h))^{-1} \sim r^{n/2} \rho'^{(n-1)/2} e^{i/\rho'} \left| \frac{dr dy d\rho' dy' d\hbar}{r \rho'^{n+1} \hbar} \right|^{1/2}.$$

This implies the kernel of $G_{(b,sc)}$ vanishes to order $(n-1)/2$ at lb_0 and to order $n/2$ at lb . \square

10. THE sc -REGIME PARAMETRIX

Let us move to the sc -regime of $M_{b,0}$. While using the sc -pseudodifferential calculus, we need perform an additional blow-up and stay briefly in $M_{sc,b,0}$. We construct a parametrix G_{sc} for P_{sc} there. Specifically,

Proposition 23 (sc -parametrix). *There is a distribution G_{sc} in the sc -regime such that*

$$P_{sc} \circ G_{sc} - \chi_{sc} \text{Id} \chi_{sc} = -E_{sc} \in e^{i/\rho'} \rho_{\text{rf}}^{(n-1)/2 - \varepsilon_r/2} \rho_{\text{bf}_0}^\infty \rho_{\text{mf}}^\infty \rho_{\text{lf}}^\infty C^\infty(M_{b,0}; {}^{b,sc}\Omega^{1/2}).$$

The kernel of G_{sc} , modulo sc -pseudodifferential operators in $\Psi_{sc}^{-2}(M_{sc,b,0}; {}^{b,sc}\Omega^{1/2})$, are Legendrian distributions lying in the class

$$\begin{aligned} I_{sc,c}^{-1/2}(M_{b,0}; (N^* \text{diag}_{M_{b,0}}, L_\pm^\circ); {}^{b,sc}\Omega^{1/2}) \\ + I_{s\Phi}^{-1/2, (n-2)/2 - \mathbf{e}; ((n-1-\mathbf{e})/2, (n-1-\mathbf{e})/2, 0)}(M_{b,0}; (L_\pm, L^\sharp); {}^{s\Phi}\Omega^{1/2}). \end{aligned}$$

Here we employ the following notations

- the diagonals in the sc -regime of $M_{b,0}$ and $M_{sc,b,0}$ are locally expressed by

$$\text{diag}_{M_{b,0}} = \{(\tilde{\rho}, \rho', y, y') \in M_{b,0} : \tilde{\rho} = 1, y = y'\}$$

$$\text{diag}_{M_{sc,b,0}} = \{(\mathbf{s}, \rho', \mathbf{y}, y') \in M_{sc,b,0} : \mathbf{s} = 0, \mathbf{y} = 0\}$$

where $\tilde{\rho} = \rho/\rho' = x'/x$, $\mathbf{s} = (\sigma - 1)/\rho'$ and $\mathbf{y} = (y - y')/\rho'$ are the projective coordinates on mf and sc ,

- $L = L_\pm$ is the forward/backward geometric Legendrian, L^\sharp is a diffractive Legendrian, and L_\pm° is the interior of L_\pm , which are explicitly given in (32) and (33),
- $\{\rho_{\text{lf}}, \rho_{\text{rf}}, \rho_{\text{mf}}\}$ denote the boundary defining functions of $\{\text{lf}, \text{rf}, \text{mf}\}$ respectively,
- the error function $\varepsilon = xe(x, y)$, with $e(x, y)$ defined in (2) and $\mathbf{e} = \sup_{(x,y) \in \mathcal{K}} \varepsilon$, produces additional oscillations near $\text{lf}, \text{rf}, L^\sharp$ for the principal symbol of G_{sc} , hence the loss \mathbf{e} of the orders of G_{sc} ,
- \bullet_l and \bullet_r denote the quantity \bullet in the left and right variables respectively,
- ρ' on $M_{b,0}$ is understood as the total boundary defining function of $\text{mf} \cup \text{rf}$.

Proof. Due to the scattering nature of P_{sc} , the proof exploits the strategy of the resolvent construction for the Laplacian on asymptotically conic manifolds (near the infinity) by Hassell-Vasy [18, Section 4]. However, we must work out the explicit oscillations near $\text{lf}, \text{rf}, L^\sharp$, caused by e , which is the essential feature of non-product conic manifolds as mentioned above.

10.1. Construction near the diagonal. We seek $G_{sc,1} \in \Psi_{sc}^{-2}(M_{sc,b,0}; {}^{b,sc}\Omega^{1/2})$ such that

$$P_{sc} \circ G_{sc,1} = \chi_{sc} \text{Id} \chi_{sc} - E_{sc,1}, \quad E_{sc,1} \in \Psi_{sc}^{-\infty}(M_{sc,b,0}; {}^{b,sc}\Omega^{1/2}).$$

This means that the remainder term $E_{sc,1}$ is smooth on $M_{sc,b,0}$ so that we have solved away the singularity along the diagonal $\text{diag}_{M_{sc,b,0}}$.

Similar to the b -regime, the standard elliptic argument applies here since the interior symbol is non-vanishing. Thus, we first choose a $\tilde{Q}_1 \in \Psi_{sc}^{-2}(M_{sc,b,0}; {}^{b,sc}\Omega^{1/2})$ with symbol $(\sigma^2(P_{sc}))^{-1}$. Then

$$P_{sc} \circ \tilde{Q}_1 = \chi_{sc} \text{Id} \chi_{sc} - \tilde{E}_1, \quad \tilde{E}_1 \in \Psi_{sc}^{-1}(M_{sc,b,0}; {}^{b,sc}\Omega^{1/2}).$$

Multiplying \tilde{Q}_1 by a finite Neumann series $(\chi_{sc} \text{Id} \chi_{sc} + \tilde{E}_1 + \cdots + \tilde{E}_1^{k-1})$ yields a remainder in $\Psi_{sc}^{-k}(M_{sc,b,0}; {}^{b,sc}\Omega^{1/2})$. Taking an asymptotic limit gives us the desired $G_{sc,1}$ and $E_{sc,1}$.

10.2. Error near the boundary of the diagonal. The sc face will be irrelevant from this stage onwards, and we thus blow down sc and live in $M_{b,0}$ hereafter. We then adopt local coordinates $(\tilde{\rho}, \rho', y, y', x)$ in this region of $M_{b,0}$. The remainder $E_{sc,1}$ has a singularity at $\partial_{sc} \text{diag}_{M_{b,0}}$,

$$\partial_{sc} \text{diag}_{M_{b,0}} = \{\rho' = 0, \tilde{\rho} = 1, y = y'\},$$

and its kernel in a neighbourhood of $\partial_{sc} \text{diag}_{M_{b,0}}$ within mf is a smooth function of ρ , $(\tilde{\rho} - 1)/\rho$, $(y - y')/\rho$, and y' . Therefore it can be expressed, in terms of the Fourier transform, as

$$E_{sc,1} = \left(\int_{\mathbb{R}^n} e^{i((y-y') \cdot \eta + (\tilde{\rho}-1)t)/\rho} a_1(\rho', y, x, \eta, t) d\eta dt \right) \nu,$$

where $\nu \in C^\infty(M_{b,0}; {}^{b,sc}\Omega^{1/2})$ and a_1 is smooth in all variables and Schwartz in (η, t) . The phase function $(y - y') \cdot \eta + (\tilde{\rho} - 1)t$ parametrizes the Legendrian

$$N^* \text{diag}_{M_{b,0}} = \{y = y', \tilde{\rho} = 1, \mu = -\mu', \tau = -\tau'\}.$$

Therefore, $E_{sc,1}$ is a Legendrian distribution of order 0 associated to $N^* \text{diag}_{M_{b,0}}$. To solve away this error inductively, we further write

$$E_{sc,1} = \sum_{k=0}^{\infty} E_{sc,1}^{(k)}, \quad E_{sc,1}^{(k)} = \left(\int_{\mathbb{R}^n} e^{i((y-y') \cdot \eta + (\tilde{\rho}-1)t)/\rho} a_1^{(k)}(\rho', y, x, \eta, t) d\eta dt \right) \nu,$$

where $E_{sc,1}^{(k)}$ is a Legendrian distribution of order k , and $a_1^{(k)}$ is smooth in all variables and Schwartz in (η, t) .

Since $\sigma_{\partial}(P_{sc}) = \tau^2 + h^{ij} \mu_i \mu_j - 1$ vanishes on a codimension one submanifolds of $N^* \text{diag}_{M_{b,0}}$, we consider its (left) Hamilton vector field V_l ,

$$V_l = 2\tau \tilde{\rho} \frac{\partial}{\partial \tilde{\rho}} + 2\tau \mu_i \frac{\partial}{\partial \mu_i} - h^{ij} \mu_i \mu_j \frac{\partial}{\partial \tau} + \left(\frac{\partial(h^{ij} \mu_i \mu_j)}{\partial \mu_i} \frac{\partial}{\partial y_j} - \frac{\partial(h^{ij} \mu_i \mu_j)}{\partial y_i} \frac{\partial}{\partial \mu_j} \right).$$

The fact that $\tau^2 + h^{ij} \mu_i \mu_j = 1$ on the characteristic variety $\Sigma(P_{sc})$ implies that either $\tau \tilde{\rho} \partial_{\tilde{\rho}}$ or $-h^{ij} \mu_i \mu_j \partial_{\tau}$ in V_l is non-vanishing. Thus V_l is transverse to $N^* \text{diag}_{M_{b,0}}$ at the intersection with $\Sigma(P_{sc})$. We define L° to be the flowout Legendrian emanating from $N^* \text{diag}_{M_{b,0}} \cap \Sigma(P_{sc})$ with respect to V_l , and L_\pm° to be the flowout in the positive and negative directions with respect to V_l .

We wish to find an intersecting Legendrian distribution

$$Q_2^{(0)} \in I_{sc,c}^{-1/2}(\tilde{M}_{b,0}; (N^* \text{diag}_{M_{b,0}}, L_+^\circ); {}^{b,sc}\Omega^{1/2}),$$




FIGURE 8. The intersecting pair of Legendrians with conic points

such that

$$(31) \quad P_{sc} \circ Q_2^{(0)} - E_{sc,1}^{(0)} \in I_{sc,c}^1(\tilde{M}_{b,0}; N^* \text{diag}_{M_{b,0}}; {}^{b,sc}\Omega^{1/2}) + I_{sc,c}^{3/2}(\tilde{M}_{b,0}; (N^* \text{diag}_{M_{b,0}}, L_+^\circ); {}^{b,sc}\Omega^{1/2}),$$

microlocally near $N^* \text{diag}_{M_{b,0}}$. Using the ellipticity of P_{sc} away from $\Sigma(P_{sc})$, we choose $Q_2^{(0)}$ with a symbol on $N^* \text{diag}_{M_{b,0}} \setminus L_+^\circ$ equal to $\sigma_\partial^0(P_{sc})^{-1} \sigma^0(E_{sc,1}^{(0)})$. Then the value of the symbol on $L_+^\circ \cap N^* \text{diag}_{M_{b,0}}$ is given by (20). The symbol of $Q_2^{(0)}$ extends to L_+ through the symbol calculus (18), which gives the transport equation

$$\left(-\imath \mathcal{L}_{H_p} - \imath(1/2 + m - n/4) \partial_\tau p + p_{\text{sub}} \right) \sigma(Q_2^{(0)}) = 0, \quad \text{for } p = \sigma_\partial^0(P_{sc}).$$

This ODE has a unique solution in a sufficiently small neighbourhood of $N^* \text{diag}_{M_{b,0}}$. Namely, the remainder $E_{sc,1}^{(0)}$ on L_+° has been solved away, and the symbol of $F_1^{(1)} = P_{sc} \circ Q_2^{(0)} - E_{sc,1}^{(0)}$ on L_+° is of order $-3/2$. Then (31) follows from the exact sequence (21).

We can inductively solve away the remainder

$$E_{sc,1}^{(k)} + F_1^{(k)} \in I_{sc,c}^k(\tilde{M}_{b,0}; N^* \text{diag}_{M_{b,0}}; {}^{b,sc}\Omega^{1/2}) + I_{sc,c}^{k+1/2}(\tilde{M}_{b,0}; (N^* \text{diag}_{M_{b,0}}, L_+^\circ); {}^{b,sc}\Omega^{1/2})$$

with an intersecting Legendrian distribution

$$Q_2^{(k)} \in I_{sc,c}^{k+1/2}(\tilde{M}_{b,0}; (N^* \text{diag}_{M_{b,0}}, L_+^\circ); {}^{b,sc}\Omega^{1/2}),$$

up to a remainder

$$F_1^{(k+1)} \in I_{sc,c}^{k+1}(\tilde{M}_{b,0}; N^* \text{diag}_{M_{b,0}}; {}^{b,sc}\Omega^{1/2}) + I_{sc,c}^{k+3/2}(\tilde{M}_{b,0}; (N^* \text{diag}_{M_{b,0}}, L_+^\circ); {}^{b,sc}\Omega^{1/2}).$$

Taking the sum $G_{sc,2} = \sum_{k=0}^\infty Q_2^{(k)}$ yields a remainder, smooth near $N^* \text{diag}_{M_{b,0}}$. But away from $\partial \text{diag}_{M_{b,0}}$, we end up with a remainder,

$$E_{sc,2} = -P_{sc} \circ G_{sc,2} + E_{sc,1} \in I_{sc,c}^{1/2}(\tilde{M}_{b,0}; L_+^\circ \setminus N^* \text{diag}_{M_{b,0}}; {}^{b,sc}\Omega^{1/2}).$$

10.3. The bicharacteristic flow. Before solving away the error on L_+° , we describe the global structure of L_+° to understand the conic singularities popping up in the closure of L_+° .

Recall that L° was defined to be the flowout from $N^*\text{diag}_{M_{b,0}} \cap \Sigma(P_{sc})$ by the Hamilton vector field V_l in left variables (ρ, y) . This is the same with the flowout generated by the Hamilton vector field V_r in right variables (x', y') .

Melrose-Zworski's calculations (with Hassell-Vasy's modifications [18, (4.7) (4.9)]) give the following explicit expressions of L° and its closure L under the local coordinates $(\tilde{\rho}, y, y', x, \tau, \tau', \mu, \mu', \xi)$ of ${}^{b,sc}T_{\text{mf}}^*M_{b,0}$.

$$L^\circ = \left\{ (\tilde{\rho}, y, y', x, \tau, \tau', \mu, \mu', 0) \left| \begin{array}{l} s, s' \in (0, \pi), \tilde{\rho} = \sin(s)/\sin(s'), \\ \tau = -\cos(s), \tau' = \cos(s'), \\ \mu = \eta(s)\sin(s), \mu' = -\eta(s')\sin(s'), \\ h^{ij}\eta_i(s)\eta_j(s) = 1, \\ (y(s), \eta(s)) \text{ and } (y'(s'), \eta'(s')) \\ \text{are geodesics in } T^*\partial\mathcal{C} \\ \text{emanating from } (y_0, \hat{\mu}_0) \end{array} \right. \right\} \cup T_\pm \cup F_\pm$$

where the vertex sets of the square $[0, \pi]^2$ are defined for $s, s' \in \{0, \pi\}$,

$$\begin{aligned} T_\pm &= \{(\tilde{\rho}, y_0, y_0, x, \pm 1, \mp 1, 0, 0, 0) : \sigma \in (0, \infty), y_0 \in \partial\mathcal{C}\}, \\ F_\pm &= \{(\tilde{\rho}, y, y', x, \mp 1, \mp 1, 0, 0, 0) : \exists \text{ geodesics of length } \pi \text{ connecting } y, y'\}. \end{aligned}$$

The closure of L° in ${}^{s\Phi}T^*M_{b,0}$ is the set,

$$(32) \quad L = \left\{ (\tilde{\rho}, y, y', x, \tau, \tau', \mu, \mu', 0) \left| \begin{array}{l} s, s' \in (0, \pi), \sin^2(s) + \sin^2(s') > 0, \\ \tilde{\rho} = \sin(s)/\sin(s'), \tau = -\cos(s), \\ \tau' = \cos(s'), \mu = \eta(s)\sin(s), \\ \mu' = -\eta(s')\sin(s'), h^{ij}\eta_i(s)\eta_j(s) = 1, \\ (y(s), \eta(s)) \text{ and } (y'(s'), \eta'(s')) \\ \text{are geodesics in } T^*\partial\mathcal{C} \\ \text{emanating from } (y_0, \hat{\mu}_0) \end{array} \right. \right\} \cup T_\pm \cup F_\pm.$$

The Legendrian L is smooth near T_\pm but has conic singularities around F_\pm . See [18, Section 4.3]

In the coordinates $(y_0, x, \hat{\mu}_0, s, s')$, the vector field V_l is given by $\sin(s')\partial_{s'}$ and V_r is given by $-\sin(s)\partial_s$. The intersection of L and $N^*\text{diag}_{M_{b,0}}$ is given by $\{s = s'\}$. Thus L_+ is given by $\{s \leq s'\}$.

There exist Legendrian submanifolds L_\pm^\sharp of ${}^{b,sc}T_{\text{mf}}^*\tilde{M}_{b,0}$, parametrized by

$$(33) \quad L_\pm^\sharp = \{(\tilde{\rho}, y, y', x, \mp 1, \mp 1, 0, 0, 0) : y, y' \in \partial X, \tilde{\rho} \in [0, \infty]\}.$$

By [18, Proposition 4.1], the Legendrian pair $\tilde{L} = (L, L_\pm^\sharp)$, intersecting at

$$L \cap L_\pm^\sharp = F_\pm,$$

forms a pair of intersecting Legendrians with conic points. See Figure 8.

Remark 24. In contrast to the double scattering n -manifolds, $M_{b,0}$ is a $(2n+1)$ -manifold such that ${}^{s\Phi}T_{\text{mf}}^*M_{b,0}$ is a $(4n+1)$ -contact manifold. However, the additional variables (x, ξ) are irrelevant to the contact forms in Section 6.3 such that $\xi \equiv 0$ on the Legendrians and x complicates nothing but acts as a smooth parameter of Legendrians. It follows that the Legendrian structures described in [18] can be employed verbatim in the present case.

10.4. Error at the bicharacteristic. We next solve away the error in the interior of L_+ . This involves solving the transport equation globally on L . What concerns us is the solution of the transport equation near the boundaries of L , as the transport equation is smooth and always solvable in the interior of L .

To begin with we examine the form of the transport equation at the boundary of L_+ . It suffices to discuss the situation near lf as it is analogous for rf.

In a neighbourhood $\{\tilde{\rho} < 2\}$ of $\text{lf} \cap \text{mf}$, we have local coordinates $(y', x, \mu', \tilde{\rho})$ near T_+ and $(y', x, \hat{\mu}', \tilde{\rho}, s)$ away from T_+ . In either set of coordinates, the Hamilton vector field in left variables $(\tilde{\rho}, y)$ thus takes the form

$$V_l = 2\tau\tilde{\rho}\partial_{\tilde{\rho}} + Z, \quad \text{for some non-vanishing vector fields } Z.$$

Recall from (19) that the subprincipal symbol of P_{sc} , is $-\imath\varepsilon\tau + o(|\mu|)$ on L_+ , where we denote

$$\varepsilon = xe(x, y).$$

By (32), $\tilde{\rho} = 0$ amounts to $\mu = 0$ in the interior of L_+ , the subprincipal symbol of P_{sc} is also of the form $-\imath\tau\varepsilon + o(\tilde{\rho})$. Therefore, the transport equation for the symbol of order $-1/2$, with the half density factor removed, takes the form

$$-\imath(\mathcal{L}_{V_l} - n\tau + \varepsilon_l\tau + o(\tilde{\rho}))q_3^{(0)} = 0.$$

Here ε_l means ε in left variables. It reduces to the following ODE,

$$-\imath\tau(\partial_{\tilde{\rho}} + o(1))(\tilde{\rho}^{-n/2+\varepsilon_l/2}q_3^{(0)}) = 0.$$

This shows that $\tilde{\rho}^{-n/2+\varepsilon_l/2}q_3^{(0)}$ is smooth near $\text{lf} \cap \text{mf}$.

Near $L^\sharp \cap L_+$, we want to understand the order of s , which measures the singularity on L^\sharp . Recall that we lift L_+ to \hat{L}_+ by blowing up $L^\sharp \cap L_+$. By [18, (4.15)], the sum of left and right Hamilton vector fields $V_l + V_r$ lifts to

$$-2s\partial_s + o(s),$$

where W is smooth and tangent to the boundary of \hat{L}_+ . To take advantage of this, we wish to combine the left and right transport equations, since the solution to the left equation also solves the right equation. The right transport operator with respect to ρ' takes the form

$$-\imath(\mathcal{L}_{V_r} - n\tau' + \varepsilon_r\tau' + o(s)),$$

where ε_r means ε in right variables, and the Hamilton vector field in right variables is

$$V_r = -2\tau'\tilde{\rho}\frac{\partial}{\partial\tilde{\rho}} + 2\tau'\mu'\frac{\partial}{\partial\mu'} - h'\frac{\partial}{\partial\tau'} + \left(\frac{\partial h'}{\partial\mu'}\frac{\partial}{\partial y'} - \frac{\partial h'}{\partial y'}\frac{\partial}{\partial\mu'}\right).$$

In addition, if we change the left variable in ρ to the right variable in ρ' , then $q_3^{(0)}$ is replaced by $\tilde{\rho}^{-1/2-n/2}q_3^{(0)}$ in the right transport equation ($\tilde{\rho}^{-1/2-n/2}$ is the transition function in (17)). It then yields an equation of the form

$$-\imath(\mathcal{L}_{V_r} - n\tau' + \varepsilon_r\tau' + o(s))(\tilde{\rho}^{-1/2-n/2}q_3^{(0)}) = 0.$$

By using the expression of the right Hamilton vector field, we reduce this equation to

$$(V_r + \tau' + \varepsilon_r\tau' + o(s))q_3^{(0)} = 0.$$

Combining it with the left transport equation and using the fact $\tau = \tau' = -1$ on L_+^\sharp gives an equation of the form

$$2\left(s\partial_s - (n-1)/2 + (\varepsilon_l + \varepsilon_r)/2 + o(s)\right)q_3^{(0)} = 0.$$

This equation may be rewritten as

$$(\partial_s + o(1))(s^{-(n-1)/2+(\varepsilon_l+\varepsilon_r)/2} q_3^{(0)}) = 0.$$

This, together with the left and right transport equations and $\tilde{\rho} = \rho_{\text{lf}}$, shows that

$$(34) \quad q_3^{(0)} \in \rho_{\text{lf}}^{n/2-\varepsilon_l/2} \rho_{\text{rf}}^{n/2-\varepsilon_r/2} s^{(n-1)/2-(\varepsilon_l+\varepsilon_r)/2} C^\infty(s^\Phi \Omega^{1/2} \otimes S^{[-1/2]}(\hat{L}_+)).$$

In light of the short exact sequence (22), the solution $q_3^{(0)}$ to the transport equation, as a principal symbol, determines a Legendrian distribution

$$Q_3^{(0)} \in I_{s^\Phi}^{-1/2, (n-2)/2-\mathbf{e}; ((n-1-\mathbf{e})/2, (n-1-\mathbf{e})/2, 0)}(M_{b,0}; \hat{L}_+; {}^s\Phi \Omega^{1/2}),$$

such that

$$\begin{aligned} P_{sc} \circ Q_3^{(0)} - E_{sc,2} &= -E_2^{(1)} \\ &\in (\log \rho_{\text{lf}} \log s)^2 I_{s^\Phi}^{3/2, n/2-\mathbf{e}; ((n+3-\mathbf{e})/2, (n-1-\mathbf{e})/2, 0)}(M_{b,0}; (L_+, L^\sharp); {}^s\Phi \Omega^{1/2}), \end{aligned}$$

where the remainder $E_2^{(1)}$ has a principal symbol in the form

$$e_2^{(1)} \in \rho_{\text{lf}}^{n/2-\varepsilon_l/2} \rho_{\text{rf}}^{n/2-\varepsilon_r/2} s^{(n-1)/2-(\varepsilon_l+\varepsilon_r)/2} (\log \rho_{\text{lf}} \log s)^2 C^\infty(s^\Phi \Omega^{1/2} \otimes S^{[3/2]}(\hat{L}_+))$$

Here we use the notation ¹⁰

$$\begin{aligned} &(\log \rho_{\text{lf}})^{k_{\text{lf}}} (\log \rho_{\text{rf}})^{k_{\text{rf}}} I_{s^\Phi}^{m; (r_{\text{lf}}, r_{\text{rf}}, r_{\text{bf}0})}(M_{b,0}; L; {}^s\Phi \Omega^{1/2}) \\ &= \left\{ u : \begin{array}{l} \sigma^m(u) = (\log \rho_{\text{lf}})^{k_{\text{lf}}} (\log \rho_{\text{rf}})^{k_{\text{rf}}} \sigma^m(\tilde{u}) \\ \tilde{u} \in I_{s^\Phi}^{m; (r_{\text{lf}}, r_{\text{rf}}, r_{\text{bf}0})}(M_{b,0}; L; {}^s\Phi \Omega^{1/2}) \end{array} \right\} \\ &(\log \rho_{\text{lf}})^{k_{\text{lf}}} (\log \rho_{\text{rf}})^{k_{\text{rf}}} (\log s)^{k_\sharp} I_{s^\Phi}^{m, p_\sharp; (r_{\text{lf}}, r_{\text{rf}}, r_{\text{bf}0})}(M_{b,0}, \tilde{L}; {}^s\Phi \Omega^{1/2}) \\ &= \left\{ u : \begin{array}{l} \sigma^m(u) = (\log \rho_{\text{lf}})^{k_{\text{lf}}} (\log \rho_{\text{rf}})^{k_{\text{rf}}} (\log s)^{k_\sharp} \sigma^m(\tilde{u}) \\ \tilde{u} \in I_{s^\Phi}^{m, p_\sharp; (r_{\text{lf}}, r_{\text{rf}}, r_{\text{bf}0})}(M_{b,0}, \tilde{L}; {}^s\Phi \Omega^{1/2}) \end{array} \right\}. \end{aligned}$$

Note that ε is not a constant but a function of (x, y) . Hence the quadratic log terms $(\log \rho_{\text{lf}} \log s)^2$ are inevitable when the second y -derivative in P_{sc} hits $\rho_{\text{lf}}^{-\varepsilon_l/2} s^{-(\varepsilon_l+\varepsilon_r)/2}$ of the symbol of $Q_3^{(0)}$.

For any $k \in \mathbb{Z}_+$, we want to find a Legendrian distribution

$$Q_3^{(k)} \in (\log \rho_{\text{lf}} \log s)^{2k} I_{s^\Phi}^{k-1/2, n/2-1-\mathbf{e}; ((n-1-\mathbf{e})/2, (n-1-\mathbf{e})/2, 0)}(M_{b,0}; \hat{L}_+; {}^s\Phi \Omega^{1/2}),$$

which solves the left equation above up to a remainder in

$$\begin{aligned} P_{sc} \circ Q_3^{(k)} - E_2^{(k)} &= -E_2^{(k+1)} \\ &\in (\log \rho_{\text{lf}} \log s)^{2(k+1)} I_{s^\Phi}^{k+3/2, n/2-\mathbf{e}; (n+3-\mathbf{e})/2, ((n-1-\mathbf{e})/2, 0)}(M_{b,0}; (L_+, L^\sharp); {}^s\Phi \Omega^{1/2}). \end{aligned}$$

This is done by solving a left transport equation of the form

$$-2i\tau \left(\tilde{\rho} \partial_{\tilde{\rho}} + (-n/2 + k) + \varepsilon_l/2 + o(\tilde{\rho}) \right) q_3^{(k)} = e_2^{(k)}.$$

The right hand side $e_2^{(k)}$ is the principal symbol of $E_2^{(k)}$. Now we want the remainder $E_2^{(k+1)}$ to be of order $k+3/2$ at L_+ and order $(n+3-\mathbf{e})/2$ at lf, this implies

$$e_2^{(k)} \in \tilde{\rho}^{n/2-\varepsilon_l/2-k+1} (\log \tilde{\rho})^{2k} C^\infty(s^\Phi \Omega^{1/2} \otimes S^{[k+3/2]}(\hat{L}_+))$$

¹⁰We of course can define the polyhomogeneity for Legendrian distributions with the index family $(\mathcal{E}_\sharp, \mathcal{E}_{\text{lf}}, \mathcal{E}_{\text{rf}})$ instead of the vanishing orders $(p_\sharp, r_{\text{lf}}, r_{\text{rf}})$ in Section 6.4. However, the log terms do not appear in the principal terms given in the main theorem but only in the lower order terms and remainders. So we would introduce Legendrian distributions with log terms only within the proof.

$$q_3^{(k)} \in \tilde{\rho}^{n/2-\varepsilon_l/2-k} (\log \tilde{\rho})^{2k} C^\infty(s^\Phi \Omega^{1/2} \otimes S^{[k-1/2]}(\hat{L}_+)).$$

By combining this with the right transport equation and using the vector field $V_l + V_r$, we have an ODE of the form

$$\begin{aligned} & -2i\tau' \left(-s\partial_s + (\varepsilon_l + \varepsilon_r)/2 + \left(-\frac{n-1}{2} + k\right) + o(s) \right) q_3^{(k)} \\ & \in s^{(n+1-\varepsilon_l-\varepsilon_r)/2-k} \rho_{\text{lf}}^{(n-\varepsilon_l)/2-k+1} \rho_{\text{rf}}^{(n-\varepsilon_r)/2-k} (\log \rho_{\text{lf}} \log s)^{2(k+1)} C^\infty(s^\Phi \Omega^{1/2} \otimes S^{[k+3/2]}(\hat{L}_+)). \end{aligned}$$

We remark that there is one more order for s in this ODE than $e_2^{(k)}$ given in the symbol map. It is because $V_l + V_r$ vanishes on L^\sharp such that the sum of the right hand side terms of the two equations lies in

$$(\log \rho_{\text{lf}} \log s)^{2k} I_{s^\Phi}^{k+1/2, n/2+1-\mathbf{e}; ((n+3-\mathbf{e})/2, (n-1-\mathbf{e})/2, 0)}(M_{b,0}; (L_+, L^\sharp); s^\Phi \Omega^{1/2}).$$

The solution obeys

$$q_3^{(k)} \in s^{(n-1-\varepsilon_l-\varepsilon_r)/2-k} \rho_{\text{lf}}^{(n-\varepsilon_l)/2-k} \rho_{\text{rf}}^{(n-\varepsilon_r)/2-k} (\log \rho_{\text{lf}} \log s)^{2k} C^\infty(s^\Phi \Omega^{1/2} \otimes S^{[k-1/2]}(\hat{L}_+))$$

as is required for the parametrix.

By asymptotically summing these correction terms, we obtain an approximation $G_{sc,3} = \sum_{k=0}^\infty Q_3^{(k)}$ to the resolvent such that

$$\begin{aligned} P_{sc} \circ G_{sc,3} - E_{sc,2} &= -E_{sc,3} \\ &\in (\log \rho_{\text{lf}} \log s)^\infty I^{\infty, n/2+1-\mathbf{e}; ((n+3-\mathbf{e})/2, (n-1-\mathbf{e})/2, 0)}(M_{b,0}, (L_+, L^\sharp); s^\Phi \Omega^{1/2}). \end{aligned}$$

This means that the error at the scattering wavefront set at L_+ has been solved away.

In addition to removing the error at L_+ , we can further make the remainder E_3 vanish to infinite order at mf . Near $\text{mf} \cap \text{rf}$, we have the boundary defining functions $\rho_{\text{mf}} = \rho$ and $\rho_{\text{rf}} = \rho'/\rho$, whilst ρ' is the total boundary defining function. By definition, the intersecting Legendrian distribution $G_{sc,3}$, in this region, takes the form

$$e^{i/\rho'} \rho'^{(n-1)/2-\varepsilon_r/2} \sum_{k \geq 0} \rho'^k (\log s)^{2k} I_{s^\Phi}^{-n/4-k+\mathbf{e}, n/4-1/2-k}(M_{b,0}; (L_{y'}, L^\sharp); s^\Phi \Omega^{1/2}).$$

Here $L_{y'}$ is the fibre Legendrian of ${}^{b,sc}T^*(F_{\text{rf}}; \text{rf})$ with smooth parameters y' . In the meantime, the remainder, also near $\text{rf} \cap \text{mf}$, takes the form

$$E_{sc,3} \sim e^{i/\rho'} e^{i/\rho} \rho^{(n+3)/2-\varepsilon_l/2} \rho'^{(n-1)/2-\varepsilon_r/2} C^\infty(M_{b,0}; s^\Phi \Omega^{1/2}).$$

On the other hand, it is obvious that the application of P_{sc} to a function in (ρ, y) will increase the vanishing order in ρ . In particular,

$$(35) \quad P_{sc}(e^{i/\rho} \rho^{(n-1)/2-\varepsilon_l/2}) = e^{i/\rho} \rho^{(n+3)/2-\varepsilon_l/2} C^\infty(M_{b,0}; s^\Phi \Omega^{1/2}).$$

Such an error can be solved away by a term of the form

$$e^{i/\rho'} e^{i/\rho} \rho^{(n+1)/2-\varepsilon_l/2} \rho'^{(n-1)/2-\varepsilon_r/2} C^\infty(M_{b,0}; s^\Phi \Omega^{1/2}).$$

We can repeat this procedure to make the remainder vanish in ρ to infinite order. Since ρ is locally the boundary defining function of mf , we have removed the error at mf .

10.5. **Error at L^\sharp .** This step is to solve away the error at L^\sharp , i.e. where $\tilde{\rho} = 0$ and $\tau = 1$, if we live near $\text{lf} \cap \text{mf}$. This amounts to solving the transport equation

$$\left(-i\tilde{\rho}\partial_{\tilde{\rho}} - i(1/2 + n/2 + k - (2n)/4) - i\varepsilon_l/2 + o(\tilde{\rho}) \right) q_4^{(k)} = e_3^{(k)}, \quad \text{for } k \in \mathbb{Z}_+,$$

as the order of $e_3^{(k)}$ at L^\sharp is $n/2 + k - \mathbf{e}$. For $k = 1$, the short exact sequence of Legendrian distributions yields the symbol $e_3^{(1)}$ of E_3 , which takes the form

$$\tilde{\rho}^{1/2+\varepsilon_l/2} \rho_{\text{mf}}^\infty C^\infty(L^\sharp).$$

Then one can find a solution to the transport equation,

$$q_4^{(1)} \in \tilde{\rho}^{-3/2+\varepsilon_l/2} \rho_{\text{mf}}^\infty C^\infty(L^\sharp),$$

such that the Legendrian distribution

$$Q_4^{(1)} \in \rho_{\text{mf}}^\infty I_{s\Phi}^{n/2+1-\mathbf{e};((n-1-\mathbf{e})/2,(n-1-\mathbf{e})/2,0)}(M_{b,0}; L^\sharp; {}^{s\Phi}\Omega^{1/2}).$$

This reduces the remainder to

$$E_3^{(2)} \in \rho_{\text{mf}}^\infty I_{s\Phi}^{n/2+2-\mathbf{e};((n+3-\mathbf{e})/2,(n-1-\mathbf{e})/2,0)}(M_{b,0}; L^\sharp; {}^{s\Phi}\Omega^{1/2}).$$

Inductively, if the remainder $E_3^{(k)}$ satisfies

$$E_3^{(k)} \in \rho_{\text{mf}}^\infty I_{s\Phi}^{n/2+k-\mathbf{e};((n+3-\mathbf{e})/2,(n-1-\mathbf{e})/2,0)}(M_{b,0}; L^\sharp; {}^{s\Phi}\Omega^{1/2}),$$

then one can find a solution to the transport equation

$$q_4^{(k)} \in \tilde{\rho}^{-1/2-k+\varepsilon_l/2} \rho_{\text{mf}}^\infty C^\infty(M_{b,0}; L^\sharp; {}^{s\Phi}\Omega^{1/2}).$$

This gives a Legendrian distribution

$$Q_4^{(k)} \in \rho_{\text{mf}}^\infty I_{s\Phi}^{n/2+k-\mathbf{e};((n-1-\mathbf{e})/2,(n-1-\mathbf{e})/2,0)}(M_{b,0}; L^\sharp; {}^{s\Phi}\Omega^{1/2}),$$

which reduces the remainder to

$$E_3^{(k+1)} \in \rho_{\text{mf}}^\infty I_{s\Phi}^{n/2+k+1-\mathbf{e};((n+3-\mathbf{e})/2,(n-1-\mathbf{e})/2,0)}(M_{b,0}; L^\sharp; {}^{s\Phi}\Omega^{1/2}).$$

Taking an asymptotic sum yields a parametrix $G_{sc,4} = \sum Q_4^{(k)}$ and a remainder $E_{sc,4}$ lying in

$$e^{i/\rho'} \rho_{\text{lf}}^{(n-1)/2-\varepsilon_l/2} \rho_{\text{rf}}^{(n-1)/2-\varepsilon_r/2} \rho_{\text{mf}}^\infty C^\infty(M_{b,0}; {}^{b,sc}\Omega^{1/2}).$$

In addition, the error at lf can be removed by an analogous argument using (35) as for mf . Then the remainder is reduced to

$$E_{sc,4} \in e^{i/\rho'} \rho_{\text{rf}}^{(n-1)/2-\varepsilon_r/2} \rho_{\text{lf}}^\infty \rho_{\text{mf}}^\infty C^\infty(M_{b,0}; {}^{b,sc}\Omega^{1/2}).$$

10.6. **Error at bf_0 .** The error at bf_0 can be solved away by a standard normal operator argument. In fact, the normal operator $\mathcal{N}_{\text{bf}_0}(P_{sc})$ is just $\Delta_{\mathfrak{C}} - (1 + i0)$. The parametrices we have constructed actually include the region near bf_0 but it might not agree with $R_{\mathfrak{C},sc}$ at $\text{bf}_0 \cap (\text{mf} \cup \text{lf} \cup \text{rf})$ due to the non-uniqueness of diffractive waves. Therefore, we further take

$$\mathcal{N}_{\text{bf}_0}(G_{sc,5}) = (R_{\mathfrak{C},sc} - \mathcal{N}_{\text{bf}_0}(G_{sc,1} + G_{sc,2} + G_{sc,3} + G_{sc,4})) \circ \mathcal{N}_{\text{bf}_0}(E_{sc,4}),$$

which is a Legendrian distribution associated only with the diffractive Legendrian L^\sharp . The restriction of the remainder $E_{sc,4}$ to bf_0 is of the form,

$$\mathcal{N}_{\text{bf}_0}(E_{sc,4}) \in e^{i/\rho'} \rho_{\text{rf}}^{(n-1)/2} \rho_{\text{lf}}^\infty \rho_{\text{mf}}^\infty C^\infty(\text{bf}_0, {}^{sc}\Omega^{1/2}).$$

Then $\mathcal{N}_{\text{bf}_0}(G_{sc,5})$ solves the normal operator equation with a remainder E_{sc} vanishing at bf_0 , whilst it lies in the space

$$I_{s\Phi}^{(n-2)/2;((n-1)/2,(n-1)/2,0)}(M_{b,0}; L^\sharp; {}^{s\Phi}\Omega^{1/2}),$$

which is even better than existing parametrices due to $\varepsilon|_{\text{bf}_0} = 0$.

Finally, the desired parametrix is just

$$G_{sc} = G_{sc,1} + G_{sc,2} + G_{sc,3} + G_{sc,4} + G_{sc,5}$$

with a remainder of the form

$$E_{sc} \in e^{i/\rho'} \rho_{\text{rf}}^{(n-1)/2-\varepsilon_r/2} \rho_{\text{bf}_0} \rho_{\text{lf}}^\infty \rho_{\text{mf}}^\infty C^\infty(M_{b,0}; {}^{b,sc}\Omega^{1/2}).$$

□

11. THE (sc, b) -REGIME PARAMETRIX

Recall that the (sc, b) -regime is a neighbourhood of $\{r' < 1\} \cup \{\rho < 1\}$ in $M_{b,0}$ and furnished with local coordinates (ρ, y, r', y', \hbar) . It has three boundary hypersurfaces $\text{rb}, \text{rb}_0, \text{bf}_0$ with boundary defining functions $r', \rho, \hbar/\rho$ respectively, where the face rb_0 is renamed lf in the sc -regime since it is defined by the left boundary defining function ρ . The compressed density ${}^{b,sc}\Omega(M_{b,0})$ locally takes the form

$$\left| \frac{d\rho dy dr' dy' d\hbar}{\rho^{n+1} r' \hbar} \right|.$$

The operator \tilde{P}_\hbar , as a left operator on $M_{b,0}$, is also a sc -operator in the (sc, b) -regime, like P_{sc} in the sc -regime. The face rb_0 in this regime is viewed as the interior of lf in the sc -regime. The sc -calculus for Legendrian distributions extends to the whole rb_0 indeed. We can construct the following parametrix for \tilde{P}_\hbar in this regime.

Proposition 25. *In the (sc, b) -regime of $M_{b,0}$, there is a parametrix $G_{(sc,b)}$ lying in*

$$\rho_{\text{rb}}^{n/2} I_{s\Phi}^{n/2+1-\mathbf{e};((n-1-\mathbf{e})/2,\infty,0)}(M_{b,0}; L^\sharp|_{\text{rb}_0}; {}^{s\Phi}\Omega^{1/2}).$$

such that

$$\tilde{P}_\hbar \circ G_{(sc,b)} - \chi_b \text{Id} \chi_{sc} = -E_{(sc,b)} \in \rho_{\text{bf}_0} \rho_{\text{rb}}^{n/2} \rho_{\text{rb}_0}^\infty C^\infty(M_{b,0}; {}^{b,sc}\Omega^{1/2}).$$

Proof. We shall first remove completely the error at rb_0 and then employ the normal operator at bf_0 to obtain the vanishing order of the parametrix at rb .

11.1. Boundary error removal. Noting that $\text{lf} \subset \text{rb}_0 = \overline{\{\rho = 0\}}$, one can extend the construction of $G_{sc,4}$ and $G_{sc,5}$ to the (sc, b) -regime near rb_0 . In fact, the face rb_0 in the (sc, b) -regime can be regarded as the interior of lf (away from mf) in the sc -regime. The only change of the argument would be the use of coordinates near the interior lf away from mf rather than those near the corner $\text{mf} \cap \text{lf}$. This yields a parametrix $G_{(sc,b)}$ which is of the form

$$I_{s\Phi}^{n/2+1-\mathbf{e};((n-1-\mathbf{e})/2,\infty,0)}(M_{b,0}; L^\sharp|_{\text{rb}_0}; {}^{s\Phi}\Omega^{1/2}).$$

Then we remove the error at rb_0 by using the computation (35) and have the remainder vanishing at rb_0 and bf_0 in exactly the same way as for lf and bf_0 in the sc -regime.

11.2. Vanishing order at rb. Since x is the boundary defining function of bf_0 , the normal operator $\mathcal{N}_{\text{bf}_0}(\tilde{P}_h)$ is equal to

$$-(\rho^2 \partial_\rho)^2 + (n-1)\rho^3 \partial_\rho + \rho^2 \Delta_{h_0} - (1 + \imath 0).$$

This is equal to $\Delta_{\mathfrak{C}} - (1 + \imath 0)$ near the infinity of the metric cone \mathfrak{C} . The resolvent of $\mathcal{N}_{\text{bf}_0}(\tilde{P}_h)$ can be calculated readily by separation of variables. Given in [14, (5.5) (5.6)], it reads

$$(\mathcal{N}_{\text{bf}_0}(\tilde{P}_h))^{-1} = \frac{\pi \imath r'}{2\rho} \sum_{j=0}^{\infty} \Pi_{\mathbb{E}_j}(y, y') J_{\nu_j}(r') \text{Ha}_{\nu_j}^{(1)}(1/\rho) \left| \frac{d\rho dy dr' dy' d\hbar}{\rho r' \hbar} \right|^{1/2}.$$

Using again the asymptotic behaviours (29) and (30) of Bessel and Hankel functions, we obtain that

$$(\mathcal{N}_{\text{bf}_0}(\tilde{P}_h))^{-1} \sim \rho^{(n-1)/2} r'^{n/2} e^{\imath/\rho} \left| \frac{d\rho dy dr' dy' d\hbar}{\rho^{n+1} r' \hbar} \right|^{1/2}.$$

This implies $G_{(b,sc)}$ is taken to vanish to order $n/2$ at rb, to match the parametrix at bf_0 as well as that in the b -regime.¹¹ \square

12. THE TRUE RESOLVENT

Through Proposition 21, Proposition 22, Proposition 23, and Proposition 25, we have constructed the parametrices G_b , $G_{(b,sc)}$, G_{sc} , and $G_{(sc,b)}$ with the remainders E_b , $E_{(b,sc)}$, E_{sc} , and $E_{(sc,b)}$ in the b -regime, the (b, sc) -regime, the sc -regime, and the (sc, b) -regime respectively. They satisfy

$$\begin{aligned} \tilde{P}_h \circ (G_b + G_{(b,sc)} + G_{sc} + G_{(sc,b)}) - \text{Id} &= -\tilde{E} = -E_b - E_{(b,sc)} - E_{sc} - E_{(sc,b)}, \\ E_b + E_{(b,sc)} &\in \rho_{\text{lb}_0}^\infty \Psi_b^{-\infty, (\infty, \hat{\mathcal{E}}_b+1, \infty, \infty)}(M_{b,0}; {}^{b,sc}\Omega^{1/2}), \quad \text{near } \{r < 1\}, \\ E_{sc} + E_{(sc,b)} &\in \rho_{\text{bf}_0}^\infty e^{\imath/\rho'} \rho_{\text{rf}}^{(n-1-\varepsilon_r)/2} \rho_{\text{rb}}^{n/2} \rho_{\text{lf}}^\infty \rho_{\text{mf}}^\infty C^\infty(M_{b,0}; {}^{b,sc}\Omega^{1/2}), \quad \text{near } \{\rho < 1\}. \end{aligned}$$

From this we obtain the true resolvent.

Theorem 26. *The resolvent kernel R of P_h is a distribution on $M_{b,0}$ consisting of*

- $R_b = \chi_b R \chi_b$ is supported near the b -regime and satisfies
 - R_b is conormal of order -2 to $\text{diag}_{M_{b,0}}$,
 - R_b is polyhomogeneous conormal to $\text{lb}, \text{rb}, \text{bf}, \text{bf}_0$ and lies in the space

$$\mathcal{A}^{(\hat{\mathcal{E}}_b+1, \check{\mathcal{E}}_b+1, \tilde{\mathcal{E}}_b+2, \mathbb{N}_0)}(M_{b,0}; {}^{b,sc}\Omega^{1/2}),$$

where the index sets $\hat{\mathcal{E}}_b, \check{\mathcal{E}}_b, \tilde{\mathcal{E}}_b$ are defined by (26), (27), (28) respectively,

- R_b vanishes to order 2 at bf , but also to order $n/2$ at lb and rb ;
- $R_{(b,sc)} = \chi_b R \chi_{sc}$ is supported near the (b, sc) -regime such that
 - $R_{(b,sc)}$ vanishes to order $n/2$ at lb and to order $(n-1)/2$ at lb_0 ;
- $R_{sc} = \chi_{sc} R \chi_{sc}$ is supported near the sc -regime and satisfies
 - R_{sc} is conormal of order -2 to $\text{diag}_{M_{sc,b,0}}$,
 - R_{sc} , near $\text{diag}_{M_{b,0}} \cap \text{mf}$, is intersecting Legendrian distributions associated with the diagonal conormal bundle and geometric Legendrian L , lying in the space

$$I_{sc,c}^{-1/2}(\tilde{M}_{b,0}; (N^* \text{diag}_{M_{b,0}}, L^\circ); {}^{b,sc}\Omega^{1/2}),$$

¹¹Be aware that one cannot solve away the error at rb!

- R_{sc} , near $(\text{lf} \cap \text{mf}) \cup (\text{rf} \cap \text{mf})$, is intersecting Legendrian distributions associated with the pair of geometric Legendrian L and diffractive Legendrian L^\sharp explicitly given in (32) and (33), lying in the space

$$I_{s\Phi}^{-1/2, (n-2)/2 - \mathbf{e}; ((n-1-\mathbf{e})/2, (n-1-\mathbf{e})/2, 0)}(M_{b,0}; (L, L^\sharp); {}^s\Phi\Omega^{1/2}),$$

where \mathbf{e} is defined in (1),

- The loss of order \mathbf{e} is caused by the additional oscillations,

$$\rho_{\text{lf}}^{-\varepsilon_l/2} \rho_{\text{rf}}^{-\varepsilon_r/2} \rho_{\sharp}^{-(\varepsilon_l + \varepsilon_r)/2}, \quad \text{near } \text{lb}_0, \text{rb}_0, L^\sharp,$$

of the principal symbol of R_{sc} in (34). It occurs (i.e. $\mathbf{e} \neq 0$) if and only if the metric is of non-product type (i.e. the boundary metric $h(x, y, dy)$ is dependent of x),

- R_{sc} vanishes to $(n-1-\mathbf{e})/2$ order at lf and rf ;
- $R_{(sc,b)} = \chi_{sc} R \chi_b$ is supported near the (sc, b) -regime such that
 - $R_{(sc,b)}$ vanishes to order $n/2$ at rb and to order $(n-1-\mathbf{e})/2$ at rb_0 .

Proof. By standard Fredholm theory, obtaining the true resolvent from parametrices will be done by understanding the inversion of $\text{Id} - \tilde{E}$ and its composition with the parametrices.

First of all, $\text{Id} - \tilde{E}$ is not necessarily invertible in the sc -regime even though \tilde{E} is a compact operator. Hassell-Vasy [18, Section 6.1] showed that this issue can be fixed by adding to the parametrix a finite rank perturbation. The remainder \tilde{E} is a Hilbert-Schmidt kernel on the weighted L^2 space of the sc -regime, say $\rho^l \chi_{sc} L^2(\mathcal{C}; {}^b, {}^{sc}\Omega^{1/2})$ uniformly in \hbar for any $l > (1 + \mathbf{e})/2$ in virtue of the stability condition (1).¹² One can choose ψ_i and ϕ_i supported near $\{\rho \leq 1\} \subset \mathcal{C}$ such that $\{\rho^l \psi_i\}$ spans the null space of $\text{Id} - \tilde{E}$ and $\{P_{sc}(\phi_i)\}$ spans the supplementary subspace to the range. Then we correct G_{sc} by adding a finite rank perturbation $G_{sc,p} = \sum_i \phi_i \langle \rho^{2l} \psi_i, \cdot \rangle$, where $\langle \cdot, \cdot \rangle$ denotes the L^2 -inner product. This results in an improved remainder E such that

$$\text{Id} - E = \text{Id} - \tilde{E} + \sum (P_{sc}(\phi_i)) \langle \rho^{2l} \psi_i, \cdot \rangle.$$

is invertible on $\rho^l \chi_{sc} L^2(\mathcal{C}; {}^b, {}^{sc}\Omega^{1/2})$. Moreover, this correction term does not change the structure of the resolvent.

Since $\text{Id} - E$ is invertible now, we can analyse the inverse

$$\text{Id} + S = (\text{Id} - E)^{-1}.$$

The identity

$$\text{Id} = (\text{Id} - E)(\text{Id} + S) = (\text{Id} + S)(\text{Id} - E)$$

implies S is also Hilbert-Schmidt and

$$S = E + E^2 + ESE.$$

Recall that the remainder E lies in the space $\mathcal{D}_b + \mathcal{D}_{sc}$, where

$$\begin{aligned} \mathcal{D}_b &= \rho_{\text{lb}_0}^\infty \mathcal{A}^{(\infty, \hat{\mathcal{E}}_b + 1, \infty, \infty)}(M_{b,0}; {}^b, {}^{sc}\Omega^{1/2}), & \text{near } \{r < 1\}, \\ \mathcal{D}_{sc} &= \rho_{\text{bf}_0}^\infty e^{i/\rho'} \rho_{\text{rb}}^{n/2} \rho_{\text{rf}}^{(n-1)/2 - \varepsilon_r/2} \rho_{\text{lf}}^\infty \rho_{\text{mf}}^\infty C^\infty(M_{b,0}; {}^b, {}^{sc}\Omega^{1/2}), & \text{near } \{\rho < 1\}. \end{aligned}$$

The term E^2 lies in the composition space

$$(\mathcal{D}_b + \mathcal{D}_{sc}) \circ (\mathcal{D}_b + \mathcal{D}_{sc}).$$

On the one hand, we see $\mathcal{D}_b \circ \mathcal{D}_b \subset \mathcal{D}_b$ and $\mathcal{D}_{sc} \circ \mathcal{D}_{sc} \subset \mathcal{D}_{sc}$ simply by the calculus of polyhomogeneous conormal distributions. On the other hand, it follows from the vanishing

¹²Recall that $\rho = \hbar/x$ and χ_{sc} is a cut-off function on \mathcal{C} supported near $\{\rho < 1\}$.

properties that the cross terms obey $\mathcal{D}_b \circ \mathcal{D}_{sc} \subset \mathcal{D}_{sc}$ and $\mathcal{D}_{sc} \circ \mathcal{D}_b \subset \mathcal{D}_b$. Therefore $-E + E^2$ also lies in $\mathcal{D}_b + \mathcal{D}_{sc}$.

To analyse ESE , it is convenient to blow down $M_{b,0}$ to M . The boundary defining functions of boundary faces of $M_{b,0}$ will be pushed forward to

$$(\beta_{b,0})_* \rho_{lf} \rho_{mf} = \langle r \rangle^{-1}, \quad (\beta_{b,0})_* \rho_{rf} \rho_{mf} = \langle r' \rangle^{-1}, \quad (\beta_{b,0})_* \rho_{rf} \rho_{lf} \rho_{bf_0} = \hbar$$

in the sc -regime and the (sc, b) -regime,

$$(\beta_{b,0})_* \rho_{lb} \rho_{bf} = r / \langle r \rangle, \quad (\beta_{b,0})_* \rho_{rb} \rho_{bf} = r' / \langle r' \rangle, \quad (\beta_{b,0})_* \rho_{lb_0} \rho_{bf_0} = \hbar$$

in the b -regime and the (b, sc) -regime, where we denote $r = x/\hbar$, $r' = x'/\hbar$ and the Japanese bracket $\langle \cdot \rangle = (1 + \cdot^2)^{1/2}$. The push-forward $\beta_{b,0*}(E)(z, z', \hbar)$ thus lies in the space

$$(36) \quad \hbar^\infty \left(\frac{r}{\langle r \rangle} \right)^\infty \langle r \rangle^{-\infty} e^{i\langle r' \rangle} \langle r' \rangle^{-(n-1)/2 + \varepsilon_r/2} \sum_{(z,k) \in \hat{\mathcal{E}}_b + 1} \left(\frac{r'}{\langle r' \rangle} \right)^z \left\langle \log \left(\frac{r'}{\langle r' \rangle} \right) \right\rangle^k C^\infty(M).$$

This distribution is weighted square integrable in the right variables, that is in the space $\langle r' \rangle^{-l} L^2(\mathcal{C})$.

Since S is Hilbert-Schmidt on $\langle r \rangle^{-l} L^2(\mathcal{C})$, the kernel of $S(z'', z''', \hbar)$ has to lie in the space

$$\langle r'' \rangle^{-l} \langle r''' \rangle^l L^2(M) \quad \text{uniformly as } \hbar \rightarrow 0,$$

where $r'' = x''/\hbar$, $r''' = x'''/\hbar$, $z'' = (x'', y'')$, $z''' = (x''', y''')$.

The term $\beta_{b,0*}(ESE)(z, z', \hbar)$ has the expression

$$\int_{\mathcal{C}^2} \left(\beta_{b,0*}(E)(z, z'', \hbar) \right) \left(S(z'', z''', \hbar) \right) \left(\beta_{b,0*}(E)(z''', z', \hbar) \right) dg(z'') dg(z''').$$

By Cauchy-Schwartz, this integral takes the form (36), i.e. ESE lies in $\mathcal{D}_b + \mathcal{D}_{sc}$. Then S in turn lies in $\mathcal{D}_b + \mathcal{D}_{sc}$.

Finally, the desired true resolvent is

$$R = G(\text{Id} + S) \quad \text{where } G = G_{sc} + G_{sc,p} + G_{(sc,b)} + G_b + G_{(b,sc)}.$$

This is distributions on $M_{b,0}$,¹³ whilst S is a distribution on M . To calculate their compositions, we introduce the projection maps from the triple space $M_{b,0} \times \mathcal{C}$,

$$\mathcal{P}_G(M_{b,0} \times \mathcal{C}) = \beta_{b,0}(M_{b,0}),$$

$$\mathcal{P}_S(M_{b,0} \times \mathcal{C}) = \beta_\hbar \circ \tilde{\mathcal{P}}_R(M_{b,0}) \times \mathcal{C},$$

where β_\hbar is the blow-down map from \mathcal{C}_b to $\mathcal{C} \times [0, 1)$ and $\tilde{\mathcal{P}}_R$ is the stretched right projection from $M_{b,0}$ to \mathcal{C}_b . In local coordinates, we write

$$(z, z'', \hbar) \in \mathcal{P}_G(M_{b,0} \times \mathcal{C}) \quad \text{and} \quad (z'', z', \hbar) \in \mathcal{P}_S(M_{b,0} \times \mathcal{C}).$$

We pull back G and S to the triple space $M_{b,0} \times \mathcal{C}$. The pullback $\mathcal{P}_G^*(G(z, z'', \hbar))$ does not change as they are irrelevant to the variable from \mathcal{C} . But $\mathcal{P}_S^*(S(z'', z', \hbar))$ will lift to a distribution of the form

$$\rho_{mf}^\infty \rho_{bf}^\infty \rho_{rb}^\infty \rho_{rb_0}^\infty e^{i\langle r' \rangle} \langle r' \rangle^{-(n-1)/2 + \varepsilon_r} \sum_{(z,k) \in \hat{\mathcal{E}}_b + 1} \left(\frac{r'}{\langle r' \rangle} \right)^z \left\langle \log \left(\frac{r'}{\langle r' \rangle} \right) \right\rangle^k C^\infty(M_{b,0} \times \mathcal{C}).$$

¹³Although the sc -pseudodifferential part of G_{sc} is defined on $M_{sc,b,0}$, it doesn't affect the composition with S , since S vanishes to infinite order at mf on $M_{b,0}$.

The composition $GS(z, z', \hbar)$ on $M_{b,0}$ has the expression

$$\int_{\mathcal{C}} \mathcal{P}_G^*(G(z, z'', \hbar)) \mathcal{P}_S^*(S(z'', z', \hbar)) dg(z'').$$

This actually makes a b -fibration (in the sense of Melrose [27]) from $M_{b,0} \times \mathcal{C}$ to $M_{b,0}$ by projecting off the middle variables of \mathcal{C}^3 (or the right variables of $M_{b,0}$). Since the polyhomogeneity and oscillations in r' agree with the behaviour of G in right variables, the push-forward theorem [27, Theorem 5] guarantees GS still lies in the space we have described in the theorem.¹⁴ \square

13. APPLICATION TO SCHRÖDINGER EQUATIONS

13.1. The Schrödinger propagator on \mathcal{C} . Now let us consider the Schrödinger propagator $e^{it\Delta_{\mathcal{C}}}$ on a non-product conic manifold \mathcal{C} with a single cone tip, which is viewed as a local neighbourhood of \mathcal{X} . By the spectral theorem of projection valued form, the Schwartz kernel of $e^{it\Delta_{\mathcal{C}}}$ reads,

$$e^{it\Delta_{\mathcal{C}}} = \int_{\text{spec}(\Delta_{\mathcal{C}})} e^{it\lambda^2} dE_{\sqrt{\Delta_{\mathcal{C}}}}(\lambda),$$

where $E_{\bullet}(\lambda)$ is the spectral projection corresponding to a self-adjoint operator \bullet , $dE_{\bullet}(\lambda)$ is called the spectral measure of \bullet , and this equation means

$$\langle e^{it\Delta_{\mathcal{C}}} \psi, \psi \rangle = \int_{\text{spec}(\Delta_{\mathcal{C}})} e^{it\lambda^2} d\langle E_{\sqrt{\Delta_{\mathcal{C}}}}(\lambda) \psi, \psi \rangle.$$

Stone's formula relates the spectral measure to the resolvent of $\Delta_{\mathcal{C}}$,

$$dE_{\sqrt{\Delta_{\mathcal{C}}}}(\lambda) = \frac{\lambda}{i\pi} \left((\Delta_{\mathcal{C}} - (\lambda + i0)^2)^{-1} - (\Delta_{\mathcal{C}} - (\lambda - i0)^2)^{-1} \right) d\lambda.$$

By combining this together with $\Delta_{\mathcal{C}} - (\lambda \pm i0)^2 = \lambda^2 P_h(\pm i0)$, we see

$$dE_{\sqrt{\Delta_{\mathcal{C}}}}(\lambda) = \frac{1}{i\pi} \left(P_h^{-1}(+i0) - P_h^{-1}(-i0) \right) \frac{d\lambda}{\lambda}.$$

For finite spectral parameters, Loya [26, Theorem 6.1] proved that $(\Delta_{\mathcal{C}} - (\lambda \pm i0)^2)^{-1}$ is meromorphic in \mathbb{C} and has only finite rank singularities. Then the spectral measure $dE_{\sqrt{\Delta_{\mathcal{C}}}}(\lambda)$ is also meromorphic and has only finite rank singularities. When the spectral parameter goes to infinity, we have constructed the resolvent $P_h^{-1}(\pm i0)$, as in Theorem 26. The subtraction in Stone's formula cancels out the diagonal singularities. Therefore the spectral measure is distributions defined on $M_{b,0}$, satisfying

- $\chi_b dE_{\sqrt{\Delta_{\mathcal{C}}}}(\lambda)$, supported in the b -regime and (b, sc) -regime, is a conormal distribution to lb and rb, vanishing at lb₀, rb₀, and bf₀;
- $\chi_{sc} dE_{\sqrt{\Delta_{\mathcal{C}}}}(\lambda)$, supported in the sc -regime and (sc, b) -regime, is intersecting Legendrian distributions with conic points, microlocally supported in the Legendrian pair \tilde{L} , vanishing at lf and rf.

Hence it is convenient to decompose the Schrödinger propagator into cut-off operators,

$$\begin{aligned} e_{\text{low}}^{it\Delta_{\mathcal{C}}} &= \int \chi_{\{\lambda < 2\}} e^{it\lambda^2} dE_{\sqrt{\Delta_{\mathcal{C}}}}(\lambda), \\ e_{\text{high}}^{it\Delta_{\mathcal{C}}} &= \int \chi_{\{\lambda > 1\}} e^{it\lambda^2} dE_{\sqrt{\Delta_{\mathcal{C}}}}(\lambda), \end{aligned}$$

¹⁴The error term GS is actually better than G . Since the infinite order vanishing at rf of $\mathcal{P}_S^* S$, we conclude that GS does not have the Legendrian structure over rf. But this is not important here.

where $\chi_{\{\lambda>1\}}$ and $\chi_{\{\lambda<2\}}$ are smooth cut-off functions supported in $(1, \infty)$ and $[0, 2)$ and satisfying $\chi_{\{\lambda<2\}} + \chi_{\{\lambda>1\}} = 1$ for any $\lambda \geq 0$.

It follows that the low energy cut-off $e_{\text{low}}^{it\Delta_c}$ is a bounded operator both from $L^2(\mathcal{C})$ to $L^2(\mathcal{C})$ and $L^1(\mathcal{C})$ to $L^\infty(\mathcal{C})$ uniformly in t .

The high energy cut-off $e_{\text{high}}^{it\Delta_c}$ is more involved. We encounter two hurdles. On the one hand, the spectral measure will have the conic singularities near $L^\# \cap L$. On the other hand, the expression of the Schrödinger propagator transforms the spectral parameter λ to the time parameter t . Unlike the resolvent and the spectral measure, one has to employ the quadratic scattering cotangent bundle to describe the Schrödinger propagator in the phase space as in [40, 19, 20].

However, understanding this finer structure does not seem necessary for our purpose. Instead, we shall adopt the classical TT^* approach and Keel-Tao's endpoint method to establish Strichartz estimates. This can be achieved by (micro)localizing the spectral measure around the diagonal as in [15, 21].¹⁵

13.2. Microlocalization and the microsupport. The microlocalization refers to the refined localization on cotangent bundles by introducing smooth cut-offs also on fibre variables not only physical variables.

The microsupport is useful to formulate the microlocalization by the conjugation. The microsupport of a distribution is defined by negating the right fibre variables of its wavefront set. Specifically, given a distribution u in the sc -regime, we say $q' = (z, z', \zeta, \zeta')$ lies in the microsupport $\text{WF}'_{sc}(u)$ if and only if $q = (z, z', \zeta, -\zeta')$ lies in the wave front set $\text{WF}_{sc}(u)$.

For a Legendrian distribution U associated with a Legendrian L , its sc -wavefront set $\text{WF}_{sc}(U)$ is contained in L . Then the sc -microsupport $\text{WF}'_{sc}(U)$ is contained in $L' = \{q' : q \in L\}$.

For a sc -pseudodifferential operator Q , its sc -wavefront set $\text{WF}_{sc}(U)$ is contained in the diagonal conormal bundle $\{(\rho, y, \rho, y, \tau, \mu, -\tau, -\mu)\}$. Since both left and right projections to the single space are diffeomorphisms,

$$\begin{aligned} \pi_L &: (\rho, y, \rho', y', \tau, \mu, \tau', \mu') \rightarrow (\rho, y, \tau, \mu) \\ \pi_R &: (\rho, y, \rho', y', \tau, \mu, \tau', \mu') \rightarrow (\rho', y', \tau', \mu'), \end{aligned}$$

we can view the sc -microsupport $\text{WF}'_{sc}(Q)$ as a subset of $\{(\rho, y, \tau, \mu)\}$ in the left space, and the sc -microsupport $\text{WF}'_{sc}(Q^*)$ as a subset of $\{(\rho', y', \tau', \mu')\}$ in the right space where the fibre variables are negated.

Conjugating Legendrian distributions U by a sc -pseudodifferential operator Q , we can microlocalize the Legendrian distributions, in terms of the microsupport of Q and U ,

$$(37) \quad \text{WF}'_{sc}(QU) \subset \pi_L^{-1}\text{WF}'_{sc}(Q) \cap \text{WF}'_{sc}(U)$$

$$(38) \quad \text{WF}'_{sc}(UQ^*) \subset \pi_R^{-1}\text{WF}'_{sc}(Q^*) \cap \text{WF}'_{sc}(U)$$

$$(39) \quad \text{WF}'_{sc}(QUQ^*) \subset \pi_L^{-1}\text{WF}'_{sc}(Q) \cap \text{WF}'_{sc}(U) \cap \pi_R^{-1}\text{WF}'_{sc}(Q^*).$$

¹⁵Apart from removing conic singularities, the microlocalization on asymptotically conic manifolds in [15, 21] also aims to avoid the potential conjugate points near the infinity. However, such an issue does not arise near the cone tip, since the geodesics are all simple here.

13.3. The conjugated spectral measure. Consider the conjugation of the spectral measure, $Q_\alpha(\lambda)dE_{\sqrt{\Delta_C}}(\lambda)Q_\alpha^*(\lambda)$, by a family of pseudodifferential operators $\{Q_\alpha(\lambda)\}_\alpha$. In light of (39), we can view this as the desired (micro)localized spectral measure by choosing the support of every $Q_\alpha(\lambda)$ sufficiently small in the physical space as well as in the phase space. Consequently, $Q_\alpha(\lambda)dE_{\sqrt{\Delta_C}}(\lambda)Q_\alpha^*(\lambda)$ will have to be supported in a neighbourhood of the diagonal with $\text{mf} \cap \text{diag}_{M_{b,0}}$ removed (which is cancelled out in Stone's formula). In particular, we can make them supported away from the conic points in the *sc*-regime. This significantly simplifies the microlocal structure in the *sc*-regime, as $Q_j(\lambda)dE_{\sqrt{\Delta_C}}(\lambda)Q_j^*(\lambda)$ is a Legendrian distribution microlocally supported only on the geometric Legendrian L .

Specifically, we shall prove that

Proposition 27. *For $\lambda > 1$, there exists a finite collection of pseudodifferential operators $\{Q_\alpha(\lambda)\}_{\alpha \in \{b, tr, 0, 1, \dots, N\}}$ such that it makes a finite partition of identity on $L^2(\mathcal{C})$,*

$$\text{Id} = Q_b(\lambda) + Q_{tr}(\lambda) + Q_0(\lambda) + \sum_{j=1}^N Q_j(\lambda),$$

and the kernel of each conjugated spectral measure on $M = \mathcal{C}^2 \times [0, 1)$ takes the form

$$Q_\alpha(\lambda)dE_{\sqrt{\Delta_C}}(\lambda)Q_\alpha^*(\lambda)(z, z') = \lambda^{n-1+\mathbf{e}/2} \left(\sum_{\pm} e^{\pm i \lambda d_{\mathcal{C}}(z, z')} a_{\pm}(z, z', \lambda) + b(z, z', \lambda) \right),$$

where \mathbf{e} is defined in (1), $d_{\mathcal{C}}(\cdot, \cdot)$ is the distance function on \mathcal{C} and the amplitudes satisfy the size estimates

$$(40) \quad |\partial_\lambda^k a_{\pm}(\lambda, z, z')| \lesssim \lambda^{-k} (1 + \lambda d_{\mathcal{C}}(z, z'))^{-(n-1)/2},$$

$$(41) \quad |\partial_\lambda^k b(\lambda, z, z')| \lesssim (1 + \lambda d_{\mathcal{C}}(z, z'))^{-K}, \quad \text{for any } K \in \mathbb{N}.$$

Proof. We apply the ideas from [15, Theorem 1.13] and its improvement [21, Proposition 1.5] to our analysis on $M_{b,0}$.

Partition of identity. Before showing the estimates, we start from elucidating how we find the partition of identity $\{Q_\alpha(\lambda)\}_\alpha$. We view these pseudodifferential operators as operators compactly supported on \mathcal{C} with a parameter $\hbar = 1/\lambda$. Then the conjugation $Q_\alpha(\lambda) \bullet Q_\alpha^*(\lambda)$ is viewed as a restriction to a neighbourhood of the diagonal in $M_{b,0}$. The diagonal of $M_{b,0}$ is a submanifold which intersects the faces $\{\text{bf}, \text{bf}_0, \text{mf}\}$. However, bf_0 is viewed as the interior since both $0 < r < \infty$ and $0 < r' < \infty$.

First of all, we divide \mathcal{C} , with a parameter $\hbar \in [0, 1)$, into three overlapping parts,

$$\begin{aligned} \mathcal{C}_b(\hbar) &= \{r = x/\hbar < \epsilon\} \\ \mathcal{C}_{tr}(\hbar) &= \{r = x/\hbar \geq \epsilon/2\} \cap \{\rho = \hbar/x \geq \epsilon/2\} \\ \mathcal{C}_{sc}(\hbar) &= \{\rho = \hbar/x < \epsilon\} \end{aligned}$$

for some sufficiently small $\epsilon > 0$.¹⁶ Then there exists a partition of identity on \mathcal{C} ,

$$\text{Id} = Q_b(\lambda) + Q_{tr}(\lambda) + Q_{sc}(\lambda),$$

with 0-th degree pseudodifferential operators $Q_b(\lambda)$, $Q_{tr}(\lambda)$, and $Q_{sc}(\lambda)$ supported in $\mathcal{C}_b(\hbar)$, $\mathcal{C}_{tr}(\hbar)$, and $\mathcal{C}_{sc}(\hbar)$ respectively. This leads to the following consequences :

- the conjugation $Q_b(\lambda) \bullet Q_b^*(\lambda)$ on $M_{b,0}$ is supported in a neighbourhood of bf ,
- the conjugation $Q_{tr}(\lambda) \bullet Q_{tr}^*(\lambda)$ on $M_{b,0}$ is supported away from the boundaries,¹⁷
- the conjugation $Q_{sc}(\lambda) \bullet Q_{sc}^*(\lambda)$ on $M_{b,0}$ is supported in a neighbourhood of mf .

¹⁶The interior $\mathcal{C} \setminus \mathcal{K}$ is thought of as a part in \mathcal{C}_{sc} .

¹⁷The interior of the face bf_0 is viewed as the interior!

Furthermore, we want to refine the decomposition near mf to make the conjugation near mf away from the diffractive Legendrian L^\sharp i.e. remove the conic singularities occur at $L^\sharp \cap L$. To accomplish this, we need to work on the phase space restricted to $\{\rho < \epsilon\}$. Recall, from (32) and (33), that the geometric Legendrian is contained in the pull-back to ${}^{sc}\Phi T^*M_{b,0}$ of the characteristic variety $\Sigma(P_{sc}) \subset {}^{sc}T^*\mathcal{C}|_{\{\rho < \epsilon\}}$,

$$\Sigma(P_{sc}) = \{(\rho, y, \tau, \mu) : \rho = 0, \tau^2 + h^{ij}\mu_i\mu_j = 1\},$$

whilst the conic singularities of the resolvent and spectral measure occur on $(\text{mf} \cap \text{lf}) \cup (\text{mf} \cap \text{rf}) \subset M_{b,0}$, in terms of microsupport, at

$$(L^\sharp)' \cap L' = \{(\tilde{\rho}, y, y', \hbar, \tau, \tau', \mu, \mu', 0) : \mu = \mu' = 0, \tau = \pm 1, \tau' = \mp 1\} \subset \text{WF}'_{sc}(dE_{\sqrt{\Delta_C}}),$$

where $\tilde{\rho} = \rho/\rho'$. Hence we decompose ${}^{sc}T^*\mathcal{C}|_{\{\rho < \epsilon\}}$ into the following overlapping subsets

$$\begin{aligned} \mathcal{I}_0 &= \{(\rho, y, \tau, \mu) : \rho < \epsilon, \tau^2 + h^{ij}\mu_i\mu_j \leq 1 - \epsilon/2 \text{ or } \tau^2 + h^{ij}\mu_i\mu_j \geq 1 + \epsilon/2\}, \\ \mathcal{I}_j &= \{(\rho, y, \tau, \mu) : \rho < \epsilon, 1 - \epsilon < \tau^2 + h^{ij}\mu_i\mu_j < 1 + \epsilon, \tau \in I_j\}, \end{aligned}$$

where we write

$$\{\tau \in \mathbb{R} : 1 - \epsilon < \tau^2 < 1 + \epsilon\} = \bigcup_{j=1}^N I_j \quad \text{with open connected intervals } |I_j| < \epsilon < 1.$$

Then we choose a partition of $Q_{sc}(\lambda)$,

$$Q_{sc}(\lambda) = Q_0(\lambda) + \sum_{j=1}^N Q_j(\lambda)$$

such that $Q_0(\lambda)$ and $Q_j(\lambda)$ are pseudodifferential operators with a microsupport contained in \mathcal{I}_0 and \mathcal{I}_j . In virtue of (39), the gain of such a decomposition is

- the conjugation $Q_0(\lambda) \bullet Q_0^*(\lambda)$ on $M_{b,0}$ is microlocally supported away from the characteristic variety of P_{sc} and thus a smooth function,
- the microsupport of the conjugation $Q_j(\lambda) \bullet Q_j^*(\lambda)$ on $M_{b,0}$ is away from $(L^\sharp)'$, as $|\tau - \tau'| \leq |I_j| < \epsilon$, i.e. $Q_j(\lambda) \bullet Q_j^*(\lambda)$ is microlocally supported away from the diffractive Legendrian L^\sharp .

Distance function on \mathcal{C} . Since \mathcal{C} is bounded and features only simple geodesics, the explicit distance function for two points $z = (x, y)$ and $z' = (x', y')$ sufficiently close in (\mathcal{C}, g) reads

$$\begin{aligned} d_{\mathcal{C}}(x, y, x', y') &= \sqrt{x^2 + x'^2 - 2xx' \cos d_Y(y, y')} \\ &= \hbar \sqrt{r^2 + r'^2 - 2rr' \cos d_Y(y, y')} \\ &= \hbar \sqrt{1/\rho^2 + 1/\rho'^2 - 2 \cos d_Y(y, y')/(\rho\rho')}, \end{aligned}$$

if we use $r = \rho^{-1} = x/\hbar$ and $r' = \rho'^{-1} = x'/\hbar$. In the projective coordinates ($\tilde{r} = r/r' = x/x', y, r' = x'/\hbar, y'$), the distance function lifts to,

$$d_b(\tilde{r}, y, r', y', \hbar) = \hbar r' \sqrt{\tilde{r}^2 + 1 - 2\tilde{r} \cos d_Y(y, y')},$$

near bf in the b -regime. In the projective coordinates ($\tilde{\rho} = \rho/\rho' = x'/x, y, \rho = \hbar/x, y'$), the distance function lifts to,

$$d_{sc}(\rho, \tilde{\rho}, y, y', \hbar) = \frac{\hbar}{\rho} \sqrt{1 + \tilde{\rho}^2 - 2\tilde{\rho} \cos d_Y(y, y')},$$

near mf in the sc -regime.

Conjugation by $Q_{tr}(\lambda)$. This conjugation $Q_{tr}(\lambda)dE_{\sqrt{\Delta_C}}(\lambda)Q_{tr}^*(\lambda)$ is localized in the interior of $M_{b,0}$,

$$\{r \geq \epsilon/2\} \cap \{r' \geq \epsilon/2\} \cap \{\rho \geq \epsilon/2\} \cap \{\rho' \geq \epsilon/2\} \subset M_{b,0}.$$

These restrictions guarantee $(1 + \lambda d_C(z, z'))^{-K}$ is bounded from below. In the meantime, the conjugated spectral measure $Q_{tr}(\lambda)dE_{\sqrt{\Delta_C}}(\lambda)Q_{tr}^*(\lambda)$ is a smooth Riemannian half density of the form

$$f|r^{n-1}r'^{n-1}drdydr'dy'|^{1/2}\frac{d\lambda}{\lambda} = \lambda^{n-1}f|x^{n-1}x'^{n-1}dxdydx'dy'|^{1/2}d\lambda,$$

for some $f \in C^\infty(M)$. So $Q_{tr}(\lambda)dE_{\sqrt{\Delta_C}}(\lambda)Q_{tr}^*(\lambda)$ is of the form $\lambda^{n-1}b(z, z', \lambda)$ and (41) holds.

Conjugation by $Q_b(\lambda)$. This conjugation $Q_b(\lambda)dE_{\sqrt{\Delta_C}}(\lambda)Q_b^*(\lambda)$ is localized in a neighbourhood of bf in the b -regime,

$$\{r < \epsilon\} \cap \{r' < \epsilon\} \subset M_{b,0}.$$

In this region, $(1 + \lambda d_C(z, z'))^{-K}$ is likewise bounded from below. The conjugated spectral measure $Q_b(\lambda)dE_{\sqrt{\Delta_C}}(\lambda)Q_b^*(\lambda)$ is a distribution on $M_{b,0}$ polyhomogeneous conormal to lb and rb. Hence it takes the form

$$\begin{aligned} \sum_{(z_l, k_l) \in \hat{\mathcal{E}}_b + 1} \rho_{lb}^{z_l} (\log \rho_{lb})^{k_l} a_{(z_l, k_l)}(\rho_{lb}, y, r', y', \hbar) \left| \frac{d\rho_{lb} dy dr' dy'}{\rho_{lb} r'} \right|^{1/2} \frac{d\lambda}{\lambda} & \quad \text{away from rb,} \\ \sum_{(z_r, k_r) \in \hat{\mathcal{E}}_b + 1} \rho_{rb}^{z_r} (\log \rho_{rb})^{k_r} a_{(z_r, k_r)}(r, y, \rho_{rb}, y', \hbar) \left| \frac{dr dy d\rho_{rb} dy'}{r \rho_{rb}} \right|^{1/2} \frac{d\lambda}{\lambda} & \quad \text{away from lb.} \end{aligned}$$

The blow down from $M_{b,0}$ to M pushes $Q_b(\lambda)dE_{\sqrt{\Delta_C}}(\lambda)Q_b^*(\lambda)$ forward to

$$\sum_{\substack{(z_l, k_l) \in \hat{\mathcal{E}}_b + 1 \\ (z_r, k_r) \in \hat{\mathcal{E}}_b + 1}} (x/\hbar)^{z_l} (\log(x/\hbar))^{k_l} (x'/\hbar)^{z_r} (\log(x'/\hbar))^{k_r} \frac{a_{(z_l, k_l, z_r, k_r)}(x, y, x', y', \hbar)}{x^{n/2} x'^{n/2}} |dg dg'|^{1/2} \frac{d\lambda}{\lambda}.$$

Then the fact $\inf \hat{\mathcal{E}}_b = \inf \check{\mathcal{E}}_b = n/2 - 1$ yields that $Q_b(\lambda)dE_{\sqrt{\Delta_C}}(\lambda)Q_b^*(\lambda)$ also takes the form $\lambda^{n-1}b(z, z', \lambda)$ with the amplitude estimates (41).

Conjugation by $Q_0(\lambda)$. This conjugation $Q_0(\lambda)dE_{\sqrt{\Delta_C}}(\lambda)Q_0^*(\lambda)$ is localized in a neighbourhood of mf in the sc -regime in the physical space $M_{b,0}$,

$$\{\rho < \epsilon\} \cap \{\rho' < \epsilon\} \subset M_{b,0}.$$

However, in the phase space level, it is microlocalized only in the elliptic region of P_{sc} ,

$$\{\rho < \epsilon, \rho' < \epsilon, \sigma_\partial(P_{sc}) \leq -\epsilon/2\} \cup \{\rho < \epsilon, \rho' < \epsilon, \sigma_\partial(P_{sc}) \geq \epsilon/2\} \subset {}^{b,sc}T^*M_{b,0},$$

where $\sigma_\partial(P_{sc}) = \tau^2 + h^{ij}\mu_i\mu_j - 1$. Therefore, $Q_0(\lambda)dE_{\sqrt{\Delta_C}}(\lambda)Q_0^*(\lambda)$ is a smooth section of the sc -half density on $M_{b,0}$.

We can write the conjugated spectral measure as

$$\begin{aligned} Q_0(\lambda)dE_{\sqrt{\Delta_C}}(\lambda)Q_0^*(\lambda) &= T_1 + T_2 + T_3 + T_4 \\ T_1 &= dE_{\sqrt{\Delta_C}}(\lambda) \\ T_2 &= -(\text{Id} - Q_0(\lambda))dE_{\sqrt{\Delta_C}}(\lambda) \\ T_3 &= -dE_{\sqrt{\Delta_C}}(\lambda)(\text{Id} - Q_0^*(\lambda)) \\ T_4 &= (\text{Id} - Q_0(\lambda))dE_{\sqrt{\Delta_C}}(\lambda)(\text{Id} - Q_0^*(\lambda)). \end{aligned}$$

Since $\text{WF}_{sc}(Q_0(\lambda)) \cap \text{WF}_{sc}(dE_{\sqrt{\Delta_C}}(\lambda)) = \emptyset$, the microlocalizing property (37) shows that $T_1 + T_2$ and $T_3 + T_4$ vanish to infinite order at lf and mf, whilst (38) proves that $T_1 + T_3$ and $T_2 + T_4$ vanish to infinite order at lf and mf. Therefore $Q_0(\lambda)dE_{\sqrt{\Delta_C}}(\lambda)Q_0^*(\lambda)$ vanishes to infinite order at lf, rf and mf. Then it is obvious that $Q_0(\lambda)dE_{\sqrt{\Delta_C}}(\lambda)Q_0^*(\lambda)$ takes the form $\lambda^{n-1}b(z, z', \lambda)$ with b satisfying (41).

Conjugation by $Q_j(\lambda)$. Every conjugation $Q_j(\lambda)dE_{\sqrt{\Delta_C}}(\lambda)Q_j^*(\lambda)$ is microlocalized in a neighbourhood of the interior of $L \setminus L^\sharp$ over mf \cup lf \cup rf in the *sc*-regime.

The geometric Legendrian is the geodesic flow emanating from the intersection of the diagonal conormal bundle $N^*\text{diag}_{M_{b,0}}$ with the characteristic variety of P_{sc} over mf,

$$\partial_0 L = \{(\tilde{\rho}, y, \rho', y, \hbar, \tau, \mu, -\tau, -\mu, 0) : \tilde{\rho} = 1, \rho' = 0, \tau^2 + \hbar^{ij}\mu_i\mu_j = 1\}.$$

As a boundary of L , $\partial_0 L$ doesn't project diffeomorphically to the base. Since the geodesics in \mathcal{C} are all simple i.e. the exponential map is a diffeomorphism, $L \setminus \partial_0 L$ projects diffeomorphically to the base. Furthermore, the proof of [15, Proposition 6.2, Lemma 6.4, Lemma 6.5] and [21, Proposition 2.6] can be used verbatim to elucidate the Legendrian structure.

Lemma 28. *The Legendrian pair $(N^*\partial_{sc}\text{diag}_{M_{b,0}}, L)$ has a clean intersection $\partial_0 L$. The projection $\pi : L \rightarrow \text{mf}$ satisfies that*

- π is a diffeomorphism on $L \setminus \partial_0 L$,
- $\det d\pi$ vanishes to order $n - 1$ at $\partial_0 L$.

It follows that the geometric Legendrian L is furnished with local coordinates,

- $(\tilde{\rho}, y, y', \hbar)$ away from $\partial_0 L$,
- $(y, \mathbf{w}_1, \mathbf{k}_2, \dots, \mathbf{k}_n, \hbar)$, in a neighbourhood of $\partial_0 L$ with $\partial_0 L$ removed, where $\mathbf{w} = (\tilde{\rho} - 1, y - y')$ and \mathbf{k} is the dual coordinates on the fibre of ${}^{b,sc}T_{\text{mf}}^*M_{b,0}$. Here we need relabel the components of \mathbf{w} and \mathbf{k} variables to ensure $d\mathbf{k}_1$ vanishes at a point in $\partial_0 L$, as $d\pi$ is not of full rank.

Consequently, the geometric Legendrian is parametrized,

- away from $\partial_0 L$, by the distance function $d_{sc}(\rho, \tilde{\rho}, y, y', \hbar)$
- in a neighbourhood of $\partial_0 L$ with $\partial_0 L$ removed, by a function $\Phi(y, \mathbf{w}, \mathbf{v})$ with $\mathbf{v} = (\mathbf{v}_2, \dots, \mathbf{v}_n) \in \mathbb{R}^{n-1}$ such that

$$(42) \quad d_{v_j}\Phi(y, \mathbf{w}, \mathbf{v}) = \mathbf{w}_j + O(\mathbf{w}_1),$$

$$(43) \quad \Phi(y, \mathbf{w}, \mathbf{v}) = \sum_{j=2}^n \mathbf{v}_j d_{v_j}\Phi(y, \mathbf{w}, \mathbf{v}) + O(\mathbf{w}_1),$$

$$(44) \quad d_{v_j v_k}^2 \Phi(y, \mathbf{w}, \mathbf{v}) = \mathbf{w}_1 A_{jk}(y, \mathbf{w}, \mathbf{v}), \quad \text{with a non-degenerate matrix } (A_{jk})_{jk},$$

$$(45) \quad \Phi(y, \mathbf{w}, \mathbf{v})/\rho = \pm \lambda d_{sc}(\rho, \tilde{\rho}, y, y', \hbar), \quad \text{when } d_{\mathbf{v}}\Phi = 0.$$

Finally, the conjugated spectral measure $Q_j(\lambda)dE_{\sqrt{\Delta_C}}(\lambda)Q_j^*(\lambda)$ is a Legendrian distribution lying in $I^{-1/2; ((n-1-\mathbf{e})/2, (n-1-\mathbf{e})/2, 0)}(M_{b,0}; L \setminus L^\sharp; {}^{b,sc}\Omega^{1/2})$ tensored with $|d\lambda/\lambda|^{1/2}$ and has the microlocal expression, away from rf,

- when the conjugation $Q_j(\lambda) \bullet Q_j^*(\lambda)$ is microlocally supported away from a neighbourhood of $\partial_0 L$,

$$(46) \quad \rho'^{-1/2+n/2} \tilde{\rho}^{(n-1-\varepsilon_l)/2} e^{i\pm \lambda d_{sc}(\rho, \tilde{\rho}, y, y', \hbar)} \tilde{a}(\tilde{\rho}, y, \rho', y', \hbar) \frac{|d\rho d\rho' dy dy'|^{1/2}}{\rho^{(n+1)/2} \rho'^{(n+1)/2}} \frac{d\lambda}{\lambda},$$

where $\tilde{a}(\tilde{\rho}, y, \rho', y', \hbar)$ is a compactly supported smooth function in the *sc*-regime,

- when the conjugation $Q_j(\lambda) \bullet Q_j^*(\lambda)$ is microlocally supported in a neighbourhood of $\partial_0 L$ with $\partial_0 L$ removed,

$$(47) \quad \rho'^{-1/2-(n-1)/2+n/2} \int_{\mathbb{R}^{n-1}} e^{i\Phi(y, \mathbf{w}, \mathbf{v})/\rho} \tilde{a}(\rho, y, \mathbf{w}, \mathbf{v}, \hbar) d\mathbf{v} \frac{|d\rho d\rho' dy dy'|^{1/2}}{\rho^{(n+1)/2} \rho'^{(n+1)/2}} \frac{d\lambda}{\lambda},$$

where $\tilde{a}(\rho, y, \mathbf{w}, \mathbf{v}, \hbar)$ is a compactly supported smooth function in the sc -regime.

Now we consider the estimates for the conjugated spectral measure of type (46). Such an expression we choose implies that we live in the sc -regime near $\text{mf} \cap \text{lf}$, i.e. $\tilde{\rho} = \rho/\rho' \rightarrow 0$. The local expression (46) further reduces to

$$\tilde{\rho}^{-\varepsilon_l/2} \rho^{(n-1)/2} e^{i\pm\lambda d_{sc}(\rho, \tilde{\rho}, y, y', \hbar)} \tilde{a}(\tilde{\rho}, y, \rho', y', \hbar) \left| \frac{d\rho d\rho' dy dy'}{\rho^{n+1} \rho'^{n+1}} \right|^{1/2} \frac{d\lambda}{\lambda}.$$

Using the local expression of the distance function d_{sc} , we see that the factor $\rho^{(n-1)/2}$ is bounded from above by

$$\rho^{(n-1)/2} \lesssim (1 + 1/\rho)^{-(n-1)/2} \lesssim (1 + \lambda d_{sc}(\rho, \tilde{\rho}, y, y', \hbar))^{-(n-1)/2}.$$

We thus take $a_{\pm} = \rho^{(n-1)/2} \tilde{a}$, which satisfies the amplitude estimates (40).

We now blow down $M_{b,0}$ to $M = \{(z, z', \hbar)\}$ and change \hbar to λ^{-1} as we need. It follows that (46) is pushed forward to,

$$\lambda^{n-1} e^{i\pm\lambda d_C(z, z')} x'^{-\varepsilon_l/2} \tilde{a}(z, z', \lambda) x^{(n-1)/2} x'^{(n-1)/2} |dx dx' dy dy'|^{1/2} d\lambda.$$

The factor $x^{(n-1)/2} x'^{(n-1)/2} |dx dx' dy dy'|^{1/2}$ gives a Riemannian half density $|dg dg'|^{1/2}$, and the stability condition (1) implies $x'^{-\varepsilon_l/2} \leq \lambda^{\mathbf{e}/2}$ in the sc -regime. It follows that we get an oscillatory function $\lambda^{n-1+\mathbf{e}/2} e^{\pm i\lambda d_C(z, z')} a_{\pm}$ up to a smooth function.

The other type (47) also takes the form $\lambda^{n-1} e^{\pm i\lambda d_C(z, z')} a_{\pm}$ but it needs a bit more work due to the complication of the Legendrian structure at the intersection with the diagonal. Recall that we now live in a small neighbourhood of $\partial_0 L$. In a small neighbourhood of $\{\mathbf{w} = 0\}$, we have $|\mathbf{w}|/\rho \sim \lambda d_C(z, z')$. In fact, it is easy to deduce that

$$\begin{aligned} \frac{|\mathbf{w}|}{\rho} &= \frac{1}{\rho} \sqrt{\left(\frac{\rho}{\rho'} - 1\right)^2 + (y - y')^2} \\ &= \sqrt{\frac{1}{\rho^2} + \frac{1}{\rho'^2} - \frac{2}{\rho\rho'} \left(1 - \frac{\rho'}{2\rho} (y - y')^2\right)} \\ &\sim \sqrt{\frac{1}{\rho^2} + \frac{1}{\rho'^2} - \frac{2}{\rho\rho'} \cos d_Y(y, y')} \\ &= \lambda d_C(z, z') \end{aligned}$$

If we blow down to M again,¹⁸ the distribution (47) is pushed forward to

$$\lambda^{n-1} \left(\int_{\mathbb{R}^{n-1}} e^{i\Phi(y, \mathbf{w}, \mathbf{v})/\rho} \tilde{a}(\rho, y, \mathbf{w}, \mathbf{v}, \lambda) d\mathbf{v} \right) x^{(n-1)/2} x'^{(n-1)/2} |dx dx' dy dy'|^{1/2} d\lambda.$$

We complete the proof of the proposition by doing estimates for the integral in the parentheses in three cases.

¹⁸Even though we no longer live in the stretched spaces, we still adopt the notations $\rho = 1/(\lambda x)$ and $\mathbf{w} = (x'/x - 1, y - y')$ for the sake of convenience.

- (a) $|\mathbf{w}| \lesssim \rho$. This condition implies $\lambda d_{\mathcal{C}}(z, z') \lesssim 1$, i.e. $(1 + \lambda d_{\mathcal{C}}(z, z'))^{-K}$ is bounded from below by a positive constant. Then the amplitude estimate for b is clearly satisfied when $k = 0$. For $k > 0$, it suffices to show

$$(\lambda \partial_{\lambda})^k \left(\int_{\mathbb{R}^{n-1}} e^{i\Phi(y, \mathbf{w}, \mathbf{v})/\rho} \tilde{a}(\rho, y, \mathbf{w}, \mathbf{v}, \lambda) dv \right) \lesssim 1.$$

If the derivative $\lambda \partial_{\lambda}$ hits the amplitude, it affects nothing. If it acts on the oscillatory term once, by (43) it will create a factor of

$$\Phi/\rho = \mathbf{v} \cdot d_{\mathbf{v}} \Phi/\rho + O(\mathbf{w}_1/\rho).$$

Noting

$$\frac{\mathbf{v} \cdot d_{\mathbf{v}} \Phi}{\rho} e^{i\Phi/\rho} = -i \mathbf{v} \cdot d_{\mathbf{v}} e^{i\Phi/\rho},$$

we perform integration by parts and get a finite integral.

- (b) $|\mathbf{w}| \gtrsim \rho$ and $|\mathbf{w}| \gtrsim |\mathbf{w}_1|$. In light of (42), we have $d_{\mathbf{v}_i} \Phi \neq 0$. Choose i such that $|d_{\mathbf{v}_i} \Phi| \gtrsim |\mathbf{w}|$. When $k = 0$, by using

$$e^{i\Phi/\rho} = \left(\frac{\rho d_{\mathbf{v}_j}}{i d_{\mathbf{v}_j} \Phi} \right)^k e^{i\Phi/\rho},$$

we make integration by parts in \mathbf{v} and obtain a finite integration, multiplied by the factor $(\rho/|\mathbf{w}|)^K$ for any $K \in \mathbb{Z}_+$. This is bounded from above by $(1 + |\mathbf{w}|/\rho)^{-K}$. If we blow down to M , this quantity is equivalent to $(1 + \lambda d_{\mathcal{C}}(z, z'))^{-K}$. When $k > 0$, the derivatives are treated similarly to Case (a).

- (c) $|\mathbf{w}| \sim |\mathbf{w}_1|$ and $|\mathbf{w}_1| \geq \rho$. It suffices to prove for $k \in \mathbb{N}$,

$$(\lambda \partial_{\lambda})^k \left(\int_{\mathbb{R}^{n-1}} e^{i\omega \tilde{\Phi}} a(z, z', \mathbf{v}, \lambda) dv \right) \lesssim \omega^{-(n-1)/2},$$

where $\omega = \lambda x |\mathbf{w}_1|$ and $\tilde{\Phi} = |\mathbf{w}_1|^{-1} (\Phi(x'/x, y, y', \mathbf{v}) \mp x^{-1} d_{\mathcal{C}}(z, z'))$. Under (45) and (44), this oscillatory integral estimates were proved by a finer integration by parts argument. See [21, 4-3].

□

13.4. Local Strichartz estimates on \mathcal{C} . The exponent pair (q, r) is said to be sharp σ -admissible if it satisfies

$$q, r \geq 2, \quad (q, r, \sigma) \neq (2, \infty, 1), \quad \frac{1}{q} + \frac{\sigma}{r} = \frac{\sigma}{2}.$$

In particular, (q, r) is said to be Schrödinger admissible if it is sharp $n/2$ -admissible as mentioned in the introduction.

Using Proposition 27, we can establish Strichartz estimates on \mathcal{C} ,

$$(48) \quad \|e^{it\Delta} f\|_{L^q([0,1]; L^{\tilde{r}}(\mathcal{C}))} \lesssim \|f\|_{L^2(\mathcal{C})},$$

for any sharp $(n/2 + \mathbf{e}/4)$ -admissible pair (q, \tilde{r}) .

This can be done by showing the following dispersive estimates and energy estimates for conjugated Schrödinger propagators from corresponding conjugated spectral measure,

Proposition 29. *Suppose $\{Q_{\alpha}(\lambda)\}_{\alpha}$ is a finite partition of identity as in Proposition 27 and write*

$$U_{\alpha}(t) = \int_{\lambda > 1} e^{it\lambda^2} Q_{\alpha}(\lambda) dE_{\sqrt{\Delta_{\mathcal{C}}}}(\lambda).$$

Then the Schrödinger propagator conjugated by $Q_\alpha(\lambda)$ features the factorization,

$$(49) \quad \int_{\lambda>1} e^{it\lambda^2} Q_\alpha(\lambda) dE_{\sqrt{\Delta_C}}(\lambda) Q_\alpha^*(\lambda) = U_\alpha(t) U_\alpha^*(t),$$

and uniformly obeys energy estimates

$$(50) \quad \|U_\alpha(t)f\|_{L^2(\mathcal{C})} \lesssim \|f\|_{L^2(\mathcal{C})},$$

as well as dispersive estimates for small times $0 < t < 1$,

$$(51) \quad \|U_\alpha(t_1)U_\alpha^*(t_2)f\|_{L^\infty(\mathcal{C})} \lesssim |t_1 - t_2|^{-n/2-\mathbf{e}/4} \|f\|_{L^1(\mathcal{C})}.$$

Given Proposition 27, the proof of (49), (50) and (51) has little to do with the conic geometry but is purely an analysis argument. We can employ the proof for [21, Proposition 5.1, Lemma 5.3, Proposition 6.1] verbatim.

Given (50) and (51) as well as the energy and dispersive estimates for $e_{\text{low}}^{it\Delta_C}$, Keel-Tao endpoint method [25, Theorem 1.2] leads to that

$$\begin{aligned} \|U_\alpha(t)f\|_{L^q([0,1];L^{\tilde{r}}(\mathcal{C}))} &\lesssim \|f\|_{L^2(\mathcal{C})} \quad \text{for each } \alpha, \\ \|e_{\text{low}}^{it\Delta_C} f\|_{L^q([0,1];L^{\tilde{r}}(\mathcal{C}))} &\lesssim \|f\|_{L^2(\mathcal{C})}. \end{aligned}$$

Since $\{Q_\alpha(\lambda)\}_\alpha$ is a finite partition of identity, (48) results from

$$e^{it\Delta_C} = e_{\text{high}}^{it\Delta_C} + e_{\text{low}}^{it\Delta_C} \quad \text{and} \quad e_{\text{high}}^{it\Delta_C} = \sum_{\alpha} U_\alpha(t).$$

Let $W^{k,p}(\mathcal{C})$ be the usual p -Sobolev space on \mathcal{C} , that is

$$W^{k,p}(\mathcal{C}) = \left\{ f \in L^p(\mathcal{C}) : \sum_{l=0}^{[k]} \|\partial^l f\|_{L^p(\mathcal{C})} + \left(\int_{\mathcal{C}^2} \frac{|f(z) - f(z')|^p}{|z - z'|^{[k]p+n}} dg(z) dg(z') \right)^{1/p} < \infty \right\}.$$

The classical Sobolev embedding theorem can be extended to \mathcal{C} (See [1, Theorem 5.4]), that is

$$W^{k_1,p_1}(\mathcal{C}) \subset W^{k_2,p_2}(\mathcal{C}), \quad \text{for} \quad \frac{1}{p_1} - \frac{k_1}{n} = \frac{1}{p_2} - \frac{k_2}{n}.$$

This allows us to rewrite (48) as Schrödinger admissible Strichartz estimates with a loss. Suppose (q, r) is Schrödinger admissible. It follows that

$$\frac{n/2 + \mathbf{e}/4}{\tilde{r}} - \frac{n/2}{r} = \frac{\mathbf{e}}{8}.$$

Then it is immediate from Sobolev embeddings and $(n/2 + \mathbf{e}/4)$ -admissibility that

$$(52) \quad \|e^{it\Delta} f\|_{L^q([0,1];W^{-\mathbf{k},r}(\mathcal{C}))} \lesssim \|f\|_{L^2(\mathcal{C})},$$

with \mathbf{k} defined in (9).

13.5. Global Strichartz estimates on \mathcal{X} .

Proof of Theorem 2. To employ local estimates (48), we choose a finite partition of unity $1 = \sum_{i=0}^m \chi_i$ on \mathcal{X} such that

- each χ_i for $i = 1, \dots, m$, is a smooth function compactly supported within \mathcal{C}_i and equal to 1 on \mathcal{K}_i ,
- χ_0 is a smooth function supported within $\mathcal{X} \setminus \bigcup_{i=1}^m \mathcal{K}_i$ and equal to 1 on $\mathcal{X} \setminus \bigcup_{i=1}^m \mathcal{C}_i$.

Now we denote $u_i = \chi_i u$ for $i = 0, 1, \dots, m$. Applying the Schrödinger operator to each u_i gives the Cauchy problem for inhomogeneous Schrödinger equations,

$$\begin{cases} i\partial_t u_i(t, z) + \Delta_{\mathcal{X}} u_i(t, z) = v_i(t, z) \\ u_i(0, z) = \chi_i f(z) \end{cases},$$

where $v_i(t, z) = [\Delta_{\mathcal{X}}, \chi_i]u(t, z)$.

We write $u_i = u'_i + u''_i$, such that

- u'_i is the solution to

$$\begin{cases} i\partial_t u'_i(t, z) + \Delta_{\mathcal{X}} u'_i(t, z) = 0 \\ u'_i(0, z) = \chi_i f(z) \end{cases};$$

- u''_i is the solution to

$$\begin{cases} i\partial_t u''_i(t, z) + \Delta_{\mathcal{X}} u''_i(t, z) = v_i(t, z) \\ u''_i(0, z) = 0 \end{cases}.$$

By (52) and Strichartz estimates on \mathbb{R}^n , we know,

$$\|e^{it\Delta_{\mathcal{X}}} u'_i\|_{L^q([0,1]; W^{-k,r}(\mathcal{X}))} \lesssim \|\chi_i f\|_{L^2(\mathcal{X})},$$

where (q, r) is Schrödinger admissible.

Then it remains to deal with u''_i . Duhamel's principle yields that

$$u''_i(t, z) = \int_0^t e^{-i(t-s)\Delta_{\mathcal{X}}} v_i(s, z) ds = e^{-i\Delta_i} \int_0^t e^{is\Delta_i} v_i(s, z) ds,$$

where $\Delta_0 = \Delta_{\mathbb{R}^n}$ and $\Delta_i = \Delta_{C_i}$ for $i = 1, \dots, m$. By (48) and Strichartz estimates on \mathbb{R}^n , it suffices to show

$$\left\| \int_0^t e^{is\Delta_i} v_i(s, z) ds \right\|_{L^2(\mathcal{X})} \lesssim \|f\|_{L^2(\mathcal{X})}.$$

Christ-Kiselev's lemma [9, Theorem 1.2] further reduces this inequality to

$$\left\| \int_0^1 e^{is\Delta_i} v_i(s, z) ds \right\|_{L^2(\mathcal{X})} \lesssim \|f\|_{L^2(\mathcal{X})}.$$

Now we need the local smoothing estimates (12) and dual local smoothing estimates

$$\left\| \int_0^1 e^{is\Delta_{\mathcal{X}}} \chi F(s, z) ds \right\|_{L^2(\mathcal{X})} \lesssim \|F\|_{L^2(\mathbb{R}, \mathcal{D}^{-1/2}(\mathcal{X}))},$$

as $[\Delta_{\mathcal{X}}, \chi_i]$ is a differential operator of order 1 supported within $\bigcup_{j=1}^m \mathcal{C}_j \setminus \mathcal{K}_j^{\circ}$. Consequently, we obtain

$$\left\| \int_{\mathbb{R}} e^{is\Delta_i} v_i(s, z) ds \right\|_{L^2(\mathcal{X})} \lesssim \|v_i\|_{L^2(\mathbb{R}, \mathcal{D}^{-1/2}(\mathcal{X}))}.$$

In addition, v_i satisfies

$$\|v_i\|_{L^2(\mathbb{R}, \mathcal{D}^{-1/2}(\mathcal{X}))} \lesssim \|u_i\|_{L^2(\mathbb{R}, \mathcal{D}^{1/2}(\mathcal{X}))}.$$

Therefore the proof of the theorem will be finished by a final application of local smoothing estimates to u_i

$$\|u_i\|_{L^2(\mathbb{R}, \mathcal{D}^{1/2}(\mathcal{X}))} \lesssim \|f\|_{L^2(\mathcal{X})}.$$

□

REFERENCES

- [1] R. A. Adams. *Sobolev spaces*. Academic Press [A subsidiary of Harcourt Brace Jovanovich, Publishers], New York-London, 1975. Pure and Applied Mathematics, Vol. 65.
- [2] P. Albin. *Analysis on non-compact manifolds*. Lecture Notes for MIT Course 18.158 in Spring 2008, available at <https://faculty.math.illinois.edu/~palbin/18158/18158May26.pdf>.
- [3] D. Baskin, J. L. Marzuola, and J. Wunsch. Strichartz estimates on exterior polygonal domains. In *Geometric and spectral analysis*, volume 630 of *Contemp. Math.*, pages 291–306. Amer. Math. Soc., Providence, RI, 2014.
- [4] D. Baskin and J. Wunsch. Resolvent estimates and local decay of waves on conic manifolds. *J. Differential Geom.*, 95(2):183–214, 2013.
- [5] M. D. Blair, G. A. Ford, S. Herr, and J. L. Marzuola. Strichartz estimates for the Schrödinger equation on polygonal domains. *J. Geom. Anal.*, 22(2):339–351, 2012.
- [6] N. Burq. Décroissance de l’énergie locale de l’équation des ondes pour le problème extérieur et absence de résonance au voisinage du réel. *Acta Math.*, 180(1):1–29, 1998.
- [7] N. Burq. Smoothing effect for Schrödinger boundary value problems. *Duke Math. J.*, 123(2):403–427, 2004.
- [8] N. Burq, P. Gérard, and N. Tzvetkov. On nonlinear Schrödinger equations in exterior domains. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 21(3):295–318, 2004.
- [9] M. Christ and A. Kiselev. Maximal functions associated to filtrations. *J. Funct. Anal.*, 179(2):409–425, 2001.
- [10] J. J. Duistermaat and L. Hörmander. Fourier integral operators. II. *Acta Math.*, 128(3-4):183–269, 1972.
- [11] G. A. Ford. The fundamental solution and Strichartz estimates for the Schrödinger equation on flat Euclidean cones. *Comm. Math. Phys.*, 299(2):447–467, 2010.
- [12] C. Guillarmou and A. Hassell. Resolvent at low energy and Riesz transform for Schrödinger operators on asymptotically conic manifolds. I. *Math. Ann.*, 341(4):859–896, 2008.
- [13] C. Guillarmou and A. Hassell. Resolvent at low energy and Riesz transform for Schrödinger operators on asymptotically conic manifolds. II. *Ann. Inst. Fourier (Grenoble)*, 59(4):1553–1610, 2009.
- [14] C. Guillarmou, A. Hassell, and A. Sikora. Resolvent at low energy III: The spectral measure. *Trans. Amer. Math. Soc.*, 365(11):6103–6148, 2013.
- [15] C. Guillarmou, A. Hassell, and A. Sikora. Restriction and spectral multiplier theorems on asymptotically conic manifolds. *Anal. PDE*, 6(4):893–950, 2013.
- [16] A. Hassell, R. Mazzeo, and R. B. Melrose. Analytic surgery and the accumulation of eigenvalues. *Comm. Anal. Geom.*, 3(1-2):115–222, 1995.
- [17] A. Hassell and A. Vasy. The spectral projections and the resolvent for scattering metrics. *J. Anal. Math.*, 79:241–298, 1999.
- [18] A. Hassell and A. Vasy. The resolvent for Laplace-type operators on asymptotically conic spaces. *Ann. Inst. Fourier (Grenoble)*, 51(5):1299–1346, 2001.
- [19] A. Hassell and J. Wunsch. The Schrödinger propagator for scattering metrics. *Ann. of Math. (2)*, 162(1):487–523, 2005.
- [20] A. Hassell and J. Wunsch. The semiclassical resolvent and the propagator for non-trapping scattering metrics. *Adv. Math.*, 217(2):586–682, 2008.
- [21] A. Hassell and J. Zhang. Global-in-time Strichartz estimates on nontrapping, asymptotically conic manifolds. *Anal. PDE*, 9(1):151–192, 2016.
- [22] P. Hintz. Resolvents and complex powers of semiclassical cone operators. *arXiv e-prints*, page arXiv:2010.01593, Oct. 2020.
- [23] P. Hintz. Semiclassical propagation through cone points. *arXiv e-prints*, page arXiv:2101.01008, Jan. 2021.
- [24] L. Hörmander. Fourier integral operators. I. *Acta Math.*, 127(1-2):79–183, 1971.
- [25] M. Keel and T. Tao. Endpoint Strichartz estimates. *Amer. J. Math.*, 120(5):955–980, 1998.
- [26] P. Loya. On the resolvent of differential operators on conic manifolds. *Comm. Anal. Geom.*, 10(5):877–934, 2002.
- [27] R. B. Melrose. Calculus of conormal distributions on manifolds with corners. *Internat. Math. Res. Notices*, (3):51–61, 1992.
- [28] R. B. Melrose. *The Atiyah-Patodi-Singer index theorem*, volume 4 of *Research Notes in Mathematics*. A K Peters, Ltd., Wellesley, MA, 1993.

- [29] R. B. Melrose. Spectral and scattering theory for the Laplacian on asymptotically Euclidian spaces. In *Spectral and scattering theory (Sanda, 1992)*, volume 161 of *Lecture Notes in Pure and Appl. Math.*, pages 85–130. Dekker, New York, 1994.
- [30] R. B. Melrose and G. A. Uhlmann. Lagrangian intersection and the cauchy problem. *Comm. Pure Appl. Math.*, 32(4):483–519, 1979.
- [31] R. B. Melrose and J. Wunsch. Propagation of singularities for the wave equation on conic manifolds. *Invent. Math.*, 156(2):235–299, 2004.
- [32] R. B. Melrose and M. Zworski. Scattering metrics and geodesic flow at infinity. *Invent. Math.*, 124(1-3):389–436, 1996.
- [33] E. Schrohe and J. Seiler. The resolvent of closed extensions of cone differential operators. *Canad. J. Math.*, 57(4):771–811, 2005.
- [34] E. Schrohe and J. Seiler. Bounded H_∞ -calculus for cone differential operators. *J. Evol. Equ.*, 18(3):1395–1425, 2018.
- [35] B.-W. Schulze. *Pseudo-differential operators on manifolds with singularities*, volume 24 of *Studies in Mathematics and its Applications*. North-Holland Publishing Co., Amsterdam, 1991.
- [36] B.-W. Schulze. *Pseudo-differential boundary value problems, conical singularities, and asymptotics*, volume 4 of *Mathematical Topics*. Akademie Verlag, Berlin, 1994.
- [37] B.-W. Schulze. *Boundary value problems and singular pseudo-differential operators*. Pure and Applied Mathematics (New York). John Wiley & Sons, Ltd., Chichester, 1998.
- [38] G. Staffilani and D. Tataru. Strichartz estimates for a Schrödinger operator with nonsmooth coefficients. *Comm. Partial Differential Equations*, 27(7-8):1337–1372, 2002.
- [39] M. E. Taylor. *Partial differential equations. II*, volume 116 of *Applied Mathematical Sciences*. Springer-Verlag, New York, 1996. Qualitative studies of linear equations.
- [40] J. Wunsch. Propagation of singularities and growth for Schrödinger operators. *Duke Math. J.*, 98(1):137–186, 1999.
- [41] J. Zhang and J. Zheng. Global-in-time Strichartz estimates and cubic Schrodinger equation on metric cone. *arXiv e-prints*, page arXiv:1702.05813, Feb. 2017.

SHANGHAI CENTER FOR MATHEMATICAL SCIENCES, FUDAN UNIVERSITY, SHANGHAI 200438, CHINA.
E-mail address: xi_chen@fudan.edu.cn

DEPARTMENT OF PURE MATHEMATICS AND MATHEMATICAL STATISTICS, UNIVERSITY OF CAMBRIDGE,
 CAMBRIDGE CB3 0WB, UK. *E-mail address:* xi.chen@dpms.cam.ac.uk