

Nonequilibrium induced by reservoirs: Physico-mathematical model and numerical tests

Rupert Klein,^{1, a)} Roya Ebrahimi Viand,¹ Felix Höfling,¹ and Luigi Delle Site^{1, b)}

Freie Universität Berlin, Institute of Mathematics, Arnimallee 6, 14195 Berlin, Germany

A recently proposed mathematical model for open many particle systems (Delle Site and Klein, J.Math.Phys. **61**, 083102 (2020)) is extended to the case of two reservoirs with different thermodynamic state points. It is assumed that (i) particles interact via short-range pair interactions and that (ii) the states of reservoir particles found within the interaction range of the open system are statistically independent of the open system state. Under these conditions, the model leads to a description of an open system out of equilibrium in which the action of the reservoirs enters as a linear sum into the Liouville-type evolution equations for the open system's statistics. This model is compared with alternative open system models that include the linearity of the action of the reservoirs as an *a priori* assumption. The possibility that nonlinear effects arise within the present model under less restrictive conditions than (i) and (ii) above is discussed together with a related numerical test. Even in such a limiting case the approximation of a linear combination of actions turns out to yield a description of the open system with sufficient accuracy for practical purposes. The model presented can be used as a guiding protocol in the development of simulation schemes for open systems out of equilibrium, thus the results of this work are not only of conceptual but also of practical interest.

^{a)}Electronic mail: rupert.klein@math.fu-berlin.de

^{b)}Electronic mail: luigi.dellesite@fu-berlin.de

I. INTRODUCTION

Theory and modeling of open systems are becoming increasingly prominent since they allow to focus on regions of relevance (where processes of interest are taking place) embedded in large complex domains while the complexity of the outside is reduced to few macroscopic variables¹. In particular, open molecular systems are of relevance because of their occurrence in many fields of current cutting edge technology, thus they require well founded numerical algorithms for efficient and accurate simulations². In this perspective physico-mathematical models of open systems represent a guideline protocol for the development of related simulation algorithms.

Two of the authors³, with such motivation, have recently derived a model of open systems following an analytic procedure based on a microscopic approach. Starting from a large system (Universe) the degrees of freedom of the particles of the reservoir, under suitable physical conditions, are analytically integrated out and an equation for the statistical evolution of the open system is derived. The original derivation considers an open system embedded in a single homogeneous reservoir. Recent numerical experiments suggest, however, that it would be interesting to treat the case of open system interfaced with two, or more, distinct and disjoint reservoirs⁴. In fact a system in contact with, for example, two reservoirs calibrated at different thermodynamic state points is subject to a situation of nonequilibrium. In fact due to the difference in temperature and/or density of the two reservoirs a thermal or mass gradient characterizes the open system. Such situations, in turn, are of special interest in the field of molecular simulation (see Ref. 4 and references therein).

For this reason we analyze in the present work the consequences of applying the strategy of Ref. 3 when two (or more) disjoint reservoirs act concurrently on an open system. The results show that the combined actions of the reservoirs in the equation of statistical evolution of the open system enters as the sum of action of each single reservoir. This result essentially summarizes the fact that, under the hypotheses of two-body short range interactions between the particles and of statistical independence of the states of reservoir particle residing close to the open system boundary, the coupling between the open system and each reservoir occurs only at the interface regions and thus the contribution of each reservoir is reduced to a surface

integral at the interface region. Other models of open systems assume instead, *a priori*, the linear contribution of the reservoirs in the equations of evolution. As an example, we compare our results with the models of Bergmann and Lebowitz^{5,6} and the model employed by Gay-Balmaz and Yoshimura in Ref. 7. Interestingly, in such models, due to the drastic *a priori* simplification of the reservoirs, whose microscopic origin is completely neglected, the possibility of considering nonlinear effects in the coupling, and in turn understanding their physical consequences is ruled out.

Instead generalizations to nonlinear and memory effects are within the scope of the present modelling approach when less restrictive conditions on the range of particle-particle interactions and on the statistics of reservoir states close to the open system boundary are adopted. Thus it is known rigorously from Ref. 8 that, just as a consequence of multidimensional wave propagation in the reservoir, a non-reflecting acoustic boundary condition *must* include memory effects. Moreover, when the single- and two-particle statistics involving reservoir particles close to the system boundary are allowed to depend on the general open system state, then nonlinear effects will arise in addition as discussed in section IV below. The inclusion of nonlinear and memory effects of the reservoirs are only discussed qualitatively here, while a detailed analysis is left for future work.

Here we focus instead on the consequences of our results for the corresponding numerical approach. To this end we test the quality of a numerical approach based on the AdResS technology^{1,9} which counts solely on a linear contribution of the reservoirs. The test consists in simulating a Lennard-Jones liquid in an open domain set up such that there is a feedback of the open system onto a sizable part of two attached reservoirs. We compare the results of our model with the results of a reference simulation of the Universe in which all particles are explicitly treated with all their degrees of freedom, but are thermalized at different temperatures in subregions equivalent to the reservoir domains of our model. With such a comparison we conclude about the effective/numerical validity of the linear approximation. Surprisingly, for a Lennard-Jones liquid at thermodynamic and gradient conditions common to a large variety of chemical-physics situations, it is shown that the linear hypothesis holds and nonlinear effects are numerically negligible. This is an encouraging result in the perspective of developing accurate and efficient simulation algorithms.




Figure 1. The partitioning of the "Universe" into the open system Ω and the reservoir $U \setminus \Omega \equiv \Omega_c$.

II. LIOUVILLE-TYPE EQUATIONS FOR AN OPEN SYSTEM

In this section we summarize the key steps of the modeling procedure adopted in Ref. 3 for an open system embedded in a single uniform reservoir which are relevant for the extension of the model to several independently acting reservoirs.

A. Topological definition of an open system

Let us consider the open system schematically illustrated in Fig.1, the total system, here called "Universe", is characterized by N (fixed) number of particles. Its spatial domain of definition for each single particle is U . Ω is the domain of definition of the open system that may contain any number $n \in \{0, \dots, N\}$ of particles, while the region of the reservoir corresponds to $U \setminus \Omega \equiv \Omega_c$ with $N - n$ particles. The phase space of the N particles in U is $S^N = \mathbb{R}^{3N} \times U^N$ (momentum space in \mathbb{R}^{3N} and position space U^N), the phase space of the open system is $S^n = \mathbb{R}^{3n} \times \Omega^n$, and the phase space of the reservoir is $S_c^{(N-n)} = \mathbb{R}^{3(N-n)} \times \Omega_c^{(N-n)}$

B. Relevant quantities that characterize U

The Universe is characterized by the Hamiltonian

$$H_N(\mathbf{p}^N, \mathbf{q}^N) = E_{\text{kin}}(\mathbf{p}^N) + V_{\text{tot}}(\mathbf{q}^N) \equiv \sum_{i=1}^N \frac{(\vec{p}_i)^2}{2M} + \frac{1}{2} \sum_{\substack{i,j=1 \\ j \neq i}}^N V(\vec{q}_j - \vec{q}_i) \quad (1)$$

where

$$(\mathbf{p}^N, \mathbf{q}^N) = (\vec{p}_1, \dots, \vec{p}_N, \vec{q}_1, \dots, \vec{q}_N) \quad (2)$$

and (\vec{p}_i, \vec{q}_i) are the momentum and position of the i th particle, respectively, M is the particle mass, assumed to be the same for all particles here, and V is the two-body interaction potential (as typical for, e.g., Molecular Dynamics). The statistical mechanics description of the system in phase space is achieved through its probability density defined as

$$\begin{aligned} F_N : \mathbb{R}^+ \times (U \times \mathbb{R}^3)^N &\rightarrow \mathbb{R} \\ (t, \mathbf{X}^N) &\mapsto F_N(t, \mathbf{X}^N) \end{aligned} \quad (3)$$

with normalization $\int_{S^N} F_N d\mathbf{X}^N = 1$. The probability density of the Universe is subject to the phase space density transport equation (Liouville equation)

$$\frac{\partial F_N}{\partial t} = \sum_{i=1}^N [\nabla_{\vec{q}_i} \cdot (\vec{v}_i F_N) + \nabla_{\vec{p}_i} \cdot (-\nabla_{q_i} V_{\text{tot}}(\mathbf{q}^N) F_N)] \quad (4)$$

where $\vec{v}_i = \vec{p}_i/M$ is the i th' particle velocity.

C. Relevant quantities that characterize Ω

The open system Ω is characterized by the Hamiltonian

$$H_n = \sum_{i=1}^n \frac{(\vec{p}_i)^2}{2M} + \frac{1}{2} \sum_{\substack{i,j=1 \\ j \neq i}}^n V(\vec{q}_j - \vec{q}_i) \quad (\vec{q}_i, \vec{q}_j \in \Omega). \quad (5)$$

The statistical mechanics description of the open system in phase space is given by the collection of all its n -particle probability densities

$$\begin{aligned} f_n : \mathbb{R}^+ \times S^n &\rightarrow \mathbb{R} \\ (t, \mathbf{X}^n) &\mapsto f_n(t, \mathbf{X}^n) \end{aligned} \quad (6)$$

for $n \in \{0, \dots, N\}$. Consistently with the fact that Ω is a subsystem of U , f_n is explicitly defined as:

$$f_n(t, \mathbf{X}^n) = \binom{N}{n} \int_{(S_c)^{N-n}} F_N(t, \mathbf{X}^n, \Xi_n^N) d\Xi_n^N \quad (7)$$

with $\Xi_n^N \equiv [\Xi_{n+1}, \dots, \Xi_N]$ where $\Xi_i = (\vec{p}_i, \vec{q}_i) \in S_c$ and normalization condition: $\sum_{n=0}^N \int_{\Omega^n} \int_{(\mathbb{R}^3)^n} f_n(t, (\mathbf{p}, \mathbf{q})) d\mathbf{p} d\mathbf{q} = 1$ (see Ref. 3).

D. Derivation of a Liouville-like equation for Ω

The starting point is Eq. 4 and the strategy to achieve a Liouville-like equation for f_n consists in the marginalization of Eq. 4 w.r.t. the degrees of freedom of the $N - n$ particles in Ω_c . The procedure can be schematized as consisting of two steps

- **Step I:** Marginalization of the term $\sum_{i=1}^N \nabla_{\vec{p}_i} \cdot (-\nabla_{q_i} V_{\text{tot}}(\mathbf{q}^N) F_N)$. Since $V_{\text{tot}}(\mathbf{q}^N) = \frac{1}{2} \sum_{i,j=1; j \neq i}^N V(\vec{q}_j - \vec{q}_i)$ one needs to analyze three specific situations:
 - (1) $i, j \in \Omega_c$; (2) $i, j \in \Omega$; (3) $i \in \Omega, j \in \Omega_c$.
- **Step II:** Marginalization of the term $\sum_{i=1}^N \nabla_{\vec{q}_i} \cdot (\vec{v}_i F_N)$, where two specific situations need to be analyzed:
 - (1) $i \in \Omega$ and (2) $i \in \Omega_c$

1. Step I: Results

1. $\forall i: n+1 \leq i \leq N; \forall j$

$$\int_{B_r(0)} \nabla_{\vec{p}_i} \cdot (\nabla_{q_i} V_{\text{tot}}(\mathbf{q}^N) F_N) d^3 p_i = \int_{\partial B_r(0)} \vec{n} \cdot (\nabla_{q_i} V_{\text{tot}}(\mathbf{q}^N)) F_N d\sigma_{p_i} \rightarrow 0 \quad (r \rightarrow \infty) \quad (8)$$

where $B_r(0)$ is the sphere of radius r in momentum space centered at the origin. It is assumed that F_N decays sufficiently rapidly for large $|\vec{p}_i|$ for the boundary integral to vanish in the limit. This is certainly true, e.g., for the Boltzmann distribution $F_N \sim \exp[-\frac{(\vec{p}_i)^2/M}{2kT}]$.

2. $\forall i, j : 1 \leq i, j \leq n$

$$\binom{N}{n} \int_{(S_c)^{N-n}} \nabla_{\vec{p}_i} \cdot \left(\nabla_{q_i} V(\vec{q}_i - \vec{q}_j) F_N(t, \mathbf{X}^n, \Xi_n^N) \right) d\Xi_n^N = \nabla_{\vec{p}_i} \cdot \left(\nabla_{q_i} V(\vec{q}_i - \vec{q}_j) f_n(t, \mathbf{X}^n) \right) \quad (9)$$

3. $\forall i : 1 \leq i \leq n; \forall j : n+1 \leq j \leq N$

$$\begin{aligned} \binom{N}{n} \int_{S_c} \int_{(S_c)^{N-n-1}} \nabla_{\vec{p}_i} \cdot \left(-\nabla_{q_i} V(\vec{q}_i - \vec{q}_j) F_N(t, \mathbf{X}^{n-1}, X_i, (\vec{p}_j, \vec{q}_j), \Xi_{n+1}^N) \right) d\Xi_{n+1}^N dp_j dq_j \\ = \nabla_{\vec{p}_i} \cdot \left(\vec{F}_{\text{av}}(\vec{q}_i) f_n(t, \mathbf{X}^{n-1}, X_i) \right) \end{aligned} \quad (10)$$

with

$$\vec{F}_{\text{av}}(\vec{q}_i) = - \int_{S_c} \nabla_{\vec{q}_i} V(\vec{q}_i - \vec{q}_j) f_2^\circ(X_j | X_i) dX_j \quad (11)$$

denoting the mean field force exerted by the outer particles onto the i th inner particle under the assumptions that

- (a) pair interactions $V(\mathbf{q}_i - \mathbf{q}_j)$ are short-ranged so that pair interactions are relevant only close to the open system's boundary,
- (b) the probability density of finding n particles in states $(\mathbf{X}^{n-1}, X_i) \in S^n$ and one other outer particle in X_j is given by $f_n(\mathbf{X}^{n-1}, X_i) f_2^\circ(X_j | X_i)$,
- (c) $f_2^\circ(X_{\text{out}} | X_{\text{in}})$ is a known or modelled conditional distribution for joint appearances of an outer particle given the state of an inner one.

Here we consider assumption (a) as a physical necessity for assumptions (b) and (c) to be justifiable in the first place, while the latter two encode the more general assumption that the statistics of the reservoir is independent of the instantaneous state of the open system for the present purposes.

2. Step II: Results

1. $\forall i \in 1, \dots, n$

$$\binom{N}{n} \int_{(S_c)^{N-n}} \nabla_{\vec{q}_i} \cdot (\vec{v}_i F_N(t, \mathbf{X}^n, \Xi_n^N)) d\Xi_n^N = \nabla_{\vec{q}_i} \cdot (\vec{v}_i f_n) \quad (12)$$

2. $\forall i : n+1 \leq i \leq N$, one of the integrals will be over $\Xi_i \in S_c$ which, after summing over the respective terms and utilizing the indistinguishability of the particles, leads to

$$\begin{aligned} & \binom{N}{n} (N-n) \int_{S_c} \int_{(S_c)^{N-n-1}} \nabla_{\vec{q}_i} \cdot (\vec{v}_i F_N(t, \mathbf{X}^n, (\vec{p}_i, \vec{q}_i), \Xi_{n+1}^N)) d\Xi_{n+1}^N d\Xi_i \\ &= -(n+1) \int_{\partial\Omega} \int_{\mathbb{R}^3} (\vec{v}_i \cdot \vec{n}) \hat{f}_{n+1}(t, \mathbf{X}^n, (\vec{p}_i, \vec{q}_i)) d^3 p_i d\sigma_i. \end{aligned} \quad (13)$$

Here, guided by the theory of characteristics, we distinguish the relevant forms of \hat{f}_{n+1} for outgoing and incoming particles as follows: Under the assumption of statistical independence of the reservoir particle states from those of the inner particles, we have

$$\hat{f}_{n+1} = \begin{cases} f_{n+1} & (\vec{v}_i \cdot \vec{n} > 0) \\ f_n f_1^\circ & (\vec{v}_i \cdot \vec{n} < 0) \end{cases}, \quad (14)$$

where f_1° is the single particle (equilibrium) density assumed for the reservoir. Alternatively, assuming a grand canonical (GC) distribution for state space trajectories that enter the open system from outside one could assume

$$\hat{f}_{n+1} = \begin{cases} f_{n+1} & (\vec{v}_i \cdot \vec{n} > 0) \\ f_{n+1}^{\text{GC}} & (\vec{v}_i \cdot \vec{n} < 0) \end{cases} \quad (15)$$

as a plausible model.

Note that we do not intend to promote the closure assumptions regarding the reservoir statistics introduced above (through the functions f_1° and f_2°) as being optimal or preferable over alternative formulations. Instead, these closures are meant to be placeholders that highlight the principal necessity of explicitly formulating assumptions on the reservoir behavior in the context of the present derivations.

E. Final Equation

Combining the results of Step I with the results of Step II one obtains a hierarchy of Liouville-type equations for f_n :

$$\frac{\partial f_n}{\partial t} + \sum_{i=1}^n \left(\nabla_{\vec{q}_i} \cdot (\vec{v}_i f_n) + \nabla_{\vec{p}_i} \cdot (\vec{F}_i f_n) \right) = \Psi_n + \Phi_n^{n+1} \quad (16)$$

with

$$\begin{aligned} \vec{F}_i &= -\frac{1}{2} \sum_{\substack{i,j=1 \\ j \neq i}}^n \nabla_{\vec{q}_i} V(\vec{q}_i - \vec{q}_j) \\ \Psi_n &= -\sum_{i=1}^n \nabla_{\vec{p}_i} \cdot \left(\vec{F}_{\text{av}}(\vec{q}_i) f_n(t, \mathbf{X}^{i-1}, X_i, \mathbf{X}_i^{n-i}) \right) \\ \Phi_n^{n+1} &= (n+1) \int \int_{\partial\Omega (\vec{p}_i \cdot \vec{n}) > 0} (\vec{v}_i \cdot \vec{n}) \left(f_{n+1}(t, \mathbf{X}^n, (\vec{p}_i, \vec{q}_i)) - f_n(t, \mathbf{X}^n) f_1^\circ(\vec{q}_i, -\vec{p}_i) \right) d^3 p_i d\sigma_i \end{aligned}$$

III. THE CASE OF TWO DISTINCT RESERVOIRS AT DIFFERENT THERMODYNAMIC CONDITIONS

Let us consider a prototype situation as that illustrated in Fig. 2 where Ω is a region that separates the Universe in two distinct (infinite) reservoirs, (Res_1 and Res_2), at different thermodynamic conditions. Straightforward physical considerations lead to the conclusion that Ω has a spatially asymmetric exchange with the Universe and may thus possess a non-equilibrium stationary state. Formally one can proceed as for the case of a single reservoir and derive an equation for f_n in this situation. The total probability distribution function F_N describes the entire Universe including the thermodynamic states of Res_1 and Res_2 . From F_N , by marginalizing w.r.t. $N - n$ degrees of freedom of particles in Ω_c , one obtains the distribution function f_n of the open system. Furthermore, the Liouville equation for F_N applies as before, and thus by marginalizing the Liouville equation for F_N w.r.t. the degrees of freedom of the particles in Ω_c one would obtain the corresponding Liouville-type equation for f_n ($n \in \{0, \dots, N\}$) as for the case of one reservoir of Ref. 3. This means an analytic derivation of the conditions of non-equilibrium induced by the concurrent action of the two




Figure 2. The partitioning of the "universe" in two large reservoirs, Res_1 and Res_2 , and the subsystem of interest Ω , with $\partial\Omega^1$ the boundary surface between Res_1 and Ω and $\partial\Omega^2$ the boundary surface between Res_2 and Ω , such that $\partial\Omega \equiv \partial\Omega^1 \cup \partial\Omega^2$. The two reservoirs are set up at different thermodynamic conditions, i.e. different densities, ρ_1, ρ_2 , and different temperatures, T_1, T_2 .

reservoirs. In the sections below we will follow the marginalization procedure adopted in the previous section and adapted to the set up illustrated in Fig. 2

A. Step I revised with Res_1 and Res_2

1. $\forall i: n+1 \leq i \leq N; \forall j$

$$\int_{B_r(0)} \nabla_{\vec{p}_i} \cdot (-\nabla_{q_i} V_{\text{tot}}(\mathbf{q}^N) F_N) d^3 p_i = - \int_{\partial B_r(0)} \vec{n} \cdot (\nabla_{q_i} V_{\text{tot}}(\mathbf{q}^N)) F_N d\sigma_{p_i} \rightarrow 0 \quad (r \rightarrow \infty). \quad (17)$$

Also in this case, in the limit of the radius of the ball tending to infinity the term will vanish because the F_N will decay sufficiently rapidly for large momenta, i.e. one can expect a decay as $\exp[-\frac{(\vec{p}_i)^2/M}{2kT_1}]$ in Res_1 and $\exp[-\frac{(\vec{p}_i)^2/M}{2kT_2}]$ for Res_2 (assuming they are both much larger than Ω).

2. $\forall i, j : 1 \leq i, j \leq n$

$$\binom{N}{n} \int_{(S_c)^{N-n}} \nabla_{\vec{p}_i} \cdot \left(\nabla_{q_i} V(\vec{q}_i - \vec{q}_j) F_N(t, \mathbf{X}^n, \Xi_n^N) \right) d\Xi_n^N = \nabla_{\vec{p}_i} \cdot \left(\nabla_{q_i} V(\vec{q}_i - \vec{q}_j) f_n(t, \mathbf{X}^n) \right) \quad (18)$$

This term remains the same as before, i.e., i and j are both particles of Ω so the marginalization w.r.t. the particles outside implies that the whole information about the particles of the reservoirs is integrated out.

3. $\forall i : 1 \leq i \leq n; \forall j : n+1 \leq j \leq N$

$$\begin{aligned} & \binom{N}{n} \int_{S_c} \int_{(S_c)^{N-n-1}} \nabla_{\vec{p}_i} \cdot \left(-\nabla_{q_i} V(\vec{q}_i - \vec{q}_j) F_N(t, \mathbf{X}^{n-1}, X_i, (\vec{p}_j, \vec{q}_j), \Xi_{n+1}^N) \right) d\Xi_{n+1}^N dp_j dq_j \\ &= \nabla_{\vec{p}_i} \cdot \left(\vec{F}_{\text{av}}(\vec{q}_i) f_n(t, \mathbf{X}^{n-1}, X_i) \right) \end{aligned} \quad (19)$$

Here emerges the first substantial difference; for $1 \leq i \leq n$ one needs to carefully consider the dependency of the potential energy on the coordinates of the particle in Ω and on the coordinates of the particles in each of the two distinct reservoirs. The expression of $\vec{F}_{\text{av}}(\vec{q}_i)$ carries the fact that the boundary on the right side has a different thermodynamic and statistical mechanics behaviour than the boundary on the left side, depending on the specific subdomain of S_c over which the integration in the variable X_j is carried out. This implies that the assumed probability density of finding n particles in states $(\mathbf{X}^{n-1}, X_i) \in S^n$ and one other outer particle in X_j is now given by $f_n(\mathbf{X}^{n-1}, X_i) f_2^{\circ, R1}(X_j|X_i)$ if $X_j \in \text{Res}_1$ and by $f_n(\mathbf{X}^{n-1}, X_i) f_2^{\circ, R2}(X_j|X_i)$ if $X_j \in \text{Res}_2$. The integration of X_j over the whole S_c results now in the sum of two integrals, i.e. the integration of X_j over the domain Res_1 and the integration of X_j over the domain Res_2 , that is $S_c \equiv \text{Res}_1 \cup \text{Res}_2$.

$$\vec{F}_{\text{av}}(\vec{q}_i) = - \left[\int_{\text{Res}_1} \nabla_{\vec{q}_i} V(\vec{q}_i - \vec{q}_j) f_2^{\circ, R1}(X_j|X_i) dX_j + \int_{\text{Res}_2} \nabla_{\vec{q}_i} V(\vec{q}_i - \vec{q}_j) f_2^{\circ, R2}(X_j|X_i) dX_j \right] \quad (20)$$

that is

$$\vec{F}_{\text{av}}(\vec{q}_i) = \vec{F}_{\text{av}}^{R1}(\vec{q}_i) + \vec{F}_{\text{av}}^{R2}(\vec{q}_i). \quad (21)$$

B. Step II revised with Res₁ and Res₂

1. $\forall i \in 1, \dots, n$

$$\binom{N}{n} \int_{(S_c)^{N-n}} \nabla_{\vec{q}_i} \cdot (\vec{v}_i F_N(t, \mathbf{X}^n, \Xi_n^N)) d\Xi_n^N = \nabla_{\vec{q}_i} \cdot (\vec{v}_i f_n)$$

This term remains the same as before because the marginalization w.r.t. the particles outside implies that any information about the reservoirs is integrated out.

2. $\forall i : n + 1 \leq i \leq N$:

the effects of the two distinct reservoirs clearly emerges in the definition of \hat{f} because one needs to define for Res₁, $\hat{f}_{n+1}^{R1} = f_n f_1^{\circ, R1}$, with $f_1^{\circ, R1}$ being the single particle (equilibrium) density in the region on the right side and equivalently $\hat{f}_{n+1}^{R2} = f_n f_1^{\circ, R2}$ for the other region. Or, equivalently, $\hat{f}_{n+1}^{R1} = f_{R1}^{GC}$ (same for R_2 .) if one makes the modeling choice of the grand canonical distribution for each reservoir. Moreover $\partial\Omega \equiv \partial\Omega^1 \cup \partial\Omega^2$ implies:

$$-(n+1) \int_{\partial\Omega} \int_{\mathbb{R}^3} (\vec{v}_i \cdot \vec{n}) \hat{f}_{n+1}(t, \mathbf{X}^n, (\vec{q}_i, \vec{p}_i)) d^3 p_i d\sigma_i = \sum_{k=1}^2 I_{\partial\Omega^k} \quad (22)$$

where

$$I_{\partial\Omega^k} = -(n+1) \int_{\partial\Omega^k} \int_{\mathbb{R}^3} (\vec{v}_i \cdot \vec{n}) \hat{f}_{n+1}(t, \mathbf{X}^n, (\vec{q}_i, \vec{p}_i)) d^3 p_i d\sigma_i. \quad (23)$$

In the case $\vec{v}_i \cdot \vec{n} < 0$ one has to replace \hat{f}^{n+1} with $f_n f_1^{\circ, R1}$ in the integral over $\partial\Omega^1$ and with $f_n f_1^{\circ, R2}$ in the integral over $\partial\Omega^2$.

C. Final equation with Res₁ and Res₂

After collecting the results from the previous sections we straightforwardly obtain:

$$\frac{\partial f_n}{\partial t} + \sum_{i=1}^n \left(\nabla_{\vec{q}_i} \cdot (\vec{v}_i f_n) + \nabla_{\vec{p}_i} \cdot (\vec{F}_i f_n) \right) = \sum_{R=R1, R2} \Psi_{n,R} + \Phi_{n,R}^{n+1}, \quad (24)$$

where

$$\begin{aligned}
\Psi_{n,R1} &= - \sum_{i=1}^n \nabla_{\vec{p}_i} \cdot \left(\vec{F}_{\text{av}}^{R1}(\vec{q}_i) f_n(t, \mathbf{X}^{i-1}, X_i, \mathbf{X}_i^{n-i}) \right) \\
\Phi_{n,R1}^{n+1} &= (n+1) \int_{\partial\Omega^1} \int_{(\vec{p}_i \cdot \vec{n}) > 0} (\vec{v}_i \cdot \vec{n}) \left(f_{n+1}(t, \mathbf{X}^n, (\vec{q}_i, \vec{p}_i)) - f_n(t, \mathbf{X}^n) f_1^{\circ, R1}(\vec{q}_i, -\vec{p}_i) \right) d^3 p_i d\sigma_i
\end{aligned} \tag{25}$$

and analogously for $r = R2$. The setup of Fig. 2 and the marginalization procedure can be straightforwardly extended to an arbitrary number of m disjoint reservoirs $R1, R2, R3 \dots Rm$, interfaced with Ω :

$$\frac{\partial f_n}{\partial t} + \sum_{i=1}^n \left(\nabla_{\vec{q}_i} \cdot (\vec{v}_i f_n) + \nabla_{\vec{p}_i} \cdot (\vec{F}_i f_n) \right) = \sum_{r=R1}^{Rm} \Psi_{n,r} + \Phi_{n,r}^{n+1}, \tag{26}$$

IV. COMPARISON WITH OTHER MODELS

The linear coupling of the open system to distinct reservoirs is the starting point of relevant mathematical models that describe exchange of matter and energy of a system with its surroundings (see e.g. Refs. 5–7 and references therein). For example the in the well established model of Bergmann-Lebowitz^{5,6}, the Liouville equation for the distribution function, $f_n(X_n, t)$ of the open system with n particles, assumes, *a priori*, the linear sum of the action of the different reservoirs:

$$\begin{aligned}
\frac{\partial f_n(X_n, t)}{\partial t} &= \{f_n(X_n, t), H(X_n)\} + \\
&+ \sum_r \sum_{n'=0}^{\infty} \int dX'_{n'} [K_{nn'}^r(X_n, X'_{n'}) f_{n'}(X'_{n'}, t) - K_{n'n}^r(X'_{n'}, X_n) f_n(X_n, t)]. \tag{27}
\end{aligned}$$

Each system-reservoir coupling is assumed to consist of an impulsive interaction formalized by a kernel, $K_{nn'}(X'_{n'}, X_n)$, i.e. a transition probability per unit time from an n -particle (open system) and phase space configuration X_n to n' particles and phase space configuration $X'_{n'}$. The overall global effect resulting from the interactions of the open system with its surrounding is assumed to be linear, i.e. : $\sum_r \sum_{n'=0}^{\infty} \int dX'_{n'} [K_{nn'}^r(X_n, X'_{n'}) f_{n'}(X'_{n'}, t) - K_{n'n}^r(X'_{n'}, X_n) f_n(X_n, t)]$, where the sum goes over different reservoirs labeled with r . The

linearity for the coupling is actually implicit in the assumption of impulsive interactions. Each system-reservoir interaction is considered discrete in time so that the open system interacts separately in time with each reservoir. Bergmann and Lebowitz also note that the model of impulsive interaction represents only an asymptotic limit which is not always realistic (see Ref. 6).

A thermodynamic perspective to justify a linear coupling is instead employed by Gay-Balmaz and Yoshimura⁷. In their model the system interacts with different “ports” each of which is a source of energy and mass and mechanical work that can be injected into/adsorbed from the system. The resulting global model is built by adding up the contribution of each port. Such a modeling perspective is justified by the application of the first principle of thermodynamics expressed in the form of time variation of the energy of the system due the action of the ports: $\frac{d}{dt}E = \sum_{n=1}^N P_{W,n}^{ext} + P_{H,n}^{ext} + P_{M,n}^{ext}$, where $P_{W,n}^{ext}$ is the power corresponding to the work done from the n -th reservoir on the system, $P_{H,n}^{ext}$ is the power corresponding to the heat transfer from the n -th reservoir to the system and $P_{M,n}^{ext}$ is the power corresponding to the matter transfer from the n -th reservoirs to the system.

By comparing the two examples above with the derivation proposed here in the previous sections it becomes clear that from the physical point of view the Bergmann-Lebowitz model and the Gay-Balmaz-Yoshimura model rely on the neglect of any feedback of the system onto the reservoirs. Our model also predicts a linear combination of the action of different reservoirs. Yet, we have restricted our analysis to short range pair interactions and an explicit assumption of statistical independence of reservoir states from the open system states. In turn, given that the extension of the region of coupling has negligible thickness, the expected surface/volume ratio necessary to define open systems without the microscopic knowledge of the reservoir is achieved.

We have not derived any nonlinear or memory effects as yet by considering long range interactions or relaxing the statistical independence assumption regarding the states of reservoir particles close to the open system boundary. Both nonlinear and memory effects are expected in general, as the following two examples demonstrate:

- Memory effects: We are dealing with thermodynamically compressible systems supporting sound waves. It is known from the theory of fluid dynamics that the modelling

of acoustic waves that exit an open domain without reflection at its boundary requires one to keep track of the system's history to properly capture the wave dynamics in the reservoir⁸.

- Nonlinearities: When the open system is at a higher pressure than the reservoir(s) initially, then the velocity statistics of outer particles near the open system boundary is clearly biased towards the outward pointing normal. This effect could be modelled by introducing single and two-particle statistics $f_{n,1}^\circ$ and $f_{n,2}^\circ$ that depend explicitly on the open system particle number n , and by coupling these functions to thermodynamic averages of the entire hierarchy of distribution functions f^ν $\nu \in \{0, \dots, N\}$. This ansatz will introduce a nonlinear effect in (20) and (23). If nonlinear effects are to arise in the presence of long range interactions, then our model would similarly be able to account for these.

Yet, our model tells something more. Even within the assumption of short range interactions, one still needs basic information about the reservoirs at the interface, e.g., one and two-particle distributions. These latter are assumed to be stationary, under the approximation that the reservoir, due to its size, fully controls the particle distribution at the surface. Such an approximation is no more valid if the range of interaction is extended because the system itself starts to significantly influence the distribution of particles near the boundary and a generally nonlinear feedback between the reservoirs is to be expected.

However, from the practical point of view an interesting question is whether the model with a linear combination of reservoir actions is sufficient for modeling systems of interest in condensed matter and thus to be used to develop suitable algorithms for numerical calculations, such as Molecular Dynamics approaches. This aspect is discussed in the next section.

V. NUMERICAL IMPLEMENTATION OF THE LINEAR COMBINATION OF RESERVOIR ACTIONS AND ITS TEST OF VALIDITY

In a recent work⁴ we have employed the Adaptive Resolution molecular dynamics algorithm (AdResS)¹⁰ to test the concept of linear combination of reservoir actions on an open




Figure 3. The AdResS set up: an atomistic region AT and an interface region Δ where molecule i interacts with molecule j through the potential $U_{ij} = U(\mathbf{q}_i - \mathbf{q}_j)$. On the left side is illustrated the large reservoir, TR , of non-interacting particles (tracers). In the $\Delta + TR$ region a thermodynamic force, $F^{th}(\mathbf{q}_i)$, acts on single particles to allow the desired thermodynamic equilibrium. Such force is derived by a self consistent procedure during the equilibration run and in practice acts by bringing the particle density at the desired value. The correspondence with the mathematical model of open system of Fig.1 and Fig.2 is illustrated by identifying each region of AdResS with the equivalent region of Fig.1 and Fig.2.

system. The AdResS algorithm allows to treat an open system at atomistic resolution coupled to a large reservoir of non interacting particles, the coupling occurs through an open interface region where molecules interact atomistically among them, but are also subject to a one particle force (thermodynamic force), determined in the course of the equilibration run (and fixed during the production run), which assure thermodynamic equilibrium, at the given conditions of density and temperature, of the atomistic subsystem (see Fig.3) .

This set up is qualitatively equivalent to the partitioning employed by our model of an open system and, in particular, it is very well suited for a numerical test of the idea of linear action. In fact AdResS can express a setup as that of Fig. 2 where the action of the two distinct reservoirs is coded in two distinct coupling conditions at the corresponding interfaces.

The coupling terms, which correspond to the thermodynamic forces, are calculated in two distinct runs; that is, the system first interacts only in the presence of Res₁ which is at temperature T_1 and density ρ_1 and one obtains the thermodynamic force needed at the interface with Res₁. Next, the system interacts only in the presence of Res₂, at $T = T_2$ and $\rho = \rho_2$ and one obtains the thermodynamic force needed at the interface with Res₂. The simulation of nonequilibrium is then achieved by running a simulation with the set up as in Fig. 3 with the corresponding thermodynamic forces applied in the corresponding interface regions.

This idea was applied to a Lennard-Jones liquid for the case of a gradient of temperature applied along an isobar. Details of the simulation can be found in Ref. 4 and its supplementary material. Here we underline only the relevant message for the current discussion. The results reported in Ref. 4 show that indeed the model reproduces very accurately data from fully atomistic reference simulations in the presence of a thermal gradient. Specifically, the study considers an isobaric setup in which the thermodynamic forces are tuned w.r.t. the temperature and density of each reservoir. Here we discuss a further step, i.e., we consider a case where the idea of linearity is pushed to its edge of validity. To this aim we numerically test what happens in the atomistic region of interest when at the interface regions one has a feedback from the rest of the system. For this purpose we show results for a nonequilibrium simulation of a system where the gradient of the temperature is applied in an isocoric setup.

In such a case the thermodynamic forces are calculated for Res₁ at $T = T_1; \rho = \rho_1 = \rho_2$ and Res₂ at $T = T_2; \rho = \rho_1 = \rho_2$. In the nonequilibrium simulation, however, the thermal gradient also induces a density gradient in AT with a feedback on a finite region of each of the two reservoirs, such that nonlinear effects of coupling at each interface arise. Fig. 4 shows that indeed in the coupling region there are effects due to this feedback, although they are not of relevance for the atomistic region, i.e., for the open system of interest.

This is a relevant conclusion in that we have predicted that in the theoretical model of Ref. 3, when some conditions are not fulfilled, nonlinear effects may arise and must be taken into account. The numerical test reported here tells us instead that, under some thermodynamic conditions corresponding to situations of interest, such effects can be ignored and the approximation of a linear coupling is actually sufficient. It must be underlined here




Figure 4. Density profile of a LJ liquid in a thermal gradient simulated with the isobaric (black line) and isochoric (green line) set-ups of non-equilibrium AdResS. Reference results from full atomistic simulations are given by red symbols. For the representation of the TR region, only the part relevant to the discussion is shown.

that this conclusion may not hold anymore, e.g., in case of extreme gradients with a much stronger feedback of the open system onto the reservoirs. Further studies of the validity of the linearization assumptions under more general conditions are deferred to future work.

VI. CONCLUSIONS

We have shown that the mathematical model of open system of Ref. 3, when extended to the case of multiple disjoint reservoirs, presents a Liouville-like equation where the combined action of the reservoirs enter as the linear sum of each single reservoir's action. The mathematical reason for such a results lies in the working hypothesis that (i) the inter-particle potential is short ranged so that the coupling terms between the open system and the reservoirs are reduced to disjoint surface integrals at each system-reservoir interface, and (ii) that the statistics of reservoir particles residing close to the open system boundary are

independent of the open system state.

We have then discussed such results considering other models of open systems in which the linear combination of the actions of the reservoir is assumed *a priori*. From such discussion emerges the necessity of considering the scenario in which nonlinear effects arise once the hypotheses of short range potentials and statistical independence of the reservoirs becomes less strict. In such a perspective, our model offers the possibility of automatically including such feedbacks between the open system and the attached reservoirs by generalizing the models pertinent models of the reservoir statistics at each interface. Such a possibility is ruled out in the other models of open system considered here. An analytic derivation of such a generalization is not provided here, however, but left to future research.

Instead we have focused on the numerical consequences of such nonlinear effects and reported a numerical test where the reference simulation automatically includes nonlinear effects while the simulation made using our model does not include the nonlinear effects. The results show that even in such conditions the approximation of linear combination of actions is numerically satisfactory for a range of systems of interest. For numerical simulations this result is a relevant information but also calls for an extension of the concepts and its numerical counterpart to nonlinear effects.

ACKNOWLEDGMENTS

This research has been funded by Deutsche Forschungsgemeinschaft (DFG) through grant CRC 1114 “Scaling Cascade in Complex Systems,” Project Number 235221301, Project C01 “Adaptive coupling of scales in molecular dynamics and beyond to fluid dynamics.”

REFERENCES

- ¹L. Delle Site and M. Praprotnik, “Molecular systems with open boundaries: Theory and simulation,” *Phys. Rep.* **693**, 1–56 (2017).
- ²G. Ciccotti and L. Delle Site, “The physics of open systems for the simulation of complex molecular environments in soft matter,” *Soft Matter* **15**, 2114 (2019).

- ³L. Delle Site and R. Klein, “Liouville-type equation for the n -particle distribution function of an open system,” *J.Math.Phys.* **61**, 083102 (2020).
- ⁴R. Ebrahimi Viand, F. Höfling, R. Klein, and L. Delle Site, “Theory and simulation of open systems out of equilibrium,” *J.Chem.Phys.* **153**, 101102 (2020).
- ⁵J. Lebowitz and P. Bergmann, “Irreversible Gibbsian ensembles,” *Ann. Phys.* **1**, 1 (1957).
- ⁶P. Bergmann and J. Lebowitz, “New approach to nonequilibrium processes,” *Phys. Rev.* **99**, 578 (1955).
- ⁷F. Gay-Balmaz and H. Yoshimura, “Dirac structures in nonequilibrium thermodynamics for simple open systems,” *J.Math.Phys.* **61**, 092701 (2020).
- ⁸B. Engquist and A. J. Majda, “Absorbing boundary conditions for the numerical evaluation of waves,” *Math. of Comput.* **31**, 629–651 (1977).
- ⁹M. Praprotnik, L. Delle Site, and K. Kremer, “Adaptive resolution molecular-dynamics simulation: Changing the degrees of freedom on the fly,” *J. Chem. Phys.* **123**, 224106 (2005).
- ¹⁰L. Delle Site, C. Krekeler, J. Whittaker, A. Agarwal, R. Klein, and F. Höfling, “Molecular dynamics of open systems: construction of a mean-field particle reservoir,” *Adv. Theory Simul.* **2**, 1900014 (2019).