AOUP in the presence of Brownian noise: a perturbative approach

David Martin

Laboratoire Matière et Systèmes Complexes, UMR 7057 CNRS/P7, Université de Paris, 10 rue Alice Domon et Léonie Duquet, 75205 Paris Cedex 13, France

E-mail: david.martin91120@gmail.com

Abstract. By working in the small-persistence-time limit, we determine the steady-state distribution of an Active Ornstein Uhlenbeck Particle (AOUP) experiencing, in addition to self-propulsion, a Gaussian white noise modelling a bath at temperature T. This allows us to derive quantitative formulas for the spatial probability density of a confined particle and for the current induced by an asymmetric periodic potential. These formulas disentangle the respective roles of the passive and active noises on the steady state of AOUPs, showing that both the correction to the Gibbs-Boltzmann distribution and the ratchet current are of order 1/T. Thus, signatures of nonequilibrium vanish in the limit of large translational diffusion. We probe the range of validity of our analytical derivations by numerical simulations. Finally, we explain how the method presented here to tackle perturbatively an Ornstein Uhlenbeck (OU) noise could be further generalized beyond the Brownian case.

Keywords: Statistical Physics, Stochastic dynamics, Active Matter, non-equilibirum processes

Submitted to: J. Stat. Mech.

1. Introduction

One of the challenges of Active Matter is to understand and predict the emerging properties of assemblies composed of individual agents able to produce mechanical work by locally dissipating energy [1]. By their very nature, such systems break detailed balance and thus lie within the realm of out-of-equilibrium physics. At a theoretical level, the modelling of the individual units of active systems reveal their nonequilibrium nature by involving non-Gaussian and colored noises differing from the familiar Wiener process of equilibrium thermal physics. On the one hand, this approach has been fruitful and numerical simulations have shown that intriguing phenomena, prohibited in equilibrium physics, arise for such noises: accumulation near walls [2,3], emergence of currents in asymmetric periodic potentials [4–7], collective motion [8, 9], motility induced phase separation [10, 11]... On the other hand, from a theoretical standpoint,

resorting to these unusual noises has the technical disadvantage of making algebraic manipulations more cumbersome. Quantifying the departure from equilibrium physics and determining the corresponding steady-state distribution is a whole research field per se [12–18]. In this paper, we focus on an Active Ornstein Uhlenbeck Particle evolving in one space dimension, subjected to an external potential $\phi(x)$ and further experiencing an additional thermal noise. Its position x(t) and self-propulsion v(t) evolve according to the following system of Langevin equations:

$$\dot{x} = -\partial_x \phi + \sqrt{2T} \,\eta_1 + v \tag{1}$$

$$\dot{v} = -\frac{v}{\tau} + \frac{\sqrt{2D}}{\tau} \eta_2 \ . \tag{2}$$

In the above dynamics (1)-(2), η_1 and η_2 are two uncorrelated Gaussian white noises of unit variance, T is the amplitude of the thermal noise while D and τ control the amplitude and the persistence of the self-propulsion respectively. When T=0, (1)-(2) correspond to the workhorse AOUP model which has been used to model transport properties of active colloids [19] as well as collective cell dynamics [20, 21]. On the theoretical side, there has been fundamental interest in its steady-state distribution, which has been characterized both in the limit of small τ [22–26] and in the limit of high τ [25–27]. However, these theoretical approaches ignore the physically relevant presence of an underlying thermal noise and the steady state distribution of (1)-(2) remains elusive for a generic combination of T and D. Indeed, such a combination of different noise sources arises in experiments and has already been used to describe a passive tracer immersed in a bath of active colloids [28]. In this paper, we aim at filling this gap by computing perturbatively the stationary probability density of an AOUP experiencing an additional thermal noise in the small persistence time limit. Note that for $\tau = 0$, the self-propulsion v falls back onto a Wiener process of amplitude D. In this particular case, the dynamics (1)-(2) is an equilibrium one with temperature T+D. Thus, intuitively, one could hope to find analytical formulas that smoothly departs from thermal equilibrium when τ is small. We develop here such a perturbative expansion and our main result is an analytical prediction of the steady-state distribution $\mathcal{P}_s(x,v)$ as a series in $\tau^{1/2}$. Building on it, we make quantitative predictions about two emerging quantities: the marginal in space of the probability density and the current in an asymmetric periodic ratchet [7, 19, 29]. We find both the correction to the Gibbs-Boltzmann distribution and the ratchet current to decay as 1/T, entailing that equilibrium physics, beyond the specific case $\tau = 0$, is also recovered in the limit $T\gg D$.

2. Systematic construction of the probability density function

We first present the derivation of the steady-state distribution $\mathcal{P}_s(x,v)$ as a series in powers of $\tau^{1/2}$. We start from the Fokker-Planck operator \mathcal{L} corresponding to (1)-(2),

which reads

$$\mathcal{L} = \partial_x(\partial_x \Phi) - v\partial_x + \partial_v \left(\frac{v}{\tau}\right) + \frac{D}{\tau^2} \partial_{vv} + T\partial_{xx} . \tag{3}$$

We now rescale v as $\tilde{v} = \sqrt{\tau}v$. Expressed in terms of the rescaled variable, $\mathcal{P}_s(x, \tilde{v})$ evolves according to

$$\tilde{\mathcal{L}}\mathcal{P}_s(x,\tilde{v}) = 0 \tag{4}$$

with the operator $\tilde{\mathcal{L}}$ being defined as :

$$\tilde{\mathcal{L}} = \partial_x(\partial_x \Phi) - \frac{\tilde{v}}{\sqrt{\tau}} \partial_x + \partial_{\tilde{v}} \left(\frac{\tilde{v}}{\tau}\right) + \frac{D}{\tau} \partial_{\tilde{v}\tilde{v}} + T \partial_{xx} . \tag{5}$$

In the remainder of this work, the tilde notation for v and \mathcal{L} will be omitted for notational simplicity. We first note that the Fokker-Planck operator (5) can be written as:

$$\mathcal{L} = \frac{1}{\tau} \mathcal{L}_1 + \frac{1}{\sqrt{\tau}} \mathcal{L}_2 + \mathcal{L}_3 . \tag{6}$$

Where \mathcal{L}_1 , \mathcal{L}_2 and \mathcal{L}_3 are given by

$$\mathcal{L}_1 = D \frac{\partial^2}{\partial^2 v} + \frac{\partial}{\partial v} v \qquad \mathcal{L}_2 = -\frac{\partial}{\partial x} v \qquad \mathcal{L}_3 = \frac{\partial}{\partial x} \frac{\partial \phi}{\partial x} + T \partial_{xx} .$$
 (7)

 \mathcal{L}_1 is the Fokker-Planck generator of the Ornstein-Uhlenbeck process, and its n^{th} eigenfunction P_n is related to the n^{th} physicists' Hermite polynomial $H_n(v) = (-1)^n e^{v^2} \partial_v^n e^{-v^2}$:

$$P_n(v) = \frac{e^{-\frac{v^2}{2D}} H_n\left(\frac{v}{\sqrt{2D}}\right)}{\sqrt{2^n n! 2\pi D}} \ . \tag{8}$$

The family $\{P_n\}$ are eigenfunctions of the operator \mathcal{L}_1 satisfying

$$\mathcal{L}_1 P_n = -n P_n \tag{9}$$

and they are further orthogonal to the family $\{H_n\}$ as

$$\delta_{k,n} = \int_{-\infty}^{+\infty} \frac{H_k\left(\frac{v}{\sqrt{2D}}\right)}{\sqrt{2^k k!}} P_n(v) dv . \tag{10}$$

We use the P_n 's to search for the solution of the stationnary distribution \mathcal{P}_s under the form of:

$$\mathcal{P}_s(x,v) = \sum_n P_n(v) A_n(x) . \tag{11}$$

Using the orthogonality property (10), the A_n 's can be obtained as

$$A_n(x) = \int \mathcal{P}_s(x, v) \frac{H_n\left(\frac{v}{\sqrt{2D}}\right)}{\sqrt{2^n n!}} dv .$$
 (12)

Injecting (11) into (4) and using (9), we find that A_n is a solution of

$$\sum_{n} P_n(v)\partial_x \left(\partial_x \phi A_n\right) + \sum_{n} P_n(v)T\partial_{xx}A_n - \sum_{n} \frac{nP_n(v)}{\tau}A_n - \sum_{n} \frac{vP_n(v)}{\sqrt{\tau}}\partial_x A_n = 0.$$
 (13)

Using the recurrence property of Hermite polynomials, $H_{n+1}(v) = 2vH_n(v) - 2nH_{n-1}(v)$, we decompose vP_n into a sum of P_{n+1} and P_{n-1}

$$vP_n = \sqrt{(n+1)D}P_{n+1} + \sqrt{nD}P_{n-1} . {14}$$

We are now in position to project equation (13) onto H_k and use the orthogonality relation (10). This leads us to the following recursion relation for the A_n 's

$$0 = -nA_n - \sqrt{\tau}\sqrt{(n+1)D}\partial_x A_{n+1} - \sqrt{\tau}\sqrt{nD}\partial_x A_{n-1} + \tau\partial_x(\partial_x \phi A_n) + \tau T\partial_{xx} A_n .$$
 (15)

We now look for the A_n 's as series in powers of $\tau^{1/2}$. We shall assume that A_{2k} contains only integer powers of τ and that its first nonzero contribution is of order τ^k . Likewise, we shall assume that A_{2k+1} contains only half-integer powers of τ and that its first nonzero contribution is of order $\tau^{k+1/2}$. We thus propose the scaling ansatz

$$A_0 = A_0^0(x) + \tau A_0^2(x) + \tau^2 A_0^4(x) + \dots$$
(16)

$$A_1 = \tau^{1/2} A_1^1(x) + \tau^{3/2} A_1^3(x) + \tau^{5/2} A_1^5(x) + \dots$$
 (17)

$$A_2 = \tau A_2^2(x) + \tau^2 A_2^4(x) + \tau^3 A_2^6(x) + \dots$$
 (18)

. . .

Let us now show that the A_i^j can be computed recursively. Looking at (15) for n = 0, we get

$$\partial_x A_1 = \sqrt{\frac{\tau}{D}} \left[\partial_x \left(\partial_x \phi A_0 \right) + T \partial_{xx} A_0 \right] . \tag{19}$$

Equating coefficients of order $\tau^{k/2}$ on both sides of (19) and integrating the x variable leads to:

$$A_1^k = \frac{1}{\sqrt{D}} \left[\partial_x \phi A_0^{k-1} + T \partial_x A_0^{k-1} \right] + b_k . \tag{20}$$

Further equating coefficients of order $\tau^{k/2}$ in (15), we obtain:

$$A_{n}^{k} = -\frac{\sqrt{(n+1)D}}{n} \partial_{x} A_{n+1}^{k-1} - \sqrt{\frac{D}{n}} \partial_{x} A_{n-1}^{k-1} + \frac{\partial_{x} \left(\partial_{x} \phi A_{n}^{k-2}\right)}{n} + \frac{T}{n} \partial_{xx} A_{n}^{k-2} . \tag{21}$$

Taking k = n in (21) and using that $A_n^j = 0$ for $j \le n$ yields the expression of A_n^n as a function of A_0^0 :

$$A_n^n = -\sqrt{\frac{D}{n}} \partial_x A_{n-1}^{n-1} = (-1)^n \frac{D^{n/2}}{\sqrt{n!}} \partial_x^n A_0^0 .$$
 (22)

Using expression (20) for k = 1 and expression (22) for n = 1, we obtain a closed equation on A_0^0 :

$$\partial_x \phi \ A_0^0 + (T+D)\partial_x A_0^0 = -b_1 \sqrt{D} \ .$$
 (23)

Since A_0^0 corresponds to the equilibrium stationary measure when $\tau = 0$ we must have

$$\int_{-\infty}^{+\infty} \mathcal{P}_s(x,v)|_{\tau=0} \ dv = A_0^0 = c_0 \ e^{-\frac{\phi}{T+D}}$$
 (24)

with c_0 fixed by normalization:

$$c_0 = \left(\int_{-\infty}^{+\infty} e^{-\frac{\phi}{T+D}} dx\right)^{-1} . \tag{25}$$

The constant b_1 is self-consistently fixed to zero such that (24) is a solution of (23). We now set out to compute the next order correction A_0^2 . Applying (21) for n = 1 and k = 3 gives:

$$A_1^3 = -\sqrt{2D}\partial_x A_2^2 - \sqrt{D}\partial_x A_0^2 + \partial_x \left(\partial_x \phi A_1^1\right) + T\partial_{xx} A_1^1 . \tag{26}$$

In (26), we can use (22) to express A_1^1 and A_2^2 as a function of A_0^0 and (20) to express A_1^3 as a function of A_0^2 . We thus obtain a differential equation for A_0^2 :

$$\frac{\partial_x \phi A_0^2}{T+D} + \partial_x A_0^2 = -\frac{D^2 \partial_x^3 A_0^0}{T+D} + \frac{D \partial_x (\partial_x \phi \partial_x A_0^0)}{T+D} + \frac{TD \partial_x^3 A_0^0}{T+D} - \frac{b_3 \sqrt{D}}{T+D} (27)$$

Using (24), we can integrate (27) and determine the expression of A_0^2

$$A_0^2 = c_0 e^{-\frac{\phi}{T+D}} D \left(-\frac{(\partial_x \phi)^2}{2(T+D)^2} + \frac{\partial_{xx} \phi}{T+D} \right) + c_2 e^{-\frac{\phi}{T+D}} - \frac{b_3 \sqrt{D}}{T+D} e^{-\frac{\phi}{T+D}} \int_0^x e^{\frac{\phi}{T+D}} dx$$
 (28)

where c_0 is defined in (25). Equation (28) involves two integration constants: c_2 and b_3 . While c_2 is found by normalization, requiring $\int_{-\infty}^{+\infty} A_0^2(x) dx = 0$, b_3 is fixed by boundary conditions on A_0^2 as we shall see in the next sections. We remark that (28) shares a common feature with the distribution of other active models [30, 31]: it is non-local. Indeed, a perturbation of the potential $\delta\phi(x)$ localized around position x will affect the steady-state at position x' located far away from x. This strongly differs from Boltzmann distribution and leads to intriguing phenomena, for example in baths of bacteria [32]. The recursion can be iterated up to an arbitrary order in τ to find both the A_0^{2k} 's and the A_i^{k} 's for i > 0. In addition to the previous constants c_{2i} and b_{2i+1} for i < k, which were determined for lower orders, A_0^{2k} generically depends on two new integration constants: c_{2k} and b_{2k+1} . The former, c_{2k} , is found by requiring the normalization of A_0^{2k} while the latter b_{2k+1} is fixed by boundary conditions for A_0^{2k} . For example, the differential equation on A_0^4 is found by applying (21) for (n=2, k=4)and (n = 1, k = 5). Its solution not only depends on c_2 and b_3 , which were previously determined upon computing A_0^2 , but also on two new integration constants: c_4 and b_5 . The constant c_4 is found by requiring normalization $\int_{-\infty}^{+\infty} A_0^4 = 0$ and b_5 is fixed by enforcing the correct boundary conditions for A_0^4 . While the explicit expressions of the A_0^{2k} rapidly become cumbersome, their systematic derivation can be implemented with a software such as Mathematica [33]. For illustration purposes, we report the complete expression of $\mathcal{P}_s(x,v)$, with its integration constants, up to the order τ^2 in (A.1).

3. Confining potential: explicit computation and numerics

The marginal in space of $\mathcal{P}_s(x,v)$ can be used to quantify how the steady-state distribution departs from the Boltzmann weight as τ increases:

$$\mathcal{P}_s(x) = \int_{-\infty}^{+\infty} \mathcal{P}_s(x, v) dv = A_0 = \sum_k A_0^{2k} \tau^k \ . \tag{29}$$

Here we consider the special case of a confining potential ϕ , and we require that, for all $k \geq 1$,

$$\lim_{x \to +\infty} A_0^{2k}(x) = 0 \tag{30}$$

$$\int_{-\infty}^{+\infty} A_0^{2k}(x) = 0. {31}$$

For a simple harmonic confinement, we note that the complete steady-state distribution $\mathcal{P}_s(x,v)$ remains gaussian and we report its expression in Appendix B. In the remainder of this paper, we will focus on the more general case of anharmonic potentials. We remark that equation (30) imposes $b_{2k+1} = 0$ for all $k \geq 1$ while (31) fixes c_{2k} for all $k \geq 1$. The function A_0^{2k} is then uniquely determined. For example, using (28) and the definition of c_0 (25), A_0^2 reads

$$A_0^2 = c_0 e^{-\frac{\phi}{T+D}} D \left(-\frac{(\partial_x \phi)^2}{2(T+D)^2} + \frac{\partial_{xx} \phi}{T+D} \right) - \frac{3 c_0^2 D}{2(T+D)} e^{-\frac{\phi}{T+D}} \int_{-\infty}^{+\infty} \partial_{xx} \phi e^{\frac{\phi}{T+D}} dx . (32)$$

When T=0 in (32), we recover the steady-state of an Active Ornstein Uhlenbeck (AOUP) particle [22]. The cumbersome expression of the full marginal in space $\mathcal{P}_s(x)$ up to order τ^2 is reported in appendix A.2. Our ansatz (16) rests on the hypothesis that $\mathcal{P}_s(x)$ is an analytic function in $\sqrt{\tau}$ which need not necessarily hold for an arbitrary potential. To check this hypothesis, we have to verify whether the series admits a finite radius of convergence. We do this for a potential $\phi(x) = x^4/4$, at fixed D and T and for two different values of τ . For $\tau = 0.01$, we show in Figure 1 that the truncation of (29) to order τ^8 is well-behaved and quantitatively agrees with the stationary distribution obtained numerically. However, for $\tau = 0.2$, Figure 1 shows the successive orders of the truncation to be typical of a divergent series: adding one order in τ increases the series by a larger amount than the sum of the previous terms, thus leading to wild oscillations. While such a result seems disappointing, it does not mean that the full series fails in capturing the steady state. Mathematically speaking, it only entails that the finite truncation yields a poor approximation of the full series and that more work should be carried out to extract physical behaviors. To regularize our diverging truncated sequence, we resort to a Padé-Borel summation method. We first introduce the Borel transform B_N associated to (29):

$$B_N(\tau) = \sum_{k=0}^{N} \frac{A_0^{2k}}{k!} \tau^k \ . \tag{33}$$

The finite-N truncation of the series (29) is exactly recovered from its N^{th} -Borel transform B_N by applying a Laplace inversion :

$$\sum_{k=0}^{N} A_0^{2k} \tau^k = \int_0^\infty B_N(\omega \tau) e^{-\omega} d\omega . \tag{34}$$

The Laplace inversion of expression (33) for B_N indeed leads back to the divergent finite truncation that we wanted to regularize. To avoid such a fate, one has to find a nonpolynomial approximation of $B_N(\tau)$ whose Taylor expansion coincides with the known terms in (33). In the Padé-Borel method, it is achieved by approximating B_N with a rational fraction $F_N = Q_N/R_N$, where Q_N and R_N are polynomials in τ of order N/2 chosen such that $B_N(\tau) = Q_N(\tau)/R_N(\tau) + o(\tau^N)$. The Borel resummation at order N of (29), B_N^r , is defined by replacing B_N in (34) by its Padé approximant F_N :

$$B_N^r = \int_0^\infty \frac{Q_N(\omega \tau)}{R_N(\omega \tau)} e^{-\omega} dt \ . \tag{35}$$




Figure 1. Steady-state distribution of (1)-(2) in a confining potential $\phi(x) = x^4/4$ Top: For $\tau = 0.01$, the finite truncation of (29) converges and agrees with the numerics (a). Its corresponding Borel resummation B_8^r also coincides with simulation data (b). Bottom: For $\tau = 0.2$, the finite truncation of (29) is rapidly diverging away (c). However, the Borel resummation B_8^r accurately follows the data (d). Parameters: D = T = 1, $dt = 10^{-4}$, $time = 10^8$.

Finally, the series (29) is formally obtained from the limit of B_N^r when $N \to \infty$. In this article, we estimate (29) while keeping N finite and we will not evaluate B_N^r beyond N=8. Interestingly, for $\tau=0.2$, while the truncated sequence of (29) is divergent, its Borel resummation B_8^r agrees quantitatively with numerical estimates of the steady-state distribution. In Figure 2, we plot the Borel resummations B_8^r and the corresponding numerics for different values of T. When $T \ll D$, the dynamics (1)-(2) is strongly out-of-equilibrium and the probability density differs significantly from the Boltzmann weight with the presence of two humps. When $T \gg D$, self-propulsion is washed out by thermal noise, the dynamics draws closer to equilibrium and the two humps of the distribution are smoothened out. Note that the Borel resummation B_8^r accurately fits the numerics without any free parameter.

4. Emerging current: analytical formula and numerics

An interesting marker of a non-equilibrium dynamics is the ratchet mechanism by which asymmetric periodic potentials may lead to steady currents. We consider here such a




Figure 2. Steady-state distribution of (1)-(2) in a confining potential $\phi(x) = x^4/4$ for different values of T. Plain curves correspond to Borel resummations B_8^r while symbols are obtained from numerical simulations of (1)-(2). In dashed lines, we plot the Gibbs-Boltzmann distributions for the two limiting cases T=0.1 and T=3.0 to highlight the activity-induced deviation. The Borel resummation B_8^r always fits the data accurately without any free parameters. Parameters : D=1, $dt=10^{-4}$, time $=10^8$.

potential ϕ of period L and we use our perturbative expansion to compute the steady-state current J, defined as

$$J = \langle \dot{x} \rangle \tag{36}$$

$$J = \int_0^L \int_{-\infty}^{\infty} \left(-\partial_x \phi + \frac{v}{\sqrt{\tau}} \right) \mathcal{P}_s(x, v) dx dv \tag{37}$$

$$J = -\sum_{k>0} \int_0^L \partial_x \phi A_0^{2k} dx + \frac{\sqrt{D}}{\sqrt{\tau}} \sum_{k>0} \int_0^L A_1^{2k+1} dx$$
 (38)

$$J = \sum_{k \ge 0} T \int_0^L \partial_x A_0^{2k} dx + L\sqrt{D} \sum_{k > 0} b_{2k+1} \tau^k . \tag{39}$$

To go from (38) to (39), we used the expression of A_1^{2k+1} in (20). We require the marginal in space $\mathcal{P}_s(x)$ to be periodic, which entails A_0^{2k} to be periodic for all $k \geq 0$. The current J then simplifies into:

$$J = L\sqrt{D} \sum_{k>0} b_{2k+1} \tau^k \ . \tag{40}$$

While the $\{b_k\}$ all vanished in the previous section as a result of confinement (30), they do not for a periodic potential. Indeed, the value of b_k is fixed upon requiring the periodicity of A_0^{k-1} . Thus, different boundary conditions lead to different distributions, highlighting once again the nonlocal nature of the steady state. We report the expression of the marginal in space $\mathcal{P}_s(x)$ for a periodic potential up to order τ^2 in (A.3)-(A.4). Using it, we find that $Lb_5\tau^2$ is the first non-vanishing contribution to the current:

$$J = \frac{DL\tau^2}{2(T+D)} \frac{\int_0^L \phi^{(1)2} \phi^{(3)} dx}{\int_0^L e^{\frac{\phi}{T+D}} dx \int_0^L e^{-\frac{\phi}{T+D}} dx} + o(\tau^2) . \tag{41}$$




Figure 3. Current J induced by a ratchet potential $\phi(x) = \sin(\pi x/2) + \sin(\pi x)$ for different values of T. Plain curves correspond to prediction (41) while dots are numerical simulations with error bars given by the standard deviation. **Left:** J as a function of τ . **Right:** J/τ^2 as a function of $1/\tau$. Parameters : D = 1, $dt = 15 \cdot 10^{-4}$, time $= 5 \cdot 10^8$.

The above formula reduces to the recently computed expression of J for an AOUP particle when T=0 [24]. It is interesting to note that, as $T\to\infty$, J vanishes as $J\propto 1/T$. Physically, when the amplitude T of the thermal noise is much bigger than the amplitude D of the self-propulsion, the particle does not "feel" the nonequilibrium nature of its dynamics and J dies out. In Figure 3, we compare our quantitative prediction (41) with the results of numerical simulations for a potential $\phi(x)=\sin(\pi x/2)+\sin(\pi x)$ and different values of T. We find quantitative agreement at small τ for $\tau<0.01$, which confirms our conclusion in the previous section for the radius of convergence of our ansatz (11). Note that J in (40) could also be regularized using Borel resummation to extend the quantitative range of agreement between theory and simulations to higher values of τ , but we leave such a regularization for futur works.

5. Conclusion

We developed theoretical insights for an AOUP submitted to an additional Brownian noise (1)-(2). First, we devised a recurrence scheme allowing us to compute its stationary ditribution at an arbitrary order in τ . We then used this result to show that activity-induced phenomena, such as spatial accumulation near walls or the emergence of current, vanish in the limit $T \gg D$ as an inverse power law 1/T. The regime of high translational diffusion thus provides a second route, besides the regime of low τ , to recover equilibrium physics. Alternatively, it is possible to derive the marginal in space $\mathcal{P}_s(x)$ up to order τ^2 by using a Markovian approximation for the evolution operator: such a method has been developed in parallel to this article [34]. We further remark that our perturbative approach can be generalized to more complex dynamics than the Brownian case. Let us consider a stochastic dynamics \mathcal{S} whose corresponding Fokker-Planck operator is $\mathcal{L}_{\mathcal{S}}$.

Adding an OU noise of amplitude D to S amounts to add $\mathcal{L}_1 + \mathcal{L}_2$ to \mathcal{L}_S . The starting point for our perturbative expansion, A_0^0 , is defined by the equation $(\mathcal{L}_S + D\partial_{xx})A_0^0 = 0$. As long as A_0^0 defined this way is known analytically, the recurrence can be carried out and the effect of the OU noise can be taken into account perturbatively in τ . For example, using our method, one could try to assess the effect of a colored noise on an underdamped Langevin dynamics.

Acknowledgments

We thanks Julien Tailleur, Frédéric van Wijland, Cesare Nardini and Thibaut Arnoulx de Pirey for insightful discussions, continuous support as well as enlightening feedback.

Appendix A. Full steady-state distribution

In this appendix, we report the steady-state probability density $\mathcal{P}_s(x,v)$ up to order τ^2 .

$$\begin{split} e^{\frac{\phi}{T+D}}\mathcal{P}_s(x,v) &= c_0 + \sqrt{\tau}P_1(v)\frac{c_0\sqrt{D}\phi^{(1)}(x)}{T+D} + \tau P_0(v) \bigg[-\frac{c_0D\phi^{(1)2}}{2(T+D)^2} + c_2 + \frac{Dc_0\phi^{(2)}}{T+D} \\ &-\frac{\sqrt{D}b_3}{T+D} \int_0^x e^{\frac{\phi(z)}{T+D}} dz \bigg] + \tau^{\frac{3}{2}} \bigg[\frac{P_3(v)c_0D^{\frac{3}{2}}}{\sqrt{6}} \bigg(\frac{\phi^{(1)3}}{(T+D)^3} - \frac{3\phi^{(1)}\phi^{(2)}}{(T+D)^2} + \frac{\phi^{(3)}}{T+D} \bigg) \\ &+ P_1(v) \bigg(\frac{b_3De^{\frac{\phi}{T+D}}}{T+D} - \frac{\phi^{(1)}b_3D}{(T+D)^2} \int_0^x e^{\frac{\phi(z)}{T+D}} dz + \frac{\sqrt{D}c_2\phi^{(1)}}{T+D} - \frac{c_0D^{\frac{3}{2}}\phi^{(1)3}}{2(T+D)^3} \\ &+ \frac{c_0\sqrt{D}(D^2-T^2)\phi^{(1)}\phi^{(2)}}{(T+D)^3} + \frac{c_0\sqrt{D}T\phi^{(3)}}{T+D} \bigg) \bigg] + \tau^2P_0(v) \bigg[\frac{c_2D\phi^{(2)}}{T+D} - \frac{c_2D\phi^{(1)2}}{2(T+D)^2} \\ &- \frac{D^{\frac{3}{2}}b_3}{(T+D)^2} \int_0^x e^{\frac{\phi(z)}{T+D}}\phi^{(2)}(z) dz + \frac{c_0D^2}{8(T+D)^4}\phi^{(1)4} - \frac{c_0D(D-T)\phi^{(1)2}\phi^{(2)}}{2(T+D)^3} \\ &+ \frac{b_3D^{\frac{3}{2}}}{(T+D)^3} \int_0^x \bigg(\int_0^s e^{\frac{\phi(z)}{T+D}} dz \bigg) \bigg(\phi^{(1)}(s)\phi^{(2)}(s) - (T+D)\phi^{(3)}(s) \bigg) \, ds + c_4 \\ &+ \frac{Dc_0}{2(T+D)} \int_0^x \phi^{(1)2}(z)\phi^{(3)}(z) dz - \frac{Dc_0(D+2T)}{(T+D)^2}\phi^{(3)}\phi^{(1)} - \frac{\sqrt{D}b_5}{T+D} \int_0^x e^{\frac{\phi(z)}{T+D}} dz \\ &+ \frac{Dc_0(D-2T)\phi^{(2)2}}{4(T+D)^2} + \frac{Dc_0(D+2T)\phi^{(4)}}{2(T+D)} \bigg] \end{split} \tag{A.1}$$

In (A.1), c_0 is defined by (25) while c_2 , c_4 , b_3 and b_5 are integration constants whose expressions must be adapted to the boundary conditions. For a confining potential, $\mathcal{P}_s(x,v)$ must vanish for $x \to \pm \infty$, and thus $b_3 = b_5 = 0$ yielding the following spatial distribution:

$$e^{\frac{\phi}{T+D}}\mathcal{P}_{s}(x) = c_{0} + \tau \left[-\frac{c_{0}D\phi^{(1)2}}{2(T+D)^{2}} + c_{2} + \frac{Dc_{0}\phi^{(2)}}{T+D} \right] + \tau^{2} \left[\frac{c_{2}D\phi^{(2)}}{T+D} - \frac{c_{2}D\phi^{(1)2}}{2(T+D)^{2}} + c_{4} \right]$$

$$+ \frac{c_{0}D^{2}}{8(T+D)^{4}}\phi^{(1)4} - \frac{c_{0}D(D-T)\phi^{(1)2}\phi^{(2)}}{2(T+D)^{3}} + \frac{Dc_{0}}{2(T+D)} \int_{0}^{x} \phi^{(1)2}(z)\phi^{(3)}(z)dz$$

$$- \frac{Dc_{0}(D+2T)}{(T+D)^{2}}\phi^{(3)}\phi^{(1)} + \frac{Dc_{0}(D-2T)\phi^{(2)2}}{4(T+D)^{2}} + \frac{Dc_{0}(D+2T)\phi^{(4)}}{2(T+D)} \right]$$
 (A.2)

 c_2 and c_4 are finally found by normalization, requiring $\int_{-\infty}^{+\infty} \mathcal{P}_s(x) dx = 1$. For a periodic potential of period L, the spatial distribution must respect $\mathcal{P}_s(x+L) = \mathcal{P}_s(x)$. This condition implies $b_3 = 0$, but $b_5 \neq 0$ and \mathcal{P}_s reads:

$$e^{\frac{\phi}{T+D}}\mathcal{P}_{s}(x) = c_{0} + \tau \left[-\frac{c_{0}D\phi^{(1)2}}{2(T+D)^{2}} + c_{2} + \frac{Dc_{0}\phi^{(2)}}{T+D} \right] + \tau^{2} \left[\frac{c_{2}D\phi^{(2)}}{T+D} - \frac{c_{2}D\phi^{(1)2}}{2(T+D)^{2}} + c_{4} \right]$$

$$+ \frac{c_{0}D^{2}}{8(T+D)^{4}}\phi^{(1)4} - \frac{c_{0}D(D-T)\phi^{(1)2}\phi^{(2)}}{2(T+D)^{3}} + \frac{Dc_{0}}{2(T+D)} \int_{0}^{x} \phi^{(1)2}(z)\phi^{(3)}(z)dz$$

$$- \frac{Dc_{0}(D+2T)}{(T+D)^{2}}\phi^{(3)}\phi^{(1)} + \frac{Dc_{0}(D-2T)\phi^{(2)2}}{4(T+D)^{2}} + \frac{Dc_{0}(D+2T)\phi^{(4)}}{2(T+D)}$$

$$- \frac{\sqrt{D}b_{5}}{T+D} \int_{0}^{x} e^{\frac{\phi(z)}{T+D}}dz \right]$$
(A.3)

$$b_5 = \frac{D}{2(T+D)} \frac{\int_0^L \phi^{(1)2} \phi^{(3)} dx}{\int_0^L e^{\frac{\phi}{T+D}} dx \int_0^L e^{-\frac{\phi}{T+D}} dx}$$
(A.4)

Once again, c_2 and c_4 are then found by normalization. Note that in expression (A.1), v corresponds to the rescaled variable \tilde{v} . To get the exact steady-state distribution associated to (1)-(2), one has thus to make the change of variable $v \to \sqrt{\tau}v$.

Appendix B. Harmonic potential

We report hereafter the steady-state distribution for the special case of a harmonic potential $\phi(x) = \kappa x^2/2$:

$$\mathcal{P}_s(x,v) = \frac{\sqrt{4ab - c^2}}{2\pi} e^{-ax^2 - bv^2 + cvx}$$
(B.1)

With the constants a, b and c defined as :

$$a = \frac{\kappa (1 + \kappa \tau)^2}{2(D + T(1 + \kappa \tau)^2)} \quad b = \frac{D(1 + \kappa \tau) + T(1 + \kappa \tau)^2}{2D(D + T(1 + \kappa \tau)^2)} \quad c = \frac{\kappa \sqrt{\tau} (1 + \kappa \tau)}{D + T(1 + \kappa \tau)^2}$$
(B.2)

Note that in expression (B.1), v corresponds to the rescaled variable \tilde{v} . To get the exact steady-state distribution associated to (1)-(2), one has thus to make the change of variable $v \to \sqrt{\tau}v$.

Appendix C. Numerical methods

To simulate dynamics (1), we used a discretized Heun scheme while dynamics (2) was integrated exactly using Gillespie's method [35]. The obtained algorithm iterates as follows:

$$\begin{array}{ll} 1 \; \mu \; = \; \exp(-\mathrm{d}t/\tau) \, ; \\ 2 \; \sigma_x \; = \; \sqrt{D(1-\mu^2)/\tau} \, ; \\ 3 \; Y_1 \; = \; \sqrt{2D\tau \, (\mathrm{d}t/\tau - 2(1-\mu) + 0.5(1-\mu^2)) - \tau D(1-\mu)^4/(1-\mu^2)} \, ; \\ 4 \; Y_2 \; = \; \sqrt{\tau D}(1-\mu)^2/\sqrt{1-\mu^2} \, ; \\ 5 \; T_1 \; = \; \sqrt{2T\mathrm{d}t} \, ; \end{array}$$

```
6 Y = x = 0;
 7 \text{ v} = \sqrt{D/\tau} * \text{normal\_distribution}(0,1);
 9 while (t < total time) {
        \eta_1 = \text{normal\_distribution}(0,1);
10
11
        \eta_2 = \text{normal\_distribution}(0,1);
12
       \eta_3 = \text{normal\_distribution}(0,1);
13
       Y = \tau * v * (1-\mu) + Y_1 * \eta_2 + Y_2 * \eta_1;
14
       \mathbf{v} = \mathbf{v} * \boldsymbol{\mu} + \boldsymbol{\sigma}_x * \boldsymbol{\eta}_1;
15
       x_1 = \mathbf{x} - \mathrm{dt} * \partial_x \phi(x) + \mathbf{Y} + T_1 * \eta_3;
       x += Y + T_1 * \eta_3 -0.5 * dt * (\partial_x \phi(x) + \partial_x \phi(x_1));
16
17
        t += dt;
```

At step (17), x(t) is stored in the variable x. The steady-state marginal in space of the distribution $\mathcal{P}_s(x)$ was then obtained by recording the particle's position recurrently into an histogram. The current J was computed using the distance travelled by the particle divided by the duration of the simulation: the error bar on J thus corresponds to the standard deviation. Such a definition for the current was heuristically found to converge faster than computing $J = \langle -\partial_x \phi + v/\sqrt{\tau} \rangle$ with recurrent recordings.

References

- Marchetti M C, Joanny J F, Ramaswamy S, Liverpool T B, Prost J, Rao M and Simha R A 2013 Reviews of Modern Physics 85 1143
- [2] Elgeti J and Gompper G 2009 EPL (Europhysics Letters) 85 38002
- [3] Tailleur J and Cates M 2009 EPL (Europhysics Letters) 86 60002
- [4] Dialynas T, Lindenberg K and Tsironis G 1997 Physical Review E 56
- [5] Lindner B, Schimansky-Geier L, Reimann P, Hänggi P and Nagaoka M 1999 Physical Review E 59 1417
- [6] Sandford C, Grosberg A Y and Joanny J F 2017 Physical Review E 96 052605
- [7] Magnasco M O 1993 Physical Review Letters 71 1477
- [8] Vicsek T, Czirók A, Ben-Jacob E, Cohen I and Shochet O 1995 Physical review letters 75 1226
- [9] Grégoire G and Chaté H 2004 Physical review letters **92** 025702
- [10] Tailleur J and Cates M 2008 Physical review letters 100 218103
- [11] Cates M E and Tailleur J 2015 Annu. Rev. Condens. Matter Phys. 6 219–244
- [12] Bianucci M 2020 arXiv preprint arXiv:2001.05809
- [13] Szamel G 2014 Physical Review E **90** 012111
- [14] Solon A P, Cates M E and Tailleur J 2015 The European Physical Journal Special Topics 224 1231–1262
- [15] Basu U, Majumdar S N, Rosso A and Schehr G 2019 Physical Review E 100 062116
- [16] Basu U, Majumdar S N, Rosso A, Sabhapandit S and Schehr G 2020 Journal of Physics A: Mathematical and Theoretical 53 09LT01
- [17] Hartmann A K, Majumdar S N, Schawe H and Schehr G 2020 Journal of Statistical Mechanics: Theory and Experiment 2020 053401
- [18] Shee A, Dhar A and Chaudhuri D 2020 Soft Matter 16 4776–4787
- [19] Koumakis N, Maggi C and Di Leonardo R 2014 Soft matter 10 5695–5701

- [20] Hakim V and Silberzan P 2017 Reports on Progress in Physics 80 076601
- [21] Deforet M, Hakim V, Yevick H, Duclos G and Silberzan P 2014 Nature communications 5 1-9
- [22] Fodor E, Nardini C, Cates M, Tailleur J, Visco P and Wijland F V 2016 *Physical Review Letters*117
- [23] Bonilla L L 2019 Physical Review E **100** 022601
- [24] Martin D, O'byrne J, Cates M E, Fodor E, Nardini C, Tailleur J and van Wijland F 2020 arXiv preprint arXiv:2008.12972v1
- [25] Woillez E, Kafri Y and Lecomte V 2020 Journal of Statistical Mechanics: Theory and Experiment 2020 063204
- [26] Jung P and Hänggi P 1987 Physical review A 35 4464
- [27] Fily Y 2019 The Journal of Chemical Physics 150 174906
- [28] Maggi C, Paoluzzi M, Pellicciotta N, Lepore A, Angelani L and Di Leonardo R 2014 *Physical review letters* 113 238303
- [29] Bartussek R, Hänggi P, Lindner B and Schimansky-Geier L 1997 Physica D: Nonlinear Phenomena 109 17–23
- [30] Van den Broeck C and Hänggi P 1984 Physical Review A 30 2730
- [31] Solon A P, Fily Y, Baskaran A, Cates M E, Kafri Y, Kardar M and Tailleur J 2015 Nature Physics 11 673–678
- [32] Galajda P, Keymer J, Chaikin P and Austin R 2007 Journal of bacteriology 189 8704–8707
- [33] Inc W R Mathematica, Version 12.1 champaign, IL, 2020 URL https://www.wolfram.com/mathematica
- [34] Private communications with Yongjoo Baek
- [35] Gillespie D T 1996 Physical review E $\mathbf{54}$ 2084