

The iron record of asteroidal processes in carbonaceous chondrites

by

A. Garenne^{1*}, P. Beck^{1,2}, G. Montes-Hernandez³, L. Bonal¹, E. Quirico¹, O. Proux⁴, J.L. Hazemann⁵.

¹Univ. Grenoble Alpes, IPAG, F-38000 Grenoble, France

CNRS, IPAG, F-38000 Grenoble, France

²Institut Universitaire de France, Paris, France.

³Univ. Grenoble Alpes / CNRS-INSU, Institut des Sciences de la Terre (IsTERRE), Grenoble, France.

⁴Observatoire des Sciences de l'Univers de Grenoble (OSUG) CNRS UMS 832, 414 rue de la piscine 38400 Saint Martin d'Hères, France

⁵Institut Néel, 25 av. des Martyrs, 38042, Grenoble, France.

*Corresponding author

Abstract:

The valence of iron has been used in terrestrial studies to trace the hydrolysis of primary silicate rocks. Here, we use a similar approach to characterize the secondary processes namely thermal metamorphism and aqueous alteration of carbonaceous chondrites. X-ray Absorption Near-Edge Spectroscopy (XANES) at the Fe-Edge was performed on a series of 36 CM, 10 CR, 9 CV, and 2 CI chondrites. While previous studies have focused on the relative distribution of Fe^0 with respect to oxidized iron ($\text{Fe}_{\text{ox}} = \text{Fe}^{2+} + \text{Fe}^{3+}$) or the iron distribution in some specific phases (e.g. Urey-Craig diagram (Urey and Craig 1953)) our measurements enable us to assess the fractions of iron in each of its three oxidation states: Fe^0 , Fe^{2+} and Fe^{3+} .

Among the 4 carbonaceous chondrites groups studied, a correlation between the Iron Oxidation Index ($\text{IOI} = ([\text{Fe}^0] + 2[\text{Fe}^{2+}] + 3[\text{Fe}^{3+}]) / [\text{Fe}_{\text{TOT}}]$) and the hydrogen content is observed. However, within the CM group, for which a progressive alteration sequence has been defined, a conversion of Fe^{3+} to Fe^{2+} is observed with increasing degree of aqueous alteration. This reduction of iron can be explained by evolution in the mineralogy of the secondary phases. In the case of the few CM chondrites that experienced some thermal metamorphism, in addition to aqueous alteration, a redox memory of the aqueous alteration is present: a significant fraction of Fe^{3+} is present, together with Fe^{2+} and sometimes Fe^0 .

From our dataset, the CR chondrites show a wider range of IOI from 1.5 to 2.5. In all considered CR chondrites, the three oxidation states of iron coexist. Even in the least altered CR chondrites, the fraction of Fe^{3+} can be high (30 % for MET 00426). This observations confirms that oxidized iron has been integrated during fine-grained amorphous material formation in the matrix (Le Guillou et al. 2015; Le Guillou and Brearley 2014; Hopp and Vollmer 2018).

Last, the IOI of CV chondrites does not reflect the reduced/oxidized classification based on metal and magnetite proportion, but is strongly correlated with petrographic types. The valence of iron in CV chondrites therefore appears to be most closely correlated related to thermal history, rather than aqueous alteration even if these processes can occurs together (Krot et al. 2004; Brearley and Krot 2013).

1. Introduction

With the occurrence of three different oxidation states in planetary environments, the valence of iron has been used as a quantitative tracer of aqueous alteration processes on Earth

(McCollom and Bach 2009; Malvoisin et al. 2012b; Malvoisin et al. 2012a). Corrosion of Fe²⁺-bearing silicates in terrestrial hydrothermal systems leads to the production of secondary Fe³⁺-bearing phases (some phyllosilicates, magnetite) coupled with the reduction of water and production of H₂ (Marcaillou et al. 2011).

Concerning chondritic meteorites, the Urey-Craig diagram, giving their relative iron contents in metal, sulfides, silicates and Fe-oxides which has been used to establish a classification (Weisberg et al. 2006; Krot et al. 2007). The initial budget of metallic iron in chondrites has been set by the compositions and relative abundances of the accreted components (type I and II chondrules, CAIs, metal/sulfides, matrix) and subsequently modified by parent body processes. The observed oxidation states in chondrites result from different processes in two distinct settings: protoplanetary disk and parent body (e.g. Lewis 1972; Rubin et al. 1988; Prinn and Fegley 1989; Zolensky et al. 1989; Bischoff 1998; Krot et al. 2000; Brearley 2006).

Primitive meteorites record the action of water (Brearley 2006) that led to significant alteration of the initially accreted phases. Although some alteration may have occurred in the solar nebula, the majority of the observed effects result from alteration on the asteroidal parent body (Bischoff 1998; Brearley 2014). In some carbonaceous chondrites, aqueous alteration can be pervasive and alteration signatures have been described in matrix, as well as in chondrule mesostasis (Brearley 2006). Meteorites from the CI and CM groups have a mineralogy that is dominated by secondary alteration products (Howard et al. 2009; King et al. 2015) and can contain up to 25 wt. % of different volatile elements in the case of Orgueil (Garenne et al. 2014). All CR chondrites have experienced aqueous alteration. The matrix appears to have been altered first, and chondrules show also aqueous alteration evidence observed as micrometric phyllosilicate rims for the least altered samples (Brearley 2006; Bonal et al. 2013; Harju et al. 2014; Le Guillou et al. 2015; Martinez and Brearley 2018). In CV chondrites, minerals resulting from low temperature metasomatism are less abundant but been unambiguously identified in some of them (e.g., Brearley 2006). These products differ between the oxidized and reduced CV subgroup (Howard et al. 2010).

The process of aqueous alteration leads to specific spectral signatures that can be searched for remotely across the small body population (e.g. small solar system bodies, such as asteroids, satellites, etc.). This includes a water-related feature around 3- μ m (Rivkin et al. 2000; Beck et al. 2010), whose shape and intensity can be used to relate meteorites to asteroids. In

addition, in the visible and near infrared (0.4-2.6 μ m), the observation of absorption features on C-type asteroids around 0.49, 0.70 and 0.90 μ m are usually related to the presence of a significant amount of Fe³⁺. These features are also observed in carbonaceous chondrites (Cloutis et al. 2011a; Cloutis et al. 2011b; Cloutis et al. 2012a ; Cloutis et al. 2012b). The valence of iron (and the mineral hosts) is the first order parameter that controls the visible-near infrared (VNIR) signature of chondrites.

In the present work, we determine the average iron oxidation index, calculated by $([Fe^0]+2[Fe^{2+}]+3[Fe^{3+}])/[Fe_{TOT}]$, in a series of primitive chondrites through synchrotron-based Fe-XANES spectroscopy . We estimate the abundance of the three iron oxidation states in bulk meteorite samples for wide range of petrologic types and chondrite groups. We focus on carbonaceous chondrites for which aqueous alteration has been documented (CM, CV, CR and CI) in the literature.

2. Methods and samples

2.1 Samples

Samples studied here are from the CM, CV, CR and CI groups of carbonaceous chondrites (Table 1). The CIs, CRs and CMs cover a range of aqueous alteration with petrologic type ranging from 1.0 to 3.0 (Rubin et al. 2007; Howard et al. 2011). Among CM samples, some are identified as thermally metamorphosed (Nakamura 2005; Alexander et al. 2012) that caused modification of the secondary products and of the bulk iron oxidation state (Beck et al. 2011). CM chondrite falls and finds are compared to determine the potential oxidation induced by terrestrial residence. Reduced (CV_{Red}) and oxidized (CV_{OxA} and CV_{OxB}) CV chondrites, that have experienced variable degrees of thermal metamorphism were selected as well. Among our 9 CV chondrites, there are 3 CV_{OxA} (Allende, Axtell, Mokoia), 3 CV_{OxB} (Bali, Grosnaja, Kaba) and 3 CV_{Red} (Efremovka, Leoville, Vigarano). Their petrologic types range from 3.1 to 3.6 (Bonal et al. 2006).

2.2 XANES spectroscopy

X-ray absorption spectroscopy is a powerful technique to study the physico-chemical properties of matter, including inter-atomic distance, coordination number, and electronic density. This method results from electronic transitions and photo-electron scattering with the neighbor atoms (single and multiple scatterings), allowing retrieval of structural and electronic information from the probed element.

X-ray Absorption Near-Edge Structure (XANES) spectra can be schematically described in three parts: (i) a pre-edge below edge-energy (E_0) corresponding to bound states transitions (mainly a 1s to 3d transition for the Fe K-edge), (ii) the absorption threshold (or edge jump) corresponding to transitions between the probed level to continuum states and/or to the last unoccupied bound levels (the edge crest, or the so-called white-line, corresponding to the 1s to 4p transition for the Fe K-edge) and (iii) the XANES part after the absorption edge corresponding to the scattering of the emitted photo-electron by the neighbor atoms.

The energy of the absorption edge will change when the core-level energies change. This absorption is specific for each element, allowing individual elements to be studied by selecting the appropriate experimental energy range parameter. X-ray absorption near-edge structure spectroscopy is very sensitive to oxidation state: as the oxidation state increases so does the absorption edge energy (Fig. 1).

2.3 Samples preparation and Fe-XANES measurements

Aliquots of 20 mg from a larger samples of meteorite powder were typically mixed with 30 mg of boron nitride (BN) and pressed in order to obtain a compact pellet with a diameter of 5 mm and a thickness of 2 mm. Each mixture of meteorite with BN was manually ground for 30 minutes in an agate mortar before pressing in order to ensure a homogeneous mixing. Each sample was analyzed for Fe-K-edge XANES at CRG-FAME beamline at the European Synchrotron Radiation Facility (Proux et al. 2005; Proux et al. 2006). The liquid nitrogen-cooled two-crystal monochromator is equipped with Si(220) crystals. Energy resolution is close to the intrinsic resolution of these crystals, i.e. 0.365 eV at the Fe K-edge. Spectra were measured in transmission mode between 7004 and 7385 eV with a typical energy step of 0.2 eV around the pre-edge and 2 eV in the further part (before 7105 eV and after 7150 eV). Energy calibration was achieved by setting the maximum of the first derivative of a Fe metal foil K-edge spectrum at 7112.0 eV.

In order to assess the homogeneity of each pellet, a scan in transmission above the iron absorption edge was performed across the sample, and three XANES spectra were measured at different locations. For each sample, the three XANES spectra were compared and averaged to have a good estimation of the iron oxidation state in the meteorite sample. The raw spectra were normalized using two polynomial curves, for the pre-edge region (linear curve) and the post-edge region (polynomial order: 2) while the edge jump was scaled by a factor to give a jump of 1. These normalizations were performed with the software ATHENA from the IFEFFIT package (Ravel and Newville 2005).

2.4 Calculation of Fe oxidation state

Several processing methods were tested on our spectral set to determine the optimum technique to derive Fe-oxidation state of the samples. In the literature, the pre-edge feature is often used: the combination of its intensity (for a normalized spectra) with its centroid position is used to infer the average valence and the site geometry for iron oxidized atoms (e.g. Wilke et al. 2001). However since significant amounts of Fe^0 are expected in our CR and CV samples, this approach was rejected.

A series of alternative methods to quantify the Fe-oxidation factor were tested, based on (i) the edge position from the derivative of the spectra, (ii) a linear combination of standards on the main part of XANES spectra (from -20 eV to +30 eV compared to E_0), (iii) a linear combination of standards of the pre-edge feature (from -15 eV to -5 eV compared to E_0). The pre-edge, expected to be less sensitive to surrounding atoms than the whole spectra, is more sensitive to the valence of iron. The method based on a linear combination of standards for the pre-edge feature was used (Beck et al. 2012). Therefore the average valence calculated for our samples is the weighted arithmetic mean of the used standards.

Our suite of standards includes iron metal (Fe^0), San Carlos olivine (Fe^{2+}), greenalite (Fe^{2+}), fayalite (Fe^{2+}), troilite (Fe^{2+}), pyrite (Fe^{2+}), cronstedtite ($\text{Fe}^{2+}+\text{Fe}^{3+}$), magnetite ($1/3 \text{Fe}^{2+} + 2/3 \text{Fe}^{3+}$), ferrihydrite (Fe^{3+}) and hematite (Fe^{3+}). For a full description of these standards see Beck et al. (2012).

3. Results

3.1 Fe-XANES spectra of bulk carbonaceous chondrites

Representative Fe-XANES spectra of CI, CM, CR and CV chondrites are compared in Figure 1a. They reveal the diversity of iron oxidation and Fe-bearing phases in these four carbonaceous chondrite groups. The Fe-XANES spectra of standards for Fe^0 , Fe^{2+} and Fe^{3+} are in Figure 1b for comparison. An increase in energy of the edge position (around intensity of 0.9) can be used as a first way to assess the average oxidation state of these samples. In contrast to the Urey-Craig diagram (Urey and Craig 1953), this study combines the information of the three oxidation states in bulk sample. The value here is then strongly dependent on the complexity of the mineralogy and the co-existing phases, possibly evolving at the same time but in different ways than the oxidation state.

As expected, the CI chondrite (Orgueil) XANES spectra reveal the presence of oxidized Fe-bearing minerals. The CV chondrite (Axtell) appears to be the most reduced, while the CM

(MET 01070) and the CR (Renazzo) appear to have similar edge positions. This position is strongly dependent on the metal contained in CRs chondrites. The energy position of the maxima of these spectra appears to increase with the abundance of metal, similarly to the behavior observed for the edge position.

The pre-edge region is valuable in deciphering the average valence of iron (Wilke et al. 2001). The spectra of the CR chondrites are distinct from the other spectra by showing an enhanced pre-edge intensity (Figure 1a), which is typical of the presence of some amount of metallic iron (Fe^0) (Figure 1b). The pre-edge region of Orgueil (CI) also shows an important pre-edge intensity, but the position (around 7114 eV) and shape of the feature is more consistent with Fe^{3+} rather than metallic iron (Figure 1b). The pre-edge features of the CM and the CV both show two peaks in this spectral region, but with different relative intensities (see inset in Fig. 1a).

Spectra obtained from CR chondrites reveal an important diversity in the intensity of the pre-edge features (Fig. 2). This feature is related to variable amounts of metal (Weisberg et al. 1993), the largest amount being in MET 00426. The spectrum of the CR1 GRO 95577 shows a much smaller pre-edge intensity with a maximum around 7114 eV, typical of the presence of Fe^{3+} . The pre-edge feature of Renazzo is intermediate between GRO 95577 and MET 00426.

Spectral variability among CMs (Fig. 3) is less obvious than for CRs. This is easily explained by the absence of significant amounts of metal (Rubin et al. 2007; Scott and Krot 2014). Still some variation is seen in terms of the edge position, the energy at the absorption maxima, and the pre-edge spectral shape. QUE 97990 (CM 2.6) seems to have the most oxidized mineralogy, MET 01070 (CM 2.0) appears intermediate, and finally PCA 02012 (heated CM) seems to have the most reduced Fe-bearing mineralogy (Fig. 3).

3.2 The average iron oxidation index

From the linear decomposition analysis of the pre-edge region, the proportion of each iron oxidation state, an average iron oxidation index and the $\text{Fe}^0/\text{Fe}_{\text{tot}}$ to $(\text{Fe}^{3+}/\text{Fe}^{2+})$ ratio were estimated (Table 1). The average “Iron Oxidation Index” (IOI) was calculated by the following equation:

$$\text{IOI} = ([\text{Fe}^0] + 2[\text{Fe}^{2+}] + 3[\text{Fe}^{3+}]) / [\text{Fe}_{\text{TOT}}]$$

This parameter is strictly equivalent to the bulk valence used for instance by Sutton et al. (2017). However, we prefer to refer to it as “iron oxidation index” since a non-integer valence is not consistent with chemical rules.

The average IOI varies between 1.60 (CR chondrite GRA 06100) and 2.61 (CI chondrite Orgueil) among our samples and increases along the sequence CR < CV < CM < CI. This pattern is consistent with the higher abundance of metal in CR chondrites (8 vol. %, Krot et al. 2007) than in CI, CM and CV chondrites. Moreover, extensively aqueously altered CI and CM chondrites have the highest valence of iron.

The bulk valence derived by Sutton et al. (2017) with a similar method are in good agreement with the present IOI. Indeed, for Orgueil we obtained an IOI of 2.61 in comparison to 2.77 (Sutton et al. 2017), and for Renazzo 1.85 in comparison to 1.91 (Sutton et al. 2017). The IOI we derived for the six CM chondrites analysed in both studies appear to be lower with an average of 2.36 against 2.49 in Sutton et al. (2017). Note that the fitting methods differ between these two studies and our standard-based fits are performed in the pre-edge region only. Indeed the higher energy part is sensitive to interaction with neighboring atoms, and thus less related to the iron valency. The values obtained for the eight samples common to the two studies show a good correlation with an R^2 value of 0.90 (slope=0.99).

3.3 $\text{Fe}^0/\text{Fe}_{\text{tot}}$ to $(\text{Fe}^{3+}/\text{Fe}^{2+})$

The matrix, being made of fine-grained material, is the petrological component most sensitive to aqueous alteration (e.g. McSween 1979; Browning et al. 1996). The abundance of matrix in each bulk sample may then affect the IOI. When comparing the metal proportion against the fraction of oxidized iron (Figure 4.b), a clear separation between the different chondrite groups appears. The CMs plot along the $\text{Fe}^{3+}/\text{Fe}^{2+}$ axis with ratios varying between 0.3 and 1.4, the CRs on the top of the diagram along $\text{Fe}^0/\text{Fe}_{\text{tot}}$ with ratios varying between 16 and 22% and the CVs are on the lower part of the diagram, closest to 0 for both ratios. CIs plot on the $\text{Fe}^{3+}/\text{Fe}^{2+}$ axis with a ratio of 1.47 and 0.70 for Orgueil and Ivuna, respectively. In contrast to the IOI, this diagram allows a clear distinction between CV and CR chondrites. This distinction results from a combination of higher proportions of Fe^{3+} and metal in CR chondrites than in CV chondrites. The $\text{Fe}^0/\text{Fe}_{\text{tot}}$ ratio is also systematically higher in heated CMs than in non-heated CMs (metal was detected in five heated chondrites) (Fig 4.b)

The variability in IOI between and within chondrite groups is discussed in the following section

4 Discussion

4.1 Group systematics, redox record and H₂ production in unheated CI, CM and CR chondrites.

The interaction between an unaltered carbonaceous chondrite and water is expected to enrich it in Fe³⁺ due to the corrosion of metallic iron. The dissolution of Fe²⁺-bearing anhydrous silicates will be followed by the precipitation of Fe²⁺- and Fe³⁺-bearing phases (e.g. (oxy)hydroxide and phyllosilicates). While Fe is oxidized, water molecules can be reduced. This reaction is a possible source of molecular hydrogen (H₂), and can be associated with hydrogen isotopic fractionation. Fingerprints of this process have been possibly identified in ordinary chondrites in the form of D/H enrichment of the IOM (Insoluble Organic Matter) (Eiler and Kitchen 2004; Alexander et al. 2007; Sutton et al. 2017).

The CI chondrites that have experienced the most intense aqueous alteration tend to have the most oxidized iron mineralogy (Fig. 4). In particular CI and CM chondrites, which are all classified as types 1 or 2 (Rubin et al. 2007; Scott and Krot 2014), have the highest average IOI (Fig 4.a). The most oxidized chondrite is Orgueil. However the two CIs, Orgueil and Ivuna have very different Fe³⁺ contents (61% and 41%, respectively) (Beck et al., 2012). The iron mineralogy of these meteorites is quite different: Orgueil contains some ferrihydrite and Ivuna does not. This difference could have partly resulted from post-terrestrial alteration (Gounelle and Zolensky 2001; Airieau et al. 2001; wiik 1956). Beck et al. (2012) found that 80% of the iron in Orgueil is present as oxides which is not the case of Ivuna which has a different mineralogy (see also Beck et al. 2014).

While several proxies and aqueous alteration schemes have been defined for carbonaceous chondrites, a convenient quantitative tracer of aqueous alteration extent is the amount of H₂O and –OH trapped in minerals (Howard et al. 2009; Howard et al. 2011; Alexander et al. 2013). IOI seems to be roughly correlated to the bulk H abundance in CI, CR and CM chondrites as measured by Alexander et al. (2012) (Figure 5): the more aqueously altered the sample is, the more oxidized the iron is and CI, CM and CR chondrites plot apart. An important spread is observed within CMs and for these samples the IOI appears to be anticorrelated with aqueous alteration. The CV chondrites are not shown on Figure 5 since H data are only available for one sample (Kaba, that falls outside of this trend). While a general increase of H -content with the valence of iron is expected, the presence of a single trend for CR, CI and CM (average) is more surprising. The slope on this diagram (Fig. 5) is related to the redox reaction leading to the production of Fe³⁺ and thus depends on the IOI of the sample prior to alteration and the nature of Fe-bearing secondary products. Since the secondary alteration products and their proportions are different among CI, CM and CR chondrites (Scott

and Krot 2014; Bonal et al. 2013; Abreu 2016), the presence of a single trend is unexpected (i.e. abundant magnetite in CIs, but rare in CMs, amorphous secondary phases in CR, higher serpentine and saponite proportion in CM and CI than CR). We cannot exclude a statistical effect due the few CI and CR chondrites in our study compared to CM chondrites. A more complete study of CIs and CRs is necessary to determine whether or not this is a single trend.

During aqueous alteration, H₂ production might occur on the asteroidal parent body. The production of H₂ can be a first step before the synthesis of complex organics through Fischer-Tropsch Type reactions. This H₂ production can also impact the evolution of asteroids since pressure build-up could lead to partial or total disruption even if a fast H₂ degassing is possible (Guo and Eiler 2007; Wilson 1999).

4.2 The reduction of Fe during aqueous alteration of CM chondrites

CM chondrites have all experienced aqueous alteration, but to various extents (e.g. Zolensky et al. 1997; Rubin et al. 2007). While it is not proven that all CMs come from a single parent body (many large CM-like asteroids are present in the main-belt, Fornasier et al. 2014; Vernazza et al. 2016), these variable degrees of aqueous alteration are usually seen as a progression, from mildly altered (unaltered mafic silicate phenocrysts in chondrules, rare metal grains preserved) to heavily altered (fully altered chondrules) (Rubin et al. 2007).

A majority of the considered CMs are Antarctic finds. The Fe valence could have been modified through terrestrial weathering (Lee and Bland 2004; Bland 2006). The average valence of iron is almost identical for CM falls (0.38, n=8) and CM finds (0.36, n=16), suggesting that terrestrial residence had a minimal impact on the Fe valence. However, we could not exclude a possible minor impact, hidden by the spread and variation of Fe-valence of these samples due to different degree of aqueous alteration.

Focusing on the CMs, the average Fe oxidation factor is compared to the hydrogen content (Alexander et al. 2012) (Fig. 5) and to the intensity of the 3- μ m band (Fig. 6) that is related to -OH and H₂O. In both cases, the average IOI appears to decrease with increasing degree of aqueous alteration or more precisely, water content. This pattern shows that iron is reduced as aqueous alteration progresses in CM chondrites. The decrease of the average IOI with the extent of aqueous alteration in CM chondrites (Fig. 6) is at first unexpected. A similar effect has been observed on CR matrix chondrites by Le Guillou et al. (2015). As stated earlier, in the case of the aqueous alteration of terrestrial ultramafic rocks, the average Fe³⁺ content increases with the degree of serpentinization (Marcaillou et al. 2011). However, CM chondrites are different from terrestrial ultramafic rocks that their protolith contained iron

metal, and quite unusual alteration products were formed, such as tochilinites (an hydroxysulfide consisting of interlayer mackinawite and brucite layers), as well as cronstedtite.

To understand the reduction of iron, it is important to consider the chemistry of iron-bearing alteration products. Cronstedtite is a Fe-rich serpentine, which contains a significant amount of Fe^{3+} in tetrahedral sites (Buseck and Hua 1993; Beck et al. 2010). Cronstedtite is abundant in the least-altered CM chondrites (Howard et al. 2011), but as aqueous alteration progresses, cronstedtite is progressively replaced by Mg-rich serpentine (Browning et al. 1996) as revealed by transmission electron microscopy observations. Mg-rich serpentine may contain Fe^{3+} , but not to the level of cronstedtite (Burns and Fisher 1994). This mineralogical evolution has been explained by the following sequence (Howard et al. 2011; Le Guillou et al. 2015): i) formation of cronstedtite from alteration of amorphous silicate and metal; ii) alteration of chondrules (Mg- and Si-rich) and increase of silica activity; iii) dissolution of cronstedtite and precipitation of Mg-rich serpentine and sulfides, iv) iron exchange from cronstedtite to serpentine in chondrules and substitution of iron by magnesium in cronstedtite.

The reduction of iron during aqueous alteration seems to be in agreement with this aqueous alteration sequence. In the least altered CM chondrites a significant fraction of Fe^{3+} can be accommodated in cronstedtite which is much less abundant in the most-altered CMs. In the latter case, the mineralogy is dominated by Mg-serpentine (which can host only a limited amount of Fe^{3+}), and Fe^{2+} -bearing sulphides (Howard et al. 2009; Howard et al. 2011; Howard et al 2016).

The fact that on average the IOI decreases with aqueous alteration requires that iron should be reduced at the expense of other elements being oxidized (if the evolution occurs in closed-system). There are many possible reducing agents, the first one being dissolved H_2 . The production of H_2 was likely significant in the first stages of aqueous alteration (from corrosion of metals) and an increase in hydrogen fugacity will destabilize cronstedtite toward Fe^{2+} -bearing serpentine greenalite (Zolotov 2014). Locally produced H_2 could therefore act as a reducing agent, under the assumption that it does not escape. A system under pressure would have facilitated the buildup of an H_2 overpressure. However, closed system alteration is an approximation and does not take into account partial escape of highly diffusing gases like H_2 (Krot et al. 2015).

An alternative scenario to explain the reduction of iron would be interaction with nebular H_2 . But this would require not only that aqueous alteration took place before nebular gas dissipation (which contradicts Mn/Cr chronology of alteration products, Fujiya et al. 2012), but also that some mechanism would permit interaction of the subsurface of the asteroid and

the surface. Therefore a reaction with nebular H_2 is unlikely. An alternative scenario is internal oxidation of different compounds on the asteroid surface reduce iron. It could be elemental sulfur, sulfides or oxysulfide into sulfates, though the sulfate amounts are likely low in carbonaceous chondrites (estimated between 0.4 and 1 wt. %; Burgess et al. 1991). Note that the sulfide abundance appears to decrease with extent of aqueous alteration (Burgess et al. 1991; Orthous-Daunay et al. 2010; Brearley 2011; Singerling and Brearley 2016). Finally, other compounds such as organic matter or dissolved CO could have acted as possible iron reducing agents in CMs.

The hydrolysis of iron metal can produce significant isotopic fractionation if the system is opened to H_2 escape (Alexander et al. 2010). In Figure 7, the oxidation factor of iron is compared to the bulk D/H measured in carbonaceous chondrites. This bulk value is the average of all H-bearing phases in carbonaceous chondrites, i.e. H_2O and -OH bearing minerals, soluble and refractory organics. In the case of CM chondrites, a positive correlation is found between bulk D/H and iron oxidation index. The simplest explanation is that this diagram reflects mixing between phyllosilicates and organics, as discussed in Alexander et al. (2012). This would mean that the D/H of water in CR and CI chondrites is distinct from the water in CM chondrites (Alexander et al. 2012). An alternative would be that this trend is related to a chemical process, which changes both the isotopic composition of water and the IOI. A possible process is the reaction between water and nebular H_2 (with a low D/H), but this seems challenging given the reasons discussed above.

4.3 Heated CM and the redox memory

Among the suite of CM chondrites studied here, some are described as heated (Table 1). The existence of such samples was initially reported following the Japanese meteorite recovery campaigns in Antarctic (Akai 1992). While the petrology points toward affinities with CM chondrites, specific mineralogical changes demonstrate that these meteorites have undergone heating, based on mineralogy, texture, oxygen isotopes or organic matter (Nakamura 2005). The most likely source of thermal metamorphism is impact heating (Nakamura 2005). When considering the IOI, the majority of heated CM samples are indistinguishable from mainstream CM chondrites. However, when considering Fe^0/Fe_{tot} , it appears that half of heated CMs contain metallic irons in contrast to unheated CMs (Table 1). Two (paired) meteorites are quite distinct by their high metal abundance, PCA 02010 and PCA 02012. PCA 02010 and PCA 02012 appear to be the most reduced CM chondrites with the highest amount of Fe^0 (7-9 at. % of Fe). These samples may have experienced more intense heating than the other heated CMs studied here

(Nakato et al. 2013; Beck et al. 2014) after they were originally hydrated (Nakamura 2005). It is difficult to determine whether the metal in heated CMs is a relic of unaltered initial iron or a product of metal formation after heating (Nozaki et al 2006). In the case of CV chondrites like Allende, the fluid-assisted thermal metamorphism seems to have oxidized the initial metal to Fe^{2+} (Krot et al. 2004; Lee et al. 1996). With the exception of the proportion of metal, the Fe^{2+} and Fe^{3+} budgets of heated CMs are similar to those in unheated CMs. It seems that for low levels of thermal alteration, the oxidized nature of iron is preserved, and an "iron oxidation state memory" of the aqueous process is present.

4.4 CV chondrites, discrepancy between iron valence and the red-ox dichotomy classification

All CV chondrites are classified as type 3, but they have experienced various degrees of aqueous alteration and thermal metamorphism. They have been classified into two subgroups, so called "oxidized" (CV_{Ox}) and "reduced" (CV_{Red}), based on the relative abundance of metal and magnetite, as well as the Ni content of metal and sulfides (Mcsween 1977). Oxidized CV chondrites were later subdivided into CV_{OxA} (Allende subgroup) and CV_{OxB} (Bali subgroup) (Weisberg et al. 1997), where CV_{OxA} typically have higher metal content and lower matrix abundance than CV_{OxB} .

Among the studied samples, Efremovka, a reduced CV, contains the highest amount of metal, and no Fe^{3+} , while Kaba, classified as a CV_{OxB} , does indeed contain a high abundance of Fe^{3+} , the highest among our series of CVs (Table 1). However, considering the Fe oxidation factor, many samples do not follow the general classification scheme of CV oxidized and reduced (Table 1). Indeed, Vigarano (CV_{Red}) seems to have a lower metal abundance and higher Fe^{3+} content than Bali (CV_{OxB}). Also, from the Fe oxidation factor perspective, Allende (CV_{OxA}) and Grosnaja (CV_{OxB}) seem to be similar. The iron oxidation in CV chondrites seems more complex than other carbonaceous chondrites. One explanation is that these materials have been altered by both aqueous and thermal metamorphism. The intensities of these geological processes are not the only parameters to take account, their chronological order too (Lee et al. 1996).

The oxidized/reduced classification is based on magnetite and metal contents only: other iron phases are not taken into account. These discrepancies have been already observed (Howard et al. 2010) and could result from iron being present in other mineral phases such as phyllosilicates. Also some of the observed variability falls within our analytical uncertainty on

the retrieval of the various proportions of iron species ($\pm 10\%$). The Fe oxidation state appears to be controlled by the thermal metamorphism and petrologic type proposed by Bonal et al. (2006).

4.5 The oxidized matrix of CR chondrites

The presence of a high abundance of presolar grains and the preservation of amorphous material in the matrix shows that some CR chondrites are amongst the most primitive meteorites in our collections (Floss and Stadermann 2009; Abreu and Brearley 2010). All of the CR chondrites in this study are classified as type 2 with the exception of GRO 95577, classified as a CR1 (Weisberg and Huber 2007; Harju et al. 2014). The CR1 sample is clearly distinct from the other studied CRs in terms of bulk H vs average iron oxidation index and falls within the array defined by CM chondrites (Fig. 5). This is not unexpected given the fact that GRO 95577 is the only CR sample that experienced extensive aqueous alteration..

The analysis of Fe-XANES spectra from our sample suite (excluding GRO 95577) shows that each CR contains iron metal, typically around 20 at. % of total iron. RBT 04133 is significantly different with a much smaller fraction of metal (5.6 at. %). This sample was previously described as anomalous by Davidson et al. (2009) from its oxygen isotope composition, with closer affinities to CV-CK than CR chondrites. It is metamorphosed (type 3.6 proposed by Bonal et al. 2016) and was proposed to be reclassified as CV_{red} (Davidson et al. 2014). With the exception of GRA 06100, all CR samples have a significant fraction of iron present as Fe³⁺. Note that GRA 06100 has been previously identified as a metamorphosed CR given its mid-infrared spectrum (Abreu and Bullock 2013; Briani et al. 2013; Beck et al. 2014).

In contrast to CVs, the CR chondrites have an important proportion of Fe³⁺, closer to CMs, probably resulting from aqueous alteration (Le Guillou et al. 2015). While this might appear surprising given that CM chondrites are generally considered more altered than CR chondrites, our measurements are probing iron, whose abundance is higher in the matrix than in chondrules. From mass balance consideration, Fe-XANES measurements are thus likely sensitive to the alteration of the matrix. Using L-edge Fe XANES spectroscopy, Le Guillou et al. (2015) were able to decipher the valence state of iron in amorphous silicates and phyllosilicates of CR chondrites. Their work showed that iron is substantially oxidized in this phase (Fe³⁺ / (Fe_{tot}) around 0.7). These values are higher than found in the present work (Fe³⁺ / (Fe_{tot}) around 0.22), but our measurements were obtained on bulk CR. The measurements obtained here include the contributions of other phases like Fe²⁺-bearing sulfides in the matrix, mixed Fe²⁺, Fe³⁺ in magnetite as well as possible Fe²⁺ phase in chondrules. With the exception

of GRA 06100, Renazzo and LAP 04156 contain the lowest Fe^{3+} proportion but also the lowest metal abundance. If the initial proportion of metal were identical in CR chondrites, this points to different oxidation conditions that produced different amount of Fe^{2+} and Fe^{3+} than other CRs.

These results confirm that iron in the matrix of CR chondrites is significantly oxidized, and water was likely the oxidative agent (Le Guillou et al. 2015). The fact that CR chondrites matrix contain a significant fraction of H_2O and $-\text{OH}$ bearing phases is now established (Alexander et al. 2013; Bonal et al. 2013; Garenne et al. 2014; Le Guillou et al. 2015, Le Guillou and Brearley 2014). Even in MET 00426, described as the most primitive CR chondrite in our dataset (the least aqueously altered) (Abreu and Brearley 2010; Le Guillou et al. 2015), the iron is significantly oxidized (Fig. 4). The pristine feature of MET 00426 matrix is debated (Bonal et al. 2013) and could have preserved those unstable phases (Scott and Krot 2005). However the two techniques (TEM and IR) are probing the samples at different scales. More generally the origin of meteorite matrices is debated, but a number of arguments seem to point toward a formation as disequilibrium condensates at least for the most primitive samples (Brearley 1993; Scott and Krot 2005, Abreu and Brearley 2010). It remains to be understood whether during the matrix formation process significant amount of Fe^{3+} may have been incorporated into the amorphous phases, together with water. Formation of hydrated amorphous silicates in the disk does not appear to be an efficient mechanism (Prinn and Fegley 1989) to explain Fe^{3+} phases in carbonaceous chondrites. The fact that GEMS grains, that have mineralogical affinities with the amorphous matrices of primitive carbonaceous chondrites are depleted in Fe^{3+} (Keller and Messenger 2012; Hopp and Vollmer 2018; Vollmer et al. 2018), argues against hydration in the disk. Therefore, the hydrated and oxidized amorphous matrix observed here in CR chondrites likely represents the very first step of aqueous alteration on the parent body (Le Guillou and Brearley 2014; Brearley 2014; Le Guillou et al. 2015).

5 Conclusion

The present study evaluates the bulk iron oxidation as well as abundance of Fe^0 , Fe^{2+} and Fe^{3+} in CM, CR, CI and CV chondrites, based on XANES spectroscopy.

-A general trend between the iron oxidation index and the hydrogen content is observed for the different groups of carbonaceous chondrites. Fe is on average more oxidized in the order of CRs < CMs < CIs

-It appears that, after the first aqueous alteration stage, extensive aqueous alteration reduces the iron oxidation state of CM chondrites. This is most likely due to the dissolution and recrystallization of cronstedtite to serpentine.

-Based on the iron oxidation, it is not possible to distinguish CM chondrites finds from falls. There is no evidence that terrestrial residence had an impact on the speciation of Fe for our set of CMs.

-The heated CMs are not clearly distinguishable from the other CMs in terms of their Fe²⁺ vs Fe³⁺ relative abundance, but they generally contain a higher proportion of metal. There is also redox memory of the aqueous alteration process.

-The CR chondrites present a wide range of oxidation of iron (this could be compared to CM chondrites) controlled by the extent of aqueous alteration. Fe³⁺ is present even in the least-altered CR chondrites.

- The iron oxidation index of CV chondrites is not consistent with the reduced/oxidized sub-classification. Rather, the iron oxidation of CVs seems strongly related to their thermal alteration.

Acknowledgment—This study was supported through funding from the H2020 European Research Council (ERC) (SOLARYS ERC-CoG2017_771691).

References

- Abreu N. M., and Brearley A. J. 2010. Early solar system processes recorded in the matrices of two highly pristine CR3 carbonaceous chondrites, MET 00426 and QUE 99177. *Geochimica et Cosmochimica Acta* 74:1146–1171.
- Abreu N. M., and Bullock E. S. 2013. Opaque assemblages in CR2 Graves Nunataks (GRA) 06100 as indicators of shock-driven hydrothermal alteration in the CR chondrite parent body. *Meteoritics & Planetary Science* 48:2406–2429.
- Abreu N. M. 2016. Why is it so difficult to classify Renazzo-type (CR) carbonaceous chondrites? - Implications from TEM observations of matrices for the sequences of aqueous alteration. *Geochim. Cosmochim. Acta* 194:91-122
- Airieu S., Farquhar J., Jackson T. L., Leshin L. A., Thiemens M. H., and Bao H. 2001. Oxygen isotope systematics of Cland CM chondritic sulphate: Implications for evolution and mobility of water in planetesimals. *Lunar Planet. Sci.* 32:#1744 (abstr.).
- Akai J. 1992. T-T-T diagram of serpentine and saponite and estimation of metamorphic heating degree of Antarctic carbonaceous chondrites. In: Proc. NIPR Symp. *Antarct. Meteorit.* 5:120–135.
- Alexander C. M. O., Fogel M., Yabuta H., and Cody G. D. 2007. The origin and evolution of chondrites recorded in the elemental and isotopic compositions of their macromolecular organic matter. *Geochimica et Cosmochimica Acta* 71:4380–4403.
- Alexander C. M. O., Newsome S. D., Fogel M. L., Nittler L. R., Busemann H., and Cody G. D. 2010. Deuterium enrichments in chondritic macromolecular material- Implications for the origin and evolution of organics, water and asteroids. *Geochimica et Cosmochimica Acta* 74:4417–4437.
- Alexander C. M. O., Bowden R., Fogel M. L., Howard K. T., Herd C. D. K., and Nittler L. R. 2012. The provenances of asteroids, and their contributions to the volatile inventories of the terrestrial planets. *Science* 337:721–723.
- Alexander C. M. O., Howard K. T., Bowden R., and Fogel M. L. 2013. The classification of CM and CR chondrites using bulk H, C and N abundances and isotopic compositions. *Geochimica et Cosmochimica Acta* 123:244-260.
- Beck P. et al. 2010. Hydrous mineralogy of CM and CI chondrites from infrared spectroscopy and their relationship with low albedo asteroids. *Geochimica et Cosmochimica Acta* 74:4881–4892.
- Beck P., De Andrade V., Orthous-Daunay F.-R., Veronesi G., Cotte M., Quirico E., and Schmitt B. 2012. The redox state of iron in the matrix of CI, CM and metamorphosed CM chondrites by XANES spectroscopy. *Geochimica & Cosmochimica Acta* 99:305–316.
- Beck P., Garenne A., Quirico E., Bonal L., Montes-Hernandez G., Moynier F., and Schmitt B. 2014. Transmission infrared spectra (2–25 μm) of carbonaceous chondrites (CI, CM, CV–CK, CR, C2 ungrouped): Mineralogy, water, and asteroidal processes. *Icarus* 229:263–277.

- Bischoff A. 1998. Aqueous alteration of carbonaceous chondrites: Evidence for preaccretionary alteration-A review. *Meteoritics & Planetary Science* 33:1113–1122.
- Bland P. 2006. Terrestrial weathering rates defined by extraterrestrial materials. *J Geochem Explor.* 88:257-261.
- Bonal L., Quirico E., Bourot-Denise M., and Montagnac G. 2006. Determination of the petrologic type of CV3 chondrites by Raman spectroscopy of included organic matter. *Geochimica et Cosmochimica Acta* 70:1849–1863.
- Bonal L., Alexander C. M. O., Huss G. R., Nagashima K., Quirico E., and Beck P. 2013. Hydrogen isotopic composition of the water in CR chondrites. *Geochimica et Cosmochimica Acta* 106:111–133.
- Bonal L., Quirico E., Flandinet L., and Montagnac G. 2016. Thermal history of type 3 chondrites from the Antarctic meteorite collection determined by Raman spectroscopy of their polyaromatic carbonaceous matter. *Geochimica et Cosmochimica Acta* 189:312–337.
- Brearley A. J. 1993. Matrix and fine-grained rims in the unequilibrated CO3 chondrite, ALHA77307 - Origins and evidence for diverse, primitive nebular dust components. *Geochimica et Cosmochimica Acta* 57:1521–1550.
- Brearley A. J. 2006. The action of water. In *Meteorites and the Early Solar System II*, edited by Binzel R. P., Lauretta D. S., and McSween Jr H. Y. Tucson: The University of Arizona Press. pp. 587–623.
- Brearley A.J. 2011. Alteration of coarse-grained Fe and Fe, Ni sulfides in the Mighei CM2 carbonaceous chondrite: Evidence for the instability of primary pyrrhotite-pentlandite grains during aqueous alteration. *Lunar Planet. Sci.* 42:#2233 (abstr.).
- Brearley A. J. and Krot A. N. 2013. Metasomatism in the early Solar System: the record from chondritic meteorites. In *Metasomatism and the Chemical Transformation of Rock* (eds. D. E. Harlov and H. Austrheim). Springer, Heidelberg, pp. 659–790.
- Brearley A. J. 2014. Nebular Versus Parent Body Processing. In *Meteorites and Cosmochemical Processes*, edited by Davis A. M. pp. 309–334.
- Briani G. et al. 2013. Short duration thermal metamorphism in CR chondrites. *Geochimica et Cosmochimica Acta* 122:267–279.
- Browning L.B., McSween Jr.H.Y.and Zolensky M.E. 1996. Correlated alteration effects in CM carbonaceous chondrites. *Geochimica et Cosmochimica Acta* 60: 2621–2633.
- Burgess R., Wright I. P., and Pillinger C. T. 1991. Determination of sulphur-bearing components in C1 and C2 carbonaceous chondrites by stepped combustion. *Meteoritics* 26:55–64.

- Burns R., and Fisher D. 1994. Nanophase mixed-valence iron minerals in meteorites identified by cryogenic Mössbauer spectroscopy. *Hyperfine Interactions* 91:571–576.
- Buseck P. R., and Hua X. 1993. Matrices of carbonaceous chondrite meteorites. *Annual Review of Earth and Planetary Sciences* 21:255–305.
- Cloutis E. A., Hudon P., Hiroi T., Gaffey M. J., and Mann P. 2011a. Spectral reflectance properties of carbonaceous chondrites: 2. CM chondrites. *Icarus* 216:309–346.
- Cloutis E. A., Hiroi T., Gaffey M. J., Alexander C. M. O., and Mann P. 2011b. Spectral reflectance properties of carbonaceous chondrites: 1. CI chondrites. *Icarus* 212:180–209.
- Cloutis E. A., Hudon P., Hiroi T., and Gaffey M. J. 2012a. Spectral reflectance properties of carbonaceous chondrites: 3. CR chondrites. *Icarus* 217:389–407.
- Cloutis E. A., Hudon P., Hiroi T., Gaffey M. J., Mann P., and Bell J. F. 2012b. Spectral reflectance properties of carbonaceous chondrites: 6. CV chondrites. *Icarus* 221:328–358.
- Davidson J., Schrader D. L., Busemann H., Franchi I. A., Connolly Jr H. C., Lauretta D. S., Alexander C. M. O'D., Verchovsky A., Gilmour M. A., Greenwood R. C. and Grady M. M. 2009. RBT 04133: A new, unusual carbonaceous chondrite. *Meteoritics and Planetary Science Supplement* 72:5141.
- Davidson J., Schrader D. L., Alexander C. M. O'D., Lauretta D. S., Busemann H., Franchi I. A., Greenwood R. C., Connolly Jr H. C., Domanik K.J. and Verchovsky A. 2014. Petrography, stable isotope compositions, microRaman spectroscopy, and presolar components of Roberts Massif 04133: A reduced CV3 carbonaceous chondrite. *Meteoritics & Planetary Science* 49:2133–2151.
- Eiler J. M., and Kitchen N. 2004. Hydrogen isotope evidence for the origin and evolution of the carbonaceous chondrites. *Geochimica et Cosmochimica Acta* 68:1395–1411.
- Floss C., and Stadermann F. 2009. Auger Nanoprobe analysis of presolar ferromagnesian silicate grains from primitive CR chondrites QUE 99177 and MET 00426. *Geochimica et Cosmochimica Acta* 73:2415–2440.
- Fornasier S., Lantz C., Barucci M. A., and Lazzarin M. 2014. Aqueous alteration on main belt primitive asteroids: Results from visible spectroscopy. *Icarus* 233:163–178.
- Fujiya W., Sugiura N., Hotta H., Ichimura K., and Sano Y. 2012. Evidence for the late formation of hydrous asteroids from young meteoritic carbonates. *Nature Communications* 3:627.
- Garenne A., Beck P., Montes-Hernandez G., Chiriac R., Toche F., Quirico E., Bonal L., and Schmitt B. 2014. The abundance and stability of “water” in type 1 and 2 carbonaceous chondrites (CI, CM and CR). *Geochimica et Cosmochimica Acta* 137:93–112.

- Gounelle M. and Zolensky M.E. 2001. A terrestrial origin for sulfate veins in CI1 chondrites. *Meteoritics & Planetary Science* 36:1321-1329.
- Guo W., and Eiler J. M. 2007. Temperatures of aqueous alteration and evidence for methane generation on the parent bodies of the CM chondrites. *Geochimica et Cosmochimica Acta* 71:5565–5575.
- Harju E. R., Rubin A. E., Ahn I., Choi B.-G., Ziegler K., and Wasson J. T. 2014. Progressive aqueous alteration of CR carbonaceous chondrites. *Geochimica et Cosmochimica Acta* 139:267–292.
- Hayatsu, R. and Anders, E. 1981. Organic compounds in meteorites and their origins. *Topics Current Chem.* 99: 1–37.
- Hopp T. and Vollmer C. 2018. Chemical composition and iron oxidation state of amorphous matrix silicates in the carbonaceous chondrite Acfer 094. *Meteorit Planet Sci.* 53: 153-166.
- Howard K. T., Benedix G. K., Bland P. A., and Cressey G. 2009. Modal mineralogy of CM2 chondrites by X-ray diffraction (PSD-XRD). Part 1: Total phyllosilicate abundance and the degree of aqueous alteration. *Geochimica et Cosmochimica Acta* 73:4576–4589.
- Howard K. T., Benedix G. K., Bland P. A., and Cressey G. 2010. Modal mineralogy of CV3 chondrites by X-ray diffraction (PSD-XRD). *Geochimica et Cosmochimica Acta* 74:5084–5097.
- Howard K. T., Benedix G. K., Bland P. A., and Cressey G. 2011. Modal mineralogy of CM chondrites by X-ray diffraction (PSD-XRD): Part 2. Degree, nature and settings of aqueous alteration. *Geochimica et Cosmochimica Acta* 75:2735–2751.
- Howard K. T., Alexander C. M. O. 'D., Schrader D. L., and Dyl K. A. 2015. Classification of hydrous meteorites (CR, CM and C2 ungrouped) by phyllosilicate fraction: PSD-XRD modal mineralogy and planetesimal environments. *Geochimica et Cosmochimica Acta* 149:206–222.
- Keller L. P., and Messenger S. 2012. Formation and Processing of Amorphous Silicates in Primitive Carbonaceous Chondrites and Cometary Dust. *Lunar Planet. Sci.* 43:#1880 (abstr.).
- King A. J., Schofield P. F., Howard K. T., and Russell S. S. 2015. Modal mineralogy of CI and CI-like chondrites by X-ray diffraction. *Geochimica et Cosmochimica Acta* 165:148–160.
- Krot A. N., Fegley B. Jr., Lodders K., and Palme H. 2000. Meteoritical and astrophysical constraints on the oxidation state of the solar nebula. *Protostars and Planets IV* 1019-1054.

- Krot A. N., Petaev I. M. and Bland P. A. 2004. Multiple formation mechanisms of ferrous olivine in CV carbonaceous chondrites during fluid-assisted metamorphism. *Antarct. Meteorit. Res.* 17:153–171.
- Krot A. N., Keil K., Scott E. R. D., Goodrich C. A., and Weisberg M. K. 2007. Classification of meteorites. In *Treatise on Geochemistry*. Elsevier. 1-52
- Krot A. N., Nagashima K., Alexander C. M. O., Ciesla F. J., Fujiya W., and Bonal L. 2015. sources of water and aqueous activity on the chondrite parent asteroids. In *Asteroids IV*. Tucson, AZ, University of Arizona Press, 635–660
- Lee M. R., Hutchinson R. and Graham A. L. 1996. Aqueous alteration in the matrix of the Vigarano (CV3) carbonaceous chondrite. *Meteoritics and Planetary Science*. 31:477-483.
- Lee M. R. and Bland P. A. 2004. Mechanisms of weathering of meteorites recovered from hot and cold deserts and the formation of phyllosilicates. *Geochim. Cosmochim. Acta* 68:893–916.
- Le Guillou C. and Brearley A. 2014. Relationships between organics, water and early stages of aqueous alteration in the pristine CR3.0 chondrite MET 00426. *Geochimica et Cosmochimica Acta*. 131:344–367.
- Le Guillou C., Changela H. G., and Brearley A. J. 2015. Widespread oxidized and hydrated amorphous silicates in CR chondrites matrices: Implications for alteration conditions and H₂ degassing of asteroids. *Earth and Planetary Science Letters* 420:162–173.
- Lewis J. S. 1972. Low temperature condensation from the solar nebula. *Icarus* 16:241–252.
- Malvoisin B., Carlut J., and Brunet F. 2012a. Serpentinization of oceanic peridotites: 1. A high-sensitivity method to monitor magnetite production in hydrothermal experiments. *Journal of Geophysical Research: Solid Earth* 117, B01104 :1-10.
- Malvoisin B., Brunet F., Carlut J., Rouméjon S., and Cannat M. 2012b. Serpentinization of oceanic peridotites: 2. Kinetics and processes of San Carlos olivine hydrothermal alteration. *Journal of Geophysical Research: Solid Earth* 117,B04102:1-13.
- Marcaillou C., Muñoz M., Vidal O., Parra T., and Harfouche M. 2011. Mineralogical evidence for H₂ degassing during serpentinization at 300 °C/300 bar. *Earth and Planetary Science Letters* 303:281–290.
- Martinez J.M. and Brearley A.J. 2018. Aqueous alteration effects in chondrules in the most pristine CR2 chondrite, Queen Alexandra Range (QUE)99177: Further insights into the earliest stages of fluid-rock interaction on the CR chondrite parent body. *Lunar Planet. Sci.* 49:#2274 (abstr.).

- McCollom T. M., and Bach W. 2009. Thermodynamic constraints on hydrogen generation during serpentinization of ultramafic rocks. *Geochimica et Cosmochimica Acta* 73:856–875.
- McSween H. Y. 1977. Petrographic variations among carbonaceous chondrites of the Vigarano type. *Geochimica et Cosmochimica Acta* 41:1777–1790.
- McSween Jr H. Y. 1979. Alteration in CM carbonaceous chondrites inferred from modal and chemical variations in matrix. *Geochimica et Cosmochimica Acta* 43:1761–1770.
- Nakamura T. 2005. Post-hydration thermal metamorphism of carbonaceous chondrites. *Journal of Mineralogical and Petrological Sciences* 100:260–272.
- Nakato A., Brearley A. J., Nakamura T., Noguchi T., Ahn I., Lee J. I., Matsuoka M., and Sasaki S. 2013. PCA 02012: A unique thermally metamorphosed carbonaceous chondrite. *Lunar Planet. Sci.* 44:#2708 (abstr.).
- Nozaki W., Nakamura T. and Noguchi T. 2006. Bulk mineralogical changes of hydrous micrometeorites during heating in the upper atmosphere at temperatures below 1000 °C. *Meteoritics and planetary science* 41:1095-1114
- Orthous-Daunay F.-R., Quirico E., Lemelle L., Beck P., deAndrade V., Simionovici A., and Derenne S. 2010. Speciation of sulfur in the insoluble organic matter from carbonaceous chondrites by XANES spectroscopy. *Earth and Planetary Science Letters* 300:321–328.
- Prinn R. G., and Fegley B. J. 1989. Solar nebula chemistry: Origin of planetary, satellite, and cometary volatile. In *Planetary and satellite atmospheres: origin and evolution.*, edited by Atreya S. K., Pollack J. B., and Mathews M. S. Tucson, AZ: The University of Arizona Press. pp. 78–136.
- . Proux O., Biquard X., Lahera E., Menthonnex J.J., Prat A., Ulrich O , Soldo Y., Trévisson P., Kapoujyan G., Perroux G., Taunier P., Grand D., Jeantet P., Deleglise M., Roux J.P. and Hazemann J.L. 2005. FAME: a new beamline for x-ray absorption investigations of very-diluted systems of environmental, material and biological interests. *Physica Scripta* T115:970-973.
- Proux O., Nassif V., Prat A., Ulrich O., Lahera E., Biquard X., Menthonnex J.-J., and Hazemann J.-L. 2006. Feedback system of a liquid-nitrogen-cooled double-crystal monochromator: design and performances. *Journal of Synchrotron Radiation* 13:59–68.
- Ravel B., and Newville M. 2005. *ATHENA , ARTEMIS , HEPHAESTUS* : data analysis for X-ray absorption spectroscopy using *IFEFFIT*. *Journal of Synchrotron Radiation* 12:537–541.
- Rivkin A. S., Howell E. S., Lebofsky L. A., Clark B. E., and Britt D. T. 2000. The Nature of M-Class Asteroids from 3- μ m Observations. *Icarus* 145:351–368.

- Rubin A. E., Fegley B., and Brett R. 1988. Oxidation state in chondrites. In *Meteorites and the Early Solar System*. Tucson, AZ, University of Arizona Press, 488–511.
- Rubin A. E., Trigo-Rodriguez J. M., Huber H., and Wasson J. T. 2007. Progressive aqueous alteration of CM carbonaceous chondrites. *Geochimica et Cosmochimica Acta* 71:2361-2382.
- Scott E. R. D., and Krot A. N. 2005. Chondritic meteorites and the high-temperature nebular origins of their components. In *Chondrites and the Protoplanetary Disk*, edited by Krot A. N., Scott E. R. D., and Reipurth B. p. 15-53.
- Scott E. R. D., and Krot A. N. 2014. Chondrites and their components. in *meteorites and cosmochemical processes*. Elsevier, Andrew M. Davis: 65–137.
- Singerling S. A. and Brearley A. J. 2016. Altered primary sulfides in CR and CM carbonaceous chondrites: formation by dissolution and pseudomorphic replacement. *Lunar Planet. Sci.* 47:#1718 (abstr.).
- Sutton S., Alexander C. M. O., Bryant A., Lanzirotti A., Newville M., and Cloutis E. A. 2017. The bulk valence state of Fe and the origin of water in chondrites. *Geochimica et Cosmochimica Acta* 211:115–132.
- Urey H. C., and Craig H. 1953. The composition of the stone meteorites and the origin of the meteorites. *Geochimica et Cosmochimica Acta* 4:36–82.
- Vernazza P., Marsset M., Beck P., Binzel R. P., Birlan M., Cloutis E. A., DeMeo F. E., Dumas C., and Hiroi T. 2016. Compositional homogeneity of CM parent bodies. *The Astronomical Journal* 152:54-64.
- Vollmer C., Leitner J., Kepaptsoglou D., Ramasse Q. M. and Hoppe P. 2018. Iron oxidation state of amorphous silicates and low-loss electron energy spectroscopy of organic matter in the pristine CM Chondrite maribo. *Meteoritical Society*. 81:#6109 (abstr.)
- Weisberg M. K., Prinz M., Clayton R. N., and Mayeda T. K. 1993. The CR (Renazzo-type) carbonaceous chondrite group and its implications. *Geochimica et Cosmochimica Acta* 57:1567–1586.
- Weisberg M. K., Prinz M., Clayton R. N., and Mayeda T. K. 1997. CV3 chondrites: three subgroups, not two. *Meteoritics and Planetary Science Supplement* 32:138.
- Weisberg M. K., McCoy T. J., and Krot A. N. 2006. Systematics and evaluation of meteorite classification. In *Meteorites and the Early Solar System II*. Tucson, University of Arizona Press : 19–52.
- Weisberg M. K., and Huber H. 2007. The GRO 95577 CR1 chondrite and hydration of the CR parent body. *Meteoritics & Planetary Science* 42:1495–1503.
- Wiik H.B. 1956. The chemical composition of some stony meteorites. *Geochim. Cosmochim. Acta* 57:1567-1586

- Wilke M., Farges F., Petit P. E., Brown G. E., and Martin F. 2001. Oxidation state and coordination of Fe in minerals; an Fe-XANES spectroscopic study. *American Mineralogist* 86:714–730.
- Wilson L., Keil K., Browning L. B., Krot A. N. and Bourcier W. 1999 Early aqueous alteration, explosive disruption, and reprocessing of asteroids. *Meteorit. Planet. Sci.* 34:541–557.
- Zolensky M. E., Bourcier W. L., and Gooding J. L. 1989. Aqueous alteration on the hydrous asteroids: Results of EQ3/6 computer simulations. *Icarus* 78:411–425.
- Zolensky M. E., Mittlefehldt D. W., Lipschutz M. E., Wang M.-S., Clayton R. N., Mayeda T. K., Grady M. M., Pillinger C., and B D. 1997. CM chondrites exhibit the complete petrologic range from type 2 to 1. *Geochimica et Cosmochimica Acta* 61:5099–5115.
- Zolotov M. Y. 2014. Formation of brucite and cronstedtite-bearing mineral assemblages on Ceres. *Icarus* 228:13–26.

Tables

Table 1: Quantification of the relative atomic proportion of the iron oxidation state for CM, CR, CI, and CV chondrites with the iron oxidation index calculated (IOI). The heated CMs as been classified by Alexander et al. (2013)

Figures:

Figure 1: a) Normalized XANES spectra of selected carbonaceous chondrites with enlarged pre-edge feature. b) XANES spectra references for Fe^0 , Fe^{2+} , and Fe^{3+} oxidation state.

Figure 2: Normalized XANES spectra of CR chondrites with inset expanded diagram of the pre-edge feature.

Figure 3: Normalized XANES spectra of CM chondrites with inset expanded diagram of the pre-edge feature.


Figure 4: a) Average of iron oxidation index for the different groups of carbonaceous chondrites. b) Diagram of $\text{Fe}^{3+}/\text{Fe}^{2+}$ in comparison to $\text{Fe}^0/\text{Fe}_{\text{tot}}$ for the different groups of carbonaceous chondrites

Figure 5: Comparison between the average oxidation factor of iron bulk H content (from Alexander et al. (2012)) for the different groups of carbonaceous chondrites.

Figure 6: Comparison between the average oxidation factor of iron with the 3- μm water band from Beck et al. (2014) for the different groups of carbonaceous chondrites.

Figure 7: Comparison between the average oxidation factor of iron with the bulk D/H ratio from Alexander et al. (2012) for the different groups of carbonaceous chondrites.

a)



b)




FIGURE 1




FIGURE 2





FIGURE 3

a)



b)




FIGURE 4




FIGURE 5




FIGURE 6

