

# Nonlinear Monte Carlo methods with polynomial runtime for high-dimensional iterated nested expectations

Christian Beck<sup>1</sup>, Arnulf Jentzen<sup>2,3</sup>, and Thomas Kruse<sup>4</sup>

<sup>1</sup> Department of Mathematics, ETH Zurich, Zürich, Switzerland, e-mail: christian.beck@math.ethz.ch

<sup>2</sup> Department of Mathematics, ETH Zurich, Zürich, Switzerland, e-mail: arnulf.jentzen@sam.math.ethz.ch

<sup>3</sup> Faculty of Mathematics and Computer Science, University of Münster, Münster, Germany, e-mail: ajentzen@uni-muenster.de

<sup>4</sup> Institute of Mathematics, University of Gießen, Gießen, Germany, e-mail: thomas.kruse@math.uni-giessen.de

December 5, 2021

## Abstract

The approximative calculation of iterated nested expectations is a recurring challenging problem in applications. Nested expectations appear, for example, in the numerical approximation of solutions of backward stochastic differential equations (BSDEs), in the numerical approximation of solutions of semilinear parabolic partial differential equations (PDEs), in statistical physics, in optimal stopping problems such as the approximative pricing of American or Bermudan options, in risk measure estimation in mathematical finance, or in decision-making under uncertainty. Nested expectations which arise in the above named applications often consist of a large number of nestings. However, the computational effort of standard nested Monte Carlo approximations for iterated nested expectations grows exponentially in the number of nestings and it remained an open question whether it is possible to approximately calculate multiply iterated high-dimensional nested expectations in polynomial time. In this article we tackle this problem by proposing and studying a new class of full-history recursive multilevel Picard (MLP) approximation schemes for iterated nested expectations. In particular, we prove under suitable assumptions that these MLP approximation schemes can approximately calculate multiply iterated nested expectations with a computational effort growing at most polynomially in the number of nestings  $K \in \mathbb{N} = \{1, 2, 3, \dots\}$ , in the problem dimension  $d \in \mathbb{N}$ , and in the reciprocal  $1/\varepsilon$  of the desired approximation accuracy  $\varepsilon \in (0, \infty)$ .

# Contents

|          |                                                                           |           |
|----------|---------------------------------------------------------------------------|-----------|
| <b>1</b> | <b>Introduction</b>                                                       | <b>2</b>  |
| <b>2</b> | <b>Full-history recursive multilevel Picard (MLP) approximations</b>      | <b>4</b>  |
| 2.1      | MLP approximation schemes for nested expectations . . . . .               | 4         |
| 2.2      | Measurability properties for MLP approximations . . . . .                 | 5         |
| 2.3      | Distribution properties for general random fields . . . . .               | 7         |
| 2.4      | Distribution properties for MLP approximations . . . . .                  | 9         |
| 2.5      | Integrability properties for MLP approximations . . . . .                 | 14        |
| <b>3</b> | <b>Error analysis for MLP approximations for nested expectations</b>      | <b>16</b> |
| 3.1      | A priori estimates for nested expectations . . . . .                      | 16        |
| 3.2      | Bias and variance estimates for MLP approximations . . . . .              | 23        |
| 3.3      | Recursive error estimates for MLP approximations . . . . .                | 28        |
| 3.4      | Full error analysis for MLP approximations . . . . .                      | 34        |
| <b>4</b> | <b>Complexity analysis for MLP approximations for nested expectations</b> | <b>39</b> |
| 4.1      | Complexity analysis for MLP approximations . . . . .                      | 40        |
| 4.2      | Exponential Euler approximations for semilinear heat equations . . . . .  | 42        |

## 1 Introduction

The approximative calculation of iterated nested expectations is a recurring challenging problem in applications. Iterated nested expectations can, for instance, arise in the numerical approximation of solutions of backward stochastic differential equations (BSDEs) (see, e.g., [7, 9, 28, 43]), in the numerical approximation of solutions of semilinear parabolic partial differential equations (PDEs) (see, e.g., [21, 44]), in statistical physics (see, e.g., [17, 33, 46, 47]), in optimal stopping problems such as the approximative pricing of American or Bermudan options (see, e.g., [2, 6, 8, 14, 16, 20, 31, 41, 48, 49]), in risk measure estimation in mathematical finance (see, e.g., [12, 13, 26, 27, 32, 40, 45]), or in decision-making under uncertainty (see, e.g., [1, 10, 11, 42]). In general, explicit solutions in closed form are not available for iterated nested expectations in the above mentioned problems and this entails a high demand for numerical approximation methods for the approximative calculation of iterated nested expectations. In the scientific literature we refer to [15, 22, 23, 24, 29, 30, 34] for results on the numerical approximation of two nested expectations. Nested expectations which appear in the numerical approximation of solutions of BSDEs, in the numerical approximation of semilinear PDEs, or in the approximative pricing of American or Bermudan options usually do not only consist of two nested expectations but do instead consist of a large number of nestings. However, the computational effort of standard nested Monte Carlo approximations for iterated nested expectations grows exponentially in the number of nestings and it remained an open question whether it is possible to approximately calculate multiply iterated high-dimensional nested expectations in polynomial time.

It is the subject of this article to attack this problem by proposing and studying a new class of full-history recursive multilevel Picard approximation schemes for iterated nested expectations (in the following we abbreviate full-history recursive multilevel Picard by MLP). MLP approximation schemes were introduced in [18, 36] for the numerical approximation of solutions of semilinear PDEs and have been shown in [3, 4, 36, 37] to overcome the curse of dimensionality in the numerical approximation of certain semilinear PDEs (cf. also [5, 19, 25, 35, 38] for numerical simulations and further mathematical results for MLP approximation schemes). In this article we

propose a new class of time-discrete MLP approximation schemes and we prove under suitable assumptions that these MLP approximation schemes can approximately calculate multiply iterated nested expectations with a computational effort growing at most polynomially in the number of nestings  $K \in \mathbb{N} = \{1, 2, 3, \dots\}$ , in the problem dimension  $d \in \mathbb{N}$ , and in the reciprocal  $1/\varepsilon$  of the desired approximation accuracy  $\varepsilon \in (0, \infty)$ . To illustrate this article's main result, Theorem 4.1 in Subsection 4.1 below, we now present in Theorem 1.1 a special case of Theorem 4.1.

**Theorem 1.1.** *Let  $L, T, \delta \in (0, \infty)$ ,  $\Theta = \bigcup_{k \in \mathbb{N}} \mathbb{Z}^k$ ,  $f \in C(\mathbb{R}, \mathbb{R})$  satisfy for all  $a, b \in \mathbb{R}$  that  $|f(a) - f(b)| \leq L|a - b|$ , let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a probability space, let  $R^\theta: \Omega \rightarrow (0, 1)$ ,  $\theta \in \Theta$ , be i.i.d. random variables, assume for all  $t \in (0, 1)$  that  $\mathbb{P}(R^0 \leq t) = t$ , let  $W^{d, \theta}: [0, T] \times \Omega \rightarrow \mathbb{R}^d$ ,  $d \in \mathbb{N}$ ,  $\theta \in \Theta$ , be i.i.d. standard Brownian motions, assume that  $(R^\theta)_{\theta \in \Theta}$  and  $(W^{d, \theta})_{(d, \theta) \in \mathbb{N} \times \Theta}$  are independent, let  $v_k^{d, K} \in C(\mathbb{R}^d, \mathbb{R})$ ,  $k \in \{0, 1, \dots, K\}$ ,  $d, K \in \mathbb{N}$ , satisfy for all  $d, K \in \mathbb{N}$ ,  $k \in \{1, 2, \dots, K\}$ ,  $x \in \mathbb{R}^d$  that  $|v_0^{d, K}(x)| \leq L$  and*

$$v_k^{d, K}(x) = \mathbb{E}[v_{k-1}^{d, K}(x + W_{T/K}^{d, 0}) + \frac{T}{K} f(v_{k-1}^{d, K}(x + W_{T/K}^{d, 0}))], \quad (1)$$

let  $V_{k, n, M}^{d, K, \theta}: \mathbb{R}^d \times \Omega \rightarrow \mathbb{R}$ ,  $\theta \in \Theta$ ,  $k \in \{0, 1, \dots, K\}$ ,  $n \in \mathbb{N}_0$ ,  $d, K, M \in \mathbb{N}$ , satisfy<sup>1</sup> for all  $d, K, M \in \mathbb{N}$ ,  $n \in \mathbb{N}_0$ ,  $k \in \{0, 1, \dots, K\}$ ,  $\theta \in \Theta$ ,  $x \in \mathbb{R}^d$  that

$$\begin{aligned} V_{k, n, M}^{d, K, \theta}(x) &= \frac{\mathbb{1}_{\mathbb{N}}(n)}{M^n} \sum_{m=1}^{M^n} \left[ v_0^{d, K}(x + W_{kT/K}^{d, (\theta, 0, -m)}) + \frac{kT}{K} f(0) \right] \\ &\quad + \sum_{j=1}^{n-1} \frac{kT}{KM^{n-j}} \sum_{m=1}^{M^{n-j}} \left[ f(V_{[kR^{(\theta, j, m)}]_{j, M}}^{d, K, (\theta, j, m)}(x + W_{kT/K}^{d, (\theta, j, m)} - W_{[kR^{(\theta, j, m)}]_{T/K}}^{d, (\theta, j, m)}) \right. \\ &\quad \left. - f(V_{[kR^{(\theta, j, m)}]_{j-1, M}}^{d, K, (\theta, j, -m)}(x + W_{kT/K}^{d, (\theta, j, m)} - W_{[kR^{(\theta, j, m)}]_{T/K}}^{d, (\theta, j, m)}) \right], \quad (2) \end{aligned}$$

and for every  $d, K, M \in \mathbb{N}$ ,  $n \in \mathbb{N}_0$  let  $\mathfrak{C}_{M, n}^{d, K} \in \mathbb{N}_0$  be the number of realizations of scalar standard normal random variables which are used to compute one realization of  $V_{K, n, M}^{d, K, 0}(0): \Omega \rightarrow \mathbb{R}$  (cf. (188) for a precise definition). Then there exist  $\mathfrak{N} = (\mathfrak{N}_\varepsilon)_{\varepsilon \in (0, 1]}: (0, 1] \rightarrow \mathbb{N}$  and  $\mathfrak{c} \in \mathbb{R}$  such that for all  $\varepsilon \in (0, 1]$ ,  $d, K \in \mathbb{N}$  it holds that  $\mathfrak{C}_{\mathfrak{N}_\varepsilon, \mathfrak{N}_\varepsilon}^{d, K} \leq \mathfrak{c} d \varepsilon^{-(2+\delta)}$  and  $(\mathbb{E}[|V_{K, \mathfrak{N}_\varepsilon, \mathfrak{N}_\varepsilon}^{d, K, 0}(0) - v_K^{d, K}(0)|^2])^{1/2} \leq \varepsilon$ .

Theorem 1.1 is an immediate consequence of Corollary 4.4 in Subsection 4.2 below. Corollary 4.4, in turn, follows from Theorem 4.1, the main result of this article. Roughly speaking, Theorem 1.1 shows for every arbitrarily small real number  $\delta \in (0, \infty)$  that MLP approximations can approximately calculate the function values  $v_K^{d, K}(0)$ ,  $d, K \in \mathbb{N}$ , which are given recursively by the iterated nested expectations in (1) above with the root mean square error bounded by  $\varepsilon \in (0, \infty)$  and the computational cost bounded by a multiple of the product of  $d \in \mathbb{N}$  and  $\varepsilon^{-(2+\delta)}$ . The real number  $T \in (0, \infty)$  and the function  $f \in C(\mathbb{R}, \mathbb{R})$  in Theorem 1.1 above describe the time horizon and the nonlinearity in the nested expectations in (1) in Theorem 1.1. The nonlinearity  $f$  in Theorem 1.1 is assumed to be Lipschitz continuous with the Lipschitz constant bounded by  $L \in (0, \infty)$ . The real number  $L \in (0, \infty)$  is also used to express a uniform boundedness assumption for the functions  $v_0^{d, K}: \mathbb{R}^d \rightarrow \mathbb{R}$ ,  $d, K \in \mathbb{N}$ , in Theorem 1.1. The functions  $v_K^{d, K}: \mathbb{R}^d \rightarrow \mathbb{R}$ ,  $d, K \in \mathbb{N}$ , are calculated approximately by means of the MLP approximation scheme in (2) in Theorem 1.1. The computational effort for evaluating the MLP approximations in (2) is measured by means of the natural numbers  $\mathfrak{C}_{M, n}^{d, K} \in \mathbb{N}$ ,  $d, K, M \in \mathbb{N}$ ,  $n \in \mathbb{N}_0$ , in Theorem 1.1 above. The specific recursion of the iterated nested expectations in (1) in Theorem 1.1 above describes just one class of examples which can be calculated approximately by the MLP approximations proposed in this

<sup>1</sup>Note that for all  $t \in \mathbb{R}$  it holds that  $\lceil t \rceil = \max((-\infty, t] \cap \mathbb{Z})$ .

paper and we refer to Theorem 4.1 in Subsection 4.1 below for our more general approximation result for iterated nested expectations.

The remainder of this article is organized as follows. In Section 2 we introduce in Subsection 2.1 MLP approximation schemes for the approximation of iterated nested expectations (see (4) in Setting 2.1 in Subsection 2.1) and we establish in Subsections 2.2, 2.4, and 2.5 basic measurability, integrability, and distribution properties for the proposed MLP approximations. In Section 3 we establish in Subsection 3.2 bias and variance estimates for MLP approximations and we provide in Subsection 3.4 a full error analysis for the proposed MLP approximation schemes. In Section 4 we combine the error analysis for the proposed MLP approximation schemes in Subsection 3.4 with a computational cost analysis for the proposed MLP approximation schemes to obtain a complexity analysis for the proposed MLP approximation schemes.

## 2 Full-history recursive multilevel Picard (MLP) approximations for iterated nested expectations

In this section we introduce in Setting 2.1 in Subsection 2.1 below MLP approximations for iterated nested expectations, we establish in Lemma 2.3 in Subsection 2.2 basic measurability properties for MLP approximations, we establish in Proposition 2.9 and Corollary 2.10 in Subsection 2.4 basic distribution properties for MLP approximations, and we establish in Lemma 2.11 in Subsection 2.5 below basic integrability properties for MLP approximations. Our proofs of Proposition 2.9, Corollary 2.10, and Lemma 2.11 are based on some general and essentially well-known distribution properties for random fields which we establish in Lemma 2.4, Lemma 2.5, Lemma 2.6, and Corollary 2.7 in Subsection 2.3 below (cf., e.g., also Hutzenthaler et al. [36, Subsection 2.2] and Hutzenthaler et al. [37, Subsection 3.3]). The MLP approximations for iterated nested expectations introduced in Subsection 2.1 below are inspired by the MLP approximations for semilinear PDEs introduced in Hutzenthaler et al. [36, Section 3].

### 2.1 MLP approximation schemes for nested expectations

**Setting 2.1.** Let  $d, K, M \in \mathbb{N}$ ,  $\Theta = \bigcup_{n \in \mathbb{N}} \mathbb{Z}^n$ ,  $\mathcal{O} \in \mathcal{B}(\mathbb{R}^d) \setminus \{\emptyset\}$ , let  $f_k: \mathcal{O} \times \mathbb{R} \rightarrow \mathbb{R}$ ,  $k \in \{0, 1, \dots, K\}$ , be  $\mathcal{B}(\mathcal{O} \times \mathbb{R})/\mathcal{B}(\mathbb{R})$ -measurable, let  $g: \mathcal{O} \rightarrow \mathbb{R}$  be  $\mathcal{B}(\mathcal{O})/\mathcal{B}(\mathbb{R})$ -measurable, let  $(S, \mathcal{S})$  be a measurable space, let  $\phi_k: \mathcal{O} \times S \rightarrow \mathcal{O}$ ,  $k \in \{0, 1, \dots, K\}$ , be  $(\mathcal{B}(\mathcal{O}) \otimes \mathcal{S})/\mathcal{B}(\mathcal{O})$ -measurable, let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a probability space, let  $W_k^\theta: \Omega \rightarrow S$ ,  $\theta \in \Theta$ ,  $k \in \{0, 1, \dots, K\}$ , be independent random variables, assume for every  $k \in \{0, 1, \dots, K\}$  that  $W_k^\theta: \Omega \rightarrow S$ ,  $\theta \in \Theta$ , are identically distributed, let  $\mathcal{R}^\theta = (\mathcal{R}_k^\theta)_{k \in \{0, 1, \dots, K\}}: \{0, 1, \dots, K\} \times \Omega \rightarrow \mathbb{N}_0$ ,  $\theta \in \Theta$ , be i.i.d. stochastic processes, assume for every  $k \in \{1, 2, \dots, K\}$  that  $\mathcal{R}_k^\theta \leq k - 1$ , let  $\mathbf{p}_{k,l} \in (0, \infty)$ ,  $k \in \{1, 2, \dots, K\}$ ,  $l \in \mathbb{N}_0$ , satisfy for all  $k \in \{1, 2, \dots, K\}$ ,  $l \in \mathbb{N}_0$  that  $\mathbf{p}_{k,l} \mathbb{P}(\mathcal{R}_k^0 = l) = |\mathbb{P}(\mathcal{R}_k^0 = l)|^2$ , assume that  $(\mathcal{R}^\theta)_{\theta \in \Theta}$  and  $(W_k^\theta)_{(\theta,k) \in \Theta \times \{0, 1, \dots, K\}}$  are independent, let  $X^{\theta,k} = (X_l^{\theta,k,x})_{(l,x) \in \{0, 1, \dots, k\} \times \mathcal{O}}: \{0, 1, \dots, k\} \times \mathcal{O} \times \Omega \rightarrow \mathcal{O}$ ,  $k \in \{0, 1, \dots, K\}$ ,  $\theta \in \Theta$ , satisfy for all  $\theta \in \Theta$ ,  $k \in \{0, 1, \dots, K\}$ ,  $l \in \{0, 1, \dots, k\}$ ,  $x \in \mathcal{O}$  that

$$X_l^{\theta,k,x} = \begin{cases} x & : l = k \\ \phi_l(X_{l+1}^{\theta,k,x}, W_l^\theta) & : l < k, \end{cases} \quad (3)$$

and let  $V_{k,n}^\theta: \mathcal{O} \times \Omega \rightarrow \mathbb{R}$ ,  $n \in \mathbb{N}_0$ ,  $k \in \{0, 1, \dots, K\}$ ,  $\theta \in \Theta$ , satisfy for all  $\theta \in \Theta$ ,  $k \in \{0, 1, \dots, K\}$ ,  $n \in \mathbb{N}_0$ ,  $x \in \mathcal{O}$  that

$$\begin{aligned} V_{k,n}^\theta(x) = & \frac{\mathbb{1}_{\mathbb{N}}(n)}{M^n} \sum_{m=1}^{M^n} \left[ g(X_0^{(\theta,0,-m),k,x}) + \sum_{l=0}^{k-1} f_l(X_l^{(\theta,0,m),k,x}, 0) \right] + \sum_{j=1}^{n-1} \frac{1}{M^{n-j}} \sum_{m=1}^{M^{n-j}} \frac{\mathbb{1}_{\mathbb{N}}(k)}{\mathfrak{p}_{k,\mathcal{R}_k^{(\theta,j,m)}}} \\ & \cdot \left[ \left( f_{\mathcal{R}_k^{(\theta,j,m)}}(X_{\mathcal{R}_k^{(\theta,j,m)}}^{(\theta,j,m),k,x}, V_{\mathcal{R}_k^{(\theta,j,m)},j}^{(\theta,j,m)}(X_{\mathcal{R}_k^{(\theta,j,m)}}^{(\theta,j,m),k,x})) - V_{\mathcal{R}_k^{(\theta,j,m)},j}^{(\theta,j,m)}(X_{\mathcal{R}_k^{(\theta,j,m)}}^{(\theta,j,m),k,x}) \right) \right. \\ & \left. - \left( f_{\mathcal{R}_k^{(\theta,j,m)}}(X_{\mathcal{R}_k^{(\theta,j,m)}}^{(\theta,j,m),k,x}, V_{\mathcal{R}_k^{(\theta,j,m)},j-1}^{(\theta,j,-m)}(X_{\mathcal{R}_k^{(\theta,j,m)}}^{(\theta,j,m),k,x})) - V_{\mathcal{R}_k^{(\theta,j,m)},j-1}^{(\theta,j,-m)}(X_{\mathcal{R}_k^{(\theta,j,m)}}^{(\theta,j,m),k,x}) \right) \right]. \end{aligned} \quad (4)$$

## 2.2 Measurability properties for MLP approximations

**Lemma 2.2.** Let  $d, K \in \mathbb{N}$ ,  $\mathcal{O} \in \mathcal{B}(\mathbb{R}^d) \setminus \{\emptyset\}$ , let  $(S, \mathcal{S})$  be a measurable space, let  $\phi_k: \mathcal{O} \times S \rightarrow \mathcal{O}$ ,  $k \in \{0, 1, \dots, K\}$ , be  $(\mathcal{B}(\mathcal{O}) \otimes \mathcal{S})/\mathcal{B}(\mathcal{O})$ -measurable, let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a probability space, let  $\mathcal{A} \subseteq \mathcal{F}$  be a sigma-algebra on  $\Omega$ , let  $W_k: \Omega \rightarrow S$ ,  $k \in \{0, 1, \dots, K\}$ , be random variables, and let  $X^k = (X_l^{k,x})_{(l,x) \in \{0,1,\dots,k\} \times \mathcal{O}}: \{0, 1, \dots, k\} \times \mathcal{O} \times \Omega \rightarrow \mathcal{O}$ ,  $k \in \{0, 1, \dots, K\}$ , satisfy for all  $k \in \{0, 1, \dots, K\}$ ,  $l \in \{0, 1, \dots, k\}$ ,  $x \in \mathcal{O}$  that

$$X_l^{k,x} = \begin{cases} x & : l = k \\ \phi_l(X_{l+1}^{k,x}, W_l) & : l < k. \end{cases} \quad (5)$$

Then

- (i) it holds for all  $k, l \in \mathbb{N}_0$  with  $l \leq k \leq K$  that  $\mathcal{O} \times \Omega \ni (x, \omega) \mapsto X_l^{k,x}(\omega) \in \mathcal{O}$  is  $(\mathcal{B}(\mathcal{O}) \otimes \sigma((W_m)_{m \in [l,k] \cap \mathbb{N}_0}))/\mathcal{B}(\mathcal{O})$ -measurable and
- (ii) it holds for all  $k, l \in \mathbb{N}_0$  with  $l \leq k \leq K$  that  $\mathcal{O} \times \Omega \ni (x, \omega) \mapsto (X_l^{k,x}(\omega), \omega) \in \mathcal{O} \times \Omega$  is  $(\mathcal{B}(\mathcal{O}) \otimes \sigma(\sigma((W_m)_{m \in [l,k] \cap \mathbb{N}_0}) \cup \mathcal{A}))/(\mathcal{B}(\mathcal{O}) \otimes \mathcal{A})$ -measurable.

*Proof of Lemma 2.2.* Throughout this proof let  $\mathbb{A}_k \subseteq \mathbb{N}_0$ ,  $k \in \{0, 1, \dots, K\}$ , satisfy for all  $k \in \{0, 1, \dots, K\}$  that

$$\mathbb{A}_k = \{l \in \{0, 1, \dots, k\} : [\forall A \in \mathcal{B}(\mathcal{O}) : (X_l^k)^{-1}(A) \in (\mathcal{B}(\mathcal{O}) \otimes \sigma((W_m)_{m \in [l,k] \cap \mathbb{N}_0}))]\}. \quad (6)$$

Observe that the fact that for all  $k \in \{0, 1, \dots, K-1\}$  it holds that  $W_k: \Omega \rightarrow S$  is  $\sigma(W_k)/\mathcal{S}$ -measurable implies that for all  $k, l \in \{0, 1, \dots, K\}$  with  $l+1 \in \mathbb{A}_k$  it holds that  $\mathcal{O} \times \Omega \ni (x, \omega) \mapsto (X_{l+1}^{k,x}(\omega), W_l(\omega)) \in \mathcal{O} \times S$  is  $(\mathcal{B}(\mathcal{O}) \otimes \sigma((W_m)_{m \in [l,k] \cap \mathbb{N}_0}))/(\mathcal{B}(\mathcal{O}) \otimes \mathcal{S})$ -measurable. The fact that for all  $k \in \{0, 1, \dots, K-1\}$  it holds that  $\phi_k: \mathcal{O} \times S \rightarrow \mathcal{O}$  is  $(\mathcal{B}(\mathcal{O}) \otimes \mathcal{S})/\mathcal{B}(\mathcal{O})$ -measurable and (5) therefore ensure that for all  $k, l \in \{0, 1, \dots, K\}$  with  $l+1 \in \mathbb{A}_k$  it holds that  $X_l^k: \mathcal{O} \times \Omega \rightarrow \mathcal{O}$  is  $(\mathcal{B}(\mathcal{O}) \otimes \sigma((W_m)_{m \in [l,k] \cap \mathbb{N}_0}))/\mathcal{B}(\mathcal{O})$ -measurable. Hence, we obtain that for all  $k, l \in \{0, 1, \dots, K\}$  with  $l+1 \in \mathbb{A}_k$  it holds that  $l \in \mathbb{A}_k$ . Moreover, observe that (5) ensures for all  $k \in \{0, 1, \dots, K\}$  that  $k \in \mathbb{A}_k$ . Induction and the fact that for all  $k, l \in \{0, 1, \dots, K\}$  with  $l+1 \in \mathbb{A}_k$  it holds that  $l \in \mathbb{A}_k$  therefore yield that for all  $k \in \{0, 1, \dots, K\}$  it holds that  $\mathbb{A}_k = \{0, 1, \dots, k\}$ . This establishes item (i). Next note that item (i) ensures that for all  $k, l \in \mathbb{N}_0$ ,  $B \in \mathcal{B}(\mathcal{O})$  with  $l \leq k \leq K$  it holds that

$$\begin{aligned} \{(x, \omega) \in \mathcal{O} \times \Omega : X_l^{k,x}(\omega) \in B\} & \in (\mathcal{B}(\mathcal{O}) \otimes \sigma((W_m)_{m \in [l,k] \cap \mathbb{N}_0})) \\ & \subseteq (\mathcal{B}(\mathcal{O}) \otimes \sigma(\sigma((W_m)_{m \in [l,k] \cap \mathbb{N}_0}) \cup \mathcal{A})). \end{aligned} \quad (7)$$

Furthermore, observe that for all  $k, l \in \mathbb{N}_0$ ,  $A \in \mathcal{A}$  with  $l \leq k \leq K$  it holds that

$$\{(x, \omega) \in \mathcal{O} \times \Omega : \omega \in A\} = (\mathcal{O} \times A) \in (\mathcal{B}(\mathcal{O}) \otimes \mathcal{A}) \subseteq (\mathcal{B}(\mathcal{O}) \otimes \sigma(\sigma((W_m)_{m \in [l,k] \cap \mathbb{N}_0}) \cup \mathcal{A})). \quad (8)$$

This and (7) imply that for all  $k, l \in \mathbb{N}_0$ ,  $B \in \mathcal{B}(\mathcal{O})$ ,  $A \in \mathcal{A}$  with  $l \leq k \leq K$  it holds that

$$\begin{aligned} & \{(x, \omega) \in \mathcal{O} \times \Omega : (X_l^{k,x}(\omega), \omega) \in B \times A\} \\ &= (\{(x, \omega) \in \mathcal{O} \times \Omega : X_l^{k,x}(\omega) \in B\} \cap (\mathcal{O} \times A)) \in (\mathcal{B}(\mathcal{O}) \otimes \sigma(\sigma((W_m)_{m \in [l,k] \cap \mathbb{N}_0}) \cup \mathcal{A})). \end{aligned} \quad (9)$$

This establishes item (ii). This completes the proof of Lemma 2.2.  $\square$

**Lemma 2.3.** *Assume Setting 2.1. Then*

(i) *it holds for all  $\theta \in \Theta$ ,  $n \in \mathbb{N}_0$  that  $\{0, 1, \dots, K\} \times \mathcal{O} \times \Omega \ni (k, x, \omega) \mapsto V_{k,n}^\theta(x, \omega) \in \mathbb{R}$  is  $(2^{\{0,1,\dots,K\}} 2^{\{0,1,\dots,K\}} \otimes \mathcal{B}(\mathcal{O}) \otimes \sigma((\mathcal{R}_s^{\theta,\vartheta})_{(s,\vartheta) \in \{1,2,\dots,K\} \times \Theta}, (W_s^{\theta,\vartheta})_{(s,\vartheta) \in \{0,1,\dots,K-1\} \times \Theta})) / \mathcal{B}(\mathbb{R})$ -measurable and*

(ii) *it holds for all  $\theta, \vartheta \in \Theta$ ,  $j \in \mathbb{N}$ ,  $k \in \{1, 2, \dots, K\}$  that*

$$\mathcal{O} \times \Omega \ni (x, \omega) \mapsto \left[ \frac{f_{\mathcal{R}_k^\theta(\omega)}(X_{\mathcal{R}_k^\theta(\omega)}^{\theta,k,x}(\omega), V_{\mathcal{R}_k^\theta(\omega),j}^\vartheta(X_{\mathcal{R}_k^\theta(\omega)}^{\theta,k,x}(\omega), \omega)) - V_{\mathcal{R}_k^\theta(\omega),j}^\vartheta(X_{\mathcal{R}_k^\theta(\omega)}^{\theta,k,x}(\omega), \omega))}{\mathfrak{p}_{k,\mathcal{R}_k^\theta(\omega)}} \right] \in \mathbb{R} \quad (10)$$

*is  $(\mathcal{B}(\mathcal{O}) \otimes \sigma((\mathcal{R}_s^\theta)_{s \in \{1,2,\dots,K\}}, (\mathcal{R}_s^{\vartheta,\eta})_{(s,\eta) \in \{1,2,\dots,K\} \times \Theta}, (W_s^\theta)_{s \in \{0,1,\dots,K-1\}}, (W_s^{\vartheta,\eta})_{(s,\eta) \in \{0,1,\dots,K-1\} \times \Theta})) / \mathcal{B}(\mathbb{R})$ -measurable.*

*Proof of Lemma 2.3.* Throughout this proof let  $\mathbb{A} \subseteq \mathbb{N}$  satisfy

$$\mathbb{A} = \left\{ n \in \mathbb{N} : \left[ \forall \theta \in \Theta, k \in \{0, 1, \dots, K\}, j \in \{0, 1, \dots, n-1\}, A \in \mathcal{B}(\mathbb{R}) : \right. \right. \\ \left. \left. (V_{k,j}^\theta)^{-1}(A) \in (\mathcal{B}(\mathcal{O}) \otimes \sigma((\mathcal{R}_s^{\theta,\vartheta})_{(s,\vartheta) \in \{1,2,\dots,K\} \times \Theta}, (W_s^{\theta,\vartheta})_{(s,\vartheta) \in \{0,1,\dots,K-1\} \times \Theta})) \right] \right\}. \quad (11)$$

Observe that the assumption that for all  $\theta \in \Theta$ ,  $k \in \{0, 1, \dots, K\}$ ,  $x \in \mathcal{O}$  it holds that  $V_{k,0}^\theta(x) = 0$  implies that  $1 \in \mathbb{A}$ . In the next step we note that item (ii) of Lemma 2.2 ensures that for all  $n \in \mathbb{A}$ ,  $\theta, \vartheta \in \Theta$ ,  $j \in \{0, 1, \dots, n-1\}$ ,  $k, m \in \{0, 1, \dots, K\}$ ,  $l \in \{0, 1, \dots, k\}$  it holds that

$$\mathcal{O} \times \Omega \ni (x, \omega) \mapsto V_{m,j}^\vartheta(X_l^{\theta,k,x}(\omega), \omega) \in \mathbb{R} \quad (12)$$

is  $(\mathcal{B}(\mathcal{O}) \otimes \sigma((W_s^\theta)_{s \in \{0,1,\dots,K-1\}}, (\mathcal{R}_s^{\vartheta,\eta})_{(s,\eta) \in \{1,2,\dots,K\} \times \Theta}, (W_s^{\vartheta,\eta})_{(s,\eta) \in \{0,1,\dots,K-1\} \times \Theta})) / \mathcal{B}(\mathbb{R})$ -measurable. The fact that  $\mathcal{B}(\mathcal{O}) \otimes \mathcal{B}(\mathbb{R}) = \mathcal{B}(\mathcal{O} \times \mathbb{R})$  and item (i) of Lemma 2.2 hence ensure that for all  $n \in \mathbb{A}$ ,  $\theta, \vartheta \in \Theta$ ,  $j \in \{0, 1, \dots, n-1\}$ ,  $k, m \in \{0, 1, \dots, K\}$ ,  $l \in \{0, 1, \dots, k\}$  it holds that

$$\mathcal{O} \times \Omega \ni (x, \omega) \mapsto (X_l^{\theta,k,x}(\omega), V_{m,j}^\vartheta(X_l^{\theta,k,x}(\omega), \omega)) \in \mathcal{O} \times \mathbb{R} \quad (13)$$

is  $(\mathcal{B}(\mathcal{O}) \otimes \sigma((W_s^\theta)_{s \in \{0,1,\dots,K-1\}}, (\mathcal{R}_s^{\vartheta,\eta})_{(s,\eta) \in \{1,2,\dots,K\} \times \Theta}, (W_s^{\vartheta,\eta})_{(s,\eta) \in \{0,1,\dots,K-1\} \times \Theta})) / (\mathcal{B}(\mathcal{O} \times \mathbb{R}))$ -measurable. Combining this with the assumption that for all  $k \in \{0, 1, \dots, K-1\}$  it holds that  $f_k : \mathcal{O} \times \mathbb{R} \rightarrow \mathbb{R}$  is  $\mathcal{B}(\mathcal{O} \times \mathbb{R}) / \mathcal{B}(\mathbb{R})$ -measurable and the fact that for all  $k, j \in \mathbb{N}_0$ ,  $\theta, \vartheta \in \Theta$ ,  $x \in \mathcal{O}$ ,  $\omega \in \Omega$  with  $1 \leq k \leq K$  it holds that

$$\frac{f_{\mathcal{R}_k^\theta(\omega)}(X_{\mathcal{R}_k^\theta(\omega)}^{\theta,k,x}(\omega), V_{\mathcal{R}_k^\theta(\omega),j}^\vartheta(X_{\mathcal{R}_k^\theta(\omega)}^{\theta,k,x}(\omega), \omega))}{\mathfrak{p}_{k,\mathcal{R}_k^\theta(\omega)}} = \sum_{l=0}^{k-1} \mathbb{1}_{\{\mathcal{R}_k^\theta=l\}}(\omega) \left[ \frac{f_l(X_l^{\theta,k,x}(\omega), V_{l,j}^\vartheta(X_l^{\theta,k,x}(\omega), \omega))}{\mathfrak{p}_{k,l}} \right] \quad (14)$$

ensures that for all  $n \in \mathbb{A}$ ,  $\theta, \vartheta \in \Theta$ ,  $k \in \{0, 1, \dots, K\}$ ,  $j \in \{0, 1, \dots, n-1\}$  it holds that

$$\mathcal{O} \times \Omega \ni (x, \omega) \mapsto f_{\mathcal{R}_k^\theta(\omega)}(X_{\mathcal{R}_k^\theta(\omega)}^{\theta,k,x}(\omega), V_{\mathcal{R}_k^\theta(\omega),j}^\vartheta(X_{\mathcal{R}_k^\theta(\omega)}^{\theta,k,x}(\omega), \omega)) \in \mathbb{R} \quad (15)$$

is  $(\mathcal{B}(\mathcal{O}) \otimes \sigma((\mathcal{R}_s^\theta)_{s \in \{1,2,\dots,K\}}, (W_s^\theta)_{s \in \{0,1,\dots,K-1\}}, (\mathcal{R}_s^{(\vartheta,\eta)})_{(s,\eta) \in \{1,2,\dots,K\} \times \Theta}, (W_s^{(\vartheta,\eta)})_{(s,\eta) \in \{0,1,\dots,K-1\} \times \Theta})) / \mathcal{B}(\mathbb{R})$ -measurable. The assumption that for all  $k \in \{0, 1, \dots, K-1\}$  it holds that  $f_k: \mathcal{O} \times \mathbb{R} \rightarrow \mathbb{R}$  is  $\mathcal{B}(\mathcal{O} \times \mathbb{R}) / \mathcal{B}(\mathbb{R})$ -measurable, the assumption that  $g: \mathcal{O} \rightarrow \mathbb{R}$  is  $\mathcal{B}(\mathcal{O}) / \mathcal{B}(\mathbb{R})$ -measurable, (4), (12), (13), and item (i) of Lemma 2.2 therefore guarantee that for all  $n \in \mathbb{A}$ ,  $\theta \in \Theta$ ,  $k \in \{0, 1, \dots, K\}$  it holds that  $V_{k,n}^\theta: \mathcal{O} \times \Omega \rightarrow \mathbb{R}$  is  $(\mathcal{B}(\mathcal{O}) \otimes \sigma((\mathcal{R}_s^{(\theta,\vartheta)})_{(s,\vartheta) \in \{1,2,\dots,K\} \times \Theta}, (W_s^{(\theta,\vartheta)})_{(s,\vartheta) \in \{0,1,\dots,K-1\} \times \Theta})) / \mathcal{B}(\mathbb{R})$ -measurable. Hence, we obtain that for all  $n \in \mathbb{A}$  it holds that  $n+1 \in \mathbb{A}$ . Induction and the fact that  $1 \in \mathbb{A}$  therefore demonstrate that  $\mathbb{A} = \mathbb{N}$ . This establishes item (i). In the next step we note that item (i) and (15) establish item (ii). This completes the proof of Lemma 2.3.  $\square$

## 2.3 Distribution properties for general random fields

**Lemma 2.4.** *Let  $(\Omega, \mathcal{F})$  be a measurable space, let  $\mathcal{G} \subseteq \mathcal{F}$  and  $\mathcal{A} \subseteq \mathcal{F}$  be sigma-algebras on  $\Omega$ , let  $(E, \mathcal{E})$ ,  $(S, \mathcal{S})$ , and  $(T, \mathcal{T})$  be measurable spaces, let  $U: S \times \Omega \rightarrow T$  be  $(\mathcal{S} \otimes \mathcal{G}) / \mathcal{T}$ -measurable, and let  $Y: E \times \Omega \rightarrow S$  be  $(\mathcal{E} \otimes \mathcal{A}) / \mathcal{S}$ -measurable. Then it holds that  $U(Y) = (E \times \Omega \ni (e, \omega) \mapsto U(Y(e, \omega), \omega) \in T)$  is  $(\mathcal{E} \otimes \mathcal{F}) / \mathcal{T}$ -measurable.*

*Proof of Lemma 2.4.* First, note that the assumption that  $\mathcal{A} \subseteq \mathcal{F}$  and the assumption that  $Y$  is  $(\mathcal{E} \otimes \mathcal{A}) / \mathcal{S}$ -measurable ensures that  $Y: E \times \Omega \rightarrow S$  is  $(\mathcal{E} \otimes \mathcal{F}) / \mathcal{S}$ -measurable. This and the assumption that  $\mathcal{G} \subseteq \mathcal{F}$  imply that  $E \times \Omega \ni (e, \omega) \mapsto (Y(e, \omega), \omega) \in S \times \Omega$  is  $(\mathcal{E} \otimes \mathcal{F}) / (\mathcal{S} \otimes \mathcal{G})$ -measurable. The assumption that  $U$  is  $(\mathcal{S} \otimes \mathcal{G}) / \mathcal{T}$ -measurable hence demonstrates that  $E \times \Omega \ni (e, \omega) \mapsto U(Y(e, \omega), \omega) \in T$  is  $(\mathcal{E} \otimes \mathcal{F}) / \mathcal{T}$ -measurable. This completes the proof of Lemma 2.4.  $\square$

**Lemma 2.5.** *Let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a probability space, let  $\mathcal{G}_i \subseteq \mathcal{F}$ ,  $i \in \{1, 2\}$ , and  $\mathcal{A}_i \subseteq \mathcal{F}$ ,  $i \in \{1, 2\}$ , be sigma-algebras on  $\Omega$ , let  $(E, \mathcal{E})$ ,  $(S, \mathcal{S})$ , and  $(T, \mathcal{T})$  be measurable spaces, let  $U_i: S \times \Omega \rightarrow T$ ,  $i \in \{1, 2\}$ , be identically distributed random fields, let  $Y_i: E \times \Omega \rightarrow S$ ,  $i \in \{1, 2\}$ , be identically distributed random fields, assume for all  $i \in \{1, 2\}$  that  $U_i$  is  $(\mathcal{S} \otimes \mathcal{G}_i) / \mathcal{T}$ -measurable, assume for all  $i \in \{1, 2\}$  that  $Y_i$  is  $(\mathcal{E} \otimes \mathcal{A}_i) / \mathcal{S}$ -measurable, and assume for all  $i \in \{1, 2\}$  that  $\mathcal{G}_i$  and  $\mathcal{A}_i$  are independent. Then*

(i) *it holds for all  $i \in \{1, 2\}$  that  $U_i(Y_i) = (E \times \Omega \ni (e, \omega) \mapsto U_i(Y_i(e, \omega), \omega) \in T)$  is  $(\mathcal{E} \otimes \mathcal{F}) / \mathcal{T}$ -measurable and*

(ii) *it holds that  $U_1(Y_1)$  and  $U_2(Y_2)$  are identically distributed random fields.*

*Proof of Lemma 2.5.* First, observe that Lemma 2.4 establishes item (i). Next we prove item (ii). For this let  $n \in \mathbb{N}$ ,  $e_1, e_2, \dots, e_n \in E$ ,  $B_1, B_2, \dots, B_n \in \mathcal{T}$ , let  $\mathbb{U}_i: S^n \times \Omega \rightarrow T^n$ ,  $i \in \{1, 2\}$ , satisfy for all  $i \in \{1, 2\}$ ,  $s = (s_1, s_2, \dots, s_n) \in S^n$ ,  $\omega \in \Omega$  that

$$\mathbb{U}_i(s, \omega) = (U_i(s_1, \omega), U_i(s_2, \omega), \dots, U_i(s_n, \omega)), \quad (16)$$

let  $\mathbb{Y}_i: \Omega \rightarrow S^n$ ,  $i \in \{1, 2\}$ , satisfy for all  $i \in \{1, 2\}$ ,  $\omega \in \Omega$  that

$$\mathbb{Y}_i(\omega) = (Y_i(e_1, \omega), Y_i(e_2, \omega), \dots, Y_i(e_n, \omega)), \quad (17)$$

and let  $\mathbb{I}: T^n \rightarrow [0, \infty)$  satisfy for all  $t = (t_1, t_2, \dots, t_n) \in T^n$  that

$$\mathbb{I}(t) = \mathbb{1}_{B_1 \times B_2 \times \dots \times B_n}(t_1, t_2, \dots, t_n). \quad (18)$$

Note that the assumption that for all  $i \in \{1, 2\}$  it holds that  $Y_i$  is  $(\mathcal{E} \otimes \mathcal{A}_i) / \mathcal{S}$ -measurable ensures that for all  $i \in \{1, 2\}$ ,  $j \in \{1, 2, \dots, n\}$  it holds that  $\Omega \ni \omega \mapsto Y_i(e_j, \omega) \in S$  is  $\mathcal{A}_i / \mathcal{S}$ -measurable. This implies that for all  $i \in \{1, 2\}$  it holds that  $\mathbb{Y}_i$  is  $\mathcal{A}_i / \mathcal{S}^{\otimes n}$ -measurable. Moreover, observe that the assumption that for all  $i \in \{1, 2\}$  it holds that  $U_i$  is  $(\mathcal{S} \otimes \mathcal{G}_i) / \mathcal{T}$ -measurable guarantees

that for all  $i \in \{1, 2\}$ ,  $j \in \{1, 2, \dots, n\}$  it holds that  $S^n \times \Omega \ni (s_1, s_2, \dots, s_n, \omega) \mapsto U_i(s_j, \omega) \in \mathcal{T}$  is  $(\mathcal{S}^{\otimes n} \otimes \mathcal{G}_i)/\mathcal{T}$ -measurable. Therefore, we obtain for all  $i \in \{1, 2\}$  that  $\mathbb{U}_i$  is  $(\mathcal{S}^{\otimes n} \otimes \mathcal{G}_i)/\mathcal{T}^{\otimes n}$ -measurable. The fact that  $\mathbb{I}$  is  $\mathcal{T}^{\otimes n}/\mathcal{B}([0, \infty))$ -measurable hence implies that for all  $i \in \{1, 2\}$  it holds that  $\mathbb{I} \circ \mathbb{U}_i$  is  $(\mathcal{S}^{\otimes n} \otimes \mathcal{G}_i)/\mathcal{B}([0, \infty))$ -measurable. Combining this, the fact that for all  $i \in \{1, 2\}$  it holds that  $\mathbb{Y}_i$  is  $\mathcal{A}_i/\mathcal{S}^{\otimes n}$ -measurable, the hypothesis that for every  $i \in \{1, 2\}$  it holds that  $\mathcal{G}_i$  and  $\mathcal{A}_i$  are independent, and Hutzenthaler et al. [36, Lemma 2.2] (applied with  $(\Omega, \mathcal{F}, \mathbb{P}) \leftarrow (\Omega, \mathcal{F}, \mathbb{P})$ ,  $\mathcal{G} \leftarrow \mathcal{G}_i$ ,  $(S, \mathcal{S}) \leftarrow (S^n, \mathcal{S}^{\otimes n})$ ,  $U \leftarrow \mathbb{I} \circ \mathbb{U}_i$ ,  $Y \leftarrow \mathbb{Y}_i$  for  $i \in \{1, 2\}$  in the notation of Hutzenthaler et al. [36, Lemma 2.2]) assure that for all  $i \in \{1, 2\}$  it holds that

$$\begin{aligned} & \mathbb{P}(U_i(Y_i(e_1)) \in B_1, U_i(Y_i(e_2)) \in B_2, \dots, U_i(Y_i(e_n)) \in B_n) \\ &= \mathbb{E}[\mathbb{1}_{B_1 \times B_2 \times \dots \times B_n}(U_i(Y_i(e_1)), U_i(Y_i(e_2)), \dots, U_i(Y_i(e_n)))] = \mathbb{E}[\mathbb{I}(\mathbb{U}_i(\mathbb{Y}_i))] \\ &= \mathbb{E}[(\mathbb{I} \circ \mathbb{U}_i)(\mathbb{Y}_i)] = \int_{S^n} \mathbb{E}[(\mathbb{I} \circ \mathbb{U}_i)(s_1, s_2, \dots, s_n)] (\mathbb{Y}_i(\mathbb{P}))_{\mathcal{S}^{\otimes n}}(d(s_1, s_2, \dots, s_n)). \end{aligned} \quad (19)$$

Moreover, note that the hypothesis that  $U_1$  and  $U_2$  are identically distributed yields that for all  $s_1, s_2, \dots, s_n \in S$  it holds that

$$\begin{aligned} & \mathbb{E}[(\mathbb{I} \circ \mathbb{U}_1)(s_1, s_2, \dots, s_n)] = \mathbb{P}(U_1(s_1) \in B_1, U_1(s_2) \in B_2, \dots, U_1(s_n) \in B_n) \\ &= \mathbb{P}(U_2(s_1) \in B_1, U_2(s_2) \in B_2, \dots, U_2(s_n) \in B_n) = \mathbb{E}[(\mathbb{I} \circ \mathbb{U}_2)(s_1, s_2, \dots, s_n)]. \end{aligned} \quad (20)$$

In addition, observe that the hypothesis that  $Y_1$  and  $Y_2$  are identically distributed ensures that  $(\mathbb{Y}_1(\mathbb{P}))_{\mathcal{S}^{\otimes n}} = (\mathbb{Y}_2(\mathbb{P}))_{\mathcal{S}^{\otimes n}}$ . This, (20), and (19) ensure that

$$\begin{aligned} & \mathbb{P}(U_1(Y_1(e_1)) \in B_1, U_1(Y_1(e_2)) \in B_2, \dots, U_1(Y_1(e_n)) \in B_n) \\ &= \int_{S^n} \mathbb{E}[(\mathbb{I} \circ \mathbb{U}_1)(s_1, s_2, \dots, s_n)] (\mathbb{Y}_1(\mathbb{P}))_{\mathcal{S}^{\otimes n}}(d(s_1, s_2, \dots, s_n)) \\ &= \int_{S^n} \mathbb{E}[(\mathbb{I} \circ \mathbb{U}_2)(s_1, s_2, \dots, s_n)] (\mathbb{Y}_2(\mathbb{P}))_{\mathcal{S}^{\otimes n}}(d(s_1, s_2, \dots, s_n)) \\ &= \mathbb{P}(U_2(Y_2(e_1)) \in B_1, U_2(Y_2(e_2)) \in B_2, \dots, U_2(Y_2(e_n)) \in B_n). \end{aligned} \quad (21)$$

Hence, we obtain that  $U_1(Y_1)$  and  $U_2(Y_2)$  are identically distributed random fields. This establishes item (ii). This completes the proof of Lemma 2.5.  $\square$

**Lemma 2.6.** *Let  $N \in \mathbb{N}$ , let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a probability space, let  $(E, \mathcal{E})$  and  $(S, \mathcal{S})$  be measurable spaces, let  $\mathcal{G}_i \subseteq \mathcal{F}$ ,  $i \in \{1, 2, \dots, N\}$ , be independent sigma-algebras on  $\Omega$ , let  $\mathcal{A}_i \subseteq \mathcal{F}$ ,  $i \in \{1, 2, \dots, N\}$ , be independent sigma-algebras on  $\Omega$ , let  $U_i: S \times \Omega \rightarrow E$ ,  $i \in \{1, 2, \dots, N\}$ , satisfy for all  $i \in \{1, 2, \dots, N\}$  that  $U_i$  is  $(\mathcal{S} \otimes \mathcal{G}_i)/\mathcal{E}$ -measurable, let  $V_i: S \times \Omega \rightarrow E$ ,  $i \in \{1, 2, \dots, N\}$ , satisfy for all  $i \in \{1, 2, \dots, N\}$  that  $V_i$  is  $(\mathcal{S} \otimes \mathcal{A}_i)/\mathcal{E}$ -measurable, and assume for all  $i \in \{1, 2, \dots, N\}$  that  $U_i$  and  $V_i$  are identically distributed random fields. Then it holds that  $(U_1, U_2, \dots, U_N) = (S \times \Omega \ni (s, \omega) \mapsto (U_1(s, \omega), U_2(s, \omega), \dots, U_N(s, \omega)) \in E^N)$  and  $(V_1, V_2, \dots, V_N) = (S \times \Omega \ni (s, \omega) \mapsto (V_1(s, \omega), V_2(s, \omega), \dots, V_N(s, \omega)) \in E^N)$  are identically distributed random fields.*

*Proof of Lemma 2.6.* Throughout this proof let  $n \in \mathbb{N}$ ,  $s_1, s_2, \dots, s_n \in S$ , let  $B_{i,j} \in \mathcal{E}$ ,  $i \in \{1, 2, \dots, N\}$ ,  $j \in \{1, 2, \dots, n\}$ , and let  $\mathbb{U}: S \times \Omega \rightarrow E^N$  and  $\mathbb{V}: S \times \Omega \rightarrow E^N$  satisfy for all  $s \in S$ ,  $\omega \in \Omega$  that  $\mathbb{U}(s, \omega) = (U_1(s, \omega), U_2(s, \omega), \dots, U_N(s, \omega))$  and  $\mathbb{V}(s, \omega) = (V_1(s, \omega), V_2(s, \omega), \dots, V_N(s, \omega))$ . Observe that the assumption that for all  $i \in \{1, 2, \dots, N\}$  it holds that  $U_i: S \times \Omega \rightarrow E$  is  $(\mathcal{S} \otimes \mathcal{G}_i)/\mathcal{E}$ -measurable implies that for all  $i \in \{1, 2, \dots, N\}$ ,  $j \in \{1, 2, \dots, n\}$  it holds that  $\Omega \ni \omega \mapsto U_i(s_j, \omega) \in E$  is  $\mathcal{G}_i/\mathcal{E}$ -measurable. This implies that for all  $i \in \{1, 2, \dots, N\}$  it holds that  $(\bigcap_{j=1}^n \{U_i(s_j) \in B_{i,j}\}) \in \mathcal{G}_i$ . The fact that  $\mathcal{G}_i$ ,  $i \in \{1, 2, \dots, N\}$ , are independent sigma-algebras on

$\Omega$  therefore ensures that

$$\begin{aligned} & \mathbb{P}\left(\mathbb{U}(s_1) \in \left[\bigtimes_{i=1}^N B_{i,1}\right], \mathbb{U}(s_2) \in \left[\bigtimes_{i=1}^N B_{i,2}\right], \dots, \mathbb{U}(s_n) \in \left[\bigtimes_{i=1}^N B_{i,n}\right]\right) \\ &= \mathbb{P}\left(\bigcap_{i=1}^N \bigcap_{j=1}^n \{U_i(s_j) \in B_{i,j}\}\right) = \prod_{i=1}^N \mathbb{P}\left(\bigcap_{j=1}^n \{U_i(s_j) \in B_{i,j}\}\right). \end{aligned} \quad (22)$$

The assumption that for all  $i \in \{1, 2, \dots, N\}$  it holds that  $U_i$  and  $V_i$  are identically distributed hence ensures that

$$\begin{aligned} & \mathbb{P}\left(\mathbb{U}(s_1) \in \left[\bigtimes_{i=1}^N B_{i,1}\right], \mathbb{U}(s_2) \in \left[\bigtimes_{i=1}^N B_{i,2}\right], \dots, \mathbb{U}(s_n) \in \left[\bigtimes_{i=1}^N B_{i,n}\right]\right) \\ &= \prod_{i=1}^N \mathbb{P}\left(\bigcap_{j=1}^n \{V_i(s_j) \in B_{i,j}\}\right). \end{aligned} \quad (23)$$

Combining this with the fact that for all  $i \in \{1, 2, \dots, N\}$  it holds that  $(\bigcap_{j=1}^n \{V_i(s_j) \in B_{i,j}\}) \in \mathcal{A}_i$  and the assumption that  $\mathcal{A}_i$ ,  $i \in \{1, 2, \dots, N\}$ , are independent sigma-algebras on  $\Omega$  yields that

$$\begin{aligned} & \mathbb{P}\left(\mathbb{U}(s_1) \in \left[\bigtimes_{i=1}^N B_{i,1}\right], \mathbb{U}(s_2) \in \left[\bigtimes_{i=1}^N B_{i,2}\right], \dots, \mathbb{U}(s_n) \in \left[\bigtimes_{i=1}^N B_{i,n}\right]\right) \\ &= \mathbb{P}\left(\bigcap_{i=1}^N \bigcap_{j=1}^n \{V_i(s_j) \in B_{i,j}\}\right) \\ &= \mathbb{P}\left(\mathbb{V}(s_1) \in \left[\bigtimes_{i=1}^N B_{i,1}\right], \mathbb{V}(s_2) \in \left[\bigtimes_{i=1}^N B_{i,2}\right], \dots, \mathbb{V}(s_n) \in \left[\bigtimes_{i=1}^N B_{i,n}\right]\right). \end{aligned} \quad (24)$$

Klenke [39, Lemma 1.42] hence ensures that  $\mathbb{U}$  and  $\mathbb{V}$  are identically distributed random fields. This completes the proof of Lemma 2.6.  $\square$

**Corollary 2.7.** *Let  $K, N \in \mathbb{N}$ , let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a probability space, let  $(S, \mathcal{S})$  be a measurable space, let  $\mathcal{G}_i \subseteq \mathcal{F}$ ,  $i \in \{1, 2, \dots, N\}$ , be independent sigma-algebras on  $\Omega$ , let  $\mathcal{A}_i \subseteq \mathcal{F}$ ,  $i \in \{1, 2, \dots, N\}$ , be independent sigma-algebras on  $\Omega$ , let  $U_i: S \times \Omega \rightarrow \mathbb{R}^K$ ,  $i \in \{1, 2, \dots, N\}$ , satisfy for all  $i \in \{1, 2, \dots, N\}$  that  $U_i$  is  $(\mathcal{S} \otimes \mathcal{G}_i)/\mathcal{B}(\mathbb{R}^K)$ -measurable, let  $V_i: S \times \Omega \rightarrow \mathbb{R}^K$ ,  $i \in \{1, 2, \dots, N\}$ , satisfy for all  $i \in \{1, 2, \dots, N\}$  that  $V_i$  is  $(\mathcal{S} \otimes \mathcal{A}_i)/\mathcal{B}(\mathbb{R}^K)$ -measurable, and assume for all  $i \in \{1, 2, \dots, N\}$  that  $U_i$  and  $V_i$  are identically distributed random fields. Then it holds that  $S \times \Omega \ni (s, \omega) \mapsto \sum_{i=1}^N U_i(s, \omega) \in \mathbb{R}^K$  and  $S \times \Omega \ni (s, \omega) \mapsto \sum_{i=1}^N V_i(s, \omega) \in \mathbb{R}^K$  are identically distributed random fields.*

*Proof of Corollary 2.7.* Observe that Lemma 2.6 (applied with  $(E, \mathcal{E}) \leftarrow (\mathbb{R}^K, \mathcal{B}(\mathbb{R}^K))$  in the notation of Lemma 2.6) proves that  $(U_1, U_2, \dots, U_N): S \times \Omega \rightarrow (\mathbb{R}^K)^N$  and  $(V_1, V_2, \dots, V_N): S \times \Omega \rightarrow (\mathbb{R}^K)^N$  are identically distributed random fields. Item (ii) of Lemma 2.5 hence implies that  $S \times \Omega \ni (s, \omega) \mapsto \sum_{i=1}^N U_i(s, \omega) \in \mathbb{R}^K$  and  $S \times \Omega \ni (s, \omega) \mapsto \sum_{i=1}^N V_i(s, \omega) \in \mathbb{R}^K$  are identically distributed random fields. This completes the proof of Corollary 2.7.  $\square$

## 2.4 Distribution properties for MLP approximations

**Lemma 2.8.** *Assume Setting 2.1. Then*

- (i) it holds for all  $\theta \in \Theta$  that  $\{(m, n) \in \mathbb{N}_0 \times \mathbb{N}_0 : n \leq m \leq K\} \times \mathcal{O} \times \Omega \ni (k, l, x, \omega) \mapsto X_l^{\theta, k, x}(\omega) \in \mathcal{O}$  and  $\{(m, n) \in \mathbb{N}_0 \times \mathbb{N}_0 : n \leq m \leq K\} \times \mathcal{O} \times \Omega \ni (k, l, x, \omega) \mapsto X_l^{0, k, x}(\omega) \in \mathcal{O}$  are identically distributed random fields and
- (ii) it holds for all  $\theta, \vartheta \in \Theta$ ,  $k, l, m \in \mathbb{N}_0$  with  $m \leq l \leq k \leq K$  that  $\mathcal{O} \times \Omega \ni (x, \omega) \mapsto X_m^{\vartheta, l, X_l^{\theta, k, x}(\omega)}(\omega) \in \mathcal{O}$  and  $\mathcal{O} \times \Omega \ni (x, \omega) \mapsto X_m^{0, k, x}(\omega) \in \mathcal{O}$  are identically distributed random fields.

*Proof of Lemma 2.8.* Throughout this proof let  $F_l^k : S^K \times \mathcal{O} \rightarrow \mathcal{O}$ ,  $l \in \{0, 1, \dots, k\}$ ,  $k \in \{0, 1, \dots, K\}$ , satisfy for all  $k \in \{0, 1, \dots, K\}$ ,  $l \in \{0, 1, \dots, k\}$ ,  $s = (s_1, s_2, \dots, s_K) \in S^K$ ,  $x \in \mathcal{O}$  that

$$F_l^k(s, x) = \begin{cases} x & : l = k \\ \phi_l(F_{l+1}^k(s, x), s_{l+1}) & : l < k. \end{cases} \quad (25)$$

Observe that induction shows that for all  $k \in \{0, 1, \dots, K\}$ ,  $l \in \{0, 1, \dots, k\}$  it holds that  $F_l^k : S^K \times \mathcal{O} \rightarrow \mathcal{O}$  is  $(\mathcal{S}^{\otimes K} \otimes \mathcal{B}(\mathcal{O}))/\mathcal{B}(\mathcal{O})$ -measurable. Moreover, note that induction, (25), and (3) ensure that for all  $k \in \{0, 1, \dots, K\}$ ,  $l \in \{0, 1, \dots, k\}$ ,  $\theta \in \Theta$ ,  $x \in \mathcal{O}$  it holds that

$$X_l^{\theta, k, x} = F_l^k(W_0^\theta, \dots, W_{K-1}^\theta, x). \quad (26)$$

Next observe that the assumption that  $W_k^\theta : \Omega \rightarrow S$ ,  $\theta \in \Theta$ ,  $k \in \{0, 1, \dots, K-1\}$ , are independent random variables and the assumption that for all  $k \in \{0, 1, \dots, K-1\}$  it holds that  $W_k^\theta : \Omega \rightarrow S$ ,  $\theta \in \Theta$ , are identically distributed implies that for all  $\theta \in \Theta$  it holds that  $(W_l^\theta)_{l \in \{0, 1, \dots, K-1\}}$  and  $(W_l^0)_{l \in \{0, 1, \dots, K-1\}}$  are identically distributed. This and (26) establish item (i). Next we prove item (ii). Note that for all  $\theta, \vartheta \in \Theta$ ,  $k, l, m \in \mathbb{N}_0$  with  $m \leq l \leq k \leq K$  it holds that  $\sigma((W_s^\vartheta)_{s \in [m, l] \cap \mathbb{N}_0})$  and  $\sigma((W_s^\theta)_{s \in [l, k] \cap \mathbb{N}_0})$  are independent. This, item (i), item (i) of Lemma 2.2, and item (ii) of Lemma 2.5 demonstrate that for all  $\theta, \vartheta \in \Theta$ ,  $k, l, m \in \mathbb{N}_0$  with  $m \leq l \leq k \leq K$  it holds that

$$\mathcal{O} \times \Omega \ni (x, \omega) \mapsto X_m^{\vartheta, l, X_l^{\theta, k, x}(\omega)}(\omega) \in \mathcal{O} \quad \text{and} \quad \mathcal{O} \times \Omega \ni (x, \omega) \mapsto X_m^{0, l, X_l^{0, k, x}(\omega)}(\omega) \in \mathcal{O} \quad (27)$$

are identically distributed random fields. Furthermore, observe that (3) and induction imply that for all  $k, l, m \in \mathbb{N}_0$ ,  $x \in \mathcal{O}$ ,  $\omega \in \Omega$  with  $m \leq l \leq k \leq K$  it holds that

$$X_m^{0, l, X_l^{0, k, x}(\omega)}(\omega) = X_m^{0, k, x}(\omega). \quad (28)$$

Combining this with (27) establishes item (ii). This completes the proof of Lemma 2.8.  $\square$

**Proposition 2.9.** Assume Setting 2.1. Then

- (i) it holds for all  $n \in \mathbb{N}_0$ ,  $\theta \in \Theta$  that  $\{0, 1, \dots, K\} \times \mathcal{O} \times \Omega \ni (k, x, \omega) \mapsto V_{k, n}^\theta(x, \omega) \in \mathbb{R}$  and  $\{0, 1, \dots, K\} \times \mathcal{O} \times \Omega \ni (k, x, \omega) \mapsto V_{k, n}^0(x, \omega) \in \mathbb{R}$  are identically distributed random fields,
- (ii) it holds for all  $n \in \mathbb{N}$ ,  $\theta, \vartheta \in \Theta$  with  $\{(\theta, \eta) \in \Theta : \eta \in \Theta\} \cap \{(\vartheta, \eta) \in \Theta : \eta \in \Theta\} = \emptyset$  that  $\{0, 1, \dots, K\} \times \mathcal{O} \times \Omega \ni (k, x, \omega) \mapsto (V_{k, n}^\theta(x, \omega), V_{k, n-1}^\vartheta(x, \omega)) \in \mathbb{R}^2$  and  $\{0, 1, \dots, K\} \times \mathcal{O} \times \Omega \ni (k, x, \omega) \mapsto (V_{k, n}^0(x, \omega), V_{k, n-1}^1(x, \omega)) \in \mathbb{R}^2$  are identically distributed random fields, and
- (iii) it holds for all  $n \in \mathbb{N}$ ,  $\theta, \vartheta \in \Theta$  with  $\{(\theta, \eta) \in \Theta : \eta \in \Theta\} \cap \{(\vartheta, \eta) \in \Theta : \eta \in \Theta\} = \emptyset$  that

$$\begin{aligned} & \{1, 2, \dots, K\} \times \mathcal{O} \times \Omega \ni (k, x, \omega) \mapsto \\ & \frac{1}{\mathbf{p}_{k, \mathcal{R}_k^\theta(\omega)}} \left[ f_{\mathcal{R}_k^\theta(\omega)}(X_{\mathcal{R}_k^\theta(\omega)}^{\theta, k, x}(\omega), V_{\mathcal{R}_k^\theta(\omega), n}^\theta(X_{\mathcal{R}_k^\theta(\omega)}^{\theta, k, x}(\omega), \omega)) - V_{\mathcal{R}_k^\theta(\omega), n}^\theta(X_{\mathcal{R}_k^\theta(\omega)}^{\theta, k, x}(\omega), \omega) \right. \\ & \left. - f_{\mathcal{R}_k^\theta(\omega)}(X_{\mathcal{R}_k^\theta(\omega)}^{\theta, k, x}(\omega), V_{\mathcal{R}_k^\theta(\omega), n-1}^\vartheta(X_{\mathcal{R}_k^\theta(\omega)}^{\theta, k, x}(\omega), \omega)) + V_{\mathcal{R}_k^\theta(\omega), n-1}^\vartheta(X_{\mathcal{R}_k^\theta(\omega)}^{\theta, k, x}(\omega), \omega) \right] \in \mathbb{R} \quad (29) \end{aligned}$$

and

$$\begin{aligned} & \{1, 2, \dots, K\} \times \mathcal{O} \times \Omega \ni (k, x, \omega) \mapsto \\ & \frac{1}{\mathfrak{p}_{k, \mathcal{R}_k^0(\omega)}} \left[ f_{\mathcal{R}_k^0(\omega)}(X_{\mathcal{R}_k^0(\omega)}^{0,k,x}(\omega), V_{\mathcal{R}_k^0(\omega),n}^0(X_{\mathcal{R}_k^0(\omega)}^{0,k,x}(\omega), \omega)) - V_{\mathcal{R}_k^0(\omega),n}^0(X_{\mathcal{R}_k^0(\omega)}^{0,k,x}(\omega), \omega) \right. \\ & \left. - f_{\mathcal{R}_k^0(\omega)}(X_{\mathcal{R}_k^0(\omega)}^{0,k,x}(\omega), V_{\mathcal{R}_k^0(\omega),n-1}^1(X_{\mathcal{R}_k^0(\omega)}^{0,k,x}(\omega), \omega)) + V_{\mathcal{R}_k^0(\omega),n-1}^1(X_{\mathcal{R}_k^0(\omega)}^{k,0}(x, \omega), \omega) \right] \in \mathbb{R} \quad (30) \end{aligned}$$

are identically distributed random fields.

*Proof of Proposition 2.9.* Throughout this proof let  $h_k: \mathcal{O} \times \mathbb{R} \rightarrow \mathbb{R}$ ,  $k \in \{0, 1, \dots, K-1\}$ , satisfy for all  $k \in \{0, 1, \dots, K-1\}$ ,  $x \in \mathcal{O}$ ,  $a \in \mathbb{R}$  that  $h_k(x, a) = f_k(x, a) - a$ , let  $\mathcal{A}_J^I \subseteq \mathcal{F}$ ,  $I, J \in 2^\Theta$ , satisfy for all  $I, J \in 2^\Theta$  that  $\mathcal{A}_J^I = \sigma((\mathcal{R}_s^\theta)_{(s,\theta) \in \{1,2,\dots,K\} \times I}, (W_s^\theta)_{(s,\theta) \in \{0,1,\dots,K-1\} \times J})$ , let  $\mathcal{X} \subseteq \mathbb{N}_0$  satisfy that

$$\mathcal{X} = \left\{ n \in \mathbb{N}_0 : \left( \begin{array}{c} \text{It holds for all } \theta \in \Theta \text{ that} \\ \{0, 1, \dots, K\} \times \mathcal{O} \times \Omega \ni (k, x, \omega) \mapsto V_{k,n}^\theta(x, \omega) \in \mathbb{R} \text{ and} \\ \{0, 1, \dots, K\} \times \mathcal{O} \times \Omega \ni (k, x, \omega) \mapsto V_{k,n}^0(x, \omega) \in \mathbb{R} \\ \text{are identically distributed random fields.} \end{array} \right) \right\}, \quad (31)$$

let  $\mathcal{Y} \subseteq \mathbb{N}$  satisfy that

$$\mathcal{Y} = \left\{ n \in \mathbb{N} : \left( \begin{array}{c} \text{It holds for all } \theta, \vartheta \in \Theta \text{ with} \\ \{(\theta, \eta) \in \Theta : \eta \in \Theta\} \cap \{(\vartheta, \eta) \in \Theta : \eta \in \Theta\} = \emptyset \\ \text{that } \{0, 1, \dots, K\} \times \mathcal{O} \times \Omega \ni (k, x, \omega) \mapsto \\ (V_{k,n}^\theta(x, \omega), V_{k,n-1}^\vartheta(x, \omega)) \in \mathbb{R}^2 \text{ and} \\ \{0, 1, \dots, K\} \times \mathcal{O} \times \Omega \ni (k, x, \omega) \mapsto (V_{k,n}^0(x, \omega), V_{k,n-1}^1(x, \omega)) \in \mathbb{R}^2 \\ \text{are identically distributed random fields.} \end{array} \right) \right\}, \quad (32)$$

and let  $\mathcal{Z} \subseteq \mathbb{N}$  satisfy that

$$\mathcal{Z} = \left\{ n \in \mathbb{N} : \left( \begin{array}{c} \text{It holds for all } \theta, \vartheta \in \Theta \text{ with} \\ \{(\theta, \eta) \in \Theta : \eta \in \Theta\} \cap \{(\vartheta, \eta) \in \Theta : \eta \in \Theta\} = \emptyset \\ \text{that } \{1, 2, \dots, K\} \times \mathcal{O} \times \Omega \ni (k, x, \omega) \mapsto \\ \frac{1}{\mathfrak{p}_{k, \mathcal{R}_k^\theta(\omega)}} [h_{\mathcal{R}_k^\theta(\omega)}(X_{\mathcal{R}_k^\theta(\omega)}^{\theta,k,x}(\omega), V_{\mathcal{R}_k^\theta(\omega),n}^\theta(X_{\mathcal{R}_k^\theta(\omega)}^{\theta,k,x}(\omega), \omega)) \\ - h_{\mathcal{R}_k^\theta(\omega)}(X_{\mathcal{R}_k^\theta(\omega)}^{\theta,k,x}(\omega), V_{\mathcal{R}_k^\theta(\omega),n-1}^\vartheta(X_{\mathcal{R}_k^\theta(\omega)}^{\theta,k,x}(\omega), \omega))] \in \mathbb{R} \\ \text{and } \{0, 1, \dots, K\} \times \mathcal{O} \times \Omega \ni (k, x, \omega) \mapsto \\ \frac{1}{\mathfrak{p}_{k, \mathcal{R}_k^0(\omega)}} [h_{\mathcal{R}_k^0(\omega)}(X_{\mathcal{R}_k^0(\omega)}^{0,k,x}(\omega), V_{\mathcal{R}_k^0(\omega),n}^0(X_{\mathcal{R}_k^0(\omega)}^{0,k,x}(\omega), \omega)) \\ - h_{\mathcal{R}_k^0(\omega)}(X_{\mathcal{R}_k^0(\omega)}^{0,k,x}(\omega), V_{\mathcal{R}_k^0(\omega),n-1}^1(X_{\mathcal{R}_k^0(\omega)}^{0,k,x}(\omega), \omega))] \in \mathbb{R} \\ \text{are identically distributed random fields.} \end{array} \right) \right\}. \quad (33)$$

Observe that the assumption that for all  $\theta \in \Theta$ ,  $k \in \{0, 1, \dots, K\}$ ,  $x \in \mathcal{O}$  it holds that  $V_{k,0}^\theta(x) = 0$  ensures that  $0 \in \mathcal{X}$ . Next we prove that  $1 \in \mathcal{X}$ . For this note that (4) yields that for all  $\theta \in \Theta$ ,  $k \in \{0, 1, \dots, K\}$ ,  $x \in \mathcal{O}$  it holds that

$$V_{k,1}^\theta(x) = \frac{1}{M} \sum_{m=1}^M \left[ g(X_0^{(\theta,0,-m),k,x}) + \sum_{l=0}^{K-1} \mathbb{1}_{[0,k)}(l) f_l(X_l^{(\theta,0,m),k,x}, 0) \right]. \quad (34)$$

Moreover, observe that item (i) of Lemma 2.8 ensures that

- (I) it holds for all  $\theta \in \Theta$ ,  $m \in \mathbb{N}$  that  $\{0, 1, \dots, K\} \times \mathcal{O} \times \Omega \ni (k, x, \omega) \mapsto g(X_0^{(\theta,0,-m),k,x}(\omega)) \in \mathbb{R}$  and  $\{0, 1, \dots, K\} \times \mathcal{O} \times \Omega \ni (k, x, \omega) \mapsto g(X_0^{(0,0,-m),k,x}(\omega)) \in \mathbb{R}$  are identically distributed random fields and

(II) it holds for all  $\theta \in \Theta$ ,  $m \in \mathbb{N}$  that  $\{1, 2, \dots, K\} \times \mathcal{O} \times \Omega \ni (k, x, \omega) \mapsto \mathbb{1}_{[0, k-1]}(l) f_l(X_l^{(\theta, 0, m), k, x}(\omega), 0) \in \mathbb{R}$  and  $\{1, 2, \dots, K\} \times \mathcal{O} \times \Omega \ni (k, x, \omega) \mapsto \mathbb{1}_{[0, k-1]}(l) f_l(X_l^{(0, 0, m), k, x}(\omega), 0) \in \mathbb{R}$  are identically distributed random fields.

Item (i) of Lemma 2.2, Corollary 2.7, and the fact that  $\sigma((W_s^\vartheta)_{s \in \{0, 1, \dots, K-1\}})$ ,  $\vartheta \in \Theta$ , are independent sigma-algebras on  $\Omega$  therefore prove that for all  $\theta \in \Theta$ ,  $m \in \mathbb{N}$  it holds that

$$\begin{aligned} \{1, 2, \dots, K\} \times \mathcal{O} \times \Omega \ni (k, x, \omega) \mapsto g(X_0^{(\theta, 0, -m), k, x}(\omega)) \\ + \sum_{l=0}^{K-1} \mathbb{1}_{[0, k-1]}(l) f_l(X_l^{(\theta, 0, m), k, x}(\omega), 0) \in \mathbb{R} \end{aligned} \quad (35)$$

and

$$\begin{aligned} \{1, 2, \dots, K\} \times \mathcal{O} \times \Omega \ni (k, x, \omega) \mapsto g(X_0^{(0, 0, -m), k, x}(\omega)) \\ + \sum_{l=0}^{K-1} \mathbb{1}_{[0, k-1]}(l) f_l(X_l^{(0, 0, m), k, x}(\omega), 0) \in \mathbb{R} \end{aligned} \quad (36)$$

are identically distributed random fields. Item (i) of Lemma 2.2, (34), Corollary 2.7, and the fact that  $\sigma((W_s^\vartheta)_{s \in \{0, 1, \dots, K-1\}})$ ,  $\vartheta \in \Theta$ , are independent sigma-algebras on  $\Omega$  hence ensure that for all  $\theta \in \Theta$  it holds that  $\{0, 1, \dots, K\} \times \mathcal{O} \times \Omega \ni (k, x, \omega) \mapsto V_{k,1}^\theta(x, \omega) \in \mathbb{R}$  and  $\{0, 1, \dots, K\} \times \mathcal{O} \times \Omega \ni (k, x, \omega) \mapsto V_{k,1}^0(x, \omega) \in \mathbb{R}$  are identically distributed random fields. Therefore, we obtain that  $1 \in \mathcal{X}$ . Next we prove that  $\{n \in \mathcal{X} : n-1 \in \mathcal{X}\} \subseteq \mathcal{Y}$ . Note that item (i) of Lemma 2.3 ensures that for all  $n \in \mathbb{N}_0$ ,  $\theta \in \Theta$  it holds that  $\{0, 1, \dots, K\} \times \mathcal{O} \times \Omega \ni (k, x, \omega) \mapsto V_{k,n}^\theta(x, \omega) \in \mathbb{R}$  is  $(2^{\{0, 1, \dots, K\}} \otimes \mathcal{B}(\mathcal{O}) \otimes \mathcal{A}_{\{(\theta, \eta) \in \Theta : \eta \in \Theta\}}^{\{(\theta, \eta) \in \Theta : \eta \in \Theta\}}) / \mathcal{B}(\mathbb{R})$ -measurable. The fact that for all  $\theta, \vartheta \in \Theta$  with  $\{(\theta, \eta) : \eta \in \Theta\} \cap \{(\vartheta, \eta) : \eta \in \Theta\} = \emptyset$  it holds that  $\mathcal{A}_{\{(\theta, \eta) : \eta \in \Theta\}}^{\{(\theta, \eta) : \eta \in \Theta\}}$  and  $\mathcal{A}_{\{(\vartheta, \eta) : \eta \in \Theta\}}^{\{(\vartheta, \eta) : \eta \in \Theta\}}$  are independent and Lemma 2.6 hence demonstrate that for all  $n \in \mathcal{X}$ ,  $\theta, \vartheta \in \Theta$  with  $\{(\theta, \eta) \in \Theta : \eta \in \Theta\} \cap \{(\vartheta, \eta) \in \Theta : \eta \in \Theta\} = \emptyset$  and  $n-1 \in \mathcal{X}$  it holds that  $\{0, 1, \dots, K\} \times \mathcal{O} \times \Omega \ni (k, x, \omega) \mapsto (V_{k,n}^\theta(x, \omega), V_{k,n-1}^\vartheta(x, \omega)) \in \mathbb{R}^2$  and  $\{0, 1, \dots, K\} \times \mathcal{O} \times \Omega \ni (k, x, \omega) \mapsto (V_{k,n}^0(x, \omega), V_{k,n-1}^1(x, \omega)) \in \mathbb{R}^2$  are identically distributed random fields. Therefore, we obtain that  $\{n \in \mathcal{X} : n-1 \in \mathcal{X}\} \subseteq \mathcal{Y}$ . Combining this with the fact that  $\{0, 1\} \subseteq \mathcal{X}$  ensures that  $1 \in \mathcal{Y}$ . Next we prove that  $\mathcal{Y} \subseteq \mathcal{Z}$ . For this note that item (i) of Lemma 2.8 and Lemma 2.5 prove that for all  $\theta \in \Theta$  it holds that  $\{1, 2, \dots, K\} \times \mathcal{O} \times \Omega \ni (k, x, \omega) \mapsto (\mathcal{R}_k^\theta(\omega), X_{\mathcal{R}_k^\theta(\omega)}^{\theta, k, x}(\omega)) \in \{0, 1, \dots, K-1\} \times \mathcal{O}$  and  $\{1, 2, \dots, K\} \times \mathcal{O} \times \Omega \ni (k, x, \omega) \mapsto (\mathcal{R}_k^0(\omega), X_{\mathcal{R}_k^0(\omega)}^{0, k, x}(\omega)) \in \{0, 1, \dots, K-1\} \times \mathcal{O}$  are identically distributed random fields. Item (i) of Lemma 2.2, item (i) of Lemma 2.3, Lemma 2.5, and the fact that for all  $\theta, \vartheta \in \Theta$  with  $\theta \notin \{(\vartheta, \eta) \in \Theta : \eta \in \Theta\}$  it holds that  $\sigma((\mathcal{R}_s^\theta)_{s \in \{1, 2, \dots, K\}}, (W_s^\theta)_{s \in \{0, 1, \dots, K-1\}})$  and  $\sigma((\mathcal{R}_s^{\vartheta, \eta})_{(s, \eta) \in \{1, 2, \dots, K\} \times \Theta}, (W_s^{\vartheta, \eta})_{(s, \eta) \in \{0, 1, \dots, K-1\} \times \Theta})$  are independent therefore demonstrate that for all  $n \in \mathcal{Y}$ ,  $\theta, \vartheta \in \Theta$  with  $\{(\theta, \eta) \in \Theta : \eta \in \Theta\} \cap \{(\vartheta, \eta) \in \Theta : \eta \in \Theta\} = \emptyset$  it holds that

$$\begin{aligned} \{1, 2, \dots, K\} \times \mathcal{O} \times \Omega \ni (k, x, \omega) \mapsto \frac{1}{\mathfrak{p}_{k, \mathcal{R}_k^\theta(\omega)}} \left[ h_{\mathcal{R}_k^\theta(\omega)}(X_{\mathcal{R}_k^\theta(\omega)}^{\theta, k, x}(\omega), V_{\mathcal{R}_k^\theta(\omega), n}^\theta(X_{\mathcal{R}_k^\theta(\omega)}^{\theta, k, x}(\omega), \omega)) \right. \\ \left. - h_{\mathcal{R}_k^\theta(\omega)}(X_{\mathcal{R}_k^\theta(\omega)}^{\theta, k, x}(\omega), V_{\mathcal{R}_k^\theta(\omega), n-1}^\vartheta(X_{\mathcal{R}_k^\theta(\omega)}^{\theta, k, x}(\omega), \omega)) \right] \in \mathbb{R} \end{aligned} \quad (37)$$

and

$$\begin{aligned} \{1, 2, \dots, K\} \times \mathcal{O} \times \Omega \ni (k, x, \omega) \mapsto \frac{1}{\mathfrak{p}_{k, \mathcal{R}_k^0(\omega)}} \left[ h_{\mathcal{R}_k^0(\omega)}(X_{\mathcal{R}_k^0(\omega)}^{0, k, x}(\omega), V_{\mathcal{R}_k^0(\omega), n}^0(X_{\mathcal{R}_k^0(\omega)}^{0, k, x}(\omega), \omega)) \right. \\ \left. - h_{\mathcal{R}_k^0(\omega)}(X_{\mathcal{R}_k^0(\omega)}^{0, k, x}(\omega), V_{\mathcal{R}_k^0(\omega), n-1}^1(X_{\mathcal{R}_k^0(\omega)}^{0, k, x}(\omega), \omega)) \right] \in \mathbb{R} \end{aligned} \quad (38)$$

are identically distributed random fields. Hence, we obtain that  $\mathcal{Y} \subseteq \mathcal{Z}$ . Combining this with the fact that  $1 \in \mathcal{Y}$  demonstrates that  $1 \in \mathcal{Z}$ . Next we prove that  $\{n \in \mathbb{N} \cap [2, \infty) : \{1, 2, \dots, n-1\} \subseteq \mathcal{Z}\} \subseteq \mathcal{X}$ . Note that item (ii) of Lemma 2.3 demonstrates that for all  $n \in \mathbb{N}$ ,  $\theta, \vartheta \in \Theta$  it holds that

$$\{1, 2, \dots, K\} \times \mathcal{O} \times \Omega \ni (k, x, \omega) \mapsto \frac{1}{\mathbf{p}_{k, \mathcal{R}_k^\theta(\omega)}} \left[ h_{\mathcal{R}_k^\theta(\omega)}(X_{\mathcal{R}_k^\theta(\omega)}^{\theta, k, x}(\omega), V_{\mathcal{R}_k^\theta(\omega), n}^\theta(X_{\mathcal{R}_k^\theta(\omega)}^{\theta, k, x}(\omega), \omega)) \right. \\ \left. - h_{\mathcal{R}_k^\theta(\omega)}(X_{\mathcal{R}_k^\theta(\omega)}^{\theta, k, x}(\omega), V_{\mathcal{R}_k^\theta(\omega), n-1}^\theta(X_{\mathcal{R}_k^\theta(\omega)}^{\theta, k, x}(\omega), \omega)) \right] \in \mathbb{R} \quad (39)$$

is  $((2^{\{0, 1, \dots, K\}} \otimes \mathcal{B}(\mathcal{O})) \otimes \mathcal{A}_{\cup_{\eta \in \Theta} \{\theta, (\theta, \eta), (\vartheta, \eta)\}}^{\cup_{\eta \in \Theta} \{\theta, (\theta, \eta), (\vartheta, \eta)\}}) / \mathcal{B}(\mathbb{R})$ -measurable. Combining this and the fact that for all  $\theta \in \Theta$ ,  $j \in \mathbb{N}$  it holds that  $\mathcal{A}_{\cup_{\eta \in \Theta} \{(\theta, j, m), (\theta, j, m, \eta), (\theta, j, -m, \eta)\}}^{\cup_{\eta \in \Theta} \{(\theta, j, m), (\theta, j, m, \eta), (\theta, j, -m, \eta)\}}$ ,  $m \in \mathbb{N}$ , are independent sigma-algebras on  $\Omega$  with Corollary 2.7 proves that for all  $n \in \{k \in [2, \infty) \cap \mathbb{N} : \{1, 2, \dots, k-1\} \subseteq \mathcal{Z}\}$ ,  $j \in \{1, 2, \dots, n-1\}$ ,  $\theta \in \Theta$  it holds that

$$\{1, 2, \dots, K\} \times \mathcal{O} \times \Omega \ni (k, x, \omega) \mapsto \sum_{m=1}^{M^{n-j}} \frac{1}{\mathbf{p}_{k, \mathcal{R}_k^{(\theta, j, m)}(\omega)}} \left[ h_{\mathcal{R}_k^{(\theta, j, m)}(\omega)}(X_{\mathcal{R}_k^{(\theta, j, m)}(\omega)}^{(\theta, j, m), k, x}(\omega), V_{\mathcal{R}_k^{(\theta, j, m)}(\omega), j}^{(\theta, j, m)}(X_{\mathcal{R}_k^{(\theta, j, m)}(\omega)}^{(\theta, j, m), k, x}(\omega), \omega)) \right. \\ \left. - h_{\mathcal{R}_k^{(\theta, j, m)}(\omega)}(X_{\mathcal{R}_k^{(\theta, j, m)}(\omega)}^{(\theta, j, m), k, x}(\omega), V_{\mathcal{R}_k^{(\theta, j, m)}(\omega), j-1}^{(\theta, j, m)}(X_{\mathcal{R}_k^{(\theta, j, m)}(\omega)}^{(\theta, j, m), k, x}(\omega), \omega)) \right] \in \mathbb{R} \quad (40)$$

and

$$\{1, 2, \dots, K\} \times \mathcal{O} \times \Omega \ni (k, x, \omega) \mapsto \sum_{m=1}^{M^{n-j}} \frac{1}{\mathbf{p}_{k, \mathcal{R}_k^{(0, j, m)}(\omega)}} \left[ h_{\mathcal{R}_k^{(0, j, m)}(\omega)}(X_{\mathcal{R}_k^{(0, j, m)}(\omega)}^{(0, j, m), k, x}(\omega), V_{\mathcal{R}_k^{(0, j, m)}(\omega), j}^{(0, j, m)}(X_{\mathcal{R}_k^{(0, j, m)}(\omega)}^{(0, j, m), k, x}(\omega), \omega)) \right. \\ \left. - h_{\mathcal{R}_k^{(0, j, m)}(\omega)}(X_{\mathcal{R}_k^{(0, j, m)}(\omega)}^{(0, j, m), k, x}(\omega), V_{\mathcal{R}_k^{(0, j, m)}(\omega), j-1}^{(0, j, m)}(X_{\mathcal{R}_k^{(0, j, m)}(\omega)}^{(0, j, m), k, x}(\omega), \omega)) \right] \in \mathbb{R} \quad (41)$$

are identically distributed random fields. This, (4), the fact that for all  $n \in \mathbb{N}$ ,  $\theta \in \Theta$  it holds that  $\{0, 1, \dots, K\} \times \mathcal{O} \times \Omega \ni (k, x, \omega) \mapsto \frac{1}{M^n} \sum_{m=1}^{M^n} [g(X_0^{(\theta, 0, -m), k, x}(\omega)) + \sum_{l=0}^{k-1} f_l(X_l^{(\theta, 0, m), k, x}(\omega), 0)] \in \mathbb{R}$  and  $\{0, 1, \dots, K\} \times \mathcal{O} \times \Omega \ni (k, x, \omega) \mapsto \frac{1}{M^n} \sum_{m=1}^{M^n} [g(X_0^{(0, 0, -m), k, x}(\omega)) + \sum_{l=0}^{k-1} f_l(X_l^{(0, 0, m), k, x}(\omega), 0)] \in \mathbb{R}$  are identically distributed random fields, item (i) of Lemma 2.2, item (i) of Lemma 2.3, and Corollary 2.7 demonstrate that for all  $n \in \{k \in [2, \infty) \cap \mathbb{N} : \{1, \dots, k-1\} \subseteq \mathcal{Z}\}$ ,  $\theta \in \Theta$  it holds that  $\{0, 1, \dots, K\} \times \mathcal{O} \times \Omega \ni (k, x, \omega) \mapsto V_{k, n}^\theta(x, \omega) \in \mathbb{R}$  and  $\{0, 1, \dots, K\} \times \mathcal{O} \times \Omega \ni (k, x, \omega) \mapsto V_{k, n}^0(x, \omega) \in \mathbb{R}$  are identically distributed random fields. Hence, we obtain that  $\{n \in [2, \infty) \cap \mathbb{N} : \{1, 2, \dots, n-1\} \subseteq \mathcal{Z}\} \subseteq \mathcal{X}$ . This, the fact that  $\{0, 1\} \subseteq \mathcal{X}$ , the fact that  $\{n \in \mathcal{X} : n-1 \in \mathcal{X}\} \subseteq \mathcal{Y}$ , the fact that  $\mathcal{Y} \subseteq \mathcal{Z}$ , and induction prove that  $\mathcal{X} = \mathbb{N}_0$ ,  $\mathcal{Y} = \mathbb{N}$ , and  $\mathcal{Z} = \mathbb{N}$ . Note that the fact that  $\mathcal{X} = \mathbb{N}_0$  establishes item (i). Moreover, observe that the fact that  $\mathcal{Y} = \mathbb{N}$  establishes item (ii). Furthermore, note that the fact that  $\mathcal{Z} = \mathbb{N}$  establishes item (iii). This completes the proof of Proposition 2.9.  $\square$

**Corollary 2.10.** Assume Setting 2.1. Then it holds for all  $k, l, n \in \mathbb{N}_0$ ,  $\theta, \vartheta \in \Theta$  with  $l \leq k \leq K$  and  $\theta \notin \{(\vartheta, \eta) \in \Theta : \eta \in \Theta\}$  that  $\mathcal{O} \times \Omega \ni (x, \omega) \mapsto (X_l^{\theta, k, x}(\omega), V_{l, n}^\vartheta(X_l^{\theta, k, x}(\omega), \omega)) \in \mathcal{O} \times \mathbb{R}$  and  $\mathcal{O} \times \Omega \ni (x, \omega) \mapsto (X_l^{0, k, x}(\omega), V_{l, n}^0(X_l^{0, k, x}(\omega), \omega)) \in \mathcal{O} \times \mathbb{R}$  are identically distributed random fields.

*Proof of Corollary 2.10.* First, observe that item (i) of Proposition 2.9 demonstrates that for all  $n \in \mathbb{N}_0$ ,  $l \in \{0, 1, \dots, K\}$ ,  $\vartheta \in \Theta$  it holds that  $\mathcal{O} \times \Omega \ni (y, \omega) \mapsto (y, V_{l, n}^\vartheta(y, \omega)) \in \mathcal{O} \times \mathbb{R}$  and  $\mathcal{O} \times \Omega \ni (y, \omega) \mapsto (y, V_{l, n}^0(y, \omega)) \in \mathcal{O} \times \mathbb{R}$  are identically distributed random fields. Moreover,

note that item (i) of Lemma 2.8 ensures that for all  $k, l \in \mathbb{N}_0$ ,  $\theta \in \Theta$  with  $l \leq k \leq K$  it holds that  $\mathcal{O} \times \Omega \ni (x, \omega) \mapsto X_l^{\theta, k, x}(\omega) \in \mathcal{O}$  and  $\mathcal{O} \times \Omega \ni (x, \omega) \mapsto X_l^{0, k, x}(\omega) \in \mathcal{O}$  are identically distributed random fields. Combining the fact that for all  $n \in \mathbb{N}_0$ ,  $l \in \{0, 1, \dots, K\}$ ,  $\vartheta \in \Theta$  it holds that  $\mathcal{O} \times \Omega \ni (y, \omega) \mapsto (y, V_{l,n}^\vartheta(y, \omega)) \in \mathcal{O} \times \mathbb{R}$  and  $\mathcal{O} \times \Omega \ni (y, \omega) \mapsto (y, V_{l,n}^0(y, \omega)) \in \mathcal{O} \times \mathbb{R}$  are identically distributed random fields, item (i) of Lemma 2.2, item (i) of Lemma 2.3, and Lemma 2.5 with the fact that for all  $\theta, \vartheta \in \Theta$  with  $\theta \notin \{(\vartheta, \eta) \in \Theta : \eta \in \Theta\}$  it holds that  $\sigma((W_s^\theta)_{s \in \{0, 1, \dots, K-1\}})$  and  $\sigma((\mathcal{R}_s^{(\vartheta, \eta)})_{(s, \eta) \in \{1, 2, \dots, K\} \times \Theta}, (W_s^{(\vartheta, \eta)})_{(s, \eta) \in \{0, 1, \dots, K-1\} \times \Theta})$  are independent hence proves that for all  $k, l, n \in \mathbb{N}_0$ ,  $\theta, \vartheta \in \Theta$  with  $l \leq k \leq K$  and  $\theta \notin \{(\vartheta, \eta) \in \Theta : \eta \in \Theta\}$  it holds that  $\mathcal{O} \times \Omega \ni (x, \omega) \mapsto (X_l^{\theta, k, x}(\omega), V_{l,n}^\vartheta(X_l^{\theta, k, x}(\omega), \omega)) \in \mathcal{O} \times \mathbb{R}$  and  $\mathcal{O} \times \Omega \ni (x, \omega) \mapsto (X_l^{0, k, x}(\omega), V_{l,n}^0(X_l^{0, k, x}(\omega), \omega)) \in \mathcal{O} \times \mathbb{R}$  are identically distributed random fields. This completes the proof of Corollary 2.10.  $\square$

## 2.5 Integrability properties for MLP approximations

**Lemma 2.11** (Integrability properties). *Assume Setting 2.1, assume for all  $k \in \{0, 1, \dots, K\}$ ,  $x \in \mathcal{O}$  that  $\mathbb{E}[|g(X_0^{0, k, x})|^2 + \sum_{l=0}^{k-1} |f_l(X_l^{0, k, x})|^2] < \infty$ , and let  $L_0, L_1, \dots, L_K \in \mathbb{R}$  satisfy for all  $k \in \{0, 1, \dots, K-1\}$ ,  $x \in \mathcal{O}$ ,  $a, b \in \mathbb{R}$  that  $|(f_k(x, a) - a) - (f_k(x, b) - b)| \leq L_k |a - b|$ . Then it holds for all  $k, l, n \in \mathbb{N}_0$ ,  $\theta, \vartheta \in \Theta$ ,  $x \in \mathcal{O}$  with  $l \leq k \leq K$  and  $\vartheta \notin \{(\theta, \eta) \in \Theta : \eta \in \Theta\}$  that  $\mathbb{E}[|V_{l,n}^\theta(X_l^{\vartheta, k, x})|^2] < \infty$ .*

*Proof of Lemma 2.11.* Throughout this proof let  $\mathbb{A} \subseteq \mathbb{N}$  satisfy that

$$\mathbb{A} = \left\{ n \in \mathbb{N} : \begin{pmatrix} \text{It holds for all } j, k, l \in \mathbb{N}_0, \theta, \vartheta \in \Theta, x \in \mathcal{O} \text{ with} \\ 0 \leq l \leq k \leq K, \vartheta \notin \{(\theta, \eta) \in \Theta : \eta \in \Theta\}, \text{ and } 0 \leq j \leq n-1 \\ \text{that } \mathbb{E}[|V_{l,j}^\theta(X_l^{\vartheta, k, x})|^2] < \infty. \end{pmatrix} \right\}. \quad (42)$$

Note that the assumption that for all  $k \in \{0, 1, \dots, K\}$ ,  $\theta \in \Theta$ ,  $x \in \mathcal{O}$  it holds that  $V_{k,0}^\theta(x) = 0$  yields that  $1 \in \mathbb{A}$ . Next observe that (4) ensures that for all  $k, l, n \in \mathbb{N}$ ,  $\theta, \vartheta \in \Theta$ ,  $x \in \mathcal{O}$  with  $l \leq k \leq K$  it holds that

$$\begin{aligned} & V_{l,n}^\theta(X_l^{\vartheta, k, x}) \\ &= \frac{1}{M^n} \sum_{m=1}^{M^n} \left[ g(X_0^{(\theta, 0, -m), l, X_l^{\vartheta, k, x}}) + \sum_{s=0}^{l-1} f_s(X_s^{(\theta, 0, m), l, X_l^{\vartheta, k, x}}, 0) \right] + \sum_{j=1}^{n-1} \frac{1}{M^{n-j}} \sum_{m=1}^{M^{n-j}} \frac{1}{\mathfrak{p}_{l, \mathcal{R}_l^{(\theta, j, m)}}} \\ & \cdot \left[ \left( f_{\mathcal{R}_l^{(\theta, j, m)}} \left( X_{\mathcal{R}_l^{(\theta, j, m)}}^{(\theta, j, m), l, X_l^{\vartheta, k, x}}, V_{\mathcal{R}_l^{(\theta, j, m)}, j}^{(\theta, j, m)}(X_{\mathcal{R}_l^{(\theta, j, m)}}^{(\theta, j, m), l, X_l^{\vartheta, k, x}}) \right) - V_{\mathcal{R}_l^{(\theta, j, m)}, j}^{(\theta, j, m)}(X_{\mathcal{R}_l^{(\theta, j, m)}}^{(\theta, j, m), l, X_l^{\vartheta, k, x}}) \right) \right. \\ & \left. - \left( f_{\mathcal{R}_l^{(\theta, j, m)}} \left( X_{\mathcal{R}_l^{(\theta, j, m)}}^{(\theta, j, m), l, X_l^{\vartheta, k, x}}, V_{\mathcal{R}_l^{(\theta, j, m)}, j-1}^{(\theta, j, -m)}(X_{\mathcal{R}_l^{(\theta, j, m)}}^{(\theta, j, m), l, X_l^{\vartheta, k, x}}) \right) - V_{\mathcal{R}_l^{(\theta, j, m)}, j-1}^{(\theta, j, -m)}(X_{\mathcal{R}_l^{(\theta, j, m)}}^{(\theta, j, m), l, X_l^{\vartheta, k, x}}) \right) \right]. \end{aligned} \quad (43)$$

The assumption that for all  $k \in \{0, 1, \dots, K-1\}$ ,  $x \in \mathcal{O}$ ,  $a, b \in \mathbb{R}$  it holds that  $|(f_k(x, a) - a) - (f_k(x, b) - b)| \leq L_k |a - b|$  hence implies that for all  $k, l, n \in \mathbb{N}$ ,  $\theta, \vartheta \in \Theta$ ,  $x \in \mathcal{O}$  with  $l \leq k \leq K$  it holds that

$$\begin{aligned} & |V_{l,n}^\theta(X_l^{\vartheta, k, x})| \leq \frac{1}{M^n} \sum_{m=1}^{M^n} \left[ \left| g(X_0^{(\theta, 0, -m), l, X_l^{\vartheta, k, x}}) \right| + \sum_{s=0}^{l-1} \left| f_s(X_s^{(\theta, 0, m), l, X_l^{\vartheta, k, x}}, 0) \right| \right] \\ & + \sum_{j=1}^{n-1} \frac{1}{M^{n-j}} \sum_{m=1}^{M^{n-j}} \frac{L_{\mathcal{R}_l^{(\theta, j, m)}}}{\mathfrak{p}_{l, \mathcal{R}_l^{(\theta, j, m)}}} \left| V_{\mathcal{R}_l^{(\theta, j, m)}, j}^{(\theta, j, m)}(X_{\mathcal{R}_l^{(\theta, j, m)}}^{(\theta, j, m), l, X_l^{\vartheta, k, x}}) - V_{\mathcal{R}_l^{(\theta, j, m)}, j-1}^{(\theta, j, -m)}(X_{\mathcal{R}_l^{(\theta, j, m)}}^{(\theta, j, m), l, X_l^{\vartheta, k, x}}) \right|. \end{aligned} \quad (44)$$

This shows that for all  $k, l, n \in \mathbb{N}$ ,  $\theta, \vartheta \in \Theta$ ,  $x \in \mathbb{R}^d$  with  $l \leq k \leq K$  it holds that

$$\begin{aligned} & |V_{l,n}^\theta(X_l^{\vartheta,k,x})|^2 \\ & \leq 3 \left[ \frac{1}{M^n} \sum_{m=1}^{M^n} \left| g(X_0^{(\theta,0,-m),l,X_l^{\vartheta,k,x}}) \right| \right]^2 + 3 \left[ \frac{1}{M^n} \sum_{m=1}^{M^n} \sum_{s=0}^{l-1} \left| f_s(X_s^{(\theta,0,m),l,X_l^{\vartheta,k,x}}, 0) \right| \right]^2 \\ & + 3 \left[ \sum_{j=1}^{n-1} \frac{1}{M^{n-j}} \sum_{m=1}^{M^{n-j}} \frac{L_{\mathcal{R}_l^{(\theta,j,m)}}}{\mathfrak{p}_{l,\mathcal{R}_l^{(\theta,j,m)}}} \left| V_{\mathcal{R}_l^{(\theta,j,m)},j}^{(\theta,j,m)}(X_{\mathcal{R}_l^{(\theta,j,m)}}^{(\theta,j,m),l,X_l^{\vartheta,k,x}}) - V_{\mathcal{R}_l^{(\theta,j,m)},j-1}^{(\theta,j,-m)}(X_{\mathcal{R}_l^{(\theta,j,m)}}^{(\theta,j,m),l,X_l^{\vartheta,k,x}}) \right| \right]^2. \end{aligned} \quad (45)$$

Jensen's inequality therefore ensures that for all  $k, l, n \in \mathbb{N}$ ,  $\theta, \vartheta \in \Theta$ ,  $x \in \mathcal{O}$  with  $l \leq k \leq K$  it holds that

$$\begin{aligned} & \mathbb{E} \left[ |V_{l,n}^\theta(X_l^{\vartheta,k,x})|^2 \right] \\ & \leq \frac{3}{M^n} \sum_{m=1}^{M^n} \mathbb{E} \left[ \left| g(X_0^{(\theta,0,-m),l,X_l^{\vartheta,k,x}}) \right|^2 \right] + \frac{3l}{M^n} \sum_{m=1}^{M^n} \sum_{s=0}^{l-1} \mathbb{E} \left[ \left| f_s(X_s^{(\theta,0,m),l,X_l^{\vartheta,k,x}}, 0) \right|^2 \right] \\ & + \sum_{j=1}^{n-1} \frac{3n}{M^{n-j}} \sum_{m=1}^{M^{n-j}} \mathbb{E} \left[ \frac{|L_{\mathcal{R}_l^{(\theta,j,m)}}|^2}{|\mathfrak{p}_{l,\mathcal{R}_l^{(\theta,j,m)}}|^2} \left| V_{\mathcal{R}_l^{(\theta,j,m)},j}^{(\theta,j,m)}(X_{\mathcal{R}_l^{(\theta,j,m)}}^{(\theta,j,m),l,X_l^{\vartheta,k,x}}) - V_{\mathcal{R}_l^{(\theta,j,m)},j-1}^{(\theta,j,-m)}(X_{\mathcal{R}_l^{(\theta,j,m)}}^{(\theta,j,m),l,X_l^{\vartheta,k,x}}) \right|^2 \right]. \end{aligned} \quad (46)$$

Item (ii) of Lemma 2.8 hence assures that for all  $k, l, n \in \mathbb{N}$ ,  $\theta, \vartheta \in \Theta$ ,  $x \in \mathcal{O}$  with  $l \leq k \leq K$  it holds that

$$\begin{aligned} & \mathbb{E} \left[ |V_{l,n}^\theta(X_l^{\vartheta,k,x})|^2 \right] \leq 3\mathbb{E} \left[ |g(X_0^{0,k,x})|^2 \right] + 3l \sum_{s=0}^{l-1} \mathbb{E} \left[ |f_s(X_s^{0,k,x}, 0)|^2 \right] \\ & + \sum_{j=1}^{n-1} \sum_{s=0}^{l-1} \frac{3n}{M^{n-j}} \sum_{m=1}^{M^{n-j}} \mathbb{E} \left[ \frac{|L_s|^2}{|\mathfrak{p}_{l,s}|^2} \left| V_{s,j}^{(\theta,j,m)}(X_s^{(\theta,j,m),l,X_l^{\vartheta,k,x}}) - V_{s,j-1}^{(\theta,j,-m)}(X_s^{(\theta,j,m),l,X_l^{\vartheta,k,x}}) \right|^2 \right] \\ & \leq 3\mathbb{E} \left[ |g(X_0^{0,k,x})|^2 \right] + 3l \sum_{s=0}^{l-1} \mathbb{E} \left[ |f_s(X_s^{0,k,x}, 0)|^2 \right] \\ & + \sum_{j=1}^{n-1} \sum_{s=0}^{l-1} \frac{6n|L_s|^2}{M^{n-j}|\mathfrak{p}_{l,s}|^2} \sum_{m=1}^{M^{n-j}} \mathbb{E} \left[ \left| V_{s,j}^{(\theta,j,m)}(X_s^{(\theta,j,m),l,X_l^{\vartheta,k,x}}) \right|^2 + \left| V_{s,j-1}^{(\theta,j,-m)}(X_s^{(\theta,j,m),l,X_l^{\vartheta,k,x}}) \right|^2 \right]. \end{aligned} \quad (47)$$

Next note that item (i) of Lemma 2.2, item (i) of Lemma 2.3, item (i) of Lemma 2.8, item (i) of Proposition 2.9, and item (ii) of Lemma 2.5 imply that for all  $k, l, m, n \in \mathbb{N}_0$ ,  $\theta, \vartheta, \xi \in \Theta$ ,  $x \in \mathcal{O}$  with  $m \leq l \leq k \leq K$  and  $\xi, \vartheta \notin \{(\theta, \eta) \in \Theta : \eta \in \Theta\}$  it holds that

$$\mathcal{O} \times \Omega \ni (x, \omega) \mapsto V_{m,n}^\theta(X_m^{\xi,l,X_l^{\vartheta,k,x}(\omega)}(\omega), \omega) \in \mathbb{R} \quad (48)$$

and

$$\mathcal{O} \times \Omega \ni (x, \omega) \mapsto V_{m,n}^0(X_m^{0,k,x}(\omega), \omega) \in \mathbb{R} \quad (49)$$

are identically distributed random fields. This and (47) imply that for all  $k, l, n \in \mathbb{N}$ ,  $\theta, \vartheta \in \Theta$ ,  $x \in \mathcal{O}$  with  $l \leq k \leq K$  and  $\vartheta \notin \{(\theta, \eta) \in \Theta : \eta \in \Theta\}$  it holds that

$$\begin{aligned} & \mathbb{E} \left[ |V_{l,n}^\theta(X_l^{\vartheta,k,x})|^2 \right] \leq 3\mathbb{E} \left[ |g(X_0^{0,k,x})|^2 \right] + 3l \sum_{s=0}^{l-1} \mathbb{E} \left[ |f_s(X_s^{0,k,x}, 0)|^2 \right] \\ & + \sum_{j=1}^{n-1} \sum_{s=0}^{l-1} \frac{12n|L_s|^2}{M^{n-j}|\mathfrak{p}_{l,s}|^2} \sum_{m=1}^{M^{n-j}} \mathbb{E} \left[ |V_{s,j}^0(X_s^{0,k,x})|^2 \right]. \end{aligned} \quad (50)$$

The fact that for all  $k \in \{0, 1, \dots, K\}$ ,  $x \in \mathcal{O}$  it holds that  $\mathbb{E}[|g(X_0^{0,k,x})|^2] + \sum_{s=0}^{k-1} \mathbb{E}[|f_s(X_s^{0,k,x}, 0)|^2] < \infty$  therefore demonstrates that for all  $n \in \mathbb{A}$  it holds that  $n + 1 \in \mathbb{A}$ . The fact that  $1 \in \mathbb{A}$  and induction hence imply that  $\mathbb{A} = \mathbb{N}$ . Therefore, we obtain that for all  $k, l, n \in \mathbb{N}_0$ ,  $\theta, \vartheta \in \Theta$ ,  $x \in \mathcal{O}$  with  $l \leq k \leq K$  and  $\vartheta \notin \{(\theta, \eta) \in \Theta : \eta \in \Theta\}$  it holds that  $\mathbb{E}[|V_{l,n}^\theta(X_l^{\vartheta,k,x})|^2] < \infty$ . This completes the proof of Lemma 2.11.  $\square$

### 3 Error analysis for MLP approximations for iterated nested expectations

In this section we provide in Proposition 3.13 in Subsection 3.4 below a full error analysis for MLP approximations for iterated nested expectations. Our proof of Proposition 3.13 is inspired by Hutzenthaler et al. [36] and is based on the idea to combine the recursive error estimate in Lemma 3.10 in Subsection 3.3, the elementary integration by parts type result in Lemma 3.11 in Subsection 3.4, the elementary Gronwall type inequality in Lemma 3.12 in Subsection 3.4, and the elementary a priori bounds for iterated nested expectations in Lemma 3.6 in Subsection 3.1. Our proof of the recursive error estimate in Lemma 3.10 in Subsection 3.3 is, roughly speaking, based on bias-variance decompositions for the approximation errors of the proposed MLP approximation schemes, the bias estimates for MLP approximations in Lemma 3.8 in Subsection 3.2, and the variance estimates for MLP approximations in Lemma 3.9 in Subsection 3.2.

#### 3.1 A priori estimates for nested expectations

**Lemma 3.1.** *Let  $d, K \in \mathbb{N}$ ,  $L \in \mathbb{R}$ ,  $\mathcal{O} \in \mathcal{B}(\mathbb{R}^d) \setminus \{\emptyset\}$ , let  $f: \mathcal{O} \times \mathbb{R} \rightarrow \mathbb{R}$  be  $\mathcal{B}(\mathcal{O} \times \mathbb{R})/\mathcal{B}(\mathbb{R})$ -measurable, assume for all  $x \in \mathcal{O}$ ,  $a, b \in \mathbb{R}$  that  $|f(x, a) - f(x, b)| \leq L|a - b|$ , let  $(S, \mathcal{S})$  be a measurable space, let  $\phi_k: \mathcal{O} \times S \rightarrow \mathcal{O}$ ,  $k \in \{0, 1, \dots, K\}$ , be  $(\mathcal{B}(\mathcal{O}) \otimes \mathcal{S})/\mathcal{B}(\mathcal{O})$ -measurable, let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a probability space, let  $W_k: \Omega \rightarrow S$ ,  $k \in \{0, 1, \dots, K\}$ , be independent random variables, for every  $k \in \{0, 1, \dots, K\}$  let  $X^k = (X_l^{k,x})_{(l,x) \in \{0,1,\dots,k\} \times \mathcal{O} : \{0,1,\dots,k\} \times \mathcal{O} \times \Omega \rightarrow \mathcal{O}}$  satisfy for all  $l \in \{0, 1, \dots, k\}$ ,  $x \in \mathcal{O}$  that*

$$X_l^{k,x} = \begin{cases} x & : l = k \\ \phi_l(X_{l+1}^{k,x}, W_l) & : l < k, \end{cases} \quad (51)$$

let  $v: \mathcal{O} \rightarrow \mathbb{R}$  be  $\mathcal{B}(\mathcal{O})/\mathcal{B}(\mathbb{R})$ -measurable, and assume for all  $x \in \mathcal{O}$  that  $\sum_{k=0}^K \mathbb{E}[|f(X_0^{k,x}, 0)|^2 + |v(X_0^{k,x})|^2] < \infty$ . Then there exists a  $\mathcal{B}(\mathcal{O})/\mathcal{B}(\mathbb{R})$ -measurable  $u: \mathcal{O} \rightarrow \mathbb{R}$  such that for all  $x \in \mathcal{O}$  it holds that  $\sum_{k=1}^K \mathbb{E}[|u(X_1^{k,x})|^2 + |f(X_0^{1,x}, v(X_0^{1,x}))|^2] < \infty$  and  $u(x) = \mathbb{E}[f(X_0^{1,x}, v(X_0^{1,x}))]$ .

*Proof of Lemma 3.1.* First, observe that the assumption that for all  $x \in \mathcal{O}$ ,  $a, b \in \mathbb{R}$  it holds that  $|f(x, a) - f(x, b)| \leq L|a - b|$  and the assumption that for all  $k \in \{0, 1, \dots, K\}$ ,  $x \in \mathcal{O}$  it holds that  $\mathbb{E}[|f(X_0^{k,x}, 0)|^2 + |v(X_0^{k,x})|^2] < \infty$  ensure that for all  $k \in \{0, 1, \dots, K\}$ ,  $x \in \mathcal{O}$  it holds that

$$\begin{aligned} \mathbb{E}[|f(X_0^{k,x}, v(X_0^{k,x}))|^2] &\leq \mathbb{E}[(|f(X_0^{k,x}, 0)| + L|v(X_0^{k,x})|)^2] \\ &\leq 2\mathbb{E}[|f(X_0^{k,x}, 0)|^2] + 2L^2\mathbb{E}[|v(X_0^{k,x})|^2] < \infty. \end{aligned} \quad (52)$$

Jensen's inequality hence proves that there exists  $u: \mathcal{O} \rightarrow \mathbb{R}$  which satisfies that for all  $x \in \mathcal{O}$  it holds that  $\mathbb{E}[|f(X_0^{1,x}, v(X_0^{1,x}))|] \leq (\mathbb{E}[|f(X_0^{1,x}, v(X_0^{1,x}))|^2])^{1/2} < \infty$  and

$$u(x) = \mathbb{E}[f(X_0^{1,x}, v(X_0^{1,x}))]. \quad (53)$$

Next note that Klenke [39, Theorem 14.16] and (53) establish that  $u$  is  $\mathcal{B}(\mathcal{O})/\mathcal{B}(\mathbb{R})$ -measurable. Moreover, observe that item (i) of Lemma 2.2 yields that for all  $k \in \{1, 2, \dots, K\}$ ,  $x \in \mathcal{O}$  it

holds that  $X_1^{k,x}: \Omega \rightarrow \mathcal{O}$  is  $\sigma((W_r)_{r \in [1,k-1] \cap \mathbb{N}_0})/\mathcal{B}(\mathcal{O})$ -measurable. Combining (51), (53), Jensen's inequality, the assumption that  $W_k$ ,  $k \in \{0, 1, \dots, K-1\}$ , are independent random variables, item (i) of Lemma 2.2, and Hutzenthaler et al. [36, Lemma 2.2] hence guarantees that for all  $k \in \{1, 2, \dots, K\}$ ,  $x \in \mathcal{O}$  it holds that

$$\begin{aligned} \mathbb{E}[|u(X_1^{k,x})|^2] &= \int_{\mathcal{O}} |u(y)|^2 (X_1^{k,x}(\mathbb{P}))(dy) \leq \int_{\mathcal{O}} \mathbb{E}[|f(X_0^{1,y}, v(X_0^{1,y}))|^2] (X_1^{k,x}(\mathbb{P}))(dy) \\ &= \mathbb{E}\left[\left|f(X_0^{1,X_1^{k,x}}, v(X_0^{1,X_1^{k,x}}))\right|^2\right] = \mathbb{E}[|f(X_0^{k,x}, v(X_0^{k,x}))|^2] < \infty. \end{aligned} \quad (54)$$

Combining this, (52), (53), and the fact that  $u$  is  $\mathcal{B}(\mathcal{O})/\mathcal{B}(\mathbb{R})$ -measurable completes the proof of Lemma 3.1.  $\square$

**Lemma 3.2.** *Let  $d, K \in \mathbb{N}$ ,  $L \in \mathbb{R}$ ,  $\mathcal{O} \in \mathcal{B}(\mathbb{R}^d) \setminus \{\emptyset\}$ , let  $f_k: \mathcal{O} \times \mathbb{R} \rightarrow \mathbb{R}$ ,  $k \in \{0, 1, \dots, K\}$ , be  $\mathcal{B}(\mathcal{O} \times \mathbb{R})/\mathcal{B}(\mathbb{R})$ -measurable, let  $g: \mathcal{O} \rightarrow \mathbb{R}$  be  $\mathcal{B}(\mathcal{O})/\mathcal{B}(\mathbb{R})$ -measurable, assume for all  $k \in \{0, 1, \dots, K\}$ ,  $x \in \mathcal{O}$ ,  $a, b \in \mathbb{R}$  that  $|f_k(x, a) - f_k(x, b)| \leq L|a - b|$ , let  $(S, \mathcal{S})$  be a measurable space, let  $\phi_k: \mathcal{O} \times S \rightarrow \mathcal{O}$ ,  $k \in \{0, 1, \dots, K\}$ , be  $(\mathcal{B}(\mathcal{O}) \otimes \mathcal{S})/\mathcal{B}(\mathcal{O})$ -measurable, let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a probability space, let  $W_k: \Omega \rightarrow S$ ,  $k \in \{0, 1, \dots, K\}$ , be independent random variables, for every  $k \in \{0, 1, \dots, K\}$  let  $X^k = (X_l^{k,x})_{(l,x) \in \{0,1,\dots,k\} \times \mathcal{O}}: \{0, 1, \dots, k\} \times \mathcal{O} \times \Omega \rightarrow \mathcal{O}$  satisfy for all  $l \in \{0, 1, \dots, k\}$ ,  $x \in \mathcal{O}$  that*

$$X_l^{k,x} = \begin{cases} x & : l = k \\ \phi_l(X_{l+1}^{k,x}, W_l) & : l < k, \end{cases} \quad (55)$$

and assume for all  $k \in \{0, 1, \dots, K\}$ ,  $x \in \mathcal{O}$  that  $\mathbb{E}[|g(X_0^{k,x})|^2 + \sum_{l=0}^{k-1} |f_l(X_l^{k,x}, 0)|^2] < \infty$ . Then there exist  $\mathcal{B}(\mathcal{O})/\mathcal{B}(\mathbb{R})$ -measurable  $v_k: \mathcal{O} \rightarrow \mathbb{R}$ ,  $k \in \{0, 1, \dots, K\}$ , such that for all  $k \in \{1, 2, \dots, K\}$ ,  $x \in \mathcal{O}$  it holds that  $\mathbb{E}[|f_{k-1}(X_{k-1}^{k,x}, v_{k-1}(X_{k-1}^{k,x}))| + \sum_{l=1}^k |v_l(X_l^{k,x})|^2] < \infty$ ,  $v_0(x) = g(x)$ , and  $v_k(x) = \mathbb{E}[f_{k-1}(X_{k-1}^{k,x}, v_{k-1}(X_{k-1}^{k,x}))]$ .

*Proof of Lemma 3.2.* Throughout this proof let  $\mathbb{A} \subseteq \{0, 1, \dots, K\}$  satisfy that

$$\mathbb{A} = \left\{ k \in \{0, 1, \dots, K\} : \begin{pmatrix} \text{There exist } \mathcal{B}(\mathcal{O})/\mathcal{B}(\mathbb{R})\text{-measurable} \\ v_l: \mathcal{O} \rightarrow \mathbb{R}, l \in \{0, 1, \dots, k\}, \text{ such that} \\ \text{for all } l \in \{1, 2, \dots, k\}, x \in \mathcal{O} \text{ it holds that} \\ \mathbb{E}[|f_{l-1}(X_{l-1}^{l,x}, v_{l-1}(X_{l-1}^{l,x}))| + \sum_{m=l}^K |v_m(X_m^{l,x})|^2] < \infty, \\ v_0(x) = g(x), \text{ and } v_l(x) = \mathbb{E}[f_{l-1}(X_{l-1}^{l,x}, v_{l-1}(X_{l-1}^{l,x}))]. \end{pmatrix} \right\}. \quad (56)$$

Note that the assumption that  $g$  is  $\mathcal{B}(\mathcal{O})/\mathcal{B}(\mathbb{R})$ -measurable ensures that  $0 \in \mathbb{A}$ . Moreover, observe that Lemma 3.1 (applied with  $d \leftarrow d$ ,  $K \leftarrow K - k$ ,  $L \leftarrow L$ ,  $\mathcal{O} \leftarrow \mathcal{O}$ ,  $f \leftarrow f_k$ ,  $(S, \mathcal{S}) \leftarrow (S, \mathcal{S})$ ,  $(\phi_l)_{l \in \{0,1,\dots,K-1\}} \leftarrow (\phi_{k+s})_{s \in \{0,1,\dots,K-k-1\}}$ ,  $(\Omega, \mathcal{F}, \mathbb{P}) \leftarrow (\Omega, \mathcal{F}, \mathbb{P})$ ,  $(W_l)_{l \in \{0,1,\dots,K-1\}} \leftarrow (W_{k+s})_{s \in \{0,1,\dots,K-k-1\}}$ ,  $v \leftarrow v_k$  for  $k \in \{0, 1, \dots, K-1\} \cap \mathbb{A}$  in the notation of Lemma 3.1) and the assumption that for all  $k \in \{0, 1, \dots, K\}$ ,  $x \in \mathcal{O}$  it holds that  $\mathbb{E}[|g(X_0^{k,x})|^2] < \infty$  ensure that for all  $k \in \{0, 1, \dots, K-1\} \cap \mathbb{A}$  there exists  $v_{k+1}: \mathcal{O} \rightarrow \mathbb{R}$  which satisfies that

- (I) it holds that  $v_{k+1}$  is  $\mathcal{B}(\mathcal{O})/\mathcal{B}(\mathbb{R})$ -measurable,
  - (II) it holds for all  $m \in \{k+1, \dots, K\}$ ,  $x \in \mathcal{O}$  that  $\mathbb{E}[|v_{k+1}(X_{k+1}^{m,x})|^2] < \infty$ , and
  - (III) it holds for all  $x \in \mathcal{O}$  that  $\mathbb{E}[|f_k(X_k^{k+1,x}, v_k(X_k^{k+1,x}))|] < \infty$  and
- $$v_{k+1}(x) = \mathbb{E}[f_k(X_k^{k+1,x}, v_k(X_k^{k+1,x}))]. \quad (57)$$

This demonstrates for all  $k \in \{0, 1, \dots, K-1\} \cap \mathbb{A}$  that  $k+1 \in \mathbb{A}$ . Induction and the fact that  $0 \in \mathbb{A}$  therefore establish that  $\mathbb{A} = \{0, 1, \dots, K\}$ . This completes the proof of Lemma 3.2.  $\square$

**Corollary 3.3.** Let  $d, K \in \mathbb{N}$ ,  $L, T \in \mathbb{R}$ , let  $t_k \in \mathbb{R}$ ,  $k \in \{0, 1, \dots, K\}$ , satisfy  $0 = t_0 < t_1 < \dots < t_K = T$ , let  $f_k: \mathbb{R}^d \times \mathbb{R} \rightarrow \mathbb{R}$ ,  $k \in \{0, 1, \dots, K\}$ , be  $\mathcal{B}(\mathbb{R}^d \times \mathbb{R})/\mathcal{B}(\mathbb{R})$ -measurable, let  $g: \mathbb{R}^d \rightarrow \mathbb{R}$  be  $\mathcal{B}(\mathbb{R}^d)/\mathcal{B}(\mathbb{R})$ -measurable, assume for all  $k \in \{0, 1, \dots, K\}$ ,  $x \in \mathbb{R}^d$ ,  $a, b \in \mathbb{R}$  that  $|f_k(x, a) - f_k(x, b)| \leq L|a - b|$ , let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a probability space, let  $W: [0, T] \times \Omega \rightarrow \mathbb{R}^d$  be a standard Brownian motion, and assume for all  $x \in \mathcal{O}$  that  $\mathbb{E}[|g(x + W_T)|^2 + \sum_{l=0}^{K-1} |f_l(x + W_{T-t_l}, 0)|^2] < \infty$ . Then there exist  $\mathcal{B}(\mathbb{R}^d)/\mathcal{B}(\mathbb{R})$ -measurable  $v_k: \mathbb{R}^d \rightarrow \mathbb{R}$ ,  $k \in \{0, 1, \dots, K\}$ , such that for all  $k \in \{1, 2, \dots, K\}$ ,  $x \in \mathcal{O}$  it holds that  $\mathbb{E}[|f_{k-1}(x + W_{t_k-t_{k-1}}, v_{k-1}(x + W_{t_k-t_{k-1}}))| + \sum_{l=1}^k |v_l(x + W_{t_k} - W_{t_l})|^2] < \infty$ ,  $v_0(x) = g(x)$ , and  $v_k(x) = \mathbb{E}[f_{k-1}(x + W_{t_k-t_{k-1}}, v_{k-1}(x + W_{t_k-t_{k-1}}))]$ .

*Proof of Corollary 3.3.* First, observe that the assumption that  $W$  is a Brownian motion and the assumption that for all  $x \in \mathbb{R}^d$  it holds that  $\mathbb{E}[|g(x + W_T)|^2] < \infty$  show that for all  $k \in \{1, 2, \dots, K\}$  it holds that

$$\begin{aligned} \mathbb{E}[|g(x + W_{t_k})|^2] &= [2\pi t_k]^{-d/2} \int_{\mathbb{R}^d} |g(x + z)|^2 \exp\left(-\frac{\|z\|^2}{2t_k}\right) dz \\ &\leq [2\pi t_k]^{-d/2} \int_{\mathbb{R}^d} |g(x + z)|^2 \exp\left(-\frac{\|z\|^2}{2T}\right) dz = \left[\frac{T}{t_k}\right]^{d/2} \mathbb{E}[|g(x + W_T)|^2] < \infty. \end{aligned} \quad (58)$$

Furthermore, note that the assumption that  $W$  is a standard Brownian motion and the assumption that  $\mathbb{E}[\sum_{k=0}^{K-1} |f_l(x + W_{T-t_l}, 0)|^2] < \infty$  show that for all  $k, l \in \{0, 1, \dots, K\}$  with  $k \geq l$  it holds that  $\mathbb{E}[|f_l(x + W_{t_k-t_l}, 0)|^2] < \infty$ . This, (58), and the assumption that  $W$  is a standard Brownian motion prove that for all  $k \in \{1, 2, \dots, K\}$ ,  $x \in \mathbb{R}^d$  it holds that  $\mathbb{E}[|g(x + W_{t_k})|^2 + \sum_{l=0}^{k-1} |f_l(x + W_{t_k} - W_{t_l}, 0)|^2] < \infty$ . Lemma 3.2 (applied with  $d \leftarrow d$ ,  $K \leftarrow K$ ,  $L \leftarrow L$ ,  $\mathcal{O} \leftarrow \mathbb{R}^d$ ,  $(f_k)_{k \in \{0, 1, \dots, K-1\}} \leftarrow (f_k)_{k \in \{0, 1, \dots, K-1\}}$ ,  $g \leftarrow g$ ,  $(S, \mathcal{S}) \leftarrow (\mathbb{R}^d, \mathcal{B}(\mathbb{R}^d))$ ,  $(\phi_k)_{k \in \{0, 1, \dots, K-1\}} \leftarrow ((\mathbb{R}^d \times \mathbb{R}^d \ni (x, w) \mapsto x + w \in \mathbb{R}^d))_{k \in \{0, 1, \dots, K-1\}}$ ,  $(\Omega, \mathcal{F}, \mathbb{P}) \leftarrow (\Omega, \mathcal{F}, \mathbb{P})$ ,  $(W_k)_{k \in \{0, 1, \dots, K-1\}} \leftarrow (W_{t_{k+1}} - W_{t_k})_{k \in \{0, 1, \dots, K-1\}}$  in the notation of Lemma 3.2) and the assumption that  $W$  is a standard Brownian motion therefore establish the claim. This completes the proof of Corollary 3.3.  $\square$

**Lemma 3.4.** Let  $d, K \in \mathbb{N}$ ,  $L \in \mathbb{R}$ ,  $\mathcal{O} \in \mathcal{B}(\mathbb{R}^d) \setminus \{\emptyset\}$ , let  $f_k: \mathcal{O} \times \mathbb{R} \rightarrow \mathbb{R}$ ,  $k \in \{0, 1, \dots, K\}$ , be  $\mathcal{B}(\mathcal{O} \times \mathbb{R})/\mathcal{B}(\mathbb{R})$ -measurable, let  $g: \mathcal{O} \rightarrow \mathbb{R}$  be  $\mathcal{B}(\mathcal{O})/\mathcal{B}(\mathbb{R})$ -measurable, assume for all  $k \in \{0, 1, \dots, K\}$ ,  $x \in \mathcal{O}$ ,  $a, b \in \mathbb{R}$  that  $|f_k(x, a) - f_k(x, b)| \leq L|a - b|$ , let  $(S, \mathcal{S})$  be a measurable space, let  $\phi_k: \mathcal{O} \times S \rightarrow \mathcal{O}$ ,  $k \in \{0, 1, \dots, K\}$ , be  $(\mathcal{B}(\mathcal{O}) \otimes \mathcal{S})/\mathcal{B}(\mathcal{O})$ -measurable, let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a probability space, let  $W_k: \Omega \rightarrow S$ ,  $k \in \{0, 1, \dots, K\}$ , be independent random variables, for every  $k \in \{0, 1, \dots, K\}$  let  $X^k = (X_l^{k,x})_{(l,x) \in \{0, 1, \dots, k\} \times \mathcal{O}}: \{0, 1, \dots, k\} \times \mathcal{O} \times \Omega \rightarrow \mathcal{O}$  satisfy for all  $l \in \{0, 1, \dots, k\}$ ,  $x \in \mathcal{O}$  that

$$X_l^{k,x} = \begin{cases} x & : l = k \\ \phi_l(X_{l+1}^{k,x}, W_l) & : l < k, \end{cases} \quad (59)$$

assume for all  $k \in \{0, 1, \dots, K\}$ ,  $x \in \mathcal{O}$  that  $\mathbb{E}[|g(X_0^{k,x})|^2 + \sum_{l=0}^{k-1} |f_l(X_l^{k,x}, 0)|^2] < \infty$ , and let  $v_k: \mathcal{O} \rightarrow \mathbb{R}$ ,  $k \in \{0, 1, \dots, K\}$ , be  $\mathcal{B}(\mathcal{O})/\mathcal{B}(\mathbb{R})$ -measurable functions which satisfy for all  $k \in \{1, 2, \dots, K\}$ ,  $x \in \mathcal{O}$  that  $v_0(x) = g(x)$  and

$$v_k(x) = \mathbb{E}[f_{k-1}(X_{k-1}^{k,x}, v_{k-1}(X_{k-1}^{k,x}))] \quad (60)$$

(cf. Lemma 3.2). Then it holds for all  $k, l \in \{0, 1, \dots, K\}$ ,  $x \in \mathcal{O}$  with  $k \geq l$  that  $\mathbb{E}[|v_l(X_l^{k,x})|] + \sum_{s=l}^{k-1} \mathbb{E}[|f_s(X_s^{k,x}, v_s(X_s^{k,x})) - v_s(X_s^{k,x})|] < \infty$  and

$$v_k(x) = \mathbb{E}[v_l(X_l^{k,x})] + \sum_{s=l}^{k-1} \mathbb{E}[f_s(X_s^{k,x}, v_s(X_s^{k,x})) - v_s(X_s^{k,x})]. \quad (61)$$

*Proof of Lemma 3.4.* Throughout this proof let  $k \in \{1, 2, \dots, K\}$ . Observe that Jensen's inequality and Lemma 3.2 prove that for all  $l \in \{0, 1, \dots, k\}$ ,  $x \in \mathcal{O}$  it holds that

$$\mathbb{E}[|v_l(X_l^{k,x})|] \leq \left(\mathbb{E}[|v_l(X_l^{k,x})|^2]\right)^{1/2} < \infty. \quad (62)$$

Combining the assumption that for all  $l \in \{0, \dots, k-1\}$ ,  $x \in \mathcal{O}$ ,  $a, b \in \mathbb{R}$  it holds that  $|f_l(x, a) - f_l(x, b)| \leq L|a - b|$  and  $\mathbb{E}[|f_l(X_l^{k,x}, 0)|^2] < \infty$  with Jensen's inequality and Minkowski's inequality hence implies that for all  $l \in \{1, 2, \dots, k\}$ ,  $x \in \mathcal{O}$  it holds that

$$\begin{aligned} \mathbb{E}[|f_{l-1}(X_{l-1}^{k,x}, v_{l-1}(X_{l-1}^{k,x}))|] &\leq \left(\mathbb{E}[|f_{l-1}(X_{l-1}^{k,x}, v_{l-1}(X_{l-1}^{k,x}))|^2]\right)^{1/2} \\ &\leq \left(\mathbb{E}[|f_{l-1}(X_{l-1}^{k,x}, 0)|^2]\right)^{1/2} + L \left(\mathbb{E}[|v_{l-1}(X_{l-1}^{k,x})|^2]\right)^{1/2} < \infty. \end{aligned} \quad (63)$$

This and (62) establish that for all  $l \in \{0, 1, \dots, k\}$ ,  $x \in \mathcal{O}$  it holds that

$$\mathbb{E}\left[|v_l(X_l^{k,x})| + \sum_{s=l}^{k-1} |f_s(X_s^{k,x}, v_s(X_s^{k,x})) - v_s(X_s^{k,x})|\right] < \infty. \quad (64)$$

In the next step we prove that for all  $l \in \{0, 1, \dots, k\}$ ,  $x \in \mathcal{O}$  it holds that

$$v_k(x) = \mathbb{E}[v_l(X_l^{k,x})] + \sum_{s=l}^{k-1} \mathbb{E}[f_s(X_s^{k,x}, v_s(X_s^{k,x})) - v_s(X_s^{k,x})]. \quad (65)$$

For this we observe that item (i) of Lemma 2.2 ensures that

- (I) for all  $l \in \{1, 2, \dots, k\}$  it holds that  $\mathcal{O} \times \Omega \ni (x, \omega) \mapsto X_{l-1}^{l,x}(\omega) \in \mathcal{O}$  is  $(\mathcal{B}(\mathcal{O}) \otimes \sigma(W_{l-1}) / \mathcal{B}(\mathcal{O}))$ -measurable and
- (II) for all  $l \in \{0, 1, \dots, k\}$ ,  $x \in \mathcal{O}$  it holds that  $\Omega \ni \omega \mapsto X_l^{k,x}(\omega) \in \mathcal{O}$  is  $\sigma((W_m)_{m \in [l, k] \cap \mathbb{N}_0}) / \mathcal{B}(\mathcal{O})$ -measurable.

This, the assumption that  $W_m: \Omega \rightarrow S$ ,  $m \in \{0, 1, \dots, K-1\}$ , are independent random variables, and Hutzenthaler et al. [36, Lemma 2.2] ensure that for all  $l \in \{1, 2, \dots, k\}$ ,  $x \in \mathcal{O}$  it holds that

$$\int_{\mathcal{O}} \mathbb{E}[|f_{l-1}(X_{l-1}^{l,y}, v_{l-1}(X_{l-1}^{l,y}))|^2] (X_l^{k,x}(\mathbb{P}))(dy) = \mathbb{E}\left[|f_{l-1}(X_{l-1}^{l, X_l^{k,x}}, v_{l-1}(X_{l-1}^{l, X_l^{k,x}}))|^2\right]. \quad (66)$$

Combining this with (59) ensures that for all  $l \in \{1, 2, \dots, k\}$ ,  $x \in \mathcal{O}$  it holds that

$$\int_{\mathcal{O}} \mathbb{E}[|f_{l-1}(X_{l-1}^{l,y}, v_{l-1}(X_{l-1}^{l,y}))|^2] (X_l^{k,x}(\mathbb{P}))(dy) = \mathbb{E}[|f_{l-1}(X_{l-1}^{k,x}, v_{l-1}(X_{l-1}^{k,x}))|^2]. \quad (67)$$

Jensen's inequality and (63) hence guarantee that for all  $l \in \{1, 2, \dots, k\}$ ,  $x \in \mathcal{O}$  it holds that

$$\begin{aligned} &\int_{\mathcal{O}} \mathbb{E}[|f_{l-1}(X_{l-1}^{l,y}, v_{l-1}(X_{l-1}^{l,y}))|] (X_l^{k,x}(\mathbb{P}))(dy) \\ &\leq \left[\int_{\mathcal{O}} \left(\mathbb{E}[|f_{l-1}(X_{l-1}^{l,y}, v_{l-1}(X_{l-1}^{l,y}))|]\right)^2 (X_l^{k,x}(\mathbb{P}))(dy)\right]^{1/2} \\ &\leq \left[\int_{\mathcal{O}} \mathbb{E}[|f_{l-1}(X_{l-1}^{l,y}, v_{l-1}(X_{l-1}^{l,y}))|^2] (X_l^{k,x}(\mathbb{P}))(dy)\right]^{1/2} \\ &= \left(\mathbb{E}[|f_{l-1}(X_{l-1}^{k,x}, v_{l-1}(X_{l-1}^{k,x}))|^2]\right)^{1/2} < \infty. \end{aligned} \quad (68)$$

The assumption that  $W_m: \Omega \rightarrow S$ ,  $m \in \{0, 1, \dots, K-1\}$ , are independent random variables, item (I), item (II), and Hutzenthaler et al. [36, Lemma 2.2] therefore ensure that for all  $l \in \{1, 2, \dots, k\}$ ,  $x \in \mathcal{O}$  it holds that

$$\begin{aligned}
& \mathbb{E} \left[ f_{l-1}(X_{l-1}^{l, X_l^{k,x}}, v_{l-1}(X_{l-1}^{l, X_l^{k,x}})) \right] \\
&= \mathbb{E} \left[ \max \left\{ f_{l-1}(X_{l-1}^{l, X_l^{k,x}}, v_{l-1}(X_{l-1}^{l, X_l^{k,x}})), 0 \right\} \right] \\
&\quad - \mathbb{E} \left[ \max \left\{ -f_{l-1}(X_{l-1}^{l, X_l^{k,x}}, v_{l-1}(X_{l-1}^{l, X_l^{k,x}})), 0 \right\} \right] \\
&= \int_{\mathcal{O}} \mathbb{E} \left[ \max \left\{ f_{l-1}(X_{l-1}^{l,y}, v_{l-1}(X_{l-1}^{l,y})), 0 \right\} \right] (X_l^{k,x}(\mathbb{P}))(dy) \\
&\quad - \int_{\mathcal{O}} \mathbb{E} \left[ \max \left\{ -f_{l-1}(X_{l-1}^{l,y}, v_{l-1}(X_{l-1}^{l,y})), 0 \right\} \right] (X_l^{k,x}(\mathbb{P}))(dy) \\
&= \int_{\mathcal{O}} \mathbb{E} \left[ f_{l-1}(X_{l-1}^{l,y}, v_{l-1}(X_{l-1}^{l,y})) \right] (X_l^{k,x}(\mathbb{P}))(dy).
\end{aligned} \tag{69}$$

Combining this with (59) yields for all  $l \in \{1, 2, \dots, k\}$ ,  $x \in \mathcal{O}$  that

$$\mathbb{E} \left[ f_{l-1}(X_{l-1}^{k,x}, v_{l-1}(X_{l-1}^{k,x})) \right] = \int_{\mathcal{O}} \mathbb{E} \left[ f_{l-1}(X_{l-1}^{l,y}, v_{l-1}(X_{l-1}^{l,y})) \right] (X_l^{k,x}(\mathbb{P}))(dy). \tag{70}$$

Note that (60) hence implies for all  $l \in \{1, 2, \dots, k\}$ ,  $x \in \mathcal{O}$  that

$$\mathbb{E} \left[ f_{l-1}(X_{l-1}^{k,x}, v_{l-1}(X_{l-1}^{k,x})) \right] = \int_{\mathbb{R}^d} v_l(y) (X_l^k(x)(\mathbb{P}))(dy) = \mathbb{E} [v_l(X_l^k(x))]. \tag{71}$$

This, (62), and (68) ensure that for all  $l \in \{1, 2, \dots, k\}$ ,  $x \in \mathcal{O}$  it holds that  $\mathbb{E}[|v_{l-1}(X_{l-1}^{k,x})| + |f_{l-1}(X_{l-1}^{k,x}, v_{l-1}(X_{l-1}^{k,x})) - v_{l-1}(X_{l-1}^{k,x})|] < \infty$  and

$$\mathbb{E} [v_l(X_l^{k,x})] = \mathbb{E} [v_{l-1}(X_{l-1}^{k,x})] + \mathbb{E} [f_{l-1}(X_{l-1}^{k,x}, v_{l-1}(X_{l-1}^{k,x})) - v_{l-1}(X_{l-1}^{k,x})]. \tag{72}$$

The fact that for all  $x \in \mathcal{O}$  it holds that  $v_k(x) = \mathbb{E}[v_k(X_k^{k,x})]$  and induction hence establish (61). This completes the proof of Lemma 3.4.  $\square$

**Lemma 3.5** (A Gronwall lemma). *Let  $\alpha \in [0, \infty)$ ,  $K \in \mathbb{N}$ , and let  $L_0, L_1, \dots, L_{K-1} \in [0, \infty)$ ,  $A_0, A_1, \dots, A_K \in [0, \infty)$  satisfy for all  $k \in \{0, 1, \dots, K\}$  that*

$$A_k \leq \alpha + \sum_{l=0}^{k-1} L_l A_l. \tag{73}$$

*Then it holds for all  $k \in \{0, 1, \dots, K\}$  that*

$$A_k \leq \alpha \prod_{l=0}^{k-1} (1 + L_l) \leq \alpha \exp \left( \sum_{l=0}^{k-1} L_l \right). \tag{74}$$

*Proof of Lemma 3.5.* Throughout this proof let  $\mathbb{A} \subseteq \{0, 1, \dots, K\}$  satisfy

$$\mathbb{A} = \left\{ k \in \{0, 1, \dots, K\} : \left( \begin{array}{c} \text{For all } s \in \{0, 1, \dots, k\} \text{ it holds that} \\ A_s \leq \alpha \prod_{l=0}^{s-1} (1 + L_s) \end{array} \right) \right\}. \tag{75}$$

Note that the fact that  $A_0 \leq \alpha = \alpha \prod_{l=0}^{-1} (1 + L_l)$  assures that  $0 \in \mathbb{A}$ . Next observe that (73) and the fact that for all  $l \in \{0, 1, \dots, K\}$  it holds that  $\prod_{s=0}^l (1 + L_s) - \prod_{s=0}^{l-1} (1 + L_s) = L_l \prod_{s=0}^{l-1} (1 + L_s)$  ensure that for all  $k \in \{0, 1, \dots, K-1\} \cap \mathbb{A}$  it holds that

$$\begin{aligned} A_{k+1} &\leq \alpha + \sum_{s=0}^k L_s \alpha \prod_{l=0}^{s-1} (1 + L_l) = \alpha + \alpha \sum_{s=0}^k \left[ \prod_{l=0}^s (1 + L_l) - \prod_{l=0}^{s-1} (1 + L_l) \right] \\ &= \alpha + \alpha \left[ \left( \prod_{l=0}^k (1 + L_l) \right) - 1 \right] = \alpha \prod_{l=0}^k (1 + L_l). \end{aligned} \quad (76)$$

This demonstrates that for all  $k \in \{0, 1, \dots, K-1\} \cap \mathbb{A}$  it holds that  $k+1 \in \mathbb{A}$ . Induction and the fact that  $0 \in \mathbb{A}$  hence demonstrate that for all  $k \in \{0, 1, \dots, K\}$  it holds that

$$A_k \leq \alpha \prod_{l=0}^{k-1} (1 + L_l). \quad (77)$$

Combining this with the fact that for all  $t \in \mathbb{R}$  it holds that  $1 + t \leq e^t$  establishes (74). This completes the proof of Lemma 3.5.  $\square$

**Lemma 3.6.** *Let  $d, K \in \mathbb{N}$ ,  $L_0, L_1, \dots, L_K \in \mathbb{R}$ ,  $\mathcal{O} \in \mathcal{B}(\mathbb{R}^d) \setminus \{\emptyset\}$ , let  $f_k: \mathcal{O} \times \mathbb{R} \rightarrow \mathbb{R}$ ,  $k \in \{0, 1, \dots, K\}$ , be  $\mathcal{B}(\mathcal{O} \times \mathbb{R})/\mathcal{B}(\mathbb{R})$ -measurable, let  $g: \mathcal{O} \rightarrow \mathbb{R}$  be  $\mathcal{B}(\mathcal{O})/\mathcal{B}(\mathbb{R})$ -measurable, assume for all  $k \in \{0, 1, \dots, K\}$ ,  $x \in \mathcal{O}$ ,  $a, b \in \mathbb{R}$  that  $|(f_k(x, a) - a) - (f_k(x, b) - b)| \leq L_k |a - b|$ , let  $(S, \mathcal{S})$  be a measurable space, let  $\phi_k: \mathcal{O} \times S \rightarrow \mathcal{O}$ ,  $k \in \{0, 1, \dots, K\}$ , be  $(\mathcal{B}(\mathcal{O}) \otimes \mathcal{S})/\mathcal{B}(\mathcal{O})$ -measurable, let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a probability space, let  $W_k: \Omega \rightarrow S$ ,  $k \in \{0, 1, \dots, K\}$ , be independent random variables, for every  $k \in \{0, 1, \dots, K\}$  let  $X^k = (X_l^{k,x})_{(l,x) \in \{0,1,\dots,k\} \times \mathcal{O}}: \{0, 1, \dots, k\} \times \mathcal{O} \times \Omega \rightarrow \mathcal{O}$  satisfy for all  $l \in \{0, 1, \dots, k\}$ ,  $x \in \mathcal{O}$  that*

$$X_l^{k,x} = \begin{cases} x & : l = k \\ \phi_l(X_{l+1}^{k,x}, W_l) & : l < k, \end{cases} \quad (78)$$

assume for all  $k \in \{0, 1, \dots, K\}$ ,  $x \in \mathcal{O}$  that  $\mathbb{E}[|g(X_0^{k,x})|^2 + \sum_{l=0}^{k-1} |f_l(X_l^{k,x}, 0)|^2] < \infty$ , and let  $v_0, v_1, \dots, v_K: \mathcal{O} \rightarrow \mathbb{R}$  be  $\mathcal{B}(\mathcal{O})/\mathcal{B}(\mathbb{R})$ -measurable functions which satisfy for all  $k \in \{1, 2, \dots, K\}$ ,  $x \in \mathcal{O}$  that  $v_0(x) = g(x)$  and

$$v_k(x) = \mathbb{E}[f_{k-1}(X_{k-1}^{k,x}, v_{k-1}(X_{k-1}^{k,x}))] \quad (79)$$

(cf. Lemma 3.2). Then it holds for all  $k \in \{0, 1, \dots, K\}$ ,  $x \in \mathcal{O}$  that

$$\left( \mathbb{E}[|v_k(X_k^{K,x})|^2] \right)^{1/2} \leq \exp \left( \sum_{l=0}^{K-1} L_l \right) \left[ \left( \mathbb{E}[|g(X_0^{K,x})|^2] \right)^{1/2} + \sum_{l=0}^{K-1} \left( \mathbb{E}[|f_l(X_l^{K,x}, 0)|^2] \right)^{1/2} \right]. \quad (80)$$

*Proof of Lemma 3.6.* First, observe that Lemma 3.2 ensures that for all  $k \in \{0, 1, \dots, K\}$ ,  $x \in \mathcal{O}$  it holds that

$$\mathbb{E}[|v_k(X_k^{K,x})|^2] < \infty. \quad (81)$$

Moreover, note that Lemma 3.4 yields that for all  $k \in \{0, 1, \dots, K\}$ ,  $x \in \mathcal{O}$  it holds that

$$v_k(x) = \mathbb{E}[g(X_0^{k,x})] + \sum_{l=0}^{k-1} \mathbb{E}[f_l(X_l^{k,x}, v_l(X_l^{k,x})) - v_l(X_l^{k,x})]. \quad (82)$$

Minkowski's inequality therefore guarantees for all  $k \in \{0, 1, \dots, K\}$ ,  $x \in \mathcal{O}$  that

$$\begin{aligned} \left( \mathbb{E}[|v_k(X_k^{K,x})|^2] \right)^{1/2} &= \left[ \int_{\mathcal{O}} |v_k(y)|^2 (X_k^{K,x}(\mathbb{P}))(dy) \right]^{1/2} \\ &\leq \left[ \int_{\mathcal{O}} |\mathbb{E}[g(X_0^{k,y})]|^2 (X_k^{K,x}(\mathbb{P}))(dy) \right]^{1/2} \\ &\quad + \sum_{l=0}^{k-1} \left[ \int_{\mathcal{O}} |\mathbb{E}[f_l(X_l^{k,y}, v_l(X_l^{k,y})) - v_l(X_l^{k,y})]|^2 (X_k^{K,x}(\mathbb{P}))(dy) \right]^{1/2}. \end{aligned} \quad (83)$$

Jensen's inequality hence yields for all  $k \in \{0, 1, \dots, K\}$ ,  $x \in \mathcal{O}$  that

$$\begin{aligned} \left( \mathbb{E}[|v_k(X_k^{K,x})|^2] \right)^{1/2} &\leq \left[ \int_{\mathcal{O}} \mathbb{E}[|g(X_0^{k,y})|^2] (X_k^{K,x}(\mathbb{P}))(dy) \right]^{1/2} \\ &\quad + \sum_{l=0}^{k-1} \left[ \int_{\mathcal{O}} \mathbb{E}[|f_l(X_l^{k,y}, v_l(X_l^{k,y})) - v_l(X_l^{k,y})|^2] (X_k^{K,x}(\mathbb{P}))(dy) \right]^{1/2}. \end{aligned} \quad (84)$$

Next observe that item (i) of Lemma 2.2 yields that for all  $k \in \{0, 1, \dots, K\}$ ,  $x \in \mathcal{O}$  it holds that  $X_k^{K,x} : \Omega \rightarrow \mathcal{O}$  is  $\sigma((W_m)_{m \in [k, K) \cap \mathbb{N}_0}) / \mathcal{B}(\mathcal{O})$ -measurable. This, the fact that for all  $k \in \{0, 1, \dots, K\}$  it holds that  $\mathcal{O} \times \Omega \ni (x, \omega) \mapsto g(X_0^{k,x}(\omega)) \in \mathbb{R}$  is  $(\mathcal{B}(\mathcal{O}) \otimes \sigma((W_m)_{m \in [0, k) \cap \mathbb{N}_0})) / \mathcal{B}(\mathbb{R})$ -measurable, the assumption that  $W_l : \Omega \rightarrow S$ ,  $l \in \{0, 1, \dots, K-1\}$ , are independent random variables, and Hutzenthaler et al. [36, Lemma 2.2] therefore demonstrate that for all  $k \in \{0, 1, \dots, K\}$ ,  $x \in \mathcal{O}$  it holds that

$$\int_{\mathcal{O}} \mathbb{E}[|g(X_0^{k,y})|^2] (X_k^{K,x}(\mathbb{P}))(dy) = \mathbb{E}\left[|g(X_0^{k, X_k^{K,x}})|^2\right]. \quad (85)$$

Moreover, note that the fact that for all  $k, l \in \mathbb{N}_0$ ,  $x \in \mathcal{O}$  with  $l \leq k \leq K$  it holds that  $X_l^{k,x} : \Omega \rightarrow \mathcal{O}$  is  $\sigma((W_m)_{m \in [l, k) \cap \mathbb{N}_0}) / \mathcal{B}(\mathcal{O})$ -measurable (cf. item (i) of Lemma 2.2), the assumption that  $W_l : \Omega \rightarrow S$ ,  $l \in \{0, 1, \dots, K-1\}$ , are independent random variables, and Hutzenthaler et al. [36, Lemma 2.2] demonstrate that for all  $k \in \{0, 1, \dots, K\}$ ,  $l \in \{0, 1, \dots, k-1\}$ ,  $x \in \mathcal{O}$  it holds that

$$\begin{aligned} &\sum_{l=0}^{k-1} \int_{\mathcal{O}} \mathbb{E}[|f_l(X_l^{k,y}, v_l(X_l^{k,y})) - v_l(X_l^{k,y})|^2] (X_k^{K,x}(\mathbb{P}))(dy) \\ &= \sum_{l=0}^{k-1} \mathbb{E}\left[|f_l(X_l^{k, X_k^{K,x}}, v_l(X_l^{k, X_k^{K,x}})) - v_l(X_l^{k, X_k^{K,x}})|^2\right]. \end{aligned} \quad (86)$$

This, (78), (84), and (85) ensure for all  $k \in \{0, 1, \dots, K\}$ ,  $x \in \mathcal{O}$  that

$$\begin{aligned} &\left( \mathbb{E}[|v_k(X_k^{K,x})|^2] \right)^{1/2} \\ &\leq \left( \mathbb{E}\left[|g(X_0^{k, X_k^{K,x}})|^2\right] \right)^{1/2} + \sum_{l=0}^{k-1} \left( \mathbb{E}\left[|f_l(X_l^{k, X_k^{K,x}}, v_l(X_l^{k, X_k^{K,x}})) - v_l(X_l^{k, X_k^{K,x}})|^2\right] \right)^{1/2} \\ &= \left( \mathbb{E}[|g(X_0^{K,x})|^2] \right)^{1/2} + \sum_{l=0}^{k-1} \left( \mathbb{E}[|f_l(X_l^{K,x}, v_l(X_l^{K,x})) - v_l(X_l^{K,x})|^2] \right)^{1/2}. \end{aligned} \quad (87)$$

Minkowski's inequality and the assumption that for all  $k \in \{0, 1, \dots, K-1\}$ ,  $a, b \in \mathbb{R}$ ,  $x \in \mathcal{O}$  it holds that  $|(f_k(x, a) - a) - (f_k(x, b) - b)| \leq L_k |a - b|$  hence guarantee that for all  $k \in \{0, 1, \dots, K\}$ ,

$x \in \mathcal{O}$  it holds that

$$\begin{aligned}
& \left( \mathbb{E}[|v_k(X_k^{K,x})|^2] \right)^{1/2} \\
& \leq \left( \mathbb{E}[|g(X_0^{K,x})|^2] \right)^{1/2} + \sum_{l=0}^{k-1} \left[ \left( \mathbb{E}[|f_l(X_l^{K,x}, 0)|^2] \right)^{1/2} + L_l \left( \mathbb{E}[|v_l(X_l^{K,x})|^2] \right)^{1/2} \right] \\
& = \left[ \left( \mathbb{E}[|g(X_0^{K,x})|^2] \right)^{1/2} + \sum_{l=0}^{K-1} \left( \mathbb{E}[|f_l(X_l^{K,x}, 0)|^2] \right)^{1/2} \right] + \sum_{l=0}^{k-1} L_l \left( \mathbb{E}[|v_l(X_l^{K,x})|^2] \right)^{1/2}.
\end{aligned} \tag{88}$$

Lemma 3.5 and (81) hence ensure for all  $k \in \{0, 1, \dots, K\}$ ,  $x \in \mathcal{O}$  that

$$\left( \mathbb{E}[|v_k(X_k^{K,x})|^2] \right)^{1/2} \leq \exp \left( \sum_{l=0}^{k-1} L_l \right) \left[ \left( \mathbb{E}[|g(X_0^{K,x})|^2] \right)^{1/2} + \sum_{l=0}^{K-1} \left( \mathbb{E}[|f_l(X_l^{K,x}, 0)|^2] \right)^{1/2} \right]. \tag{89}$$

This establishes (80). This completes the proof of Lemma 3.6.  $\square$

### 3.2 Bias and variance estimates for MLP approximations

**Lemma 3.7.** Assume Setting 2.1, let  $c, L_0, L_1, \dots, L_K \in [0, \infty)$  satisfy for all  $k \in \{0, 1, \dots, K\}$ ,  $x \in \mathcal{O}$ ,  $a, b \in \mathbb{R}$  that  $|(f_k(x, a) - a) - (f_k(x, b) - b)| \leq L_k|a - b|$ , assume for all  $k, l \in \mathbb{N}_0$  with  $l < k \leq K$  that  $L_l \leq c\mathbb{P}(\mathcal{R}_k^0 = l)$ , and assume for all  $k \in \{0, 1, \dots, K\}$ ,  $x \in \mathcal{O}$  that  $\mathbb{E}[|g(X_0^{0,k,x})|^2 + \sum_{l=0}^{k-1} |f_l(X_l^{0,k,x}, 0)|^2] < \infty$ . Then it holds for all  $k \in \{1, 2, \dots, K\}$ ,  $n \in \mathbb{N}$ ,  $x \in \mathcal{O}$  that  $\mathbb{E}[|g(X_0^{0,k,x})| + \sum_{l=0}^{k-1} |f_l(X_l^{0,k,x}, V_{l,n-1}^0(X_l^{0,k,x})) - V_{l,n-1}^0(X_l^{0,k,x})|] < \infty$  and

$$\mathbb{E}[V_{k,n}^0(x)] = \mathbb{E}[g(X_0^{0,k,x})] + \sum_{l=0}^{k-1} \mathbb{E}[f_l(X_l^{0,k,x}, V_{l,n-1}^0(X_l^{0,k,x})) - V_{l,n-1}^0(X_l^{0,k,x})]. \tag{90}$$

*Proof of Lemma 3.7.* First, observe that Lemma 2.11 guarantees that for all  $k, l, n \in \mathbb{N}_0$ ,  $x \in \mathcal{O}$  with  $l \leq k \leq K$  it holds that  $\mathbb{E}[|V_{l,n}^0(X_l^{0,k,x})|^2] < \infty$ . This, the assumption that for all  $k \in \{0, 1, \dots, K-1\}$ ,  $x \in \mathcal{O}$ ,  $a, b \in \mathbb{R}$  it holds that  $|(f_k(x, a) - a) - (f_k(x, b) - b)| \leq L_k|a - b|$ , the assumption that for all  $k \in \{0, 1, \dots, K\}$ ,  $x \in \mathcal{O}$  it holds that  $\mathbb{E}[|g(X_0^{0,k,x})|^2 + \sum_{l=0}^{k-1} |f_l(X_l^{0,k,x}, 0)|^2] < \infty$ , and (4) ensure that for all  $k \in \{1, 2, \dots, K\}$ ,  $n \in \mathbb{N}$ ,  $x \in \mathcal{O}$  it holds that  $\mathbb{E}[|g(X_0^{0,k,x})| + \sum_{l=0}^{k-1} |f_l(X_l^{0,k,x}, V_{l,n-1}^0(X_l^{0,k,x})) - V_{l,n-1}^0(X_l^{0,k,x})|] < \infty$  and

$$\begin{aligned}
\mathbb{E}[V_{k,n}^0(x)] &= \frac{1}{M^n} \sum_{m=1}^{M^n} \mathbb{E} \left[ g(X_0^{(0,0,-m),k,x}) + \sum_{l=0}^{k-1} f_l(X_l^{(0,0,m),k,x}, 0) \right] + \sum_{j=1}^{n-1} \sum_{m=1}^{M^{n-j}} \frac{1}{M^{n-j}} \\
&\cdot \left[ \mathbb{E} \left[ \frac{1}{\mathbf{p}_{k, \mathcal{R}_k^{(0,j,m)}}} \left( f_{\mathcal{R}_k^{(0,j,m)}} \left( X_{\mathcal{R}_k^{(0,j,m)}}^{(0,j,m),k,x}, V_{\mathcal{R}_k^{(0,j,m)},j}^{(0,j,m)}(X_{\mathcal{R}_k^{(0,j,m)}}^{(0,j,m),k,x}) \right) - V_{\mathcal{R}_k^{(0,j,m)},j}^{(0,j,m)}(X_{\mathcal{R}_k^{(0,j,m)}}^{(0,j,m),k,x}) \right) \right. \right. \\
&\left. \left. - \frac{1}{\mathbf{p}_{k, \mathcal{R}_k^{(0,j,m)}}} \left( f_{\mathcal{R}_k^{(0,j,m)}} \left( X_{\mathcal{R}_k^{(0,j,m)}}^{(0,j,m),k,x}, V_{\mathcal{R}_k^{(0,j,m)},j-1}^{(0,j,-m)}(X_{\mathcal{R}_k^{(0,j,m)}}^{(0,j,m),k,x}) \right) - V_{\mathcal{R}_k^{(0,j,m)},j-1}^{(0,j,-m)}(X_{\mathcal{R}_k^{(0,j,m)}}^{(0,j,m),k,x}) \right) \right] \right].
\end{aligned} \tag{91}$$

Next note that the fact that for all  $\theta \in \Theta$ ,  $k \in \{1, 2, \dots, K\}$  it holds that  $\mathbb{P}(\mathcal{R}_k^\theta \in \{0, 1, \dots, k-1\}) =$

1 ensures that for all  $n, m \in \mathbb{N}$ ,  $k \in \{1, 2, \dots, K\}$ ,  $j \in \{1, 2, \dots, n-1\}$ ,  $x \in \mathcal{O}$  it holds that

$$\begin{aligned}
& \mathbb{E} \left[ \frac{1}{\mathbf{p}_{k, \mathcal{R}_k^{(0,j,m)}}} \left( f_{\mathcal{R}_k^{(0,j,m)}}(X_{\mathcal{R}_k^{(0,j,m)}}^{(0,j,m),k,x}, V_{\mathcal{R}_k^{(0,j,m)},j}^{(0,j,m)}(X_{\mathcal{R}_k^{(0,j,m)}}^{(0,j,m),k,x})) - V_{\mathcal{R}_k^{(0,j,m)},j}^{(0,j,m)}(X_{\mathcal{R}_k^{(0,j,m)}}^{(0,j,m),k,x}) \right) \right] \\
& - \mathbb{E} \left[ \frac{1}{\mathbf{p}_{k, \mathcal{R}_k^{(0,j,m)}}} \left( f_{\mathcal{R}_k^{(0,j,m)}}(X_{\mathcal{R}_k^{(0,j,m)}}^{(0,j,m),k,x}, V_{\mathcal{R}_k^{(0,j,m)},j-1}^{(0,j,-m)}(X_{\mathcal{R}_k^{(0,j,m)}}^{(0,j,m),k,x})) - V_{\mathcal{R}_k^{(0,j,m)},j-1}^{(0,j,-m)}(X_{\mathcal{R}_k^{(0,j,m)}}^{(0,j,m),k,x}) \right) \right] \\
& = \sum_{l=0}^{k-1} \mathbb{E} \left[ \frac{\mathbb{1}_{\{\mathcal{R}_k^{(0,j,m)}=l\}}}{\mathbf{p}_{k,l}} \left( f_l(X_l^{(0,j,m),k,x}, V_{l,j}^{(0,j,m)}(X_l^{(0,j,m),k,x})) - V_{l,j}^{(0,j,m)}(X_l^{(0,j,m),k,x}) \right) \right] \\
& - \sum_{l=0}^{k-1} \mathbb{E} \left[ \frac{\mathbb{1}_{\{\mathcal{R}_k^{(0,j,m)}=l\}}}{\mathbf{p}_{k,l}} \left( f_l(X_l^{(0,j,m),k,x}, V_{l,j-1}^{(0,j,-m)}(X_l^{(0,j,m),k,x})) - V_{l,j-1}^{(0,j,-m)}(X_l^{(0,j,m),k,x}) \right) \right]. \tag{92}
\end{aligned}$$

Item (i) of Lemma 2.2, item (i) of Lemma 2.3, and the fact that for all  $n, m \in \mathbb{N}$ ,  $j \in \{1, 2, \dots, n-1\}$  it holds that  $\sigma((\mathcal{R}_s^{(0,j,m)})_{s \in \{1,2,\dots,K\}})$  and  $\sigma((\mathcal{R}_s^{(0,j,m,\vartheta)})_{(s,\vartheta) \in \{1,2,\dots,K\} \times \Theta}, (\mathcal{R}_s^{(0,-j,m,\vartheta)})_{(s,\vartheta) \in \{1,2,\dots,K\} \times \Theta}, (W_s^{(0,j,m,\vartheta)})_{(s,\vartheta) \in \{0,1,\dots,K-1\} \times \Theta}, (W_s^{(0,j,-m,\vartheta)})_{(s,\vartheta) \in \{0,1,\dots,K-1\} \times \Theta}, (W_s^{(0,j,m)})_{s \in \{0,1,\dots,K-1\}})$  are independent therefore ensure that for all  $n, m \in \mathbb{N}$ ,  $k \in \{1, 2, \dots, K\}$ ,  $j \in \{1, 2, \dots, n-1\}$ ,  $x \in \mathcal{O}$  it holds that

$$\begin{aligned}
& \mathbb{E} \left[ \frac{1}{\mathbf{p}_{k, \mathcal{R}_k^{(0,j,m)}}} \left( f_{\mathcal{R}_k^{(0,j,m)}}(X_{\mathcal{R}_k^{(0,j,m)}}^{(0,j,m),k,x}, V_{\mathcal{R}_k^{(0,j,m)},j}^{(0,j,m)}(X_{\mathcal{R}_k^{(0,j,m)}}^{(0,j,m),k,x})) - V_{\mathcal{R}_k^{(0,j,m)},j}^{(0,j,m)}(X_{\mathcal{R}_k^{(0,j,m)}}^{(0,j,m),k,x}) \right) \right] \\
& - \mathbb{E} \left[ \frac{1}{\mathbf{p}_{k, \mathcal{R}_k^{(0,j,m)}}} \left( f_{\mathcal{R}_k^{(0,j,m)}}(X_{\mathcal{R}_k^{(0,j,m)}}^{(0,j,m),k,x}, V_{\mathcal{R}_k^{(0,j,m)},j-1}^{(0,j,-m)}(X_{\mathcal{R}_k^{(0,j,m)}}^{(0,j,m),k,x})) - V_{\mathcal{R}_k^{(0,j,m)},j-1}^{(0,j,-m)}(X_{\mathcal{R}_k^{(0,j,m)}}^{(0,j,m),k,x}) \right) \right] \\
& = \sum_{l=0}^{k-1} \frac{\mathbb{P}(\mathcal{R}_k^{(0,j,m)} = l)}{\mathbf{p}_{k,l}} \left[ \mathbb{E} \left[ f_l(X_l^{(0,j,m),k,x}, V_{l,j}^{(0,j,m)}(X_l^{(0,j,m),k,x})) - V_{l,j}^{(0,j,m)}(X_l^{(0,j,m),k,x}) \right] \right. \\
& \left. - \mathbb{E} \left[ f_l(X_l^{(0,j,m),k,x}, V_{l,j-1}^{(0,j,-m)}(X_l^{(0,j,m),k,x})) - V_{l,j-1}^{(0,j,-m)}(X_l^{(0,j,m),k,x}) \right] \right]. \tag{93}
\end{aligned}$$

The fact that for all  $\theta \in \Theta$ ,  $k \in \{1, 2, \dots, K\}$ ,  $l \in \{0, 1, \dots, k-1\}$  with  $\mathbb{P}(\mathcal{R}_k^\theta = l) > 0$  it holds that  $\mathbb{P}(\mathcal{R}_k^\theta = l) = \mathbf{p}_{k,l}$  and the fact that for all  $\theta \in \Theta$ ,  $k \in \{1, 2, \dots, K\}$ ,  $l \in \{0, 1, \dots, k-1\}$  with  $\mathbb{P}(\mathcal{R}_k^\theta = l) = 0$  it holds that  $L_l = 0$  hence yield that for all  $n, m \in \mathbb{N}$ ,  $k \in \{1, 2, \dots, K\}$ ,  $j \in \{1, 2, \dots, n-1\}$ ,  $x \in \mathcal{O}$  it holds that

$$\begin{aligned}
& \mathbb{E} \left[ \frac{1}{\mathbf{p}_{k, \mathcal{R}_k^{(0,j,m)}}} \left( f_{\mathcal{R}_k^{(0,j,m)}}(X_{\mathcal{R}_k^{(0,j,m)}}^{(0,j,m),k,x}, V_{\mathcal{R}_k^{(0,j,m)},j}^{(0,j,m)}(X_{\mathcal{R}_k^{(0,j,m)}}^{(0,j,m),k,x})) - V_{\mathcal{R}_k^{(0,j,m)},j}^{(0,j,m)}(X_{\mathcal{R}_k^{(0,j,m)}}^{(0,j,m),k,x}) \right) \right] \\
& - \mathbb{E} \left[ \frac{1}{\mathbf{p}_{k, \mathcal{R}_k^{(0,j,m)}}} \left( f_{\mathcal{R}_k^{(0,j,m)}}(X_{\mathcal{R}_k^{(0,j,m)}}^{(0,j,m),k,x}, V_{\mathcal{R}_k^{(0,j,m)},j-1}^{(0,j,-m)}(X_{\mathcal{R}_k^{(0,j,m)}}^{(0,j,m),k,x})) - V_{\mathcal{R}_k^{(0,j,m)},j-1}^{(0,j,-m)}(X_{\mathcal{R}_k^{(0,j,m)}}^{(0,j,m),k,x}) \right) \right] \\
& = \sum_{l=0}^{k-1} \left[ \mathbb{E} \left[ f_l(X_l^{(0,j,m),k,x}, V_{l,j}^{(0,j,m)}(X_l^{(0,j,m),k,x})) - V_{l,j}^{(0,j,m)}(X_l^{(0,j,m),k,x}) \right] \right. \\
& \left. - \mathbb{E} \left[ f_l(X_l^{(0,j,m),k,x}, V_{l,j-1}^{(0,j,-m)}(X_l^{(0,j,m),k,x})) - V_{l,j-1}^{(0,j,-m)}(X_l^{(0,j,m),k,x}) \right] \right]. \tag{94}
\end{aligned}$$

This and Corollary 2.10 assure that for all  $n, m \in \mathbb{N}$ ,  $k \in \{1, 2, \dots, K\}$ ,  $j \in \{0, 1, \dots, n-1\}$ ,  $x \in \mathcal{O}$

it holds that

$$\begin{aligned}
& \mathbb{E} \left[ \frac{1}{\mathbf{p}_{k, \mathcal{R}_k^{(0,j,m)}}} \left( f_{\mathcal{R}_k^{(0,j,m)}}(X_{\mathcal{R}_k^{(0,j,m)}}^{(0,j,m),k,x}, V_{\mathcal{R}_k^{(0,j,m)},j}^{(0,j,m)}(X_{\mathcal{R}_k^{(0,j,m)}}^{(0,j,m),k,x})) - V_{\mathcal{R}_k^{(0,j,m)},j}^{(0,j,m)}(X_{\mathcal{R}_k^{(0,j,m)}}^{(0,j,m),k,x}) \right) \right] \\
& - \mathbb{E} \left[ \frac{1}{\mathbf{p}_{k, \mathcal{R}_k^{(0,j,m)}}} \left( f_{\mathcal{R}_k^{(0,j,m)}}(X_{\mathcal{R}_k^{(0,j,m)}}^{(0,j,m),k,x}, V_{\mathcal{R}_k^{(0,j,m)},j-1}^{(0,j,m)}(X_{\mathcal{R}_k^{(0,j,m)}}^{(0,j,m),k,x})) - V_{\mathcal{R}_k^{(0,j,m)},j-1}^{(0,j,m)}(X_{\mathcal{R}_k^{(0,j,m)}}^{(0,j,m),k,x}) \right) \right] \\
& = \sum_{l=0}^{k-1} \mathbb{E}[f_l(X_l^{0,k,x}, V_{l,j}^0(X_l^{0,k,x})) - V_{l,j}^0(X_l^{0,k,x})] \\
& - \sum_{l=0}^{k-1} \mathbb{E}[f_l(X_l^{0,k,x}, V_{l,j-1}^0(X_l^{0,k,x})) - V_{l,j-1}^0(X_l^{0,k,x})].
\end{aligned} \tag{95}$$

This, (91), and item (i) of Lemma 2.8 imply that for all  $k \in \{1, 2, \dots, K\}$ ,  $n \in \mathbb{N}$ ,  $x \in \mathcal{O}$  it holds that

$$\begin{aligned}
\mathbb{E}[V_{k,n}^0(x)] &= \mathbb{E}[g(X_0^{0,k,x})] + \sum_{l=0}^{k-1} \mathbb{E}[f_l(X_l^{0,k,x}, 0)] \\
&+ \sum_{j=1}^{n-1} \sum_{l=0}^{k-1} \mathbb{E}[f_l(X_l^{0,k,x}, V_{l,j}^0(X_l^{0,k,x})) - V_{l,j}^0(X_l^{0,k,x})] \\
&- \sum_{j=1}^{n-1} \sum_{l=0}^{k-1} \mathbb{E}[f_l(X_l^{0,k,x}, V_{l,j-1}^0(X_l^{0,k,x})) - V_{l,j-1}^0(X_l^{0,k,x})] \\
&= \mathbb{E}[g(X_0^{0,k,x})] + \sum_{l=0}^{k-1} \mathbb{E}[f_l(X_l^{0,k,x}, 0)] \\
&+ \sum_{l=0}^{k-1} \mathbb{E}[f_l(X_l^{0,k,x}, V_{l,n-1}^0(X_l^{0,k,x})) - V_{l,n-1}^0(X_l^{0,k,x})] \\
&- \sum_{l=0}^{k-1} \mathbb{E}[f_l(X_l^{0,k,x}, V_{l,0}^0(X_l^{0,k,x})) - V_{l,0}^0(X_l^{0,k,x})].
\end{aligned} \tag{96}$$

The fact that for all  $k \in \{0, 1, \dots, K\}$ ,  $x \in \mathcal{O}$  it holds that  $V_{k,0}^0(x) = 0$  therefore implies that for all  $k \in \{1, 2, \dots, K\}$ ,  $n \in \mathbb{N}$ ,  $x \in \mathcal{O}$  it holds that

$$\mathbb{E}[V_{k,n}^0(x)] = \mathbb{E}[g(X_0^{0,k,x})] + \sum_{l=0}^{k-1} \mathbb{E}[f_l(X_l^{0,k,x}, V_{l,n-1}^0(X_l^{0,k,x})) - V_{l,n-1}^0(X_l^{0,k,x})]. \tag{97}$$

This establishes (90). This completes the proof of Lemma 3.7.  $\square$

**Lemma 3.8** (Bias estimate). *Assume Setting 2.1, let  $c, L_0, L_1, \dots, L_K \in [0, \infty)$  satisfy for all  $k \in \{0, 1, \dots, K\}$ ,  $x \in \mathcal{O}$ ,  $a, b \in \mathbb{R}$  that  $|(f_k(x, a) - a) - (f_k(x, b) - b)| \leq L_k|a - b|$ , assume for all  $k, l \in \mathbb{N}_0$  with  $l < k \leq K$  that  $L_l \leq c\mathbb{P}(\mathcal{R}_k^0 = l)$ , assume for all  $k \in \{0, 1, \dots, K\}$ ,  $x \in \mathcal{O}$  that  $\mathbb{E}[|g(X_0^{0,k,x})|^2 + \sum_{l=0}^{k-1} |f_l(X_l^{0,k,x}, 0)|^2] < \infty$ , and let  $v_k: \mathcal{O} \rightarrow \mathbb{R}$ ,  $k \in \{0, 1, \dots, K\}$ , be  $\mathcal{B}(\mathcal{O})/\mathcal{B}(\mathbb{R})$ -measurable functions which satisfy for all  $k \in \{1, 2, \dots, K\}$ ,  $x \in \mathcal{O}$  that  $v_0(x) = g(x)$  and*

$$v_k(x) = \mathbb{E}[f_{k-1}(X_{k-1}^{0,k,x}, v_{k-1}(X_{k-1}^{0,k,x}))] \tag{98}$$

(cf. Lemma 3.2). Then it tholds for all  $k \in \{1, 2, \dots, K\}$ ,  $n \in \mathbb{N}$ ,  $x \in \mathcal{O}$  that

$$|\mathbb{E}[V_{k,n}^0(x)] - v_k(x)|^2 \leq \left( \sum_{l=0}^{k-1} L_l \right) \left( \sum_{l=0}^{k-1} L_l \mathbb{E}[|V_{l,n-1}^0(X_l^{0,k,x}) - v_l(X_l^{0,k,x})|^2] \right). \tag{99}$$

*Proof of Lemma 3.8.* First, observe that Lemma 3.4 ensures for all  $k \in \{1, 2, \dots, K\}$ ,  $x \in \mathcal{O}$  that

$$v_k(x) = \mathbb{E}[g(X_0^{0,k,x})] + \sum_{l=0}^{k-1} \mathbb{E}[f_l(X_l^{0,k,x}, v_l(X_l^{0,k,x})) - v_l(X_l^{0,k,x})]. \quad (100)$$

Moreover, note that Lemma 3.7 proves that for all  $k \in \{1, 2, \dots, K\}$ ,  $n \in \mathbb{N}$ ,  $x \in \mathcal{O}$  it holds that

$$\mathbb{E}[V_{k,n}^0(x)] = \mathbb{E}[g(X_0^{0,k,x})] + \sum_{l=0}^{k-1} \mathbb{E}[f_l(X_l^{0,k,x}, V_{l,n-1}^0(X_l^{0,k,x})) - V_{l,n-1}^0(X_l^{0,k,x})]. \quad (101)$$

This and (100) ensure that for all  $k \in \{1, 2, \dots, K\}$ ,  $n \in \mathbb{N}$ ,  $x \in \mathcal{O}$  it holds that

$$\begin{aligned} & \mathbb{E}[V_{k,n}^0(x)] - v_k(x) \\ &= \sum_{l=0}^{k-1} \mathbb{E}[(f_l(X_l^{0,k,x}, V_{l,n-1}^0(X_l^{0,k,x})) - V_{l,n-1}^0(X_l^{0,k,x})) - (f_l(X_l^{0,k,x}, v_l(X_l^{0,k,x})) - v_l(X_l^{0,k,x}))]. \end{aligned} \quad (102)$$

The assumption that for all  $k \in \{0, 1, \dots, K-1\}$ ,  $a, b \in \mathbb{R}$ ,  $x \in \mathcal{O}$  it holds that  $|(f_k(x, a) - a) - (f_k(x, b) - b)| \leq L_k |a - b|$  hence yields that for all  $k \in \{1, 2, \dots, K\}$ ,  $n \in \mathbb{N}$ ,  $x \in \mathcal{O}$  it holds that

$$|\mathbb{E}[V_{k,n}^0(x)] - v_k(x)| \leq \sum_{l=0}^{k-1} L_l \mathbb{E}[|V_{l,n-1}^0(X_l^{0,k,x}) - v_l(X_l^{0,k,x})|]. \quad (103)$$

The Cauchy-Schwarz inequality therefore shows that for all  $k \in \{0, 1, \dots, K\}$ ,  $n \in \mathbb{N}$ ,  $x \in \mathcal{O}$  it holds that

$$\begin{aligned} |\mathbb{E}[V_{k,n}^0(x)] - v_k(x)|^2 &\leq \left[ \sum_{l=0}^{k-1} L_l^{1/2} \left( L_l^{1/2} \mathbb{E}[|V_{l,n-1}^0(X_l^{0,k,x}) - v_l(X_l^{0,k,x})|] \right) \right]^2 \\ &\leq \left( \sum_{l=0}^{k-1} L_l \right) \left( \sum_{l=0}^{k-1} L_l \mathbb{E}[|V_{l,n-1}^0(X_l^{0,k,x}) - v_l(X_l^{0,k,x})|]^2 \right) \\ &\leq \left( \sum_{l=0}^{k-1} L_l \right) \left( \sum_{l=0}^{k-1} L_l \mathbb{E}[|V_{l,n-1}^0(X_l^{0,k,x}) - v_l(X_l^{0,k,x})|^2] \right). \end{aligned} \quad (104)$$

This establishes (99). This completes the proof of Lemma 3.8.  $\square$

**Lemma 3.9** (Variance estimate). *Assume Setting 2.1, let  $L_0, L_1, \dots, L_K \in [0, \infty)$  satisfy for all  $k \in \{0, 1, \dots, K\}$ ,  $x \in \mathcal{O}$ ,  $a, b \in \mathbb{R}$  that  $|(f_k(x, a) - a) - (f_k(x, b) - b)| \leq L_k |a - b|$ , and assume for all  $k \in \{0, 1, \dots, K\}$ ,  $x \in \mathcal{O}$  that  $\mathbb{E}[|g(X_0^{0,k,x})|^2 + \sum_{l=0}^{k-1} |f_l(X_l^{0,k,x}, 0)|^2] < \infty$ . Then it holds for all  $k \in \{1, 2, \dots, K\}$ ,  $n \in \mathbb{N}$ ,  $x \in \mathcal{O}$  that*

$$\begin{aligned} \text{Var}[V_{k,n}^0(x)] &\leq \frac{1}{M^n} \mathbb{E}[|g(X_0^{0,k,x})|^2 + |\sum_{l=0}^{k-1} f_l(X_l^{0,k,x}, 0)|^2] \\ &\quad + \sum_{j=1}^{n-1} \frac{1}{M^{n-j}} \left[ \max_{l \in \{0, 1, \dots, k-1\}} \frac{L_l}{\mathbf{p}_{k,l}} \right] \left( \sum_{l=0}^{k-1} L_l \mathbb{E}[|V_{l,j}^0(X_l^{0,k,x}) - V_{l,j-1}^1(X_l^{0,k,x})|^2] \right). \end{aligned} \quad (105)$$

*Proof of Lemma 3.9.* First, observe that item (i) of Lemma 2.2, item (ii) of Lemma 2.3, and the fact that for all  $k \in \{1, 2, \dots, K\}$  it holds that  $\{W_s^\theta, \mathcal{R}_k^\theta: \theta \in \Theta, s \in \{0, 1, \dots, K-1\}\}$  is a family of

independent random variables ensure that for all  $k \in \{1, 2, \dots, K\}$ ,  $n \in \mathbb{N}$ ,  $x \in \mathcal{O}$  it holds that

$$\begin{aligned} \text{Var}[V_{k,n}^0(x)] &= \sum_{m=1}^{M^n} \text{Var}\left[\frac{g(X_0^{(0,0,-m),k,x})}{M^n}\right] + \sum_{m=1}^{M^n} \text{Var}\left[\frac{\sum_{l=0}^{k-1} f_l(X_l^{(0,0,m),k,x}, 0)}{M^n}\right] \\ &+ \sum_{j=1}^{n-1} \sum_{m=1}^{M^{n-j}} \text{Var}\left[\left(\frac{f_{\mathcal{R}_k^{(0,j,m)}}(X_{\mathcal{R}_k^{(0,j,m)}}^{(0,j,m),k,x}, V_{\mathcal{R}_k^{(0,j,m),j}}^{(0,j,m)}(X_{\mathcal{R}_k^{(0,j,m)}}^{(0,j,m),k,x})) - V_{\mathcal{R}_k^{(0,j,m),j}}^{(0,j,m)}(X_{\mathcal{R}_k^{(0,j,m)}}^{(0,j,m),k,x})}{M^{n-j}}\right.\right. \\ &\quad \left.\left.- \frac{f_{\mathcal{R}_k^{(0,j,m)}}(X_{\mathcal{R}_k^{(0,j,m)}}^{(0,j,m),k,x}, V_{\mathcal{R}_k^{(0,j,m),j-1}}^{(0,j,-m)}(X_{\mathcal{R}_k^{(0,j,m)}}^{(0,j,m),k,x})) - V_{\mathcal{R}_k^{(0,j,m),j-1}}^{(0,j,-m)}(X_{\mathcal{R}_k^{(0,j,m)}}^{(0,j,m),k,x})}{M^{n-j}}\right) \cdot \frac{1}{\mathbf{p}_{k,\mathcal{R}_k^{(0,j,m)}}}\right]. \end{aligned} \quad (106)$$

Item (i) of Lemma 2.8 and item (iii) of Proposition 2.9 therefore ensure that for all  $k \in \{1, 2, \dots, K\}$ ,  $n \in \mathbb{N}$ ,  $x \in \mathcal{O}$  it holds that

$$\begin{aligned} \text{Var}[V_{k,n}^0(x)] &= \frac{1}{M^n} \left( \text{Var}[g(X_0^{0,k,x})] + \text{Var}\left[\sum_{l=0}^{k-1} f_l(X_l^{0,k,x}, 0)\right] \right) \\ &+ \sum_{j=1}^{n-1} \frac{1}{M^{n-j}} \text{Var}\left[\frac{1}{\mathbf{p}_{k,\mathcal{R}_k^0}} (f_{\mathcal{R}_k^0}(X_{\mathcal{R}_k^0}^{0,k,x}, V_{\mathcal{R}_k^0,j}^0(X_{\mathcal{R}_k^0}^{0,k,x})) - V_{\mathcal{R}_k^0,j}^0(X_{\mathcal{R}_k^0}^{0,k,x})) \right. \\ &\quad \left. - (f_{\mathcal{R}_k^0}(X_{\mathcal{R}_k^0}^{0,k,x}, V_{\mathcal{R}_k^0,j-1}^1(X_{\mathcal{R}_k^0}^{0,k,x})) - V_{\mathcal{R}_k^0,j-1}^1(X_{\mathcal{R}_k^0}^{0,k,x})) \right]. \end{aligned} \quad (107)$$

Combining this with the fact that for every random variable  $\mathcal{X}: \Omega \rightarrow \mathbb{R}$  with  $\mathbb{E}[|\mathcal{X}|] < \infty$  it holds that  $\text{Var}[\mathcal{X}] \leq \mathbb{E}[|\mathcal{X}|^2] \in [0, \infty]$  and Lemma 2.11 yields that for all  $k \in \{0, 1, \dots, K\}$ ,  $n \in \mathbb{N}$ ,  $x \in \mathcal{O}$  it holds that

$$\begin{aligned} \text{Var}[V_{k,n}^0(x)] &\leq \frac{1}{M^n} \mathbb{E}\left[|g(X_0^{0,k,x})|^2 + \left|\sum_{l=0}^{k-1} f_l(X_l^{0,k,x}, 0)\right|^2\right] \\ &+ \sum_{j=1}^{n-1} \frac{1}{M^{n-j}} \mathbb{E}\left[\left|\frac{1}{\mathbf{p}_{k,\mathcal{R}_k^0}} (f_{\mathcal{R}_k^0}(X_{\mathcal{R}_k^0}^{0,k,x}, V_{\mathcal{R}_k^0,j}^0(X_{\mathcal{R}_k^0}^{0,k,x})) - V_{\mathcal{R}_k^0,j}^0(X_{\mathcal{R}_k^0}^{0,k,x})) \right.\right. \\ &\quad \left.\left.- (f_{\mathcal{R}_k^0}(X_{\mathcal{R}_k^0}^{0,k,x}, V_{\mathcal{R}_k^0,j-1}^1(X_{\mathcal{R}_k^0}^{0,k,x})) - V_{\mathcal{R}_k^0,j-1}^1(X_{\mathcal{R}_k^0}^{0,k,x}))\right|^2\right]. \end{aligned} \quad (108)$$

Moreover, note that item (ii) of Lemma 2.2, item (i) of Lemma 2.3, the fact that  $\sigma((\mathcal{R}_s^0)_{s \in \{1,2,\dots,K\}})$  and  $\sigma((\mathcal{R}_s^{(0,\eta)})_{(s,\eta) \in \{1,2,\dots,K\} \times \Theta}, (W_s^{(0,\eta)})_{(s,\eta) \in \{0,1,\dots,K-1\} \times \Theta}, (W_s^0)_{s \in \{0,1,\dots,K-1\}})$  are independent sigma-algebras, and the fact that  $\sigma((\mathcal{R}_s^0)_{s \in \{1,2,\dots,K\}})$  and  $\sigma((\mathcal{R}_s^{(1,\eta)})_{(s,\eta) \in \{1,2,\dots,K\} \times \Theta}, (W_s^{(1,\eta)})_{(s,\eta) \in \{0,1,\dots,K-1\} \times \Theta}, (W_s^0)_{s \in \{0,1,\dots,K-1\}})$  are independent sigma-algebras on  $\Omega$  ensure that for all  $k \in \{1, 2, \dots, K\}$ ,  $n \in \mathbb{N}$ ,

$j \in \{1, 2, \dots, n-1\}$ ,  $x \in \mathcal{O}$  it holds that

$$\begin{aligned}
& \mathbb{E} \left[ \left| \frac{1}{\mathfrak{p}_{k, \mathcal{R}_k^0}} (f_{\mathcal{R}_k^0}(X_{\mathcal{R}_k^0}^{0,k,x}, V_{\mathcal{R}_k^0, j}^0(X_{\mathcal{R}_k^0}^{0,k,x})) - V_{\mathcal{R}_k^0, j}^0(X_{\mathcal{R}_k^0}^{0,k,x})) \right. \right. \\
& \quad \left. \left. - (f_{\mathcal{R}_k^0}(X_{\mathcal{R}_k^0}^{0,k,x}, V_{\mathcal{R}_k^0, j-1}^1(X_{\mathcal{R}_k^0}^{0,k,x})) - V_{\mathcal{R}_k^0, j-1}^1(X_{\mathcal{R}_k^0}^{0,k,x})) \right|^2 \right] \\
&= \sum_{l=0}^{k-1} \mathbb{E} \left[ \frac{\mathbb{1}_{\{\mathcal{R}_k^0=l\}}}{|\mathfrak{p}_{k, l}|^2} \left| (f_l(X_l^{0,k,x}, V_{l, j}^0(X_l^{0,k,x})) - V_{l, j}^0(X_l^{0,k,x})) \right. \right. \\
& \quad \left. \left. - (f_l(X_l^{0,k,x}, V_{l, j-1}^1(X_l^{0,k,x})) - V_{l, j-1}^1(X_l^{0,k,x})) \right|^2 \right] \\
&= \sum_{l=0}^{k-1} \frac{1}{\mathfrak{p}_{k, l}} \mathbb{E} \left[ \left| (f_l(X_l^{0,k,x}, V_{l, j}^0(X_l^{0,k,x})) - V_{l, j}^0(X_l^{0,k,x})) \right. \right. \\
& \quad \left. \left. - (f_l(X_l^{0,k,x}, V_{l, j-1}^1(X_l^{0,k,x})) - V_{l, j-1}^1(X_l^{0,k,x})) \right|^2 \right].
\end{aligned} \tag{109}$$

This and (108) yield that for all  $k \in \{1, 2, \dots, K\}$ ,  $n \in \mathbb{N}$ ,  $x \in \mathcal{O}$  it holds that

$$\begin{aligned}
\text{Var}[V_{k, n}^0(x)] &\leq \frac{1}{M^n} \mathbb{E} \left[ |g(X_0^{0,k,x})|^2 + \left| \sum_{l=0}^{k-1} f_l(X_l^{0,k,x}, 0) \right|^2 \right] \\
&+ \sum_{j=1}^{n-1} \frac{1}{M^{n-j}} \sum_{l=0}^{k-1} \frac{1}{\mathfrak{p}_{k, l}} \mathbb{E} \left[ \left| (f_l(X_l^{0,k,x}, V_{l, j}^0(X_l^{0,k,x})) - V_{l, j}^0(X_l^{0,k,x})) \right. \right. \\
& \quad \left. \left. - (f_l(X_l^{0,k,x}, V_{l, j-1}^1(X_l^{0,k,x})) - V_{l, j-1}^1(X_l^{0,k,x})) \right|^2 \right].
\end{aligned} \tag{110}$$

The assumption that for all  $k \in \{0, 1, \dots, K-1\}$ ,  $a, b \in \mathbb{R}$ ,  $x \in \mathcal{O}$  it holds that  $|(f_k(x, a) - a) - (f_k(x, b) - b)| \leq L_k |a - b|$  therefore implies that for all  $k \in \{1, 2, \dots, K\}$ ,  $n \in \mathbb{N}$ ,  $x \in \mathcal{O}$  it holds that

$$\begin{aligned}
\text{Var}[V_{k, n}^0(x)] &\leq \frac{1}{M^n} \mathbb{E} \left[ |g(X_0^{0,k,x})|^2 + \left| \sum_{l=0}^{k-1} f_l(X_l^{0,k,x}, 0) \right|^2 \right] \\
&+ \sum_{j=1}^{n-1} \frac{1}{M^{n-j}} \sum_{l=0}^{k-1} \frac{L_l^2}{\mathfrak{p}_{k, l}} \mathbb{E} \left[ |V_{l, j}^0(X_l^{0,k,x}) - V_{l, j-1}^1(X_l^{0,k,x})|^2 \right].
\end{aligned} \tag{111}$$

Hölder's inequality hence ensures for all  $k \in \{1, 2, \dots, K\}$ ,  $n \in \mathbb{N}$ ,  $x \in \mathcal{O}$  that

$$\begin{aligned}
\text{Var}[V_{k, n}^0(x)] &\leq \frac{1}{M^n} \mathbb{E} \left[ |g(X_0^{0,k,x})|^2 + \left| \sum_{l=0}^{k-1} f_l(X_l^{0,k,x}, 0) \right|^2 \right] \\
&+ \sum_{j=1}^{n-1} \frac{1}{M^{n-j}} \left[ \max_{l \in [0, k) \cap \mathbb{N}_0} \frac{L_l}{\mathfrak{p}_{k, l}} \right] \left( \sum_{l=0}^{k-1} L_l \mathbb{E} \left[ |V_{l, j}^0(X_l^{0,k,x}) - V_{l, j-1}^1(X_l^{0,k,x})|^2 \right] \right).
\end{aligned} \tag{112}$$

This establishes (105). This completes the proof of Lemma 3.9.  $\square$

### 3.3 Recursive error estimates for MLP approximations

**Lemma 3.10.** Assume Setting 2.1, let  $c \in [0, \infty)$ , let  $\alpha_k, \beta_k, L_k \in [0, \infty)$ ,  $k \in \{0, 1, \dots, K\}$ , assume for all  $k \in \{0, 1, \dots, K\}$ ,  $x \in \mathcal{O}$  that  $\mathbb{E}[|g(X_0^{0,k,x})|^2 + \sum_{l=0}^{k-1} |f_l(X_l^{0,k,x}, 0)|^2] < \infty$ , assume for all  $k \in \{0, 1, \dots, K-1\}$ ,  $x \in \mathcal{O}$ ,  $a, b \in \mathbb{R}$  that  $|(f_k(x, a) - a) - (f_k(x, b) - b)| \leq L_k |a - b|$ , assume for all  $k, l \in \mathbb{N}_0$  with  $l < k \leq K$  that  $L_l \leq c \mathbb{P}(\mathcal{R}_k^0 = l)$ , assume for all  $l \in \{0, 1, \dots, K-1\}$

that  $L_l \sum_{k=l+1}^K \alpha_k \leq \beta_l \sum_{j=0}^{K-1} L_j$ , and let  $v_k: \mathcal{O} \rightarrow \mathbb{R}$ ,  $k \in \{0, 1, \dots, K\}$ , be  $\mathcal{B}(\mathcal{O})/\mathcal{B}(\mathbb{R})$ -measurable functions which satisfy for all  $k \in \{1, 2, \dots, K\}$ ,  $x \in \mathcal{O}$  that  $v_0(x) = g(x)$  and

$$v_k(x) = \mathbb{E}[f_{k-1}(X_{k-1}^{0,k,x}, v_{k-1}(X_{k-1}^{0,k,x}))] \quad (113)$$

(cf. Lemma 3.2). Then it holds for all  $n \in \mathbb{N}$ ,  $x \in \mathcal{O}$  that

$$\begin{aligned} & \left( \sum_{k=0}^K \alpha_k \mathbb{E} \left[ |V_{k,n}^0(X_k^{0,K,x}) - v_k(X_k^{0,K,x})|^2 \right] \right)^{1/2} \\ & \leq \frac{1}{\sqrt{M^n}} \left[ \sum_{k=0}^K \alpha_k \right]^{1/2} \left[ \left( \mathbb{E} \left[ |g(X_0^{0,K,x})|^2 \right] \right)^{1/2} + \sum_{l=0}^{K-1} \left( \mathbb{E} \left[ |f_l(X_l^{0,K,x}, 0)|^2 \right] \right)^{1/2} \right] \\ & + 2 \left[ \max_{k,l \in [0,K] \cap \mathbb{N}_0, k > l} \left( \frac{L_l}{\mathfrak{p}_{k,l}} \right) \right] \left[ \sum_{j=0}^{n-1} \frac{1}{\sqrt{M^{n-j-1}}} \left( \sum_{k=0}^{K-1} \beta_k \mathbb{E} \left[ |V_{k,j}^0(X_k^{0,K,x}) - v_k(X_k^{0,K,x})|^2 \right] \right)^{1/2} \right]. \end{aligned} \quad (114)$$

*Proof of Lemma 3.10.* First, note that Minkowski's inequality guarantees that for all  $n \in \mathbb{N}_0$ ,  $x \in \mathcal{O}$  it holds that

$$\begin{aligned} & \left( \sum_{k=0}^K \alpha_k \mathbb{E} \left[ |V_{k,n}^0 - v_k|^2(X_k^{0,K,x}) \right] \right)^{1/2} \\ & \leq \left( \sum_{k=0}^K \alpha_k \mathbb{E} \left[ |V_{k,n}^0 - \mathbb{E}[V_{k,n}^0]|^2(X_k^{0,K,x}) \right] \right)^{1/2} + \left( \sum_{k=0}^K \alpha_k \mathbb{E} \left[ |\mathbb{E}[V_{k,n}^0] - v_k|^2(X_k^{0,K,x}) \right] \right)^{1/2}. \end{aligned} \quad (115)$$

Moreover, note that Lemma 3.8 ensures that for all  $k \in \{1, 2, \dots, K\}$ ,  $n \in \mathbb{N}$ ,  $x \in \mathcal{O}$  it holds that

$$|\mathbb{E}[V_{k,n}^0(x)] - v_k(x)|^2 \leq \left( \sum_{l=0}^{k-1} L_l \right) \left( \sum_{l=0}^{k-1} L_l \mathbb{E} \left[ |V_{l,n-1}^0(X_l^{0,k,x}) - v_l(X_l^{0,k,x})|^2 \right] \right). \quad (116)$$

In addition, observe that item (ii) of Lemma 2.2 and item (i) of Lemma 2.3 ensure that for all  $k, l, n \in \mathbb{N}_0$  with  $l \leq k \leq K$  it holds that  $\mathcal{O} \times \Omega \ni (x, \omega) \mapsto V_{l,n}^0(X_l^{0,k,x}(\omega), \omega) \in \mathbb{R}$  is  $(\mathcal{B}(\mathcal{O}) \otimes \sigma((\mathcal{R}_s^{(0,\vartheta)})_{(s,\vartheta) \in \{1,2,\dots,K\} \times \Theta}, (W_s^{(0,\vartheta)})_{(s,\vartheta) \in \{0,1,\dots,K-1\} \times \Theta}, (W_s^0)_{s \in [l,k] \cap \mathbb{N}_0})) / \mathcal{B}(\mathbb{R})$ -measurable. Combining this and the fact that for all  $k, l \in \mathbb{N}_0$  with  $l \leq k \leq K$  it holds that  $\sigma((W_s^0)_{s \in [k,K-1] \cap \mathbb{N}_0})$  and  $\sigma((\mathcal{R}_s^{(0,\vartheta)})_{(s,\vartheta) \in \{1,2,\dots,K\} \times \Theta}, (W_s^0)_{s \in [l,k] \cap \mathbb{N}_0}, (W_s^{(0,\vartheta)})_{(s,\vartheta) \in \{0,1,\dots,K-1\} \times \Theta})$  are independent sigma-algebras on  $\Omega$  with Lemma 3.8 and Hutzenthaler et al. [36, Lemma 2.2] ensures that for all  $k \in \{1, 2, \dots, K\}$ ,  $n \in \mathbb{N}$ ,  $x \in \mathcal{O}$  it holds that

$$\begin{aligned} & \mathbb{E} \left[ |\mathbb{E}[V_{k,n}^0] - v_k|^2(X_k^{0,K,x}) \right] = \int_{\mathcal{O}} |\mathbb{E}[V_{k,n}^0(y)] - v_k(y)|^2 \left[ (X_k^{0,K,x}(\mathbb{P}))(dy) \right] \\ & \leq \int_{\mathcal{O}} \left( \sum_{l=0}^{k-1} L_l \right) \left( \sum_{l=0}^{k-1} L_l \mathbb{E} \left[ |V_{l,n-1}^0(X_l^{0,k,y}) - v_l(X_l^{0,k,y})|^2 \right] \right) \left[ (X_k^{0,K,x}(\mathbb{P}))(dy) \right] \\ & = \left( \sum_{l=0}^{k-1} L_l \right) \left( \sum_{l=0}^{k-1} L_l \int_{\mathcal{O}} \mathbb{E} \left[ |V_{l,n-1}^0(X_l^{0,k,y}) - v_l(X_l^{0,k,y})|^2 \right] \left[ (X_k^{0,K,x}(\mathbb{P}))(dy) \right] \right) \\ & = \left( \sum_{l=0}^{k-1} L_l \right) \left( \sum_{l=0}^{k-1} L_l \mathbb{E} \left[ |V_{l,n-1}^0(X_l^{0,k,X_k^{0,K,x}}) - v_l(X_l^{0,k,X_k^{0,K,x}})|^2 \right] \right) \\ & = \left( \sum_{l=0}^{k-1} L_l \right) \left( \sum_{l=0}^{k-1} L_l \mathbb{E} \left[ |V_{l,n-1}^0(X_l^{0,K,x}) - v_l(X_l^{0,K,x})|^2 \right] \right). \end{aligned} \quad (117)$$

This and the fact that for all  $n \in \mathbb{N}$ ,  $x \in \mathcal{O}$  it holds that  $V_{0,n}^0(x) = g(x)$  demonstrate that for all  $n \in \mathbb{N}$ ,  $x \in \mathcal{O}$  it holds that

$$\begin{aligned}
& \sum_{k=0}^K \alpha_k \mathbb{E} \left[ \left| \mathbb{E}[V_{k,n}^0] - v_k \right|^2 (X_k^{0,K,x}) \right] \\
& \leq \sum_{k=1}^K \alpha_k \left( \sum_{l=0}^{k-1} L_l \right) \left( \sum_{l=0}^{k-1} L_l \mathbb{E} \left[ \left| V_{l,n-1}^0(X_l^{0,K,x}) - v_l(X_l^{0,K,x}) \right|^2 \right] \right) \\
& = \sum_{l=0}^{K-1} \sum_{k=l+1}^K \alpha_k \left( \sum_{j=0}^{k-1} L_j \right) L_l \mathbb{E} \left[ \left| V_{l,n-1}^0(X_l^{0,K,x}) - v_l(X_l^{0,K,x}) \right|^2 \right] \\
& \leq \left( \sum_{j=0}^{K-1} L_j \right) \sum_{l=0}^{K-1} L_l \left( \sum_{k=l+1}^K \alpha_k \right) \mathbb{E} \left[ \left| V_{l,n-1}^0(X_l^{0,K,x}) - v_l(X_l^{0,K,x}) \right|^2 \right].
\end{aligned} \tag{118}$$

Next note that the fact that for all  $l \in \{0, 1, \dots, K-1\}$  with  $\mathbb{P}(\mathcal{R}_K^0 = l) > 0$  it holds that  $\mathbb{P}(\mathcal{R}_K^0 = l) = \mathbf{p}_{K,l}$ , the fact that for all  $l \in \{0, 1, \dots, K-1\}$  with  $\mathbb{P}(\mathcal{R}_K^0 = l) = 0$  it holds that  $L_l = 0$ , and the fact that  $\sum_{l=0}^{K-1} \mathbb{P}(\mathcal{R}_K^0 = l) = 1$  ensure that

$$\begin{aligned}
\sum_{l=0}^{K-1} L_l &= \sum_{l=0}^{K-1} L_l \mathbb{1}_{(0,\infty)}(\mathbb{P}(\mathcal{R}_K^0 = l)) \\
&= \sum_{l=0}^{K-1} \frac{L_l}{\mathbf{p}_{K,l}} \mathbf{p}_{K,l} \mathbb{1}_{(0,\infty)}(\mathbb{P}(\mathcal{R}_K^0 = l)) = \sum_{l=0}^{K-1} \frac{L_l}{\mathbf{p}_{K,l}} \mathbb{P}(\mathcal{R}_K^0 = l) \mathbb{1}_{(0,\infty)}(\mathbb{P}(\mathcal{R}_K^0 = l)) \\
&\leq \left[ \max_{l \in \{0,1,\dots,K-1\}} \frac{L_l}{\mathbf{p}_{K,l}} \right] \left[ \sum_{l=0}^{K-1} \mathbb{P}(\mathcal{R}_K^0 = l) \mathbb{1}_{(0,\infty)}(\mathbb{P}(\mathcal{R}_K^0 = l)) \right] \\
&= \left[ \max_{l \in \{0,1,\dots,K-1\}} \frac{L_l}{\mathbf{p}_{K,l}} \right] \left[ \sum_{l=0}^{K-1} \mathbb{P}(\mathcal{R}_K^0 = l) \right] = \left[ \max_{l \in \{0,1,\dots,K-1\}} \frac{L_l}{\mathbf{p}_{K,l}} \right].
\end{aligned} \tag{119}$$

The assumption that for all  $l \in \{0, 1, \dots, K-1\}$  it holds that  $L_l \sum_{k=l+1}^K \alpha_k \leq \beta_l \sum_{j=0}^{K-1} L_j$  and (118) therefore ensure that for all  $n \in \mathbb{N}$ ,  $x \in \mathcal{O}$  it holds that

$$\begin{aligned}
& \sum_{k=0}^K \alpha_k \mathbb{E} \left[ \left| \mathbb{E}[V_{k,n}^0] - v_k \right|^2 (X_k^{0,K,x}) \right] \\
& \leq \left[ \sum_{j=0}^{K-1} L_j \right]^2 \left( \sum_{l=0}^{K-1} \beta_l \mathbb{E} \left[ \left| V_{l,n-1}^0(X_l^{0,K,x}) - v_l(X_l^{0,K,x}) \right|^2 \right] \right) \\
& \leq \left[ \max_{j \in [0,K) \cap \mathbb{N}_0} \frac{L_j}{\mathbf{p}_{K,j}} \right]^2 \left( \sum_{l=0}^{K-1} \beta_l \mathbb{E} \left[ \left| V_{l,n-1}^0(X_l^{0,K,x}) - v_l(X_l^{0,K,x}) \right|^2 \right] \right).
\end{aligned} \tag{120}$$

In the next step we note that item (i) of Lemma 2.3 ensures that for all  $n \in \mathbb{N}_0$ ,  $k \in \{0, 1, \dots, K\}$  it holds that

$$\mathcal{O} \times \Omega \ni (x, \omega) \mapsto \left| V_{k,n}^0(x, \omega) - \int_{\Omega} V_{k,n}^0(x, \xi) \mathbb{P}(d\xi) \right|^2 \in \mathbb{R} \tag{121}$$

is  $(\mathcal{B}(\mathcal{O}) \otimes \sigma((\mathcal{R}_s^{(0,\vartheta)})_{(s,\vartheta) \in \{1,2,\dots,K\} \times \Theta}, (W_s^{(0,\vartheta)})_{(s,\vartheta) \in \{0,1,\dots,K-1\} \times \Theta})) / \mathcal{B}(\mathbb{R})$ -measurable. The fact that  $\sigma((\mathcal{R}_s^{(0,\vartheta)})_{(s,\vartheta) \in \{1,2,\dots,K\} \times \Theta}, (W_s^{(0,\vartheta)})_{(s,\vartheta) \in \{0,1,\dots,K-1\} \times \Theta})$  and  $\sigma((W_s^0)_{s \in \{0,1,\dots,K-1\}})$  are independent, item

(i) of Lemma 2.2, and Hutzenthaler et al. [36, Lemma 2.2] hence show that for all  $k \in \{0, 1, \dots, K\}$ ,  $n \in \mathbb{N}$ ,  $x \in \mathcal{O}$  it holds that

$$\mathbb{E}\left[|V_{k,n}^0 - \mathbb{E}[V_{k,n}^0]|^2(X_k^{0,K,x})\right] = \int_{\mathcal{O}} \mathbb{E}\left[|V_{k,n}^0(y) - \mathbb{E}[V_{k,n}^0(y)]|^2\right] (X_k^{0,K,x}(\mathbb{P}))(dy). \quad (122)$$

Lemma 3.9 hence guarantees for all  $k \in \{1, 2, \dots, K\}$ ,  $n \in \mathbb{N}$ ,  $x \in \mathcal{O}$  that

$$\begin{aligned} & \mathbb{E}\left[|V_{k,n}^0 - \mathbb{E}[V_{k,n}^0]|^2(X_k^{0,K,x})\right] \\ & \leq \int_{\mathcal{O}} \frac{1}{M^n} \mathbb{E}\left[|g(X_0^{0,k,y})|^2 + \left|\sum_{l=0}^{k-1} f_l(X_l^{0,k,y}, 0)\right|^2\right] (X_k^{0,K,x}(\mathbb{P}))(dy) \\ & + \int_{\mathcal{O}} \sum_{j=1}^{n-1} \frac{\max_{l \in \{0,1,\dots,k-1\}} \left(\frac{L_l}{p_{k,l}}\right)}{M^{n-j}} \left(\sum_{l=0}^{k-1} L_l \mathbb{E}\left[|V_{l,j}^0(X_l^{0,k,y}) - V_{l,j-1}^1(X_l^{0,k,y})|^2\right]\right) (X_k^{0,K,x}(\mathbb{P}))(dy). \end{aligned} \quad (123)$$

Combining this with item (i) of Lemma 2.2 and Hutzenthaler et al. [36, Lemma 2.2] yields that for all  $k \in \{1, 2, \dots, K\}$ ,  $n \in \mathbb{N}$ ,  $x \in \mathcal{O}$  it holds that

$$\begin{aligned} & \mathbb{E}\left[|V_{k,n}^0 - \mathbb{E}[V_{k,n}^0]|^2(X_k^{0,K,x})\right] \\ & \leq \frac{1}{M^n} \mathbb{E}\left[|g(X_0^{0,k,X_k^{0,K,x}})|^2 + \left|\sum_{l=0}^{k-1} f_l(X_l^{0,k,X_k^{0,K,x}}, 0)\right|^2\right] \\ & + \int_{\mathcal{O}} \sum_{j=1}^{n-1} \frac{\max_{l \in \{0,1,\dots,k-1\}} \left(\frac{L_l}{p_{k,l}}\right)}{M^{n-j}} \left(\sum_{l=0}^{k-1} L_l \mathbb{E}\left[|V_{l,j}^0(X_l^{0,k,y}) - V_{l,j-1}^1(X_l^{0,k,y})|^2\right]\right) (X_k^{0,K,x}(\mathbb{P}))(dy). \end{aligned} \quad (124)$$

Next observe that item (ii) of Lemma 2.2 and item (i) of Lemma 2.3 show that for all  $k, l \in \mathbb{N}_0$ ,  $n \in \mathbb{N}$  with  $l \leq k \leq K$  it holds that  $\mathcal{O} \times \Omega \ni (x, \omega) \mapsto |V_{l,n}^0(X_l^{0,k,x}(\omega), \omega) - V_{l,n-1}^1(X_l^{0,k,x}(\omega), \omega)|^2 \in [0, \infty)$  is  $(\mathcal{B}(\mathcal{O}) \otimes \sigma((\mathcal{R}_s^{(0,\eta)})_{(s,\eta) \in \{1,2,\dots,K\} \times \Theta}, (\mathcal{R}_s^{(1,\eta)})_{(s,\eta) \in \{1,2,\dots,K\} \times \Theta}, (W_s^{(0,\eta)})_{(s,\eta) \in \{0,1,\dots,K-1\} \times \Theta}, (W_s^0)_{s \in [l,k) \cap \mathbb{N}_0}, (W_s^{(1,\eta)})_{(s,\eta) \in \{0,1,\dots,K-1\} \times \Theta})) / \mathcal{B}([0, \infty))$ -measurable. This, the fact that for all  $k, l \in \mathbb{N}_0$  with  $l \leq k \leq K$  it holds that  $\sigma((W_s^0)_{s \in [k,K) \cap \mathbb{N}_0})$  and  $\sigma((\mathcal{R}_s^{(0,\eta)})_{(s,\eta) \in \{1,2,\dots,K\} \times \Theta}, (\mathcal{R}_s^{(1,\eta)})_{(s,\eta) \in \{1,2,\dots,K\} \times \Theta}, (W_s^0)_{s \in [l,k) \cap \mathbb{N}_0}, (W_s^{(0,\eta)})_{(s,\eta) \in \{0,1,\dots,K-1\} \times \Theta}, (W_s^{(1,\eta)})_{(s,\eta) \in \{0,1,\dots,K-1\} \times \Theta})$  are independent, and Hutzenthaler et al. [36, Lemma 2.2] demonstrate that for all  $k \in \{1, 2, \dots, K\}$ ,  $n \in \mathbb{N}$ ,  $x \in \mathcal{O}$  it holds that

$$\begin{aligned} & \mathbb{E}\left[|V_{k,n}^0 - \mathbb{E}[V_{k,n}^0]|^2(X_k^{0,K,x})\right] \\ & \leq \frac{1}{M^n} \mathbb{E}\left[|g(X_0^{0,k,X_k^{0,K,x}})|^2 + \left|\sum_{l=0}^{k-1} f_l(X_l^{0,k,X_k^{0,K,x}}, 0)\right|^2\right] \\ & + \sum_{j=1}^{n-1} \frac{1}{M^{n-j}} \left[\max_{l \in \{0,1,\dots,k-1\}} \frac{L_l}{p_{k,l}}\right] \left(\sum_{l=0}^{k-1} L_l \mathbb{E}\left[|V_{l,j}^0(X_l^{0,k,X_k^{0,K,x}}) - V_{l,j-1}^1(X_l^{0,k,X_k^{0,K,x}})|^2\right]\right). \end{aligned} \quad (125)$$

This and (3) ensure for all  $k \in \{1, 2, \dots, K\}$ ,  $n \in \mathbb{N}$ ,  $x \in \mathcal{O}$  that

$$\begin{aligned} & \mathbb{E}\left[|V_{k,n}^0 - \mathbb{E}[V_{k,n}^0]|^2(X_k^{0,K,x})\right] \leq \frac{1}{M^n} \mathbb{E}\left[|g(X_0^{0,K,x})|^2 + \left|\sum_{l=0}^{k-1} f_l(X_l^{0,K,x}, 0)\right|^2\right] \\ & + \sum_{j=1}^{n-1} \frac{1}{M^{n-j}} \left[\max_{l \in \{0,1,\dots,k-1\}} \frac{L_l}{p_{k,l}}\right] \left(\sum_{l=0}^{k-1} L_l \mathbb{E}\left[|V_{l,j}^0(X_l^{0,K,x}) - V_{l,j-1}^1(X_l^{0,K,x})|^2\right]\right). \end{aligned} \quad (126)$$

The fact that for all  $n \in \mathbb{N}$ ,  $x \in \mathcal{O}$  it holds that  $V_{0,n}^0(x) = g(x)$  hence guarantees that for all  $n \in \mathbb{N}$ ,  $x \in \mathcal{O}$  it holds that

$$\begin{aligned}
& \sum_{k=0}^K \alpha_k \mathbb{E} \left[ |V_{k,n}^0 - \mathbb{E}[V_{k,n}^0]|^2 (X_k^{0,K,x}) \right] \\
& \leq \frac{1}{M^n} \left[ \sum_{k=0}^K \alpha_k \right] \mathbb{E} \left[ |g(X_0^{0,K,x})|^2 + \left( \sum_{l=0}^{K-1} |f_l(X_l^{0,K,x}, 0)| \right)^2 \right] \\
& + \sum_{k=1}^K \sum_{j=1}^{n-1} \frac{\alpha_k}{M^{n-j}} \left[ \max_{l \in \{0,1,\dots,k-1\}} \frac{L_l}{\mathfrak{p}_{k,l}} \right] \left( \sum_{l=0}^{k-1} L_l \mathbb{E} \left[ |V_{l,j}^0(X_l^{0,K,x}) - V_{l,j-1}^1(X_l^{0,K,x})|^2 \right] \right).
\end{aligned} \tag{127}$$

This yields for all  $n \in \mathbb{N}$ ,  $x \in \mathcal{O}$  that

$$\begin{aligned}
& \sum_{k=0}^K \alpha_k \mathbb{E} \left[ |V_{k,n}^0 - \mathbb{E}[V_{k,n}^0]|^2 (X_k^{0,K,x}) \right] \\
& \leq \frac{1}{M^n} \left[ \sum_{k=0}^K \alpha_k \right] \mathbb{E} \left[ |g(X_0^{0,K,x})|^2 + \left( \sum_{l=0}^{K-1} |f_l(X_l^{0,K,x}, 0)| \right)^2 \right] \\
& + \left[ \max_{k,l \in [0,K] \cap \mathbb{N}_0, k>l} \frac{L_l}{\mathfrak{p}_{k,l}} \right] \sum_{j=1}^{n-1} \frac{1}{M^{n-j}} \sum_{l=0}^{K-1} \left[ L_l \left( \sum_{k=l+1}^K \alpha_k \right) \right] \mathbb{E} \left[ |V_{l,j}^0(X_l^{0,K,x}) - V_{l,j-1}^1(X_l^{0,K,x})|^2 \right].
\end{aligned} \tag{128}$$

The assumption that for all  $l \in \{0, 1, \dots, K-1\}$  it holds that  $L_l \sum_{k=l+1}^K \alpha_k \leq \beta_l \sum_{j=0}^{K-1} L_j$  therefore demonstrates that for all  $n \in \mathbb{N}$ ,  $x \in \mathcal{O}$  it holds that

$$\begin{aligned}
& \sum_{k=0}^K \alpha_k \mathbb{E} \left[ |V_{k,n}^0 - \mathbb{E}[V_{k,n}^0]|^2 (X_k^{0,K,x}) \right] \\
& \leq \frac{1}{M^n} \left[ \sum_{k=0}^K \alpha_k \right] \mathbb{E} \left[ |g(X_0^{0,K,x})|^2 + \left( \sum_{l=0}^{K-1} |f_l(X_l^{0,K,x}, 0)| \right)^2 \right] \\
& + \left[ \max_{k,l \in [0,K] \cap \mathbb{N}_0, k>l} \frac{L_l}{\mathfrak{p}_{k,l}} \right] \left[ \sum_{l=0}^{K-1} L_l \right] \left[ \sum_{j=1}^{n-1} \frac{1}{M^{n-j}} \left( \sum_{l=0}^{K-1} \beta_l \mathbb{E} \left[ |V_{l,j}^0(X_l^{0,K,x}) - V_{l,j-1}^1(X_l^{0,K,x})|^2 \right] \right) \right].
\end{aligned} \tag{129}$$

This and (119) ensure that for all  $n \in \mathbb{N}$ ,  $x \in \mathcal{O}$  it holds that

$$\begin{aligned}
& \sum_{k=0}^K \alpha_k \mathbb{E} \left[ |V_{k,n}^0 - \mathbb{E}[V_{k,n}^0]|^2 (X_k^{0,K,x}) \right] \\
& \leq \frac{1}{M^n} \left[ \sum_{k=0}^K \alpha_k \right] \mathbb{E} \left[ |g(X_0^{0,K,x})|^2 + \left( \sum_{l=0}^{K-1} |f_l(X_l^{0,K,x}, 0)| \right)^2 \right] \\
& + \left[ \max_{k,l \in [0,K] \cap \mathbb{N}_0, k>l} \left( \frac{L_l}{\mathfrak{p}_{k,l}} \right) \right]^2 \left[ \sum_{j=1}^{n-1} \frac{1}{M^{n-j}} \left( \sum_{l=0}^{K-1} \beta_l \mathbb{E} \left[ |V_{l,j}^0(X_l^{0,K,x}) - V_{l,j-1}^1(X_l^{0,K,x})|^2 \right] \right) \right].
\end{aligned} \tag{130}$$

The fact that for all  $m \in \mathbb{N}$ ,  $a_1, a_2, \dots, a_m \in [0, \infty)$  it holds that  $\sqrt{\sum_{i=1}^m a_i} \leq \sum_{i=1}^m \sqrt{a_i}$ , and

Minkowski's inequality therefore prove that for all  $n \in \mathbb{N}$ ,  $x \in \mathcal{O}$  it holds that

$$\begin{aligned}
& \left( \sum_{k=0}^K \alpha_k \mathbb{E} \left[ |V_{k,n}^0 - \mathbb{E}[V_{k,n}^0]|^2 (X_k^{0,K,x}) \right] \right)^{1/2} \\
& \leq \frac{1}{\sqrt{M^n}} \left[ \sum_{k=0}^K \alpha_k \right]^{1/2} \left[ \left( \mathbb{E} \left[ |g(X_0^{0,K,x})|^2 \right] \right)^{1/2} + \sum_{l=0}^{K-1} \left( \mathbb{E} \left[ |f_l(X_l^{0,K,x}, 0)|^2 \right] \right)^{1/2} \right] \\
& + \left[ \max_{k,l \in [0,K] \cap \mathbb{N}_0, k>l} \left( \frac{L_l}{\mathfrak{p}_{k,l}} \right) \right] \left[ \sum_{j=1}^{n-1} \frac{1}{\sqrt{M^{n-j}}} \left( \sum_{l=0}^{K-1} \beta_l \mathbb{E} \left[ |V_{l,j}^0(X_l^{0,K,x}) - v_l(X_l^{0,K,x})|^2 \right] \right)^{1/2} \right] \\
& + \left[ \max_{k,l \in [0,K] \cap \mathbb{N}_0, k>l} \left( \frac{L_l}{\mathfrak{p}_{k,l}} \right) \right] \left[ \sum_{j=1}^{n-1} \frac{1}{\sqrt{M^{n-j}}} \left( \sum_{l=0}^{K-1} \beta_l \mathbb{E} \left[ |V_{l,j-1}^1(X_l^{0,K,x}) - v_l(X_l^{0,K,x})|^2 \right] \right)^{1/2} \right]. \tag{131}
\end{aligned}$$

Item (i) of Lemma 2.2, item (i) of Lemma 2.3, the assumption that  $\sigma((W_s^{(1,\eta)})_{(s,\eta) \in \{0,1,\dots,K-1\} \times \Theta}, (\mathcal{R}_s^{(1,\eta)})_{(s,\eta) \in \{1,2,\dots,K\} \times \Theta})$  and  $\sigma((W_s^0)_{s \in \{0,1,\dots,K-1\}})$  are independent, item (i) of Lemma 2.8, item (i) of Proposition 2.9, the assumption that  $\sigma((W_s^{(0,\eta)})_{(s,\eta) \in \{0,1,\dots,K-1\} \times \Theta}, (\mathcal{R}_s^{(0,\eta)})_{(s,\eta) \in \{1,2,\dots,K\} \times \Theta})$  and  $\sigma((W_s^0)_{s \in \{0,1,\dots,K-1\}})$  are independent, and Hutzenthaler et al. [36, Lemma 2.2] hence yield that for all  $n \in \mathbb{N}$ ,  $x \in \mathcal{O}$  it holds that

$$\begin{aligned}
& \left( \sum_{k=0}^K \alpha_k \mathbb{E} \left[ |V_{k,n}^0 - \mathbb{E}[V_{k,n}^0]|^2 (X_k^{0,K,x}) \right] \right)^{1/2} \\
& \leq \frac{1}{\sqrt{M^n}} \left[ \sum_{k=0}^K \alpha_k \right]^{1/2} \left[ \left( \mathbb{E} \left[ |g(X_0^{0,K,x})|^2 \right] \right)^{1/2} + \sum_{l=0}^{K-1} \left( \mathbb{E} \left[ |f_l(X_l^{0,K,x}, 0)|^2 \right] \right)^{1/2} \right] \\
& + \left[ \max_{k,l \in [0,K] \cap \mathbb{N}_0, k>l} \left( \frac{L_l}{\mathfrak{p}_{k,l}} \right) \right] \left[ \sum_{j=1}^{n-1} \frac{1}{\sqrt{M^{n-j}}} \left( \sum_{l=0}^{K-1} \beta_l \mathbb{E} \left[ |V_{l,j}^0(X_l^{0,K,x}) - v_l(X_l^{0,K,x})|^2 \right] \right)^{1/2} \right] \\
& + \left[ \max_{k,l \in [0,K] \cap \mathbb{N}_0, k>l} \left( \frac{L_l}{\mathfrak{p}_{k,l}} \right) \right] \left[ \sum_{j=1}^{n-1} \frac{1}{\sqrt{M^{n-j}}} \left( \sum_{l=0}^{K-1} \beta_l \mathbb{E} \left[ |V_{l,j-1}^0(X_l^{0,K,x}) - v_l(X_l^{0,K,x})|^2 \right] \right)^{1/2} \right]. \tag{132}
\end{aligned}$$

This, (115), and (120) therefore demonstrate that for all  $n \in \mathbb{N}$ ,  $x \in \mathcal{O}$  it holds that

$$\begin{aligned}
& \left( \sum_{k=0}^K \alpha_k \mathbb{E} \left[ |V_{k,n}^0 - v_k|^2 (X_k^{0,K,x}) \right] \right)^{1/2} \\
& \leq \frac{1}{\sqrt{M^n}} \left[ \sum_{k=0}^K \alpha_k \right]^{1/2} \left[ \left( \mathbb{E} \left[ |g(X_0^{0,K,x})|^2 \right] \right)^{1/2} + \sum_{l=0}^{K-1} \left( \mathbb{E} \left[ |f_l(X_l^{0,K,x}, 0)|^2 \right] \right)^{1/2} \right] \\
& + 2 \left[ \max_{k,l \in [0,K] \cap \mathbb{N}_0, k>l} \left( \frac{L_l}{\mathfrak{p}_{k,l}} \right) \right] \left[ \sum_{j=0}^{n-1} \frac{1}{\sqrt{M^{n-j-1}}} \left( \sum_{l=0}^{K-1} \beta_l \mathbb{E} \left[ |V_{l,j}^0(X_l^{0,K,x}) - v_l(X_l^{0,K,x})|^2 \right] \right)^{1/2} \right]. \tag{133}
\end{aligned}$$

This establishes (114). This completes the proof of Lemma 3.10.  $\square$

### 3.4 Full error analysis for MLP approximations

**Lemma 3.11.** *Let  $K \in \mathbb{N}$ ,  $L_0, L_1, \dots, L_{K-1} \in [0, \infty)$  satisfy  $\sum_{j=0}^{K-1} L_j > 0$ , let  $\alpha_k^{(q)} \in [0, \infty)$ ,  $q \in \mathbb{N}_0$ ,  $k \in \{0, 1, \dots, K\}$ , and assume for all  $k \in \{0, 1, \dots, K-1\}$ ,  $q \in \mathbb{N}$  that*

$$\alpha_K^{(0)} = 1, \quad \alpha_k^{(0)} = 0, \quad \text{and} \quad \alpha_k^{(q)} = \frac{\left[\sum_{j=k}^{K-1} L_j\right]^q - \left[\sum_{j=k+1}^{K-1} L_j\right]^q}{q! \left[\sum_{j=0}^{K-1} L_j\right]^q}. \quad (134)$$

Then it holds for all  $k \in \{0, 1, \dots, K-1\}$ ,  $q \in \mathbb{N}_0$  that  $\alpha_k^{(q+1)} \sum_{j=0}^{K-1} L_j \geq L_k \sum_{j=k+1}^K \alpha_j^{(q)}$ .

*Proof of Lemma 3.11.* First, note that for all  $k \in \{0, 1, \dots, K-1\}$  it holds that  $\sum_{i=k+1}^K \alpha_i^{(0)} = 1$ . This and (134) ensure for all  $k \in \{0, 1, \dots, K-1\}$  that

$$L_k \left[ \sum_{i=k+1}^K \alpha_i^{(0)} \right] = L_k = \left[ \sum_{j=k}^{K-1} L_j \right] - \left[ \sum_{j=k+1}^{K-1} L_j \right] = \alpha_k^{(1)} \left[ \sum_{j=0}^{K-1} L_j \right]. \quad (135)$$

In the next step we observe that the fact that for all  $q \in \mathbb{N}_0$ ,  $a, b \in \mathbb{R}$  it holds that  $a^{q+1} - b^{q+1} = (a-b) \sum_{i=0}^q a^i b^{q-i}$  implies that for all  $q \in \mathbb{N}_0$ ,  $a, b \in \mathbb{R}$  with  $a \geq b$  it holds that

$$a^{q+1} - b^{q+1} = (a-b) \sum_{i=0}^q a^i b^{q-i} \geq (a-b) \sum_{i=0}^q b^q = (q+1)(a-b)b^q. \quad (136)$$

This and (134) ensure that for all  $k \in \{0, 1, \dots, K-1\}$ ,  $q \in \mathbb{N}$  it holds that

$$\begin{aligned} & (q+1)! L_k \left[ \sum_{j=k+1}^K \alpha_j^{(q)} \right] \left[ \sum_{j=0}^{K-1} L_j \right]^{q+1} \\ &= (q+1) L_k \left[ \sum_{j=k+1}^{K-1} \left( \left[ \sum_{l=j}^{K-1} L_l \right]^q - \left[ \sum_{l=j+1}^{K-1} L_l \right]^q \right) \right] \left[ \sum_{j=0}^{K-1} L_j \right] \\ &= (q+1) L_k \left[ \sum_{j=k+1}^{K-1} L_j \right]^q \left[ \sum_{j=0}^{K-1} L_j \right] \leq \left( \left[ \sum_{j=k}^{K-1} L_j \right]^{q+1} - \left[ \sum_{j=k+1}^{K-1} L_j \right]^{q+1} \right) \left[ \sum_{j=0}^{K-1} L_j \right] \\ &= (q+1)! \alpha_k^{(q+1)} \left[ \sum_{j=0}^{K-1} L_j \right]^{q+2}. \end{aligned} \quad (137)$$

This and (135) establish that for all  $k \in \{0, 1, \dots, K-1\}$ ,  $q \in \mathbb{N}_0$  it holds that  $\alpha_k^{(q+1)} \sum_{j=0}^{K-1} L_j \geq L_k \sum_{j=k+1}^K \alpha_j^{(q)}$ . This completes the proof of Lemma 3.11.  $\square$

**Lemma 3.12.** *Let  $M \in \mathbb{N}$ ,  $N \in \mathbb{N}_0$ ,  $\alpha, \beta, \kappa \in [0, \infty)$ , let  $e_{n,q} \in [0, \infty)$ ,  $n, q \in \mathbb{N}_0$ , satisfy for all  $n \in \{0, 1, \dots, N\}$ ,  $q \in \mathbb{N}_0$  that*

$$e_{n,q} \leq \frac{\alpha \sqrt{\kappa^q}}{\sqrt{q!} M^n} + \beta \sum_{l=0}^{n-1} \frac{e_{l,q+1}}{\sqrt{M^{n-1-l}}}. \quad (138)$$

Then it holds that

$$e_{N,0} \leq \frac{\alpha \exp(\frac{\kappa M}{2})}{M^{N/2}} (1 + \beta)^N. \quad (139)$$

*Proof of Lemma 3.12.* Throughout this proof assume w.l.o.g. that  $N \in \mathbb{N}$  and let  $\varepsilon_n \in [0, \infty)$ ,  $n \in \{0, 1, \dots, N\}$ , satisfy for all  $n \in \{0, 1, \dots, N\}$  that

$$\varepsilon_n = \sup \left\{ \frac{e_{n,q}}{\sqrt{M^k}} : k, q \in \mathbb{N}_0, n + k + q = N \right\}. \quad (140)$$

Observe that (138) ensures that for all  $n, k, q \in \mathbb{N}_0$  with  $n + k + q = N$  it holds that

$$\begin{aligned} \frac{e_{n,q}}{\sqrt{M^k}} &\leq \frac{\alpha \sqrt{\kappa^q}}{\sqrt{q! M^{n+k}}} + \beta \sum_{l=0}^{n-1} \frac{e_{l,q+1}}{\sqrt{M^{n-1-l+k}}} = \frac{\alpha \sqrt{\kappa^q}}{\sqrt{q! M^{N-q}}} + \beta \sum_{l=0}^{n-1} \frac{e_{l,q+1}}{\sqrt{M^{N-(q+1)-l}}} \\ &\leq \frac{\alpha \sqrt{\kappa^q}}{\sqrt{q! M^{N-q}}} + \beta \sum_{l=0}^{n-1} \varepsilon_l. \end{aligned} \quad (141)$$

The fact that for all  $q \in \mathbb{N}_0$  it holds that  $\frac{\kappa^q M^q}{q!} \leq e^{\kappa M}$  hence implies that for all  $n, k, q \in \mathbb{N}_0$  with  $n + k + q = N$  it holds that

$$\frac{e_{n,q}}{\sqrt{M^k}} \leq \frac{\alpha \exp(\frac{\kappa M}{2})}{M^{N/2}} + \beta \sum_{l=0}^{n-1} \varepsilon_l. \quad (142)$$

This and (140) yield for all  $n \in \{0, 1, \dots, N\}$  that

$$\varepsilon_n = \sup \left\{ \frac{e_{n,q}}{\sqrt{M^k}} : k, q \in \mathbb{N}_0, n + k + q = N \right\} \leq \frac{\alpha \exp(\frac{\kappa M}{2})}{M^{N/2}} + \beta \sum_{l=0}^{n-1} \varepsilon_l. \quad (143)$$

Lemma 3.5 hence ensures for all  $n \in \{0, 1, \dots, N\}$  that

$$\varepsilon_n \leq \frac{\alpha \exp(\frac{\kappa M}{2})}{M^{N/2}} (1 + \beta)^n. \quad (144)$$

This establishes (139). This completes the proof of Lemma 3.12.  $\square$

**Proposition 3.13.** *Let  $d, K, M \in \mathbb{N}$ ,  $\Theta = \bigcup_{n \in \mathbb{N}} \mathbb{Z}^n$ ,  $\mathcal{O} \in \mathcal{B}(\mathbb{R}^d) \setminus \{\emptyset\}$ ,  $c, L_0, L_1, \dots, L_K \in \mathbb{R}$ , let  $f_k: \mathcal{O} \times \mathbb{R} \rightarrow \mathbb{R}$ ,  $k \in \{0, 1, \dots, K\}$ , be  $\mathcal{B}(\mathcal{O} \times \mathbb{R})/\mathcal{B}(\mathbb{R})$ -measurable, let  $g: \mathcal{O} \rightarrow \mathbb{R}$  be  $\mathcal{B}(\mathcal{O})/\mathcal{B}(\mathbb{R})$ -measurable, assume for all  $k \in \{0, 1, \dots, K\}$ ,  $x \in \mathcal{O}$ ,  $a, b \in \mathbb{R}$  that  $|(f_k(x, a) - a) - (f_k(x, b) - b)| \leq L_k |a - b|$ , let  $(S, \mathcal{S})$  be a measurable space, let  $\phi_k: \mathcal{O} \times S \rightarrow \mathcal{O}$ ,  $k \in \{0, 1, \dots, K\}$ , be  $(\mathcal{B}(\mathcal{O}) \otimes \mathcal{S})/\mathcal{B}(\mathcal{O})$ -measurable, let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a probability space, let  $W^\theta = (W_k^\theta)_{k \in \{0, 1, \dots, K\}}: \{0, 1, \dots, K\} \times \Omega \rightarrow S$ ,  $\theta \in \Theta$ , be i.i.d. stochastic processes, assume for all  $\theta \in \Theta$  that  $W_0^\theta, W_1^\theta, \dots, W_{K-1}^\theta$  are independent, let  $\mathcal{R}^\theta = (\mathcal{R}_k^\theta)_{k \in \{0, 1, \dots, K\}}: \{0, 1, \dots, K\} \times \Omega \rightarrow \mathbb{N}_0$ ,  $\theta \in \Theta$ , be i.i.d. stochastic processes, assume for every  $k \in \{1, 2, \dots, K\}$  that  $\mathcal{R}_k^\theta \leq k - 1$ , let  $\mathbf{p}_{k,l} \in (0, \infty)$ ,  $k \in \{1, 2, \dots, K\}$ ,  $l \in \mathbb{N}_0$ , satisfy for all  $k \in \{1, 2, \dots, K\}$ ,  $l \in \mathbb{N}_0$  that  $\mathbf{p}_{k,l} \mathbb{P}(\mathcal{R}_k^0 = l) = |\mathbb{P}(\mathcal{R}_k^0 = l)|^2$ , assume for all  $k, l \in \mathbb{N}_0$  with  $l < k \leq K$  that  $L_l \leq c \mathbb{P}(\mathcal{R}_k^0 = l)$ , assume that  $(\mathcal{R}^\theta)_{\theta \in \Theta}$  and  $(W_k^\theta)_{(\theta, k) \in \Theta \times \{0, 1, \dots, K\}}$  are independent, let  $X^{\theta, k} = (X_l^{\theta, k, x})_{(l, x) \in \{0, 1, \dots, k\} \times \mathcal{O}}: \{0, 1, \dots, k\} \times \mathcal{O} \times \Omega \rightarrow \mathcal{O}$ ,  $k \in \{0, 1, \dots, K\}$ ,  $\theta \in \Theta$ , satisfy for all  $\theta \in \Theta$ ,  $k \in \{0, 1, \dots, K\}$ ,  $l \in \{0, 1, \dots, k\}$ ,  $x \in \mathcal{O}$  that*

$$X_l^{\theta, k, x} = \begin{cases} x & : l = k \\ \phi_l(X_{l+1}^{\theta, k, x}, W_l^\theta) & : l < k, \end{cases} \quad (145)$$

assume for all  $k \in \{0, 1, \dots, K\}$ ,  $x \in \mathcal{O}$  that  $\mathbb{E}[|g(X_0^{0, k, x})|^2 + \sum_{l=0}^{k-1} |f_l(X_l^{0, k, x}, 0)|^2] < \infty$ , let  $V_{k,n}^\theta: \mathcal{O} \times \Omega \rightarrow \mathbb{R}$ ,  $n \in \mathbb{N}_0$ ,  $k \in \{0, 1, \dots, K\}$ ,  $\theta \in \Theta$ , satisfy for all  $\theta \in \Theta$ ,  $k \in \{0, 1, \dots, K\}$ ,  $n \in \mathbb{N}_0$ ,  $x \in \mathcal{O}$  that

$$\begin{aligned} V_{k,n}^\theta(x) &= \frac{\mathbb{1}_{\mathbb{N}}(n)}{M^n} \sum_{m=1}^{M^n} \left[ g(X_0^{(\theta, 0, -m), k, x}) + \sum_{l=0}^{k-1} f_l(X_l^{(\theta, 0, m), k, x}, 0) \right] + \sum_{j=1}^{n-1} \frac{1}{M^{n-j}} \sum_{m=1}^{M^{n-j}} \frac{\mathbb{1}_{\mathbb{N}}(k)}{\mathbf{p}_{k, \mathcal{R}_k^{(\theta, j, m)}}} \\ &\quad \cdot \left[ \left( f_{\mathcal{R}_k^{(\theta, j, m)}}(X_{\mathcal{R}_k^{(\theta, j, m)}}^{(\theta, j, m), k, x}, V_{\mathcal{R}_k^{(\theta, j, m)}, j}^{(\theta, j, m)}(X_{\mathcal{R}_k^{(\theta, j, m)}}^{(\theta, j, m), k, x})) - V_{\mathcal{R}_k^{(\theta, j, m)}, j}^{(\theta, j, m)}(X_{\mathcal{R}_k^{(\theta, j, m)}}^{(\theta, j, m), k, x}) \right) \right. \\ &\quad \left. - \left( f_{\mathcal{R}_k^{(\theta, j, m)}}(X_{\mathcal{R}_k^{(\theta, j, m)}}^{(\theta, j, m), k, x}, V_{\mathcal{R}_k^{(\theta, j, m)}, j-1}^{(\theta, j, -m)}(X_{\mathcal{R}_k^{(\theta, j, m)}}^{(\theta, j, m), k, x})) - V_{\mathcal{R}_k^{(\theta, j, m)}, j-1}^{(\theta, j, -m)}(X_{\mathcal{R}_k^{(\theta, j, m)}}^{(\theta, j, m), k, x}) \right) \right], \end{aligned} \quad (146)$$

and let  $v_0, v_1, \dots, v_K: \mathcal{O} \rightarrow \mathbb{R}$  be  $\mathcal{B}(\mathcal{O})/\mathcal{B}(\mathbb{R})$ -measurable functions which satisfy for all  $k \in \{1, 2, \dots, K\}$ ,  $x \in \mathcal{O}$  that  $v_0(x) = g(x)$  and

$$v_k(x) = \mathbb{E}[f_{k-1}(X_{k-1}^{0,k,x}, v_{k-1}(X_{k-1}^{0,k,x}))] = \mathbb{E}[f_{k-1}(\phi_{k-1}(x, W_{k-1}^0), v_{k-1}(\phi_{k-1}(x, W_{k-1}^0)))] \quad (147)$$

(cf. Lemma 3.2). Then it holds for all  $N \in \mathbb{N}_0$ ,  $x \in \mathcal{O}$  that

$$\begin{aligned} (\mathbb{E}[|V_{K,N}^0(x) - v_K(x)|^2])^{1/2} &\leq \frac{\exp(\frac{M}{2} + \sum_{j=0}^{K-1} L_j)}{M^{N/2}} \left[ 1 + 2 \max_{k,l \in \mathbb{N}_0, l < k \leq K} \left( \frac{L_l}{\mathfrak{p}_{k,l}} \right) \right]^N \\ &\quad \cdot \left[ \left( \mathbb{E}[|g(X_0^{0,K,x})|^2] \right)^{1/2} + \sum_{l=0}^{K-1} \left( \mathbb{E}[|f_l(X_l^{0,K,x}, 0)|^2] \right)^{1/2} \right]. \end{aligned} \quad (148)$$

*Proof of Proposition 3.13.* We first prove (148) in the case that  $\sum_{k=0}^{K-1} L_k > 0$ . For this let  $x \in \mathcal{O}$ ,  $N \in \mathbb{N}_0$ , let  $e_{n,q} \in [0, \infty)$ ,  $n, q \in \mathbb{N}_0$ , let  $\alpha_k^{(q)} \in [0, \infty)$ ,  $q \in \mathbb{N}_0$ ,  $k \in \{0, 1, \dots, K\}$ , satisfy for all  $k \in \{0, 1, \dots, K-1\}$ ,  $q \in \mathbb{N}$  that

$$\alpha_K^{(0)} = 1, \quad \alpha_k^{(0)} = 0 = \alpha_K^{(q)}, \quad \text{and} \quad \alpha_k^{(q)} = \frac{\left[ \sum_{j=k}^{K-1} L_j \right]^q - \left[ \sum_{j=k+1}^{K-1} L_j \right]^q}{q! \left[ \sum_{j=0}^{K-1} L_j \right]^q}, \quad (149)$$

and assume for all  $n, q \in \mathbb{N}_0$  that

$$e_{n,q}^2 = \sum_{k=0}^K \alpha_k^{(q)} \mathbb{E}[|V_{k,n}^0(X_k^{0,K,x}) - v_k(X_k^{0,K,x})|^2]. \quad (150)$$

Observe that (149), (150), Lemma 3.10, and Lemma 3.11 ensure that for all  $n \in \mathbb{N}$ ,  $q \in \mathbb{N}_0$  it holds that

$$\begin{aligned} e_{n,q} &\leq \frac{1}{\sqrt{M^n}} \left[ \sum_{k=0}^K \alpha_k^{(q)} \right]^{1/2} \left[ \left( \mathbb{E}[|g(X_0^{0,K,x})|^2] \right)^{1/2} + \sum_{l=0}^{K-1} \left( \mathbb{E}[|f_l(X_l^{0,K,x}, 0)|^2] \right)^{1/2} \right] \\ &\quad + 2 \left[ \max_{k,l \in \mathbb{N}_0, l < k \leq K} \left( \frac{L_l}{\mathfrak{p}_{k,l}} \right) \right] \sum_{j=0}^{n-1} \frac{1}{\sqrt{M^{n-j-1}}} \left( \sum_{k=0}^{K-1} \alpha_k^{(q+1)} \mathbb{E}[|V_{k,j}^0(X_k^{0,K,x}) - v_k(X_k^{0,K,x})|^2] \right)^{1/2}. \end{aligned} \quad (151)$$

The fact that for all  $q \in \mathbb{N}_0$  it holds that  $\sum_{k=0}^K \alpha_k^{(q)} = \frac{1}{q!}$  and (150) therefore ensure that for all  $n \in \mathbb{N}$ ,  $q \in \mathbb{N}_0$  it holds that

$$\begin{aligned} e_{n,q} &\leq \frac{1}{\sqrt{q! M^n}} \left[ \left( \mathbb{E}[|g(X_0^{0,K,x})|^2] \right)^{1/2} + \sum_{l=0}^{K-1} \left( \mathbb{E}[|f_l(X_l^{0,K,x}, 0)|^2] \right)^{1/2} \right] \\ &\quad + 2 \left[ \max_{k,l \in \mathbb{N}_0, l < k \leq K} \left( \frac{L_l}{\mathfrak{p}_{k,l}} \right) \right] \sum_{j=0}^{n-1} \frac{e_{j,q+1}}{\sqrt{M^{n-j-1}}}. \end{aligned} \quad (152)$$

Moreover, note that Lemma 3.6 ensures that

$$\begin{aligned} &\sup_{k \in \{0, 1, \dots, K\}} \left( \mathbb{E}[|v_k(X_k^{0,K,x})|^2] \right)^{1/2} \\ &\leq \exp \left( \sum_{j=0}^{K-1} L_j \right) \left[ \left( \mathbb{E}[|g(X_0^{0,K,x})|^2] \right)^{1/2} + \sum_{l=0}^{K-1} \left( \mathbb{E}[|f_l(X_l^{0,K,x}, 0)|^2] \right)^{1/2} \right]. \end{aligned} \quad (153)$$

This, the fact that for all  $q \in \mathbb{N}_0$  it holds that  $\sum_{k=0}^K \alpha_k^{(q)} = \frac{1}{q!}$ , and (150) ensure that for all  $q \in \mathbb{N}_0$  it holds that

$$\begin{aligned} e_{0,q} &= \left[ \sum_{k=0}^K \alpha_k^{(q)} \mathbb{E}[|v_k(X_k^{0,K,x})|^2] \right]^{1/2} \leq \left[ \sum_{k=0}^K \alpha_k^{(q)} \right]^{1/2} \left[ \sup_{k \in \{0,1,\dots,K\}} \left( \mathbb{E}[|v_k(X_k^{0,K,x})|^2] \right)^{1/2} \right] \\ &\leq \frac{1}{\sqrt{q!}} \exp\left(\sum_{j=0}^{K-1} L_j\right) \left[ \left( \mathbb{E}[|g(X_0^{0,K,x})|^2] \right)^{1/2} + \sum_{l=0}^{K-1} \left( \mathbb{E}[|f_l(X_l^{0,K,x}, 0)|^2] \right)^{1/2} \right]. \end{aligned} \quad (154)$$

Combining this with (152) proves that for all  $n, q \in \mathbb{N}_0$  it holds that

$$\begin{aligned} e_{n,q} &\leq \frac{\exp(\sum_{j=0}^{K-1} L_j)}{\sqrt{q!} M^n} \left[ \left( \mathbb{E}[|g(X_0^{0,K,x})|^2] \right)^{1/2} + \sum_{l=0}^{K-1} \left( \mathbb{E}[|f_l(X_l^{0,K,x}, 0)|^2] \right)^{1/2} \right] \\ &\quad + 2 \left[ \max_{k,l \in \mathbb{N}_0, l < k \leq K} \left( \frac{L_l}{\mathfrak{p}_{k,l}} \right) \right] \sum_{j=0}^{n-1} \frac{e_{j,q+1}}{\sqrt{M^{n-j-1}}}. \end{aligned} \quad (155)$$

Lemma 3.12, (149), and (150) hence demonstrate that

$$\begin{aligned} (\mathbb{E}[|V_{K,N}^0(x) - v_K(x)|^2])^{1/2} &= e_{N,0} \leq \left[ 1 + 2 \max_{k,l \in \mathbb{N}_0, l < k \leq K} \left( \frac{L_l}{\mathfrak{p}_{k,l}} \right) \right]^N \\ &\quad \cdot \exp\left(\sum_{j=0}^{K-1} L_j\right) \left[ \left( \mathbb{E}[|g(X_0^{0,K,x})|^2] \right)^{1/2} + \sum_{l=0}^{K-1} \left( \mathbb{E}[|f_l(X_l^{0,K,x}, 0)|^2] \right)^{1/2} \right] \frac{e^{M/2}}{M^{N/2}}. \end{aligned} \quad (156)$$

This establishes (148) in the case that  $\sum_{k=0}^{K-1} L_k > 0$ . Next we prove (148) in the case that  $\sum_{k=0}^{K-1} L_k \geq 0$ . For this, we note that (148) in the case that  $\sum_{k=0}^{K-1} L_k > 0$  ensures that for all  $N \in \mathbb{N}_0$ ,  $x \in \mathcal{O}$ ,  $\eta \in (0, \infty)$  it holds that

$$\begin{aligned} (\mathbb{E}[|V_{K,N}^0(x) - v_K(x)|^2])^{1/2} &\leq \exp\left(K\eta + \sum_{j=0}^{K-1} L_j\right) \frac{e^{M/2}}{M^{N/2}} \left[ 1 + 2 \max_{k,l \in \mathbb{N}_0, l < k \leq K} \left( \frac{L_l + \eta}{\mathfrak{p}_{k,l}} \right) \right]^N \\ &\quad \cdot \left[ \left( \mathbb{E}[|g(X_0^{0,K,x})|^2] \right)^{1/2} + \sum_{l=0}^{K-1} \left( \mathbb{E}[|f_l(X_l^{0,K,x}, 0)|^2] \right)^{1/2} \right]. \end{aligned} \quad (157)$$

Taking the infimum over  $\eta \in (0, \infty)$  hence establishes (148). This completes the proof of Proposition 3.13.  $\square$

**Corollary 3.14.** *Let  $d, K, M \in \mathbb{N}$ ,  $\Theta = \bigcup_{k \in \mathbb{N}} \mathbb{Z}^k$ ,  $L, T, t_0, t_1, \dots, t_K \in \mathbb{R}$  satisfy  $0 = t_0 < t_1 < \dots < t_K = T$ , let  $f \in C([0, T] \times \mathbb{R}^d \times \mathbb{R}, \mathbb{R})$ , let  $v_k: \mathbb{R}^d \rightarrow \mathbb{R}$ ,  $k \in \{0, 1, \dots, K\}$ , be  $\mathcal{B}(\mathbb{R}^d)/\mathcal{B}(\mathbb{R})$ -measurable, let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a probability space, let  $W^\theta: [0, T] \times \Omega \rightarrow \mathbb{R}^d$ ,  $\theta \in \Theta$ , be i.i.d. standard Brownian motions, let  $R^\theta: \Omega \rightarrow (0, 1)$ ,  $\theta \in \Theta$ , be i.i.d. random variables, assume for all  $r \in (0, 1)$  that  $\mathbb{P}(R^0 \leq r) = r$ , assume that  $(R^\theta)_{\theta \in \Theta}$  and  $(W^\theta)_{\theta \in \Theta}$  are independent, let  $\mathcal{R}^\theta = (\mathcal{R}_k^\theta)_{k \in \{0,1,\dots,K\}}: \{0, 1, \dots, K\} \times \Omega \rightarrow \mathbb{N}_0$ ,  $\theta \in \Theta$ , satisfy for all  $\theta \in \Theta$ ,  $k \in \{0, 1, \dots, K\}$  that  $\mathcal{R}_k^\theta = \max\{n \in \mathbb{N}_0: t_n \leq t_k R^\theta\}$ , assume for all  $k \in \{1, 2, \dots, K\}$ ,  $t \in [0, T]$ ,  $x \in \mathbb{R}^d$ ,  $a, b \in \mathbb{R}$  that  $|f(t, x, a) - f(t, x, b)| \leq L|a - b|$ ,  $\mathbb{E}[|v_0(x + W_T^0)|^2 + \sum_{l=0}^{K-1} |f(t_l, x + W_{T-t_l}^0, 0)|^2] < \infty$ , and*

$$v_k(x) = \mathbb{E}[v_{k-1}(x + W_{t_k - t_{k-1}}^0) + (t_k - t_{k-1})f(t_{k-1}, x + W_{t_k - t_{k-1}}^0, v_{k-1}(x + W_{t_k - t_{k-1}}^0))] \quad (158)$$

(cf. Corollary 3.3), and let  $V_{k,n}^\theta: \mathbb{R}^d \times \Omega \rightarrow \mathbb{R}$ ,  $n \in \mathbb{N}_0$ ,  $k \in \{0, 1, \dots, K\}$ ,  $\theta \in \Theta$ , satisfy for all  $\theta \in \Theta$ ,  $k \in \{0, 1, \dots, K\}$ ,  $n \in \mathbb{N}_0$ ,  $x \in \mathbb{R}^d$  that

$$\begin{aligned} V_{k,n}^\theta(x) &= \frac{\mathbb{1}_{\mathbb{N}}(n)}{M^n} \sum_{m=1}^{M^n} \left[ v_0(x + W_{t_k}^{(\theta,0,-m)}) + \sum_{l=0}^{k-1} (t_{l+1} - t_l) f(t_l, x + W_{t_k}^{(\theta,0,m)} - W_{t_l}^{(\theta,0,m)}, 0) \right] \\ &+ \sum_{j=1}^{n-1} \frac{t_k}{M^{n-j}} \sum_{m=1}^{M^{n-j}} \left[ f(t_{\mathcal{R}_k^{(\theta,j,m)}}, x + W_{t_k}^{(\theta,j,m)} - W_{t_{\mathcal{R}_k^{(\theta,j,m)}}}^{(\theta,j,m)}, V_{\mathcal{R}_k^{(\theta,j,m)},j}^{(\theta,j,m)}(x + W_{t_k}^{(\theta,j,m)} - W_{t_{\mathcal{R}_k^{(\theta,j,m)}}}^{(\theta,j,m)})) \right. \\ &\left. - f(t_{\mathcal{R}_k^{(\theta,j,m)}}, x + W_{t_k}^{(\theta,j,m)} - W_{t_{\mathcal{R}_k^{(\theta,j,m)}}}^{(\theta,j,m)}, V_{\mathcal{R}_k^{(\theta,j,m)},j-1}^{(\theta,j,-m)}(x + W_{t_k}^{(\theta,j,m)} - W_{t_{\mathcal{R}_k^{(\theta,j,m)}}}^{(\theta,j,m)})) \right]. \end{aligned} \quad (159)$$

Then it holds for all  $N \in \mathbb{N}_0$ ,  $x \in \mathbb{R}^d$  that

$$\begin{aligned} \left( \mathbb{E} \left[ |V_{K,N}^0(x) - v_K(x)|^2 \right] \right)^{1/2} &\leq \frac{\exp(LT + \frac{M}{2})[1 + 2LT]^N}{M^{N/2}} \\ &\cdot \left[ \left( \mathbb{E} \left[ |v_0(x + W_T^0)|^2 \right] \right)^{1/2} + \sum_{l=0}^{K-1} (t_{l+1} - t_l) \left( \mathbb{E} \left[ |f(t_l, x + W_{T-t_l}^0, 0)|^2 \right] \right)^{1/2} \right]. \end{aligned} \quad (160)$$

*Proof of Corollary 3.14.* Throughout this proof let  $\|\cdot\|: \mathbb{R}^d \rightarrow [0, \infty)$  be the standard norm on  $\mathbb{R}^d$  and let  $\mathbf{p}_{k,l} \in [0, \infty)$ ,  $k \in \{1, 2, \dots, K\}$ ,  $l \in \mathbb{N}_0$ , satisfy for all  $k \in \{1, 2, \dots, K\}$  that

$$\mathbf{p}_{k,l} = \begin{cases} \frac{t_{l+1} - t_l}{t_k} & : l < k \\ 1 & : l \geq k. \end{cases} \quad (161)$$

Observe that the assumption that  $W^0$  is a Brownian motion and the assumption that for all  $x \in \mathbb{R}^d$  it holds that  $\mathbb{E}[|v_0(x + W_T^0)|^2 + \sum_{l=0}^{K-1} |f(t_l, x + W_{T-t_l}^0, 0)|^2] < \infty$  show that for all  $k \in \{0, 1, \dots, K\}$ ,  $x \in \mathbb{R}^d$  it holds that  $\mathbb{E}[|v_0(x + W_{t_k}^0)|^2 + \sum_{l=0}^{k-1} |f(t_l, x + W_{t_k}^0 - W_{t_l}^0, 0)|^2] < \infty$ . In addition, note that (158) and the fact that  $W^0: [0, T] \times \Omega \rightarrow \mathbb{R}^d$  is a standard Brownian motion ensure that for all  $k \in \{1, 2, \dots, K\}$ ,  $x \in \mathbb{R}^d$  it holds that

$$\begin{aligned} v_k(x) &= \mathbb{E}[v_{k-1}(x + W_{t_k}^0 - W_{t_{k-1}}^0) \\ &+ (t_k - t_{k-1}) f(t_{k-1}, x + W_{t_k}^0 - W_{t_{k-1}}^0, v_{k-1}(x + W_{t_k}^0 - W_{t_{k-1}}^0))]. \end{aligned} \quad (162)$$

Moreover, note that Proposition 3.13 (applied with  $d \leftarrow d$ ,  $K \leftarrow K$ ,  $M \leftarrow M$ ,  $\Theta \leftarrow \Theta$ ,  $\mathcal{O} \leftarrow \mathbb{R}^d$ ,  $(f_k)_{k \in \{0,1,\dots,K\}} \leftarrow ((\mathbb{R}^d \times \mathbb{R} \ni (x, a) \mapsto a + (t_{\min\{k+1,K\}} - t_k) f(t_k, x, a) \in \mathbb{R}))_{k \in \{0,1,\dots,K\}}$ ,  $g \leftarrow v_0$ ,  $c \leftarrow LT$ ,  $(L_k)_{k \in \{0,1,\dots,K\}} \leftarrow ((t_{\min\{k+1,K\}} - t_k) L)_{k \in \{0,1,\dots,K\}}$ ,  $(S, \mathcal{S}) \leftarrow (\mathbb{R}^d, \mathcal{B}(\mathbb{R}^d))$ ,  $(\phi_k)_{k \in \{0,1,\dots,K\}} \leftarrow (\mathbb{R}^d \times \mathbb{R}^d \ni (x, w) \mapsto x + w \in \mathbb{R}^d)_{k \in \{0,1,\dots,K\}}$ ,  $(\Omega, \mathcal{F}, \mathbb{P}) \leftarrow (\Omega, \mathcal{F}, \mathbb{P})$ ,  $(W_k^\theta)_{(\theta,k) \in \Theta \times \{0,1,\dots,K\}} \leftarrow (W_{t_{\min\{k+1,K\}}}^\theta - W_{t_k}^\theta)_{(\theta,k) \in \Theta \times \{0,1,\dots,K\}}$ ,  $(\mathcal{R}^\theta)_{\theta \in \Theta} \leftarrow (\mathcal{R}^\theta)_{\theta \in \Theta}$ ,  $(\mathbf{p}_{k,l})_{(k,l) \in \{1,2,\dots,K\} \times \mathbb{N}_0} \leftarrow (\mathbf{p}_{k,l})_{(k,l) \in \{1,2,\dots,K\} \times \mathbb{N}_0}$  in the notation of Proposition 3.13) and the fact that  $W^0$  is a standard Brownian motion therefore demonstrate that for all  $N \in \mathbb{N}_0$ ,  $x \in \mathbb{R}^d$  it holds that

$$\begin{aligned} \left( \mathbb{E} \left[ |V_{K,N}^0(x) - v_K(x)|^2 \right] \right)^{1/2} &\leq \frac{e^{M/2}}{M^{N/2}} \exp \left( \sum_{l=0}^{K-1} (t_{l+1} - t_l) L \right) \left[ 1 + 2 \max_{k,l \in \{0,1,\dots,K\}, k > l} \frac{(t_{l+1} - t_l) L}{\mathbf{p}_{k,l}} \right]^N \\ &\cdot \left[ \left( \mathbb{E} \left[ |v_0(x + W_T^0)|^2 \right] \right)^{1/2} + \sum_{l=0}^{K-1} \left( \mathbb{E} \left[ |(t_{l+1} - t_l) f(t_l, x + W_T^0 - W_{t_l}^0, 0)|^2 \right] \right)^{1/2} \right] \\ &= \frac{e^{M/2} e^{LT} [1 + 2LT]^N}{M^{N/2}} \left[ \left( \mathbb{E} \left[ |v_0(x + W_T^0)|^2 \right] \right)^{1/2} + \sum_{l=0}^{K-1} (t_{l+1} - t_l) \left( \mathbb{E} \left[ |f(t_l, x + W_{T-t_l}^0, 0)|^2 \right] \right)^{1/2} \right]. \end{aligned} \quad (163)$$

This establishes (160). This completes the proof of Corollary 3.14.  $\square$

**Corollary 3.15.** Let  $d, K, M \in \mathbb{N}$ ,  $\Theta = \bigcup_{n \in \mathbb{N}} \mathbb{Z}^n$ ,  $L, T \in [0, \infty)$ ,  $f \in C(\mathbb{R}, \mathbb{R})$  satisfy for all  $a, b \in \mathbb{R}$  that  $|(f(a) - a) - (f(b) - b)| \leq L|a - b|$ , let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a probability space, let  $W^\theta: [0, T] \times \Omega \rightarrow \mathbb{R}^d$ ,  $\theta \in \Theta$ , be i.i.d. standard Brownian motions, let  $R^\theta: \Omega \rightarrow (0, 1)$ ,  $\theta \in \Theta$ , be i.i.d. random variables, assume for all  $r \in (0, 1)$  that  $\mathbb{P}(R^\theta \leq r) = r$ , assume that  $(R^\theta)_{\theta \in \Theta}$  and  $(W^\theta)_{\theta \in \Theta}$  are independent, let  $\mathcal{R}_k^\theta: \Omega \rightarrow \{0, 1, \dots, K\}$ ,  $\theta \in \Theta$ ,  $k \in \{0, 1, \dots, K\}$ , satisfy for all  $\theta \in \Theta$ ,  $k \in \{0, 1, \dots, K\}$  that  $\mathcal{R}_k^\theta = \max\{n \in \mathbb{N}_0: n \leq kR^\theta\}$ , let  $v_k: \mathbb{R}^d \rightarrow \mathbb{R}$ ,  $k \in \{0, 1, \dots, K\}$ , satisfy for all  $k \in \{1, 2, \dots, K\}$ ,  $x \in \mathbb{R}^d$  that  $v_0 \in C(\mathbb{R}^d, \mathbb{R})$ ,  $\mathbb{E}[|v_0(x + W_T^0)|^2] < \infty$ , and

$$v_k(x) = \mathbb{E}[f(v_{k-1}(x + W_{T/K}^0))] \quad (164)$$

(cf. Corollary 3.3), and let  $V_{k,n}^\theta: \mathbb{R}^d \times \Omega \rightarrow \mathbb{R}$ ,  $n \in \mathbb{N}_0$ ,  $k \in \{0, 1, \dots, K\}$ ,  $\theta \in \Theta$ , satisfy for all  $\theta \in \Theta$ ,  $k \in \{0, 1, \dots, K\}$ ,  $n \in \mathbb{N}_0$ ,  $x \in \mathbb{R}^d$  that

$$\begin{aligned} V_{k,n}^\theta(x) &= \frac{\mathbb{1}_{\mathbb{N}}(n)}{M^n} \sum_{m=1}^{M^n} \left[ v_0(x + W_{kT/K}^{(\theta,0,-m)}) + kf(0) \right] + \sum_{j=1}^{n-1} \frac{k}{M^{n-j}} \\ &\cdot \sum_{m=1}^{M^{n-j}} \left[ \left( f(V_{\mathcal{R}_k^{(\theta,j,m)},j}^{(\theta,j,m)}(x + W_{kT/K}^{(\theta,j,m)} - W_{\mathcal{R}_k^{(\theta,j,m)}T/K}^{(\theta,j,m)})) - V_{\mathcal{R}_k^{(\theta,j,m)},j}^{(\theta,j,m)}(x + W_{kT/K}^{(\theta,j,m)} - W_{\mathcal{R}_k^{(\theta,j,m)}T/K}^{(\theta,j,m)})) \right) \right. \\ &\quad \left. - \left( f(V_{\mathcal{R}_k^{(\theta,j,m)},j-1}^{(\theta,j,-m)}(x + W_{kT/K}^{(\theta,j,m)} - W_{\mathcal{R}_k^{(\theta,j,m)}T/K}^{(\theta,j,m)})) - V_{\mathcal{R}_k^{(\theta,j,m)},j-1}^{(\theta,j,-m)}(x + W_{kT/K}^{(\theta,j,m)} - W_{\mathcal{R}_k^{(\theta,j,m)}T/K}^{(\theta,j,m)})) \right) \right]. \end{aligned} \quad (165)$$

Then it holds for all  $N \in \mathbb{N}_0$ ,  $x \in \mathbb{R}^d$  that

$$\left( \mathbb{E}[|V_{K,N}^0(x) - v_K(x)|^2] \right)^{1/2} \leq \frac{e^{M/2+KL} [1 + 2KL]^N}{M^{N/2}} \left[ (\mathbb{E}[|v_0(x + W_T^0)|^2])^{1/2} + K|f(0)| \right]. \quad (166)$$

*Proof of Corollary 3.15.* Observe that Corollary 3.14 (applied with  $d \leftarrow d$ ,  $K \leftarrow K$ ,  $M \leftarrow M$ ,  $L \leftarrow \frac{KL}{T}$ ,  $T \leftarrow T$ ,  $\Theta \leftarrow \Theta$ ,  $(t_k)_{k \in \{0,1,\dots,K\}} \leftarrow (\frac{kT}{K})_{k \in \{0,1,\dots,K\}}$ ,  $f \leftarrow ([0, T] \times \mathbb{R}^d \times \mathbb{R} \ni (t, x, a) \mapsto \frac{K}{T}(f(a) - a) \in \mathbb{R})$ ,  $(v_k)_{k \in \{0,1,\dots,K\}} \leftarrow (v_k)_{k \in \{0,1,\dots,K\}}$ ,  $(\Omega, \mathcal{F}, \mathbb{P}) \leftarrow (\Omega, \mathcal{F}, \mathbb{P})$ ,  $(W^\theta)_{\theta \in \Theta} \leftarrow (W^\theta)_{\theta \in \Theta}$ ,  $(R^\theta)_{\theta \in \Theta} \leftarrow (R^\theta)_{\theta \in \Theta}$  in the notation of Corollary 3.14) ensures that for all  $N \in \mathbb{N}_0$ ,  $x \in \mathbb{R}^d$  it holds that

$$\begin{aligned} & \left( \mathbb{E}[|V_{K,N}^0(x) - v_K(x)|^2] \right)^{1/2} \leq \frac{e^{M/2}}{M^{N/2}} [1 + 2(\frac{KL}{T})T]^N \\ & \cdot e^{\frac{KL}{T}T} \left[ \left( \mathbb{E}[|v_0(x + W_T^0)|^2] \right)^{1/2} + \sum_{l=0}^{K-1} \frac{T}{K} \left( \mathbb{E}[|\frac{K}{T}(f(0) - 0)|^2] \right)^{1/2} \right] \\ & = \frac{e^{M/2}}{M^{N/2}} e^{KL} (1 + 2KL)^N \left[ (\mathbb{E}[|v_0(x + W_T^0)|^2])^{1/2} + K|f(0)| \right]. \end{aligned} \quad (167)$$

This establishes (166). This completes the proof of Corollary 3.15.  $\square$

## 4 Complexity analysis for MLP approximations for iterated nested expectations

In this section we combine the error analysis for the proposed MLP approximation schemes in Proposition 3.13 in Subsection 3.4 with a computational cost analysis for the proposed MLP approximation schemes to obtain in Theorem 4.1 in Subsection 4.1 below a complexity analysis for the proposed MLP approximation schemes. In Corollary 4.4 in Subsection 4.2 below we illustrate Theorem 4.1 by means of a sample application to exponential Euler approximations.

## 4.1 Complexity analysis for MLP approximations

**Theorem 4.1.** Let  $\alpha, \gamma, c, \kappa, p \in [0, \infty)$ ,  $\Theta = \bigcup_{n \in \mathbb{N}} \mathbb{Z}^n$ , let  $\xi_d \in \mathbb{R}^d$ ,  $d \in \mathbb{N}$ , let  $L_k^{d,K} \in \mathbb{R}$ ,  $k \in \{0, 1, \dots, K\}$ ,  $d, K \in \mathbb{N}$ , and  $f_k^{d,K} \in C(\mathbb{R}^d \times \mathbb{R}, \mathbb{R})$ ,  $k \in \{0, 1, \dots, K\}$ ,  $d, K \in \mathbb{N}$ , satisfy for all  $d, K \in \mathbb{N}$ ,  $k \in \{0, 1, \dots, K\}$ ,  $x \in \mathbb{R}^d$ ,  $a, b \in \mathbb{R}$  that  $|(f_k^{d,K}(x, a) - a) - (f_k^{d,K}(x, b) - b)| \leq L_k^{d,K}|a - b|$ , for every  $d, K \in \mathbb{N}$ ,  $k \in \{0, 1, \dots, K\}$  let  $\phi_k^{d,K} : \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}^d$  be  $(\mathcal{B}(\mathbb{R}^d \times \mathbb{R}^d))/\mathcal{B}(\mathbb{R}^d)$ -measurable, let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a probability space, for every  $d, K \in \mathbb{N}$  let  $W^{d,K,\theta} = (W_k^{d,K,\theta})_{k \in \{0,1,\dots,K\}} : \{0, 1, \dots, K\} \times \Omega \rightarrow \mathbb{R}^d$ ,  $\theta \in \Theta$ , be i.i.d. stochastic processes, assume for all  $d, K \in \mathbb{N}$ ,  $\theta \in \Theta$  that  $W_0^{d,K,\theta}, W_1^{d,K,\theta}, \dots, W_{K-1}^{d,K,\theta}$  are independent, for every  $d, K \in \mathbb{N}$  let  $\mathcal{R}^{d,K,\theta} = (\mathcal{R}_k^{d,K,\theta})_{k \in \{0,1,\dots,K\}} : \{0, 1, \dots, K\} \times \Omega \rightarrow \mathbb{N}_0$ ,  $\theta \in \Theta$ , be i.i.d. stochastic processes, let  $\mathfrak{p}_{k,l}^{d,K} \in (0, \infty)$ ,  $l \in \mathbb{N}_0$ ,  $k \in \{1, 2, \dots, K\}$ ,  $d, K \in \mathbb{N}$ , assume for all  $d, K \in \mathbb{N}$ ,  $k \in \{1, 2, \dots, K\}$ ,  $l \in \{0, 1, \dots, k-1\}$  that  $\mathcal{R}_k^{d,K,\theta} \leq k-1$ ,  $\mathfrak{p}_{k,l}^{d,K} \mathbb{P}(\mathcal{R}_k^{d,K,\theta} = l) = |\mathbb{P}(\mathcal{R}_k^{d,K,\theta} = l)|^2$ , and  $L_l^{d,K} \leq c\mathbb{P}(\mathcal{R}_k^{d,K,\theta} = l)$ , assume for all  $d, K \in \mathbb{N}$  that  $(R^{d,K,\theta})_{\theta \in \Theta}$  and  $(W^{d,K,\theta})_{\theta \in \Theta}$  are independent, for every  $d, K \in \mathbb{N}$ ,  $k \in \{0, 1, \dots, K\}$ ,  $\theta \in \Theta$  let  $X^{d,K,\theta,k} = (X_l^{d,K,\theta,k,x})_{(l,x) \in \{0,1,\dots,k\} \times \mathbb{R}^d} : \{0, 1, \dots, k\} \times \mathbb{R}^d \times \Omega \rightarrow \mathbb{R}^d$  satisfy for all  $l \in \{0, 1, \dots, k\}$ ,  $x \in \mathbb{R}^d$  that

$$X_l^{d,K,\theta,k,x} = \begin{cases} x & : l = k \\ \phi_l^{d,K}(X_{l+1}^{d,K,\theta,k,x}, W_l^{d,K,\theta}) & : l < k, \end{cases} \quad (168)$$

for every  $d, K \in \mathbb{N}$  let  $v_k^{d,K} : \mathbb{R}^d \rightarrow \mathbb{R}$ ,  $k \in \{0, 1, \dots, K\}$ , be  $\mathcal{B}(\mathbb{R}^d)/\mathcal{B}(\mathbb{R})$ -measurable, assume for all  $d, K \in \mathbb{N}$ ,  $k \in \{1, 2, \dots, K\}$ ,  $x \in \mathbb{R}^d$  that  $\mathbb{E}[|v_0^{d,K}(X_0^{d,K,0,k,x})|^2 + \sum_{l=0}^{k-1} |f_l^{d,K}(X_l^{d,K,0,k,x}, 0)|^2] < \infty$ ,  $(\mathbb{E}[|v_0^{d,K}(X_0^{d,K,0,K,\xi_d})|^2])^{1/2} + \sum_{l=0}^{K-1} (\mathbb{E}[|f_l^{d,K}(X_l^{d,K,0,K,\xi_d}, 0)|^2])^{1/2} \leq \kappa$ , and

$$v_k^{d,K}(x) = \mathbb{E}[f_{k-1}^{d,K}(X_{k-1}^{d,K,0,k,x}, v_{k-1}^{d,K}(X_{k-1}^{d,K,0,k,x}))] \quad (169)$$

(cf. Lemma 3.2), let  $V_{k,n,M}^{d,K,\theta} : \mathbb{R}^d \times \Omega \rightarrow \mathbb{R}$ ,  $\theta \in \Theta$ ,  $k \in \{0, 1, \dots, K\}$ ,  $n \in \mathbb{N}_0$ ,  $d, K, M \in \mathbb{N}$ , satisfy for all  $d, K, M \in \mathbb{N}$ ,  $n \in \mathbb{N}_0$ ,  $\theta \in \Theta$ ,  $k \in \{0, 1, \dots, K\}$ ,  $x \in \mathbb{R}^d$  that

$$\begin{aligned} & V_{k,n,M}^{d,K,\theta}(x) \\ &= \frac{\mathbb{1}_{\mathbb{N}}(n)}{M^n} \sum_{m=1}^{M^n} \left[ v_0^{d,K}(X_0^{d,K,(\theta,0,-m),k,x}) + \sum_{l=0}^{k-1} f_l^{d,K}(X_l^{d,K,(\theta,0,m),k,x}, 0) \right] + \sum_{j=1}^{n-1} \frac{1}{M^{n-j}} \sum_{m=1}^{M^{n-j}} \frac{\mathbb{1}_{\mathbb{N}}(k)}{\mathfrak{p}_{k,\mathcal{R}_k^{d,K,(\theta,j,m)}}^{d,K}} \\ & \cdot \left[ f_{\mathcal{R}_k^{d,K,(\theta,j,m)}}^{d,K}(X_{\mathcal{R}_k^{d,K,(\theta,j,m)}}^{d,K,(\theta,j,m),k,x}, V_{\mathcal{R}_k^{d,K,(\theta,j,m)},j,M}^{d,K,(\theta,j,m)}(X_{\mathcal{R}_k^{d,K,(\theta,j,m)}}^{d,K,(\theta,j,m),k,x})) - V_{\mathcal{R}_k^{d,K,(\theta,j,m)},j,M}^{d,K,(\theta,j,m)}(X_{\mathcal{R}_k^{d,K,(\theta,j,m)}}^{d,K,(\theta,j,m),k,x}) \right. \\ & \left. - f_{\mathcal{R}_k^{d,K,(\theta,j,m)}}^{d,K}(X_{\mathcal{R}_k^{d,K,(\theta,j,m)}}^{d,K,(\theta,j,m),k,x}, V_{\mathcal{R}_k^{d,K,(\theta,j,m)},j-1,M}^{d,K,(\theta,j,m)}(X_{\mathcal{R}_k^{d,K,(\theta,j,m)}}^{d,K,(\theta,j,m),k,x})) + V_{\mathcal{R}_k^{d,K,(\theta,j,m)},j-1,M}^{d,K,(\theta,j,m)}(X_{\mathcal{R}_k^{d,K,(\theta,j,m)}}^{d,K,(\theta,j,m),k,x}) \right], \end{aligned} \quad (170)$$

and let  $\mathfrak{C}_{M,n}^{d,K} \in \mathbb{R}$ ,  $d, K, M, n \in \mathbb{N}_0$ , satisfy for all  $d, K, M \in \mathbb{N}$ ,  $n \in \mathbb{N}_0$  that

$$\mathfrak{C}_{M,n}^{d,K} \leq \gamma K^\alpha d^p M^n + \sum_{j=1}^{n-1} M^{n-j} (1 + \gamma K^\alpha d^p + \mathfrak{C}_{M,j}^{d,K} + \mathfrak{C}_{M,j-1}^{d,K}). \quad (171)$$

Then

(i) it holds for all  $d, K, M \in \mathbb{N}$ ,  $N \in \mathbb{N}_0$  that

$$\left( \mathbb{E}[|V_{K,N,M}^{d,K,0}(\xi_d) - v_K^{d,K}(\xi_d)|^2] \right)^{1/2} \leq \kappa(1 + 2c)^N M^{-N/2} \exp(M/2 + c) \quad (172)$$

and

(ii) there exist  $\mathfrak{N} = (\mathfrak{N}_\varepsilon)_{\varepsilon \in (0,1]} : (0, 1] \rightarrow \mathbb{N}$  and  $\mathfrak{c} = (\mathfrak{c}_\delta)_{\delta \in (0,1]} : (0, 1] \rightarrow [0, \infty)$  such that for all  $d, K \in \mathbb{N}$ ,  $\delta, \varepsilon \in (0, 1]$  it holds that  $\mathfrak{C}_{\mathfrak{N}_\varepsilon, \mathfrak{N}_\varepsilon}^{d,K} \leq \mathfrak{c}_\delta K^\alpha d^p \varepsilon^{-(2+\delta)}$  and

$$\left( \mathbb{E} [|V_{K, \mathfrak{N}_\varepsilon, \mathfrak{N}_\varepsilon}^{d,K,0}(\xi_d) - v_K^{d,K}(\xi_d)|^2] \right)^{1/2} \leq \varepsilon. \quad (173)$$

*Proof of Theorem 4.1.* First, observe that Proposition 3.13 (applied with  $d \leftarrow d$ ,  $K \leftarrow K$ ,  $M \leftarrow M$ ,  $\Theta \leftarrow \Theta$ ,  $\mathcal{O} \leftarrow \mathbb{R}^d$ ,  $(f_k)_{k \in \{0,1,\dots,K\}} \leftarrow (f_k^{d,K})_{k \in \{0,1,\dots,K\}}$ ,  $g \leftarrow v_0^{d,K}$ ,  $c \leftarrow c$ ,  $(L_k)_{k \in \{0,1,\dots,K\}} \leftarrow (L_k^{d,K})_{k \in \{0,1,\dots,K\}}$ ,  $(\phi_k)_{k \in \{0,1,\dots,K\}} \leftarrow (\phi_k^{d,K})_{k \in \{0,1,\dots,K\}}$ ,  $(\mathcal{R}^\theta)_{\theta \in \Theta} \leftarrow (\mathcal{R}^{d,K,\theta})_{\theta \in \Theta}$ ,  $(S, \mathcal{S}) \leftarrow (\mathbb{R}^d, \mathcal{B}(\mathbb{R}^d))$ ,  $(\Omega, \mathcal{F}, \mathbb{P}) \leftarrow (\Omega, \mathcal{F}, \mathbb{P})$ ,  $(\mathfrak{p}_{k,l})_{(k,l) \in \{1,2,\dots,K\} \times \mathbb{N}_0} \leftarrow (\mathfrak{p}_{k,l}^{d,K})_{(k,l) \in \{1,2,\dots,K\} \times \mathbb{N}_0}$ ,  $(W^\theta)_{\theta \in \Theta} \leftarrow (W^{d,K,\theta})_{\theta \in \Theta}$  for  $d, K, M \in \mathbb{N}$  in the notation of Proposition 3.13) ensures that for all  $d, K, M \in \mathbb{N}$ ,  $N \in \mathbb{N}_0$ ,  $x \in \mathbb{R}^d$  it holds that

$$\begin{aligned} & \left( \mathbb{E} [|V_{K,N,M}^{d,K,0}(x) - v_K^{d,K}(x)|^2] \right)^{1/2} \\ & \leq \frac{e^{M/2}}{M^{N/2}} \exp \left( \sum_{j=0}^{K-1} L_j^{d,K} \right) \left[ 1 + 2 \max_{k,j \in [0,K] \cap \mathbb{N}_0, k > j} \left( \frac{L_j^{d,K}}{\mathfrak{p}_{k,j}^{d,K}} \right) \right]^N \\ & \quad \cdot \left[ \left( \mathbb{E} [|v_0^{d,K}(X_0^{d,K,0,K,x})|^2] \right)^{1/2} + \sum_{l=0}^{K-1} \left( \mathbb{E} [|f_l^{d,K}(X_l^{d,K,0,K,x}, 0)|^2] \right)^{1/2} \right]. \end{aligned} \quad (174)$$

The fact that for all  $d, K \in \mathbb{N}$ ,  $k \in \{1, 2, \dots, K\}$ ,  $l \in \{0, 1, \dots, k-1\}$  it holds that  $L_l^{d,K} \leq c \mathbb{P}(\mathcal{R}_k^{d,K,0} = l) \leq c \mathfrak{p}_{k,l}^{d,K}$  hence implies that for all  $d, K, M \in \mathbb{N}$ ,  $N \in \mathbb{N}_0$ ,  $x \in \mathbb{R}^d$  it holds that

$$\begin{aligned} & \left( \mathbb{E} [|V_{K,N,M}^{d,K,0}(x) - v_K^{d,K}(x)|^2] \right)^{1/2} \\ & \leq \frac{e^{M/2}}{M^{N/2}} e^c (1 + 2c)^N \left[ \left( \mathbb{E} [|v_0^{d,K}(X_0^{d,K,0,K,x})|^2] \right)^{1/2} + \sum_{l=0}^{K-1} \left( \mathbb{E} [|f_l^{d,K}(X_l^{d,K,0,K,x}, 0)|^2] \right)^{1/2} \right]. \end{aligned} \quad (175)$$

Combining this with the assumption that for all  $d, K \in \mathbb{N}$  it holds that  $(\mathbb{E} [|v_0^{d,K}(X_0^{d,K,0,K,\xi_d})|^2])^{1/2} + \sum_{l=0}^{K-1} (\mathbb{E} [|f_l^{d,K}(X_l^{d,K,0,K,\xi_d}, 0)|^2])^{1/2} \leq \kappa$  shows that for all  $d, K, M \in \mathbb{N}$ ,  $N \in \mathbb{N}_0$  it holds that

$$\left( \mathbb{E} [|V_{K,N,M}^{d,K,0}(\xi_d) - v_K^{d,K}(\xi_d)|^2] \right)^{1/2} \leq \frac{e^{M/2}}{M^{N/2}} e^c (1 + 2c)^N \kappa. \quad (176)$$

This establishes item (i). Next we prove item (ii). Observe that [3, Lemma 3.14] (applied with  $\alpha \leftarrow \gamma K^\alpha d^p$ ,  $\beta \leftarrow 1 + \gamma K^\alpha d^p$ ,  $M \leftarrow M$ ,  $(C_n)_{n \in \mathbb{N}_0} \leftarrow (\max\{\mathfrak{C}_{M,n}^{d,K}, 0\})_{n \in \mathbb{N}_0}$  for  $d, K, M \in \mathbb{N}$  in the notation of [3, Lemma 3.14]) and (171) show that for all  $d, K, M, n \in \mathbb{N}$  it holds that

$$\mathfrak{C}_{M,n}^{d,K} \leq \frac{2\gamma K^\alpha d^p + (1 + \gamma K^\alpha d^p)}{2} (3M)^n \leq \left[ \frac{1 + 3\gamma}{2} \right] d^p K^\alpha (3M)^n. \quad (177)$$

In addition, note that [4, Lemma 4.2] (applied with  $\alpha \leftarrow 3$ ,  $\beta \leftarrow 2$ ,  $c \leftarrow \kappa + 1$ ,  $\kappa \leftarrow \sqrt{e}$ ,  $\rho \leftarrow 1$ ,  $K \leftarrow 0$ ,  $(\gamma_n)_{n \in \mathbb{N}} \leftarrow ((3n)^n)_{n \in \mathbb{N}}$ ,  $(\epsilon_{n,r})_{n \in \mathbb{N}, r \in [\rho, \infty)} \leftarrow (\kappa e^c [\sqrt{e}(1 + 2c)]^n n^{-n/2})_{n \in \mathbb{N}, r \in [1, \infty)}$ ,  $L \leftarrow ((0, \infty) \ni r \mapsto c \in [0, \infty))$ ,  $\varrho \leftarrow (\mathbb{N} \ni n \mapsto n \in (0, \infty))$  in the notation of [4, Lemma 4.2]) ensures that there exist  $\mathfrak{N} = (\mathfrak{N}_\varepsilon)_{\varepsilon \in (0,1]} : (0, 1] \rightarrow \mathbb{N}$  and  $\mathfrak{c} = (\mathfrak{c}_\delta)_{\delta \in (0,1]} : (0, 1] \rightarrow [0, \infty)$  which satisfy for all  $\delta, \varepsilon \in (0, 1]$  that

$$(3\mathfrak{N}_\varepsilon)^{\mathfrak{N}_\varepsilon} \leq \mathfrak{c}_\delta \varepsilon^{-(2+\delta)} \quad \text{and} \quad \kappa e^c [\sqrt{e}(1 + 2c)]^{\mathfrak{N}_\varepsilon} \mathfrak{N}_\varepsilon^{-\mathfrak{N}_\varepsilon/2} \leq \varepsilon. \quad (178)$$

This, (176), and (177) imply that for all  $\delta \in (0, \infty)$ ,  $\varepsilon \in (0, 1]$ ,  $d, K \in \mathbb{N}$  it holds that

$$\mathfrak{C}_{\mathfrak{N}_\varepsilon, \mathfrak{N}_\varepsilon}^{d,K} \leq \mathfrak{c}_\delta \left[ \frac{1 + 3\gamma}{2} \right] d^p K^\alpha \varepsilon^{-(2+\delta)} \quad \text{and} \quad \left( \mathbb{E} [|V_{K, \mathfrak{N}_\varepsilon, \mathfrak{N}_\varepsilon}^{d,K,0}(\xi_d) - v_K^{d,K}(\xi_d)|^2] \right)^{1/2} \leq \varepsilon. \quad (179)$$

This establishes item (ii). This completes the proof of Theorem 4.1.  $\square$

## 4.2 Exponential Euler approximations for semilinear heat equations

**Lemma 4.2.** *Let  $d \in \mathbb{N}$ ,  $p \in (0, 2)$ ,  $r, T \in [0, \infty)$ , let  $\|\cdot\|: \mathbb{R}^d \rightarrow [0, \infty)$  be the standard norm on  $\mathbb{R}^d$ , let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a probability space, let  $W: [0, T] \times \Omega \rightarrow \mathbb{R}^d$  be a standard Brownian motion, and let  $h \in C(\mathbb{R}^d, \mathbb{R})$  satisfy that  $\mathbb{E}[|h(W_T)|^2] < \infty$ . Then  $\sup_{x \in \mathbb{R}^d, \|x\| \leq r} \mathbb{E}[|h(x + W_T)|^p] < \infty$ .*

*Proof of Lemma 4.2.* Throughout this proof assume without loss of generality that  $\min\{r, T\} > 0$ , let  $\alpha \in (0, 1)$  satisfy that  $(1 - \alpha)^2 > \frac{p}{2}$ , and let  $R = \frac{r}{\alpha}$ . Note that the assumption that  $h \in C(\mathbb{R}^d, \mathbb{R})$  and the fact that  $\{z \in \mathbb{R}^d: \|z\| \leq r\}$  is compact ensure that

$$\sup_{x \in \mathbb{R}^d, \|x\| \leq r} \mathbb{E}[\mathbb{1}_{\{\|W_T\| \leq 2R\}} |h(x + W_T)|^p] < \infty. \quad (180)$$

Next note that the triangle inequality proves that for all  $x, z \in \mathbb{R}^d$  with  $\|x\| \leq r$  and  $\|z - x\| \geq 2R$  it holds

(i) that  $\|z\| \geq 2R - \|x\| \geq 2R - r = 2R - \alpha R > R$ ,

(ii) that  $\|x\| \leq r = \alpha R \leq \alpha \|z\|$ , and

(iii) that  $\|z - x\| \geq \|z\| - \|x\| \geq (1 - \alpha) \|z\|$ .

Observe that item (iii) implies that for all  $x \in \mathbb{R}^d$  with  $\|x\| \leq r$  it holds that

$$\begin{aligned} & \mathbb{E}[\mathbb{1}_{\{\|W_T\| > 2R\}} |h(x + W_T)|^p] \\ &= (2\pi T)^{-d/2} \int_{\mathbb{R}^d} \mathbb{1}_{\{z \in \mathbb{R}^d: \|z\| > 2R\}}(y) |h(x + y)|^p \exp(-\frac{\|y\|^2}{2T}) dy \\ &= (2\pi T)^{-d/2} \int_{\mathbb{R}^d} \mathbb{1}_{\{z \in \mathbb{R}^d: \|z\| > 2R\}}(y - x) |h(y)|^p \exp(-\frac{\|y - x\|^2}{2T}) dy \\ &\leq (2\pi T)^{-d/2} \int_{\mathbb{R}^d} |h(y)|^p \exp(-(1 - \alpha)^2 \frac{\|y\|^2}{2T}) dy. \end{aligned} \quad (181)$$

The fact that  $(1 - \alpha)^2 > \frac{p}{2}$  and Hölder's inequality therefore show that for all  $x \in \mathbb{R}^d$  with  $\|x\| \leq r$  it holds that

$$\begin{aligned} & \mathbb{E}[\mathbb{1}_{\{\|W_T\| > 2R\}} |h(x + W_T)|^p] \\ &\leq (2\pi T)^{-d/2} \int_{\mathbb{R}^d} \left[ |h(y)|^2 \exp(-\frac{\|y\|^2}{2T}) \right]^{p/2} \exp(-((1 - \alpha)^2 - \frac{p}{2}) \frac{\|y\|^2}{2T}) dy \\ &\leq (2\pi T)^{-d/2} (\mathbb{E}[|h(W_T)|^2])^{p/2} \left[ \int_{\mathbb{R}^d} \exp(-\frac{(1 - \alpha)^2 - p/2}{1 - p/2} \frac{\|y\|^2}{2T}) dy \right]^{1 - p/2}. \end{aligned} \quad (182)$$

Hence, we obtain that  $\sup_{x \in \mathbb{R}^d, \|x\| \leq r} \mathbb{E}[\mathbb{1}_{\{\|W_T\| \geq 2R\}} |h(x + W_T)|^p] < \infty$ . Combining this with (180) shows that  $\sup_{x \in \mathbb{R}^d, \|x\| \leq r} \mathbb{E}[|h(x + W_T)|^p] < \infty$ . This completes the proof of Lemma 4.2.  $\square$

**Lemma 4.3.** *Let  $d \in \mathbb{N}$ ,  $f \in C(\mathbb{R}^d \times \mathbb{R}, \mathbb{R})$ ,  $g \in C(\mathbb{R}^d, \mathbb{R})$ ,  $L, T \in [0, \infty)$ , let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a probability space, let  $W: [0, T] \times \Omega \rightarrow \mathbb{R}^d$  be a standard Brownian motion, assume for all  $x \in \mathbb{R}^d$ ,  $a, b \in \mathbb{R}$  that  $|f(x, a) - f(x, b)| \leq L|a - b|$  and  $\mathbb{E}[|f(x + W_T, 0)|^2 + |g(x + W_T)|^2] < \infty$ , and let  $u: \mathbb{R}^d \rightarrow \mathbb{R}$  satisfy for all  $x \in \mathbb{R}^d$  that  $u(x) = \mathbb{E}[f(x + W_T, g(x + W_T))]$  (cf. Lemma 3.1). Then  $u \in C(\mathbb{R}^d, \mathbb{R})$ .*

*Proof of Lemma 4.3.* Throughout this proof let  $\|\cdot\|: \mathbb{R}^d \rightarrow [0, \infty)$  be the standard norm on  $\mathbb{R}^d$  and let  $\mathbf{x}_n \in \mathbb{R}^d$ ,  $n \in \mathbb{N}_0$ , satisfy  $\limsup_{n \rightarrow \infty} \|\mathbf{x}_n - \mathbf{x}_0\| = 0$ . The assumption that  $f \in C(\mathbb{R}^d \times \mathbb{R}, \mathbb{R})$  and the assumption that  $g \in C(\mathbb{R}^d, \mathbb{R})$  ensure that for all  $\omega \in \Omega$  it holds that

$$\limsup_{n \rightarrow \infty} |f(\mathbf{x}_n + W_T(\omega), g(\mathbf{x}_n + W_T(\omega))) - f(\mathbf{x}_0 + W_T(\omega), g(\mathbf{x}_0 + W_T(\omega)))| = 0. \quad (183)$$

Moreover, note that the assumption that for all  $x \in \mathbb{R}^d$ ,  $a, b \in \mathbb{R}$  it holds that  $|f(x, a) - f(x, b)| \leq L|a - b|$  and the assumption that for all  $x \in \mathbb{R}^d$  it holds that  $\mathbb{E}[|f(x + W_T, 0)|^2 + |g(x + W_T)|^2] < \infty$  imply that

$$\mathbb{E}[|f(x + W_T, g(x + W_T))|^2] \leq 2\mathbb{E}[|f(x + W_T, 0)|^2] + 2\mathbb{E}[|g(x + W_T)|^2] < \infty. \quad (184)$$

Lemma 4.2 hence ensures that for all  $p \in (1, 2)$  it holds that  $\sup_{n \in \mathbb{N}} \mathbb{E}[|f(\mathbf{x}_n + W_T, g(\mathbf{x}_n + W_T))|^p] < \infty$ . Vitali's convergence theorem (see, e.g., Klenke [39, Theorem 6.25]), the de la Vallée Poussin theorem (see, e.g., Klenke [39, Theorem 6.19]), and (183) therefore demonstrate that

$$\limsup_{n \rightarrow \infty} \mathbb{E}[|f(\mathbf{x}_n + W_T, g(\mathbf{x}_n + W_T)) - f(\mathbf{x}_0 + W_T, g(\mathbf{x}_0 + W_T))|] = 0. \quad (185)$$

This completes the proof of Lemma 4.3.  $\square$

**Corollary 4.4.** Let  $L, T \in \mathbb{R}$ ,  $\Theta = \bigcup_{k \in \mathbb{N}} \mathbb{Z}^k$ ,  $f \in C(\mathbb{R}, \mathbb{R})$  satisfy for all  $a, b \in \mathbb{R}$  that  $|f(a) - f(b)| \leq L|a - b|$ , let  $\xi_d \in \mathbb{R}^d$ ,  $d \in \mathbb{N}$ , for every  $K \in \mathbb{N}$  let  $t_0^K, t_1^K, \dots, t_K^K \in \mathbb{R}$  satisfy  $0 = t_0^K < t_1^K < \dots < t_K^K = T$ , let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a probability space, let  $R^\theta: \Omega \rightarrow (0, 1)$ ,  $\theta \in \Theta$ , be i.i.d. random variables, assume for all  $r \in (0, 1)$  that  $\mathbb{P}(R^0 \leq r) = r$ , let  $W^{d, \theta}: [0, T] \times \Omega \rightarrow \mathbb{R}^d$ ,  $d \in \mathbb{N}$ ,  $\theta \in \Theta$ , be i.i.d. standard Brownian motions, assume that  $(R^\theta)_{\theta \in \Theta}$  and  $(W^{d, \theta})_{(d, \theta) \in \mathbb{N} \times \Theta}$  are independent, let  $v_k^{d, K} \in C(\mathbb{R}^d, \mathbb{R})$ ,  $k \in \{0, 1, \dots, K\}$ ,  $d, K \in \mathbb{N}$ , satisfy for all  $d, K \in \mathbb{N}$ ,  $k \in \{1, 2, \dots, K\}$ ,  $x \in \mathbb{R}^d$  that  $\mathbb{E}[|v_0^{d, K}(\xi_d + W_T^{d, 0})|^2] \leq L$ ,  $\mathbb{E}[|v_0^{d, K}(x + W_T^{d, 0})|^2] < \infty$ , and

$$v_k^{d, K}(x) = \mathbb{E}\left[v_{k-1}^{d, K}(x + W_{t_k^K - t_{k-1}^K}^{d, 0}) + (t_k^K - t_{k-1}^K)f(v_{k-1}^{d, K}(x + W_{t_k^K - t_{k-1}^K}^{d, 0}))\right] \quad (186)$$

(cf. Lemma 4.3), let  $\mathcal{R}^{K, \theta} = (\mathcal{R}_k^{K, \theta})_{k \in \{0, 1, \dots, K\}}: \{0, 1, \dots, K\} \times \Omega \rightarrow \mathbb{N}_0$ ,  $K \in \mathbb{N}$ ,  $\theta \in \Theta$ , satisfy for all  $K \in \mathbb{N}$ ,  $\theta \in \Theta$ ,  $k \in \{0, 1, \dots, K\}$  that  $\mathcal{R}_k^{K, \theta} = \max\{n \in \mathbb{N}_0: t_n^K \leq t_k^K R^\theta\}$ , let  $V_{k, n, M}^{d, K, \theta}: \mathbb{R}^d \times \Omega \rightarrow \mathbb{R}$ ,  $\theta \in \Theta$ ,  $k \in \{0, 1, \dots, K\}$ ,  $n \in \mathbb{N}_0$ ,  $d, K, M \in \mathbb{N}$ , satisfy for all  $\theta \in \Theta$ ,  $d, K, M \in \mathbb{N}$ ,  $n \in \mathbb{N}_0$ ,  $k \in \{0, 1, \dots, K\}$ ,  $x \in \mathbb{R}^d$  that

$$\begin{aligned} V_{k, n, M}^{d, K, \theta}(x) &= \frac{\mathbb{1}_{\mathbb{N}}(n)}{M^n} \sum_{m=1}^{M^n} \left[ v_0^{d, K}(x + W_{t_k^K}^{d, (\theta, 0, -m)}) + t_k^K f(0) \right] \\ &\quad + \sum_{j=1}^{n-1} \frac{t_k^K}{M^{n-j}} \sum_{m=1}^{M^{n-j}} \left[ f(V_{\mathcal{R}_k^{K, (\theta, j, m)}, j, M}^{d, K, (\theta, j, m)}(x + W_{t_k^K}^{d, (\theta, j, m)} - W_{\mathcal{R}_k^{K, (\theta, j, m)}}^{d, (\theta, j, m)})) \right. \\ &\quad \left. - f(V_{\mathcal{R}_k^{K, (\theta, j, m)}, j-1, M}^{d, K, (\theta, j, m)}(x + W_{t_k^K}^{d, (\theta, j, m)} - W_{\mathcal{R}_k^{K, (\theta, j, m)}}^{d, (\theta, j, m)})) \right], \end{aligned} \quad (187)$$

and let  $\mathfrak{C}_{M, n}^{d, K} \in \mathbb{R}$ ,  $d, K, M, n \in \mathbb{N}_0$ , satisfy for all  $d, K, M \in \mathbb{N}$ ,  $n \in \mathbb{N}_0$  that

$$\mathfrak{C}_{M, n}^{d, K} \leq dM^n + \sum_{j=1}^{n-1} M^{n-j}(d+1 + \mathfrak{C}_{M, j}^{d, K} + \mathfrak{C}_{M, j-1}^{d, K}). \quad (188)$$

Then there exist  $\mathfrak{N} = (\mathfrak{N}_\varepsilon)_{\varepsilon \in (0, 1]}: (0, 1] \rightarrow \mathbb{N}$  and  $\mathfrak{c} = (\mathfrak{c}_\delta)_{\delta \in (0, 1]}: (0, 1] \rightarrow \mathbb{R}$  such that for all  $\delta, \varepsilon \in (0, 1]$ ,  $d, K \in \mathbb{N}$  it holds that  $\mathfrak{C}_{\mathfrak{N}_\varepsilon, \mathfrak{N}_\varepsilon}^{d, K} \leq \mathfrak{c}_\delta d \varepsilon^{-(2+\delta)}$  and  $(\mathbb{E}[|V_{K, \mathfrak{N}_\varepsilon, \mathfrak{N}_\varepsilon}^{d, K, 0}(\xi_d) - v_K^{d, K}(\xi_d)|^2])^{1/2} \leq \varepsilon$ .

*Proof of Corollary 4.4.* Throughout this proof let  $\mathfrak{p}_{k,l}^K \in (0, \infty)$ ,  $k \in \{1, 2, \dots, K\}$ ,  $l \in \mathbb{N}_0$ ,  $K \in \mathbb{N}$ , satisfy for all  $K \in \mathbb{N}$ ,  $k \in \{1, 2, \dots, K\}$ ,  $l \in \mathbb{N}_0$  that

$$\mathfrak{p}_{k,l}^K = \begin{cases} \frac{t_{l+1}^K - t_l^K}{t_k^K} & : l < k \\ 1 & : l \geq k. \end{cases} \quad (189)$$

Observe that item (ii) of Theorem 4.1 (applied with  $\alpha \leftarrow 0$ ,  $\gamma \leftarrow 1$ ,  $c \leftarrow LT$ ,  $\kappa \leftarrow L + T|f(0)|$ ,  $p \leftarrow 1$ ,  $\Theta \leftarrow \Theta$ ,  $\xi_d \leftarrow \xi_d$ ,  $(L_k^{d,K})_{k \in \{0,1,\dots,K\}} \leftarrow (L(t_{\min\{k+1,K\}}^K - t_k^K))_{k \in \{0,1,\dots,K\}}$ ,  $(f_k^{d,K})_{k \in \{0,1,\dots,K\}} \leftarrow ((\mathbb{R}^d \times \mathbb{R} \ni (x, a) \mapsto a + (t_{\min\{k+1,K\}}^K - t_k^K)f(a) \in \mathbb{R}))_{k \in \{0,1,\dots,K\}}$ ,  $(\phi_k^{d,K})_{k \in \{0,1,\dots,K\}} \leftarrow (\mathbb{R}^d \times \mathbb{R}^d \ni (x, w) \mapsto x + w \in \mathbb{R}^d)_{k \in \{0,1,\dots,K\}}$ ,  $(\Omega, \mathcal{F}, \mathbb{P}) \leftarrow (\Omega, \mathcal{F}, \mathbb{P})$ ,  $(\mathfrak{C}_{M,n}^{d,K})_{(M,n) \in \mathbb{N} \times \mathbb{N}_0} \leftarrow (\mathfrak{C}_{M,n}^{d,K})_{(M,n) \in \mathbb{N} \times \mathbb{N}_0}$ ,  $(\mathcal{R}^{d,K,\theta})_{\theta \in \Theta} \leftarrow (\mathcal{R}^{K,\theta})_{\theta \in \Theta}$ ,  $(W_k^{d,K,\theta})_{(k,\theta) \in \{0,1,\dots,K\} \times \Theta} \leftarrow (W_{t_{\min\{k+1,K\}}^K}^{d,\theta} - W_{t_k^K}^{d,\theta})_{(k,\theta) \in \{0,1,\dots,K\} \times \Theta}$ ,  $(\mathfrak{p}_{k,l}^{d,K})_{(k,l) \in \{1,2,\dots,K\} \times \mathbb{N}_0} \leftarrow (\mathfrak{p}_{k,l}^K)_{(k,l) \in \{1,2,\dots,K\} \times \mathbb{N}_0}$ ,  $(v_k^{d,K})_{k \in \{0,1,\dots,K\}} \leftarrow (v_k^{d,K})_{k \in \{0,1,\dots,K\}}$  for  $d, K \in \mathbb{N}$  in the notation of Theorem 4.1) ensures that there exist  $\mathfrak{N} = (\mathfrak{N}_\varepsilon)_{\varepsilon \in (0,1]} : (0, 1] \rightarrow \mathbb{N}$  and  $\mathfrak{c} = (\mathfrak{c}_\delta) : (0, 1] \rightarrow [0, \infty)$  such that for all  $\delta, \varepsilon \in (0, 1]$ ,  $d, K \in \mathbb{N}$  it holds that  $\mathfrak{C}_{\mathfrak{N}_\varepsilon, \mathfrak{N}_\varepsilon}^{d,K} \leq \mathfrak{c}_\delta d \varepsilon^{-(2+\delta)}$  and

$$\left( \mathbb{E} \left[ |V_{K, \mathfrak{N}_\varepsilon, \mathfrak{N}_\varepsilon}^{d,K,0}(\xi_d) - v_K^{d,K}(\xi_d)|^2 \right] \right)^{1/2} \leq \varepsilon. \quad (190)$$

This completes the proof of Corollary 4.4.  $\square$

## Acknowledgements

The second author acknowledges funding by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy EXC 2044-390685587, Mathematics Muenster: Dynamics-Geometry-Structure.

## References

- [1] ADES, A., LU, G., AND CLAXTON, K. Expected value of sample information calculations in medical decision modeling. *Medical decision making* 24, 2 (2004), 207–227.
- [2] ANDERSEN, L. B. A simple approach to the pricing of bermudan swaptions in the multi-factor libor market model. *Available at SSRN 155208* (1999).
- [3] BECK, C., GONON, L., AND JENTZEN, A. Overcoming the curse of dimensionality in the numerical approximation of high-dimensional semilinear elliptic partial differential equations. *arXiv:2003.00596* (2020), 50 pages.
- [4] BECK, C., HORNING, F., HUTZENTHALER, M., JENTZEN, A., AND KRUSE, T. Overcoming the curse of dimensionality in the numerical approximation of Allen–Cahn partial differential equations via truncated full-history recursive multilevel Picard approximations. *Accepted by the Journal of Numerical Mathematics, arXiv:1907.06729* (2019), 31 pages.
- [5] BECKER, S., BRAUNWARTH, R., HUTZENTHALER, M., JENTZEN, A., AND VON WURSTEMBERGER, P. Numerical simulations for full history recursive multilevel picard approximations for systems of high-dimensional partial differential equations. *arXiv preprint arXiv:2005.10206* (2020), 21 pages.
- [6] BELOMESTNY, D. Pricing Bermudan options by nonparametric regression: optimal rates of convergence for lower estimates. *Finance Stoch.* 15, 4 (2011), 655–683.

- [7] BENDER, C., GÄRTNER, C., AND SCHWEIZER, N. Pathwise dynamic programming. *Math. Oper. Res.* 43, 3 (2018), 965–995.
- [8] BENDER, C., SCHWEIZER, N., AND ZHUO, J. A primal-dual algorithm for BSDEs. *Math. Finance* 27, 3 (2017), 866–901.
- [9] BOUCHARD, B., AND TOUZI, N. Discrete-time approximation and Monte-Carlo simulation of backward stochastic differential equations. *Stochastic Process. Appl.* 111, 2 (2004), 175–206.
- [10] BRATVOLD, R. B., BICKEL, J. E., LOHNE, H. P., ET AL. Value of information in the oil and gas industry: past, present, and future. *SPE Reservoir Evaluation & Engineering* 12, 04 (2009), 630–638.
- [11] BRENNAN, A., KHARROUBI, S., O’HAGAN, A., AND CHILCOTT, J. Calculating partial expected value of perfect information via Monte Carlo sampling algorithms. *Medical Decision Making* 27, 4 (2007), 448–470.
- [12] BROADIE, M., DU, Y., AND MOALLEMI, C. C. Efficient risk estimation via nested sequential simulation. *Management Science* 57, 6 (2011), 1172–1194.
- [13] BROADIE, M., DU, Y., AND MOALLEMI, C. C. Risk estimation via regression. *Oper. Res.* 63, 5 (2015), 1077–1097.
- [14] BROADIE, M., AND GLASSERMAN, P. Pricing American-style securities using simulation. *J. Econom. Dynam. Control* 21, 8-9 (1997), 1323–1352. Computational financial modelling.
- [15] BUJOK, K., HAMBLY, B. M., AND REISINGER, C. Multilevel simulation of functionals of Bernoulli random variables with application to basket credit derivatives. *Methodol. Comput. Appl. Probab.* 17, 3 (2015), 579–604.
- [16] CARRIERE, J. F. Valuation of the early-exercise price for options using simulations and nonparametric regression. *Insurance Math. Econom.* 19, 1 (1996), 19–30.
- [17] DIMITS, A. M., COHEN, B. I., CAFLISCH, R. E., ROSIN, M. S., AND RICKETSON, L. F. Higher-order time integration of Coulomb collisions in a plasma using Langevin equations. *J. Comput. Phys.* 242 (2013), 561–580.
- [18] E, W., HUTZENTHALER, M., JENTZEN, A., AND KRUSE, T. Multilevel Picard iterations for solving smooth semilinear parabolic heat equations. *arXiv:1607.03295* (2016), 19 pages.
- [19] E, W., HUTZENTHALER, M., JENTZEN, A., AND KRUSE, T. On multilevel Picard numerical approximations for high-dimensional nonlinear parabolic partial differential equations and high-dimensional nonlinear backward stochastic differential equations. *J. Sci. Comput.* 79, 3 (2019), 1534–1571.
- [20] EGLOFF, D. Monte Carlo algorithms for optimal stopping and statistical learning. *Ann. Appl. Probab.* 15, 2 (2005), 1396–1432.
- [21] EL KAROUI, N., KAPOUDJIAN, C., PARDOUX, E., PENG, S., AND QUENEZ, M. C. Reflected solutions of backward SDE’s, and related obstacle problems for PDE’s. *Ann. Probab.* 25, 2 (1997), 702–737.
- [22] GILES, M. B. MLMC for Nested Expectations. In *Contemporary Computational Mathematics-A Celebration of the 80th Birthday of Ian Sloan*. Springer, 2018, pp. 425–442.

- [23] GILES, M. B., AND GODA, T. Decision-making under uncertainty: using MLMC for efficient estimation of EVPPI. *Statistics and Computing* (2018), 1–13.
- [24] GILES, M. B., AND HAJI-ALI, A.-L. Multilevel nested simulation for efficient risk estimation. *arXiv:1802.05016* (2018), 28 pages.
- [25] GILES, M. B., JENTZEN, A., AND WELTI, T. Generalised multilevel Picard approximations. *arXiv:1911.03188* (2019), 61 pages.
- [26] GLASSERMAN, P., HEIDELBERGER, P., AND SHAHABUDDIN, P. Variance reduction techniques for estimating Value-at-Risk. *Management Science* 46, 10 (2000), 1349–1364.
- [27] GLASSERMAN, P., HEIDELBERGER, P., AND SHAHABUDDIN, P. Portfolio Value-at-Risk with heavy-tailed risk factors. *Mathematical Finance* 12, 3 (2002), 239–269.
- [28] GOBET, E., LEMOR, J.-P., AND WARIN, X. A regression-based Monte Carlo method to solve backward stochastic differential equations. *Ann. Appl. Probab.* 15, 3 (2005), 2172–2202.
- [29] GODA, T., HIRONAKA, T., AND IWAMOTO, T. Multilevel Monte Carlo estimation of expected information gains. *Stochastic Analysis and Applications* 38, 4 (2020), 581–600.
- [30] GODA, T., MURAKAMI, D., TANAKA, K., AND SATO, K. Decision-theoretic sensitivity analysis for reservoir development under uncertainty using multilevel quasi-Monte Carlo methods. *Computational Geosciences* 22, 4 (2018), 1009–1020.
- [31] GOLDBERG, D. A., AND CHEN, Y. Beating the curse of dimensionality in options pricing and optimal stopping. *arXiv:1807.02227* (2018), 62 pages.
- [32] GORDY, M. B., AND JUNEJA, S. Nested simulation in portfolio risk measurement. *Management Science* 56, 10 (2010), 1833–1848.
- [33] HAJI-ALI, A.-L., AND TEMPONE, R. Multilevel and multi-index Monte Carlo methods for the McKean-Vlasov equation. *Stat. Comput.* 28, 4 (2018), 923–935.
- [34] HIRONAKA, T., GILES, M. B., GODA, T., AND THOM, H. Multilevel Monte Carlo estimation of the expected value of sample information. *arXiv:1909.00549* (2019), 26 pages.
- [35] HUTZENTHALER, M., JENTZEN, A., AND KRUSE, T. Overcoming the curse of dimensionality in the numerical approximation of parabolic partial differential equations with gradient-dependent nonlinearities. *arXiv:1912.02571* (2019), 33 pages.
- [36] HUTZENTHALER, M., JENTZEN, A., KRUSE, T., NGUYEN, T. A., AND VON WURSTEMBERGER, P. Overcoming the curse of dimensionality in the numerical approximation of semilinear parabolic partial differential equations. *Accepted by Proc. Roy. Soc. London A*, *arXiv:1807.01212* (2018), 30 pages.
- [37] HUTZENTHALER, M., JENTZEN, A., AND VON WURSTEMBERGER, P. Overcoming the curse of dimensionality in the approximative pricing of financial derivatives with default risks. *Electronic Journal of Probability* 25 (2020).
- [38] HUTZENTHALER, M., AND KRUSE, T. Multilevel Picard Approximations of High-Dimensional Semilinear Parabolic Differential Equations with Gradient-Dependent Nonlinearities. *SIAM J. Numer. Anal.* 58, 2 (2020), 929–961.

- [39] KLENKE, A. *Probability theory*, second ed. Universitext. Springer, London, 2014. A comprehensive course.
- [40] KORN, R., KORN, E., AND KROISANDT, G. *Monte Carlo methods and models in finance and insurance*. Chapman & Hall/CRC Financial Mathematics Series. CRC Press, Boca Raton, FL, 2010.
- [41] LONGSTAFF, F. A., AND SCHWARTZ, E. S. Valuing American options by simulation: a simple least-squares approach. *The review of financial studies* 14, 1 (2001), 113–147.
- [42] NAKAYASU, M., GODA, T., TANAKA, K., SATO, K., ET AL. Evaluating the value of single-point data in heterogeneous reservoirs with the expectation-maximization algorithm. *SPE Economics & Management* 8, 01 (2016), 1–10.
- [43] PARDOUX, É., AND PENG, S. Adapted solution of a backward stochastic differential equation. *Systems Control Lett.* 14, 1 (1990), 55–61.
- [44] PARDOUX, E., AND PENG, S. Backward stochastic differential equations and quasilinear parabolic partial differential equations. In *Stochastic partial differential equations and their applications (Charlotte, NC, 1991)*, vol. 176 of *Lect. Notes Control Inf. Sci.* Springer, Berlin, 1992, pp. 200–217.
- [45] PENG, S. Nonlinear expectations, nonlinear evaluations and risk measures. In *Stochastic methods in finance*, vol. 1856 of *Lecture Notes in Math.* Springer, Berlin, 2004, pp. 165–253.
- [46] ROSIN, M., RICKETSON, L., DIMITS, A. M., CAFLISCH, R. E., AND COHEN, B. I. Multilevel Monte Carlo simulation of Coulomb collisions. *Journal of Computational Physics* 274 (2014), 140–157.
- [47] SZPRUCH, L., TAN, S., AND TSE, A. Iterative particle approximation for McKean-Vlasov sdes with application to multilevel Monte Carlo estimation. *arXiv preprint arXiv:1706.00907* (2017).
- [48] TSITSIKLIS, J. N., AND VAN ROY, B. Optimal stopping of Markov processes: Hilbert space theory, approximation algorithms, and an application to pricing high-dimensional financial derivatives. *IEEE Trans. Automat. Control* 44, 10 (1999), 1840–1851.
- [49] TSITSIKLIS, J. N., AND VAN ROY, B. Regression methods for pricing complex american-style options. *IEEE Transactions on Neural Networks* 12, 4 (2001), 694–703.