Quantum vacuum effects in braneworlds on AdS bulk

A. A. Saharian^{*}

Department of Physics, Yerevan State University, 1 Alex Manoogian Street, 0025 Yerevan, Armenia

January 13, 2022

Abstract

We review the results of investigations for brane-induced effects on the local properties of quantum vacuum in background of AdS spacetime. Two geometries are considered: a brane parallel to the AdS boundary and a brane intersecting the AdS boundary. For both these cases the contribution in the vacuum expectation value (VEV) of the energy-momentum tensor is separated explicitly and its behavior in various asymptotic regions of the parameters is studied. It is shown that the influence of the gravitational field on the local properties of the quantum vacuum is essential at distance from the brane larger than the AdS curvature radius. In the geometry with a brane parallel to the AdS boundary the VEV of the energy-momentum tensor is considered for scalar field with the Robin boundary condition, for Dirac field with the bag boundary condition and for the electromagnetic field. In the latter case two types of boundary conditions are discussed. The first one is a generalization of the perfect conductor boundary condition and the second one corresponds to the confining boundary condition used in QCD for gluons. For the geometry of a brane intersecting the AdS boundary, the case of a scalar field is considered. The corresponding energy-momentum tensor, apart from the diagonal components, has nonzero off-diagonal component. As a consequence of the latter, in addition to the normal component, the Casimir force acquires a component parallel to the brane.

1 Introduction

Quantum field theory in anti-de Sitter (AdS) spacetime is an active area of research. This activity is motivated by several reasons. First of all, the corresponding geometry is maximally symmetric and a sufficiently large number of problems are exactly solvable on its background. This helps to give an idea of the influence of the classical gravitational field on quantum phenomena in more complicated geometries. Qualitatively new features in the dynamics of quantum fields on the AdS background are related to the lack of global hyperbolicity and to the existence of both regular and irregular modes. In particular, boundary conditions on propagating fields need to be imposed at timelike infinity to prevent loss of unitarity. The different boundary conditions define different theories in the bulk. Another new feature, that has no analogue in Minkowskian field theories, is related to the possibility of regularization for infrared divergences in interacting field theories without reducing the symmetries [1]. This is closely related to the natural length scale of the AdS spacetime. The high interest to the AdS geometry is also related to its natural appearance as a ground state in supergravity and as the near horizon geometry for extremal black holes, black strings and domain walls.

The AdS spacetime plays a fundamental role in two exciting developments of contemporary theoretical physics. The first one, the AdS/CFT correspondence (for reviews see [2, 3, 4]), establishes

^{*}E-mail: saharian@ysu.am

duality between string theories or supergravity in the AdS bulk and a conformal field theory localized on the AdS boundary. This duality provides an interesting possibility for the investigation of nonperturbative effects in both sides of the correspondence by using the weak coupling regime of the dual theory. The recent developments include applications in various condensed matter systems such as holographic superconductors and topological insulators [5, 6]. The second development is related to various types of braneworld models with large extra dimensions [7]. In the corresponding setup the standard model fields are localized on a brane embedded in a higher dimensional AdS spacetime. Braneworlds naturally appear in string/M-theory context and have been initially proposed for a geometrical resolution of the hierarcy problem between the electroweak and Planck energy scales. They provide an alternative framework to address the problems in particle physics and cosmology from different perspectives.

An inherent feature of field theories in AdS/CFT correspondence and in braneworld models is the need to impose boundary conditions on fileds propagating in the AdS bulk. They include the conditions on the AdS boundary and the conditions on the branes in braneworld scenario. In particular, in braneworld models of the Randall-Sundrum type the boundary conditions on the branes are dictated by the Z_2 -symmetry. In quantum field theory, the boundary conditions modify the spectrum of the zero-point fluctuations and, as a consequence, the vacuum expectation values (VEVs) are shifted by an amount that depends on the bulk and boundary geometries and also on the boundary conditions. This is the well-known Casimir effect [8]-[11]. In braneworld models the Casimir forces acting on the branes may provide a mechanism for stabilization of the brane location. In particular, this stabilization is required to prevent the variations of physical constants on the branes. In addition, the quantum effects of bulk fields generate a cosmological constant on the brane. Motivated by these points, the Casimir effect in braneworld models on the AdS bulk has been investigated in the large number of papers (see, for instance, references given in [12, 13]).

The main part of the papers on the Casimir effect in the AdS bulk consider global quantities, such as the Casimir energy and the forces acting on the branes. The local quantities carry more detailed information about the properties of the quantum vacuum. In particular, the expectation value of the energy-momentum tensor is of special interest. It appears as the source in the semiclassical Einstein equations and therefore plays an important role in modelling self-consistent dynamics involving the gravitational field. The VEV of the energy-momentum tensor for a conformally coupled scalar field in confromally-flat geometries has been investigated in [14]. Massive scalar fields with general curvature coupling in the geometry of two branes on AdS bulk were considered in [15, 16]. The Casimir densities for a Z_2 -symmetric thick brane for the general case of static plane symmetric interior structure have been discussed in [17]. The VEVs of the energy-momentum tensor for Dirac spinor field and for the electromagnetic field are investigated in [18, 19] and [13, 20, 21, 22]. The geometry of a brane intersecting the AdS boundary has been considered in [23]. For a scalar field with general curvature coupling, the background AdS geometry with an additional compact subspace is discussed in [24, 25] (for the Casimir energy in those models see [26, 27]).

In the present paper we review the results for the VEV of the energy-momentum tensor in the geometry of a single brane on background of (D + 1)-dimensional AdS spacetime. The cases of scalar, Dirac and electromagnetic fields will be considered. The organization of the paper is as follows. In section 2, for a planar brane parallel to the AdS boundary, we consider complete sets of orthonormalized mode functions for both the regions between the AdS boundary and the brane and between the brane and the AdS horizon. The VEVs of the energy-momentum tensors for scalar, Dirac and electromagnetic fields are investigated in section 3. The behavior in the asymptotic regions of the parameters is discussed in detail. Section 4 is devoted to the study of the vacuum energy-momentum tensor for a scalar field in the geometry with a brane perpendicular to the AdS boundary. The main results are summarized in section 5.

2 Mode functions in the geometry with a brane parallel to the AdS boundary

In Poincaré coordinates $(x^0 = t, x^1, \dots, x^{D-1}, y)$, with $-\infty < x^i, y < +\infty, i = 0, 1, \dots, D-1$, the line element for the (D+1)-dimensional AdS spacetime is presented as

$$ds^{2} = e^{-2y/a} \left[\left(dx^{0} \right)^{2} - \left(dx^{1} \right)^{2} - \dots - \left(dx^{D-1} \right)^{2} \right] - dy^{2}, \tag{2.1}$$

where a is the curvature radius of the background geometry sourced by a negative cosmological constant $\Lambda = -D(D-1)/(2a^2)$. For the curvature scalar and the Ricci tensor one has $R = -D(D+1)/a^2$ and $R_{\mu\rho} = -Dg_{\mu\rho}/a^2$, with the metric tensor $g_{\mu\rho}$ defined from (2.1). Introducing a new coordinate z, $0 \leq z < \infty$, in accordance with $z = ae^{y/a}$, the line element is written in conformally flat form

$$ds^{2} = g_{\mu\rho}dx^{\mu}dx^{\rho} = (a/z)^{2}\eta_{\mu\rho}dx^{\mu}dx^{\rho}, \qquad (2.2)$$

where $x^D = z$ and $\eta_{\mu\rho} = \text{diag}(1, -1, \dots, -1)$ is the metric tensor for (D+1)-dimensional Minkowski spacetime. The hypersurfaces z = 0 and $z = \infty$ correspond to the AdS boundary and horizon, respectively. In what follows we will work in the coordinate system defined by (2.2).

We are interested in the effects on the local properties of the quantum vacuum induced by a codimension one brane. First we consider the case where the brane is parallel to the AdS boundary and is located at $z = z_0$. It divides the space into two subspaces: the region between the AdS boundary and the brane, $0 \le z \le z_0$ (L-region) and the region between the brane and AdS horizon, $z_0 \le z < \infty$ (R-region). The brane has a nonzero extrinsic curvature tensor and, as a consequence, the properties of the vacuum state in the L- and R-regions are different. The evaluation of the VEVs for local physical observables in those regions requires different procedures and we will discuss them separately. The VEVs are presented in the form of mode-sums over complete set of mode functions for quantum fields and we start by considering the modes for scalar, Dirac and electromagnetic fields.

2.1 Scalar field

First we consider a scalar field $\varphi(x)$ with the mass m. Assuming a general curvature coupling with the parameter ξ , the field equation reads

$$\left(g^{\mu\rho}\nabla_{\mu}\nabla_{\rho} + m^2 + \xi R\right)\varphi(x) = 0, \qquad (2.3)$$

where ∇_{μ} is the covariant derivative operator. The most important special cases correspond to minimally ($\xi = 0$) and conformally ($\xi = \xi_D = (D-1)/(4D)$) coupled fields. Let $\varphi_{\sigma}^{(\pm)}(x)$ be a complete set of positive and negative energy mode functions specified by the set of quantum numbers σ . In accordance with the problem symmetry, they can be presented in the form

$$\varphi_{\sigma}(x) = e^{i\mathbf{k}\mathbf{x}\mp i\omega t} f(z), \qquad (2.4)$$

where $\mathbf{x} = (x^1, x^2, \dots, x^{D-1})$, $\mathbf{k} = (k_1, k_2, \dots, k_{D-1})$, and $\mathbf{k}\mathbf{x} = \sum_{l=1}^{D-1} k_l x^l$. Plugging into the field equation we get an ordinary differential equation for the function f(z):

$$z^{D+1}\partial_z \left[z^{1-D}\partial_z f(z) \right] + \left[\lambda^2 z^2 + \left(\xi D(D+1) - a^2 m^2 \right) \right] f(z) = 0,$$
(2.5)

where the energy ω is expressed in terms of λ and $k^2 = \sum_{i=1}^{D-1} k_i^2$ as $\omega = \sqrt{\lambda^2 + k^2}$. The general solution of (2.5) is expressed in terms of the Bessel and Neumann functions $J_{\nu}(\lambda z)$ and $Y_{\nu}(\lambda z)$:

$$f(z) = z^{D/2} \left[c_1 J_{\nu}(\lambda z) + c_2 Y_{\nu}(\lambda z) \right], \qquad (2.6)$$

where c_1 and c_2 are constants and

$$\nu = \sqrt{D^2/4 - D(D+1)\xi + m^2 a^2}.$$
(2.7)

In order to have a stable vacuum we assume the values of the parameters in (2.7) for which $\nu \ge 0$ (see [28, 29]). For a conformally coupled massless field one has $\nu = 1/2$ and the mode functions (2.4) with (2.6) are conformally related to the modes in the Minkowski bulk. The scalar modes are normalized by the condition

$$\int d^D x \, g^{00} \sqrt{|g|} \varphi_{\sigma}^{(s)}(x) \varphi_{\sigma'}^{(s')*}(x) = \frac{\delta_{ss'}}{2\omega} \delta_{\lambda\lambda'} \delta(\mathbf{k} - \mathbf{k}'), \tag{2.8}$$

where g is the determinant of the metric tensor. Here $\delta_{\lambda\lambda'}$ is the Kronecker delta in the problems with discrete eigenvalues of the quantum number λ and $\delta_{\lambda\lambda'} = \delta(\lambda - \lambda')$ in the problems with continuous spectrum for λ .

We are interested in the effects of a codimension one brane, localized at $z = z_0$, on the local properties of the scalar vacuum. The Robin boundary condition will be imposed for the field operator on the brane:

$$(\beta n^{\mu} \nabla_{\mu} + 1) \varphi(x) = 0, \ z = z_0,$$
(2.9)

where n^{μ} is the inward pointing normal to the brane and β is a constant. The latter encodes the properties of the brane. In the special cases $\beta = 0$ and $\beta = \infty$ the condition (2.9) is reduced to the Dirichlet and Neumann boundary conditions, respectively. For the normal in (2.9) one has $n^{\mu} = \delta_{(J)} \delta^{\mu}_{D} z/a$, where J=L, $\delta_{(L)} = -1$ in the L-region and J=R, $\delta_{(R)} = 1$ in the R-region. In general, the values of the constant β could be different for those regions.

First let us consider the modes in the R-region. From the boundary condition (2.9) it follows that $c_2/c_1 = -\bar{J}_{\nu}(\lambda z_0)/\bar{Y}_{\nu}(\lambda z_0)$ for the coefficients in (2.6). Here and below, for a given function F(x), we use the notation with the bar defined in accordance with

$$\bar{F}(x) = B_0 x F'(x) + A_0 F(x), \qquad (2.10)$$

with the coefficients

$$A_0 = 1 + \delta_{(J)} \frac{D\beta}{2a}, \quad B_0 = \delta_{(J)} \frac{\beta}{a}.$$
 (2.11)

The mode functions in the R-region obeying the boundary condition (2.9) are presented as

$$\varphi_{(\mathrm{R})\sigma}^{(\pm)}(x) = C_{(\mathrm{R})\sigma} z^{D/2} g_{\nu}(\lambda z_0, \lambda z) e^{i\mathbf{k}\mathbf{x}\mp i\omega t}, \qquad (2.12)$$

with the function

$$g_{\nu}(u,v) = J_{\nu}(v)\bar{Y}_{\nu}(u) - \bar{J}_{\nu}(u)Y_{\nu}(v).$$
(2.13)

The spectrum for λ is continuous and from the normalization condition (2.8) with $\delta_{\lambda\lambda'} = \delta(\lambda - \lambda')$ we get

$$|C_{(R)\sigma}|^{2} = \frac{(2\pi a)^{1-D} \lambda}{2\omega \left[\bar{J}_{\nu}^{2}(\lambda z_{0}) + \bar{Y}_{\nu}^{2}(\lambda z_{0})\right]}.$$
(2.14)

The modes are specified by the set $\sigma = (\mathbf{k}, \lambda)$ with $-\infty < k_i < +\infty$, $i = 1, \ldots, D-1$, and $0 \le \lambda < \infty$. The analog of the mode functions (2.12) in the region between two parallel branes on the AdS bulk has been considered in [16].

Note that we could also have modes with purely imaginary λ , $\lambda = i|\lambda|$. For those modes $f(z) = c_1 z^{D/2} K_{\nu}(|\lambda|z)$, where $K_{\nu}(x)$ is the Macdonald function (the modes with the modified Bessel function $I_{\nu}(|\lambda|z)$ are not normalizable). From the boundary condition (2.9) we get the equation for the allowed values of $|\lambda|$: $\bar{K}_{\nu}(|\lambda|z_0) = 0$. The energy corresponding to these modes is given by $\omega = \sqrt{k^2 - |\lambda|^2}$ and it becomes imaginary for $k < |\lambda|$. This leads to the instability of the vacuum state. In order to exclude the unstable modes we restrict the allowed values for the Robin coefficient in the region

where the equation $\bar{K}_{\nu}(|\lambda|z_0) = 0$ has no roots. It can be seen that for non-Dirichlet boundary conditions $(\beta \neq 0)$ the corresponding condition is expressed as $a/\beta < \nu - D/2$ (for more detailed discussion in models with compact dimensions see [30]). For $a/\beta > \nu - D/2$ there is a single root. For the special value $\beta/a = 1/(\nu - D/2)$ there exists a mode with $\lambda = 0$ and with the mode functions $\varphi_{(R)\sigma}^{(\pm)}(x) = C_{(R)\sigma}z^{D/2-\nu}e^{i\mathbf{kx}\mp ikt}$. For a minimally coupled massless scalar field one has $\nu = D/2$ and this special value corresponds to the Neumann boundary condition. The corresponding mode functions do not depend on z.

In the L-region the integration over z in (2.8) goes over the interval $z \in [0, z_0]$. For the solutions (2.4) with (2.6) and $c_2 \neq 0$, the z-integral diverges at the lower limit z = 0 in the range $\nu \geq 1$. Hence, in that range, from the normalizability condition for the mode functions it follows that we should take $c_2 = 0$. In the range $0 \leq \nu < 1$, the solution (2.6) with $c_2 \neq 0$ is normalizable and in order to uniquely define the mode functions an additional boundary condition at the AdS boundary is required [28, 31]. The general class of allowed boundary conditions has been discussed in [32, 33]. In particular, they include the Dirichlet and Neumann boundary conditions, the most frequently used in the literature. Here, for the values of the parameters corresponding to the range $0 \leq \nu < 1$ we will choose the Dirichlet condition which gives $c_2 = 0$. With this choice, the mode functions in the L-region are specified as

$$\varphi_{(\mathrm{L})\sigma}^{(\pm)}(x) = C_{(\mathrm{L})\sigma} z^{D/2} J_{\nu}(\lambda z) e^{i\mathbf{k}\mathbf{x}\mp i\omega t}.$$
(2.15)

From the boundary condition (2.9) on the brane we get the equation for the eigenvalues of the quantum number λ :

$$\bar{J}_{\nu}(\lambda z_0) = 0. \tag{2.16}$$

If we denote by $x = \lambda_{\nu,n}$ the positive zeros of the functions $\bar{J}_{\nu}(x)$, numerated by n = 1, 2, ..., then the eigenvalues are given by $\lambda = \lambda_{\nu,n}/z_0$. Note that the roots $\lambda_{\nu,n}$ do not depend on the location of the brane. From the normalization condition (2.8), with $\delta_{\lambda\lambda'} = \delta_{nn'}$ and with the z-integartion over $[0, z_0]$, one finds

$$|C_{\sigma}|^{2} = \frac{(2\pi a)^{1-D} \lambda_{\nu,n} T_{\nu}(\lambda_{\nu,n})}{z_{0} \sqrt{k^{2} z_{0}^{2} + \lambda_{\nu,n}^{2}}},$$
(2.17)

with the function $T_{\nu}(x) = x[x^2 J_{\nu}'^2(x) + (x^2 - \nu^2) J_{\nu}^2(x)]^{-1}.$

Similar to the R-region, the stability condition for the vacuum state in the L-region imposes restrictions to the allowed values of the Robin coefficient β . That condition excludes the presence of purely imaginary roots $\lambda = i|\lambda|$ for the equation (2.16). It can be shown that there are no such modes for $a/\beta < D/2 + \nu$ and there is a single mode for $a/\beta > D/2 + \nu$. As seen, the stability condition in the L-region is less restrictive than that for the R-region. In the special case $\beta/a = 1/(D/2 + \nu)$ one has a mode with $\lambda = 0$ with the mode functions $\varphi_{(L)\sigma}^{(\pm)}(x) = C_{(L)}z^{D/2+\nu}e^{i\mathbf{kx}\mp ikt}$.

2.2 Dirac field

Now we turn to a massive Dirac field $\psi(x)$. The dynamics of the field is governed by the Dirac equation

$$[i\gamma^{\mu}\left(\partial_{\mu}+\Gamma_{\mu}\right)-m]\psi(x)=0, \qquad (2.18)$$

where Γ_{μ} is the spin connection. The curved spacetime Dirac matrices γ^{μ} are expressed in terms of the flat spacetime matrices $\gamma^{(b)}$ by the relation $\gamma^{\mu} = e^{\mu}_{(b)}\gamma^{(b)}$, with $e^{\mu}_{(b)}$ being the tetrad fields. In the coordinate system corresponding to (2.2) the latter can be taken as $e^{\mu}_{(b)} = (z/a)\delta^{\mu}_{b}$. For the components of the spin connection this gives $\Gamma_{D} = 0$ and $\Gamma_{i} = \eta_{il}\gamma^{(D)}\gamma^{(l)}/(2z)$ for $i = 0, \ldots, D-1$. In an irreducible representation of the Clifford algebra the matrices $\gamma^{(b)}$ are $N \times N$ matrices, where $N = 2^{[(D+1)/2]}$ and the square brackets in the exponent mean the integer part. Up to a similarity transformation, the irreducible representation is unique in odd numbers of spatial dimension D. For even values of D one has two inequivalent irreducible representations. We use the flat spacetime gamma matrices in the representation

$$\gamma^{(0)} = \begin{pmatrix} 0 & \chi_0 \\ \chi_0^{\dagger} & 0 \end{pmatrix}, \ \gamma^{(l)} = \begin{pmatrix} 0 & \chi_l \\ -\chi_l^{\dagger} & 0 \end{pmatrix}, \ l = 1, 2, \dots, D - 1,$$
(2.19)

and $\gamma^{(D)} = si \operatorname{diag}(1, -1)$, where $s = \pm 1$. In odd spatial dimensions the two values of the parameter s correspond to two inequivalent representations. The commutation relations for the $N/2 \times N/2$ matrices χ_l and for their hermitian conjugate matrices χ_l^{\dagger} are obtained from those for the Dirac matrices $\gamma^{(l)}$. They are reduced to the relations $\chi_0 \chi_l^{\dagger} = \chi_l \chi_0^{\dagger}$, $\chi_0^{\dagger} \chi_l = \chi_l^{\dagger} \chi_0$, $\chi_0^{\dagger} \chi_0 = 1$ and $\chi_l \chi_i^{\dagger} + \chi_i \chi_l^{\dagger} = 2\delta_{li}$, $\chi_l^{\dagger} \chi_i + \chi_i^{\dagger} \chi_l = 2\delta_{li}$, for $l, i = 1, 2, \ldots, D - 1$. The representation (2.19) for the construction of the gamma matrices in AdS spacetime has been considered in [34]. Another representation is taken in [19].

Assuming the dependence on the coordinates (t, \mathbf{x}) in the form $e^{i\mathbf{kx}\mp i\omega t}$ and decomposing the spinor $\psi(x)$ into the upper and lower components, in the representation (2.19) the corresponding equations are separated. The dependence of those components on the z-coordinate is expressed in terms of the function $c_1 J_{ma\pm s/2}(\lambda z) + c_2 Y_{ma\pm s/2}(\lambda z)$, where the upper and lower signs correspond to the upper and lower components. The coefficients are determined by the normalization condition and by the boundary condition on the brane at $z = z_0$. The positive and negative energy fermionic modes $\psi_{\sigma}^{(\pm)}(x)$, specified by the set of quantum numbers σ , are normalized by the condition

$$\int d^D x \, (a/z)^D \psi_{\sigma}^{(\pm)\dagger} \psi_{\sigma'}^{(\pm)} = \delta_{\sigma\sigma'}.$$
(2.20)

As in the case of a scalar field, here $\delta_{\sigma\sigma'}$ is understood as the Dirac delta function for the continuous components of σ and the Kronecker delta for discrete ones. On the brane at $z = z_0$ we impose the bag boundary condition

$$(1+i\gamma^{\mu}n_{\mu})\psi(x) = 0, \ z = z_0, \tag{2.21}$$

where $n_{\mu} = -\delta_{(J)}\delta^{D}_{\mu}a/z$, J=R,L, with $\delta_{(R)} = -\delta_{(L)} = 1$ for the L- and R-regions.

In the R-region, $z_0 \leq z < \infty$, from the boundary condition (2.21), for the ratio of the coefficients in the linear combination of the Bessel and Neumann functions one finds $c_2/c_1 = -J_{ma+1/2}(\lambda z_0)/Y_{ma+1/2}(\lambda z_0)$. The positive and negative energy mode functions, obeying the boundary condition (2.21), are expressed as

$$\psi_{(\mathrm{R})\sigma}^{(+)}(x) = C_{(\mathrm{R})\sigma}^{(+)} z^{\frac{D+1}{2}} e^{i\mathbf{k}\mathbf{x}-i\omega t} \left(\begin{array}{c} \frac{\mathbf{k}\chi\chi_{0}^{\dagger}+i\lambda-\omega}{\omega} g_{ma+1/2,ma+s/2}(\lambda z_{0},\lambda z)w^{(\gamma)} \\ i\chi_{0}^{\dagger} \frac{\mathbf{k}\chi\chi_{0}^{\dagger}+i\lambda+\omega}{\omega} g_{ma+1/2,ma-s/2}(\lambda z_{0},\lambda z)w^{(\gamma)} \end{array} \right),$$

$$\psi_{(\mathrm{R})\sigma}^{(-)}(x) = C_{(\mathrm{R})\sigma}^{(-)} z^{\frac{D+1}{2}} e^{i\mathbf{k}\mathbf{x}+i\omega t} \left(\begin{array}{c} i\chi_{0} \frac{\mathbf{k}\chi^{\dagger}\chi_{0}-i\lambda+\omega}{\omega} g_{ma+1/2,ma+s/2}(\lambda z_{0},\lambda z)w^{(\gamma)} \\ \frac{\mathbf{k}\chi^{\dagger}\chi_{0}-i\lambda-\omega}{\omega} g_{ma+1/2,ma-s/2}(\lambda z_{0},\lambda z)w^{(\gamma)} \end{array} \right), \quad (2.22)$$

where $\mathbf{k}\chi = \sum_{l=1}^{D-1} k_l \chi_l$ and

$$g_{\mu,\rho}(x,u) = J_{\mu}(x)Y_{\rho}(u) - J_{\rho}(u)Y_{\mu}(x).$$
(2.23)

In (2.22), the one-column matrices $w^{(\gamma)}$, $\gamma = 1, \ldots, N/2$, are introduced with the elements $w_l^{(\sigma)} = \delta_{l\sigma}$ and having N/2 rows. The normalization constant is determined from the condition (2.20):

$$\left|C_{(\mathrm{R})\sigma}^{(\pm)}\right|^{2} = \lambda \frac{\left[J_{ma+1/2}^{2}(\lambda z_{0}) + Y_{ma+1/2}^{2}(\lambda z_{0})\right]^{-1}}{4\left(2\pi\right)^{D-1}a^{D}}.$$
(2.24)

The set of quantum numbers σ is specified as $\sigma = (\mathbf{k}, \lambda, \gamma)$.

In the L-region and for $ma \ge 1/2$ from the normalizability condition we get $c_2 = 0$. In the range of the mass corresponding to ma < 1/2 the modes with $c_2 \ne 0$ are normalizable as well and an additional boundary condition is required to uniquely define the mode functions. Here we consider a special case that corresponds to the choice $c_2 = 0$ for all values of the mass. The fermionic modes are given by the expressions

$$\psi_{(\mathrm{L})\sigma}^{(+)}(x) = C_{(\mathrm{L})\sigma}^{(+)} z^{\frac{D+1}{2}} e^{i\mathbf{k}\mathbf{x}-i\omega t} \begin{pmatrix} \frac{\mathbf{k}\chi\chi_{0}^{\dagger}+i\lambda-\omega}{\omega} J_{ma+s/2}(\lambda z)w^{(\gamma)} \\ i\chi_{0}^{\dagger}\frac{\mathbf{k}\chi\chi_{0}^{\dagger}+i\lambda+\omega}{\omega} J_{ma-s/2}(\lambda z)w^{(\gamma)} \end{pmatrix},$$

$$\psi_{(\mathrm{L})\sigma}^{(-)}(x) = C_{(\mathrm{L})\sigma}^{(-)} z^{\frac{D+1}{2}} e^{i\mathbf{k}\mathbf{x}+i\omega t} \begin{pmatrix} i\chi_{0}\frac{\mathbf{k}\chi^{\dagger}\chi_{0}-i\lambda+\omega}{\omega} J_{ma+s/2}(\lambda z)w^{(\gamma)} \\ \frac{\mathbf{k}\chi^{\dagger}\chi_{0}-i\lambda-\omega}{\omega} J_{ma-s/2}(\lambda z)w^{(\gamma)} \end{pmatrix}.$$
(2.25)

The allowed values of λ are determined from the boundary condition (2.21) on the brane. They are roots of the equation

$$J_{ma-1/2}(\lambda z_0) = 0, (2.26)$$

and are expressed as $\lambda = \lambda_{ma-1/2,n}/z_0$. The normalization coefficient is obtained from (2.20) and is given by

$$|C_{(\mathrm{L})\sigma}^{(\pm)}|^{2} = \frac{J_{ma+1/2}^{-2}(\lambda_{ma-1/2,n})}{2(2\pi)^{D-1}a^{D}z_{0}^{2}}.$$
(2.27)

Note that the eigenvalues for λ are the same for both the representations of the Clifford algebra.

2.3 Electromagnetic field

For the electromagnetic field with the vector potential $A_{\mu}(x)$, $\mu = 0, 1, \ldots, D$, the field equation reads

$$\nabla_{\rho}F^{\mu\rho} = 0, \qquad (2.28)$$

where $F_{\mu\rho} = \partial_{\mu}A_{\rho} - \partial_{\rho}A_{\mu}$ is the field strength tensor. In order to find a complete set of mode functions $A_{\sigma\mu}(x)$ for the vector potential we impose the Lorentz condition $\nabla_{\mu}A^{\mu} = 0$ and an additional gauge condition $A^{D} = 0$. For the positive energy modes, presenting the dependence on the coordinates $(t, x^{1}, \ldots, x^{D-1})$ in the form $e^{i\mathbf{kx}-i\omega t}$, from the field equation we can see that

$$A_{\sigma\mu}(x) = \epsilon_{(\gamma)\mu} z^{D/2-1} \left[c_1 J_{D/2-1}(\lambda z) + c_2 Y_{D/2-1}(\lambda z) \right] e^{i\mathbf{kx} - i\omega t}.$$
 (2.29)

Here, $\gamma = 1, \ldots, D-1$ correspond to different polarizations specified by the polarization vector $\epsilon_{(\sigma)\mu}$. For the latter one has the normalization condition $\eta^{\mu\rho}\epsilon_{(\gamma)\mu}\epsilon_{(\gamma')\rho} = -\delta_{\gamma\gamma'}$ and the constraints $\epsilon_{(\gamma)D} = 0$ and $\eta^{\mu\rho}k_{\mu}\epsilon_{(\gamma)\rho} = 0$. The latter two relations follow from the gauge conditions. The modes (2.29) are normalized by the condition

$$\int d^D x \sqrt{|g|} [A^*_{\sigma'\mu} \nabla^0 A^\mu_\sigma - (\nabla^0 A^*_{\sigma'\mu}) A^\mu_\sigma] = 4i\pi \delta_{\sigma\sigma'}, \qquad (2.30)$$

where the set of quantum numbers is given by $\sigma = (\mathbf{k}, \lambda, \gamma)$.

The coefficients c_1 and c_2 in (2.29) are determined from (2.30) and from the boundary condition on the brane. We will consider two types of gauge invariant constraints. The first condition is the analog of the perfect conductor boundary condition in D = 3 Maxwell electrodynamics and is given by

$$n^{\mu_1 *} F_{\mu_1 \cdots \mu_{D-1}} = 0, \ z = z_0, \tag{2.31}$$

with $F_{\mu_1\cdots\mu_{D-1}}$ being the dual of the field tensor and n^{μ} is the normal vector to the boundary. The second boundary conditions is expressed as

$$n^{\mu}F_{\mu\rho} = 0, \ z = z_0. \tag{2.32}$$

This type of condition has been used in quantum chromodynamics to confine the gluons. In the gauge under consideration and for the mode functions (2.29) the boundary condition (2.31) yields $A_{\sigma D}|_{z=z_0} = 0$, whereas the condition (2.32) gives $\partial_D A_{\sigma l}|_{z=z_0} = 0$.

In the R-region, from the boundary condition on the brane we get $c_2/c_1 = -J_{\nu}(\lambda z_0)/Y_{\nu}(\lambda z_0)$, where

$$\nu = \begin{cases} D/2 - 1, & \text{for } (2.31), \\ D/2 - 2, & \text{for } (2.32). \end{cases}$$
(2.33)

The corresponding mode functions for the vector potential are presented as

$$A_{\sigma\mu} = C_{(\mathrm{R})\sigma} \epsilon_{(\gamma)\mu} z^{D/2-1} g_{\nu,D/2-1}(\lambda z_0, \lambda z) e^{i\mathbf{k}\mathbf{x} - i\omega t}, \qquad (2.34)$$

where the function $g_{\mu,\rho}(x,u)$ is given by (2.23). The normalization constant is found from (2.30):

$$C_{(\mathrm{R})\sigma}^{2} = \lambda \frac{\left[J_{\nu}^{2}(\lambda z_{0}) + Y_{\nu}^{2}(\lambda z_{0})\right]^{-1}}{(2\pi)^{D-2} a^{D-3} \sqrt{k^{2} + \lambda^{2}}},$$
(2.35)

with $0 \leq \lambda < \infty$.

In the L-region from the normalizability of the mode functions it follows that in (2.29) $c_2 = 0$ for $D \ge 4$. For D = 3 an additional condition on the AdS boundary is required in order to uniquely define the modes. Here we consider a special case with $c_2 = 0$ for D = 3 as well. For the mode functions we obtain

$$A_{\sigma\mu}(x) = C_{(\mathrm{L})\sigma}\epsilon_{(\gamma)\mu}z^{D/2-1}J_{D/2-1}(\lambda z)e^{i\mathbf{k}\mathbf{x}-i\omega t}.$$
(2.36)

The allowed values for λ are determined by the boundary condition on the brane and they are the roots of the equation

$$J_{\nu}(\lambda z_0) = 0, \qquad (2.37)$$

with ν from (2.33). Hence, we have $\lambda = \lambda_{\nu,n}/z_0$. For the normalization coefficient one gets

$$C_{(\mathrm{L})\sigma}^{2} = \frac{2 (2\pi)^{2-D} a^{3-D}}{z_{0} \sqrt{k^{2} z_{0}^{2} + \lambda_{\nu,n}^{2}} J_{\nu}^{\prime 2}(\lambda_{\nu,n})}.$$
(2.38)

The mode functions in the region between two branes have been recently considered in [13].

2.4 Boundary conditions in Z_2 -symmetric braneworlds

An example of the Z_2 -symmetric braneworld is provided by the Randall-Sundrum model with a single brane (RSII model) formulated in background of (4+1)-dimensional AdS spacetime (see [35, 36] and the review [7] for the RSI and RSII models). For an arbitrary number of spatial dimensions the line element is given by (2.1) with $e^{-2y/a}$ replaced by $e^{-2|y|/a}$. The regions $-\infty < y < 0$ and $0 < y < +\infty$ are identified by the Z_2 -symmetry. The brane is located at y = 0. Hence, in the corresponding setup two copies of the R-region are employed with $z_0 = a$. The boundary conditions on the bulk fields at the location of the brane are obtained by integrating the field equations about y = 0 (see, e.g., the discussions in [16, 34, 37, 38, 39, 40]).

For scalar fields even under the reflection with respect to the brane, the Robin boundary condition is obtaned with the coefficient $\beta = -2/(c_b + 4D\xi/a)$, where c_b is the brane mass term. The latter appears in the part of the action located on the brane, $S_b = -c_b \int d^D x dy \sqrt{|g|} \delta(y) \varphi^2/2$. For odd scalar fields the Dirichlet boundary condition is obtained. For fermionic fields two types of boundary conditions are obtained. The first one is reduced to the bag boundary condition (2.21) and the second one is obtained from (2.21) by the change of the sign in the term containing the normal to the brane. For vector fields even under the reflection with respect to the brane, the boundary condition is reduced to (2.32) and for odd fields the condition (2.31) is obtained. For fermionic fields with the boundary condition obtained from (2.21) by the change of the sign in the second term, the VEV of the energy-momentum tensor is evaluated in a way similar to that we have demonstrated for the bag boundary condition. The corresponding mode functions are obtained from (2.22) by the replacement $ma + 1/2 \rightarrow ma - 1/2$ in the first index of the functions $g_{\mu,\rho}(x,u)$, $g_{ma+1/2,ma+s/2}(\lambda z_0, \lambda z) \rightarrow g_{ma-1/2,ma+s/2}(\lambda z_0, \lambda z)$, and in the expression (2.24) for the normalization coefficient.

Hence, we conclude that the VEVs of the energy-momentum tensors for scalar, Dirac and electromagnetic fields in Z_2 -symmetric braneworlds with a single brane are obtained from the results given above by an appropriate choice of the boundary conditions on the brane. The only difference is that an additional factor 1/2 should be added. The latter is related to the presence of two copies of the R-region.

3 Vacuum energy-momentum tensor

3.1 General properties

Having the mode functions for quantum fields we can investigate the VEVs of the local characteristics of the vacuum state. Among the most important characteristics is the VEV of the energy-momentum tensor. In particular, it determines the distribution of the vacuum energy density and the forces acting on the boundaries (the Casimir forces). For a free field $\Psi(x)$ (the only interaction is that with background gravitational field) the operator of the energy-momentum tensor is a bilinear form in the field operator:

$$T_{\mu\rho} = T_{\mu\rho} \{\Psi(x), \Psi(x)\}.$$
(3.1)

The corresponding expressions of the bilinear form in the right-hand side for scalar, Dirac and vector fields can be found, for example, [41, 42]. Expanding the field opertor in terms of a complete set $(\Psi_{\sigma}^{(+)}(x), \Psi_{\sigma}^{(-)}(x))$ of the positive and negative energy mode functions, specified by quantum numbers σ , using the commutation relations for the creation and annihilation operators and the definition of the vacuum state $|0\rangle$, for the VEV of the energy-momentum tensor the following sum over the modes is obtained:

$$\langle 0|T_{\mu\rho}|0\rangle \equiv \langle T_{\mu\rho}\rangle = \frac{1}{2} \sum_{\sigma} \sum_{s=\pm} T_{\mu\rho} \left\{ \Psi_{\sigma}^{(s)}(x), \Psi_{\sigma}^{(s)}(x) \right\}.$$
(3.2)

Here, \sum_{σ} is understood as summation for discrete components of σ and as integration over the continuous components. The expression in the right-hand side of (3.2) is divergent and a regularization procedure is required. The regularization can be made by point splitting, by using the local zeta function technique or by introducing a cutoff function [41]-[46]. For definiteness, we will assume that a cutoff function is introduced in (3.2) without writing it explicitly. The details of the evaluation do not depend on the specific regularization scheme.

In the geometry with a brane, we will decompose the vacuum energy-momentum tensor into two contributions:

$$\langle T_{\mu\rho} \rangle = \langle T_{\mu\rho} \rangle_0 + \langle T_{\mu\rho} \rangle_{\rm b} \,, \tag{3.3}$$

where $\langle T_{\mu\rho} \rangle_0$ is the VEV in the absence of the brane and the part $\langle T_{\mu\rho} \rangle_b$ is induced by the brane. The VEV in the brane-free AdS geometry has been widely discussed in the literature (for recent discussion see, for example, [47]) and here we are mainly interested in the brane-induced effects. From the maximal symmetry of the AdS spacetime it follows that $\langle T_{\mu\rho} \rangle_0 = \text{const} \cdot g_{\mu\rho}$. On the base of the symmetry of the problem with a brane parallel to the AdS boundary, we expect that the VEV $\langle T_{\mu\rho} \rangle_b$ is diagonal. From the Lorentz invariance in the subspace $(t, x^1, \ldots, x^{D-1})$ one concludes that the stresses in the directions parallel to the brane are equal to the energy density:

$$\langle T_0^0 \rangle_{\mathbf{b}} = \langle T_1^1 \rangle_{\mathbf{b}} = \dots = \langle T_{D-1}^{D-1} \rangle_{\mathbf{b}}.$$
 (3.4)

An additional relation between the components of the brane-induced VEV is obtained from the covariant continuity equation $\nabla_{\mu} \langle T^{\mu}_{\rho} \rangle_{\rm b} = 0$. The latter is reduced to

$$z^{D+1}\partial_z \left(z^{-D} \left\langle T_D^D \right\rangle_{\mathbf{b}} \right) + D \left\langle T_0^0 \right\rangle_{\mathbf{b}} = 0.$$
(3.5)

As it will be shown below, the brane-induced contribution in the VEVs depend on the coordinate z and on the location of the brane through the ratio z/z_0 . This property is a consequence of the maximal symmetry of the AdS spacetime and of the vacuum state we consider here. The procedure for the evaluation of the brane-induced contribution in the VEV of the energy-momentum tensor follows similar steps for scalar, fermion and electromagnetic fields and we illustrate it on the example of the fermionic field.

The geometry given by (2.2) is conformally flat and for conformally coupled fields (conformally coupled massless scalar field ($\xi = \xi_D$, m = 0), massless Dirac field, the electromagnetic field in D = 3 spatial dimensions) the problem under consideration is conformally related to the corresponding problem in the Minkowski bulk. Denoting by $\langle T^{\rho}_{\mu} \rangle_{\rm b}^{({\rm M})}$ the boundary-induced VEV in the Minkowskian problem, we have the following relation

$$\left\langle T^{\rho}_{\mu} \right\rangle_{\mathbf{b}} = (z/a)^{D+1} \left\langle T^{\rho}_{\mu} \right\rangle_{\mathbf{b}}^{(\mathbf{M})}. \tag{3.6}$$

For the R-region, the problem with a brane in AdS bulk parallel to the AdS boundary is conformally related to the problem in the Minkowski bulk with the line element $ds_M^2 = \eta_{\mu\rho} dx^{\mu} dx^{\rho}$, $x^D = z$, and with a single Robin boundary at $z = z_0$. For the L-region the Minkowskian counterpart contains two boundaries. The first one is located at $z = z_0$ and is the conformal image of the brane and the second one is located at z = 0 and is the conformal image of the AdS boundary. The boundary conditions on z = 0 for fields in the Minkowskian problem are related to the special boundary conditions we have imposed on the AdS boundary.

3.2 R-region

We will describe the procedure for the evaluation of the brane-induced energy-momentum tensor on the example of the Dirac field. The procedure for scalar and electromagnetic fields follows similar steps.

3.2.1 Dirac field

By using the expression for the energy-momentum tensor of the Dirac field and the corresponding mode functions from the previous section, the diagonal components in the R-region are presented in the form (here and in what follows in the expressions for $\langle T^{\mu}_{\mu} \rangle$ no summation is assumed over μ)

$$\langle T^{\mu}_{\mu} \rangle^{(\mathrm{f})} = \frac{Na^{-D-1}(z/z_0)^{D+2}}{2(4\pi)^{(D-1)/2}\Gamma((D-1)/2)} \int_0^\infty du \, u^{D-2} \\ \times \int_0^\infty dx \frac{x}{\sqrt{x^2 + u^2}} \frac{f^{(\mu)}_{(\mathrm{R})}(x, xz/z_0)}{J^2_{ma+1/2}(x) + Y^2_{ma+1/2}(x)},$$
(3.7)

where

$$f_{(R)}^{(0)}(x,y) = -(x^{2}+u^{2}) \left[g_{ma+1/2,ma+s/2}^{2}(x,y) + g_{ma+1/2,ma-s/2}^{2}(x,y) \right],$$

$$f_{(R)}^{(D)}(x,y) = x^{2} \left[g_{ma+1/2,ma+s/2}^{2}(x,y) + g_{ma+1/2,ma-s/2}^{2}(x,y) - \frac{2ma}{y} g_{ma+1/2,ma+s/2}(x,y) g_{ma+1/2,ma-s/2}(x,y) \right].$$
(3.8)

As seen from (3.8), the VEVs for s = +1 and s = -1 coincide. For the separation of the brane-induced part we use the relation

$$\frac{g_{\mu,\rho'}(x,u)g_{\mu,\rho'}(x,u)}{J_{\mu}^{2}(x)+Y_{\mu}^{2}(x)} = J_{\rho}(u)J_{\rho'}(u) - \frac{1}{2}\sum_{s=1,2}\frac{J_{\mu}(x)}{H_{\mu}^{(s)}(x)}H_{\rho}^{(s)}(u)H_{\rho'}^{(s)}(u),$$
(3.9)

where $H_{\rho}^{(s)}(u)$, s = 1, 2, are the Hankel functions. The part in the VEV coming from the first term in the right-hand side of (3.9) is the vacuum energy-momentum tensor in the geometry where the brane is absent (the part $\langle T_{\mu\rho} \rangle_0$ in (3.3)). In the parts with the last term in (3.9) we rotate the integration contour over x by the angle $\pi/2$ for s = 1 terms and by the angle $-\pi/2$ for s = -1 terms. Introducing the modified Bessel functions $I_{\mu}(x)$ and $K_{\mu}(x)$, we get

$$\langle T^{\mu}_{\mu} \rangle_{\rm b}^{\rm (f)} = -\frac{2^{-D}N}{\pi^{D/2}\Gamma(D/2)a^{D+1}} \int_0^\infty dx \, x^{D+1} \frac{I_{ma+1/2}(xz_0/z)}{K_{ma+1/2}(xz_0/z)} F^{(\mu)}_{\rm (R)}(x), \tag{3.10}$$

with the notations

$$F_{(R)}^{(0)}(x) = \frac{1}{D} \left[K_{ma+1/2}^2(x) - K_{ma-1/2}^2(x) \right],$$

$$F_{(R)}^{(D)}(x) = K_{ma-1/2}^2(x) - K_{ma+1/2}^2(x) + \frac{2ma}{x} K_{ma+1/2}(x) K_{ma-1/2}(x).$$
(3.11)

For massive fields both the energy density $\langle T_0^0 \rangle_{\rm b}^{\rm (f)}$ and the normal stress $\langle T_D^D \rangle_{\rm b}^{\rm (f)}$ are negative. As seen from (3.10), the brane-induced contribution depends on the coordinates z and z_0 through the ratio $z/z_0 = e^{(y-y_0)/a}$, where $y_0 = a \ln(z_0/a)$ is the location of the brane in terms of the coordinate y. Note that $y - y_0$ is the proper distance from the brane. For a massless field the brane-induced contribution vanishes, $\langle T_{\mu}^{\mu} \rangle_{\rm b} = 0$. The massless fermionic field is conformally invariant and this result could be directly obtained from the corresponding result for a single boundary in the Minkowski bulk by using the relation (3.6).

The Minkowskian result for a massive field is obtained from (3.10) in the limit $a \to \infty$ for fixed y (see (2.1)). From the relation $z = ae^{y/a}$ it follows that $z \approx a + y$ and the values for the coordinate z are large. By using the uniform asymptotic expansions for the modified Bessel functions for large values of the order, the leading term in the expansion of the components with $\mu = 0, 1, \ldots, D-1$ coincides with the VEV for a planar boundary in the Minkowski bulk:

$$\langle T^{\mu}_{\mu} \rangle_{\rm b}^{({\rm Mf})} = -\frac{2^{-D}Nm}{\pi^{D/2}D\Gamma(D/2)} \int_{m}^{\infty} dx \, (x^2 - m^2)^{D/2} \frac{e^{-2x(y-y_0)}}{x+m},$$
 (3.12)

and the normal stress vanishes, $\langle T_D^D \rangle_{\rm b}^{({\rm Mf})} = 0$.

The general formula (3.10) for the brane-induced contribution in the energy-momentum tensor is simplified in the region near the brane and at large distances from it (near-horizon limit). For large values x the integrand in (3.10) behaves as $x^{D-1}e^{-2x(1-z_0/z)}$ and the integral is divergent for points on the brane. These surface divergences are well-known in quantum field theory with boundaries and have been widely discussed in the literature for different geometries of boundaries (see, for instance, [8]). Near the brane, assuming that $z/z_0 - 1 \ll 1$, the contribution of large values of x dominate in the integral and by making use of the corresponding asymptotic formulas for the modified Bessel functions, in the leading order we obtain

$$\langle T_0^0 \rangle_{\rm b}^{\rm (f)} \approx -\frac{Nm\Gamma((D+1)/2)}{(4\pi)^{(D+1)/2}D(y-y_0)^D}, \ \langle T_D^D \rangle_{\rm b}^{\rm (f)} \approx -\frac{DNm\Gamma((D-1)/2)}{2(4\pi)^{(D+1)/2}Da(y-y_0)^{D-1}}.$$
 (3.13)

In terms of the coordinate y, these asymptotics are valid under the conditions $y - y_0 \ll a, m^{-1}$. The leading terms for the energy density and for the stresses parallel to the brane coincide with those

for a boundary in the Minkowski bulk. This is related to the fact that near the brane the dominant contribution to the VEV comes from the vacuum fluctuations with wavelengths smaller than the curvatures radius and the influence of the gravitational field on those fluctuations is weak. The effects of gravity are essential at distances from the brane larger than the curvature radius, $z/z_0 \gg 1$ or $y - y_0 \gg a$. In that region the leading term in the asymptotic expansion for the energy density is expressed as

$$\langle T_0^0 \rangle_{\rm b}^{\rm (f)} \approx -Nm \frac{\exp[-(2m+1/a)(y-y_0)]}{2^{D+2ma+1}\pi^{(D-1)/2}a^D} \\ \times \frac{\Gamma(ma+(D+3)/2)\Gamma(2ma+D/2+1)}{(2ma+1)\Gamma^2(ma+1/2)\Gamma(ma+D/2+2)},$$
(3.14)

and for the normal stress we get $\langle T_D^D \rangle_{\rm b}^{\rm (f)} \approx D \langle T_0^0 \rangle_{\rm b}^{\rm (f)} / (D + 2ma + 1)$. Note that for a boundary in the Minkowski bulk and for $y - y_0 \gg 1/m$ the VEV decays like $e^{-2m(y-y_0)}$. In the AdS spacetime, the decay of the boundary-induced contribution, as a function of the proper distance from the boundary, is stronger.

The left panel of figure 1 presents the dependence of the brane-induced contributions to the energy density and the normal stress on the ratio z/z_0 for fixed value ma = 1. On the right panel we have displayed the brane-induced parts as functions of the field mass for fixed $z/z_0 = 2$. For both panels the full and dashed curves correspond to D = 3 and D = 4, respectively.



Figure 1: The brane-induced contributions in the VEVs of the energy density and the normal stress as functions of z/z_0 (left panel) and of ma (right panel). For the left panel we have taken ma = 1 and for the right panel $z/z_0 = 2$. The full and dashed curves correspond to D = 3 and D = 4.

3.2.2 Scalar field

In a similar way, for the brane-induced contribution in the case of a scalar field with the boundary condition (2.9) one obtains

$$\langle T^{\mu}_{\mu} \rangle_{\rm b}^{\rm (s)} = -\frac{2^{-D}}{\pi^{D/2} \Gamma(D/2) a^{D+1}} \int_0^\infty dx \, x^{D+1} \frac{\bar{I}_{\nu}(x z_0/z)}{\bar{K}_{\nu}(x z_0/z)} S^{(\mu)} \left[K_{\nu}(x) \right], \tag{3.15}$$

where the notation with the bar is defined as (2.10) and

$$S^{(0)}[g(x)] = -\xi_1 \left[g^{\prime 2}(x) + \frac{D + 4\xi/\xi_1}{x} g(x)g^{\prime}(x) + \left(1 + \frac{2}{D\xi_1} + \frac{\nu^2}{x^2}\right)g^2(x) \right],$$

$$S^{(D)}[g(x)] = -g^{\prime 2}(x) + \frac{D\xi_1}{x} g(x)g^{\prime}(x) + \left(1 + \frac{2m^2a^2 - \nu^2}{x^2}\right)g^2(x),$$
(3.16)

with

$$\xi_1 = 4\xi - 1. \tag{3.17}$$

For a conformally coupled massless scalar field the brane-induced contribution (3.15) vanishes. In the limit $a \to \infty$, with fixed y and y_0 , from (3.15) we get the vacuum energy-momentum tensor for a Robin boundary in the Minkowski bulk:

$$\langle T^{\mu}_{\mu} \rangle_{\rm b}^{\rm (Ms)} = \frac{2^{-D} \pi^{-D/2}}{\Gamma(D/2)} \int_{m}^{\infty} dx \, \left(x^2 - m^2\right)^{D/2 - 1} \left[\frac{m^2}{D} - 4\left(\xi - \xi_D\right) x^2\right] \frac{\beta x + 1}{\beta x - 1} e^{-2x(y - y_0)}, \qquad (3.18)$$

for $\mu = 0, 1, 2, \dots, D - 1$ and $\langle T_D^D \rangle_{\rm b}^{({\rm Ms})} = 0.$

For points near the brane, $y - y_0 \ll a, m^{-1}, |\beta|$, the leading term in the asymptotic expansion of the energy density is given by the expression

$$\langle T_0^0 \rangle_{\rm b}^{\rm (s)} \approx \pm \frac{D\Gamma\left((D+1)/2\right)\left(\xi - \xi_D\right)}{2^D \pi^{(D+1)/2} (y - y_0)^{D+1}},$$
(3.19)

where the upper and lower signs correspond to $\beta = 0$ (Dirichlet boundary condition) and $\beta \neq 0$ (non-Dirichlet boundary conditions), respectively. The leading term in (3.19) coincides with that for the Minkwoski bulk. For the normal stress we find $\langle T_D^D \rangle_{\rm b}^{(\rm s)} \approx \langle T_0^0 \rangle_{\rm b}^{(\rm s)}(y-y_0)/a$. In the case of a conformally coupled field, the leading terms vanish and the divergences on the brane are weaker. At large distances from the brane and for $\nu > 0$, the energy density is approximated as

$$\langle T_0^0 \rangle_{\rm b}^{\rm (s)} \approx \frac{(2\nu - 1)e^{-2\nu(y-y_0)/a}}{2^{2\nu + D - 1}\pi^{(D-1)/2}a^{D+1}} \frac{1 + \beta\nu}{1 - \beta\nu} \left(4\xi - \frac{D + 2\nu}{D + 2\nu + 1} \right) \\ \times \frac{\Gamma(D/2 + \nu + 1)\Gamma(D/2 + 2\nu)}{\Gamma(\nu + 1)\Gamma(D/2 + 1/2 + \nu)}.$$

$$(3.20)$$

For the normal stress we get $\langle T_D^D \rangle_{\rm b}^{({\rm s})} \approx D \langle T_0^0 \rangle_{\rm b}^{({\rm s})}/(D+2\nu)$. Note that for $\nu > 0$ the decay of the brane-induced VEV is exponential for both massless and massive fields. In the Minkowski bulk, the boundary-induced contribution decays as $e^{-2m(y-y_0)}$ for massive fields and as power-law $(y-y_0)^{-D-1}$ for non-conformally coupled massless fields.

On the left panel of figure 2 we display the brane-induced energy density in the R-region as a function of z/z_0 for D = 4 minimally coupled scalar field with ma = 1. The graphs are plotted for the Dirichlet and Neumann boundary conditions and for the Robin boundary conditions with $\beta/a = -0.1, -0.15, -0.5$. The graphs for the Robin boundary conditions are located between the curves corresponding to the Dirichlet and Neumann conditions and for them $\langle T_0^0 \rangle_b^{(s)}$ increases with increasing $|\beta|$. The normal stress displays a similar behavior as a function of z/z_0 . The right panel in figure 2 presents the brane-induced contributions to the energy density (full curves) and normal stress (dashed curves) for fixed $z/z_0 = 2$ as functions of ma for D = 4 minimally coupled field. The graphs are plotted for the Dirichlet and Neumann boundary conditions and for the Robin boundary condition with $\beta/a = -0.15$. In the latter case the energy density changes the sign at $ma \approx 3.13$ and the normal stress changes the sign at $ma \approx 3.34$.

Figure 3 presents similar results for D = 4 conformally coupled field ($\xi = 3/16$). Note that in this case the stability condition $a/\beta < \nu - D/2$ on the Robin coefficient is stronger. For the left panel we have taken ma = 2. With this value, for all the boundary conditions with $\beta \leq 0$ the vacuum is stable. The right panel is plotted for $z/z_0 = 2$. There is a range for the values of ma in which the Neumann boundary condition for a conformally coupled field leads to the vacuum instability.



Figure 2: The brane-induced contributions in the VEVs of the energy density and the normal stress for D = 4 minimally coupled scalar field. For the left panel ma = 1 and for the right panel $z/z_0 = 2$. The full and dashed curves on the right panel correspond to the energy density and normal stress, respectively (for the boundary conditions chosen see the text).



Figure 3: The same as in figure 2 for D = 4 conformally coupled scalar field. The left and right panels are plotted for ma = 2 and $z/z_0 = 2$, respectively.

3.2.3 Electromagnetic field

For the electromagnetic field with the boundary conditions (2.31) and (2.32) on the brane, the braneinduced VEV has the form

$$\langle T^{\mu}_{\mu} \rangle_{\rm b}^{\rm (v)} = -\frac{\delta_{\nu} \left(D-1\right)}{2^{D} \pi^{D/2} \Gamma(D/2) a^{D+1}} \int_{0}^{\infty} dx \, x^{D+1} \frac{I_{\nu}(x z_{0}/z)}{K_{\nu}(x z_{0}/z)} V^{(\mu)}_{\rm (R)}(x), \tag{3.21}$$

where $\delta_{\nu} = 1$ for $\nu = D/2 - 1$ (boundary condition (2.31)), $\delta_{\nu} = -1$ for $\nu = D/2 - 2$ (boundary condition (2.32)), and

$$V_{(R)}^{(0)}(x) = \left(1 - \frac{4}{D}\right) K_{D/2-1}^2(x) + \left(1 - \frac{2}{D}\right) K_{D/2-2}^2(x),$$

$$V_{(R)}^{(D)}(x) = K_{D/2-1}^2(x) - K_{D/2-2}^2(x).$$
(3.22)

Note that the VEV has different signs for the boundary conditions (2.31) and (2.32). For D = 3 the electromagnetic field is conformally invariant and the VEV (3.21) is zero. One has $\langle T^{\mu}_{\mu} \rangle_{\rm b}^{(v)} < 0$ for the boundary condition (2.31) and $\langle T^{\mu}_{\mu} \rangle_{\rm b}^{(v)} > 0$ for the condition (2.31). The VEVs for a plate in the Minkowski bulk are obtained from (3.21) in the limit $a \to \infty$:

$$\langle T^{\mu}_{\mu} \rangle^{(M_{V})}_{b} = \mp \frac{(D-1)(D-3)\Gamma((D+1)/2)}{2(4\pi)^{(D+1)/2}(y-y_{0})^{D+1}},$$
(3.23)

for $\mu = 0, 1, 2, \ldots, D-1$ and $\langle T_D^D \rangle_{\rm b}^{({\rm Mv})} = 0$. The upper and lower signs in (3.23) correspond to the conditions (2.31) and (2.32), respectively.

For D > 3 the VEV (3.21) diverges on the brane. The leading terms in the asymptotic expansions for the energy density and for the normal stress over the distance from the brane are expressed as

$$\langle T_0^0 \rangle_{\rm b}^{\rm (v)} \approx -\delta_{\nu} \frac{(D-1) (D-3) \Gamma((D+1)/2)}{2 (4\pi)^{(D+1)/2} (y-y_0)^{D+1}}, \ \langle T_D^D \rangle_{\rm b}^{\rm (v)} \approx \frac{y-y_j}{a} \langle T_0^0 \rangle.$$
(3.24)

The leading term for the energy density coincides with the exact result for a boundary in the Minkowski bulk, given by (3.23). At large distances from the brane, assuming that $z_0/z \ll 1$, for $\nu > 0$ to the leading order one gets

$$\langle T^{\mu}_{\mu} \rangle_{\rm b}^{\rm (v)} \approx -\frac{2^{1-D-2\nu} \delta_{\nu} \left(D-1\right) e^{-2\nu(y-y_0)}}{\pi^{D/2} \Gamma(D/2) \nu \Gamma^2(\nu) a^{D+1}} \int_0^\infty dx \, x^{D+2\nu+1} V^{(\mu)}_{\rm (R)}(x), \tag{3.25}$$

where the integral is expressed in terms of the product of the gamma functions (see [49]). For D = 4 and for the boundary condition (2.32) one has $\nu > 0$ and at large distances the brane-induced part in the energy-momentum tensor falls as $1/(y - y_0)$.

In figure 4 we present the brane-induced energy density and the normal stress versus z/z_0 for D = 4 (left panel) and D = 5 (right panel) electromagnetic fields. The full and dashed curves correspond to the boundary conditions (2.31) and (2.32), respectively, and near the curves the value of the index μ is given for $\langle T^{\mu}_{\mu} \rangle_{\rm b}^{(\rm v)}$ from (3.21).

3.3 L-region

In the L-region the eigenvalues for the quantum number λ are discrete. Again, we illustrate the evaluation procedure for the example of the Dirac field. The VEVs for scalar and electromagnetic fields are evaluated in a similar way.



Figure 4: The brane-induced VEVs of the energy density and the normal stress as functions of z/z_0 for D = 4 (left panel) and D = 5 (right panel) electromagnetic fields. The full and dashed curves correspond to the conditions (2.31) and (2.32).

3.3.1 Dirac field

By using the mode functions (2.25), the mode sum for the energy-momentum tensor is presented in the form

$$\langle T^{\mu}_{\mu} \rangle^{(f)} = -\frac{(4\pi)^{(1-D)/2} N z^D}{\Gamma\left((D-1)/2\right) a^{D+1} z_0} \int_0^\infty dk \, k^{D-2} \sum_{n=1}^\infty \frac{f^{(\mu)}_{(L)}(\lambda_{ma-1/2,n} z/z_0)}{\sqrt{k^2 z_0^2 + \lambda_{ma-1/2,n}^2} J^2_{ma+1/2}(\lambda_{ma-1/2,n})}, \quad (3.26)$$

with the functions

$$f_{(L)}^{(0)}(x) = -(k^2 z^2 + x^2) \left[J_{ma+s/2}^2(x) + J_{ma-s/2}^2(x) \right],$$

$$f_{(L)}^{(D)}(x) = x^2 \left[J_{ma+s/2}^2(x) + J_{ma-s/2}^2(x) - \frac{2ma}{x} J_{ma+s/2}(x) J_{ma-s/2}(x) \right].$$
(3.27)

By taking into account that the eigenvalues $\lambda_{ma-1/2,n}$ do not depend on s, we conclude that, similar to the R-region, the VEV of the energy-momentum tensor is the same for both inequivalent representations of the Clifford algebra.

For the summation of the series over the roots $\lambda_{ma-1/2,n}$ we use the Abel-Plana type formula [50, 51]

$$\sum_{n=1}^{\infty} \frac{f(\lambda_{ma-1/2,n})/\lambda_{ma-1/2,n}}{J_{ma+1/2}^{2}(\lambda_{ma-1/2,n})} = \frac{1}{2} \int_{0}^{\infty} dx f(x) - \frac{1}{2\pi} \int_{0}^{\infty} dx \frac{K_{ma-1/2}(x)}{I_{ma-1/2}(x)} \times \left[e^{(1/2-ma)\pi i} f(ix) + e^{(ma-1/2)\pi i} f(-ix) \right].$$
(3.28)

The part of the VEV corresponding to the first term in the right-hand side of (3.28) gives the VEV in the geometry without the brane. For the brane-induced contribution coming from the second integral in (3.28) we get

$$\langle T^{\mu}_{\mu} \rangle_{\rm b}^{\rm (f)} = -\frac{2^{-D}N}{\pi^{D/2}\Gamma(D/2)a^{D+1}} \int_0^\infty dx \, x^{D+1} \frac{K_{ma-1/2}(xz_0/z)}{I_{ma-1/2}(xz_0/z)} F^{(\mu)}_{\rm (L)}(x), \tag{3.29}$$

where the functions in the integrand are defined as

$$F_{(L)}^{(0)}(x) = \frac{1}{D} \left[I_{ma-1/2}^{2}(x) - I_{ma+1/2}^{2}(x) \right],$$

$$F_{(L)}^{(D)}(x) = I_{ma+1/2}^{2}(x) - I_{ma-1/2}^{2}(x) + \frac{2ma}{x} I_{ma+1/2}(x) I_{ma-1/2}(x).$$
(3.30)

The energy density and the parallel stresses corresponding to (3.29) are negative, $\langle T^{\mu}_{\mu} \rangle_{\rm b}^{\rm (f)} < 0, \mu = 0, 1, \ldots, D-1$, whereas the normal stress is positive. Comparing with the results for the R-region, we see that the energy densities in the R- and L-regions have the same sign and the normal stresses have opposite signs.

For a massless field one obtains $\langle T_D^D \rangle_{\rm b}^{\rm (f)} = -D \langle T_0^0 \rangle_{\rm b}^{\rm (f)}$ and the expression for the energy density is simplified to

$$\langle T_0^0 \rangle_{\rm b}^{\rm (f)} = -\left(\frac{z}{az_0}\right)^{D+1} \frac{N\Gamma((D+1)/2)}{(4\pi)^{(D+1)/2}} (1-2^{-D})\zeta(D+1),$$
 (3.31)

where $\zeta(x)$ is the Riemann zeta function. In this case the brane-induced part is traceless. The massless fermionic field is conformally invariant and (3.31) corresponds to the conformal relation (3.6). In the Minkowskian counterpart, z_0 is the separation between two parallel planar boundaries. For a massive field and in the Minkowskian limit $a \to 0$ we obtain the expression (3.12) for the energy density and parallel stresses, with the replacement $y - y_0 \to y_0 - y$, and the normal stress vanishes.

The VEV (3.29) depends on z and z_0 in the form of the ration z/z_0 . Let us consider its behavior in the asymptotic regions. The limit $z/z_0 \to 1$ corresponds to the points on the brane. The brane-induced contribution (3.29) diverges on the brane. The leading terms in the asymptotic expansions over the distance from the brane are obtained from the corresponding expressions (3.13) for the R-region by the replacement $y - y_0 \to y_0 - y$ and by an additional change of the sign for the normal stress. For $z/z_0 \ll 1$ we use the small argument asymptotics for the functions $F_{(L)}^{(\mu)}(x)$. To the leading order this gives

$$\langle T_0^0 \rangle_{\rm b}^{\rm (f)} \approx -\frac{2^{-D-2ma} \pi^{-D/2} N(z/z_0)^{D+2ma+1}}{\Gamma(D/2+1) \Gamma^2 (ma+1/2) a^{D+1}} \int_0^\infty dx \, x^{D+2ma} \frac{K_{ma-1/2}(x)}{I_{ma-1/2}(x)} dx^{D+2ma} \frac{K_{ma-1/2}(x)}{I_{ma-1/2}(x)}$$

for the energy density and

$$\langle T_D^D \rangle_{\rm b}^{\rm (f)} \approx -\frac{D}{2ma+1} \langle T_0^0 \rangle_{\rm b}^{\rm (f)}$$

for the normal stress. In particular, the brane-induced VEV tends to zero on the AdS boundary as $z^{D+2ma+1}$.

The brane-induced contributions for the energy density and the normal stress in the L-region, $\langle T^{\mu}_{\mu} \rangle^{(f)}_{\rm b}$, $\mu = 0, D$, for D = 3 (full curves) and D = 4 (dashed curves) Dirac fields are presented in figure 5 as functions of z/z_0 and of ma. The left and right panels are plotted for ma = 1 and $z/z_0 = 0.6$.



Figure 5: The same as in figure 1 for the L-region. The right panel is plotted for $z/z_0 = 0.6$.

3.3.2 Scalar field

For a scalar field the mode-sum formula for the VEV of the energy-momentum tensor contains series over the zeros of the function $\bar{J}_{\nu}(x) = B_0 x J'_{\nu}(x) + A_0 J_{\nu}(x)$. The corresponding summation formula can be found in [50, 51]. The brane-induced contribution for a scalar field is presented as

$$\langle T^{\mu}_{\mu} \rangle_{\rm b}^{\rm (s)} = -\frac{2^{-D}}{\pi^{D/2} \Gamma(D/2) a^{D+1}} \int_0^\infty dx \, x^{D+1} \frac{\bar{K}_{\nu}(xz_0/z)}{\bar{I}_{\nu}(xz_0/z)} S^{(\mu)} \left[I_{\nu}(x) \right], \tag{3.32}$$

where the functions $S^{(\mu)}[g(x)]$ are given by (3.16). For a conformally coupled massless field one has $\nu = 1/2$ and $S^{(0)}[I_{1/2}(x)] = 2/(\pi Dx)$, $S^{(D)}[I_{1/2}(x)] = -2/(\pi x)$. For the energy density we get

$$\langle T_0^0 \rangle_{\rm b}^{\rm (s)} = -\frac{(z/a)^{D+1} z_0^{-D-1}}{2^{2D+1} \pi^{D/2} \Gamma(D/2+1)} \int_0^\infty dx \frac{x^D}{\frac{2a/\beta + 1 - D - x}{2a/\beta + 1 - D + x}} e^x - 1, \tag{3.33}$$

and for the normal stress $\langle T_D^D \rangle_{\rm b}^{\rm (s)} = -D \langle T_0^0 \rangle_{\rm b}^{\rm (s)}$. Note that one has the conformal relation $\varphi(x) = (z/a)^{(D-1)/2} \varphi_{\rm M}(x)$ between the fields in the AdS and Minkowski bulk. From here it follows that the conformal image of the boundary condition (2.9) is the condition $(\beta_{\rm M} n_{\rm M}^{\mu} \nabla_{\mu} + 1) \varphi_{\rm M}(x) = 0, z = z_0$, with the relation between the Robin coefficients

$$\frac{z_0}{\beta_{\rm M}} = \frac{a}{\beta} - \frac{D-1}{2}.$$
(3.34)

By taking into account (3.34), we can see that (3.33) corresponds to the conformal relation (3.6) with the Minkowskian problem of two boundaries with the Dirichlet boundary condition on the plate z = 0(conformal image of the AdS boundary) and the Robin condition with the coefficient $\beta_{\rm M}$ on the plate at $z = z_0$. In the Minkowskian limit the result (3.18) is obtained for the energy density and parallel stresses and the normal stress becomes zero.

Let us consider the behavior of the brane-induced VEV (3.32) near the brane and near the AdS boundary. Near the brane one has $1 - z/z_0 \ll 1$ and the contribution of large x dominates in the integral of (3.32). For non-conformally coupled fields the leading term in the expansion of the energy density is given by (3.19) with the replacement $y - y_0 \rightarrow y_0 - y$ and the relation between the energy density and the normal stress remain the same. In particular, we see that near the brane the energy density has the same sign in the R- and L-regions, whereas the normal stresses have opposite signs. For points near the AdS boundary one has $z/z_0 \ll 1$ and the leading term in the asymptotic expansion is given by

$$\langle T_0^0 \rangle_{\rm b}^{(\rm s)} \approx \frac{(z/z_0)^{D+2\nu}}{\pi^{D/2}\Gamma(D/2)} \frac{(D+2\nu+1)\xi_1+1}{2^{D+2\nu}\Gamma(\nu)\Gamma(\nu+1)a^{D+1}} \int_0^\infty dx x^{D+2\nu-1} \frac{\bar{K}_{\nu}(x)}{\bar{I}_{\nu}(x)}.$$
 (3.35)

In the same order, the normal stress is found from $\langle T_D^D \rangle_{\rm b}^{\rm (s)} \approx -D \langle T_0^0 \rangle_{\rm b}^{\rm (s)}/(2\nu)$. Hence, the braneinduced VEVs vanish on the AdS boundary like $z^{D+2\nu}$. Note that the factor $(D+2\nu+1)\xi_1+1$ is negative for both minimally and conformally coupled field and for the Dirichlet boundary condition the energy density is negative near the AdS boundary. In general, depending on the Robin coefficient, the energy density can be either negative or positive.

The brane-induced energy density for D = 4 minimally coupled scalar field is presented on the left panel of figure 6 as a function of z/z_0 . The graphs are plotted for ma = 1, for the Dirichlet and Neumann boundary conditions (the letters near the curves), and for the Robin boundary conditions with $\beta/a = -0.5, -0.15, -0.1$. For the latter cases the energy density $\langle T_0^0 \rangle_b^{(s)}$ increases with increasing $|\beta/a|$. On the right panel we display the energy density and the normal stress as functions of the mass for the Dirichlet and Neumann boundary conditions, and for the Robin condition with $\beta/a = -0.15$. The graphs are plotted for $z/z_0 = 0.6$. For the Robin boundary condition the brane-induced VEVs may change the sign as functions of the mass.



Figure 6: The same as in figure 2 for a minimally coupled scalar field in the L-region. The right panel is plotted for $z/z_0 = 0.6$.

3.3.3 Electromagnetic field

For the electromagnetic field the eigenmodes of the quantum number λ are roots of the equation (2.37), where ν is given by (2.33) for the boundary conditions (2.31) and (2.32). The corresponding summation formula for the series in the mode-sum of the energy-momentum tensor is obtained from (3.28) by the replacement $ma - 1/2 \rightarrow \nu$. The brane-induced contribution is expressed as

$$\langle T^{\mu}_{\mu} \rangle_{\rm b}^{\rm (v)} = -\frac{\delta_{\nu} \left(D-1\right)}{2^{D} \pi^{D/2} \Gamma(D/2) a^{D+1}} \int_{0}^{\infty} dx \, x^{D+1} \frac{K_{\nu}(x z_{0}/z)}{I_{\nu}(x z_{0}/z)} V^{(\mu)}_{\rm (L)}(x), \tag{3.36}$$

where we have defined the functions

$$V_{(L)}^{(0)}(x) = \left(1 - \frac{4}{D}\right) I_{D/2-1}^2(x) + \left(1 - \frac{2}{D}\right) I_{D/2-2}^2(x),$$

$$V_{(L)}^{(D)}(x) = I_{D/2-1}^2(x) - I_{D/2-2}^2(x).$$
(3.37)

For $D \ge 4$ the energy density is negative for the boundary condition (2.31) and positive for the condition (2.32). For the normal stress one has $\langle T_D^D \rangle_{\rm b}^{({\rm v})} > 0$ in the case of the condition (2.31) and $\langle T_D^D \rangle_{\rm b}^{({\rm v})} < 0$ for (2.32). In the Minkowskian limit $a \to \infty$, with fixed y, y_0 , we obtain the result (3.23).

Unlike the R-region, in the L-region the brane-induced contribution is different from zero for D = 3. For the boundary condition (2.31) one gets

$$\langle T^{\rho}_{\mu} \rangle^{(v)}_{b} = -\frac{\pi^2}{720} \left(\frac{z}{az_0}\right)^4 \text{diag}(1, 1, 1, -3).$$
 (3.38)

This determines the Casimir force per unit surface of the brane equal to $-\pi^2 a^{-4}/240$. The latter is attractive with respect to the AdS boundary and does not depend on the location of the brane. In the case of the boundary condition (2.32) we obtain the expression

$$\langle T^{\rho}_{\mu} \rangle^{(v)}_{b} = -\frac{7\pi^{2}}{5760} \left(\frac{z}{az_{0}}\right)^{4} \operatorname{diag}(1,1,1,-3).$$
 (3.39)

The corresponding force per unit surface of the brane is given by $7\pi^2 \alpha^{-4}/1920$ and is repulsive with respect to the AdS boundary.

The near-brane asymptotics in the L-region are given by (3.24) with the replacement $y-y_0 \rightarrow y_0-y$ in the expression for the energy density. Near the AdS boundary one has $z/z_0 \ll 1$ and by using the expression of the modified Bessel function for small arguments, in the leading order we get

$$\langle T_0^0 \rangle_{\rm b}^{(\rm v)} \approx -\frac{8\delta_{\nu} \left(D-1\right) \left(z/2z_0\right)^{2D-2}}{\pi^{D/2} D \Gamma^3 (D/2-1) a^{D+1}} \int_0^\infty dx \, x^{2D-3} \frac{K_{\nu}(x)}{I_{\nu}(x)},$$

$$\langle T_D^D \rangle_{\rm b}^{(\rm v)} \approx -\frac{D}{D-2} \langle T_0^0 \rangle_{\rm b}^{(\rm v)}.$$

$$(3.40)$$

In spatial dimension D = 3 the integral is equal to $\pi^5/240$ for the boundary condition (2.31) and $7\pi^5/1920$ for (2.32). In this special case the asymptotics (3.40) coincide with the exact results (3.38) and (3.39).

For D = 4 and D = 5 electromagnetic fields, the brane contributions to the vacuum energy density and the normal stress are plotted on the left and right panels of figure 7 (the values of the index μ for $\langle T^{\mu}_{\mu} \rangle_{\rm b}^{(v)}$ are displayed near the curves). The full and dashed curves correspond to the boundary conditions (2.31) and (2.32), respectively.



Figure 7: The same as in figure 4 for the L-region.

4 Geometry with a brane perpendicular to the AdS boundary

In a number of recent developments of the AdS/CFT correspondence, branes intersecting the AdS boundary are considered. They include the extensions of the correspondence for conformal field theories with boundaries (AdS/BCFT correspondence) [52, 53] and the geometric procedure for the evaluation of the entanglement entropy for a bounded region in CFT [54, 55]. In this section, based on [23], we will consider the effects on the scalar vacuum induced by a brane perpendicular to the AdS boundary.

The background geometry is described by the line element (2.2) and the brane is located at $x^1 = 0$. The problem is symmetric with respect to the brane and we will consider the region $x^1 \ge 0$. The scalar field $\varphi(x)$ obeys the field equation (2.3) and the boundary condition

$$(\beta \partial_1 + 1) \varphi(x) = 0, \tag{4.1}$$

on the brane. Introducing the notations $\mathbf{x} = (x^2, \dots, x^{D-1})$ and $\mathbf{k} = (k_2, \dots, k_{D-1})$, the normalized mode functions obeying the boundary condition has the form

$$\varphi_{\sigma}^{(\pm)}(x) = \frac{\sqrt{2k_1/\omega}}{(2\pi a)^{(D-1)/2}} z^{D/2} J_{\nu}(\lambda z) \cos[k_1 x^1 + \alpha_0(k_1)] e^{i\mathbf{k}\mathbf{x}\mp i\omega t}, \tag{4.2}$$

where $0 \le k_1, \lambda < \infty$, the energy is given by $\omega = \sqrt{k^2 + k_1^2 + \lambda^2}$ with $k^2 = \sum_{i=2}^{D} k_i^2$ and ν is defined by (2.7). The function $\alpha_0(k_1)$ is defined by the relation

$$e^{2i\alpha_0(k_1)} = \frac{i\beta k_1 - 1}{i\beta k_1 + 1}.$$
(4.3)

Note that in the case $0 \le \nu < 1$ we impose the same boundary condition on the AdS boundary as that in section 2 for the L-region.

For positive values of the Robin parameter β in (4.1), in addition to (4.2), there is a mode for which the part depending on the coordinate x^1 is expressed in terms of the exponential function $e^{-x^1/\beta}$. The corresponding energy is given by $\omega = \sqrt{k^2 + \lambda^2 - 1/\beta^2}$. In the subspace $k^2 + \lambda^2 < 1/\beta^2$ of the quantum numbers the energy is imaginary and the vacuum state is unstable. Note that for a Robin plate in the Minkowski bulk the energy for the corresponding bound states is positive in the range $\beta > 1/m$. In order to have a stable vacuum in the AdS bulk we will assume that $\beta \leq 0$.

With the mode functions (4.2), the VEV of the energy-momentum tensor is evaluated by using the mode-sum formula (3.2). We write the parts containing the products of the trigonometric functions with the arguments $k_1x^1 + \alpha_0(k_1)$ in the form of the sum of exponents. After rotating the integration contour over k_1 for separate exponents by the angles $\pi/2$ and $-\pi/2$, the VEV of the energy-momentum tensor is presented in the decomposed form (3.3). For the diagonal components of the brane-induced part one gets the representation

$$\langle T^{\mu}_{\mu} \rangle_{\rm b}^{\rm (s)} = -\frac{(4\pi)^{(1-D)/2}}{2^{2\nu+1}a^{D+1}} \int_{0}^{\infty} dx \, x e^{-2xx^{1}/z} \frac{\beta x/z+1}{\beta x/z-1} \\ \times \left[A_{\mu}x^{D+2\nu} F_{\nu}^{D/2+1}(x) + \hat{B}_{\mu}(x)x^{D+2\nu} F_{\nu}^{D/2}(x) \right],$$

$$(4.4)$$

where $A_1 = 0$, $A_D = (1 - D)/2$, $A_l = 1/2$ for l = 0, 2, ..., D - 1. The operators $\hat{B}_{\mu}(x)$ are defined as

$$\hat{B}_{l}(x) = \frac{\xi_{1}}{4}\hat{B}(x) + \frac{\xi}{x}\left(\partial_{x} - \frac{D}{x}\right) - \xi_{1}\delta_{1l}, \ l = 0, 1, 2, \dots, D - 1,
\hat{B}_{D}(x) = \frac{1}{4}\hat{B}(x) - D\frac{\xi}{x}\left(\partial_{x} - \frac{D}{x}\right) - \frac{m^{2}a^{2}}{x^{2}} + \xi_{1},$$
(4.5)

with $\hat{B}(x) = \partial_x^2 - ((D-1)/x)\partial_x + 4$ and ξ_1 is given by (3.17). In (4.4) we have introduced the function

$$F_{\nu}^{\mu}(x) = \frac{2^{2\nu+1}x^{-2\nu}}{\Gamma(\mu-1/2)} \int_{0}^{1} du \, u(1-u^{2})^{\mu-3/2} J_{\nu}^{2}(xu)$$

$$= \frac{{}_{1}F_{2}\left(\nu+1/2;\nu+\mu+1/2,2\nu+1;-x^{2}\right)}{\Gamma(\nu+\mu+1/2)\Gamma(\nu+1)}, \qquad (4.6)$$

where ${}_{1}F_{2}(a; b, c; y)$ is the hypergeometric function. The diagonal components in the region $x^{1} < 0$ are given by the expression (4.4) with x^{1} replaced by $|x^{1}|$.

An important difference from the geometry with a brane parallel to the AdS boundary is the presence of the off-diagonal component of the vacuum energy-momentum tensor:

$$\langle T_D^1 \rangle_{\rm b}^{\rm (s)} = -\frac{(4\pi)^{(1-D)/2}}{2^{2\nu+2}a^{D+1}} \int_0^\infty dx \, e^{-2xx^1/z} \frac{\beta x/z+1}{\beta x/z-1} \left(\xi_1 x \partial_x + 4\xi\right) x^{D+2\nu} F_\nu^{D/2}(x). \tag{4.7}$$

This expression for the off-diagonal component in the region $x^1 < 0$ is obtained from (4.7) changing the sign and replacing $x^1 \to |x^1|$. Note that $\langle T_D^1 \rangle_{\rm b}^{({\rm s})} = \langle T_1^D \rangle_{\rm b}^{({\rm s})}$. Due to the nonzero off-diagonal component $\langle T_D^1 \rangle_{\rm b}^{({\rm s})}$, the Casimir force acting on the brane has two components. The first one is determined by the stress $\langle T_1^1 \rangle_{\rm b}^{({\rm s})}$ and corresponds to the component normal to the brane. The second one is obtained from

 $\langle T_D^1 \rangle_{\rm b}^{\rm (s)}$ and is directed along the z-axis. It corresponds to the shear force. Of course, because of the surface divergences in the local VEVs, both these components require an additional renormalization.

From the expressions (4.4) and (4.7) it follows that for the Dirichlet and Neumann boundary conditions the brane-induced contributions to the VEV of the energy-momentum tensor differ by the signs. In these special cases the corresponding expressions are further simplified as

$$\langle T^{\mu}_{\mu} \rangle_{\rm b}^{\rm (s)} = \pm \frac{a^{-D-1}}{2^{D/2+\nu+1}\pi^{D/2}} [\hat{C}_{\mu}(u) - D\xi] f_{\nu}(u),$$

$$\langle T^{1}_{D} \rangle_{\rm b}^{\rm (s)} = \pm \frac{2a^{-D-1}x^{1/2}}{2^{D/2+\nu+1}\pi^{D/2}} \left[\xi_{1} \left(u - 1 \right) \partial_{u}^{2} + \left(2\xi - 1 \right) \partial_{u} \right] f_{\nu}(u),$$
 (4.8)

where the upper and lower signs correspond to the Dirichlet and Neumann boundary conditions, respectively, and $u = 1+2(x^1/z)^2$. The operators $\hat{C}_{\mu}(u)$ in the expressions for the diagonal components are given by

$$\hat{C}_{l}(u) = \xi_{1} \left(u^{2} - 1 \right) \partial_{u}^{2} + \left[4\xi - 2 + \left(\frac{D+1}{2} \xi_{1} - \frac{1}{2} \right) (u-1) \right] \partial_{u},$$
(4.9)

for l = 0, 2, ..., D - 1, and

$$\hat{C}_{1}(u) = \xi_{1} (u-1)^{2} \partial_{u}^{2} + \left(\frac{D+1}{2}\xi_{1} - \frac{1}{2}\right) (u-1) \partial_{u},$$

$$\hat{C}_{D}(u) = 2\xi_{1} (u-1) \partial_{u}^{2} + \left[4\xi - 2 + \frac{D}{2}\xi_{1} (u-1)\right] \partial_{u}.$$
(4.10)

The function $f_{\nu}(u)$ is expressed in terms of the hypergeometric function $_{2}F_{1}(a,b;c;x)$:

$$f_{\nu}(u) = \frac{\Gamma(\nu + D/2)}{\Gamma(\nu + 1)u^{\nu + D/2}} {}_{2}F_{1}\left(\frac{D + 2\nu + 2}{4}, \frac{D + 2\nu}{4}; \nu + 1; \frac{1}{u^{2}}\right).$$
(4.11)

For a conformally coupled massless field one has $\nu = 1/2$ and the problem under consideration is conformally related to the problem in the Minkowski spacetime with the line element $ds^2 = \eta_{\mu\rho} dx^{\mu} dx^{\rho}$, $x^D = z$, and with planar codimension one boundaries located at z = 0 (the conformal image of the AdS boundary) and $x^1 = 0$ (the conformal image of the brane). The Minkowskian field obeys the Dirichlet boundary condition at z = 0 and the Robin boundary condition (4.1) at $x^1 = 0$. The Dirichlet boundary condition at z = 0 is a consequence of the condition we have imposed on the AdS boundary. Taking $\nu = 1/2$, from the results given above we can obtain the VEV of the energy-momentum tensor for a conformally coupled massless field in the geometry of perpendicular planar boundaries in the Minkowski bulk by using the relation $\langle T^{\rho}_{\mu} \rangle_{\rm b}^{({\rm M})} = (a/z)^{D+1} \langle T^{\rho}_{\mu} \rangle_{\rm b}$ (see (3.6)). In the Minkowskian limit we get the result 3.18 (with the replacement $y - y_0 \to x^1$) for the components $\mu = 0, 2, \ldots, D$.

Near the brane and for non-Dirichlet boundary conditions, $x^1 \ll z, |\beta|$, the leading term in the asymptotic expansions for diagonal components with $\mu = 0, 2, ..., D$ is given by

$$\langle T^{\mu}_{\mu} \rangle^{(s)}_{b} \approx -\frac{D\Gamma((D+1)/2)\left(\xi - \xi_{D}\right)}{2^{D}\pi^{(D+1)/2}\left(ax^{1}/z\right)^{D+1}}.$$
(4.12)

This coincides with the Minkowskian result where the distance from the boundary is replaced by the ratio ax^1/z (compare with (3.19)). Note that, in accordance with (2.2), the latter is the proper distance from the brane measured by an observer at rest with respect to the brane. Note that the proper distance ax^1/z is different from the geodesic distance $\sigma(x, x')$. The latter between the spacetime points $x = (t, 0, \mathbf{x}, z)$ and $x' = (t, x^1, \mathbf{x}, z)$ is given as $\cosh(\sigma(x, x')/a) = 1 + (x^1/z)^2/2$. For the normal stress and for the off-diagonal component near the brane one gets

$$\langle T_1^1 \rangle_{\rm b}^{\rm (s)} \approx -\frac{\left(x^1/z\right)^2}{D-1} \langle T_0^0 \rangle_{\rm b}^{\rm (s)}, \ \langle T_D^1 \rangle_{\rm b}^{\rm (s)} \approx \frac{x^1}{z} \langle T_0^0 \rangle_{\rm b}^{\rm (s)}.$$
 (4.13)

The leading terms for the Dirichlet boundary condition differ from the ones given above by the sign. The expressions (4.12) and (4.13) describe also the asymptotic behavior of the brane-induced VEV near the AdS horizon (large values of z for fixed given x^1).

Now let us consider the asymptotics at large distances from the brane, $x^1 \gg z$. For non-Neumann boundary conditions, additionally assuming that $x^1 \gg |\beta|$, for the components $\mu = 0, \ldots, D-1$ the leading term is given by

$$\langle T^{\mu}_{\mu} \rangle_{\rm b}^{\rm (s)} \approx \frac{(\xi - 1/4) (D + 2\nu) + \xi}{\pi^{D/2} \Gamma(\nu) a^{D+1} (2x^{1}/z)^{D+2\nu}} \Gamma(D/2 + \nu).$$
 (4.14)

The asymptotics for the remaining components are expressed as

$$\langle T_D^D \rangle_{\rm b}^{\rm (s)} \approx -\frac{D}{2\nu} \langle T_0^0 \rangle_{\rm b}^{\rm (s)}, \ \langle T_D^1 \rangle_{\rm b}^{\rm (s)} \approx \frac{D/2 + \nu}{\nu x^1/z} \langle T_0^0 \rangle_{\rm b}^{\rm (s)}.$$
 (4.15)

As seen, the leading order terms for $0 < |\beta| < \infty$ coincide with those for the Dirichlet boundary condition. For a scalar field with the Neumann boundary condition $(\beta = 0)$, the leading terms differ from those for the Dirichlet condition by the signs. Hence, the Dirichlet boundary condition is the attractor in a class of Robin boundary conditions with $\beta \neq 0$. At large distances, the brane-induced contributions, considered as functions of the proper distance from the brane, exhibit a power-law fall-off for both massless and massive fields. This behavior is in clear contrast with the case of the Minkowski bulk, where the boundary-induced VEVs decay exponentially for massive fields, like e^{-2mx^1} (see (3.18)). Note that for large x^1/z one has the relation $(x^1/z)^2 = \exp[\sigma(x, x')/a]$, with $\sigma(x, x')$ being the geodesic distance. For fixed x^1 , the asymptotic formulas (4.14) and (4.15) describe the behavior of the VEVs near the AdS boundary. The diagonal components decay as $z^{D+2\nu}$, whereas the off-diagonal component tends to zero like $z^{D+2\nu+1}$. The qualitative behavior of the brane-induced energy density, as a function of the distance from the brane, is similar to that we have described in the previous section for a scalar field with the Robin boundary condition on the brane parallel to the AdS boundary.

5 Summary

We have considered the influence of a brane in AdS bulk on the properties of quantum vacuum. Two geometries are discussed: (i) a brane parallel to the AdS boundary and (ii) a brane perpendicular to the AdS boundary. In the first geometry, as a local characteristic of the vacuum state, the VEV of the energy-momentum tensor is investigated for scalar, Dirac and electromagnetic fields. For scalar field a general Robin boundary condition is considered and the Dirac field is constrained by the bag boundary condition. In the case of the electromagnetic field two types of boundary conditions are discussed. The first one corresponds to the perfect conductor boundary conditions in 3D electrodynamics and the second one is the analog of the boundary condition used in bag models of hadrons to confine the gluons. The VEV of the energy-momentum tensor is expressed as a mode-sum over complete set of mode functions and for all these cases the corresponding sets are given. The brane divides the background geometry into two regions: the region between the brane and AdS horizon (R-region) and the region between the brane and AdS boundary (L-region). Though the AdS spacetime is homogeneous, the brane has a nonzero extrinsic curvature tensor and the properties of the quantum vacuum in those regions are different. In particular, the spectrum of the quantum number λ , corresponding to the momentum along the direction normal to the AdS boundary, is continuous in the R-region and discrete in the L-region. In the latter region the eigenvalues are zeros of cylinder functions. The mode-sum for the VEV of the energy-momentum tensor contains series over those zeros and for the summation we have employed the generalized Abel-Plana formula. That allowed to extract from the VEV the part corresponding to the geometry without a brane and to present the brane-induced contribution in terms of integral, exponentially convergent for points away form the brane. A similar decomposition is provided for the R-region.

Near the brane, the leading terms in the asymptotic expansions for the energy density and parallel stresses coincide with the corresponding expressions for a single boundary in the Minkowski bulk, where the distance from the boundary is replaced by the proper distance from the brane on the AdS bulk. For those VEVs the effect of gravity is weak. This is related to the fact that near the brane the main contribution to the corresponding VEVs come from the vacuum fluctuations with the wavelengths smaller than the curvature radius of the background geometry and influence of the gravitational field on those modes is weak. For a boundary in the Minkowski bulk the normal stress is zero. The nonzero normal stress in the geometry of a brane on the AdS bulk is a purely gravitational effect. The effect of gravity on the brane-induced VEVs is essential at distances from the brane larger than the curvature radius. In particular, for the R-region, at large distances the decay of the brane-induced contribution in the vacuum energy-momentum tensor, as a function of the proper distance, is exponential for both massless and massive fields. For the Minkowski bulk and for massless fields the fall-off of the boundaryinduced contribution is as power-law. On the AdS boundary, the brane-induced contributions tend to zero like $z^{D+\beta}$, where $\beta = 2\nu$ for scalar field (with ν given by (2.7)), $\beta = 2ma + 1$ for the Dirac field, and $\beta = D - 2$ for the electromagnetic field. Near the AdS boundary one has a simple relation between the energy density and the normal stress, given by $\langle T_D^D \rangle_{\rm b} \approx -(D/\beta) \langle T_0^0 \rangle_{\rm b}$. This correspond to the barotropic equation of state for the vacuum pressure $-\langle T_D^D \rangle_{\rm b}$ along the z-direction and vacuum energy density. Note that for the pressures along the directions parallel to the brane the equation of state is of the cosmological constant type.

The brane-induced effects on the quantum vacuum for the geometry (ii) we have considered on the example of a scalar field with general curvature coupling parameter. For the Robin boundary condition the mode functions have the form (4.2). The diagonal components of the brane-induced energy-momentum tensor are given by the expressions (4.4). An important difference from the problem with a brane parallel to the AdS boundary is the presence of nonzero off-diagonal component (4.7) of the vacuum energy-momentum tensor. As a consequence, the Casimir force acting on the brane, in addition to the normal component, contains a component directed parallel to the brane (shear force). At large distances from the brane, the decay of the brane-induced contribution to the energymomentum tensor, as a function of the proper distance from the brane, is as power-law for both massive and massless field. As it has been mentioned above, in the Minkowski bulk the decay for massive fields is exponential.

In the discussion above we have considered the VEV of the bulk energy-momentum tensor. In manifolds with boundaries, in addition to the latter a surface energy-momentum tensor may present which is localized on the boundaries. In the general case of bulk and boundary geometries, the expression of the surface energy-momentum tensor for a scalar field with general curvature coupling has been obtained in [56] by using the standard variational procedure. The VEV of the surface energy-momentum tensor for branes parallel to the AdS boundary is investigated in [57] by using the generalized zeta function technique. It has been shown that in the geometry of two branes the surface energy density induced on the visible brane by the presence of the hidden brane gives rise to naturally suppressed cosmological constant. The corresponding surface densities in models with additional compact dimensions are discussed in [58, 59] for neutral and charged scalar fields. In the latter case, the value of the induced cosmological constant on the brane is controlled by tuning the magnetic flux enclosed by compact dimensions.

For charged fields, another important local characteristic of the vacuum state is the expectation value of the current density. This VEV in models with local AdS geometry and with toroidally compact spatial dimensions, in the presence of single and two branes has been investigated in [30, 60] and [12, 34] for charged scalar and fermionic fields. The vacuum currents have nonzero components along the compact dimensions only. They are periodic functions of the magnetic flux with the period equal to the flux quantum. Depending on the boundary conditions imposed on the fields at the locations of the branes, the brane-induced effects lead to increase or decrease of the current density. Applications were discussed to Randall-Sundrum type braneworld models and also to curved graphene

tubes.

Acknowledgments

The author is grateful to Stefano Bellucci, Eugenio Bezerra de Mello, Emilio Elizalde, Sergei Odintsov, Mohammad Setare and Valery Vardanyan for collaboration.

References

- Callan, C.G.; Wilczek, F. Infrared behavior at negative curvature. Nucl. Phys. B 1990, 340, 366-386.
- [2] Aharony, O.; Gubser, S.S.; Maldacena, J.; Ooguri, H.; Oz, Y. Large N field theories, string theory and gravity. *Phys. Rep.* 2000, 323, 183-386.
- [3] Năstase, H. Introduction to AdS/CFT correspondence; Cambridge University Press: Cambridge, UK, 2015.
- [4] Ammon, M.; Erdmenger, J. Gauge/Gravity Duality: Foundations and Applications; Cambridge University Press: Cambridge, UK, 2015.
- [5] Pires, A.S.T. AdS/CFT Correspondence in Condensed Matter; Morgan & Claypool Publishers: USA, 2014.
- [6] Zaanen, J.; Sun, Y.-W.; Liu, Y.; Schalm, K. Holographic Duality in Condensed Matter Physics; Cambridge University Press: Cambridge, UK, 2015.
- [7] Maartens, R.; Koyama, K. Brane-World Gravity. Living Rev. Relativity, 2010, 13, 1.
- [8] Mostepanenko, V.M.; Trunov, N.N. The Casimir Effect and Its Applications; Clarendon Press: Oxford, UK, 1997.
- [9] Milton, K.A. The Casimir Effect: Physical Manifestation of Zero-Point Energy; World Scientific: Singapore, 2002.
- [10] Bordag, M.; Klimchitskaya, G.L.; Mohideen, U.; Mostepanenko, V.M. Advances in the Casimir Effect; Oxford University Press: New York, USA, 2009.
- [11] Dalvit, D.; Milonni, P.; Roberts, D.; da Rosa, F. (Eds.) Casimir Physics; Lecture Notes in Physics; Springer-Verlag: Berlin, Germany, 2011; Volume 834.
- [12] Bellucci, S.; Saharian, A.A.; Sargsyan, H.G.; Vardanyan, V.V. Fermionic vacuum currents in topologically nontrivial braneworlds: Two-brane geometry. *Phys. Rev. D* 2020, 101, 045020.
- [13] Saharian, A.A.; Kotanjyan, A.S.; Sargsyan, H.G. Electromagnetic field correlators and the Casimir effect for planar boundaries in AdS spacetime with application in braneworlds; arXiv:2009.02072.
- [14] Saharian, A.A.; Setare, M.R. The Casimir effect on background of conformally flat brane-world geometries. *Phys. Lett. B* 2003, 552, 119-126.
- [15] Knapman, A.; Toms, D.J. Stress-energy tensor for a quantized bulk scalar field in the Randall-Sundrum brane model. *Phys. Rev. D* 2004, 69, 044023.

- [16] Saharian, A.A. Wightman function and Casimir densities on AdS bulk with application to the Randall-Sundrum braneworld. Nucl. Phys. B 2005, 712, 196-228.
- [17] Saharian, A.A.; Mkhitaryan, A.L. Wightman function and vacuum densities for a Z₂-symmetric thick brane in AdS spacetime. J. High Energy Phys. 2007, 08, 063.
- [18] Shao, S.-H.; Chen, P.; Gu, J.-A. Stress-energy tensor induced by a bulk Dirac spinor in the Randall-Sundrum model. *Phys. Rev. D* 2010, *81*, 084036.
- [19] Elizalde, E.; Odintsov, S.D.; Saharian, A.A. Fermionic Casimir densities in anti-de Sitter spacetime. Phys. Rev. D 2013, 87, 084003.
- [20] Saharian, A.A.; Kotanjyan, A.S.; Saharyan, A.A. Proceedings of the Yerevan State University. Physical and Mathematical Sciences 2016, 3, 37-41.
- [21] Kotanjyan, A.S.; Saharian, A.A. Electromagnetic quantum effects in anti-de Sitter spacetime. *Physics of Atomic Nuclei* 2017, 80, 562-571.
- [22] Kotanjyan, A.S.; Saharian, A.A.; Saharyan, A.A. Electromagnetic Casimir Effect in AdS Spacetime. *Galaxies* 2017, 5, 102.
- [23] Bezerra de Mello, E.R.; Saharian, A.A.; Setare, M. R. Vacuum densities for a brane intersecting the AdS boundary. *Phys. Rev. D* 2015, *92*, 104005.
- [24] Saharian, A.A. Wightman function and vacuum fluctuations in higher dimensional brane models. *Phys. Rev. D* 2006, 73, 044012.
- [25] Saharian, A.A. Bulk Casimir densities and vacuum interaction forces in higher dimensional brane models. *Phys. Rev. D* 2006, 73, 064019.
- [26] Flachi, A.; Garriga, J.; Pujolàs, O.; Tanaka, T. Moduli stabilization in higher dimensional brane models. J. High Energy Phys. 2003, 08, 053.
- [27] Flachi, A.; Pujolàs, O. Quantum self-consistency of AdS×Σ brane models. Phys. Rev. D 2003, 68, 025023.
- [28] Breitenlohner, P.; Freedman, D.Z. Stability in gauged extended supergravity. Ann. Phys. (NY) 1982, 144 (2), 249-281.
- [29] Mezincescu, L.; Townsend, P.K. Stability at a local maximum in higher dimensional anti-deSitter space and applications to supergravity. Ann. Phys. (NY) 1985, 160 (2), 406-419.
- [30] Bellucci, S.; Saharian, A.A.; Vardanyan, V. Vacuum currents in braneworlds on AdS bulk with compact dimensions. J. High Energy Phys. 2015, 11, 092.
- [31] Avis, S.J.; Isham, C.J.; Storey, D. Quantum field theory in anti-de Sitter space-time. Phys. Rev. D 1978, 18, 3565-3576.
- [32] Ishibashi, A.; Wald, R.M. Dynamics in non-globally-hyperbolic static spacetimes: III. Anti-de Sitter spacetime. Class. Quantum Grav. 2004, 21, 2981.
- [33] Morley, T.; Taylor, P.; Winstanley, E. Quantum field theory on global anti-de Sitter space-time with Robin boundary conditions; arXiv:2004.02704.
- [34] Bellucci, S.; Saharian, A.A.; Simonyan, D.H.; Vardanyan, V. Fermionic currents in topologically nontrivial braneworlds. *Phys. Rev. D* 2018, 98, 085020.

- [35] Randall, L.; Sundrum, R. Large mass hierarchy from a small extra dimension. Phys. Rev. Lett. 1999, 83, 3370-3373.
- [36] Randall, L.; Sundrum, R. An alternative to compactification. Phys. Rev. Lett. 1999, 83, 4690-4693.
- [37] Gherghetta, T.; Pomarol, A. Bulk fields and supersymmetry in a slice of AdS. Nucl. Phys. B 2000, 586, 141-162.
- [38] Flachi, A.; Toms, D.J. Quantized bulk scalar fields in the Randall-Sundrum brane-model. Nucl. Phys. B 2001, 610, 144-168.
- [39] Flachi, A.; Moss, I.G.; Toms, D.J. Quantized bulk fermions in the Randall-Sundrum brane model. *Phys. Rev. D* 2001, 64, 105029.
- [40] Chan, S.; Park, S.Ch.; Song, J. Kaluza-Klein masses of bulk fields with general boundary conditions in AdS₅ space. *Phys. Rev. D* 2005, 71, 106004.
- [41] Grib, A.A.; Mamayev, S.G.; Mostepanenko, V.M. Vacuum Quantum Effects in Strong Fields; Friedmann Laboratory Publishing: St. Petersburg, 1994.
- [42] Birrell, N.D.; Davies, P.C.W. Quantum Fields in Curved Space; Cambridge University Press: Cambridge, England, 1982.
- [43] Buchbinder, I.L., Odintsov, S.D.; Shapiro, I.L. *Effective Action in Quantum Gravity*; Taylor & Francis: BocaRaton, 1992.
- [44] Elizalde, E.; Odintsov, S.D.; Romeo, A.; Bytsenko, A.A.; Zerbini, S. Zeta Regularization Techniques with Applications; World Scientific: Singapore, 1994.
- [45] Kirsten, K. Spectral Functions in Mathematics and Physics; CRC Press: Boca Raton, FL, 2001.
- [46] Bytsenko, A.A.; Cognola, G.; Elizalde, E.; Moretti, V.; Zerbini, S. Analytic Aspects of Quantum Fields; World Scientific: Singapore, 2003.
- [47] Kent, C.; Winstanley, E. Hadamard renormalized scalar field theory on anti-de Sitter spacetime. *Phys. Rev. D* 2015, 91, 044044.
- [48] Ambrus, V.E.; Winstanley, E. Renormalised fermion vacuum expectation values on anti-deSitter space-time. *Phys. Lett. B* 2015, 749, 597-602.
- [49] Prudnikov, A.P.; Brychkov, Yu.A.; Marichev, O.I. Integrals and series; Gordon and Breach: New York, 1986, Vol.2.
- [50] Saharian, A.A. A generalized Abel-Plana formula. Applications to cylindrical functions. Izv. Akad. Nauk Arm. SSR Mat. 1987, 22, 166-179 [Sov. J. Contemp. Math. Anal. 1987, 22, 70-86].
- [51] Saharian, A.A. The Generalized Abel-Plana Formula with Applications to Bessel Functions and Casimir Effect; Yerevan State University Publishing House: Yerevan, 2008; Report No. ICTP/2007/082; arXiv:0708.1187.
- [52] Takayanagi, T. Holographic Dual of a Boundary Conformal Field Theory. Phys. Rev. Lett. 2011, 107, 101602.
- [53] Fujita, M.; Takayanagi, T.; Tonni, E. Aspects of AdS/BCFT. J. High Energy Phys. 2011, 11, 043.

- [54] Ryu, S; Takayanagi, T. Holographic Derivation of Entanglement Entropy from the anti-de Sitter Space/Conformal Field Theory Correspondence. *Phys. Rev. Lett.* 2006, *96*, 181602.
- [55] Ryu, S; Takayanagi, T. Aspects of holographic entanglement entropy. J. High Energy Phys. 2006, 08, 045.
- [56] Saharian, A.A. Energy-momentum tensor for a scalar field on manifolds with boundaries. *Phys. Rev. D* 2004, 69, 085005.
- [57] Saharian, A.A. Surface Casimir densities and induced cosmological constant on parallel branes in AdS spacetime *Phys. Rev. D* 2004, 70, 064026.
- [58] Saharian, A.A. Surface Casimir densities and induced cosmological constant in higher dimensional braneworlds. *Phys. Rev. D* 2006, 74, 124009.
- [59] Saharian, A.A.; Sargsyan, H. G. Induced Cosmological Constant in Brane Models with a Compact Dimension. Astrophysics 2018, 61, 375–390.
- [60] Bellucci, S.; Saharian, A.A.; Vardanyan, V. Hadamard function and the vacuum currents in braneworlds with compact dimensions: Two-brane geometry. *Phys. Rev. D* 2016, *93*, 084011.