

On an extension of the generalized BGW tau-function

Di Yang, Chunhui Zhou

Dedicated to the memory of Boris Anatol'evich Dubrovin, with admiration

Abstract

For an arbitrary solution to the Burgers-KdV hierarchy, we define the tau-vector (τ_1, τ_2) of the solution. We then introduce certain linear maps and derive an explicit formula for the product $\tau_1\tau_2$. Applications to an extension of the generalized BGW tau-function and to the open partition function are given.

1 Introduction

The Brézin–Gross–Witten (BGW) partition function was introduced in [9, 19]. Recently, Norbury [27] constructed a new cohomology class on the moduli space of closed Riemann surfaces. It has been proved [27] that the partition function of the intersection numbers of the Norbury class coupling ψ -classes coincides with the BGW partition function. Recalling another recent construction about the moduli space of open Riemann surfaces [28] (cf. also [11]), we consider in this article an extension of the BGW partition function, that should serve as the partition function of intersection numbers of certain extended version of the Norbury class coupling the descendants on the open and closed Riemann surfaces. Considering that there is a one-parameter generalization of the BGW partition function given in [1], we will perform the extension for an arbitrary value of the parameter.

Denote by $\mathcal{A}_{(w,\rho)}$ the ring of polynomials of $w_x, \rho_x, w_{xx}, \rho_{xx}, \dots$ with coefficients being smooth functions of w, ρ . The Burgers-KdV integrable hierarchy [10] is defined as the following pairwise commuting system of non-linear PDEs:

$$\frac{\partial w}{\partial q_{2n-1}} = \partial_x K_{2n-1}, \quad \frac{\partial w}{\partial q_{2n}} = 0, \quad n \geq 1, \quad (1.1)$$

$$\frac{\partial \rho}{\partial q_{2n-1}} = \partial_x R_{2n-1}, \quad \frac{\partial \rho}{\partial q_{2n}} = \partial_x R_{2n}, \quad n \geq 1, \quad (1.2)$$

where $K_n, R_n \in \mathcal{A}_{(w,\rho)}$ are defined recursively by

$$K_1 = w, \quad K_2 = 0, \quad R_1 = \rho, \quad R_2 = \rho^2 + \rho_x + 2w, \quad (1.3)$$

$$\partial_x K_{2n+1} = \left(2w\partial_x + w_x + \frac{1}{4}\partial_x^3 \right) K_{2n-1}, \quad (1.4)$$

$$R_{2n+1} = (\partial_x^2 + 2\rho\partial_x + \rho^2 + \rho_x + 2w) R_{2n-1} + \left(\rho + \frac{3}{2}\partial_x \right) K_{2n-1}, \quad (1.5)$$

$$R_{2n+2} = (\partial_x^2 + 2\rho\partial_x + \rho^2 + \rho_x + 2w) R_{2n}, \quad n \geq 0. \quad (1.6)$$

The $n = 1$ flow reads

$$\frac{\partial w}{\partial q_1} = w_x \quad \frac{\partial \rho}{\partial q_1} = \rho_x.$$

Therefore, we can identify the variable q_1 with x . Note that equations (1.1) are the KdV integrable hierarchy. Hence, the Burgers-KdV hierarchy can be viewed as an integrable extension of (1.1); for more details about the Burgers-KdV hierarchy see [10].

Recall that the canonical tau structure [17, 16] of the KdV hierarchy is defined as the unique family of elements $(\Omega_{i,j})_{i,j \geq 1} \in \mathcal{A}_w$ satisfying

$$\Omega_{1,1} = w, \quad \Omega_{i,j} = \Omega_{j,i}, \quad \frac{\partial \Omega_{i,j}}{\partial q_{2k-1}} = \frac{\partial \Omega_{i,k}}{\partial q_{2j-1}}, \quad \forall i, j, k \geq 1 \quad (1.7)$$

along with the normalization condition $\Omega_{i,j}|_{w_{kx}=0, k \geq 0} = 0$. Here, $\mathcal{A}_w \subset \mathcal{A}_{(w,\rho)}$ denotes the ring of polynomials of w_x, w_{xx}, \dots with coefficients being smooth functions in w . For the existence of $\Omega_{i,j}$ see e.g. [17, 16]. Inspired by the open and closed intersection numbers [7, 10, 11, 28] and by the tau-structure of the Burgers hierarchy [15], we will define the tau-vector structure for the Burgers-KdV hierarchy. Indeed, observe that

$$\frac{\partial R_i}{\partial q_j} = \frac{\partial R_j}{\partial q_i}, \quad \forall i, j \geq 1. \quad (1.8)$$

Combining with (1.7) we arrive at the following lemma.

Lemma 1.1 *For an arbitrary solution (w, ρ) in $\mathbb{C}[[\mathbf{q}]]^2$ to the Burgers-KdV hierarchy, there exists $(\tau_1, \tau_2) \in \mathbb{C}[[\mathbf{q}]]^2$, such that*

$$\frac{\partial^2 \log \tau_1}{\partial q_{2i-1} \partial q_{2j-1}} = \Omega_{i,j}, \quad \frac{\partial^2 \log \tau_1}{\partial q_{2i} \partial q_j} = 0, \quad \forall i, j \geq 1, \quad (1.9)$$

$$\frac{\partial \log \tau_2}{\partial q_i} = R_i, \quad \forall i \geq 1. \quad (1.10)$$

Moreover, the vector (τ_1, τ_2) is uniquely determined by (w, ρ) up to the freedom described by

$$(\tau_1, \tau_2) \mapsto \left(\tau_1 e^{\beta_0 + \sum_{n \geq 1} \beta_n q_n}, \tau_2 e^{\alpha_0} \right), \quad (1.11)$$

where $\beta_n, n \geq 0$ and α_0 are an arbitrary constant.

We call (τ_1, τ_2) the *tau-vector* of the solution (w, ρ) to the Burgers-KdV hierarchy, and call the product $\tau_1 \cdot \tau_2 =: \tau_{\mathbb{E}}$ the tau-function for the Burgers-KdV hierarchy. Equations (1.9)–(1.10) for $i = j = 1$ read

$$w = \partial_x^2 \log \tau_1, \quad \rho = \partial_x \log \tau_2. \quad (1.12)$$

As it was proven in [5], the Dubrovin-Zhang type tau-function τ_1 for the KdV hierarchy is also a Sato type tau-function; vice versa. To proceed we first recall some notations. Denote by \mathbb{Y} the set of all the partitions. For each partition $\lambda \in \mathbb{Y}$, denote by $\ell(\lambda)$ the length of λ , by $|\lambda|$ the weight, and by $(m_1, \dots, m_r | n_1, \dots, n_r)$ the Frobenius notation (cf. [25]). The Schur polynomial associated to λ is defined as follows:

$$s_\lambda := \det (h_{\lambda_i - i + j})_{1 \leq i, j \leq \ell(\lambda)}, \quad (1.13)$$

where $\{h_k\}_{k \geq 0}$ are polynomials in $\mathbb{C}[\mathbf{q}]$ defined by the generating function

$$\sum_{k \geq 0} h_k z^k = \exp \left(\sum_{n \geq 1} q_n z^n \right). \quad (1.14)$$

According to the Sato theory, the component τ_1 , can be expressed as the linear combination of the Schur polynomials with coefficients being the Plücker coordinates, i.e.,

$$\tau_1 = c_0 \sum_{\lambda \in \mathbb{Y}} \pi_\lambda s_\lambda. \quad (1.15)$$

Here, c_0 is a nonzero constant that is irrelevant of the study and is often taken as 1, and

$$\pi_\lambda := (-1)^{\sum_{i=1}^r n_i} \det (A_{m_i, n_j})_{1 \leq i, j \leq r} \quad (1.16)$$

with $A_{m, n}$, $m, n \geq 0$ being the *affine coordinates* [18, 32, 7, 33] for the point of Sato Grassmanian corresponding to τ_1 . In this paper, we will derive explicit expressions for $\tau_1 \tau_2$.

Before presenting the results, we introduce further some notations. Note that the Schur polynomials form a basis of $\mathbb{C}[[\mathbf{q}]]$, and define a sequence of linear operators $T_n : \mathbb{C}[[\mathbf{q}]] \rightarrow \mathbb{C}[[\mathbf{q}]]$ in the following way:

$$T_n(s_\lambda) = \begin{cases} (-1)^i s_{(\lambda_1-1, \dots, \lambda_{i-1}-1, n+i, \lambda_{i+1}, \dots, \lambda_{\ell(\lambda)})}, & \text{if } \lambda_i > n+i \geq \lambda_{i+1}, \text{ for some } i, \\ 0, & \text{otherwise.} \end{cases} \quad (1.17)$$

Here we set $\lambda_0 = +\infty$ and $\lambda_{\ell(\lambda)+1} = 0$. The operator T_0 has the alternative expression:

$$T_0(s_{(m_1, \dots, m_r | n_1, \dots, n_r)}) = \begin{cases} (-1)^r s_{(m_1-1, \dots, m_r-1 | n_1+1, \dots, n_r+1)} & m_r \geq 1, \\ 0 & m_r = 0. \end{cases} \quad (1.18)$$

Note that for each $n \geq 0$, the linear operator T_n induces the linear transformation on $\text{Span}_{\mathbb{C}}(\mathbb{Y})$, denoted again by T_n . The first result of the paper is given by the following proposition.

Proposition 1.2 *Let (w, ρ) be an arbitrary solution in $\mathbb{C}[[\mathbf{q}]]^2$ to the Burgers-KdV hierarchy (1.1)–(1.2), and $(\tau_1, \tau_2) \in \mathbb{C}[[\mathbf{q}]]^2$ its tau-vector with $\tau_2(\mathbf{0})$ chosen as 1. There exists a unique sequence of numbers b_1, b_2, b_3, \dots , such that*

$$\tau_1 \cdot \tau_2 = Q \circ T_0 \circ Q^{-1}(\tau_1), \quad (1.19)$$

where $Q = e^{\sum_{k \geq 1} k b_k q_k}$ denotes the multiplication operator on $\mathbb{C}[[\mathbf{q}]]$.

The proof is in Section 3.

Noticing that $Q \circ T_0 \circ Q^{-1}$ is a linear operator and using (1.15), we will simplify (1.19) via computing the action of $Q \circ T_0 \circ Q^{-1}$ on each single Schur polynomial. Let us present the results here, and leave the details of calculations to Section 3. For $\lambda \in \mathbb{Y}$, define $\pi_{-\lambda} = -\pi_\lambda$. Let $c_{\lambda\mu}^\nu$, $\lambda, \mu, \nu \in \mathbb{Y}$ denote the Littlewood-Richardson coefficients [24]:

$$s_\lambda \cdot s_\mu =: \sum_{\nu \in \mathbb{Y}} c_{\lambda\mu}^\nu s_\nu. \quad (1.20)$$

For $\lambda, \mu, \nu \in \mathbb{Y}$, we set $c_{\lambda, -\mu}^\nu := -c_{\lambda, \mu}^\nu$. Define a sequence of numbers $g_0, g_1, g_2, g_3, \dots$ by

$$\sum_{n \geq 0} g_n z^{-n} := e^{\sum_{k \geq 1} b_k z^{-k}}. \quad (1.21)$$

Proposition 1.3 *The tau-function τ_E has the expression*

$$\tau_E = \sum_{\lambda \in \mathbb{Y}} s_\lambda \sum_{\substack{0 \leq k \leq |\lambda| \\ \mu \in T_k^{-1}(\lambda)}} g_k \pi_\mu. \quad (1.22)$$

Alternatively, denoting $\mathbf{b} := (b_1, b_2, b_3, \dots)$, we have

$$\tau_E = \sum_{\lambda \in \mathbb{Y}} s_\lambda \sum_{\substack{\alpha, \mu, \nu \in \mathbb{Y} \\ \beta \in \mathbb{Y} \setminus \text{Ker}(T_0)}} \pi_\alpha c_{\mu\alpha}^\beta c_{\nu, T_0(\beta)}^\lambda s_\mu(-\mathbf{b}) s_\nu(\mathbf{b}). \quad (1.23)$$

Observe that for a hook partition λ with the Frobenius notation $(m|n)$,

$$T_k^{-1}(\lambda) = \begin{cases} \{\emptyset\}, & n = 0, k = m + 1, \\ \{(0|n-1)\}, & n \geq 1, k = m + 1, \\ \{-(m+1|n-1)\}, & n \geq 1, k = 0, \\ \emptyset, & \text{otherwise.} \end{cases} \quad (1.24)$$

Then from (1.22) we see that the coefficient of $s_{(m|n)}$ in τ_E , denoted by $c_{m,n}$, has the more explicit expression:

$$c_{m,n} = \begin{cases} g_{m+1}, & n = 0, \\ (-1)^n (g_{m+1} A_{0,n-1} - A_{m+1,n-1}), & n \geq 1. \end{cases} \quad (1.25)$$

Together with Theorem B.1 and the formula similar to (1.15) for the KP hierarchy, we have

Corollary 1.4 *The tau-function τ_E has the following explicit expression:*

$$\tau_E = \sum_{\lambda \in \mathbb{Y}} \pi'_\lambda s_\lambda. \quad (1.26)$$

Here for a partition $\lambda = (m_1, \dots, m_r | n_1, \dots, n_r)$, $\pi'_\lambda = \det(c_{m_i, n_j})_{1 \leq i, j \leq r}$.

We proceed with defining the extension of the generalized BGW tau-function. Consider the solution to the Burgers-KdV hierarchy (1.1)-(1.2) specified by the initial condition

$$w|_{q_2=q_3=\dots=0} = \frac{C}{(1-x)^2}, \quad \rho|_{q_2=q_3=\dots=0} = 0, \quad (1.27)$$

where C is an arbitrarily given constant ($N = \sqrt{\frac{1}{4} - 2C}$ corresponds to the parameter in [3]). We call this solution the generalized BGW solution to the Burgers-KdV hierarchy, denoted by $(w^{\Theta(C)}, \rho^{\Theta(C)})$. Let $(\tau_1^{\Theta(C)}, \tau_2^{\Theta(C)})$ be the tau-vector of $(w^{\Theta(C)}, \rho^{\Theta(C)})$. As in [3, 6, 14, 16] we normalize the first component $\tau_1^{\Theta(C)}$ by

$$\sum_{k \geq 0} (2k+1)(q_{2k+1} - \delta_{k,0}) \frac{\partial \tau_1^{\Theta(C)}}{\partial q_{2k+1}} + C \tau_1^{\Theta(C)} = 0. \quad (1.28)$$

and by requiring $\partial \tau_1^{\Theta(C)} / \partial q_{2n} = 0$, call it the generalized BGW tau-function. We normalize the second component $\tau_2^{\Theta(C)}$ by requiring

$$\tau_2^{\Theta(C)}(\mathbf{0}) = 1. \quad (1.29)$$

We note that when the parameter C is taken to be $1/8$, the first component $\tau_1^{\Theta(C)}$ becomes the BGW tau-function. Norbury [27] proves the following identity:

$$\tau_1^{\Theta(\frac{1}{8})} = \exp\left(\sum_{n \geq 1} \sum_{i_1, \dots, i_n \geq 0} \frac{1}{n!} \prod_{k=1}^n (2i_k + 1)!! q_{2i_k+1} \int_{\mathcal{M}_{g,n}} \Theta_{g,n} \psi_1^{i_1} \cdots \psi_n^{i_n}\right). \quad (1.30)$$

Here $\Theta_{g,n}$ are certain cohomology classes [27]. Our next result is given by the following theorem.

Theorem 1.5 *Define a sequence of numbers $g_n^{\Theta(C)}$ by*

$$g_0^{\Theta(C)} = 1, \quad g_n^{\Theta(C)} = (n-1)! \sum_{k=1}^n \frac{1}{k!(k-1)!} \prod_{i=1}^k \left(C + \frac{i(i-1)}{2}\right), \quad n \geq 1. \quad (1.31)$$

Define b_1, b_2, b_3, \dots via $\sum_{k \geq 1} b_k z^{-k} = \log \sum_{n \geq 0} g_n^{\Theta(C)} z^{-n}$, and denote by $Q = e^{\sum_{k \geq 1} k b_k q_k}$ the multiplication operator. The following identity holds true:

$$\tau_2^{\Theta(C)} = \frac{1}{\tau_1^{\Theta(C)}} Q \circ T_0 \circ Q^{-1} (\tau_1^{\Theta(C)}). \quad (1.32)$$

Moreover, $\tau_E^{\Theta(C)} := \tau_1^{\Theta(C)} \tau_2^{\Theta(C)}$ satisfies the following linear equation:

$$\sum_{n \geq 1} n \tilde{q}_n \frac{\partial \tau_E^{\Theta(C)}}{\partial q_n} + C \tau_E^{\Theta(C)} = 0, \quad \tilde{q}_n = q_n - \delta_{n,1}. \quad (1.33)$$

Recall that the explicit expression for the affine coordinates $A_{m,n}^{\Theta(C)}$ of the point of the Sato Grassmannian corresponding to $\tau_1^{\Theta(C)}$ was obtained in [16]:

$$A_{m,n}^{\Theta(C)} = (-1)^m \sum_{\substack{r \geq m+1, s \geq 0 \\ r+s=m+n+1}} \frac{r-s}{r+s} a_r a_s, \quad a_k := \frac{(-1)^k}{k!} \prod_{i=1}^k \left(C + \binom{i}{2}\right). \quad (1.34)$$

Define $\pi_\lambda^{\Theta(C)}$ via (1.16) with $A_{m,n}$ replaced by $A_{m,n}^{\Theta(C)}$. Combining with (1.22) we arrive at

Corollary 1.6 *The tau-function $\tau_E^{\Theta(C)}$ has the form*

$$\tau_E^{\Theta(C)} = \sum_{\lambda \in \mathbb{Y}} s_\lambda \sum_{\substack{0 \leq k \leq |\lambda| \\ \mu \in T_k^{-1}(\lambda)}} g_k^{\Theta(C)} \pi_\mu^{\Theta(C)}. \quad (1.35)$$

For the reader's convenience we list the first few coefficients of $\tau_E^{\Theta(C)}$ in Appendix A.

Propositions 1.2 and 1.3 also apply to the topological solution to the Burgers-KdV hierarchy, denoted by $(w^{\text{WK}}, \rho^{\text{WK}})$, which is specified by the initial condition

$$w^{\text{WK}}|_{q_2=q_3=\dots=0} = x, \quad \rho^{\text{WK}}|_{q_2=q_3=\dots=0} = 0. \quad (1.36)$$

This solution governs the open and closed intersection numbers [28, 10, 11]. Let τ_1^{WK} be the tau-function of the solution w^{WK} to the KdV hierarchy, normalized by the string equation:

$$\sum_{n \geq 0} (2n+1) (q_{2n+1} - \delta_{n,1}) \frac{\partial \tau_1^{\text{WK}}}{\partial q_{2n+1}} + \frac{x^2}{2} \tau_1^{\text{WK}} = 0. \quad (1.37)$$

The explicit expression for the affine coordinates $A_{m,n}^{\text{WK}}$ for the point of the Sato Grassmannian corresponding to τ_1^{WK} was first obtained in [32] and later re-proved in [4]:

$$\begin{aligned} A_{3m-1,3n}^{\text{WK}} &= A_{3m-3,3n+2}^{\text{WK}} \\ &= \frac{(-1)^n (6m+1)!!}{36^{m+n} (2m+2n)!} \prod_{j=0}^{n-1} (m+j) \prod_{j=1}^n (2m+2j-1) \left(B(n,m) + \frac{2^n (6n+1)!!}{(6m+1)(2n)!} \right), \end{aligned} \quad (1.38)$$

$$\begin{aligned} A_{3m-2,3n+1}^{\text{WK}} &= \frac{(-1)^{n+1} (6m+1)!!}{36^{m+n} (2m+2n)!} \prod_{j=0}^{n-1} (m+j) \prod_{j=1}^n (2m+2j-1) \left(B(n,m) + \frac{2^n (6n+1)!!}{(6m-1)(2n)!} \right), \end{aligned} \quad (1.39)$$

where $n \geq 0$, $m \geq 1$ and $B(n,m)$ are defined as

$$B(n,m) := \frac{1}{6} \sum_{j=1}^n 108^j \frac{2^{n-j} (6n-6j+1)!!}{(2n-2j)!} \frac{\Gamma(m+n+1)}{\Gamma(m+n+2-j)}. \quad (1.40)$$

Recall from [11] that the values g_n for the topological solution are given by

$$g_n^{\text{WK}} = \begin{cases} \sum_{i=0}^m \frac{3^i (6m-6i)!}{288^{m-i} (2m-2i)! (3m-3i)!} \prod_{j=1}^i (m + \frac{1}{2} - j), & n = 3m, \\ 0, & \text{otherwise.} \end{cases} \quad (1.41)$$

Define π_λ^{WK} via (1.16) with $A_{m,n}$ replaced by $A_{m,n}^{\text{WK}}$. Using (1.22) we then arrive at

Corollary 1.7 *The power series τ_E^{WK} has the following expression:*

$$\tau_E^{\text{WK}} = \sum_{\lambda \in \mathbb{Y}} s_\lambda \sum_{\substack{0 \leq k \leq |\lambda| \\ \mu \in T_k^{-1}(\lambda)}} g_k^{\text{WK}} \pi_\mu^{\text{WK}}. \quad (1.42)$$

Organization of the paper In Section 2, we will recall about some basics about the KdV hierarchy. In Section 3, we prove Propositions 1.2 and 1.3. In Section 4, we prove Theorem 1.5. Besides, we give a new proof of a result of Alexandrov in Appendix B.

Acknowledgement We are grateful to Si-Qi Liu and Youjin Zhang for very helpful suggestions and encouragements. D.Y. is grateful to Boris Dubrovin and Don Zagier for their advice. The work receives supports from a national research grant.

2 A brief review of the KdV hierarchy

In this section, we recall some basics about the KdV hierarchy (1.1). It is well known that this hierarchy can be written into the Lax form

$$\frac{\partial L}{\partial q_n} = \left[\left(L^{\frac{n}{2}} \right)_+, L \right], \quad n \geq 1, \quad (2.1)$$

where $L := \partial_x^2 + 2w(x)$. Here, for the meaning of $L^{n/2}$ see e.g. [13]. For any solution $w \in \mathbb{C}[[\mathbf{q}]]$ to the KdV hierarchy¹, there exists [13, 16] a pseudo-differential operator $P = \sum_{i \geq 0} \phi_i \partial_x^{-i} \in$

¹Usually the even flows are not included. Since these flows vanish and are therefore trivial, we have added them in (1.1). In other words, we understand in the way that (1.1) provide solutions for the KP hierarchy.

$\mathbb{C}[[\mathbf{q}]][[\partial_x^{-1}]]$, $\phi_0 := 1$ satisfying the followings:

$$L = P \circ \partial_x^2 \circ P^{-1}, \quad (2.2)$$

$$\frac{\partial P}{\partial q_n} = -(L^{\frac{n}{2}})_- \circ P, \quad \forall n \geq 1. \quad (2.3)$$

The formal adjoint operator of P is defined by

$$P^* := \sum_{i \geq 0} (-\partial_x)^{-i} \circ \phi_i =: \sum_{n \geq 1} \phi_i^* \partial_x^{-i}. \quad (2.4)$$

For each solution w , define the wave function ψ and its dual by

$$\psi := \psi(\mathbf{q}; z) = \sum_{i \geq 0} \phi_i z^{-i} e^{\sum_{n \geq 1} q_n z^n}, \quad (2.5)$$

$$\psi^* := \psi^*(\mathbf{q}; z) = \sum_{i \geq 0} \phi_i^* z^{-i} e^{-\sum_{n \geq 1} q_n z^n}. \quad (2.6)$$

Then ψ and ψ^* satisfy the following linear equations:

$$\mathcal{L}\psi = z^2\psi, \quad \mathcal{L}\psi^* = z^2\psi^*, \quad (2.7)$$

$$\frac{\partial \psi}{\partial q_n} = \left(\mathcal{L}^{\frac{n}{2}}\right)_+ \psi, \quad \frac{\partial \psi^*}{\partial q_n} = \left((\mathcal{L}^*)^{\frac{n}{2}}\right)_+ \psi^*, \quad \forall n \geq 1. \quad (2.8)$$

The following lemma was proven in e.g. [12].

Lemma 2.1 ([12]) *The following bilinear identity holds true:*

$$\text{res}_{z=\infty} \psi(\mathbf{q}; z) \psi^*(\mathbf{q}'; z) dz = 0, \quad \forall \mathbf{q} = (q_1, q_2, \dots), \mathbf{q}' = (q'_1, q'_2, \dots). \quad (2.9)$$

According to the Sato theory [29, 12, 13], there exists a (unique up to a multiplicative constant) formal series $\tau(\mathbf{q})$ satisfying

$$\psi(\mathbf{q}; z) = \frac{\tau(\mathbf{q} - [z^{-1}])}{\tau(\mathbf{q})} e^{\sum_{n \geq 1} q_n z^n}, \quad \psi^*(\mathbf{q}; z) = \frac{\tau(\mathbf{q} + [z^{-1}])}{\tau(\mathbf{q})} e^{-\sum_{n \geq 1} q_n z^n}. \quad (2.10)$$

where $[z^{-1}] := (\frac{1}{z}, \frac{1}{2z^2}, \frac{1}{3z^3}, \dots)$. The series $\tau(\mathbf{q})$ is called the Sato type tau-function of the solution w for the KdV hierarchy, which will play an important role in the later sections. Observe that the above ϕ_i and ϕ_i^* , $i \geq 0$ do not depend on q_2, q_4, q_6, \dots . Therefore, if we view q_2, q_4, q_6, \dots as constants, then as it was proved in [16] the ψ and ψ^* defined by (2.5)–(2.6) satisfy the pair condition of [16]. This statement is applied in several places of this paper.

3 Proofs of Propositions 1.2 and 1.3

The goal of this section is to prove Propositions 1.2 and 1.3.

The following lemma, that will play a fundamental rule, generalizes a result of Buryak [11].

Lemma 3.1 *Let $(w, \rho) \in \mathbb{C}[[\mathbf{q}]]^2$ be an arbitrary solution to the Burgers-KdV hierarchy, and let $(\tau_1, \tau_2) \in \mathbb{C}[[\mathbf{q}]]^2$ be its tau-vector with a fixed choice (1.11). There exists a unique series*

$$g(z) = \sum_{n \geq 0} g_n z^{-n} \in \mathbb{C}[[z^{-1}]], \quad g_n \in \mathbb{C}, \quad g_0 \neq 0, \quad (3.1)$$

such that the component τ_2 has the expression

$$\tau_2 = -\operatorname{res}_{z=\infty} g(z) \frac{\tau_1(\mathbf{q} - [z^{-1}])}{\tau_1(\mathbf{q})} e^{\sum_{n \geq 1} q_n z^n} \frac{dz}{z}. \quad (3.2)$$

Proof According to [5], the Dubrovin–Zhang type tau-function τ_1 of the solution w for the KdV hierarchy defined by (1.9) is also a Sato type tau-function of w , i.e., the ψ defined by

$$\psi(\mathbf{q}; z) := \frac{\tau_1(\mathbf{q} - [z^{-1}])}{\tau_1(\mathbf{q})} e^{\sum_{n \geq 1} q_n z^n}$$

is a wave function (2.8) of the KP hierarchy. Let us define a sequence of number g_n , $n \geq 0$

$$g_0 = \tau_2(\mathbf{0}), \quad (3.3)$$

$$g_n = (\partial_x^n \tau_2)(\mathbf{0}) + \sum_{k=0}^n \sum_{i=0}^{\min\{k, n-1\}} \operatorname{res}_{z=\infty} \frac{g_i dz}{z^{k-i+1}} \partial_x^{n-k} \frac{\tau_1(x - \frac{1}{z}, -\frac{1}{2z^2}, \dots)}{\tau_1(x, 0, \dots)} \Big|_{x=0}, \quad n \geq 1. \quad (3.4)$$

Define $g(z) = \sum_{n \geq 0} g_n z^{-n}$, and define $\tilde{\tau}_2 \in \mathbb{C}[[\mathbf{q}]]$ by $\tilde{\tau}_2 = -\operatorname{res}_{z=\infty} g(z) \psi(\mathbf{q}; z) \frac{dz}{z}$. From the definition of g_n we have

$$\tilde{\tau}_2(x, 0, \dots) = \tau_2(x, 0, \dots). \quad (3.5)$$

Using (2.8) we know that $\forall n \geq 1$,

$$\frac{\partial \tilde{\tau}_2}{\partial q_n} = \left(L^{\frac{n}{2}} \right)_+ (\tilde{\tau}_2), \quad (3.6)$$

where $L = \partial_x^2 + w$. According (1.10) and [10], the second component τ_2 satisfies

$$\frac{\partial \tau_2}{\partial q_n} = \left(L^{\frac{n}{2}} \right)_+ (\tau_2), \quad \forall n \geq 1. \quad (3.7)$$

Combining the formulae (3.5)–(3.7), we obtain $\tau_2 = \tilde{\tau}_2$. The uniqueness follows from the requirement of (3.2) at $q_2 = q_3 = \dots$ identically in q_1 . The lemma is proved. \square

Lemma 3.2 *For each $n \geq 1$ and $\lambda \in \mathbb{Y}$, we have*

$$T_n(s_\lambda) = \det M_{n, \lambda_1, \dots, \lambda_\ell(\lambda)}. \quad (3.8)$$

where M_{k_1, \dots, k_m} denotes the matrix $(h_{k_i - i + j})_{1 \leq i, j \leq m}$ for $m \geq 1$, $k_1, \dots, k_m \in \mathbb{Z}$.

The proof of this lemma is elementary and we omit its details.

Lemma 3.3 *The transformation T_0 defined by (1.18) satisfies*

$$T_0(f) = -\operatorname{res}_{z=\infty} f(\mathbf{q} - [z^{-1}]) e^{\sum_{n \geq 1} q_n z^n} \frac{dz}{z}, \quad \forall f \in \mathbb{C}[[\mathbf{q}]]. \quad (3.9)$$

Proof Due to the linearity and the fact that Schur polynomials form a basis of $\mathbb{C}[[\mathbf{q}]]$, it suffices to show (3.9) for $f = s_\lambda$, $\lambda \in \mathbb{Y}$. By Lemma 3.2 we have

$$T_0(s_\lambda) = \det M_{0,\lambda_1,\dots,\lambda_{\ell(\lambda)}}. \quad (3.10)$$

Hence it suffices to prove that the right-hand side of (3.9) is equal to $\det M_{0,\lambda_1,\dots,\lambda_{\ell(\lambda)}}$. Indeed, we obtain from (1.14) that

$$\sum_{k \geq 0} h_k(\mathbf{q} - [z^{-1}]) y^k = \exp\left(\sum_{k \geq 1} \left(q_k - \frac{1}{kz^k}\right) y^k\right) = \exp\left(\sum_{k \geq 1} q_k y^k\right) \left(1 - \frac{y}{z}\right). \quad (3.11)$$

It follows that

$$h_k(\mathbf{q} - [z^{-1}]) = h_k(\mathbf{q}) - \frac{1}{z} h_{k-1}(\mathbf{q}), \quad \forall k \geq 1. \quad (3.12)$$

Therefore

$$s_\lambda(\mathbf{q} - [z^{-1}]) = \det \left(h_{\lambda_i - i + j} - \frac{1}{z} h_{\lambda_i - i + j - 1} \right)_{1 \leq i, j \leq \ell(\lambda)} =: \sum_{k=0}^{\ell(\lambda)} M^{[k]} z^{-k}, \quad (3.13)$$

where $M^{[k]} := M^{[k]}(\mathbf{q})$ is just the $(1, k+1)$ -th cofactor of the matrix $M_{0,\lambda_1,\dots,\lambda_{\ell(\lambda)}}$. Now the right-hand side of (3.9) can be simplified to

$$- \operatorname{res}_{z=\infty} \left(\sum_{k=0}^{\ell(\lambda)} M^{[k]} z^{-k} \right) \left(\sum_{k \geq 0} h_k z^k \right) \frac{dz}{z} = \sum_{k=0}^{\ell(\lambda)} h_k M^{[k]} = \det M_{0,\lambda_1,\dots,\lambda_{\ell(\lambda)}}. \quad (3.14)$$

The lemma is proved. \square

We notice that similarly with (3.9), we have the following formula:

$$\begin{aligned} - \operatorname{res}_{z=\infty} s_{(m_1, \dots, m_r | n_1, \dots, n_r)}(\mathbf{q} + [z^{-1}]) e^{-\sum_{n \geq 1} q_n z^n} \frac{dz}{z} \\ = (-1)^r s_{(m_1+1, \dots, m_r+1 | n_1-1, \dots, n_r-1)}(\mathbf{q}), \end{aligned} \quad (3.15)$$

if $n_r \geq 1$; and the left-hand side vanishes, if $n_r = 0$.

Now we are ready to prove Proposition 1.2 given in the introduction.

Proof of Proposition 1.2 By using Lemma 3.1, we can find a series $g(z) \in \mathbb{C}[[z^{-1}]]$ such that

$$\tau_1 \tau_2 = - \operatorname{res}_{z=\infty} g(z) \tau_1(\mathbf{q} - [z^{-1}]) e^{\sum_{n \geq 1} q_n z^n} \frac{dz}{z}. \quad (3.16)$$

Consider the expansion

$$\log g(z) =: b_0 + \sum_{k \geq 1} b_k z^{-k}. \quad (3.17)$$

and denote $Q := e^{\sum_{k \geq 1} k b_k q_k}$. By using the identity (3.16) and Lemma 3.3, we have

$$Q^{-1}(\tau_1) \cdot \tau_2 = - \operatorname{res}_{z=\infty} (Q^{-1} \tau_1)(\mathbf{q} - [z^{-1}]) e^{\sum_{n \geq 1} q_n z^n} \frac{dz}{z} = T_0 \circ Q^{-1}(\tau_1).$$

The proposition is proved. \square

Lemma 3.4 Let $Q := e^{\sum_{k \geq 1} k b_k q_k}$ denote the multiplication operator. We have

$$Q \circ T_0 \circ Q^{-1}(s_\lambda) = \sum_{n \geq 0} h_n(\mathbf{b}) T_n(s_\lambda). \quad (3.18)$$

Proof By using Lemma 3.3 and Lemma 3.2 we have

$$\begin{aligned} Q \circ T_0 \circ Q^{-1}(s_\lambda) &= -\operatorname{res}_{z=\infty} e^{\sum_{k \geq 1} b_k z^{-k}} s_\lambda(\mathbf{q} - [z^{-1}]) e^{\sum_{n \geq 1} q_n z^n} \frac{dz}{z} \\ &= \sum_{n, m \geq 0} h_n(\mathbf{b}) h_m(\mathbf{q}) z^{m-n} \sum_{k=0}^{\ell(\lambda)} M^{[k]} z^{-k} \\ &= \sum_{n \geq 0} h_n(\mathbf{b}) \det M_{n, \lambda_1, \dots, \lambda_{\ell(\lambda)}}. \end{aligned}$$

The lemma is proved. \square

We are now ready to prove Proposition 1.3.

Proof of Proposition 1.3 By using Proposition 1.2 and Lemma 3.4, it is easy to see that

$$\tau_E = \sum_{\lambda \in \mathbb{Y}} \pi_\lambda \cdot Q \circ T \circ Q^{-1}(s_\lambda) = \sum_{\lambda \in \mathbb{Y}} \pi_\lambda \sum_{n \geq 0} h_n(\mathbf{b}) T_n(s_\lambda), \quad (3.19)$$

which is nothing but an equivalent form of (1.22). Next we are to show (1.23). It is known [25] that the following identity holds true:

$$e^{\sum_{k \geq 1} k q_k q'_k} = \sum_{\lambda \in \mathbb{Y}} s_\lambda(\mathbf{q}) s_\lambda(\mathbf{q}'). \quad (3.20)$$

Then we have

$$\begin{aligned} Q \circ T_0 \circ Q^{-1}(s_\lambda) &= \sum_{\nu \in \mathbb{Y}} s_\nu(\mathbf{b}) s_\nu(\mathbf{q}) T_0 \left(\sum_{\mu \in \mathbb{Y}} c_{\alpha\mu}^\beta s_\mu(-\mathbf{b}) s_\beta(\mathbf{q}) \right) \\ &= \sum_{\substack{\lambda, \mu \in \mathbb{Y} \\ \beta \in \mathbb{Y} \setminus \operatorname{Ker}(T_0)}} c_{\alpha\mu}^\beta c_{\nu, T_0(\beta)}^\lambda s_\mu(-\mathbf{b}) s_\nu(\mathbf{b}) s_\lambda(\mathbf{q}). \end{aligned}$$

Formula (1.23) is then obtained in a similar way with (3.19). The proposition is proved. \square

For the reader's convenience we provide in below a few Littlewood-Richardson coefficients $c_{\alpha\mu}^\beta$ that appear in the above proof:

$$\begin{aligned} s(3)s(3) &= s(6) + s(5,1) + s(4,2) + s(3,3), \\ s(3)s(2,1) &= s(5,1) + s(4,2) + s(4,1,1) + s(3,2,1), \\ s(3)s(1,1,1) &= s(4,1,1) + s(3,1,1,1), \\ s(2,1)s(2,1) &= s(4,2) + s(4,1,1) + s(3,3) + 2s(3,2,1) + s(3,1,1,1) + s(2,2,2) + s(2,2,1,1), \\ s(2,1)s(1,1,1) &= s(3,2,1) + s(3,1,1,1) + s(2,2,1,1) + s(2,1,1,1,1), \\ s(1,1,1)s(1,1,1) &= s(2,2,2) + s(2,2,1,1) + s(2,1,1,1,1) + s(1,1,1,1,1,1). \end{aligned}$$

For the case of the topological solution to the Burgers–KdV hierarchy, it was conjectured by Witten [31] and proved by Kontsevich the following identity [23]:

$$\tau_1^{\text{WK}} = \exp \left(\sum_{n \geq 1} \sum_{i_1, \dots, i_n \geq 0} \frac{1}{n!!} \prod_{k=1}^n (2i_k + 1)!! q_{2i_k+1} \int_{\mathcal{M}_{g,n}} \psi_1^{i_1} \cdots \psi_n^{i_n} \right), \quad (3.21)$$

where τ_1^{WK} is defined by (1.36)–(1.37). The second component τ_2^{WK} in the tau-vector of the topological solution plays the role of the partition function for the open intersection numbers [28, 10, 11]. From Corollary 1.7 and a straightforward computation, we obtain the first few terms of τ_E^{WK} as follows:

$$\begin{aligned} \tau_E^{\text{WK}} = & 1 + \frac{41}{24} s^{(3)} - \frac{5}{24} s^{(2,1)} - \frac{7}{24} s^{(1,1,1)} + \frac{9241}{1152} s^{(6)} - \frac{385}{1152} s^{(5,1)} - \frac{455}{1152} s^{(4,1,1)} + \frac{205}{576} s^{(3,3)} \\ & + \frac{287}{576} s^{(3,2,1)} + \frac{25}{1152} s^{(3,1,1,1)} - \frac{35}{576} s^{(2,2,2)} - \frac{385}{1152} s^{(2,1,1,1,1)} - \frac{455}{1152} s^{(1,1,1,1,1,1)} + \cdots, \end{aligned} \quad (3.22)$$

which agree with the computations in [22]. We hope that Corollary 1.7 could be helpful in understanding open and closed intersection numbers.

4 The extended generalized BGW tau-function

In this section, following [11] and [7] we prove Theorem 1.5. We also prove that the extended generalized BGW tau-function with some particularly chosen value of C is a polynomial.

Let $(w^{\Theta(C)}, \rho^{\Theta(C)})$ and $(\tau_1^{\Theta(C)}, \tau_2^{\Theta(C)})$ be defined in Introduction (cf. (1.27)–(1.29)). According to [5], the series $\tau_1^{\Theta(C)}$ is also the Sato type tau-function of the solution $(w^{\Theta(C)}, \rho^{\Theta(C)})$ for the KdV hierarchy. Therefore, as mentioned in Section 2, the $\psi_{\Theta(C)} := \psi_{\Theta(C)}(\mathbf{q}; z)$ and $\psi_{\Theta(C)}^* := \psi_{\Theta(C)}^*(\mathbf{q}; z)$ defined by

$$\psi_{\Theta(C)} = \frac{\tau_1^{\Theta(C)}(\mathbf{q} - [z^{-1}])}{\tau_1^{\Theta(C)}(\mathbf{q})} e^{\sum_{n \geq 1} q_n z^n}, \quad \psi_{\Theta(C)}^* = \frac{\tau_1^{\Theta(C)}(\mathbf{q} + [z^{-1}])}{\tau_1^{\Theta(C)}(\mathbf{q})} e^{-\sum_{n \geq 1} q_n z^n}, \quad (4.1)$$

form a pair of wave and dual wave functions of $w^{\Theta(C)}$ for the KdV hierarchy. To proceed we introduce the notations:

$$K_{\Theta(C)} := \tau_1^{\Theta(C)} \psi_{\Theta(C)}, \quad f_{\Theta(C)}(x, z) := \psi_{\Theta(C)} \Big|_{q_2=q_3=\dots=0}, \quad (4.2)$$

$$K_{\Theta(C)}^* := \tau_1^{\Theta(C)} \psi_{\Theta(C)}^*, \quad f_{\Theta(C)}^*(x, z) := \psi_{\Theta(C)}^* \Big|_{q_2=q_3=\dots=0}, \quad (4.3)$$

$$L_0^{\text{Ext}} := \sum_{n \geq 1} n \tilde{q}_n \frac{\partial}{\partial q_n} + C, \quad \tilde{q}_n := q_n - \delta_{n,1}. \quad (4.4)$$

Lemma 4.1 *The following formulae hold true:*

$$L_0^{\text{Ext}} K_{\Theta(C)} = (z \partial_z - z) K_{\Theta(C)}, \quad (4.5)$$

$$L_0^{\text{Ext}} K_{\Theta(C)}^* = (z \partial_z + z) K_{\Theta(C)}^*, \quad (4.6)$$

Proof Performing a shift $\mathbf{q} \mapsto (\mathbf{q} - [z^{-1}])$ to the identity (1.28), we obtain

$$\sum_{k \geq 0} \left((2k+1)\tilde{q}_{2k+1} - \frac{1}{z^{2k+1}} \right) \frac{\partial \tau_1^{\Theta(C)}}{\partial q_{2k+1}} (\mathbf{q} - [z^{-1}]) + C \tau_1^{\Theta(C)} (\mathbf{q} - [z^{-1}]) = 0. \quad (4.7)$$

Then the left-hand side of (4.5) is equal to

$$\begin{aligned} & e^{\sum_{n \geq 1} q_n z^n} \left(\sum_{k \geq 0} (2k+1)\tilde{q}_{2k+1} \frac{\partial \tau_1^{\Theta(C)} (\mathbf{q} - [z^{-1}])}{\partial q_{2k+1}} + \tau_1^{\Theta(C)} (\mathbf{q} - [z^{-1}]) \left(\sum_{n \geq 1} n \tilde{q}_n z^n + C \right) \right) \\ &= e^{\sum_{n \geq 1} q_n z^n} \sum_{k \geq 0} z^{-2k-1} \frac{\partial \tau_1^{\Theta(C)} (\mathbf{q} - [z^{-1}])}{\partial q_{2k+1}} + K_{\Theta(C)} \sum_{n \geq 1} n \tilde{q}_n z^n \\ &= (z \partial_z - z) K_{\Theta(C)}. \end{aligned} \quad (4.8)$$

The proof for (4.6) is similar. The lemma is proved. \square

The next lemma was obtained in [16], and we will provide a new proof of it using the method given in [11].

Lemma 4.2 ([16]) *The following formulae are true:*

$$f_{\Theta(C)} = e^{-xz} \sum_{n \geq 0} \frac{\theta_n}{z^n (1-x)^n}, \quad f_{\Theta(C)}^* = e^{xz} \sum_{n \geq 0} \frac{(-1)^n \theta_n}{z^n (1-x)^n}, \quad (4.9)$$

where $\theta_n := (-1)^n \prod_{i=0}^{n-1} (C + \frac{i+1}{2}) / n!$, $n \geq 0$.

Proof Setting $q_2 = q_3 = \dots = 0$ in (4.5) we have

$$(x-1) \partial_x f_{\Theta(C)} = (z \partial_z - z) f_{\Theta(C)}. \quad (4.10)$$

Then there exists a formal series $\theta(z) \in \mathbb{C}[[z^{-1}]]$ such that $f_{\Theta(C)} = e^{-xz} \theta(z(1-x))$. Setting $q_2 = q_3 = \dots = 0$ in formula (2.7) we have

$$\partial_x^2 f_{\Theta(C)} + (2C - z^2) f_{\Theta(C)} = 0. \quad (4.11)$$

It follows that $\theta(z)$ satisfies the ODE:

$$(z^2 \partial_z^2 - 2z^2 \partial_z + 2C) \theta(z) = 0. \quad (4.12)$$

The rest of the proof can be found in [16]. The proof of the second equality is similar. \square

We are ready to prove Theorem 1.5 using the method of [11] improved in [7].

Proof of Theorem 1.5 Due to Lemma 3.1, there exists a series $g(z)$ of the form (3.1) satisfying

$$\tau_2^{\Theta(C)} = -\operatorname{res}_{z=\infty} g(z) \frac{\tau_1^{\Theta(C)} (\mathbf{q} - [z^{-1}])}{\tau_1^{\Theta(C)} (\mathbf{q})} e^{\sum_{n \geq 1} q_n z^n} \frac{dz}{z}. \quad (4.13)$$

Using (1.29) we find $g_0 = 1$. Using Lemma 4.1 we have

$$L_0^{\text{Ext}}(\tau_E^{\Theta(C)}) = \tau_1^{\Theta(C)} \operatorname{res}_{z=\infty} (g'(z) + g(z)) \psi_{\Theta(C)} dz. \quad (4.14)$$

It follows from (1.28), (1.12) and (1.27) that

$$L_0^{\text{Ext}}(\tau_E^{\Theta(C)})|_{q_2=q_3=\dots=0} \equiv 0. \quad (4.15)$$

Thus taking $q_2 = q_3 = \dots = 0$ on the both sides of (4.14) we find

$$\text{res}_{z=\infty} (g'(z) + g(z)) f_{\Theta(C)}(x, z) dz \equiv 0. \quad (4.16)$$

Since $\psi_{\Theta(C)}$ satisfies equations (2.8), it follows from (4.18) that

$$\text{res}_{z=\infty} (g'(z) + g(z)) \psi_{\Theta(C)}(\mathbf{q}; z) dz \equiv 0. \quad (4.17)$$

Following [7] we employ the bilinear identity (2.9) with $\mathbf{q} = (x, \mathbf{0})$, $\mathbf{q}' = (x', \mathbf{0})$, and we find

$$\text{res}_{z=\infty} f_{\Theta(C)}^*(x', z) f_{\Theta(C)}(x, z) dz \equiv 0. \quad (4.18)$$

Further taking $x' = 0$ and using Lemma 4.2 we find that

$$\text{res}_{z=\infty} \theta(-z) f_{\Theta(C)}(x, z) dz \equiv 0. \quad (4.19)$$

Comparing (4.16) and (4.19), and using the uniqueness nature we find that $g(z)$ must satisfy

$$g'(z) + g(z) = \theta(-z). \quad (4.20)$$

Obviously, solution to (4.20) of the form (3.1) is unique. By solving (4.20) we finally obtain that $g(z) = 1 + \sum_{n \geq 1} g_n^{\Theta(C)} z^{-n}$ with $g_n^{\Theta(C)}$ being given by (1.31). The identities (1.32) and (1.33) are proved. \square

It was shown in [3] that if $C = -m(m+1)/2$ for some non-negative integer m , the generalized BGW tau-function $\tau_1^{\Theta(C)}$ is a polynomial. We find in the next proposition that when m is even, the tau-function $\tau_E^{\Theta(C)}$ is also a polynomial.

Proposition 4.3 *For $C = -m(m+1)/2$, $m \in \mathbb{Z}_{\geq 0}$, if m is even, then $\tau_E^{\Theta(C)}$ is a polynomial.*

Proof Define $g(z) = \sum_{k \geq 0} g_k^{\Theta(C)} z^{-k}$. Using Theorem 1.5 we find

$$g(z) = \sum_{k=0}^{m-1} \frac{g_k^{\Theta(C)}}{z^k} + a_m \sum_{k \geq m} \frac{(k-1)!}{z^k}, \quad (4.21)$$

where

$$a_m = \sum_{k=1}^m \frac{\prod_{i=0}^{k-1} \left(-\frac{m(m+1)}{2} + \frac{i(i+1)}{2} \right)}{k!(k-1)!}. \quad (4.22)$$

One can show that the numbers a_m satisfy

$$a_m = \frac{1}{m!} f^{(m)}(0), \quad f(z) := -z(1+z^2)^{-\frac{3}{2}}. \quad (4.23)$$

Since m is even, we find a_m is zero. Then from (4.21) it follows that $g(z)$ is a polynomial in $1/z$. Combining this with the polynomiality property of the component $\tau_1^{\Theta(C)}$, one can see

$$g(z) \tau_1^{\Theta(C)}(\mathbf{q} - [z^{-1}]) \in \mathbb{C}[\mathbf{q}][z^{-1}]. \quad (4.24)$$

Recall the formula (1.32) and rewrite it into

$$\tau_E^{\Theta(C)} = -\text{res}_{z=\infty} g(z) \tau_1^{\Theta(C)}(\mathbf{q} - [z^{-1}]) e^{\sum_{n \geq 1} q_n z^n} \frac{dz}{z}. \quad (4.25)$$

We conclude from (4.24) and (4.25) that $\tau_E^{\Theta(C)}$ is a polynomial. \square

A The first few coefficients of $\tau_E^{\Theta(C)}$

λ	Coefficient of s_λ
(1)	C
(2)	$C(C+3)/2$
(1,1)	$C(C-1)/2$
(3)	$C(C+3)(C+7)/6$
(2,1)	$C(C+3)(2C-1)/6$
(1,1,1)	$C(C+1)(C-3)/6$
(4)	$C(C+3)(C+9)(C+10)/24$
(3,1)	$C(C+3)(C^2+7C-2)/8$
(2,2)	$C^2(C+1)(C+3)/12$
(2,1,1)	$C(C+1)(C+3)(C-2)/8$
(1,1,1,1)	$C(C+1)(C+3)(C-6)/24$
(5)	$C(C+3)(C+10)(C^2+27C+186)/120$
(4,1)	$C(C+3)(C+10)(2C^2-3C-30)/60$
(3,2)	$C^2(C+1)(C+3)(C+8)/24$
(3,1,1)	$C(C+1)(C+3)(C^2+6C-10)/20$
(2,2,1)	$C^3(C+1)(C+3)/24$
(2,1,1,1)	$C(C+1)(C+3)(2C^2-3C-30)/60$
(1,1,1,1,1)	$C(C+1)(C+3)(C+6)(C-10)/120$
(6)	$C(C+3)(C+10)(C+21)(C^2+31C+270)/720$
(5,1)	$C(C+3)(C+10)(C^3+28C^2+201C-18)/144$
(4,2)	$C^2(C+1)(C+3)(C+10)(C+11)/80$
(4,1,1)	$C(C+1)(C+3)(C+10)(C^2+9C-9)/72$
(3,3)	$C^2(C+1)(C+3)^2C(C+10)/144$
(3,2,1)	$C^2(C+1)(C+3)(4C^2+34C+15)/180$
(3,1,1,1)	$C(C+1)(C+3)(C^3+7C^2-15C-90)/72$
(2,2,2)	$C^2(C+1)(C+3)(C^2+C+6)/144$
(2,2,1,1)	$C^2(C+1)(C+3)(C^2+C-10)/144$
(2,1,1,1,1)	$C(C+1)(C+3)(C+6)(C^2-5C-30)/144$
(1,1,1,1,1,1)	$C(C+1)(C+3)(C+6)(C+10)(C-15)/720$

B A novel proof of a result of Alexandrov

In this section, we give a new proof of a result of Alexandrov [2].

Let $\mathbb{C}((z^{-1}))$ be the linear space of formal series with finitely many terms of positive powers, and consider the Sato Grassmannian GM as defined originally in [29]. A point $W \in GM$ is

a subspace of $\mathbb{C}((z^{-1}))$ which can be written as a linear span of a set of basis vectors. A particular useful choice of basis of W is of the following form:

$$W = \text{Span}_{\mathbb{C}} \left\{ z^l + \sum_{k \geq 0} A_{k,l} z^{-k-1} \right\}_{l \geq 0}. \quad (\text{B.1})$$

Here the collection $(A_{k,l})_{k,l \geq 0}$ is called the *affine coordinate* of W . Note that such a choice is not always possible: existence of basis of form (B.1) characterizes the *big cell* (See [30] for the details). Given a point $W \in GM$, the affine coordinate, if exist, must be unique.

Define a family of numbers as follows:

$$\pi_{(m_1, \dots, m_r | n_1, \dots, n_r)}(W) := (-1)^{\sum_{i=1}^r n_i} \det (A_{m_i, n_j})_{1 \leq i, j \leq r}. \quad (\text{B.2})$$

Then according to [18, 4], the formal series defined by

$$\tau_W := \sum_{\lambda \in \mathbb{Y}} \pi_{\lambda}(W) s_{\lambda} \quad (\text{B.3})$$

is a Sato type tau-function of the KP hierarchy, i.e., it satisfies the following bilinear identity:

$$\text{res}_{z=\infty} \tau_W(\mathbf{q} - [z^{-1}]) \tau_W(\mathbf{q}' + [z^{-1}]) e^{\sum_{n \geq 1} (q_n - q'_n) z^n} dz = 0. \quad (\text{B.4})$$

Theorem B.1 ([2]) *If τ is a Sato type tau-function of the KP hierarchy, so is*

$$\tau' := -\text{res}_{z=\infty} f(z) \tau(\mathbf{q} - [z^{-1}]) e^{\sum_{n \geq 1} q_n z^n} \frac{dz}{z}, \quad (\text{B.5})$$

where $f(z) = 1 + \sum_{n \geq 1} f_n z^{-n} \in \mathbb{C}[[z^{-1}]]$.

Proof By using Lemma 3.3, we can rewrite (B.5) as

$$\tau' = Q \circ T_0 \circ Q^{-1}(\tau), \quad (\text{B.6})$$

where $Q = e^{-\sum_{k \geq 1} b_k z^{-k}}$ denotes the multiplication operator with b_1, b_2, \dots being numbers such that $\log f(z) = \sum_{k \geq 1} b_k z^{-k}$.

Let τ be the Sato type tau-function given by (B.3). We first show $T_0(\tau)$ is a Sato type tau-function. Define a point in GM as

$$W' := \text{Span}_{\mathbb{C}} \left\{ z^l - \sum_{k \geq 0} A_{k+1, l-1} z^{-k-1} \right\}_{l \geq 1} \quad (\text{B.7})$$

It is obvious that the affine coordinates $(A'_{k,l})_{k,l \geq 0}$ of W' are given by

$$(A'_{k,l})_{k,l \geq 0} = \begin{pmatrix} \vdots & \vdots & \vdots & \vdots \\ 0 & -A_{20} & -A_{21} & \cdots \\ 0 & -A_{10} & -A_{11} & \cdots \end{pmatrix}. \quad (\text{B.8})$$

Recall the expression (1.18) of T_0 . Then by using (B.3) and observing that

$$\det(A'_{m_i, n_j})_{1 \leq i, j \leq r} = \begin{cases} (-1)^r \det(A_{m_i+1, n_j-1})_{1 \leq i, j \leq r}, & n_r \geq 1, \\ 0, & n_r = 0, \end{cases} \quad (\text{B.9})$$

one can obtain that

$$T_0(\tau) = \tau_{\mathcal{W}'}$$

Hence $T_0(\tau)$ is Sato type tau-function of the KP hierarchy. In other words, the transformation T_0 has the property that it maps an arbitrary Sato type tau-function of the KP hierarchy to another one. Obviously, the operator Q also has this property. Combining with (B.6), we therefore conclude that the formal series τ' is also a Sato type tau-function. The theorem is proved. \square

Let us give a direct consequence of Proposition 1.3 and Corollary 1.4. For arbitrary numbers $A_{m,n}$, $m, n \geq 0$ and g_k , $k \geq 0$ with $g_0 = 1$, define $c_{m,n}$ by (1.25) and π_λ by (1.16). Then for any partition $\lambda = (m_1, \dots, m_r | n_1, \dots, n_r)$, the following identity holds true:

$$\det(c_{m_i, n_j})_{1 \leq i, j \leq r} = \sum_{\substack{0 \leq k \leq |\lambda| \\ \mu \in T_k^{-1}(\lambda)}} g_k \pi_\mu = \sum_{\substack{\alpha, \mu, \nu \in \mathbb{Y} \\ \beta \in \mathbb{Y} \setminus \text{Ker}(T_0)}} \pi_\alpha c_{\mu\alpha}^\beta c_{\nu, T_0(\beta)}^\lambda s_\mu(-\mathbf{b}) s_\nu(\mathbf{b}), \quad (\text{B.10})$$

where $\mathbf{b} := (b_1, b_2, \dots)$ are defined via $e^{\sum_{k \geq 1} b_k z^{-k}} = \sum_{n \geq 0} g_n z^{-n}$.

References

- [1] Alexandrov, A., Open intersection numbers, matrix models and MKP hierarchy, *J. High Energy Phys.*, **3** (2015), 1–14.
- [2] Alexandrov, A., Open intersection numbers, Kontsevich-Penner model and cut-and-join operators, *J. High Energy Phys.*, **8** (2015), 1–25.
- [3] Alexandrov, A., Cut-and-join description of generalized Brezin-Gross-Witten model, *Adv. Theor. Math. Phys.*, **22** (2018), 1347–1399.
- [4] Balogh, F., Yang, D., Geometric interpretation of Zhou’s explicit formula for the Witten–Kontsevich tau function, *Lett. Math. Phys.*, **107** (2017), 1837–1857.
- [5] Bertola, M., Dubrovin, B., Yang, D., Correlation functions of the KdV hierarchy and applications to intersection numbers over $\overline{\mathcal{M}}_{g,n}$, *Physica D Nonlinear Phenomena*, **327** (2016), 30–57.
- [6] Bertola, M., Ruzza, G., Brezin-Gross-Witten tau-function and isomonodromic deformations, *Communications in Number Theory and Physics*, **13** (2019), 827–883.
- [7] Bertola, M., Yang, D., The partition function of the extended r -reduced Kadomtsev-Petviashvili hierarchy, *Journal of Physics A Mathematical and Theoretical*, **48** (2015).
- [8] Borot, G., Bouchard V., Chidambaram N.K., Creutzig T., Noshchenko D., Higher Airy structures, \mathcal{W} algebras and topological recursion, arXiv:1812.08738.

- [9] Brezin, E., Gross, D.J., The external field problem in the large N limit of QCD, *Physics Letters B*, **97** (1980), 120–124.
- [10] Buryak, A., Equivalence of the open KdV and the open Virasoro equations for the moduli space of Riemann surfaces with boundary, *Lett. Math. Phys.*, **105** (2015), 1427–1448.
- [11] Buryak, A., Open intersection numbers and the wave function of the KdV hierarchy, *Mosc. Math. J.*, **16** (2016), 27–44.
- [12] Date, E., Jimbo, M., Kashiwara, M., Miwa, T., Transformation groups for soliton equations, *Proceedings of RIMS*, 1981.
- [13] Dickey, L.A., *Soliton equations and Hamiltonian systems*, World Scientific, Singapore (2003).
- [14] Do, N., Norbury, P., Topological recursion on the Bessel curve, *Commun. Number Theory Phys.*, **12** (2018), 53–73.
- [15] Dubrovin, B., Yang, D., Remarks on intersection numbers and integrable hierarchies. I. Quasi-triviality, arXiv:1905.08106.
- [16] Dubrovin, B., Yang, D., Zagier, D., On tau-functions for the KdV hierarchy, arXiv:1812.08488.
- [17] Dubrovin, B., Zhang, Y., Normal forms of hierarchies of integrable PDEs, Frobenius manifolds and Gromov–Witten invariants, arXiv: math/0108160.
- [18] Enolski, V.Z., Harnad, J., Schur function expansions of Kadomtsev-Petviashvili τ -functions associated with algebraic curves, *Uspekhi Mat. Nauk*, **66** (2011), 137–178.
- [19] Gross, D.J., Witten, E., Possible third-order phase transition in the large-N lattice gauge theory, *Physical review D: Particles and fields*, **21** (1980), 446–453.
- [20] Itzykson, C., Zuber, J.-B., Combinatorics of the modular group. II. The Kontsevich integrals, *Internat. J. Modern Phys. A*, **7** (1992), 5661–5705.
- [21] Kac, V., Schwarz, A., Geometric interpretation of the partition function of 2D gravity, *Physics Letters B* **257** (1991), 329–334.
- [22] Ke, H.-Z., On a geometric solution to open KdV and Virasoro, *Adv. Math. (China)*, **46** (2017), 91–96.
- [23] Kontsevich, M., Intersection theory on the moduli space of curves and the matrix airy function, *Comm. Math. Phys.*, **147** (1992), 1–23.
- [24] Littlewood, D.E., Richardson, A.R., (1934), *Group Characters and Algebra*, *Philosophical Transactions of the Royal Society of London. Series A*, **233** (1934), 99–141.
- [25] Macdonald, I.G., *Symmetric functions and hall polynomials*, *Oxford Classic Texts in the Physical Sciences*. Clarendon Press, Oxford (1998).

- [26] Mironov, A., Morozov, A., Semenoff, G.W., Unitary matrix integrals in the framework of generalized Kontsevich model, *Internat. J. Modern Phys. A*, **11** (1996), 5031–5080.
- [27] Norbury, P., A new cohomology class on the moduli space of curves. arXiv:1712.03662.
- [28] Pandharipande, R., Solomon, J.P., Tessler, R.J., Intersection theory on moduli of disks, open KdV and Virasoro, arXiv:1409.2191.
- [29] Sato, M., Soliton equations as dynamical systems on infinite dimensional Grassmann manifolds, *Res. Inst. Math. Sci. Kokyuroku*, **439** (1981), 30–46.
- [30] Segal, G., Wilson, G., Loop groups and equations of KdV type, *Surv. Differ. Geom*, **4** (1998), 403–466.
- [31] Witten, E., Two-dimensional gravity and intersection theory on moduli space, *Surveys in differential geometry* (Cambridge, MA, 1990), 243-310, Lehigh Univ., Bethlehem, PA, 1991.
- [32] Zhou, J., Explicit formula for Witten-Kontsevich tau-function, arXiv:1306.5429.
- [33] Zhou, J., Emergent geometry and mirror symmetry of a point, arXiv:1507.01679.

School of Mathematical Sciences, USTC, Hefei 230026, P.R. China
diyang@ustc.edu.cn, zhouch@ustc.edu.cn