

# Gravitational Waves in Higher Order Teleparallel Gravity

Salvatore Capozziello<sup>1,2,3</sup>, Maurizio Capriolo<sup>4</sup>, and Loredana Caso<sup>4</sup>

<sup>1</sup>*Dipartimento di Fisica "E. Pancini", Università di Napoli "Federico II", Compl. Univ. di Monte S. Angelo, Edificio G, Via Cinthia, I-80126, Napoli, Italy,*

<sup>2</sup>*INFN Sezione di Napoli, Compl. Univ. di Monte S. Angelo, Edificio G, Via Cinthia, I-80126, Napoli, Italy,*

<sup>3</sup>*Tomsk State Pedagogical University, ul. Kievskaya, 60, 634061 Tomsk, Russia,*

<sup>4</sup>*Dipartimento di Matematica Università di Salerno, via Giovanni Paolo II, 132, Fisciano, SA I-84084, Italy.*

February 16, 2022

## Abstract

The teleparallel equivalent of higher order Lagrangians like  $L_{\square R} = -R + a_0 R^2 + a_1 R \square R$  can be obtained by means of the boundary term  $B = 2\nabla_\mu T^\mu$ . In this perspective, we derive the field equations in presence of matter for higher-order teleparallel gravity considering, in particular, sixth-order theories where the  $\square$  operator is linearly included. In the weak field approximation, gravitational wave solutions for these theories are derived. Three states of polarization are found: the two standard  $+$  and  $\times$  polarizations, namely 2-helicity massless transverse tensor polarizations, and a 0-helicity massive, with partly transverse and partly longitudinal scalar polarization. Moreover, these gravitational waves exhibit four oscillation modes related to four degrees of freedom: the two classical  $+$  and  $\times$  tensor modes of frequency  $\omega_1$ , related to the standard Einstein waves with  $k_1^2 = 0$ ; two mixed longitudinal-transverse scalar modes for each frequencies  $\omega_2$  and  $\omega_3$ , related to two different 4-wave vectors,  $k_2^2 = M_2^2$  and  $k_3^2 = M_3^2$ . The four degrees of freedom are the amplitudes of each individual mode, i.e.  $\hat{\epsilon}^{(+)}(\omega_1)$ ,  $\hat{\epsilon}^{(\times)}(\omega_1)$ ,  $\hat{B}_2(\mathbf{k})$ , and  $\hat{B}_3(\mathbf{k})$ . To describe a general teleparallel gravity model of order  $(2p + 2)$ , we used the teleparallel Lagrangian  $L_{\square^k T}^{neq} = e(T + a_0 T^2 + \sum_{k=1}^p a_k T \square^k T)$  which demonstrates to be not equivalent to  $L_{\square^k R} = \sqrt{-g}(R + \sum_{k=0}^p a_k R \square^k R)$ . By varying its action, the related field equations in the presence of matter are derived. Hence we obtain the gravitational waves for these teleparallel gravity models, i.e.  $L_{\square^k T}^{neq}$ , that are exactly the Einstein gravitational waves as in  $f(T)$  teleparallel gravity. In conclusion, the boundary term  $B$  generates extra polarizations and additional modes beyond the standard ones. In particular  $B^2$  and  $B \square B$  terms generate both the additional scalar polarization and the extra scalar modes.

PACS numbers: 04.50.Kd, 04.30.-w, 98.80.-k

Keywords: Teleparallel gravity; modified gravity; gravitational waves.

# 1 Introduction

In affine geometry, spacetime can be endowed with a connection whose curvature and torsion do not vanish. Connections are rules that tell us how to derive geometrical objects on a manifold and then how to parallel transport them. Furthermore, it is possible to add a metric to the spacetime whose connection is generally non-metric compatible (metric-affine geometry). However, if we want lengths and angles between vectors to be preserved under parallel transport, we have to impose that the connection is metric-compatible, that is  $\nabla g = 0$ . After turning off both curvature and non-metricity in the connection, gravity appears as a manifestation of the only torsion as in the Teleparallel Equivalent of General Relativity. In this framework, the autoparallel curves (straight lines), i.e. the affine geodesics, related to the connection, and the extremal curves (shortest lines), i.e. the metric geodesics or simply the geodesics, related to metric, are not the same. Autoparallel and extremal curves are the same for light rays or massless wave traveling at the speed of light, while the particle trajectories follow the geodesics as in General Relativity [1, 2]. Finally, in the Weitzenböck spacetime where both curvature and non-metricity are zero, we have that the local Lorentz invariance is violated and the spin connection can vanish. In a nutshell, teleparallelism is a dynamical theory of the vierbein  $\{e_a\}$ , based on the notion of distance or absolute parallelism  $\nabla e_a = 0$ , with respect to the curvature-free torsion-based connection used in the Weitzenböck spacetime. So gravity is completely described by the torsion two-form  $T^a = de^a$  where the one-forms  $\{e^a\}$  are a local basis on the cotangent space [3].

Historically, teleparallelism was proposed by Einstein himself, in 1928, who tried to unify General Relativity with Electromagnetism. The approach was based on replacing the Levi-Civita connection with a non-symmetric connection, the Weitzenböck connection. The attempt was a failure because the components of the Electromagnetic field, identified with the additional six component of the tetrad, could be eliminated by imposing the local Lorentz invariance. However, the idea of modifying the spacetime geometry by using the Weitzenböck connection allowed to formulate a theory of gravity, based on teleparallelism, equivalent to General Relativity, i.e. the Teleparallel Equivalent of General Relativity (TEGR) [4, 5, 6]. Teleparallelism and General Relativity describe the same physics because they give the same field equations coming from the Hilbert-Einstein Lagrangian, linear in the Ricci curvature scalar  $R$  [7, 8], and the Teleparallel Lagrangian, linear in the scalar torsion  $T$ . The two Lagrangian differ by a four-divergence as we will discuss below.

The above theories can be extended in view of Quantum Gravity, considering more general effective Lagrangians. In fact, the issue to renormalize and regularize General Relativity can be improved by adding higher-order curvature invariants and related derivatives, minimally or non-minimally coupled with scalar and vector fields in the Hilbert-Einstein action. Improving the actions with higher-order terms can make the theory more regular and renormalizable at least at lower loop level [9]. Specifically, such contributions appear when the scale lengths approach the Planck scale as one-loop or higher-loop quantum corrections in the effective action. A straightforward extension of the Hilbert-Einstein action can be realized by assuming analytical functions of Ricci curvature scalar  $R$ , the so called  $f(R)$  gravity where General Relativity is recovered for  $f(R) \rightarrow R$  [10, 11, 12, 13].

In the same way, it is possible to extend TEGR considering analytical functions  $f(T)$  which can be non-linear with respect to  $T$  and, as standard TEGR, exhibit the violation of local Lorentz invariance [14, 15]. Extended teleparallel gravity  $f(T)$  differs from extended gravity  $f(R)$  essentially because the former leads to second-order field equations while the latter leads to fourth-order field equations in metric formalism. Then  $f(R)$  gravity is not equivalent to  $f(T)$  gravity but, if we introduce the boundary terms  $B = 2\nabla_\mu (T^\mu)$ , the equivalence can be restored being  $f(T, B) \rightarrow f(R)$

gravity [16].

Extended teleparallel theories  $f(T)$  can be adopted, for example, to explain the accelerated expansion of the Universe at the present time [17]. If we want to study higher-order teleparallel theories, we cannot limit to  $f(T)$ , because, as said above, it always produces second-order dynamical equations. To compare extended teleparallel theories with the analogous in curvature invariants, we have to introduce the boundary term  $B$  with its derivatives and terms like  $\square T$ ,  $\square^k T$  in the teleparallel Lagrangian [18, 19]. In order to develop our considerations, we can therefore start from Lagrangians in terms of  $R$  and  $\square R$  and find their teleparallel equivalent, using the boundary term  $B$ . Otherwise we can generally start directly from a theory of gravity of order  $n$  in teleparallel geometry, without the boundary term  $B$  but only with  $\square^k T$ . We can study, in particular, the field equations associated with the Lagrangian  $L_{\square^k T} = e(T + a_0 T^2 + \sum_{k=1}^p a_k T \square^k T)$ .

All the mentioned theories of gravity, both in Riemannian and teleparallel frameworks, can be studied in the weak field limit to be linearized. This allows us to derive gravitational waves (GWs) for each theory and to analyze their properties. These theories differ in their predictions for propagation speed and polarizations of gravitational radiation. GW polarization measurements, implemented by interferometers, are a powerful tool to select viable theories of gravity. On the other hand, GW detection can contribute to define the effective number of degrees of freedom (d.of.) of a given theory of gravity [16, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30].

In this paper, we want to investigate higher-order teleparallel gravity in view of a classification of GW solutions. Furthermore, we want to compare such theories with the analogous in curvature invariants to establish, eventually, if they are equivalent or not.

The layout of the paper is the following. In Sec.2, we obtain the extended teleparallel Lagrangian  $L_{\square T}^e$  equivalent to the curvature Lagrangian  $L_{\square R} = -R + a_0 R^2 + a_1 R \square R$  by means of a boundary term  $B$ . As discussed in [31, 32, 33], this theory is important because it is related to one-loop corrections of Quantum Gravity and it gives a double-inflationary evolution in early cosmology because it can be reduced to a double-scalar field theory by conformal transformations.

In Sec.3, we obtain the field equations of sixth-order teleparallel gravity. Hence, in Sec.4, linearized equations are derived in the weak field limit. In Sec.5, adopting the method of distributions, we get GW solutions in sixth-order teleparallel gravity. Properties of GWs, as polarization and helicity, are investigated by geodesic deviation and Newmann-Penrose formalism in Sec.6. Sec.7 is devoted to the study of teleparallel gravity at order  $(2p + 2)$ . We point out the non-equivalence with extended curvature  $R$ -based theory considering a flat geometry with torsion in a zero spin connection. Varying the action  $L_{\square^k T}$  with respect to the tetrad  $e^a_\rho$  and imposing that it vanishes with its derivatives on the boundary of the  $\Omega$  domain, we obtain the teleparallel higher-order field equations in presence of matter. After linearization, the related GW solutions are obtained and studied. Conclusions are drawn in Sec.8. In Appendices, the differential identities and the variations are reported.

## 2 Teleparallel Equivalent of Sixth Order Curvature Gravity

Let us consider sixth-order curvature gravity in view of obtaining the teleparallel equivalent of Lagrangian  $L_{\square R} = -R + a_0 R^2 + a_1 R \square R$ , expressed in terms of the Levi Civita connection  $\overset{\circ}{\Gamma}$ , replacing it with the curvature-free and torsion-based Weitzenböck connection  $\tilde{\Gamma}$ . In teleparallelism, we adopt the sixteen components of the tetrad basis  $\{e_a\}$  and its dual basis  $\{e^a\}$  to describe spacetime manifold. These dynamical variables form a local orthonormal basis for the tangent

space at each point  $\{x_\mu\}$  of the spacetime. The components of the *vierbein* satisfy the relations [34, 35, 36, 37, 38, 39]

$$e^a{}_\mu e_a{}^\nu = \delta_\mu^\nu, \quad \text{and} \quad e^a{}_\mu e_b{}^\mu = \delta_a^b, \quad (2.1)$$

$$\eta_{ab} = g_{\mu\nu} e_a{}^\mu e_b{}^\nu, \quad \text{and} \quad g_{\mu\nu} = \eta_{ab} e_a{}^\mu e_b{}^\nu, \quad (2.2)$$

where the signature of the metric  $g_{\mu\nu}$  is  $(+, -, -, -)$ . The Weitzenböck connection is defined to parallel-transport the tetrads on the manifold, that is

$$\tilde{\Gamma}^\rho{}_{\mu\nu} = e_a{}^\rho \partial_\nu e^a{}_\mu, \quad (2.3)$$

and the related torsion tensor is

$$T^\nu{}_{\rho\mu} = \tilde{\Gamma}^\nu{}_{\mu\rho} - \tilde{\Gamma}^\nu{}_{\rho\mu}. \quad (2.4)$$

Subtracting the Levi-Civita connection from the Weitzenböck connection, we obtain the contortion tensor

$$K^\nu{}_{\rho\mu} = \tilde{\Gamma}^\nu{}_{\rho\mu} - \overset{\circ}{\Gamma}^\nu{}_{\rho\mu} = \frac{1}{2} (T_\rho{}^\nu{}_\mu + T_\mu{}^\nu{}_\rho - T^\nu{}_{\rho\mu}). \quad (2.5)$$

The superpotential torsion tensor  $S^{\rho\mu\nu}$  is

$$S^{\rho\mu\nu} = \frac{1}{2} (K^{\mu\nu\rho} - g^{\rho\nu} T^\sigma{}_\mu{}^\rho + g^{\rho\mu} T^\sigma{}_\nu{}^\rho). \quad (2.6)$$

Contracting the torsion tensor with the superpotential, we get the scalar torsion  $T$

$$T = T_{\rho\mu\nu} S^{\rho\mu\nu}. \quad (2.7)$$

On the other hand, the torsion vector is

$$T^\sigma = T^{\nu\sigma}{}_\nu. \quad (2.8)$$

The Weitzenböck connection curvature vanishes  $R[\tilde{\Gamma}] = 0$  but we can express the Levi Civita connection curvature  $R[\overset{\circ}{\Gamma}]$  in terms of the scalar torsion  $T$  and the vector torsion  $T^\sigma$ , that is

$$-R = T + \frac{2}{e} \partial_\sigma (e T^{\nu\sigma}{}_\nu), \quad (2.9)$$

and, in more compact form,<sup>1</sup>

$$-R = T + B, \quad (2.10)$$

where  $e = \det(e^a{}_\rho)$  and the boundary term is

$$B = \frac{2}{e} \partial_\sigma (e T^{\nu\sigma}{}_\nu). \quad (2.11)$$

---

<sup>1</sup>For the signature of the boundary term, see the discussion in [65].

With the above positions, we can express the sixth-order curvature Lagrangian in the teleparallel equivalent form by the transformation

$$\begin{aligned}
- R + a_0 R^2 + a_1 R \square R &= \underbrace{T + a_0 T^2 + a_1 T \square T}_{\text{First term}} \\
&+ \underbrace{a_0 \left( \frac{2}{e} \partial_\sigma (e T^{\nu\sigma}{}_\nu) \right) \left( 2T + \frac{2}{e} \partial_\sigma (e T^{\nu\sigma}{}_\nu) \right)}_{\text{Second Term}} - \underbrace{a_1 \nabla^\mu \left( \frac{2}{e} \partial_\sigma (e T^{\nu\sigma}{}_\nu) \right) \nabla_\mu \left( 2T + \frac{2}{e} \partial_\sigma (e T^{\nu\sigma}{}_\nu) \right)}_{\text{Third term}} \\
&+ \underbrace{\frac{1}{e} \partial_\sigma \left\{ a_1 h \nabla^\sigma \left[ \frac{2}{e} \partial_\eta (e T^{\nu\eta}{}_\nu) T + \frac{1}{2} \left( \frac{2}{e} \partial_\eta (e T^{\nu\eta}{}_\nu) \right)^2 \right] + 2e T^{\nu\sigma}{}_\nu \right\}}_{\text{Fourth term}}, \quad (2.12)
\end{aligned}$$

or, more explicitly, via the boundary term  $B$

$$\begin{aligned}
- R + a_0 R^2 + a_1 R \square R &= T + a_0 T^2 + a_1 T \square T + a_0 B (2T + B) - a_1 \nabla^\mu B \nabla_\mu (2T + B) \\
&+ \frac{1}{e} \partial_\sigma \left\{ a_1 e \nabla^\sigma \left[ B T + \frac{1}{2} B^2 \right] \right\}, \quad (2.13)
\end{aligned}$$

where  $\nabla_\mu$  is the covariant derivative with respect to the Levi-Civita connection  $\overset{\circ}{\Gamma}$ . The fourth term of Eq. (2.12) is a four-divergence which vanishes if the tetrad with its derivatives vanish on the boundary of the domain; the first term of Eq. (2.12) is the component we need and the second term exists because it depends on the  $\partial^2 e_a^\rho$  of the tetrad and guarantees fourth-order equations in the absence of  $\square$  operator. Otherwise, considering only the scalar  $T$  depending on the first derivatives of the tetrad  $\partial e_a^\rho$ , we cannot go above the second order (because  $f(T)$  theories are described by second order equations and not by fourth-order equations as  $f(R)$  which, instead, depend on the second derivatives of the metric). The third term of Eq. (2.12) depends on third derivatives of tetrad  $\partial^3 e_a^\rho$  insuring sixth-order field equations. The possibility of finding an equivalent teleparallel Lagrangian is guaranteed by the boundary term  $B$  which is not incorporated into a four-divergence. The TEGR, whose Lagrangian depends linearly on the scalar torsion  $T$ , is equivalent to the Hilbert-Einstein Lagrangian because the term  $B$  is incorporated into the four-divergence that vanishes on the boundary while the teleparallel extended theories, to be equivalent to extended theories like  $f(R)$  or  $f(R, \square R)$ , must include boundary terms. Another form of sixth-order teleparallel Lagrangian, up to four-divergence, is

$$\mathcal{L}_{\square T}^{eq} = \underbrace{T + a_0 B^2 + a_1 B \square B}_{\text{terms that generate the first order}} + a_0 T^2 + a_1 T \square T + 2a_0 B T - 2a_1 \nabla^\mu B \nabla_\mu T - \frac{1}{2} a_1 \square B^2, \quad (2.14)$$

where we pointed out parts containing first-order terms generating GWs as we are going to discuss in the next sections.

### 3 Field Equations of Sixth-Order Teleparallel Equivalent Gravity

Let us take into account now the sixth-order teleparallel equivalent Lagrangian up to a four-divergence (2.13), which can vanish imposing appropriate conditions for the fields on the boundary,

that is

$$L_{\square T}^{eq} = e \mathcal{L}_{\square T}^{eq} = \frac{e}{2\kappa^2} [T + a_0 T^2 + a_1 T \square T + a_0 B (2T + B) - a_1 \nabla^\mu B \nabla_\mu (2T + B)] , \quad (3.1)$$

where  $\kappa^2 = 8\pi G/c^4$  is the gravitational coupling. The following action with the material term  $L_m = e \mathcal{L}_m$

$$I = \int_{\Omega} d^4x [\mathcal{L}_{\square T} + \mathcal{L}_m] e , \quad (3.2)$$

can be examined. Varying with respect to the tetrad  $e^a_\rho$  with the variation of the tetrad field vanishing on the boundary, we obtain the following sixth-order field equations:

$$\boxed{\begin{aligned} & \frac{4}{e} \partial_\sigma [e (1 + 2a_0 T + 2a_1 \square T) S_a^{\rho\sigma}] - 4 (1 + 2a_0 T + 2a_1 \square T) T^\mu_{\nu a} S_\mu^{\nu\rho} + (T + a_0 T^2 + a_1 T \square T) e_a^\rho \\ & - a_1 \left\{ \left[ e_a^\nu T (\nabla^\rho \nabla_\nu + \nabla_\nu \nabla^\rho) T - \left( e_a^\eta e_b^\rho \partial^\nu e^b_\nu + T_a^\eta{}^\rho - e_a^\eta T^\rho - g^{\eta\rho} T_a \right) T \partial_\eta T \right] \right. \\ & + \frac{1}{e} \partial_\sigma [e (e_a^\rho g^{\eta\sigma} - e_a^\sigma g^{\eta\rho} - e_a^\eta g^{\rho\sigma}) T \partial_\eta T] \Big\} - [a_0 B^2 - a_1 B \square (T + B) + a_1 \partial^\mu B \partial_\mu (2T + B)] e_a^\rho \\ & + \frac{4}{e} \partial_\lambda \{ e [a_0 (T + B) + a_1 \square (T + B)] \} \left( T^{\rho\lambda}_a + e_a^\lambda T^\rho + g^{\lambda\rho} T_a - T^\lambda e_a^\rho \right) \\ & + \frac{4}{e} \partial_\sigma \{ e \partial_\lambda [a_0 (T + B) + a_1 \square (T + B)] (e_a^\rho g^{\lambda\sigma} - e_a^\sigma g^{\lambda\rho}) \} \\ & - 8 [a_0 B + a_1 \square B] (T^\mu_{\nu a} S_\mu^{\nu\rho}) + \frac{8}{e} \partial_\sigma [e (a_0 B + a_1 \square B) S_a^{\rho\sigma}] = 2\kappa^2 \mathcal{T}^{(m)}_a{}^\rho \end{aligned}} , \quad (3.3)$$

with  $T^\lambda = T^{\nu\lambda}_\nu$  or alternatively:

$$\begin{aligned} & \frac{4}{e} \partial_\sigma [e (1 + 2a_0 T + 2a_1 \square T) S_a^{\rho\sigma}] - 4 (1 + 2a_0 T + 2a_1 \square T) T^\mu_{\nu a} S_\mu^{\nu\rho} + (T + a_0 T^2 + a_1 T \square T) e_a^\rho \\ & - a_1 \left\{ \left[ e_a^\nu T (\nabla^\rho \nabla_\nu + \nabla_\nu \nabla^\rho) T - \left( e_a^\eta e_b^\rho \partial^\nu e^b_\nu + T_a^\eta{}^\rho - e_a^\eta T^\rho - g^{\eta\rho} T_a \right) T \partial_\eta T \right] \right. \\ & + \frac{1}{e} \partial_\sigma [e (e_a^\rho g^{\eta\sigma} - e_a^\sigma g^{\eta\rho} - e_a^\eta g^{\rho\sigma}) T \partial_\eta T] \Big\} - [a_0 B^2 - a_1 B \square (T + B) + a_1 \partial^\mu B \partial_\mu (2T + B)] e_a^\rho \\ & + \frac{4}{e} \partial_\lambda (eG) \left( T^{\rho\lambda}_a + e_a^\lambda T^\rho + g^{\lambda\rho} T_a - T^\lambda e_a^\rho \right) + \frac{4}{e} \partial_\sigma [e \partial_\lambda G (e_a^\rho g^{\lambda\sigma} - e_a^\sigma g^{\lambda\rho})] \\ & - 8 [a_0 B + a_1 \square B] (T^\mu_{\nu a} S_\mu^{\nu\rho}) + \frac{8}{e} \partial_\sigma [e (a_0 B + a_1 \square B) S_a^{\rho\sigma}] = 2\kappa^2 \mathcal{T}^{(m)}_a{}^\rho , \quad (3.4) \end{aligned}$$

where  $G = a_0 (T + B) + a_1 \square (T + B)$ . The energy-momentum tensor of matter  $\mathcal{T}^{(m)}_a{}^\rho$  is:

$$\mathcal{T}^{(m)}_a{}^\rho = - \frac{1}{e} \frac{\delta (e \mathcal{L}_m)}{\delta e^a_\rho} . \quad (3.5)$$

For details on differential identities and variations, see Appendices A and B.

## 4 Weak Field Limit of Sixth-Order Teleparallel Equivalent Gravity

Let us perturb the tetrad field around the background metric with the hypothesis that we are sufficiently far from massive bodies. We can limit our considerations to a flat background described by the trivial tetrad  $e^a{}_\mu = \delta^a{}_\mu$  as follows

$$e^a{}_\mu = \delta^a{}_\mu + E^a{}_\mu , \quad (4.1)$$

where  $|E^a{}_\mu| \ll 1$ . We therefore expand the metric tensor  $g_{\mu\nu}$  to first order in  $E^a{}_\mu$  and we get

$$g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu} + \mathcal{O}(h^2) = \eta_{\mu\nu} + \eta_{\mu a} E^a{}_\nu + \eta_{\nu a} E^a{}_\mu + \mathcal{O}(E^2) . \quad (4.2)$$

Perturbations are related each other by the relation

$$h_{\mu\nu} = \eta_{\mu a} E^a{}_\nu + \eta_{\nu a} E^a{}_\mu . \quad (4.3)$$

To first order in  $E^a{}_\mu$ , the Weitzenböck connection becomes

$$\tilde{\Gamma}^{\rho(1)}_{\mu\nu} = \delta_a{}^\rho \partial_\nu E^a{}_\mu . \quad (4.4)$$

The zero-order covariant derivative  $\nabla_\mu$  and the covariant d'Alembert operator  $\square = g^{\mu\nu} \nabla_\mu \nabla_\nu$  become

$$\nabla_\mu^{(0)} = \partial_\mu , \quad (4.5)$$

$$\square^{(0)} = \eta^{\mu\nu} \partial_\nu \partial_\mu = \partial^\mu \partial_\mu . \quad (4.6)$$

The torsion tensor  $T^\mu{}_{\nu\rho}$ , its contraction  $T^{\mu\nu}{}_\mu$  and the contortion tensor  $K^\rho_{\mu\nu}$ , to first order in tetrad perturbation  $E^a{}_\mu$  can be written as

$$T^{\mu(1)}_{\nu\rho} = \delta_a{}^\mu (\partial_\nu E^a{}_\rho - \partial_\rho E^a{}_\nu) , \quad (4.7)$$

$$T^{\rho\sigma(1)}_\rho = \delta_a{}^\mu \eta^{\sigma\nu} (\partial_\nu E^a{}_\mu - \partial_\mu E^a{}_\nu) , \quad (4.8)$$

$$K^{\rho(1)}_{\mu\nu} = \eta_{\mu\lambda} \delta_a{}^\lambda \partial^\rho E^a{}_\nu - \delta_a{}^\rho \partial_\mu E^a{}_\nu . \quad (4.9)$$

The superpotential  $S_\rho{}^{\mu\nu}$ , the scalar torsion  $T$  and the boundary term  $B$ , perturbed at the lowest order, become

$$\begin{aligned} 2S_\rho{}^{\mu\nu(1)} = & \delta_a{}^\nu \partial^\mu E^a{}_\rho - \delta_a{}^\mu \partial^\nu E^a{}_\rho - \delta_\rho{}^\nu (\delta_a{}^\sigma \partial^\mu E^a{}_\sigma - \eta^{\alpha\mu} \delta_a{}^\alpha \partial_\sigma E^\sigma{}_a) \\ & + \delta_\rho{}^\mu (\delta_a{}^\sigma \partial^\nu E^a{}_\sigma - \eta^{\alpha\nu} \delta_a{}^\alpha \partial_\sigma E^\sigma{}_a) , \end{aligned} \quad (4.10)$$

$$T^{(2)} = T^{\mu\nu\rho(1)} S_{\mu\nu\rho}^{(1)} , \quad (4.11)$$

$$B^{(1)} = \left( \frac{2}{e} \partial_\sigma (e T^\sigma) \right)^{(1)} = 2 \delta_a{}^\nu [\square E^a{}_\nu - \partial^\mu \partial_\nu E^a{}_\mu] . \quad (4.12)$$

The lowest order where the scalar torsion  $T$  does not vanish is the second order in the tetrad perturbation  $E^a{}_\mu$ . The Ricci curvature  $R$ , to first order in  $E^a{}_\mu$ , gets the following form

$$R^{(1)} = -B^{(1)} , \quad (4.13)$$

that is, to first order, only the boundary term  $B$  contributes to the Ricci curvature and not the torsion  $T$ . Finally we obtain the useful relation

$$2\partial_\nu S_\rho^{\mu\nu(1)} = \delta_a^\nu \partial_\nu \partial^\mu E_\rho^a - \delta_a^\mu \square E_\rho^a - \delta_a^\sigma \partial_\rho \partial^\mu E_\sigma^a + \eta^{\rho\mu} \delta_a^\sigma \partial_\rho \partial_\sigma E_\rho^a + \delta_\rho^\mu \delta_a^\sigma \square E_\sigma^a - \delta_\rho^\mu \delta_a^\sigma \eta^{\alpha\nu} \partial_\nu \partial_\sigma E_\alpha^a . \quad (4.14)$$

The first-order tetrad perturbation  $E_\mu^a$  is not symmetric because the sixth-order teleparallel equivalent gravity is not invariant under a local Lorentz transformation [40, 41]

$$\eta_{\mu a} E_\nu^a \neq \eta_{\nu a} E_\mu^a , \quad (4.15)$$

and then, we can decompose the tetrad perturbation  $E_{\mu\nu}$  into symmetric and antisymmetric parts

$$E_{\mu\nu} = E_{(\mu\nu)} + E_{[\mu\nu]} . \quad (4.16)$$

However, the antisymmetric term  $E_{[\mu\nu]}$  has no physical meaning because it is not involved into the Lagrangian  $L_{\square T}^{eq}$  and the field equations: in fact, they depend on the symmetric term  $E_{(\mu\nu)}$  by means of  $T$  and  $B$ . Hence, we can set to zero the antisymmetric part  $E_{[\mu\nu]}$

$$E_{[\mu\nu]} = 0 , \quad (4.17)$$

and the metric perturbation becomes

$$h_{\mu\nu} = 2\eta_{\mu a} E_\nu^a . \quad (4.18)$$

We now linearize the field equations of sixth-order teleparallel equivalent gravity, Eq.(3.3), keeping, at most, first order terms in tetrad perturbation  $E_\mu^a$ . We therefore obtain

$$4\partial_\sigma S_\tau^{\rho\sigma(1)} + 4(\eta^{\lambda\sigma} \delta_\tau^\rho - \eta^{\lambda\rho} \delta_\tau^\sigma) \partial_\sigma \partial_\lambda (a_0 B^{(1)} + a_1 \square B^{(1)}) = 2\kappa^2 \mathcal{T}_\tau^{\rho(0)} . \quad (4.19)$$

The harmonic gauge, to the first order in tetrad perturbation, becomes

$$\partial_\mu \bar{E}^{\mu\nu} = \partial_\mu \left( E^{\mu\nu} - \frac{1}{2} \eta^{\mu\nu} E \right) = 0 , \quad (4.20)$$

where we set

$$E_{\mu\nu} = \eta_{\mu a} E_\nu^a , \quad E = \delta_a^\mu E_\mu^a , \quad (4.21)$$

and

$$\bar{E}_{\mu\nu} = E_{\mu\nu} - \frac{1}{2} \eta_{\mu\nu} E . \quad (4.22)$$

The boundary term to first order  $B^{(1)}$  and Eq.(4.14) in the harmonic gauge take the form

$$B^{(1)} = -\square \bar{E} , \quad (4.23)$$

$$2\partial_\nu S_\rho^{\mu\nu(1)} = -\square \bar{E}^\mu_\rho . \quad (4.24)$$

Substituting Eqs.(4.23) and (4.24) into (4.19), we obtain the linearized field equations in presence of matter (4.19) as

$$\boxed{\square \bar{E}^\rho_\tau + 2a_0 (\delta_\tau^\rho \square^2 - \partial_\tau \partial^\rho \square) \bar{E} + 2a_1 (\delta_\tau^\rho \square^3 - \partial_\tau \partial^\rho \square^2) \bar{E} = -\kappa^2 \mathcal{T}_\tau^{\rho(0)}} . \quad (4.25)$$



In a more compact form

$$\boxed{\square \bar{E}^\rho{}_\tau + 2 \sum_{k=0}^1 a_k (\delta^\rho{}_\tau \square^{k+2} - \partial_\tau \partial^\rho \square^{k+1}) \bar{E} = -\kappa^2 \mathcal{T}^\rho{}_{(0)}}, \quad (4.26)$$

where the trace is

$$\square \bar{E} + 6 \sum_{k=0}^1 a_k \square^{k+2} \bar{E} = -\kappa^2 \mathcal{T}^{(0)}. \quad (4.27)$$

Setting  $l = k + 2$  and defining  $c_l$  as

$$c_l := \begin{cases} 1 & \text{if } l = 1 \\ 6a_{l-2} & \text{if } l > 1 \end{cases},$$

we get the trace equation of Eq. (4.26) as

$$\boxed{\sum_{l=1}^3 c_l \square^l \bar{E} = -\kappa^2 \mathcal{T}^{(0)}}. \quad (4.28)$$

We have now all the ingredient to search for GW solutions in sixth-order teleparallel gravity.

## 5 Gravitational Waves in Sixth-Order Teleparallel Equivalent Gravity

The solution of Eq.(4.28) is an object more general than a function.  $\bar{E}(x)$  is a distribution in  $\mathbb{R}^4$  [42] or more precisely a *tempered distribution*.

Let us now give a precise definition of the notion of distribution. Let  $\phi$  be a complex-valued function on  $\mathbb{R}^4$ . The closure of the set of points  $\{x \in \mathbb{R}^4 : \phi(x) \neq 0\}$  is called the support of  $\phi$  and it is denoted by  $\text{supp}(\phi)$ . A function  $\phi : \mathbb{R}^4 \rightarrow \mathbb{C}$  is called a  $C^\infty$  function if all its partial derivatives  $D^k \phi$  exist and are continuous. A  $C^\infty$  function with compact support is called a *test function* and the space of the test functions on  $\mathbb{R}^4$  is denoted by  $\mathcal{D}(\mathbb{R}^4)$  that is, the space of functions with compact support. A distribution on  $\mathbb{R}^4$  is a continuous complex-valued linear functional  $T$  defined on  $\mathcal{D}(\mathbb{R}^4)$  and the space of all distributions  $T$  is denoted by  $\mathcal{D}'(\mathbb{R}^4)$ . The space  $\mathcal{D}(\mathbb{R}^4)$  is however not closed under Fourier transforms thus we replace it with a new space, closed under Fourier transforms, the Schwartz space  $\mathcal{S}(\mathbb{R}^4)$  of rapidly decreasing functions. The space of Schwartz functions is defined as the space of infinitely differentiable functions whose derivatives decay faster than any polynomial at infinity, that is, formally,  $\mathcal{S}(\mathbb{R}^4) = \{\phi \in C^\infty : \|\phi\|_{(N,\alpha)} < \infty \forall N, \alpha\}$  where we have defined the norm  $\|\phi\|_{(N,\alpha)} = \sup_{x \in \mathbb{R}^4} (1 + |x|^N) |\partial^\alpha \phi(x)|$  for any non-negative integer  $N$  and any multi-index  $\alpha$ . The tempered distribution is a continuous linear functional on  $\mathcal{S}(\mathbb{R}^4)$  and the space of all tempered distributions on  $\mathbb{R}^4$  is denoted by  $\mathcal{S}'(\mathbb{R}^4)$ , that is the dual space of the space of Schwartz functions  $\mathcal{S}(\mathbb{R}^4)$ .

The equation of trace (4.28) in vacuum, in  $k$ -space, becomes

$$\left( \sum_{l=1}^3 c_l (-1)^l k^{2l} \right) \hat{A}(k) = 0, \quad (5.1)$$

where  $k^2 = \omega^2 - \mathbf{k} \cdot \mathbf{k} = \omega^2 - q^2$  with the four-wavevector  $k^\mu = (\omega, \mathbf{k})$ .  $\hat{A}(k) \in \mathcal{S}'(\mathbb{R}^4)$  is the Fourier transformation of  $\bar{E}(x) \in \mathcal{S}'(\mathbb{R}^4)$

$$\hat{A}(k) = \int \frac{d^4 k}{(2\pi)^2} \bar{E}(x) e^{-ik^\alpha x_\alpha} . \quad (5.2)$$

We solve Eq.(5.1) as an algebraic equation in the space of distributions with  $\hat{A}(k) \in \mathcal{S}'(\mathbb{R}^4)$

$$(6a_1 k^6 - 6a_0 k^4 + k^2) \hat{A}(k) = 0 , \quad (5.3)$$

then

$$\hat{A}(k) = f(k) [\delta(6a_1 k^6 - 6a_0 k^4 + k^2)] , \quad (5.4)$$

where we used the  $\delta$ -distribution with a suitable complex function  $f(k)$ . We use now the following properties of  $\delta$ -distribution

$$\delta(F(x)) = \sum_{h=1}^n \frac{\delta(x - x_h)}{|F'(x_h)|} , \quad (5.5)$$

such as  $F(x_h) = 0$  for  $h = 1, \dots, n$  and

$$f(x) \delta(x - a) = f(a) \delta(x - a) . \quad (5.6)$$

The solutions of the related algebraic equation

$$6a_1 k^6 - 6a_0 k^4 + k^2 = 0 , \quad (5.7)$$

are

$$k^2 = 0 \quad k^2 = M_2^2 \neq 0 \quad k^2 = M_3^2 \neq 0 , \quad (5.8)$$

where

$$M_{2,3}^2 = \frac{-3a_0 \mp \sqrt{3a_0^2 - a_1}}{6a_1} . \quad (5.9)$$

Keeping  $\mathbf{k}$  fixed and varying the time-component  $k^0$  of four-wavevector  $k^\mu$ , we obtain

$$k^2 = 0 \rightarrow (k^0)^2 = |\mathbf{k}|^2 \rightarrow k^0 = \pm |\mathbf{k}| = \pm \omega_1 , \quad (5.10)$$

$$k^2 = M_2^2 \rightarrow (k^0)^2 - |\mathbf{k}|^2 = M_2^2 \rightarrow k^0 = \pm \sqrt{M_2^2 + |\mathbf{k}|^2} = \pm \omega_2 , \quad (5.11)$$

$$k^2 = M_3^2 \rightarrow (k^0)^2 - |\mathbf{k}|^2 = M_3^2 \rightarrow k^0 = \pm \sqrt{M_3^2 + |\mathbf{k}|^2} = \pm \omega_3 . \quad (5.12)$$

From (5.5), it is

$$\begin{aligned} \delta(6a_1 k^6 - 6a_0 k^4 + k^2) = & \frac{1}{2\omega_1 |18a_1 \omega_1^4 - 12a_0 \omega_1^2 + 1|} [\delta(k^0 - \omega_1) + \delta(k^0 + \omega_1)] \\ & + \frac{1}{2\omega_2 |18a_1 \omega_2^4 - 12a_0 \omega_2^2 + 1|} [\delta(k^0 - \omega_2) + \delta(k^0 + \omega_2)] \\ & + \frac{1}{2\omega_3 |18a_1 \omega_3^4 - 12a_0 \omega_3^2 + 1|} [\delta(k^0 - \omega_3) + \delta(k^0 + \omega_3)] , \end{aligned} \quad (5.13)$$

and we have

$$\hat{A}(k) = f(k^0, \mathbf{k}) \sum_{m=1}^3 \frac{\delta(k^0 - \omega_m) + \delta(k^0 + \omega_m)}{2\omega_m |18a_1\omega_m^4 - 12a_0\omega_m^2 + 1|} , \quad (5.14)$$

and, by (5.6), we get

$$\hat{A}(k) = \sum_{m=1}^3 \frac{\delta(k^0 - \omega_m) f(\omega_m, \mathbf{k}) + \delta(k^0 + \omega_m) f(-\omega_m, \mathbf{k})}{2\omega_m |18a_1\omega_m^4 - 12a_0\omega_m^2 + 1|} . \quad (5.15)$$

Imposing that the function  $\bar{E}(x)$  is real, we obtain

$$f^*(k) = f(-k) , \quad (5.16)$$

that is

$$f^*(\omega_m, \mathbf{k}) = f(-\omega_m, -\mathbf{k}) \rightarrow f^*(\omega_m, -\mathbf{k}) = f(-\omega_m, \mathbf{k}) . \quad (5.17)$$

After setting

$$f(\omega_m, \mathbf{k}) = Q_m(\mathbf{k}) , \quad (5.18)$$

from (5.17) and (5.18), we obtain, from Eq. (5.15),

$$\hat{A}(k) = \sum_{m=1}^3 \frac{\delta(k^0 - \omega_m) Q_m(\mathbf{k}) + \delta(k^0 + \omega_m) Q_m^*(-\mathbf{k})}{2\omega_m |18a_1\omega_m^4 - 12a_0\omega_m^2 + 1|} . \quad (5.19)$$

In a more compact form, it is

$$\hat{A}(k) = \sqrt{2\pi} \sum_{m=1}^3 \left[ \delta(k^0 - \omega_m) \hat{B}_m(\mathbf{k}) + \delta(k^0 + \omega_m) \hat{B}_m^*(-\mathbf{k}) \right] , \quad (5.20)$$

with

$$\hat{B}_m(\mathbf{k}) = \frac{Q_m(\mathbf{k})}{2\sqrt{2\pi}\omega_m |18a_1\omega_m^4 - 12a_0\omega_m^2 + 1|} . \quad (5.21)$$

Let us now perform the inverse Fourier transform of  $\hat{A}(k)$ . It is

$$\bar{E}(x) = \int \frac{d^4k}{(2\pi)^2} \hat{A}(k) e^{ik^\alpha x_\alpha} , \quad (5.22)$$

and, by (5.20), we get

$$\bar{E}(x) = \frac{1}{(2\pi)^{3/2}} \int d^3\mathbf{k} \int dk^0 e^{i(k^0 x^0 - \mathbf{k} \cdot \mathbf{x})} \sum_{m=1}^3 \left[ \delta(k^0 - \omega_m) \hat{B}_m(\mathbf{k}) + \delta(k^0 + \omega_m) \hat{B}_m^*(-\mathbf{k}) \right] . \quad (5.23)$$

Finally, we obtain the solution

$$\bar{E}(x) = \frac{1}{(2\pi)^{3/2}} \sum_{m=1}^3 \int d^3\mathbf{k} \left( \hat{B}_m(\mathbf{k}) e^{ik_m^\alpha x_\alpha} + c.c. \right) , \quad (5.24)$$

with  $k_m^\mu = (\omega_m, \mathbf{k})$ . Let us rename  $M_2$  as  $k_2$  and  $M_3$  as  $k_3$ , then we have, from (5.7),

$$\begin{cases} k_1^2 = 0 & \text{if } m = 1 \\ \sum_{l=0}^1 a_l (-1)^{l+2} k_m^{2(l+1)} = \frac{1}{6} & \text{if } m = 2, 3 \end{cases} . \quad (5.25)$$

Therefore, from Eqs.(4.26) and (5.25), we get, in vacuum,

$$\square \bar{E}_{\rho\tau}(x) = \sum_{m=2}^3 \int \frac{d^3\mathbf{k}}{(2\pi)^{3/2}} \left\{ \left( -\frac{k_m^2}{3} \right) \left[ \eta_{\rho\tau} - \frac{(k_m)_\rho (k_m)_\tau}{k_m^2} \right] \right\} \left( \hat{B}_m(\mathbf{k}) e^{ik_m^\alpha x_\alpha} + c.c. \right) . \quad (5.26)$$

The general solution of (4.26), in vacuum, considering the homogeneous plus a particular solution is

$$\begin{aligned} \bar{E}_{\rho\tau}(x) &= \int \frac{d^3\mathbf{k}}{(2\pi)^{3/2}} \hat{C}_{\rho\tau}(\mathbf{k}) e^{ik_1^\alpha x_\alpha} \\ &+ \sum_{m=2}^3 \int \frac{d^3\mathbf{k}}{(2\pi)^{3/2}} \left\{ \left( -\frac{1}{3} \right) \left[ \eta_{\rho\tau} - \frac{(k_m)_\rho (k_m)_\tau}{k_m^2} \right] \right\} \hat{B}_m(\mathbf{k}) e^{ik_m^\alpha x_\alpha} + c.c. . \end{aligned} \quad (5.27)$$

From Eq.(4.22), we derive the GWs in vacuum for sixth-order teleparallel equivalent gravity, that is

$$\begin{aligned} E_{\rho\tau}(x) &= \int \frac{d^3\mathbf{k}}{(2\pi)^{3/2}} C_{\rho\tau}(\mathbf{k}) e^{ik_1^\alpha x_\alpha} \\ &+ \sum_{m=2}^3 \int \frac{d^3\mathbf{k}}{(2\pi)^{3/2}} \left\{ \frac{1}{3} \left[ \frac{\eta_{\rho\tau}}{2} + \frac{(k_m)_\rho (k_m)_\tau}{k_m^2} \right] \right\} \hat{B}_m(\mathbf{k}) e^{ik_m^\alpha x_\alpha} + c.c. . \end{aligned} \quad (5.28)$$

For  $f(T, B)$  teleparallel gravity and for higher-order metric gravity see [16, 20].

## 6 Polarizations and helicity

Let us consider the wave traveling along the  $+z$  direction in a local proper reference frame with a separation vector  $\vec{\chi} = (x^1, x^2, x^3)$  which connects two nearby geodesics. The geodesic deviation is

$$\ddot{x}^i = -R^i_{0k0} x^k , \quad (6.1)$$

where the Latin index ranges over the set  $\{1, 2, 3\}$  and  $R^i_{0k0}$  are the so-called "electric" components of the Riemann tensor, the only measurable components. Linearized electric components of the Riemann tensor  $R^{(1)}_{i0j0}$ , expressed in terms of the tetrad perturbation  $E_{\mu\nu}$ , restricted to the Hilbert space of square integrable functions denoted by  $L^2(\mathbb{R}^4)$ , are

$$R^{(1)}_{i0j0} = (E_{i0,0j} + E_{0j,i0} - E_{ij,00} - E_{00,ij}) . \quad (6.2)$$

Replacing them into Eq.(6.1), we obtain

$$\begin{cases} \ddot{x}(t) = -(xE_{11,00} + yE_{12,00}) \\ \ddot{y}(t) = -(xE_{12,00} + yE_{11,00}) \\ \ddot{z}(t) = (2E_{03,03} - E_{33,00} - E_{00,33})z \end{cases} . \quad (6.3)$$

From the general solution (5.28), the plane wave propagating in the  $+\hat{z}$  direction with  $k_1^2 = 0$ , keeping  $\mathbf{k}$  fixed and  $k_1^\mu = (\omega_1, 0, 0, k_z)$  can be written as

$$E_{\mu\nu}^{(k_1)}(t, z) = \sqrt{2} \left[ \hat{\epsilon}^{(+)}(\omega_1) \epsilon_{\mu\nu}^{(+)} + \hat{\epsilon}^{(\times)}(\omega_1) \epsilon_{\mu\nu}^{(\times)} \right] e^{i\omega_1(t-z)} + c.c. , \quad (6.4)$$

where

$$\epsilon_{\mu\nu}^{(+)} = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} , \quad (6.5)$$

$$\epsilon_{\mu\nu}^{(\times)} = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} . \quad (6.6)$$

Always from Eq.(5.28), the  $m$ -th plane wave propagating in the  $+\hat{z}$  direction, for  $k_m^2 \neq 0$  with  $m \in \{2, 3\}$ , keeping  $\mathbf{k}$  fixed, and  $k_m^\mu = (\omega_m, 0, 0, k_z)$  can be written as

$$E_{\mu\nu}^{(k_m)}(t, z) = \left[ \left( \frac{1}{2} + \frac{\omega_m^2}{k_m^2} \right) \epsilon_{\mu\nu}^{(TT)} - \frac{\sqrt{2}\omega_m k_z}{k_m^2} \epsilon_{\mu\nu}^{(TS)} - \frac{1}{\sqrt{2}} \epsilon_{\mu\nu}^{(b)} + \left( -\frac{1}{2} + \frac{k_z^2}{k_m^2} \right) \epsilon_{\mu\nu}^{(l)} \right] \frac{\hat{B}_m(k_z)}{3} e^{i(\omega_m t - k_z z)} + c.c. , \quad (6.7)$$

where

$$\epsilon_{\mu\nu}^{(TT)} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} , \quad \epsilon_{\mu\nu}^{(TS)} = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix} , \quad (6.8)$$

$$\epsilon_{\mu\nu}^{(b)} = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} , \quad \epsilon_{\mu\nu}^{(l)} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} . \quad (6.9)$$

In more compact form, the linear perturbation of tetrad  $E_{\mu\nu}$ , traveling in the  $+\hat{z}$  direction and keeping  $\mathbf{k}$  fixed, may be expressed as

$$E_{\mu\nu}(t, z) = \sqrt{2} \left[ \hat{\epsilon}^{(+)}(k_z) \epsilon_{\mu\nu}^{(+)} + \hat{\epsilon}^{(\times)}(k_z) \epsilon_{\mu\nu}^{(\times)} \right] e^{i\omega_1(t-z)} + \hat{\epsilon}_{\mu\nu}^{(s_2)}(k_z) e^{i(\omega_2 t - k_z z)} + \hat{\epsilon}_{\mu\nu}^{(s_3)}(k_z) e^{i(\omega_3 t - k_z z)} + c.c. , \quad (6.10)$$

where  $\hat{\epsilon}_{\mu\nu}^{(s_m)}$  is the polarization tensor associated to the scalar mode with  $m \in \{2, 3\}$

$$\hat{\epsilon}_{\mu\nu}^{(s_m)}(k_z) = \left[ \left( \frac{1}{2} + \frac{\omega_m^2}{k_m^2} \right) \epsilon_{\mu\nu}^{(TT)} - \frac{\sqrt{2}\omega_m k_z}{k_m^2} \epsilon_{\mu\nu}^{(TS)} - \frac{1}{\sqrt{2}} \epsilon_{\mu\nu}^{(b)} + \left( -\frac{1}{2} + \frac{k_z^2}{k_m^2} \right) \epsilon_{\mu\nu}^{(l)} \right] \frac{\hat{B}_m(k_z)}{3} . \quad (6.11)$$

The two d.o.f.  $\hat{\epsilon}^{(+)}$  and  $\hat{\epsilon}^{(\times)}$  produce the two tensor modes associated to the polarization tensors  $\epsilon_{\mu\nu}^{(+)}$  and  $\epsilon_{\mu\nu}^{(\times)}$  while the two d.o.f.  $\hat{B}_2$  and  $\hat{B}_3$  produce two scalar modes associated to the polarization tensors  $\hat{\epsilon}_{\mu\nu}^{(s_2)}$  and  $\hat{\epsilon}_{\mu\nu}^{(s_3)}$ . Indeed the polarization tensor  $\hat{\epsilon}_{\mu\nu}^{(s)}$ , restricted to the spatial components  $\hat{\epsilon}_{i,j}^{(s)}$ , is provided by

$$\hat{\epsilon}_{i,j}^{(s_m)} = -\frac{1}{3\sqrt{2}}\hat{B}_m(k_z)\epsilon_{i,j}^{(b)} + \frac{1}{3}\left(-\frac{1}{2} + \frac{k_z^2}{k_m^2}\right)\hat{B}_m(k_z)\epsilon_{i,j}^{(l)}, \quad (6.12)$$

where  $(i, j)$  range over  $(1, 2, 3)$ . Hence, the scalar mode is a combination of the longitudinal scalar mode and the transverse breathing scalar mode.

More explicitly, for a massless plane wave  $E_{\mu\nu}^{(k_1)}$ , Eq.(6.3) gives

$$\begin{cases} \ddot{x}(t) = \omega_1^2 [\hat{\epsilon}^{(+)}(\omega_1)x + \hat{\epsilon}^{(\times)}(\omega_1)y] e^{i\omega_1(t-z)} + c.c. \\ \ddot{y}(t) = \omega_1^2 [\hat{\epsilon}^{(\times)}(\omega_1)x - \hat{\epsilon}^{(+)}(\omega_1)y] e^{i\omega_1(t-z)} + c.c. \\ \ddot{z}(t) = 0 \end{cases}, \quad (6.13)$$

where we obtain the two standard GW polarizations of General Relativity, the purely transverse  $+$  and  $\times$  polarizations and the two-helicity massless tensor modes.

Instead, for the  $m$ -th massive plane wave,  $E_{\mu\nu}^{(k_m)}$ , where  $m = 2, 3$ , with  $k_m^2 = M_m^2 = \omega_m^2 - k_z^2$  equal to the square of the  $m$ -th mass of the scalar field, Eqs.(6.3) give

$$\begin{cases} \ddot{x}(t) = -\frac{1}{6}\omega_m^2\hat{B}_m(k_z)x e^{i(\omega_m t - k_z z)} + c.c. \\ \ddot{y}(t) = -\frac{1}{6}\omega_m^2\hat{B}_m(k_z)y e^{i(\omega_m t - k_z z)} + c.c. \\ \ddot{z}(t) = -\frac{1}{6}M_m^2\hat{B}_m(k_z)z e^{i(\omega_m t - k_z z)} + c.c. \end{cases}. \quad (6.14)$$

This system of equations can be integrated assuming small  $E_{\mu\nu}(t, z)$  and, hence, we have

$$\begin{cases} x(t) = x(0) + \frac{1}{6}\hat{B}_m(k_z)x(0)e^{i(\omega_m t - k_z z)} + c.c. \\ y(t) = y(0) + \frac{1}{6}\hat{B}_m(k_z)y(0)e^{i(\omega_m t - k_z z)} + c.c. \\ z(t) = z(0) + \frac{1}{6\omega_m^2}M_m^2\hat{B}_m(k_z)z(0)e^{i(\omega_m t - k_z z)} + c.c. \end{cases}, \quad (6.15)$$

where we obtain two further mixed massive scalar modes, zero-helicity, partially longitudinal and partially transverse breathing, that is each one with the same mixed scalar polarization. To visualize GW polarizations, we use the geodesic deviations when a  $m$ -th plane GW of frequency  $\omega_m$  strikes a sphere of freely falling particles of radius  $r = \sqrt{x^2(0) + y^2(0) + z^2(0)}$ . The displacement of a given particle from the center of the ring  $\vec{\chi}$  is given by the solution of the geodesic deviation equation (6.15). The sphere will be distorted into an ellipsoid described by

$$\left(\frac{x}{\rho_{1m}(t)}\right)^2 + \left(\frac{y}{\rho_{1m}(t)}\right)^2 + \left(\frac{z}{\rho_{2m}(t)}\right)^2 = r^2, \quad (6.16)$$

where both  $\rho_{1m}(t) = 1 + \frac{1}{3}\hat{B}_m(k_z)\cos(\omega_m t - k_z z)$  and  $\rho_{2m}(t) = 1 + \frac{M_m^2}{3\omega_m^2}\hat{B}_m(k_z)\cos(\omega_m t - k_z z)$  vary between their maximum and minimum value. Each swinging ellipsoid represents an additional scalar mode with zero-helicity which is partly longitudinal and partly transverse [43]. The d.o.f. of the sixth-order teleparallel equivalent gravity are four: two of these,  $\hat{\epsilon}^{(+)}$  and  $\hat{\epsilon}^{(\times)}$ , generate the  $+$  and  $\times$  tensor modes and the two d.o.f  $\hat{B}_2$  and  $\hat{B}_3$  generate two mixed longitudinal-transverse scalar modes. Finally our theory of gravity shows three polarizations, two tensor and one mixed scalar polarizations, and four modes, two tensor and two mixed scalar modes for each frequency  $\omega_2$  and  $\omega_3$ .

The same results can be obtained adopting the Newman-Penrose (NP) formalism for massless and massive waves propagating along either null or nearly non-null geodesics [44, 45]. To determine the *little group*<sup>2</sup>  $E(2)$  classification of massive GWs<sup>3</sup>, we linearize the vacuum field equations in the limit of nearly massless plane waves (observer far from the source). We introduce a quasinormal local null tetrad basis  $\{\hat{e}_a\} = (k, l, m, \bar{m})$  as

$$k = \frac{1}{\sqrt{2}}(\partial_t + \partial_z) , \quad l = \frac{1}{\sqrt{2}}(\partial_t - \partial_z) , \quad (6.17)$$

$$m = \frac{1}{\sqrt{2}}(\partial_x + i\partial_y) , \quad \bar{m} = \frac{1}{\sqrt{2}}(\partial_x - i\partial_y) , \quad (6.18)$$

which satisfies the relations

$$\begin{aligned} k \cdot l &= -m \cdot \bar{m} = 1 , \\ k \cdot k &= l \cdot l = m \cdot m = \bar{m} \cdot \bar{m} = 0 , \\ k \cdot m &= k \cdot \bar{m} = l \cdot m = l \cdot \bar{m} = 0 . \end{aligned} \quad (6.19)$$

Knowing that

$$\hat{e}_a = e_a^\mu \partial_\mu \quad \theta^a = \theta_\mu^a dx^\mu , \quad (6.20)$$

$$g_{\mu\nu} = \tilde{\eta}_{ab} \theta_\mu^a \theta_\nu^b \quad \tilde{\eta}_{ab} = e_a^\mu e_b^\nu g_{\mu\nu} , \quad (6.21)$$

with metric  $\tilde{\eta}_{ab}$

$$\tilde{\eta}_{ab} = \tilde{\eta}^{ab} = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & -1 & 0 \end{pmatrix} , \quad (6.22)$$

it is

$$g^{\mu\nu} = 2k^{(\mu} l^{\nu)} - 2m^{(\mu} \bar{m}^{\nu)} , \quad (6.23)$$

where  $\{\theta^a\}$  is the dual tetrad of  $\{\hat{e}_a\}$ . Now we express the fifteen NP scalars in terms of the four-dimensional Weyl tensor  $C_{\mu\nu\rho\sigma}$  in a null tetrad basis defined as

$$C_{\mu\nu\rho\sigma} = R_{\mu\nu\rho\sigma} - 2g_{[\mu[\rho} R_{\sigma]\nu]} + \frac{1}{3}g_{\mu[\rho} g_{\sigma]\nu} R . \quad (6.24)$$

<sup>2</sup>This is the subgroup of Lorentz transformations leaving the wave vector  $\mathbf{k}$  invariant.

<sup>3</sup>It is worth noticing that this classification is strictly valid only for massless waves.

The five complex Weyl-NP scalars, classified with spin weight  $s$ , from the Weyl tensor expressed in a null tetrad basis, are

$$\begin{aligned}
s = +2 \quad \Psi_0 &\equiv C_{kmkm} , \\
s = +1 \quad \Psi_1 &\equiv C_{klkm} , \\
s = 0 \quad \Psi_2 &\equiv C_{km\bar{m}l} , \\
s = -1 \quad \Psi_3 &\equiv C_{kl\bar{m}l} , \\
s = -2 \quad \Psi_4 &\equiv C_{\bar{m}l\bar{m}l} ,
\end{aligned} \tag{6.25}$$

while the ten Ricci-NP scalars, classified with spin weight  $s$ , from the Ricci tensor expressed in a null tetrad basis, are

$$\begin{aligned}
s = 2 \quad \Phi_{02} &\equiv -\frac{1}{2}R_{mm} , \\
s = 1 \quad \begin{cases} \Phi_{01} \equiv -\frac{1}{2}R_{km} \\ \Phi_{12} \equiv -\frac{1}{2}R_{lm} \end{cases} , \\
s = 0 \quad \begin{cases} \Phi_{00} \equiv -\frac{1}{2}R_{kk} \\ \Phi_{11} \equiv -\frac{1}{4}(R_{kl} + R_{m\bar{m}}) \\ \Phi_{22} \equiv -R_{ll} \end{cases} , \\
s = -1 \quad \begin{cases} \Phi_{10} \equiv -\frac{1}{2}R_{k\bar{m}} = \Phi_{01}^* \\ \Phi_{21} \equiv -\frac{1}{2}R_{l\bar{m}} = \Phi_{12}^* \end{cases} , \\
s = -2 \quad \Phi_{20} &\equiv -\frac{1}{2}R_{m\bar{m}} = \Phi_{02}^* , \\
\Lambda &= \frac{R}{24} .
\end{aligned}$$

However, for plane waves, they become

$$\Psi_0 = \Psi_1 = 0 , \tag{6.26}$$

$$\Psi_2 = \frac{1}{6}R_{lklk} , \tag{6.27}$$

$$\Psi_3 = -\frac{1}{2}R_{lkl\bar{m}} , \tag{6.28}$$

$$\Psi_4 = -R_{l\bar{m}l\bar{m}} , \tag{6.29}$$

$$\Phi_{00} = \Phi_{01} = \Phi_{10} = \Phi_{02} = \Phi_{20} = 0 , \tag{6.30}$$

$$\Phi_{22} = -R_{lml\bar{m}} , \tag{6.31}$$

$$\Phi_{11} = \frac{3}{2}\Psi_2 , \tag{6.32}$$

$$\Phi_{12} = \Phi_{21} = \bar{\Psi}_3 , \tag{6.33}$$

where we can choose the set  $\{\Psi_2(u), \Psi_3(u), \Psi_4(u), \Phi_{22}(u)\}$  as independent amplitudes to describe GWs propagating in the  $+z$  direction with the retarded time  $u = t - z$ . Taking into account that the spatial components of driving-force matrix  $S(t)$  of an ideal detector are the electric components



of the Riemann tensor  $R_{i0j0}$  that is [46, 47, 48, 49, 50, 51, 52]

$$S_{ij}(t) = R_{i0j0}(\tilde{u}) , \quad (6.34)$$

we express the response matrix  $S(t)$  in terms of the six new basis polarization matrices  $W_A(\mathbf{e}_z)$  belonging to the wave direction  $\mathbf{k} = \mathbf{e}_z$  with  $A$  ranging over  $\{1, 2, 3, 4, 5, 6\}$ , that is [53, 54]

$$S(t) = \sum_A p_A(\mathbf{e}_z, t) W_A(\mathbf{e}_z) , \quad (6.35)$$

where

$$\begin{aligned} W_1(\mathbf{e}_z) &= -6 \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} , & W_2(\mathbf{e}_z) &= -2 \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix} , \\ W_3(\mathbf{e}_z) &= 2 \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} , & W_4(\mathbf{e}_z) &= -\frac{1}{2} \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix} , \\ W_5(\mathbf{e}_z) &= \frac{1}{2} \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} , & W_6(\mathbf{e}_z) &= -\frac{1}{2} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix} . \end{aligned} \quad (6.36)$$

The six polarization amplitudes  $p_A(\mathbf{e}_z, t)$  can be expressed in terms of the six NP scalars, each with their own helicity as [55, 56, 57, 58, 59, 60]

$$p_1^{(l)} \equiv \Psi_2 \quad s = 0 , \quad (6.37)$$

$$p_2^{(x)} \equiv \text{Re } \Psi_3 \quad s = 1 , \quad (6.38)$$

$$p_3^{(y)} \equiv \text{Im } \Psi_3 \quad s = -1 , \quad (6.39)$$

$$p_4^{(+)} \equiv \text{Re } \Psi_4 \quad s = 2 , \quad (6.40)$$

$$p_5^{(\times)} \equiv \text{Im } \Psi_4 \quad s = -2 , \quad (6.41)$$

$$p_6^{(b)} \equiv \Phi_{22} \quad s = 0 . \quad (6.42)$$

$$(6.43)$$

The six polarizations modes are: the longitudinal mode  $p_1^{(l)}$ , the vector- $x$  mode  $p_2^{(x)}$ , the vector- $y$  mode  $p_3^{(y)}$ , the  $+$  mode  $p_4^{(+)}$ , the  $\times$  mode  $p_5^{(\times)}$  and the breathing mode  $p_6^{(b)}$ . Hence  $\Psi_2$  and  $\phi_{22}$  are both scalar modes, respectively purely transverse and purely longitudinal, the complex  $\Psi_3$  are vector modes mixed and the complex  $\Psi_4$  are the tensor modes purely transverse.

We conclude that the GWs of sixth-order teleparallel equivalent gravity are composed by both massless plane waves  $k_1^2 = 0$ , moving along null-geodesics, and massive plane waves  $k_2^2, k_3^2 \neq 0$ , moving along non null-geodesics. In the hypothesis of slightly massive waves, we can use the NP-formalism for weak, plane, nearly null GWs [44]. They have a wave 4-vector  $k_m^\mu$ , normal to surfaces of constant retarded time  $\tilde{u}$ , that is

$$(k_m)_\mu = -\tilde{u}_{,\mu} . \quad (6.44)$$

The GW propagation speed  $|\mathbf{v}_g|$  is less than  $c$  for massive waves and  $c$  for massless waves

$$\mathbf{v}_g = \frac{d\omega(\mathbf{k})}{d\mathbf{k}} = \frac{\mathbf{k}}{\omega(\mathbf{k})} \quad |\mathbf{v}_g| \leq c . \quad (6.45)$$

One can define a parameter  $\epsilon$  as the speed difference between electromagnetic waves and GWs in a local proper reference frame, that is

$$\epsilon = \left( \frac{c}{v_g} \right)^2 - 1 . \quad (6.46)$$

in this local observer frame, we may expand  $k_m^\mu$  in terms of the null-tetrad local basis  $(k, l, m, \bar{m})$ , that is

$$k_m^\mu = k^\mu (1 + \epsilon_k) + \epsilon_l l^\mu + \epsilon_m m^\mu + \bar{\epsilon}_m \bar{m}^\mu , \quad (6.47)$$

where  $\{\epsilon_k, \epsilon_l, \epsilon_m\} \sim \mathcal{O}(\epsilon)$  and  $\epsilon$  vanishes for null wave because  $|\mathbf{v}_g| = c$ . The only NP quantities which are not  $\mathcal{O}(\epsilon_n)$  are the four scalar  $\{\Psi_2, \Psi_3, \Psi_4, \Phi_{22}\}$ , with  $\Psi_3$  and  $\Psi_4$  complex, that is

$$\Psi_2 = -\frac{1}{6} R_{lklk} + \mathcal{O}(\epsilon_l R) , \quad (6.48)$$

$$\Psi_3 = -\frac{1}{2} R_{lkl\bar{m}} + \mathcal{O}(\epsilon_l R) , \quad (6.49)$$

$$\Psi_4 = -R_{l\bar{m}l\bar{m}} + \mathcal{O}(\epsilon_l R) , \quad (6.50)$$

$$\Phi_{22} = -R_{lm\bar{l}\bar{m}} + \mathcal{O}(\epsilon_l R) . \quad (6.51)$$

The linearized harmonic gauge in the perturbation  $E_{\mu\nu}$  for our plane wave propagating in the  $+z$  direction becomes

$$\begin{cases} \partial_0 E_0^0 + \partial_3 E_0^3 = \frac{1}{2} \partial_0 E \\ \partial_0 E_1^0 + \partial_3 E_1^3 = 0 \\ \partial_0 E_2^0 + \partial_3 E_2^3 = 0 \\ \partial_0 E_3^0 + \partial_3 E_3^3 = \frac{1}{2} \partial_3 E \end{cases} , \quad (6.52)$$

where  $E$  is the trace of  $E_{\mu\nu}$ . After simple algebraic and differential calculations, we have

$$\begin{cases} \partial_0 E_{01} = \partial_3 E_{31} \\ \partial_0 E_{02} = \partial_3 E_{32} \\ (\partial_{00} - \partial_{33})(E_{00} + E_{33}) = -(\partial_{00} + \partial_{33})(E_{11} + E_{22}) \\ (\partial_{00} - \partial_{33})E_{30} = -\partial_{03}(E_{11} + E_{22}) \end{cases} , \quad (6.53)$$

which, after derivations, give

$$\begin{cases} E_{01} = -\frac{k_z}{\omega} E_{31} \\ E_{02} = -\frac{k_z}{\omega} E_{32} \\ E_{11} + E_{22} = -\frac{\omega^2 - k_z^2}{\omega^2 + k_z^2} (E_{00} + E_{33}) \\ E_{30} = \frac{\omega k_z}{\omega^2 - k_z^2} (E_{11} + E_{22}) \end{cases} . \quad (6.54)$$

Then the four NP-scalars become

$$\Psi_2 = \frac{1}{6} \ddot{E}_{kk} + \mathcal{O}(\epsilon_l R) = -\frac{1}{6} \left[ \frac{\omega^2 - k_z^2}{\omega^2 + k_z^2} (\omega^2 E_{33} - k_z^2 E_{00}) \right] + \mathcal{O}(\epsilon_l R) , \quad (6.55)$$

$$\Psi_3 = \frac{1}{2} \ddot{E}_{k\bar{m}} + \mathcal{O}(\epsilon_l R) = -\frac{1}{2} (\omega^2 - k_z^2) E_{13} + \frac{1}{2} i (\omega^2 - k_z^2) E_{23} + \mathcal{O}(\epsilon_l R) , \quad (6.56)$$

$$\Psi_4 = \ddot{E}_{m\bar{m}} + \mathcal{O}(\epsilon_l R) = \left( \frac{\omega^2 - k_z^2}{\omega^2 + k_z^2} \right) \omega^2 (E_{00} + E_{33}) + 2\omega^2 E_{22} + 2i\omega^2 E_{12} + \mathcal{O}(\epsilon_l R) , \quad (6.57)$$

$$\Phi_{22} = \ddot{E}_{m\bar{m}} + \mathcal{O}(\epsilon_l R) = \left( \frac{\omega^2 - k_z^2}{\omega^2 + k_z^2} \right) \omega^2 (E_{00} + E_{33}) + \mathcal{O}(\epsilon_l R) , \quad (6.58)$$

where  $\ddot{E} = E_{,\tilde{u}\tilde{u}}$  and  $\tilde{u} = t - v_g z$ .

Hence by Eqs.(6.4) and (6.7), we get, for massless modes  $\omega_1$  associated to the plane wave  $E_{\mu\nu}^{k_1}$ , the following NP-quantities

$$\Psi_2 = \Psi_3 = \Phi_{22} = 0 , \quad (6.59)$$

$$0 \neq \Psi_4 = -2\omega_1^2 \hat{\epsilon}^{(+)}(\omega_1) + 2i\omega_1^2 \hat{\epsilon}^{(\times)}(\omega_1) \left( e^{i\omega_1(t-z)} + c.c. \right) , \quad (6.60)$$

and amplitudes

$$\begin{aligned} p_1^{(l)}(\mathbf{e}_z, t) &= p_2^{(x)}(\mathbf{e}_z, t) = p_3^{(y)}(\mathbf{e}_z, t) = p_6^{(b)}(\mathbf{e}_z, t) = 0 , \\ p_4^{(+)}(\mathbf{e}_z, t) &= -2\omega_1^2 \hat{\epsilon}^{(+)}(\omega_1) e^{i\omega_1(t-z)} + c.c. , \\ p_5^{(\times)}(\mathbf{e}_z, t) &= 2\omega_1^2 \hat{\epsilon}^{(\times)}(\omega_1) e^{i\omega_1(t-z)} + c.c. , \end{aligned} \quad (6.61)$$

that is, for frequency  $\omega_1$ , we obtain the two standard  $+$  and  $\times$  transverse modes with helicity 2 and the  $E(2)$  classification is  $N_2$ . More precisely the  $+$  mode is generated by the d.o.f.  $\hat{\epsilon}^{(+)}(\omega_1)$  while the  $\times$  mode is generated by the d.o.f.  $\hat{\epsilon}^{(\times)}(\omega_1)$ .

On the other hand, by Eqs.(6.4) and (6.7) for massive modes  $\omega_2$  and  $\omega_3$  associated to the plane wave  $E_{\mu\nu}^{k_m}$  with  $m$  ranging over  $\{2, 3\}$ , we get the following NP-quantities

$$\Psi_3 = \Psi_4 = 0 , \quad (6.62)$$

$$0 \neq \Psi_2 = \frac{1}{36} (\omega_m^2 - k_z^2) \hat{B}_m(k_z) e^{i(\omega_m t - k_z z)} + \mathcal{O}(\epsilon_l R) + c.c. , \quad (6.63)$$

$$0 \neq \Phi_{22} = \frac{\omega_m^2}{3} \hat{B}_m(k_z) e^{i(\omega_m t - k_z z)} + \mathcal{O}(\epsilon_l R) + c.c. , \quad (6.64)$$

and amplitudes

$$\begin{aligned} p_1^{(l)}(\tilde{\mathbf{e}}_z, t) &= \frac{1}{36} (\omega_m^2 - k_z^2) \hat{B}_m(k_z) e^{i(\omega_m t - k_z z)} + \mathcal{O}(\epsilon_l R) + c.c. , \\ p_2^{(x)}(\tilde{\mathbf{e}}_z, t) &= p_3^{(y)}(\mathbf{e}_z, t) = p_4^{(+)}(\mathbf{e}_z, t) = p_5^{(\times)}(\mathbf{e}_z, t) = 0 , \\ p_6^{(b)}(\tilde{\mathbf{e}}_z, t) &= \frac{\omega_m^2}{3} \hat{B}_m(k_z) e^{i(\omega_m t - k_z z)} + \mathcal{O}(\epsilon_l R) + c.c. , \end{aligned} \quad (6.65)$$

that is for each frequency  $\omega_2$  and  $\omega_3$ , we obtain one mixed longitudinal-transverse scalar mode with helicity 0 and the  $E(2)$  classification is  $II_6$ . More specifically both scalars  $\Psi_2$  and  $\Phi_{22}$  are driven by one d.o.f.  $B_m$  for each frequency  $\omega_m$ . Specifically, this fact means that both longitudinal and breathing modes, related to each  $\omega_m$ , show themselves always coupled in a single mixed scalar mode. In conclusion, we obtained three polarizations, two tensor polarizations and one mixed scalar polarization, and four modes, two tensor modes and two scalar modes. This result is because the two mixed scalar modes, associated with each frequency, exhibit the same polarization. If we go up with the order of the theory equivalent to polynomial curvature Lagrangian  $f(R, \square R, \dots, \square^k R)$ , for every two orders, a single mixed longitudinal-transverse scalar mode has to be added as shown in Table 1.

## 7 Higher-Order Teleparallel Gravity not equivalent to Higher-Order Curvature Gravity

Finally let us investigate a generic teleparallel Lagrangian of order  $n$  which does not contain the boundary term  $B$ . To get the field equations of any order  $n$ , we have to introduce the linear differential terms  $\square^k T$ , in analogy with the Lagrangian  $L_{\square^k R} = \sqrt{-g} (R + \sum_{k=0}^p a_k R \square^k R)$ , that is

$$L_{\square^k T}^{neq} = \frac{e}{2\kappa^2} \left( T + a_0 T^2 + \sum_{k=1}^p a_k T \square^k T \right). \quad (7.1)$$

By varying the Lagrangian  $L_{\square^k T}^{neq}$  plus the material Lagrangian term  $L_m$  with respect to a tetrad basis  $e^a_\rho$ , we obtain the  $(2p+2)$ -order equations:

$$\begin{aligned} & \frac{4}{e} \partial_\sigma \left[ e \left( 1 + 2a_0 T + 2 \sum_{k=1}^p a_k \square^k T \right) S_a^{\rho\sigma} \right] - 4 \left( 1 + 2a_0 T + 2 \sum_{k=1}^p a_k \square^k T \right) T^\mu_{\nu a} S_\mu^{\nu\rho} \\ & + \left( T + a_0 T^2 + \sum_{k=1}^p a_k T \square^k T \right) e_a^\rho - e_a^\rho \sum_{k=1}^p a_k \sum_{h=1}^k \square^{h-1} T (\nabla^\rho \nabla_\nu + \nabla_\nu \nabla^\rho) \square^{k-h} T \\ & + \sum_{k=1}^p a_k \sum_{h=1}^k \square^{h-1} T \partial_\eta \square^{k-h} T \left( e_a^\eta e_b^\rho \partial^\nu e^b_\nu + T_a^{\eta\rho} - e_a^\eta T^\rho - g^{\eta\rho} T_a \right) \\ & - \sum_{k=1}^p a_k \sum_{h=1}^k \partial_\sigma \left[ (e_a^\rho g^{\eta\sigma} - e_a^\sigma g^{\eta\rho} - e_a^\eta g^{\sigma\rho}) \square^{h-1} T \partial_\eta \square^{k-h} T \right] = 2\kappa^2 \mathcal{T}^{(m)}_a{}^\rho. \end{aligned} \quad (7.2)$$

Thus, we linearize field Eqs. (7.2) keeping up to first order terms in  $E^a_\mu$ . We get

$$2\partial_\sigma S_\tau^{\rho\sigma(1)} = \kappa^2 \mathcal{T}_\tau^{\rho(0)}. \quad (7.3)$$

Adopting the approximation (4.24), one obtains

$$\square \bar{E}^\mu_\rho = -\kappa^2 \mathcal{T}_\tau^{\rho(0)}. \quad (7.4)$$

Solving Eq.(7.4) in vacuum we get the solutions

$$E_{\rho\tau}(x) = \int \frac{d^3 \mathbf{k}}{(2\pi)^{3/2}} \hat{C}_{\rho\tau}(\mathbf{k}) e^{ik^\alpha x_\alpha} + c.c., \quad (7.5)$$

with  $k^2 = 0$  which, for a wave propagating along the  $+\hat{z}$  direction, becomes

$$E_{\mu\nu}(t, z) = \int \frac{d^3\mathbf{k}}{(2\pi)^{3/2}} \sqrt{2} \left[ \hat{\epsilon}^{(+)}(\mathbf{k}) \epsilon_{\mu\nu}^{(+)} + \hat{\epsilon}^{(\times)}(\mathbf{k}) \epsilon_{\mu\nu}^{(\times)} \right] e^{i\omega(t-z)} + c.c. , . \quad (7.6)$$

This results means that we achieve the Einstein GWs with two transverse tensor polarizations and helicity 2. In other words, the absence of boundary term  $B$  in the Lagrangian generates only the two standard  $+$  and  $\times$  polarizations. Further modes emerge if we introduce the boundary term  $B$  that excites extra polarizations. This result generalizes the one obtained in [30] for  $f(T)$  gravity. In other words, higher-order teleparallel gravity without boundary terms is equivalent to General Relativity from the point of view of GWs.

A summary of polarizations and helicity states is reported in Table 1 for several extended teleparallel theories.

| Lagrangian              | Order       | Frequency                                    | Polarization | Type                              | Modes<br>d.o.f. | State  | Helicity | Mass      |
|-------------------------|-------------|--|--------------|-----------------------------------|-----------------|--|----------|-----------|
| $L_{f(T)}$              | 2th         | $\omega_1$                                   | 2            | transverse tensor                 | 2               | $\epsilon_{\mu\nu}^{(+)}, \epsilon_{\mu\nu}^{(\times)}$  | 2        | 0         |
| $L_{f(T,B)}$            | 4th         | $\omega_1, \omega_2$                         | 3            | transverse tensor<br>mixed scalar | 3               | $\epsilon_{\mu\nu}^{(+)}, \epsilon_{\mu\nu}^{(\times)}$<br>$\epsilon_{\mu\nu}^{(s_2)}$   | 2, 0     | $M_2$     |
| $L_{\square T}^{eq}$    | 6th         | $\omega_1, \omega_2$<br>$\omega_3$           | 3            | transverse tensor<br>mixed scalar | 4               | $\epsilon_{\mu\nu}^{(+)}, \epsilon_{\mu\nu}^{(\times)}$<br>$\epsilon_{\mu\nu}^{(s_2)}, \epsilon_{\mu\nu}^{(s_3)}$                                | 2, 0     | $M_3$     |
| $L_{\square^2 T}^{eq}$  | 8th         | $\omega_1, \omega_2$<br>$\omega_3, \omega_4$ | 3            | transverse tensor<br>mixed scalar | 5               | $\epsilon_{\mu\nu}^{(+)}, \epsilon_{\mu\nu}^{(\times)}$<br>$\epsilon_{\mu\nu}^{(s_2)}, \epsilon_{\mu\nu}^{(s_3)}$<br>$\epsilon_{\mu\nu}^{(s_4)}$ | 2, 0     | $M_4$     |
| $\vdots$                | $\vdots$    | $\vdots$                                     | $\vdots$     | $\vdots$                          | $\vdots$        | $\vdots$   | $\vdots$ |           |
| $L_{\square^p T}^{eq}$  | $2(p+2)$ th | $\omega_1, \dots$<br>$\dots, \omega_{p+2}$   | 3            | transverse tensor<br>mixed scalar | $p+3$           | $\epsilon_{\mu\nu}^{(+)}, \epsilon_{\mu\nu}^{(\times)}$<br>$\epsilon_{\mu\nu}^{(s_2)}, \dots$<br>$\dots, \epsilon_{\mu\nu}^{(s_{p+2})}$          | 2, 0     | $M_{p+2}$ |
| $L_{\square^k T}^{neq}$ | $2(p+1)$ th | $\omega = q$                                 | 2            | tensor                            | 2               | $\epsilon_{\mu\nu}^{(+)}, \epsilon_{\mu\nu}^{(\times)}$  | 2        | 0         |

Table 1: Classification of Gravitational Waves for Extended Teleparallel Theories of Gravity

## 8 Conclusions

Teleparallelism is an approach to gravity equivalent to General Relativity, if torsion and curvature scalars are linear into the gravitational Lagrangian. Considering  $f(R)$  and  $f(T)$  theories, or, more generally, higher-order theories such as  $f(R, \square R, \dots, \square^k R)$  and  $f(T, \square T, \dots, \square^k T)$  the non-equivalence emerges because the involved d.o.f. are different. To describe the same physics, one has to take into account the boundary term  $B$  with its derivatives. The equivalence is restored according to the relation (2.10).

As shown above, we derived the teleparallel Lagrangian equivalent to  $L_{\square R} = -R + a_0 R^2 + a_1 R \square R$  considering the boundary term  $B$  and its derivatives. The linearized field equations, in the low energy regime, have been solved in vacuum adopting a distribution theory approach. Thus, we have derived the GWs of sixth-order equivalent teleparallel gravity and investigated their properties. GWs show three polarizations and four oscillation modes related to the four d.o.f. More in detail, in addition to the two tensor modes of General Relativity of frequency  $\omega_1$ , both transversely polarized, massless, with helicity 2 and propagation speed  $c$ , we obtained two further scalar modes associated with each frequency  $\omega_2$  and  $\omega_3$ . Every individual scalar mode is partly transversally polarized and partly longitudinally polarized because governed by a single d.o.f., respectively  $\tilde{B}_2(k_z)$  and  $\tilde{B}_3(k_z)$ : they form a single mixed transverse-longitudinal massive scalar mode with helicity 0 and propagation speed less than  $c$ . These results can be both demonstrated adopting the geodesic deviation equation and the Newmann-Penrose formalism, generalized to slightly massive waves. It is important to point out that in every higher-order teleparallel theory, equivalent to the polynomial Lagrangian  $f(R, \square R, \dots, \square^k R)$ , every two higher-orders excite a mixed scalar mode. On the other hand, considering theories like  $f(T, \square T, \dots, \square^k T)$  is completely equivalent to General Relativity from the point of view of GWs. We stress again the crucial relevance of the boundary term  $B$  to improve the number of GW modes and polarizations.

This means that precise polarization measurements cannot only invalidate or promote a given theory of gravity [60, 61, 62, 63, 64, 65, 66] but also suggest if metric or teleparallel representation better represent the gravitational interaction. See also [8] for a discussion on this point.

## Acknowledgements

SC is supported in part by the INFN sezione di Napoli, *iniziative specifiche* MOONLIGHT2 and QGSKY.

## A Appendix: Differential Identities

Let  $f$  and  $g$  be two scalar functions and  $A^\mu$  a contravariant vector, the following differential identities hold:

$$f \square g = \nabla^\nu (f \nabla_\nu g - g \nabla_\nu f) + g \square f , \quad (\text{A.1})$$

$$\square (fg) = f \square g + g \square f + 2 \nabla^\mu f \nabla_\mu g , \quad (\text{A.2})$$

$$f \square^k g = \nabla^\nu (\dots)_\nu + g \square^k f , \quad (\text{A.3})$$

$$\partial_\mu (e A^\mu) = e \nabla_\mu A^\mu . \quad (\text{A.4})$$

## B Appendix: Main Variations

Here a summary of main variations adopted along the text:

$$\delta e = e e_a{}^\rho \delta e^a{}_\rho = -e e_a{}_\rho \delta e^a{}^\rho , \quad (\text{B.1})$$

$$\delta e_a{}^\rho = -e_a{}^\sigma e_b{}^\rho \delta e^b{}_\sigma , \quad (\text{B.2})$$

$$\delta g^{\mu\nu} = (-g^{\nu\rho} e_a{}^\mu - g^{\mu\rho} e_a{}^\nu) \delta e^a{}_\rho , \quad (\text{B.3})$$

$$\delta T = -4T^\mu_{\nu a} S_\mu^{\nu\rho} \delta e^a_\rho - 4S_a^{\rho\sigma} \delta (\partial_\sigma e^a_\rho) , \quad (\text{B.4})$$

$$\delta \square T = \nabla_\mu \nabla_\nu T \delta g^{\mu\nu} + \square \delta T - g^{\mu\nu} \left( \delta \tilde{\Gamma}^\eta_{\nu\mu} - \delta K^\eta_{\nu\mu} \right) \partial_\eta T , \quad (\text{B.5})$$

$$\begin{aligned} \delta \square T = & \left[ -e_a^\nu (\nabla^\rho \nabla_\nu + \nabla_\nu \nabla^\rho) T + (e_a^\eta e_b^\rho \partial^\nu e^b_\nu + T_a^\eta{}^\rho - e_a^\eta T^\rho - g^{\eta\rho} T_a) \partial_\eta T \right] \delta e^a_\rho \\ & + (e_a^\rho g^{\eta\sigma} - e_a^\sigma g^{\eta\rho} - e_a^\eta g^{\rho\sigma}) \partial_\eta T \delta (\partial_\sigma e^a_\rho) + \square \delta T , \end{aligned} \quad (\text{B.6})$$

$$T \delta \square^k T = -2e_a^\nu \sum_{h=1}^k \square^{h-1} T \nabla^{(\rho} \nabla_{\nu)} \square^{k-h} T \delta e^a_\rho - e \sum_{h=1}^k \square^{h-1} T \partial_\eta \square^{k-h} T g^{\mu\nu} \delta \overset{\circ}{\Gamma}^\eta_{\nu\mu} + e \square^k T \delta T , \quad (\text{B.7})$$

$$\delta T^\nu{}_\lambda = - \left( T^{\rho\lambda}{}_a + e_a^\lambda T^\rho + g^{\lambda\rho} T_a \right) \delta e^a_\rho + (e_a^\rho g^{\lambda\sigma} - e_a^\sigma g^{\lambda\rho}) \delta (\partial_\sigma e^a_\rho) , \quad (\text{B.8})$$

$$\delta B = \delta \left[ \frac{2}{e} \partial_\sigma (e T^{\nu\sigma}{}_\nu) \right] = -\frac{B}{e} \delta e + \frac{2}{e} \partial_\mu (T^\mu \delta e + e \delta T^\mu) . \quad (\text{B.9})$$

## References

- [1] F. W. Hehl, P. von der Heyde, G. D. Kerlick, and J. M. Nester, *General relativity with spin and torsion: Foundations and prospects* Rev. Mod. Phys. **48**, 393 (1976).
- [2] J. B. Fonseca-Neto, C. Romero and S. P. G. Martinez, *Scalar torsion and a new symmetry of general relativity*, Gen. Relativ. Gravit. **45**, 1579-1601 (2013).
- [3] K. Hayashi and T. Shirafuji, *New general relativity* Phys. Rev. D **19**, 3524 (1979).
- [4] J. W. Maluf, *The teleparallel equivalent of general relativity*, Ann. Phys. **525**, 339 (2013).
- [5] L. Combi, G. E. Romero, *Is Teleparallel Gravity Really Equivalent to General Relativity?*, Ann. Phys. **530**, 1700175 (2018).
- [6] G. Kofinas and E. N. Saridakis, *Teleparallel equivalent of Gauss-Bonnet gravity and its modifications*, Phys. Rev. D **90**, 084044 (2014).
- [7] R. Aldrovandi, J. G. Pereira, *Teleparallel Gravity: An Introduction*. Fundamental Theories of Physics, vol. **173** (Springer, Dordrecht, 2013).
- [8] Y. Cai, S. Capozziello, M. De Laurentis, E. N. Saridakis, *f(T) teleparallel gravity and cosmology*, Rept. Prog. Phys. **79**, 106901 (2016).
- [9] N. D. Birrell and P. C. W. Davies, *Quantum Fields in Curved Space*, Cambridge Univ. Press, Cambridge (1984).
- [10] S. Capozziello, M. De Laurentis, *Extended Theories of Gravity*, Phys. Rept. **509**, 167 (2011).
- [11] S. Nojiri and S. D. Odintsov, *Unified cosmic history in modified gravity: from F(R) theory to Lorentz non-invariant models*, Phys. Rept. **505** (2011) 59
- [12] S. Nojiri, S. D. Odintsov and V. K. Oikonomou, *Modified Gravity Theories on a Nutshell: Inflation, Bounce and Late-time Evolution*, Phys. Rept. **692** (2017) 1

- [13] A. Petrov, *Introduction to Modified Gravity*, Springer Briefs in Physics, (2020) Cham Switzerland.
- [14] R. Ferraro and F. Fiorini, *Non-trivial frames for  $f(T)$  theories of gravity and beyond*, Phys. Lett. B **702**, 75 (2011).
- [15] R. Ferraro,  *$f(R)$  and  $f(T)$  theories of modified gravity*, AIP Conf.Proc. **1471**, 103-110 (2012) arXiv:1204.6273 [gr-qc].
- [16] S. Capozziello, M. Capriolo and L. Caso, *Weak field limit and gravitational waves in  $f(T, B)$  teleparallel gravity*, Eur. Phys. J. C **80**, 156 (2020).
- [17] J. B. Dent, S. Dutta and E. N. Saridakis,  *$f(T)$  gravity mimicking dynamical dark energy. Background and perturbation analysis*, JCAP **1101** (2011) 009
- [18] G. Otalora, E. N. Saridakis, *Modified teleparallel gravity with higher-derivative torsion terms*, Phys. Rev. D **94**, 084021 (2016).
- [19] S. Bahamonde, S. Capozziello, *Noether symmetry approach in  $f(T, B)$  teleparallel cosmology*, Eur. Phys. J. C **77**, 107 (2017).
- [20] S. Capozziello, M. Capriolo and L. Caso, *Weak field limit and gravitational waves in higher order gravity*, Int. J. Geom. Methods Mod. Phys. **16** No.03, 1950047 (2019).
- [21] Y. F. Cai, C. Li, E. N. Saridakis, and L.Q. Xue,  *$f(T)$  gravity after GW170817 and GRB170817A*, Phys. Rev. D **97**, 103513 (2018).
- [22] R. C. Nunes, S. Pan and E. N. Saridakis, *New observational constraints on  $f(T)$  gravity through gravitational-wave astronomy*, Phys. Rev. D **98**, 104055 (2018).
- [23] M. Hohmann, C. Pfeifer, J. L. Said and U. Ualikhanova, *Propagation of gravitational waves in symmetric teleparallel gravity theories*, Phys. Rev. D **99** 024009 (2019).
- [24] M. Hohmann, M. Krssak, C. Pfeifer and U. Ualikhanova, *Propagation of gravitational waves in teleparallel gravity theories*, Phys. Rev. D **98**, (2018) 124004 (2018).
- [25] M. Hohmann, *Polarization of gravitational waves in general teleparallel theories of gravity*, Astron. Rep. **62**, 890 (2018).
- [26] D. Liang, Y. Gong, S. Hou and Y. Liu, *Polarizations of gravitational waves in  $f(R)$  gravity*, Phys. Rev. D **95**, 104034 (2017).
- [27] T. Katsuragawa, T. Nakamura, T. Ikeda and S. Capozziello, *Gravitational Waves in  $F(R)$  Gravity: Scalar Waves and the Chameleon Mechanism*, Phys. Rev. D **99**, 124050 (2019).
- [28] S. Capozziello, R. Cianci, M. De Laurentis and S. Vignolo, *Testing metric-affine  $f(R)$ -gravity by relic scalar gravitational waves* Eur. Phys. J. C **70**, 341 (2010).
- [29] H. R. Kausar, L. Philippoz, and P. Jetzer, *Gravitational wave polarization modes in  $f(R)$  theories*, Phys. Rev. D **93**, 124071 (2016).
- [30] K. Bamba, S. Capozziello, M. De Laurentis, S. Nojiri and D. Saez-Gomez, *No further gravitational wave modes in  $F(T)$  gravity*, Phys. Lett. B **727**, 194 (2013).



- [31] S. Gottloeber, H. J. Schmidt and A. A. Starobinsky, *Sixth Order Gravity and Conformal Transformations*, Class. Quant. Grav. **7** (1990) 893.
- [32] A. Battaglia Mayer and H. J. Schmidt, *The de Sitter space-time as attractor solution in eighth order gravity*, Class. Quant. Grav. **10** (1993) 2441.
- [33] L. Amendola, A. Battaglia Mayer, S. Capozziello, F. Occhionero, S. Gottloeber, V. Muller and H. J. Schmidt, *Generalized sixth order gravity and inflation*, Class. Quant. Grav. **10** (1993) L43.
- [34] L. D. Landau and E. M. Lifshitz, *The Classical Theory of Fields*, Course of Theoretical Physics Vol. 2 4th ed, Butterworth- Heinemann, Oxford (1975).
- [35] S. M. Carroll, *Spacetime And Geometry*, Addison Wesley, San Francisco (2004).
- [36] R. M. Wald, *General Relativity*, The University of Chicago Press, Chicago (1984).
- [37] R. Aldrovandi and J. G. Pereira, *An Introduction to Geometrical Physics*, World Scientifics, Singapore (1995) **77**, 107 (2017).
- [38] N. Straumann, *General Relativity*, Springer-Verlag, Dordrecht (2013).
- [39] M. Nakahara, *Geometry, topology and physics* IOP Publishing, London (2003).
- [40] H. Abedi and S. Capozziello, *Gravitational waves in modified teleparallel theories of gravity*, Eur. Phys. J. C **78**, 474 (2018).
- [41] Yu. N. Obukhov, J. G. Pereira, *Teleparallel origin of the Fierz picture for spin-2 particle*, Phys. Rev. D **67**, 044008 (2003).
- [42] V. D. Gerrit, *Distribution Theory. Convolution, Fourier Transform, and Laplace Transform*, De Gruyter, Berlin (2013).
- [43] E. Poisson and C. M. Will, *Gravity Newtonian, Post-Newtonian, Relativistic*, Cambridge University Press, Cambridge (2014). doi:10.1017/CBO9781139507486
- [44] C. M. Will, *Theory and Experiment in Gravitational Physics*, Cambridge University Press, Cambridge (2018). doi:10.1017/9781316338612.
- [45] E. Newman and R. Penrose, *An approach to gravitational radiation by a method of spin coefficients*, J. Math. Phys. **3**, 566 (1962).
- [46] Y. H. Hyun, Y. Kim, and S. Lee, *Exact amplitudes of six polarization modes for gravitational waves*, Phys. Rev. D **99**, 124002 (2019).
- [47] D. Bessada and O. D. Miranda, *CMB polarization and theories of gravitation with massive gravitons*, Classical Quantum Gravity **26**, 045005 (2009).
- [48] M. E. S. Alves, O. D. Miranda, and J. C. N. de Araujo, *Extra polarization states of cosmological gravitational waves in alternative theories of gravity*, Classical Quantum Gravity **27**, 145010 (2010).

- [49] W. L. S. de Paula, O. D. Miranda, and R. M. Marinho, *Polarization states of gravitational waves with a massive graviton*, Classical Quantum Gravity **21**, 4595 (2004).
- [50] R. V. Wagoner, *Scalar-Tensor Theory and Gravitational Waves*, Phys. Rev. D **1**, 3209 (1970).
- [51] G. Farrugia, J. L. Said, V. Gakis, and E. N. Saridakis, *Gravitational waves in modified teleparallel theories*, Phys. Rev. D **97**, 124064 (2018).
- [52] Y. Gong and S. Hou, *Gravitational Wave Polarizations in  $f(R)$  Gravity and Scalar- Tensor Theory*, EPJ Web Conf. **168**, 01003 (2018).
- [53] D. M. Eardley, D. L. Lee, and A. P. Lightman, *Gravitational-Wave Observations as a Tool for Testing Relativistic Gravity*, Phys. Rev. D **8**, 3308 (1973).
- [54] D. M. Eardley, D. L. Lee, A. P. Lightman, R. V. Wagoner and C. M. Will, *Gravitational-Wave Observations as a Tool for Testing Relativistic Gravity*, Phys. Rev. Lett. **30**, 884 (1973).
- [55] M. Maggiore and A. Nicolis, *Detection strategies for scalar gravitational waves with interferometers and resonant spheres*, Phys. Rev. D **62**, 024004 (2000).
- [56] C. M. Will, *The confrontation between general relativity and experiment*, Living Rev. Relativity **17**, 4 (2014).
- [57] M. E. S. Alves, O. D. Miranda, and J. C. N. de Araujo, *Probing the  $f(R)$  formalism through gravitational wave polarizations*, Phys. Lett. B **679**, 401 (2009).
- [58] Y. S. Myung, *Propagating degrees of freedom in  $f(R)$  gravity*, Adv. High Energy Phys. **2016**, 3901734 (2016).
- [59] S. Capozziello, C. Corda, and M. De Laurentis, *Massive gravitational waves from  $f(R)$  theories of gravity: Potential detection with LISA*, Phys. Lett. B **669**, 255 (2008).
- [60] C. Bogdanos, S. Capozziello, M. De Laurentis and S. Nesseris, *Massive, massless and ghost modes of gravitational waves from higher-order gravity*, Astropart. Phys. **34**, 236 (2010).
- [61] B. P. Abbott et al., LIGO Scientific and Virgo Collaborations, *GW170817: Observation of Gravitational Waves from a Binary Neutron Star Inspiral*, Phys.Rev.Lett. **119**,161101 (2017).
- [62] B. P. Abbott et al., LIGO Scientific and Virgo Collaborations, *Gravitational Waves and Gamma-rays from a Binary Neutron Star Merger: GW170817 and GRB 170817A*, ApJ Lett. **848**, L13 (2017).
- [63] B. P. Abbott et al., LIGO Scientific and Virgo Collaborations, *Search for Tensor, Vector, and Scalar Polarizations in the Stochastic Gravitational-Wave Background*, Phys. Rev. Lett. **120**, 201102 (2018).
- [64] L. Lombriser and A. Taylor, *Breaking a Dark Degeneracy with Gravitational Waves*, JCAP **03** (2016) 031.
- [65] S. Capozziello, M. Capriolo and M. Transirico, *The gravitational energy-momentum pseudotensor: The cases of  $f(R)$  and  $f(T)$  Gravity*, Int. J. Geom. Methods Mod. Phys. **15** No.sup01, 1850164 (2018).

- [66] S. Capozziello, M. Capriolo and M. Transirico, *The gravitational energy-momentum pseudo-tensor of higher-order theories of gravity*, Ann. Phys. **525**, 1600376 (2017).