

PRODUCT THEOREM ON DELTA INVARIANTS VIA ADDING A GENERAL BOUNDARY

CHUYU ZHOU

ABSTRACT. It's well-known that adding a general boundary would create K-stability. As an application, we reprove product theorem for delta invariants of Fano varieties.

CONTENTS

1. Introduction	1
Acknowledgement	2
2. Preliminaries	2
3. Product theorem for K-stability	3
References	4

1. INTRODUCTION

It's well known that a K-unstable Fano manifold cannot admit Kähler-Einstein (KE) metrics. However, it can admit conic KE metrics along some smooth divisors. This inspires the concept of twisted K-stability. In [Der16], Dervan introduced the concept of twisted K-stability and later it is algebraically reformulated using twisted generalized Futaki invariants in [BLZ19], which indicates that K-unstable Fano manifolds can be twisted K-stable. For example, if X is a given K-unstable Fano manifold and $H \in |-lK_X|$ is a general smooth divisor on X , where $-lK_X$ is a very ample line bundle, then the pair $(X, \frac{1-\beta}{l}H)$ is uniformly K-stable for sufficiently small $0 < \beta \ll 1$. Thus there exists a conic KE metric $w(\beta)$ such that the following equation holds:

$$\text{Ric}(w(\beta)) = \beta w(\beta) + (1 - \beta)[H].$$

The following theorem proved by [BL18] are natural products of this phenomenon, that is, adding a general boundary would create K-stability.

Theorem 1.1. ([BL18, Section 7]) *Let (X, Δ) be a log Fano pair, then the following statements are equivalent.*

- (1) $\delta(X, \Delta) \geq \mu$ for some positive $0 < \mu \leq 1$,
- (2) For any rational $0 < \epsilon < \mu$, there exists an element $D \in |-K_X - \Delta|_{\mathbb{Q}}$ such that the pair $(X, \Delta + (1 - \epsilon)D)$ is K-semistable,
- (3) For any rational $0 < \epsilon < \mu$, there exists an element $D \in |-K_X - \Delta|_{\mathbb{Q}}$ such that the pair $(X, \Delta + (1 - \epsilon)D)$ is uniformly K-stable.

As an application of this principle, we reprove the following product theorem for delta invariants of Fano varieties, which is originally proved by [Zhu20].

Theorem 1.2. *Let $(X_i, \Delta_i), i = 1, 2$, be two log Fano pairs.*

- (1) *Both pairs are K -semistable if and only if $(X_1 \times X_2, p_1^* \Delta_1 + p_2^* \Delta_2)$ is K -semistable,*
(2) *If one of the two pairs is K -unstable, then*

$$\delta(X_1 \times X_2, p_1^* \Delta_1 + p_2^* \Delta_2) = \min\{\delta(X_1, \Delta_1), \delta(X_2, \Delta_2)\}.$$

- (3) *In general, we have following inequality,*

$$\delta(X_1 \times X_2, p_1^* \Delta_1 + p_2^* \Delta_2) \geq \min\{\delta(X_1, \Delta_1), \delta(X_2, \Delta_2), 1\}.$$

Acknowledgement. The author would like to thank Yuchen Liu and Ziquan Zhuang for beneficial comments. The author is supported by grant European Research Council (ERC-804334).

2. PRELIMINARIES

In this section, we present some necessary preliminaries. Throughout the note we work over complex number field \mathbb{C} . A log pair (X, Δ) consists of a normal projective variety X and an effective \mathbb{Q} -divisor Δ on X such that $K_X + \Delta$ is \mathbb{Q} -Cartier. We say a log pair (X, Δ) is log Fano if the pair admits klt singularities and $-K_X - \Delta$ is ample. The \mathbb{Q} -linear system $|-K_X - \Delta|_{\mathbb{Q}}$ is defined as follows:

$$|-K_X - \Delta|_{\mathbb{Q}} := \{D \geq 0 \mid D \sim_{\mathbb{Q}} -K_X - \Delta\}.$$

By the works of [FO18, BJ20], we can give the following definition for delta invariants of log Fano pairs.

Definition 2.1. Let (X, Δ) be a log Fano pair. We define

$$\delta_m(X, \Delta) := \inf_E \frac{A_{X, \Delta}(E)}{S_m(E)} \quad \text{and} \quad \delta(X, \Delta) := \inf_E \frac{A_{X, \Delta}(E)}{S_{X, \Delta}(E)},$$

where E runs over all prime divisors over X . Here

$$A_{X, \Delta}(E) = \text{ord}_E(K_Y - f^*(K_X + \Delta)) + 1$$

for some log resolution $f : Y \rightarrow X$ such that $E \subset Y$; and

$$S_m(E) = \sup_{D_m} \text{ord}_E(D_m),$$

where D_m is of the form $\frac{\sum_j \{s_j=0\}}{m \dim H^0(X, -m(K_X + \Delta))}$ and $\{s_j\}_j$ is a complete basis of the vector space $H^0(X, -m(K_X + \Delta))$; and

$$S_{X, \Delta}(E) = \frac{1}{\text{vol}(-K_X - \Delta)} \int_0^\infty \text{vol}(-f^*(K_X + \Delta) - tE) dt.$$

The following result is now well-known.

Theorem 2.2. ([FO18, BJ20]) *Let (X, Δ) be a log Fano pair, then*

- (1) $\lim_{m \rightarrow \infty} \delta_m(X, \Delta) = \delta(X, \Delta)$,
(2) (X, Δ) is K -semistable if and only if $\delta(X, \Delta) \geq 1$,
(3) (X, Δ) is uniformly K -stable if and only if $\delta(X, \Delta) > 1$.

One can just put this theorem as the definition of K -semistability (resp. uniform K -stability) of (X, Δ) .

3. PRODUCT THEOREM FOR K-STABILITY

In this section, we will prove Theorem 1.2. We first show the following result on K-stability of product varieties.

Theorem 3.1. *Let $(X_i, \Delta_i), i = 1, 2$, be two log Fano pairs, then $(X_1 \times X_2, p_1^* \Delta_1 + p_2^* \Delta_2)$ is K-semistable if and only if both pairs are K-semistable.*

Proof. The only if direction is easy. Suppose $(X_1 \times X_2, p_1^* \Delta_1 + p_2^* \Delta_2)$ is K-semistable. For any m -basis type divisor $D_m^{(i)}$ for (X_i, Δ_i) , we see that $p_1^* D_m^{(1)} + p_2^* D_m^{(2)}$ is an m -basis type divisor for $(X_1 \times X_2, p_1^* \Delta_1 + p_2^* \Delta_2)$. Denote $\delta_m := \delta_m(X_1 \times X_2, p_1^* \Delta_1 + p_2^* \Delta_2)$, then the pair

$$(X_1 \times X_2, p_1^* \Delta_1 + p_2^* \Delta_2 + \delta_m(p_1^* D_m^{(1)} + p_2^* D_m^{(2)}))$$

is log canonical, so are $(X_1, \Delta_1 + \delta_m D_m^{(1)})$ and $(X_2, \Delta_2 + \delta_m D_m^{(2)})$. Thus $\delta_m(X_i, \Delta_i) \geq \delta_m$, which implies that both $(X_i, \Delta_i), i = 1, 2$, are K-semistable.

For the converse direction, we just apply Theorem 1.1 to the case $\mu = 1$. For any fixed rational $0 < \epsilon < 1$, there exist $D_i \in |-K_{X_i} - \Delta_i|_{\mathbb{Q}}, i = 1, 2$ such that the pairs

$$(X_1, \Delta_1 + (1 - \epsilon)D_1) \quad \text{and} \quad (X_2, \Delta_2 + (1 - \epsilon)D_2)$$

are both uniformly K-stable by Theorem 1.1 (3). By [LTW19], one can construct KE metrics on the pairs $(X_i, \Delta_i + (1 - \epsilon)D_i), i = 1, 2$. Therefore, by taking product, the pair

$$(X_1 \times X_2, p_1^* \Delta_1 + p_2^* \Delta_2 + (1 - \epsilon)(p_1^* D_1 + p_2^* D_2))$$

also admits a KE metric, hence is K-semistable (even K-polystable) by [Ber16]. Again by Theorem 1.1 (2), we see the pair $(X_1 \times X_2, p_1^* \Delta_1 + p_2^* \Delta_2)$ is K-semistable. The proof is finished. \square

If we assume one of the two pairs is K-unstable, then we have following more precise result on delta invariant of the product variety.

Theorem 3.2. *Let $(X_i, \Delta_i), i = 1, 2$, be two log Fano pairs. Suppose one of the two pairs is K-unstable, then*

$$\delta(X_1 \times X_2, p_1^* \Delta_1 + p_2^* \Delta_2) = \min\{\delta(X_1, \Delta_1), \delta(X_2, \Delta_2)\}.$$

Proof. We assume $\delta(X_1, \Delta_1) \leq \delta(X_2, \Delta_2)$. We first show

$$\delta(X_1 \times X_2, p_1^* \Delta_1 + p_2^* \Delta_2) \geq \delta(X_1, \Delta_1).$$

We apply Theorem 1.1 (3). For any rational $0 < \epsilon < \delta(X_1, \Delta_1)$, one can find an element $D_1 \in |-K_{X_1} - \Delta_1|_{\mathbb{Q}}$ and an element $D_2 \in |-K_{X_2} - \Delta_2|_{\mathbb{Q}}$ such that both $(X_i, \Delta_i + (1 - \epsilon)D_i)$ are uniformly K-stable. By Theorem 3.1, the pair

$$(X_1 \times X_2, p_1^* \Delta_1 + p_2^* \Delta_2 + (1 - \epsilon)(p_1^* D_1 + p_2^* D_2))$$

is K-semistable. By Theorem 1.1 (2) we see that

$$\delta(X_1 \times X_2, p_1^* \Delta_1 + p_2^* \Delta_2) \geq \delta(X_1, \Delta_1).$$

We next show $\delta(X_1 \times X_2, p_1^* \Delta_1 + p_2^* \Delta_2) \leq \delta(X_1, \Delta_1)$. It suffices to show that

$$\delta_m(X_1 \times X_2, p_1^* \Delta_1 + p_2^* \Delta_2) \leq \delta_m(X_1, \Delta_1).$$

Similar to the proof of Theorem 3.1, we arbitrarily choose m -basis type divisors $D_m^{(i)}$ for (X_i, Δ_i) , then $p_1^* D_m^{(1)} + p_2^* D_m^{(2)}$ is an m -basis type divisor for $(X_1 \times X_2, p_1^* \Delta_1 + p_2^* \Delta_2)$. Denote $\delta_m := \delta_m(X_1 \times X_2, p_1^* \Delta_1 + p_2^* \Delta_2)$, we see the pair

$$(X_1 \times X_2, p_1^* \Delta_1 + p_2^* \Delta_2 + \delta_m(p_1^* D_m^{(1)} + p_2^* D_m^{(2)}))$$

is log canonical, so are the pairs $(X_i, \Delta_i + \delta_m D_m^{(i)})$, $i = 1, 2$. Therefore we have

$$\delta_m(X_1, \Delta_1) \geq \delta_m.$$

Taking the limit we have

$$\delta(X_1, \Delta_1) \geq \delta(X_1 \times X_2, p_1^* \Delta_1 + p_2^* \Delta_2).$$

The proof is finished. □

Proof of Theorem 1.2. The proof is just a combination of Theorem 3.1 and Theorem 3.2. □

REFERENCES

- [Ber16] Robert J. Berman, *K-polystability of \mathbb{Q} -Fano varieties admitting Kähler-Einstein metrics*, Invent. Math. **203** (2016), no. 3, 973–1025. MR3461370 ↑3
- [BJ20] Harold Blum and Mattias Jonsson, *Thresholds, valuations, and K-stability*, Adv. Math. **365** (2020), 107062, 57. MR4067358 ↑2
- [BL18] Harold Blum and Yuchen Liu, *Openness of uniform K-stability in families of \mathbb{Q} -Fano varieties*, arXiv e-prints (2018Aug), arXiv:1808.09070, available at 1808.09070. ↑1
- [BLZ19] Harold Blum, Yuchen Liu, and Chuyu Zhou, *Optimal destabilization of K-unstable Fano varieties via stability thresholds*, arXiv e-prints (2019Jul), arXiv:1907.05399, available at 1907.05399. ↑1
- [Der16] Ruadhaí Dervan, *Uniform stability of twisted constant scalar curvature Kähler metrics*, Int. Math. Res. Not. IMRN **15** (2016), 4728–4783. MR3564626 ↑1
- [FO18] Kento Fujita and Yuji Odaka, *On the K-stability of Fano varieties and anticanonical divisors*, Tohoku Math. J. (2) **70** (2018), no. 4, 511–521. MR3896135 ↑2
- [LTW19] Chi Li, Gang Tian, and Feng Wang, *The uniform version of Yau-Tian-Donaldson conjecture for singular Fano varieties*, arXiv e-prints (2019Mar), arXiv:1903.01215, available at 1903.01215. ↑3
- [Zhu20] Ziquan Zhuang, *Product theorem for K-stability*, Adv. Math. **371** (2020), 107250, 18. MR4108221 ↑1

ÉCOLE POLYTECHNIQUE FÉDÉRALE DE LAUSANNE (EPFL), MA C3 615, STATION 8, 1015 LAUSANNE, SWITZERLAND

Email address: chuyu.zhou@epfl.ch