# ALGEBRAIC CURVES ADMITTING INNER AND OUTER GALOIS POINTS

SATORU FUKASAWA

ABSTRACT. There are two purposes in this article. One is to present a criterion for the existence of a birational embedding into a projective plane with inner and outer Galois points for algebraic curves. Another is to classify plane curves of degree d admitting an inner Galois point P and an outer Galois point Q with  $G_PG_Q = G_P \rtimes G_Q$  or  $G_P \ltimes G_Q$ , under the assumption that the characteristic pis zero or p does not divide d - 1.

## 1. INTRODUCTION

In 1996, Hisao Yoshihara introduced the notion of a Galois point for a plane curve  $C \subset \mathbb{P}^2$  over an algebraically closed field k of characteristic  $p \geq 0$  ([1, 11, 13]): a point  $P \in \mathbb{P}^2$  is called a Galois point, if the extension  $k(C)/\pi_P^*k(\mathbb{P}^1)$  of function fields induced by the projection  $\pi_P$  from P is a Galois extension. The associated Galois group is denoted by  $G_P$ . Furthermore, a Galois point P is said to be inner (resp. outer), if  $P \in C \setminus \operatorname{Sing}(C)$  (resp. if  $P \in \mathbb{P}^2 \setminus C$ ). In this article, we investigate (possibly singular) plane curves C in the case where there exist an inner Galois point P and an outer Galois point Q.

Since we do not assume that the smoothness of C, sometimes we start with a (reduced, irreducible) smooth projective curve X. In this case, C is the image of some morphism  $\varphi : X \to \mathbb{P}^2$ , which is birational onto its image.

In [3], the present author presented a criterion for the existence of a birational embedding with two Galois points P and Q. In this criterion, it is assumed that both points P and Q are inner, or outer. In this article, we consider the case where P is inner and Q is outer.

**Theorem 1.** Let  $G_1$  and  $G_2$  be finite subgroups of Aut(X), and let  $P \in X$ . Then the two conditions

1

<sup>2010</sup> Mathematics Subject Classification. 14H50, 14H05, 14H37.

Key words and phrases. Galois point, plane curve, Galois group, automorphism group

The author was partially supported by JSPS KAKENHI Grant Number JP19K03438.

- (a)  $X/G_1 \cong \mathbb{P}^1$  and  $X/G_2 \cong \mathbb{P}^1$ , and
- (b) there exists  $\eta \in G_2$  such that  $P + \sum_{\sigma \in G_1} \sigma(\eta(P)) = \sum_{\tau \in G_2} \tau(P)$

are satisfied, if and only if there exists a birational embedding  $\varphi : X \to \mathbb{P}^2$  of degree  $|G_2|$  and an outer Galois point  $Q \in \mathbb{P}^2 \setminus \varphi(X)$  exists for  $\varphi(X)$  such that  $\varphi(P)$  is an inner Galois point, and  $G_{\varphi(P)} = G_1$  and  $G_Q = G_2$ .

In [6], the present author and Speziali classified plane curves admitting two outer Galois points  $P_1$  and  $P_2$  with  $G_{P_1}G_{P_2} = G_{P_1} \rtimes G_{P_2}$ . The second purpose in this article is to classify C admitting an inner Galois point P and an outer Galois point Q with  $G_P G_Q = G_P \rtimes G_Q$  or  $G_P \ltimes G_Q$ . The case where  $G_P G_Q = G_P \rtimes G_Q$  is determined as follows.

**Theorem 2.** Let C be a plane curve of degree  $d \ge 3$ . Assume that p = 0 or d - 1 is prime to p. Then the following conditions are equivalent.

- (a) There exist an inner Galois point P and an outer Galois point Q such that  $G_P G_Q = G_P \rtimes G_Q;$
- (b) There exist an inner Galois point P and an outer Galois point Q such that  $G_P G_Q = G_P \times G_Q;$
- (c) C is in one of the following cases:
  - -p = 0 or d is prime to p, and C is projectively equivalent to the plane curve defined by  $x^{d-1} + y^d + c = 0$ , where c = 0 or 1;
  - p > 0, d is divisible by p, and C is projectively equivalent to the plane curve defined by  $x^{d-1} + (\sum_{m \mid p^i-1} \alpha_i y^{p^i})^m + c = 0$ , where  $d = p^e m$ , m is not divisible by p,  $\alpha_e \alpha_0 \neq 0$ , and c = 0 or 1.

On the other hand, there exist examples of plane curves such that  $G_P G_Q = G_P \ltimes G_Q$  and  $G_P G_Q \neq G_P \times G_Q$ . In this case, such curves are classified as follows.

**Theorem 3.** Let C be a plane curve of degree  $d \ge 3$ . Assume that p = 0 or d(d-1) is prime to p. Then there exist an inner Galois point P and an outer Galois point Q such that  $G_PG_Q = G_P \ltimes G_Q$  and  $G_PG_Q \neq G_P \times G_Q$ , if and only if C is in one of the following cases:

(a)  $p \neq 2, 3, d = 3$  and C is projectively equivalent to the curve defined by

$$y^{2}x + (x+1)^{2}(x-8) = 0.$$

(b)  $p \neq 2, 3, d = 4$  and C is projectively equivalent to the curve defined by

$$y^{3}x + (x+1)^{3}(x+9) = 0.$$

For the proof of the if-part of Theorem 3, we prove the existence of triples  $(G_1, G_2, P)$  with conditions (a) and (b) in Theorem 1 such that  $G_1G_2 \cong S_3$  or  $A_4$ , where  $S_3$  (resp.  $A_4$ ) is the symmetric (resp. alternative) group of degree three (resp. of degree four). For the case where d is divisible by p, the following holds.

**Proposition 1.** Let p > 0 and let d be divisible by p. Then there exist an inner Galois point P and an outer Galois point Q such that  $G_PG_Q = G_P \ltimes G_Q$ ,  $G_PG_Q \neq$  $G_P \times G_Q$  and  $\overline{PQ}$  is not the tangent line at P, if and only if d is a power of p, and C is projectively equivalent to the curve defined by

$$y^{d-1}x + (x+1)^d = 0.$$

## 2. Proof of Theorem 1

Proof of Theorem 1. We consider the if-part. Since  $\varphi(P)$  and Q are Galois points with  $G_{\varphi(P)} = G_1$  and  $G_Q = G_2$ , it follows that fixed fields  $k(X)^{G_{\varphi(P)}} = k(X)^{G_1}$  and  $k(X)^{G_Q} = k(X)^{G_2}$  are rational. Condition (a) is satisfied. Let  $\ell$  be the line passing through points  $\varphi(P)$  and Q, and let  $D = \varphi^* \ell$ . Since Q is an outer Galois point and  $G_Q = G_2$ ,

$$D = (\pi_Q \circ \varphi)^* (\pi_Q(\varphi(P))) = \sum_{\tau \in G_2} \tau(P)$$

(see [12, III.7.1, III.7.2, III.8.2]). Let  $\eta \in G_2 \setminus \{1\}$ . Then  $\varphi(\eta(P)) \in \ell$ . To prove that condition (b) is satisfied, we would like to prove that

$$D - P = \sum_{\sigma \in G_1} \sigma(\eta(P)).$$

Assume that  $\eta(P) \neq P$ . Then  $\varphi(\eta(P)) \in \ell \setminus \{\varphi(P)\}$ . Since  $\varphi(P)$  is an inner Galois point and  $G_{\varphi(P)} = G_1$ , it follows that

$$D - P = (\pi_{\varphi(P)} \circ \varphi)^* (\pi_{\varphi(P)} \circ \varphi(\eta(P))) = \sum_{\sigma \in G_1} \sigma(\eta(P)).$$

The claim follows. Assume that  $\eta(P) = P$ . Then  $\ell$  is the tangent line at  $\varphi(P)$ . Since  $\varphi(P)$  is an inner Galois point, it follows that  $P \in \operatorname{supp}(\pi_{\varphi(P)} \circ \varphi)^*(\pi_{\varphi(P)} \circ \varphi(P))$  and

$$D - P = (\pi_{\varphi(P)} \circ \varphi)^* (\pi_{\varphi(P)} \circ \varphi(P)) = \sum_{\sigma \in G_1} \sigma(P).$$

It follows from the condition  $P = \eta(P)$  that the claim follows.

We consider the only-if part. Let  $\eta \in G_2$  be an automorphism as in condition (b). By condition (a), there exists a function  $f \in k(X)$  such that

$$k(X)^{G_1} = k(f), \ (f)_{\infty} = \sum_{\sigma \in G_1} \sigma(\eta(P))$$

(see [12, III.7.1, III.7.2, III.8.2]). Similarly, there exists  $g \in k(X)$  such that

$$k(X)^{G_2} = k(g), \ (g)_{\infty} = \sum_{\tau \in G_2} \tau(P).$$

Considering condition (b), we take a divisor

$$D := P + \sum_{\sigma \in G_1} \sigma(\eta(P)) = \sum_{\tau \in G_2} \tau(P).$$

Then  $f, g \in \mathcal{L}(D)$  and the sublinear system of |D| corresponding to a linear space  $\langle f, g, 1 \rangle$  is base-point-free. Since orders  $|G_1|$  and  $|G_2|$  are coprime, it follows that  $G_1 \cap G_2 = \{1\}$ . By this condition, the induced morphism

$$\varphi: X \to \mathbb{P}^2; \ (f:g:1)$$

is birational onto its image (see [3, Proposition 1]), and of degree  $|G_2|$ . Then the projection from the point  $\varphi(P) = (0:1:0)$  coincides with the morphism (f:1) and the projection from the point Q := (1:0:0) coincides with the morphism (g:1). Therefore,  $\varphi(P)$  is an inner Galois point with  $G_{\varphi(P)} = G_1$  and Q is an outer Galois point with  $G_Q = G_2$ .

## 3. Proof of Theorem 2

Let  $C \subset \mathbb{P}^2$  be an irreducible curve of degree  $d \geq 3$ , and let  $\varphi : X \to C$  be the normalization.

**Lemma 1.** Let  $P \in C$  be an inner Galois point, let  $Q \in \mathbb{P}^2 \setminus C$  be an outer Galois point and let  $\ell$  be the line passing through P and Q. Then the following hold.

- (a) The number of points of  $\operatorname{supp}(\varphi^*\ell)$  is 1 or d.
- (b) If  $G_P G_Q = G_P \rtimes G_Q$  or  $G_P \ltimes G_Q$ , then  $G_P$  fixes P. In particular, if  $\sigma \in G_P \setminus \{1\}$  fixes a point  $R \in \operatorname{supp}(\varphi^* \ell)$ , then R = P.
- (c) If  $G_P G_Q = G_P \rtimes G_Q$ , then  $\varphi^* \ell = dP$ .

*Proof.* We consider assertion (a). Let

$$\varphi^* \ell = \sum_{R \in \operatorname{supp}(\varphi^* \ell)} n(R) R.$$

Since Q is an outer Galois point and  $\ell \ni R$ , it follows from [12, III.7.2] that n(R) = n(R') for any  $R, R' \in \text{supp}(\varphi^* \ell)$ . If  $\ell$  is not a tangent line at P, then n(R) = n(P) = 1. This implies that the number of points of  $\text{supp}(\varphi^* \ell)$  is equal to d. Assume that  $\ell$  is the tangent line at P. Then the divisor

$$\varphi^* \ell - P = (n(P) - 1)P + \sum_{R \in \operatorname{supp}(\varphi^* \ell), \varphi(R) \neq P} n(R)R$$

coincides with  $(\pi_P \circ \varphi)^*(\pi_P(P))$ . Assume that  $R \in \operatorname{supp}(\varphi^*\ell)$  with  $\varphi(R) \neq P$  exists. Since P is a Galois point, it follows that n(R) = n(P) - 1. This is a contradiction. Therefore, in this case,  $\varphi^*\ell = dP$ .

We consider (b). By (a), the number of points of  $\operatorname{supp}(\varphi^*\ell)$  is 1 or d. For the former case, it follows from [12, III.8.2] that the assertion holds. We can assume that  $\operatorname{supp}(\varphi^*\ell)$  consists of d points. Let  $\sigma \in G_P \setminus \{1\}$ . By the assumption, there exist  $\sigma' \in G_P \setminus \{1\}$  and  $\tau, \tau' \in G_Q \setminus \{1\}$  such that  $\tau \sigma = \sigma' \tau'$ . Then  $\tau \sigma(P) = \sigma' \tau'(P) \in \varphi^{-1}(\ell)$ . Since  $\tau^{-1}(\varphi^{-1}(\ell)) \subset \varphi^{-1}(\ell)$ , it follows that  $\sigma(P) \in T_P C \cap \ell = \{P\}$ . Therefore,  $\sigma(P) = P$  follows.

We consider (c). Let  $\tau \in G_Q$  and let  $\sigma \in G_P \setminus \{1\}$ . By (b),  $\tau \sigma \tau^{-1}$  fixes  $\tau(P)$ . Since  $G_P G_Q = G_P \rtimes G_Q$ ,  $\tau \sigma \tau^{-1} \in G_P \setminus \{1\}$ . By (b),  $\tau(P) = P$ . It follows from [12, III.8.2] that the ramification index at P for  $\pi_Q$  is d. The assertion follows.

Proof of Theorem 2. We prove (a)  $\Rightarrow$  (b). By condition (a) and Lemma 1, the group  $G_P \rtimes G_Q$  fixes P. It follows from [9, Lemma 11.44] that if p = 0 or d(d-1) is prime to p, then  $G_P \rtimes G_Q$  is a cyclic group. Therefore,  $G_P \rtimes G_Q = G_P \times G_Q$ . Assume that p > 0 and p divides d. Let  $d = p^e m$ , where m is not divisible by p. It follows from [9, Lemma 11.44] that  $G_P \cong \mathbb{Z}/(d-1)\mathbb{Z}$  and there exist subgroups N, H such that  $G_Q = N \rtimes H$ ,  $|N| = p^e$  and  $H \cong (\mathbb{Z}/m\mathbb{Z})$ . Since  $|G_P \rtimes G_Q| = (d-1)p^e m$ , the subgroup N is a normal subgroup of  $G_P \rtimes G_Q$ , and hence  $G_P \rtimes N = G_P \times N$ . Since  $|G_P \rtimes H| = (d-1)m$  is not divisible by p, the group  $G_P \rtimes H$  is a cyclic group. Therefore,  $G_P \rtimes H = G_P \times H$ . As a consequence,  $G_P G_Q = G_P \times G_Q$  follows.

We prove (b)  $\Rightarrow$  (c). By assertion (b) in Lemma 1,  $\varphi^*T_PC = dP$ . By the assumption,  $G_Q$  acts on the subfield  $k(X)^{G_P}$ , which is rational. Assume that p = 0 or d(d-1) is prime to p. Since  $G_Q$  is totally ramified at  $\pi_P(P)$  and d is prime to p, it follows that  $G_Q$  is a cyclic group and there exists a ramification point  $R \in X/G_P$  different from  $\pi_P(P)$  (see, for example, [9, Theorem 11.91]). The line corresponding to R is denoted by  $\ell_Q$ . By the assumption that p = 0 or d-1 is prime to p, a similar argument can be applied to the morphism  $X \to X/G_Q$ . It is inferred that  $G_P$  is a

cyclic group and there exists a ramification point  $S \in X/G_Q$  different from  $\pi_Q(P)$ . The line corresponding to S is denoted by  $\ell_P$ . The linear system corresponding to  $\varphi: X \to C \subset \mathbb{P}^2$  is determined by three lines  $\ell, \ell_Q$  and  $\ell_P$ . For a suitable system of coordinates, we can assume that such lines are defined by Z = 0, Y = 0 and X = 0, respectively. Then the covering maps  $X \to X/G_P$  and  $X/G_P \to X/\langle G_P, G_Q \rangle$  are represented by  $(X:Y:Z) \mapsto (Y:Z)$  and  $(Y:Z) \mapsto (Y^d:Z^d)$ . When we represent the covering map  $X \to X/G_Q$  by  $(X:Y:Z) \mapsto (X:Z)$ , we infer that the covering map  $X/G_Q \to X/\langle G_Q, G_P \rangle$  is represented by  $(X:Z) \mapsto (aX^{d-1} + bZ^{d-1}:Z^{d-1})$  for some  $a, b \in k$ , by considering the branch points of the covering map  $X \to X/\langle G_P, G_Q \rangle$ . Therefore,  $y^d = ax^{d-1} + b$  holds in k(X). This gives the defining equation of C. For a suitable system of coordinates, we can assume that a = -1 and  $b \in \{-1, 0\}$ .

Assume that p > 0 and d is divisible by p. Let  $d = p^e m$ , where m is not divisible by p. Since  $G_Q$  is totally ramified at  $\pi_P(P)$ , it follows that m divides  $p^e - 1$  and there exist subgroups N, H such that  $G_Q = N \rtimes H, N \cong (\mathbb{Z}/p\mathbb{Z})^{\oplus e}$  and  $H \cong \mathbb{Z}/m\mathbb{Z}$  (see, for example, [9, Theorem 11.91]). Then there exists a ramification point  $R \in X/G_P$ different from  $\pi_P(P)$ . The line corresponding to R is denoted by  $\ell_Q$ . Considering the assumption that d-1 is prime to p, we infer that  $G_P$  is a cyclic group and there exists a ramification point  $S \in X/G_Q$  different from  $\pi_Q(P)$ . The line corresponding to S is denoted by  $\ell_P$ . The linear system corresponding to  $\varphi : X \to C \subset \mathbb{P}^2$  is determined by three lines  $\ell, \ell_Q$  and  $\ell_P$ . For a suitable system of coordinates, we can assume that such lines are defined by Z = 0, Y = 0 and X = 0, respectively. According to Lemma 2 below, the covering maps  $X \to X/G_P$  and  $X/G_P \to X/\langle G_P, G_Q \rangle$  are represented by  $(X : Y : Z) \mapsto (Y : Z)$  and  $(Y : Z) \mapsto (f(Y/Z)Z^d : Z^d)$  for a polynomial

$$f(y) = \left(\sum_{m \mid p^{i-1}} \alpha_{i} y^{p^{i}}\right)^{m},$$

where  $\alpha_e \ldots, \alpha_0 \in k$  and  $\alpha_e \alpha_0 \neq 0$ . When we represent the covering map  $X \to X/G_Q$  by  $(X : Y : Z) \mapsto (X : Z)$ , we infer that the covering map  $X/G_Q \to X/\langle G_Q, G_P \rangle$  is represented by  $(X : Z) \mapsto (aX^{d-1} + bZ^{d-1} : Z^{d-1})$  for some  $a, b \in k$ , by considering the branch points of the covering map  $X \to X/\langle G_P, G_Q \rangle$ . Therefore,  $(\sum_{m \mid p^i - 1} \alpha_i y^{p^i})^m = ax^{d-1} + b$  holds in k(X). This gives the defining equation of C. For a suitable system of coordinates, we can assume that a = -1 and  $b \in \{-1, 0\}$ . **Lemma 2.** Let p > 0, e > 0,  $m \mid p^e - 1$  and let  $\psi : \mathbb{P}^1 \to \mathbb{P}^1$  be a Galois covering of degree d such that the Galois group G is described as  $G = N \rtimes H$  for a p-subgroup N of order  $p^e$  and a cyclic group H of order m. Assume that  $\psi(1:0) = (1:0)$  and  $\psi(0:1) = (0:1)$ . If G fixes (1:0) and H fixes (0:1), then  $\psi(Y:Z) = (f(Y/Z)Y^d : Z^d)$  for some polynomial

$$f(y) = c \left(\sum_{m \mid p^{i}-1} \alpha_{i} y^{p^{i}}\right)^{m},$$

where  $\alpha_e, \ldots, \alpha_0, c \in k$  with  $\alpha_e \alpha_0 \neq 0$ .

Proof. The covering  $\psi$  is the composite map of the Galois coverings  $\psi_1 : \mathbb{P}^1 \to \mathbb{P}^1/N$ and  $\psi_2 : \mathbb{P}^1/N \to \mathbb{P}^1/(G/N)$ , where  $G/N \cong H$ . We can assume that  $\psi_1(1:0) =$ (1:0) and  $\psi_1(0:1) = (0:1)$ . Since N fixes (1:0) and is of degree  $p^e$ ,  $\psi_1 =$  $(g(Y/Z)Z^{p^e}:Z^{p^e})$  for some polynomial g(y) and there exists an additive subgroup  $S \subset k$  such that  $N = \{(Y:Z) \mapsto (Y + \alpha Z:Z) \mid \alpha \in S\}$ . It follows from the assumption  $\psi_1(0:1) = (0:1)$  that g(0) = 0. Since g(y) is invariant under the action of N, it follows from [8, Proposition 1.1.5 and Theorem 1.2.1] that

$$g(y) = \alpha_e y^{p^e} + \dots + \alpha_1 y^p + \alpha_0 y$$

for some  $\alpha_e, \ldots, \alpha_0 \in k$  with  $\alpha_e \alpha_0 \neq 0$ . Since the cyclic covering  $\psi_2$  is ramified at points (1 : 0) and (0 : 1),  $\psi_2(Y : Z) = (cY^m : Z^m)$  for some  $c \in k$ . Then  $\psi = (c(g(y))^m : 1)$ . Let  $\zeta$  be a primitive *m*-th root of unity. Since  $m \mid p^e - 1$ , it follows that  $g(\zeta y) = \zeta g(y)$ . Then  $\alpha_i \neq 0$  implies that  $(\zeta)^{p^i} = \zeta$ . We obtain the required form.  $\Box$ 

**Remark 1.** The curves defined by  $x^{d-1}+y^d+1=0$  or  $x^{d-1}+(\sum_{m \mid p^i-1} \alpha_i y^{p^i})^m+1=0$  are smooth. For these curves, points (1:0:0) and (0:1:0) are all Galois points. This was proved in [2, Theorem 3].

For the curve  $x^{d-1} + y^d = 0$ , the same holds. It is not difficult to prove it, since (0:0:1) is a cusp and (1:0:0) is a unique total inflection point.

**Remark 2.** (1) When p > 0 and d - 1 is a power of p,  $G_P G_Q = G_P \rtimes G_Q$ does not imply  $G_P G_Q = G_P \times G_Q$ . For (the plane model of) the Giulietti– Korchmáros curve

$$x^{q^3} + x - (x^q + x)^{q^2 - q + 1} - y^{q^3 + 1} = 0$$

(see [7]), where q is a power of p, P = (1 : 0 : 0) is an inner Galois point and Q = (0 : 1 : 0) is an outer Galois point ([5]). However,  $G_Q$  is not a normal subgroup of  $G_P G_Q = G_P \rtimes G_Q$ .

(2) Assume that p > 0 and p divides d - 1. In this case also, we can prove that  $G_Q \cong \mathbb{Z}/d\mathbb{Z}$  and the defining equation of C is of the form  $y^d + h(x) = 0$ , where h(x) is a polynomial of degree at most d, up to a projective equivalence.

## 4. Proof of Theorem 3

In this section, we consider plane curves admitting an inner Galois point P and an outer Galois point Q with  $G_P G_Q = G_P \ltimes G_Q$  and  $G_P G_Q \neq G_P \times G_Q$ .

**Lemma 3.** If d(d-1) is prime to p, then the number of points of  $\operatorname{supp}(\varphi^* \ell)$  is d.

Proof. By Lemma 1 (a), the number of points of  $\operatorname{supp}(\varphi^*\ell)$  is 1 or d. Assume the former. Note that  $G_P \ltimes G_Q$  fixes P. It follows from the assumption on the degree and [9, Lemma 11.44] that  $G_P \ltimes G_Q$  is a cyclic group. This implies that  $G_P G_Q = G_P \times G_Q$ . This is a contradiction.

Hereafter, we assume that  $\operatorname{supp}(\varphi^*\ell)$  consists of d points.

**Lemma 4.** For any  $\sigma \in G_P \setminus \{1\}$  and  $\tau \in G_Q \setminus \{1\}$ ,  $\sigma \tau \neq \tau \sigma$ .

*Proof.* Assume that  $\sigma \tau = \tau \sigma$ . By Lemma 1 (b), it follows that

$$\sigma\tau(P) = \tau\sigma(P) = \tau(P).$$

If  $\tau(P) \neq P$ , then by Lemma 1 (b),  $\sigma(\tau(P)) \neq \tau(P)$ . Therefore,  $\tau(P) = P$ . Since  $\operatorname{supp}(\varphi^*\ell)$  consists of d points, it follows that  $\tau = 1$ . This is a contradiction.  $\Box$ 

We assume that d-1 is prime to p. In this case, the covering map  $X/G_Q \to X/G_P G_Q$  is tame and totally ramified at  $\pi_Q(P)$ . Similarly to the proof of Theorem 2, it follows that  $G_P$  has an iclusion  $G_P \hookrightarrow PGL(3, k)$ ,  $G_P$  is a cyclic group of order d-1, and there exists a line  $\ell_P \subset \mathbb{P}^2 \setminus \{P\}$  with  $\ell_P \ni Q$  such that

$$\ell_P \cup \{P\} = \{R \in \mathbb{P}^2 \mid \sigma(R) = R \text{ for all } \sigma \in G_P\}.$$

For a suitable system of coordinates, P = (0 : 1 : 0), Q = (1 : 0 : 0), lines  $\ell_P$  and  $T_PC$  are defined by Y = 0 and X = 0 respectively, and C is defined by

$$y^{d-1}x + g(x) = 0,$$

where q(x) is a polynomial of degree d.

**Lemma 5.** The number of points of  $(C \setminus \text{Sing}(C)) \cap \ell_P$  is zero or one.

*Proof.* Assume that there exist two different smooth points  $R_1$  and  $R_2 \in \ell_P$ . Then there exists  $\tau \in G_Q \setminus \{1\}$  such that  $\tau(R_1) = R_2$ . Let  $\sigma \in G_P \setminus \{1\}$ . Then

$$\tau^{-1}\sigma^{-1}\tau\sigma(R_1) = \tau^{-1}\sigma^{-1}\tau(R_1) = \tau^{-1}\sigma^{-1}(R_2) = \tau^{-1}(R_2) = R_1$$

Note that  $\sigma^{-1}\tau\sigma \in G_Q$ , since  $G_PG_Q = G_P \ltimes G_Q$ . Since  $\pi_Q$  is not ramified at  $R_1$ ,  $\tau^{-1}\sigma^{-1}\tau\sigma = 1$ . Therefore,  $\tau\sigma = \sigma\tau$ . This is a contradiction to Lemma 4.

**Lemma 6.** Assume that  $d \ge 4$  and there exists a ramification point  $S \in C \setminus \ell_P$  of index  $m \ge 2$  for  $\pi_Q$ . Let n be the number of points of  $C \cap \ell_P$ . Then the following holds.

- (a) If  $d \ge 5$ , then  $n \ge \frac{m-1}{m}d$ .
- (b) m = 2.
- (c) There does not exist a point  $R \in C \cap \ell_P$  such that the multiplicity of R is two.

*Proof.* We consider (a). On the line  $\overline{QS}$ , there exist d/m ramification points of index m. Since any automorphism in  $G_P$  is a linear transformation, fixes Q and does not fix any point of  $C \cap \overline{QS}$ , it follows that there exist  $(d-1) \times (d/m)$  ramification points of index m. It follows from the Riemann-Hurwitz formula that for  $\pi_Q$ ,

$$2g - 2 \ge -2d + (d - 1) \times \frac{d}{m}(m - 1).$$

For  $\pi_P$ ,

$$2g - 2 \le -2(d - 1) + (d - 2) + (d - 2)n.$$

With these inequalities used, it follows that

$$n \ge \frac{m-1}{m}d - \frac{1}{m}\left(1 + \frac{2}{d-2}\right).$$

If  $d \ge 5$ , then  $\frac{1}{m} \left( 1 + \frac{2}{d-2} \right) < 1$ . Therefore,

$$n \ge \frac{m-1}{m}d.$$

We consider (b). Assume that  $d \ge 5$  and  $m \ge 3$ . Since the number of singular points on  $\ell_P$  is at most d/2, by (a), the number of smooth points on  $\ell_P$  is at least

$$\frac{m-1}{m}d - \frac{d}{2}.$$

If  $d \ge 7$  or  $m \ge 4$ , then this value is strictly more than 1. This is a contradiction to Lemma 5. Assume that d = 6 and m = 3. In this case,  $n \ge 4$  and there exists a smooth point on  $\ell_P$ . Therefore, the number of singular points on  $\ell_P$  is at most two, and hence, there exist two smooth points on  $\ell_P$ . This is a contradiction to Lemma 5. Assume that d = 4. Then m = 2 or 4. Assume that m = 4. Then by the proof of (a),

$$n \ge \frac{3}{4}d - \frac{1}{4}\left(1 + \frac{2}{d-2}\right) = \frac{5}{2}.$$

Therefore,  $n \geq 3$ . Then there exist two smooth points on  $\ell_P$ . This is a contradiction to Lemma 5.

We consider (c). By (b), d is even. Assume that  $R \in \ell_P$  is a point of multiplicity two. Note that  $\overline{RP} \cap C = \{R, P\}$ , by considering the defining equation. If  $\varphi^{-1}(R)$ consists of two points, then the degree of  $\pi_P$  is divisible by two. Since d is even and d-1 is odd, this is a contradiction. Therefore,  $\varphi^{-1}(R)$  consists of a unique point. Such a point also is denoted by R. Then the ramification index of R for  $\pi_P$  is d-1, and that of R for  $\pi_Q$  is two. The stabilizer subgroup  $G_Q(R)$  for  $G_Q$  consists of a unique involution  $\tau$ . Let  $\sigma \in G_P \setminus \{1\}$ . Since  $\tau$  and  $\sigma$  fix R,  $\sigma^{-1}\tau\sigma$  fixes R. It follows from the condition  $G_P G_Q = G_P \ltimes G_Q$  that  $\sigma^{-1}\tau\sigma \in G_Q(R)$ . Since  $|G_Q(R)| = 2$ ,  $\sigma^{-1}\tau\sigma = \tau$ . This is a contradiction to Lemma 4.

**Lemma 7.** Assume that all ramification points for  $\pi_Q$  are contained in the line  $\ell_P$ . Then the following holds.

- (a) p > 0.
- (b) The order of each element of  $G_Q \setminus \{1\}$  is equal to p. In particular,  $d = p^e$  for some e > 0.

*Proof.* It follows from the Riemann–Hurwitz formula that there exist a wild ramification. Therefore, p > 0. In this case, there exists an element  $\tau \in G_Q$  of order p. By Lemma 4, it follows that

$$G_Q = \{ \sigma^{-1} \tau \sigma \mid \sigma \in G_P \} \cup \{1\}$$

It follows from Sylow's theorem that  $d = p^e$  for some e > 0.

Proof of Theorem 3. We consider the only-if part. By Lemma 7, there exists a ramification point  $S \in C \setminus \ell_P$  for  $\pi_Q$ . We prove that  $d \leq 4$ . Assume that  $d \geq 5$ . By Lemma 6 (a) and (b),  $n \geq d/2$ . The number of smooth points on  $\ell_P$  is zero or one, by Lemma 5. If the number is zero, then, by Lemma 6 (c), there exist d/2 points of

multiplicity at least three on  $\ell_P$ . This is obviously a contradiction. Therefore, the number of smooth points on  $\ell_P$  is one. By Lemma 6 (c), there exist (d/2) - 1 points of multiplicity at least three on  $\ell_P$ . Then we have an inequality

$$1 + 3 \times \left(\frac{d}{2} - 1\right) \le d.$$

This implies that  $d \leq 4$ .

Let d = 4. By the assumption on the degree, it follows that  $p \neq 2, 3$ . By Lemmas 5 and 6, there exists a singular point of multiplicity 3 on  $\ell_P$ . In particular, C is rational. We can assume that for a suitable system of coordinates, P = (0 : 1 : 0), Q = (1 : 0 : 0) and C is defined by

$$y^{3}x + (x+1)^{3}(x+a) = 0,$$

where  $a \in k \setminus \{0\}$ . Let  $y = \beta$  be a branch point of  $\pi_Q$ , where  $\beta \in k \setminus \{0\}$ . Then there exist  $c, d \in k$  such that

$$\beta^3 x + (x+1)^3 (x+a) = (x^2 + cx + d)^2.$$

Comparing the coefficients of both sides, we have relations

$$a + 3 = 2c$$
,  $3a + 3 = c^2 + 2d$ ,  $3a + \beta^3 + 1 = 2cd$ ,  $a = d^2$ .

By these equations, we have

$$d^{4} - 6d^{2} + 8d - 3 = (d - 1)^{3}(d + 3) = 0.$$

If d = 1, then  $\beta = 0$ . This is a contradiction. It follows that d = -3, a = 9, c = 6and  $\beta^3 = -64$ . Then the curve C is defined by

$$y^{3}x + (x+1)^{3}(x+9) = y^{3}x + 64x + (x^{2} + 6x - 3)^{2} = 0.$$

Let d = 3. By the assumption on the degree, it follows that  $p \neq 2, 3$ . By Lemma 5, there exists a singular point on  $\ell_P$ . Then C is rational. Note that the index of all ramification points contained in  $C \setminus \ell_P$  for  $\pi_Q$  is three. We can assume that for a suitable system of coordinates, P = (0:1:0), Q = (1:0:0) and C is defined by

$$y^{2}x + (x+1)^{2}(x+a) = 0,$$

where  $a \in k \setminus \{0\}$ . Let  $y = \beta$  be a branch point of  $\pi_Q$ , where  $\beta \in k \setminus \{0\}$ . Then there exists  $c \in k$  such that

$$\beta^2 x + (x+1)^2 (x+a) = (x+c)^3.$$

Comparing the coefficients of both sides, we have relations

$$a + 2 = 3c$$
,  $2a + \beta^2 + 1 = 3c^2$ ,  $a = c^3$ .

By these equations, we have

$$c^{3} - 3c + 2 = (c - 1)^{2}(c + 2) = 0.$$

If c = 1, then  $\beta = 0$ . This is a contradiction. It follows that c = -2, a = -8 and  $\beta^2 = 27$ . Then the curve C is defined by

$$y^{2}x + (x+1)^{2}(x-8) = y^{2}x - 27x + (x-2)^{3} = 0.$$

To prove the if-part, using Theorem 1, we prove the existence of plane rational curves of d = 3 or 4 with required properties. We consider subgroups of  $\operatorname{Aut}(\mathbb{P}^1) \cong$ PGL(2,k). It is well known that  $A_4$  is a subgroup of PGL(2,k) (see, for example, [9, Theorem 11.91]). It is also well known that there exist subgroups  $G_1, G_2$  of  $A_4$ such that  $G_1$  is a cyclic group of order three,  $G_2$  is a normal subgroup of  $A_4$  with  $G_2 \cong (\mathbb{Z}/2\mathbb{Z})^{\oplus 2}$ , and  $G_1G_2 = G_1 \ltimes G_2 = A_4$ . Let  $P \in \mathbb{P}^1$  be a point such that  $G_1$ fixes P. We prove that conditions (a) and (b) in Theorem 1 are satisfied for the triple  $(G_1, G_2, P)$ . By Lüroth's theorem, condition (a) is satisfied. With the action of  $A_4$  considered, there exist three points  $P_2, P_3, P_4$  such that  $A_4P = \{P, P_2, P_3, P_4\}$ . Let  $\eta \in G_2 \setminus \{1\}$ . Then,

$$P + \sum_{\sigma \in G_1} \sigma(\eta(P)) = P + P_2 + P_3 + P_4 = \sum_{\tau \in G_2} \tau(P).$$

This implies condition (b) in Theorem 1. By Theorem 1, the required rational plane curve of d = 4 is obtained. For d = 3, a similar argument can be applied.

Proof of Proposition 1. We consider the only-if part. We prove that all ramification points for  $\pi_Q$  are contained in  $\ell_P$ . When  $d \geq 5$ , according to the first paragraph of the proof of the only-if part of Theorem 3, the claim follows. Assume by contradiction that there exists a ramification point in  $S \setminus \ell_P$  for  $\pi_Q$ . By Lemmas 5 and 6, if  $d \leq 4$ , then C is rational. If d = 4, then p = 2 and  $\pi_Q$  is wildly ramified at 6 ramification points contained in  $C \setminus \ell_P$ , by Lemma 6 (b). According to the Riemann-Hurwitz formula for  $\pi_Q$ , this is a contradiction. If d = 3, then p = 3 and  $\pi_Q$  is wildly ramified at 2 ramification points. This is a contradiction. Therefore, the claim follows for all  $d \geq 3$ . By Lemma 7 (a) and (b), p > 0 and  $d = p^e$  for some integer e > 0. Let  $R \in \varphi^{-1}(\ell_P)$  and let  $G_Q(R)$  be the stabilizer subgroup of  $G_Q$  of order  $p^k$ . By the assumption, k > 0. Let  $G_P(R)$  be the stabilizer subgroup of  $G_P$ . For any  $\sigma \in G_P(R)$ ,  $\sigma^{-1}G_Q(R)\sigma = G_Q(R)$ . It follows from Lemma 4 that  $|G_P(R)| + 1 \leq |G_Q(R)|$ . With the action of  $G_P$  on  $\varphi^{-1}(\ell_P)$  considered, the set  $\varphi^{-1}(\ell_P)$  contains

$$\frac{|G_P|}{|G_P(R)|} \ge \frac{p^e - 1}{p^k - 1}$$

points. With the action of  $G_Q$  considered, it follows that

$$p^{e} = |G_Q| \ge |G_Q(R)| \times \frac{p^{e} - 1}{p^{k} - 1} = p^{k} \times \frac{p^{e} - 1}{p^{k} - 1}$$

It follows that

$$p^{e-k} \le 1.$$

Therefore, k = e, that is,  $G_Q(R) = G_Q$ . This implies that  $C \cap \ell_P = \{\varphi(R)\}$ . For a suitable system of coordinates, we can assume that  $\varphi(R) = (-1:0:1)$ . Then C is defined by

$$y^{d-1}x + (x+1)^d = 0.$$

We consider the if-part. The curve C is projectively equivalent to the curve defined by

$$x - y^q = 0,$$

where p > 0 and q is a power of p. Let P = (0:0:1) and let  $Q \in \{Z = 0\} \setminus \{(1:0:0), (0:1:0)\}$ . According to [4], P is an inner Galois point and Q is an outer Galois point. Note that there exists an inclusion  $G_P \hookrightarrow PGL(3, k)$  and  $G_P$  fixes Q. It is inferred that  $G_PG_Q = G_P \ltimes G_Q$ . Since the line passing through P and Q intersects C at q points, it follows that  $G_PG_Q \neq G_P \rtimes G_Q$ , by Lemma 1.

**Remark 3.** For the curve of degree 4 in Theorem 3, the point P is a unique inner Galois point. This is proved easily, because two flexes are needed for each inner Galois points and there exists  $G_P \hookrightarrow PGL(3, k)$ . For the curve described in Proposition 1, the set of Galois points is determined in [4].

**Remark 4.** According to a classification of finite subgroups G of PGL(2, k) (see, for example, [9, Theorem 11.91]), a finite non-abelian subgroup G containing subgroups  $H_1, H_2 \subset G$  such that  $G = H_1 \ltimes H_2$ ,  $|H_2| = |H_1| + 1 \ge 4$  and G has a short orbit of length  $|H_2|$  must be one of the following:

(a)  $A_4$  with  $p \neq 2, 3$ ;

(b) the semidirect product of an elementary abelian *p*-group of order q with a cyclic group of order q - 1.

The cases (a) and (b) are described in Theorem 3 (b) and Proposition 1.

**Remark 5.** If  $\operatorname{supp}(\varphi^*\ell)$  consists of d points, then the action of the group  $G_P \ltimes G_Q$ on  $\operatorname{supp}(\varphi^*\ell)$  is doubly transitive.

**Remark 6.** Assume that  $\operatorname{supp}(\varphi^*\ell)$  consists of d points. Let  $P_1 = P$ , let  $P_2 = \tau(P)$  and let  $G_2 = \tau G_P \tau^{-1}$ , where  $\tau \in G_Q \setminus \{1\}$ . It is inferred that conditions in [3, Theorem 1] are satisfied for the 4-tuple  $(G_1, G_2, P_1, P_2)$ . Therefore, X has a birational embedding with two inner Galois points. Note that  $G_1$  fixes  $P_1$  and  $G_2$  fixes  $P_2$ . In this case, the group  $G = \langle G_{P_1}, G_{P_2} \rangle (= G_P \ltimes G_Q)$  has been classified by Korchmáros, Lia, and Timpanella [10]. It is noted that this birational embedding is different from the one given by the triple  $(G_P, G_Q, P)$  in the case of Theorem 3 (b). In fact, the embedding of d = 4 described in Theorem 3 does not admit two inner Galois points.

**Remark 7.** Assume that p > 0 and p divides d - 1. In this case also, we can prove that the defining equation of C is of the form  $(\sum_{m \mid p^i - 1} \alpha_i x^{p^i})^m + h(y) = 0$ , where  $d - 1 = p^e m$ , m is not divisible by p,  $\alpha_e \alpha_0 \neq 0$ , and h(y) is a polynomial of degree d, up to a projective equivalence.

## Acknowledgements

The author is grateful to Doctor Kazuki Higashine for helpful discussions.

#### References

- S. Fukasawa, Galois points for a plane curve in arbitrary characteristic, Proceedings of the IV Iberoamerican Conference on Complex Geometry, Geom. Dedicata 139 (2009), 211–218.
- [2] S. Fukasawa, On the number of Galois points for a plane curve in positive characteristic, III, Geom. Dedicata 146 (2010), 9–20.
- [3] S. Fukasawa, A birational embedding of an algebraic curve into a projective plane with two Galois points, J. Algebra 511 (2018), 95–101.
- [4] S. Fukasawa and T. Hasegawa, Singular plane curves with infinitely many Galois points, J. Algebra 323 (2010), 10–13.
- [5] S. Fukasawa and K. Higashine, Galois lines for the Giulietti–Korchmáros curve, Finite Fields Appl. 57 (2019), 268–275.
- [6] S. Fukasawa and P. Speziali, Plane curves possessing two outer Galois points, preprint, arXiv:1801.03198.

14

- [7] M. Giulietti and G. Korchmáros, A new family of maximal curves over a finite field, Math. Ann. 343 (2009), 229–245.
- [8] D. Goss, Basic structures of function fields arithmetic, Springer, Berlin, 1996.
- [9] J. W. P. Hirschfeld, G. Korchmáros and F. Torres, Algebraic curves over a finite field, Princeton Univ. Press, Princeton, 2008.
- [10] G. Korchmáros, S. Lia and M. Timpanella, Curves with more than one inner Galois point, J. Algebra 566 (2021), 374–404.
- [11] K. Miura and H. Yoshihara, Field theory for function fields of plane quartic curves, J. Algebra 226 (2000), 283–294.
- [12] H. Stichtenoth, Algebraic function fields and codes, Universitext, Springer-Verlag, Berlin, 1993.
- [13] H. Yoshihara, Function field theory of plane curves by dual curves, J. Algebra 239 (2001), 340–355.

DEPARTMENT OF MATHEMATICAL SCIENCES, FACULTY OF SCIENCE, YAMAGATA UNIVERSITY, KOJIRAKAWA-MACHI 1-4-12, YAMAGATA 990-8560, JAPAN

Email address: s.fukasawa@sci.kj.yamagata-u.ac.jp