# A Superdeterministic Toy Model

Sandro Donadi, Sabine Hossenfelder

Frankfurt Institute for Advanced Studies Ruth-Moufang-Str. 1, D-60438 Frankfurt am Main, Germany

#### **Abstract**

A superdeterministic toy model for quantum mechanics is introduced and discussed. It is demonstrated that, when averaged over the hidden variables, the model produces the same predictions as quantum mechanics. In the model considered here, the dynamics depends only on the settings of the measurement device at the detection time, not how those settings were chosen. This provides a counter-example to the claim that superdeterminism is fine-tuned and unscientific.

### 1 Introduction

Superdeterminism [1] is the only known consistent, local, and deterministic completion of quantum mechanics. Despite having surfaced in literature already in the 1950s as a solution to the measurement problem, the option never received much attention, largely because it is widely believed to be a "conspiracy" that requires "fine-tuning". Those arguments were already addressed in [1, 2]. The purpose of this paper is to enrich the discussion with a simple toy model that demonstrates just why these fine-tuning arguments are misleading. What we mean by "fine-tuning" here is that the model requires one to specify such a large amount of detail to make a prediction that it becomes scientifically useless.

Several superdeterministic toy models have previously been put forward in the literature [3, 4, 5, 6, 7, 8] (for a review, see [9]). The one introduced here has the advantage that it can be applied to any kind of quantum mechanical system with a finite dimensional Hilbert space. This new model should not be taken too seriously as a viable description of nature. It is neither pretty nor does it make sense as a fundamental theory for reasons that will be discussed later. However, it will serve as a proof of principle to show that superdeterministic theories are not necessarily fine-tuned and that Bell-type tests cannot rule them out.

Before we get to the equations, here is a general argument for why superdeterminism does not need to be fine-tuned: The hallmark of a superdeterministic model is that the outcome of a quantum measurement depends on the detector settings at the time of measurement<sup>1</sup>. The typical argument against superdeterminism is that there are many ways to choose detector settings,

<sup>&</sup>lt;sup>1</sup>If you find this confusing, you should start with the accompanying paper "Superdeterminism: A Guide for the Perplexed" [2].

and the superdeterministic model must work for all of them. This, it is then argued, requires that the model specifies what happens for a huge number of different cases and must still reproduce quantum mechanics on the average.

The problem with this argument is that there is no reason why, in a superdeterministic model, the measurement outcome should depend on how the detector settings were chosen any more than this is the case in quantum mechanics. The settings at the time of measurement alone are enough to get the correct average predictions. What happens in the brain of the experimenter (or with any other mechanism used to determine detector settings [10, 11, 12]) does not matter; the only thing that matters is the setting of the measurement device at the time of measurement. It follows that the average outcome is insensitive to the details, hence not fine-tuned.

We already know this is possible because it's what we do in quantum mechanics itself: We predict average measurement outcomes from the detector settings alone. The next section will support this idea with a concrete example. We will use units in which  $c = \hbar = 1$ .

### 2 The Toy Model

Suppose we have a quantum mechanical system described by a wave-function  $|\Psi\rangle$  and a Hamiltonian  $\hat{H}$  in a N-dimensional Hilbert space. We want to measure an observable with N eigenstates  $|i\rangle$ ,  $i\in\{1,...,N\}$ , each one corresponding to a different state of the detector at the time of detection<sup>2</sup>. The system could be a composite system and the detector could be distributed over different locations; the following will not depend on these details. If there are multiple detectors, we choose the space-time slicing so that the measurement takes place at the same coordinate time for all detectors.<sup>3</sup>

We then postulate that all the states in the Hilbert-space except the eigenstates of the measurement observable are unstable under any kind of disturbance that comes from the hidden variables. That is, we will introduce a deviation from quantum mechanics quantified by a parameter,  $\kappa > 0$ , which collapses arbitrary initial states into detector eigenstates. The measurement outcome will then be determined by the initial state of the system and the value of the hidden variables. When we average over these hidden variables, only the dependency on the initial state remains and the probabilistic outcome agrees with that of quantum mechanics. The reader can think of these variables as encoding the detailed degrees of freedom of the detector because that is the most minimal possibility, but it could be more complicated.

In the next subsection, we will start with the case N=2 to demonstrate the general idea, and then generalize to larger N.

<sup>&</sup>lt;sup>2</sup>Strictly speaking, of course, detection does not happen in a single moment but during a finite time-interval. This distinction, however, does not matter for the purposes of this toy model as will become clear later.

<sup>&</sup>lt;sup>3</sup>If there is no space-like slice that can accommodate all the detectors, then they are causally related, in which case the later detectors can be ignored because the state is already "collapsed".

#### **2.1** N = 2

For N = 2 we have

$$|\Psi(t)\rangle := a_1(t)|1\rangle + a_2(t)|2\rangle \quad , \tag{1}$$

with complex factors  $a_1(t)$  and  $a_2(t)$  that fulfill  $|a_1(t)|^2 + |a_2(t)|^2 = 1$ . We will denote the time of system preparation with  $t_p$  and that of measurement with  $t_d$ . Let us further denote with  $|\Psi^*(t)\rangle$  the solution to the normal Hamiltonian evolution

$$|\Psi^*(t)\rangle := \exp(-i\hat{H}(t-t_p))|\Psi(t_p)\rangle. \tag{2}$$

It will also be handy to define the coefficients of the eigenvectors at the time of measurement under the usual Hamiltonian evolution:

$$\alpha_I := \langle I | \Psi^*(t_{\rm d}) \rangle \ . \tag{3}$$

We know that in quantum mechanics the probability of measuring the I-th outcome is  $|\alpha_I|^2$ , so the superdeterministic theory must reproduce this on the average. These numbers can be calculated from  $\hat{H}$  alone, so the underlying dynamics that we are about to construct is unnecessary to make the probabilistic predictions, but this isn't the point. The point of having a superdeterministic model is to give a proper statistical interpretation to Born's rule and to get rid of the collapse postulate.

As hidden variables we use complex numbers that are uniformly distributed inside the complex unit circle. The radius of the distribution could be chosen differently for each of these variables, but this would just add unnecessary parameters. We will denote the random variable as  $\lambda_2$ , where the index 2 on  $\lambda_2$  refers to N=2.

With that, we construct, as usual, the density matrix  $\rho(t) := |\Psi(t)\rangle\langle\Psi(t)|$  but change the dynamical law to

$$\partial_t \rho(t) = -\mathrm{i} \left[ \hat{H}, \rho(t) \right] + \kappa \left( L_{21} \rho(t) L_{21}^{\dagger} - \frac{1}{2} \{ \rho_t, L_{21}^{\dagger} L_{21} \} \right) . \tag{4}$$

Here,  $\kappa$  is a constant of dimension energy (for the interpretation, see Section 3), the curly brackets denote the anti-commutator and

$$L_{21} := \theta(\sigma_2)|2\rangle\langle 1| + \theta(-\sigma_2)|1\rangle\langle 2|, \qquad (5)$$

$$L_{21} := \theta(\sigma_2)|2\rangle\langle 1| + \theta(-\sigma_2)|1\rangle\langle 2|,$$

$$\sigma_2 := |\lambda_2 \alpha_2|^2 - (1 - |\lambda_2|^2)|\alpha_1|^2.$$
(5)

Again, the indices on  $L_{21}$  and  $\sigma_2$  refer to the case N=2 and are merely there to make it easier to later generalize to higher N.  $\theta(\cdot)$ , as usual, denotes the Heaviside-function which we define as  $\theta(x) = 0$  for x < 0 and  $\theta(x) = 1$  otherwise. It must be emphasized that the  $\rho(t)$  in Eq. (4) is not the density matrix of quantum mechanics. To get the density matrix of quantum mechanics, we have to average over the hidden variable  $\lambda_2$ .

The dynamical law (4) has the Lindblad-form [13, 14]. It is the master-equation for one of the most common examples of decoherence, that of amplitude damping in a two-level system [15]. We will in the following work in the limit where  $\kappa$  is much larger than the largest eigenvalue of  $\hat{H}$ , which we will denote as  $E_{\text{max}}$ . In this limit,  $\kappa \gg E_{\text{max}}$ , it is then easy to explicitly solve (4). This solution (for details, see Section 4) has the following asymptotic behavior

$$\lim_{\kappa t \to \infty} \rho(t) = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \quad \text{for } \sigma_2 > 0 ,$$

$$\lim_{\kappa t \to \infty} \rho(t) = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \quad \text{for } \sigma_2 \le 0 .$$
(8)

$$\lim_{\kappa t \to \infty} \rho(t) = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \quad \text{for } \sigma_2 \le 0 \ . \tag{8}$$

This toy model, importantly, is not a model of spontaneous collapse [16] or, more generally a model of a stochastic process [17], because there is no stochasticity in the dynamics. The dynamics is determined by the time-independent random variable and not a random walk. If one selects a specific  $\lambda_2$ , then one knows for certain what  $\sigma_2$  is and one knows what the outcome of the evolution is.

We now have to show that the probabilities for the state to evolve to  $|1\rangle$  and  $|2\rangle$  come out correctly. As we just saw, in the limit where  $\kappa \gg E_{\text{max}}$  an initial state  $|\Psi(t)\rangle$  will go to  $|2\rangle$  if  $\sigma_2 > 0$ . We therefore have

$$P(|2\rangle) = P(\sigma_2 > 0) , \qquad (9)$$

where the P here and in the following denotes a probability. In terms of the random variable  $\lambda_2$ , we can write the condition  $\sigma_2 > 0$  as

$$\begin{aligned} |\lambda_{2}\alpha_{2}|^{2} &> (1 - |\lambda_{2}|^{2})|\alpha_{1}|^{2} \\ \Leftrightarrow & |\lambda_{2}\alpha_{2}|^{2} &> (1 - |\lambda_{2}|^{2})(1 - |\alpha_{2}|^{2}) \\ \Leftrightarrow & |\lambda_{2}|^{2} &> 1 - |\alpha_{2}|^{2} . \end{aligned}$$
(10)

Since  $\lambda_2$  is uniformly distributed in the unit circle, this means we are asking for the area of the ring between the radii  $\sqrt{1-|\alpha_2|^2}$  and 1 relative to the area of the whole disk. The area of the ring and the disk scale with the square of the radius, so the fraction is  $1 - 1 + |\alpha_2|^2 = |\alpha_2|^2$ , and we have

$$P(|2\rangle) = |\alpha_2|^2 \ . \tag{11}$$

It follows that the probability of the state to evolve to  $|1\rangle$  is  $|\alpha_1|^2$ . This is exactly Born's rule. It may appear irrelevant that the random variable is complex-valued, but note that if it was not, the probabilities would not scale with the area, hence not give Born's rule.

#### **2.2** N > 2

To generalize from N=2 to larger N, it helps to have a physical description of the process that the dynamical law (4) is commonly used to describe. For the case  $\sigma_2 > 0$ , the state  $|1\rangle$  is unstable and decays to  $|2\rangle$ , while for  $\sigma_2 \leq 0$ , it is the other way around,  $|2\rangle$  decays to  $|1\rangle$ . The constant  $\kappa$  determines the decay time. Provided that the contribution from the Hamiltonian can be neglected, this leads asymptotically to the limits in (8).

For N > 2 one then iteratively adds more random vectors  $\lambda_N$ , each of which is independently uniformly distributed on the unit disk and defines

$$\sigma_N := |\alpha_N \lambda_N|^2 - \sum_{I=1}^{N-1} |\alpha_I|^2 (1 - |\lambda_N|^2) .$$
 (12)

With that, we construct the Lindblad-operators recursively for higher N by drawing on the comparison with decaying states. In the step from N-1 to N, we introduce N-1 new operators

$$L_{NM} := \theta(\sigma_N)|N\rangle\langle M| + \theta(-\sigma_N)|M\rangle\langle N| \quad \text{for} \quad M \in \{1..N-1\} . \tag{13}$$

This means that we have in total N(N-1)/2 Lindblad-operators  $L_{KM}$  for N dimensions, and by convention we have labelled them so that the second index is always strictly smaller than the first. With that, the master equation can be written as

$$\partial_{t} \rho(t) = -\mathrm{i} \left[ \hat{H}, \rho(t) \right] + \kappa \sum_{K > M = 1}^{N} \left( L_{KM} \rho(t) L_{KM}^{\dagger} - \frac{1}{2} \{ \rho(t), L_{KM}^{\dagger} L_{KM} \} \right) . \tag{14}$$

As above for the case N = 2, we obtain the density matrix of quantum mechanics by averaging the  $\rho$  from Eq. (14) over the hidden variables.


Before we move on, let us briefly interpret the structure of the Lindblad-operators that we have constructed. The N-1 new operators that we add in each step N-1 to N have the property that they either take any initial state  $|K\rangle$  with K < N and asymptotically convert it to  $|N\rangle$  (if  $\sigma_N > 0$ ) or they take the initial state  $|N\rangle$  and distribute it evenly on the states  $|K\rangle$  with K < N (if  $\sigma_N \le 0$ ). This is illustrated in Fig. 1.

So, if  $\sigma_N > 0$ , we know that the result will be  $|N\rangle$ . Let us then look at the case  $\sigma_N \le 0$ . Once the state  $|N\rangle$  is cleared out and distributed over the subspace spanned by  $|1\rangle...|N-1\rangle$ , the same thing happens again for the remaining states, but this time depending on the sign of  $\sigma_{N-1}$ : If  $\sigma_{N-1} > 0$ , then all remaining states go to  $|N-1\rangle$ , if  $\sigma_{N-1} \le 0$ , then they will be distributed over the subspace spanned by  $|1\rangle...|N-1\rangle$ , and so on.

Having said that, we can now show that the generalization to N > 2 still fulfills Born's rule. This is straight-forward to see for the eigenstate  $|N\rangle$ . As just explained, if  $\sigma_N > 0$ , then the operators  $L_{NK}$  will convert any initial state  $|K\rangle$  with K < N into the state  $|N\rangle$ . One can calculate the probability for this the same way as above in (10) and easily finds

$$P(|N\rangle) = P(\sigma_N > 0) = |\alpha_N|^2.$$
(15)

Correspondingly, the probability that the state goes to any one of the other N-1 eigenstates is  $1-|\alpha_N|^2$ .



**Figure 1:** The dynamics induced by the Lindblad terms involving the operators  $L_{NM}$  with M = 1, 2, ...N - 1. For  $\sigma_N > 0$  the system will collapse to the state  $|N\rangle$ . For  $\sigma_N < 0$ , if the system is in the state  $|N\rangle$  will evolve evenly into the states  $|1\rangle, ... |N-1\rangle$ .

To obtain the probability that an initial state asymptotically goes to  $|N-1\rangle$ , one then needs to calculate the probability for

$$P(|N\rangle) = P(\sigma_N < 0 \land \sigma_{N-1} > 0) = P(\sigma_N < 0)P(\sigma_{N-1} > 0)$$
, (16)

because the random variables are by assumption uncorrelated.

For this, one has to take into account that for N-1 basis states  $\sum_{i=1}^{N-1} |\alpha_i|^2 \le 1$ , and therefore the probability for  $\sigma_{N-1}$  to be larger or smaller than one has to be divided by this sum. One sees this by rewriting the requirement  $\sigma_{N-1} > 0$  as

$$|\lambda_{N-1}\alpha_{N-1}|^{2} > (1 - |\lambda_{N-1}|^{2}) \sum_{i=1}^{N-2} |\alpha_{i}|^{2}$$

$$\Leftrightarrow |\lambda_{N-1}|^{2} \frac{|\alpha_{N-1}|^{2}}{\sum_{i=1}^{N-1} |\alpha_{i}|^{2}} > (1 - |\lambda_{N-1}|^{2}) \left(1 - \frac{|\alpha_{N-1}|^{2}}{\sum_{i=1}^{N-1} |\alpha_{i}|^{2}}\right)$$

$$\Leftrightarrow |\lambda_{N-1}|^{2} > 1 - \frac{|\alpha_{N-1}|^{2}}{\sum_{i=1}^{N-1} |\alpha_{i}|^{2}}.$$
(17)

So, by the same argument as before, the probability for  $\sigma_{N-1} > 0$  is now  $|\alpha_{N-1}|^2 / \sum_{i=1}^{N-1} |\alpha_i|^2$  (instead of just  $|\alpha_{N-1}|^2$ , as it was for N-1 states). Hence we have

$$P(|N-1\rangle) = (1 - |\alpha_N|^2) \frac{|\alpha_{N-1}|^2}{\sum_{i=1}^{N-1} |\alpha_i|^2} = |\alpha_{N-1}|^2 , \qquad (18)$$

which is what Born's rule requires.

It works the same way for all remaining eigenstates. For example, the probability of an initial state to go to  $|N-2\rangle$  is given by

$$P(|N-2\rangle) = P(\sigma_N \le 0 \land \sigma_{N-1} \le 0 \land \sigma_{N-2} > 0)$$
  
=  $P(\sigma_N \le 0)P(\sigma_{N-1} \le 0)P(\sigma_{N-2} > 0)$ , (19)

which is

$$(1 - |\alpha_N|^2) \left( 1 - \frac{|\alpha_{N-1}|^2}{\sum_{i=1}^{N-1} |\alpha_i|^2} \right) \frac{|\alpha_{N-2}|^2}{\sum_{i=1}^{N-2} |\alpha_i|^2} = |\alpha_{N-2}|^2.$$
 (20)

In general, then, the probability for any initial state to asymptotically come out as  $|K\rangle$  is

$$P(|K\rangle) = P(\sigma_N \le 0 \land \dots \land \sigma_{K+1} \le 0 \land \sigma_K > 0)$$

$$= (1 - |\alpha_N|^2) \prod_{J=N-1}^{K+1} \left(1 - \frac{|\alpha_J|^2}{\sum_{I=1}^J |\alpha_I|^2}\right) \left(\frac{|\alpha_K|^2}{\sum_{I=1}^K |\alpha_I|^2}\right) = |\alpha_K|^2.$$
 (21)

This completes our proof that Born's rule is fulfilled for all *N*.

The way we have defined the *N*-dimensional case depends on the order of the basis-states. Since the order is arbitrary, this seems unphysical. It does not really matter for the purposes of this model because the resulting probability distribution is independent of the ordering. However, one could make the model explicitly independent of the ordering by summing over all possible permutations.

One may note that this distribution fulfils the requirements for Born's rule laid out in [18].

# 3 Finetuning, Absence Of

One should not think of this model as a viable description of nature because the way that the random variables enter the dynamics has no good motivation. It is a toy model for an effective limit of a more fundamental theory. The parameter  $\kappa$ , for example, defines a time-scale like for decoherence and should be understood as induced by the interaction with a larger system that has been integrated out. That is,  $\kappa$  scales with the total size of the system and in the limit when the prepared state has no further interaction,  $\kappa=0$ , one just has normal quantum mechanics.

This is also why the model is not Lorenz-invariant: The environment defines a preferred frame, yet the environment does not explicitly appear. The model further violates energy conservation. Again, this is because it stands in for an effective description that, among other things, does not take into account the recoil (and resulting entanglement with) parts of the experimental equipment. For the same reason, as mentioned in footnoote 2, the time of measurement is not really *the* time of measurement, but a time at which we would for all practical purposes say the measurement has been completed.

Furthermore, since the operators  $L_{NK}$  are defined in terms of the eigenstates of the observable measured at the time of detection, we have still used an external definition of detection that is not included in the definition of the model. Now, the pointer basis defining these operators can be identified using the einselection methods introduced by Zurek [19] when one includes an environment, so this is not a problem per so. However, a more fundamental model is needed to explain how these operators come to couple to the prepared state the way that they do.

All these issues are resolvable in principle, but given that this model is not intended to make a lot of sense, putting more effort into it seems not a good time-investment. The reason for this little exercise was merely to demonstrate that there is nothing to fine-tune here. For a Hilbert-space of dimension N, we have N-1 uniformly distributed, complex, random variables. Picking a set of specific values for these variables will in the limit  $\kappa t \to \infty$  lead to one particular detector eigenstate: No collapse postulate necessary. But if we do not know what the values of

the random variables are, we average over all possible values and get Born's rule. The model has one free parameter,  $\kappa$ , but its value doesn't matter as long as it's  $\gg E_{\text{max}}$ . You may not like this construction for one reason or the other, but clearly there is no huge number of details to be specified here.

This works regardless of what the Hamiltonian is, how large the system is, how many detectors there are, or what observable is being measured. It will work for the double-slit<sup>4</sup>, for Stern-Gerlach, and for EPR-type experiments, no matter how the settings were chosen. In addition, the dynamical law that we have made up here is both deterministic and local: It's not like the prepared state actually interacts with the detector before measurement, it just has the required information already in the dynamical law. It appears that we have done the seemingly impossible: We have found a local, deterministic, hidden variable model that reproduces quantum mechanics on the average without a hint of conspiracy. How can this possibly be?

The reason this works is that the dynamical law explicitly depends on the measurement settings at the time of measurement. This may appear as if it goes against the arrow of time, but note that one could express the measurement settings at the time of measurement by the settings at the time of preparation, it's just that the relation between the two might be very difficult. Causality is not violated because there is no time-like direction of communication singled out in this setup, there is merely a correlation. The consequence of this, however, is that while the distribution of the hidden variables,  $\lambda_J$ , does not depend on the settings, the distribution of the  $\sigma_J$ , which accounts for the outcomes of the experiment, depends on the initial state of the system, the  $\lambda_S$ , and the detector settings at the time of the measurement. So, the model violates Statistical Independence; it is "superdeterministic" to use the common term. Therefore, one of the assumptions for Bell's theorem is not fulfilled and there is nothing contradictory about this hidden variable theory reproducing quantum mechanics.

Now, for what the economy of this model is concerned, one may debate whether replacing the collapse postulate with random variables is a good trade-off. It certainly is a conceptual advantage, because it gives a well-defined probabilistic interpretation to quantum mechanics. However, if we merely count axioms, our toy model has no advantage over quantum mechanics. This is not surprising: If one believes in "shut up and calculate" and a probabilistic prediction is all one wants, then one can as well stick with quantum mechanics.

But the collapse postulate in quantum mechanics is ambiguous, as emphasized already by Schödinger [22] and later stressed by Bell [23]: The collapse postulate has it that a measurement induces wave-function collapse, but it does not define just what a "measurement" is to begin with.

Let us also recall that re-deriving quantum mechanics was not the reason for discussing this model. The reason for discussing this model was to demonstrate that superdeterminism is a viable completion of quantum mechanics, and so it may open doors to the development of a

<sup>&</sup>lt;sup>4</sup>Double slit experiments are typically described using infinite Hilbert-spaces, which we did not consider here. However, in reality one never actually measures the particle position with infinite precision; one always has a finite grid of positions. Therefore, while it may be mathematically more convenient to use an infinite-dimensional Hilbert-space, the system can in principle be described using a finite Hilbert space.

<sup>&</sup>lt;sup>5</sup>It is not only difficult, it is also unnecessary, as we just saw.

more fundamental theory in which the variables, rather than being random, are determined by a to-be-found law of nature.

Our toy model is local in the sense of respecting "continuity of action" as defined in [20]. It avoids non-local interactions by hard-coding the dependence on the detector settings into the evolution law. This is another reasons why one should not take this model too seriously: A good, fundamental, model should allow us to *derive* that the effective law for the prepared state depends on the detector settings. This requires that the to-be-found fundamental model includes the detectors and the environment and possibly other transformation devices that are part of the experimental setup. This has to be the case because otherwise we would lack information to define what the detector eigenstates are.<sup>6</sup>

# 4 Superluminal Signalling, Absence Of

It has been argued in the literature that superdeterministic models require fine-tuning to prevent that entanglement can be exploited for superluminal signalling [21, 24, 25, 26]. Given that there is no finetuning in our model, we will therefore now have a close look at this issue. Ahead, one may notice that chances are good the toy model proposed here does not allow for superluminal signalling because the master equation (14) is linear in the density matrix. The model hence does not suffer from the problem discussed in [27, 28]<sup>7</sup>. However, we wish to study this more explicitly using the common example of spin entanglement.

We will concretely look at a singlet state, shared between two – potentially very distant – observers, Alice (A) and Bob (B), who measure the spin of a particle, which is either up ( $\uparrow$ ) or down ( $\downarrow$ ) in their basis. We will chose the weights of the entangled state to be q and  $\sqrt{1-q^2}$ , respectively, (instead of using  $1/\sqrt{2}$  for both) to make sure cancellations between terms do not occur because of the symmetry of the state.

The total dimension of the Hilbert space is  $N = 2^2 = 4$ . To connect the notation of the previous section with the common notation, we define:

$$|1\rangle := |\downarrow_A\rangle \otimes |\uparrow_B\rangle \equiv |\downarrow_A\uparrow_B\rangle \quad , \quad |2\rangle := |\uparrow_A\rangle \otimes |\downarrow_B\rangle \equiv |\uparrow_A\downarrow_B\rangle ,$$
  
$$|3\rangle := |\downarrow_A\rangle \otimes |\downarrow_B\rangle \equiv |\downarrow_A\downarrow_B\rangle \quad , \quad |4\rangle := |\uparrow_A\rangle \otimes |\uparrow_B\rangle = |\uparrow_A\uparrow_B\rangle . \tag{22}$$

The prepared state at time  $t = t_p$  is

$$|\Psi(t_{\rm p})\rangle = q|\uparrow_A\downarrow_B\rangle - \sqrt{1 - q^2}|\downarrow_A\uparrow_B\rangle = q|1\rangle - \sqrt{1 - q^2}|2\rangle , \qquad (23)$$

and has the density matrix

$$\rho(t_{\rm p}) = q^2 |1\rangle\langle 1| + (1 - q^2)|2\rangle\langle 2| - q\sqrt{1 - q^2} (|1\rangle\langle 2| + |2\rangle\langle 1|) . \tag{24}$$

<sup>&</sup>lt;sup>6</sup>Whatever the underlying model, it most likely will have a future-input dependence [20]. For more about this, see [2].

<sup>&</sup>lt;sup>7</sup>It should be mentioned that [27, 28] were concerned with a different question than the one investigated here, namely whether non-linear generalizations of the Schrödinger-equation allow for superluminal signalling. Their answer is "Yes" for deterministic evolutions and "in some cases" for stochastic dynamics. But the equation we use here is neither non-linear nor is it a Schrödinger-equation.

Our task is now to solve the master-equation (14) for the density matrix with the initial value (24) and then take the average over the hidden variables to get

$$\overline{\rho}(t) := \int \rho(t) \prod_{I=2}^{N} d\lambda_{I} . \tag{25}$$

For a 4-dimensional Hilbert-space, we have 6 Lindblad-operators, one for each entry above the diagonal. For the singlet state under consideration,  $\alpha_1^2 = q^2$ ,  $\alpha_2 = 1 - q^2$ ,  $\alpha_3^2 = \alpha_4^2 = 0$ , so we have

$$\sigma_2 = |\lambda_2|^2 - q^2, \ \sigma_3 = |\lambda_3|^2 - 1, \ \sigma_4 = |\lambda_4|^2 - 1.$$
 (26)

Since the  $\lambda$ 's all lie within the unit circle, we see that  $\sigma_3$  and  $\sigma_4$  are both  $\leq 0$ . This reflects the fact that the probability to end up in the state  $|3\rangle$  or  $|4\rangle$  should be zero.

One can then integrate the master-equation (see Appendix), using the initial state (24), from which one obtains

$$\rho_{11}(t) = \theta(\sigma_{2})q^{2}e^{-\kappa t} + \theta(-\sigma_{2})\left(1 - \left(1 - q^{2}\right)e^{-\kappa t}\right), 
\rho_{22}(t) = \theta(\sigma_{2})(1 - q^{2}e^{-\kappa t}) + \theta(-\sigma_{2})\left(1 - q^{2}\right)e^{-\kappa t}, 
\rho_{12}(t) = \rho_{22}(t) = q\sqrt{1 - q^{2}}e^{-\kappa t/2}.$$
(27)

It has the property that  $Tr(\rho) = 1$  for all times.

We already know that

$$P(\sigma_2 < 0) = q^2, P(\sigma_2 > 0) = 1 - q^2.$$
 (28)

So for the average of the density matrix we have

$$\begin{split} \overline{\rho}_{11}(t) &= (1-q^2)q^2e^{-\kappa t} + q^2\left(1 - \left(1 - q^2\right)e^{-\kappa t}\right) = q^2 \;, \\ \overline{\rho}_{22}(t) &= (1-q^2)(1-q^2e^{-\kappa t}) + q^2\left(1 - q^2\right)e^{-\kappa t} = 1 - q^2 \;, \\ \overline{\rho}_{12}(t) &= \overline{\rho}_{22}(t) &= q\sqrt{1 - q^2}e^{-\kappa t/2} \;. \end{split} \tag{29}$$

This means the only thing the Lindblad-operators do is to exponentially suppress the off-diagonal terms in the density matrix. We can now take the partial traces

$$\overline{\rho}_A = \operatorname{Tr}_B(\overline{\rho}) = q^2 |\uparrow_A\rangle \langle \uparrow_A | + (1 - q^2) |\downarrow_A\rangle \langle \downarrow_A |, \qquad (30)$$

$$\overline{\rho}_{B} = \operatorname{Tr}_{A}(\overline{\rho}) = q^{2} |\downarrow_{B}\rangle\langle\downarrow_{B}| + (1 - q^{2}) |\uparrow_{B}\rangle\langle\uparrow_{B}|. \tag{31}$$

This is exactly the same result as in quantum mechanics. We thus see that if one treats the random variables in the above discussed toy model as truly random then superluminal signalling is not possible.

However, one may object that if the hidden variables are really random, then the model is not superdeterministic because it is not deterministic in the first place. Indeed, in a superdeterministic model, the randomness should not be fundamental but merely a consequence of lacking

detailed information – as one expects in a 'hidden variables' theory. Thus, the distribution of the hidden variables should in certain circumstances deviate from being completely uniform. In this case, taking the average over the hidden variables will not give the same result as in quantum mechanics. This is apparent from Eqs. (27) which, when averaged with probabilities other than (28) will not give (29), but have correction terms in whatever is the deviation of the distribution of the  $\lambda$ 's from uniformity.

This per se does not constitute an opportunity for signalling, because to exploit this channel one of the observers would have to be able to skew the distribution of hidden variables so that the measurement outcome for the other observer would be correlated with information they desire to send. What it takes to do that is, again, a question this toy model cannot address because it does not specify the origin of the randomness. However, that so far no experiment has revealed any deviations from Born's rule suggests it is not easy to prepare a system in a state where the distribution of hidden variables does not reproduce quantum mechanics.

#### 5 Conclusion

This toy model demonstrates superdeterminism is a scientifically sound option for resolving the quantum measurement problem. We have shown that fine-tuning, which has frequently been used as an argument against superdetermism, it unnecessary: To make predictions with this model one does not need to introduce a huge number of delicately chosen parameters.

The model, when averaged over the uniformly distributed hidden variables,  $\lambda$ , reproduces the quantum mechanical predictions. If the variables were not uniformly distributed, for example because the sample space is not large enough, this model would give predictions that differ from quantum mechanics. That is, even in an experiment where one expects to have a good enough sampling to reproduce Born's rule with only small statistical deviations, the observed distribution could still be heavily skewed in the space of hidden variables, hence not approximate Born's rule as expected. This makes clear that superdeterminism is not an interpretation of quantum mechanics; it is a concrete possibility for replacing quantum mechanics with a better theory and more effort should be made to experimentally test it.

For example, following [29], consider the case in which the values of the hidden variables are determined by the degrees of freedom of the measurement device (other than the measurement setting itself). Then, if we perform multiple measurements of non-commuting observables using the same setup, but the degrees of freedoms of the device which determine the  $\lambda s$  only change very slowly between the measurements, we might end up sampling always the same region of the parameter space of the  $\lambda s$ , hence observing results which are more strongly correlated than quantum mechanics predicts.

Exactly when the deviations from quantum mechanics become non-negligible depends on what the hidden variables are; the above introduced toy model cannot answer this question. The toy model is therefore, strictly speaking, untestable, because it does not specify where the distribution of hidden variables comes from. But, as pointed out above and in more detail in [29, 1], if the hidden variables are the degrees of freedom of the detector, it is reasonable to

expect that minimizing the variation in the detectors' degrees of freedom between consecutive measurements will reveal deviations from Born's rule which cannot be detected by Bell-type experiments [10, 11, 12], no matter how ingeniously conceived and executed.

### Acknowledgements

We gratefully acknowledge support from the Fetzer Franklin Fund. SH thanks Tim Palmer for helpful discussion.

# **Appendix: General Solution for** N = 4

As previously, we work in the limit  $\kappa \gg E_{\text{max}}$  and therefore set H = 0. We begin with rewriting Eq. (14) more explicitly inserting Eq. (13):

$$\partial_{t}\rho(t) = \kappa \sum_{M>K=1}^{4} \left( L_{MK}\rho(t)L_{MK}^{\dagger} - \frac{1}{2} \{\rho(t), L_{MK}^{\dagger}L_{MK}\} \right) 
= \kappa \sum_{M>K=1}^{4} \left[ \theta(\sigma_{M}) \left( |M\rangle\langle K|\rho(t)|K\rangle\langle M| - \frac{1}{2} \{\rho(t), |K\rangle\langle K|\} \right) 
+ \theta(-\sigma_{M}) \left( |K\rangle\langle M|\rho(t)|M\rangle\langle K| - \frac{1}{2} \{\rho(t), |M\rangle\langle M|\} \right) \right].$$
(32)

We now introduce the notation  $\rho_{IJ}(t) := \langle I | \rho(t) | J \rangle$  by use of which the master-equation takes the form

$$\partial_{t}\rho_{IJ}(t) = \kappa \sum_{M>K=1}^{4} \left[ \theta(\sigma_{M}) \left( \delta_{IM} \delta_{MJ} \rho_{KK}(t) - \frac{1}{2} \delta_{KJ} \rho_{IK}(t) - \frac{1}{2} \delta_{IK} \rho_{KJ}(t) \right) + \theta(-\sigma_{M}) \left( \delta_{IK} \delta_{KJ} \rho_{MM}(t) - \frac{1}{2} \rho_{IM}(t) \delta_{MJ} - \frac{1}{2} \delta_{IM} \rho_{MJ}(t) \right) \right].$$
(33)

To apply the Kronecker delta, we recall that the double sum can be equivalently written as:

$$\sum_{M>K}^{4} = \sum_{M=2}^{4} \sum_{K=1}^{M-1} = \sum_{K=1}^{3} \sum_{M=K+1}^{4} . \tag{34}$$

Using this, we obtain for I = J

$$\partial_{t}\rho_{JJ}(t)/\kappa = \sum_{M=2}^{4} \sum_{K=1}^{M-1} \theta(\sigma_{M}) \delta_{JM} \delta_{MJ} \rho_{KK}(t) - \frac{1}{2} \sum_{K=1}^{3} \sum_{M=K+1}^{4} \theta(\sigma_{M}) \delta_{KJ} \rho_{JK}(t) 
- \frac{1}{2} \sum_{K=1}^{3} \sum_{M=K+1}^{4} \theta(\sigma_{M}) \delta_{JK} \rho_{KJ}(t) + \sum_{K=1}^{3} \sum_{M=K+1}^{4} \theta(-\sigma_{M}) \delta_{JK} \delta_{KJ} \rho_{MM}(t) 
- \frac{1}{2} \sum_{M=2}^{4} \sum_{K=1}^{M-1} \theta(-\sigma_{M}) \rho_{JM}(t) \delta_{MJ} - \frac{1}{2} \sum_{M=2}^{4} \sum_{K=1}^{M-1} \theta(-\sigma_{M}) \delta_{JM} \rho_{MJ}(t) 
= (1 - \delta_{J1}) \theta(\sigma_{J}) \sum_{K=1}^{J-1} \rho_{KK}(t) - \frac{1}{2} (1 - \delta_{J4}) \rho_{JJ}(t) \sum_{M=J+1}^{4} \theta(\sigma_{M}) 
- \frac{1}{2} (1 - \delta_{J4}) \rho_{JJ}(t) \sum_{M=J+1}^{4} \theta(\sigma_{M}) + (1 - \delta_{J4}) \sum_{M=J+1}^{4} \theta(-\sigma_{M}) \rho_{MM}(t) 
- \frac{1}{2} (1 - \delta_{J1}) \sum_{K=1}^{J-1} \theta(-\sigma_{J}) \rho_{JJ}(t) - \frac{1}{2} (1 - \delta_{J1}) \sum_{K=1}^{J-1} \theta(-\sigma_{J}) \rho_{JJ}(t) 
= (1 - \delta_{J1}) \left( \theta(\sigma_{J}) \sum_{K=1}^{J-1} \rho_{KK}(t) - \sum_{K=1}^{J-1} \theta(-\sigma_{J}) \rho_{JJ}(t) \right) 
+ (1 - \delta_{J4}) \left( \sum_{M=J+1}^{4} \theta(-\sigma_{M}) \rho_{MM}(t) - \rho_{JJ}(t) \sum_{M=J+1}^{4} \theta(\sigma_{M}) \right) 
= (1 - \delta_{J1}) \left[ \theta(\sigma_{J}) \sum_{K=1}^{J-1} \rho_{KK}(t) - \theta(-\sigma_{J}) \rho_{JJ}(t) (J-1) \right] 
+ (1 - \delta_{J4}) \sum_{M=J+1}^{4} \left[ \theta(-\sigma_{M}) \rho_{MM}(t) - \theta(\sigma_{M}) \rho_{JJ}(t) \right] .$$
(35)

This can be written explicitly as

$$\partial_{t}\rho_{11}(t)/\kappa = \theta(-\sigma_{2})\rho_{22}(t) + \theta(-\sigma_{3})\rho_{33}(t) + \theta(-\sigma_{4})\rho_{44}(t) 
- \sum_{M=2}^{4} \theta(\sigma_{M})\rho_{11}(t) ,$$
(36)
$$\partial_{t}\rho_{22}(t)/\kappa = \theta(\sigma_{2})\rho_{11}(t) - (\theta(-\sigma_{2}) + \theta(\sigma_{3}) + \theta(\sigma_{4}))\rho_{22}(t) 
+ \theta(-\sigma_{3})\rho_{33}(t) + \theta(-\sigma_{4})\rho_{44}(t) ,$$
(37)
$$\partial_{t}\rho_{33}(t)/\kappa = \theta(\sigma_{3})\rho_{11}(t) + \theta(\sigma_{3})\rho_{22}(t) 
- (2\theta(-\sigma_{3}) + \theta(\sigma_{4}))\rho_{33}(t) + \theta(-\sigma_{4})\rho_{44}(t) ,$$
(38)
$$\partial_{t}\rho_{44}(t)/\kappa = \theta(\sigma_{4})\rho_{11}(t) + \theta(\sigma_{4})\rho_{22}(t) + \theta(\sigma_{4})\rho_{33}(t) - 3\theta(-\sigma_{4})\rho_{44}(t) .$$
(39)

For  $I \neq J$  Eq. (33) becomes:

$$\partial_{t}\rho_{IJ}(t)/\kappa = \sum_{K=1}^{3} \sum_{M=K+1}^{4} \theta(\sigma_{M}) \left( -\frac{1}{2} \delta_{KJ}\rho_{IK}(t) - \frac{1}{2} \delta_{IK}\rho_{KJ}(t) \right) 
+ \sum_{M=2}^{4} \sum_{K=1}^{M-1} \theta(-\sigma_{M}) \left( -\frac{1}{2} \rho_{IM}(t) \delta_{MJ} - \frac{1}{2} \delta_{IM}\rho_{MJ}(t) \right) 
= -\frac{1}{2} \left( \sum_{M=J+1}^{4} \theta(\sigma_{M})(1 - \delta_{J4})\rho_{IJ}(t) + \sum_{M=I+1}^{4} (1 - \delta_{I4})\theta(\sigma_{M})\rho_{IJ}(t) \right) 
- \frac{1}{2} \left( (1 - \delta_{J1}) \sum_{K=1}^{J-1} \theta(-\sigma_{J})\rho_{IJ}(t) + (1 - \delta_{I1}) \sum_{K=1}^{I-1} \theta(-\sigma_{I})\rho_{IJ}(t) \right) 
= -\frac{1}{2} \left[ \left( (1 - \delta_{J4}) \sum_{M=J+1}^{4} \theta(\sigma_{M}) + (1 - \delta_{I4}) \sum_{M=I+1}^{4} \theta(\sigma_{M}) \right) 
+ \left( (1 - \delta_{J1}) \sum_{K=1}^{J-1} \theta(-\sigma_{J}) + (1 - \delta_{I1}) \sum_{K=1}^{I-1} \theta(-\sigma_{I}) \right) \right] \rho_{IJ}(t) 
= -\frac{1}{2} \left[ \left( (1 - \delta_{J4}) \sum_{M=J+1}^{4} \theta(\sigma_{M}) + (1 - \delta_{I4}) \sum_{M=I+1}^{4} \theta(\sigma_{M}) \right) 
+ \left( (1 - \delta_{J1})(J - 1)\theta(-\sigma_{J}) + (1 - \delta_{I1})(I - 1)\theta(-\sigma_{J}) \right) \right] \rho_{IJ}(t) . \tag{40}$$

With that, we can write for  $I \neq J$ 

$$\partial_t \rho_{IJ}(t) = -\frac{\kappa}{2} \Lambda_{IJ} \rho_{IJ}(t) , \qquad (41)$$

where

$$\Lambda_{IJ} := \left( (1 - \delta_{J4}) \sum_{M=J+1}^{4} \theta(\sigma_{M}) + (1 - \delta_{I,4}) \sum_{M=I+1}^{4} \theta(\sigma_{M}) \right) 
+ \left( (1 - \delta_{J1})(J - 1)\theta(-\sigma_{J}) + (1 - \delta_{I1})(I - 1)\theta(-\sigma_{I}) \right) .$$
(42)

Eq. (41) can be integrated straight-forwardly to

$$\rho_{IJ}(t) = \rho_{IJ}(0) e^{-\frac{\kappa}{2}\Lambda_{IJ}t}.$$
(43)

For the singlet state we always have  $\sigma_{3.4} \le 0$  so Eqs. (36) - (39) simplify to

$$\partial_t \rho_{11}(t) = \kappa [\theta(-\sigma_2)\rho_{22}(t) + \rho_{33}(t) + \rho_{44}(t) - \theta(\sigma_2)\rho_{11}(t)], \qquad (44)$$

$$\partial_t \rho_{22}(t) = \kappa [\theta(\sigma_2)\rho_{11}(t) - \theta(-\sigma_2)\rho_{22}(t) + \rho_{33}(t) + \rho_{44}(t)],$$
 (45)

$$\partial_t \rho_{33}(t) = -2\kappa \rho_{33}(t) + \kappa \rho_{44}(t) ,$$
 (46)

$$\partial_t \rho_{44}(t) = -3\kappa \rho_{44}(t) . \tag{47}$$

We solve these equations starting from the last one

$$\rho_{44}(t) = \rho_{44}(0)e^{-3\kappa t} . (48)$$

This can be inserted into the third equation and gives

$$\rho_{33}(t) = -\rho_{44}(0)e^{-3\kappa t} + (\rho_{33}(0) + \rho_{44}(0))e^{-2\kappa t}. \tag{49}$$

Inserting the initial state (24) now reveals that the solution to Eqs. (48) and (49) is simply  $\rho_{33}(t) = \rho_{44}(t) = 0$ .

Then the first two equations are

$$\partial_t \rho_{11}(t) = \kappa \theta(-\sigma_2) \rho_{22}(t) - \kappa \theta(\sigma_2) \rho_{11}(t) , \qquad (50)$$

$$\partial_t \rho_{22}(t) = \kappa \theta(\sigma_2) \rho_{11}(t) - \kappa \theta(-\sigma_2) \rho_{22}(t) , \qquad (51)$$

which can be integrated to

$$\rho_{11}(t) = \theta(\sigma_2)\rho_{11}(0)e^{-\kappa t} + \theta(-\sigma_2)\left(\rho_{11}(0) + \rho_{22}(0)\left(1 - e^{-\kappa t}\right)\right), \tag{52}$$

$$\rho_{22}(t) = \theta(\sigma_2) \left( \rho_{22}(0) + \rho_{11}(0) \left( 1 - e^{-\kappa t} \right) \right) + \theta(-\sigma_2) \rho_{22}(0) e^{-\kappa t} . \tag{53}$$

After inserting the initial state  $\rho_{11}(0) = q^2$  and  $\rho_{22}(0) = 1 - q^2$  we get for the diagonal elements

$$\rho_{11}(t) = \theta(\sigma_2)q^2e^{-\kappa t} + \theta(-\sigma_2)\left[1 - (1 - q^2)e^{-\kappa t}\right], \tag{54}$$

$$\rho_{22}(t) = \theta(\sigma_2) \left( 1 - q^2 e^{-\kappa t} \right) + \theta(-\sigma_2) \left( 1 - q^2 \right) e^{-\kappa t} . \tag{55}$$

For the off-diagonal elements of the density matrix, we note that all contributions to  $\Lambda_{IJ}$  for I, J equal to 3 or 4 vanish and we only have one non-zero matrix element which is

$$\Lambda_{12} = \theta(\sigma_2) + \theta(-\sigma_2) = 1. \tag{56}$$

then

$$\rho_{12}(t) = -q\sqrt{1 - q^2} e^{-\frac{\kappa}{2}t} . \tag{57}$$

Eqs. (54), (55), and (57) are used in (27).

#### References

- [1] S. Hossenfelder, T. Palmer, "Rethinking Superdeterminism," Front. Phys. 8, 139 (2020) [arXiv:1912.06462 [quant-ph]].
- [2] S. Hossenfelder, "Superdeterminism: A Guide for the Perplexed", arXiv:
- [3] C. H. Brans, "Bell's theorem does not eliminate fully causal hidden variables," International Journal of Theoretical Physics 27, 219 (1988).

- [4] T. N. Palmer, "A local deterministic model of quantum spin measurement" Proc. Roy. Soc. A, 451, 585-608 (1995) [arXiv:quant-ph/9505025].
- [5] J. Degorre, S. Laplante, J. Roland, "Simulating quantum correlations as a distributed sampling problem, Phys. Rev. A 72, 062314 (2005) [arXiv:quant-ph/0507120].
- [6] T. N. Palmer, "The Invariant Set Postulate: a new geometric framework for the foundations of quantum theory and the role played by gravity." Proc. Roy. Soc. A, 465 (2009) [arXiv:0812.1148 [quant-ph]].
- [7] M. J. W. Hall, "Local deterministic model of singlet state correlations based on relaxing measurement independence, Phys. Rev. Lett. 105:250404 (2010) [arXiv:1007.5518 [quant-ph]].
- [8] G. S. Ciepielewski, E. Okon, D. Sudarsky, "On Superdeterministic Rejections of Settings Independence," arXiv:2008.00631 [quant-ph].
- [9] M. J. W. Hall, "The significance of measurement independence for Bell inequalities and locality," in At the Frontier of Spacetime, 189-204, (2016) [arXiv:1511.00729 [quant-ph]].
- [10] C. Leung, A. Brown, H. Nguyen, A. S. Friedman, D. I. Kaiser, J. Gallicchio, "Astronomical random numbers for quantum foundations experiments," Phys. Rev. A **97**, no. 4, 042120 (2018) [arXiv:1706.02276 [quant-ph]].
- [11] The BIG Bell Test Collaboration, "Challenging local realism with human choices," Nature, 557, 212-216 (2018) [arXiv:1805.04431 [quant-ph]].
- [12] A. S. Friedman, A. H. Guth, M. J. W. Hall, D. I. Kaiser, J. Gallicchio, "Relaxed Bell Inequalities with Arbitrary Measurement Dependence for Each Observer," Phys. Rev. A **99**, no. 1, 012121 (2019) [arXiv:1809.01307 [quant-ph]].
- [13] G. Lindblad, "On the generators of quantum dynamical semigroups," Commun. Math. Phys. 119, 48 (1976).
- [14] V. Gorini, A. Kossakowski, E. C. Sudarsahan, "Completely positive semigroups of n-level systems," J. Math. Phys. 17, 821 (1976).
- [15] M. A. Nielsen, I. L. Chuang, "Quantum Computation and Quantum Information," Cambridge University Press (2010).
- [16] A. Bassi and G. C. Ghirardi, "Dynamical reduction models," Phys. Rept. **379**, 257 (2003) [arXiv:quant-ph/0302164 [quant-ph]].
- [17] S.L. Adler, D. C. Brody, T. A. Brun, L. P. Hughston, "Martingale Models for Quantum State Reduction," J. Phys. A 34, 8795-8820 (2001) [arXiv:quant-ph/0107153].
- [18] S. Hossenfelder, "Born's rule from almost nothing," [arXiv:2006.14175 [quant-ph]].
- [19] W. H. Zurek, "Pointer basis of quantum apparatus: Into what mixture does the wave packet collapse?". Phys. Rev. D **24.6**, 1516 (1981).
- [20] K. B. Wharton, N. Argaman, "Bell's Theorem and Spacetime-Based Reformulations of Quantum Mechanics,", arXiv:1906.04313 [quant-ph].

- [21] C. J. Wood, R. W. Spekkens, "The lesson of causal discovery algorithms for quantum correlations: Causal explanations of Bell-inequality violations require fine-tuning, New J. Phys. 17, 033002 (2015) [arXiv:1208.4119 [quant-ph]].
- [22] E. Schrödinger, *Die gegenwärtige Situation in der Quantenmechanik*, Naturwissenschaften **23**, 823–828 (1935).
- [23] J. S. Bell, Speakable and unspeakable in quantum mechanics: Collected papers on quantum philosophy, Cambridge university press (2004).
- [24] A. Bendersky, G. Senno, G. de la Torre, S. Figueira, A. Acin, "Non-signaling deterministic models for non-local correlations have to be uncomputable," PRL 118, 130401 (2017), [arXiv:1606.08403 [quant-ph]].
- [25] I. Sen, A. Valentini, "Superdeterministic hidden-variables models I: nonequilibrium and signalling" arXiv:2003.11989 [quant-ph]
- [26] I. Sen, A. Valentini, "Superdeterministic hidden-variables models II: conspiracy" arXiv:2003.12195 [quant-ph].
- [27] N. Gisin, "Stochastic quantum dynamics and relativity," Helv. Phys. Acta 62, 363-371 (1989).
- [28] A. Bassi and K. Hejazi, "No-faster-than-light-signaling implies linear evolution. A re-derivation," Eur. J. Phys. **36**, no.5, 055027 (2015) [arXiv:1411.1768 [quant-ph]].
- [29] S. Hossenfelder, "Testing super-deterministic hidden variables theories," Found. Phys. **41**, 1521 (2011) [arXiv:1105.4326 [quant-ph]].