

Period change of binary pulsars due to the accretion of dark matter particles

Ebrahim Hassani ^{1*}, Hossein Ebadi ², Reza Pazhouhesh ³

Abstract

As binary systems move inside galaxies, they interact with the dark matter halo. This interaction leads to the accretion of dark matter particles inside binary components. The accretion of dark matter particles increases the mass of the binary components and then, the total mass of the system. Increased mass by this way can affect other physical parameters of the systems, like orbital periods of the systems. In this work, we estimated the period change of some known compact binary systems due to the accretion of dark matter particles inside them. Our conclusion is that, for WIMP particles with masses in the range $\simeq 10 \text{ GeV} \cdot c^{-2}$ and dark matter density as high as the dark matter density around the sun, period change due to accretion of dark matter particles is negligible compared to the period change due to the emission of gravitational waves from the systems.

Keywords

Dark Matter, Binary Pulsars, White dwarfs, Neutron stars

¹ Department of Physics, University of Birjand, Iran

*Corresponding author: ebrahim.hassani@birjand.ac.ir

² Department of Theoretical Physics and Astrophysics,
University of Tabriz, PO Box 51664, Tabriz, Iran, e-mail: hosseinebadi@tabrizu.ac.ir

³ Department of Physics, Faculty of Sciences,
University of Birjand, Birjand, Iran, e-mail: rpazhouhesh@birjand.ac.ir

Contents

1	Introduction	2
2	Capture rate by white dwarfs	2
3	Capture rate by neutron stars	4
4	Capture rate by compact binary systems	4
5	Results and Conclusions	5
	Acknowledgments	7

1. Introduction

Rotation curves of galaxies reveal the non-uniform distribution of dark matter (DM) inside galaxies.¹ From this point, one can infer that all stars that are distributed inside galaxies are immersed inside DM. Then, it is logical to suppose that the DM must affect structure and evolutionary courses of stars. For the first time Steigman used DM supposition to solve the solar neutrino problem². Simulation of dwarf galaxies also supports the interaction between DM and stars³. In addition, DM effect on stars can be used to solve the paradox of youth problem for stars that are located near the Galactic massive black hole⁴. In addition to normal stars, DM effect on other celestial bodies like the moon^{5,6}, planets⁷, neutron stars (NS)^{8–35}, white dwarf stars (WD),^{36–44} black holes^{8,16,45,46} and binary star systems⁴⁷ are also investigated in the literature.

For the first time, Press and Spergel estimated the rate at which DM particles (WIMP particles, exactly speaking) accrete into the sun⁴⁸. In the next step, Gould obtained a general relation for capture rate (CR) of DM particles by other round bodies like planets and stars⁴⁹. Kouvaris used Press and Spergel relation to derive CR relation for NSs¹³ (see section 3 for more details). Hurst et al used Gould relation to obtain CR relation for WDs⁴³ (see section 2 for more details).

As a compact binary system moves inside the DM halo, it feels dynamical friction by the DM halo. Induced dynamical friction by the DM halo can alter the orbital parameters (e.g. period and semi-major axis) of these systems. In a series of studies, researchers tried to estimate the period change of pulsar binary systems due to this dynamical friction and compare the results with the period change due to the gravitational wave emission.^{50–58}

In this work we estimated the period change of compact binary systems due to the accretion of DM particles (instead of due to the dynamical friction) inside them. Accretion of DM particles inside compact binary systems increases the mass of the binary components. Then, increased mass leads to the change in the orbital periods of these system. To the best of my knowledge, there is not any previous study that estimates the period change of binary systems due to the accretion of DM particles.

The rest of this paper is structured as follows. In sections 2 and 3 we presented CR relations for WDs and NSs, separately. In section 4 we presented an approximate relation for the CR by compact binary systems. Finally, section 5 devoted to the results and discussions.

2. Capture rate by white dwarfs

Capture Rate of DM particles by white dwarfs can be calculated using the equation⁴³:

$$CR_{WD} = \sum_i CR_{WD,i} = \sqrt{\frac{3}{2}} \frac{\rho_x}{m_x} \sigma_{\chi,i} \frac{v_{esc,RWD}}{\bar{v}} N_i \langle \hat{\phi} \rangle \frac{erf(\eta)}{\eta} \quad (1)$$

like equation 8 for neutron stars, ρ_x , m_x and \bar{v} are the density of DM halo around white dwarf (or around compact binary system), mass of WIMP particles and velocity dispersion of WIMPs (which we supposed

to be $\bar{v} = 220 \text{ km} \cdot \text{sec}^{-1}$ for all stars in this study), respectively. R_{WD} is the radius of the white dwarf and $v_{esc, R_{WD}}$ is the escape velocity at the surface of the white dwarf. η is the ratio of star's velocity through its halo to the local velocity dispersion of the DM halo.

$\sigma_{\chi, i}$ is the scattering cross section of WIMP particles from nuclear species i . As WDs consumed almost all of their hydrogen then, we just consider the spin-independent part of the scattering cross section and neglect the spin-dependent part. So, the spin-independent scattering cross section for all elements except hydrogen is^{43, 59, 60} :

$$\sigma_{\chi, i} = \sigma_{\chi, SI} A_i^2 \left(\frac{m_{\chi} m_{n, i}}{m_{\chi} + m_{n, i}} \right)^2 \left(\frac{m_{\chi} + m_{proton}}{m_{\chi} m_{proton}} \right)^2 \quad (2)$$

where $\sigma_{\chi, SI}$ is the spin-independent scattering cross-section which is determined through experimental direct DM detection methods. In this study we considered it to be $\sigma_{\chi, SI} = 10^{-44} \text{ cm}^2$ (as mentioned again in the section 3 too). A_i is atomic number of species i , $m_{n, i}$ in nuclear mass of species i and m_{proton} is the mass of a proton. To simplify calculations, we supposed that WDs composed of just carbon atoms with $m_{n, c} = 12 u$ where u is the atomic mass unit (amu) and atomic number $A_c = 6$. The similar supposition had done in the similar work⁴³.

In equation 1, $\hat{\phi}$ is dimensionless potential for stellar nucleons and defined as⁴³ :

$$\hat{\phi} = \frac{v_{esc}^2(r)}{v_{esc}^2(R)} \quad (3)$$

The quantity $\langle \hat{\phi} \rangle$ is the average of $\hat{\phi}$ over all nucleons in the star. To calculate $\langle \hat{\phi} \rangle$, we supposed that, like the case for NSs, average density of all WDs is constant. So, for a typical WD (e.g. Sirius B WD star with $M_{SiriusB} = 1.018 M_{\odot}$ mass⁶¹ and radius⁶² $M_{SiriusB} = 0.0084 R_{\odot}$) we have:

$$\bar{\rho}_{WD} = \frac{M_{WD}}{(4/3)\pi R_{WD}^3} = \frac{1.018 M_{\odot}}{(4/3)\pi (0.0084 R_{\odot})^3} = 2.38 \times 10^9 \left(\frac{Kg}{m^3} \right) \quad (4)$$

So, the approximate mass-radius relation for white dwarfs becomes :

$$R_{WD} = \left(\frac{M_{WD}}{(4/3)\pi \bar{\rho}_{WD}} \right)^{\frac{1}{3}} = 4.65 \times 10^{-4} \times M_{WD}^{1/3} \text{ (m)}. \quad (5)$$

(We need R_{WD} to calculate $v_{esc}(R)$ for WDs with different masses too). Now, $\langle \hat{\phi} \rangle$ becomes:

$$\langle \hat{\phi} \rangle = \frac{1}{R} \int_0^R \frac{v_{esc}^2(r)}{v_{esc}^2(R)} dr = \frac{1}{3} \quad (6)$$

In equation 1, N_i is the total number of nucleons of species i in the WD star. In the case of carbon WDs it becomes:

$$N_c = \frac{M_{WD}}{m_c} = \frac{M_{WD}}{12 u} \quad (7)$$

3. Capture rate by neutron stars

Capture Rate by Neutron stars, considering the general relativity corrections, can be calculated using the below equation¹³ :

$$CR_{NS} = 4\pi^2 \frac{\rho_x}{m_x} \left(\frac{3}{2\pi\bar{v}^2} \right)^{3/2} \frac{(2GM_{NS}R_{NS})}{1 - \frac{2GM_{NS}}{c^2 R_{NS}}} \times \min\left(\frac{1}{3}\bar{v}^2, E_0\right) \times f \quad (8)$$

where ρ_x is the density of the DM halo around the neutron star, m_x is the mass of the WIMP particles, \bar{v} is the the velocity dispersion of the DM particles (in this study we took it to be $\bar{v} = 220 \text{ km} \cdot \text{sec}^{-1}$ for all stars in the study), G is the gravitational constant, c is the speed of light, M_{NS} is the mass of the neutron star and R_{NS} is the radius of the neutron star.

To estimate the radius of a neutron stars R_{NS} (to use in equation 8), we supposed that the average density of the all neutron stars is constant. So, the average density of a typical neutron star with $M_{NS} = 1.44 M_\odot$ and radius $R_{NS} = 10.6 \text{ km}$ (see table 1 of the paper⁶³ for these quantities) is :

$$\bar{\rho}_{NS} = \frac{M_{NS}}{(4/3)\pi R_{NS}^3} = \frac{1.44 M_\odot}{(4/3)\pi (10.6 \times 10^3)^3} = 5.74 \times 10^{17} \left(\frac{\text{Kg}}{\text{m}^3} \right) \quad (9)$$

So, we obtain an approximate mass-radius relation for neutron stars:

$$R_{NS} = \left(\frac{M_{NS}}{(4/3)\pi \bar{\rho}_{NS}} \right)^{1/3} = 7.46 \times 10^{-7} \times M_{NS}^{1/3} \text{ (m)}. \quad (10)$$

In equation 8, E_0 is a constant that parametrizes the maximum kinetic energy per mass of the WIMPs at an asymptotically large distance from the star in order for the WIMPS to be captured by the star. In the case of neutron stars and WIMPs with masses of the order GeV, $E_0 \gg (1/3)\bar{v}^2$. Then, in equation 8, $(1/3)\bar{v}^2$ should be taken always as minimum (for more detailed dicussion about this selection see the section 2 of the paper¹³).

Not all the WIPMs that enter a neutron star will be captured by it. In equation 8, f represents the fraction of the WIPMs that, after one or several scattering from neclear matter, lose enough energy to be captured by a typical neutron star. For WIMP-necleon scattering cross sections $\sigma_\chi > 10^{-45} \text{ cm}^2$, $f = 1$ (as in the case of this study) ; otherwise f is given by¹³ :

$$f = 0.45 \frac{\sigma_\chi}{\sigma_{crit}} \quad (11)$$

which $\sigma_{crit} = m_n R^2 / M \simeq 6 \times 10^{-46} \text{ cm}^2$. In this study we supposed $\sigma_\chi = 10^{-44} \text{ cm}^2$ (then, $f = 1$) which is the maximum magnitude that are determined through the experimental DM detection experiment.⁶⁴

4. Capture rate by compact binary systems

According to the Kepler's third law, square of the period in binary systems is proportional to the cube of the semi-major axis of the orbit⁶⁵ :

$$P^2 = \left(\frac{4\pi^2}{GM} \right) a^3 \quad (12)$$

which $M = M_1 + M_2$. If we suppose that all binary systems that are located inside the galaxies immersed inside the DM halos then, it is logical to suppose that the both components of the systems accumulate DM particles while they are orbiting around each other. By this way, by passing the time, the total mass of the systems will increase. In the next, we will estimate how much the increased mass will affect the other parameters of the systems. Taking differential of the both sides of the equation 12 yealds:

$$\begin{aligned}
 2PdP &= \frac{4\pi^2}{G} d\left(\frac{a^3}{M}\right) \\
 &= \frac{4\pi^2}{G} \frac{3a^2 da}{(M_1 + M_2)} - \frac{4\pi^2 a^3}{G} \frac{dM_1}{(M_1 + M_2)^2} - \frac{4\pi^2 a^3}{G} \frac{dM_2}{(M_1 + M_2)^2} \\
 &= \underbrace{\frac{4\pi^2 a^3}{G(M_1 + M_2)}}_{=P^2} \left[\frac{3}{a} da - \frac{dM_1 + dM_2}{M_1 + M_2} \right]
 \end{aligned} \tag{13}$$

which simplifies to :

$$\frac{dP}{P} = \frac{3}{2} \frac{da}{a} - \frac{dM}{2M} = \frac{3}{2} \frac{da}{a} - \frac{dM_1 + dM_2}{2(M_1 + M_2)} \tag{14}$$

multiplying both sides of the equation 14 to $1/dt$ yealds :

$$\frac{\dot{P}}{P} = \frac{3}{2} \frac{\dot{a}}{a} - \frac{1}{2} \frac{\dot{M}}{M} = \frac{3}{2} \frac{\dot{a}}{a} - \frac{1}{2} \frac{\dot{M}_1 + \dot{M}_2}{(M_1 + M_2)} \tag{15}$$

where $\dot{P} = dp/dt$, $\dot{a} = da/dt$, $\dot{M} = dM/dt$, $\dot{M}_1 = dM_1/dt$ and $\dot{M}_2 = dM_2/dt$. \dot{M}_1 and \dot{M}_2 signifies the accretion of DM particles into compact stars. To calculate \dot{M}_1 and \dot{M}_2 it is enough to multiply equation 8 (in the case of neutron stars) or equation 1 (in the case of white dwarf stars) to the mass of the DM particles m_χ (e.g. $\dot{M}_1 = CR_{NS} \times m_\chi$ if the M_1 star is a neutron star). In table 1 we estimated \dot{M}/M (which $\dot{M} = \dot{M}_1 + \dot{M}_2$) for some known compact binary systems.

5. Results and Conclusions

Using equation 15, it is possible to estimate the accretion of DM particles in compact binary systems. In our calculations, we supposed stars with masses lower than the chandrasekhar mass ($1.4M_\odot$) to be white dwarfs⁷⁴. Stars with masses higher than $1.4M_\odot$ considered to be neutron stars. The mass of the WIMP particles and density of DM halo around binary systems considered to be $m_\chi = 10 \text{ GeV}.c^{-2}$ and $\rho_\chi = 8.16 \times 10^{-3} M_\odot/pc^3$ (which is the estimated DM density around solar neibourhoud⁷), respectively. In equation 15, it is possible to measure \dot{P}/P term through observations. But the precision of the current observational technology is not enough to measure the \dot{a}/a term in the right hand side of the equation. Paying attention to the amounts of \dot{M}/M and \dot{P}/P in table 1, we see that the amounts of \dot{M}/M are about 12-14 orders of magnitude smaller than the amounts of \dot{P}/P (for binaries that at least one of the components is a neutron star). For binaries that both of the components are white dwarfs (e.g. J0737-3039, B1913+16

Table 1. Compact binary system parameters that we used in this study to test our results. All parameters in the table are obtained through observations, except the \dot{M}/M column which is obtained theoretically using the assumptions that are explained in the text.

Name	M_1 (M_\odot)	M_2 (M_\odot)	P_d (days)	\dot{P}^*	\dot{P}/P (sec^{-1})	\dot{M}/M (sec^{-1})	Ref. ^{**}
J1903+0327	1.0	1.7	95	-6.4e-11	-7.8e-18	1.9e-32	66
J0737-3039	1.2	1.3	0.10	-4.0e-15	-4.6e-19	3.5e-45	67
B1913+16	1.4	1.4	0.32	5.0e-15	1.8e-19	3.6e-45	67
J1012+5307	0.16	1.6	0.60	-1.8e-14	-3.5e-19	2.5e-32	68,69
J1614-2230	0.49	1.9	8.7	3.4e-13	4.5e-19	2.6e-32	68
J1909-3744	0.21	1.5	1.5	-4.0e-15	-3.1e-20	2.3e-32	68,69
J0751+1807	0.16	1.6	0.26	-4.6e-14	-2.1e-18	2.5e-32	69
J1910+1256	0.19	1.6	58	-2.0e-11	-4.0e-18	2.5e-32	70
J2016+1948	0.29	1.0	635	-1.0e-9	-1.8e-17	2.7e-45	70
J0348+0432	0.17	2.0	0.10	-1.1e-14	-1.3e-18	3.2e-32	71
J1713+0747	0.29	1.3	68	3.0e-14	5.1e-21	2.9e-45	72
J0613-0200	0.12	1.2	1.2	2.7e-14	2.6e-19	2.5e-45	68,69
J1738+0333	0.19	1.5	0.35	2.0e-15	6.6e-20	2.4e-32	73

* \dot{P} is a dimensionless quantity.

** References for all parameters except \dot{M}/M , which is calculated in this study.

or J2016+1948 systems) the results are more dramatic and the amounts of \dot{M}/M are about 24-26 orders of magnitude smaller than the amounts of \dot{P}/P . Then, we can say that the accretion of DM particles is not the main reason of period change in compact binaries. In order to interpret the period changes \dot{P}/P , other physical sources that affects systems parameters, e.g. gravitaional emission, mass transfer, dynamical friction etc., must be token into account.

As an overall result, our calculations show that the accretion of DM particles in compact binary systems do not change the systems parameters significantly. In addition, accretion of DM particles can not be considered the main source of period changes in binary pulsars.

Acknowledgments

The author would like to express his special thanks to Prof. Joakim Edsjö from the University of Stockholm, Sweden, and Dr. Amin Rezaei Akbarieh from the University of Tabriz, Iran and Prof. Gianfranco Bertone from the University of Amsterdam, Netherlands and Dr. Marco Taoso from the Istituto Nazionale di Fisica Nucleare (INFN), Italy because of their worthwhile discussions during the research.

References

- [1] Y. Sofue and V. Rubin, “Rotation curves of spiral galaxies,” *Annual Review of Astronomy and Astrophysics*, vol. 39, pp. 137–174, sep 2001.
- [2] G. Steigman, H. Quintana, C. L. Sarazin, and J. Faulkner, “Dynamical interactions and astrophysical effects of stable heavy neutrinos,” *The Astronomical Journal*, vol. 83, p. 1050, sep 1978.
- [3] J. I. Read, M. G. Walker, and P. Steger, “Dark matter heats up in dwarf galaxies,” *Monthly Notices of the Royal Astronomical Society*, vol. 484, no. 1, pp. 1401–1420, 2019.
- [4] E. Hassani, R. Pazhouhesh, and H. Ebadi, “The effect of dark matter on stars at the Galactic center: The paradox of youth problem,” *International Journal of Modern Physics D*, vol. 29, p. 2050052, jun 2020.
- [5] M. H. Chan and C. M. Lee, “Constraining the spin-independent elastic scattering cross section of dark matter using the Moon as a detection target and the background neutrino data,” *Physical Review D*, vol. 102, no. 2, p. 23024, 2020.
- [6] R. Garani and P. Tinyakov, “Constraints on dark matter from the Moon,” *Physics Letters, Section B: Nuclear, Elementary Particle and High-Energy Physics*, vol. 804, p. 135403, may 2020.
- [7] D. Hooper and J. H. Steffen, “Dark matter and the habitability of planets,” *Journal of Cosmology and Astroparticle Physics*, vol. 2012, no. 7, p. 46, 2012.

- [8] S. D. McDermott, H. B. Yu, and K. M. Zurek, “Constraints on scalar asymmetric dark matter from black hole formation in neutron stars,” *Physical Review D - Particles, Fields, Gravitation and Cosmology*, vol. 85, p. 23519, jan 2012.
- [9] M. RAZEIRA, A. MESQUITA, C. A. Z. VASCONCELLOS, and R. O. GOMES, “Accretion of Dark Matter in Neutron Stars,” *International Journal of Modern Physics E*, vol. 20, no. supp02, pp. 109–116, 2011.
- [10] M. A. Perez-Garcia, J. Silk, and J. R. Stone, “Dark matter, neutron stars, and strange quark matter,” *Physical Review Letters*, vol. 105, no. 14, p. 141101, 2010.
- [11] A. De Lavallaz and M. Fairbairn, “Neutron stars as dark matter probes,” *Physical Review D - Particles, Fields, Gravitation and Cosmology*, vol. 81, p. 123521, jun 2010.
- [12] G. Bertone and M. Fairbairn, “Compact stars as dark matter probes,” *Physical Review D - Particles, Fields, Gravitation and Cosmology*, vol. 77, p. 43515, feb 2008.
- [13] C. Kouvaris, “WIMP annihilation and cooling of neutron stars,” *Physical Review D - Particles, Fields, Gravitation and Cosmology*, vol. 77, p. 23006, jan 2008.
- [14] N. F. Bell, G. Busoni, S. Robles, and M. Virgato, “Improved treatment of dark matter capture in neutron stars,” *Journal of Cosmology and Astroparticle Physics*, vol. 2020, no. 09, pp. 028–028, 2020.
- [15] A. Joglekar, N. Raj, P. Tanedo, and H.-B. Yu, “Kinetic Heating from Contact Interactions with Relativistic Targets: Electrons Capture Dark Matter in Neutron Stars,” 2020.
- [16] G. L. Lin and Y. H. Lin, “Analysis on the black hole formations inside old neutron stars by isospin-violating dark matter with self-interaction,” *Journal of Cosmology and Astroparticle Physics*, vol. 2020, no. 8, pp. 022–022, 2020.
- [17] A. Herrero, M. A. Pérez-García, J. Silk, and C. Albertus, “Dark matter and bubble nucleation in old neutron stars,” *Physical Review D*, vol. 100, no. 10, p. 103019, 2019.
- [18] N. F. Bell, G. Busoni, and S. Robles, “Capture of leptophilic dark matter in neutron stars,” *Journal of Cosmology and Astroparticle Physics*, vol. 2019, p. 54, jun 2019.
- [19] M. Fabbrichesi and A. Urbano, “Charged neutron stars and observational tests of a dark force weaker than gravity,” *Journal of Cosmology and Astroparticle Physics*, vol. 2020, no. 6, p. 7, 2020.
- [20] N. F. Bell, G. Busoni, and S. Robles, “Heating up neutron stars with inelastic dark matter,” *Journal of Cosmology and Astroparticle Physics*, vol. 2018, no. 9, p. 18, 2018.

- [21] J. Kopp, R. Laha, T. Opferkuch, and W. Shepherd, “Cuckoo’s eggs in neutron stars: can LIGO hear chirps from the dark sector?,” *Journal of High Energy Physics*, vol. 2018, no. 11, p. 96, 2018.
- [22] M. Cermeño, M. Á. Pérez-García, and J. Silk, “Fermionic Light Dark Matter Particles and the New Physics of Neutron Stars,” *Publications of the Astronomical Society of Australia*, vol. 34, p. e043, 2017.
- [23] X. Huang, J.-F. Liu, W.-H. Wang, Q. Cheng, and X.-P. Zheng, “Bondi accretion of dark matter by neutron stars,” *arXiv e-prints*, p. arXiv:1705.08864, 2017.
- [24] M. Cermeño, M. A. Pérez-García, and R. A. Lineros, “Enhanced Neutrino Emissivities in Pseudoscalar-mediated Dark Matter Annihilation in Neutron Stars,” *The Astrophysical Journal*, vol. 863, no. 2, p. 157, 2018.
- [25] M. Baryakhtar, J. Bramante, S. W. Li, T. Linden, and N. Raj, “Dark Kinetic Heating of Neutron Stars and an Infrared Window on WIMPs, SIMPs, and Pure Higgsinos,” *Physical Review Letters*, vol. 119, no. 13, p. 131801, 2017.
- [26] M. A. Perez-Garcia, “Inhomogeneous seeding of quark bubbles in Neutron Stars,” in *Compact Stars in the QCD Phase Diagram IV*, 2015.
- [27] M. Pérez-García and J. Silk, “Dark matter annihilation and local warming in the core of a neutron star,” in *20th International Conference on Particles and Nuclei*, pp. 365–368, 2014.
- [28] H. Zheng, K. J. Sun, and L. W. Chen, “Old neutron stars as probes of isospin-violating dark matter,” *Astrophysical Journal*, vol. 800, no. 2, p. 141, 2015.
- [29] M. Tachibana and M. Ruggieri, “Comment on Dark Matter Capture in Neutron Stars with Exotic Phases,” in *arXiv:1312.5802 [hep-ph]*, 2013.
- [30] M. A. Perez-Garcia, “Deconfinement of quark content in neutron stars due to cosmic external agents,” *arXiv e-prints*, p. arXiv:1306.0148, 2013.
- [31] M. Á. Pérez-García, F. Daigne, and J. Silk, “Short gamma-ray bursts and dark matter seeding in neutron stars,” *Astrophysical Journal*, vol. 768, no. 2, p. 145, 2013.
- [32] N. F. Bell, A. Melatos, and K. Petraki, “Realistic neutron star constraints on bosonic asymmetric dark matter,” *Physical Review D - Particles, Fields, Gravitation and Cosmology*, vol. 87, p. 123507, jun 2013.
- [33] J. Bramante, K. Fukushima, and J. Kumar, “Constraints on bosonic dark matter from observation of old neutron stars,” *Physical Review D - Particles, Fields, Gravitation and Cosmology*, vol. 87, p. 55012, mar 2013.

- [34] A. P. Serebrov, O. M. Zhrebtssov, A. K. Fomin, and M. S. Onegin, “Ultracold-neutron trap as a tool to search for dark matter interacting with long-range forces,” *Physical Review C - Nuclear Physics*, vol. 86, no. 2, p. 25802, 2012.
- [35] M. Á. Pérez-García, J. Silk, and J. R. Stone, “Dark matter seeding in neutron stars,” in *AIP Conference Proceedings* (S. G. Steadman, G. S. Stephans, and F. E. Taylor, eds.), vol. 1441 of *American Institute of Physics Conference Series*, pp. 525–527, 2012.
- [36] K. W. Wong, *Physical Properties of White Dwarf with a Dark Matter Core*. PhD thesis, Hong Kong, Chinese U., 2011.
- [37] B. Dasgupta, A. Gupta, and A. Ray, “Dark matter capture in celestial objects: improved treatment of multiple scattering and updated constraints from white dwarfs,” *Journal of Cosmology and Astroparticle Physics*, vol. 2019, pp. 018–018, aug 2019.
- [38] M. Cermeño and M. A. Pérez-García, “Gamma rays from dark mediators in white dwarfs,” *Physical Review D*, vol. 98, no. 6, p. 63002, 2018.
- [39] P. W. Graham, R. Janish, V. Narayan, S. Rajendran, and P. Riggins, “White dwarfs as dark matter detectors,” *Physical Review D*, vol. 98, no. 11, p. 115027, 2018.
- [40] J. S. Niu, T. Li, W. Zong, H. F. Xue, and Y. Wang, “Probing the dark matter-electron interactions via hydrogen-atmosphere pulsating white dwarfs,” *Physical Review D*, vol. 98, no. 10, p. 103023, 2018.
- [41] J.-S. Niu, W. Zong, and H.-F. Xue, “Constrain the Dark Matter Electron Cross Section from Pulsating White Dwarfs,” *arXiv e-prints*, p. arXiv:1703.10104, 2017.
- [42] P. Amaro-Seoane, J. Casanellas, R. Schödel, E. Davidson, and J. Cuadra, “Probing dark matter crests with white dwarfs and IMBHs,” *Monthly Notices of the Royal Astronomical Society*, vol. 459, no. 1, pp. 695–700, 2016.
- [43] T. J. Hurst, A. R. Zentner, A. Natarajan, and C. Badenes, “Indirect probes of dark matter and globular cluster properties from dark matter annihilation within the coolest white dwarfs,” *Physical Review D - Particles, Fields, Gravitation and Cosmology*, vol. 91, no. 10, p. 103514, 2015.
- [44] S. C. Leung, M. C. Chu, L. M. Lin, and K. W. Wong, “Dark-matter admixed white dwarfs,” *Physical Review D - Particles, Fields, Gravitation and Cosmology*, vol. 87, p. 123506, jun 2013.
- [45] H. Umeda, N. Yoshida, K. Nomoto, S. Tsuruta, M. Sasaki, and T. Ohkubo, “Early black hole formation by accretion of gas and dark matter,” *Journal of Cosmology and Astroparticle Physics*, vol. 2009, no. 8, p. 24, 2009.

- [46] K. M. Belotsky, A. E. Dmitriev, E. A. Esipova, V. A. Gani, A. V. Grobov, M. Y. Khlopov, A. A. Kirillov, S. G. Rubin, and I. V. Svadkovsky, “Signatures of primordial black hole dark matter,” *Modern Physics Letters A*, vol. 29, p. 1440005, dec 2014.
- [47] E. Hassani, H. Ebadi, and R. Pazhouhesh, “Capture rate of weakly interacting massive particles (WIMPs) in binary star systems,” sep 2020.
- [48] W. H. Press and D. N. Spergel, “Capture by the sun of a galactic population of weakly interacting, massive particles,” *The Astrophysical Journal*, vol. 296, p. 679, sep 1985.
- [49] A. Gould, “Resonant enhancements in weakly interacting massive particle capture by the earth,” *The Astrophysical Journal*, vol. 321, p. 571, oct 1987.
- [50] L. G. Gómez and J. A. Rueda, “Dark matter dynamical friction versus gravitational wave emission in the evolution of compact-star binaries,” *Physical Review D*, vol. 96, no. 6, p. 63001, 2017.
- [51] J. Peñarrubia, A. D. Ludlow, J. Chanamé, and M. G. Walker, “Wide binaries in ultrafaint galaxies: A window on to dark matter on the smallest scales,” *Monthly Notices of the Royal Astronomical Society: Letters*, vol. 461, pp. L72–L76, may 2016.
- [52] D. Blas, D. L. Nacir, and S. Sibiryakov, “Secular effects of ultralight dark matter on binary pulsars,” *Physical Review D*, vol. 101, p. 063016, mar 2020.
- [53] L. G. Gómez, “Constraining light fermionic dark matter with binary pulsars,” *Physics of the Dark Universe*, vol. 26, p. 100343, 2019.
- [54] D. Blas, D. L. Nacir, and S. Sibiryakov, “Ultralight Dark Matter Resonates with Binary Pulsars,” *Physical Review Letters*, vol. 118, jun 2017.
- [55] P. Pani, “Binary pulsars as dark-matter probes,” *Physical Review D - Particles, Fields, Gravitation and Cosmology*, vol. 92, no. 12, p. 123530, 2015.
- [56] J. M. Armaleo, D. L. Nacir, and F. R. Urban, “Binary pulsars as probes for spin-2 ultralight dark matter,” *Journal of Cosmology and Astroparticle Physics*, vol. 2020, p. 053, jan 2020.
- [57] A. Caputo, J. Zavala, and D. Blas, “Binary pulsars as probes of a Galactic dark matter disk,” *Physics of the Dark Universe*, vol. 19, pp. 1–11, mar 2018.
- [58] J. Yoo, J. Chaname, and A. Gould, “The End of the MACHO Era: Limits on Halo Dark Matter from Stellar Halo Wide Binaries,” *The Astrophysical Journal*, vol. 601, pp. 311–318, jan 2004.
- [59] G. Jungman, M. Kamionkowski, and K. Griest, “Supersymmetric dark matter,” *Physics Report*, vol. 267, pp. 195–373, mar 1996.

- [60] I. Lopes, J. Casanellas, and D. Eugénio, “The capture of dark matter particles through the evolution of low-mass stars,” *Physical Review D - Particles, Fields, Gravitation and Cosmology*, vol. 83, p. 63521, mar 2011.
- [61] H. E. Bond, G. H. Schaefer, R. L. Gilliland, J. B. Holberg, B. D. Mason, I. W. Lindenblad, M. Seitz-McLeese, W. D. Arnett, P. Demarque, F. Spada, P. A. Young, M. A. Barstow, M. R. Burleigh, and D. Gudehus, “The Sirius System and Its Astrophysical Puzzles: Hubble Space Telescope and Ground-based Astrometry,” *The Astrophysical Journal*, vol. 840, p. 70, may 2017.
- [62] J. B. Holberg, M. A. Barstow, F. C. Bruhweiler, A. M. Cruise, and A. J. Penny, “Sirius B: A New, More Accurate View,” *The Astrophysical Journal*, vol. 497, pp. 935–942, apr 1998.
- [63] T. Güver, A. E. Erkoca, M. H. Reno, and I. Sarcevic, “On the capture of dark matter by neutron stars,” *Journal of Cosmology and Astroparticle Physics*, vol. 2014, pp. 013–013, jan 2014.
- [64] E. Behnke, J. Behnke, S. J. Brice, D. Broemmelsiek, J. I. Collar, P. S. Cooper, M. Crisler, C. E. Dahl, D. Fustin, J. Hall, J. H. Hinnefeld, M. Hu, I. Levine, E. Ramberg, T. Shepherd, A. Sonnenschein, and M. Szydagis, “Improved limits on spin-dependent WIMP-proton interactions from a two liter CF3I bubble chamber,” *Physical Review Letters*, vol. 106, p. 21303, jan 2011.
- [65] R. W. Hilditch, *An Introduction to Close Binary Stars*. Cambridge University Press, mar 2001.
- [66] P. C. Freire, C. G. Bassa, N. Wex, I. H. Stairs, D. J. Champion, S. M. Ransom, P. Lazarus, V. M. Kaspi, J. W. Hessels, M. Kramer, J. M. Cordes, J. P. Verbiest, P. Podsiadlowski, D. J. Nice, J. S. Deneva, D. R. Lorimer, B. W. Stappers, M. A. McLaughlin, and F. Camilo, “On the nature and evolution of the unique binary pulsar J1903+0327,” *Monthly Notices of the Royal Astronomical Society*, vol. 412, pp. 2763–2780, apr 2011.
- [67] N. Wex, “Testing relativistic gravity with radio pulsars,” in *Applications and Experiments*, DE GRUYTER, aug 2014.
- [68] Z. Arzoumanian, A. Brazier, S. Burke-Spolaor, S. Chamberlin, S. Chatterjee, B. Christy, J. M. Cordes, N. J. Cornish, F. Crawford, H. T. Cromartie, K. Crowter, M. E. DeCesar, P. B. Demorest, T. Dolch, J. A. Ellis, R. D. Ferdman, E. C. Ferrara, E. Fonseca, N. Garver-Daniels, P. A. Gentile, D. Halmrast, E. A. Huerta, F. A. Jenet, C. Jessup, G. Jones, M. L. Jones, D. L. Kaplan, M. T. Lam, T. J. W. Lazio, L. Levin, A. Lommen, D. R. Lorimer, J. Luo, R. S. Lynch, D. Madison, A. M. Matthews, M. A. McLaughlin, S. T. McWilliams, C. Mingarelli, C. Ng, D. J. Nice, T. T. Pennucci, S. M. Ransom, P. S. Ray, X. Siemens, J. Simon, R. Spiewak, I. H. Stairs, D. R. Stinebring, K. Stovall, J. K. Swiggum, S. R. Taylor, M. Vallisneri, R. van Haasteren, S. J. Vigeland, and W. Zhu, “The NANOGrav 11-year Data Set: High-precision Timing of 45 Millisecond Pulsars,” *The Astrophysical Journal Supplement Series*, vol. 235, p. 37, apr 2018.

- [69] G. Desvignes, R. N. Caballero, L. Lentati, J. P. Verbiest, D. J. Champion, B. W. Stappers, G. H. Janssen, P. Lazarus, S. Osłowski, S. Babak, C. G. Bassa, P. Brem, M. Burgay, I. Cognard, J. R. Gair, E. Graikou, L. Guillemot, J. W. Hessels, A. Jessner, C. Jordan, R. Karuppusamy, M. Kramer, A. Lassus, K. Lazaridis, K. J. Lee, K. Liu, A. G. Lyne, J. McKee, C. M. Mingarelli, D. Perrodin, A. Petiteau, A. Possenti, M. B. Purver, P. A. Rosado, S. Sanidas, A. Sesana, G. Shaifullah, R. Smits, S. R. Taylor, G. Theureau, C. Tiburzi, R. Van Haasteren, and A. Vecchio, “High-precision timing of 42 millisecond pulsars with the European Pulsar Timing Array,” *Monthly Notices of the Royal Astronomical Society*, vol. 458, pp. 3341–3380, mar 2016.
- [70] M. E. Gonzalez, I. H. Stairs, R. D. Ferdman, P. C. Freire, D. J. Nice, P. B. Demorest, S. M. Ransom, M. Kramer, F. Camilo, G. Hobbs, R. N. Manchester, and A. G. Lyne, “High-precision timing of five millisecond pulsars: Space velocities, binary evolution, and equivalence principles,” *Astrophysical Journal*, vol. 743, dec 2011.
- [71] J. Antoniadis, P. C. Freire, N. Wex, T. M. Tauris, R. S. Lynch, M. H. Van Kerkwijk, M. Kramer, C. Bassa, V. S. Dhillon, T. Driebe, J. W. Hessels, V. M. Kaspi, V. I. Kondratiev, N. Langer, T. R. Marsh, M. A. McLaughlin, T. T. Pennucci, S. M. Ransom, I. H. Stairs, J. Van Leeuwen, J. P. Verbiest, and D. G. Whelan, “A massive pulsar in a compact relativistic binary,” *Science*, vol. 340, p. 448, apr 2013.
- [72] W. W. Zhu, G. Desvignes, N. Wex, R. N. Caballero, D. J. Champion, P. B. Demorest, J. A. Ellis, G. H. Janssen, M. Kramer, A. Krieger, L. Lentati, D. J. Nice, S. M. Ransom, I. H. Stairs, B. W. Stappers, J. P. Verbiest, Z. Arzoumanian, C. G. Bassa, M. Burgay, I. Cognard, K. Crowter, T. Dolch, R. D. Ferdman, E. Fonseca, M. E. Gonzalez, E. Graikou, L. Guillemot, J. W. Hessels, A. Jessner, G. Jones, M. L. Jones, C. Jordan, R. Karuppusamy, M. T. Lam, K. Lazaridis, P. Lazarus, K. J. Lee, L. Levin, K. Liu, A. G. Lyne, J. W. McKee, M. A. McLaughlin, S. Osłowski, T. Pennucci, D. Perrodin, A. Possenti, S. Sanidas, G. Shaifullah, R. Smits, K. Stovall, J. Swiggum, G. Theureau, and C. Tiburzi, “Tests of gravitational symmetries with pulsar binary J1713+0747,” *Monthly Notices of the Royal Astronomical Society*, vol. 482, no. 3, pp. 3249–3260, 2019.
- [73] P. C. Freire, N. Wex, G. Esposito-Farèse, J. P. Verbiest, M. Bailes, B. A. Jacoby, M. Kramer, I. H. Stairs, J. Antoniadis, and G. H. Janssen, “The relativistic pulsar-white dwarf binary PSR J1738+0333 - II. The most stringent test of scalar-tensor gravity,” *Monthly Notices of the Royal Astronomical Society*, vol. 423, pp. 3328–3343, jul 2012.
- [74] R. Ellis, *An Introduction to the Theory of Stellar Structure and Evolution*, 2nd edn., by Dina Prialnik, vol. 52. jul 2011.