

symODE2: Symbolic analysis of second-order ordinary differential equations with polynomial coefficients

Tolga Birkandan*

Department of Physics, Istanbul Technical University, 34469 Istanbul, Turkey.

October 6, 2020

Abstract

An open-source package for symbolic analysis of second-order ordinary differential equations with polynomial coefficients is proposed. The approach is mainly based on the singularity structure of the equation and the routines are written under the open-source computer algebra system SageMath. The code is able to obtain the singularity structure, indices and recurrence relations associated with the regular singular points, and symbolic solutions of the hypergeometric equation, Heun equation, and their confluent forms.

Keywords: Ordinary differential equations, symbolic analysis, special functions

1 Introduction

Mathematical analysis of physical problems generally requires the methods of solving ordinary differential equations (ODEs). Numerical solutions of the initial and boundary value problems are often sufficient to give a concrete idea about the behavior of the system, whereas the analytic solution of an ODE in closed form, especially in terms of special functions may have more importance than constituting an exact solution. In some systems, the type of solution has the potential to reveal the symmetries of the system. For example, the emergence of the hypergeometric function in the solutions may indicate the conformal symmetry [1–4].

The applications of the general and confluent hypergeometric equation have dominated the 20th-century [5]. Although the mathematical theory of the Heun equation and its confluent forms are far from complete, these functions have been known and employed by the experts in the area for many years [6, 7]. Besides, the number of their applications increased substantially after the implementation of the Heun functions in the computer algebra system Maple [8] in 2005 [9–12]. The implementation of the Heun type functions in Mathematica in 2020 is another big leap for the Heun community [13, 14].

Most of the free and open-source computer algebra systems and packages involve the solutions of the hypergeometric equation, its confluent form, and related equations at least numerically [15–18]. However, the symbolic solutions of the Heun type equations are defined only in Maple and Mathematica which are commercial systems. The numerical evaluation of the general Heun and singly confluent Heun functions are studied by Motygin using the freely available GNU Octave language [19–21].

A historical review on computer algebra systems is given in 22 along with gravitational applications. The general methods for obtaining symbolic solutions to ODEs and a review of comprehensive literature before the year 2000 are given in [23]. Among the papers released prior to 2000, we should cite the seminal paper of Kovacic [24] and Duval

*E-mail: birkandant@itu.edu.tr

and Loday-Richaud’s work in which the hypergeometric and Heun type equations are studied in particular [25]. Among the papers on the solutions of ODEs in terms of special functions that are published after 2000, we can cite Bronstein and Lafaille [26], Chan and Cheb-Terrab [27], and van Hoeij with his collaborators [28–30].

SageMath is a free and open-source general-purpose computer algebra system licensed under the GPL [15]. SageMath offers a Python-based language and it is built on many open-source packages such as Maxima, SciPy, NumPy, and matplotlib. A variety of modules are present for many areas such as differential geometry and tensor calculus [31, 32] that enable calculations on quantum field theory and general relativity [33].

We will focus on the singularity analysis and symbolic solutions of the hypergeometric and Heun type equations using the SageMath system.

This paper is organized in the following way: In the second section, we present the code structure of our package with the functions defined in the code. In the third, fourth, and fifth sections, we briefly explain the singularity structure, series solutions, and change of variables for a second order ODE with polynomial coefficients, respectively. In the sixth and seventh sections, we give the results of our code for the standard forms and for the applications of the hypergeometric and Heun type equations. Section eight involves our conclusions and we describe the standard forms of the equations in the appendix.

All equations that are *not* ending with a comma or a period are computer-generated.

2 The code structure of the symODE2 package

The symODE2 package is written under SageMath 9.1 using a laptop computer with Intel(R) Core(TM) i7-6500U CPU @ 2.50GHz and 8 GB memory. The operating system is Windows 10 Enterprise ver.1909.

The package consists of two main parts: `ode2analyzer.sage` for the general analysis and `hypergeometric_heun.sage` for the symbolic solutions of the equations. `hypergeometric_heun.sage` calls the routines defined in `ode2analyzer.sage` when needed. We suggest the user put these two files in the same directory. A sample SageMath session is also given on the GitHub page [34].

The parts of the package and a sample session can be downloaded from the address [34]:

<https://github.com/tbirkandan/symODE2>

2.1 General analysis: ode2analyzer

The first part, `ode2analyzer` contains the routines,

- `find_singularities(diffeqn,y,z)`: Finds the singularity structure of the input ODE. The output is an array that involves the locations of the singularities, indices of the regular singularities and the ranks of the irregular singularities.
- `find_indices_recurrence(diffeqn,y,z,point,index,operation)`: Finds the indices and/or the recurrence relation with respect to a regular singular point using the θ -operator method which will be defined below.
- `ode_change_of_variable(diffeqn,y,z,transformation)`: Performs a change of variables.
- `normal_form_ode2(diffeqn,y,z)`: Finds the normal form of a second order ODE. For an input in the form (1), the output is in the form (4).

Here, the input arguments are

- `diffeqn`: A 2nd order ODE with polynomial coefficients.

- **y**: The dependent variable of the ODE.
- **z**: The independent variable of the ODE.
- **point**: The regular singular point for which the indices and/or the recurrence relation will be found.
- **index**: The name of the parameter that will be used for the characteristic exponent.
- **operation**: The result of the routine (indices, recurrence, full (both)).
- **transformation**: The transformation function that defines the change of variables.

2.2 Symbolic solutions of special ODEs: `hypergeometric_heun`

The second part, `hypergeometric_heun` contains the routines,

- `find_special_ode(diffeqn,y,z)`: Finds the type of the ODE using its singularity structure and solves it using the routines defined below.
- `ode_finder_bruteforce(diffeqn,y,z)`: Uses a change of variables list in order to bring the input ODE into a special form that is recognized in the package.
- `find_2F1(diffeqn,y,z)`: Solves a hypergeometric equation.
- `find_1F1(diffeqn,y,z)`: Solves a confluent hypergeometric equation.
- `find_HG(diffeqn,y,z)`: Solves a general Heun equation.
- `find_HC(diffeqn,y,z)`: Solves a (singly) confluent Heun equation.
- `find_HD(diffeqn,y,z)`: Solves a double confluent Heun equation.
- `find_HB(diffeqn,y,z)`: Solves a biconfluent Heun equation.
- `find_HT(diffeqn,y,z)`: Solves a triconfluent Heun equation.

The input arguments (`diffeqn,y,z`) are the same as defined in the first part.

3 Singularity analysis

A second order ODE can be written in the form,

$$f_1(x)\frac{d^2y(x)}{dx^2} + f_2(x)\frac{dy(x)}{dx} + f_3(x)y(x) = 0. \quad (1)$$

From this point on, the coefficient functions $f_{1,2,3}(x)$ will be regarded as polynomials in the parameter x , and the x -dependence of the function y will be omitted.

We can denote $p(x) = f_2(x)/f_1(x)$ and $q(x) = f_3(x)/f_1(x)$ to obtain,

$$\frac{d^2y}{dx^2} + p(x)\frac{dy}{dx} + q(x)y = 0. \quad (2)$$

If the functions $p(x)$ and $q(x)$ are analytic at a point $x = x_0$, then x_0 is an “ordinary point” for this ODE.

The points that make $p(x)$ or $q(x)$ divergent are called the singular points or singularities of the ODE. If x_* is a singular point and if $(x - x_*)p(x)$ and $(x - x_*)^2q(x)$ are both analytic at $x = x_*$, then x_* is called a “regular singular

point”. Otherwise, the singular point is “irregular” [35]. The singularity behavior at $x \rightarrow \infty$ can be analyzed by performing the transformation $\tilde{x} = 1/x$ and checking the behavior at $\tilde{x} = 0$. If all the singular points of an ODE are regular, then the ODE is said to be a “Fuchsian equation”.

If the singularity at $x = x_*$ is irregular but $(x - x_*)^k p(x)$ and $(x - x_*)^{2k} q(x)$ are analytic where k is the least integer satisfying this condition, then the irregular singular point at $x = x_*$ has a rank $(k - 1)$ [35]. Consequently, a regular singular point is of rank-0 as $k = 1$.

We can write the equation (2) in normal form in which the coefficient of the first derivative vanishes. We define $y(x) = g(x)w(x)$ and for

$$g(x) = e^{-\frac{1}{2} \int^x p(x') dx'}, \quad (3)$$

we obtain

$$\frac{d^2 w}{dx^2} + q'(x)w = 0, \quad (4)$$

where

$$q'(x) = q(x) - \frac{1}{2} \frac{dp(x)}{dx} - \frac{p(x)^2}{4}. \quad (5)$$

Our code transforms the input equation into the normal form in order to deal with the singular points of only one function, namely $q'(x)$. Although the package involves a routine for general change of variables, the results of the transformation $\tilde{x} = 1/x$ is included in the function that finds the singularity behavior as the analysis of $x \rightarrow \infty$ case is inevitable.

4 Series solution around a regular singular point

An indicial equation can be defined for a regular singular point [7]. For a finite regular singular point x_* we have,

$$r(r - 1) + p_* r + q_* = 0, \quad (6)$$

where p_* and q_* are the residues of $p(x)$ and $(x - x_*)q(x)$ at $x = x_*$, respectively. For the regular singularity at infinity, one can write

$$r(r + 1) - p_\infty r + q_\infty = 0, \quad (7)$$

where p_∞ and q_∞ are the residues of $p(x)$ and $xq(x)$ at $x \rightarrow \infty$, respectively.

The roots $r = r_1$ and $r = r_2$ of the indicial equation are called the “indices” or “characteristic exponents”, or “Frobenius exponents” of the corresponding regular singularity [7]. The sum of the all $(2n)$ indices corresponding to all (n) regular singular points in a Fuchsian equation should be equal to $n - 2$ [36].

One can find at least one series solution, the “Frobenius solution” of the form,

$$y = \sum_{n=0}^{\infty} C_n (x - x_*)^{n+r}, \quad (8)$$

near a finite regular singular point x_* , r being a characteristic exponent associated with x_* . The details on the second solution and the solution around infinity can be found in [7]. We substitute the solution (8) into equation (1) to obtain a recurrence relation among the coefficients C_n . For example, the hypergeometric equation admits a two-term recurrence relation which connects C_n with C_{n-1} , while the Heun equation has a three-term recurrence relation connecting C_n , C_{n-1} and C_{n-2} .

The θ -operator method yields the indicial equation and the recurrence relation for a regular singular point with less effort than the formal Frobenius series calculation. Following [37], we define $D = \frac{d}{dx}$ and,

$$\theta = x \frac{d}{dx} = xD, \quad (9)$$

$$\theta(\theta - 1) = x^2 D^2. \quad (10)$$

Similarly, we have

$$x^n D^n = (-1)^n (-\theta)_n, \quad (11)$$

where

$$(-\theta)_n = \prod_{j=0}^{n-1} (-\theta + j), \quad (12)$$

is the generalized factorial notation. A general n^{th} order ODE can be written as

$$[a_0(x)D^n + a_1(x)D^{n-1} + \dots + a_n(x)]y = 0. \quad (13)$$

Using eq.(11), we can write the equation in the θ -form, namely

$$[A_0(\theta) + xA_1(\theta) + x^2A_2(\theta) + \dots + x^m A_m(\theta)]y = 0, \quad (14)$$

if the coefficients of the original equation are polynomials in x . Here, $A_{0,1,\dots,m}(\theta)$ are polynomials in θ .

Let us assume that $x = 0$ is a regular singular point and seek a series solution in the form (8) for $x_* = 0$. It is known that any polynomial expression in θ operating on x^n yields the same polynomial in n times x^n , thus $P(\theta)x^n = P(n)x^n$ [37]. Using this property, we get

$$\sum_{n=0}^{\infty} C_n [A_0(n+r)x^{n+r} + A_1(n+r)x^{n+r+1} + \dots + A_m(n+r)x^{n+r+m}] = 0. \quad (15)$$

In order to have an arbitrary C_0 , the indicial equation is obtained as $A_0(r) = 0$. Let us shift the indices in the sum,

$$\sum_{n=0}^{\infty} [C_n A_0(n+r) + C_{n-1} A_1(n+r-1) + \dots + C_{n-m} A_m(n+r-m)] x^{n+r} = 0, \quad (16)$$

and the recurrence relation reads

$$C_n A_0(n+r) + C_{n-1} A_1(n+r-1) + \dots + C_{n-m} A_m(n+r-m) = 0. \quad (17)$$

For a regular singular point x_* other than zero, one should make the transformation $x' = x - x_*$ and do the calculation for x' . For the details of the θ -operator method, proofs and examples, we refer the reader to [37].

In our code, the function that finds the indices and the recurrence relation for a given regular singular point is based on the θ -operator method.

Let us reset the SageMath session, load the file containing the routines and declare our functions,

```

1 # Start the session
2 reset()
3 load('ode2analyzer.sage')
4 #####
5 u = function('u')(x)
6 u(x)=u
7 f1 = function('f1')(x)
8 f2 = function('f2')(x)
9 f3 = function('f3')(x)
10 #####

```

Here, the line $u(x)=u$ is written in order to avoid a “Deprecation Warning” as explained in page 162 of [38]. The variables and functions that define the hypergeometric equation are given as

```

1 # Hypergeometric equation:
2 var('a,b,c');
3 f1(x)=x*(1-x);

```

```

4 f2(x)=(c-(a+b+1)*x);
5 f3(x)=-a*b;
6
7 myeqn=f1(x)*diff(u(x),x,x)+f2(x)*diff(u(x),x)+f3(x)*u(x)
8 show(myeqn)

```

The hypergeometric equation as given in eq. (78) is thus defined and shown as

$$-abu(x) - (x-1)x \frac{\partial^2}{(\partial x)^2} u(x) - ((a+b+1)x-c) \frac{\partial}{\partial x} u(x) \quad (18)$$

The singularity structure of the equation is found by the command

```

1 # Singularity structure
2 mystructure=find_singularities(diffeqn=myeqn,y=u,z=x)
3 show(mystructure)

```

as

$$([1, 0, +\infty], [[-a-b+c, 0], [-c+1, 0], [b, a]], [], []) \quad (19)$$

There are four sub-arrays in the resulting array. The first one contains the locations of the regular singular points, the second one contains the indices (characteristic exponents) associated with the regular singular points (respectively), the third one contains the locations of the irregular singular points and the last array contains the ranks of the irregular singular points (respectively).

Here, the singularity structure of the general hypergeometric equation is found correctly [39]. We can run our code for an ODE with irregular singularities as in the following example. The double confluent Heun equation 100 can be defined as

```

1 # Double confluent Heun:
2 var('alpha,gamma,delta,epsilon,q');
3 f1(x)=1;
4 f2(x)=(delta/x)+(gamma/(x^2))+epsilon;
5 f3(x)=(alpha*x-q)/(x^2)
6
7 myeqn=f1(x)*diff(u(x),x,x)+f2(x)*diff(u(x),x)+f3(x)*u(x)
8 show(myeqn)

```

$$\left(\epsilon + \frac{\delta}{x} + \frac{\gamma}{x^2}\right) \frac{\partial}{\partial x} u(x) + \frac{(\alpha x - q)u(x)}{x^2} + \frac{\partial^2}{(\partial x)^2} u(x) \quad (20)$$

and the singularity structure can be found by the same command as

$$([], [], [0, +\infty], [1, 1]) \quad (21)$$

Here, we find two rank-1 irregular singular points located at $(0, \infty)$ [39]. The (singly) confluent Heun equation (92) case can be analyzed as

```

1 # (Singly) Confluent Heun:
2 var('alpha,gamma,delta,epsilon,q');
3 f1(x)=1;
4 f2(x)=(gamma/x)+(delta/(x-1))+epsilon;
5 f3(x)=(alpha*x-q)/(x*(x-1))
6
7 myeqn=f1(x)*diff(u(x),x,x)+f2(x)*diff(u(x),x)+f3(x)*u(x)
8 show(myeqn)
9
10 mystructure=find_singularities(diffeqn=myeqn,y=u,z=x)
11 show(mystructure)

```

$$\left(\epsilon + \frac{\delta}{x-1} + \frac{\gamma}{x}\right) \frac{\partial}{\partial x} u(x) + \frac{(\alpha x - q)u(x)}{(x-1)x} + \frac{\partial^2}{(\partial x)^2} u(x) \quad (22)$$

$$([1, 0], [[-\delta + 1, 0], [-\gamma + 1, 0]], [+ \infty], [1]) \quad (23)$$

with two regular singularities at $(0, 1)$ and one irregular singularity at (∞) of rank-1 [7, 39].

Another routine of our code is able to find the indices and the recurrence relation for a given regular singular point of a second-order ODE with polynomial coefficients. An input parameter `operation` can take the values: "indices", "recurrence", and "full", yielding the corresponding results ("full" gives both the indices and the recurrence relation).

For the hypergeometric equation, around $x = 0$ we have

```

1 # Hypergeometric equation
2 var('a,b,c'); f1(x)=x*(1-x); f2(x)=(c-(a+b+1)*x); f3(x)=-a*b;
3 myeqn=f1(x)*diff(u(x),x,x)+f2(x)*diff(u(x),x)+f3(x)*u(x)
4
5 var('r')
6 myindicesandrecurrence=find_indices_recurrence(diffeqn=myeqn,y=u,z=x,point=0,index=r,
7         operation="full")
8 myindices=find_indices_recurrence(myeqn,u,x,0,r,"indices")
9 myrecurrence=find_indices_recurrence(myeqn,u,x,0,r,"recurrence")
10
11 show(myindicesandrecurrence)
12 show(myindices)
13 show(myrecurrence)

```

Here, "r" is the parameter associated with the characteristic exponent. The results of the commands with parameters "indices" and "recurrence" are

$$[-c + 1, 0] \quad (24)$$

$$\begin{aligned} & ((a-1)b + (a+b-2)n + n^2 + (a+b+2n-2)r + r^2 - a + 1)C(n-1) \\ & - ((c-1)n + n^2 + (c+2n-1)r + r^2)C(n) \end{aligned} \quad (25)$$

We obtain the recurrence relation corresponding to $r = 0$ by substituting this value in the recurrence relation and solving it for $C(n)$. The result will be in the form $C(n) = (\dots)C(n-1)$ as expected, namely

```

1 # The recurrence relation
2 solve(myrecurrence==0,C(n))[0].subs(r=0).rhs().full_simplify().collect_common_factors()

```

$$C(n) = \frac{((a-1)b + (a+b-2)n + n^2 - a + 1)C(n-1)}{(c+n-1)n} \quad (26)$$

which agrees with the one given in [7].

The hypergeometric function is implemented in SageMath as `hypergeometric([a,b],[c],x)`. Therefore we can verify our recurrence relation graphically using the code below. The code adds new terms to the summation if the new term is greater than the "machine epsilon" of the system.

```

1 # Plot of 2F1
2 def hypfromrecurrence(a,b,c,x):
3     machineepsilon=abs((7./3)-(4./3)-1)
4     Cn=1
5     n=0
6     result=1
7     while True:
8         coeff=(a*b + (a+b)*n + n^2)*(Cn)/((c+1)*n + n^2 + c)

```

```

9     newterm=coeff*(x^(n+1))
10    if abs(newterm)<machineepsilon:
11        break
12    else:
13        result=result+newterm
14        Cn=coeff
15        n=n+1
16    return result
17
18    # https://dlmf.nist.gov/15.3.F2.mag
19    a=5
20    b=-10
21    c=1
22
23    hypsage(x)=hypergeometric([a,b],[c],x)
24
25    p1=plot(hypsage,(x,-0.023,0.7),legend_label='SageMath',axes_labels=[r'$x$', r'$_2F_1$
26        ([5,-10],[1],x)'])
27    p2=plot(hypfromrecurrence(a,b,c,x),(x,-0.023,0.7),legend_label='Recurrence relation',
28        plot_points=10, linestyle='', marker='.',color='red')
29    (p1+p2).show()

```

We take the parameters a, b, c such that our result coincides with the plot given in [39] as Figure (15.3.2). We present the result of the SageMath and our recurrence relation on the same plot in Figure (1).




Figure 1: Plot of the hypergeometric function found by our recurrence relation result and the internal command of SageMath.

We should note that finding a solution using the recurrence relation generally requires more effort than presented here. Methods such as analytic continuation should be carefully applied in order to deal with the circle of convergence of the series solution [20, 21].

For the general Heun equation we have,

```

1    # Heun equation
2    var('alpha,beta,gamma,delta,epsilon,q,a1');
3    epsilon=alpha+beta+1-gamma-delta;

```



```

4 f1(x)=1;f2(x)=(gamma/x)+(delta/(x-1))+(epsilon/(x-a1));f3(x)=(alpha*beta*x-q)/(x*(x-1)*(
  x-a1))
5
6 myeqn=f1(x)*diff(u(x),x,x)+f2(x)*diff(u(x),x)+f3(x)*u(x)
7
8 var('r')
9 myrecurrence=find_indices_recurrence(myeqn,u,x,0,r,"recurrence").subs(r=0)
10 show(myrecurrence)
11
12 # The recurrence relation given in \protect\vrule width0pt\protect\href{http://dlmf.nist
  .gov/31.3.E3}{http://dlmf.nist.gov/31.3.E3}
13 # with an overall minus sign
14 Rn=a1*(n+1)*(n+gamma)
15 Pn=(n-1+alpha)*(n-1+beta)
16 Qn=n*((n-1+gamma)*(1+a1)+a1*delta+epsilon)
17 recurrenceDLMF=-((Rn*C(n+1)-(Qn+q)*C(n)+Pn*C(n-1))
18 # We need to shift n by 1 to check with the DLMF result
19 myrecurrence=myrecurrence.subs(n=n+1)
20
21 # Their difference should be zero
22 shouldbezero=(myrecurrence-recurrenceDLMF).expand().factor().full_simplify()
23 show(shouldbezero)

```

We obtain the recurrence relation as

$$\begin{aligned}
& ((a_1 + 1)n^2 - (a_1 - 1)\delta - a_1\gamma + ((a_1 - 1)\delta + a_1\gamma - 3a_1 + \alpha + \beta - 2)n + 2a_1 - \alpha - \beta + q + 1)C(n - 1) \\
& - ((\alpha - 2)\beta + (\alpha + \beta - 4)n + n^2 - 2\alpha + 4)C(n - 2) \\
& - (a_1n^2 + (a_1\gamma - a_1)n)C(n)
\end{aligned} \tag{27}$$

We also enter the recurrence relation given in [39] with an overall minus sign. The difference of the two results is zero, verifying the result of our code.

We plot the series solution for the general Heun equation in Figure (2) using a similar code with an array of plots and compare it with the plot given in the Wolfram Blog post [14] to see that they are similar. The details of this analysis can be found in the sample SageMath session [34].




Figure 2: Plot of the general Heun function similar to the one given in [14].

5 Change of variables for a second order ODE

For an ODE of the form (1), we can change the independent variable from x to $z(x)$ using

$$\frac{dy}{dx} = \frac{dz}{dx} \frac{dy}{dz}, \quad (28)$$

$$\frac{d^2y}{dx^2} = \left(\frac{dz}{dx}\right)^2 \frac{d^2y}{dz^2} + \frac{d^2z}{dx^2} \frac{dy}{dz}. \quad (29)$$

As an example, let us make the change $x \rightarrow \sqrt{z}$. Then $z = x^2$ and

$$\frac{dy}{dx} = 2\sqrt{z} \frac{dy}{dz}, \quad (30)$$

$$\frac{d^2y}{dx^2} = 4z \frac{d^2y}{dz^2} + 2 \frac{dy}{dz}, \quad (31)$$

and the ODE becomes

$$4zf_1(z) \frac{d^2y(z)}{dz^2} + 2[f_1(z) + \sqrt{z}f_2(z)] \frac{dy(z)}{dz} + f_3(z)y(z) = 0. \quad (32)$$

We can work with this transformation for a general 2nd order ODE using the `ode_change_of_variable` routine of our code.

```

1 # Change of variables
2 reset()
3 load('ode2analyzer.sage')
4 #####
5 u = function('u')(x)
6 u(x)=u
7 f1 = function('f1')(x)
8 f2 = function('f2')(x)
9 f3 = function('f3')(x)
10 #####
11 myeqn=f1(x)*diff(u(x),x,x)+f2(x)*diff(u(x),x)+f3(x)*u(x)
12 show(myeqn)
13 myresult=ode_change_of_variable(diffeqn=myeqn,y=u,z=x,transformation=sqrt(x))
14 show(myresult)

```

which yields

$$myeqn = f_3(x)u(x) + f_2(x) \frac{\partial}{\partial x}u(x) + f_1(x) \frac{\partial^2}{(\partial x)^2}u(x) \quad (33)$$

$$myresult = 4xf_1(\sqrt{x}) \frac{\partial^2}{(\partial x)^2}u(x) + f_3(\sqrt{x})u(x) + 2(\sqrt{x}f_2(\sqrt{x}) + f_1(\sqrt{x})) \frac{\partial}{\partial x}u(x) \quad (34)$$

The user should be careful as the variable name is not changed. The routine `ode_change_of_variable` used here is actually intended for the intrinsic use of other functions in our code where the unchanged variable name is more appropriate as the explicit declaration of a new variable is required in SageMath system.

6 Hypergeometric and Heun-type equations

The code attempts to find symbolic solutions of some special ODEs, namely, the hypergeometric equation, the Heun equation, and their confluent forms. The analysis of the equations is based on the singularity structure. Locations of the singularities and corresponding characteristic exponents play a major role in the method. Using particular

substitutions and transformations, the input equation is brought in a standard form that can be recognized by the routines.

A basic example of our approach can be given by using the hypergeometric equation. The Riemann P -symbol for the standard form of the hypergeometric equation (78) is

$$P \left\{ \begin{array}{ccc} 0 & 1 & \infty \\ 0 & 0 & a \\ 1-c & c-a-b & b \end{array} \begin{array}{c} x \\ \\ \end{array} \right\}. \quad (35)$$

Here, the locations of the singular points are given in the first row and each column exhibits the characteristic exponents of the corresponding singular points as found by our code in (19).

Let us study a more general second order ODE with three regular singular points (x_1, x_2, x_3) and corresponding indices $(c_{i1}, c_{i2}, i = 1, 2, 3)$, namely,

$$P \left\{ \begin{array}{ccc} x_1 & x_2 & x_3 \\ c_{11} & c_{21} & c_{31} \\ c_{12} & c_{22} & c_{32} \end{array} \begin{array}{c} x \\ \\ \end{array} \right\}. \quad (36)$$

The substitution,

$$u(x) \rightarrow \begin{cases} \left(\frac{x-x_1}{x-x_3}\right)^{c_{11}} \left(\frac{x-x_2}{x-x_3}\right)^{c_{21}} u(x), & \text{if } x_3 \neq \infty, \\ (x-x_1)^{c_{11}} (x-x_2)^{c_{21}} u(x), & \text{if } x_3 = \infty, \end{cases} \quad (37)$$

brings the P -symbol in the form

$$P \left\{ \begin{array}{ccc} x_1 & x_2 & x_3 \\ 0 & 0 & c_{31} + c_{11} + c_{21} \\ c_{12} - c_{11} & c_{22} - c_{21} & c_{32} + c_{11} + c_{21} \end{array} \begin{array}{c} x \\ \\ \end{array} \right\}. \quad (38)$$

The transformation,

$$x \rightarrow \frac{(x_2 - x_3)(x - x_1)}{(x_2 - x_1)(x - x_3)}, \quad (39)$$

moves the locations of the singular points from (x_1, x_2, x_3) to $(0, 1, \infty)$ as in the standard form of the hypergeometric equation [40]. Now we have,

$$P \left\{ \begin{array}{ccc} 0 & 1 & \infty \\ 0 & 0 & c_{31} + c_{11} + c_{21} \\ c_{12} - c_{11} & c_{22} - c_{21} & c_{32} + c_{11} + c_{21} \end{array} \begin{array}{c} x \\ \\ \frac{(x_2-x_3)(x-x_1)}{(x_2-x_1)(x-x_3)} \end{array} \right\}, \quad (40)$$

which corresponds to the standard form of the hypergeometric equation. We note that the sum of the indices is not changed. A similar analysis of the general Heun equation can be found in [40].

Our approach is similar to this example for other equations: we change the indices of the regular singular points and move the locations of the singular points in order to obtain a standard form. After reaching the standard singularity structure of an equation that can be recognized by the code, the parameters are read either from the characteristic exponents or by matching the final form of the input equation with the standard equation in their normal forms.

The polynomial coefficients of the normal forms are matched in the confluent cases. The parameters of these ODEs can be found by solving single equations, i.e. the code finds some parameters by solving algebraic equations that depend only on one parameter. The rest of the parameters are found by substitution. For the Fuchsian ODEs, the parameters are read from the characteristic exponents, e.g. the non-zero exponent of the singular point at zero in the hypergeometric equation yields $1 - c$, etc.

We find the parameters with this method and use the Maple or Mathematica forms of the solutions to substitute these parameters. The standard forms of the equations are given in the Appendix.

The hypergeometric equation has three pairs of Frobenius solutions around its three regular singular points and these solutions can be transformed into other solutions via specific transformations [39]. The number of all solutions of the hypergeometric equation is 24. The number of total solutions is 192 for the general Heun equation [41]. The user of our code may need to use some transformations or function identities in order to obtain the desired form of the solution [36, 39, 41].

The results of the hypergeometric and confluent hypergeometric equations are numerically usable as these functions are defined in SageMath. However, the numerical solutions of the Heun-type functions are not defined. The numerical solutions of the general Heun and (singly) confluent Heun functions are defined by Motygin for GNU Octave/MATLAB [20, 21]. GNU Octave/MATLAB commands can be run in a SageMath session. However, this procedure is not straightforward and it is beyond the scope of this work.

Regarding our code, although the results of our limited number of tests seem proper, it should be emphasized that the solutions should be taken with great care as such routines should be tested heavily before using them for new problems. The user should approach these routines as an experimental effort and follow the updates from the GitHub page [34] regularly.

The outputs of the following examples are obtained using the following code structure,

```

1 # Standard equations
2 reset()
3 load('hypergeometric_heun.sage') #Routines for the symbolic solutions
4 #####
5 u = function('u')(x)
6 u(x)=u
7 f1 = function('f1')(x)
8 f2 = function('f2')(x)
9 f3 = function('f3')(x)
10 #####
11
12 # Definition of the variables , f1, f2, and f3.
13
14 myeqn=f1(x)*diff(u(x),x,x)+f2(x)*diff(u(x),x)+f3(x)*u(x)
15 show(myeqn)
16 mysingularities=find_singularities(diffeqn=myeqn,y=u,z=x)
17 show(mysingularities)
18 myresult=find_special_ode(diffeqn=myeqn,y=u,z=x)

```

The singularity structures and the solutions for the corresponding equations are found by the code as in the following sections.

6.1 Hypergeometric equation

$$-abu(x) - (x-1)x \frac{\partial^2}{(\partial x)^2} u(x) - ((a+b+1)x-c) \frac{\partial}{\partial x} u(x) \quad (41)$$

$$([1, 0, +\infty], [[-a-b+c, 0], [-c+1, 0], [b, a]], [], []) \quad (42)$$

$$\frac{C_2 x {}_2F_1\left(\begin{matrix} a-c+1, b-c+1 \\ -c+2 \end{matrix}; x\right) + C_1 x^c {}_2F_1\left(\begin{matrix} a, b \\ c \end{matrix}; x\right)}{x^c} \quad (43)$$

The result is numerically usable as `hypergeometric([a,b],[c],x)` functions are defined within SageMath.

6.2 Confluent hypergeometric equation

$$-au(x) + (b-x)\frac{\partial}{\partial x}u(x) + x\frac{\partial^2}{(\partial x)^2}u(x) \quad (44)$$

$$([0], [[-b+1, 0]], [+∞], [1]) \quad (45)$$

$$C_1M(a, b, x) + C_2U(a, b, x) \quad (46)$$

The result is numerically usable as `hypergeometric_M(a,b,x)` and `hypergeometric_U(a,b,x)` functions are defined within SageMath.

6.3 (General) Heun equation

$$-\left(\frac{\alpha + \beta - \delta - \gamma + 1}{a_1 - x} - \frac{\delta}{x-1} - \frac{\gamma}{x}\right)\frac{\partial}{\partial x}u(x) - \frac{(\alpha\beta x - q)u(x)}{(a_1 - x)(x-1)x} + \frac{\partial^2}{(\partial x)^2}u(x) \quad (47)$$

$$([a_1, 1, 0, +∞], [[-\alpha - \beta + \delta + \gamma, 0], [-\delta + 1, 0], [-\gamma + 1, 0], [\beta, \alpha]], [], []) \quad (48)$$

$$C_2x^{-\gamma+1}\text{HG}(a_1, -(a_1\delta + \alpha + \beta - \delta - \gamma + 1)(\gamma - 1) + q, \alpha - \gamma + 1, \beta - \gamma + 1, -\gamma + 2, \delta, x) \\ + C_1\text{HG}(a_1, q, \alpha, \beta, \gamma, \delta, x) \quad (49)$$

6.4 (Singly) Confluent Heun equation

$$\left(\frac{\nu}{x-1} + \frac{\mu}{x}\right)u(x) + \left(\alpha + \frac{\gamma+1}{x-1} + \frac{\beta+1}{x}\right)\frac{\partial}{\partial x}u(x) + \frac{\partial^2}{(\partial x)^2}u(x) \quad (50)$$

$$([1, 0], [[-\gamma, 0], [-\beta, 0]], [+∞], [1]) \quad (51)$$

$$C_1\text{HC}\left(\alpha, \beta, \gamma, -\frac{1}{2}\alpha\beta - \frac{1}{2}\alpha\gamma - \alpha + \mu + \nu, \frac{1}{2}(\alpha-1)\beta - \frac{1}{2}(\beta+1)\gamma + \frac{1}{2}\alpha - \mu, x\right) \\ + \frac{C_2\text{HC}\left(\alpha, -\beta, \gamma, -\frac{1}{2}\alpha\beta - \frac{1}{2}\alpha\gamma - \alpha + \mu + \nu, \frac{1}{2}(\alpha-1)\beta - \frac{1}{2}(\beta+1)\gamma + \frac{1}{2}\alpha - \mu, x\right)}{x^\beta} \quad (52)$$

6.5 Double confluent Heun equation

$$\left(\epsilon + \frac{\delta}{x} + \frac{\gamma}{x^2}\right)\frac{\partial}{\partial x}u(x) + \frac{(\alpha x - q)u(x)}{x^2} + \frac{\partial^2}{(\partial x)^2}u(x) \quad (53)$$

$$([], [], [0, +∞], [1, 1]) \quad (54)$$

$$C_2x^{-\delta+2}\text{HD}(\delta + q - 2, \alpha - 2\epsilon, -\gamma, -\delta + 4, -\epsilon, x)e^{(-\epsilon x + \frac{\gamma}{x})} + C_1\text{HD}(q, \alpha, \gamma, \delta, \epsilon, x) \quad (55)$$

6.6 Biconfluent Heun equation

$$\left(\epsilon x + \delta + \frac{\gamma}{x}\right)\frac{\partial}{\partial x}u(x) + \frac{(\alpha x - q)u(x)}{x} + \frac{\partial^2}{(\partial x)^2}u(x) \quad (56)$$

$$([0], [[-\gamma + 1, 0]], [+∞], [2]) \quad (57)$$

$$C_2x^{-\gamma+1}\text{HB}(\delta(\gamma-1) + q, -\epsilon(\gamma-1) + \alpha, -\gamma + 2, \delta, \epsilon, x) + C_1\text{HB}(q, \alpha, \gamma, \delta, \epsilon, x) \quad (58)$$

6.7 Triconfluent Heun equation

$$(\alpha x - q)u(x) + (\epsilon x^2 + \delta x + \gamma)\frac{\partial}{\partial x}u(x) + \frac{\partial^2}{(\partial x)^2}u(x) \quad (59)$$

$$([], [], [+∞], [3]) \quad (60)$$

$$C_2\text{HT}(\delta + q, \alpha - 2\epsilon, -\gamma, -\delta, -\epsilon, x)e^{(-\frac{1}{3}\epsilon x^3 - \frac{1}{2}\delta x^2 - \gamma x)} + C_1\text{HT}(q, \alpha, \gamma, \delta, \epsilon, x) \quad (61)$$

7 Applications with symODE2

We will study some examples in the literature in order to test our code. For further applications, the user should keep in mind that some transformations or identities may be required to obtain the same result with the literature.

7.1 Hypergeometric equation: Equation (15) of [42]

The equation (15) of Nasheeha et. al [42],

$$4(ax+1)(bx+1)\frac{\partial^2}{(\partial x)^2}u(x) - \left(\left(a(2\alpha+1)^2 - 2a+b\right)b + ab - b^2\right)u(x) + 2(a-b)\frac{\partial}{\partial x}u(x) \quad (62)$$

can be solved using our code,

```

1 # The equation (15) of Nasheeha et. al
2 reset()
3 load('hypergeometric_heun.sage') #Routines for the symbolic solutions
4 #####
5 u = function('u')(x)
6 u(x)=u
7 f1 = function('f1')(x)
8 f2 = function('f2')(x)
9 f3 = function('f3')(x)
10 #####
11 var('a,b,k,alpha');
12 f1(x)=4*(1+a*x)*(1+b*x);
13 f2(x)=2*(a-b);
14 f3(x)=((b^2)-a*b-k*b)
15
16 myeqn=(f1(x)*diff(u(x),x,x)+f2(x)*diff(u(x),x)+f3(x)*u(x))
17 myresult=find_special_ode(diffeqn=myeqn,y=u,z=x)

```

as

$$\frac{\sqrt{ax+1}C_2\sqrt{b}{}_2F_1\left(-\alpha-\frac{1}{2},\alpha+\frac{1}{2};-\frac{abx+b}{a-b}\right) + C_1\sqrt{-a+b}{}_2F_1\left(-\alpha-1,\alpha;-\frac{abx+b}{a-b}\right)}{\sqrt{-a+b}} \quad (63)$$

Here, the code found the change of variables itself. If we use the transformation given in the article, we find the same result as in the article, namely,

```

1 # Change of variables
2 myeqn2=ode_change_of_variable(diffeqn=myeqn,y=u,z=x,transformation=((a-b)*x-a)/(a*b))
3 myeqn2=myeqn2.subs(k==a*(2*alpha+1)^2-2*a+b)
4 myresult=find_special_ode(diffeqn=myeqn2,y=u,z=x)

```

and

$$C_2 x^{\frac{3}{2}} {}_2F_1\left(-\alpha + \frac{1}{2}, \alpha + \frac{3}{2}; x\right) + C_1 {}_2F_1\left(-\alpha - 1, \alpha; -\frac{1}{2}; x\right) \quad (64)$$

7.2 Confluent hypergeometric equation: Whittaker equation

The Whittaker equation [39],

$$\frac{d^2 u}{dz^2} + \left(-\frac{1}{4} + \frac{\kappa}{x} + \frac{\frac{1}{4} - \mu^2}{x^2}\right) u = 0, \quad (65)$$

can be obtained from the confluent hypergeometric equation. Hence the solutions can be given in terms of the confluent hypergeometric functions M and U . We start by the substitution $u(x) = e^{-x/2} u(x)$ and the code

```

1 # The Whittaker equation
2 var('kappa , mu '); f1(x)=1; f2(x)=0; f3(x)=(-1/4)+(kappa/x)+((1/4)-(mu^2))/(x^2)
3 myeqn=f1(x)*diff(u(x),x,x)+f2(x)*diff(u(x),x)+f3(x)*u(x)
4 myeqn=myeqn.substitute_function(u(x),exp(-x/2)*u(x))
5 myresult=find_special_ode(diffeqn=myeqn,y=u,z=x)

```

yields

$$\left(C_1 M\left(-\kappa + \mu + \frac{1}{2}, 2\mu + 1, x\right) + C_2 U\left(-\kappa + \mu + \frac{1}{2}, 2\mu + 1, x\right)\right) x^{\mu + \frac{1}{2}} \quad (66)$$

as the same result given in [39]. The code finds the factor $x^{\mu + \frac{1}{2}}$ itself.

7.3 General Heun equation: Equation (10) of [43]

The equation (10) of Petroff [43] is

$$-(x^2 - 1)(A - x) \frac{\partial^2}{(\partial x)^2} u(x) - 4Au(x) - (5Ax - 4x^2 - 1) \frac{\partial}{\partial x} u(x) \quad (67)$$

The author applies a change of variables as $z = (1 - x)/2$. Let us use this transformation and solve the resulting equation with our code,

```

1 # The equation (10) of Petroff
2 var('A '); f1(x)=(x-A)*((x^2)-1); f2(x)=4*(x^2)-5*A*x+1; f3(x)=-4*A;
3
4 myeqn=(f1(x)*diff(u(x),x,x)+f2(x)*diff(u(x),x)+f3(x)*u(x))
5 show(myeqn)
6 myeqn2=ode_change_of_variable(diffeqn=myeqn,y=u,z=x,transformation=1-2*x)
7 myresult=find_special_ode(diffeqn=myeqn2,y=u,z=x)

```

that yields

$$C_1 \text{HG}\left(-\frac{1}{2}A + \frac{1}{2}, -2A, 3, 0, \frac{5}{2}, \frac{5}{2}, x\right) + \frac{C_2 \text{HG}\left(-\frac{1}{2}A + \frac{1}{2}, -\frac{1}{8}A - \frac{3}{8}, \frac{3}{2}, -\frac{3}{2}, -\frac{1}{2}, \frac{5}{2}, x\right)}{x^{\frac{3}{2}}} \quad (68)$$

which is the same with the reference.

7.4 Confluent Heun equation: Equation (26) of [44]

The ODE (26) in Sakallı et al. [44],

$$-(x - 1)x \frac{\partial^2}{(\partial x)^2} u(x) - \left(L_1 m^2 r(x - 1) - L_1^2 \omega^2 \left(\frac{1}{x} - 1\right) - L_1(L_2 + 1)L_2\right) u(x) - (2x - 1) \frac{\partial}{\partial x} u(x) \quad (69)$$

needs the introduction of the factor $x^{iL_1\omega}$ given in eq. (27) of this article in order to obtain it in the standard form of the (singly) confluent Heun equation. Our code finds this function automatically. Keeping in mind that L_1 is L and L_2 is ℓ , we run

```

1 # The ODE (26) in Sakalli et al.
2 var('L1,L2,omega,m,r');
3 f1(x)=x*(1-x);
4 f2(x)=1-2*x;
5 f3(x)=((L1^2)*(omega^2)*((1/x)-1))+((m^2)*(L1)*r*(1-x))+(L2*(L2+1)*L1)
6
7 myeqn=(f1(x)*diff(u(x),x,x)+f2(x)*diff(u(x),x)+f3(x)*u(x))
8 myresult=find_special_ode(diffeqn=myeqn,y=u,z=x)

```

and we get

$$\begin{aligned}
& C_2 x^{iL_1\omega} x^{-2iL_1\omega} \text{HC}(0, -2iL_1\omega, 0, L_1 m^2 r, -L_1 m^2 r - L_1 L_2^2 - L_1 L_2, x) \\
& + C_1 x^{iL_1\omega} \text{HC}(0, 2iL_1\omega, 0, L_1 m^2 r, -L_1 m^2 r - L_1 L_2^2 - L_1 L_2, x)
\end{aligned} \tag{70}$$

which agrees with the original article.

7.5 Biconfluent Heun equation: Equation (10) of [45]

The equation (10) of Vitória et. al [45] can be solved in terms of biconfluent Heun functions. We need the substitution $u(x) \rightarrow x^\gamma e^{-\frac{x^2}{2}} u(x)$ and the transformation $x \rightarrow -x$. Using the code

```

1 # The equation (10) of Vitoria et. al
2 var('gamma,beta,m,omega,delta')
3 f1(x)=1
4 f2(x)=1/x
5 f3(x)=(beta/(m*omega))+(delta/x)-((gamma^2)/(x^2))-x^2
6 myeqn=f1(x)*diff(u(x),x,x)+f2(x)*diff(u(x),x)+f3(x)*u(x)
7 myeqn=myeqn.substitute_function(u(x),exp(-(x^2)/2)*(x^gamma)*u(x))
8 myeqn=(myeqn/(exp(-(x^2)/2)*(x^gamma))).full_simplify().expand().collect(u).collect(diff
9 (u,x))
10 show(myeqn)
11 myeqn=ode_change_of_variable(diffeqn=myeqn,y=u,z=x,transformation=-x)
12 myresult=find_special_ode(diffeqn=myeqn,y=u,z=x)

```

we obtain the equation,

$$\frac{\partial}{\partial x} u(x) - \frac{(m\omega x^4 + \gamma^2 m\omega + \delta m\omega x - \beta x^2)u(x)}{m\omega x^2} + \frac{\partial^2}{(\partial x)^2} u(x) \tag{71}$$

and the solution,

$$C_1 \text{HB}\left(\delta, -2\gamma + \frac{\beta}{m\omega} - 2, 2\gamma + 1, 0, -2, x\right) e^{(-\frac{1}{2}x^2)} + \frac{C_2 \text{HB}\left(\delta, 2\gamma + \frac{\beta}{m\omega} - 2, -2\gamma + 1, 0, -2, x\right) e^{(-\frac{1}{2}x^2)}}{x^{2\gamma}} \tag{72}$$

as given in the original article after employing the correspondence given in equations (104)-(108).

7.6 Double confluent Heun equation: Equation (11) of [46]

As explained in the appendix, the singularity structure of the Maple-based approach of the double confluent Heun equation is different from the Mathematica version. The symODE2 package uses the Mathematica form as it involves more parameters. However, this choice makes it difficult to verify our analysis with the examples in literature.

The equation (11) of Vieira [46],

$$\left(A_1 + \frac{A_2}{x^2} + \frac{A_3}{x^4}\right)u(x) + \frac{\partial}{\partial x}u(x) + \frac{\partial^2}{(\partial x)^2}u(x) \quad (73)$$

has the singularity structure $([], [], [0, +\infty], [1, 1])$. The equation can be solved with the code

```

1 # The equation (11) of Vieira
2 var('A1, A2, A3'); f1(x)=1; f2(x)=1/x; f3(x)=A1+(A2/x^2)+(A3/x^4)
3
4 myeqn=f1(x)*diff(u(x), x, x)+f2(x)*diff(u(x), x)+f3(x)*u(x)
5 myresult=find_special_ode(diffeqn=myeqn, y=u, z=x)
6 show(myresult)

```

as

$$C_2\text{HD} \left(-A_2 - 2\sqrt{-A_1}\sqrt{-A_3} - \frac{1}{4}, -2\sqrt{-A_1}, -2\sqrt{-A_3}, 2, -2\sqrt{-A_1}, x \right) e^{\left(\frac{1}{3}\sqrt{-A_1}x^3 + \frac{1}{4}x^2 - 2\sqrt{-A_1}x + \sqrt{-A_3}x + \frac{2\sqrt{-A_3}}{x}\right)}$$

$$+ C_1\text{HD} \left(-A_2 - 2\sqrt{-A_1}\sqrt{-A_3} - \frac{1}{4}, 2\sqrt{-A_1}, 2\sqrt{-A_3}, 2, 2\sqrt{-A_1}, x \right) e^{\left(\frac{1}{3}\sqrt{-A_1}x^3 + \frac{1}{4}x^2 + \sqrt{-A_3}x\right)} \quad (74)$$

If we apply the transformation (12) $(x \rightarrow \frac{x^2+1}{x^2-1})$ as given in the article, we find the form of the equation (13) in the article that can be related with the Maple form, namely,

$$\frac{(A_1 + A_2 + A_3)x^2 + 2(A_1 - A_3)x + A_1 - A_2 + A_3}{x^6 - 3x^4 + 3x^2 - 1}u(x) + \frac{2(x^5 - 2x^3 + x)}{(x+1)^3(x-1)^3} \frac{\partial}{\partial x}u(x) + \frac{\partial^2}{(\partial x)^2}u(x) \quad (75)$$

The singularity structure of this equation is $([], [], [-1, 1], [1, 1])$ as used by Maple.

7.7 Triconfluent Heun equation: Equation (5) of [47]

The equation (5) of Dong et. al [47],

$$-\frac{\sqrt{2}(2bx^2 + a) \frac{\partial}{\partial x}u(x)}{\sqrt{b}} + \frac{(a^2 + 4b\epsilon - 4\sqrt{2}\sqrt{b^3x})u(x)}{2b} + \frac{\partial^2}{(\partial x)^2}u(x) \quad (76)$$

needs the change of coordinates $z = \sqrt[6]{\frac{8b}{9}}x$ in order to bring it in a specific form of the triconfluent Heun equation. Let us apply this transformation and solve the resulting equation with the code

```

1 # The equation (5) of Dong et. al
2 var('a, b, epsilon');
3 f1(x)=1
4 f2(x)=-sqrt(2)*(a+2*b*x^2)/sqrt(b)
5 f3(x)=((a^2)-4*sqrt(2*b^3)*x+4*b*epsilon)/(2*b)
6
7 myeqn=(f1(x)*diff(u(x), x, x)+f2(x)*diff(u(x), x)+f3(x)*u(x))
8 show(myeqn)
9 show(find_singularities(diffeqn=myeqn, y=u, z=x))
10 myeqn2=ode_change_of_variable(diffeqn=myeqn, y=u, z=x, transformation=x*((8*b)/9)^-(1/6))
11 myresult=find_special_ode(diffeqn=myeqn2, y=u, z=x)

```

which yields

$$C_1\text{HT} \left(-\frac{3^{\frac{2}{3}}a^2}{4b^{\frac{4}{3}}} - \frac{3^{\frac{2}{3}}\epsilon}{b^{\frac{1}{3}}}, 3, \frac{3^{\frac{1}{3}}a}{b^{\frac{2}{3}}}, 0, 3, x \right) e^{\left(x^3 + \frac{3^{\frac{1}{3}}ax}{b^{\frac{2}{3}}}\right)} + C_2\text{HT} \left(-\frac{3^{\frac{2}{3}}a^2}{4b^{\frac{4}{3}}} - \frac{3^{\frac{2}{3}}\epsilon}{b^{\frac{1}{3}}}, -3, -\frac{3^{\frac{1}{3}}a}{b^{\frac{2}{3}}}, 0, -3, x \right) \quad (77)$$

that agrees with the original paper using the correspondence of the parameters given in equations (112-116).

7.8 Further applications

The analysis of other ODEs will be in the same manner, therefore we will just give a list of the other cases that our code succeeded. The corresponding analyses can be found in the sample SageMath session given in [34].

- General Heun: Lamé equation as given in [48].
- Biconfluent Heun: The quantum-mechanical doubly anharmonic oscillator potential as given in [49].
- Triconfluent Heun: The classical anharmonic oscillator equation as given in [50]. It should be noted that our code finds the parameter ϵ using a quadratic equation favoring the positive root. The Mathematica solution has the negative root.
- Confluent Heun: Equation (57) of [51].
- Hypergeometric: Legendre equation as given in [39].
- Confluent hypergeometric: Hermite equation as given in [5]. The equations (7.2) and (7.3) of this paper can be obtained from the solution of the Hermite equation given in the same paper. The identity (13.8(iii)) in [39] should be used in the calculation.

8 Conclusion

We proposed an open-source package under SageMath for symbolic analysis of second-order ordinary differential equations with polynomial coefficients. Our approach was based on the singularity structure, namely, the locations of the singularities, corresponding characteristic exponents, and the ranks of the irregular singular points of the equation.

The singularity structure, indices, and recurrence relations associated with the regular singular points, and symbolic solutions of the hypergeometric equation, Heun equation, and their confluent forms could be found using the package.

Using particular substitutions and transformations, the singularity structure of the input equation was brought in a standard form that could be recognized by the routines. After being reached the standard singularity structure of an equation, the parameters were obtained either using the characteristic exponents or by matching the final form of the input equation with the normal form of the standard equation.

As they were defined in SageMath, the results of the hypergeometric and confluent hypergeometric equations were numerically usable, unlike the Heun-type functions.

We presented that our code worked properly with a limited number of tests. However, we emphasized that the results should be taken with great care as such routines should be tested heavily before using them for new problems. We also noted that some transformations, substitutions, or identities might be needed in order to reach the results of the literature.

Acknowledgement

The author would like to thank Prof. Mahmut Hortaçsu for stimulating discussions. The author is also indebted to Maplesoft and Wolfram Research Inc. for providing excellent online resources for the Heun community.

A Appendix: Hypergeometric and Heun type equations in DLMF, Maple, Mathematica, and symODE2

The standard forms of the equations may be defined differently in the literature and in the computer algebra systems. Here, we consider DLMF, a well-known library of mathematical functions [39], and two computer algebra systems, Maple and Mathematica which can work with hypergeometric and Heun type functions. We give lists of correspondence of the parameters in these programs. We also note the standard forms of the equations used in the symODE2 code.

A.1 Hypergeometric equation

In DLMF [39],

$$x(1-x)\frac{d^2y}{dx^2} + [c - (a+b+1)y]\frac{dy}{dx} - aby = 0. \quad (78)$$

In Maple [52],

$$x(x-1)\frac{d^2y}{dx^2} + [(a+b+1)x - c]\frac{dy}{dx} + aby = 0, \quad (79)$$

with solution `hypergeom([a,b],[c],x)`.

In Mathematica [53],

$$-\left(x(x-1)\frac{d^2y}{dx^2} + [(a+b+1)x - c]\frac{dy}{dx} + aby\right) = 0, \quad (80)$$

with solution `Hypergeometric2F1[a,b,c,z]`.

All equations coincide and they have three regular singularities located at $\{0, 1, \infty\}$. symODE2 uses this form of the equation as well.

A.2 Confluent hypergeometric equation

In DLMF [39],

$$x\frac{d^2y}{dx^2} + (b-x)\frac{dy}{dx} - ay = 0. \quad (81)$$

In Maple [52],

$$x\frac{d^2y}{dx^2} + (c-x)\frac{dy}{dx} - ay = 0, \quad (82)$$

with solution `hypergeom([a],[c],x)`.

In Mathematica [54],

$$x\frac{d^2y}{dx^2} + (b-x)\frac{dy}{dx} - ay = 0, \quad (83)$$

with solution `HypergeometricU[a,b,x]`.

All have one regular singularity located at $\{0\}$ and one irregular singularity of rank-1 at $\{\infty\}$. The equations coincide and symODE2 also uses this form of the equation.

A.3 (General) Heun equation

In DLMF [39],

$$\frac{d^2y}{dx^2} + \left(\frac{\gamma}{x} + \frac{\delta}{x-1} + \frac{\epsilon}{x-a}\right)\frac{dy}{dx} + \frac{\alpha\beta x - q}{x(x-1)(x-a)}y = 0, \quad (84)$$

and in Maple [55],

$$\frac{d^2y}{dx^2} + \left(\frac{\gamma}{x} + \frac{\delta}{x-1} + \frac{\epsilon}{x-a} \right) \frac{dy}{dx} + \frac{\alpha\beta x - q}{x(x-1)(x-a)} y = 0, \quad (85)$$

where $\epsilon = \alpha + \beta + 1 - \gamma - \delta$ with solution $\text{HeunG}(a, q, \alpha, \beta, \gamma, \delta, x)$.

In Mathematica [56],

$$\frac{d^2y}{dx^2} + \left(\frac{\gamma}{x} + \frac{\delta}{x-1} + \frac{\alpha + \beta + 1 - \gamma - \delta}{x-a} \right) \frac{dy}{dx} + \frac{\alpha\beta x - q}{x(x-1)(x-a)} y = 0, \quad (86)$$

with solution $\text{HeunG}(a, q, \alpha, \beta, \gamma, \delta, x)$.

All have four regular singularities located at $\{0, 1, a, \infty\}$. symODE2 uses the same structure of the equation.

A.4 (Singly) Confluent Heun equation

In DLMF [39],

$$\frac{d^2y}{dx^2} + \left(\frac{\gamma}{x} + \frac{\delta}{x-1} + \epsilon \right) \frac{dy}{dx} + \frac{\alpha x - q}{x(x-1)} y = 0. \quad (87)$$

In Maple [55],

$$\frac{d^2y}{dx^2} - \frac{-x^2\alpha + (-\beta + \alpha - \gamma - 2)x + \beta + 1}{x(x-1)} \frac{dy}{dx} - \frac{[(-\beta - \gamma - 2)\alpha - 2\delta]x + (\beta + 1)\alpha + (-\gamma - 1)\beta - 2\eta - \gamma}{2x(x-1)} y = 0, \quad (88)$$

with solution $\text{HeunC}(\alpha, \beta, \gamma, \delta, \eta, x)$. This form can be transformed into

$$\frac{d^2y}{dx^2} + \left(\frac{\beta + 1}{x} + \frac{\gamma + 1}{x-1} + \alpha \right) \frac{dy}{dx} + \left(\frac{\mu}{x} + \frac{\nu}{x-1} \right) y = 0, \quad (89)$$

where

$$\delta = \mu + \nu - \alpha \frac{\beta + \gamma + 2}{2}, \quad (90)$$

$$\eta = \frac{(\alpha - \gamma)(\beta + 1) - \beta}{2} - \mu. \quad (91)$$

In Mathematica [56],

$$\frac{d^2y}{dx^2} + \left(\frac{\gamma}{x} + \frac{\delta}{x-1} + \epsilon \right) \frac{dy}{dx} + \frac{\alpha x - q}{x(x-1)} y = 0, \quad (92)$$

with solution $\text{HeunC}(q, \alpha, \gamma, \delta, \epsilon, x)$.

We can define the correspondence,

$$\gamma_{Mat} = \beta_{Map} + 1, \quad (93)$$

$$\delta_{Mat} = \gamma_{Map} + 1, \quad (94)$$

$$\epsilon_{Mat} = \alpha_{Map}, \quad (95)$$

$$\alpha_{Mat} = \mu_{Map} + \nu_{Map}, \quad (96)$$

$$q_{Mat} = \mu_{Map}, \quad (97)$$

where “*Mat*” and “*Map*” stand for Mathematica and Maple, respectively.

All have two regular singularities located at $\{0, 1\}$ and an irregular singularity of rank-1 at $\{\infty\}$. symODE2 uses the Maple form.

A.5 Double confluent Heun equation

In DLMF [39],

$$\frac{d^2y}{dx^2} + \left(\frac{\delta}{x^2} + \frac{\gamma}{x} + 1 \right) \frac{dy}{dx} + \frac{\alpha x - q}{x^2} y = 0. \quad (98)$$

This equation has two irregular singular points of rank-1 located at $\{0, \infty\}$.

In Maple [55],

$$\frac{d^2y}{dx^2} - \frac{\alpha x^4 - 2x^5 + 4x^3 - \alpha - 2x}{(x-1)^3(x+1)^3} \frac{dy}{dx} - \frac{-x^2\beta + (-\gamma - 2\alpha)x - \delta}{(x-1)^3(x+1)^3} y = 0, \quad (99)$$

with solution `HeunD`($\alpha, \beta, \gamma, \delta, x$). This equation has two irregular singular points of rank-1 located at $\{-1, 1\}$.

In Mathematica [56],

$$\frac{d^2y}{dx^2} + \left(\frac{\gamma}{x^2} + \frac{\delta}{x} + \epsilon \right) \frac{dy}{dx} + \frac{\alpha x - q}{x^2} y = 0, \quad (100)$$

with solution `HeunD`($q, \alpha, \gamma, \delta, \epsilon, x$). This equation has two irregular singular points of rank-1 located at $\{0, \infty\}$. `symODE2` uses the Mathematica form.

We will not write a correspondence between the Maple and Mathematica parameters as the singularity structures of the equations are not the same.

A.6 Biconfluent Heun equation

In DLMF [39],

$$\frac{d^2y}{dx^2} - \left(\frac{\gamma}{x} + \delta + x \right) \frac{dy}{dx} + \frac{\alpha x - q}{z} y = 0. \quad (101)$$

In Maple [55],

$$\frac{d^2y}{dx^2} - \frac{\beta x + 2x^2 - \alpha - 1}{x} \frac{dy}{dx} - \frac{(2\alpha - 2\gamma + 4)x + \beta\alpha + \beta + \delta}{2x} y = 0, \quad (102)$$

with solution `HeunB`($\alpha, \beta, \gamma, \delta, x$).

In Mathematica [56],

$$\frac{d^2y}{dx^2} + \left(\frac{\gamma}{x} + \delta + \epsilon x \right) \frac{dy}{dx} + \frac{\alpha x - q}{x} y = 0, \quad (103)$$

with solution `HeunB`($q, \alpha, \gamma, \delta, \epsilon, x$).

All have one regular singularity located at $\{0\}$ and one irregular singularity of rank-2 at $\{\infty\}$. `symODE2` uses the Mathematica form.

We can define the correspondence,

$$\gamma_{Mat} = \alpha_{Map} + 1, \quad (104)$$

$$\delta_{Mat} = -\beta_{Map}, \quad (105)$$

$$\epsilon_{Mat} = -2, \quad (106)$$

$$\alpha_{Mat} = \gamma_{Map} - \alpha_{Map} - 2, \quad (107)$$

$$q_{Mat} = \frac{\beta_{Map}(\alpha_{Map} + 1) + \delta_{Map}}{2}, \quad (108)$$

where “*Mat*” and “*Map*” stand for Mathematica and Maple, respectively.

A.7 Triconfluent Heun equation

In DLMF [39],

$$\frac{d^2y}{dx^2} + (\gamma + x)x \frac{dy}{dx} + (\alpha x - q)y = 0. \quad (109)$$

In Maple [55],

$$\frac{d^2y}{dx^2} - (3x^2 + \gamma) \frac{dy}{dx} + [(\beta - 3)x + \alpha]y = 0, \quad (110)$$

with solution `HeunT`(α, β, γ, x).

In Mathematica [56],

$$\frac{d^2y}{dx^2} + (\gamma + \delta x + \epsilon x^2) \frac{dy}{dx} + (\alpha x - q)y = 0, \quad (111)$$

with solution `HeunT`($q, \alpha, \gamma, \delta, \epsilon, x$).

All have one irregular singularity of rank-3 at $\{\infty\}$. `symODE2` uses the Mathematica form.

We can define the correspondence,

$$\epsilon_{Mat} = -3, \quad (112)$$

$$\delta_{Mat} = 0, \quad (113)$$

$$\gamma_{Mat} = -\gamma_{Map}, \quad (114)$$

$$\alpha_{Mat} = \beta_{Map} - 3, \quad (115)$$

$$q_{Mat} = -\alpha_{Map}, \quad (116)$$

where “*Mat*” and “*Map*” stand for Mathematica and Maple, respectively.

References

- [1] M. Hortacsu, “Explicit examples on conformal invariance,” *Int. J. Theor. Phys.* **42**, 49 (2003). [arXiv:hep-th/0106080 [hep-th]].
- [2] T. Birkandan and M. Cvetič, “Conformal Invariance and Near-extreme Rotating AdS Black Holes,” *Phys. Rev. D* **84**, 044018 (2011). [arXiv:1106.4329 [hep-th]].
- [3] T. Birkandan and M. Cvetič, “Addendum to: Conformal Invariance and Near-extreme Rotating AdS Black Holes,” *Phys. Rev. D* **90**, no.6, 067504 (2014). [arXiv:1406.5208 [hep-th]].
- [4] T. Birkandan and M. Cvetič, “An analysis of the wave equation for the $U(1)^2$ gauged supergravity black hole,” *Class. Quant. Grav.* **32**, no.8, 085007 (2015). [arXiv:1501.03144 [hep-th]].
- [5] J. Dereziński, “Hypergeometric type functions and their symmetries”, *Ann. Henri Poincare* **15**, 1569 (2014). [arXiv:1305.3113 [math.CA]].
- [6] A. Ronveaux (ed.), *Heun’s Differential Equations*, Oxford University Press, New York (1995).
- [7] S.Yu. Slavyanov and W. Lay, *Special Functions, A Unified Theory Based on Singularities*, Oxford University Press, New York (2000).
- [8] Maple 2020, Maplesoft, a division of Waterloo Maple Inc., Waterloo, Ontario (2020).
- [9] M. Hortacsu, “Heun functions and some of their applications in physics”, *Advances in High Energy Physics* **2018**, 8621573 (2018). [arXiv:1101.0471 [math-ph]].

- [10] T. Birkandan and M. Hortacsu, “Dirac Equation in the Background of the Nutku Helicoid Metric,” *J. Math. Phys.* **48**, 092301 (2007). [arXiv:0706.2543 [gr-qc]].
- [11] T. Birkandan and M. Hortacsu, “Examples of Heun and Mathieu functions as solutions of wave equations in curved spaces,” *J. Phys. A* **40**, 1105-1116 (2007). [arXiv:gr-qc/0607108 [gr-qc]].
- [12] P.P. Fiziev, “Exact solutions of Regge-Wheeler equation and quasi-normal modes of compact objects,” *Class. Quant. Grav.* **23**, 2447-2468 (2006). [arXiv:gr-qc/0509123 [gr-qc]].
- [13] Mathematica Version 12.1, Wolfram Research, Inc., Champaign, IL (2020).
- [14] T. Ishkhanyan, “From Sine to Heun: 5 New Functions for Mathematics and Physics in the Wolfram Language”, Accessed 13 September 2020, <https://blog.wolfram.com/2020/05/06/from-sine-to-heun-5-new-functions-for-mathematics-and-physics-in-the-wolfram-language/>
- [15] SageMath, the Sage Mathematics Software System (Version 9.1), The Sage Developers, <https://www.sagemath.org> (2020).
- [16] Maxima, a Computer Algebra System, Version 5.44.0, <http://maxima.sourceforge.net/> (2020).
- [17] P. Virtanen, R. Gommers, T. E. Oliphant, M. Haberland, T. Reddy, D. Cournapeau, E. Burovski, P. Peterson, W. Weckesser, J. Bright, S. J. van der Walt, M. Brett, J. Wilson, K. J. Millman, N. Mayorov, A. R. J. Nelson, E. Jones, R. Kern, E. Larson, C.J. Carey, Í. Polat, Y. Feng, E. W. Moore, J. VanderPlas, D. Laxalde, J. Perktold, R. Cimrman, I. Henriksen, E.A. Quintero, C. R. Harris, A. M. Archibald, A. H. Ribeiro, F. Pedregosa, P. van Mulbregt, and SciPy 1.0 Contributors, “SciPy 1.0: Fundamental Algorithms for Scientific Computing in Python”, *Nature Methods* **17**(3), 261-272 (2020).
- [18] REDUCE, a portable general-purpose computer algebra system, <https://reduce-algebra.sourceforge.io/> (2020).
- [19] GNU Octave Version 5.2.0, Scientific Programming Language, <https://www.gnu.org/software/octave/> (2020).
- [20] O.V. Motygin, “On numerical evaluation of the Heun functions”, *Proceedings of the 2015 Days on Diffraction (DD)*, IEEE (2015). [arXiv:1506.03848 [math.NA]].
- [21] O.V. Motygin, “On evaluation of the confluent Heun functions”, *Proceedings of the 2018 Days on Diffraction (DD)*, IEEE (2018). [arXiv:1804.01007 [math.NA]].
- [22] M.A.H. MacCallum, “Computer algebra in gravity research,” *Living Rev. Rel.* **21**, no.1, 6 (2018).
- [23] J. Grabmeier, E. Kaltrofen, U. Weispfennig (eds.), *Computer Algebra Handbook*, Springer, Berlin (2003).
- [24] J.J. Kovacic, “An algorithm for solving second order linear homogeneous differential equations”, *J. Symbolic Computation* **2**, 3, (1986).
- [25] A. Duval and M. Loday-Richaud, “Kovacic’s algorithm and its application to some families of special functions”, *AAECC* **3**, 211 (1992).
- [26] M. Bronstein and S. Lafaille, “Solutions of linear ordinary differential equations in terms of special functions”, *Proceedings of the 2002 International Symposium on Symbolic and algebraic computation*, 23 (2002).
- [27] L. Chan and E.S. Cheb-Terrab, “Non-liouvillian solutions for second order Linear ODEs”, *Proceedings of the 2004 International Symposium on Symbolic and Algebraic Computation*, 80 (2004). [arXiv:math-ph/0402063].
- [28] R. Debeerst, M. van Hoeij, and W. Koepf, “Solving differential equations in terms of Bessel functions”, *Proceedings of the twenty-first international symposium on Symbolic and algebraic computation*, 39 (2008).
- [29] V. Kunwar and M. van Hoeij, “Second order differential equations with hypergeometric solutions of degree three”, *Proceedings of the twenty-first international symposium on Symbolic and algebraic computation*, 235 (2008).

- [30] E. Imamoglu and M. van Hoeij, “Computing hypergeometric solutions of second order linear differential equations using quotients of formal solutions and integral bases”, *Journal of Symbolic Computation* **83**, 254 (2017). [arXiv:1606.01576 [cs.SC]].
- [31] E. Gourgoulhon, M. Bejger, M. Mancini, “Tensor calculus with open-source software: the SageManifolds project”, *J. Phys. Conf. Ser.* **600**, 012002 (2015). [arXiv:1412.4765 [gr-qc]].
- [32] E. Gourgoulhon, M. Mancini, “Symbolic tensor calculus on manifolds: a SageMath implementation”, *Les cours du CIRM* **6**, I (2018). [arXiv:1804.07346 [gr-qc]].
- [33] T. Birkandan, C. Güzelgün, E. Şirin and M. C. Uslu, “Symbolic and numerical analysis in general relativity with open source computer algebra systems,” *Gen. Rel. Grav.* **51**, no.1, 4 (2019). [arXiv:1703.09738 [gr-qc]].
- [34] “The symODE2 package”, Accessed 27 September 2020, <https://github.com/tbirkandan/symODE2>
- [35] F.W.J. Olver, *Asymptotics and Special Functions*, Academic Press, New York (1974).
- [36] R.S. Maier, “On reducing the Heun equation to the hypergeometric equation”, *J. Differential Equations* **213**, 171 (2005). [arXiv:math/0203264 [math.CA]].
- [37] G. Allen, “Some efficient methods for obtaining infinite series solutions of nth-order linear ordinary differential equations”, NASA Technical Report (NASA TR-R-390) (1972).
- [38] R.A. Mezei, *An Introduction to SAGE Programming*, Wiley, Hoboken (2016).
- [39] F.W.J. Olver, A.B. Olde Daalhuis, D.W. Lozier, B.I. Schneider, R.F. Boisvert, C.W. Clark, B.R. Miller, B.V. Saunders, H.S. Cohl, and M.A. McClain (eds.), *NIST Digital Library of Mathematical Functions*, <http://dlmf.nist.gov/>, Release 1.0.27 of 2020-06-15.
- [40] G. Kristensson, *Second Order Differential Equations: Special Functions and Their Classification*, Springer, New York (2010).
- [41] R.S. Maier, “The 192 Solutions of the Heun Equation”, *Math. Comp.* **76**, 811 (2007). [arXiv:math/0408317 [math.CA]].
- [42] R.N. Nasheeha, S. Thirukkanesh and F.C. Ragel, “Anisotropic generalization of isotropic models via hypergeometric equation,” *Eur. Phys. J. C* **80**, no.1, 6 (2020).
- [43] D. Petroff, “Slowly rotating homogeneous stars and the Heun equation,” *Class. Quant. Grav.* **24**, 1055-1068 (2007). [arXiv:gr-qc/0701081 [gr-qc]].
- [44] İ. Sakallı, K. Jusufi and A. Övgün, “Analytical Solutions in a Cosmic String Born-Infeld-dilaton Black Hole Geometry: Quasinormal Modes and Quantization,” *Gen. Rel. Grav.* **50**, no.10, 125 (2018). [arXiv:1803.10583 [gr-qc]].
- [45] R.L.L. Vitória, C. Furtado and K. Bakke, “On a relativistic particle and a relativistic position-dependent mass particle subject to the Klein–Gordon oscillator and the Coulomb potential,” *Annals Phys.* **370**, 128-136 (2016). [arXiv:1511.05072 [quant-ph]].
- [46] H.S. Vieira, “Resonant frequencies of the hydrodynamic vortex,” *Int. J. Mod. Phys. D* **26**, no.04, 1750035 (2016). [arXiv:1510.08298 [gr-qc]].
- [47] Q. Dong, G.H. Sun, M.A. Aoki, C.Y. Chen and S.H. Dong, “Exact solutions of a quartic potential,” *Mod. Phys. Lett. A* **34**, no.26, 1950208 (2019).
- [48] ‘HeunG’, Accessed 27 September 2020, <https://reference.wolfram.com/language/ref/HeunG.html>.
- [49] ‘HeunB’, Accessed 27 September 2020, <https://reference.wolfram.com/language/ref/HeunB.html>.
- [50] ‘HeunT’, Accessed 27 September 2020, <https://reference.wolfram.com/language/ref/HeunT.html>.

- [51] M. S. Cunha and H. R. Christiansen, “Confluent Heun functions in gauge theories on thick braneworlds,” *Phys. Rev. D* **84**, 085002 (2011). [arXiv:1109.3486 [hep-th]].
- [52] “Solving Some Second Order Linear ODEs that Admit Hypergeometric 2F1, 1F1, and 0F1 Function Solutions”, Accessed 12 September 2020, <https://www.maplesoft.com/support/help/Maple/view.aspx?path=dsolve/hyper3>.
- [53] “Hypergeometric2F1”, Accessed 12 September 2020, <https://functions.wolfram.com/HypergeometricFunctions/Hypergeometric2F1/13/01/01/01/>.
- [54] “HypergeometricU”, Accessed 12 September 2020, <https://reference.wolfram.com/language/ref/HypergeometricU.html>.
- [55] “The five Second Order Linear Heun equations and the corresponding Heun function solutions”, Accessed 12 September 2020, <https://www.maplesoft.com/support/help/Maple/view.aspx?path=Heun>.
- [56] “Heun and Related Functions”, Accessed 12 September 2020, <https://reference.wolfram.com/language/guide/HeunAndRelatedFunctions.html>